

GLUING DETERMINANTAL CREMONA MAPS

RÉMI BIGNALET-CAZALET

ABSTRACT. We study determinantal Cremona maps, i.e. birational maps whose base ideal is the maximal minors ideal of a given matrix Φ , via the resolution of the polynomials systems defined by Φ . Using convex geometry, this approach leads in particular to describe the projective degrees of some almost linear determinantal maps.

INTRODUCTION

In this note, we consider rational maps $f = (f_0 : \dots : f_n) : \mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^n$ defined by $n + 1$ homogeneous polynomials $f_0, \dots, f_n \in \mathbb{R} = k[x_0, \dots, x_n]$ without common factor, that are determinantal i.e. that the *base ideal* $I_f = (f_0, \dots, f_n) \subset \mathbb{R}$ of f is the n -minors ideal of its minimal presenting matrix Φ_f .

Our guiding remark is that the so-called *standard Cremona map* (see [GSP06] for more references about the standard Cremona maps)

$$\tau_n = (x_1 \cdots x_n : x_0 x_2 \cdots x_n : \dots : x_0 \cdots x_{n-1}) : \mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^n$$

can be interpreted as the gluing of n standard Cremona maps $\tau_1^{(1)} = (x_1 : x_0) : \mathbb{P}_k^1 \dashrightarrow \mathbb{P}_k^1$, $\tau_1^{(2)} = (x_2 : x_1) : \mathbb{P}_k^1 \dashrightarrow \mathbb{P}_k^1$, \dots , $\tau_1^{(n)} = (x_n : x_{n-1}) : \mathbb{P}_k^1 \dashrightarrow \mathbb{P}_k^1$. Indeed, I_{τ_n} is the n -minors ideal of the $(n + 1) \times n$ matrix

$$\Phi_{\tau_n} = \begin{pmatrix} x_0 & 0 & \dots & \dots & 0 \\ -x_1 & x_1 & \ddots & & \vdots \\ 0 & -x_2 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ \vdots & & \ddots & \ddots & x_{n-1} \\ 0 & \dots & \dots & 0 & -x_n \end{pmatrix}$$

which is the concatenation of the presenting matrices $\Phi_{\tau_1^{(1)}} = \begin{pmatrix} x_0 \\ -x_1 \end{pmatrix}$, \dots , $\Phi_{\tau_1^{(n)}} = \begin{pmatrix} x_{n-1} \\ -x_n \end{pmatrix}$ respectively of $I_{\tau_1^{(1)}}$, \dots , $I_{\tau_1^{(n)}}$ completed with zeros entries.

This work takes its roots in trying to generalize this previous construction and, given two integers $m, m' \geq 1$, to understand the projective degrees of such an outputted glued determinantal map $[g|g'] : \mathbb{P}_k^{m+m'} \dashrightarrow \mathbb{P}_k^{m+m'}$ via the properties

Date: January 1, 2022.

2010 Mathematics Subject Classification. 13D02, 14E05,

Key words and phrases. determinantal Cremona maps, multidegree/projective degrees of a rational map, mixed volumes of polytopes, Bernstein's theorem on sparse polynomial systems, glued determinantal Cremona map.

The author was funded by the European Union's Horizon 2020 research and innovation program as a Marie Skłodowska-Curie fellow of the Istituto Nazionale di Alta Matematica "Francesco Severi" grant No. 713485.

of the inputted determinantal maps $g : \mathbb{P}_k^m \dashrightarrow \mathbb{P}_k^m$ and $g' : \mathbb{P}_k^{m'} \dashrightarrow \mathbb{P}_k^{m'}$. Recall that given any map $f = (f_0 : \dots : f_n) : \mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^n$ and assuming that the base field k is algebraically closed, the projective degrees $d_0(f), d_1(f), \dots, d_n(f)$ of f are the quantities such that for any $k \in \{0, \dots, n\}$, $d_k(f)$ is the cardinal $\#(H_x^k \cap f^{-1}(H_y^{n-k}))$ of the intersection between a general linear space H_x^k of codimension k in the source space of f with the preimage of a general linear space H_y^{n-k} of codimension $n - k$ in the target space of f , see [Har92, Ex. 19.4] or below for more developments about the projective degrees of a map. Given this latter definition, let us state a first result describing, in some situations, the projective degree of $[g|g']$.

Proposition 1. *Let $m, m' \geq 1$, $R_m = k[x_0, \dots, x_m]$, $R_{m'} = k[x_m, \dots, x_{m+m']$ and $R = k[x_0, \dots, x_{m+m']$ and let $g : \mathbb{P}_k^m \dashrightarrow \mathbb{P}_k^m$ (resp. $g' : \mathbb{P}_k^{m'} \dashrightarrow \mathbb{P}_k^{m'}$) be a determinantal map such that I_g (resp. $I_{g'}$) is the m -minors ideal of a matrix $\Phi_g \in R_m^{(m+1) \times m}$ (resp. the m' -minors ideal of $\Phi_{g'} \in R_{m'}^{(m'+1) \times m'}$). Put*

$$\Phi_{[g|g']} = \left\{ \left(\begin{array}{c|c} \overbrace{\Phi_g}^m & \overbrace{0_{m \times m'}}^{m'} \\ \hline \overbrace{0_{m' \times m}}^{m'} & \overbrace{\Phi_{g'}}^{m'} \end{array} \right) \right\} \begin{array}{l} m+1 \\ \\ \\ \\ m' \end{array} \in R^{(m+m'+1) \times (m+m')}$$

and assume that the ideal of $(m + m')$ -minors of $\Phi_{[g|g']}$ is the base ideal of a map $[g|g'] : \mathbb{P}_k^{m+m'} \dashrightarrow \mathbb{P}_k^{m+m'}$.

Suppose that the tensor product $\mathbb{F}_g \otimes \mathbb{F}_{g'}$ of a free resolution \mathbb{F}_g of the graph of g and a free resolution $\mathbb{F}_{g'}$ of the graph of g' provides a free resolution of the graph of $[g|g']$. Then, given any $k \in \{0, \dots, m + m'\}$

$$(1) \quad d_k([g|g']) = \sum_{p=0}^k d_p(g) d_{k-p}(g')$$

with the convention that $d_p(g) = 0$ (resp. $d_p(g') = 0$) if $p > m$ (resp. $p > m'$).

As we will explain, Proposition 1 applies in particular in the previous example of the standard Cremona map.

Actually, other definitions of a glued determinantal map $[g|g']$ can be given depending on the convention adopted to concatenate the matrices Φ_g and $\Phi_{g'}$. For instance, in the case where each entry of any given column of Φ_g and $\Phi_{g'}$ is a general combination of given polynomials, a pseudo concatenate matrix $\Phi_{[g|g']}$ and an associated glued map $[g|g']$ can still be defined. In this direction:

Proposition 2.14. *Let $m, d \geq 1$ and let $\Phi_{[g|g']} = (\phi_{ij})_{\substack{0 \leq i \leq m+2 \\ 1 \leq j \leq m+2}}$ be such that:*

- for all $l \in \{1, \dots, m\}$, all the entries ϕ_{il} of the l -th column of $\Phi_{[g|g']}$ are general linear combinations of x_0, \dots, x_m ,
- all the entries $\phi_{i, m+1}$ of the $m + 1$ -th column of $\Phi_{[g|g']}$ are general linear combinations of x_{m+1} and x_{m+2} ,

- all the entries $\phi_{i,m+2}$ of the $m + 2$ -th column of $\Phi_{[g|g']}$ are general linear combinations of the generators of the ideal

$$(x_m, x_{m+1}, x_{m+2}) \cdot (x_{m+1}, x_{m+2})^{d-1} = (x_m x_{m+1}^{d-1}, x_m x_{m+2}^{d-1}) + (x_{m+1}, x_{m+2})^d.$$

Then the map $[g|g'] : \mathbb{P}_k^{m+2} \dashrightarrow \mathbb{P}_k^{m+2}$ whose base ideal $I_{[g|g']}$ is the $(m+2)$ -minors ideal of $\Phi_{[g|g']}$ is a determinantal map and moreover:

$$\forall k \in \{0, \dots, m+2\}, d_k([g|g']) = \binom{m}{m-k} + (d+1) \binom{m}{m-k+1} + \binom{m}{m-k+2}$$

with the convention that $\binom{j}{i} = 0$ if $i < 0$ or $i > j$. In particular, the projective degrees' vector $d([g|g']) = (d_0([g|g']), \dots, d_{m+2}([g|g']))$ of $[g|g']$ is palindromic.

Our goal behind this glued construction is to approach combinatorially the projective degrees of the *almost linear determinantal maps*, i.e. the determinantal map f defined by *almost linear matrix* Φ_f for which all but one column are filled with linear polynomials (see [KPU11] for development about maps defined by almost linear matrices), in order to eventually extend in greater dimension previous works such as in [DH17].

The proof of Proposition 2.14 relies on the following observations. When considering a map $f : \mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^n$ of base ideal I_f with presenting matrix Φ_f , the scheme $\mathbb{P}(I_f) = \text{Proj}(\text{Sym}(I_f))$ provides an approximation of the projective degrees of f . This is now a classical approach, see [RS01], [BCJ09], [BCRS20] and [CR21] for developments and applications in broader contexts. If f is in addition *Koszul-determinantal* i.e. $\mathbb{P}(I_f)$ is a complete intersection in its embedding in $\mathbb{P}_k^n \times \mathbb{P}_k^n$ defined by Φ_f (a condition on f stronger than being determinantal and that we will explicit below), one can use the resolution of zero-dimensional complete intersection defined by sparse polynomials to compute, in some situations, the projective degrees of f . Indeed, Bernstein's theorem states that, under conditions, the mixed volume associated to a polynomial system (E) compute the solutions of (E) with non zero coordinates, see [CLO05, 7. Th. 5.4] or Theorem 2.7 below. Proposition 2.14 then follows from convex geometry, and in particular the projection formula in [ST10, Lemma 6] decomposing some mixed volumes as the product of mixed volumes in smaller dimension.

Contents of the paper. In Section 1, we precise the definition of the projective degrees of a map f , emphasizing how they are related to (bi-graded) free resolutions of the ideal of the graph Γ_f of f . It gives the background of our proof of Proposition 1. In Section 2, we precise the approximations of the projective degrees of a Koszul-determinantal map given by the mixed volumes associated to the presentation matrix its the base ideal. We then give another definition of a glued determinantal map $[g|g'] : \mathbb{P}_k^{m+m'} \dashrightarrow \mathbb{P}_k^{m+m'}$ via general properties of two initial determinantal maps $g : \mathbb{P}_k^m \dashrightarrow \mathbb{P}_k^m$ and $g' : \mathbb{P}_k^{m'} \dashrightarrow \mathbb{P}_k^{m'}$. Under assumptions on g and g' , the projective degrees of such a glued map $[g|g']$ are then described by a Kunneth-like formula in the projective degrees of g and g' , see Proposition 2.14 and its proof.

The explicit computations given in this paper were made using basic functions of the software systems POLYMAKE and MACAULAY2 and the CREMONA package [Sta17] associated. The corresponding codes are available on request.

Acknowledgements. I thank Adrien Dubouloz and Daniele Faenzi for their comments about a first version of this manuscript. I also warmly thank Alessandro de Stefani for its help on some arguments of the first section.

Notation 2. We let k be any field in Section 1 and $k = \mathbb{C}$ in Section 2.

When recalling generalities about rational maps, we consider a positive integer $n \in \mathbb{N}^*$, $R = k[x_0, \dots, x_n]$, $\mathbb{P}_k^n = \text{Proj}(R)$ and a rational map $f : \mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^n$.

When considering glued maps, we let $m, m' \geq 1$, $R_m = k[x_0, \dots, x_m]$, $R_{m'} = k[x_m, \dots, x_{m+m']$, $R = k[x_0, \dots, x_{m+m']$ and we always consider that R_m and $R_{m'}$ are embedded in R . Let also $g : \mathbb{P}_k^m \dashrightarrow \mathbb{P}_k^m$ (resp. $g' : \mathbb{P}_k^{m'} \dashrightarrow \mathbb{P}_k^{m'}$), where $\mathbb{P}_k^m = \text{Proj}(R_m)$ (resp. $\mathbb{P}_k^{m'} = \text{Proj}(R_{m'})$). The associated glued map is denoted $[g|g'] : \mathbb{P}_k^{m+m'} \dashrightarrow \mathbb{P}_k^{m+m'}$ where $\mathbb{P}_k^{m+m'} = \text{Proj}(R_{m+m'})$.

1. FREE RESOLUTIONS OF THE GRAPH OF GLUED DETERMINANTAL MAPS

Our motivation in describing a free resolution of the ideal of the graph of a map f (free resolution of the graph for short) lies on the fact it gives an insight on the projective degrees of f and let us briefly outline why.

As explained in [MS05, Chapter 8], the theory of multigraded Hilbert series is the fine context for defining and studying the multidegree of subschemes in multi-projective spaces so we adapt here previous expositions as in [MS05] or the more recent one in [CR21] to the following bi-graded context. Let $n \geq 1$ be an integer, k be any field (if no additional specific assumptions), $R = k[x_0, \dots, x_n]$ and $S = R[y_0, \dots, y_n]$ be the \mathbb{Z}^2 -graded polynomial ring with the standard graduation $\deg(x_i) = (1, 0)$ and $\deg(y_i) = (0, 1)$ for all $i \in \{0, \dots, n\}$. Given a bi-graded algebra B , we let $\text{BiProj}(B)$ be the set of bi-graded prime ideals of B . Thus, a subscheme \mathbb{X} of $\text{BiProj}(S) \simeq \mathbb{P}_k^n \times \mathbb{P}_k^n$ is defined by a bi-graded ideal $\mathcal{A} \subset S$ and one associates to $\mathbb{X} = \text{BiProj}(S/\mathcal{A})$ its bi-variate Hilbert series

$$H_{S/\mathcal{A}}(T_0, T_1) = \sum_{(n_0, n_1) \in \mathbb{Z}^2} (\text{length}(S/\mathcal{A})_{(n_0, n_1)} T_0^{n_0} T_1^{n_1}).$$

Then, writing $H_{S/\mathcal{A}}(T_0, T_1) = \frac{\text{Num}_{S/\mathcal{A}}(T_0, T_1)}{(1-T_0)^c (1-T_1)^c}$ and assuming that $\mathbb{X} = \text{BiProj}(S/\mathcal{A})$ has codimension c , the coefficients of the homogeneous component of $\text{Num}_{S/\mathcal{A}}(1 - T_0, 1 - T_1)$ of total degree c define the multidegree of \mathbb{X} .

Definition 1.1 (multidegree of a subscheme in $\mathbb{P}_k^n \times \mathbb{P}_k^n$). Let \mathcal{A} be a bi-graded ideal of S and assume that $\mathbb{X} = \text{BiProj}(S/\mathcal{A})$ has codimension c in $\text{BiProj}(S) \simeq \mathbb{P}_k^n \times \mathbb{P}_k^n$.

Given $k \in \{0, \dots, c\}$, define $\deg_{\mathbb{P}}^{c-k, k} \mathbb{X}$ as the coefficient of the monomial of bi-degree $(c-k, k)$ of $\text{Num}_{S/\mathcal{A}}(1 - T_0, 1 - T_1)$.

The *multidegree* $\deg_{\mathbb{P}} \mathbb{X}$ of \mathbb{X} is the $c+1$ -uple

$$\deg_{\mathbb{P}} \mathbb{X} = (\deg_{\mathbb{P}}^{c,0} \mathbb{X}, \dots, \deg_{\mathbb{P}}^{0,c} \mathbb{X}).$$

Remark 1.2. As explained for instance in [Har92, Proposition 7.16] or [CR21, Theorem 4.7], the multidegree $\deg_{\mathbb{P}} \mathbb{X} = (\deg_{\mathbb{P}}^{c,0} \mathbb{X}, \dots, \deg_{\mathbb{P}}^{0,c} \mathbb{X})$ of a c -dimensional subscheme $\mathbb{X} \subset \mathbb{P}_k^n \times \mathbb{P}_k^n$ has the following geometric interpretation. Assuming the base field k is algebraically closed and letting $k \in \{0, \dots, c\}$, we have:

$$\deg_{\mathbb{P}}^{c-k, k} \mathbb{X} = \text{length}(H_{\mathbf{x}}^k \cap \mathbb{X} \cap H_{\mathbf{y}}^{c-k})$$

where $H_{\mathbf{x}}^k = \mathbb{V}(l_{1,0}, \dots, l_{k,0})$ is the zero locus of k general linear forms $l_{1,0}, \dots, l_{k,0}$ in the \mathbf{x} -variables (that is $\deg(l_{i,0}) = (1, 0)$ for all $i \in \{0, \dots, k\}$) and $H_{\mathbf{y}}^k =$

$\mathbb{V}(l_{0,1}, \dots, l_{0,c-k})$ is the zero locus of $c-k$ general linear forms $l_{0,1}, \dots, l_{0,c-k}$ in the y -variables.

Definition 1.3 (projective degrees of a map). Let $f : \mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^n$, regular on a dense open subset $U \subset \mathbb{P}_k^n$, and let $\Gamma_f = \overline{\{(x, f(x)), x \in U\}} \subset \mathbb{P}_k^n \times \mathbb{P}_k^n$ be the graph of f . It is an n -fold so, for $k \in \{0, \dots, n\}$, the k -th projective degree of f , written $d_k(f)$, is defined by

$$d_k(f) := \deg_{\mathbb{P}}^{n-k, k} \Gamma_f.$$

In the following, our convention is also that $d_k(f) = 0$ if $k > n$.

By Remark 1.2, note that $d_0(f) = 1$ if and only if f is an isomorphism between two dense open subsets of \mathbb{P}_k^n in which case f is said to be a *Cremona map*.

Following for instance [EH00, III.3.6], a bigraded free resolution of S/I_{Γ_f} provides the numerator $\text{Num}_{S/I_{\Gamma_f}}(T_0, T_1)$ of the Hilbert series of the coordinate ring S/I_{Γ_f} of Γ_f and thus provides the projective degrees of f .

Now let us clarify in which embedding we consider the graphs Γ_g and $\Gamma_{g'}$ of, respectively, a map $g : \mathbb{P}_k^m \dashrightarrow \mathbb{P}_k^m$ and $g' : \mathbb{P}_k^{m'} \dashrightarrow \mathbb{P}_k^{m'}$ see Notation 2. Denote by $S_m = R_m[y_0, \dots, y_m]$, $S_{m'} = R_{m'}[y_m, \dots, y_{m+m'}]$, $S = R[y_0, \dots, y_{m+m'}]$ and $I_{\Gamma_g} \subset S$ (resp. $I_{\Gamma_{g'}} \subset S$) be the ideal of the graph $\Gamma_g \subset \mathbb{P}_k^{m+m'} \times \mathbb{P}_k^{m+m'}$ of g (resp. $\Gamma_{g'} \subset \mathbb{P}_k^{m+m'} \times \mathbb{P}_k^{m+m'}$ of g') where $\Gamma_g \subset \mathbb{P}_k^{m+m'} \times \mathbb{P}_k^{m+m'} = \text{BiProj}(S)$. Geometrically, Γ_g (resp. $\Gamma_{g'}$) is thus the cone of vertex $\Gamma_g \cap \mathbb{P}_k^m \times \mathbb{P}_k^m$ (resp. $\Gamma_{g'} \cap \mathbb{P}_k^{m'} \times \mathbb{P}_k^{m'}$) where $\mathbb{P}_k^m \times \mathbb{P}_k^m = \text{BiProj}(S_m)$ (resp. $\mathbb{P}_k^{m'} \times \mathbb{P}_k^{m'} = \text{BiProj}(S_{m'})$). Proposition 1 follows from the following results about $\Gamma_g \cap \Gamma_{g'}$.

Lemma 1.4. *Assume that the tensor product $\mathbb{F}_g \otimes \mathbb{F}_{g'}$, of a bigraded free resolution \mathbb{F}_g of Γ_g with a bigraded free resolution $\mathbb{F}_{g'}$ of $\Gamma_{g'}$, is a free resolution of $\Gamma_g \cap \Gamma_{g'}$. Then for any $k \in \{0, \dots, m+m'\}$:*

$$d_k([g|g']) = \sum_{p=0}^k d_p(g) d_{k-p}(g').$$

Proof. By assumptions, the numerator $\text{Num}_{S/I_{\Gamma_g \cap \Gamma_{g'}}}(T_0, T_1)$ of the Hilbert series of $S/I_{\Gamma_g \cap \Gamma_{g'}}$ is the product $\text{Num}_g(T_0, T_1) \text{Num}_{g'}(T_0, T_1)$ of the numerators of the Hilbert series of respectively S/I_g and $S/I_{g'}$.

Focusing on the homogeneous component of total degree $m+m'$ of $\text{Num}_{[g|g']}(1-T_0, 1-T_1)$, we thus have:

$$\begin{aligned} & (\text{Num}_{S/I_{\Gamma_g \cap \Gamma_{g'}}}(1-T_0, 1-T_1))_{m+m'} \\ &= (\text{Num}_g(1-T_0, 1-T_1) \text{Num}_{g'}(1-T_0, 1-T_1))_{m+m'} \\ &= (\text{Num}_g(1-T_0, 1-T_1))_m (\text{Num}_{g'}(1-T_0, 1-T_1))_{m'} \end{aligned}$$

the last equality holds because $(\text{Num}_g(1-T_0, 1-T_1))_m$ (resp. $(\text{Num}_{g'}(1-T_0, 1-T_1))_{m'}$) is also the homogeneous component of smallest total degree of $\text{Num}_g(1-T_0, 1-T_1)$ (resp. $\text{Num}_{g'}(1-T_0, 1-T_1)$) as Γ_g , being the cone over the graph of g , is irreducible of codimension m (resp. $\Gamma_{g'}$ is irreducible of codimension m'). \square

Given these preliminaries and before proving Proposition 1, let us briefly underline why, when building a glued map $[g|g']$, we restrict to inputted determinantal maps g and g' .

Definition 1.5 (Hilbert-Burch matrix of a determinantal map). Given that the base ideal $I_f = (f_0 : \dots : f_n) \subset R$ of f verifies $\text{codim}(\mathbb{V}(I_f)) \geq 2$ and that I_f is the n -minors ideal of a $(n+1) \times n$ -matrix, Hilbert-Burch theorem [Eis95, Theorem 20.15] states that I_f is the n -minors ideal of its presentation matrix Φ_f , i.e. the presentation of I_f reads:

$$0 \rightarrow R^n \xrightarrow{\Phi_f} R^{n+1} \xrightarrow{(f_0 \dots f_n)} I_f \rightarrow 0.$$

It motivates why the matrix Φ_f is called the *Hilbert-Burch matrix* of f in the following, see for instance [KPU11], [KPU13] and [KM20] for recent works about maps defined by Hilbert-Burch matrices.

When gluing two determinantal maps g and g' as in Proposition 1, the $(m+m')$ -minors ideal of the concatenated matrix $\Phi_{[g|g']}$ always defines a map $\mathbb{P}_k^{m+m'} \dashrightarrow \mathbb{P}_k^{m+m'}$ and, if $\text{codim} \mathbb{V}(I_{m+m'}(\Phi_{[g|g']})) = 2$, the base ideal $I_{[g|g']}$ of $[g|g']$ is equal to $I_{m+m'}(\Phi_{[g|g']})$. Let us mention here that a glued map $[g|g']$ can also be defined when g and g' are not necessarily determinantal however in this case $[g|g']$ is not necessarily a map from $\mathbb{P}_k^{m+m'}$ to $\mathbb{P}_k^{m+m'}$.

Now recall that, algebraically, the graph Γ_f of a map $f = (f_0 : \dots : f_n) : \mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^n$ is the Proj of the Rees algebra $\mathcal{R}(I_f) = \bigoplus_{k \geq 0} I_f^k t^k$ of the base ideal I_f of f and the embedding of Γ_f in $\mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^n$ is defined by the surjection $S \twoheadrightarrow \mathcal{R}(I_f)$ sending the variables y_i to $f_i t$ ($S = R[y_0, \dots, y_n]$). In practice, the kernel J of the latter surjection may be difficult to compute and an approximation of J is more accessible by considering the symmetric algebra of I_f , this is a classical approach that we now summarize briefly in our context of rational maps, see [Vas05] for an introduction in a broader context and pointers to references about this procedure. The presentation matrix Φ_f of I_f , by definition verifying the following short exact sequence:

$$R^{n'} \xrightarrow{\Phi_f} R^{n+1} \xrightarrow{(f_0 \dots f_n)} I_f \rightarrow 0,$$

defines an embedding of $\mathbb{P}(I_f) = \text{Proj}(\text{Sym}(I_f))$ in $\mathbb{P}_k^n \times \mathbb{P}_k^n$ whose ideal J_1 is generated by the n' entries of the line matrix $(y_0 \dots y_n)\Phi_f$ (here we consider that Φ_f is a matrix both in R and S). Since the natural surjection $\text{Sym}(I_f) \twoheadrightarrow \mathcal{R}(I_f)$ factorizing the maps $I_f^{\otimes k} \twoheadrightarrow I_f^k$ defines an embedding of Γ_f in $\mathbb{P}(I_f)$, one has that $J_1 \subset J$, i.e. J_1 provides some equations of Γ_f . The ideal I_f is said to be of *linear type* when $J_1 = J$.

When f is determinantal of Hilbert-Burch matrix $\Phi_f \in R^{(n+1) \times n}$, the kernel of $\text{Sym}(I_f) \twoheadrightarrow \mathcal{R}(I_f)$ is the R -torsion of $\text{Sym}(I_f)$ [Mic64] and are described by the Fitting ideals of I_f , i.e. by the ideals $I_k(\Phi_f)$ for $k \in \{1, \dots, n-1\}$ of k -minors ideals of the presentation matrix Φ_f of I_f , see [Vas05, Prop. 1.1 and below] and [BCJ09]. Hence, the irreducible components of $\mathbb{P}(I_f)$ are the graph Γ_f of f and eventual additional pieces lying above closed strict subschemes of the source space \mathbb{P}_k^n of f .

With all these facts, let us now show Proposition 1.

Proof of Proposition 1. Assume that $\text{codim} \mathbb{V}(I_{m+m'}(\Phi_{[g|g']})) = 2$ so the base ideal $I_{[g|g']}$ of the glued map $[g|g'] : \mathbb{P}_k^{m+m'} \dashrightarrow \mathbb{P}_k^{m+m'}$ is the $(m+m')$ -minors ideal of the matrix $\Phi_{[g|g']}$. It is clear that $\Gamma_{[g|g']}$ is included in $\Gamma_g \cap \Gamma_{g'}$. Indeed, by definition, $\Gamma_{[g|g']} \subset \mathbb{P}(I_{[g|g']}) = \mathbb{P}(I_g) \cap \mathbb{P}(I_{g'})$ so $\Gamma_{[g|g']} \subset \mathbb{P}(I_g)$ (resp. $\Gamma_{[g|g']} \subset \mathbb{P}(I_{g'})$) and

since $\Gamma_{[g|g']}$ cannot be included in a component of $\mathbb{P}(I_g)$ (resp. $\mathbb{P}(I_{g'})$) lying above a closed strict subscheme of $\mathbb{P}_k^{m+m'}$, it is necessarily included in Γ_g (resp. $\Gamma_{g'}$).

Now, since, by assumption, $\mathbb{F}_g \otimes \mathbb{F}_{g'}$ is a resolution of $\Gamma_{[g|g']}$ and $\Gamma_{[g|g']} \subset \Gamma_g \cap \Gamma_{g'}$, one has $\Gamma_{[g|g']} = \Gamma_g \cap \Gamma_{g'}$.

The conclusion of Proposition 1 follows then from Lemma 1.4. \square

Note that, as set, $\Gamma_{[g|g']}$ and $\Gamma_g \cap \Gamma_{g'}$ are not always equal as illustrated by the following example.

Example 1.6. Let $\Phi_g = \begin{pmatrix} x_2 & 0 \\ x_1 & x_0x_2 \\ 0 & x_1^2 \end{pmatrix}$ and $\Phi_{g'} = \begin{pmatrix} x_3 & 0 \\ x_4 & x_2x_3 \\ 0 & x_4^2 \end{pmatrix}$ and $\Phi_{[g|g']}$ be the concatenate matrix of Φ_g and $\Phi_{g'}$, as in Proposition 1. Then, as it can be checked by direct computation on a computer algebra system such as MACAULAY2, one has that $\text{codim } \mathbb{V}(\Phi_{[g|g']}) \geq 2$ and that $d([g|g']) = (1, 6, 10, 6, 1)$, $d(g) = d(g') = (1, 3, 1)$ and

$$d_2([g|g']) = 10 < 11 = d_0(g)d_2(g') + d_1(g)d_1(g') + d_2(g)d_0(g').$$

In this example $\mathbb{F}_g \otimes \mathbb{F}_{g'}$ provides a bigraded free resolution of $\Gamma_g \cap \Gamma_{g'}$ and $\Gamma_{[g|g']} \subsetneq \Gamma_g \cap \Gamma_{g'}$.

Let us also mention that there exist situations where the ideal of $\Gamma_g \cap \Gamma_{g'}$ decomposes as the intersection of the ideal of $\Gamma_{[g|g']}$ and the ideal of embedded components, so that $\Gamma_{[g|g']}$ and $\Gamma_g \cap \Gamma_{g'}$ are not scheme-theoretically equal but have the same projective degrees.

Even if it might seem a bit artificial (since the results about the standard Cremona maps are well known, in particular its projective degrees, and computable by other means, see [GSP06]), we illustrate an application of Proposition 1 in the example of the standard Cremona maps $\tau_{m+m'} = [\tau_m | \tau_{m'}]$.

Proposition 1.7. *Following Notation 2, let*

$$\begin{aligned} \tau_m &= (x_1 \cdots x_m : \dots : x_0 \cdots x_{m-1}) : \mathbb{P}_k^m \dashrightarrow \mathbb{P}_k^m \\ \tau_{m'} &= (x_{m+1} \cdots x_{m+m'} : \dots : x_m \cdots x_{m+m'-1}) : \mathbb{P}_k^{m'} \dashrightarrow \mathbb{P}_k^{m'} \end{aligned}$$

with associated Hilbert-Burch matrices $\Phi_{\tau_m} \in \mathbb{R}_m^{(m+1) \times m}$ and $\Phi_{\tau_{m'}} \in \mathbb{R}_{m'}^{(m'+1) \times m'}$. Let also $\Phi_{[\tau_m | \tau_{m'}]} \in \mathbb{R}^{(m+m'+1) \times (m+m')}$ be as in Proposition 1. Then:

- (1) $\text{codim } \mathbb{V}(I_{m+m'}(\Phi_{[\tau_m | \tau_{m'}]})) = 2$,
- (2) the ideal of $\mathbb{P}(I_{[\tau_m | \tau_{m'}]})$ is minimally resolved by $\mathbb{F}_{\tau_m} \otimes \mathbb{F}_{\tau_{m'}}$, which is the Koszul complex on the entries of the line matrix $(y_0 \ \dots \ y_{m+m'}) \Phi_{[\tau_m | \tau_{m'}]}$.
- (3) $\mathbb{P}(I_{[\tau_m | \tau_{m'}]}) = \Gamma_{[\tau_m | \tau_{m'}]}$ and, consequently, for any $k \in \{0, \dots, m+m'\}$

$$(1.1) \quad d_k(\tau_{m+m'}) = \sum_{p=0}^k d_p(\tau_m) d_{k-p}(\tau_{m'}) = \binom{m+m'}{k}.$$

Proof. Let $m, m' \geq 1$ and

$$(1.2) \quad \Phi_{\tau_{m+m'}} = \begin{pmatrix} x_0 & 0 & \dots & \dots & 0 \\ -x_1 & x_1 & \ddots & & \vdots \\ 0 & -x_2 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ \vdots & & \ddots & \ddots & \vdots \\ 0 & \dots & \dots & 0 & \begin{matrix} x_{m+m'-1} \\ -x_{m+m'} \end{matrix} \end{pmatrix}$$

Since

$$\mathbf{I}_{m+m'}(\Phi_{[\tau_m|\tau_{m'}]}) = (x_1 \cdots x_{m+m'}, \dots, x_0 \cdots x_{m+m'-1}),$$

one has that $\text{codim } \mathbb{V}(\mathbf{I}_{m+m'}(\Phi_{[\tau_m|\tau_{m'}]})) = 2$ (the generators of $\mathbf{I}_{m+m'}(\Phi_{[\tau_m|\tau_{m'}]})$ do not share a common factor) and thus Item (1) is verified.

The proof of Item (2) and Item (3) relies on an induction on m and m' . The inductive step consists in first showing that the tensor product $\mathbb{F}_{\tau_m} \otimes \mathbb{F}_{\tau_{m'}}$ of a free resolution \mathbb{F}_{τ_m} of $\mathbb{P}(\mathbf{I}_{\tau_m})$ and a free resolution $\mathbb{F}_{\tau_{m'}}$ of $\mathbb{P}(\mathbf{I}_{\tau_{m'}})$ provides a free resolution of $\mathbb{P}(\mathbf{I}_{\tau_m}) \cap \mathbb{P}(\mathbf{I}_{\tau_{m'}}) = \mathbb{P}(\mathbf{I}_{[\tau_m|\tau_{m'}]})$. The second step consists in showing that $\mathbb{P}(\mathbf{I}_{[\tau_m|\tau_{m'}]})$ is irreducible or, in other words, that $\mathbf{I}_{\mathbb{P}(\mathbf{I}_{\tau_m})} + \mathbf{I}_{\mathbb{P}(\mathbf{I}_{\tau_{m'}})}$ is prime. This latter property insures that

$$\Gamma_{\tau_m} \cap \Gamma_{\tau_{m'}} = \mathbb{P}(\mathbf{I}_{\tau_m}) \cap \mathbb{P}(\mathbf{I}_{\tau_{m'}}) = \mathbb{P}(\mathbf{I}_{[\tau_m|\tau_{m'}]}) = \Gamma_{[\tau_m|\tau_{m'}]}$$

and Equation (1.1) follows then from a direct application of Proposition 1.

- Initial case: Item (2) and Item (3) are verified in the case $m = 1$, this follows from the fact that $\mathbb{P}(\mathbf{I}_{\tau_1}) = \mathbb{V}(x_0y_0 - x_1y_1)$ whose minimal free resolution is the Koszul complex on the single irreducible polynomial $x_0y_0 - x_1y_1 \in \mathbb{k}[x_0, x_1, y_0, y_1]$.
- Inductive step: let $m, m' \geq 1$ and assume that Item (2) and Item (3) hold for τ_m and $\tau_{m'}$. In particular the ideal \mathcal{I}_m of $\mathbb{P}(\mathbf{I}_{\tau_m}) = \Gamma_{\tau_m}$ (resp. the ideal $\mathcal{I}_{m'}$ of $\mathbb{P}(\mathbf{I}_{\tau_{m'}}) = \Gamma_{\tau_{m'}}$) is minimally resolved by the Koszul complex \mathbb{F}_{τ_m} on the entries of the line matrix $(y_0 \ \dots \ y_m) \Phi_{\tau_m}$ (resp. by the Koszul complex $\mathbb{F}_{\tau_{m'}}$ on the entries of the line matrix $(y_m \ \dots \ y_{m'}) \Phi_{\tau_{m'}}$). We show Item (2) and Item (3) via classical methods involving Gröbner bases and we refer to [AL94] for the associated definitions associated to this argument. Remark that $\mathbb{F}_{\tau_m} \otimes \mathbb{F}_{\tau_{m+m'}}$ is the Koszul complex on the entries of $(y_0 \ \dots \ y_{m+m'}) \Phi_{[\tau_m|\tau_{m'}]}$ and one has that $\mathbb{F}_{\tau_m} \otimes \mathbb{F}_{\tau_{m'}}$ resolves the ideal of $\mathbb{P}(\mathbf{I}_{[\tau_m|\tau_{m'}]})$ if and only if

$$\forall i \geq 1, \text{Tor}^i(\mathbb{S}/\mathcal{I}_m, \mathbb{S}/\mathcal{I}_{m'}) = 0,$$

\mathbb{S} being the polynomial ring $\mathbb{k}[x_0, \dots, x_{m+m'}, y_0, \dots, y_{m+m'}]$. Actually, by Tor rigidity, the latter conditions are verified if and only if

$$\text{Tor}^1(\mathbb{S}/\mathcal{I}_m, \mathbb{S}/\mathcal{I}_{m'}) = \mathcal{I}_m \cap \mathcal{I}_{m'} / \mathcal{I}_m \cdot \mathcal{I}_{m'} = 0.$$

To show this last condition, i.e. to show that

$$\mathcal{I}_m \cap \mathcal{I}_{m'} = \mathcal{I}_m \cdot \mathcal{I}_{m'},$$

we compute $\mathcal{I}_m \cap \mathcal{I}_{m'}$ by eliminating t in the ideal $t\mathcal{I}_m + (1-t)\mathcal{I}_{m'} \subset \mathbb{S}[t]$ see [AL94, Prop. 2.3.5] for this standard use of Gröbner bases. Using the

- for $i \in \{m+2, \dots, m+m'+1\}$, $\Phi_{ij} = 0$.
- for any $j \in \{m+1, \dots, m+m'\}$, let $k_j \geq 2$ and $\lambda_{1,j}, \dots, \lambda_{k_j,j} \in \mathbb{R}_{m'} = \mathbb{k}[x_m, \dots, x_{m+m'}] \subset \mathbb{R}$ and:
 - for $i \in \{1, \dots, m\}$, $\Phi_{ij} = 0$,
 - for $i \in \{m+1, \dots, m+m'+1\}$, let $\Phi_{ij} \in |\lambda_{1,j}, \dots, \lambda_{k_j,j}|$ be a general linear combination of $\lambda_{1,j}, \dots, \lambda_{k_j,j}$.

Let also $\Phi_g = (\Phi_{ij})_{\substack{0 \leq i \leq m+1 \\ 1 \leq j \leq m}} \in \mathbb{R}_m^{(m+1) \times m}$ and $\Phi_{g'} = (\Phi_{ij})_{\substack{m+1 \leq i \leq m+m'+1 \\ m+1 \leq j \leq m+m'}} \in \mathbb{R}_n^{(m'+1) \times m'}$ and $[g|g'] : \mathbb{P}^{m+m'} \dashrightarrow \mathbb{P}^{m+m'}$, $g : \mathbb{P}^m \dashrightarrow \mathbb{P}^m$ and $g' : \mathbb{P}^{m'} \dashrightarrow \mathbb{P}^{m'}$ be the determinantal maps defined by $\Phi_{[g|g']}$, Φ_g and $\Phi_{g'}$ (where $\mathbb{P}^m = \text{Proj}(\mathbb{R}_m) \subset \text{Proj}(\mathbb{R}) = \mathbb{P}^{m+m'}$ and $\mathbb{P}^{m'} = \text{Proj}(\mathbb{R}_{m'}) \subset \text{Proj}(\mathbb{R}) = \mathbb{P}^{m+m'}$).

Then for all $i \in \{0, \dots, m+m'\}$:

$$(1.3) \quad d_k([g|g']) = \sum_{p=0}^k d_p(g) d_{k-p}(g').$$

2. PROJECTIVE DEGREES VS MIXED VOLUMES OF KOSZUL-DETERMINANTAL MAPS

For the rest of the paper, we let $\mathbb{k} = \mathbb{C}$. Given a map $f : \mathbb{P}_{\mathbb{k}}^n \dashrightarrow \mathbb{P}_{\mathbb{k}}^n$ of base ideal I_f , the inclusion $\Gamma_f \subset \mathbb{P}(I_f)$ of the graph Γ_f of f in the Proj of the symmetric algebra of I_f , see below Definition 1.5 for more details, provides the following estimations:

$$(2.1) \quad \forall k \in \{0, \dots, n\}, \quad d_k(f) \leq \deg_{\mathbb{P}}^{n-k, k} \mathbb{P}(I_f).$$

Let us now present a specificity when computing the projective degrees of the Koszul-determinantal map f we are now going to define. Recall Definition 1.5 that, assuming that f is determinantal, the presentation matrix Φ_f of I_f is called the Hilbert-Burch matrix of f . In this latter case, letting

$$0 \rightarrow \bigoplus_{k=1}^n \mathbb{R}(-d_k) \xrightarrow{\Phi_f} \mathbb{R}^{n+1} \xrightarrow{(f_0 \dots f_n)} I_f(d) \rightarrow 0$$

be a graded free resolution of the base ideal of f , we call the positive integer $d_1, \dots, d_n \geq 1$ the *Hilbert-Burch degrees* of f .

Remark from the equations $(y_0 \dots y_n)\Phi_f$ defining the embedding of $\mathbb{P}(I_f)$ in $\mathbb{P}_{\mathbb{k}}^n \times \mathbb{P}_{\mathbb{k}}^n$ that the conditions:

$$(2.2) \quad \forall k \in \{1, \dots, n-1\}, \quad \text{codim } \mathbb{V}(I_k(\Phi_f)) \geq n+1-k$$

are the required conditions in order that $\mathbb{P}(I_f)$ has codimension n in which case $\mathbb{P}(I_f)$ is a complete intersection and a minimal free resolution of its coordinate ring is the Koszul complex on the entries of $(y_0 \dots y_n)\Phi_f$.

Definition 2.1 (Koszul-determinantal map). A *Koszul-determinantal map* $f = (f_0 : \dots : f_n) : \mathbb{P}_{\mathbb{k}}^n \dashrightarrow \mathbb{P}_{\mathbb{k}}^n$ is a map such that the following two conditions are verified:

- a minimal presentation of $I_f = (f_0, \dots, f_n)$ reads:

$$0 \rightarrow \mathbb{R}^n \xrightarrow{\Phi_f} \mathbb{R}^{n+1} \xrightarrow{(f_0 \dots f_n)} I_f \rightarrow 0.$$

- the conditions (2.2) are satisfied.

Proposition 2.2. *Let $f = (f_0 : \dots : f_n) : \mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^n$ be a Koszul-determinantal map of Hilbert-Burch degree $d_1, \dots, d_n \geq 1$, then for all $k \in \{0, \dots, n\}$,*

$$(2.3) \quad d_k(f) \leq \sigma_{n-k,n}(d_1, \dots, d_n) = \sum_{\{i_1, \dots, i_k\} \subset \{1, \dots, n\}} d_{i_1} \dots d_{i_k}$$

where $\sigma_{k,n}(u_1, \dots, u_n)$ is the k -th symmetric polynomial in n variables.

Proof. Letting Φ_f be the $(n+1) \times n$ Hilbert-Burch matrix of f , the ideal J_1 of $\mathbb{P}(\mathbb{I}_f)$ in $\mathbb{P}_k^n \times \mathbb{P}_k^n$ is generated by the n entries ϕ_1, \dots, ϕ_n of the line matrix $(y_0 \dots y_n)\Phi_f$. Since conditions (2.2) are moreover satisfied, $\mathbb{P}(\mathbb{I}_f)$ has codimension n in $\mathbb{P}_k^n \times \mathbb{P}_k^n$ so $\mathbb{P}(\mathbb{I}_f)$ is a complete intersection and the Koszul complex on $\phi_1, \dots, \phi_n \in \mathbb{S}$

$$0 \rightarrow \mathbb{S}(-\sum_{k=1}^n d_k, -n) \rightarrow \dots \rightarrow \wedge^2 \left(\bigoplus_{k=1}^n \mathbb{S}(-d_k, -1) \right) \rightarrow \bigoplus_{k=1}^n \mathbb{S}(-d_k, -1) \rightarrow \mathbb{S}$$

provides a minimal bi-graded free resolution of the coordinate ring of $\mathbb{P}(\mathbb{I}_f)$ from which we extract the component of total degree n of $\text{Num}_{\mathbb{S}/J_1}(1 - T_0, 1 - T_1)$ (see [EH00, III.3.6]):

$$\forall k \in \{0, \dots, n\}, \quad \deg_{\mathbb{S}_{\mathbb{P}}}^{n-k,k} \mathbb{P}(\mathbb{I}_f) = s_{n-k,n}(d_1, \dots, d_n) = \sum_{\{i_1, \dots, i_k\} \subset \{1, \dots, n\}} d_{i_1} \dots d_{i_k}.$$

The conclusion of the proposition follows then from (2.1). \square

We point out that the conclusion of Proposition 2.2 is classical and had been established in more general context of maps $f : \mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^{n'}$ with $n \leq n'$, see for instance [CR21, Theorem 5.7]. It explains however why, when interested in determinantal Cremona maps, one has also to consider base ideal not of linear type or else, if the base ideal is of linear type, the entries of the presentation matrix can only contain linear polynomials (case $d_1 = \dots = d_n = 1$ of the previous proposition). In other words, one has to consider the kernel of the surjection $\text{Sym}(\mathbb{I}_f) \rightarrow \mathcal{R}(\mathbb{I}_f)$ in order to understand the level of approximations of the inequalities (2.3). As we previously explained in Section 1, this kernel is described by the ideals $\mathbb{I}_k(\Phi_f)$ for $k \in \{1, \dots, n-1\}$ of k -minors ideals of the presentation matrix Φ_f of \mathbb{I}_f which, in our context of determinantal base ideal \mathbb{I}_f , have an expected codimension:

$$\forall k \in \{1, \dots, n\}, \quad \text{codim } \mathbb{V}(\mathbb{I}_k(\Phi_f)) \geq n + 2 - k,$$

ensuring, if these latter conditions are verified, that \mathbb{I}_f is of linear type [RS01, Subsection 2.1]. Hence the conditions (2.2) are the first step to consider when interested in determinantal Cremona maps.

2.1. Bernstein theorem's bound on the number of solutions of a zero dimensional polynomial system. When interested in the topological degree $d_0(f)$ of a determinantal map $f = (f_0 : \dots : f_n) : \mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^n$, remark that two polynomial systems can be considered. An initial one, in the end the main object of this work, is defined by f itself: given $\mathbf{y} = (y_0 : \dots : y_n) \in \mathbb{P}_k^n$ in the target space of f , one wants to find the pre-images $\mathbf{x} \in \mathbb{P}_k^n$ of \mathbf{y} , i.e. wants to solve the polynomial system

$$(2.4) \quad f(\mathbf{x}) = \mathbf{y} \Leftrightarrow \begin{cases} y_0 &= f_0(x_0, \dots, x_n) \\ &\vdots \\ y_n &= f_n(x_0, \dots, x_n) \end{cases}$$

Expressing \mathbf{y} as the intersection of n hyperplans in the target space of f and assuming that \mathbf{y} is general enough, Bézout theorem then asserts that $d_0(f) = \#f^{-1}(\{\mathbf{y}\}) \leq d^n$ where $d = d_{n-1}(f)$ is the common degree of the polynomials f_i generating the base ideal $\mathbf{I}_f = (f_0, \dots, f_n)$ of f . An intermediate polynomial system is moreover defined by a presentation matrix Φ_f of \mathbf{I}_f . Indeed, by definition of Φ_f , one has $(f_0 \dots f_n)\Phi_f = 0$ so the polynomial system

$$(2.5) \quad (y_0 \dots y_n)\Phi_f = 0$$

whose number of equations is the number of columns of Φ_f , contains by definition $f^{-1}(\{\mathbf{y}\})$. Now consider the situation where $\Phi_f = (\phi_{ij})_{\substack{0 \leq i \leq n \\ 1 \leq j \leq n}}$ is a $(n+1) \times n$ matrix where each entry ϕ_{ij} of the j -th column of Φ_f has degree $d_j \geq 1$ and in which case $d = d_{n-1}(f) = d_1 + \dots + d_n$ by Hilbert-Burch theorem. Then, provided that $\text{codim } \mathbb{V}(\mathbf{I}_k(\Phi_f)) \geq n+1-k$ for all $k \in \{1, \dots, n\}$, the system (2.5) is 0-dimensional in \mathbb{P}_k^n and, by Bézout theorem, has $d_1 \cdot \dots \cdot d_n$ solutions. Moreover, in case the entries of Φ_f are sparse polynomials that all verify the same algebraic constraints on their coefficients, one can refine the bound on the number of solutions of (2.5) by using Bernstein's theorem on sparse polynomials. It then gives a combinatorial translation, via the computation of *mixed volumes* associated to the polynomial entries of Φ_f , to the problem of detecting determinantal Cremona maps of given Hilbert-Burch degree (d_1, \dots, d_n) among all determinantal maps of Hilbert-Burch degree (d_1, \dots, d_n) .

This idea structures our approach of Proposition 2.14, see [TV05] or [LM09] for developments in more general contexts and for pointers to the related literature. We refer to [CLO05, Chapter 7] for the background in convex geometry and all the material we use for the actual computations. Following [CLO05, 7.4], a set $C \subset \mathbb{R}^n$ is *convex* if it contains any segments between two points in C and the *convex hull* $\text{Conv}(S)$ of a subset $S \subset \mathbb{R}^n$ is the smallest convex set containing S . A *polytope* is the convex hull $\text{Conv}(A)$ of a finite set $A \subset \mathbb{R}^n$ and the polytopes which are the convex hull of points with integer coordinates are called *lattice polytopes*.

Given $\phi = \sum_{\alpha \in \mathbb{N}^n} c_\alpha x^\alpha \in k[x_1, \dots, x_n]$ where $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$, the *Newton polytope* of ϕ , denoted $\text{NP}(\phi)$, is the lattice polytope $\text{NP}(\phi) = \text{Conv}\{\alpha \in \mathbb{N}^n, c_\alpha \neq 0\}$.

A polytope $P \subset \mathbb{R}^n$ has an n -dimensional volume $\text{Vol}_n(P)$. Given two polytopes $P, Q \subset \mathbb{R}^n$ and a real number $\lambda \geq 0$, the *Minkowski sum* of P and Q , denoted $P+Q$ is the set

$$P+Q := \{p+q, p \in P, q \in Q\}$$

where $p+q$ denotes the usual vector sum in \mathbb{R}^n and the λP stands for the polytope $\{\lambda p, p \in P\}$ where λp is the usual scalar multiplication in \mathbb{R}^n .

Given any collection $P_1, \dots, P_r \subset \mathbb{R}^n$ and r non negative scalar $\lambda_1, \dots, \lambda_r \in \mathbb{R}$, then $\text{Vol}_n(\lambda_1 P_1 + \dots + \lambda_r P_r)$ is a homogeneous polynomial of degree n in the λ_i [CLO05, 7. Prop.4.9].

Definition 2.3 (mixed volume of a collection of polytopes). The n -dimensional *mixed volume* $\text{MV}_n(P_1, \dots, P_n)$ of given polytopes P_1, \dots, P_n is the coefficient of the monomial $\lambda_1, \dots, \lambda_n$ in $\text{Vol}_n(P_1, \dots, P_n)$.

Theorem 2.4. [CLO05, 7. Th.4.12]

- (i) *The mixed volume $\text{MV}_n(P_1, \dots, P_n)$ is invariant if the P_i are replaced by their images under a volume-preserving transformation of \mathbb{R}^n .*

- (ii) $MV_n(P_1, \dots, P_n)$ is symmetric and linear in each variable (multilinearity of the mixed volume).
- (iii) $MV_n(P_1, \dots, P_n) \geq 0$ and $MV_n(P_1, \dots, P_n) = 0$ if one of the P_i has dimension zero (i.e. if P_i consists of a single point).
 $MV_n(P_1, \dots, P_n) > 0$ if every P_i has dimension n .
- (iv) $MV_n(P_1, \dots, P_n) = \sum_{k=1}^n (-1)^{n-k} \sum_{\substack{I \subset \{1, \dots, n\} \\ |I|=k}} \text{Vol}_n(\sum_{i \in I} P_i)$ where $\sum_{i \in I} P_i$ is the Minkowski sum of polytope.

In addition to Theorem 2.4, our work relies also on the following toolbox.

Lemma 2.5. [ST10, Lemma 6] *Let $n, n' \in \mathbb{N}^*$ and let P_1, \dots, P_n be polytopes in $\mathbb{R}^{n+n'}$ and $P_{n+1}, \dots, P_{n+n'}$ be polytopes in $\mathbb{R}^n \times \{0_{\mathbb{R}^{n'}}\} \subset \mathbb{R}^{n+n'}$. Then:*

$$MV_{n+n'}(P_1, \dots, P_{n+n'}) = MV_n(P_1, \dots, P_n) MV_{n'}(\pi_n(P_{n+1}), \dots, \pi_n(P_{n+n'}))$$

where $\pi_{n'} : \mathbb{R}^{n+n'} \rightarrow \mathbb{R}^{n'}$ stands for the projection on the last n' coordinates.

Following [CLO05, 7 Section 5], let us now define the genericity of a polynomial with respect to a polytope.

Definition 2.6 (genericity with respect to a polytope). Given finite set $A \subset \mathbb{Z}^n$, put $L(A) := \{ \sum_{\alpha \in A} c_\alpha x^\alpha \in k[x_1, \dots, x_n] \}$ and remark that each $L(A)$ can be considered as an affine space $k^{\#A}$ with the coordinate c_α as coordinates.

A polynomial $\sum_{\alpha \in A} c_\alpha x^\alpha$ is said to be *generic with respect to $L(A)$* if its coefficients are generic in $L(A)$.

Given $k \geq 1$ and finite sets $A_1, \dots, A_k \in \mathbb{Z}^n$, a property is said to *hold generically* for polynomials $(\phi_1, \dots, \phi_n) \in L(A_1) \times \dots \times L(A_k)$ if there is a non zero polynomial in the coefficients of the ϕ_i such that the property holds for all ϕ_1, \dots, ϕ_n for which the polynomial is non vanishing, in particular if every ϕ_1, \dots, ϕ_n is generic with respect to its own polytope $L(A_1), \dots, L(A_k)$.

Theorem 2.7 (Bernstein theorem). *Given polynomials $\phi_1, \dots, \phi_n \in k[x_1, \dots, x_n]$ with finitely many common zeroes in $(k^*)^n$, let $P_i = \text{NP}(\phi_i)$. Then the number of common zeroes in $(\mathbb{C}^*)^n$ is bounded above by the mixed volume $MV_n(P_1, \dots, P_n)$. Moreover if each ϕ_i is generic with respect to P_i , the number of common zero solutions is exactly $MV_n(P_1, \dots, P_n)$.*

See [CLO05, Proof of 7. Th. 5.4] and the references therein for highlights about Bernstein theorem.

2.2. Projective degrees of Koszul-determinantal maps defined by sparse polynomials. Given a homogeneous polynomial $\phi \in R = k[x_0, \dots, x_n]$, the Newton polytope $\text{NP}(\phi) \subset \mathbb{R}^n$ associated to ϕ is the Newton polytope of the de-homogenization of ϕ with respect to a variable x_i (omitted when irrelevant) and, reciprocally, given a polytope $P \subset \mathbb{R}^n$, we denote $\phi \in P$ for a homogeneous polynomial $\phi \in R$ whose de-homogenization with respect to the variable x_i is in P . Furthermore, we consider more general Newton polytopes $P \subset \mathbb{R}^n \times \mathbb{R}^n$ corresponding to bi-homogeneous polynomials $\phi \in S = R[y_0, \dots, y_n]$, that is the bi-de-homogenization of ϕ with respect to one fixed variable x_i and one fixed variable y_j ($i, j \in \{0, \dots, n\}$) is in P .

Let us also denote by $S_n = \text{Conv}\{(0, \dots, 0), (1, \underbrace{0, \dots, 0}_{n-1}), \dots, (\underbrace{0, \dots, 0}_{n-1}, 1)\} \subset \mathbb{R}^n$ the unit simplex of \mathbb{R}^n and let $S_n^x := S_n \times \{0_{\mathbb{R}^n}\} \subset \mathbb{R}^n \times \mathbb{R}^n$ and $S_n^y := \{0_{\mathbb{R}^n}\} \times S_n \subset \mathbb{R}^n \times \mathbb{R}^n$.

Proposition 2.8. *Let $f : \mathbb{P}_k^n \rightarrow \mathbb{P}_k^n$ be a determinantal map of Hilbert-Burch matrix $\Phi_f = (\phi_{ij})_{\substack{0 \leq i \leq n \\ 1 \leq j \leq n}}$ of Hilbert-Burch degree (d_1, \dots, d_n) . Then for all $k \in \{0, \dots, n\}$:*

$$d_k(f) \leq \text{MV}_{2n}(\underbrace{S_n^x, \dots, S_n^x}_k, P_1^y, \dots, P_n^y, \underbrace{S_n^y, \dots, S_n^y}_{n-k}) \leq \sigma_{n-k,n}(d_1, \dots, d_n)$$

where for $l \in \{1, \dots, n\}$, $P_l^y \subset \mathbb{P}_k^n \times \mathbb{P}_k^n$ is the Newton polytope of the l -th entry of the matrix $(y_0 \dots y_n)\Phi_f$ and $\sigma_{k,n}$ is the k -th symmetric polynomial in n -variables.

Proof. As explained in Remark 1.2, for all $k \in \{0, \dots, n\}$, $\deg_{\mathbb{P}}^{n-k,k} \mathbb{P}(I_f)$ is the number of common zero solutions in $\mathbb{P}_k^n \times \mathbb{P}_k^n$ of the polynomials

$$(2.6) \quad l_{1,0}, \dots, l_{k,0}, \phi_1, \dots, \phi_n, l_{0,1}, \dots, l_{0,n-k}$$

where $l_{1,0}, \dots, l_{k,0}$ (resp. $l_{0,1}, \dots, l_{0,n-k}$) are generic with respect to S_n^x (resp. generic with respect to S_n^y) and $\phi_1, \dots, \phi_n \in \mathbb{S} = \mathbb{R}[y_0, \dots, y_n]$ are the entries of the matrix $(y_0 \dots y_n)\Phi_f$ (the associated polynomial system being 0-dimensional by definition of a determinantal map, see Definition 1.5).

In this setting, by Theorem 2.7 in the generic case, the quantity

$$\text{MV}_{2n}(\underbrace{S_n^x, \dots, S_n^x}_k, P_1^y, \dots, P_n^y, \underbrace{S_n^y, \dots, S_n^y}_{n-k})$$

is the number of solutions of (2.6) whose coordinates are all non zero which shows the right hand side inequality of Proposition 2.8 since the number of common zero of (2.6) is equal to $\sigma_{n-k,n}(d_1, \dots, d_n)$.

The set of common zero of (2.6) contains moreover the set

$$H_x^k \cap \Gamma_f \cap H_y^{n-k} \subset \mathbb{P}_k^n \times \mathbb{P}_k^n$$

where $H_x^k = \mathbb{V}(l_{1,0}, \dots, l_{k,0})$ is the zero locus of $l_{1,0}, \dots, l_{k,0}$, Γ_f is the graph of f and $H_y^{n-k} = \mathbb{V}(l_{0,1}, \dots, l_{0,n-k})$. However, by the genericity assumptions, remark that the points of $H_x^k \cap \Gamma_f \cap H_y^{n-k}$ have all non zero coordinates which shows the left hand side inequality of Proposition 2.8. \square

Let us now isolate two technical facts that rely on Lemma 2.5:

Lemma 2.9. *Let $P_1, \dots, P_n \subset \mathbb{R}^n$ be n polytopes and for $j \in \{1, \dots, n\}$ put $P_j^y := P_j \times \{0_{\mathbb{R}^n}\} + \{0_{\mathbb{R}^n}\} \times S_n^y \subset \mathbb{R}^n \times \mathbb{R}^n$.*

Then

$$\text{MV}_{2n}(P_1^y, \dots, P_n^y, \underbrace{S_n^y, \dots, S_n^y}_n) = \text{MV}_n(P_1, \dots, P_n).$$

Proof. It suffices to apply the multilinearity of the mixed volume (Theorem 2.4. Item (ii)), the projection formula (Lemma 2.5) and the vanishing condition (Theorem 2.4. Item (iii)) on the left member of the equality to obtain:

$$\begin{aligned} \text{MV}_{2n}(P_1^y, \dots, P_n^y, \underbrace{S_n^y, \dots, S_n^y}_n) &= \text{MV}_n(P_1, \dots, P_n) \text{MV}_n(\underbrace{S_n, \dots, S_n}_n) \\ &= \text{MV}_n(P_1, \dots, P_n). \end{aligned}$$

□

Lemma 2.10. *Let $m, m' \in \mathbb{N}^*$ and:*

- $P_1, \dots, P_m \subset \mathbb{R}^m \times \{0_{\mathbb{R}^{m'}}\} \subset \mathbb{R}^m \times \mathbb{R}^{m'}$,
 - $P_{m+1}, \dots, P_{m+m'} \subset \{0_{\mathbb{R}^m}\} \times \mathbb{R}^{m'} \subset \mathbb{R}^m \times \mathbb{R}^{m'}$,
- be $m + m'$ polytopes. For $j \in \{1, \dots, m + m'\}$ put:

$$P_j^y := P_j \times \{0_{\mathbb{R}^{m+m'}}\} + \{0_{\mathbb{R}^{m+m'}}\} \times S_{m+m'}^y \subset \mathbb{R}^{m+m'} \times \mathbb{R}^{m+m'}.$$

Then for all $k \in \{0, \dots, m + m'\}$,

$$\begin{aligned} & \text{MV}_{2(m+m')}(\underbrace{S_{m+m'}^x, \dots, S_{m+m'}^x}_{m+m'-k}, P_1^y, \dots, P_{m+m'}^y, \underbrace{S_{m+m'}^y, \dots, S_{m+m'}^y}_k) \\ &= \\ & \sum_{p=0}^k \left[\left(\sum_{\substack{\{l_1, \dots, l_p\} \subset \\ \{1, \dots, m'\}}} \text{MV}_{m'}(\underbrace{S_{m'}, \dots, S_{m'}}_{m'-p}, P_{l_1}, \dots, P_{l_p}) \right) \times \right. \\ & \left. \left(\sum_{\substack{\{l_1, \dots, l_{k-p}\} \subset \\ \{m'+1, \dots, m+m'\}}} \text{MV}_m(\underbrace{S_m, \dots, S_m}_{m+p-k}, P_{l_1}, \dots, P_{l_{k-p}}) \right) \right] \end{aligned}$$

Proof. The formula follows from by decomposing first $S_{m+m'}^x \subset \mathbb{R}^{m+m'} \times \mathbb{R}^{m+m'}$ (resp. $S_{m+m'}^y$) as the sum $S_m^x \times \{0_{\mathbb{R}^{m'}}\} \times \{0_{\mathbb{R}^{m+m'}}\} + \{0_{\mathbb{R}^m}\} \times S_{m'}^x \times \{0_{\mathbb{R}^{m+m'}}\} \subset \mathbb{R}^{m+m'} \times \mathbb{R}^{m+m'}$ (resp. as the sum $\{0_{\mathbb{R}^{m+m'}}\} \times S_m^y \times \{0_{\mathbb{R}^{m'}}\} + \{0_{\mathbb{R}^{m+m'}}\} \times \{0_{\mathbb{R}^m}\} \times S_{m'}^y \times \{0_{\mathbb{R}^{m+m'}}\}$) and then applying the multilinearity of the mixed volume (Theorem 2.4. Item (ii)), the projection formula (Lemma 2.5), the vanishing condition (Theorem 2.4. Item (iii)) and eventually Lemma 2.9. □

Proposition 2.11. *Let $f : \mathbb{P}_k^n \dashrightarrow \mathbb{P}_k^n$ be a Koszul-determinantal map of Hilbert-Burch matrix $\Phi_f = (\phi_{ij})_{\substack{0 \leq i \leq n \\ 1 \leq j \leq n}}$. Then:*

$$\begin{aligned} \forall k \in \{0, \dots, n\}, d_k(f) &= \text{MV}_{2n}(\underbrace{S_n^x, \dots, S_n^x}_k, P_1^y, \dots, P_n^y, \underbrace{S_n^y, \dots, S_n^y}_{n-k}) \\ &\updownarrow \\ & \overline{\mathbb{P}(\text{I}_f) \setminus \Gamma_f} \subset \mathbb{V}(\prod_{i=0}^n x_i) \subset \mathbb{P}_k^n \times \mathbb{P}_k^n \end{aligned}$$

where for $l \in \{1, \dots, n\}$, $P_l^y \subset \mathbb{R}^n \times \mathbb{R}^n$ is the Newton polytope of the l -th entry of the matrix $(y_0 \dots y_n)\Phi_f$.

Proof. Assume that $\overline{\mathbb{P}(\text{I}_f) \setminus \Gamma_f} \subset \mathbb{V}(\prod_{i=0}^n x_i) \subset \mathbb{P}_k^n \times \mathbb{P}_k^n$ so $\mathbb{P}(\text{I}_f)$ and Γ_f coincide away the coordinate axis defined by the x_i . Now for any $k \in \{0, \dots, n\}$,

$$(2.7) \quad \text{MV}_{2n}(\underbrace{S_n^x, \dots, S_n^x}_k, P_1^y, \dots, P_n^y, \underbrace{S_n^y, \dots, S_n^y}_{n-k})$$

is the number of point with non zero coordinate in the \mathbf{x} -variable of the reduced scheme $H_{\mathbf{x}}^k \cap \mathbb{P}(\text{I}_f) \cap H_{\mathbf{y}}^{n-k}$ where $H_{\mathbf{x}}^k$ is the zero locus of k general linear forms in the \mathbf{x} -variables and $H_{\mathbf{y}}^{n-k}$ is the zero locus of $c - k$ general linear forms in the \mathbf{y} -variables. So (2.7) is also equal to the number of point with non zero coordinate in the \mathbf{x} -variable of the reduced scheme $H_{\mathbf{x}}^k \cap \Gamma_f \cap H_{\mathbf{y}}^{n-k}$. But since $H_{\mathbf{x}}^k$ and $H_{\mathbf{y}}^{n-k}$

are general linear subspace, this last quantity is equal to $d_k(f)$ which show the indirect implication in Proposition 2.11.

Now assume that $\overline{\mathbb{P}(\mathbb{I}_f) \setminus \Gamma_f} \not\subset \mathbb{V}(\prod_{i=0}^n x_i)$. This implies that there exists a codimension n irreducible component \mathbb{T} of $\mathbb{P}(\mathbb{I}_f)$, distinct from Γ_f , that is not included in $\mathbb{V}(\prod_{i=0}^n x_i)$. Since \mathbb{T} lies above the zero locus of some Fitting ideals of Φ_f , it not included neither in $\mathbb{V}(\prod_{i=0}^n y_i)$. Hence, there is a $k \in \{0, \dots, n\}$ such that for any $H_{\mathbf{x}}^k$ which is the zero locus of k general linear forms in the \mathbf{x} -variables and any $H_{\mathbf{y}}^k$ which is the zero locus of $c - k$ general linear forms in the \mathbf{y} -variables, the scheme $H_{\mathbf{x}}^k \cap \mathbb{T} \cap H_{\mathbf{y}}^{n-k}$ contain a point with all non zero coordinate. Since this point is taken in account in the mixed volume (2.7), one has the strict inequality $d_k(f) < (2.7)$. \square

Hence, to know the actual term $d_k(f)$ of the projective degrees of a determinantal map f , the computation of the associated mixed volume has to be completed by a preliminary control on the support of the successive ideal of minors of Φ_f . We illustrate such a control in the next subsection.

2.3. Gluing determinantal maps subject to general conditions. Let us now focus on another construction of a *glued determinantal map*

$$[g|g'] : \mathbb{P}_{\mathbf{k}}^{m+m'} \dashrightarrow \mathbb{P}_{\mathbf{k}}^{m+m'},$$

where $m, m' \in \mathbb{N}^*$, starting from two determinantal Cremona maps $g : \mathbb{P}_{\mathbf{k}}^m \dashrightarrow \mathbb{P}_{\mathbf{k}}^m$ and $g' : \mathbb{P}_{\mathbf{k}}^{m'} \dashrightarrow \mathbb{P}_{\mathbf{k}}^{m'}$. In the following, given polynomials $\phi, \psi_1, \dots, \psi_l$, we denote by $\phi \in |\psi_1, \dots, \psi_l|$ the condition that $\phi = \sum_{k=1}^l \lambda_k \psi_k$ is a general linear combination of ψ_1, \dots, ψ_l .

Proposition-Definition 2.12 (glued map). Given any $j \in \{1, \dots, m + m'\}$, let $l_j \in \mathbb{N}$ and let:

- $\psi_0^{(j)}, \dots, \psi_{l_j}^{(j)} \in \mathbb{R}_m = \mathbb{k}[x_0, \dots, x_m]$ if $j \in \{1, \dots, m\}$
- $\psi_0^{(j)}, \dots, \psi_{l_j}^{(j)} \in \mathbb{R}_{m'} = \mathbb{k}[x_m, \dots, x_{m+m'}]$ if $j \in \{m + 1, \dots, m + m'\}$.

Let also $\Phi_g = (\phi_{ij}^{(g)})_{\substack{0 \leq i \leq m \\ 1 \leq j \leq m}} \in \mathbb{R}_m^{(m+1) \times m}$, $\Phi_{g'} = (\phi_{ij}^{(g')})_{\substack{0 \leq i \leq m' \\ m+1 \leq j \leq m+m'}} \in \mathbb{R}_{m'}^{(m'+1) \times m'}$ and $\Phi_{[g|g']} = (\phi_{ij})_{\substack{0 \leq i \leq m+m' \\ 1 \leq j \leq m+m'}} \in \mathbb{R}_{m+m'}^{(m+m'+1) \times (m+m')}$ where $\mathbb{R}_{m+m'} = \mathbb{k}[x_0, \dots, x_{m+m'}]$ be such that:

- for any $j \in \{1, \dots, m\}$, each entry $\phi_{ij}^{(g)}$ of the j -th column of Φ_g is such that $\phi_{ij}^{(g)} \in |\psi_0^{(j)}, \dots, \psi_{l_j}^{(j)}|$.
- for any $j \in \{1, \dots, m'\}$, each entry $\phi_{ij}^{(g')}$ of the j -th column of $\Phi_{g'}$ is such that $\phi_{ij}^{(g')} \in |\psi_0^{(j+m)}, \dots, \psi_{l_j}^{(j+m)}|$.
- for any $j \in \{1, \dots, m + m'\}$, each entry ϕ_{ij} of the j -th column of $\Phi_{[g|g']}$ is such that $\phi_{ij} \in |\psi_0^{(j)}, \dots, \psi_{l_j}^{(j)}|$.

Assume that $\text{codim } \mathbb{V}(\mathbb{I}_m(\Phi_g)) = \text{codim } \mathbb{V}(\mathbb{I}_{m'}(\Phi_{g'})) = 2$, then

$$\text{codim } \mathbb{V}(\mathbb{I}_{m+m'}(\Phi_{[g|g']})) = 2$$

and define the *glued map* $[g|g'] : \mathbb{P}_k^{m+m'} \dashrightarrow \mathbb{P}_k^{m+m'}$ as the map whose base locus $I_{[g|g']}$ is the $(m+m')$ -minors ideal of $\Phi_{[g|g']}$.

In the following, under the notation of Proposition-Definition 2.12 and under the assumption that $\text{codim } \mathbb{V}(I_m(\Phi_g)) = \text{codim } \mathbb{V}(I_{m'}(\Phi_{g'})) = 2$, we let $g : \mathbb{P}_k^m \dashrightarrow \mathbb{P}_k^m$ (resp. $g' : \mathbb{P}_k^{m'} \dashrightarrow \mathbb{P}_k^{m'}$) be the map whose base locus I_g is equal to $I_m(\Phi_g)$ (resp. $I_{g'} = I_{m'}(\Phi_{g'})$).

Proof. Assume that $\text{codim } \mathbb{V}(I_m(\Phi_g)) = \text{codim } \mathbb{V}(I_{m'}(\Phi_{g'})) = 2$ and suppose by contradiction that $\text{codim } \mathbb{V}(I_{m+m'}(\Phi_{[g|g']})) < 2$. Then there is a common factor to each $m+m'$ minors of $\Phi_{[g|g']}$, so, after operation on columns, one column of $\Phi_{[g|g']}$ have all its entries sharing a common factor. But it is impossible under the genericity assumption on the entries of $\Phi_{[g|g']}$. \square

In the following example, we illustrate however that the glued map $[g|g']$ of two Koszul-determinantal maps g and g' , though determinantal by Proposition-Definition 2.12, may not be Koszul-determinantal.

Example 2.13. Put $m = n = 2$ and let $l_1 = l_2 = l_3 = l_4 = 2$ with:

- $\psi_0^{(1)} = \psi_0^{(2)} = x_1, \psi_1^{(1)} = \psi_1^{(2)} = x_2$
- $\psi_0^{(3)} = \psi_0^{(4)} = x_2, \psi_1^{(3)} = \psi_1^{(4)} = x_3$.

Then, the maps $g : \mathbb{P}_k^2 \dashrightarrow \mathbb{P}_k^2$ and $g' : \mathbb{P}_k^2 \dashrightarrow \mathbb{P}_k^2$ are Koszul-determinantal but $[g|g'] : \mathbb{P}_k^4 \dashrightarrow \mathbb{P}_k^4$ is such that $\text{codim } \mathbb{V}(I_1(\Phi_{[g|g']})) = 3 < 4$ ($I_1(\Phi_{[g|g']}) = (x_1, x_2, x_3)$).

Let us now describe the projective degrees of a glued determinantal map in the almost linear setting.

Proposition 2.14. *Let $m, d \geq 1$, put $l_1 = \dots = l_m = m+1, l_{m+1} = 1, l_{m+2} = m+3$*

- $\forall j \in \{1, \dots, m\}, k \in \{0, \dots, m\}$, let $\psi_k^{(j)} = x_k$
- $\psi_0^{(m+1)} = x_0, \psi_1^{(m+1)} = x_1$
- $\forall k \in \{1, \dots, 3d\}$, $\psi_k^{(m+2)}$ is the k -th generator of the product

$$(x_m, x_{m+1}, x_{m+2}) \cdot (x_{m+1}, x_{m+2})^{d-1} = (x_m x_{m+1}^{d-1}, x_m x_{m+2}^{d-1}) + (x_{m+1}, x_{m+2})^d.$$

Let $[g|g'] : \mathbb{P}_k^{m+2} \dashrightarrow \mathbb{P}_k^{m+2}$ be the glued map associated to the previous data as defined in Proposition-Definition 2.12 and whose base ideal $I_{[g|g']}$ is the $(m+2)$ -minors ideal of $\Phi_{[g|g']}$.

Then $[g|g']$ is a determinantal map and moreover:

$$\forall k \in \{0, \dots, m+2\}, d_k([g|g']) = \binom{m}{m-k} + (d+1) \binom{m}{m-k+1} + \binom{m}{m-k+2}$$

with the convention that $\binom{j}{i} = 0$ if $i < 0$ or $i > j$.

Proof. First, under our general assumptions, $\text{codim } I_{m+2}(\Phi_{[g|g']}) = 2$ and for all $k \in \{1, \dots, n\}$, $\text{codim } I_k(\Phi_{[g|g']}) \geq 3 + m - k$, hence $[g|g']$ is Koszul-determinantal.

Moreover $\overline{\mathbb{P}(I_{[g|g']})} \setminus \Gamma_{[g|g']} \subset \mathbb{V}(\prod_{i=0}^n x_i) \subset \mathbb{P}_k^{m+2} \times \mathbb{P}_k^{m+2}$ hence, by Proposition 2.11,

we can use the mixed volumes of the polytopes $P_j = \text{NP}(\psi_1^{(j)}) + \dots + \text{NP}(\psi_{l_j}^{(j)})$ ($j \in \{1, \dots, m+2\}$) to compute the projective degrees of $[g|g']$.

By applying Lemma 2.10, given any $k \in \{0, \dots, m+2\}$, one has the formula:

$$\begin{aligned}
& \text{MV}_{2(m+2)}(\underbrace{S_{m+2}^{\mathbf{x}}, \dots, S_{m+2}^{\mathbf{x}}}_{m+2-k}, P_1^{\mathbf{y}}, \dots, P_{m+2}^{\mathbf{y}}, \underbrace{S_{m+2}^{\mathbf{y}}, \dots, S_{m+2}^{\mathbf{y}}}_k) \\
&= \\
& \sum_{p=0}^k \left[\left(\sum_{\substack{\{l_1, \dots, l_{k-p}\} \subset \\ \{1, \dots, m\}}} \text{MV}_m(S_{m_1}, \dots, S_{m_{k-p}}, P_{l_1}, \dots, P_{l_{k-p}}) \right) \times \right. \\
& \quad \left. \left(\sum_{\substack{\{l_1, \dots, l_p\} \subset \\ \{m+1, m+2\}}} \text{MV}_2(S_{2_1}, \dots, S_{2_p}, P_{l_1}, \dots, P_{l_p}) \right) \right] \\
& \quad \updownarrow \\
& d_k(g|g') = \sum_{p=0}^k [d_p(g)d_{k-p}(g')]
\end{aligned}$$

with the convention that $d_p(g) = 0$ if $p > m$ and $d_p(g') = 0$ if $p > 2$.

The result of Proposition 2.14 follows from the fact that:

- $d(g) = (1, \binom{m}{1}, \dots, \binom{m}{m-1}, \binom{m}{m})$ as g is general determinantal map, see [GSP06, Theorem 2]
- $d(g') = (d_0(g'), d_1(g'), d_2(g')) = (1, d+1, 1)$, this follows for instance from [BCRS20, Th. 5.14].

□

REFERENCES

- [AL94] W.W. Adams and P. Lounstaunau. *An introduction to Gröbner bases*, volume 3 of *Graduate studies in math.* AMS, 1994. [↑8](#), [↑9](#)
- [BCJ09] L. Busé, M. Chardin, and J.-P. Jouanolou. Torsion of the symmetric algebra and implicitization. *Proc. Amer. Math. Soc.*, 137:1855–1865, 2009. [↑3](#), [↑6](#)
- [BCRS20] L. Busé, Y. Cid-Ruiz, and A. Simis. Degree and birationality of multi-graded rational maps. *Proc. of the London Math. Soc.*, 121(4):743–787, 2020. [↑3](#), [↑18](#)
- [CLO05] D.A. Cox, J. Little, and D. O’Shea. *Using Algebraic Geometry*. Graduate Texts in Mathematics. Springer-Verlag New York, 2005. [↑3](#), [↑12](#), [↑13](#)
- [CR21] Y. Cid-Ruiz. Mixed multiplicities and projective degrees of rational maps. *J. of Algebra*, 566:136–162, 2021. [↑3](#), [↑4](#), [↑11](#)
- [DH17] J. Déserti and F. Han. Quarto-quartic birational maps of $\mathbb{P}^3(\mathbb{C})$. *Internat. J. Math.*, 28, 2017. [↑3](#)
- [EH00] D. Eisenbud and J. Harris. *The Geometry of Schemes*. Graduate Texts in Mathematics. Springer-Verlag New York, 2000. [↑5](#), [↑11](#)
- [Eis95] D. Eisenbud. *Commutative algebra, with a view toward algebraic geometry*. Graduate Texts in Mathematics. Springer, 1995. [↑6](#)
- [GSP06] G. Gonzalez-Sprinberg and I. Pan. On characteristic classes of determinantal Cremona transformations. *Mathematische Annalen*, 335:479–487, 2006. [↑1](#), [↑7](#), [↑18](#)
- [Har92] J. Harris. *Algebraic Geometry*. Graduate Texts in Mathematics. Springer, 1992. [↑2](#), [↑4](#)
- [KM20] Y. Kim and V. Mukundan. Equations defining certain graphs. *Michigan Math. J.*, page 36 pages, 2020. [↑6](#)
- [KPU11] A. Kustin, C. Polini, and B. Ulrich. Rational normal scrolls and the defining equations of Rees algebras. *J. reine angew. Math.*, 650:23–65, 2011. [↑3](#), [↑6](#)
- [KPU13] A. Kustin, C. Polini, and B. Ulrich. The bi-graded structure of Symmetric Algebras with applications to Rees rings. *J. of Algebra*, 469:188–250, 2013. [↑6](#)
- [LM09] R. Lazarsfeld and Mustață. Convex bodies associated to linear series. *Annales scient. de l’ENS*, 42(5):783–835, 2009. [↑12](#)

- [Mic64] A. Micali. Sur les algèbre universelles. *Ann. Inst. Fourier*, 14:33–87, 1964. [↑6](#)
- [MS05] E. Miller and B. Sturmfels. *Combinatorial Commutative Algebra*, volume 227 of *Graduate Texts in Mathematics*. Springer-Verlag, 2005. [↑4](#)
- [RS01] F. Russo and A. Simis. On birational maps and jacobian matrices. *Compositio Math.*, 126:335–358, 2001. [↑3](#), [↑9](#), [↑11](#)
- [ST10] R. Steffens and T. Theobald. Mixed volume techniques for embeddings of Laman graphs. *Computational Geo.*, 43:84–93, 2010. [↑3](#), [↑13](#)
- [Sta17] G. Stagliano. A Macaulay2 package for computations with rational maps. *J. of Software for Alg. and Geo.*, 8(1):61–70, 2017. [↑3](#)
- [TV05] N. V. Trung and J. Verma. Mixed multiplicities of ideals versus mixed volumes of polytopes. *Transactions of the American Mathematical Society*, 359:4711–4727, 2005. [↑12](#)
- [Vas05] W. Vasconcelos. *Integral closure, Rees Algebras, Multiplicities, Algorithms*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2005. [↑6](#)

UNIVERSITÀ DEGLI STUDI DI GENOVA, DIPARTIMENTO DI MATEMATICA, VIA DODECANESO 35,
16146 GENOVA (GE), ITALY

Email address: `bignalet@dima.unige.fr`