

# RATIONAL POINTS OF LATTICE IDEALS ON A TORIC VARIETY AND TORIC CODES

MESUT ŞAHİN

ABSTRACT. We show that the number of rational points of a subgroup inside a toric variety over a finite field defined by a homogeneous lattice ideal can be computed via Smith normal form of the matrix whose columns constitute a basis of the lattice. This generalizes and yields a concise toric geometric proof of the same fact proven purely algebraically by Lopez and Villarreal for the case of a projective space and a standard homogeneous lattice ideal of dimension one. We also prove a Nullstellensatz type theorem over a finite field establishing a one to one correspondence between subgroups of the dense split torus and certain homogeneous lattice ideals. As application, we compute the main parameters of generalized toric codes on subgroups of the torus of Hirzebruch surfaces, generalizing the existing literature.

## 1. INTRODUCTION

Let  $\Sigma$  be a complete simplicial fan with rays generated by the lattice vectors  $\mathbf{v}_1, \dots, \mathbf{v}_r \in \mathbb{R}^n$ . Each cone  $\sigma \in \Sigma$  defines an affine toric variety  $U_\sigma$  over a field  $\mathbb{K}$  whose ring of regular functions is the semigroup ring  $\mathbb{K}[\check{\sigma} \cap \mathbb{Z}^n]$ . Gluing these affine pieces in a standard way, we obtain the split toric variety  $X_\Sigma$  as an abstract variety over the field  $\mathbb{K}$ . Our main motivation comes from the immediate applications to error-correcting codes defined on toric varieties over a finite field  $\mathbb{K} = \mathbb{F}_q$ . In this regard, we assume that the class group  $\text{Cl } X_\Sigma$  is free throughout, which does not harm the generality, as the codes on a singular toric variety survive when the singularity is resolved and smooth complete toric varieties have a torsion-free class group, by [6, Proposition 4.2.5].

The Cox ring  $S = \mathbb{K}[x_1, \dots, x_r]$  of  $X_\Sigma$  is graded naturally by the group  $\text{Cl } X_\Sigma$ , i.e. we have the decomposition  $S = \bigoplus_{\alpha \in \text{Cl } X_\Sigma} S_\alpha$ , where  $S_\alpha$  is the  $\mathbb{K}$ -vector space spanned by the monomials of degree  $\alpha$ . For any  $\alpha$  and any subset  $Y = \{P_1, \dots, P_N\}$  of  $X_\Sigma$ , we have the following *evaluation map* for  $\mathbb{K} = \mathbb{F}_q$ :

$$(1.1) \quad \text{ev}_Y : S_\alpha \rightarrow \mathbb{K}^N, \quad F \mapsto (F(P_1), \dots, F(P_N)).$$

The image  $\mathcal{C}_{\alpha, Y} = \text{ev}_Y(S_\alpha)$  is a linear code, called a *generalized toric code*. The three basic parameters of  $\mathcal{C}_{\alpha, Y}$  are block-length which is  $N$ , the dimension which is  $K = \dim_{\mathbb{F}_q}(\mathcal{C}_{\alpha, Y})$ , and the minimum distance  $\delta = \delta(\mathcal{C}_{\alpha, Y})$  which is the minimum of the numbers of nonzero components of nonzero vectors in  $\mathcal{C}_{\alpha, Y}$ .

When  $Y = T_X$  is the maximal torus of  $X = X_\Sigma$  we get the classical toric code examined for the first time by Hansen in [9] and studied further in e.g. [11, 21, 2, 22, 13, 4] showcasing some champion codes. Although the block-length of  $\mathcal{C}_{\alpha, Y}$  equals

---

2020 *Mathematics Subject Classification*. Primary 14M25; 14G05 ; Secondary 94B27 ; 11T71.  
The author is supported by TÜBİTAK Project No:119F177.

$N = |T_X| = (q - 1)^n$  in this classical case, its computation depends on the way  $Y$  is presented in the general case. For instance, if  $Y = Y_Q$  is parameterised by the monomials whose power is a column of an integer matrix  $Q$ , then some algorithms are given for computing the length  $N = |Y|$ , see [20, 1]. The vanishing ideal  $I(Y)$  of  $Y$  which is generated by homogeneous polynomials vanishing on  $Y$  is relevant in the computation of the other parameters. Namely, the kernel of  $\text{ev}_Y$  is just the subspace  $I_\alpha(Y) := I(Y) \cap S_\alpha$ , and hence the dimension  $K = \dim_{\mathbb{F}_q}(\mathcal{C}_{\alpha,Y})$  is the value  $H_Y(\alpha)$  of the multigraded Hilbert function  $H_Y$  of  $Y$ , see [24]. The minimum distance function of  $I(Y)$  is introduced and is related to the minimum distance of  $\mathcal{C}_{\alpha,Y}$  in [16] for the case of the projective space  $X = \mathbb{P}^n$ . The ideal  $I(Y_Q)$  is a lattice ideal  $I_L$  for a unique lattice  $L \subseteq \mathbb{Z}^r$  of rank  $n$  as shown in [20, 7, 1]. Moreover, every subgroup of  $T_X$  is not only of the form  $Y_Q$  but also is the common zeroes  $V_X(I_L)$  of binomials generating a lattice ideal  $I_L$ , see [23, Section 3].

The present paper aims mainly to develop methods for computing parameters of generalized toric codes  $\mathcal{C}_{\alpha,Y}$  from subgroups  $Y = V_X(I_L)$  of  $T_X$ . Before we state our main results, we need to introduce a little more notation.

Let  $\mathcal{B}$  be an  $r \times n$  matrix whose columns constitute a basis for  $L$ ; let  $\mathcal{A}$  and  $\mathcal{C}$  be unimodular matrices of sizes  $r$  and  $n$ , respectively, so that

$$\mathcal{D} = \mathcal{A}\mathcal{B}\mathcal{C} = [d_1\mathbf{e}_1 | \cdots | d_n\mathbf{e}_n]$$

is the Smith-Normal form of  $\mathcal{B}$ , where  $\mathbf{e}_1, \dots, \mathbf{e}_n \in \mathbb{Z}^r$ . These  $d_i$ 's are known as the *invariant factors* having the property that  $d_i$  divides  $d_{i+1}$  for each  $i \in [n-1] := \{1, \dots, n-1\}$  and that the order of the torsion part  $T(\mathbb{Z}^r/L)$  of the group  $\mathbb{Z}^r/L$  is  $d_1 \cdots d_n$ , by the fundamental structure theorem for finitely generated abelian groups, see [10, pp. 187–188].

We first observe in Proposition 3.3 that the common solutions  $V_X(I_L)(\mathbb{K})$  of polynomials in  $I_L$  lie in the torus  $T_X$ , for any field  $\mathbb{K}$ .

Then, we count characters to see the number of  $\mathbb{K}$ -rational points is finite:  $|V_X(I_L)(\mathbb{K})| \leq d_1 \cdots d_n$ . Furthermore, it meets this upper bound if and only if  $\mathbb{K}$  has all the  $d_i$ -th roots of unity, for every  $i \in [n]$ , see Theorem 3.7.

Let  $Y$  be a subgroup of  $T_X(\mathbb{F}_q)$ . Then,  $I(Y) = I_L$  for a unique lattice  $L$  with  $(q-1)L_\beta \subseteq L \subseteq L_\beta$ , see Lemma 4.2. Moreover, the number of  $\mathbb{F}_q$ -rational points in  $Y$  is given in Theorem 4.4 by

$$|Y| = |V_X(I_L)(\mathbb{F}_q)| = |L_\beta/L| = d_1 \cdots d_n.$$

Therefore, the length of the code  $\mathcal{C}_\alpha(Y)$  will be  $d_1 \cdots d_n$ .

We also prove a Nullstellensatz type result over the finite field  $\mathbb{F}_q$  saying that  $I(V_X(I_L)) = I_L$  establishing a 1–1 correspondence between the subgroups  $V_X(I_L)$  and the lattice ideals  $I_L$ , where  $(q-1)L_\beta \subseteq L \subseteq L_\beta$ , see Theorem 5.2. This result will eliminate the need for an algorithm to compute the vanishing ideal of the subgroup  $V_X(I_L)$ , which will save time in search of codes with good parameters.

If  $I_L$  is a homogeneous ideal of dimension  $d = r - n$ , then its degree is just the constant Hilbert polynomial, which is computed to be  $\deg(I_L) = |L_\beta/L| = d_1 \cdots d_n$ , in Theorem 6.3, generalizing the main result of [14]. This will be nothing but the mixed volume of the polytopes of binomials generating  $I_L$  by Corollary 6.5.

Finally, we uncover the structure and give two descriptions for the points of the subgroup cut out by the lattice ideal  $I_L$ . More precisely, we have

$$V_X(I_L)(\mathbb{K}) = \langle [P_1] \rangle \times \cdots \times \langle [P_n] \rangle,$$

for the point  $[P_i] = [\eta_i^{a_{i1}} : \cdots : \eta_i^{a_{ir}}]$ , where  $\eta_i$  is a primitive  $d_i$ -th root of unity in  $\mathbb{K}$ , for each  $i \in [n]$ . In addition,

$$V_X(I_L)(\mathbb{F}_q) = Y_Q = \{[\mathbf{t}^{\mathbf{q}_1} : \cdots : \mathbf{t}^{\mathbf{q}_r}] \mid \mathbf{t} \in (\mathbb{F}_q^*)^s\}$$

is parameterized by the columns of the matrix  $Q = [\mathbf{q}_1 \mathbf{q}_2 \cdots \mathbf{q}_r]$ , whose  $i$ -th row is  $(q-1)/d_i$  times the  $i$ -th row of the matrix  $\mathcal{A}$ , see Theorem 7.1.

As application, we compute the main parameters of generalized toric codes on subgroups of the torus of Hirzebruch surfaces, generalizing the existing literature, see Theorem 8.4. The parameterization above is especially useful in computing the minimum distance.

## 2. PRELIMINARIES

In this section we fix our notation and recall standard facts from toric geometry relying on the wonderful book [6], although there are other excellent books.

Let  $\Sigma$  be a complete simplicial fan with rays generated by the lattice vectors  $\mathbf{v}_1, \dots, \mathbf{v}_r \in \mathbb{R}^n$ . The fan gives the following short exact sequence:

$$(2.1) \quad \mathfrak{P} : 0 \longrightarrow \mathbb{Z}^n \xrightarrow{\phi} \mathbb{Z}^r \xrightarrow{\beta} \mathbb{Z}^d \longrightarrow 0,$$

where  $\phi$  denotes the matrix  $[\mathbf{v}_1 \cdots \mathbf{v}_r]^T$  and  $d = r - n$  is the rank of  $\text{Cl } X_\Sigma$ . Denote by  $L_\beta$  the key sublattice of  $\mathbb{Z}^r$  which is isomorphic to  $\mathbb{Z}^n$  via  $\phi$ , whose basis is given by the columns  $\mathbf{u}_1, \dots, \mathbf{u}_n$  of  $\phi$ . We are interested in sublattices  $L$  of  $L_\beta$  and the  $L_\beta$ -homogeneous *lattice ideals* they define in the Cox ring  $\mathbb{K}[x_1, \dots, x_r]$  of  $X(\mathbb{K})$ :

$$I_L = \langle \mathbf{x}^{\mathbf{m}^+} - \mathbf{x}^{\mathbf{m}^-} \mid \mathbf{m} = \mathbf{m}^+ - \mathbf{m}^- \in L \rangle.$$

Applying  $\text{Hom}(-, \mathbb{K}^*)$  functor to  $\mathfrak{P}$  we get the following dual short exact sequence since  $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}^d, \mathbb{K}^*) = 1$ :

$$(2.2) \quad \mathfrak{P}^* : 1 \longrightarrow G \xrightarrow{i} (\mathbb{K}^*)^r \xrightarrow{\pi} (\mathbb{K}^*)^n \longrightarrow 1,$$

where  $\pi : P \mapsto (\mathbf{x}^{\mathbf{u}_1}(P), \dots, \mathbf{x}^{\mathbf{u}_n}(P))$  is expressed using the notation

$$\mathbf{x}^{\mathbf{a}}(P) = p_1^{a_1} \cdots p_r^{a_r} \text{ for } P = (p_1, \dots, p_r) \in (\mathbb{K}^*)^r \text{ and for } \mathbf{a} = (a_1, \dots, a_r) \in \mathbb{Z}^r.$$

Notice also that the algebraic group  $G = \ker(\pi)$  is the zero locus in  $(\mathbb{K}^*)^r$  of the toric (prime lattice) ideal  $I_{L_\beta}$ :

$$\begin{aligned} V(I_{L_\beta}) \cap (\mathbb{K}^*)^r &:= \{P \in (\mathbb{K}^*)^r \mid (\mathbf{x}^{\mathbf{m}^+} - \mathbf{x}^{\mathbf{m}^-})(P) = 0 \text{ for all } \mathbf{m} \in L_\beta\} \\ &= \{P \in (\mathbb{K}^*)^r \mid \mathbf{x}^{\mathbf{m}}(P) = 1 \text{ for all } \mathbf{m} \in L_\beta\}. \end{aligned}$$

As proved by Cox in [5], we see that the toric variety  $X(\mathbb{K})$  is the geometric quotient  $[\mathbb{K}^r \setminus V(B)]/G$ , where  $B$  is the monomial ideal in  $\mathbb{K}[x_1, \dots, x_r]$  generated by the monomials  $\mathbf{x}^{\hat{\sigma}} = \prod_{\rho_i \notin \sigma} x_i$  corresponding to cones  $\sigma \in \Sigma$ . Hence, points of  $X(\mathbb{K})$  are orbits  $[P] := G \cdot P$ , for  $P \in \mathbb{K}^r \setminus V(B)$  and that  $T_X(\mathbb{K}) = (\mathbb{K}^*)^r/G$ .

As the elements of the algebraic torus  $(\mathbb{K}^*)^r$  in  $\mathfrak{P}^*$  is already identified with the characters  $\rho : \mathbb{Z}^r \rightarrow \mathbb{K}^*$ , we take the approach to investigate characters in order to compute the number of  $\mathbb{K}$ -rational points of the lattice ideal  $I_L$  on the toric variety  $X(\mathbb{K})$ .

## 3. SUBGROUPS DEFINED BY LATTICE IDEALS OVER ANY FIELD

In this section, we study subgroups of the torus  $T_X$  defined by lattice ideals inside  $\mathbb{K}[x_1, \dots, x_r]$ , which is an  $L_\beta$ -graded ring via  $\deg(x_i) = \beta_i$ , where  $\beta_i$  is the  $i$ -th column of the matrix  $\beta$  in (2.1).

First of all,  $\text{rank } L_\beta = n$  since  $L_\beta$  is isomorphic to  $\mathbb{Z}^n$  by the sequence  $\mathfrak{P}$  in (2.1). On the other hand, it is well-known that  $\dim J = r - \text{height}(J)$ , where  $\dim J$  is the Krull dimension of the quotient ring  $\mathbb{K}[x_1, \dots, x_r]/J$ . So, we have

$$\text{rank } L = n = \text{height}(I_L) \iff \dim I_L = d = r - n.$$

As the vanishing ideals of subgroups of the torus  $T_X$  are  $L_\beta$ -homogeneous lattice ideals of Krull-dimension  $d$  by [23, Proposition 2.6], and  $I_L$  is  $L_\beta$ -homogeneous if and only if  $L \subseteq L_\beta$  by [23, Proposition 2.3], we focus on lattices  $L$  satisfying  $L \subseteq L_\beta$  and  $\text{rank } L = n$ .

**Lemma 3.1.** *Let  $L \subseteq L_\beta$  and  $\text{rank } L = n$ . Then,  $\ell L_\beta \subseteq L$  for the order  $\ell = |L_\beta/L|$ .*

*Proof.* Since  $L_\beta/L$  is a finite group of order  $\ell$ , we have  $\ell(\mathbf{m} + L) = L$ , for every  $\mathbf{m} \in L_\beta$ . This means that  $\ell\mathbf{m} \in L$  for all  $\mathbf{m} \in L_\beta$ .  $\square$

**Lemma 3.2.**  *$L \subseteq L'$  if and only if  $I_L \subseteq I_{L'}$ .*

*Proof.* ( $\Rightarrow$ ): If  $L \subseteq L'$ , then every  $\mathbf{m} \in L$  lies in  $L'$ . Therefore, every binomial generator  $\mathbf{x}^{\mathbf{m}^+} - \mathbf{x}^{\mathbf{m}^-}$  of  $I_L$  lies in  $I_{L'}$ . So,  $I_L \subseteq I_{L'}$ .

( $\Leftarrow$ ): Assume that  $I_L \subseteq I_{L'}$  and  $\mathbf{m} \in L$ . Then, the binomial  $\mathbf{x}^{\mathbf{m}^+} - \mathbf{x}^{\mathbf{m}^-}$  of  $I_L$  lies in  $I_{L'}$ . It follows from [14, Lemma 3.2] that  $\mathbf{m} \in L'$ .  $\square$

**Proposition 3.3.** *Let  $L \subseteq L_\beta$  and  $\text{rank } L = n$ . Then,  $V_X(I_L)$  is a subgroup of  $T_X$  for  $X = X(\mathbb{K})$ .*

*Proof.* By Lemma 3.1, we have  $\ell L_\beta \subseteq L \subseteq L_\beta$  for the order  $\ell = |L_\beta/L|$ . As smaller lattices correspond to smaller lattice ideals by Lemma 3.2, it follows that  $I_{\ell L_\beta} \subseteq I_L \subseteq I_{L_\beta}$ . Thus, we have the inclusions below:

$$\{[1]\} = V_X(I_{L_\beta}) \subseteq V_X(I_L) \subseteq V_X(I_{\ell L_\beta}).$$

So as to prove that  $V_X(I_L)$  is a subset of the torus  $T_X$ , it suffices to reveal that  $V_X(I_{\ell L_\beta}) \subseteq T_X$ . Given any point  $[P] \in V_X(I_{\ell L_\beta})$ , we have  $P \in V(I_{\ell L_\beta}) \setminus V(B)$ . By definition,  $P \in V(I_{\ell L_\beta})$  exactly when  $\mathbf{x}^{\ell\mathbf{m}^+}(P) = \mathbf{x}^{\ell\mathbf{m}^-}(P)$  for all  $\mathbf{m} = \mathbf{m}^+ - \mathbf{m}^- \in L_\beta$ . Since  $\mathbf{x}^{\ell\mathbf{m}^+}(P) = \mathbf{x}^{\ell\mathbf{m}^-}(P)$  is equivalent to  $\mathbf{x}^{\mathbf{m}^+}(P^\ell) = \mathbf{x}^{\mathbf{m}^-}(P^\ell)$ , it follows that  $P \in V(I_{\ell L_\beta})$  if and only if  $P^\ell \in V(I_{L_\beta})$ . As  $B$  is a monomial ideal,  $P \notin V(B)$  if and only if  $P^\ell \notin V(B)$ . Thus,  $P \in V(I_{\ell L_\beta}) \setminus V(B)$  implies that  $P^\ell \in V(I_{L_\beta}) \setminus V(B) = G$ . As  $G$  lies in the torus  $(\mathbb{K}^*)^r$ , we have  $P \in (\mathbb{K}^*)^r$  yielding  $[P] \in T_X$ . Hence,  $V_X(I_L) \subseteq T_X$ .

If  $[P], [Q] \in V_X(I_L)$ , then we have the following:

$$\mathbf{x}^{\mathbf{m}^+}(P \cdot Q^{-1}) = \mathbf{x}^{\mathbf{m}^+}(P)\mathbf{x}^{\mathbf{m}^+}(Q)^{-1} = \mathbf{x}^{\mathbf{m}^-}(P)\mathbf{x}^{\mathbf{m}^-}(Q)^{-1} = \mathbf{x}^{\mathbf{m}^-}(P \cdot Q^{-1}).$$

Thus,  $[P] \cdot [Q]^{-1}$  lies in  $V_X(I_L)$  demonstrating that it is indeed a subgroup.  $\square$

**Remark 3.4.** *It is easy to see that  $V_X(I_L) \cap T_X$  is a subgroup of  $T_X$ , when we only have  $L \subseteq L_\beta$ , see also [23, Corollary 2.4]. But, if  $\text{rank } L < n$ , then  $V_X(I_L)$  need not lie inside the torus  $T_X$ , see [23, Example 3.6].*

For any  $\mathbf{m} = \mathbf{m}^+ - \mathbf{m}^- \in L$ , the binomial  $\mathbf{x}^{\mathbf{m}^+} - \mathbf{x}^{\mathbf{m}^-} \in I_L$ , hence we have  $\mathbf{x}^{\mathbf{m}^+}(P) = \mathbf{x}^{\mathbf{m}^-}(P)$  for a point  $[P] \in V_X(I_L)$ . If we further assume that  $L \subseteq L_\beta$  and  $\text{rank } L = n$ , then  $\mathbf{x}^{\mathbf{m}}(P) = 1$  as  $V_X(I_L) \subseteq T_X$ . Thus, a point  $[P] \in V_X(I_L)$ , defines a character  $\chi_P : L_\beta/L \rightarrow \mathbb{K}^*$  via  $\chi_P(\mathbf{m} + L) := \mathbf{x}^{\mathbf{m}}(P) = p_1^{m_1} \cdots p_r^{m_r}$ , for all  $\mathbf{m} = (m_1, \dots, m_r) \in L$  under these circumstances.

**Lemma 3.5.** *Let  $L \subseteq L_\beta$  and  $\text{rank } L = n$ . If  $X = X(\mathbb{K})$ , then the groups  $V_X(I_L)$  and  $\text{Hom}(L_\beta/L, \mathbb{K}^*)$  are isomorphic.*

*Proof.* Let  $\psi : V_X(I_L) \rightarrow \text{Hom}(L_\beta/L, \mathbb{K}^*)$  be defined by  $[P] \rightarrow \chi_P$ .

- The map  $\psi$  is well-defined, for if  $g \in G = V(I_{L_\beta}) \cap (\mathbb{K}^*)^r$ , then  $\chi_{gP} = \chi_P$ , since for all  $\mathbf{m} \in L_\beta$ , we have

$$\mathbf{x}^{\mathbf{m}}(gP) = g^{\mathbf{m}} \mathbf{x}^{\mathbf{m}}(P) = \mathbf{x}^{\mathbf{m}}(P).$$

- $\psi$  is a homomorphism, since  $\chi_{P \cdot P'} = \chi_P \cdot \chi_{P'}$ .
- It is injective, as  $\chi_P(\mathbf{m} + L) = \mathbf{x}^{\mathbf{m}}(P) = 1$ , for all  $\mathbf{m} \in L_\beta$ , implies that  $P \in G$  that is  $[P] = [1]$ .
- It is surjective, as we prove now. Let  $\{\mathbf{u}_1, \dots, \mathbf{u}_n\}$  be the  $\mathbb{Z}$ -basis for the lattice  $L_\beta$ . For a given  $\chi : L_\beta/L \rightarrow \mathbb{K}^*$ , let  $t_i = \chi(\mathbf{u}_i + L)$  for each  $i \in [n]$ . Since  $\mathfrak{P}^*$  is exact, there is a point  $P \in (\mathbb{K}^*)^r$  such that  $\pi(P) = (\mathbf{x}^{\mathbf{u}_1}(P), \dots, \mathbf{x}^{\mathbf{u}_n}(P)) = (t_1, \dots, t_n)$ . Every element  $\mathbf{m} \in L_\beta$  is written as  $\mathbf{m} = c_1 \mathbf{u}_1 + \dots + c_n \mathbf{u}_n$  for some  $(c_1, \dots, c_n) \in \mathbb{Z}^n$ . Thus, we have

$$\chi(\mathbf{m} + L) = t_1^{c_1} \cdots t_n^{c_n} = (\mathbf{x}^{\mathbf{u}_1}(P))^{c_1} \cdots (\mathbf{x}^{\mathbf{u}_n}(P))^{c_n} = \mathbf{x}^{\mathbf{m}}(P) = \chi_P(\mathbf{m} + L).$$

If  $\mathbf{m} \in L$ , then it follows that  $\mathbf{x}^{\mathbf{m}}(P) = \chi(\mathbf{m} + L) = 1$  for all  $\mathbf{m} \in L$  and hence  $[P] \in V_X(I_L)$ . Hence, we have  $\chi = \chi_P$ .

These complete the proof.  $\square$

**Lemma 3.6.** *Let  $L \subseteq L_\beta$  and  $\text{rank } L = n$ . If  $d_i$  is an invariant factor of the group  $L_\beta/L$  and  $C_i := \{c \in \mathbb{K}^* : c^{d_i} = 1\}$  is the group of  $d_i$ -th roots of unity lying inside  $\mathbb{K}$ , for each  $i \in [n]$ , then the groups  $\text{Hom}(L_\beta/L, \mathbb{K}^*)$  and  $C_1 \times \cdots \times C_n$  are isomorphic.*

*Proof.* Let  $\{\mathbf{m}_1, \dots, \mathbf{m}_n\}$  be a  $\mathbb{Z}$ -basis for the lattice  $L_\beta$  such that  $\{d_1 \mathbf{m}_1, \dots, d_n \mathbf{m}_n\}$  is a  $\mathbb{Z}$ -basis of  $L$ .

We regard elements in  $\text{Hom}(L_\beta/L, \mathbb{K}^*)$  as (partial) characters  $\rho : L_\beta \rightarrow \mathbb{K}^*$  whose restriction to  $L$  is the identity  $1_L$ . So, we have  $\rho(\mathbf{m}_i)^{d_i} = \rho(d_i \mathbf{m}_i) = 1$ , for each  $i \in [n]$ . Since  $\rho$  is determined by the values  $\rho(\mathbf{m}_1), \dots, \rho(\mathbf{m}_n)$ , the map  $\Theta : \text{Hom}(L_\beta/L, \mathbb{K}^*) \rightarrow C_1 \times \cdots \times C_n$ , defined by  $\Theta(\rho) = (\rho(\mathbf{m}_1), \dots, \rho(\mathbf{m}_n))$  is well-defined.

- $\Theta$  is a homomorphism as  $\Theta(\rho \cdot \rho') = \Theta(\rho) \cdot \Theta(\rho')$ .
- It is injective, since  $\Theta(\rho) = (\rho(\mathbf{m}_1), \dots, \rho(\mathbf{m}_n)) = (1, \dots, 1) \implies \rho = 1_{L_\beta}$ .
- $\Theta$  is surjective as we demonstrate below:

Every choice of  $(c_1, \dots, c_n) \in C_1 \times \cdots \times C_n$  determines an element

$$\rho : L_\beta \rightarrow \mathbb{K}^* \quad \text{via} \quad \rho(k_1 \mathbf{m}_1 + \cdots + k_n \mathbf{m}_n) = c_1^{k_1} \cdots c_n^{k_n}.$$

Furthermore, if  $\mathbf{m} = k_1 \mathbf{m}_1 + \cdots + k_n \mathbf{m}_n \in L$  then  $d_i$  divides  $k_i$  and thus  $\rho(\mathbf{m}) = 1$ , as  $c_1^{k_1} = \cdots = c_n^{k_n} = 1$ . Hence,  $\rho \in \text{Hom}(L_\beta/L, \mathbb{K}^*)$ .

Thus,  $\Theta$  is the required isomorphism, completing the proof.  $\square$

**Theorem 3.7.** *Let  $X = X(\mathbb{K})$ ,  $L \subseteq L_\beta$  and  $\text{rank } L = n$ . If  $d_i$  is an invariant factor of the group  $L_\beta/L$  and  $C_i := \{c \in \mathbb{K}^* : c^{d_i} = 1\}$  is the group of  $d_i$ -th roots of unity lying inside  $\mathbb{K}$ , for each  $i \in [n]$ , then  $V_X(I_L) \cong C_1 \times \cdots \times C_n$ , and thus the number of  $\mathbb{K}$ -rational points  $|V_X(I_L)|$  is  $|C_1| \cdots |C_n| \leq d_1 \cdots d_n$ . Moreover,  $|V_X(I_L)| = d_1 \cdots d_n \iff \mathbb{K}$  has all the  $d_i$ -th roots of unity, for every  $i \in [n]$ .*

*Proof.* It follows from Lemma 3.5 that  $V_X(I_L) \cong \text{Hom}(L_\beta/L, \mathbb{K}^*)$ . We also have  $\text{Hom}(L_\beta/L, \mathbb{K}^*) \cong C_1 \times \cdots \times C_n$  by Lemma 3.6, whence  $V_X(I_L) \cong C_1 \times \cdots \times C_n$ . Since  $|C_i| \leq d_i$  in general, we have  $|V_X(I_L)| = |C_1| \cdots |C_n| \leq d_1 \cdots d_n$ .

The second part follows, since  $|C_i| = d_i \iff \mathbb{K}$  has all the  $d_i$ -th roots of unity, for every  $i \in [n]$ .  $\square$

#### 4. SUBGROUPS DEFINED BY LATTICE IDEALS OVER A FINITE FIELD

In this section, we compute the number of  $\mathbb{F}_q$ -rational points of a subgroup of the algebraic group  $T_X$ . By the virtue of Theorem 3.7, this amounts to computing the order of the cyclic subgroup  $C_i$  of  $d_i$ -th roots of unity lying inside the cyclic group  $\mathbb{F}_q^*$  of order  $q - 1$ .

**Theorem 4.1.** *Let  $X = X(\mathbb{F}_q)$ . If  $L \subseteq L_\beta$  and  $\text{rank } L = n$ , then the number of  $\mathbb{F}_q$ -rational points is given by  $|V_X(I_L)| = (d_1, q - 1) \cdots (d_n, q - 1)$ , where  $(d_i, q - 1)$  is the greatest common divisor of  $q - 1$  with an invariant factor  $d_i$ .*

*Proof.* Fix  $i \in [n]$ . Recall that  $C_i = \{c \in \mathbb{K}^* : c^{d_i} = 1\}$  is a cyclic subgroup of  $\mathbb{F}_q^*$  of order  $|C_i|$ , and so  $|C_i|$  divides  $q - 1$ . The order  $|C_i|$  of a generator of  $C_i$  divides  $d_i$ , as well. Hence,  $|C_i|$  divides  $(d_i, q - 1)$ . On the other hand, as  $(d_i, q - 1)$  divides  $q - 1$ , there is a cyclic subgroup of  $\mathbb{F}_q^*$  of order  $(d_i, q - 1)$ , which is clearly contained in  $C_i$ , since  $(d_i, q - 1)$  divides  $d_i$ . Therefore,  $|C_i| = (d_i, q - 1)$ .

The proof now follows directly from Theorem 3.7.  $\square$

Before going further, let us single out lattices corresponding to subgroups of  $T_X$ .

**Lemma 4.2.** *Let  $X = X(\mathbb{F}_q)$ . If  $Y$  is a subgroup of  $T_X$  then  $I(Y) = I_L$  for a unique lattice  $L$  with  $(q - 1)L_\beta \subseteq L \subseteq L_\beta$ . Therefore, subgroups of  $T_X$  are of the form  $V_X(I_L)$  for lattices with  $(q - 1)L_\beta \subseteq L \subseteq L_\beta$ .*

*Proof.* By [23, Theorem 2.9]  $I(Y) = I_L$  for a unique lattice  $L$  with  $L \subseteq L_\beta$ . Since  $Y \subseteq T_X$  implies  $I(T_X) \subseteq I(Y)$  and we have  $I(T_X) = I_{(q-1)L_\beta}$  by [23, Corollary 4.14 (ii)], the first claim follows from Lemma 3.2.

If  $Y$  is a subgroup of  $T_X$ , then  $Y = V_X(I(Y))$  by [23, Lemma 2.8] which is nothing but  $V_X(I_L)$  by above for a unique lattice  $L$  with  $(q - 1)L_\beta \subseteq L \subseteq L_\beta$ .  $\square$

Therefore, in order to compute the order of a subgroup of the torus  $T_X$  of  $X = X(\mathbb{F}_q)$ , it is enough to focus on lattices with  $(q - 1)L_\beta \subseteq L \subseteq L_\beta$ .

**Remark 4.3.** *Let  $L \subseteq L_\beta$  and  $\text{rank } L = n$ . Then,  $(q - 1)L_\beta \subseteq L$  if and only if  $\exp(L_\beta/L) = d_n$  divides  $q - 1$ . Recall that the exponent  $\exp(H)$  of a finite group  $H$  is the least common multiple of the orders of all elements in  $H$ .*

*Proof.* Recall that  $L_\beta$  has a basis  $\{\mathbf{m}_1, \dots, \mathbf{m}_n\}$  such that  $\{d_1\mathbf{m}_1, \dots, d_n\mathbf{m}_n\}$  is a basis of  $L$ , where  $d_i$ 's are the invariant factors of  $L_\beta/L$ . By the structure theorem for finite abelian groups  $L_\beta/L$  and  $\mathbb{Z}_{d_1} \oplus \cdots \oplus \mathbb{Z}_{d_n}$  are isomorphic. Since  $d_i$  divides  $d_{i+1}$  for each  $i \in [n - 1]$ , the exponent of  $L_\beta/L$  is  $d_n$ .

Therefore, if  $(q-1)L_\beta \subseteq L$ , then  $(q-1)\mathbf{m}_i \in L$ . Since  $\{d_1\mathbf{m}_1, \dots, d_n\mathbf{m}_n\}$  is a basis of  $L$ , it follows immediately that  $d_i$  divides  $q-1$ , for every  $i \in [n]$ . Thus,  $\exp(L_\beta/L)$  divides  $q-1$ . Conversely, if the exponent divides  $q-1$ , then each  $d_i$  divides  $q-1$ , and hence  $(q-1)\mathbf{m}_i \in L$  yielding to the inclusion  $(q-1)L_\beta \subseteq L$ .  $\square$

We are now ready to prove the main result of the section yielding to an efficient method to count the number of  $\mathbb{F}_q$ -rational points of the subgroup  $V_X(I_L)$  of  $T_X$  using a basis of  $L$ .

**Theorem 4.4.** *Let  $X = X(\mathbb{F}_q)$ . If  $(q-1)L_\beta \subseteq L \subseteq L_\beta$ , then the number of  $\mathbb{F}_q$ -rational points is given by  $|V_X(I_L)| = |L_\beta/L| = d_1 \cdots d_n$ .*

*Proof.* By Remark 4.3, it follows that  $d_i$  divides  $q-1$ , and hence  $(d_i, q-1) = d_i$ , for every  $i \in [n]$ . Therefore,  $|V_X(I_L)| = d_1 \cdots d_n$ , by Theorem 4.1.  $\square$

**Remark 4.5.** *Theorem 4.4 follows from Lemma 3.5 together with a more general fact which is communicated in private to us by Laurence Barker: given a finite group  $A$  and a finite field  $\mathbb{F}_q$ ,  $\text{Hom}(A, \mathbb{F}_q^*)$  is isomorphic to  $A$  if and only if  $\exp(A)$  divides  $q-1$ .*

*Proof.* The usual dual of  $A$ , sometimes called the Pontryagin dual, is defined to be  $A^* = \text{Hom}(A, \mathbb{Q}/\mathbb{Z})$ . The group  $\mathbb{Q}/\mathbb{Z}$ , is isomorphic to the torsion subgroup of  $\mathbb{C}^*$ . So  $A^*$  can be identified with the group of irreducible complex characters of  $A$ .

The standard result that  $A$  is non-canonically isomorphic to  $A^*$  is plain when  $A$  is cyclic. It is easy to see that  $(A_1 \times A_2)^* \cong A_1^* \times A_2^*$ . So the general case follows by decomposing  $A$  as a direct product of cyclic groups.

Viewing finite cyclic groups as embedded in each other wherever possible, then  $\mathbb{Q}/\mathbb{Z}$  is the union of all those groups, in other words, the colimit of the diagram of embeddings. In particular,  $A^*$  can be identified with  $\text{Hom}(A, C)$  for any cyclic group  $C$  that is sufficiently large in the sense where  $C' \leq C$  provided  $|C'|$  divides  $|C|$ . The exact criterion for  $C$  to be sufficiently large is that every homomorphism  $A \rightarrow \mathbb{Q}/\mathbb{Z}$  has image contained in the subgroup isomorphic to  $C$ . That is equivalent to saying that the exponent of  $A$  divides  $|C|$ . Taking  $C = \mathbb{F}_q^*$  completes the proof.  $\square$

**Corollary 4.6.** *Let  $X = X(\mathbb{F}_q)$ . If  $(q-1)L_\beta \subseteq L \subseteq L_\beta$ , then  $|V_X(I_L)| = |\det \mathbf{ML}|$ , where  $\mathbf{ML}$  is the matrix with entries  $b_{ij} \in \mathbb{Z}$  such that the  $j$ -th basis element of  $L$  is written as  $b_{1j}\mathbf{u}_1 + \cdots + b_{nj}\mathbf{u}_n$  in terms of the basis  $\{\mathbf{u}_1, \dots, \mathbf{u}_n\}$  of  $L_\beta$ .*

*Proof.* By Theorem 4.4, we have  $|V_X(I_L)| = |L_\beta/L|$ . Recall that the map  $\phi$  of (2.1) is an isomorphism between  $\mathbb{Z}^n$  and  $L_\beta$ . If  $\Lambda$  is the sublattice of  $\mathbb{Z}^n$  which is isomorphic to  $L$  under  $\phi$ , then the order  $|L_\beta/L| = |\mathbb{Z}^n/\Lambda|$ . Let  $\mathbf{ML} := \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$  be a subset of  $\mathbb{Z}^n$  constituting a  $\mathbb{Z}$ -basis for the lattice  $\Lambda$ . By abusing the notation, let us denote by  $\mathbf{ML}$  the matrix whose columns are the elements of the set  $\mathbf{ML}$ .

It is known that the volume of the fundamental domain  $\Pi(\mathbf{ML})$  of  $\mathbf{ML}$  is  $|\det \mathbf{ML}|$ , where

$$\Pi(\mathbf{ML}) := \left\{ \sum_{i=1}^n \lambda_i \mathbf{b}_i : 0 \leq \lambda_i < 1 \right\}.$$

It is also known that  $|\det \mathbf{ML}|$  is the number  $|\Pi(\mathbf{ML}) \cap \mathbb{Z}^n|$  of lattice points inside  $\Pi(\mathbf{ML})$  as well as the index  $[\mathbb{Z}^n : \Lambda] = |\mathbb{Z}^n/\Lambda|$ .

Altogether, we have  $|V_X(I_L)| = |L_\beta/L| = |\mathbb{Z}^n/\Lambda| = |\det \mathbf{ML}|$ .  $\square$

**Corollary 4.7.** *If  $\{c_1\mathbf{u}_1, \dots, c_n\mathbf{u}_n\}$  constitute a basis for  $L$ , for positive integers  $c_i$  dividing  $q-1$ , then  $|V_X(I_L)| = c_1 \cdots c_n$ .*

*Proof.* Since  $\{\mathbf{u}_1, \dots, \mathbf{u}_n\} \subset \mathbb{Z}^r$  form a basis of  $L_\beta$  and  $\phi(\mathbf{e}_i) = \mathbf{u}_i$ , we observe that  $\phi(c_i\mathbf{e}_i) = c_i\mathbf{u}_i$ , for all  $i \in [n]$ . So,  $\mathbf{ML} = \{c_1\mathbf{e}_1, \dots, c_n\mathbf{e}_n\}$  is a basis for the lattice  $\Lambda$  that appeared in the proof of Corollary 4.6 and  $|\det(\mathbf{ML})| = c_1 \cdots c_n$ . Therefore, it follows that  $\ell = |L_\beta/L| = c_1 \cdots c_n$  and that  $V_X(L)$  has  $c_1 \cdots c_n$  elements by Corollary 4.6.  $\square$

If  $(q-1)L_\beta \subseteq L \subseteq L_\beta$ , then obviously  $\ell = |L_\beta/L|$  is coprime to  $p = \text{char}(\mathbb{F}_q)$ . On the other hand, whenever  $\ell = |L_\beta/L|$  is not divisible by a prime  $p$ , there are finite fields of characteristic  $p$  verifying this condition as we demonstrate next.

**Proposition 4.8.** *Let  $L \subseteq L_\beta$  and  $\text{rank } L = n$ . If  $\ell = |L_\beta/L|$  is not divisible by a prime  $p$ , then there is a finite field  $\mathbb{F}_q$  of characteristic  $p$  such that  $\exp(L_\beta/L)$  divides  $q-1$ . Thus, we have  $(q-1)L_\beta \subseteq L \subseteq L_\beta$  in this case.*

*Proof.* Let  $\mathbb{F}_q$  be the splitting field of  $x^\ell - 1$  over  $\mathbb{F}_p$ . As  $p$  does not divide  $\ell$ ,  $x^\ell - 1$  and its derivative  $\ell x^{\ell-1}$  have no roots in common. So, roots of  $x^\ell - 1$  are all distinct. It is easy to see that they form a cyclic subgroup of  $\mathbb{F}_q^*$  of order  $\ell$ . Thus,  $\ell$  divides  $q-1$  whence so does  $\exp(L_\beta/L)$ .

The second part follows from Remark 4.3.  $\square$

We close the section with an interesting example illustrating the results.

**Example 4.9.** *Consider the Hirzebruch surface  $X = \mathcal{H}_2$  whose fan in  $\mathbb{R}^2$  have rays generated by*

$$\mathbf{v}_1 = (1, 0), \mathbf{v}_2 = (0, 1), \mathbf{v}_3 = (-1, 2), \text{ and } \mathbf{v}_4 = (0, -1).$$




FIGURE 1. The fan for the Hirzebruch surface.

The exact sequence in (2.1) becomes:

$$\mathfrak{P} : 0 \longrightarrow \mathbb{Z}^2 \xrightarrow{\phi} \mathbb{Z}^4 \xrightarrow{\beta} \mathbb{Z}^2 \longrightarrow 0,$$

where

$$\phi = \begin{bmatrix} 1 & 0 & -1 & 0 \\ 0 & 1 & 2 & -1 \end{bmatrix}^T \quad \text{and} \quad \beta = \begin{bmatrix} 1 & -2 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}.$$



Thus, the class group is  $\mathcal{A} = \mathbb{Z}^2$  grading the Cox ring  $S = \mathbb{K}[x_1, x_2, x_3, x_4]$  of  $X$  via

$$\deg_{\mathcal{A}}(x_1) = \deg_{\mathcal{A}}(x_3) = (1, 0), \deg_{\mathcal{A}}(x_2) = (-2, 1), \deg_{\mathcal{A}}(x_4) = (0, 1).$$

Clearly, a  $\mathbb{Z}$ -basis of  $L_{\beta}$  is given by  $\mathbf{u}_1 = (1, 0, -1, 0)$  and  $\mathbf{u}_2 = (0, 1, 2, -1)$ .

Consider the lattice  $L$  spanned over  $\mathbb{Z}$  by  $3\mathbf{u}_1$  and  $5\mathbf{u}_2$ . Since  $B = \{3\mathbf{e}_1, 5\mathbf{e}_2\}$  is a basis for the lattice  $\Lambda$  that appeared in the proof of Corollary 4.6 we have  $\ell = |L_{\beta}/L| = \det B = 15$  and so  $15L_{\beta} \subset L \subset L_{\beta}$  by Lemma 3.1. Hence, if 15 divides  $q - 1$ , which is the case for instance when  $q = 16$  or  $q = 31$ , then the subgroup  $V_X(I_L)$  has 15 members of  $T_X$  over the finite field  $\mathbb{F}_q$ .

## 5. FINITE NULLSTELLENSATZ FOR LATTICE IDEALS

In this section, we prove a Nullstellensatz type result over  $\mathbb{F}_q$  establishing a one to one correspondence between the subgroups  $V_{X(\mathbb{F}_q)}(I_L)$  and lattice ideals  $I_L$ , where  $(q - 1)L_{\beta} \subseteq L \subseteq L_{\beta}$ .

**Lemma 5.1.** *If  $(q - 1)L_{\beta} \subseteq L \subseteq L_{\beta}$ , then the subgroups  $V_{X(\mathbb{K})}(I_L)$  and  $V_{X(\mathbb{F}_q)}(I_L)$  are the same, for any field  $\mathbb{K}$  which is an extension of  $\mathbb{F}_q$ .*

*Proof.* Since  $\mathbb{F}_q^* \subset \mathbb{K}^*$ , we have the inclusion  $V_{X(\mathbb{F}_q)}(I_L) \subseteq V_{X(\mathbb{K})}(I_L)$ . By Theorem 4.4,  $|V_{X(\mathbb{F}_q)}(I_L)| = d_1 \cdots d_n$ , and so  $\mathbb{F}_q^*$  has all the  $d_i$ -th roots of unity. Thus,  $\mathbb{K}^*$  has all the  $d_i$ -th roots of unity and hence  $|V_{X(\mathbb{K})}(I_L)| = d_1 \cdots d_n$  as well, by Theorem 3.7. Having the same orders,  $V_{X(\mathbb{K})}(I_L) = V_{X(\mathbb{F}_q)}(I_L)$  for any field  $\mathbb{K}$  which is an extension of  $\mathbb{F}_q$ .  $\square$

It is now time to demonstrate that the necessary condition suggested by the Lemma 4.2 for a lattice to correspond to a vanishing ideal of a subgroup is indeed sufficient.

**Theorem 5.2.** *If  $X$  is the toric variety  $X(\mathbb{F}_q)$  over  $\mathbb{F}_q$  and  $(q - 1)L_{\beta} \subseteq L \subseteq L_{\beta}$ , then  $I(V_X(I_L)) = I_L$ .*

*Proof.* By [23, Theorem 5.1(iii)], we have the following primary decomposition for the vanishing ideal of  $V_X(I_L)$  in  $S = \mathbb{F}_q[x_1, \dots, x_r]$ :

$$I(V_X(I_L)) = \bigcap_{[P] \in V_X(I_L)} I([P]).$$

Suppose now that  $\mathfrak{p} \subset S$  is a minimal prime of the radical ideal  $I_L \subseteq I(V_X(I_L))$ . Then, as  $\mathfrak{p} \supseteq I_L$ , we have  $V_{X(\overline{\mathbb{F}}_q)}(\mathfrak{p}) \subseteq V_{X(\overline{\mathbb{F}}_q)}(I_L)$ . As  $V_{X(\overline{\mathbb{F}}_q)}(\mathfrak{p})$  cannot be empty over the algebraically closed field  $\overline{\mathbb{F}}_q$ , it has a point  $[P] \in V_{X(\overline{\mathbb{F}}_q)}(I_L) = V_X(I_L)$ , by the virtue of Lemma 5.1. Therefore,  $[P] \in V_X(\mathfrak{p})$  so that  $\mathfrak{p} \subseteq I(V_X(\mathfrak{p})) \subseteq I([P]) \subset S$ . So,  $\text{height}(\mathfrak{p}) \leq \text{height}(I([P])) = n$ . Since we also have  $\mathfrak{p} \supseteq I_L$ , it follows that  $\text{height}(\mathfrak{p}) \geq \text{height}(I_L) = n$ . Thus,  $\text{height}(\mathfrak{p}) = \text{height}(I([P])) = n$  and these primes must coincide:  $\mathfrak{p} = I([P])$ . Therefore, the two ideals  $I_L \subseteq I(V_X(I_L))$  having the same minimal primary decomposition must coincide:  $I_L = I(V_X(I_L))$ .  $\square$

Recall that subgroups of  $T_X$  are of the form  $Y_Q = \{[\mathbf{t}^{\mathbf{q}_1} : \cdots : \mathbf{t}^{\mathbf{q}_r}] \mid \mathbf{t} \in (\mathbb{F}_q^*)^s\}$  for a matrix  $Q = [\mathbf{q}_1 \mathbf{q}_2 \cdots \mathbf{q}_r] \in M_{s \times r}(\mathbb{Z})$  by [23, Theorem 3.2 and Corollary 3.7]. In [19], a handier description of the lattice of the ideal  $I(Y_Q)$  is given, in terms of  $Q$  and  $\beta$ , under a condition on the lattice  $\mathcal{L} = QL_{\beta} = \{Q\mathbf{m} \mid \mathbf{m} \in L_{\beta}\}$ . Before stating this result, let us remind that  $\mathcal{L} : (q - 1) = \{\mathbf{m} \in \mathbb{Z}^s \mid (q - 1)\mathbf{m} \in \mathcal{L}\}$ .

**Theorem 5.3** ([19]). *Let  $L = (L_Q \cap L_\beta) + (q - 1)L_\beta$ . Then  $I_L \subseteq I(Y_Q)$ . The equality holds if and only if  $\mathcal{L} = \mathcal{L} : (q - 1)$ .*

If  $L_Q \subseteq L_\beta$ , then  $I_{L_Q}$  is an  $L_\beta$ -homogeneous toric ideal whose zero locus inside the torus is denoted  $V_Q := V_X(I_{L_Q}) \cap T_X$ . The following consequence of Theorem 5.3 ensures that  $V_Q$  coincides with the subgroup  $Y_Q$  if the condition  $\mathcal{L} = \mathcal{L} : (q - 1)$  holds, following [20, Proposition 4.3] and [20, Corollary 4.4].

**Theorem 5.4** ([19]). *If  $L_Q \subseteq L_\beta$  and  $L = L_Q + (q - 1)L_\beta$ , then we have the following*

- (1)  $V_Q = V_X(I_L)$ ,
- (2)  $V_Q = Y_Q$ , if the condition  $\mathcal{L} = \mathcal{L} : (q - 1)$  holds,
- (3)  $I(V_X(I_L)) = I_L$ , if the condition  $\mathcal{L} = \mathcal{L} : (q - 1)$  holds.

Using Theorem 5.2, we are now able to prove that  $V_Q$  coincides with the subgroup  $Y_Q$  if and only if the condition  $\mathcal{L} = \mathcal{L} : (q - 1)$  holds.

**Corollary 5.5.** *If  $L = (L_Q \cap L_\beta) + (q - 1)L_\beta$ , then  $Y_Q = V_X(I_L)$  if and only if  $\mathcal{L} : (q - 1) = \mathcal{L}$ .*

*Proof.* If  $\mathcal{L} : (q - 1) = \mathcal{L}$  then by Theorem 5.3 we have  $I(Y_Q) = I_L$  implying that  $Y_Q = V_X(I(Y_Q)) = V_X(I_L)$ . For the converse it suffices to note that the lattice  $L = (L_Q \cap L_\beta) + (q - 1)L_\beta$  satisfies the condition  $(q - 1)L_\beta \subseteq L \subseteq L_\beta$  of Theorem 5.2, and hence  $I(Y_Q) = I(V_X(I_L)) = I_L$ . Applying Theorem 5.3 again, we get  $\mathcal{L} : (q - 1) = \mathcal{L}$ .  $\square$

We close this section by sharing a procedure for computing the vanishing ideal of  $V_X(I_L) \cap T_X$  for lattices  $L \subseteq L_\beta$  not satisfying the condition  $(q - 1)L_\beta \subseteq L$ .

**Corollary 5.6.** *If  $L \subseteq L_\beta$ , then the vanishing ideal of  $V_X(I_L) \cap T_X$  is  $I_{L'}$ , for the lattice  $L' = L + (q - 1)L_\beta$ . Moreover, the order of  $V_X(I_L) \cap T_X$  is  $|L_\beta/L'|$ .*

*Proof.* If  $L' = L + (q - 1)L_\beta$ , then  $I_{L'} = I_L + I_{(q-1)L_\beta}$ . Since  $I(T_X) = I_{(q-1)L_\beta}$  it follows that we have

$$V_X(I_{L'}) = V_X(I_L) \cap V_X(I_{(q-1)L_\beta}) = V_X(I_L) \cap T_X.$$

As  $(q - 1)L_\beta \subseteq L' \subseteq L_\beta$ , Theorem 5.2 implies that  $I_{L'} = I(V_X(I_{L'}))$ . Hence, the  $B$ -saturated ideal corresponding to  $Y = V_X(I_L) \cap T_X$  is  $I(Y) = I_{L'}$ . The second claim follows directly from Theorem 4.4.  $\square$

## 6. DEGREE OF A LATTICE IDEAL

In this section, we assume that the  $n$ -dimensional toric variety  $X$  is smooth and projective. Let  $S$  be the Cox ring of  $X$  and  $M$  an  $S$ -graded module. Denote by  $H_M(\alpha)$  the multigraded Hilbert function of  $M$ , that is,  $H_M(\alpha) = \dim_{\mathbb{K}}(M_\alpha)$ . Maclagan and Smith proved in [15, Proposition 2.10] that there is a polynomial  $P_M(t_1, \dots, t_d) \in \mathbb{Q}[t_1, \dots, t_d]$  called the *multigraded Hilbert polynomial* of  $M$  such that  $P_M(\alpha) = H_M(\alpha)$  for a non-empty subset of  $\mathbb{N}\beta$ . They also showed that for a multigraded ideal  $I$ , the Hilbert polynomials of  $S/I$  and that of  $S/(I : B^\infty)$  are the same, where  $(I : B^\infty)$  is the  $B$ -saturation of  $I$ , see [15, Lemma 2.13]. Therefore, if  $I$  is the  $B$ -saturated ideal corresponding to a finite set of points then Hilbert polynomial is constant: the cardinality of  $V_X(I)$ , see [15, Example 4.12]. Inspired from these, we make the following

**Definition 6.1.** Let  $I$  be the  $B$ -saturated homogeneous ideal corresponding to a finite set of points. The degree of  $I$  is the constant Hilbert polynomial  $P_{S/I}$ .

**Lemma 6.2.** If  $I_L$  is an  $L_\beta$ -homogeneous ideal, then its (multigraded) Hilbert function defined by  $H_{I_L}(\alpha) := \dim_{\mathbb{K}}(S_\alpha/I_L)$  is independent of the field  $\mathbb{K}$ .

*Proof.* Recall that the vector space  $S_\alpha$  is spanned by the monomials

$$\mathbf{x}^{\mathbf{a}} = x_1^{a_1} \cdots x_r^{a_r} \text{ with } \deg(\mathbf{x}^{\mathbf{a}}) = \sum_{j=1}^r a_j \beta_j = \alpha.$$

By [17, Theorem 7.3], the quotient ring  $S/I_L$  is isomorphic to the semigroup ring of the semigroup  $\mathbb{N}^r / \sim_L$ , where  $\mathbf{a}_1 \sim_L \mathbf{a}_2 \iff \mathbf{a}_1 - \mathbf{a}_2 \in L$ . If  $I_L$  is  $L_\beta$ -homogeneous, then  $L \subseteq L_\beta$ . Therefore,  $\mathbf{a}_1 - \mathbf{a}_2 \in L$  implies  $\mathbf{a}_1 - \mathbf{a}_2 \in L_\beta$ , i.e. if  $\mathbf{x}^{\mathbf{a}_1} + I_L = \mathbf{x}^{\mathbf{a}_2} + I_L$  in  $S/I_L$ , then  $\deg(\mathbf{x}^{\mathbf{a}_1}) = \deg(\mathbf{x}^{\mathbf{a}_2})$  meaning that  $\mathbf{x}^{\mathbf{a}_1} + I_L = \mathbf{x}^{\mathbf{a}_2} + I_L$  in  $S_\alpha/I_L$ . Hence,  $H_{I_L}(\alpha) = \dim_{\mathbb{K}}(S_\alpha/I_L)$  is the number of equivalence classes  $\mathbf{x}^{\mathbf{a}} + I_L$  of monomials  $\mathbf{x}^{\mathbf{a}}$  of degree  $\alpha \in \mathbb{N}_\beta$ , which is also the number of equivalence classes of vectors  $\mathbf{a} \in \mathbb{N}^r$  with respect to the equivalence relation  $\sim_L$  such that  $\sum_{j=1}^r a_j \beta_j = \alpha$ . This number is independent of the field, completing the proof.  $\square$

**Theorem 6.3.** If  $I_L$  is an  $L_\beta$ -homogeneous ideal of dimension  $d = r - n$ , then its degree is  $\deg(I_L) = |L_\beta/L| = d_1 \cdots d_n$ , where  $d_i$ 's are the invariant factors of  $L_\beta/L$ .

*Proof.* By Lemma 6.2, the multigraded Hilbert polynomial of  $I_L$  is independent of the field. Under the hypothesis  $\text{rank } L = n$  and  $L \subseteq L_\beta$ , so by Proposition 4.8 there is a finite field  $\mathbb{F}_q$  such that  $(q-1)L_\beta \subseteq L \subseteq L_\beta$ . Thus, the number of  $\mathbb{F}_q$ -rational points is  $|V_X(I_L)(\mathbb{F}_q)| = d_1 \cdots d_n$  by Theorem 4.4. On the other hand, the  $B$ -saturated ideal corresponding to  $V_X(I_L)$  is  $I(V_X(I_L)) = I_L$  by Theorem 5.2. Thus, the Hilbert polynomial of  $I_L$  will be  $d_1 \cdots d_n$ , which is the degree of  $I_L$ .  $\square$

The following reveals that Theorem 6.3 generalizes the main result of the nice paper [14] by Lopez and Villarreal.  $X$  is the projective space  $\mathbb{P}^n$  in this case. The matrix  $\beta = [1 \cdots 1]$ ,  $L_\beta = \mathbb{Z}\{e_1 - e_r, \dots, e_{r-1} - e_r\}$  and hence an ideal is  $L_\beta$ -homogeneous with respect to the standard grading:  $\deg(x_i) = 1$ , for each  $i \in [r]$ , where  $r = n + 1$ .

**Corollary 6.4.** [14, Theorem 3.13] If  $I_L$  is an  $L_\beta$ -homogeneous ideal of dimension 1, then its degree is  $\deg(I_L) = |T(\mathbb{Z}^r/L)|$ , where  $T(\mathbb{Z}^r/L)$  is the torsion part of the group  $\mathbb{Z}^r/L$ .

*Proof.* It is well known from the fundamental structure theorem for finitely generated abelian groups that the order of the torsion part  $T(\mathbb{Z}^r/L)$  of the group  $\mathbb{Z}^r/L$  is  $d_1 \cdots d_n$ , where  $d_i$ 's are the invariant factors of the matrix whose columns span the lattice  $L$ , see [10, pp. 187–188]. So, Theorem 6.3 completes the proof.  $\square$

If an ideal  $I = \langle F_1, \dots, F_n \rangle$  is a complete intersection generated in semi-ample degrees  $\alpha_1, \dots, \alpha_n$ , then its Hilbert Polynomial is the Bernstein—Kushnirenko bound, which is the mixed volume  $n!V(P_{\alpha_1}, \dots, P_{\alpha_n})$  of the polytopes  $P_{\alpha_1}, \dots, P_{\alpha_n}$  by the proof of [24, Theorem 3.16]. The following reveals the same is true in the special case that  $I = I_L$  is a lattice ideal, without the assumption that the degrees  $\alpha_1, \dots, \alpha_n$  are semi-ample.

**Corollary 6.5.** *Assume that  $\{d_1\mathbf{m}_1, \dots, d_n\mathbf{m}_n\}$  is a basis of the lattice  $L$  such that  $\{\mathbf{m}_1, \dots, \mathbf{m}_n\}$  is a basis of  $L_\beta$ . If  $\alpha_i = \deg(\mathbf{x}^{d_i\mathbf{m}_i^+} - \mathbf{x}^{d_i\mathbf{m}_i^-})$ , for  $i \in [n]$ , then the degree of  $I_L$  is the mixed volume  $n!V(P_{\alpha_1}, \dots, P_{\alpha_n})$ .*

*Proof.* By Proposition 4.8, there is a finite field  $\mathbb{K} = \mathbb{F}_q$  containing all the  $d_i$ -th roots of unity, for each  $i \in [n]$ . We know that  $V_X(I_L)(\mathbb{K})$  lies inside the torus  $T_X(\mathbb{K})$  by Proposition 3.3. So, localizing to the torus does not change the order of  $V_X(I_L)(\mathbb{K})$ . This amounts to passing to the Laurent polynomial ring

$$\mathbb{K}[\mathbf{x}^\mp] = \mathbb{K}[x_1, \dots, x_r]_{x_1 \cdots x_r} = \mathbb{K}[x_1, \dots, x_r, x_1^{-1}, \dots, x_r^{-1}].$$

By [8, Theorem 2.1], the ideal  $\mathbb{K}[\mathbf{x}^\mp] \cdot I_L$  is a local complete intersection generated by the Laurent polynomials  $\mathbf{x}^{d_1\mathbf{m}_1} - 1, \dots, \mathbf{x}^{d_n\mathbf{m}_n} - 1$  forming a regular sequence, even if  $I_L$  is not the (global) complete intersection ideal

$$\langle \mathbf{x}^{d_1\mathbf{m}_1^+} - \mathbf{x}^{d_1\mathbf{m}_1^-}, \dots, \mathbf{x}^{d_n\mathbf{m}_n^+} - \mathbf{x}^{d_n\mathbf{m}_n^-} \rangle.$$

Thus,  $|V_X(I_L)(\overline{\mathbb{K}})|$  meets the Bernstein–Kushnirenko bound, see [3, 12]. Since  $\mathbb{K}$  has all the  $d_i$ -th roots of unity, for each  $i \in [n]$ , we have  $V_X(I_L)(\overline{\mathbb{K}}) = V_X(I_L)(\mathbb{K})$ . Thus,  $n!V(P_{\alpha_1}, \dots, P_{\alpha_n}) = d_1 \cdots d_n$ .

As the Hilbert polynomial of  $I_L$  is the constant  $d_1 \cdots d_n$  by Theorem 6.3, the claim follows.  $\square$

**Example 6.6.** *We revisit Example 4.9. Consider the fan in  $\mathbb{R}^2$  with rays generated by  $\mathbf{v}_1 = (1, 0)$ ,  $\mathbf{v}_2 = (0, 1)$ ,  $\mathbf{v}_3 = (-1, 2)$ , and  $\mathbf{v}_4 = (0, -1)$ . Then the corresponding toric variety is the Hirzebruch surface  $X = \mathcal{H}_2$ . Recall that a  $\mathbb{Z}$ -basis of  $L_\beta$  is given by  $\mathbf{u}_1 = (1, 0, -1, 0)$  and  $\mathbf{u}_2 = (0, 1, 2, -1)$ . Consider the lattice  $L$  spanned over  $\mathbb{Z}$  by  $3\mathbf{u}_1$  and  $5\mathbf{u}_2$ . Then, clearly  $15L_\beta \subset L \subset L_\beta$ . So, the degree of  $I_L$  is just the order  $|L_\beta/L| = 15$ . One can check that  $I_L$  is a  $B$ -saturated radical ideal over any field whose characteristic is coprime to 15, where  $B = \langle x_1x_2, x_2x_3, x_3x_4, x_4x_1 \rangle$ .*

*For instance,  $I_L = I(V_X(I_L))$  is the  $B$ -saturated ideal of  $V_X(I_L)$ , by Theorem 5.2 over the field  $\mathbb{F}_{16}$ . Thus,  $V_X(I_L)$  has  $15 = \deg(I_L)$  members of  $T_X$  over  $\mathbb{F}_{16}$ .*

*However, if  $q = 11$  then  $I_L$  is not the  $B$ -saturated ideal of  $V_X(I_L)$ . Indeed, we have to replace  $L$  by  $L' = L + (q-1)L_\beta$  so as to obtain the correct lattice. This is because, we have  $I_{L'} = I_L + I_{(q-1)L_\beta}$  and that*

$$V_X(I_{L'}) = V_X(I_L) \cap V_X(I_{(q-1)L_\beta}) = V_X(I_L) \cap T_X = V_X(I_L)$$

*whenever  $L' = L + (q-1)L_\beta$ . Therefore,  $I_{L'} = I(V_X(I_L))$  is the  $B$ -saturated ideal of  $V_X(I_L)$ , by Theorem 5.2 as  $10L_\beta \subset L' \subset L_\beta$ . A basis for the lattice  $L'$  is given by  $\mathbf{u}_1$  and  $5\mathbf{u}_2$ , since  $\mathbf{u}_1 = -3(3\mathbf{u}_1) + 10\mathbf{u}_1 \in L'$ . The Smith-normal form of the matrix with columns  $\mathbf{u}_1$  and  $5\mathbf{u}_2$  is  $[\mathbf{e}_1 | 5\mathbf{e}_2]$ , where  $\mathbf{e}_i$  is a standard basis vector of  $\mathbb{Z}^4$ . Thus,  $V_X(I_{L'})$  has  $5 = \deg(I_{L'})$  members of  $T_X$  over the finite field  $\mathbb{F}_{11}$ .*

## 7. PARAMETERIZATION OF THE SUBGROUP CUT OUT BY A LATTICE IDEAL

In this section, we describe the points of the subgroup  $V_X(I_L)$  of the torus  $T_X$  defined by the lattice ideal  $I_L$ .

Let  $\mathcal{B}$  be an  $r \times n$  matrix whose columns constitute a basis for  $L \subseteq L_\beta$ ; let  $\mathcal{A}$  and  $\mathcal{C}$  be unimodular matrices of sizes  $r$  and  $n$ , respectively, so that

$$\mathcal{D} = \mathcal{A}\mathcal{B}\mathcal{C} = [d_1\mathbf{e}_1 | \cdots | d_n\mathbf{e}_n]$$

is the Smith-Normal form of  $\mathcal{B}$ , where  $\mathbf{e}_1, \dots, \mathbf{e}_n \in \mathbb{Z}^r$ . Recall that  $d_i$  divides  $d_{i+1}$ , for all  $i \in [n-1]$ . If  $\mathcal{A}^{-1} = [\mathbf{m}_1 | \cdots | \mathbf{m}_r]$  is the inverse of  $\mathcal{A}$ , then its columns span

$\mathbb{Z}^r$ . Thus, the columns  $\{d_1 \mathbf{m}_1, \dots, d_n \mathbf{m}_n\}$  of  $\mathcal{A}^{-1} \mathcal{D} = \mathcal{BC}$  provide us with another basis of the lattice  $L$  such that  $\{\mathbf{m}_1, \dots, \mathbf{m}_n\}$  is a basis of  $L_\beta$ . Since  $\text{rank } L = n$ , we have  $V_X(I_L) \subseteq T_X$  by Proposition 3.3.

**Theorem 7.1.** *Let  $L \subseteq L_\beta$  be a lattice of rank  $n$ . Then, the following hold:*

- (i) *If  $[P_i] = [\eta_i^{a_{i1}} : \dots : \eta_i^{a_{ir}}]$ , where  $\eta_i$  is a generator for the cyclic subgroup of  $d_i$ -th roots of unity in  $\mathbb{K}$ , for all  $i \in [n]$ , then*

$$V_X(I_L)(\mathbb{K}) = \langle [P_1] \rangle \times \dots \times \langle [P_n] \rangle.$$

- (ii) *If  $d_i$  divides  $q - 1$ , for all  $i \in [n]$ , and  $Q$  is the matrix whose  $i$ -th row is  $[a_{i1}(q-1)/d_i | \dots | a_{ir}(q-1)/d_i]$  which is  $(q-1)/d_i$  times the  $i$ -th row of the matrix  $\mathcal{A}$ , then  $V_X(I_L)(\mathbb{F}_q) = Y_Q$ .*

*Proof.* The algebraic isomorphism induced by the change of basis map  $\mathbb{Z}^r \rightarrow \mathbb{Z}^r$  defined via multiplication by the matrix  $\mathcal{A} = [\mathbf{a}_1 | \dots | \mathbf{a}_r]$  is as follows:

$$(7.1) \quad \Phi_{\mathcal{A}} : (\mathbb{K}^*)^r \rightarrow (\mathbb{K}^*)^r, \quad \Phi_{\mathcal{A}}(y_1, \dots, y_r) = (\mathbf{y}^{\mathbf{a}_1}, \dots, \mathbf{y}^{\mathbf{a}_r}).$$

with the following inverse isomorphism:

$$(7.2) \quad \Phi_{\mathcal{A}}^{-1} : (\mathbb{K}^*)^r \rightarrow (\mathbb{K}^*)^r, \quad \Phi_{\mathcal{A}}^{-1}(x_1, \dots, x_r) = (\mathbf{x}^{\mathbf{m}_1}, \dots, \mathbf{x}^{\mathbf{m}_r}).$$

Recall that  $G$  is the subgroup  $V(I_{L_\beta}) \cap (\mathbb{K}^*)^r$  and thus  $(x_1, \dots, x_r) \in G$  if and only if  $\mathbf{x}^{\mathbf{m}_1} = \dots = \mathbf{x}^{\mathbf{m}_n} = 1$ , by the virtue of [23, Lemma 3.3] saying that

$$V(I_{L_\beta}) \cap (\mathbb{K}^*)^r = \bigcap_{i=1}^n V(\mathbf{x}^{\mathbf{m}_i^+} - \mathbf{x}^{\mathbf{m}_i^-}) \cap (\mathbb{K}^*)^r.$$

Therefore, if  $(y_1, \dots, y_r) = \Phi_{\mathcal{A}}^{-1}(x_1, \dots, x_r)$ , then it follows that  $(x_1, \dots, x_r) \in G$  if and only if  $y_1 = \dots = y_n = 1$ . Thus, the group  $G' := \Phi_{\mathcal{A}}^{-1}(G)$  is described by

$$G' = \{(1, \dots, 1, y_{n+1}, \dots, y_r) : y_i \in (\mathbb{K}^*)^r\}.$$

On the other hand, since a basis for  $L$  is given by the set  $\{d_1 \mathbf{m}_1, \dots, d_n \mathbf{m}_n\}$ , we similarly have the following:

$$(x_1, \dots, x_r) \in V(I_L) \cap (\mathbb{K}^*)^r \iff \mathbf{x}^{d_1 \mathbf{m}_1} = \dots = \mathbf{x}^{d_n \mathbf{m}_n} = 1.$$

Finally, as the elements  $[x_1 : \dots : x_r] \in T_X$  are orbits  $G \cdot (x_1, \dots, x_r)$ , it follows that

$$[x_1 : \dots : x_r] \in T_X \iff [y_1 : \dots : y_n : 1 : \dots : 1] \in (\mathbb{K}^*)^r / G'.$$

Thus,  $\Phi_{\mathcal{A}}^{-1}$  induces an isomorphism between  $T_X$  and the torus  $(\mathbb{K}^*)^r / G'$  identifying the subgroup  $V_X(I_L)$  of  $T_X$  with the following subgroup:

$$\mathcal{G}_L := \{[y_1 : \dots : y_n : 1 : \dots : 1] \in (\mathbb{K}^*)^r / G' : y_1^{d_1} = \dots = y_n^{d_n} = 1\}.$$

If  $C_i = \{y_i \in \mathbb{K}^* : y_i^{d_i} = 1\}$ , for all  $i \in [n]$ , then,  $\mathcal{G}_L$  is isomorphic to  $C_1 \times \dots \times C_n$ , via

$$[y_1 : \dots : y_n : 1 : \dots : 1] \rightarrow (y_1, \dots, y_n).$$

If  $C_i$  is the cyclic subgroup of  $\mathbb{K}^*$  generated by  $\eta_i$ , then the elements of  $\mathcal{G}_L$  are of the form

$$[y_1 : \dots : y_n : 1 : \dots : 1] = [\eta_1^{k_1} : \dots : \eta_n^{k_n} : 1 : \dots : 1],$$

which bijectively correspond to the  $n$ -tuples  $(k_1, \dots, k_n)$  with  $0 \leq k_i \leq |C_i| - 1$ . Let  $[P] = [x_1 : \dots : x_r]$  be a point with  $x_i = \mathbf{y}^{\mathbf{a}_i} = y_1^{a_{i1}} \dots y_r^{a_{ri}}$ . Then, we have

$$[P] \in V_X(I_L) \iff \Phi_{\mathcal{A}}^{-1}(P) \in \mathcal{G}_L.$$

Therefore,  $[P] \in V_X(I_L) \iff$  there is an  $n$ -tuple  $(k_1, \dots, k_n)$  with  $0 \leq k_i \leq |C_i| - 1$  such that

$$(7.3) \quad [P] = [\eta_1^{a_{11}k_1} \dots \eta_n^{a_{n1}k_n} : \dots : \eta_1^{a_{1r}k_1} \dots \eta_n^{a_{nr}k_n}]$$

$$(7.4) \quad = [\eta_1^{a_{11}} : \dots : \eta_1^{a_{1r}}]^{k_1} \dots [\eta_n^{a_{n1}} : \dots : \eta_n^{a_{nr}}]^{k_n} = [P_1]^{k_1} \dots [P_n]^{k_n}.$$

This proves the first part (i):

$$V_X(I_L)(\mathbb{K}) = \langle [P_1] \rangle \times \dots \times \langle [P_n] \rangle.$$

When  $\mathbb{K} = \mathbb{F}_q$  and  $d_i$  divides  $q - 1$ ,  $C_i$  becomes the cyclic subgroup of  $\mathbb{F}_q^* = \langle \eta \rangle$  of order  $d_i$ , generated by  $\eta_i = \eta^{q-1/d_i}$ , for all  $i \in [n]$ . Letting  $t_i = \eta^{k_i}$  for all  $i \in [n]$ , we observe that the point in (7.3) becomes

$$\begin{aligned} [P] &= [\eta^{a_{11}k_1(q-1)/d_1} \dots \eta^{a_{n1}k_n(q-1)/d_n} : \dots : \eta^{a_{1r}k_1(q-1)/d_1} \dots \eta^{a_{nr}k_n(q-1)/d_n}] \\ &= [t_1^{a_{11}(q-1)/d_1} \dots t_n^{a_{n1}(q-1)/d_n} : \dots : t_1^{a_{1r}(q-1)/d_1} \dots t_n^{a_{nr}(q-1)/d_n}]. \end{aligned}$$

If  $Q$  is the matrix whose  $i$ -th row is  $[a_{i1}(q-1)/d_i \dots |a_{ir}(q-1)/d_i]$ , then it follows that  $[P] \in V_X(I_L)$  implies  $[P] \in Y_Q$ .

In order to prove the converse inclusion, take  $[P] \in Y_Q$ . Then, there are  $t_i \in \mathbb{F}_q^*$  such that

$$[P] = [t_1^{a_{11}(q-1)/d_1} \dots t_n^{a_{n1}(q-1)/d_n} : \dots : t_1^{a_{1r}(q-1)/d_1} \dots t_n^{a_{nr}(q-1)/d_n}].$$

Since  $\mathbb{F}_q^* = \langle \eta \rangle$ , we have  $\ell_i$  between 0 and  $q - 2$  for which  $t_i = \eta^{\ell_i}$ , for all  $i \in [n]$ . Then, there is a unique  $k_i$  between 0 and  $d_i - 1$  such that  $\ell_i \cong k_i$  modulo  $d_i$ . Thus,  $t_i^{(q-1)/d_i} = \eta^{\ell_i(q-1)/d_i} = \eta^{k_i(q-1)/d_i}$ , yielding  $[P] \in V_X(I_L)$ .  $\square$

**Corollary 7.2.** *If  $V_X(I_L)(\mathbb{F}_q)$  is a cyclic subgroup, then  $V_X(I_L)(\mathbb{F}_q) = Y_Q$  for a row matrix  $Q = [a_{n1}(q-1)/d_n \dots |a_{nr}(q-1)/d_n]$  which is  $(q-1)/d_n$  times the  $n$ -th row of the matrix  $\mathcal{A}$ .*

*Proof.* By Theorem 4.4,  $V_X(I_L)(\mathbb{F}_q)$  has  $d_1 \dots d_n$  elements. Since it is cyclic,  $d_1 = \dots = d_{n-1} = 1$  and  $|V_X(I_L)(\mathbb{F}_q)| = d_n$ . By Theorem 7.1, it follows that  $V_X(I_L)(\mathbb{F}_q) = Y'_Q$ , where  $Q'$  is the matrix whose  $i$ -th row is  $(q-1)/d_i$  times the  $i$ -th row of the matrix  $\mathcal{A}$ . When  $d_i = 1$ , the following point

$$[P] = [t_1^{a_{11}(q-1)/d_1} \dots t_n^{a_{n1}(q-1)/d_n} : \dots : t_1^{a_{1r}(q-1)/d_1} \dots t_n^{a_{nr}(q-1)/d_n}]$$

of  $Y'_Q$  becomes

$$[P] = [t_n^{a_{n1}(q-1)/d_n} : \dots : t_n^{a_{nr}(q-1)/d_n}]$$

due to the fact that  $t_i^{a_{ij}(q-1)/d_i} = 1$ , for all  $i \in [n-1]$  and  $j \in [r]$ . Thus,  $[P]$  is a point of  $Y_Q$ . As the converse is also true, the claim follows.  $\square$

**Remark 7.3.** *Although a parameterization  $Y_Q$  of the subgroup  $V_X(I_L) \cap T_X$  is given in [23, Proposition 3.4], the size of the matrix  $Q$  for which  $V_X(I_L) \cap T_X = Y_Q$  was always  $r \times r$ , even if the subgroup is cyclic. This is in contrast with the matrices provided in Theorem 7.1 and Corollary 7.2 that are more natural and succinct.*

8. EVALUATION CODES ON SUBGROUPS OF THE TORUS  $T_X(\mathbb{F}_q)$ 

In this section, we apply the main results of the paper to compute basic parameters of toric codes on subgroups  $Y = V_X(I_L)$  of the torus  $T_X(\mathbb{F}_q)$ , where a basis for  $L$  is given by  $\{c_1 \mathbf{u}_1, \dots, c_n \mathbf{u}_n\}$  for positive integers  $c_i$  dividing  $q - 1$ .

We first recall the evaluation code defined on a subset  $Y = \{[P_1], \dots, [P_N]\}$  of  $X(\mathbb{F}_q)$ . Let  $\mathbb{N}\beta$  be the subsemigroup of  $\mathbb{N}^n$  generated (not necessarily minimally) by  $\beta_1, \dots, \beta_r$ . Recall that the Cox ring  $S = \bigoplus_{\alpha \in \mathbb{N}\beta} S_\alpha$  of  $X$  is multigraded by  $\mathbb{N}\beta$  via

$\deg(x_i) = \beta_i$  for  $i = 1, \dots, r$ . For any multidegree  $\alpha \in \mathbb{N}\beta$ , we have the following evaluation map

$$(8.1) \quad \text{ev}_Y : S_\alpha \rightarrow \mathbb{F}_q^N, \quad F \mapsto (F(P_1), \dots, F(P_N)).$$

The image  $\mathcal{C}_{\alpha,Y} = \text{ev}_Y(S_\alpha)$  is a linear code, called the *generalized toric code*. The three basic parameters of  $\mathcal{C}_{\alpha,Y}$  are block-length which is  $N$ , the dimension which is  $K = \dim_{\mathbb{F}_q}(\mathcal{C}_{\alpha,Y})$ , and the minimum distance  $\delta = \delta(\mathcal{C}_{\alpha,Y})$  which is the minimum of the numbers of nonzero components of nonzero vectors in  $\mathcal{C}_{\alpha,Y}$ .

It is clear that the kernel of the linear map  $\text{ev}_Y$  equals the homogeneous piece  $I(Y)_\alpha$  of the vanishing ideal  $I(Y)$  in degree  $\alpha$ . Therefore, the code  $\mathcal{C}_{\alpha,Y}$  is isomorphic to the  $\mathbb{K}$ -vector space  $S_\alpha/I(Y)_\alpha$ . Thus, the dimension of  $\mathcal{C}_{\alpha,Y}$  is the multigraded Hilbert function  $H_Y(\alpha) := \dim_{\mathbb{F}_q} S_\alpha - \dim_{\mathbb{F}_q} I(Y)_\alpha$  of  $I(Y)$ . As  $\mathcal{C}_{\alpha,Y}$  is a subspace of  $\mathbb{F}_q^N$ , we have  $H_Y(\alpha) \leq N$ . When we have the equality  $K = H_Y(\alpha) = N$ , the code becomes trivial, i.e.  $\delta = \delta(\mathcal{C}_{\alpha,Y}) = 1$ , by the Singleton bound  $\delta \leq N + 1 - K$ . A related algebraic notion useful to eliminate these trivial codes is the so-called *multigraded regularity* defined by

$$\text{reg}(Y) := \{\alpha \in \mathbb{N}\beta \quad : \quad H_Y(\alpha) = |Y|\} \subseteq \mathbb{N}^d.$$

The values of the Hilbert function  $H_Y(\alpha)$  can also be used to detect (monomially) equivalent codes. Indeed, by [24, Proposition 4.3], the codes  $\mathcal{C}_{\alpha,Y}$  and  $\mathcal{C}_{\alpha',Y}$  are equivalent if  $H_Y(\alpha) = H_Y(\alpha')$  and  $\alpha - \alpha' \in \mathbb{N}\beta$ .

The first application of our results is that the length of the code  $\mathcal{C}_{\alpha,Y}$  is given by  $N = |Y| = c_1 \cdots c_n$ , due to Corollary 4.7, where  $Y = V_X(I_L)$  and a basis for  $L$  is given by  $\{c_1 \mathbf{u}_1, \dots, c_n \mathbf{u}_n\}$  for positive integers  $c_i$  dividing  $q - 1$ .

In order to compute the dimension and minimum distance of the code  $\mathcal{C}_{\alpha,Y}$ , we need to determine a minimal generating set for the vanishing ideal  $I(Y)$ . We apply our Theorem 5.2 together with a characterization of complete intersection lattice ideals given by Morales and Thoma [18] using the following concept.

**Definition 8.1.** *Let  $A$  be a matrix whose entries are all integers.  $A$  is called mixed if there is a positive and a negative entry in every column. If no square submatrix of  $A$  is mixed, it is called dominating.*

**Theorem 8.2.** [18, Theorem 3.9] *Let  $L \subseteq \mathbb{Z}^r$  be a lattice with the property that  $L \cap \mathbb{N}^r = 0$ . Then,  $I_L$  is complete intersection  $\iff$   $L$  has a basis  $\mathbf{m}_1, \dots, \mathbf{m}_k$  such that the matrix  $[\mathbf{m}_1 \cdots \mathbf{m}_k]$  is mixed dominating. In the affirmative case, we have  $I_L = \langle \mathbf{x}^{\mathbf{m}_1^+} - \mathbf{x}^{\mathbf{m}_1^-}, \dots, \mathbf{x}^{\mathbf{m}_k^+} - \mathbf{x}^{\mathbf{m}_k^-} \rangle$ .  $\square$*

Applying our Theorem 5.2 to  $Y = V_X(I_L)$  we immediately obtain  $I(Y) = I_L$ . Using then Theorem 8.2, one can confirm when  $I(Y_Q) = I_L$  is a complete intersection by looking at a basis of the lattice  $L$ . The following is an instance where everything works very well.

**Proposition 8.3.** *Let  $X$  be the Hirzebruch surface  $\mathcal{H}_\ell$ . If  $\{c_1\mathbf{u}_1, c_2\mathbf{u}_2\}$  constitute a basis for  $L$ , and  $Y = V_X(I_L)$  then  $I(Y)$  is a complete intersection generated minimally by  $x_3^{c_1} - x_1^{c_1}$  and  $x_4^{c_2} - x_2^{c_2}x_3^{\ell c_2}$ . Furthermore,  $(c_1 + c_2\ell, c_2) + \mathbb{N}^2 \subseteq \text{reg}(Y)$ .*

*Proof.* By Theorem 5.2, we get  $I(Y) = I_L$ . Since the columns of  $\mathbf{M}L$  constitute a basis of  $L$ , where

$$\mathbf{M}L = [c_1\mathbf{u}_1 \quad c_2\mathbf{u}_2] = \begin{bmatrix} c_1 & 0 & -c_1 & 0 \\ 0 & c_2 & \ell c_2 & -c_2 \end{bmatrix}^T$$

and  $\mathbf{M}L$  is mixed dominating, it follows from Theorem 8.2 that

$$I(Y) = \langle x_3^{c_1} - x_1^{c_1}, x_4^{c_2} - x_2^{c_2}x_3^{\ell c_2} \rangle.$$

There is a geometrically significant subsemigroup  $\mathcal{K}$  contained in the semigroup  $\mathbb{N}\beta = \mathbb{N}^2$  whose elements are semiample degrees corresponding to the classes of numerically effective line bundles on  $X$ , which is given by

$$\mathcal{K} = \bigcap_{\sigma \in \Sigma} \mathbb{N}\check{\sigma} = \mathbb{N}(1, 0) + \mathbb{N}(\ell, 1),$$

where  $\mathbb{N}\check{\sigma}$  is the semigroup generated by  $\beta_i$  corresponding to rays  $\rho_i \notin \sigma$ , for details see [6, Theorem 6.3.12]. Thus, the degrees  $c_1(1, 0)$  and  $c_2(\ell, 1)$  of the generators are semiample. Therefore,  $(c_1 + c_2\ell, c_2) + \mathbb{N}^2 \subseteq \text{reg}(Y)$  by [24, Theorem 3.16].  $\square$

It is now time to compute the other main parameters of the toric code  $\mathcal{C}_{\alpha, Y}$ , where  $\alpha = (c, d) \in \mathbb{N}\beta$ . Hansen computed these parameters for the case  $Y = T_X$ ,  $c < q - 1$  and  $d = b$ , where  $b$  is to be defined below, see [9]. More recently, the conditions  $c < q - 1$  and  $d = b$  is relaxed in [1] again for the torus  $Y = T_X$ .

**Theorem 8.4.** *Let  $\{c_1\mathbf{u}_1, c_2\mathbf{u}_2\}$  be a basis for  $L$ , and  $Y = V_X(I_L)$  be the subgroup of the torus  $T_X$  of the Hirzebruch surface  $\mathcal{H}_\ell$  over  $\mathbb{K} = \mathbb{F}_q$  for which  $c_1$  and  $c_2$  divides  $q - 1$ . If  $\alpha = (c, d) \in \mathbb{N}\beta$ , then the dimension  $K$  of the toric code  $\mathcal{C}_{\alpha, Y}$  is given by*

$$K = \begin{cases} (b+1)[c+1-\ell b/2], & \text{if } c < c_1 \text{ and } b \leq c_2 - 1 \\ c_2[c+1-(c_2-1)\ell/2] & \text{if } c < c_1 \text{ and } b' < c_2 - 1 < b \\ c_1(b'+1) + (b-b')[c+1-\ell(b+b'+1)/2], & \text{if } c \geq c_1 \text{ and } b \leq c_2 - 1 \\ c_1(b'+1) + (c_2-1-b')[c+1-\ell(c_2+b')/2], & \text{if } c \geq c_1 \text{ and } b' < c_2 - 1 < b \\ c_1c_2, & \text{if } c \geq c_1 \text{ and } b' \geq c_2 - 1, \end{cases}$$

and the minimum distance equals

$$\delta(\mathcal{C}_{\alpha, Y}) = \begin{cases} c_2[c_1 - c], & \text{if } c < c_1 \text{ and } b \leq c_2 - 1 \\ c_2[c_1 - c], & \text{if } c < c_1 \text{ and } b' < c_2 - 1 < b \\ c_2 - b', & \text{if } c \geq c_1 \text{ and } b \leq c_2 - 1 \\ c_2 - b', & \text{if } c \geq c_1 \text{ and } b' < c_2 - 1 < b \\ 1, & \text{if } c \geq c_1 \text{ and } b' \geq c_2 - 1. \end{cases}$$

where  $b$  (respectively  $b'$ ) is the greatest non-negative integer with the property that  $c - b\ell \geq 0$  and  $d - b \geq 0$  (respectively  $c - b'\ell \geq c_1 - 1$  and  $d - b' \geq 0$ ).



*Proof.* Theorem 7.1 gives a parameterization which will be crucial in computing the minimum distance below. So, we first see that  $Q = \begin{bmatrix} 0 & 0 & (q-1)/c_1 & 0 \\ 0 & 0 & 0 & (q-1)/c_2 \end{bmatrix}$  parameterizes the subgroup  $Y = V_X(I_L)$ . Let  $\eta_1 = \eta^{q-1/c_1}$  and  $\eta_2 = \eta^{q-1/c_2}$  where  $\eta$  is a generator for the cyclic group  $\mathbb{K}^*$ . Then,  $Y_Q$  consists of the following  $c_1 c_2$  points  $P_{i,j} = [1 : 1 : \eta_1^i : \eta_2^j] \in T_X$ , for  $i \in [c_1]$  and  $j \in [c_2]$ . Since these points satisfy the equations:

$$x_3^{c_1} - x_1^{c_1} = 0 \text{ and } x_4^{c_2} - x_2^{c_2} x_3^{\ell c_2} = 0$$

it follows that they belong to  $Y$ , yielding  $Y_Q \subseteq Y$ . By Corollary 4.7,  $|Y| = c_1 c_2$  forcing the equality  $Y_Q = Y$ .

Let us find a  $\mathbb{K}$ -basis for  $S_\alpha$  for any  $\alpha = (c, d) \in \mathbb{N}\beta$  where  $\beta = \begin{bmatrix} 1 & 0 & 1 & \ell \\ 0 & 1 & 0 & 1 \end{bmatrix}$ .

Since  $b$  is the greatest non-negative integer with  $\alpha = (c, d) = b(\ell, 1) + (a, a')$  for some non-negative integers  $a = c - b\ell \geq 0$  and  $a' = d - b \geq 0$ , we have  $0 \leq a_4 \leq b$  if  $\deg(\mathbf{x}^{\mathbf{a}}) = \beta \mathbf{a} = \alpha$ . following set  $B_\alpha := \{\mathbf{x}^{\mathbf{a}} \mid \deg(\mathbf{x}^{\mathbf{a}}) = \beta \mathbf{a} = \alpha, 0 \leq a_4 \leq b\}$  is a  $\mathbb{K}$ -basis for  $S_\alpha$ . For a fixed  $a_4$ , and fixed  $0 \leq a_3 \leq c - \ell a_4$  the powers  $a_2 = d - a_4$  and  $a_1 = c - \ell a_4 - a_3$  in  $\mathbf{x}^{\mathbf{a}} = x_1^{a_1} x_2^{a_2} x_3^{a_3} x_4^{a_4}$  are fixed too. So,

$$B_\alpha = \{\mathbf{x}^{\mathbf{a}} \mid 0 \leq a_4 \leq b, 0 \leq a_3 \leq c - \ell a_4, a_2 = d - a_4 \geq 0, a_1 = c - \ell a_4 - a_3 \geq 0\}.$$

Let  $\mu_1$  be the minimum of  $c$  and  $c_1 - 1$ , and  $\mu_2$  be the minimum of  $b$  and  $c_2 - 1$ . Since  $x_3^{c_1} = x_1^{c_1}$  and  $x_4^{c_2} = x_2^{c_2} x_3^{\ell c_2}$  in the ring  $S/I(Y)$ , a basis for  $S_\alpha/I_\alpha(Y)$  is

$$\bar{B}_\alpha = \{\mathbf{x}^{\mathbf{a}} \mid 0 \leq a_4 \leq \mu_2, 0 \leq a_3 \leq \min\{c - \ell a_4, \mu_1\}, d - a_4 \geq 0, c - \ell a_4 - a_3 \geq 0\}.$$

If  $b' \geq c_2 - 1$ , then  $c \geq c_1$  as otherwise  $b' = 0$ . Hence, we have  $\mu_1 = c_1 - 1$  and  $\mu_2 = c_2 - 1$ . So,  $a_4 \leq c_2 - 1 \leq b'$  implying  $c - \ell a_4 \geq c_1 - 1$ , and hence  $0 \leq a_3 \leq c_1 - 1$ . So, for every choice of  $a_4$  there are  $c_1$  choice for  $a_3$ . Thus, we have

$$(8.2) \quad \dim_{\mathbb{K}} \mathcal{C}_{\alpha, Y} = |\bar{B}_\alpha| = c_1 c_2 = N$$

Since the dimension reaches its maximum value, the code  $\mathcal{C}_{\alpha, Y}$  is trivial, that is,  $\delta(\mathcal{C}_{\alpha, Y}) = 1$ . Therefore, we assume  $b' < c_2 - 1$  from now on.

By the definition of  $b'$ , we have  $\min\{c - \ell a_4, c_1 - 1\} = c - \ell a_4$  for  $b' < a_4 \leq \mu_2$  and  $\min\{c - \ell a_4, c_1 - 1\} = c_1 - 1$  for  $0 \leq a_4 \leq b'$ . So, if  $0 \leq a_4 \leq b'$  then  $0 \leq a_3 \leq \mu_1$  but if  $a_4 > b'$  then  $0 \leq a_3 \leq c - \ell a_4$ , yielding the formula

$$(8.3) \quad \begin{aligned} \dim_{\mathbb{K}} \mathcal{C}_{\alpha, Y} &= |\bar{B}_\alpha| = (\mu_1 + 1)(b' + 1) + \sum_{a_4=b'+1}^{\mu_2} (c - \ell a_4 + 1) \\ &= (\mu_1 + 1)(b' + 1) + (\mu_2 - b')[c + 1 - (\mu_2 + b' + 1)\ell/2]. \end{aligned}$$

Notice that if  $a \geq c_1 - 1$ , then  $c - \ell b = a > c_1 - 1$ , and thus  $b' = b$ . Otherwise,

$$b' = \lfloor (c - c_1 + 1)/\ell \rfloor = b - \lfloor (c_1 - 1 - a)/\ell \rfloor.$$

**Case I:** Let  $c = a + b\ell < c_1$ . Then,  $\mu_1 = c$  and  $b' = 0$ .

If  $b \leq c_2 - 1$ , then  $\mu_2 = b$  and by (8.3), we have

$$\dim_{\mathbb{K}} \mathcal{C}_{\alpha, Y} = (b + 1)(c + 1) - b(b + 1)\ell/2 = (b + 1)[c + 1 - \ell b/2].$$

If  $b' < c_2 - 1 < b$ , then  $\mu_2 = c_2 - 1$  and by (8.3), we have

$$\dim_{\mathbb{K}} \mathcal{C}_{\alpha, Y} = c_2(c + 1) - (c_2 - 1)c_2/2 = c_2[c + 1 - (c_2 - 1)\ell/2].$$

**Case II:** Let  $c \geq c_1$ .

If  $b \leq c_2 - 1$ , then  $\mu_1 = c_1 - 1$  and  $\mu_2 = b$  in (8.3), yielding the formula

$$\dim_{\mathbb{K}} \mathcal{C}_{\alpha, Y} = c_1(b' + 1) + (b - b')[c + 1 - (b + b' + 1)\ell/2].$$

If  $b' < c_2 - 1 < b$ , then  $\mu_1 = c_1 - 1$  and  $\mu_2 = c_2 - 1$  in (8.3), yielding the formula

$$\dim_{\mathbb{K}} \mathcal{C}_{\alpha, Y} = c_1(b' + 1) + (c_2 - 1 - b')[c + 1 - (c_2 + b')\ell/2].$$

Let  $(c, d) = (a + \ell b, a' + b)$ . Since the set  $\bar{B}_\alpha$  is independent of  $a'$ , it follows that  $H_Y(\alpha) = |\bar{B}_\alpha| = H_Y(a + \ell b, b)$ .

If  $a \geq c_1 - 1$ , then  $c - \ell b = a \geq c_1 - 1$ . So,  $b' = b$ . If  $b' = b \geq c_2 - 1$ , then  $H_Y(\alpha) = |\bar{B}_\alpha| = |Y| = N$ , by (8.2). If  $b' = b < c_2 - 1$ , then by **Case II** above  $H_Y(\alpha) = |\bar{B}_\alpha| = c_1(b + 1)$ . It follows that the codes  $\mathcal{C}_{\alpha', Y}$  with  $a > c_1 - 1$  are all equivalent to the code  $\mathcal{C}_{\alpha, Y}$  with  $a = c_1 - 1$ , by [24, Proposition 4.3].

Let us compute the minimum distance now. Let  $\eta_1 = \eta^{q-1/c_1}$  and  $\eta_2 = \eta^{q-1/c_2}$  where  $\eta$  is a generator for the cyclic group  $\mathbb{K}^*$ . Then,  $Y = V_X(I_L)$  consists of the following points  $P_{i,j} = [1 : 1 : \eta_1^i : \eta_2^j] \in T_X$ , for  $i \in [c_1]$  and  $j \in [c_2]$ .

Fix  $F \in S_\alpha$  and set  $J = \{j \in [c_2] : x_4 - \eta_2^j x_2 x_3^\ell \text{ divides } F\}$ . Then, we first prove that the number of zeroes of  $F$  in  $Y$  satisfies

$$(8.4) \quad |V_X(F) \cap Y| \leq c_1|J| + (c_2 - |J|) \deg_{x_3} F.$$

It is clear that  $F(P_{i,j}) = 0$  whenever  $j \in J$  and there are  $c_1|J|$  many such points in  $Y$ . Moreover, the polynomial  $F(1, 1, x_3, \eta_2^j) \in \mathbb{K}[x_3] \setminus \{0\}$  if  $j \notin J$ . Thus, it can have at most  $\deg_{x_3} F$  many zeroes for each  $j \notin J$ . So, we can have at most  $(c_2 - |J|) \deg_{x_3} F$  many such roots. Therefore, altogether  $F$  can have at most  $c_1|J| + (c_2 - |J|) \deg_{x_3} F$  zeroes in  $Y$ .

As a second step, we elaborate on this upper bound depending on whether  $|J| \leq b'$  or  $|J| > b'$ .

Suppose that  $0 \leq |J| \leq b'$ . Since  $\deg_{x_3} F \leq \mu_1$ , it follows from (8.4) that

$$\begin{aligned} |V_X(F) \cap Y| &\leq c_1|J| + (c_2 - |J|)\mu_1 = c_2\mu_1 + (c_1 - \mu_1)|J| \\ &\leq c_2\mu_1 + (c_1 - \mu_1)b'. \end{aligned}$$

Since  $b' = 0$  and  $\mu_1 = c$  when  $c < c_1$ , and  $\mu_1 = c_1 - 1$  when  $c \geq c_1$ , we have

$$(8.5) \quad |V_X(F) \cap Y| \leq \begin{cases} c_2c, & \text{if } c < c_1 \\ c_2(c_1 - 1) + b', & \text{if } c \geq c_1. \end{cases}$$

Suppose now that  $|J| > b'$ . In this case, we have  $c - \ell|J| < c_1 - 1$ , by definition of  $b'$ , which will be crucial below. Since  $\deg_{x_3} F \leq c - \ell|J|$ , it follows from (8.4) that

$$\begin{aligned} |V_X(F) \cap Y| &\leq c_1|J| + (c_2 - |J|)(c - \ell|J| - c_1 + c_1) \\ &\leq c_1c_2 - c_2 + c_2 + (c_2 - |J|)(c - \ell|J| - c_1) \\ &\leq (c_1 - 1)c_2 + c_2 + (c_2 - b')(c - \ell|J| - c_1). \end{aligned}$$

Since the difference

$$b' - [c_2 + (c_2 - b')(c - \ell|J| - c_1)] = (c_2 - b')(\ell|J| + c_1 - c - 1) > 0$$

we have that

$$(8.6) \quad |V_X(F) \cap Y| < (c_1 - 1)c_2 + b'.$$

**Case I:** Let  $c < c_1$ . Then, by (8.5),  $F$  can have at most  $c_2c$  many zeroes in  $Y$ . On the other hand, the following polynomial

$$F = x_2^d \prod_{i=1}^c (x_3 - \eta_1^i x_1) \in S_\alpha$$

vanishes exactly at the  $c_2c$  points  $P_{i,j} = [1 : 1 : \eta_1^i : \eta_2^j] \in Y$ , where  $1 \leq i \leq c$  and  $1 \leq j \leq c_2$ . Thus, there is a codeword  $\text{ev}_{\alpha,Y}(F)$  with weight  $c_1c_2 - c_2c$ . Hence,

$$\delta(\mathcal{C}_{\alpha,Y}) = c_1c_2 - c_2c = c_2(c_1 - c).$$

**Case II:** Let  $c \geq c_1$ . Then, by (8.5) and (8.6),  $F$  can have at most  $(c_1 - 1)c_2 + b'$  many zeroes in  $Y$ . On the other hand, the following polynomial

$$F = x_1^{c-\ell b'-(c_1-1)} x_2^{d-b'} \prod_{i=1}^{c_1-1} (x_3 - \eta_1^i x_1) \prod_{j=1}^{b'} (x_4 - \eta_2^j x_1^\ell x_2)$$

vanishes exactly at  $c_1b' + (c_2 - b')(c_1 - 1) = c_2(c_1 - 1) + b'$ , and thus there is a codeword with weight  $c_2 - b'$ . This shows that  $\delta(\mathcal{C}_{\alpha,Y}) = c_2 - b'$ .  $\square$

**Corollary 8.5.** *Let  $\{c_1\mathbf{u}_1, c_2\mathbf{u}_2\}$  be a basis for  $L$ , and  $Y = V_X(I_L)$  be the subgroup of the torus  $T_X$  of the Hirzebruch surface  $\mathcal{H}_\ell$  over  $\mathbb{K} = \mathbb{F}_q$  for which  $c_1$  and  $c_2$  divides  $q - 1$ . If  $\ell \geq c_1$ , then the parameters of the toric code  $\mathcal{C}_{\alpha,Y}$  are  $N = c_1c_2$ ,*

$$\begin{aligned} K &= a + 1 \text{ and } \delta = c_2(c_1 - a), \text{ if } c < c_1 \\ K &= c_1b + 1 + a \text{ and } \delta = c_2 - b + 1, \text{ if } a < c_1 - 1 < c \\ K &= c_1(b + 1) \text{ and } \delta = c_2 - b, \text{ if } a = c_1 - 1 < c. \end{aligned}$$

Furthermore,  $(c_1 + c_2\ell, c_2) + \mathbb{N}^2 = \text{reg}(Y)$ .

*Proof.* Let  $\ell \geq c_1$ . Since  $a \leq c_1 - 1$ , it follows that  $b' = b$  if  $a = c_1 - 1$  and  $b' = b - 1$  if  $a < c_1 - 1$ . So, if  $b \geq c_2$ , then  $b' \geq c_2 - 1$  and thus the code is trivial by (8.2). Thus, non-equivalent codes occur only for  $\alpha = a(1, 0) + b(\ell, 1)$  where  $0 \leq a \leq c_1 - 1$  and  $0 \leq b \leq c_2 - 1$ .

**Case I:** If  $c = a + \ell b < c_1 \leq \ell$ , then  $0 \leq a < \ell(1 - b)$  implying that  $b = 0$ . Hence,  $\dim_{\mathbb{K}} \mathcal{C}_{\alpha,Y} = H_Y(a, 0) = a + 1$  by the proof of Theorem 8.4. It follows from Theorem 8.4 that  $\delta(\mathcal{C}_{\alpha,Y}) = c_2(c_1 - a)$ , since  $c = a$ .

**Case II:** Let  $c \geq c_1$ .

If  $a = c_1 - 1$ , then  $b' = b$  and  $\dim_{\mathbb{K}} \mathcal{C}_{\alpha,Y} = H_Y(\alpha) = c_1(b + 1)$ . Furthermore, we have  $\delta(\mathcal{C}_{\alpha,Y}) = c_2 - b' = c_2 - b$ .

If  $a < c_1 - 1$ , then  $b' = b - 1$  and  $\dim_{\mathbb{K}} \mathcal{C}_{\alpha,Y} = H_Y(\alpha) = c_1b + c + 1 - \ell b = c_1b + 1 + a$ . Moreover, we have  $\delta(\mathcal{C}_{\alpha,Y}) = c_2 - b' = c_2 - b + 1$ .

The final claim follows easily.  $\square$

**Example 8.6.** *Let  $X = \mathcal{H}_2$  over  $\mathbb{F}_{13}$ . Let  $c_1 = 2$ ,  $c_2 = 6$ . Then the parameters of the code are given in Table 1.*

Table 1: Parameters of Toric Codes on  $X = \mathcal{H}_2$ .

$(a, b)$	$\alpha$	$[N, K, \delta]$	$(a, b)$	$\alpha$	$[N, K, \delta]$
$(0, 0)$	$(0, 0)$	$[12, 1, 12]$	$(1, 0)$	$(1, 0)$	$[12, 2, 6]$
$(0, 1)$	$(2, 1)$	$[12, 3, 6]$	$(1, 1)$	$(3, 1)$	$[12, 4, 5]$
$(0, 2)$	$(4, 2)$	$[12, 5, 5]$	$(1, 2)$	$(5, 2)$	$[12, 6, 4]$

$(0, 3)$	$(6, 3)$	$[12, 7, 4]$	$(1, 3)$	$(7, 3)$	$[12, 8, 3]$
$(0, 4)$	$(8, 4)$	$[12, 9, 3]$	$(1, 4)$	$(9, 4)$	$[12, 10, 2]$
$(0, 5)$	$(10, 5)$	$[12, 11, 2]$	$(1, 5)$	$(11, 5)$	$[12, 12, 1]$

## REFERENCES

- [1] Esmā Baran and Mesut Şahin. On parameterised toric codes. *Appl. Algebra Engrg. Comm. Comput.*, 2021.
- [2] Peter Beelen and Diego Ruano. The order bound for toric codes. In *Applied algebra, algebraic algorithms, and error-correcting codes*, volume 5527 of *Lecture Notes in Comput. Sci.*, pages 1–10. Springer, Berlin, 2009.
- [3] D. N. Bernstein. The number of roots of a system of equations. *Funkcional. Anal. i Priložen.*, 9(3):1–4, 1975.
- [4] Gavin Brown and Alexander M. Kasprzyk. Seven new champion linear codes. *LMS J. Comput. Math.*, 16:109–117, 2013.
- [5] David A. Cox. The homogeneous coordinate ring of a toric variety. *J. Algebraic Geom.*, 4(1):17–50, 1995.
- [6] David A. Cox, John B. Little, and Henry K. Schenck. *Toric varieties*, volume 124 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2011.
- [7] Eduardo Dias and Jorge Neves. Codes over a weighted torus. *Finite Fields Appl.*, 33:66–79, 2015.
- [8] David Eisenbud and Bernd Sturmfels. Binomial ideals. *Duke Math. J.*, 84(1):1–45, 1996.
- [9] Johan P. Hansen. Toric varieties Hirzebruch surfaces and error-correcting codes. *Appl. Algebra Engrg. Comm. Comput.*, 13(4):289–300, 2002.
- [10] Nathan Jacobson. *Basic algebra. I*. W. H. Freeman and Company, New York, second edition, 1985.
- [11] David Joyner. Toric codes over finite fields. *Appl. Algebra Engrg. Comm. Comput.*, 15(1):63–79, 2004.
- [12] A. G. Kušnirenko. Newton polyhedra and Bezout’s theorem. *Funkcional. Anal. i Priložen.*, 10(3):82–83, 1976.
- [13] John B. Little. Remarks on generalized toric codes. *Finite Fields Appl.*, 24:1–14, 2013.
- [14] Hiram H. López and Rafael H. Villarreal. Computing the degree of a lattice ideal of dimension one. *J. Symbolic Comput.*, 65:15–28, 2014.
- [15] Diane Maclagan and Gregory G. Smith. Uniform bounds on multigraded regularity. *J. Algebraic Geom.*, 14(1):137–164, 2005.
- [16] José Martínez-Bernal, Yuriko Pitones, and Rafael H. Villarreal. Minimum distance functions of graded ideals and Reed-Muller-type codes. *J. Pure Appl. Algebra*, 221(2):251–275, 2017.
- [17] Ezra Miller and Bernd Sturmfels. *Combinatorial Commutative Algebra*. Cambridge Studies in Advanced Mathematics. Springer-Verlag New York, 2005.
- [18] Marcel Morales and Apostolos Thoma. Complete intersection lattice ideals. *J. Algebra*, 284(2):755–770, 2005.
- [19] Esmā Baran Özkan. Vanishing ideals of parameterized subgroups in a toric variety, 2021.
- [20] Carlos Rentería-Márquez, Aron Simis, and Rafael H. Villarreal. Algebraic methods for parameterized codes and invariants of vanishing ideals over finite fields. *Finite Fields Appl.*, 17(1):81–104, 2011.
- [21] Diego Ruano. On the structure of generalized toric codes. *J. Symbolic Comput.*, 44(5):499–506, 2009.
- [22] Ivan Soprunov. Toric complete intersection codes. *J. Symbolic Comput.*, 50:374–385, 2013.
- [23] Mesut Şahin. Toric codes and lattice ideals. *Finite Fields Appl.*, 52:243–260, 2018.
- [24] Mesut Şahin and Ivan Soprunov. Multigraded Hilbert functions and toric complete intersection codes. *J. Algebra*, 459:446–467, 2016.

DEPARTMENT OF MATHEMATICS, HACETTEPE UNIVERSITY, ANKARA, TURKEY  
 Email address: mesut.sahin@hacettepe.edu.tr