

Solving Tolman-Oppenheimer-Volkoff equations in $f(T)$ gravity: a novel approach

H.G.M. Fortes,^{a,1} and J.C.N. de Araujo^a

^aDivisão de Astrofísica, Instituto Nacional de Pesquisas Espaciais,
Avenida dos Astronautas 1758, São José dos Campos, SP 12227-010, Brazil

E-mail: hemily.gomes@inpe.br, jcarlos.dearaujo@inpe.br

Abstract. The torsion models have stood out among the proposals for an alternative description of gravity. The simplest of them, the Teleparallel theory, is equivalent to General Relativity and there are many studies that seek to study its extension to more general functions of the torsion T . The purpose of our study is to consider a family of $f(T)$ models and apply their corresponding Tolman-Oppenheimer-Volkof equations to compact objects such as neutron stars. Consequently, through a numerical analysis, calculate, among other things, the maximum mass allowed by the model for a neutron star, which would also allow us to evaluate which models are in accordance with observations. In the present paper, the first in the series, we show explicitly the set of equations that must be solved, and how to solve it, in order to model compact stars in $f(T)$ gravity without the need to adopt any particular form for the metric functions or consider any perturbative approach, as has been done in some works in the literature.

¹Corresponding author.

Contents

1	Introduction	1
2	The basic equations of $f(T)$ gravity for spherically symmetric metric	2
3	Compact stars on $f(T)$ models	4
4	Numerical examples with polytropics	7
5	Final Remarks	8
A	Recovering TOV GR	10

1 Introduction

The theory of General Relativity (GR) with a cosmological constant has offered an explanation for the accelerated expansion of the universe in the context of the Λ CDM model, where it is also assumed the existence of the so-called dark matter whose properties are still unknown. In addition to the lack of knowledge about dark matter, there is also the problem of the cosmological constant whose value is very small compared to the predictions of quantum field theory.

In an attempt to circumvent the theoretical impasses and offer an alternative explanation for the observed acceleration, it is necessary to make changes in the fundamental theory. In this sense, there have been several proposals for alternative theories of gravitation based on modifications of the GR, for example, adding to the Lagrangian terms of mass or non-linear terms in curvature, among others.

The so-called Teleparallel theory of Gravity [1]-[3] is known to be equivalent to GR, with the basic difference of having in its formulation the torsion scalar T instead of the curvature scalar R .

Thus, we can consider changes in both GR and Teleparallel theory in order to build alternative models of gravitation. An important family of modifications of the Einstein-Hilbert action are the theories $f(R)$ where we have a function of the scalar of curvature as Lagrangian density. However, the field equations of these models are in general fourth order differential equations, which makes their analysis difficult. It is possible, then, to make equivalent modifications in the Teleparallel theory considering the Lagrangian density as a function of the torsion scalar. These models, in turn, have the advantage of taking us to second order field equations, therefore, simpler than those obtained in $f(R)$. It is worth mentioning that the theory $f(T)$ has presented interesting cosmological and astrophysical solutions, providing alternative interpretations for the acceleration phases of the universe [4]-[7].

Tolman-Oppenheimer-Volkoff (TOV) [8] equations are derived from the theory of gravity and can be used to model the structure of a spherically symmetrical object that is in hydrostatic equilibrium. From the Einstein-Hilbert action of GR, these equations are obtained and can be used to describe, for example, neutron stars. With this, it is possible to determine, among other things, the maximum mass allowed for this type of star for a given equation

of state. In this sense, considering different gravity models could, in principle, lead us to different maximum masses, see [9]. This would be important, for example, to explain one of the components involved in the event GW190814 [10] concerning the coalescence involving a black hole of 22.2–24.3 M_{\odot} and a compact object of mass 2.50–2.67 M_{\odot} , the latter being a relatively high-mass neutron star or a low-mass black hole. Therefore, it is evident the great relevance of considering alternative models of gravitation, such as the $f(T)$ models in the description of compact objects.

The goal of our study is to propose a specific family of $f(T)$ models and consider their applications in some astrophysical contexts, namely, the description of compact objects such as neutron stars. Such models must present solutions for these objects with a mass-radius relation that is in agreement with the observations. For this, it is necessary to obtain the TOV equations corresponding to the model in question and look for solutions of these equations. The proposal is to obtain these solutions for neutron stars from a more general approach than those already presented in the literature [11]–[13].

The TOV equations for $f(T)$ models will depend on the form assumed for this function and, together with the equation of state of neutron star, lead us to a system of coupled differential equations of non-trivial resolution. Thus, many authors choose to consider perturbative cases, approximations, assume conditions on the parameters of the theory or restrict themselves to very simple forms for $f(T)$, which is avoided with the approach presented in this paper.

It is worth mentioning that in [14], the authors seem to be among the few who solve the problem appropriately for a particular $f(T)$, although they do not explicitly show the basic equations and how to solve them.

In the present paper, the first in the series, we explicitly show the set of equations that must be solved, and how to solve it, in order to model compact stars in $f(T)$ gravity without the need to adopt any particular form for the metric functions or consider any perturbative approach. This is an essential procedure which allows the readers to extend more easily the approach to other $f(T)$. Section 2 considers the basic equations, in Section 3 we present the equations to be solved to model compact stars in $f(T)$ gravity, in Section 4 a numerical example is presented and the final remarks are considered in Section 5.

2 The basic equations of $f(T)$ gravity for spherically symmetric metric

Theories with torsion are formulated from an alternative dynamical variable $h^a{}_{\mu}$, the tetrad, which is a set of four vectors that define an inertial frame at each point [3]. The metric $g_{\mu\nu}$, usually adopted in models with curvature, and the tetrad $h^a{}_{\mu}$ are related as follows:

$$g_{\mu\nu} = h^a{}_{\mu} h^b{}_{\nu} \eta_{ab} , \quad (2.1)$$

where $\eta_{ab} = \text{diag}(+1, -1, -1, -1)$ is the Minkowski metric of the tangent space.

Gravity theories constructed from the metric will always be invariant under local Lorentz transformations. On the other hand, torsion models will not necessarily be, which will generally depend on a choice of tetrads and appropriate spin connection. In addition, a non-invariant theory will be sensitive to the choice of the tetrad and may lead to different solutions. Thus, this choice is fundamental for the good behavior of the $f(T)$ models, which has been extensively studied in recent years [3, 15] and is not always taken into account when working with models with torsion.

This was so till the seminal paper [16], in which a covariant $f(T)$ gravity is formulated. Thus, the problems related to the violation of local Lorentz invariance and frame dependence could be resolved. To obtain a covariant $f(T)$ gravity they also considered the inertial spin connection as a dynamical variable. Recall that in the usual formulation (hereafter referred to original formulation) only the tetrad can be considered as dynamical variable.

It is also worth mentioning that in the covariant $f(T)$ formulation by [16] one is allowed to adopt an arbitrary tetrad and then obtain its corresponding spin connection. Thus, one does not need to be worried about bad or good tetrad, since all them lead to the same field equations. An interesting issue is how to obtain the spin connection, but since this is out of the scope of the present article, we refer the reader to [16] for detail. It is noteworthy that although the covariant formulation is mentioned here, it is not used in any derivation in the present paper.

For the purpose of this article, we consider the $f(T)$ models in spherically symmetric spacetime [17]-[24] derived from the original formulation. A suitable tetrad in this formulation reads [15]:

$$h_{\mu}^a = \begin{pmatrix} e^{A/2} & 0 & 0 & 0 \\ 0 & e^{B/2} \sin \theta \cos \phi & 0 & 0 \\ 0 & -r (\cos \theta \cos \phi \sin \gamma + \sin \phi \cos \gamma) & 0 & 0 \\ 0 & r \sin \theta (\sin \phi \sin \gamma - \cos \theta \cos \phi \cos \gamma) & 0 & 0 \\ & & e^{B/2} \sin \theta \sin \phi & e^{B/2} \cos \theta \\ & & r (\cos \phi \cos \gamma - \cos \theta \sin \phi \sin \gamma) & r \sin \theta \sin \gamma \\ & & -r \sin \theta (\cos \theta \sin \phi \cos \gamma + \cos \phi \sin \gamma) & r \sin^2 \theta \cos \gamma \end{pmatrix}, \quad (2.2)$$

where A , B , θ , ϕ are those from the spherically symmetric metric $ds^2 = e^{A(t,r)} dt^2 - e^{B(t,r)} dr^2 - r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2$, and $\gamma = \gamma(r)$ is one of the Euler's angles [15, 25].

Since [15] was one of the first papers to obtain the correct field equations for spherically symmetrical spacetimes, we will adopt it to obtain the basic equations used here to model compact stars in $f(T)$ gravity.

The torsion tensor is built from the antisymmetry part of the connection only,

$$T^{\sigma}{}_{\mu\nu} = \Gamma^{\sigma}{}_{\nu\mu} - \Gamma^{\sigma}{}_{\mu\nu} = h_i^{\sigma} (\partial_{\mu} h^i{}_{\nu} - \partial_{\nu} h^i{}_{\mu}), \quad (2.3)$$

since the spin connection is null in the original formulation.

By using the tetrad (2.2), we can write $T(r)$ as:

$$T(r) = \frac{2e^{-B}}{r^2} \left[1 + e^B + 2e^{B/2} \sin \gamma + 2e^{B/2} r \gamma' \cos \gamma + r A' \left(1 + e^{B/2} \sin \gamma \right) \right], \quad (2.4)$$

where the prime corresponds to derivative with respect to r .

From the action for the $f(T)$ theory,

$$S \propto \int d^4x \det(h_{\mu}^a) f(T), \quad (2.5)$$

we can obtain the following field equations:

$$4\pi P = -\frac{f}{4} + \frac{f_T e^{-B}}{4r^2} (2 - 2e^B + r^2 e^B T - 2r A') \quad (2.6)$$

$$4\pi\rho = \frac{f}{4} - \frac{f_T e^{-B}}{4r^2} (2 - 2e^B + r^2 e^B T - 2r B') - \frac{f_{TT} T' e^{-B}}{r} (1 + e^{B/2} \sin \gamma) \quad (2.7)$$

$$f_{TT} T' \cos \gamma = 0 \quad (2.8)$$

$$f_T \dot{B} = 0 \quad (2.9)$$

$$\begin{aligned} \dot{B} \left[2 f_{TT} (1 + e^{B/2} \sin \gamma) (2 - 2e^B + r^2 e^B T + 2r A') + e^B r^2 f_T \right] + \\ - 4r f_{TT} \dot{A}' (1 + e^{B/2} \sin \gamma)^2 = 0 \end{aligned} \quad (2.10)$$

$$\begin{aligned} f_{TT} \left[-4e^A r T' - \dot{B}^2 (2 - 2e^B + r^2 e^B T) - 2r A' (e^A r T' + \dot{B}^2) \right] + \\ + 4r \dot{B} \dot{A}' (1 + e^{B/2} \sin \gamma) \left. \right] + f_T \left[4e^A - 4e^A e^B - e^A r^2 A'^2 + 2e^A r B' + \right. \\ \left. + e^A r A' (2 + r B') - 2r^2 e^A A'' - e^B r^2 \dot{A} \dot{B} + e^B r^2 \dot{B}^2 + 2e^B r^2 \ddot{B} \right] = 0 \end{aligned} \quad (2.11)$$

where $f_T = \frac{\partial f}{\partial T}$ and $f_{TT} = \frac{\partial^2 f}{\partial T^2}$ and the overdots correspond to differentiation with respect to t . The source term is given by an isotropic perfect fluid, with pressure P and energy density ρ . Additionally, the well-known conservation equation reads [25]:

$$A' = -2 \frac{P'}{P + \rho}, \quad (2.12)$$

which will be very useful for our purposes later on.

3 Compact stars on $f(T)$ models

Since we are interested to model compact stars in hydrostatic equilibrium in $f(T)$ gravity, A and B , and consequently all other quantities such as P and ρ , do not depend on time. As a result, equations (2.9) and (2.10) are identically null, and several terms in equation (2.11) are taken equal to zero. The resulting equations are complicated anyway and it does not seem possible to get a set of equations similar in structure to the TOV equations that come from General Relativity.

In fact, there are papers in the literature related to compact stars in $f(T)$ gravity [11–13, 26] in which very complicated equations are obtained for different tetrads, as well as for some *ansatz* for B and or T , for example.

The main objective here is, then, to obtain the system of differential equations adequate for this type of analysis and, without assuming restrictions for the functions $A(r)$, $B(r)$, etc., find their solutions without approximations or perturbative calculations.

In principle, we are going to consider, as an example, $f(T)$ given by the following form

$$f(T) = T + \xi T^2, \quad (3.1)$$

where ξ is an arbitrary real. However, the procedure could be repeated later on for even more general forms of $f(T)$.

With this specific choice for $f(T)$, taking $\gamma = -\pi/2^1$ in order to get rid of the restriction on $f(T)$ from (2.8) and by considering the hydrostatic equilibrium ($\dot{B} = 0$, $\dot{A} = 0$), equations (2.7-2.11) can be rewritten as:

$$\left\{ \frac{(1 - e^B)}{2} - 4\pi P r^2 e^B + \frac{\xi}{r^2} (6 + 3e^{-B} - e^B - 8e^{-B/2}) + \frac{1}{2r} \left[r^2 + 12\xi (1 + e^{-B} - 2e^{-B/2}) \right] A' + \xi (3e^{-B} - 4e^{B/2} + 1) A'^2 \right\} e^{-B} = 0 \quad (3.2)$$

$$\left\{ -\frac{(1 - e^B)}{2r^2} - 4\pi \rho e^B + \frac{\xi}{r^4} (18 + e^B + 5e^{-B} - 8e^{B/2} - 16e^{-B/2}) + \frac{1}{2r^3} \left[r^2 + 12\xi (1 + e^{-B} - 2e^{-B/2}) \right] B' - \frac{4\xi}{r^2} (1 + e^{-B} - 2e^{-B/2}) A'' + \frac{2\xi}{r^2} (1 + 3e^{-B} - 4e^{-B/2}) B' A' - \frac{\xi}{r^2} (1 + e^{-B} - 2e^{-B/2}) A'^2 \right\} e^{-B} = 0 \quad (3.3)$$

$$\left\{ -2 \left[r^2 + 4\xi (1 + 3e^{-B} - 4e^{-B/2}) \right] A'' + \frac{2}{r} \left[r^2 + 4\xi (1 + 3e^{-B} - 5e^{-B/2}) \right] B' + \frac{4}{r^2} \left[r^2 (1 - e^B) + 4\xi (2 - e^B + 3e^{-B} + 2e^{B/2} - 6e^{-B/2}) \right] + \left[16\xi r e^{-B/2} (1 - e^{-B/2}) A'' + \left\{ r^2 + 4\xi (1 + 9e^{-B} - 8e^{-B/2}) \right\} B' + \frac{2}{r} \left\{ r^2 + 4\xi (1 + 7e^{-B} + 2e^{B/2} - 10e^{-B/2}) \right\} A' + \left[4\xi r e^{-B/2} (3e^{-B/2} - 2) B' - \left\{ r^2 + 4\xi (1 - 3e^{-B} + 2e^{-B/2}) \right\} A'^2 + -4\xi r e^{-B/2} (e^{-B/2} - 1) A'^3 \right\} e^A = 0 \quad (3.4)$$

Notice that the equations obtained are not simple since we have nontrivial coupled differential equations which mix all the functions and their derivatives, namely, B , B' , A' , and A'' . Some authors have preferred to consider some approximations, restriction of the functions or perturbative calculations even before getting the system of equations. So, equations (3.2)-(3.4) had never been considered explicitly in the literature so far.

The next step is to obtain an analytical solution which gives us explicitly P and other functions, such as A' or B' , so that they can be numerically integrated. After some manipulations and with the help of mathematical software in order to deal with the complex expressions, we were able to reach the following solution:

$$P = \frac{e^{-B}}{8\pi r^4} \left\{ r^2 (1 - e^B) + r^3 A' + 12r (e^{-1/2 B} - 1)^2 \xi A' + 2e^B (e^{-1/2 B} - 1)^3 (3e^{-1/2 B} + 1) \xi + 2r^2 \xi (e^{-1/2 B} - 1) (3e^{-1/2 B} - 1) A'^2 \right\}, \quad (3.5)$$

¹The specific choice $\gamma = -\pi/2$ will allow us to reproduce the TOV GR equations in the $\xi = 0$ limit.

$$\begin{aligned}
B' = & -\frac{e^B}{r} \left\{ r^4 (1 - e^{-B} - 8\pi r^2 \rho) - 32\pi r^4 \rho (e^{-1/2B} - 1)(3e^{-1/2B} - 1)\xi + \right. \\
& -2r^2 (e^{-1/2B} - 1)^2 (9e^{-B} + 10e^{-1/2B} - 11)\xi - 8(e^{-1/2B} - 1)^4 (9e^{-B} + 8e^{-1/2B} - 9)\xi^2 + \\
& -8r^3 e^{-1/2B} (e^{-1/2B} - 1) \left[8\pi r^2 \rho + (e^{-1/2B} - 1)(2e^{-1/2B} + 1) \right] A'\xi + \\
& -16r e^{-1/2B} (e^{-1/2B} - 1)^3 (9e^{-1/2B} - 5) A'\xi^2 + 2r^4 e^{-B} (e^{-1/2B} - 1)^2 A'^2 \xi + \\
& \left. -8r^2 e^{-B} (e^{-1/2B} - 1)^3 (9e^{-1/2B} + 1) A'^2 \xi^2 \right\} / \left\{ r^4 + 8r^2 (e^{-1/2B} - 1)(2e^{-1/2B} - 1)\xi + \right. \\
& +16(e^{-1/2B} - 1)^3 (3e^{-1/2B} - 1)\xi^2 + 16r^3 e^{-1/2B} (e^{-1/2B} - 1) A'\xi + \\
& \left. +32r e^{-1/2B} (e^{-1/2B} - 1)^2 (3e^{-1/2B} - 2) A'\xi^2 + 48r^2 e^{-B} (e^{-1/2B} - 1)^2 A'^2 \xi^2 \right\}, \quad (3.6)
\end{aligned}$$

and

$$\begin{aligned}
A'' = & e^B \left\{ r^4 (8\pi r^2 \rho + 3e^{-B} - 3) - 2r^2 (64e^{-1/2B} \pi r^2 \rho - 48e^{-B} \pi r^2 \rho - 16\pi r^2 \rho + \right. \\
& -48e^{-1/2B} - 25e^{-2B} + 40e^{-3/2B} + 14e^{-B} + 19)\xi + 8(-13 - 76e^{-5/2B} + \\
& +60e^{-1/2B} + 81e^{-2B} + 21e^{-3B} - 89e^{-B} + 16e^{-3/2B})\xi^2 + \left[\frac{1}{2} r^5 (8\pi r^2 \rho + 3e^{-B} - 1) + \right. \\
& -r^3 (128e^{-1/2B} \pi r^2 \rho - 144e^{-B} \pi r^2 \rho - 16\pi r^2 \rho - 64e^{-1/2B} + 34e^{-B} + \\
& -77e^{-2B} + 96e^{-3/2B} + 11)\xi + 4r(111e^{-3B} - 344e^{-5/2B} + 88e^{-1/2B} + \\
& -199e^{-B} + 305e^{-2B} + 48e^{-3/2B} - 9)\xi^2 \Big] A' + \left[-\frac{1}{2} e^{-B} r^6 - 4r^4 (2e^{-B} + \right. \\
& +8e^{-1/2B} \pi r^2 \rho - 12e^{-B} \pi r^2 \rho - e^{-1/2B} - 5e^{-2B} + 4e^{-3/2B})\xi + \\
& +4r^2 (10e^{-1/2B} + 174e^{-2B} + 93e^{-3B} - 242e^{-5/2B} + 16e^{-3/2B} - 51e^{-B})\xi^2 \Big] A'^2 + \\
& + \left[-r^5 (7e^{-2B} - 8e^{-3/2B} + e^{-B})\xi + 4r^3 (24e^{-2B} + 21e^{-3B} - 44e^{-5/2B} - e^{-B})\xi^2 \right] A'^3 + \\
& \left. -12r^4 (e^{-3B} + e^{-2B} - 2e^{-5/2B})\xi^2 A'^4 \right\} / r^2 \left\{ r^4 + (8r^2 - 24e^{-1/2B} r^2 + 16e^{-B} r^2)\xi + \right. \\
& + (16 - 96e^{-1/2B} + 192e^{-B} - 160e^{-3/2B} + 48e^{-2B})\xi^2 + \left[16r^3 (e^{-B} - e^{-1/2B})\xi + \right. \\
& \left. -32r(2e^{-1/2B} - 7e^{-B} + 8e^{-3/2B} - 3e^{-2B})\xi^2 \right] A' + 48r^2 (e^{-B} - 2e^{-3/2B} + e^{-2B})\xi^2 A'^2 \Big\}. \quad (3.7)
\end{aligned}$$

Notice that, although we have obtained the expression just above for A'' in terms of the independent variables, one does not necessitate to use it, since it will be more useful to consider alternatively the conservation equation (2.12) which relates A' , P' and ρ instead, since the latter is a much simpler expression.

If ρ is related to the mass as in the TOV RG, then one would have:

$$\frac{dm}{dr} = 4\pi\rho r^2. \quad (3.8)$$

It is worth mentioning, however, that equation (3.8) may not apply to $f(T)$. This is still an open problem in such a theory (see, e.g., [14] and [22]). Although, in a recent paper on

boson stars in $f(T)$ [23], the authors also adopted equation (3.8) to obtain the mass of such stars.

Having in mind that it is not an observable, the total rest mass M_0 is an interesting quantity to calculate, since it can be used to make comparisons between different theories. For example, for a given central density and a given equation of state, different theories must provide different values of M_0 .

Recall that the total rest mass M_0 is obtained by solving

$$\frac{dm_0}{dr} = 4\pi\rho_0 e^{B/2}r^2, \quad (3.9)$$

where ρ_0 is the rest mass density and $4\pi e^{B/2}r^2 dr$ is the proper volume element.

One clearly sees that for $f(T) = T$, one should recover the TOV equations that come from General Relativity (see A for detail).

In summary, to model stars on the particular $f(T)$ gravity adopted here for a given equation of state (EOS), one needs to integrate the following system of differential equations: (2.12), (3.5), (3.6) and (3.8).

As we are interested in modelling neutron stars, we will need to consider their EOS, which can be quite complex. For this reason, it is common to start by considering the so-called polytropic equations instead of the realistic ones. In general, the polytropic offers a very consistent description of these objects [27]. For more information and examples of neutron star state equations, see [28]-[30].

4 Numerical examples with polytropics

The main proposal here is to show that, despite the apparent complexity of the equations obtained in last section, we can apply them to cases of physical interest.

As an practical example, we consider an specific polytropic equation of state [27], since it can offer a simpler and satisfactory description for neutron stars, for example.

Recall that the polytropic reads

$$P = k \rho_o^\gamma, \quad (4.1)$$

where ρ_o is the rest-mass density, k is the polytropic gas constant and n defined by $\gamma \equiv 1+1/n$ is the polytropic index. From the first law of thermodynamics one obtains the mass-energy density, ρ , namely $\rho = \rho_o + n P$ (see, e.g., [31]).

We can proceed by solving equations (2.12), (3.5), (3.6) and (3.8) to model a compact star for the $n = 1$ polytropic equation of state, $P = \rho_o^2$, where we have set all quantities in nondimensional form, following [31]. In practice, this is equivalent to set $k = G = c = 1$. This is a useful procedure since our aim here is to compare how different $f(T)$ is as compared to General Relativity regarding compact objects.

In order to perform the numerical integration, we have written a numerical code in Python. The procedure is essentially to integrate numerically the differential equations for $m(r)$ and $P(r)$ for a given EOS. Then, it is necessary to choose a central density, ρ_c , and using the EOS one obtains the central pressure P_c . In addition, providing the central boundary conditions

$$m = 0 \quad \text{and} \quad P = P_c \quad \text{at} \quad r = 0, \quad (4.2)$$

one obtains $m(r)$, $P(r)$ and $\rho(r)$, i.e., the structure of the star. The radius of the star, R , is given by the value of r for which $P(r) = 0$. That is, one starts the integration of the set of

differential equations at $r = 0$ and continues it till the value of r for which $P(r) = 0$. Finally, the mass of the star is then given by $M = m(R)$.

Here, to model a compact star one has an additional equation in comparison to General Relativity, namely, the differential equation for $B(r)$. Then, one needs to provide a central boundary condition for B . One can convince oneself that $B_c = 0$.

In figure 1, we can compare different models of gravity from $f(T) = T + \xi T^2$. For a range of values of ρ_c one obtains the corresponding masses and radii. Consequently, one obtains the “Mass \times Radius” and “Mass $\times \rho_c$ ” curves, from which one can obtain, for example, the maximum mass allowed for a given EOS. For $\xi = 0$, we have $f(T) = T$, which is nothing but the TEGR, whose TOV equations after a simple manipulation are reduced to TOV GR.

Note that for $\xi \geq 0$ there are maximum masses. In addition, the greater ξ is, the lower the maximum mass. Considering now that ξ could have negative values, one clearly see that, as compared to GR, the masses for a given ρ_c increases for decreasing values of ξ .

Notice that this very polytropic was studied by [14] and a comparison with the models we have in common shows that the results are in perfect agreement. For example: for $\xi = 0.05$, which corresponds to the model a = 0.1 shown in Table 1 of Ref. [14], we obtain $R = 0.820$ and $M_0 = 0.163$, whereas they obtain $R = 0.819$ and $M_0 = 0.165$, respectively. It is noteworthy that these authors adopted the covariant formulation and wrote the set of differential equations to be solved differently from what we did here, so our approaches are not the same, although they are equivalent. Consequently, comparing our results is very timely, as it validates both approaches.

In figure 2 are shown “Mass \times Radius” and “Mass $\times \rho_c$ ” sequences, respectively, now for a polytropic with $n = 3/2$, i.e., $P = \rho_0^{5/3}$ (non-relativistic Fermi gas). As can be seen, the behaviour is qualitatively the same as those for $n = 1$ ($P = \rho_0^2$) polytropic.

It is worth mentioning that we consider TOV in $f(T) = T + \xi T^2$ in two other articles. We refer the reader to these papers for further models in this particular $f(T)$. In one of them we study other polytropics [32]. In the other one, we study some realistic equations of state in modeling neutron stars [33].

5 Final Remarks

Modeling spherical stars has been a much-addressed topic in the recent literature. At the same time, alternative descriptions of gravity have been highly pursued due the open problems in this area. In view of that, it is always productive offer different approaches do deal with the problem and even more useful if the procedure is quite general and able to be reproduced for other proposals. The novel approach presented here, despite the considerable equations that appear, can be exactly repeated for other models. Additionally, one of our concerns was to explicit each step of the method in order to facilitate future works with the same approach, in contrast of some authors who have adopted effective strategies but do not detail the procedure sufficiently.

As we showed in Section 3, for the particular case in which $f(T) = T + \xi T^2$, the equations to be solved are very complicated. Nevertheless, the correspondent TOV equations could be obtained without any restrictions. It is worth stressing that we adopted here a particular form for $f(T)$ as an example, but since the procedure considered here is quite general, any $f(T)$ can be in principle considered. Obviously, one is also free to choose the equation of state.




Figure 1. Right (left): sequences of mass-energy M vs. central mass-energy density ρ_c (radius R) for $P = \rho_0^2$ and different values of ξ .




Figure 2. The same as figure 1 now for $P = \rho_0^{5/3}$.

For the present choice of $f(T)$, one can then solve equations (2.12), (3.5), (3.6) and (3.8)², to model a compact star for a given equation of state. Notice that in the way these equations are obtained, one does not need to assume beforehand any particular form for A , B and T , as some papers in the literature do.

One could ask oneself why this particular set of equations has been chosen. Notice also that in the present context, by setting $\xi = 0$ into equations (3.5) and (3.6), one obtains (A.1) and (A.2), respectively. Therefore, one could think of equations (3.5) and (3.6) as a kind of generalisation of (A.1) and (A.2). It is worth mentioning that, if we use (2.12), equation (3.7) is no longer necessary since it does not contain any new information. Moreover, it can well be obtained by a combination of the other equations that in fact will be resolved.

Another interesting issue concerning the equations is that the combination of equations (2.12) and (3.5) is a generalisation of equation (A.7). It was precisely this fact that has allowed the resolution of the system in the current approach without any further complications.

Also, it is worth noticing that $e^{-B(r)} = 1 - 2m(r)/r$ in TOV GR, whereas here one has to numerically integrate equation (3.6) to obtain $B(r)$. This integration can be performed without any kind of approximation.

Once the equation of state has been chosen, the first numerical test to be performed would be with $\xi = 0$, since, for this value, we should retrieve results already known from the RG TOV equations, for example, the maximum mass allowed for a neutron star for the chosen equation of state. Then, different values of ξ should be taken, leading to different $f(T)$ gravity models applied to neutron stars.

Then, using the most appropriate Python integrator, we can implement a program that, starting from a given central density for the neutron star and the specific equation of state, could lead us to the structure of the star, i.e., P , ρ and m as a function of r , as was done in the previous section for the $n = 1$ and $3/2$ polytropics. This allows us, among other things, to obtain the value of r for which the pressure P vanishes, which would correspond to the external limit of the star, i.e., its radius R and, consequently, the mass $M = m(R)$ is also obtained. Repeating the procedure for different central densities, it is possible to obtain a curve of the type “*Mass vs. Radius*”, which would indicate the maximum mass allowed by the model for a given equation of state.

Due to the complexity of the differential equations, all this process of finding the best integrator and performing the numerical analysis is quite elaborate, but it has been shown perfectly possible to obtain its numerical solution, as we have seen from the examples considered previously. Analysis for more general equations of state will be presented in future works to appear elsewhere, either for polytropic and realistic equations of state.

A Recovering TOV GR

Although this discussion appears in related studies, it is worth showing explicitly that one can recover TOV GR by setting $f(T) = T$. As is well known, such a particular case is nothing but the teleparallel equivalent of GR.

By substituting $f(T) = T$ into equations (2.6) and (2.7), we obtain

$$8\pi pr^2 = -1 + e^{-B} + r A' e^{-B} \tag{A.1}$$

²We are considering exactly the same mass-energy equation of TOV GR, having in mind that this is still an open problem in $f(T)$ gravity.

and

$$8\pi\rho r^2 = 1 - e^{-B} + r B' e^{-B} \quad (\text{A.2})$$

Integrating equation (A.2) one obtains

$$e^{-B} = 1 - \frac{2m}{r}, \quad (\text{A.3})$$

where

$$\frac{dm}{dr} = 4\pi\rho r^2. \quad (\text{A.4})$$

Using these last two equations in (A.1), one obtains

$$A' = 2 \frac{4\pi\rho r^3 + m}{r^2 - 2mr} \quad (\text{A.5})$$

Since from the *Conservation Equation* (see, e.g., Ref. [25]) one has

$$A' = -2 \frac{p'}{p + \rho}, \quad (\text{A.6})$$

the equation for the hydrostatic equilibrium reads

$$p' = -(p + \rho) \frac{4\pi\rho r^3 + m}{r^2 - 2mr}. \quad (\text{A.7})$$

Equations (A.3) - (A.7), as expected, are the equations one has in general relativity, namely, TOV GR.

Acknowledgments

J.C.N.A. thanks FAPESP (2013/26258 – 4) and CNPq (308367/2019-7) for partial financial support. The authors would like to thank Rafael da Costa Nunes for discussions related to the $f(T)$ theory.

References

- [1] R. Aldrovandi and J. G. Pereira, *Teleparallel Gravity: An Introduction*, Springer, Dordrecht - Heidelberg - New York - London (2013).
- [2] S. Bahamonde et al. *Teleparallel Gravity: From Theory to Cosmology* (2021) arXiv:2106.13793 [gr-qc].
- [3] Y. Cai et al., *$f(T)$ teleparallel gravity and cosmology*, *Rept. Prog. Phys.* **79** (2016), 106901.
- [4] R. Ferraro, F. Fiorini, *Modified teleparallel gravity: Inflation without inflaton*, *Phys. Rev. D* **75**, (2007) 084031 [gr-qc/0610067].
- [5] E. V. Linder, *Einstein's Other Gravity and the Acceleration of the Universe*, *Phys. Rev. D* **81**, (2010) 127301 [arXiv:1005.3039].
- [6] R. Myrzakulov, *Accelerating universe from $F(T)$ gravity*, *Eur. Phys. J. C* **71** (2011) 1752 [arXiv:1006.1120].
- [7] K. Karami, A. Abdolmaleki, *$f(T)$ modified teleparallel gravity models as an alternative for holographic and new agegraphic dark energy models*, *Res. Astron. Astrophys.* **13** (2013) 757.

- [8] C. M. Will, *Theory and experiment in gravitational physics*, Cambridge University Press (1993).
- [9] R. C. Nunes, J. G. Coelho, J. C. N. de Araujo, *Weighing massive neutron star with screening gravity: A look on PSR J0740+6620 and GW190814 secondary component*, *Eur. Phys. J. C* **80** (2020) 11115 [arXiv:2008.10395].
- [10] R. Abbott et al., *GW190814: Gravitational Waves from the Coalescence of a 23 Solar Mass Black Hole with a 2.6 Solar Mass Compact Object*, *The Astrophysical Journal* **896** (2020) L44 [arXiv:2006.12611].
- [11] M. G. Ganiou et al., *Strong magnetic field effects on neutron stars within $f(T)$ theory of gravity*, *Eur. Phys. J. Plus* **132** (2017) 250.
- [12] A. V. Kpadonou, M. J. S. Houndjo, M. E. Rodrigues, *Tolman-Oppenheimer-Volkoff Equations and their implications for the structures of relativistic Stars in $f(T)$ gravity*, *Astrophys. Space Sci.* **361** (2016) 244.
- [13] M. Pace, J. L. Said, *A Perturbative Approach to Neutron Stars in $f(T, T)$ -Gravity*, *Eur. Phys. J. C* **77** (2017) 283.
- [14] S. Ilijić, M. Sossich, *Compact stars in $f(T)$ extended theory of gravity*, *Phys. Rev. D* **98** (2018) 064047.
- [15] N. Tamanini and C. G. Böhmer, *Good and bad tetrads in $f(T)$ gravity*, *Phys. Rev. D* **86** (2012) 044009.
- [16] M. Krššák, E.N. Saridakis, *The covariant formulation of $f(T)$ gravity*, *Classical and Quantum Gravity* **33** (2016) 115009.
- [17] X.-h. Meng, Y.-b. Wang, *Birkhoff's theorem in $f(T)$ gravity*, *Eur. Phys. J. C* **71** (2011) 1755.
- [18] H. Dong, Y.-b. Wang, X.-h. Meng, *Extended Birkhoff's theorem in $f(T)$ gravity*, *Eur. Phys. J. C* **72** (2012) 2002.
- [19] H. Dong, Y.-b. Wang, X.-h. Meng, *Birkhoff's Theorem in $f(T)$ Gravity up to the Perturbative Order*, *Eur. Phys. J. C* **72** (2012) 2201.
- [20] C. G. Böhmer, A. Mussa, N. Tamanini, *Existence of relativistic stars in $f(T)$ gravity*, *Classical Quantum Gravity* **28** (2011) 245020.
- [21] R. Ferraro, F. Fiorini, *Spherically symmetric static spacetimes in vacuum $f(T)$ gravity*, *Phys. Rev. D* **84** (2011) 083518.
- [22] G.J. Olmo, D. Rubiera-Garcia, A. Wojnar, *Stellar structure models in modified theories of gravity: Lessons and challenges*, *Phys. Report* **876** (2020) 1.
- [23] S. Ilijić, M. Sossich, *Boson stars in $f(T)$ extended theory of gravity*, *Phys. Rev. D* **102** (2020) 084019.
- [24] L. Iorio, E. N. Saridakis, *Solar system constraints on $f(T)$ gravity*, *Mon. Not. Roy. Astron. Soc.* **427** (2012) 1555.
- [25] C. G. Böhmer, A. Mussa and N. Tamanini, *Existence of relativistic stars in $f(T)$ gravity*, *Class. Quantum Grav.* **28** (2011) 245020.
- [26] M. Pace, J. L. Said, *Quark stars in $f(T, T)$ -gravity*, *Eur. Phys. J. C* **77** (2017) 62.
- [27] L. Ferrari, P. C. R. Rossi, M. Malheiro, *A polytropic approach to neutron stars*, *International Journal of Modern Physics D* **19** (2010) 1569.
- [28] J. M. Lattimer, M. Prakash, *Neutron Star Structure and the Equation of State*, *Astrophys. J.* **550** (2001) 426.
- [29] J. M. Lattimer, M. Prakash, *Equation of state, neutron stars and exotic phases*, *Nucl. Phys. A* **777** (2006) 479.

- [30] J. S. Read et al., *Constraints on a phenomenologically parametrized neutron-star equation of state*, *Physical Review D* **79** (2009) 124032.
- [31] T.W. Baumgarte, S. L. Shapiro, "Numerical Relativity: Solving Einstein's Equations on the Computer", Cambridge University Press, 2010.
- [32] H.G.M. Fortes, J.C.N. de Araujo, *Solving Tolman-Oppenheimer-Volkoff equations in $f(T)$ gravity: a novel approach applied to polytropic equations of state*, arxiv:2105.09118.
- [33] H.G.M. Fortes, J.C.N. de Araujo, *Solving Tolman-Oppenheimer-Volkoff equations in $f(T)$ gravity: a novel approach applied to some realistic equations of state*, arxiv:2109.01155.