

# Exact Recovery in the General Hypergraph Stochastic Block Model

Qiaosheng Zhang, Vincent Y. F. Tan, *Senior Member, IEEE*

## Abstract

This paper investigates fundamental limits of exact recovery in the general  $d$ -uniform hypergraph stochastic block model ( $d$ -HSBM), wherein  $n$  nodes are partitioned into  $k$  disjoint communities with relative sizes  $(p_1, \dots, p_k)$ . Each subset of nodes with cardinality  $d$  is generated independently as an order- $d$  hyperedge with a certain probability that depends on the ground-truth communities that the  $d$  nodes belong to. The goal is to exactly recover the  $k$  hidden communities based on the observed hypergraph. We show that there exists a sharp threshold such that exact recovery is achievable above the threshold and impossible below the threshold (apart from a small regime of parameters that will be specified precisely). This threshold is represented in terms of a quantity which we term as the generalized Chernoff-Hellinger divergence between communities. Our result for this general model recovers prior results for the standard SBM and  $d$ -HSBM with two symmetric communities as special cases. En route to proving our achievability results, we develop a polynomial-time two-stage algorithm that meets the threshold. The first stage adopts a certain hypergraph spectral clustering method to obtain a coarse estimate of communities, and the second stage refines each node individually via local refinement steps to ensure exact recovery.

## Index Terms

Community detection, hypergraph stochastic block model (HSBM), exact recovery, hypergraph spectral clustering methods.

## I. INTRODUCTION

The stochastic block model (SBM) [1] is a celebrated random graph model that has been widely studied for the community detection problem, and the objective therein is to partition  $n$  nodes into  $k$  disjoint communities (a.k.a., clusters) based on the randomly generated graph. The recent *award-winning*<sup>1</sup> papers [2], [3] discovered a phase transition phenomenon for *exact recovery* (i.e., all the nodes are required to be classified correctly) in the SBM with two symmetric communities. That is, there is a sharp threshold such that exact recovery is achievable above the threshold, and impossible below the threshold. This phase transition phenomenon was later extended to the *general SBM* with  $k \geq 2$  communities and without imposing symmetric structures [4], [5]. Popularized by these breakthroughs, community detection in the SBM and its variants have then received significant attention, and many works have progressively contributed to this field by considering the information-theoretic limits of some variants of the SBM [6]–[9], efficient algorithms with theoretical guarantees [10]–[17], the effect of side information [18]–[22], etc. We refer the readers to [23] for a comprehensive survey.

While most prior works focused on community detection on graphs, it is also of keen interest to study community detection on *hypergraphs*. This is because *higher-order* relational information among multiple nodes, which can naturally be captured by

Qiaosheng Zhang is with the Department of Electrical and Computer Engineering, National University of Singapore (e-mail: elezqiao@nus.edu.sg).

Vincent Y. F. Tan is with the Department of Electrical and Computer Engineering and Department of Mathematics, National University of Singapore (e-mail: vtan@nus.edu.sg).

<sup>1</sup>The Information Theory Society Paper Award, 2020.

hypergraphs, is ubiquitous in many applications. For example, friendships between users in social networks can be captured by graphs, but chat groups are usually represented by hyperedges in hypergraphs. Authors in co-authorship networks can also be connected by hyperedges. There are also applications in computer vision (such as object recognition and image registration) that are concerned with the *point-set matching* problem [24], [25], which aims to find strongly connected components in a uniform hypergraph. Motivated by these applications, in recent years some efforts have been expended to advance our understanding of community detection on hypergraphs. In particular, Ghoshdastidar and Dukkipati [26] first proposed a random hypergraph model called the *d-uniform hypergraph stochastic block model (d-HSBM)*, in which each subset of nodes with cardinality  $d$  is generated independently as an *order-d hyperedge* with a certain probability that depends on the communities that the  $d$  nodes belong to. Subsequent researchers further investigated the recovery limits of  $d$ -HSBMs by developing various hypergraph clustering algorithms (such as spectral clustering methods [27]–[33], semidefinite programming-based methods [34]–[36], tensor decomposition-based methods [37], approximate-message passing algorithms [38], [39], etc) with theoretical guarantees, and characterizing the minimax misclassification ratio [40]–[42] as well as the exact recovery criterion for the special case of two symmetric communities [34].

Although an information-theoretic limit for exact recovery has been derived in [34], their setting considering only two equal-sized communities and symmetric hyperedge generation probabilities is rather restrictive for real-world applications, and a general theory for exact recovery in the  $d$ -HSBM is still lacking. Motivated by this gap in the literature, in this work we consider the exact recovery criterion in the *general d-HSBM*. Our problem setting and the distinctions compared to other works are summarized as follows.

- 1) Nodes are partitioned into  $k \geq 2$  non-overlapping communities, in which each node is assigned to one of these  $k$  communities with probabilities  $\{p_i\}_{i=1}^k$ . This generalizes the setting in [30], [34] in which  $k = 2$  and  $p_1 = p_2$ , and the setting in [31], [40]–[42] wherein the  $k$  communities are of equal or approximately equal sizes (i.e.,  $p_i \approx p_j$  for all  $1 \leq i < j \leq k$ ).
- 2) The probability that an order- $d$  hyperedge appears depends on the number of nodes in each of the  $k$  communities (which is quantified by a length- $k$  vector  $(T_1, T_2, \dots, T_k)$  with  $\sum_i T_i = d$ , where  $T_i$  is the number of nodes in community  $i$ ). In contrast, many prior works with theoretical guarantees consider more restrictive assumptions on the probability that a hyperedge is present. For example, in [30], [31], [33]–[35] the hyperedge probability can only take *two* values depending on whether all the  $d$  nodes belong to the same community. Although other works such as [40]–[42] relax the restrictions in [30], [31], [33]–[35], they are nevertheless particularizations of our general HSBM model. We are aware that the assumption on hyperedge probabilities made in [26]–[29] is similar to ours, but their focus is to characterize the performance of a hypergraph spectral clustering method contained therein in terms of the fraction of misclassified nodes, whereas we aim to quantify necessary and sufficient conditions for *exact recovery* (see Definition 1 for details).
- 3) Based on the observed  $d$ -uniform hypergraph, the learner is tasked to achieve exact recovery of the hidden partition, i.e., all the  $n$  nodes should be assigned to the ground-truth communities that they belong to with high probability as the size of the hypergraph grows. We are also interested in deriving an algorithm-agnostic impossibility result that matches the performance guarantee of the learner’s algorithm.

### Main Contributions

The main contributions and key technical challenges of this work are summarized as follows.

- 1) We establish a phase transition for the general  $d$ -HSBM (see Theorems 1 and 2), apart from a small subset of  $d$ -HSBMs that contains communities whose so-called *second-order degree profiles* are identical (to be specified in Section III-A). That is, there is a sharp threshold such that exact recovery is possible above the threshold, and impossible below the threshold. This threshold is represented in terms of a quantity which we term as the *generalized Chernoff-Hellinger (GCH) divergence* between different communities, and is a generalization of the CH-divergence discovered in [4] for the SBM. Our result also recovers the exact recovery criteria for the SBM [2], [3] and  $d$ -HSBM with two symmetric communities [34] as special cases.
- 2) We develop a polynomial-time algorithm that meets the information-theoretic limits. This implies that there is no information-computation gap for exact recovery in the general  $d$ -HSBM (apart from the aforementioned small regime of parameters). Our two-stage algorithm consists of a *hypergraph spectral clustering* step in the first stage to ensure *almost exact recovery* (see Definition 2). It then performs local refinement steps for each of the  $n$  nodes in the second stage to ensure exact recovery. To circumvent the problem that conditioned on the success of the first stage certain *a priori* independent random variables become dependent, we adopt a *hypergraph splitting* technique to split the hypergraph into two sub-hypergraphs (see Section IV-A), such that the two stages can be run on the two independent sub-hypergraphs respectively, preserving the independence of the two stages to facilitate the analysis. Although this technique is not new, our analytical method is different from previous analyses (such as [4]). We prove that with high probability over splitting of the given hypergraph into two sub-hypergraphs, the good properties of the resultant sub-hypergraphs are preserved. This guarantees the success of the two stages.
- 3) A main technical challenge lies in the development and analysis of an efficient algorithm that leads to almost exact recovery (for the first stage) for the general  $d$ -HSBM. To the best of our knowledge, such an algorithm with accompanying guarantees is lacking in the literature. Thus, the hypergraph spectral clustering method developed here and its analysis may be of independent interest. We are aware that various clustering algorithms have been developed. However, theoretical guarantees (on the fraction of misclassified nodes) are usually restricted to special classes of HSBMs and they do not readily apply to the general  $d$ -HSBM. For example, the performance of the spectral clustering method in [40] depends on the  $k$ -th largest singular value of a specific matrix, but bounding this value turns out to be non-trivial for general  $d$ -HSBMs. The semidefinite programming-based method in [34] is only applicable to symmetric settings. Our new algorithm overcomes these stumbling blocks by leveraging and judiciously combining various ideas from prior works for the SBM [9], [10] and the HSBM [40]. Our theoretical result (Theorem 3) shows that, with probability approaching one, all but a vanishing fraction of the  $n$  nodes can be assigned to their true communities (i.e., almost exact recovery is achieved) in the general  $d$ -HSBM.

## Organization

We describe the general  $d$ -HSBM and the exact recovery criterion in Section II, and provide our main results (with accompanying discussions) in Section III. Our computationally efficient two-stage algorithm is introduced in Section IV, and its theoretical guarantee is formally established in Section V. The converse part is proved in Section VI. Section VII concludes this work and proposes several directions that are fertile avenues for future research.

## II. PRELIMINARIES AND PROBLEM STATEMENT

### A. Notation

For any integer  $m \geq 1$ , let  $[m]$  represent the set of integers  $\{1, 2, \dots, m\}$ , and  $\mathcal{S}_m$  be the set of all permutations from  $[m]$  to  $[m]$ . Random variables and their realizations are respectively denoted by upper-case and lower-case letters, while vectors, matrices, and tensors are denoted by boldface letters. For a length- $n$  vector  $\mathbf{x}$ , let  $x_i$  denote its  $i$ -th element, and  $\mathbf{x}_{\sim i}$  denote the length- $(n-1)$  sub-vector that excludes  $x_i$ . For a matrix  $\mathbf{A}$ , its operator norm and Frobenius norm are respectively represented by  $\|\mathbf{A}\|_{\text{op}}$  and  $\|\mathbf{A}\|_{\text{F}}$ , and its  $v$ -th column is denoted by  $\mathbf{A}_v$ .

### B. The $d$ -uniform hypergraph stochastic block model ( $d$ -HSBM)

Let  $n \in \mathbb{N}$  be the number of nodes, and  $k \geq 2$  be the number of non-overlapping communities. Each node  $v \in [n]$  belongs to one of the  $k$  communities, and is associated with a latent random variable  $Z_v$  on  $[k]$  with prior distribution  $\mathbf{p} = (p_1, p_2, \dots, p_k)$ , where  $\sum_{i \in [k]} p_i = 1$ . That is, if node  $v$  belongs to community  $i$ , then  $Z_v = i$ . The length- $n$  vector  $\mathbf{Z} = (Z_1, Z_2, \dots, Z_n)$  thus represents the *ground-truth community vector* of the  $n$  nodes. Furthermore, we define  $\mathcal{V}_i \triangleq \{v \in [n] : Z_v = i\}$  as the collection of nodes that belong to the community  $i$  (for  $i \in [k]$ ).

Let  $d \geq 2$  be the *order* of the hyperedges (i.e., the number of nodes contained in each hyperedge), and  $\mathcal{W}$  be the set of all order- $d$  hyperedges on  $[n]$  (where  $|\mathcal{W}| = \binom{n}{d}$ ). It is assumed that the hypergraph considered in this work only contains order- $d$  hyperedges; thus it is referred to as a  *$d$ -uniform hypergraph*. The generation process (underlying statistical model) of our random  $d$ -uniform hypergraph  $G = ([n], \mathcal{E})$  is as follows. For each  $e \in \mathcal{W}$ , the probability that it appears in the hypergraph  $G$  (i.e.,  $e \in \mathcal{E}$ ) depends on the number of nodes in each community  $\{\mathcal{V}_i\}_{i=1}^k$ . Formally, let

$$\mathcal{T} \triangleq \{\mathbf{T} \in \mathbb{N}^k : T_1 + T_2 + \dots + T_k = d\} \quad (1)$$

be the collection of length- $k$  vectors such that each vector  $\mathbf{T} \in \mathcal{T}$  (with  $T_i$  representing the number of nodes in  $\mathcal{V}_i$ ) represents a possible *community assignment* of  $d$  nodes, where ‘‘community assignment’’ is referred to as the number of nodes contained in each community. The generation of the hyperedges in  $G$  is fully characterized by a set of numbers  $\{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}} \subset \mathbb{R}_+$ . The probability of a hyperedge  $e \in \mathcal{W}$  appearing is  $Q_{\mathbf{T}(e)} \frac{\log n}{n^{d-1}}$ , where  $\mathbf{T}(e) \in \mathcal{T}$  denotes the community assignment of the  $d$  nodes in hyperedge  $e$ . That is,  $\mathbf{T}(e)$  is the length- $k$  vector whose  $i$ -th entry represents the number of nodes in the hyperedge  $e$  that belongs to the  $i$ -th community.

**Example 1.** Suppose  $d = 3$ ,  $k = 4$ ,  $n = 8$ , and  $\mathcal{V}_1 = \{1, 2\}$ ,  $\mathcal{V}_2 = \{3, 4\}$ ,  $\mathcal{V}_3 = \{5, 6\}$ ,  $\mathcal{V}_4 = \{7, 8\}$ . We list three different order- $d$  hyperedges  $e_1, e_2, e_3 \in \mathcal{W}$ , as well as their community assignments in the table below. Although  $e_1 \neq e_2$ , the probabilities that  $e_1 \in \mathcal{E}$  and  $e_2 \in \mathcal{E}$  are the same since they have the same community assignment. On the other hand, the probability that  $e_3 \in \mathcal{E}$  is in general different from that for  $e_1$  and  $e_2$ .

Hyperedges	Community assignments	Hyperedge probabilities
$e_1 = (1, 4, 8)$	$\mathbf{T}(e_1) = (1, 1, 0, 1)$	$Q_{(1,1,0,1)}(\log n)/n^2$
$e_2 = (1, 3, 7)$	$\mathbf{T}(e_2) = (1, 1, 0, 1)$	$Q_{(1,1,0,1)}(\log n)/n^2$
$e_3 = (2, 5, 7)$	$\mathbf{T}(e_3) = (1, 0, 1, 1)$	$Q_{(1,0,1,1)}(\log n)/n^2$

The reason why we consider the  $\Theta(\frac{\log n}{n^{d-1}})$ -regime for the connectivity probability is that it ensures the average degree of each node is  $\Theta(\log n)$  and it was shown [33], [34], [40] that phase transition for exact recovery occurs in this *logarithmic*

average degrees regime. Furthermore, we define  $Q_{\max} \triangleq \max_{\mathbf{T} \in \mathcal{T}} Q_{\mathbf{T}}$  and  $Q_{\min} \triangleq \min_{\mathbf{T} \in \mathcal{T}} Q_{\mathbf{T}}$ , and it is assumed that the parameters  $Q_{\max}, Q_{\min}, k, d$  and  $\{p_i\}_{i \in [k]}$  do not scale with  $n$ .

Similar to the adjacency matrices for graphs, any  $d$ -uniform hypergraph  $G$  can be represented by a  $d$ -dimensional  $n \times \dots \times n$  adjacency tensor  $\mathbf{A} = [A_{\mathbf{b}}]$ , where  $\mathbf{b} = [b_1, \dots, b_d] \in [n]^d$  is the access index of the element in the tensor. Here,  $A_{\mathbf{b}} \in \{0, 1\}$ , and  $A_{\mathbf{b}} = 1$  means the presence of the hyperedge corresponding to the  $d$  nodes in  $\mathbf{b}$ . In particular,  $A_{\mathbf{b}} = 0$  if the  $d$  elements in  $\mathbf{b}$  are *not* distinct (since each hypergraph must contain  $d$  nodes), and  $A_{\mathbf{b}} = A_{\mathbf{b}'}$  if there exists a permutation  $\pi$  such that  $(b'_{\pi(1)}, b'_{\pi(2)}, \dots, b'_{\pi(d)}) = (b_1, b_2, \dots, b_d)$ .

### C. Objective

Given the observation of the hypergraph  $G$  (or the adjacency tensor  $\mathbf{A}$ ), the learner aims to use an estimator  $\phi = \phi(G)$  to recover the *partition* of the  $n$  nodes into  $k$  communities. The output of the estimator  $\phi$  is denoted by  $\widehat{\mathbf{Z}} = (\widehat{Z}_1, \widehat{Z}_2, \dots, \widehat{Z}_n)$ . We measure the accuracy of  $\widehat{\mathbf{Z}}$  in terms of the *misclassification proportion*  $l(\widehat{\mathbf{Z}}, \mathbf{Z})$ , which is defined as

$$l(\widehat{\mathbf{Z}}, \mathbf{Z}) \triangleq \min_{\pi \in \mathcal{S}_k} \frac{1}{n} \sum_{v \in [n]} \mathbb{1} \left\{ \widehat{Z}_v \neq \pi(Z_v) \right\}. \quad (2)$$

**Definition 1** (Exact recovery). *An estimator  $\phi$  is said to achieve exact recovery if it ensures that with probability  $1 - o(1)$ , the misclassification proportion  $l(\widehat{\mathbf{Z}}, \mathbf{Z}) = 0$ .*

**Definition 2** (Almost exact recovery). *An estimator  $\phi$  is said to achieve almost exact recovery if it ensures that with probability  $1 - o(1)$ , the misclassification proportion  $l(\widehat{\mathbf{Z}}, \mathbf{Z}) \rightarrow 0$  as  $n$  tends to infinity.*

## III. MAIN RESULTS AND DISCUSSIONS

We first introduce several notations that are useful for stating our main results. Let

$$\mathcal{M} \triangleq \{\mathbf{m} \in \mathbb{N}^k : m_1 + m_2 + \dots + m_k = d - 1\} \quad (3)$$

be the collection of length- $k$  vectors such that the sum of the  $k$  elements equals  $d - 1$ . Each element in  $\mathcal{M}$  represents one possible community assignment of  $d - 1$  nodes. For each  $\mathbf{m} \in \mathcal{M}$ , we define

$$R_{\mathbf{m}} \triangleq \prod_{s=1}^k \binom{np_s}{m_s} \quad \text{and} \quad R'_{\mathbf{m}} \triangleq \frac{R_{\mathbf{m}}}{n^{d-1}} \quad (4)$$

as the expected number (and normalized expected number) of combinations of  $d - 1$  nodes that have community assignment  $\mathbf{m}$ . Note that  $R_{\mathbf{m}} = \Theta(n^{d-1})$  since  $\binom{np_s}{m_s} = \Theta(n^{m_s})$  and  $\sum_{s \in [k]} m_s = d - 1$ , and thus  $R'_{\mathbf{m}} = \Theta(1)$ . Suppose a node belongs to  $\mathcal{V}_i$  (where  $i \in [k]$ ) and other  $d - 1$  nodes have community assignment  $\mathbf{m} = (m_1, \dots, m_i, \dots, m_k) \in \mathcal{M}$ , then the joint community assignment of these  $d$  nodes is denoted by

$$\mathbf{m} \oplus i \triangleq (m_1, \dots, m_{i-1}, m_i + 1, m_{i+1}, \dots, m_k) \in \mathcal{T}. \quad (5)$$

For instance, when  $d = 3, k = 4, \mathbf{m} = (1, 1, 0, 0)$  and  $i = 3$ , we have  $\mathbf{m} \oplus i = (1, 1, 1, 0)$ .

### A. Separation between communities

1) *Degree profile*: For each community  $\mathcal{V}_i$  (where  $i \in [k]$ ), we define  $\mu_{\mathbf{m} \oplus i} \triangleq R'_{\mathbf{m}} Q_{\mathbf{m} \oplus i}$  for each  $\mathbf{m} \in \mathcal{M}$ . The interpretation of  $\mu_{\mathbf{m} \oplus i}$  is as follows: for any node  $v \in \mathcal{V}_i$ , the expected number of hyperedges that contain  $v$  and have community assignment  $\mathbf{m} \oplus i$  is  $R_{\mathbf{m}} \cdot \frac{Q_{\mathbf{m} \oplus i} \log n}{n^{d-1}}$ , thus  $\mu_{\mathbf{m} \oplus i}$  is the corresponding normalized quantity which scales as  $\Theta(1)$ . We then refer to the

collection  $\{\mu_{\mathbf{m} \oplus i}\}_{\mathbf{m} \in \mathcal{M}}$  as the *degree profile* of community  $i$ . Intuitively, two communities are easier to be separated if the degree profiles of these two communities are further apart. The discrepancy between any two communities in the  $d$ -HSBM can be measured in terms of the *generalized Chernoff-Hellinger divergence (GCH-divergence)* defined below, which generalizes the CH-divergence for the SBM that was first discovered by Abbe and Sandon [4, Eqn. (3)].

**Definition 3** (GCH-divergence). *For any  $i, j \in [k]$  such that  $i \neq j$ , we define the GCH-divergence between  $i$  and  $j$  as*

$$D_+(i, j) \triangleq \max_{t \in [0, 1]} \sum_{\mathbf{m} \in \mathcal{M}} t \mu_{\mathbf{m} \oplus i} + (1 - t) \mu_{\mathbf{m} \oplus j} - \mu_{\mathbf{m} \oplus i}^t \mu_{\mathbf{m} \oplus j}^{1-t}, \quad (6)$$

where  $D_+(i, j)$  is a function of  $\mathbf{p}$  and  $\{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}$ .

Note that  $D_+(i, j) = 0$  if and only if the degree profiles of communities  $\mathcal{V}_i$  and  $\mathcal{V}_j$  are exactly the same, in which case the two communities are statistically indistinguishable.

2) *Second-order degree profile*: For each community  $\mathcal{V}_i$  (where  $i \in [k]$ ), we define its *second-order degree profile* as  $\{\sum_{\mathbf{m} \in \mathcal{M}: m_s \geq 1} m_s \mu_{\mathbf{m} \oplus i}\}_{s \in [k]}$ , where each element  $\sum_{\mathbf{m} \in \mathcal{M}: m_s \geq 1} m_s \mu_{\mathbf{m} \oplus i}$  represents the normalized expected number of hyperedges that contain two fixed nodes belonging to  $\mathcal{V}_i$  and  $\mathcal{V}_s$  respectively. When the second-order degree profile of two communities are exactly the same, our analysis also shows that there may be some inherent difficulties in distinguishing them. Formally, we define  $\Xi$  as the subset of model parameters  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}})$  such that there exist two communities having the same second-order degree profiles, i.e.,

$$\Xi \triangleq \left\{ (\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) : \exists i \neq j \text{ such that } \sum_{\mathbf{m} \in \mathcal{M}: m_s \geq 1} m_s \mu_{\mathbf{m} \oplus i} = \sum_{\mathbf{m} \in \mathcal{M}: m_s \geq 1} m_s \mu_{\mathbf{m} \oplus j} \text{ for all } s \in [k] \right\}. \quad (7)$$

## B. Main Results

**Theorem 1** (Converse). *It is impossible to achieve exact recovery when the model parameters  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}})$  satisfy*

$$\min_{i, j \in [k]: i \neq j} D_+(i, j) < 1. \quad (8)$$

**Theorem 2** (Achievability). *Assume that the model parameters  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \notin \Xi$ . Then the polynomial-time two-stage algorithm (Algorithm 1) achieves exact recovery when the model parameters  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}})$  satisfy*

$$\min_{i, j \in [k]: i \neq j} D_+(i, j) > 1. \quad (9)$$

Some remarks on Theorems 1 and 2 are in order.

- 1) For community detection in the  $d$ -HSBM, most of the settings considered in prior works, such as the one that  $\{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}$  can only take two values depending on whether  $d$  nodes belong to the same community [30]–[35], satisfy  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \notin \Xi$ . Thus, our result is a strict generalization of these existing works. An example for the case that  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \in \Xi$  is as follows. Suppose  $k = 2$ ,  $d = 3$ ,  $\mathbf{p} = (p_1, p_2)$ , and  $\{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}} = \{Q_{(3,0)}, Q_{(2,1)}, Q_{(1,2)}, Q_{(0,3)}\}$ . When  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}})$  satisfies  $p_1 = p_2 = 1/2$ ,  $Q_{(3,0)} = Q_{(1,2)}$  and  $Q_{(2,1)} = Q_{(0,3)}$ , one can check that the two communities  $\mathcal{V}_1$  and  $\mathcal{V}_2$  have the same second-order degree profile, thus  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \in \Xi$ .
- 2) When  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \in \Xi$  and  $\min_{i, j \in [k]: i \neq j} D_+(i, j) > 1$ , it remains open whether exact recovery is possible. However, we would like to point out that this scenario does not apply to the SBM (equivalently, the 2-HSBM), since the condition  $\min_{i, j \in [k]: i \neq j} D_+(i, j) > 1$  immediately implies that  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \notin \Xi$  in this setting. Our sharp threshold is applicable to all model parameters when  $d = 2$ .

- 3) Algorithm 1 contains two stages. The first stage utilizes a hypergraph spectral clustering method to obtain an initial estimate  $\widehat{\mathbf{Z}}^{(0)}$  that satisfies almost exact recovery, while the second stage involves performing local refinement steps to each node individually to correct the misclassified nodes in  $\widehat{\mathbf{Z}}^{(0)}$ .
- 4) When  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \notin \Xi$ , the first stage ensures almost exact recovery (as shown in Theorem 3), and the condition  $\min_{i,j \in [k]: i \neq j} D_+(i,j) > 1$  is the criterion for Stage 2 to succeed. Roughly speaking, performing a local refinement step for each node is equivalent to performing a hypothesis test with independent but non-identically distributed samples. The corresponding error probability can be represented by a variant of the *Chernoff information* [43, Chapter 11.9], and this further reduces to  $n^{-\min_{i,j \in [k]: i \neq j} D_+(i,j)}$  which is in the form of the GCH-divergence. Thus, when  $\min_{i,j \in [k]: i \neq j} D_+(i,j) > 1$ , taking a union bound over the  $n$  nodes results in a vanishing error probability (i.e., exact recovery is achieved).
- 5) Our analysis of the first stage is not able to handle the case in which  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \in \Xi$  because the key step in the hypergraph spectral clustering method is to map the order- $d$  adjacency tensor  $\mathbf{A}$  to an  $n \times n$  matrix  $\mathbf{L}$  (defined in (12) below), and the subsequent clustering algorithm critically relies on the discrepancy between the columns of  $\mathbf{L}$  (which corresponds to the second-order degree profiles of communities). We conjecture that this issue may be circumvented if one directly applies clustering algorithms on the adjacency tensor (such as the method proposed in [37]), and the exact recovery threshold  $\min_{i,j \in [k]: i \neq j} D_+(i,j) = 1$  holds without the assumption that the second-order degree profiles of any two communities are distinct.
- 6) Both the GCH-divergence and CH-divergence [4] can be considered as generalizations of the *Hellinger divergence* and *Chernoff divergence*. We refer the readers to [4, Section 2.3] for a detailed discussion.

### C. Recovering prior results from Theorems 1 and 2

To the best of our knowledge, the sharp threshold established by Theorems 1 and 2 is the most general result for exact recovery in the SBM/HSBM literature. As discussed below, several problem settings investigated in prior works are subsumed by our result, and the thresholds derived in the prior works can be recovered from Theorems 1 and 2.

1) *Exact recovery in the SBM [4]*: The SBM considered in [4] corresponds to our  $d$ -HSBM with  $d = 2$ . In [4], the prior distribution of each node is also  $\mathbf{p} = (p_1, p_2, \dots, p_k)$ , and the edge probabilities are characterized by  $\{Q_{i,j} \frac{\log n}{n}\}_{i,j \in [k]}$ , where  $Q_{i,j}$  corresponds to edges that contain nodes in  $\mathcal{V}_i$  and  $\mathcal{V}_j$ . The authors of [4] showed that the threshold for exact recovery is

$$\min_{i,j \in [k]: i \neq j} \max_{t \in [0,1]} \sum_{s \in [k]} p_s [tQ_{s,i} + (1-t)Q_{s,j} - Q_{s,i}^t Q_{s,j}^{1-t}] = 1. \quad (10)$$

In our setting with  $d = 2$ , the set  $\mathcal{M} = \{\mathbf{m}_1, \mathbf{m}_2, \dots, \mathbf{m}_k\}$  contains  $k$  distinct length- $k$  vectors, where each  $\mathbf{m}_s \triangleq (0, \dots, 0, 1, 0, \dots, 0)$  contains a single one which is at the  $s$ -th location. By noting that  $\mu_{\mathbf{m} \oplus i} = p_s Q_{s,i}$  (resp.  $\mu_{\mathbf{m} \oplus j} = p_s Q_{s,j}$ ) in [4] and  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \notin \Xi$  when  $\min_{i,j \in [k]: i \neq j} D_+(i,j) > 1$ , we recover their threshold stated in Eqn. (10) from Theorems 1 and 2.

2) *Exact recovery in the  $d$ -HSBM with two symmetric communities [34]*: The model considered in [34] is a special  $d$ -HSBM with two equal-sized communities that have symmetric structures. It corresponds to our general  $d$ -HSBM with  $k = 2$ ,  $p_1 = p_2 = 1/2$ , and  $\{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}} = \{Q_1, Q_2\}$  (where each hyperedge appears with probability  $Q_1$  when  $d$  nodes are in the same communities, and  $Q_2$  otherwise). Kim, Bandeira, and Goemans [34] showed that the threshold for exact recovery is

$$\frac{1}{2^{d-1}} \left( \sqrt{\frac{Q_1}{(d-1)!}} - \sqrt{\frac{Q_2}{(d-1)!}} \right)^2 = 1. \quad (11)$$

Specializing our result to this symmetric setting, we note that the set  $\mathcal{M} = \{(d-1, 0), (d-2, 1), \dots, (0, d-1)\}$  contains  $d$  distinct length-2 vectors, and the values of  $\mu_{\mathbf{m} \oplus 1}$  and  $\mu_{\mathbf{m} \oplus 2}$  for all  $\mathbf{m} \in \mathcal{M}$  can then be calculated. By noting that the model parameters  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \notin \Xi$  and  $t = 1/2$  maximizes the GCH-divergence in (6) for symmetric SBMs and HSBMs, we recover the threshold stated in Eqn. (11) from Theorems 1 and 2. Furthermore, we also recover the celebrated exact recovery threshold  $|\sqrt{Q_1} - \sqrt{Q_2}| = \sqrt{2}$  for the SBM with two symmetric communities [2], since it is a special case of [34] for  $d = 2$ .

#### IV. THE TWO-STAGE ALGORITHM FOR EXACT RECOVERY

In this section, we present our polynomial-time algorithm that is used to achieve the information-theoretic limit shown in Theorem 2. As mentioned in Section III-B, our algorithm consists of two stages such that the first stage achieves almost exact recovery via the hypergraph spectral clustering method and the second stage achieves exact recovery via local refinement steps. This “*from global to local*” principle has been employed in many contexts, such as community detection in the SBM [4], [9]–[11], [44] and HSBM [33], [34], [40]–[42], matrix completion [45]–[47], etc. It is also worth noting that when analyzing two-stage algorithms, random variables that are initially independent may become dependent conditioned on the success of a preceding stage. To ameliorate this problem, we adopt the graph splitting technique (as described in Subsection IV-A) which is inspired by prior works on community detection [4], [11], [33], [48]. Our algorithm is described in detail in Algorithm 1.

##### A. Graph Splitting

Let  $F = ([n], \mathcal{W})$  be the *complete  $d$ -uniform hypergraph* on node set  $[n]$ , and the hyperedge set  $\mathcal{W}$  contains all the  $\binom{n}{d}$  order- $d$  hyperedges (as defined in Section II-B). We randomly split  $F$  into two sub-hypergraph  $F_1 = ([n], \mathcal{W}_1)$  and  $F_2 = ([n], \mathcal{W}_2)$ . Each hyperedge in  $\mathcal{W}$  is sampled to  $\mathcal{W}_1$  with probability  $\gamma_n / \log n$ , and to  $\mathcal{W}_2$  with probability  $1 - (\gamma_n / \log n)$ , where  $\gamma_n$  belongs to  $\omega(1) \cap o(\log n)$ . Note that  $\mathcal{W}_2$  is the complement of  $\mathcal{W}_1$ . This splitting process is independent of the generation of the hypergraph  $G = ([n], \mathcal{E})$  (which is generated according to  $\mathbf{p}$  and  $\{Q_{\mathbf{T}}(\log n) / n^{d-1}\}_{\mathbf{T} \in \mathcal{T}}$ ). We then define  $G_1 = ([n], \mathcal{E}_1)$  as the *sub-HSBM* that is generated on the hyperedge set  $\mathcal{W}_1$  of  $F_1$ , where  $\mathcal{E}_1 = \mathcal{E} \cap \mathcal{W}_1$  is the intersection of the hyperedge sets of the HSBM  $G$  and the sub-hypergraph  $F_1$ . Similarly, we define  $G_2 = ([n], \mathcal{E}_2)$  as the sub-HSBM that is generated on the hyperedge set  $\mathcal{W}_2$  of  $F_2$ , where  $\mathcal{E}_2 = \mathcal{E} \cap \mathcal{W}_2$ .

##### B. Almost exact recovery via a hypergraph spectral clustering method (Stage 1)

The main focus of this subsection is the sub-HSBM  $G_1 = ([n], \mathcal{E}_1)$ . We apply a hypergraph spectral clustering method on  $G_1$  to obtain an initial estimate of the ground-truth community vector  $\mathbf{Z}$ , denoted by  $\widehat{\mathbf{Z}}^{(0)} = (\widehat{Z}_1^{(0)}, \dots, \widehat{Z}_n^{(0)})$ .

Let  $\mathbf{H} = [H_{ve}]$  be the  $n \times \binom{n}{d}$  binary incidence matrix corresponding to  $G_1$  such that each entry  $H_{ve} = 1$  if the hyperedge  $e \in \mathcal{E}_1$  and  $e$  contains node  $v$ , and  $H_{ve} = 0$  otherwise. Note that there is an one-to-one mapping between  $\mathbf{H}$  and the observed adjacency tensor  $\mathbf{A}$ . For each node  $v \in [n]$ , its degree (in  $G_1$ ) is denoted by  $d_v \triangleq \sum_{e \in \mathcal{E}} H_{ve}$ . Let  $\mathbf{D} = \text{diag}(d_1, \dots, d_n)$  be an  $n \times n$  diagonal matrix that represents the degrees of the  $n$  nodes. We then define the *hypergraph Laplacian* as

$$\mathbf{L} \triangleq \mathbf{H}\mathbf{H}^\top - \mathbf{D}, \quad (12)$$

where  $\mathbf{L}$  is an  $n \times n$  matrix and the  $(i, j)$ -entry represents the number of hyperedges that contain both node  $i$  and node  $j$ . To ensure a good performance of the hypergraph spectral clustering method, one typically needs to remove a small fractions of nodes that have significantly higher degrees [40] than the average. Thus, we define the set of “good” nodes that have degree no larger than a certain threshold  $\tau$  as

$$\Gamma \triangleq \{v \in [n] : d_v \leq \tau\}, \quad (13)$$



---

**Algorithm 1: THE TWO-STAGE ALGORITHM**


---

- 1 **Input:** Hypergraphs  $G_1, G_2$ , number of communities  $k$ , radius  $r = \frac{\gamma_n^2}{n \log(\gamma_n)}$ ,  $\widehat{\mathcal{V}}_0^{(0)} = \emptyset$ ;
  - 2 **Stage 1: Almost exact recovery via the hypergraph spectral clustering method**
  - 3  $\mathbf{L} \leftarrow \mathbf{H}\mathbf{H}^\top - \mathbf{D}$ ;  $\mathbf{L}_\Gamma \leftarrow$  trim the rows and columns in  $\mathbf{L}$  that correspond to  $v \notin \Gamma$ ;
  - 4  $\mathbf{L}_\Gamma^{(k)} \leftarrow$  rank- $k$  approximation of  $\mathbf{L}_\Gamma$ ;
  - 5  $\Psi \leftarrow$  a subset of nodes that contains  $\lceil \log n \rceil$  random samples (with replacement) from  $\Gamma$ ;
  - 6  $\mathcal{B}_v \leftarrow \{u \in \Gamma : \|(\mathbf{L}_\Gamma^{(k)})_u - (\mathbf{L}_\Gamma^{(k)})_v\|_2^2 \leq r\}$ , for all  $v \in \Psi$ ;
  - 7 **for**  $j = 1$  **to**  $k$  **do**
  - 8  $v_j^* \leftarrow \arg \max_{v \in \Psi} |\mathcal{B}_v \setminus (\cup_{l=0}^{j-1} \widehat{\mathcal{V}}_l^{(0)})|$ ;
  - 9  $\widehat{\mathcal{V}}_j^{(0)} \leftarrow \mathcal{B}_{v_j^*} \setminus (\cup_{l=0}^{j-1} \widehat{\mathcal{V}}_l^{(0)})$
  - 10 **end**
  - 11 **for**  $v \in \Gamma \setminus (\cup_{j=1}^k \widehat{\mathcal{V}}_j^{(0)})$  **do**
  - 12  $j_* \leftarrow \arg \min_j \|(\mathbf{L}_\Gamma^{(k)})_v - (\mathbf{L}_\Gamma^{(k)})_{v_j^*}\|_2^2$ ;
  - 13  $\widehat{\mathcal{V}}_{j_*}^{(0)} \leftarrow \widehat{\mathcal{V}}_{j_*}^{(0)} \cup \{v\}$
  - 14 **end**
  - 15 Randomly assign each  $v \in [n] \setminus \Gamma$  to one community in  $\{\widehat{\mathcal{V}}_i^{(0)}\}_{i \in [k]}$
  - 16 Output the initial estimate  $\widehat{\mathbf{Z}}^{(0)}$  based on  $\{\widehat{\mathcal{V}}_i^{(0)}\}_{i \in [k]}$
  - 17 **Stage 2: Local refinement steps**
  - 18 **for**  $v \in [n]$  **do**
  - 19  $\widehat{Z}_v \triangleq \arg \max_{i \in [k]} \mathbb{P}(Z_v = i | G_2 = g_2, \widehat{\mathbf{Z}}_{\sim v}^{(0)} = \widehat{\mathbf{z}}_{\sim v}^{(0)})$ ;
  - 20 **end**
- Output:** Final estimate  $\widehat{\mathbf{Z}} = (\widehat{Z}_1, \widehat{Z}_2, \dots, \widehat{Z}_n)$
- 

where  $\tau$  is set to be  $CQ_{\max} \gamma_n$  for some large constant  $C > 0$ , such that  $\tau$  is much larger than the expected degree of every node.

We apply Stage 1 of Algorithm 1 (lines 2 – 16) to obtain an almost exact recovery of the  $k$  communities. Initially, we calculate the hypergraph Laplacian  $\mathbf{L}$ , and then “trim” the rows and columns in  $\mathbf{L}$  that correspond to nodes that do not belong to  $\Gamma$ . Specifically, if node  $v \notin \Gamma$ , we replace all the entries in the  $v$ -th row and  $v$ -th column of  $\mathbf{L}$  by 0. This yields the *trimmed hypergraph Laplacian*  $\mathbf{L}_\Gamma$ . In addition, we also perform an singular value decomposition (SVD) on  $\mathbf{L}_\Gamma$  to obtain the optimal rank- $k$  approximation  $\mathbf{L}_\Gamma^{(k)}$ .

We then perform a clustering algorithm (lines 5 – 16) on the columns of  $\mathbf{L}_\Gamma^{(k)}$ , i.e., the set of column vectors  $\{(\mathbf{L}_\Gamma^{(k)})_v\}_{v \in \Gamma}$ . An example of our clustering algorithm is illustrated in Fig. 1. We first randomly select  $\lceil \log n \rceil$  nodes from  $\Gamma$  (with replacement) as *reference nodes*, and it can be shown (in Lemma 5 below) that each community contains at least one reference node with high probability. This set of reference nodes is denoted by  $\Psi$ . For each node  $v \in \Psi$ , we construct a ball  $\mathcal{B}_v$  with center  $v$  and radius  $r \triangleq \frac{\gamma_n^2}{n \log(\gamma_n)}$  which includes all the neighboring nodes (i.e., the nodes in  $\mathcal{B}_v$ ). Among  $\{\mathcal{B}_v\}_{v \in \Psi}$ , we find the one that has the largest cardinality, declare  $v_1^* \triangleq \arg \max_{v \in \Psi} |\mathcal{B}_v|$ , and set the largest community  $\widehat{\mathcal{V}}_1^{(0)}$  to be  $\mathcal{B}_{v_1^*}$ . To find the second largest community, we remove all the nodes in  $\widehat{\mathcal{V}}_1^{(0)}$  and then follow a similar procedure to find the ball with the largest cardinality. That is, we declare  $v_2^* \triangleq \arg \max_{v \in \Psi} |\mathcal{B}_v \setminus \widehat{\mathcal{V}}_1^{(0)}|$ , and set the second largest community  $\widehat{\mathcal{V}}_2^{(0)}$  to be  $\mathcal{B}_{v_2^*} \setminus \widehat{\mathcal{V}}_1^{(0)}$ . By repeating




Fig. 1: An example with 3 communities. Each node corresponds to a column of  $\mathbf{L}_\Gamma^{(k)}$ .

this procedure for  $2 \leq j \leq k$ , we obtain  $k$  estimated communities  $\hat{\mathcal{V}}_1^{(0)}, \hat{\mathcal{V}}_2^{(0)}, \dots, \hat{\mathcal{V}}_k^{(0)}$  (lines 7 – 10). Furthermore, we assign the nodes belonging to  $\Gamma \setminus (\cup_{j \in [k]} \hat{\mathcal{V}}_j^{(0)})$  to their nearest communities (lines 11 – 14), and the nodes that do not belong to  $\Gamma$  to each community randomly (line 15). Finally, for each node  $v \in \hat{\mathcal{V}}_j^{(0)}$  (for all  $j \in [k]$ ), we set  $\hat{Z}_v^{(0)} = j$ .

The high-level intuition of the analysis of Stage 1 (when  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \notin \Xi$ ) is as follows. Let  $\mathbf{M} \triangleq \mathbb{E}(\mathbf{L}_\Gamma)$ , and note that  $\mathbf{M}$  is an  $n \times n$  matrix of rank  $k$ . If nodes  $u$  and  $v$  are in the same cluster, we have  $\mathbf{M}_u = \mathbf{M}_v$ ; otherwise we show in Lemma 3 that  $\|\mathbf{M}_u - \mathbf{M}_v\|_2^2 = \Omega(\gamma_n^2/n)$ . As a result, one can prove that, by running Stage 1, each node  $v$  will be misclassified only if the distance  $\|(\mathbf{L}_\Gamma^{(k)})_v - \mathbf{M}_v\|_2^2 = \Omega(\gamma_n^2/n)$ . On the other hand, one can prove that the sum of the distances satisfies  $\sum_{v \in \Gamma} \|(\mathbf{L}_\Gamma^{(k)})_v - \mathbf{M}_v\|_2^2 = \|\mathbf{L}_\Gamma^{(k)} - \mathbf{M}\|_F^2 = \mathcal{O}(\gamma_n)$  with high probability. Therefore, the output of Stage 1 results in a misclassification of at most  $\mathcal{O}(n/\gamma_n)$  nodes, and the misclassified proportion tends to zero as  $n$  tends to infinity (since  $\gamma_n = \omega(1)$ ). The theoretical guarantee of Stage 1 is formally established in Section V-A.

**Remark 1.** *The number of randomly select reference nodes is set to be  $\lceil \log n \rceil$  such that it is large enough to ensure that each community contains at least one reference node (with high probability). Since the radius  $r = \gamma_n^2/(n \log \gamma_n)$  is also large enough, one can use the reference node in each community (denoted by  $v$ ) to find most of its community members, via the ball  $\mathcal{B}_v$  with center  $v$  and radius  $r$ . As a result, we only need to compute  $\Theta(n \log n)$  pairwise distances  $\|(\mathbf{L}_\Gamma^{(k)})_u - (\mathbf{L}_\Gamma^{(k)})_v\|_2^2$ ; while some related works [40] use all the  $n$  nodes as reference nodes and thus  $\Theta(n^2)$  pairwise distances need to be computed.*

### C. Local refinements (Stage 2)

After obtaining the initial estimate  $\hat{\mathbf{Z}}^{(0)}$ , we refine the label of each node  $v \in [n]$  based on the observation of the hypergraph  $G_2 = g_2$  as well as the estimated labels  $\{\hat{\mathbf{Z}}_{\sim v}^{(0)}\}$  for the remaining nodes. For each node  $v \in [n]$ , we perform a local *maximum a posteriori (MAP)* estimation step as follows:

$$\hat{Z}_v(g_2, \hat{\mathbf{z}}_{\sim v}^{(0)}) \triangleq \arg \max_{i \in [k]} \mathbb{P} \left( Z_v = i \mid G_2 = g_2, \hat{\mathbf{Z}}_{\sim v}^{(0)} = \hat{\mathbf{z}}_{\sim v}^{(0)} \right). \quad (14)$$

This leads to the final estimate  $\hat{\mathbf{Z}} = (\hat{Z}_1, \dots, \hat{Z}_n)$  of the ground-truth community vector. A detailed analysis of Stage 2 is provided in Section V-B.

## V. THEORETICAL GUARANTEES OF ALGORITHM 1 (THEOREM 2)

In this section, we prove that as long as  $\min_{i,j \in [k]: i \neq j} D_+(i, j) > 1$  and  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \notin \Xi$ , applying Algorithm 1 on the observed hypergraph  $G$  ensures  $l(\mathbf{Z}, \hat{\mathbf{Z}}) = 0$  with high probability for sufficiently large  $n$  (i.e., exact recovery).

First, we note that with high probability, the size of each community  $\mathcal{V}_j = \{v \in [n] : Z_v = j\}$  is close to  $np_j$  for all  $j \in [k]$ . This is stated in Lemma 1 below and can be proved by applying the Chernoff bound.

**Lemma 1.** *Fix a constant  $\delta \in (0, 1)$  which can be chosen to be arbitrarily small. We say that the length- $n$  vector  $\mathbf{Z} \in \mathcal{A}_Z$  (where  $\mathcal{A}_Z$  is referred to as the typical set of  $\mathbf{Z}$ ) if the communities  $\{\mathcal{V}_j\}_{j \in [k]}$  associated with  $\mathbf{Z}$  satisfy*

$$\left(1 - n^{-\frac{1}{2} + \frac{\delta}{2}}\right) p_j n \leq |\mathcal{V}_j| \leq \left(1 + n^{-\frac{1}{2} + \frac{\delta}{2}}\right) p_j n, \quad \text{for all } j \in [k]. \quad (15)$$

Then, we have  $\mathbb{P}(\mathbf{Z} \in \mathcal{A}_Z) \geq 1 - \exp(-\Theta(n^\delta))$ .

Therefore, one can focus on typical ground-truth community vectors  $\mathbf{z} \in \mathcal{A}_Z$  in the following analysis.

#### A. Theoretical guarantees of Stage 1

For a fixed ground-truth community vector  $\mathbf{z} \in \mathcal{A}_Z$ , we first introduce an artificial  $d$ -HSBM  $\tilde{G}_1$  which is generated with respect to the ground-truth community vector  $\mathbf{z}$  and hyperedge probabilities  $\{\frac{Q_{\mathbf{T}} \gamma_n}{n^{d-1}}\}_{\mathbf{T} \in \mathcal{T}}$ . Note that the generation process of  $\tilde{G}_1$  is equivalent to first generating a sub-hypergraph  $F_1$  (with splitting parameter  $\gamma_n / \log n$ ) and then generating a sub-HSBM  $G_1$  on the hyperedge set of  $F_1$  (with hyperedge probabilities  $\{\frac{Q_{\mathbf{T}} \log n}{n^{d-1}}\}_{\mathbf{T} \in \mathcal{T}}$ ). Thus, we investigate the misclassification proportion  $l(\mathbf{z}, \hat{\mathbf{Z}}^{(0)})$  based on the random hypergraph  $\tilde{G}_1$ .

**Theorem 3** (Theoretical guarantee of Stage 1). *Suppose the model parameters  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \notin \Xi$ . For any fixed  $\mathbf{z} \in \mathcal{A}_Z$ , there exist vanishing sequences  $\{\epsilon_n\}$  and  $\{\eta_n\}$  (which depend on  $\{\gamma_n\}$ ) such that with probability at least  $1 - \epsilon_n$  over the generation of  $\tilde{G}_1$ , running Stage 1 of Algorithm 1 ensures that  $l(\mathbf{z}, \hat{\mathbf{Z}}^{(0)}) \leq \eta_n$ , i.e., almost exact recovery is achieved.*

Theorem 3 is proved in the rest of this section, and in the following we assume that  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \notin \Xi$ .

Note that with high probability over the generation of  $\tilde{G}_1$ , the degrees of most nodes in  $[n]$  are smaller than the threshold  $\tau$ , thus only a vanishing fraction of nodes (at most  $\mathcal{O}(n/\gamma_n)$  nodes) is trimmed. This result is adapted from [40, Lemma D.3] and stated below.

**Lemma 2** (Adapted from Lemma D.3 in [40]). *There exists a large constant  $C > 0$  such that if we set  $\tau = CQ_{\max} \gamma_n$ , then with probability at least  $1 - \exp(-C'n)$  (for some constant  $C' > 0$ ), the cardinality of the set  $\Gamma$  satisfies  $|\Gamma| \geq n(1 - (1/\tau))$ .*

Furthermore, let  $\mathcal{V}_j^\Gamma \triangleq \mathcal{V}_j \cap \Gamma$  for all  $j \in [k]$ . Lemma 2 also implies that with high probability,

$$(1 - o(1))p_j n \leq |\mathcal{V}_j^\Gamma| \leq (1 + o(1))p_j n, \quad \text{for all } j \in [k]. \quad (16)$$

We then focus on the remaining nodes in the set  $\Gamma$ . First recall that  $\mathbf{M} = \mathbb{E}(\mathbf{L}_\Gamma)$  is a rank- $k$  matrix such that  $\mathbf{M}_u = \mathbf{M}_v$  if  $u$  and  $v$  belong to the same community (where  $u, v \in \Gamma$ ). Lemma 3 below shows that the distance between the two columns  $\mathbf{M}_u$  and  $\mathbf{M}_v$  scales as  $\Omega(\gamma_n^2/n)$  if  $u$  and  $v$  belong to different communities.

**Lemma 3.** *Suppose  $u, v \in \Gamma$ . Then we have*

$$\|\mathbf{M}_u - \mathbf{M}_v\|_2^2 = \begin{cases} 0, & \text{if } u \text{ and } v \text{ belong to the same cluster;} \\ \Omega(\gamma_n^2/n), & \text{otherwise.} \end{cases} \quad (17)$$

The proof of Lemma 3 can be found in Appendix A. Lemma 4 below shows that with high probability, both the operator norm and the Frobenius norm of the difference between  $\mathbf{L}_\Gamma$  and  $\mathbf{M}$  can be appropriately upper bounded.

**Lemma 4.** For any constant  $C'_1 > 0$ , there exists some constant  $C_1 > 0$  such that with probability at least  $1 - n^{-C'}$ ,

$$\|\mathbf{L}_\Gamma - \mathbf{M}\|_{\text{op}} \leq C_1 \cdot \left[ \sqrt{\gamma_n Q_{\max}} + \sqrt{\tau} + \frac{\gamma_n Q_{\max}}{\sqrt{\gamma_n Q_{\max}} + \sqrt{\tau}} \right]. \quad (18)$$

As a result, the Frobenius norm between  $\mathbf{M}$  and  $\mathbf{L}_\Gamma^{(k)}$  (the rank- $k$  approximation of  $\mathbf{L}_\Gamma$ ) can be upper-bounded as

$$\|\mathbf{L}_\Gamma^{(k)} - \mathbf{M}\|_{\text{F}}^2 \leq 8kC_1^2 \cdot \left[ \sqrt{\gamma_n Q_{\max}} + \sqrt{\tau} + \frac{\gamma_n Q_{\max}}{\sqrt{\gamma_n Q_{\max}} + \sqrt{\tau}} \right]^2 = \mathcal{O}(\gamma_n). \quad (19)$$

*Proof of Lemma 4.* The proof of (18) can be adapted from [40, Lemma D.4], so our main focus is on the proof of (19). Note that

$$\|\mathbf{L}_\Gamma^{(k)} - \mathbf{M}\|_{\text{F}}^2 \leq 2k \cdot \|\mathbf{L}_\Gamma^{(k)} - \mathbf{M}\|_{\text{op}}^2 \quad (20)$$

$$\leq 4k \cdot \|\mathbf{L}_\Gamma^{(k)} - \mathbf{L}_\Gamma\|_{\text{op}}^2 + 4k \cdot \|\mathbf{L}_\Gamma - \mathbf{M}\|_{\text{op}}^2 \quad (21)$$

$$\leq 4k \cdot \sigma_{k+1}^2 + 4k \cdot \|\mathbf{L}_\Gamma - \mathbf{M}\|_{\text{op}}^2 \quad (22)$$

$$\leq 8k \cdot \|\mathbf{L}_\Gamma - \mathbf{M}\|_{\text{op}}^2, \quad (23)$$

where  $\sigma_{k+1}$  is the  $(k+1)$ -th largest singular value of  $\mathbf{L}_\Gamma$ . Eqn. (20) holds since the rank of  $\mathbf{L}_\Gamma^{(k)} - \mathbf{M}$  is at most  $2k$ , and  $\|\mathbf{X}\|_{\text{F}}^2 \leq r\|\mathbf{X}\|_{\text{op}}^2$  for any matrix  $\mathbf{X}$  of rank  $r$ . Eqn. (21) holds since  $\|\mathbf{L}_\Gamma^{(k)} - \mathbf{M}\|_{\text{op}} \leq \|\mathbf{L}_\Gamma^{(k)} - \mathbf{L}_\Gamma\|_{\text{op}} + \|\mathbf{L}_\Gamma - \mathbf{M}\|_{\text{op}}$  and  $2\|\mathbf{L}_\Gamma^{(k)} - \mathbf{L}_\Gamma\|_{\text{op}} \cdot \|\mathbf{L}_\Gamma - \mathbf{M}\|_{\text{op}} \leq \|\mathbf{L}_\Gamma^{(k)} - \mathbf{L}_\Gamma\|_{\text{op}}^2 + \|\mathbf{L}_\Gamma - \mathbf{M}\|_{\text{op}}^2$ . Eqn. (22) follows from the *Eckart–Young–Mirsky theorem* [49], while Eqn. (23) is due to *Weyl's inequality*<sup>2</sup> [50]. Combining Eqns (18) and (23), it is then clear that

$$\|\mathbf{L}_\Gamma^{(k)} - \mathbf{M}\|_{\text{F}}^2 \leq 2k \cdot \|\mathbf{L}_\Gamma^{(k)} - \mathbf{M}\|_{\text{op}}^2 \leq 8kC_1^2 \cdot \left[ \sqrt{\gamma_n Q_{\max}} + \sqrt{\tau} + \frac{\gamma_n Q_{\max}}{\sqrt{\gamma_n Q_{\max}} + \sqrt{\tau}} \right]^2 = \mathcal{O}(\gamma_n). \quad (24)$$

This completes the proof of Lemma 4.  $\square$

**Analysis of Stage 1:** In the following, we show that when Lemmas 2–4 hold, running Stage 1 ensures that  $l(\mathbf{z}, \widehat{\mathbf{Z}}^{(0)}) \rightarrow 0$  with high probability. Our analysis is inspired by [9] for SBMs, but is adapted to our general  $d$ -HSBM problem. We first partition the nodes in  $\Gamma$  as follows. Recall that the radius  $r$  is set to be  $\gamma_n^2/(n \log(\gamma_n))$ , and let

$$\mathcal{I}_j^{\text{in}} \triangleq \{v \in \mathcal{V}_j^\Gamma : \|(\mathbf{L}_\Gamma^{(k)})_v - \mathbf{M}_v\|_2^2 \leq r/4\}, \quad \text{for all } j \in [k], \quad (25)$$

$$\mathcal{I}_j^{\text{out}} \triangleq \{v \in \mathcal{V}_j^\Gamma : \|(\mathbf{L}_\Gamma^{(k)})_v - \mathbf{M}_v\|_2^2 \leq 4r\}, \quad \text{for all } j \in [k], \quad (26)$$

and the remaining nodes in  $\Gamma$  belong to  $\mathcal{U} \triangleq \Gamma \setminus \{\cup_{j \in [k]} \mathcal{I}_j^{\text{out}}\}$ . Note that these sets have the following properties:

(i) For all  $j \in [k]$ , most of the nodes  $v \in \mathcal{V}_j^\Gamma$  are such that  $(\mathbf{L}_\Gamma^{(k)})_v$  is  $\frac{r}{4}$ -close to its expectation  $\mathbf{M}_v$ , since

$$|\mathcal{V}_j^\Gamma \setminus \mathcal{I}_j^{\text{in}}| \leq \frac{\|\mathbf{L}_\Gamma^{(k)} - \mathbf{M}\|_{\text{F}}^2}{r/4} = \mathcal{O}\left(\frac{\log(\gamma_n)}{\gamma_n} n\right) = o(n). \quad (27)$$

Thus,  $|\mathcal{I}_j^{\text{in}}| \geq |\mathcal{V}_j^\Gamma \cap \mathcal{I}_j^{\text{in}}| = |\mathcal{V}_j^\Gamma| - |\mathcal{V}_j^\Gamma \setminus \mathcal{I}_j^{\text{in}}| \geq p_j n(1 - o(1))$ .

(ii) Similar to (i), we have  $|\mathcal{U}| \leq |\Gamma \setminus (\cup_{j \in [k]} \mathcal{I}_j^{\text{in}})| \leq \|\mathbf{L}_\Gamma^{(k)} - \mathbf{M}\|_{\text{F}}^2/(r/4) = \mathcal{O}\left(\frac{\log(\gamma_n)}{\gamma_n} n\right) = o(n)$ .

(iii) If node  $v \in \mathcal{U} \cap \Psi$ , then  $\mathcal{B}_v \cap (\cup_{j \in [k]} \mathcal{I}_j^{\text{in}}) = \emptyset$ , and as a result of (i), we have  $|\mathcal{B}_v| = \mathcal{O}\left(\frac{\log(\gamma_n)}{\gamma_n} n\right) = o(n)$ .

(iv) If node  $v \in \mathcal{I}_j^{\text{in}} \cap \Psi$ , then  $\mathcal{I}_j^{\text{in}} \subseteq \mathcal{B}_v$ .

(v) If nodes  $u \in \mathcal{I}_i^{\text{out}}$  and  $v \in \mathcal{I}_j^{\text{out}}$  such that  $i \neq j$ , then  $T_u \cap \mathcal{B}_v = \emptyset$ .

(vi) For all  $j \in [k]$ ,  $|\mathcal{I}_j^{\text{out}}| \leq p_j n(1 + o(1))$ . This is because  $|\mathcal{I}_j^{\text{in}}| \geq p_j n(1 - o(1))$  from (i), and thus

$$|\mathcal{I}_j^{\text{out}}| \leq n - |\Gamma| - \sum_{j' \neq j} |\mathcal{I}_{j'}^{\text{out}}| \leq n - \sum_{j' \neq j} |\mathcal{I}_{j'}^{\text{in}}| \leq n - (p_{j'} n(1 - o(1))) \leq p_j n(1 + o(1)).$$

<sup>2</sup>Let  $\mathbf{A}$  and  $\mathbf{B}$  be  $n \times n$  matrices, and their singular values are respectively denoted by  $\{\sigma_i(\mathbf{A})\}_{i \in [n]}$  and  $\{\sigma_i(\mathbf{B})\}_{i \in [n]}$ , both in decreasing orders. Weyl's inequality states that for every  $i \in [n]$ ,  $|\sigma_i(\mathbf{A}) - \sigma_i(\mathbf{B})| \leq \|\mathbf{A} - \mathbf{B}\|_{\text{op}}$ .

Lemma 5 below states that with high probability, each community contains at least one reference node.

**Lemma 5.** *With probability  $1 - o(1)$  over the selection of  $\Psi$ , we have  $|\mathcal{I}_j^{\text{in}} \cap \Psi| \geq 1$  for all  $j \in [k]$ .*

*Proof.* For any  $j \in [k]$ , the probability that a randomly selected reference node does not belong to  $\mathcal{I}_j^{\text{in}}$  is  $1 - (|\mathcal{I}_j^{\text{in}}|/n)$ , thus the probability that there exists at least one reference node belongs to  $\mathcal{I}_j^{\text{in}}$  is

$$1 - (1 - (|\mathcal{I}_j^{\text{in}}|/n))^{\log n} = 1 - \exp(-\Theta(n/\log n)). \quad (28)$$

Taking a union bound over all  $j \in [k]$ , we complete the proof.  $\square$

With these properties, we are able to show that running Stage 1 yields almost exact recovery. For ease of presentation, in the following we focus mainly on the case when  $p_1 > p_2 > \dots > p_k$  (such that the community sizes satisfy  $|\mathcal{V}_1^\Gamma| > |\mathcal{V}_2^\Gamma| > \dots > |\mathcal{V}_k^\Gamma|$  with high probability). The analysis can be easily generalized to the case when  $p_i = p_j$  for some  $i \neq j$ , but it requires more cumbersome notations (such as permutations) which makes the subsequent analysis more difficult to understand.

The “for loop” in lines 7 – 10: When  $j = 1$ , we will show that  $v_1^* \in \mathcal{I}_1^{\text{out}}$  and  $|\mathcal{B}_{v_1^*}| \geq np_1(1 - o(1))$ . From Lemma 5, there exists a node  $v_1 \in \mathcal{I}_1^{\text{in}} \cap \Psi$ , and its corresponding set  $\mathcal{B}_{v_1}$  is a superset of  $\mathcal{I}_1^{\text{in}}$  (due to (iv)). Thus, we have  $|\mathcal{B}_{v_1}| \geq |\mathcal{I}_1^{\text{in}}| \geq p_1 n(1 - o(1))$ , where the last inequality is due to (i). As  $|\mathcal{B}_{v_1^*}|$  is at least  $|\mathcal{B}_{v_1}|$ , we obtain that  $|\mathcal{B}_{v_1^*}| \geq p_1 n(1 - o(1))$ . To prove that  $v_1^* \in \mathcal{I}_1^{\text{out}}$ , one can verify that

- 1) for any  $v \in \mathcal{U} \cap \Psi$ ,  $|\mathcal{B}_v| = o(n) < |\mathcal{B}_{v_1^*}|$  by (iii);
- 2) for any  $v \in \mathcal{I}_j^{\text{out}} \cap \Psi$  where  $j \neq 1$ ,  $|\mathcal{B}_v| \leq |\mathcal{I}_j^{\text{out}}| + |\mathcal{U}| \leq np_j(1 + o(1)) + o(n) < |\mathcal{B}_{v_1^*}|$  (due to (vi) and the fact that  $p_1 > p_j$ ).

For  $2 \leq j \leq k$ , we will show that  $v_j^* \in \mathcal{I}_j^{\text{out}}$  and  $|\mathcal{B}_{v_j^*}| \geq np_j(1 - o(1))$ . Similar to the analysis above, there exists a node  $v_j \in \mathcal{I}_j^{\text{in}} \cap \Psi$  and  $\mathcal{B}_{v_j} \supseteq \mathcal{I}_j^{\text{in}}$ . From (v) we know that  $|\mathcal{B}_{v_j} \setminus (\cup_{l=0}^{j-1} \widehat{\mathcal{V}}_l^{(0)})| = |\mathcal{B}_{v_j}|$ , thus  $|\mathcal{B}_{v_j} \setminus (\cup_{l=0}^{j-1} \widehat{\mathcal{V}}_l^{(0)})| \geq |\mathcal{I}_j^{\text{in}}| \geq np_j(1 - o(1))$ . As  $|\mathcal{B}_{v_j^*} \setminus (\cup_{l=0}^{j-1} \widehat{\mathcal{V}}_l^{(0)})|$  is at least  $|\mathcal{B}_{v_j} \setminus (\cup_{l=0}^{j-1} \widehat{\mathcal{V}}_l^{(0)})|$ , we obtain that  $|\mathcal{B}_{v_j^*}| \geq np_j(1 - o(1))$ . To prove that  $v_j^* \in \mathcal{I}_j^{\text{out}}$ , one can verify that

- 1) for any  $v \in \mathcal{U} \cap \Psi$ ,  $|\mathcal{B}_v \setminus (\cup_{l=0}^{j-1} \widehat{\mathcal{V}}_l^{(0)})| = o(n)$ ;
- 2) for any  $v \in \mathcal{I}_{j'}^{\text{out}} \cap \Psi$  where  $j' > j$ ,  $|\mathcal{B}_v \setminus (\cup_{l=0}^{j-1} \widehat{\mathcal{V}}_l^{(0)})| \leq |\mathcal{B}_v| \leq |\mathcal{I}_{j'}^{\text{out}}| + |\mathcal{U}| \leq np_{j'}(1 + o(1)) + o(n) < |\mathcal{B}_{v_j^*}|$  (due to (vi) and the fact that  $p_j > p_{j'}$ );
- 3) for any  $v \in \mathcal{I}_{j'}^{\text{out}} \cap \Psi$  where  $j' < j$ , we have  $|\mathcal{B}_v \setminus (\cup_{l=0}^{j-1} \widehat{\mathcal{V}}_l^{(0)})| \leq |\mathcal{B}_v \setminus \widehat{\mathcal{V}}_{j'}^{(0)}| \leq |\mathcal{I}_{j'}^{\text{out}}| + |\mathcal{U}| - |\mathcal{B}_{v_{j'}^*}| = o(n)$ , since  $|\mathcal{B}_{v_{j'}^*}| \geq np_{j'}(1 - o(1))$ .

Therefore, under the assumption  $p_1 > p_2 > \dots > p_k$ , we have  $v_j^* \in \mathcal{I}_j^{\text{out}}$  for all  $j \in [k]$ .<sup>3</sup> This implies that all the elements in  $\{v_j^*\}_{j \in [k]}$  are far from each other—specifically, every pair of nodes  $(v_i^*, v_j^*)$  satisfies

$$\begin{aligned} \|(\mathbf{L}_\Gamma^{(k)})_{v_i^*} - (\mathbf{L}_\Gamma^{(k)})_{v_j^*}\|_2^2 &\geq \frac{1}{2} \|\mathbf{M}_{v_i^*} - (\mathbf{L}_\Gamma^{(k)})_{v_j^*}\|_2^2 - \|\mathbf{M}_{v_i^*} - (\mathbf{L}_\Gamma^{(k)})_{v_i^*}\|_2^2 \\ &\geq \frac{1}{2} \left( \frac{1}{2} \|\mathbf{M}_{v_j^*} - \mathbf{M}_{v_i^*}\|_2^2 - \|\mathbf{M}_{v_j^*} - (\mathbf{L}_\Gamma^{(k)})_{v_j^*}\|_2^2 \right) - \|\mathbf{M}_{v_i^*} - (\mathbf{L}_\Gamma^{(k)})_{v_i^*}\|_2^2 = \Omega(\gamma_n^2/n), \end{aligned} \quad (29)$$

since  $\|(\mathbf{L}_\Gamma^{(k)})_{v_i^*} - \mathbf{M}_{v_i^*}\|_2^2 \leq 4r$ ,  $\|(\mathbf{L}_\Gamma^{(k)})_{v_j^*} - \mathbf{M}_{v_j^*}\|_2^2 \leq 4r$ , and  $\|\mathbf{M}_{v_j^*} - \mathbf{M}_{v_i^*}\|_2^2 = \Omega(\gamma_n^2/n)$  by Lemma 3. If a node  $v \in \mathcal{V}_i^\Gamma$  is misclassified to  $\widehat{\mathcal{V}}_j^{(0)}$  (where  $i \neq j$ ) in the first “for loop”, then it must be close to the center of  $\widehat{\mathcal{V}}_j^{(0)}$ , i.e.,

<sup>3</sup>Without this assumption, we have that there exists a permutation  $\pi$  on  $[k]$  such that  $v_j^* \in \mathcal{I}_{\pi(j)}^{\text{out}}$  for all  $j \in [k]$ .

$\|(\mathbf{L}_\Gamma^{(k)})_v - (\mathbf{L}_\Gamma^{(k)})_{v_j^*}\|_2^2 \leq r$ . Thus, we have

$$\begin{aligned} \|(\mathbf{L}_\Gamma^{(k)})_v - \mathbf{M}_v\|_2^2 &\geq \frac{1}{2}\|\mathbf{M}_v - \mathbf{M}_{v_j^*}\|_2^2 - \|(\mathbf{L}_\Gamma^{(k)})_v - \mathbf{M}_{v_j^*}\|_2^2 \\ &\geq \frac{1}{2}\|\mathbf{M}_v - \mathbf{M}_{v_j^*}\|_2^2 - 2\|(\mathbf{L}_\Gamma^{(k)})_v - (\mathbf{L}_\Gamma^{(k)})_{v_j^*}\|_2^2 - 2\|(\mathbf{L}_\Gamma^{(k)})_{v_j^*} - \mathbf{M}_{v_j^*}\|_2^2 \\ &= \Omega(\gamma_n^2/n). \end{aligned} \quad (30)$$

The ‘‘for loop’’ in lines 11 – 14: Consider a specific node  $v \in \Gamma \setminus (\cup_{j=1}^k \widehat{\mathcal{V}}_j^{(0)})$ . If  $v \in \mathcal{V}_i^\Gamma$  is misclassified to  $\widehat{\mathcal{V}}_j^{(0)}$  (where  $i \neq j$ ) in the second ‘‘for loop’’, then it must be closer to the center of  $\widehat{\mathcal{V}}_j^{(0)}$  than the center of  $\widehat{\mathcal{V}}_i^{(0)}$ , i.e.,  $\|(\mathbf{L}_\Gamma^{(k)})_v - (\mathbf{L}_\Gamma^{(k)})_{v_j^*}\|_2^2 \leq \|(\mathbf{L}_\Gamma^{(k)})_v - (\mathbf{L}_\Gamma^{(k)})_{v_i^*}\|_2^2$ . Since the two centers are far from each other, one can show that  $v$  is far from the center of  $\widehat{\mathcal{V}}_i^{(0)}$ , i.e.,

$$\Omega(\gamma_n^2/n) \leq \|(\mathbf{L}_\Gamma^{(k)})_{v_i^*} - (\mathbf{L}_\Gamma^{(k)})_{v_j^*}\|_2^2 \leq 2\|(\mathbf{L}_\Gamma^{(k)})_{v_i^*} - (\mathbf{L}_\Gamma^{(k)})_v\|_2^2 + 2\|(\mathbf{L}_\Gamma^{(k)})_{v_j^*} - (\mathbf{L}_\Gamma^{(k)})_v\|_2^2 \leq 4\|(\mathbf{L}_\Gamma^{(k)})_{v_i^*} - (\mathbf{L}_\Gamma^{(k)})_v\|_2^2. \quad (31)$$

Thus, node  $v$  must satisfy

$$\|(\mathbf{L}_\Gamma^{(k)})_v - \mathbf{M}_v\|_2^2 \geq \frac{1}{2}\|(\mathbf{L}_\Gamma^{(k)})_v - (\mathbf{L}_\Gamma^{(k)})_{v_i^*}\|_2^2 - \|\mathbf{M}_v - (\mathbf{L}_\Gamma^{(k)})_{v_i^*}\|_2^2 = \Omega(\gamma_n^2/n). \quad (32)$$

Combining Eqns. (30) and (32), we conclude that for any node  $v \in \Gamma$ , if it is misclassified to another cluster, it must satisfy  $\|(\mathbf{L}_\Gamma^{(k)})_v - \mathbf{M}_v\|_2^2 = \Omega(\gamma_n^2/n)$ . Since  $\sum_{v \in \Gamma} \|(\mathbf{L}_\Gamma^{(k)})_v - \mathbf{M}_v\|_2^2 = \|\mathbf{L}_\Gamma^{(k)} - \mathbf{M}\|_F^2 = \mathcal{O}(\gamma_n)$  (by Lemma 4), we know that the number of misclassified nodes in  $\Gamma$  is at most  $\mathcal{O}(n/\gamma_n)$ . Taking into account the number of nodes that do not belong to  $\Gamma$  (which also scales as  $\mathcal{O}(n/\gamma_n)$  by Lemma 2), we complete the proof of Theorem 3.

## B. Theoretical guarantees of Stage 2

From Theorem 3 we know that for a fixed ground-truth community vector  $\mathbf{z} \in \mathcal{A}_Z$ , running Stage 1 on  $\widetilde{G}_1$  ensures that  $l(\mathbf{z}, \widehat{\mathbf{Z}}^{(0)}) \leq \eta_n$  with probability at least  $1 - \epsilon_n$ . In the following, we show that the hypergraph spectral clustering method does not only work well on  $\widetilde{G}_1$ , but also works well on the sub-HSBM  $G_1$  (which is generated on the fixed sub-hypergraph  $F_1$ ) with high probability over the graph splitting process.

**Definition 4.** Let  $(f_1, f_2)$  be the realizations of the sub-HSBMs  $(F_1, F_2)$ , and consider a fixed ground-truth community vector  $\mathbf{z}$ .

- We say  $f_1$  is a good realization of the first sub-hypergraph with respect to  $\mathbf{z}$  (denoted by  $f_1 \in \mathcal{G}_1^z$ ) if the probability that ‘‘running Stage 1 on  $G_1$  (which depends on  $f_1$ ) ensures  $l(\mathbf{z}, \widehat{\mathbf{Z}}^{(0)}) \leq \eta_n$ ’’ is at least  $1 - \sqrt{\epsilon_n}$ , i.e.,  $\mathbb{P}_{G_1}(l(\mathbf{z}, \widehat{\mathbf{Z}}^{(0)}) \leq \eta_n) \geq 1 - \sqrt{\epsilon_n}$ .
- We say  $f_2$  is a good realization of the second sub-hypergraph with respect to  $\mathbf{z}$  (denoted by  $f_2 \in \mathcal{G}_2^z$ ) if for every node  $v \in [n]$  and every community assignment  $\mathbf{m} \in \mathcal{M}$ , the number of hyperedges in  $f_2$  that includes node  $v$  and other  $d - 1$  nodes with community assignment  $\mathbf{m}$  (denoted by  $D_{v,\mathbf{m}}$ ) satisfies  $(1 - (3\gamma_n/\log n))R_{\mathbf{m}} \leq D_{v,\mathbf{m}} \leq (1 + n^{-\frac{1}{2} + \frac{\delta}{2}})R_{\mathbf{m}}$ .

**Lemma 6.** Suppose  $\mathbf{z} \in \mathcal{A}_Z$ . With probability at least  $1 - 2\sqrt{\epsilon_n}$ , the randomly generated sub-hypergraphs  $(F_1, F_2)$  satisfy  $F_1 \in \mathcal{G}_1^z$  and  $F_2 \in \mathcal{G}_2^z$  simultaneously.

*Proof.* See Appendix B. □

Due to Lemma 6, one can then focus on a specific typical ground-truth community vector  $\mathbf{z} \in \mathcal{A}_Z$  and good realizations of the sub-hypergraphs  $f_1 \in \mathcal{G}_1^z$  and  $f_2 \in \mathcal{G}_2^z$  in the following. Also, one can suppose that the initial estimate  $\widehat{\mathbf{z}}^{(0)}$  (after Stage 1) satisfies  $l(\mathbf{z}, \widehat{\mathbf{z}}^{(0)}) \leq \eta_n$ , since running Stage 1 on  $G_1$  (which depends on the good realization  $f_1$ ) ensures  $l(\mathbf{z}, \widehat{\mathbf{z}}^{(0)}) \leq \eta_n$  with high probability. Without loss of generality, we further assume that the minimum value of  $l(\mathbf{z}, \widehat{\mathbf{z}}^{(0)}) = \min_{\pi \in \mathcal{S}_k} \frac{1}{n} \sum_{v \in [n]} \mathbb{1}\{\widehat{z}_v^{(0)} \neq \pi(v)\}$

$\pi(z_v)$  is achieved by the identity permutation in  $\mathcal{S}_k$ , and this assumption helps us to remove the presence of the permutation in the following analysis.

Note that the local MAP estimation in (14) for each node  $v \in [n]$  can be alternatively represented as

$$\widehat{Z}_v(g_2, \widehat{\mathbf{z}}_{\sim v}^{(0)}) = \arg \max_{i \in [k]} \mathbb{P} \left( Z_v = i | G_2 = g_2, \widehat{\mathbf{Z}}_{\sim v}^{(0)} = \widehat{\mathbf{z}}_{\sim v}^{(0)} \right) \quad (33)$$

$$= \arg \max_{i \in [k]} \mathbb{P} \left( G_2 = g_2 | Z_v = i, \widehat{\mathbf{Z}}_{\sim v}^{(0)} = \widehat{\mathbf{z}}_{\sim v}^{(0)} \right) \cdot p_i. \quad (34)$$

Thus, for a fixed node  $v \in \mathcal{V}_i$  (i.e.,  $z_v = i$ ) for some  $i \in [k]$ , the probability that it is misclassified to a different community can be bounded as follows:

$$\mathbb{P} \left( \widehat{Z}_v(G_2, \widehat{\mathbf{z}}_{\sim v}^{(0)}) \neq z_v \right) = \mathbb{P}_{G_2} \left( \arg \max_{i' \in [k]} \mathbb{P} \left( G_2 | Z_v = i', \widehat{\mathbf{z}}_{\sim v}^{(0)} \right) \cdot p_{i'} \neq i \right) \quad (35)$$

$$= \mathbb{P}_{G_2} \left( \exists j \neq i : \mathbb{P} \left( G_2 | Z_v = j, \widehat{\mathbf{z}}_{\sim v}^{(0)} \right) \cdot p_j \geq \mathbb{P} \left( G_2 | Z_v = i, \widehat{\mathbf{z}}_{\sim v}^{(0)} \right) \cdot p_i \right) \quad (36)$$

$$\leq \sum_{j \neq i} \mathbb{P}_{G_2} \left( \mathbb{P} \left( G_2 | Z_v = j, \widehat{\mathbf{z}}_{\sim v}^{(0)} \right) \cdot p_j \geq \mathbb{P} \left( G_2 | Z_v = i, \widehat{\mathbf{z}}_{\sim v}^{(0)} \right) \cdot p_i \right) \quad (37)$$

$$= \sum_{j \neq i} \sum_{g_2} \mathbb{P} \left( g_2 | Z_v = i, \mathbf{z}_{\sim v} \right) \times \mathbb{1} \left\{ \mathbb{P} \left( g_2 | Z_v = j, \widehat{\mathbf{z}}_{\sim v}^{(0)} \right) \cdot p_j \geq \mathbb{P} \left( g_2 | Z_v = i, \widehat{\mathbf{z}}_{\sim v}^{(0)} \right) \cdot p_i \right\}. \quad (38)$$

Note that the above error probability depends only on the hyperedges in  $G_2$  (which is generated on the sub-hypergraph  $f_2$ ) that contains node  $v$ . Recall that the number of hyperedges in  $f_2$  that comprise node  $v$  and other  $d-1$  nodes with community assignment  $\mathbf{m}$  is  $D_{v,\mathbf{m}}$  (as defined in Definition 4), thus these hyperedges can be represented by Bernoulli random variables  $\{X_a^{\mathbf{m}}\}_{a=1}^{D_{v,\mathbf{m}}}$ , where  $X_a^{\mathbf{m}} = 1$  means the presence of this hyperedge in  $G_2$ . Taking all possible  $\mathbf{m} \in \mathcal{M}$  into account, we know that the hyperedges in  $f_2$  that contains node  $v$  can be represented by the collection of Bernoulli random variables  $\{\{X_a^{\mathbf{m}}\}_{a=1}^{D_{v,\mathbf{m}}}\}_{\mathbf{m} \in \mathcal{M}}$ . We also denote  $D_v \triangleq \sum_{\mathbf{m} \in \mathcal{M}} D_{v,\mathbf{m}}$  as the total number of hyperedges in  $f_2$  that contains  $v$ , where  $D_v$  scales as  $\Theta(n^{d-1})$  by Lemma 6.

To further upper bound (38), we would like to substitute the terms  $\mathbb{P}(g_2 | Z_v = i, \widehat{\mathbf{z}}_{\sim v}^{(0)})$  and  $\mathbb{P}(g_2 | Z_v = j, \widehat{\mathbf{z}}_{\sim v}^{(0)})$  in the indicator function by  $\mathbb{P}(g_2 | Z_v = i, \mathbf{z}_{\sim v})$  and  $\mathbb{P}(g_2 | Z_v = j, \mathbf{z}_{\sim v})$  respectively; thus we can get rid of the fact that  $\widehat{\mathbf{z}}_{\sim v}^{(0)}$  is not exactly the same as  $\mathbf{z}_{\sim v}$ . We now consider the ratio between  $\mathbb{P}(G_2 | Z_v = j, \widehat{\mathbf{z}}_{\sim v}^{(0)})$  and  $\mathbb{P}(G_2 | Z_v = j, \mathbf{z}_{\sim v})$  for an arbitrary  $j \in [k]$ . Due to the independence of hyperedges  $\{\{X_a^{\mathbf{m}}\}_{a=1}^{D_{v,\mathbf{m}}}\}_{\mathbf{m} \in \mathcal{M}}$  in  $G_2$ , we have

$$\frac{\mathbb{P} \left( G_2 | Z_v = j, \widehat{\mathbf{z}}_{\sim v}^{(0)} \right)}{\mathbb{P} \left( G_2 | Z_v = j, \mathbf{z}_{\sim v} \right)} = \prod_{\mathbf{m} \in \mathcal{M}} \prod_{a=1}^{D_{v,\mathbf{m}}} \frac{\mathbb{P} \left( X_a^{\mathbf{m}} | Z_v = j, \widehat{\mathbf{z}}_{\sim v}^{(0)} \right)}{\mathbb{P} \left( X_a^{\mathbf{m}} | Z_v = j, \mathbf{z}_{\sim v} \right)} \quad (39)$$

Conditioned on  $Z_v = j$  and the ground truth  $\mathbf{z}_{\sim v}$ , the hyperedge  $X_a^{\mathbf{m}}$  includes node  $v \in \mathcal{V}_j$  and  $d-1$  other nodes with community assignment  $\mathbf{m}$ , thus  $\mathbb{P}(X_a^{\mathbf{m}} = 1 | Z_v = j, \mathbf{z}_{\sim v}) = \frac{Q_{\mathbf{m} \oplus j} \log n}{n^{d-1}}$ . On the other hand, conditioned on  $Z_v = j$  and the estimated labels  $\widehat{\mathbf{z}}_{\sim v}^{(0)}$ , the hyperedge  $X_a^{\mathbf{m}}$  is considered to include node  $v \in \mathcal{V}_j$ , but the community assignment of the other  $d-1$  nodes, denoted by  $\mathbf{m}' \in \mathcal{M}$ , may not be equal to  $\mathbf{m}$ , and we have  $\mathbb{P}(X_a^{\mathbf{m}} = 1 | Z_v = j, \widehat{\mathbf{z}}_{\sim v}^{(0)}) = \frac{Q_{\mathbf{m}' \oplus j} \log n}{n^{d-1}}$ . Here,  $\mathbf{m}' = \mathbf{m}$  if all the  $d-1$  nodes (except for  $v$ ) in this hyperedge are not misclassified in  $\widehat{\mathbf{z}}_{\sim v}^{(0)}$ , while  $\mathbf{m}' \neq \mathbf{m}$  otherwise. For hyperedges such that their community assignments  $\mathbf{m}'$  (under  $\widehat{\mathbf{z}}_{\sim v}^{(0)}$ ) equal  $\mathbf{m}$ , they are cancelled out in (39) since the distributions are invariant conditioned on either  $\widehat{\mathbf{z}}_{\sim v}^{(0)}$  or  $\mathbf{z}_{\sim v}$ . It then remains to focus on the hyperedges that contain at least one misclassified node in  $\widehat{\mathbf{z}}_{\sim v}^{(0)}$ . Since  $l(\mathbf{z}, \widehat{\mathbf{z}}^{(0)}) \leq \eta_n$ , the number of such hyperedges is at most  $\eta_n \cdot \binom{n-2}{d-2} \triangleq D'_v$ , where  $D'_v$  scales as  $\mathcal{O}(\eta_n n^{d-1})$ . For ease of presentation, these  $D'_v$  Bernoulli random variables in  $\{\{X_a^{\mathbf{m}}\}_{a=1}^{D_{v,\mathbf{m}}}\}_{\mathbf{m} \in \mathcal{M}}$  are alternatively relabelled as  $\{Y_a\}_{a=1}^{D'_v}$ .

**Definition 5.** Let  $\{c_n\}$  be a vanishing sequence that satisfy  $c_n \log(c_n/\eta_n) = \omega(1)$ . We say  $G_2 \in \mathcal{G}_v$  if the random variables  $\{Y_a\}_{a=1}^{D'_v}$  satisfies  $\sum_{a=1}^{D'_v} Y_a \leq c_n \log n$ , and  $G_2 \notin \mathcal{G}_v$  otherwise.

Lemma 7 below states that  $G_2 \in \mathcal{G}_v$  with high probability (conditioned on the ground truth  $Z_v = i$  and  $\mathbf{z}_{\sim v}$ ), and for every realization  $g_2 \in \mathcal{G}_v$ , the ratio between  $\mathbb{P}(g_2|Z_v = j, \widehat{\mathbf{z}}_{\sim v}^{(0)})$  and  $\mathbb{P}(g_2|Z_v = j, \mathbf{z}_{\sim v})$  is bounded for any  $j \in [k]$ .

**Lemma 7.** *Suppose the initial estimate  $\widehat{\mathbf{z}}^{(0)}$  satisfies  $l(\mathbf{z}, \widehat{\mathbf{z}}^{(0)}) \leq \eta_n$ . The probability that  $G_2$  does not belong to  $\mathcal{G}_v$  is at most  $n^{-\omega(1)}$ , i.e.,*

$$\mathbb{P}(G_2 \notin \mathcal{G}_v | Z_v = i, \mathbf{z}_{\sim v}) = \mathbb{P}\left(\sum_{a=1}^{D'_v} Y_a > c_n \log n | Z_v = i, \mathbf{z}_{\sim v}\right) \leq n^{-\omega(1)}. \quad (40)$$

Furthermore, there exist  $L_h = n^{o(1)}$  and  $L_l = n^{-o(1)}$  such that for all  $g_2 \in \mathcal{G}_v$  and  $j \in [k]$ ,

$$L_l \leq \frac{\mathbb{P}(g_2 | Z_v = j, \widehat{\mathbf{z}}_{\sim v}^{(0)})}{\mathbb{P}(g_2 | Z_v = j, \mathbf{z}_{\sim v})} \leq L_h. \quad (41)$$

*Proof.* See Appendix C. □

We now return to (38). When  $g_2 \in \mathcal{G}_v$  and  $\mathbb{P}(g_2|Z_v = j, \widehat{\mathbf{z}}_{\sim v}^{(0)}) \cdot p_j \geq \mathbb{P}(g_2|Z_v = i, \widehat{\mathbf{z}}_{\sim v}^{(0)}) \cdot p_i$ , we have  $\mathbb{P}(g_2|Z_v = i, \mathbf{z}_{\sim v}) \leq (p_j L_u / p_i L_l) \mathbb{P}(g_2|Z_v = j, \mathbf{z}_{\sim v})$  according to Lemma 7. Thus, one can upper-bound (38) as

$$\begin{aligned} & \sum_{j \neq i} \sum_{g_2 \in \mathcal{G}_v} \mathbb{P}(g_2 | Z_v = i, \mathbf{z}_{\sim v}) \mathbb{1} \left\{ \mathbb{P}(g_2 | Z_v = j, \widehat{\mathbf{z}}_{\sim v}^{(0)}) \cdot p_j \geq \mathbb{P}(g_2 | Z_v = i, \widehat{\mathbf{z}}_{\sim v}^{(0)}) \cdot p_i \right\} + \sum_{g_2 \notin \mathcal{G}_v} \mathbb{P}(g_2 | Z_v = i, \mathbf{z}_{\sim v}) \\ & \leq \sum_{j \neq i} \sum_{g_2 \in \mathcal{G}_v} \frac{p_j L_u}{p_i L_l} \min \{ \mathbb{P}(g_2 | Z_v = i, \mathbf{z}_{\sim v}), \mathbb{P}(g_2 | Z_v = j, \mathbf{z}_{\sim v}) \} + n^{-\omega(1)} \end{aligned} \quad (42)$$

$$\leq \sum_{j \neq i} \frac{p_j L_u}{p_i L_l} \sum_{g_2} \min \{ \mathbb{P}(g_2 | Z_v = i, \mathbf{z}_{\sim v}), \mathbb{P}(g_2 | Z_v = j, \mathbf{z}_{\sim v}) \} + n^{-\omega(1)}. \quad (43)$$

Lemma 8 below bounds the term  $\sum_{g_2} \min \{ \mathbb{P}(g_2 | Z_v = i, \mathbf{z}_{\sim v}), \mathbb{P}(g_2 | Z_v = j, \mathbf{z}_{\sim v}) \}$  for any  $i \neq j$  in terms of the GCH-divergence  $D_+(i, j)$  between communities  $\mathcal{V}_i$  and  $\mathcal{V}_j$ .

**Lemma 8.** *When the ground-truth community vector  $\mathbf{z} \in \mathcal{A}_Z$  and the sub-hypergraphs  $f_1 \in \mathcal{G}_1^{\mathbf{z}}$  and  $f_2 \in \mathcal{G}_2^{\mathbf{z}}$ , we have*

$$\sum_{g_2} \min \{ \mathbb{P}(g_2 | Z_v = i, \mathbf{z}_{\sim v}), \mathbb{P}(g_2 | Z_v = j, \mathbf{z}_{\sim v}) \} \leq n^{-D_+(i, j) + o(1)}, \quad (44)$$

where  $D_+(i, j) = \max_{t \in [0, 1]} \sum_{\mathbf{m} \in \mathcal{M}} t \mu_{\mathbf{m} \oplus i} + (1 - t) \mu_{\mathbf{m} \oplus j} - \mu_{\mathbf{m} \oplus i}^t \mu_{\mathbf{m} \oplus j}^{1-t}$ .

*Proof of Lemma 8.* As  $G_2$  is equivalent to the collection of random variables  $\{ \{X_a^{\mathbf{m}}\}_{a=1}^{D_{v, \mathbf{m}}} \}_{\mathbf{m} \in \mathcal{M}}$ , we have

$$\begin{aligned} & \sum_{g_2} \min \{ \mathbb{P}(g_2 | Z_v = i, \mathbf{z}_{\sim v}), \mathbb{P}(g_2 | Z_v = j, \mathbf{z}_{\sim v}) \} \\ & = \sum_{\mathbf{m} \in \mathcal{M}} \sum_{\{x_a^{\mathbf{m}}\}_{a=1}^{D_{v, \mathbf{m}}}} \min \left\{ \prod_{\mathbf{m} \in \mathcal{M}} \prod_{a=1}^{D_{v, \mathbf{m}}} \mathbb{P}(x_a^{\mathbf{m}} | Z_v = i, \mathbf{z}_{\sim v}), \prod_{\mathbf{m} \in \mathcal{M}} \prod_{a=1}^{D_{v, \mathbf{m}}} \mathbb{P}(x_a^{\mathbf{m}} | Z_v = j, \mathbf{z}_{\sim v}) \right\} \end{aligned} \quad (45)$$

$$\leq \sum_{\mathbf{m} \in \mathcal{M}} \sum_{\{x_a^{\mathbf{m}}\}_{a=1}^{D_{v, \mathbf{m}}}} \left( \prod_{\mathbf{m} \in \mathcal{M}} \prod_{a=1}^{D_{v, \mathbf{m}}} \mathbb{P}(x_a^{\mathbf{m}} | Z_v = i, \mathbf{z}_{\sim v}) \right)^t \left( \prod_{\mathbf{m} \in \mathcal{M}} \prod_{a=1}^{D_{v, \mathbf{m}}} \mathbb{P}(x_a^{\mathbf{m}} | Z_v = j, \mathbf{z}_{\sim v}) \right)^{1-t} \quad (46)$$

$$= \prod_{\mathbf{m} \in \mathcal{M}} \prod_{a=1}^{D_{v, \mathbf{m}}} \sum_{x_a^{\mathbf{m}} \in \{0, 1\}} \mathbb{P}(x_a^{\mathbf{m}} | Z_v = i, \mathbf{z}_{\sim v})^t \mathbb{P}(x_a^{\mathbf{m}} | Z_v = j, \mathbf{z}_{\sim v})^{1-t} \quad (47)$$

$$= \prod_{\mathbf{m} \in \mathcal{M}} \left[ \left( \frac{Q_{\mathbf{m} \oplus i} \log n}{n^{d-1}} \right)^t \left( \frac{Q_{\mathbf{m} \oplus j} \log n}{n^{d-1}} \right)^{1-t} + \left( 1 - \frac{Q_{\mathbf{m} \oplus i} \log n}{n^{d-1}} \right)^t \left( 1 - \frac{Q_{\mathbf{m} \oplus j} \log n}{n^{d-1}} \right)^{1-t} \right]^{D_{v, \mathbf{m}}}, \quad (48)$$



where (46)-(48) hold for any  $t \in [0, 1]$ . By applying a Taylor series expansion, we have

$$\begin{aligned} & \left(1 - \frac{Q_{\mathbf{m} \oplus i} \log n}{n^{d-1}}\right)^t \left(1 - \frac{Q_{\mathbf{m} \oplus j} \log n}{n^{d-1}}\right)^{1-t} \\ &= \exp \left\{ \log \left\{ \frac{(n^{d-1} - Q_{\mathbf{m} \oplus i} \log n)^t (n^{d-1} - Q_{\mathbf{m} \oplus j} \log n)^{1-t}}{n^{d-1}} \right\} \right\} \end{aligned} \quad (49)$$

$$= \exp \left\{ t \log \left( n^{d-1} \left(1 - \frac{Q_{\mathbf{m} \oplus i} \log n}{n^{d-1}}\right) \right) + (1-t) \log \left( n^{d-1} \left(1 - \frac{Q_{\mathbf{m} \oplus j} \log n}{n^{d-1}}\right) \right) - \log(n^{d-1}) \right\} \quad (50)$$

$$= \exp \left\{ -t \frac{Q_{\mathbf{m} \oplus i} \log n}{n^{d-1}} - (1-t) \frac{Q_{\mathbf{m} \oplus j} \log n}{n^{d-1}} - \frac{1}{2} (tQ_{\mathbf{m} \oplus i}^2 + (1-t)Q_{\mathbf{m} \oplus j}^2) \frac{(\log n)^2}{n^{2d-2}} + \mathcal{O} \left( \frac{(\log n)^3}{n^{3d-3}} \right) \right\} \quad (51)$$

$$= 1 - t \frac{Q_{\mathbf{m} \oplus i} \log n}{n^{d-1}} - (1-t) \frac{Q_{\mathbf{m} \oplus j} \log n}{n^{d-1}} - \mathcal{O} \left( \frac{(\log n)^2}{n^{2d-2}} \right), \quad (52)$$

where in (52) one needs to be aware that the lower order term has a negative coefficient. Thus, one can rewrite (48) as

$$\begin{aligned} & \exp \left\{ \sum_{\mathbf{m} \in \mathcal{M}} D_{v, \mathbf{m}} \cdot \log \left[ 1 - t \frac{Q_{\mathbf{m} \oplus i} \log n}{n^{d-1}} - (1-t) \frac{Q_{\mathbf{m} \oplus j} \log n}{n^{d-1}} + \left( \frac{Q_{\mathbf{m} \oplus i} \log n}{n^{d-1}} \right)^t \left( \frac{Q_{\mathbf{m} \oplus j} \log n}{n^{d-1}} \right)^{1-t} - \mathcal{O} \left( \frac{(\log n)^2}{n^{2d-2}} \right) \right] \right\} \\ &= \exp \left\{ -(\log n) \sum_{\mathbf{m} \in \mathcal{M}} \frac{D_{v, \mathbf{m}}}{n^{d-1}} \cdot \left( tQ_{\mathbf{m} \oplus i} + (1-t)Q_{\mathbf{m} \oplus j} - Q_{\mathbf{m} \oplus i}^t Q_{\mathbf{m} \oplus j}^{1-t} + \mathcal{O} \left( \frac{\log n}{n^{d-1}} \right) \right) \right\} \end{aligned} \quad (53)$$

$$\leq n^{-\sum_{\mathbf{m} \in \mathcal{M}} R'_{\mathbf{m}} (tQ_{\mathbf{m} \oplus i} + (1-t)Q_{\mathbf{m} \oplus j} - Q_{\mathbf{m} \oplus i}^t Q_{\mathbf{m} \oplus j}^{1-t}) + \mathcal{O}(\gamma_n / \log n)}, \quad (54)$$

where (54) is due to the fact that  $(1 - (3\gamma_n / \log n)) R_{\mathbf{m}} \leq D_{v, \mathbf{m}} \leq (1 + n^{-\frac{1}{2} + \frac{\delta}{2}}) R_{\mathbf{m}}$ . Since (54) is valid for any  $t \in [0, 1]$  and recall that  $\mu_{\mathbf{m} \oplus i} = R'_{\mathbf{m}} Q_{\mathbf{m} \oplus i}$ , we eventually obtain that

$$\sum_{g_2} \min \{ \mathbb{P}(g_2 | Z_v = i, \mathbf{z}_{\sim v}), \mathbb{P}(g_2 | Z_v = j, \mathbf{z}_{\sim v}) \} \leq n^{-\max_{t \in [0, 1]} \sum_{\mathbf{m} \in \mathcal{M}} t\mu_{\mathbf{m} \oplus i} + (1-t)\mu_{\mathbf{m} \oplus j} - \mu_{\mathbf{m} \oplus i}^t \mu_{\mathbf{m} \oplus j}^{1-t} + o(1)} \quad (55)$$

$$= n^{-D_+(i, j) + o(1)}. \quad (56)$$

This completes the proof of Lemma 8.  $\square$

Since  $\min_{i, j \in [k]: i \neq j} D_+(i, j) > 1$ , there exists an  $\varepsilon > 0$  such that  $D_+(i, j) > 1 + \varepsilon$  for all  $i \neq j$ . By also noting that  $L_u/L_l = n^{o(1)}$ , we have that  $\sum_{j \neq i} (p_j L_u / p_i L_l) \sum_{g_2} \min \{ \mathbb{P}(g_2 | Z_v = i, \mathbf{z}_{\sim v}), \mathbb{P}(g_2 | Z_v = j, \mathbf{z}_{\sim v}) \} \leq n^{-(1+\varepsilon)+o(1)}$ . Thus, one can bound the error probability for node  $v$  from above as

$$\mathbb{P}(\widehat{Z}_v \neq z_v) \leq n^{-(1+\varepsilon)+o(1)} + n^{-\omega(1)}. \quad (57)$$

Note that the above analysis is also valid for nodes that belong to any other communities (not necessarily community  $\mathcal{V}_i$ ), thus one can take a union bound over all the  $n$  nodes to obtain that

$$\mathbb{P}(\exists v \in [n] : \widehat{Z}_v \neq z_v) = \mathbb{P}(\widehat{\mathbf{Z}} \neq \mathbf{z}) \leq n^{-\varepsilon/2} \quad (58)$$

when  $n$  is sufficiently large. This means that all the nodes can be recovered correctly with probability at least  $1 - n^{-\varepsilon/2}$ .

### C. The Overall Success Probability

Let  $\mathcal{E}_{\text{suc}}$  be the event that  $l(\mathbf{z}, \widehat{\mathbf{Z}}) = 0$ . From the analysis of Stage 2, we know that for all  $\mathbf{z} \in \mathcal{A}_Z$ ,  $f_1 \in \mathcal{G}_1^z$ ,  $f_2 \in \mathcal{G}_2^z$ , and  $\widehat{\mathbf{z}}^{(0)}$  satisfying  $l(\mathbf{z}, \widehat{\mathbf{z}}) \leq \eta_n$ ,

$$\mathbb{P}(\mathcal{E}_{\text{suc}} | \mathbf{z}, f_1, f_2, \widehat{\mathbf{z}}^{(0)}) \geq 1 - n^{-\varepsilon/2}, \quad (59)$$

Therefore, the overall success probability is

$$\mathbb{P}\left(l(\mathbf{Z}, \widehat{\mathbf{Z}}) = 0\right) = \sum_{\mathbf{z}} \mathbb{P}(\mathbf{z}) \sum_{f_1, f_2} \mathbb{P}(f_1, f_2) \sum_{\widehat{\mathbf{z}}^{(0)}} \mathbb{P}(\widehat{\mathbf{z}}^{(0)} | \mathbf{z}, f_1, f_2) \cdot \mathbb{P}(\mathcal{E}_{\text{suc}} | \mathbf{z}, f_1, f_2, \widehat{\mathbf{z}}^{(0)}) \quad (60)$$

$$\geq \sum_{\mathbf{z} \in \mathcal{A}_Z} \mathbb{P}(\mathbf{z}) \sum_{f_1 \in \mathcal{G}_1^Z, f_2 \in \mathcal{G}_2^Z} \mathbb{P}(f_1, f_2) \sum_{\widehat{\mathbf{z}}^{(0)}: l(\widehat{\mathbf{z}}^{(0)}, \mathbf{z}) \leq \eta_n} \mathbb{P}(\widehat{\mathbf{z}}^{(0)} | \mathbf{z}, f_1, f_2) \cdot \mathbb{P}(\mathcal{E}_{\text{suc}} | \mathbf{z}, f_1, f_2, \widehat{\mathbf{z}}^{(0)}) \quad (61)$$

$$\geq (1 - \exp(-\Theta(n^\delta))) \cdot (1 - 2\sqrt{\epsilon_n}) \cdot (1 - \sqrt{\epsilon_n}) \cdot (1 - n^{-\epsilon/2}) \quad (62)$$

$$= 1 - o(1), \quad (63)$$

where inequality (62) follows from Lemma 1, Lemma 6, the definition of good realization  $f_1$  in Definition 4, and Eqn. (59). This means that exact recovery is achievable.

## VI. PROOF OF CONVERSE (THEOREM 1)

In this section, we show that when the model parameters  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}})$  satisfy  $\min_{i, j \in [k]: i \neq j} D_+(i, j) < 1$ , exact recovery is impossible. This converse proof is inspired by that for the SBM [4], but is adapted to the  $d$ -HSBM setting.

First, we recall from Lemma 1 that with high probability the number of nodes in each community  $\mathcal{V}_j$  is tightly concentrated around the expectation  $np_j$  (i.e., the ground-truth community vector  $\mathbf{Z} \in \mathcal{A}_Z$ ). Hence, we consider a fixed  $\mathbf{z} \in \mathcal{A}_Z$  from now on. Let  $S$  be a random set that contains  $n/(\log n)^3$  randomly selected nodes from  $[n]$ . By applying the Chernoff bound, we can show that with probability  $1 - \exp(-\Theta(n^\delta))$  over the selection process, the number of nodes in both  $\mathcal{V}_j$  and  $S$ , denoted by  $\mathcal{V}_j^S$ , satisfies

$$(1 - n^{-\frac{1}{2} + \frac{\delta}{2}})^2 \frac{np_j}{(\log n)^3} \leq |\mathcal{V}_j^S| \leq (1 + n^{-\frac{1}{2} + \frac{\delta}{2}})^2 \frac{np_j}{(\log n)^3}, \quad \forall j \in [k], \quad (64)$$

and thus the number of nodes in both  $\mathcal{V}_j$  and  $S^c$ , denoted by  $\mathcal{V}_j^{S^c}$ , satisfies

$$np_j \left(1 - \frac{1}{(\log n)^3} - 2n^{-\frac{1}{2} + \frac{\delta}{2}}\right) \leq |\mathcal{V}_j^{S^c}| \leq np_j \left(1 - \frac{1}{(\log n)^3} + 2n^{-\frac{1}{2} + \frac{\delta}{2}}\right), \quad \forall j \in [k] \quad (65)$$

for sufficiently large  $n$ . We then consider a fixed set  $S$  that satisfies (64) and (65). Let

$$f_{i,j}(t) \triangleq \sum_{\mathbf{m} \in \mathcal{M}} t\mu_{\mathbf{m} \oplus i} + (1-t)\mu_{\mathbf{m} \oplus j} - \mu_{\mathbf{m} \oplus i}^t \mu_{\mathbf{m} \oplus j}^{1-t}, \quad (66)$$

and note that  $D_+(i, j) = \max_{t \in [0, 1]} f_{i,j}(t)$ . To obtain the maximizer of  $f_{i,j}(t)$ , we set  $f'_{i,j}(t) = 0$  and this implies

$$\sum_{\mathbf{m} \in \mathcal{M}} \mu_{\mathbf{m} \oplus i}^t \mu_{\mathbf{m} \oplus j}^{1-t} \log \left( \frac{\mu_{\mathbf{m} \oplus i}}{\mu_{\mathbf{m} \oplus j}} \right) = \sum_{\mathbf{m} \in \mathcal{M}} \mu_{\mathbf{m} \oplus i} - \mu_{\mathbf{m} \oplus j}. \quad (67)$$

Let  $\tau_{\mathbf{m}} \triangleq \lfloor \mu_{\mathbf{m} \oplus i}^{t^*} \mu_{\mathbf{m} \oplus j}^{1-t^*} \log n \rfloor$  for each  $\mathbf{m} \in \mathcal{M}$ , where  $t^*$  is set to satisfy (67).

**Definition 6.** For each node  $v \in S$ , let  $N_v^{\mathbf{m}}$  denote the number of hyperedges that contains  $v$  and other  $d-1$  nodes from  $[n] \setminus S$  that have community assignment  $\mathbf{m} \in \mathcal{M}$ . A node  $v \in S$  is said to be ambiguous if  $N_v^{\mathbf{m}} = \tau_{\mathbf{m}}$  for all  $\mathbf{m} \in \mathcal{M}$ .

In the following, we show that if  $D_+(i, j) < 1$ , there is at least one ambiguous node in  $\mathcal{V}_i^S$  and one ambiguous node in  $\mathcal{V}_j^S$ . For a node  $v \in \mathcal{V}_i^S$ ,  $N_v^{\mathbf{m}}$  equals the sum of  $\widetilde{R}_{\mathbf{m}}$  i.i.d. Bernoulli random variables  $\{B_r^{\mathbf{m}}\}_{r=1}^{\widetilde{R}_{\mathbf{m}}}$  with expectation  $Q_{\mathbf{m} \oplus i} \frac{\log n}{n^{d-1}}$ , where  $\widetilde{R}_{\mathbf{m}} = \prod_{s \in [k]} \binom{|\mathcal{V}_s^{S^c}|}{m_s}$ . Due to (64) and (65), one can show that

$$\left(1 - \frac{1}{(\log n)^3} - 2n^{-\frac{1}{2} + \frac{\delta}{2}}\right) \leq \frac{\widetilde{R}_{\mathbf{m}}}{R_{\mathbf{m}}} \leq \left(1 - \frac{1}{(\log n)^3} + 2n^{-\frac{1}{2} + \frac{\delta}{2}}\right). \quad (68)$$

Following [51, Exercise 2.2], one can show that the probability of  $N_v^{\mathbf{m}} = \tau_{\mathbf{m}}$  is

$$\mathbb{P}\left(\sum_{r=1}^{\tilde{R}_{\mathbf{m}}} B_r^{\mathbf{m}} = \tau_{\mathbf{m}}\right) = \exp\left\{-\tilde{R}_{\mathbf{m}}\mathbb{D}\left(\frac{R_{\mathbf{m}} Q_{\mathbf{m}\oplus i}^{t^*} Q_{\mathbf{m}\oplus j}^{1-t^*} \log n}{\tilde{R}_{\mathbf{m}} n^{d-1}} \left\| \frac{Q_{\mathbf{m}\oplus i} \log n}{n^{d-1}}\right.\right) - \frac{1}{2} \log(2\pi\tilde{R}_{\mathbf{m}}) - \frac{1}{2} \log\left(1 - \frac{R_{\mathbf{m}} Q_{\mathbf{m}\oplus i}^{t^*} Q_{\mathbf{m}\oplus j}^{1-t^*} \log n}{\tilde{R}_{\mathbf{m}} n^{d-1}}\right) - \frac{1}{2} \log\left(\frac{R_{\mathbf{m}} Q_{\mathbf{m}\oplus i}^{t^*} Q_{\mathbf{m}\oplus j}^{1-t^*} \log n}{\tilde{R}_{\mathbf{m}} n^{d-1}}\right) - C_2\right\}, \quad (69)$$

where  $C_2 > 0$  is a constant. By using a Taylor series expansion and the fact that  $\tilde{R}_{\mathbf{m}}/R_{\mathbf{m}}$  is bounded (as shown in (68)), we have

$$\begin{aligned} & \tilde{R}_{\mathbf{m}}\mathbb{D}\left(\frac{R_{\mathbf{m}} Q_{\mathbf{m}\oplus i}^{t^*} Q_{\mathbf{m}\oplus j}^{1-t^*} \log n}{\tilde{R}_{\mathbf{m}} n^{d-1}} \left\| \frac{Q_{\mathbf{m}\oplus i} \log n}{n^{d-1}}\right.\right) \\ &= R_{\mathbf{m}} \frac{Q_{\mathbf{m}\oplus i}^{t^*} Q_{\mathbf{m}\oplus j}^{1-t^*} \log n}{n^{d-1}} \log\left(\frac{R_{\mathbf{m}}}{\tilde{R}_{\mathbf{m}}}\left(\frac{Q_{\mathbf{m}\oplus j}}{Q_{\mathbf{m}\oplus i}}\right)^{1-t^*}\right) + \tilde{R}_{\mathbf{m}}\left(1 - \frac{R_{\mathbf{m}} Q_{\mathbf{m}\oplus i}^{t^*} Q_{\mathbf{m}\oplus j}^{1-t^*} \log n}{\tilde{R}_{\mathbf{m}} n^{d-1}}\right) \\ & \quad \times \log\left(1 + \frac{Q_{\mathbf{m}\oplus i} \log n - \frac{R_{\mathbf{m}} Q_{\mathbf{m}\oplus i}^{t^*} Q_{\mathbf{m}\oplus j}^{1-t^*} \log n}{\tilde{R}_{\mathbf{m}}}}{n^{d-1} - Q_{\mathbf{m}\oplus i} \log n}\right) \end{aligned} \quad (70)$$

$$= R'_{\mathbf{m}} Q_{\mathbf{m}\oplus i}^{t^*} Q_{\mathbf{m}\oplus j}^{1-t^*} (\log n) (1-t^*) \log\left(\frac{Q_{\mathbf{m}\oplus j}}{Q_{\mathbf{m}\oplus i}}\right) + \frac{\tilde{R}_{\mathbf{m}}}{R_{\mathbf{m}}}\left(R'_{\mathbf{m}} Q_{\mathbf{m}\oplus i} \log n - \frac{R_{\mathbf{m}}}{\tilde{R}_{\mathbf{m}}} R'_{\mathbf{m}} Q_{\mathbf{m}\oplus i}^{t^*} Q_{\mathbf{m}\oplus j}^{1-t^*} \log n + \mathcal{O}(1/(\log n)^2)\right) \quad (71)$$

$$= (\log n) \left[ \mu_{\mathbf{m}\oplus i}^{t^*} \mu_{\mathbf{m}\oplus j}^{1-t^*} (1-t^*) \log\left(\frac{\mu_{\mathbf{m}\oplus j}}{\mu_{\mathbf{m}\oplus i}}\right) + \mu_{\mathbf{m}\oplus i} - \mu_{\mathbf{m}\oplus i}^{t^*} \mu_{\mathbf{m}\oplus j}^{1-t^*} + \mathcal{O}(1/(\log n)^3) \right]. \quad (72)$$

Since  $\tilde{R}_{\mathbf{m}} = \Theta(n^{d-1})$ , we also have

$$\frac{1}{2} \log(2\pi\tilde{R}_{\mathbf{m}}) + \frac{1}{2} \log\left(1 - \frac{R_{\mathbf{m}} Q_{\mathbf{m}\oplus i}^{t^*} Q_{\mathbf{m}\oplus j}^{1-t^*} \log n}{\tilde{R}_{\mathbf{m}} n^{d-1}}\right) + \frac{1}{2} \log\left(\frac{R_{\mathbf{m}} Q_{\mathbf{m}\oplus i}^{t^*} Q_{\mathbf{m}\oplus j}^{1-t^*} \log n}{\tilde{R}_{\mathbf{m}} n^{d-1}}\right) \quad (73)$$

$$= \frac{d-1}{2} \log n - \frac{d-1}{2} \log n + \mathcal{O}(\log \log n) = \mathcal{O}(\log \log n). \quad (74)$$

Combining (69), (72), and (74), we then have

$$\mathbb{P}(N_v^{\mathbf{m}} = \tau_{\mathbf{m}}) = n^{-\left[(1-t^*)\mu_{\mathbf{m}\oplus i}^{t^*} \mu_{\mathbf{m}\oplus j}^{1-t^*} \log(\mu_{\mathbf{m}\oplus j}/\mu_{\mathbf{m}\oplus i}) + \mu_{\mathbf{m}\oplus i} - \mu_{\mathbf{m}\oplus i}^{t^*} \mu_{\mathbf{m}\oplus j}^{1-t^*} + o(1)\right]}. \quad (75)$$

Taking all  $\mathbf{m} \in \mathcal{M}$  into account, we obtain the probability that a node  $v \in \mathcal{V}_i^S$  is ambiguous as follows:

$$\mathbb{P}(v \in \mathcal{V}_i^S \text{ is ambiguous}) = \mathbb{P}(\forall \mathbf{m} \in \mathcal{M} : N_v^{\mathbf{m}} = \tau_{\mathbf{m}}) \quad (76)$$

$$= n^{-\left[\sum_{\mathbf{m} \in \mathcal{M}} (1-t^*)\mu_{\mathbf{m}\oplus i}^{t^*} \mu_{\mathbf{m}\oplus j}^{1-t^*} \log(\mu_{\mathbf{m}\oplus j}/\mu_{\mathbf{m}\oplus i}) + \mu_{\mathbf{m}\oplus i} - \mu_{\mathbf{m}\oplus i}^{t^*} \mu_{\mathbf{m}\oplus j}^{1-t^*} + o(1)\right]} \quad (77)$$

$$= n^{-[D_+(i,j)+o(1)]}, \quad (78)$$

where (77) is due to the independence of  $\{N_v^{\mathbf{m}}\}_{\mathbf{m} \in \mathcal{M}}$ , and (78) holds since  $t^*$  satisfies (67). Similarly, one can also show that for a node  $v \in \mathcal{V}_j^S$ ,

$$\mathbb{P}(v \in \mathcal{V}_j^S \text{ is ambiguous}) = \mathbb{P}(\forall \mathbf{m} \in \mathcal{M} : N_v^{\mathbf{m}} = \tau_{\mathbf{m}}) \quad (79)$$

$$= n^{-\left[\sum_{\mathbf{m} \in \mathcal{M}} t^* \mu_{\mathbf{m}\oplus i}^{t^*} \mu_{\mathbf{m}\oplus j}^{1-t^*} \log(\mu_{\mathbf{m}\oplus i}/\mu_{\mathbf{m}\oplus j}) + \mu_{\mathbf{m}\oplus j} - \mu_{\mathbf{m}\oplus i}^{t^*} \mu_{\mathbf{m}\oplus j}^{1-t^*} + o(1)\right]} \quad (80)$$

$$= n^{-[D_+(i,j)+o(1)]}. \quad (81)$$

By noting that  $D_+(i,j) < 1$  and the cardinalities of both  $\mathcal{V}_i^S$  and  $\mathcal{V}_j^S$  scale as  $\Theta(n/(\log n)^3)$ , one can show that with probability  $1 - o(1)$ , there is at least one ambiguous node in  $\mathcal{V}_i^S$  (denoted by  $v_1$ ) and also one ambiguous node in  $\mathcal{V}_j^S$  (denoted by  $v_2$ ).

In addition, we prove that with high probability, node  $v_1$  (resp.  $v_2$ ) is not connected to any node in  $S$ . This is because the number of hyperedges that contains  $v_1$  (resp.  $v_2$ ) and another node in  $S$  is at most  $|S|\binom{n}{d-2} \leq n^{d-1}/(\log n)^3$ , and the probability of each hyperedge is at most  $Q_{\max}(\log n)/n^{d-1}$ , thus the probability that  $v_1$  (resp.  $v_2$ ) does not have any connection with other nodes in  $S$  is at least

$$\left(1 - Q_{\max} \frac{\log n}{n^{d-1}}\right)^{\frac{n^{d-1}}{(\log n)^3}} \geq e^{-\frac{2Q_{\max}}{(\log n)^2}} \geq 1 - \frac{4Q_{\max}}{(\log n)^2}, \quad (82)$$

for sufficiently large  $n$ . Finally, note that both  $v_1$  and  $v_2$  are not connected to any node in  $S$ , and both of them are ambiguous (i.e., have the same number of hyperedges  $\{N_v^{\mathbf{m}}\}_{\mathbf{m} \in \mathcal{M}}$  outside  $S$ ), thus it is impossible to distinguish them and to achieve exact recovery.

## VII. CONCLUSION AND FUTURE DIRECTIONS

This paper establishes a sharp phase transition for exact recovery in the general  $d$ -HSBM, apart from a small subset of generative distributions such that there exists two communities with the same second-order degree profiles. We also develop a polynomial-time algorithm (with theoretical guarantees) that achieves the information-theoretic limit, showing that there is no information-computation gap. Our two-stage algorithm is based on hypergraph spectral clustering and local refinement steps. We put forth several promising directions for future work.

- 1) Our algorithm fails if the parameters belong to  $\Xi$  because we apply the hypergraph spectral clustering method to the processed hypergraph Laplacian  $\mathbf{L}$  (rather than the observed adjacency tensor  $\mathbf{A}$ ). This pre-processing step from  $\mathbf{A}$  to  $\mathbf{L}$  annihilates some salient information for distinguishing two communities with the same second-order degree profile. We conjecture that this issue may be circumvented if one directly applies clustering algorithms to the adjacency tensor  $\mathbf{A}$  (such as the method proposed in [37]). However, theoretical guarantees of such algorithms in the general  $d$ -HSBM may be non-trivial to develop. Nonetheless, in future work, one would expect to validate our conjecture and show that the exact recovery threshold  $\min_{i,j \in [k]: i \neq j} D_+(i,j) = 1$  holds even without the restriction that the second-order degree profiles of two distinct communities are different.
- 2) It would also be interesting to extend our theory to even more general settings and other variants of the HSBM, such as the non-uniform HSBM (as proposed in [27]–[29]), HSBM with overlapping communities, weighted or labelled HSBMs, HSBM with side information, etc.

## APPENDIX A

### PROOF OF LEMMA 3

Without loss of generality, we assume nodes  $u$  and  $v$  respectively belong to communities  $\mathcal{V}_i$  and  $\mathcal{V}_j$ . Since we require  $(\mathbf{p}, \{Q_{\mathbf{T}}\}_{\mathbf{T} \in \mathcal{T}}) \notin \Xi$ , there must exist a  $s \in [k]$  such that

$$\sum_{\mathbf{m} \in \mathcal{M}: m_s \geq 1} m_s \mu_{\mathbf{m} \oplus i} \neq \sum_{\mathbf{m} \in \mathcal{M}: m_s \geq 1} m_s \mu_{\mathbf{m} \oplus j}. \quad (83)$$

Let  $R_{\mathbf{m},s} \triangleq \binom{np_s}{m_s-1} \cdot \prod_{a \in [k] \setminus \{s\}} \binom{np_a}{m_a}$ . Thus, we have

$$\|\mathbf{M}_u - \mathbf{M}_v\|_2^2 \geq \sum_{w \in \mathcal{V}_s} (M_{w,u} - M_{w,v})^2 \quad (84)$$

$$= \sum_{w \in \mathcal{V}_s} \left( \sum_{\mathbf{m} \in \mathcal{M}: m_s \geq 1} R_{\mathbf{m},s} Q_{\mathbf{m} \oplus i} \frac{\gamma_n}{n^{d-1}} - \sum_{\mathbf{m} \in \mathcal{M}: m_s \geq 1} R_{\mathbf{m},s} Q_{\mathbf{m} \oplus j} \frac{\gamma_n}{n^{d-1}} \right)^2 \quad (85)$$

$$= \sum_{w \in \mathcal{V}_s} \left( \frac{\gamma_n}{np_s} \left[ \sum_{\mathbf{m} \in \mathcal{M}: m_s \geq 1} m_s \mu_{\mathbf{m} \oplus i} - m_s \mu_{\mathbf{m} \oplus j} \right] \right)^2 \quad (86)$$

$$= \Omega(\gamma_n^2/n), \quad (87)$$

where (86) holds since  $R_{\mathbf{m},s} = m_s R_{\mathbf{m}} / (np_s)$  and  $R_{\mathbf{m}} Q_{\mathbf{m} \oplus i} / n^{d-1} = \mu_{\mathbf{m} \oplus i}$ , and (87) follows from (83) and  $|\mathcal{V}_s| \approx np_s$ .

## APPENDIX B

### PROOF OF LEMMA 6

For any realization  $F_1 = f_1$ , let  $P_{\text{suc}}(f_1, \mathbf{z})$  be the probability that running a hypergraph spectral clustering method on  $G_1$  (which depends on  $f_1$ ) ensures  $l(\mathbf{z}, \widehat{\mathbf{Z}}^{(0)}) \leq \eta_n$ . From Theorem 3, we have

$$\sum_{f_1} \mathbb{P}(F_1 = f_1) P_{\text{suc}}(f_1, \mathbf{z}) \geq 1 - \epsilon_n. \quad (88)$$

We now prove Lemma 6 by contradiction. Suppose the probability that  $F_1 \in \mathcal{G}_1^{\mathbf{z}}$  is less than  $1 - \sqrt{\epsilon_n}$ , then we have

$$\sum_{f_1} \mathbb{P}(F_1 = f_1) P_{\text{suc}}(f_1, \mathbf{z}) < \sum_{f_1 \in \mathcal{G}_1^{\mathbf{z}}} \mathbb{P}(F_1 = f_1) + \sum_{f_1 \notin \mathcal{G}_1^{\mathbf{z}}} \mathbb{P}(F_1 = f_1) (1 - \sqrt{\epsilon_n}) \quad (89)$$

$$= \sum_{f_1 \in \mathcal{G}_1^{\mathbf{z}}} \mathbb{P}(F_1 = f_1) + (1 - \sqrt{\epsilon_n}) \left( 1 - \sum_{f_1 \notin \mathcal{G}_1^{\mathbf{z}}} \mathbb{P}(F_1 = f_1) \right) \quad (90)$$

$$< 1 - \sqrt{\epsilon_n} + (1 - \sqrt{\epsilon_n}) \cdot \sqrt{\epsilon_n} \quad (91)$$

$$= 1 - \epsilon_n, \quad (92)$$

where (89) follows from the fact that  $P_{\text{suc}}(f_1, \mathbf{z}) < 1 - \sqrt{\epsilon_n}$  for  $f_1 \notin \mathcal{G}_1^{\mathbf{z}}$  (see Definition 4), and (91) is due to our assumption. Since Eqns. (89)-(92) contradict with the fact in (88), we obtain that  $\mathbb{P}(F_1 \in \mathcal{G}_1^{\mathbf{z}}) \geq 1 - \sqrt{\epsilon_n}$ .

Let  $N_{\mathbf{m}} \triangleq \prod_{s=1}^k \binom{|\mathcal{V}_s|}{m_s}$  for each  $\mathbf{m} \in \mathcal{M}$ . For each node  $v \in [n]$ , the expected number of hyperedges in  $F_2$  that contain node  $v$  and other  $d-1$  nodes with community assignment  $\mathbf{m}$  is  $\mathbb{E}(D_{v,\mathbf{m}}) = N_{\mathbf{m}} \left( 1 - \frac{\gamma_n}{\log n} \right)$ . By applying the Chernoff bound, we have

$$\mathbb{P} \left( D_{v,\mathbf{m}} \leq N_{\mathbf{m}} \left( 1 - \frac{2\gamma_n}{\log n} \right) \right) \leq \mathbb{P} \left( D_{v,\mathbf{m}} \leq \left( 1 - \frac{\gamma_n}{\log n} \right) \mathbb{E}(D_{v,\mathbf{m}}) \right) \leq \exp \left( -\frac{1}{3} \frac{\gamma_n^2}{(\log n)^2} \mathbb{E}(D_{v,\mathbf{m}}) \right) \quad (93)$$

$$= \exp \left( -\Theta(n^{d-1} \gamma_n^2 / (\log n)^2) \right). \quad (94)$$

Taking a union bound over all  $\mathbf{m} \in \mathcal{M}$  and all the  $n$  nodes, we have that with probability at least  $1 - \exp \left( -\Theta(n^{d-1} \gamma_n^2 / (\log n)^2) \right)$ , every node  $v \in [n]$  satisfies

$$(1 - (2\gamma_n / \log n)) N_{\mathbf{m}} \leq D_{v,\mathbf{m}} \leq N_{\mathbf{m}}, \quad \forall \mathbf{m} \in \mathcal{M}. \quad (95)$$

Combining the fact that  $(1 - n^{-\frac{1}{2} + \frac{\delta}{2}}) R_{\mathbf{m}} \leq N_{\mathbf{m}} \leq (1 + n^{-\frac{1}{2} + \frac{\delta}{2}}) R_{\mathbf{m}}$  (since  $\mathbf{z} \in \mathcal{A}_Z$ ), we have

$$(1 - (3\gamma_n / \log n)) R_{\mathbf{m}} \leq D_{v,\mathbf{m}} \leq (1 + n^{-\frac{1}{2} + \frac{\delta}{2}}) R_{\mathbf{m}}, \quad \forall \mathbf{m} \in \mathcal{M}. \quad (96)$$

Thus,  $\mathbb{P}(F_2 \in \mathcal{G}_2^{\mathbf{z}}) \geq 1 - \exp \left( -\Theta(n^{d-1} \gamma_n^2 / (\log n)^2) \right)$ . This completes the proof.

APPENDIX C  
PROOF OF LEMMA 7

Note that  $\mathbb{E}(\sum_{a=1}^{D'_v} Y_a) \leq D'_v \cdot Q_{\max}(\log n)/n^{d-1} = \mathcal{O}(\eta_n \log n)$ , since the success probability of each Bernoulli random variable is at most  $Q_{\max}(\log n)/n^{d-1}$ . In the following, we show that the probability that  $\sum_{a=1}^{D'_v} Y_a \geq c_n \log n$  is at most  $n^{-\omega(1)}$ , where  $c_n \log n = \omega(\eta_n \log n)$ . Note that

$$\mathbb{P}\left(\sum_{a=1}^{D'_v} Y_a \geq c_n \log n\right) = \sum_{\theta=c_n \log n}^{D'_v} \mathbb{P}\left(\sum_{a=1}^{D'_v} Y_a = \theta\right) \quad (97)$$

$$= \sum_{\theta=c_n \log n}^{D'_v} \binom{D'_v}{\theta} \left(\frac{Q_{\max} \log n}{n^{d-1}}\right)^\theta \left(1 - \frac{Q_{\max} \log n}{n^{d-1}}\right)^{D'_v - \theta} \quad (98)$$

$$\leq D'_v \cdot \left(\frac{e D'_v}{c_n \log n}\right)^{c_n \log n} \left(\frac{Q_{\max} \log n}{n^{d-1}}\right)^{c_n \log n} \quad (99)$$

$$= D'_v \cdot n^{-c_n \log(c_n/\eta_n)} = n^{-\omega(1)}, \quad (100)$$

where (99) follows from the facts that  $\binom{n}{k} \leq (en/k)^k$  and  $\theta = c_n \log n$  maximizes the terms in (98).

We then prove the second part. For random variables  $\{Y_a\}_{a=1}^{D'_v}$ , we have

$$\frac{Q_{\min}}{Q_{\max}} \leq \frac{\mathbb{P}(Y_a = 1 | Z_v = j, \hat{\mathbf{z}}_{\sim v}^{(0)})}{\mathbb{P}(Y_a = 1 | Z_v = j, \mathbf{z}_{\sim v})} \leq \frac{Q_{\max}}{Q_{\min}}, \quad \text{and} \quad (101)$$

$$1 - \frac{(Q_{\max} - Q_{\min}) \log n}{n^{d-1}} \leq \frac{\mathbb{P}(Y_a = 0 | Z_v = j, \hat{\mathbf{z}}_{\sim v}^{(0)})}{\mathbb{P}(Y_a = 0 | Z_v = j, \mathbf{z}_{\sim v})} \leq 1 + \frac{(Q_{\max} - Q_{\min}) \log n}{n^{d-1}}. \quad (102)$$

Since  $\sum_{a=1}^{D'_v} y_a \leq c_n \log n$  for  $g_2 \in \mathcal{G}_v$ , we have

$$\frac{\mathbb{P}(g_2 | Z_v = j, \hat{\mathbf{z}}_{\sim v}^{(0)})}{\mathbb{P}(g_2 | Z_v = j, \mathbf{z}_{\sim v})} = \frac{\prod_{a=1}^{D'_v} \mathbb{P}(y_a | Z_v = j, \hat{\mathbf{z}}_{\sim v}^{(0)})}{\prod_{a=1}^{D'_v} \mathbb{P}(y_a | Z_v = j, \mathbf{z}_{\sim v})} \leq \left(\frac{Q_{\max}}{Q_{\min}}\right)^{c_n \log n} \left(1 + \frac{(Q_{\max} - Q_{\min}) \log n}{n^{d-1}}\right)^{D'_v - c_n \log n} \triangleq L_h, \quad (103)$$

$$\frac{\mathbb{P}(g_2 | Z_v = j, \hat{\mathbf{z}}_{\sim v}^{(0)})}{\mathbb{P}(g_2 | Z_v = j, \mathbf{z}_{\sim v})} = \frac{\prod_{a=1}^{D'_v} \mathbb{P}(y_a | Z_v = j, \hat{\mathbf{z}}_{\sim v}^{(0)})}{\prod_{a=1}^{D'_v} \mathbb{P}(y_a | Z_v = j, \mathbf{z}_{\sim v})} \geq \left(\frac{Q_{\min}}{Q_{\max}}\right)^{c_n \log n} \left(1 - \frac{(Q_{\max} - Q_{\min}) \log n}{n^{d-1}}\right)^{D'_v - c_n \log n} \triangleq L_l, \quad (104)$$

and note that  $L_h = n^{o(1)}$  and  $L_l = n^{-o(1)}$ .

REFERENCES

- [1] P. W. Holland, K. B. Laskey, and S. Leinhardt, "Stochastic blockmodels: First steps," *Social networks*, vol. 5, no. 2, pp. 109–137, 1983.
- [2] E. Abbe, A. S. Bandeira, and G. Hall, "Exact recovery in the stochastic block model," *IEEE Transactions on Information Theory*, vol. 62, no. 1, pp. 471–487, 2015.
- [3] E. Mossel, J. Neeman, and A. Sly, "Consistency thresholds for the planted bisection model," in *Proceedings of the forty-seventh annual ACM symposium on Theory of computing*, 2015, pp. 69–75.
- [4] E. Abbe and C. Sandon, "Community detection in general stochastic block models: Fundamental limits and efficient algorithms for recovery," in *IEEE 56th Annual Symposium on Foundations of Computer Science*, 2015, pp. 670–688.
- [5] —, "Recovering communities in the general stochastic block model without knowing the parameters," *arXiv preprint arXiv:1506.03729*, 2015.
- [6] B. Hajek, Y. Wu, and J. Xu, "Information limits for recovering a hidden community," *IEEE Transactions on Information Theory*, vol. 63, no. 8, pp. 4729–4745, 2017.
- [7] G. Reeves, V. Mayya, and A. Volfovsky, "The geometry of community detection via the mmse matrix," in *2019 IEEE International Symposium on Information Theory (ISIT)*, 2019, pp. 400–404.
- [8] V. Jog and P.-L. Loh, "Information-theoretic bounds for exact recovery in weighted stochastic block models using the Renyi divergence," *arXiv preprint arXiv:1509.06418*, 2015.
- [9] S.-Y. Yun and A. Proutiere, "Optimal cluster recovery in the labeled stochastic block model," *Advances in Neural Information Processing Systems*, vol. 29, pp. 965–973, 2016.
- [10] —, "Accurate community detection in the stochastic block model via spectral algorithms," *arXiv preprint arXiv:1412.7335*, 2014.

- [11] P. Chin, A. Rao, and V. Vu, “Stochastic block model and community detection in sparse graphs: A spectral algorithm with optimal rate of recovery,” in *Conference on Learning Theory*, 2015, pp. 391–423.
- [12] B. Hajek, Y. Wu, and J. Xu, “Achieving exact cluster recovery threshold via semidefinite programming,” *IEEE Transactions on Information Theory*, vol. 62, no. 5, pp. 2788–2797, 2016.
- [13] —, “Achieving exact cluster recovery threshold via semidefinite programming: Extensions,” *IEEE Transactions on Information Theory*, vol. 62, no. 10, pp. 5918–5937, 2016.
- [14] A. Montanari and S. Sen, “Semidefinite programs on sparse random graphs and their application to community detection,” in *Proceedings of the forty-eighth annual ACM symposium on Theory of Computing*, 2016, pp. 814–827.
- [15] F. Caltagirone, M. Lelarge, and L. Miolane, “Recovering asymmetric communities in the stochastic block model,” *IEEE Transactions on Network Science and Engineering*, vol. 5, no. 3, pp. 237–246, 2017.
- [16] N. Agarwal, A. S. Bandeira, K. Koiliaris, and A. Kolla, “Multisection in the stochastic block model using semidefinite programming,” in *Compressed Sensing and its Applications*, 2017, pp. 125–162.
- [17] A. Perry and A. S. Wein, “A semidefinite program for unbalanced multisection in the stochastic block model,” in *2017 International Conference on Sampling Theory and Applications (SampTA)*, 2017, pp. 64–67.
- [18] A. R. Asadi, E. Abbe, and S. Verdú, “Compressing data on graphs with clusters,” in *IEEE Int. Symp. Inf. Theory (ISIT)*, 2017, pp. 1583–1587.
- [19] H. Saad and A. Nosratinia, “Community detection with side information: Exact recovery under the stochastic block model,” *IEEE Journal of Selected Topics in Signal Processing*, vol. 12, no. 5, pp. 944–958, 2018.
- [20] —, “Recovering a single community with side information,” *IEEE Transactions on Information Theory*, vol. 66, no. 12, pp. 7939–7966, 2020.
- [21] —, “Exact recovery in community detection with continuous-valued side information,” *IEEE Signal Processing Letters*, vol. 26, no. 2, pp. 332–336, 2018.
- [22] V. Mayya and G. Reeves, “Mutual information in community detection with covariate information and correlated networks,” in *57th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, 2019, pp. 602–607.
- [23] E. Abbe, “Community detection and stochastic block models: recent developments,” *The Journal of Machine Learning Research*, vol. 18, no. 1, pp. 6446–6531, 2017.
- [24] M. Chertok and Y. Keller, “Efficient high order matching,” *IEEE Transactions on Pattern Analysis and Machine Intelligence*, vol. 32, no. 12, pp. 2205–2215, 2010.
- [25] H. Liu, L. Latecki, and S. Yan, “Robust clustering as ensembles of affinity relations,” *Advances in neural information processing systems*, vol. 23, pp. 1414–1422, 2010.
- [26] D. Ghoshdastidar and A. Dukkipati, “Consistency of spectral partitioning of uniform hypergraphs under planted partition model,” *Advances in Neural Information Processing Systems*, vol. 27, pp. 397–405, 2014.
- [27] —, “A provable generalized tensor spectral method for uniform hypergraph partitioning,” in *International Conference on Machine Learning*, 2015, pp. 400–409.
- [28] —, “Spectral clustering using multilinear svd: Analysis, approximations and applications,” in *Proceedings of the AAAI Conference on Artificial Intelligence*, vol. 29, no. 1, 2015.
- [29] —, “Consistency of spectral hypergraph partitioning under planted partition model,” *Annals of Statistics*, vol. 45, no. 1, pp. 289–315, 2017.
- [30] S. Pal and Y. Zhu, “Community detection in the sparse hypergraph stochastic block model,” *arXiv preprint arXiv:1904.05981*, 2019.
- [31] S. Cole and Y. Zhu, “Exact recovery in the hypergraph stochastic block model: A spectral algorithm,” *Linear Algebra and its Applications*, vol. 593, pp. 45–73, 2020.
- [32] K. Ahn, K. Lee, and C. Suh, “Information-theoretic limits of subspace clustering,” in *2017 IEEE International Symposium on Information Theory (ISIT)*, 2017, pp. 2473–2477.
- [33] —, “Hypergraph spectral clustering in the weighted stochastic block model,” *IEEE Journal of Selected Topics in Signal Processing*, vol. 12, no. 5, pp. 959–974, 2018.
- [34] C. Kim, A. S. Bandeira, and M. X. Goemans, “Stochastic block model for hypergraphs: Statistical limits and a semidefinite programming approach,” *arXiv preprint arXiv:1807.02884*, 2018.
- [35] J. Lee, D. Kim, and H. W. Chung, “Robust hypergraph clustering via convex relaxation of truncated mle,” *IEEE Journal on Selected Areas in Information Theory*, 2020.
- [36] C. Kim, A. S. Bandeira, and M. X. Goemans, “Community detection in hypergraphs, spiked tensor models, and sum-of-squares,” in *2017 International Conference on Sampling Theory and Applications (SampTA)*, 2017, pp. 124–128.
- [37] Z. T. Ke, F. Shi, and D. Xia, “Community detection for hypergraph networks via regularized tensor power iteration,” *arXiv preprint arXiv:1909.06503*, 2019.

- [38] M. C. Angelini, F. Caltagirone, F. Krzakala, and L. Zdeborová, “Spectral detection on sparse hypergraphs,” in *2015 53rd Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, 2015, pp. 66–73.
- [39] T. Lesieur, L. Miolane, M. Lelarge, F. Krzakala, and L. Zdeborová, “Statistical and computational phase transitions in spiked tensor estimation,” in *2017 IEEE International Symposium on Information Theory (ISIT)*, 2017, pp. 511–515.
- [40] I. E. Chien, C.-Y. Lin, and I.-H. Wang, “On the minimax misclassification ratio of hypergraph community detection,” *IEEE Transactions on Information Theory*, vol. 65, no. 12, pp. 8095–8118, 2019.
- [41] C.-Y. Lin, I. E. Chien, and I.-H. Wang, “On the fundamental statistical limit of community detection in random hypergraphs,” in *2017 IEEE International Symposium on Information Theory (ISIT)*, 2017, pp. 2178–2182.
- [42] I. Chien, C.-Y. Lin, and I.-H. Wang, “Community detection in hypergraphs: Optimal statistical limit and efficient algorithms,” in *International Conference on Artificial Intelligence and Statistics*, 2018, pp. 871–879.
- [43] T. M. Cover and J. A. Thomas, *Elements of Information Theory*. John Wiley & Sons, 2012.
- [44] C. Gao, Z. Ma, A. Y. Zhang, and H. H. Zhou, “Achieving optimal misclassification proportion in stochastic block models,” *The Journal of Machine Learning Research*, vol. 18, no. 1, pp. 1980–2024, 2017.
- [45] P. Jain, P. Netrapalli, and S. Sanghavi, “Low-rank matrix completion using alternating minimization,” in *STOC*, 2013, pp. 665–674.
- [46] R. H. Keshavan, A. Montanari, and S. Oh, “Matrix completion from a few entries,” *IEEE Trans. Inf. Theory*, vol. 56, no. 6, pp. 2980–2998, 2010.
- [47] Q. Zhang, G. Suh, C. Suh, and V. Y. Tan, “Mc2g: An efficient algorithm for matrix completion with social and item similarity graphs,” *arXiv preprint arXiv:2006.04373*, 2020.
- [48] V. Vu, “A simple svd algorithm for finding hidden partitions,” *arXiv preprint arXiv:1404.3918*, 2014.
- [49] C. Eckart and G. Young, “The approximation of one matrix by another of lower rank,” *Psychometrika*, vol. 1, no. 3, pp. 211–218, 1936.
- [50] H. Weyl, “Das asymptotische verteilungsgesetz der eigenwerte linearer partieller differentialgleichungen (mit einer anwendung auf die theorie der hohlraumstrahlung),” *Mathematische Annalen*, vol. 71, no. 4, pp. 441–479, 1912.
- [51] I. Csiszár and J. Körner, *Information theory: coding theorems for discrete memoryless systems*. Cambridge University Press, 2011.