

Position-space representation of charged particles' propagators in a constant magnetic field as an expansion over Landau levels

Iablokov S. N.* and Kuznetsov A. V.

P.G. Demidov Yaroslavl State University, Yaroslavl, Russia

(Dated: January 8, 2022)

We obtained propagators in the position space as an expansion over Landau levels for the charged scalar particle, fermion and massive vector boson in a constant external magnetic field. The summation terms in the resulting expressions consisted of two factors, one being rotationally invariant in the 2-dimensional Euclidean space perpendicular to the direction of the field, and the other being Lorentz-invariant in the 1+1-dimensional space-time. The discussed methods, namely, the Fock-Schwinger method, the modified Fock-Schwinger method and the "sum over solutions" approach demonstrated consistent results, thus, contributing to the development of the position-space techniques in QFT.

I. INTRODUCTION

In the vast majority of QFT problems, the momentum-space paradigm is adopted due to the simplicity of calculations. However, in the case of a (partially) broken translational invariance, e.g. in the presence of external electromagnetic field, the explicit position-space expressions might be useful for the thorough analysis of the underlying problem. When studying propagators, these expressions allow to investigate the corresponding time-like, space-like and light-cone behaviour. Unfortunately, the position-space methods of QFT are far less developed than the momentum-space techniques, with a moderate number of analytic formulae known for the propagators of quantum fields.

Besides the case of a free field [1, 2], there exist another canonical QFT problem, namely a charged particle in a constant magnetic field, with known exact analytic expression for the respective propagator in the position space. This expression is obtained using the Fock-Schwinger method [3, 4] and is represented as an integral over the proper-time [5] parameter. Two other known analytic representations of charged particles' propagators are in the momentum space and are written either via the proper-time or as an expansion over Landau levels [6–13].

In our previous paper [14] we obtained a "missing" position-space Landau-levels representation of a scalar charged particle propagator in a constant magnetic field. Here, we further develop position-space techniques and compare three different methods for solving this problem, namely, the original Fock-Schwinger (FS) method, the modified Fock-Schwinger (MFS) method [15, 16] and the "sum over solutions" approach. Additionally, we obtain the expressions for the propagators of fermions and massive vector bosons in the position space as an expansion over Landau levels.

Throughout the calculations we adhere to the West-Coast metric convention $g_{\mu\nu} = (+, -, -, -)$. The constant magnetic field B is assumed to be directed along

the z -axis. This naturally leads to the decomposition of the spacetime vectors into the parallel (\parallel) and perpendicular (\perp) components

$$p_{\perp}^{\mu} = (0, p_1, p_2, 0), \quad p_{\parallel}^{\mu} = (p_0, 0, 0, p_3), \quad (1)$$

which belong to the Euclidean $\{1, 2\}$ -subspace and the Minkowski $\{0, 3\}$ -subspace, correspondingly. Then, for the arbitrary four-vectors p_{μ}, q_{μ} one has

$$(pq)_{\perp} = p_1 q_1 + p_2 q_2, \quad (pq)_{\parallel} = p_0 q_0 - p_3 q_3, \quad (2)$$

with the full scalar product written as

$$(pq) = (pq)_{\parallel} - (pq)_{\perp}. \quad (3)$$

II. CHARGED SCALAR PARTICLE PROPAGATOR

A. Proper-time representation

In this section we outline the derivation of the proper-time position-space representation of a charged scalar particle propagator in a constant magnetic field using the original Fock-Schwinger (FS) approach.

Briefly, we are to solve the following propagator equation:

$$H(\partial_X, X) G(X, X') = \delta^{(4)}(X - X'). \quad (4)$$

The FS method consists in representing the unknown function $G(X, X')$ as an integral

$$G(X, X') = (-i) \int_{-\infty}^0 d\tau U(X, X'; \tau), \quad (5)$$

where $U(X, X'; \tau)$ satisfies a Schrödinger-type equation:

$$i \partial_{\tau} U(X, X'; \tau) = H(\partial_X, X) U(X, X'; \tau) \quad (6)$$

subject to the following boundary conditions:

$$U(X, X'; 0) = \delta^{(4)}(X - X'), \quad U(X, X'; -\infty) = 0. \quad (7)$$

* physics@iablockov.ru

The solution then reads:

$$U(X, X'; \tau) = e^{-i\tau H(\partial_X, X) + \varepsilon\tau \delta^{(4)}(X - X')}. \quad (8)$$

The action of the exponential operator could be evaluated directly (as will be discussed later), however, the original FS method follows a different strategy. First, to ease notation, let us rewrite (8) as follows:

$$U(X, X'; \tau) = \langle X | e^{-i\tau H} | X' \rangle, \quad (9)$$

where we also suppressed the $i\varepsilon$, which will be restored in the end of calculations. Next, we once again consider the differential equation for $U(X, X'; \tau)$, this time written in new terms:

$$i\partial_\tau U(X, X'; \tau) = \langle X | H e^{-i\tau H} | X' \rangle. \quad (10)$$

Finally, if one is able to factor a scalar function out of the matrix element in (10), such that

$$\langle X | H e^{-i\tau H} | X' \rangle = F(X, X'; \tau) \langle X | e^{-i\tau H} | X' \rangle, \quad (11)$$

then substituting (9) in the right-hand side of (11) gives us a differential equation

$$i\partial_\tau U(X, X'; \tau) = F(X, X'; \tau) U(X, X'; \tau) \quad (12)$$

with the obvious solution:

$$U(X, X'; \tau) = C(X, X') \exp^{-i \int^\tau d\tau' F(X, X'; \tau')}. \quad (13)$$

Here, the result of integration on the lower bound is incorporated into $C(X, X')$, which is to be determined later.

For the scalar charged particle in a constant magnetic field (directed along the z -axis) the corresponding $H(\partial_X, X)$ operator has the following form:

$$H(\partial_X, X) = (\Pi^\mu \Pi_\mu - m^2), \quad (14)$$

where $\Pi^\mu = i\partial^\mu - eQA^\mu(X)$. Here, Q is a dimensionless charge of the particle and $e > 0$ is the elementary charge.

In complete analogy with the proof of Erenfest's theorem, Heisenberg-picture operators are introduced:

$$\Pi^\mu(\tau) = e^{i\tau H} \Pi^\mu e^{-i\tau H}, \quad (15)$$

$$X^\mu(\tau) = e^{i\tau H} X^\mu e^{-i\tau H}, \quad (16)$$

$$H(\tau) = e^{i\tau H} H e^{-i\tau H} = H. \quad (17)$$

They satisfy the following equations

$$\frac{d\Pi^\mu(\tau)}{d\tau} = i[H(\tau), \Pi^\mu(\tau)] = -2eQF_{\nu}^{\mu}\Pi^\nu(\tau), \quad (18)$$

$$\frac{dX^\mu(\tau)}{d\tau} = i[H(\tau), X^\mu(\tau)] = -2\Pi^\mu(\tau). \quad (19)$$

Solving for $\Pi(\tau)$ in terms of $X(\tau)$, one obtains:

$$\begin{aligned} \Pi^\mu(\tau)\Pi_\mu(\tau) &= \frac{1}{4\tau^2} [X^\mu(\tau) - X^\mu(0)]_\parallel^2 \\ &\quad - \frac{\beta^2}{4\sin^2(\beta\tau)} [X^\mu(\tau) - X^\mu(0)]_\perp^2. \end{aligned} \quad (20)$$

With the aid of the commutation relation

$$[X^\mu(\tau), X_\mu(0)] = -4i\tau - 2i\beta^{-1} \sin(2\beta\tau) \quad (21)$$

we are able to factorize a scalar function out of the matrix element in Eq. (10):

$$F(X, X', \tau) = \frac{(X - X')_\parallel^2}{4\tau^2} - \frac{i}{\tau} - m^2 - \frac{\beta^2(X - X')_\perp^2}{4\sin^2(\beta\tau)} - \frac{i\beta}{\tan(\beta\tau)}. \quad (22)$$

After evaluating the integral in (13), we obtain the following expression:

$$U(X, X'; \tau) = \frac{C(X, X')}{\tau \sin(\beta\tau)} e^{i \left[\frac{(X - X')_\parallel^2}{4\tau} - \frac{\beta(X - X')_\perp^2}{4 \tan(\beta\tau)} + m^2\tau \right]}. \quad (23)$$

To determine $C(X, X')$ we consider the equality

$$[i\partial_X^\mu - eQA^\mu(X)] \langle X | e^{-i\tau H} | X' \rangle = \langle X | e^{-i\tau H} \Pi^\mu(\tau) | X' \rangle,$$

which gives us the differential equation:

$$\left[i\partial_X^\mu - eQA^\mu - \frac{eQ}{2} F_{\nu}^{\mu}(X - X')_\perp^\nu \right] C(X, X') = 0. \quad (24)$$

Integrating (24) leads to:

$$C(X, X') = C(X') e^{-ieQ \int_{X'}^X d\xi^\mu [A_\mu(\xi) + \frac{1}{2} F_{\mu\nu}(\xi - X')^\nu]}. \quad (25)$$

The remaining constant $C(X')$ is determined from (4) and is equal to $-i\beta/(4\pi)^2$. To further simplify the result, we notice that the curl of the expression in brackets in (25) vanishes, making the integral independent of the integration path. Using the linear parametrization between X and X' , we eliminate the second term, which is a contraction of an antisymmetric and symmetric tensors.

Restoring the $i\varepsilon$ prescription, we provide the final expression for the proper-time position-space representation of the charged scalar particle propagator in a constant magnetic field:

$$G(X, X') = -\frac{\beta}{(4\pi)^2} e^{i\Phi(X, X')} \int_{-\infty}^0 d\tau \frac{1}{\tau \sin(\beta\tau)} \exp\left\{ i \left[\frac{Z_\parallel^2}{4\tau} - \frac{\beta Z_\perp^2}{4 \tan(\beta\tau)} + (m^2 - i\varepsilon)\tau \right] \right\}, \quad (26)$$

where $Z^\mu = X^\mu - X'^\mu$ and

$$\Phi(X, X') = -eQ \int_{X'}^X d\xi^\mu \left[A_\mu(\xi) + \frac{1}{2} F_{\mu\nu}(\xi - X')^\nu \right]. \quad (27)$$

B. Landau-levels representation

In this section we switch from the proper-time representation (26) to the sum over Landau levels.

One can notice that the integral in (26) resembles the well-known identity (see, e.g., [17]) for the modified Bessel functions of the second kind:

$$K_\nu(z) = \frac{1}{2} \left(\frac{z}{2}\right)^\nu \int_0^\infty \frac{dt}{t^{\nu+1}} e^{-t - \frac{z^2}{4t}}, \quad (28)$$

which is valid for $\text{Re } z^2 > 0$ and $|\arg z| < \pi/4$.

To use (28), we transform (26) according to the following recipe. First, a change of integration variable is performed:

$$t \equiv -i(m^2 - i\varepsilon)\tau. \quad (29)$$

This effectively rotates the integration contour by the angle of $\approx -\pi/2 - \varepsilon/m^2$ and leads to the following formula:

$$G(X, X') = \frac{i\beta}{(4\pi)^2} \int_{(i+\varepsilon)\infty}^0 \frac{dt}{t} e^{-\frac{(-Z_\parallel^2)(m^2 - i\varepsilon)}{4t} - t} \times \left([\sinh(\beta t/m^2)]^{-1} e^{-\frac{\beta Z_\perp^2}{4} \coth(\beta t/m^2)} \right). \quad (30)$$

The expression in brackets can be written as:

$$\begin{aligned} [\sinh(\beta t/m^2)]^{-1} e^{-\frac{\beta Z_\perp^2}{4} \coth(\beta t/m^2)} &= \\ &= 2 \frac{e^{-b}}{1-c} e^{a \frac{c+1}{c-1}} = 2 \frac{e^{-b} e^{-a}}{1-c} e^{-\frac{2ac}{1-c}}, \end{aligned} \quad (31)$$

where $a = \beta Z_\perp^2/4$, $b = t\beta/m^2$ and $c = e^{-2b}$.

The final expression for the position-space Landau-levels representation of the propagator reads:

$$G(X, X') = \frac{-i\beta}{4\pi^2} e^{i\Phi} \sum_{n=0}^{\infty} L_n e^{-\beta Z_\perp^2/4} \left[K_0 - \frac{i\pi}{2} H_0^{(2)} \right], \quad (36)$$

with the notations: $L_n = L_n\left(\frac{\beta Z_\perp^2}{2}\right)$, $K_0 = K_0\left(M_n \sqrt{-Z_\parallel^2 + i\varepsilon}\right) \theta(-Z_\parallel^2)$ and $H_0^{(2)} = H_0^{(2)}\left(M_n \sqrt{Z_\parallel^2 - i\varepsilon}\right) \theta(Z_\parallel^2)$.

The light-cone behaviour of K_0 and $H_0^{(2)}$ is discussed, e.g., in [18].

C. Sum over solutions

An alternative way to obtain analytic expression for the propagator is to consider a time-ordered sum over the solutions of the corresponding wave-equation:

$$H(\partial_X, X)\psi(X) = 0. \quad (37)$$

The H operator can be written as:

$$\begin{aligned} H(\partial_X, X) &= \Pi^\mu \Pi_\mu - m^2 = \\ &= (i\partial_\eta)_\parallel^2 + \beta [d_\eta^2 - \eta^2] - m^2, \end{aligned} \quad (38)$$

where $\eta = \sqrt{\beta} \left(x - Q \frac{p_y}{\beta}\right)$ and $\beta = eB$. It leads to the following set of solutions:

$$\psi_n^{(\pm)}(X) = \frac{\beta^{1/4}}{\sqrt{2p_0 L_y L_z}} e^{\mp i p_0 t + i p_y y + i p_z z} V_n(\eta), \quad (39)$$

Transition to the sum over Landau levels is performed via the following formula for the generating function of Laguerre polynomials L_n , which is valid for $|c| < 1$:

$$\frac{1}{1-c} e^{-\frac{2ac}{1-c}} = \sum_{n=0}^{\infty} L_n(2a) c^n. \quad (32)$$

We additionally change the contour of integration from $((i+\varepsilon)\infty, 0)$ to $(0, \infty)$, where the condition $c = e^{-2b} = e^{-2t\beta/m^2} < 1$ is satisfied.

After the straightforward rearrangements in the exponential factors in (30) and the following substitutions

$$\frac{t}{m^2} = \frac{s}{M_n^2}, \quad M_n^2 = m^2 + (2n+1)\beta, \quad (33)$$

we use (28) and obtain for $Z_\parallel^2 < 0$:

$$\begin{aligned} G(X, X') &= -\frac{i\beta}{4\pi^2} e^{i\Phi} e^{-\beta Z_\perp^2/4} \times \\ &\sum_{n=0}^{\infty} K_0 \left(M_n \sqrt{-Z_\parallel^2 + i\varepsilon} \right) L_n \left(\frac{\beta Z_\perp^2}{2} \right). \end{aligned} \quad (34)$$

For the case $Z_\parallel^2 > 0$, we transform K_0 using a standard relation for Bessel functions:

$$K_\nu(z) = -\frac{i\pi}{2} e^{-i\pi\nu/2} H_\nu^{(2)} \left(z e^{-i\pi/2} \right), \quad (35)$$

where $H_\nu^{(2)}$ are the Hankel functions of the second kind.

where

$$p_0 = \sqrt{p_z^2 + m^2 + (2n+1)\beta} \quad (40)$$

and L_y/L_z are the volume normalization factors. The functions V_n are the quantum harmonic oscillator (QHO) eigenvectors

$$V_n(\eta) = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} e^{-\eta^2/2} H_n(\eta), \quad (41)$$

satisfying

$$[d_\eta^2 - \eta^2] V_n(\eta) = -(2n+1)V_n(\eta), \quad (42)$$

with H_n being the Hermite polynomials.

Performing standard calculations in the canonical quantization scheme, we obtain the propagator

$$G(X, X') = (-i) \langle 0 | T \{ \psi(X) \psi^*(X') \} | 0 \rangle \quad (43)$$

as the "sum over solutions":

$$G(X, X') = \sqrt{\beta} \sum_{n=0}^{\infty} \int \frac{d^2 p_{\parallel} dp_y}{(2\pi)^3} \times \quad (44)$$

$$\times \frac{e^{-i(p(X-X')_{\parallel} + ip_y(y-y'))}}{p_{\parallel}^2 - M_n^2 + i\epsilon} V_n(\eta) V_n(\eta'),$$

where $\eta' = \sqrt{\beta} \left(x' - Q \frac{p_y}{\beta} \right)$. This form of the propagator, however, is not symmetric with respect to x, y coordinates and, therefore, does not reflect the internal symmetry of the problem. To symmetrize (44), we should perform the p_y -integration:

$$I_{n,n'} = \int dp_y e^{ip_y(y-y')} V_n(\eta) V_{n'}(\eta'). \quad (45)$$

First, we make a change of the integration variable:

$$u = -Q \frac{p_y}{\sqrt{\beta}} + \frac{\sqrt{\beta}}{2} [(x+x') + iQ(y-y')]. \quad (46)$$

This leads to:

$$I_{n,n'} = \frac{e^{i\Phi(X,X')}}{\sqrt{2^{n+n'} n! n'! \pi}} \sqrt{\beta} e^{-\frac{\beta}{4}(X-X')_{\perp}^2} \tilde{I}_{n,n'}, \quad (47)$$

where

$$\Phi(X, X') = \frac{Q\beta}{2} (x+x')(y-y'), \quad (48)$$

$$\tilde{I}_{n,n'} = \int_{-\infty}^{\infty} du e^{-u^2} H_n(u+a) H_{n'}(u+b), \quad (49)$$

with the following substitutions:

$$a = \frac{\sqrt{\beta}}{2} [(x-x') - iQ(y-y')],$$

$$b = -\frac{\sqrt{\beta}}{2} [(x-x') + iQ(y-y')]. \quad (50)$$

The phase (48) can be shown to be in agreement with the one in Eq. (27), see e.g. Ref. [12]. Secondly, according to Ref. [17], the $\tilde{I}_{n,n'}$ integral evaluates to:

$$\tilde{I}_{n,n'} = 2^{n'} \sqrt{\pi} n! b^{n'-n} L_n^{(n'-n)}(-2ab) =$$

$$= 2^{n'} \sqrt{\pi} n! b^{n'-n} L_n^{(n'-n)} \left(\frac{\beta}{2} Z_{\perp}^2 \right) \quad (51)$$

for $n \leq n'$. Finally, we obtain the symmetrized representation of the propagator:

$$G(X, X') = \frac{\beta}{2\pi} e^{i\Phi} \sum_{n=0}^{\infty} L_n \left(\frac{\beta Z_{\perp}^2}{2} \right) e^{-\beta Z_{\perp}^2 / 4}$$

$$\times \int \frac{d^2 p_{\parallel}}{(2\pi)^2} \frac{e^{-i(pZ)_{\parallel}}}{p_{\parallel}^2 - M_n^2 + i\epsilon}. \quad (52)$$

We observe that, aside from the overall non-invariant phase factor $e^{i\Phi}$, the summation terms in (73) decompose into two factors. The first one depends only on the

x, y -coordinates and is invariant with respect to rotations in the x, y -plane which is perpendicular to the direction of the magnetic field. The second factor (the Fourier integral)

$$J_{\parallel} = \int \frac{d^2 p_{\parallel}}{(2\pi)^2} \frac{e^{-i(pZ)_{\parallel}}}{p_{\parallel}^2 - M_n^2 + i\epsilon} \quad (53)$$

depends only on the t, z -coordinates and allows for further simplification. In the case $Z_{\parallel}^2 < 0$ it evaluates to:

$$J_{\parallel} = \frac{-i}{2\pi} K_0 \left(M_n \sqrt{-Z_{\parallel}^2 + i\epsilon} \right). \quad (54)$$

For $Z_{\parallel}^2 > 0$, we once again apply (35), which gives the same final answer (36) for the propagator.

D. Modified Fock-Schwinger method

Yet another approach for finding propagators of charged particles in external electromagnetic fields, modified Fock-Schwinger (MFS) method [15, 16], consists in the direct evaluation of (8). First, one should choose an appropriate representation of the δ -function. For the considered problem, the following decomposition is particularly convenient:

$$\delta^{(4)}(X - X') = \sqrt{\beta} \sum_{n=0}^{\infty} \int \frac{d^3 p_{\parallel, y}}{(2\pi)^3} e^{-i(pZ)_{\parallel, y}} V_n(\eta) V_n(\eta'). \quad (55)$$

Next, using (38) and (42), we write (5) as:

$$G(X, X') = -i\sqrt{\beta} \sum_{n=0}^{\infty} \int \frac{d^3 p_{\parallel, y}}{(2\pi)^3} e^{-i(pZ)_{\parallel, y}} \quad (56)$$

$$\int_{-\infty}^0 d\tau e^{-i\tau(p_{\parallel}^2 - M_n^2 + i\epsilon)} V_n V_n',$$

which after integration over τ gives us (44). The rest of calculations are identical to (45) – (54).

For the case of a scalar particle, MFS method gives no obvious advantage as compared to the calculations performed in the canonical quantization scheme. However, for particles with spin it significantly simplifies the notation and provides additional representations of the propagator. In the following chapters we focus our attention exclusively on the calculations using the MFS method.

III. CHARGED FERMION PROPAGATOR

The $H(\partial_X, X)$ operator in Eq. (4) for a charged fermion in a constant magnetic field reads:

$$H(\partial_X, X) = \Pi_{\mu} \gamma^{\mu} - m. \quad (57)$$

Squaring the operator by the following substitution

$$G(X, X') = (\Pi_{\mu} \gamma^{\mu} + m) S(X, X') \quad (58)$$

leads to the equation on the new unknown function $S(X, X')$:

$$H(\partial_X, X) S(X, X') = \delta^{(4)}(X - X'), \quad (59)$$

Provided that $\Sigma_3 = i\gamma_1\gamma_2 = \text{diag}(+1, -1, +1, -1)$ commutes with the unit matrix I , we are able to separate the exponential operator in (8) into two exponents:

$$S(X, X') = (-i)\sqrt{\beta} \sum_{n=0}^{\infty} \int \frac{d^2 p_{||} dp_y}{(2\pi)^3} \int_{-\infty}^0 d\tau e^{-i\tau Q\beta\Sigma_3} e^{-i\tau[p_{||}^2 - \beta(2n+1) - m^2 + i\varepsilon]} e^{-i(p(X-X'))_{||,y}} V_n(\eta) V_n(\eta'). \quad (61)$$

Next, we evaluate $\exp(-i\tau Q\beta\Sigma_3)$ and combine it with the second exponent, followed by the shift of the summation index where appropriate. This allows us to perform the proper-time integration and obtain the following expression:

$$S(X, X') = \sqrt{\beta} \sum_{n=0}^{\infty} \int \frac{d^2 p_{||} dp_y}{(2\pi)^3} \frac{e^{-i(p(X-X'))_{||,y}}}{p_{||}^2 - M_n^2 + i\varepsilon} \left[V_{n-1} V'_{n-1} \Pi_{n-1}^{(Q)} + V_n V'_n \Pi_n^{(Q)} \right], \quad (62)$$

where

$$\begin{aligned} \Pi_{n-1}^{(Q)} &= \frac{1-Q}{2} \Pi_+ + \frac{1+Q}{2} \Pi_-, & \Pi_n^{(Q)} &= \frac{1+Q}{2} \Pi_+ + \frac{1-Q}{2} \Pi_-, \\ \Pi_+ &= \text{diag}(1, 0, 1, 0), & \Pi_- &= \text{diag}(0, 1, 0, 1) \end{aligned} \quad (63)$$

and $M_n^2 = m^2 + 2\beta n$. Following the same techniques as in the case of a scalar particle and using the corollary from (24) and (25), namely,

$$\left(i\partial_\mu - eQA_\mu \right) [e^{i\Phi} f] = e^{i\Phi} \left(i\partial_\mu + \frac{eQ}{2} F_{\mu\nu} (X - X')^\nu \right) f, \quad (64)$$

we apply Eq. (58) and arrive at the following result:

$$G(X, X') = \frac{-i\beta}{4\pi^2} e^{i\Phi} \left[\left(i\partial_\mu + \frac{eQ}{2} F_{\mu\nu} Z^\nu \right) \gamma^\mu + m \right] \sum_{n=0}^{\infty} e^{-\beta Z_\perp^2 / 4} \left[L_{n-1} \Pi_{n-1}^{(Q)} + L_n \Pi_n^{(Q)} \right] \left(K_0 - \frac{i\pi}{2} H_0^{(2)} \right). \quad (65)$$

This representation of the propagator is of practical importance by itself, and can be used in calculations involving integrals of $G(X, X')$, where the derivative can be integrated by parts. It is a straightforward task to take the derivative by applying standard relations for the special functions. The resulting expression is rather cumbersome and is not provided.

IV. CHARGED MASSIVE VECTOR BOSON PROPAGATOR

Lastly, let us consider the charged vector boson. The corresponding propagator equation is given by:

$$\begin{aligned} H^\mu_\nu G^\nu_\rho(X, X') &= \delta^\mu_\rho \delta^4(X - X'), & (66) \\ H^\mu_\nu &= (H_0)^\mu_\nu + (H_F)^\mu_\nu + (H_\xi)^\mu_\nu, \end{aligned}$$

where

$$(H_0)^\mu_\nu = (\text{III} - m^2) \delta^\mu_\nu, \quad (67)$$

$$(H_F)^\mu_\nu = -2ieQF^\mu_\nu, \quad (68)$$

$$(H_\xi)^\mu_\nu = \left(\frac{1}{\xi} - 1 \right) \Pi^\mu \Pi_\nu. \quad (69)$$

We proceed with calculations exactly as in [16]. Briefly, we first note that $[H_0 + H_F, H_\xi] = 0$ and

$[H_0, H_F] = 0$. This allows for a step-by-step separation of $e^{-i\tau(H_0 + H_F + H_\xi)}$, namely:

$$\begin{aligned} e^{-i\tau(H_0 + H_F + H_\xi)} &= e^{-i\tau H_\xi} e^{-i\tau(H_0 + H_F)} = \\ &= e^{-i\tau H_\xi} e^{-i\tau H_F} e^{-i\tau H_0}. \end{aligned} \quad (70)$$

Expanding the exponents

$$[e^{-i\tau H_F}]^\mu_\nu = [e^{-2Q\beta\tau\varphi}]^\mu_\nu = \quad (71)$$

$$= \delta^\mu_\nu + \frac{e^{i2\beta\tau}}{2} (\delta^\mu_{\perp\nu} + iQ\varphi^\mu_\nu) + \frac{e^{-i2\beta\tau}}{2} (\delta^\mu_{\perp\nu} - iQ\varphi^\mu_\nu)$$

and

$$[e^{-i\tau H_\xi}]^\mu_\nu = e^{-i\tau(\frac{1}{\xi} - 1)\Pi^\mu \Pi_\nu} = \quad (72)$$

$$= \delta^\mu_\nu + \Pi^\mu \frac{e^{-i\tau(\frac{1}{\xi} - 1)\text{III}} - 1}{\text{III}} \Pi_\nu,$$

we shift the summation index (as for the fermion case) and evaluate the corresponding τ -integral. Performing

the transformations (45) – (51), we obtain the symmetrized representation of the propagator:

$$G^\mu_\nu(X, X') = \frac{\beta}{2\pi} e^{i\Phi} \sum_{n=-1}^{\infty} \int \frac{d^2 p_\parallel}{(2\pi)^2} \frac{e^{-i(pZ)_\parallel} e^{-\beta Z_\perp^2/4}}{p_\parallel^2 - M_n^2 + i\varepsilon} \left(d^\mu_\nu + \frac{\xi - 1}{p_\parallel^2 - \tilde{M}_n^2 + i\varepsilon} f^\mu_\nu \right), \quad (73)$$

$$d^\mu_\nu = \delta^\mu_{\parallel\nu} L_n + \frac{1}{2} \delta^\mu_{\perp\nu} (L_{n+1} + L_{n-1}) - \frac{iQ}{2} \varphi^\mu_\nu (L_{n+1} - L_{n-1}), \quad (74)$$

$$f^\mu_\nu = \left[p_\parallel^\mu p_{\parallel\nu} + \frac{\beta Q}{2} (p_\parallel^\mu (Z\varphi)_\nu + (\varphi Z)^\mu p_{\parallel\nu}) - \left((2n+1)\beta - \frac{\beta^2}{4} Z_\perp^2 \right) \delta^\mu_{\perp\nu} + \frac{iQ\beta}{2} \varphi^\mu_\nu \right] L_n + \frac{i\beta}{2} \left[(p_\parallel^\mu Z_{\perp\nu} + Z_\perp^\mu p_{\parallel\nu}) - i\delta^\mu_{\perp\nu} - \frac{Q\beta}{2} Z_\perp^2 \varphi^\mu_\nu \right] (L_n^{(1)} + L_{n-1}^{(1)}) - \beta^2 (\varphi Z)^\mu (Z\varphi)_\nu L_{n-1}^{(2)}. \quad (75)$$

In Eqs. (73) – (75), $M_n^2 = m^2 + (2n+1)\beta$, $\tilde{M}_n^2 = \xi m^2 + (2n+1)\beta$ and all the Laguerre polynomials $L_n^{(m)}$ have $\beta Z_\perp^2/2$ as their arguments: $L_n^{(m)} = L_n^{(m)}(\beta Z_\perp^2/2)$.

Next, we have to consider both J_\parallel and a similar integral

$$\tilde{J}_\parallel = \int \frac{d^2 p_\parallel}{(2\pi)^2} \frac{e^{-i(pZ)_\parallel}}{p_\parallel^2 - M_n^2 + i\varepsilon} \frac{1}{p_\parallel^2 - \tilde{M}_n^2 + i\varepsilon} = \frac{i}{2\pi} \frac{1}{m^2(\xi - 1)} \left[K_0 - \frac{i\pi}{2} H_0^{(2)} - \tilde{K}_0 + \frac{i\pi}{2} \tilde{H}_0^{(2)} \right], \quad (76)$$

where $\tilde{K}_0 = K_0 \left(\tilde{M}_n \sqrt{-Z_\parallel^2 + i\varepsilon} \right) \theta(-Z_\parallel^2)$ and $\tilde{H}_0^{(2)} = H_0^{(2)} \left(\tilde{M}_n \sqrt{Z_\parallel^2 - i\varepsilon} \right) \theta(Z_\parallel^2)$. For the ground-state level, one can notice that the large field values lead to the vacuum instability [19].

Replacing p_\parallel by $i\partial_\parallel$ in (75), we finally obtain the position-space representation of the charged massive vector boson propagator as an expansion over Landau levels:

$$G^\mu_\nu(X, X') = \frac{-i\beta}{4\pi^2} e^{i\Phi} e^{-\beta Z_\perp^2/4} \sum_{n=-1}^{\infty} \left\{ P^\mu_\nu - \frac{1}{m^2} Q^\mu_\nu \right\}, \quad (77)$$

$$P^\mu_\nu = \left[\delta^\mu_{\parallel\nu} L_n + \frac{1}{2} \delta^\mu_{\perp\nu} (L_{n+1} + L_{n-1}) - \frac{iQ}{2} \varphi^\mu_\nu (L_{n+1} - L_{n-1}) \right] \left[K_0 - \frac{i\pi}{2} H_0^{(2)} \right], \quad (78)$$

$$Q^\mu_\nu = \left[\left\{ -\partial_\parallel^\mu \partial_{\parallel\nu} + \frac{i\beta Q}{2} (\partial_\parallel^\mu (Z\varphi)_\nu + (\varphi Z)^\mu \partial_{\parallel\nu}) - \left((2n+1)\beta - \frac{\beta^2}{4} Z_\perp^2 \right) \delta^\mu_{\perp\nu} + \frac{iQ\beta}{2} \varphi^\mu_\nu \right\} L_n - \frac{\beta}{2} \left\{ (\partial_\parallel^\mu Z_{\perp\nu} + Z_\perp^\mu \partial_{\parallel\nu}) - \delta^\mu_{\perp\nu} + \frac{iQ\beta}{2} Z_\perp^2 \varphi^\mu_\nu \right\} (L_n^{(1)} + L_{n-1}^{(1)}) - \beta^2 (\varphi Z)^\mu (Z\varphi)_\nu L_{n-1}^{(2)} \right] \times \left[K_0 - \frac{i\pi}{2} H_0^{(2)} - \tilde{K}_0 + \frac{i\pi}{2} \tilde{H}_0^{(2)} \right]. \quad (79)$$

V. CONCLUSION

We considered three methods for obtaining the position-space Landau-levels representation of a charged scalar particle's propagator in a constant magnetic field, namely, the Fock-Schwinger method (FS), the modified Fock-Schwinger method (MFS) and the "sum over solutions" approach. All of them showed the consistent results and provided various useful intermediate expressions of the propagator.

In the expansion over Landau levels, each summation term decomposed into two factors. The first one depends only on the coordinates of the plane perpendicular to the

direction of magnetic field and is rotationally invariant, thus, emphasizing the underlying symmetry of the problem. Being a product of a Laguerre polynomial and a damping exponential, it localizes the propagation in the x, y -plane. The second factor describes propagation in a 1+1 – dimensional space-time, and is invariant with respect to Lorentz transformations in this subspace. Similarly to the case of a free field, it contains both a time-like oscillatory term $H_0^{(2)}$ and a space-like damping term K_0 .

Additionally, we obtained the respective propagators for the charged fermions and charged massive vector bosons. These expressions could be useful in the calculations of processes in external fields, and the discussed

methods are likely to contribute to the development of the position-space techniques in QFT.

ACKNOWLEDGMENTS

We are grateful to A.Ya. Parkhomenko and D. A. Romyantsev for useful remarks.

The reported study was funded by RFBR, project number 19-32-90137.

-
- [1] N. Bogoliubov and D. Shirkov, *Introduction to the theory of quantized fields* (1959).
- [2] W. Greiner and J. Reinhardt, *Quantum electrodynamics* (Springer-Verlag Berlin Heidelberg, 2009).
- [3] J. Schwinger, Phys. Rev. **82**, 664 (1951).
- [4] C. Itzykson and J. Zuber, *Quantum Field Theory*, International Series In Pure and Applied Physics (McGraw-Hill, New York, 1980).
- [5] V. Fock, Phys. Z. Sowjetunion **12**, 404 (1937).
- [6] A. Erdas and G. Feldman, Nuclear Physics B **343**, 597 (1990).
- [7] A. Chodos, K. Everding, and D. A. Owen, Phys. Rev. D **42**, 2881 (1990).
- [8] A. Erdas and C. Isola, Physics Letters B **494**, 262 (2000).
- [9] T.-K. Chyi, C.-W. Hwang, W. F. Kao, G.-L. Lin, K.-W. Ng, and J.-J. Tseng, Phys. Rev. D **62**, 105014 (2000).
- [10] A. Ayala, A. Sánchez, G. Piccinelli, and S. Sahu, Phys. Rev. D **71**, 023004 (2005).
- [11] A. V. Kuznetsov and A. A. Okrugin, International Journal of Modern Physics A **26**, 2725 (2011), <https://doi.org/10.1142/S0217751X11053535>.
- [12] A. Kuznetsov and N. Mikheev, *Electroweak Processes in External Active Media* (Springer, 2013).
- [13] A. V. Kuznetsov, A. A. Okrugin, and A. M. Shitova, International Journal of Modern Physics A **30**, 1550140 (2015), <https://doi.org/10.1142/S0217751X15501407>.
- [14] S. N. Iablokov and A. V. Kuznetsov, J. Phys. Conf. Ser. **1690**, 012087 (2020), arXiv:2010.05195 [hep-th].
- [15] S. N. Iablokov and A. V. Kuznetsov, Journal of Physics: Conference Series **1390**, 012078 (2019).
- [16] S. N. Iablokov and A. V. Kuznetsov, Phys. Rev. D **102**, 096015 (2020), arXiv:2008.07890 [hep-th].
- [17] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products, 8th ed.* (Academic Press, 2015).
- [18] H.-H. Zhang, K.-X. Feng, S.-W. Qiu, A. Zhao, and X.-S. Li, Chin. Phys. C **34**, 1576 (2010), arXiv:0811.1261 [math-ph].
- [19] N. K. Nielsen and P. Olesen, Nucl. Phys. B **144**, 376 (1978).