

Type-3/2 Seesaw Mechanism

Durmuş Demir,¹ Canan Karahan,^{2,*} and Ozan Sargin³

¹*Sabancı University, Faculty of Engineering and Natural Sciences, 34956 Tuzla İstanbul, Turkey*

²*Physics Engineering Department, İstanbul Technical University, 34469 Maslak İstanbul, Turkey*

³*Izmir Institute of Technology, Department of Physics, 35430, Izmir, Turkey*

(Dated: March 16, 2022)

Type-I seesaw provides a natural explanation for the tiny neutrino masses. The right-handed neutrino masses it requires are, however, too large to keep the Higgs boson mass at its measured value. Here we show that vector-spinors, singlet leptons like the right-handed neutrinos, generate the tiny neutrino masses naturally by the exchange of its spin-1/2 and spin-3/2 components. This one-step seesaw mechanism, which we call Type-3/2 seesaw, keeps the Higgs boson mass unchanged at one loop, and gives cause therefore to no fine-tuning problem. If the on-shell vector-spinor is a pure spin-3/2 particle then it becomes a potential candidate for hidden dark matter diluting due only to the expansion of the Universe. The Type-3/2 seesaw provides a natural framework for the neutrino, Higgs boson and dark matter sectors, with overall agreement with current experiments and observations.

Introduction— Neutrino oscillations [1, 2] are a proof that the active neutrinos are massive. Neutrino mass, whose nature is still an open question, is a proof that there is new physics beyond the Standard Model (SM) [3–8].

Even though neutrino physics has come of age in the past two decades [22], there is still no telling if neutrinos are Dirac ($\nu \neq \nu^c$) or Majorana ($\nu = \nu^c$) fermions [8–11]. The Dirac masses [12–19] conserve lepton number. The Dirac neutrinos acquire their masses via the electroweak breaking as all the other fermions, albeit with an unnaturally small Yukawa coupling. The Majorana masses, on the other hand, break lepton number, and arise via the electroweak breaking, with naturally heavy SM-singlet right-handed neutrinos [3–6].

The new physics that generates the tiny neutrino masses has been modeled variously by invoking various fields, symmetries and scales [20–22]. The Type-I seesaw mechanism, the first and the foremost of all [4], leads to small active neutrino masses [4] via the dimension-5 Weinberg operator [23] induced by the heavy right-handed neutrino mediation. There are also Type-II [24–26] and Type-III [27] seesaw mechanisms, which are mediated, respectively, by the triplet scalars and triplet fermions. In addition to these tree-level mechanisms, various models [28–39] have been constructed to generate the small neutrino masses radiatively.

In this Letter we further the Type-I seesaw. We do this by replacing the right-handed neutrino with a vector-spinor field [40] involving both spin-3/2 and spin-1/2 components when it is off-shell and only a spin-3/2 component when it is on-shell. We call the resulting neutrino mass generation mechanism “Type-3/2 seesaw

mechanism”. As will be shown in the sequel, this new mediator leads to important new results for the neutrino, Higgs and dark matter sectors.

Below, we first summarize the Type-I seesaw, and discuss its implications for leptogenesis, dark matter and Higgs boson mass. Next, we turn to the vector-spinor and construct the Type-3/2 seesaw, with a detailed discussion of its implications for leptogenesis, dark matter and the Higgs boson mass. Finally, we conclude the Letter by contrasting the salient features and implications of the Type-I and Type-3/2 seesaw mechanisms.

Type-I Seesaw— The right-handed neutrino N , an SM-singlet spin-1/2 neutral fermion [3, 4], gives a simple model of the tiny neutrino masses. It couples to the active neutrinos ν_L through the left-handed lepton doublet $L = (\nu_L, \ell_L)$ as

$$y_N \bar{L} H N + \frac{M_N}{2} \bar{N} N + \text{H.C.} \quad (1)$$

so that an active neutrino acquires the Majorana mass

$$m_\nu = \frac{y_N^2 \langle H \rangle^2}{2M_N} \quad (2)$$

via the Feynman diagram in Fig. 1. This induced mass, resulting from the Weinberg operator [23], agrees with the experimental data [1, 2] for right-handed neutrino masses $M_N \approx 10^{14}$ GeV, Higgs vacuum expectation value $\langle H \rangle \approx 246$ GeV, and Yukawa coupling $y_N \approx \mathcal{O}(1)$. This dynamical mechanism, the Type-I seesaw [3–6], generates neutrino Majorana masses m_ν naturally with naturally heavy SM-singlet right-handed neutrinos [7]. Neutrino mixings [21, 22] are realized with two or more right-handed neutrinos.

The interactions in (1) give cause for not only the neutrino masses as in (2) but also the Higgs mass shift [41]

$$(\delta m_h^2)_N = \frac{m_\nu M_N^3}{2\pi^2 \langle H \rangle^2} \log \frac{Q}{M_N} \quad (3)$$

* ckarahan@itu.edu.tr




FIG. 1. Type-I seesaw: Heavy right-handed neutrinos N lead to the neutrino Majorana masses m_ν .

at the renormalization scale $Q \gtrsim M_N$ via the Feynman diagram in Fig. 2. This mass correction, evaluated in dimensional regularization in which quadratic (and quartic) corrections all vanish identically [42], exceeds the Higgs boson mass itself unless $M_N \lesssim 10^7$ GeV [43–45]. This bound is in clear contradiction with the value $M_N \approx 10^{14}$ GeV required by the neutrino masses. This contradiction shows that the right-handed neutrinos generate the active neutrino masses at the expense of an immense fine-tuning of the model parameters entering the Higgs boson mass [46]. This is a serious naturalness problem because the logarithmic correction (3) survives to impede the Type-I seesaw [44] even in the supersymmetry [47].

The right-handed neutrinos can decay and annihilate [48] via their interactions with the SM fields in (1), and can facilitate therefore, for instance, the leptogenesis [49, 50]. Thermal leptogenesis requires the lightest right-handed neutrino N_1 to have a mass $M_{N_1} \gtrsim 2 \times 10^9$ GeV in order to produce the requisite asymmetry in the lepton sector. To be able to produce such a massive N_1 thermally, the reheat temperature T_{rh} after the inflation must have a value $T_{rh} > M_{N_1}$ [51–53]. It is clear that this leptogenesis value of M_{N_1} , too, is in contradiction with the bound $M_{N_1} \lesssim 10^7$ GeV imposed by the Higgs boson mass correction in (3). Even though it is not possible to suppress all the radiative corrections, sometimes gravity-mediated softly broken supersymmetry is incorporated into the thermal leptogenesis to reduce the quadratic corrections from heavy right-handed neutrinos to the milder logarithmic ones. This attempt leads, however, to the well-known gravitino problem [54, 55]. It turns out that for gravitinos of masses below a few TeV the reheat temperature of the Universe should not exceed 10^5 GeV [56, 57].




FIG. 2. The active neutrino/right-handed neutrino loop that gives cause to the Higgs boson mass shift in (3).

The Type-I seesaw does not offer a unique dark matter candidate though, as an analogous low-energy extension, one can consider incorporating the sterile neutrinos into the setup [58].

Type-3/2 Seesaw— Having discussed the Type-I seesaw and its physics implications, we now develop a new approach in which we envision the right-handed neutrino as spin-1/2 component of an SM-singlet, neutral vector-spinor ψ_μ of mass M_ψ . This Rarita-Schwinger field [40] decomposes as $[(1, \frac{1}{2}) \oplus (0, \frac{1}{2})] \oplus [(\frac{1}{2}, 1) \oplus (\frac{1}{2}, 0)]$ under the Lorentz group. Namely, it involves spin-3/2 and spin-1/2 components. As was already pointed out in [59], it directly couples to the Higgs and lepton doublets via the Lagrangian

$$y_\psi \bar{L} H \gamma^\mu \psi_\mu + y_\psi \bar{\psi}_\mu \gamma^\mu H^\dagger L + \bar{\psi}_\mu \Lambda^{\mu\nu} \psi_\nu \quad (4)$$

in which the kinetic operator [40]

$$\Lambda^{\mu\nu} = \eta^{\mu\nu} (\not{p} - M_\psi) - (\gamma^\mu p^\nu + p^\mu \gamma^\nu) + \gamma^\mu \not{p} \gamma^\nu + M_\psi \gamma^\mu \gamma^\nu \quad (5)$$

differs from the Dirac operator by the terms in the second line. The Lagrangian (4) leads to the equation of motion

$$\Lambda_{\mu\nu} \psi^\nu = y_\psi \gamma_\mu H^\dagger L \quad (6)$$

in agreement with the vector-spinor description in [60, 61]. For a free field, that is, for an on-shell vector-spinor $\psi_{(free)}^\mu(p)$ satisfying $p^2 = M_\psi^2$ one gets the equation of motion $\Lambda_{\mu\nu} \psi_{(free)}^\nu = 0$. This homogeneous equation can be consistently split into three distinct parts

$$(\not{p} - M_\psi) \psi_{(free)}^\mu = 0, \quad (7)$$

$$\gamma_\mu \psi_{(free)}^\mu = 0, \quad (8)$$

$$p_\mu \psi_{(free)}^\mu = 0 \quad (9)$$

as follows from the kinetic structure in (5) as particular choices for its individual terms. Here the point is that the equations (8) and (9) eliminate the spin-1/2 component of the vector-spinor $\psi_{(free)}^\mu$ to yield an on-shell pure spin-3/2 field.

For an off-shell vector-spinor $\psi^\mu(p)$ satisfying $p^2 \neq M_\psi^2$ the description is given by its propagator $S^{\mu\nu}$ [60, 61]

$$S^{\mu\nu} = -\frac{i\eta^{\mu\nu}}{\not{p} - M_\psi} + \frac{i}{\not{p} - M_\psi} \left(\frac{\gamma^\mu \gamma^\nu}{3} + \frac{(\gamma^\mu p^\nu - p^\mu \gamma^\nu)}{3M_\psi} + \frac{2p^\mu p^\nu}{3M_\psi^2} \right) \quad (10)$$

whose second line stands for deviation from the spin-1/2 propagator in parallel with (5). This propagator is nothing but the inverse of the kinetic structure in (5) namely $S^{\mu\alpha} \Lambda_{\alpha\nu} = \delta_\nu^\mu$. This propagator holds in the entire momentum and spin space [62]. In other words, this vector-spinor propagator involves propagations of both spin-3/2 and spin-1/2 components. It certainly relates ψ_μ to its source in (6) but it does not satisfy the individual motion equations in (7), (8) and (9).




FIG. 3. Type-3/2 seesaw: Heavy vector-spinor neutrinos ψ_μ lead to the neutrino Majorana masses m_ν .

It proves useful to contrast the vector-spinor above with the gravitino [63, 64] – the gauge field of supergravity which acquires mass by swallowing the goldstino field [65]. The gravitino is a spin-3/2 field and therefore obeys the equations (7), (8) and (9) when it is on-shell and off-shell. The gravitino propagator is constructed in [63] by imposing (7), (8) and (9) (with (8) and (9) being kind of gauge conditions) so that as a propagator it satisfies (7), (8) and (9).

The vector-spinor ψ_μ is different than the gravitino. Its propagator in (10) propagates both spin-3/2 and spin-1/2 components, and does not obey the motion equations (7), (8) and (9). It can be reduced to a spin-3/2 particle when it is on-shell, that is, when it obeys the individual motion equations (7), (8) and (9). Its electric neutrality ensures that there arises no problem with local causality [66, 67].

In similarity to the right-handed neutrinos, the ψ_μ couplings in (4) lead to the active neutrino masses

$$m_\nu = \frac{2y_\psi^2 \langle H \rangle^2}{9M_\psi} \quad (11)$$

via the Feynman diagram in Fig. 3 with the exchange of spin-3/2 and spin-1/2 components. This result, derived by using the ψ_μ propagator in (10), agrees with the experimental data [1, 2] for $M_\psi \approx 10^{14}$ GeV and $y_\psi \approx \mathcal{O}(1)$. This new mechanism, which is what we term as Type-3/2 seesaw, generates the neutrino Majorana masses naturally with a naturally heavy SM-singlet vector-spinor ψ_μ . Neutrino mixings [21, 22] can be realized with two or more ψ_μ fields.

If the vector-spinor ψ_μ obeys the three equations of motion (7), (8) and (9) then it becomes a spin-3/2 field when it is on-shell and in this case, in contrast to the right-handed neutrinos, it can neither decay nor annihilate despite its couplings in (4) to the SM fields. The reason is that in such processes ψ_μ is on its mass-shell as an isolated physical particle and for an on-shell ψ_μ namely for a ψ_μ satisfying (7), (8) and (9) the H - L - ψ_μ vertex vanishes identically ([60, 61]). (This

vertex vanishes for both on-shell and off-shell gravitino [63].) This means that the scattering processes with on-shell ψ_μ (its decays, annihilations and productions) all vanish. This on-shell nullity of ψ_μ has three important implications. Firstly, ψ_μ , unlike the right-handed neutrinos, cannot facilitate leptogenesis simply because there exists no decay channel to transfer lepton number in ψ_μ to active neutrinos and charged leptons [68]. Needless to say, baryogenesis can occur via some other mechanism such as Affleck-Dine [69] baryogenesis, and this sets the theory exempt from the gravitino problem [70] in gravity-mediated softly-broken supersymmetry [65].

Secondly, if on-shell ψ_μ is a spin-3/2 particle obeying (7), (8) and (9) then it is an everlasting particle. It was around, is around and will be around to participate in certain processes in a hidden or invisible way [59]. Its density falls with the volume of the Universe as it dilutes due only to the expansion of the Universe. It can therefore form dark matter [71] if it leads to flat rotation curves, structure formation and other relevant phenomena. The setup in (4) can be modified to obtain a detectable and more conventional dark matter candidate. One possibility is to invoke higher-dimension operators [72] but the H - L - ψ_μ coupling in (4) must still be taken into account when ψ_μ is off-shell.




FIG. 4. The vector-spinor/active neutrino loop that does not lead to any Higgs boson mass shift.

Thirdly, when it is off-shell ψ_μ reveals itself via its spin-3/2 and spin-1/2 components [59] by inducing the neutrino Majorana masses as in (11), altering certain SM scattering amplitudes such as $hh \rightarrow \bar{\nu}_L \nu_L$ [59], and facilitating loop corrections to certain SM masses and couplings. Its loop with the active neutrinos, depicted in Fig. 4, is expected to shift the Higgs boson mass as in (3) but, to one's surprise, it gives actually zero contribution

$$(\delta m_h^2)_{\psi_\mu} = 0 \quad (12)$$

at one loop. This follows from the fact that the Feynman diagram in Fig. 4 evaluates to zero

$$\begin{aligned}
& -i \frac{y_\psi^2}{2} \int \frac{d^4 p}{(2\pi)^4} \frac{1}{p^2(p^2 - M_\psi^2)} \text{Tr} \left[P_R \gamma_\alpha (\not{p} + M_\psi) \left\{ \eta^{\alpha\beta} - \frac{\gamma^\alpha \gamma^\beta}{3} - \frac{2p^\alpha p^\beta}{3M_\psi^2} - \frac{(\gamma^\alpha p^\beta - p^\alpha \gamma^\beta)}{3M_\psi} \right\} \gamma_\beta \not{p} P_R \right] \\
& = -i y_\psi^2 M_\psi^2 \lim_{\epsilon \rightarrow 0} Q^\epsilon \left(\frac{M_\psi^2}{4\pi} \right)^{2-\frac{\epsilon}{2}} \left[\Gamma \left(-1 + \frac{\epsilon}{2} \right) - \left(-2 + \frac{\epsilon}{2} \right) \Gamma \left(-2 + \frac{\epsilon}{2} \right) \right] = 0
\end{aligned} \tag{13}$$

in confirmation of the nullity of $\delta^4(0)$ in dimensional regularization [42]. This zero Higgs mass shift, parametrized by the ψ_μ momentum p_α , right projector $P_R = (1 + \gamma_5)/2$, and the renormalization scale Q , is evaluated in the same regularization scheme as the Higgs mass shift in (3). It ensures that the active neutrinos and the Higgs boson can acquire their measured masses with no contradiction concerning the scale of $M_{3/2}$. The naturalness (fine-tuning) problem impeding the Type-I seesaw does simply not exist in the Type-3/2 seesaw, owing to (12).

Conclusion— In this Letter we have shown that the SM-singlet vector-spinor [40] gives rise to a new one-step seesaw mechanism. This new mechanism, the Type-3/2 seesaw, exhibits physically important features not found in the Type-I seesaw. The salient features are:

- Tiny neutrino Majorana masses arise naturally (in agreement with the Type-I seesaw),
- Higgs boson acquires its mass without fine-tuning (in disagreement with the Type-I seesaw),
- If the on-shell vector-spinor is a spin-3/2 field then baryogenesis is not sourced by leptogenesis (in disagreement with the Type-I seesaw), and this makes the model impervious to the gravitino problem in the Type-I seesaw, and finally
- If the on-shell vector-spinor is a spin-3/2 field dark matter can exist as an everlasting field hidden in

the SM spectrum (in disagreement with Type-I seesaw).

These points are tabulated in Table I in a comparative fashion. In view of them, one can conclude that the Type-3/2 seesaw proposed in this Letter can open up a novel approach to neutrino and dark sector phenomenology, with its inherent naturalness in both the neutrino and Higgs sectors. The minimal structure considered in this Letter can be extended to make the spin-3/2 dark matter detectable. This can be done in various ways such as using higher-dimension operators as in [72, 73] or adding new fields, such as a scalar field coupling to $\psi_\mu \psi^\mu$ [74].

ACKNOWLEDGMENTS

The work of CK is supported by İTÜ BAP grant TAB-2020-42312.

TABLE I. A comparative summary of the implications of the Type-I and Type-3/2 seesaw mechanisms. The label ^(a) is a warning that the Type-I seesaw does not bring up any distinctive dark matter candidate. The label ^(b), on the other hand, is a reminder that baryogenesis must occur via mechanisms other than leptogenesis.

	Type-I Seesaw	Type-3/2 Seesaw
ν -oscillations	Yes	Yes
Stable Higgs mass	No	Yes
Dark matter	No ^(a)	Yes
Leptogenesis	Yes	No ^(b)

-
- [1] Q. R. Ahmad *et al.* [SNO], Phys. Rev. Lett. **87**, 071301 (2001) [arXiv: 0106015 [nucl-ex]].
- [2] Y. Fukuda *et al.* [Super-Kamiokande], Phys. Rev. Lett. **81**, 1562-1567 (1998) [arXiv: 9807003 [hep-ex]].
- [3] P. Minkowski, Phys. Lett. B **67**, 421-428 (1977).
- [4] M. Gell-Mann, P. Ramond and R. Slansky, Conf. Proc. C **790927**, 315-321 (1979) [arXiv: 1306.4669 [hep-th]].
- [5] T. Yanagida, in *Workshop on the Unified Theories and the Baryon Number in the Universe*, Ed. O. Sawada and A. Sugamoto (Tsukuba, Japan, 1979).
- [6] R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
- [7] R. N. Mohapatra and A. Y. Smirnov, Ann. Rev. Nucl. Part. Sci. **56**, 569-628 (2006) [arXiv: 0603118 [hep-ph]].
- [8] P. Langacker, Int. J. Mod. Phys. A **20**, 5254-5265 (2005) [arXiv: 0411116 [hep-ph]].
- [9] M. Agostini *et al.* [GERDA], Phys. Rev. Lett. **125**, no.25, 252502 (2020) [arXiv:2009.06079 [nucl-ex]].
- [10] M. J. Dolinski, A. W. P. Poon and W. Rodejohann, Ann. Rev. Nucl. Part. Sci. **69**, 219-251 (2019) [arXiv: 1902.04097 [nucl-ex]].
- [11] M. Czakon, M. Zralek and J. Gluza, Acta Phys. Polon. B **30**, 3121-3138 (1999) [arXiv: 9910357 [hep-ph]].
- [12] D. A. Demir, L. L. Everett and P. Langacker, Phys. Rev. Lett. **100**, 091804 (2008) [arXiv: 0712.1341 [hep-ph]].
- [13] C. Bonilla and J. W. F. Valle, Phys. Lett. B **762**, 162-165

- (2016) [arXiv: 1605.08362 [hep-ph]].
- [14] W. Wang and Z. L. Han, JHEP **04**, 166 (2017) [arXiv: 1611.03240 [hep-ph]].
- [15] W. Wang, R. Wang, Z. L. Han and J. Z. Han, Eur. Phys. J. C **77**, no.12, 889 (2017) [arXiv: 1705.00414 [hep-ph]].
- [16] C. Y. Yao and G. J. Ding, Phys. Rev. D **97**, no.9, 095042 (2018) [arXiv: 1802.05231 [hep-ph]].
- [17] J. Calle, D. Restrepo, C. E. Yaguna and Ó. Zapata, Phys. Rev. D **99**, no.7, 075008 (2019) [arXiv: 1812.05523 [hep-ph]].
- [18] S. Saad, Nucl. Phys. B **943**, 114636 (2019) [arXiv: 1902.07259 [hep-ph]].
- [19] S. Jana, V. P. K. and S. Saad, Eur. Phys. J. C **79**, no.11, 916 (2019) [arXiv: 1904.07407 [hep-ph]].
- [20] S. F. King, Rept. Prog. Phys. **67**, 107-158 (2004) [arXiv: 0310204 [hep-ph]].
- [21] R. N. Mohapatra, S. Antusch, K. S. Babu, G. Barenboim, M. C. Chen, S. Davidson, A. de Gouvêa, P. de Holanda, B. Dutta and Y. Grossman, *et al.* Rept. Prog. Phys. **70**, 1757-1867 (2007) [arXiv: 0510213 [hep-ph]].
- [22] A. Baha Balantekin and B. Kayser, Ann. Rev. Nucl. Part. Sci. **68**, 313-338 (2018) [arXiv: 1805.00922 [hep-ph]].
- [23] S. Weinberg, Physica A **96**, no.1-2, 327-340 (1979).
- [24] W. Konetschny and W. Kummer, Phys. Lett. B **70**, 433 (1977).
- [25] T. P. Cheng and L. F. Li, Phys. Rev. D **22**, 2860 (1980).
- [26] J. Schechter and J. W. F. Valle, Phys. Rev. D **22**, 2227 (1980).
- [27] R. Foot, H. Lew, X. G. He and G. C. Joshi, Z. Phys. C. **44**, 441 (1989).
- [28] W. S. Hou and G. G. Wong, Phys. Lett. B **339**, 109-113 (1994) [arXiv: 9405395 [hep-ph]].
- [29] T. Nomura, H. Okada and Y. Uesaka, JHEP **01**, 016 (2021) [arXiv: 2005.05527 [hep-ph]].
- [30] K. Bamba, C. Q. Geng and S. H. Ho, JCAP **09**, 001 (2008) [arXiv: 0806.0952 [hep-ph]].
- [31] E. Ma and D. Suematsu, Mod. Phys. Lett. A **24**, 583-589 (2009) [arXiv: 0809.0942 [hep-ph]].
- [32] E. Ma, Phys. Rev. D **80**, 013013 (2009) [arXiv: 0904.4450 [hep-ph]].
- [33] K. S. Babu and J. Julio, Phys. Rev. D **85**, 073005 (2012) [arXiv: 1112.5452 [hep-ph]].
- [34] D. Hehn and A. Ibarra, Phys. Lett. B **718**, 988-991 (2013) [arXiv: 1208.3162 [hep-ph]].
- [35] A. Hriche, K. L. McDonald and S. Nasri, JHEP **10**, 167 (2014) [arXiv: 1404.5917 [hep-ph]].
- [36] T. Nomura, H. Okada and Y. Orikasa, Eur. Phys. J. C **77**, no.2, 103 (2017) [arXiv: 1602.08302 [hep-ph]].
- [37] A. Hriche, A. Jueid and S. Nasri, Phys. Rev. D **97**, no.9, 095012 (2018) [arXiv: 1710.03824 [hep-ph]].
- [38] Y. Cai, J. Herrero-García, M. A. Schmidt, A. Vicente and R. R. Volkas, Front. in Phys. **5**, 63 (2017) [arXiv: 1706.08524 [hep-ph]].
- [39] C. Klein, M. Lindner and S. Ohmer, JHEP **03**, 018 (2019) [arXiv: 1901.03225 [hep-ph]].
- [40] W. Rarita and J. Schwinger, Phys. Rev. **60**, 61 (1941).
- [41] F. Vissani, Phys. Rev. D **57** (1998) 7027 [arXiv: 9709409 [hep-ph]].
- [42] G. Leibbrandt, Rev. Mod. Phys. **47**, 849 (1975).
- [43] P. Chattopadhyay and K. M. Patel, Nucl. Phys. B **921** (2017) 487 [arXiv: 1703.09541 [hep-ph]].
- [44] F. Bazzocchi and M. Fabbrichesi, Phys. Rev. D **87**, no.3, 036001 (2013) [arXiv: 1212.5065 [hep-ph]].
- [45] G. Bambhaniya, P. S. Bhupal Dev, S. Goswami, S. Khan and W. Rodejohann, Phys. Rev. D **95**, no.9, 095016 (2017) [arXiv: 1611.03827 [hep-ph]].
- [46] J. A. Casas, J. R. Espinosa and I. Hidalgo, JHEP **11**, 057 (2004) [arXiv: 0410298 [hep-ph]].
- [47] H. Baer and X. Tata, Weak Scale Supersymmetry: From Superfields to Scattering Events, (Cambridge University Press, 2006).
- [48] M. Drewes, Int. J. Mod. Phys. E **22**, 1330019 (2013) [arXiv: 1303.6912 [hep-ph]].
- [49] M. A. Luty, Phys. Rev. D **45**, 455-465 (1992).
- [50] W. Buchmuller, R. D. Peccei and T. Yanagida, Ann. Rev. Nucl. Part. Sci. **55**, 311-355 (2005) [arXiv: 0502169 [hep-ph]].
- [51] W. Buchmüller, P. Di Bari and M. Plumacher, Nucl. Phys. B **643** 367 (2002) and Erratum-ibid,B793 362 (2008).
- [52] W. Buchmüller, P. Di Bari and M. Plumacher, Annals Phys. **315**, 305 (2005).
- [53] W. Buchmüller, P. Di Bari and M. Plumacher, New J. Phys. **6**, 105 (2004).
- [54] M. Khlopov and A. Linde, Phys. Lett. B **138**, 265 (1984).
- [55] J.R. Ellis, J.E. Kim, D.V. Nanopoulos, Phys. Lett. B **145**, 181 (1984).
- [56] M. Kawasaki, K. Kohri, T. Moroi, A. Yotsuyanagi, Phys. Rev. D **78** (6), 065011 (2008).
- [57] R.H. Cyburt, J.R. Ellis, B.D. Fields, K.A. Olive, Phys. Rev. D **67**, 103521 (2003).
- [58] A. Boyarsky, M. Drewes, T. Lasserre, S. Mertens and O. Ruchayskiy, Prog. Part. Nucl. Phys. **104**, 1-45 (2019) [arXiv:1807.07938 [hep-ph]].
- [59] D. Demir, C. Karahan, B. Korutlu and O. Sargin, Eur. Phys. J. C **77**, no.9, 593 (2017) [arXiv: 1708.07956 [hep-ph]].
- [60] V. Pascalutsa, Phys. Lett. B **503**, 85-90 (2001) [arXiv: 0008026 [hep-ph]].
- [61] T. Pilling, Int. J. Mod. Phys. A **20**, 2715-2742 (2005) [arXiv: 0404131 [hep-th]].
- [62] H. Haberkzettel, [arXiv:nucl-th/9812043 [nucl-th]].
- [63] P. Van Nieuwenhuizen, Phys. Rept. **68**, 189-398 (1981)
- [64] V. S. Rychkov and A. Strumia, Phys. Rev. D **75**, 075011 (2007) [arXiv:hep-ph/0701104 [hep-ph]].
- [65] L. J. Hall, J. D. Lykken and S. Weinberg, Phys. Rev. D **27**, 2359-2378 (1983).
- [66] K. Johnson and E. C. G. Sudarshan, Annals Phys.**13**, 126-145 (1961).
- [67] Giorgio Velo and Daniel Zwanziger, Phys. Rev. **186**, 1337 (1969).
- [68] O. Sargin, “Leptogenesis via spin-3/2 field”, work in progress (2021).
- [69] I. Affleck and M. Dine, Nucl. Phys. B **249**, 361-380 (1985).
- [70] S. Weinberg. Phys. Rev. Lett. **48**, 1303 (1982).
- [71] V. C. Rubin, W. K. Ford, Jr., N. Thonnard, and M. Roberts, Astron. J. **81**, 687 (1976).
- [72] C. F. Chang, X. G. He and J. Tandean, Phys. Rev. D **96**, no.7, 075026 (2017) [arXiv: 1704.01904 [hep-ph]].
- [73] M. A. G. Garcia, Y. Mambrini, K. A. Olive and S. Verner, Phys. Rev. D **102**, no.8, 083533 (2020) [arXiv: 2006.03325 [hep-ph]].
- [74] Canan Karahan and Ozan Sargin, “Making the spin-3/2 dark matter visible”, work in progress (2021).