

**Comment on “An axially symmetric spacetime with  
causality violation [Phys. Scr. 96 (07), 075208 (2021)],  
arXiv:2004.13769[gr-qc]”**

Faizuddin Ahmed<sup>1</sup>

National Academy Gauripur, Assam, India, 783331

**Abstract**

Recently, an axially symmetric spacetime with causality violation is appeared in Ref. [1]. There, author presented a type III metric with vanishing expansion, shear and twist. The matter-energy represents a pure radiation field with a negative cosmological constant. The spacetime is asymptotically anti-de Sitter space in the radial direction. Actually, this work with different form of the metric was already published in Ref. [4]. Therefore, the current paper Ref. [1] is a duplicate one of the previous work.

Recently, an axially symmetric metric with causality violation is appeared in Ref. [1]. The metric in cylindrical coordinates  $(t, r, \phi, z)$  is given by

$$ds^2 = \frac{dr^2}{\alpha^2 r^2} + r^2 (dz^2 - 2 dt d\phi - t d\phi^2) + \frac{\beta z}{r^2} dr d\phi, \quad (1)$$

where  $\alpha > 0, \beta > 0$  are non-zero real numbers. The coordinates are in the ranges  $-\infty < t < \infty, 0 \leq r < \infty, -\infty < z < \infty$  and  $\phi$  is a periodic.

The scalar curvature invariant constructed from the Riemann tensor are

$$R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} = 24 \alpha^4. \quad (2)$$

with the scalar curvature

$$R = R^\mu{}_\mu = -12 \alpha^2. \quad (3)$$

---

<sup>1</sup>faizuddinahmed15@gmail.com ; faiz4U.enter@rediffmail.com

The space-time (1) satisfy the Field Equations

$$R^{\mu\nu} = \Lambda g^{\mu\nu} + \rho k^\mu k^\nu \quad (4)$$

with the stress-energy tensor the pure radiation field. The radiation energy density with a negative Cosmology constant is given by

$$\Lambda = -3\alpha^2 \quad , \quad \rho = \frac{\alpha^2 \beta^2}{8r^8}, \quad (5)$$

where the null vector for the metric (1)

$$k^\mu = (1, 0, 0, 0) \quad (6)$$

satisfies null condition  $k_\mu k^\mu = 0$ . Therefore, the energy-density of radiation field satisfies the null energy condition (NEC).

One can easily show that the null vector (6) is a covariantly constant null vector field, that is,

$$k_{\mu;\nu} = 0 \quad (7)$$

To classify the metric (1) according to the Petrov classification scheme, author constructed a set of tetrad vectors  $(\mathbf{k}, \mathbf{l}, \mathbf{m}, \bar{\mathbf{m}})$ . Using the set of tetrad vectors, one can calculate the five Weyl scalars and these are

$$\Psi_0 = \Psi_1 = 0 = \Psi_2 \quad , \quad \Psi_3 = \frac{i\alpha^2\beta}{4\sqrt{2}r^2} \quad , \quad \Psi_4 = -\frac{\alpha\beta(i+2rz\alpha)}{8r^2}. \quad (8)$$

Therefore the metric represented by (1) is of type III in the Petrov classification scheme Ref. [2].

One can easily show that the metric (1) is non-expanding, non-twisting and shear-free which are as follows

$$\begin{aligned} \Theta &= \frac{1}{2} k^\mu{}_{;\mu} = 0, \\ \omega &= \sqrt{\frac{1}{2} k_{[\mu;\nu]} k^{\mu;\nu}} = 0, \\ |\sigma| &= \sqrt{\frac{1}{2} k_{(\mu;\nu)} k^{\mu;\nu} - \Theta^2} = 0. \end{aligned} \quad (9)$$

The study space-time admits closed time-like curves (CTCs). Consider an azimuthal curve  $\gamma$  defined by  $\{t = \text{const.}, r = \text{const.}, z = \text{const.}\}$ , and  $\phi$  is a periodic coordinate. From the metric (1), we have

$$ds^2 = -t r^2 d\phi^2. \quad (10)$$

The integral curve with  $(t, r, z)$  fixed is time-like provided  $t > 0$  and thus closed time-like curves are formed at an instant of time Ref. [3].

The original metric with different form was already obtained by other author in Ref. [4]. The original metric in cylindrical coordinate is given by

$$ds^2 = 4r^2 dr^2 + e^{2\alpha r^2} (dz^2 - t d\phi^2 - 2 dt d\phi) + 4\beta z r e^{-\alpha r^2} dr d\phi, \quad (11)$$

where  $\alpha, \beta$  are non-zero real numbers. We do the following transformation

$$r \rightarrow \sqrt{\varrho} \quad (12)$$

into the original metric (11), one will get

$$ds^2 = d\varrho^2 + e^{2\alpha\varrho} (dz^2 - t d\phi^2 - 2 dt d\phi) + 2\beta z e^{-\alpha\varrho} d\varrho d\phi. \quad (13)$$

Finally doing another transformation as follows

$$\varrho \rightarrow \frac{1}{\alpha} \ln r' \quad (14)$$

into the metric (13), we get

$$ds^2 = \frac{dr'^2}{\alpha^2 r'^2} + r'^2 (dz^2 - 2 dt d\phi - t d\phi^2) + \frac{2\beta z}{\alpha r'^2} dr' d\phi, \quad (15)$$

which is very similar to the metric (1). Therefore, the results obtained in the current paper Ref. [1] are same with the previous work Ref. [4]. So, the current paper Ref. [1] is a duplicate one of the previous work.

## References

- [1] Bidyut Bikash Hazarika, An axially symmetric spacetime with causality violation, Phys. Scr. **96** (07), 075208 (2021), arXiv : 2004.13769v1 [gr-qc].
- [2] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers and E. Herlt, Exact Solutions to Einstein's Field Equations, Cambridge University Press, Cambridge (2005).
- [3] F. Ahmed, A type N radiation field solution with  $\Lambda < 0$  in a curved space-time and closed time-like curves, Eur. Phys. J. C **78**, 385 (2018).
- [4] F. Ahmed, Type III Spacetime with Closed Timelike Curves, PROGRESS IN PHYSICS (PiP) Volume **12**, Issue 4 (October), pp. 329-331 (2016), <http://www.ptep-online.com>.