

ABELIAN COVERS AND SECOND FUNDAMENTAL FORM

PAOLA FREDIANI

ABSTRACT. We give some conditions on a family of abelian covers of \mathbb{P}^1 of genus g curves, that ensure that the family yields a subvariety of \mathbf{A}_g which is not totally geodesic, hence it is not Shimura. As a consequence, we show that for any abelian group G , there exists an integer M which only depends on G such that if $g > M$, then the family yields a subvariety of \mathbf{A}_g which is not totally geodesic. We prove then analogous results for families of abelian covers of $\tilde{C}_t \rightarrow \mathbb{P}^1 = \tilde{C}_t/\tilde{G}$ with an abelian Galois group \tilde{G} of even order, proving that under some conditions, if $\sigma \in \tilde{G}$ is an involution, the family of Pryms associated with the covers $\tilde{C}_t \rightarrow C_t = \tilde{C}_t/\langle\sigma\rangle$ yields a subvariety of \mathbf{A}_p^δ which is not totally geodesic. As a consequence, we show that if $\tilde{G} = (\mathbb{Z}/N\mathbb{Z})^m$ with N even, and σ is an involution in \tilde{G} , there exists an integer $M(N)$ which only depends on N such that, if $\tilde{g} = g(\tilde{C}_t) > M(N)$, then the subvariety of the Prym locus in \mathbf{A}_p^δ induced by any such family is not totally geodesic (hence it is not Shimura).

1. INTRODUCTION

In this paper we study families of abelian covers of \mathbb{P}^1 , in relation with the Coleman-Oort conjecture and with an analogue of this conjecture for the Prym maps of (possibly ramified) double covers.

Given a family of Galois covers of genus g curves, $C_t \rightarrow C_t/G \cong \mathbb{P}^1$, the associated family of Jacobians gives a subvariety of \mathbf{A}_g and we ask under which conditions this subvariety is a Shimura subvariety of \mathbf{A}_g . Coleman-Oort conjecture predicts that for g sufficiently high, there should not exist positive dimensional Shimura subvarieties of \mathbf{A}_g generically contained in the Torelli locus. In [9] (and in [15], [16] in the ramified case) a similar question was posed about the existence of positive dimensional Shimura subvarieties of the moduli space of polarised abelian varieties of a fixed dimension, which are generically contained in the Prym loci, that is in the closure of the image of the Prym maps of (possibly ramified) double covers. Both in the case of the Torelli map and in the case of Prym maps, there are examples of such Shimura subvarieties for low values of g . These constructions all use families of Galois covers (see [12], [40], [31], [32], [13], [18], [21], [14] for the Torelli case, [9], [15], [16] for the Prym case).

Let us first explain the main result of this paper in the case of the Torelli morphism. Fix a family of abelian covers of \mathbb{P}^1 , $C_t \rightarrow C_t/G \cong \mathbb{P}^1$, where C_t has genus g . We show that there exists an integer M which only depends on G such that if $g > M$, then the family of Jacobians of the curves C_t yields a subvariety of \mathbf{A}_g which is not totally geodesic, hence it is not Shimura. The moduli space \mathbf{A}_g of principally polarised abelian varieties of dimension g is endowed with an orbifold metric which is induced by the symmetric metric on the Siegel space \mathbf{H}_g and a Shimura subvariety of \mathbf{A}_g is totally geodesic with respect to this metric ([33], [30]).

More precisely, assume we have a family of Galois covers of \mathbb{P}^1 with Galois group G . With the notation used in [13, Section 2], denote by $\mathbf{M}_g(\mathbf{m}, G, \theta)$ the corresponding subvariety of \mathbf{M}_g . Here $\theta : \Gamma_r = \langle \gamma_1, \dots, \gamma_r \mid \prod_{i=1}^r \gamma_i = 1 \rangle \rightarrow G$ denotes the monodromy of the cover ($\mathbf{m} := (m_1, \dots, m_r)$ is such that m_i is the order of $\theta(\gamma_i)$ in G). Consider $j : \mathbf{M}_g \rightarrow \mathbf{A}_g$ the Torelli morphism, which is an orbifold immersion outside the hyperelliptic locus ([37]). The second fundamental form of the Torelli map with respect to the Siegel metric has been studied in [10], [8], [19]. Its dual at a point $[C] \in \mathbf{M}_g$ corresponding to a non hyperelliptic curve C is a map

$$(1.1) \quad \rho : I_2(K_C) \rightarrow S^2 H^0(C, K_C^{\otimes 2})$$

where $I_2(K_C)$ is the kernel of the multiplication map $S^2 H^0(C, K_C) \rightarrow H^0(C, K_C^2)$, that is, the space of quadrics containing the canonical image of C .

In [31] in the cyclic case and in [13, Theorem 3.9] for any group G , it is proven that if the following condition holds

$$(*) \quad \dim(S^2 H^0(C, K_C))^G = \dim H^0(C, K_C^2)^G,$$

for a general member $[C] \in \mathbf{M}_g(\mathbf{m}, G, \theta)$, then the family of Galois covers yields a Shimura (hence totally geodesic) subvariety of \mathbf{A}_g . Moonen proved in [31], using techniques in positive characteristic,

that condition (*) is also necessary for a family of cyclic covers of \mathbb{P}^1 to give a Shimura subvariety of \mathbf{A}_g . This was then generalised in [29], still using positive characteristic techniques, for 1-dimensional families of abelian covers of \mathbb{P}^1 . Using this sufficient condition (*) and computer computations, many examples of Shimura subvarieties of \mathbf{A}_g generically contained in the Torelli locus have been constructed in [31], [32], [13], [18]. Recently in [11] the authors proved that if $g \leq 100$ the examples found in [13], that are all in genus $g \leq 7$, are the only ones satisfying condition (*), thus giving a strong evidence for the Coleman-Oort conjecture (at least for Shimura subvarieties obtained via families of Galois covers of \mathbb{P}^1). In [14] it is shown that the only families of Galois covers of curves of genus $g' > 0$ satisfying condition (*) are the ones found in [18] that are all in low genus $g \leq 4$. In [8, Proposition 5.4] it is proven that if a family of cyclic covers of \mathbb{P}^1 does not satisfy (*) and another condition on the dimension of the eigenspaces for the representation of G on $H^0(C, K_C)$ is satisfied, then the family of cyclic covers gives a subvariety of \mathbf{A}_g which is not even totally geodesic (hence it is not Shimura). Here we generalise this result to families of abelian covers of \mathbb{P}^1 in Theorem 3.1. Let $C \rightarrow C/G \cong \mathbb{P}^1$ be a general element of a family of abelian covers. Since G is abelian, we have an isomorphism of G with the group of characters $G^* = \text{Hom}(G, \mathbb{C}^*)$, hence we identify an element n of G with the corresponding character $\chi \in G^*$ and we denote by $d_n := \dim H^0(C, K_C)_\chi$ (see Section 2).

We have the following

Theorem 1.1. (*Theorem 3.1*)

- (1) *If there exists an element $n \in G \subseteq (\mathbb{Z}/N\mathbb{Z})^m$ such that $n \neq -n$, $d_n \geq 2$, $d_{-n} \geq 2$, then the subvariety of \mathbf{A}_g induced by the family of abelian covers of genus $g \geq 4$ is not totally geodesic (hence it is not Shimura).*
- (2) *If there exists an element $n \in G \subseteq (\mathbb{Z}/N\mathbb{Z})^m$ of order 2 such that $d_n \geq 3$, then the subvariety of \mathbf{A}_g induced by the family of abelian covers of genus $g \geq 4$ is not totally geodesic (hence it is not Shimura).*

As a consequence we prove the following

Theorem 1.2. (*Corollaries 3.2, 3.3, 3.4*)

Assume $G \subseteq (\mathbb{Z}/N\mathbb{Z})^m$, $N \geq 3$, set $d := \#G$, $g \geq 4$. Assume that we have a family of G -covers of \mathbb{P}^1 yielding a totally geodesic subvariety of \mathbf{A}_g . Then $r \leq 2Nm$ and $g \leq 1 + d(m(N-1) - 1)$.

If $G = (\mathbb{Z}/2\mathbb{Z})^m$ ($g \geq 4$) and we have a family of G -covers of \mathbb{P}^1 yielding a totally geodesic subvariety of \mathbf{A}_g , then $m \leq 6$, $r \leq 6m \leq 36$, and $g \leq 1 + 2^{m-1}(3m - 2) \leq 513$.

If $G = (\mathbb{Z}/p\mathbb{Z})^m$, with p a prime number, $p \geq 3$, $g \geq 4$. Assume that we have a family of G -covers of \mathbb{P}^1 yielding a totally geodesic subvariety of \mathbf{A}_g . Then $m \leq 2p$, $r \leq 4p^2$ and $g \leq 1 + p^{2p}(2p(p-1) - 1)$.

Note that in [27, Theorem 5.2] it is proven, using characteristic p techniques, that if there exist $n \in G$ such that $\{d_n, d_{-n}\} \neq \{0, r-2\}$, where r is the number of branch points of the cover, and $d_n + d_{-n} \geq r-2$, then the subvariety of \mathbf{A}_g induced by the family of abelian covers is not Shimura.

In the second part of the paper we show similar statements for families of Galois covers yielding subvarieties generically contained in the Prym loci of possibly ramified double covers.

Denote by $\mathbf{R}_{g,b}$ the moduli space of isomorphism classes of triples $[(C, \alpha, B)]$ where C is a smooth complex projective curve of genus g , B is a reduced effective divisor of degree b on C and $\alpha \in \text{Pic}(C)$ is such that $\alpha^2 = \mathcal{O}_C(B)$. A point $[(C, \alpha, B)] \in \mathbf{R}_{g,b}$ determines a double cover of C , $f: \tilde{C} \rightarrow C$ branched on B , with $\tilde{C} = \text{Spec}(\mathcal{O}_C \oplus \alpha^{-1})$.

The Prym variety $P(C, \alpha, B)$ (also denoted by $P(\tilde{C}, C)$) associated to $[(C, \alpha, B)]$ is the connected component containing the origin of the kernel of the norm map $\text{Nm}_f: J\tilde{C} \rightarrow JC$. When $b > 0$, $\ker \text{Nm}_f$ is connected. The variety $P(C, \alpha, B)$ is a polarised abelian variety of dimension $g - 1 + \frac{b}{2}$. In fact, if we denote by Ξ the restriction to $P(\tilde{C}, C)$ of the principal polarisation on $J\tilde{C}$, if $b = 0$, Ξ is twice a principal polarisation, hence we consider on $P(\tilde{C}, C)$ this principal polarisation. If $b > 0$, we endow $P(\tilde{C}, C)$ with the polarisation Ξ which is of type $\delta = (1, \dots, 1, \underbrace{2, \dots, 2}_{g \text{ times}})$.

Denote by $\mathbf{A}_{g-1+\frac{b}{2}}^\delta$ the moduli space of abelian varieties of dimension $g - 1 + \frac{b}{2}$ with a polarization of type δ , then we consider the Prym map

$$\mathbf{P}_{g,b}: \mathbf{R}_{g,b} \longrightarrow \mathbf{A}_{g-1+\frac{b}{2}}^\delta, \quad [(C, \alpha, B)] \longmapsto [(P(C, \alpha, B), \Xi)].$$

The dual of the differential of the Prym map $\mathbf{P}_{g,b}$ at a generic point $[(C, \alpha, B)]$ is given by the multiplication map

$$(1.2) \quad (d\mathbf{P}_{g,b})^* : S^2 H^0(C, \omega_C \otimes \alpha) \rightarrow H^0(C, \omega_C^2 \otimes \mathcal{O}_C(B))$$

The multiplication map is surjective if $\dim \mathbf{R}_{g,b} \leq \dim \mathbf{A}_{g-1+\frac{b}{2}}^\delta$, ([24]). So the Prym map $\mathbf{P}_{g,b}$ is generically finite, if and only if $\dim \mathbf{R}_{g,b} \leq \dim \mathbf{A}_{g-1+\frac{b}{2}}^\delta$, that is if either $b \geq 6$ and $g \geq 1$, or $b = 4$ and $g \geq 3$, $b = 2$ and $g \geq 5$, $b = 0$ and $g \geq 6$.

For $b = 0$ the Prym map is generically injective for $g \geq 7$ ([20], [23]). If $b > 0$, in [25], [26], [35], it is proven that $\mathbf{P}_{g,b}$ is generically injective in all the cases except for $b = 4$, $g = 3$, when the degree is 3 ([34], [1]). Moreover, a global Prym-Torelli theorem was recently proved for all g and $b \geq 6$ ([22] for $g = 1$ and [36] for all g).

Assume we have a family of Galois covers $\psi_t : \tilde{C}_t \rightarrow \mathbb{P}^1 = \tilde{C}_t/\tilde{G}$, where \tilde{G} is a group containing a central involution σ . Then we have an exact sequence $0 \rightarrow \langle \sigma \rangle \rightarrow \tilde{G} \rightarrow G \rightarrow 0$, and a commutative diagram

$$(1.3) \quad \begin{array}{ccc} \tilde{C}_t & \xrightarrow{\varphi_t} & C_t = \tilde{C}_t/\langle \sigma \rangle \\ & \searrow \psi_t & \swarrow \pi_t \\ & \mathbb{P}^1 & \end{array}$$

For a general element of the family \tilde{C}_t , denote by $V := H^0(\tilde{C}_t, K_{\tilde{C}_t}) \cong V_+ \oplus V_-$, where V_+ is the set of σ -invariants elements, and V_- is the set of σ -anti-invariants elements. The double cover $\psi_t : \tilde{C}_t \rightarrow C_t = \tilde{C}_t/\langle \sigma \rangle$ corresponds to a triple (C_t, α_t, B_t) , where B_t is a reduced effective divisor of degree b on C_t and α_t is a line bundle on C_t such that $\alpha_t^2 = \mathcal{O}_{C_t}(B_t)$. Then $V_+ \cong H^0(C_t, K_{C_t})$ and $V_- \cong H^0(C_t, K_{C_t} \otimes \alpha_t) \cong H^{1,0}(P(\tilde{C}_t, C_t))$.

For the precise construction of the subvarieties of $\mathbf{A}_{g-1+\frac{b}{2}}^\delta$ generically contained in the Prym loci given by families of Galois covers, see [16, §3]. Given a family of Galois covers as above, observe that multiplication map $m : S^2 V \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})$, which is the dual of the differential of the Torelli map $\tilde{j} : \mathbf{M}_{\tilde{g}} \rightarrow \mathbf{A}_{\tilde{g}}$ at the point $[\tilde{C}_t] \in \mathbf{M}_{\tilde{g}}$, is \tilde{G} -equivariant, hence m maps $(S^2 V)^{\tilde{G}}$ to $H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$. We have the following isomorphism, $(S^2 V)^{\tilde{G}} = (S^2 V_+)^{\tilde{G}} \oplus (S^2 V_-)^{\tilde{G}}$.

The codifferential of the restriction of Prym map to the subvariety of $\mathbf{R}_{g,b}$ given by our family of Galois covers at the point $[(C_t, \alpha_t, B_t)]$ is the restriction of the multiplication map m to $(S^2 V_-)^{\tilde{G}}$, that we still denote by $m : (S^2 V_-)^{\tilde{G}} \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$ (see [15], [16]).

In [15, Theorem 3.2], which is a generalisation of Theorems 3.2 and 4.2 in [9], it is shown that if the map $m : (S^2 V_-)^{\tilde{G}} \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$ is an isomorphism, then the family of Pryms yields a Shimura subvariety of $\mathbf{A}_{g-1+\frac{b}{2}}^\delta$. Using this criterion, in [9], [15], [16] many examples of Shimura subvarieties generically contained in the Prym loci have been constructed.

A natural question is to ask whether the above condition is also necessary for such families of abelian covers to yield Shimura subvarieties of $\mathbf{A}_{g-1+\frac{b}{2}}^\delta$ generically contained in the Prym loci.

We give a partial answer to this question in the case of abelian covers in Theorem 4.2, which is an analogue of Theorem 3.1 in the Prym case. Notice that Theorem 4.2 improves the results obtained in [28] by different techniques.

As a consequence we show the following

Theorem 1.3. (Corollaries 4.4, 4.5, and Remark 4.6)

Assume $\tilde{G} = (\mathbb{Z}/N\mathbb{Z})^m$ with N even, and let σ be an involution in \tilde{G} . Consider a family of abelian covers of \mathbb{P}^1 , $\tilde{C}_t \rightarrow \tilde{C}_t/\tilde{G} \cong \mathbb{P}^1$, with $g(\tilde{C}_t) = \tilde{g} \geq 4$. Assume that the covers $\tilde{C}_t \rightarrow C_t = \tilde{C}_t/\langle \sigma \rangle$ have b ramification points and denote by g the genus of C_t . If the multiplication map $m : (S^2 V_-)^{\tilde{G}} \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$ is surjective at a general element of the family of covers, then there exists an integer $M(N)$ which only depends on N such that, if $\tilde{g} > M(N)$, then the subvariety of the Prym locus in $\mathbf{A}_{g-1+\frac{b}{2}}^\delta$ induced by any such family is not totally geodesic (hence it is not Shimura).

We remark that the condition on the surjectivity of the multiplication map $m : (S^2 V_-)^{\tilde{G}} \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$ at a general element of the family of covers is automatically satisfied if $b \geq 6$ thanks to the Prym-Torelli theorem proved in [36], [22].

The technique used to prove the main Theorems 3.1 and 4.2 is the computation of the second fundamental forms of the Torelli map and of the Prym maps on some particular quadrics invariant under the action of the Galois group. The fact that the Galois group is abelian allows to explicitly construct such quadrics. Then, the techniques developed in [10], [8] for the second fundamental form of the Torelli map and in [5], [6] for the second fundamental form of the Prym map, allow to compute these second fundamental forms on the quadrics that we have constructed and show that they do not vanish. A similar technique in the case of cyclic groups, has been used in [19] to show that the bielliptic and the bihyperelliptic loci are not totally geodesic.

The structure of the paper is as follows. In section 2 we recall the construction of abelian covers of \mathbb{P}^1 following [29], [41]. In section 3 we prove the main results in the case of the Torelli map and we finish giving one example. In section 4 we prove the main results in the case of the Prym maps and we conclude giving some examples.

ACKNOWLEDGEMENTS

I would like to thank Alessandro Ghigi for useful discussions and Roberto Pignatelli for suggestions that improved the estimates given in Corollaries 3.2, 3.4.

2. ABELIAN COVERS OF \mathbb{P}^1

In this section, we recall the construction of abelian covers of \mathbb{P}^1 following [29], [41] (see also [38], [16]).

Consider an $m \times r$ matrix $A = (r_{ij})$ whose entries r_{ij} are in $\mathbb{Z}/N\mathbb{Z}$ for some $N \geq 2$ such that the sum of the columns of A is zero in $(\mathbb{Z}/N\mathbb{Z})^m$ and all the columns are different from zero. Denote by \tilde{r}_{ij} the lift of r_{ij} to $\mathbb{Z} \cap [0, N)$ and take $t_1, \dots, t_r \in \mathbb{C}$, $t_i \neq t_j$, $\forall i \neq j$. Let $\overline{\mathbb{C}(x)}$ be the algebraic closure of $\mathbb{C}(x)$. For each $i = 1, \dots, m$, choose a function $w_i \in \overline{\mathbb{C}(x)}$ with

$$(2.1) \quad w_i^N = \prod_{j=1}^r (x - t_j)^{\tilde{r}_{ij}} \quad \text{for } i = 1, \dots, m,$$

in $\mathbb{C}(x)[w_1, \dots, w_m]$. The equations (2.1) give in general a singular affine curve and we take its normalization. Notice that the cover given by (2.1) is not ramified over the infinity and the local monodromy around the branch point t_j is given by the column vector $(r_{1j}, \dots, r_{mj})^t$. Hence the order of ramification over t_j is $\frac{N}{\gcd(N, \tilde{r}_{1j}, \dots, \tilde{r}_{mj})}$. The covering has Galois group which is given by the subgroup G of $(\mathbb{Z}/N\mathbb{Z})^m$ generated by the columns of A . So by Riemann-Hurwitz formula, we compute the genus of the curve as follows:

$$(2.2) \quad g = 1 + d \left(\frac{r-2}{2} - \frac{1}{2N} \sum_{j=1}^r \gcd(N, \tilde{r}_{1j}, \dots, \tilde{r}_{mj}) \right),$$

where d is the cardinality of G . Denote by $G^* = \text{Hom}(G, \mathbb{C}^*)$ the group of the characters of G . Consider a Galois covering $\pi : C \rightarrow \mathbb{P}^1$ with Galois group G . The group G acts on the sheaves $\pi_*(\mathcal{O}_C)$ and $\pi_*(\mathbb{C})$ via its characters and we get corresponding eigenspace decompositions $\pi_*(\mathcal{O}_C) = \bigoplus_{\chi \in G^*} \pi_*(\mathcal{O}_C)_\chi$ and $\pi_*(\mathbb{C}) = \bigoplus_{\chi \in G^*} \pi_*(\mathbb{C})_\chi$. Set $L_\chi^{-1} = \pi_*(\mathcal{O}_C)_\chi$ and let $\mathbb{C}_\chi = \pi_*(\mathbb{C})_\chi$ denote the eigensheaves corresponding to the character χ . Then L_χ is a line bundle and \mathbb{C}_χ is a local system of rank 1 outside of the branch locus of π .

Denote by l_j be the j -th column of the matrix A . The group G is the subgroup of $(\mathbb{Z}/N\mathbb{Z})^m$ generated by the columns of A , hence we see the column l_j as an element in G . Since G is a finite abelian group, then the character group $G^* = \text{Hom}(G, \mathbb{C}^*)$ is isomorphic to G . In fact, if $G = \mathbb{Z}/N\mathbb{Z}$ is cyclic, then we can fix an isomorphism between $\mathbb{Z}/N\mathbb{Z}$ and the group of N -th roots of unity in \mathbb{C}^* via $1 \mapsto e^{\frac{2\pi i}{N}}$. Therefore G^* is isomorphic to the group of N -th roots of unity via $\chi \mapsto \chi(1)$. In general, the abelian group G is a product of finite cyclic groups, so this isomorphism extends to an isomorphism $G \xrightarrow{\sim} G^*$. Fix a character χ of G , then $\chi(l_j) \in \mathbb{C}^*$ and since G is finite, $\chi(l_j)$ is a root of unity. Hence there exists a unique integer $\alpha_j \in [0, N)$ such that $\chi(l_j) = e^{\frac{2\pi i \alpha_j}{N}}$. Equivalently, we can obtain α_j as follows: let $n \in G \subseteq (\mathbb{Z}/N\mathbb{Z})^m$ be the element corresponding to χ under the above isomorphism. We see n a row vector. Then $(\alpha_1, \dots, \alpha_r) = n \cdot A$.

Denote by \tilde{n} the lift of n to $(\mathbb{Z} \cap [0, N))^m$ and set $\tilde{n} \cdot \tilde{A} = (\tilde{\alpha}_1, \dots, \tilde{\alpha}_r)$, where \tilde{A} is the matrix with entries given by the \tilde{r}_{ij} 's. This means that $\tilde{\alpha}_j = \sum_{i=1}^m n_i \tilde{r}_{ij} \in \mathbb{Z}$ (notice that $\tilde{\alpha}_j$ is not necessarily in $\mathbb{Z} \cap [0, N)$).

Let us denote by K_C the canonical sheaf of C . The sheaf $\pi_*(K_C)_\chi$ decomposes according to the action of G . Let χ be the character associated to an element $n \in G$. Then we have: $L_\chi = \mathcal{O}_{\mathbb{P}^1}(\sum_{j=1}^r \langle \frac{\alpha_j}{N} \rangle)$, where $\langle x \rangle$ is the fractional part of the real number x and

$$(2.3) \quad \pi_*(K_C)_\chi = K_{\mathbb{P}^1} \otimes L_{\chi^{-1}} = \mathcal{O}_{\mathbb{P}^1} \left(-2 + \sum_{j=1}^r \left\langle -\frac{\alpha_j}{N} \right\rangle \right).$$

For a proof of this, see [16, Lemma 4.2], [38, Proposition 1.2]. Consider now an abelian group $G \subseteq (\mathbb{Z}/N\mathbb{Z})^m$ and a G -abelian cover given by the equations (2.1). Let $n \in G$ be the element $(n_1, \dots, n_m) \in (\mathbb{Z}/N\mathbb{Z})^m$ under the inclusion $G \subseteq (\mathbb{Z}/N\mathbb{Z})^m$ with $n_i \in \mathbb{Z} \cap [0, N)$. By (2.3), we have

$$(2.4) \quad d_n := \dim H^0(C, K_C)_n = -1 + \sum_{j=1}^r \left\langle -\frac{\alpha_j}{N} \right\rangle.$$

A basis for $H^0(C, K_C)$ is given by the forms

$$(2.5) \quad \omega_{n,\nu} = x^\nu w_1^{n_1} \cdots w_m^{n_m} \prod_{j=1}^r (x - t_j)^{\lfloor -\frac{\bar{\alpha}_j}{N} \rfloor} dx.$$

Here $\bar{\alpha}_j$ is as above and $0 \leq \nu \leq d_n - 1 = -2 + \sum_{j=1}^r \langle -\frac{\alpha_j}{N} \rangle$. The fact that these elements give a basis of $H^0(C, K_C)$ is proven [29], proof of Lemma 5.1.

To construct a family of abelian covers of \mathbb{P}^1 , we fix a matrix A as above and we let the points (t_1, \dots, t_r) vary in $Y_r := \{(t_1, \dots, t_r) \in (\mathbb{A}_{\mathbb{C}}^1)^r \mid t_i \neq t_j, \forall i \neq j\}$. Over this affine open set we define a family of abelian covers of \mathbb{P}^1 by the equation (2.1). Clearly the branch points are $(t_1, \dots, t_r) \in Y_r$. Varying the branch points we get a family $f : \tilde{C} \rightarrow Y_r$ of smooth projective curves whose fibers C_t are the abelian covers of \mathbb{P}^1 introduced above. Denote by $M_g(G, A) \subset M_g$ the corresponding subvariety in M_g . For more details on the construction of families of Galois covers see e.g. [13], [2], [3], [16].

3. TOTALLY GEODESIC SUBVARIETIES IN THE TORELLI LOCUS

Assume we have a family of Galois covers of \mathbb{P}^1 with Galois group G . With the notation used in [13, Section 2], denote by $M_g(\mathbf{m}, G, \theta)$ the corresponding subvariety of M_g . Here $\theta : \Gamma_r = \langle \gamma_1, \dots, \gamma_r \mid \prod_{i=1}^r \gamma_i = 1 \rangle \rightarrow G$ denotes the monodromy of the cover ($\mathbf{m} := (m_1, \dots, m_r)$ is such that m_i is the order of $\theta(\gamma_i)$ in G). Consider $j : M_g \rightarrow A_g$ the Torelli morphism, which is an orbifold immersion outside the hyperelliptic locus. We endow A_g with the Siegel metric, that is the orbifold metric induced on A_g by the symmetric metric on the Siegel space H_g . The second fundamental form of the Torelli map with respect to the Siegel metric has been studied in [10], [8], [19]. Its dual at a point $[C] \in M_g$ corresponding to a non hyperelliptic curve C is a map

$$(3.1) \quad \rho : I_2(K_C) \rightarrow S^2 H^0(C, K_C^{\otimes 2})$$

where $I_2(K_C)$ is the kernel of the multiplication map $S^2 H^0(C, K_C) \rightarrow H^0(C, K_C^{\otimes 2})$. In [31] in the cyclic case and in [13, Theorem 3.9] for any group G , it is proven that if, for a general curve C of our family of Galois covers, the following condition holds

$$(*) \quad \dim(S^2 H^0(C, K_C))^G = \dim H^0(C, K_C^{\otimes 2})^G,$$

then the family of Galois covers yields a Shimura (hence totally geodesic) subvariety of A_g . We want to show if $(*)$ does not hold, the group G is abelian, and under some assumptions on the dimension of eigenspaces of the action of G on $H^0(K_C)$, the family of abelian covers is not totally geodesic.

So we assume that the Galois group is an abelian group $G \subseteq (\mathbb{Z}/N\mathbb{Z})^m$ of cardinality d , and the family has equations given by (2.1) given by an $m \times r$ matrix $A = (r_{ij})$ whose entries are in $\mathbb{Z}/N\mathbb{Z}$ for some $N \geq 2$. For an element $n \in G$ denote by $V_n := H^0(C, K_C)_n$ and by $d_n = \dim V_n = -1 + \sum_{j=1}^r \langle -\frac{\alpha_j}{N} \rangle$ (see (2.4)). We have the following

- Theorem 3.1.** (1) *If there exists an element $n \in G \subseteq (\mathbb{Z}/N\mathbb{Z})^m$ such that $n \neq -n$, $d_n \geq 2$, $d_{-n} \geq 2$, then the subvariety of A_g induced by the family of abelian covers of genus $g \geq 4$ is not totally geodesic (hence it is not Shimura).*
- (2) *If there exists an element $n \in G \subseteq (\mathbb{Z}/N\mathbb{Z})^m$ of order 2 such that $d_n \geq 3$, then the subvariety of A_g induced by the family of abelian covers of genus $g \geq 4$ is not totally geodesic (hence it is not Shimura).*

Proof. Assume first that the generic curve C of the family of abelian covers is not hyperelliptic. In case (1), by assumption, and by (2.5) there exists subspaces $\langle \omega_1, \omega_2 = x\omega_1 \rangle \subseteq V_n$, $\langle \omega_3, \omega_4 = x\omega_3 \rangle \subseteq V_{-n}$. Then the quadric $Q := \omega_1 \odot \omega_4 - \omega_2 \odot \omega_3$ clearly lies in $(I_2(K_C))^G$. Consider the $d : 1$ cover $\pi : C \rightarrow C/G \cong \mathbb{P}^1$ and take a generic fibre $\pi^{-1}(t) = \{p_1, \dots, p_d\}$ such that $p_i \neq p_j, \forall i \neq j$ and such that $\forall i, p_i$ does not belong to the base locus of the pencil given by $\langle \omega_1, \omega_3 \rangle$, nor to the ramification locus of the map given by the pencil $\langle \omega_1, \omega_3 \rangle$.

Consider the vector $v = \sum_{i=1}^d \xi_{p_i} \in H^1(T_C)^G$, where ξ_{p_i} denotes the Schiffer variation at p_i (for a definition of it see e.g. [8, section 2.2]). Then using [10, Theorem 3.1], or [8, Theorem 2.2] and the fact that ρ is G -equivariant, we have

$$(3.2) \quad \rho(Q)(v \odot v) = \sum_{i \neq j} \rho(Q)(\xi_{p_i} \odot \xi_{p_j}) + \sum_{i=1}^d \rho(Q)(\xi_{p_i} \odot \xi_{p_i}) = -4\pi i \sum_{i \neq j} Q(p_i, p_j) \cdot \eta_{p_i}(p_j) + d\rho(Q)(\xi_{p_1} \odot \xi_{p_1}).$$

For $i \neq j$, we have $Q(p_i, p_j) = 0$, since $x(p_i) = x(p_j) = t$, while $\rho(Q)(\xi_{p_1} \odot \xi_{p_1}) = -2\pi i \cdot \mu_2(Q)(p_1)$, where $\mu_2 : I_2(K_C) \rightarrow H^0(C, K_C^{\otimes 4})$ is the second Gaussian map of the canonical line bundle K_C .

So we have

$$(3.3) \quad \rho(Q)(v \odot v) = k\mu_2(Q)(p_1),$$

where k is a non zero constant. Assume that in a local coordinate around p_1 , $\omega_i = f_i(z)dz$, with $f_2(z) = x(z)f_1(z)$, $f_4(z) = x(z)f_3(z)$. Then we have

$$\begin{aligned} \mu_2(Q) &= (f_1'(z) \cdot f_4'(z) - f_2'(z) \cdot f_3'(z))(dz)^4 = \\ &= (f_1' \cdot (x'(z)f_3(z) + x(z)f_3'(z)) - (x'(z)f_1(z) + x(z)f_1'(z)) \cdot f_3'(z))(dz)^4 = \\ &= x'(z) \cdot (f_1'(z)f_3(z) - f_1(z)f_3'(z))(dz)^4 = x'(z)dz \cdot \mu_1(\omega_1 \wedge \omega_3), \end{aligned}$$

where $\mu_1 : \Lambda^2 H^0(C, K_C) \rightarrow H^0(C, K_C^{\otimes 3})$ denotes the first Gaussian map of the canonical bundle K_C . So $\mu_2(Q)$ vanishes exactly in the base locus of the linear system $\langle \omega_1, \omega_3 \rangle$, in the ramification points of the map given by $\langle \omega_1, \omega_3 \rangle$ and in the ramification points of the cover $\pi : C \rightarrow \mathbb{P}^1$. Hence by our choice of t , we get $\mu_2(Q)(p_1) \neq 0$, so $\rho(Q)(v \odot v) \neq 0$, so the variety given by the family of abelian covers is not totally geodesic.

In case (2) by assumption, there exist a subspace $\langle \omega_1, \omega_2 = x\omega_1, \omega_3 = x^2\omega_1 \rangle \subseteq V_n$, with $n = -n$. Thus we can take the quadric $Q := \omega_1 \odot \omega_3 - \omega_2 \odot \omega_2 \in I_2(K_C)^G$. For a general fibre $\pi^{-1}(t) = \{p_1, \dots, p_d\}$, take $v = \sum_{i=1}^d \xi_{p_i} \in H^1(T_C)^G$ as above, then

$$(3.4) \quad \rho(Q)(v \odot v) = -4\pi i \sum_{i \neq j} Q(p_i, p_j) \cdot \eta_{p_i}(p_j) + d\rho(Q)(\xi_{p_1} \odot \xi_{p_1}) = k\mu_2(Q)(p_1).$$

If we write in local coordinates as above $\omega_i = f_i(z)dz$, $i = 1, 2, 3$, we have $f_2(z) = x(z)f_1(z)$, $f_3(z) = (x(z))^2 f_1(z)$. So we compute

$$\begin{aligned} \mu_2(Q) &= f_1'(z) \cdot (2x(z)x'(z)f_1(z) + (x(z))^2 f_1'(z)) - (x'(z)f_1(z) + x(z)f_1'(z)) \cdot (x'(z)f_1(z) + x(z)f_1'(z)) \\ &= -(x'(z))^2 (f_1(z))^2. \end{aligned}$$

Therefore if we take t generic as above, we have $\mu_2(Q)(p_1) \neq 0$, hence $\rho(Q)(v \odot v) \neq 0$.

It remains to consider the case in which the family of abelian covers is contained in the hyperelliptic locus. The Torelli map restricted to the hyperelliptic locus $\mathbf{HE}_g, j_h : \mathbf{HE}_g \rightarrow \mathbf{A}_g$, is an orbifold immersion ([37]) and in [4, Prop. 5.1], [19, Section 6] the second fundamental form of the restriction of the Torelli map to the hyperelliptic locus has been studied.

We have the following tangent bundle exact sequence

$$(3.5) \quad \begin{array}{ccccccc} & & 0 & & & & \\ & & \downarrow & & & & \\ 0 & \longrightarrow & T_{\text{HE}_g} & \longrightarrow & T_{A_g|_{\text{HE}_g}} & \longrightarrow & N_{\text{HE}_g/A_g} \longrightarrow 0 \\ & & \downarrow & & \downarrow = & & \downarrow \\ & & T_{M_g|_{\text{HE}_g}} & \longrightarrow & T_{A_g|_{\text{HE}_g}} & \longrightarrow & N_{M_g/A_g|_{\text{HE}_g}} \longrightarrow 0 \end{array}$$

Denote by

$$(3.6) \quad \rho_{HE} : N_{\text{HE}_g|_{A_g}}^* \rightarrow S^2 T_{\text{HE}_g}$$

the dual of the second fundamental form of j_h .

At a point $[C] \in \text{HE}_g$, the dual of (3.5) is

$$(3.7) \quad \begin{array}{ccccccc} 0 & \longrightarrow & I_2(K_C) & \longrightarrow & S^2 H^0(K_C) & \xrightarrow{m} & H^0(K_C^{\otimes 2}) \\ & & \downarrow = & & \downarrow = & & \downarrow \\ 0 & \longrightarrow & I_2(K_C) & \longrightarrow & S^2 H^0(K_C) & \xrightarrow{m} & H^0(K_C^{\otimes 2})^+ \longrightarrow 0 \end{array}$$

where $H^0(C, K_C^{\otimes 2})^+$ denotes the subspace of $H^0(C, K_C^{\otimes 2})$ of the element which are invariant under the hyperelliptic involution τ and $I_2(K_C)$ can be identified with the set of quadrics containing the rational normal curve.

We have $\forall Q \in I_2(K_C)$, $\forall v, w \in H^1(T_C)^+$, $\rho_{HE}(Q)(v \odot w) = \rho(Q)(v \odot w)$ (see [4, Prop. 5.1]). Here we denote by $\rho(Q)$ the section $Q \cdot \hat{\eta}$, seen as an element in $S^2 H^0(K_C^{\otimes 2})$ as in [8, Theorem 3.7].

So, if the hyperelliptic involution is contained in the Galois group G of the cover, the tangent vector $v = \sum_{i=1}^d \xi_{p_i}$ is G -invariant, hence τ -invariant, i.e. $v \in H^1(T_C)^+$. Therefore if we take a quadric Q as above (both in case (1), and (2)), we have:

$$\begin{aligned} \rho_{HE}(Q)(v \odot v) &= \rho(Q)(v \odot v) = \sum_{i \neq j} \rho(Q)(\xi_{p_i} \odot \xi_{p_j}) + \sum_{i=1}^d \rho(Q)(\xi_{p_i} \odot \xi_{p_i}) = \\ &= -4\pi i \sum_{i \neq j} Q(p_i, p_j) \cdot \eta_{p_i}(p_j) + d\rho(Q)(\xi_{p_1} \odot \xi_{p_1}) = -2\pi i d \mu_2(Q)(p_1) \neq 0 \end{aligned}$$

for a generic choice of the point $t \in \mathbb{P}^1$.

If the hyperelliptic involution τ is not contained in G , we can consider the subgroup \tilde{G} of $\text{Aut}(C)$ generated by G and τ . Since τ is central in $\text{Aut}(C)$, the group \tilde{G} is abelian and, substituting G with \tilde{G} we can assume that the hyperelliptic involution is contained in the Galois group of the cover. By this we mean that we take the quadrics Q as above, we consider the map $h : C \rightarrow C/\tilde{G} \cong \mathbb{P}^1$, we take a generic point $t \in \mathbb{P}^1$ and the vector $v = \sum_{i=1}^{2d} \xi_{q_i} \in H^1(T_C)^{\tilde{G}} \subset H^1(T_C)^+$, where $h^{-1}(t) = \{q_1, \dots, q_{2d}\}$. Then we have

$$\begin{aligned} \rho_{HE}(Q)(v \odot v) &= \rho(Q)(v \odot v) = \sum_{i \neq j} \rho(Q)(\xi_{q_i} \odot \xi_{q_j}) + \sum_{i=1}^d \rho(Q)(\xi_{q_i} \odot \xi_{q_i}) = \\ &= -4\pi i \sum_{i \neq j} Q(q_i, q_j) \cdot \eta_{q_i}(q_j) + d\rho(Q)(\xi_{q_1} \odot \xi_{q_1}) = -2\pi i d \mu_2(Q)(q_1) \neq 0, \end{aligned}$$

for t general. This concludes the proof. \square

Corollary 3.2. *Under the above assumptions, if $G = (\mathbb{Z}/2\mathbb{Z})^m$ ($g \geq 4$) and the family is totally geodesic, then*

- (1) $r \leq 6m$, and $g \leq 1 + 2^{m-1}(3m - 2)$,
- (2) $m \leq 6$, hence $r \leq 36$, $g \leq 513$.

Proof. By Theorem 3.1 (2), if the family is totally geodesic, for every $n \in G$, with $n \neq 0$, we must have $d_n \leq 2$. Consider the i -th row (r_{i1}, \dots, r_{ir}) of the matrix A that gives the monodromy of the cover $\pi : C \rightarrow C/G \cong \mathbb{P}^1$. Denote by \tilde{r}_{ij} the lift of r_{ij} in $\mathbb{Z} \cap [0, N)$. Set $e_i = (0, \dots, 0, 1, 0, \dots, 0)^t \in G$ and denote by β_i the number of nonzero entries in the row $(r_{i1}, \dots, r_{ir}) = e_i \cdot A$. Then

$$2 \geq d_{e_i} = -1 + \sum_{j=1}^r \left\langle \frac{-\tilde{r}_{ij}}{2} \right\rangle = -1 + \frac{\beta_i}{2},$$

hence $\beta_i \leq 6$, $\forall i = 1, \dots, m$. So if $r > 6m$ there must be a column of A which is zero, a contradiction. Recall that the columns of A give the monodromy of the cover, hence they have order 2. So $r \leq 6m$ and by Riemann Hurwitz formula we have:

$$2g - 2 = -2 \cdot 2^m + r \cdot \frac{2^m}{2},$$

so

$$g = 1 + 2^{m-1} \left(\frac{r}{2} - 2 \right) \leq 1 + 2^{m-1} (3m - 2).$$

It remains to show that $m \leq 6$. Since $G = (\mathbb{Z}/2\mathbb{Z})^m$ is generated by the columns of the matrix A , the set of the columns of A contains a basis of the vector space $(\mathbb{Z}/2\mathbb{Z})^m$. Hence, composing with an automorphism of $(\mathbb{Z}/2\mathbb{Z})^m$, we can assume that the canonical basis e_1, \dots, e_m is a subset of the set of the columns of A . So, consider the functional on G that does not vanish on e_1, \dots, e_m , that is the element $n := (1, \dots, 1)^t$, seen as a character $\chi : G \rightarrow \{\pm 1\} \subset \mathbb{C}^*$. Then $(1, \dots, 1)A = (\alpha_1, \dots, \alpha_r)$ has at least m non zero entries (corresponding to the columns given by e_1, \dots, e_m). So one immediately computes $d_n \geq -1 + \frac{m}{2}$, and by Theorem 3.1 (2), we must have $m \leq 6$. Thus $r \leq 6m \leq 36$ and $g \leq 1 + 2^{m-1} (3m - 2) \leq 513$. □

Corollary 3.3. *Assume $G \subseteq (\mathbb{Z}/N\mathbb{Z})^m$, $N \geq 3$, set $d := \#G$, $g \geq 4$. Assume that we have a family of G -covers of \mathbb{P}^1 yielding a totally geodesic subvariety of A_g . Then $r \leq 2Nm$ and $g \leq 1 + d(m(N-1) - 1)$.*

Proof. Consider as above the element $e_i = (0, \dots, 0, 1, 0, \dots, 0)^t$, then $e_i^t \cdot A = (r_{i1}, \dots, r_{ir})$ is the i^{th} -row of A . By Theorem 3.1, for each row $e_i^t \cdot A$ of the matrix A we must have $d_{e_i} \leq 2$, if $e_i^t \cdot A$ has order 2, otherwise either $d_{e_i} \leq 1$, or $d_{-e_i} \leq 1$.

We claim that in the first case we obtain $\beta_i \leq 6$, while in the second case $\beta_i \leq 2N$.

In fact if $e_i^t \cdot A$ has order 2, then all its nonzero entries are equal to $N/2$, hence $2 \geq d_{e_i} = -1 + \sum_{j=1}^r \left\langle \frac{-\tilde{r}_{ij}}{N} \right\rangle = -1 + \frac{\beta_i}{2}$, so $\beta_i \leq 6$.

If $e_i^t \cdot A$ has order bigger than 2, we have either

$$1 \geq d_{e_i} = -1 + \sum_{j=1}^r \left\langle \frac{-\tilde{r}_{ij}}{N} \right\rangle = -1 + \sum_{j|\tilde{r}_{ij} \neq 0} \left(1 - \frac{\tilde{r}_{ij}}{N} \right) \geq -1 + \beta_i - \beta_i \frac{N-1}{N},$$

(since $\tilde{r}_{ij} \leq N-1$), $\forall i, j$, thus $\beta_i \leq 2N$,

or

$$1 \geq d_{-e_i} = -1 + \sum_{j=1}^r \left\langle \frac{\tilde{r}_{ij}}{N} \right\rangle = -1 + \sum_{j|\tilde{r}_{ij} \neq 0} \frac{\tilde{r}_{ij}}{N} \geq -1 + \frac{\beta_i}{N},$$

thus again we have $\beta_i \leq 2N$.

Denote by p the number of rows of A of order 2, and by q the number of rows of A of order greater than 2, then we must have $r \leq \sum_{i=1}^m \beta_i \leq 6p + 2Nq \leq 2N(p+q) = 2Nm$.

By Riemann Hurwitz formula we have:

$$2g - 2 = d \left(-2 + \sum_{i=1}^r \left(1 - \frac{1}{m_i} \right) \right) \leq -2d + dr \left(1 - \frac{1}{N} \right) \leq -2d + 2dNm \left(1 - \frac{1}{N} \right) = 2d(m(N-1) - 1),$$

since $m_i \leq N$, $\forall i$ and $r \leq 2Nm$. Hence $g \leq 1 + d(m(N-1) - 1)$. □

Corollary 3.4. *Assume $G = (\mathbb{Z}/p\mathbb{Z})^m$, with p a prime number, $p \geq 3$, $g \geq 4$. Assume that we have a family of G -covers of \mathbb{P}^1 yielding a totally geodesic subvariety of A_g . Then $m \leq 2p$, $r \leq 4p^2$ and $g \leq 1 + p^{2p}(2p(p-1) - 1)$.*

Proof. By Corollary 3.3 we know that $r \leq 2pm$ and $g \leq 1 + p^m(m(p-1) - 1)$. Then it suffices to show that $m \leq 2p$.

Since the columns of A generate the vector space $G = (\mathbb{Z}/p\mathbb{Z})^m$, a subset of the set of the columns of A gives a basis of G . Hence, applying an automorphism of G we can assume that the canonical basis e_1, \dots, e_m is a subset of the set of the columns of A . Consider the element $n = (1, \dots, 1)^t \in G$, set $n^t A = (\alpha_1, \dots, \alpha_r)$. Then, by Theorem 3.1 (1), either

$$1 \geq d_n = -1 + \sum_{j=1}^r \left\langle \frac{-\alpha_j}{p} \right\rangle \geq -1 + m\left(1 - \frac{1}{p}\right),$$

or

$$1 \geq d_{-n} = -1 + \sum_{j=1}^r \left\langle \frac{\alpha_j}{p} \right\rangle \geq -1 + \frac{m}{p}.$$

Hence $m \leq 2p$. □

So this concludes the proof of Theorem 1.2.

Clearly the above estimates are not sharp, as one can see by the following example.

Example. $r = 8, g = 13$. $G = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} = \langle g_1 \rangle \times \langle g_2 \rangle \subseteq \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$, $g_1 \mapsto (2, 0)^t$, $g_2 \mapsto (0, 1)^t$. The family of covers is given by the matrix

$$A = \begin{pmatrix} 2 & 2 & 2 & 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix},$$

so the equations can be written as:

$$w_1^2 = \prod_{i=1}^4 (x - t_i)$$

$$w_2^4 = \prod_{i=5}^8 (x - t_i).$$

By Riemann Hurwitz one immediately computes $g = 13$. The non zero elements of G , with the identification of G with its group of characters explained in section 2 are:

$$g_1 = (10)A, g_2 = (01)A, g_1 + g_2 = (11)A, g_1 + 2g_2 = (12)A, g_1 + 3g_2 = (13)A, 2g_2 = (02)A, 3g_2 = (03)A.$$

So we have:

$$d_{g_1} = 1, d_{g_2} = 2, d_{g_1+g_2} = 4, d_{g_1+2g_2} = 3, d_{g_1+3g_2} = 2, d_{2g_2} = 1, d_{3g_2} = 0.$$

Since $d_{g_1+g_2} = 4$, $-(g_1 + g_2) = g_1 + 3g_2$ and $d_{g_1+3g_2} = 2$, we can apply Theorem 3.1(1) and conclude that this family gives a subvariety of A_{13} contained in the Torelli locus which is not totally geodesic. We could also apply Theorem 3.1(2) to conclude, since $d_{g_1+2g_2} = 3$ and $d_{g_1+2g_2}$ has order 2.

4. TOTALLY GEODESIC SUBVARIETIES IN THE PRYM LOCI

Denote by $\mathbf{R}_{g,b}$ the moduli space of isomorphism classes of triples $[(C, \alpha, B)]$ where C is a smooth complex projective curve of genus g , B is a reduced effective divisor of degree b on C and $\alpha \in \text{Pic}(C)$ is such that $\alpha^{\otimes 2} = \mathcal{O}_C(B)$. A point $[(C, \alpha, B)] \in \mathbf{R}_{g,b}$ determines a double cover of C , $f : \tilde{C} \rightarrow C$ branched on B , with $\tilde{C} = \text{Spec}(\mathcal{O}_C \oplus \alpha^{-1})$.

The Prym variety $P(C, \alpha, B)$ (also denoted by $P(\tilde{C}, C)$) associated to $[(C, \alpha, B)]$ is the connected component containing the origin of the kernel of the norm map $\text{Nm}_f : J\tilde{C} \rightarrow JC$. When $b > 0$, $\ker \text{Nm}_f$ is connected. The variety $P(C, \alpha, B)$ is a polarised abelian variety of dimension $g - 1 + \frac{b}{2}$. In fact, if we denote by Ξ the restriction to $P(\tilde{C}, C)$ of the principal polarisation on $J\tilde{C}$, if $b = 0$, Ξ is twice a principal polarisation, hence we consider on $P(\tilde{C}, C)$ this principal polarisation. If $b > 0$, we endow $P(\tilde{C}, C)$ with the polarisation Ξ which is of type $\delta = (1, \dots, 1, \underbrace{2, \dots, 2}_{g \text{ times}})$.

If we denote by $\mathbf{A}_{g-1+\frac{b}{2}}^\delta$ the moduli space of abelian varieties of dimension $g - 1 + \frac{b}{2}$ with a polarization of type δ , then we consider the Prym map

$$\mathbf{P}_{g,b} : \mathbf{R}_{g,b} \longrightarrow \mathbf{A}_{g-1+\frac{b}{2}}^\delta, \quad [(C, \alpha, B)] \longmapsto [(P(C, \alpha, B), \Xi)].$$

The dual of the differential of the Prym map $P_{g,b}$ at a generic point $[(C, \alpha, B)]$ is given by the multiplication map

$$(4.1) \quad (dP_{g,b})^* : S^2 H^0(C, \omega_C \otimes \alpha) \rightarrow H^0(C, \omega_C^2 \otimes \mathcal{O}_C(B))$$

The multiplication map is surjective if $\dim R_{g,b} \leq \dim A_{g-1+\frac{b}{2}}^\delta$, ([24]). Assume we have a family of Galois covers $\psi_t : \tilde{C}_t \rightarrow \mathbb{P}^1 = \tilde{C}_t/\tilde{G}$, where \tilde{G} is a finite group containing a central involution σ . Then we have an exact sequence $0 \rightarrow \langle \sigma \rangle \rightarrow \tilde{G} \rightarrow G \rightarrow 0$, and a commutative diagram

$$(4.2) \quad \begin{array}{ccc} \tilde{C}_t & \xrightarrow{\varphi_t} & C_t = \tilde{C}_t/\langle \sigma \rangle \\ & \searrow \psi_t & \swarrow \pi_t \\ & \mathbb{P}^1 & \end{array}$$

For a general element \tilde{C} of the family, denote by $V := H^0(\tilde{C}, K_{\tilde{C}}) \cong V_+ \oplus V_-$, where V_+ is the set of σ -invariant elements, and V_- is the set of σ -anti-invariant elements. The double cover $\varphi : \tilde{C} \rightarrow C = \tilde{C}/\langle \sigma \rangle$ corresponds to a triple (C, α, B) , where B is a reduced effective divisor of degree b on C and α is a line bundle on C such that $\alpha^{\otimes 2} = \mathcal{O}_C(B)$. Then $V_+ \cong H^0(C, K_C)$ and $V_- \cong H^0(C, K_C \otimes \alpha) \cong H^{1,0}(P(\tilde{C}, C))$.

For the precise construction of the subvarieties of $A_{g-1+\frac{b}{2}}^\delta$ generically contained in the Prym loci given by families of Galois covers, see [16, §3]. Given a family of Galois covers as above, consider the multiplication map $m : S^2 V \rightarrow W = H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})$, which is the dual of the differential of the Torelli map $\tilde{j} : M_{\tilde{g}} \rightarrow A_{\tilde{g}}$ at the point $[\tilde{C}] \in M_{\tilde{g}}$. The map m is \tilde{G} -equivariant, hence it maps $(S^2 V)^{\tilde{G}}$ to $H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$. We have the following isomorphism: $(S^2 V)^{\tilde{G}} = (S^2 V_+)^{\tilde{G}} \oplus (S^2 V_-)^{\tilde{G}}$.

The dual of the differential of the restriction of Prym map to the subvariety of $R_{g,b}$ given by our family of Galois covers at the point $[(C, \alpha, B)]$ is the restriction of the multiplication map m to $(S^2 V_-)^{\tilde{G}}$. We still denote by

$$(4.3) \quad m : (S^2 V_-)^{\tilde{G}} \longrightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$$

this restriction (see [15], [16]).

In [15, Theorem 3.2], which is a generalisation of Theorems 3.2 and 4.2 in [9], it is shown that if the map $m : (S^2 V_-)^{\tilde{G}} \longrightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$ is an isomorphism, then the family of Pryms yields a Shimura subvariety of $A_{g-1+\frac{b}{2}}^\delta$. Using this criterion, in [9], [15], [16] many examples of Shimura subvarieties generically contained in the Prym loci have been constructed.

We prove that in the case of abelian covers, under some assumptions, the above condition is also necessary for such families of abelian covers to yield a Shimura subvariety of $A_{g-1+\frac{b}{2}}^\delta$ generically contained in the Prym loci.

So, consider now an abelian group $\tilde{G} \subseteq (\mathbb{Z}/N\mathbb{Z})^m$ containing an involution σ as above and a \tilde{G} -abelian cover given by the equations (2.1). Let $n \in \tilde{G}$ be the element $(n_1, \dots, n_m) \in (\mathbb{Z}/N\mathbb{Z})^m$ under the inclusion $\tilde{G} \subseteq (\mathbb{Z}/N\mathbb{Z})^m$ with $n_i \in \mathbb{Z} \cap [0, N)$. By (2.4), $\dim H^0(\tilde{C}, K_{\tilde{C}})_n = -1 + \sum_{j=1}^r \langle -\frac{\alpha_j}{N} \rangle$. A basis for the \mathbb{C} -vector space $H^0(\tilde{C}, K_{\tilde{C}})$ is given by the forms in (2.5): $\omega_{n,\nu} = x^\nu w_1^{n_1} \cdots w_m^{n_m} \prod_{j=1}^r (x-t_j)^{\lfloor -\frac{\alpha_j}{N} \rfloor} dx$, where $\tilde{\alpha}_j$ is as introduced above and $0 \leq \nu \leq d_n - 1 = -2 + \sum_{j=1}^r \langle -\frac{\alpha_j}{N} \rangle$.

The action of σ is given by $w_i \mapsto -w_i$ for some subset of $\{1, \dots, m\}$, and $w_j \mapsto w_j$ for j in the complement of this subset. We may then, without loss of generality, assume that $\sigma(w_i) = -w_i$ for $i \in \{1, \dots, k\}$ for some $k \leq m$ and $\sigma(w_i) = w_i$ for the $i > k$. We recall now [28, Lemma 2.5].

Lemma 4.1. *The group \tilde{G} acts on $H^0(\tilde{C}, K_{\tilde{C}})_-$ and for $n \in \tilde{G}$, $n = (n_1, \dots, n_m)^t$, we have $H^0(\tilde{C}, K_{\tilde{C}})_{-,n} = H^0(\tilde{C}, K_{\tilde{C}})_n$ if $n_1 + \dots + n_k$ is odd and $H^0(\tilde{C}, K_{\tilde{C}})_{-,n} = 0$ otherwise. Similar equalities hold for $H^1(\tilde{C}, \mathbb{C})_{-,n}$.*

Under these hypotheses, we have the following

Theorem 4.2. *Assume that the multiplication map $m : (S^2 V_-)^{\tilde{G}} \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$ is surjective at the generic point of the family, then*

- (1) *If there exists $n \in \tilde{G}$ of order greater than 2, $\dim(V_-)_n = d_n \geq 2$ and $\dim(V_-)_{(-n)} = d_{-n} \geq 2$, then the family of Pryms gives a subvariety of $A_{g-1+\frac{b}{2}}^\delta$ which is not totally geodesic.*

(2) If there exists $n \in \tilde{G}$ such that $n = -n$ and $\dim(V_-)_n = d_n \geq 3$, then the family of Pryms gives a subvariety of $\mathbf{A}_{g-1+\frac{1}{2}}^\delta$ which is not totally geodesic.

Proof. Assume that $\tilde{G} \subseteq (\mathbb{Z}/N\mathbb{Z})^m$ and the family of \tilde{G} -covers is given by a matrix $A = (r_{ij})$ whose entries are in $\mathbb{Z}/N\mathbb{Z}$ for some $N \geq 2$. Then the equations for the cover \tilde{G} are

$$(4.4) \quad w_i^N = \prod_{j=1}^r (x - t_j)^{\tilde{r}_{ij}} \quad \text{for } i = 1, \dots, m,$$

and we can assume that the involution σ is given by $\sigma(w_i) = -w_i$, $\forall i = 1, \dots, k$, $\sigma(w_i) = w_i$, $\forall i \geq k+1$. Then, denoting as above by $V = H^0(\tilde{C}, K_{\tilde{C}})$ and by $V_- = H^0(\tilde{C}, K_{\tilde{C}}^-) \cong H^0(C, K_C \otimes \alpha) \cong H^{1,0}(P(\tilde{C}, C))$, by Lemma 4.1, we have $(V_-)_n = 0$, if $n_1 + \dots + n_k$ is even, while $(V_-)_n = V_n$ if $n_1 + \dots + n_k$ is odd.

Hence if we are in case (1), there exist subspaces $\langle \omega_1, \omega_2 = x\omega_1 \rangle \subseteq (V_-)_n =$, and $\langle \omega_3, \omega_4 = x\omega_3 \rangle \subseteq (V_-)_{(-n)}$. So the quadric $Q := \omega_1 \odot \omega_4 - \omega_2 \odot \omega_3$ is \tilde{G} -invariant and it belongs to the kernel $I_2(K_C \otimes \alpha)^{\tilde{G}}$ of the multiplication map $m : (S^2 V_-)^{\tilde{G}} \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$. The assumption on the surjectivity of the map m is equivalent to say that the differential of the restriction of the Prym map to the subvariety \mathbf{X} of $\mathbf{R}_{g,b}$ given by our family of abelian covers is injective. So it is possible to study the second fundamental form of the immersion of \mathbf{X} in $\mathbf{A}_{g-1+\frac{1}{2}}^\delta$ given by the restriction of the Prym map to \mathbf{X} .

In [6, section 2] it is proven that if

$$\rho_P : I_2(K_C \otimes \alpha)^{\tilde{G}} \rightarrow (S^2 H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2}))^{\tilde{G}}$$

is the dual of the second fundamental form of the restriction of the Prym map to \mathbf{X} we have $\rho_P(Q)(v \odot v) = \tilde{\rho}(Q)(v \odot v)$, $\forall v \in H^1(T_{\tilde{C}})^{\tilde{G}}$, where $\tilde{\rho}$ is the dual of the second fundamental form of the Torelli map of the family of covers $\psi : \tilde{C} \rightarrow \tilde{C}/\tilde{G} = \mathbb{P}^1$. As in the proof of Theorem 3.1, if the family of covers $\tilde{C} \rightarrow \tilde{C}/\tilde{G}$ is contained in the hyperelliptic locus then $\tilde{\rho}(Q)$ is the section $\tilde{Q} \cdot \tilde{\eta}$, seen as an element in $S^2 H^0(K_{\tilde{C}}^{\otimes 2})$ as in [8, Theorem 3.7].

Let $t \in \mathbb{P}^1$ be a general point, then $\psi^{-1}(t) = \{p_1, \dots, p_d\}$, where d is the order of \tilde{G} and $p_i \neq p_j$, $\forall i \neq j$. Set $v := \sum_{i=1}^d \xi_{p_i}$. Then clearly $v \in H^1(T_{\tilde{C}})^{\tilde{G}}$ and we have:

$$\begin{aligned} \rho_P(Q)(v \odot v) &= \tilde{\rho}(Q)(v \odot v) = \sum_{i \neq j} \tilde{\rho}(Q)(\xi_{p_i} \odot \xi_{p_j}) + \sum_{i=1}^d \tilde{\rho}(Q)(\xi_{p_i} \odot \xi_{p_i}) \\ &= -4\pi i \sum_{i \neq j} Q(p_i, p_j) \tilde{\eta}_{p_i}(p_j) + d \tilde{\rho}(Q)(\xi_{p_1} \odot \xi_{p_1}) = -2\pi i d \mu_2(Q)(p_1), \end{aligned}$$

where $\mu_2 : I_2(K_{\tilde{C}}) \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 4})$ is the second Gaussian map of the canonical line bundle $K_{\tilde{C}}$. Here we used again the \tilde{G} -equivariance of $\tilde{\rho}$ and the fact that $Q(p_i, p_j) = 0$, for all i, j , since $x(p_i) = x(p_j)$. Hence if we take $t \in \mathbb{P}^1$ generic as in the proof of Theorem 3.1 we have $\mu_2(Q)(p_1) \neq 0$. This concludes the proof of case (1).

Case (2) is very similar: by assumption there exist a subspace $\langle \omega_1, \omega_2 = x\omega_1, \omega_3 = x^2\omega_1 \rangle \subseteq (V_-)_n = V_n$, with $n = -n$. Thus we can take the quadric $Q := \omega_1 \odot \omega_3 - \omega_2 \odot \omega_2 \in I_2(K_C \otimes \alpha)^{\tilde{G}}$. For a general fibre $\psi^{-1}(q) = \{p_1, \dots, p_d\}$, we have again

$$(4.5) \quad \rho_P(Q)(v \odot v) = -4\pi i \sum_{i \neq j} Q(p_i, p_j) \cdot \eta_{p_i}(p_j) + d \rho(Q)(\xi_{p_1} \odot \xi_{p_1}) = -2\pi i d \mu_2(Q)(p_1) \neq 0,$$

for t generic. □

Remark 4.3. *The assumption on the surjectivity of the multiplication map $m : (S^2 V_-)^{\tilde{G}} \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$ at the generic point of the family is automatically satisfied if $b \geq 6$, thanks to the Prym-Torelli theorem proved in [22], [36].*

Corollary 4.4. *With the above notation, assume that the multiplication map $m : (S^2 V_-)^{\tilde{G}} \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$ is surjective, $\tilde{G} = (\mathbb{Z}/2\mathbb{Z})^m$, and $\sigma(w_i) = -w_i$, for $i = 1, \dots, m$. If the family of Pryms yields a totally geodesic subvariety, then $r \leq 6m$ and $\tilde{g} \leq 1 + 2^{m-1}(3m - 2)$.*

Proof. Since $\sigma(w_i) = -w_i$, $\forall i = 1, \dots, m$, setting as usual $e_i = (0, \dots, 0, 1, 0, \dots, 0)^t \in \tilde{G}$, by Lemma 4.1 we have $V_{e_i} = (V_-)_{e_i}$, for all $i = 1, \dots, m$. Hence by Theorem 4.2 (2), if the family is totally geodesic, we must have $d_{e_i} \leq 2$, $\forall i = 1, \dots, m$. So we conclude as in Corollary 3.2. \square

Corollary 4.5. *Assume that the multiplication map $m : (S^2V_-)^{\tilde{G}} \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}}$ is surjective, $\tilde{G} \subseteq (\mathbb{Z}/N\mathbb{Z})^m$, $N \geq 3$, $\sigma(w_i) = -w_i$, for $i = 1, \dots, m$. Set $d := \#\tilde{G}$. If the family of Pryms yields a totally geodesic subvariety, then $r \leq 2Nm$ and $\tilde{g} \leq 1 + d(-1 + m(N-1))$.*

Proof. Since $\sigma(w_i) = -w_i$, $\forall i = 1, \dots, m$, by Lemma 4.1 we have $V_{e_i} = (V_-)_{e_i}$, for all $i = 1, \dots, m$. Hence by Theorem 4.2, for each row $e_i^t \cdot A$, $i = 1, \dots, m$, we must have $d_{e_i} \leq 2$, if $e_i \cdot A$ has order 2, otherwise either $d_{e_i} \leq 1$, or $d_{-e_i} \leq 1$.

As in Corollary 3.3, we show that in the first case we obtain $\beta_i \leq 6$, while in the second case $\beta_i \leq 2N$, $i = 1, \dots, m$ and we conclude as in Corollary 3.3. \square

Remark 4.6. *Notice that in the case $\tilde{G} = (\mathbb{Z}/N\mathbb{Z})^m$ we can always assume that $\sigma(w_i) = -w_i$, for $i = 1, \dots, m$, that is, $\sigma = (\frac{N}{2}, \dots, \frac{N}{2})^t$, since, given any two elements σ_1, σ_2 of order 2, there always exists an automorphism of \tilde{G} sending σ_1 to σ_2 .*

Corollaries 4.4, 4.5 and Remark 4.6 prove Theorem 1.3.

We give now some examples where the assumptions of Theorem 4.2 are satisfied.

Clearly the above estimates are not sharp, as one can see by example 2.

Example 1 $r = 8$, $b = 8$, $\tilde{g} = 5$, $g = 1$. $\tilde{G} = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. The family of covers is given by the matrix

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix},$$

so the equations are:

$$\begin{aligned} w_1^2 &= (x-t_1)(x-t_2)(x-t_3)(x-t_4) \\ w_2^2 &= (x-t_1)(x-t_2)(x-t_3)(x-t_4)(x-t_5)(x-t_6)(x-t_7)(x-t_8). \end{aligned}$$

By Riemann Hurwitz formula, one immediately computes $\tilde{g} = 5$. The involution $\sigma = (1, 1)^t$ acts as follows: $\sigma(w_1) = -w_1$, $\sigma(w_2) = -w_2$. The map $\tilde{C} \rightarrow C = \tilde{C}/\langle \sigma \rangle$ ramifies over t_i , $i = 1, \dots, 4$, hence it has 8 ramification points, so by Riemann Hurwitz we see that $g = g(C) = 1$.

We compute:

$d_{(1,0)} = -1 + \frac{4}{2} = 1$, and $(V_-)_{(1,0)} = \langle \alpha_1 = w_1 \frac{dx}{(x-t_1)(x-t_2)(x-t_3)(x-t_4)} = \frac{dx}{w_1} \rangle$,
 $d_{(0,1)} = -1 + \frac{8}{2} = 3$, and $(V_-)_{(0,1)} = \langle \alpha_2 = w_2 \frac{dx}{\prod_{i=1}^8 (x-t_i)} = \frac{dx}{w_2}, \alpha_3 = x \frac{dx}{w_2}, \alpha_4 = x^2 \frac{dx}{w_2} \rangle$. Hence $V_- = \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4 \rangle$, and

$$(S^2V_-)^{\tilde{G}} = \langle \alpha_1 \odot \alpha_1, \alpha_2 \odot \alpha_2, \alpha_2 \odot \alpha_3, \alpha_2 \odot \alpha_4, \alpha_3 \odot \alpha_3, \alpha_3 \odot \alpha_4, \alpha_4 \odot \alpha_4 \rangle \cong \mathbb{C}^7.$$

Since the differential of the Prym map $P_{1,8} : R_{1,8} \rightarrow A_4^\delta$ is injective ([36]), the multiplication map

$$m : (S^2V_-)^{\tilde{G}} \cong \mathbb{C}^7 \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}} \cong \mathbb{C}^5$$

is surjective, and $\dim(V_-)_{(0,1)} = 3$, so we can apply Theorem 4.2 (2) and conclude that the family of Pryms is not totally geodesic.

One can also explicitly compute the kernel of the multiplication map as follows: We have $\alpha_1^2 = \frac{dx^2}{(x-t_1)(x-t_2)(x-t_3)(x-t_4)}$, $\alpha_2^2 = \frac{dx^2}{\prod_{i=1}^8 (x-t_i)}$, $\alpha_2\alpha_3 = x\alpha_2^2$, $\alpha_2\alpha_4 = x^2\alpha_2^2$, $\alpha_3^2 = x^2\alpha_2^2$, $\alpha_3\alpha_4 = x^3\alpha_2^2$, $\alpha_4^2 = x^4\alpha_2^2$. We can assume that $t_5 = 0$, $t_6 = 1$, $t_7 = -1$, so one easily computes that

$$a_1\alpha_1^2 + a_2\alpha_2^2 + a_3\alpha_2\alpha_3 + a_4\alpha_2\alpha_4 + a_5\alpha_3^2 + a_6\alpha_3\alpha_4 + a_7\alpha_4^2 = 0,$$

if and only if $a_7 = -a_1$, $a_6 = t_8 a_1$, $a_5 = -a_4 + a_1$, $a_3 = -t_8 a_1$, $a_2 = 0$. Hence the kernel of the multiplication map $m : (S^2V_-)^{\tilde{G}} \cong \mathbb{C}^7 \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}} \cong \mathbb{C}^5$ has dimension 2 and it is generated by the quadrics $Q_1 = \alpha_2 \odot \alpha_4 - \alpha_3 \odot \alpha_3$, $Q_2 = \alpha_1 \odot \alpha_1 - t_8 \alpha_2 \odot \alpha_3 + \alpha_3 \odot \alpha_3 + t_8 \alpha_3 \odot \alpha_4 - \alpha_4 \odot \alpha_4$. Since

$\dim(S^2V_-)^{\tilde{G}} = 7$ and $\dim H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}} = 8 - 3 = 5$, we conclude that m is surjective.

Example 2 $r = 8, b = 0, \tilde{g} = 33, g = 17$. $\tilde{G} = \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$. The family of covers is given by the matrix

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix},$$

so the equations are:

$$\begin{aligned} w_1^4 &= (x - t_1)(x - t_2)(x - t_3)(x - t_4) \\ w_2^4 &= (x - t_5)(x - t_6)(x - t_7)(x - t_8). \end{aligned}$$

By Riemann Hurwitz formula one immediately computes $\tilde{g} = 33$. The involution $\sigma = (1, 1)^t$ acts as follows: $\sigma(w_1) = -w_1, \sigma(w_2) = -w_2$. The map $\tilde{C} \rightarrow C = \tilde{C}/\langle \sigma \rangle$ is étale so $g = 17$.

We compute:

$$d_{(1,0)} = -1 + 4 \cdot \frac{3}{4} = 2 = d_{(0,1)},$$

$$d_{(3,0)} = -1 + 4 \cdot \frac{1}{4} = 0 = d_{(0,3)},$$

$$d_{(1,2)} = -1 + 4 \cdot \frac{3}{4} + 4 \cdot \frac{1}{2} = 4 = d_{(2,1)}, \quad d_{(3,2)} = -1 + 4 \cdot \frac{1}{4} + 4 \cdot \frac{1}{2} = 2 = d_{(2,3)}.$$

So $(S^2V_-)^{\tilde{G}} = ((V_-)_{(1,2)} \otimes (V_-)_{(3,2)}) \oplus ((V_-)_{(2,1)} \otimes (V_-)_{(2,3)})$ and since $d_{(1,2)} = 4$ and $d_{(3,2)} = 2$, Theorem 4.2 (1) applies, provided that the multiplication map $m : (S^2V_-)^{\tilde{G}} \cong \mathbb{C}^{16} \rightarrow H^0(\tilde{C}, K_{\tilde{C}}^{\otimes 2})^{\tilde{G}} \cong \mathbb{C}^5$ is surjective.

We show that the restriction of m to $((V_-)_{(1,2)} \otimes (V_-)_{(3,2)})$ has rank 5, hence m is surjective. A basis of $(V_-)_{(1,2)}$ is $\{\beta_1 = \frac{w_1 w_2^2 dx}{\prod_{i=1}^8 (x-t_i)}, \beta_2 = x\beta_1, \beta_3 = x^2\beta_1, \beta_4 = x^3\beta_1\}$. A basis of $(V_-)_{(3,2)}$ is $\{\gamma_1 = \frac{w_1^3 w_2^2 dx}{\prod_{i=1}^8 (x-t_i)}, \gamma_2 = x\gamma_1\}$. So a basis of $((V_-)_{(1,2)} \otimes (V_-)_{(3,2)})$ is given by $\{\beta_i \odot \gamma_j \mid i = 1, \dots, 4, j = 1, 2\}$.

$$\sum_{i,j} a_{ij} \beta_i \gamma_j = 0$$

if and only if

$$\beta_1 \gamma_1 (a_{11} + x a_{21} + x^2 a_{31} + x^3 a_{41} + x a_{12} + x^2 a_{22} + x^3 a_{32} + x^4 a_{42}) = 0.$$

So we have $a_{11} = a_{42} = 0, a_{21} = -a_{12}, a_{31} = -a_{22}, a_{41} = -a_{32}$. So The kernel of the restriction of m to $((V_-)_{(1,2)} \otimes (V_-)_{(3,2)})$ has dimension 3, hence m is surjective and the family of Pryms is not totally geodesic.

REFERENCES

- [1] Bardelli, F., Ciliberto, C., Verra, A., Curves of minimal genus on a general abelian variety, *Compos. Math.* 96 (1995), no. 2, 115–147.
- [2] F. Catanese, M. Lönne, and F. Perroni. Irreducibility of the space of dihedral covers of the projective line of a given numerical type. *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl.*, 22(3):291–309, 2011.
- [3] F. Catanese, M. Lönne, and F. Perroni. Genus stabilization for the components of moduli spaces of curves with symmetries. *Algebr. Geom.*, 3(1):23–49, 2016.
- [4] E. Colombo and P. Frediani. Siegel metric and curvature of the moduli space of curves. *Trans. Amer. Math. Soc.*, 362(3):1231–1246, 2010.
- [5] Colombo, E., Frediani P., A bound on the dimension of a totally geodesic submanifold in the Prym locus. *Collectanea Mathematica*, 70, n.1, (2019), 51-57. DOI: 10.1007/s13348-018-0215-0.
- [6] E. Colombo and P. Frediani. Second fundamental form of the Prym map in the ramified case. "Galois Covers, Grothendieck-Teichmueller Theory and Dessins d'Enfants - Interactions between Geometry, Topology, Number Theory and Algebra", *Springer Proceedings in Mathematics & Statistics*, Vol.330, (2020).
- [7] E. Colombo and P. Frediani. On the dimension of totally geodesic submanifolds in the Prym loci. arXiv:2101.05189. To appear in *Bollettino U. M. I.* DOI: 10.1007/s40574-021-00287-4.
- [8] E. Colombo, P. Frediani, and A. Ghigi. On totally geodesic submanifolds in the Jacobian locus. *International Journal of Mathematics*, 26 (2015), no. 1, 1550005 (21 pages).
- [9] Colombo, E., Frediani, P., Ghigi, A., Penegini M., Shimura curves in the Prym locus. *Communications in Contemporary Mathematics*, Vol. 21, N0. 2 (2019) 1850009 (34 pages). DOI: 10.1142/S0219199718500098.
- [10] E. Colombo, G. P. Pirola, and A. Tortora. Hodge-Gaussian maps. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 30(1):125–146, 2001.
- [11] D. Conti, A. Ghigi, R. Pignatelli. Some evidence for the Coleman-Oort conjecture. arXiv:math/2102.12349.
- [12] J. de Jong and R. Noot. Jacobians with complex multiplication. In *Arithmetic algebraic geometry (Texel, 1989)*, volume 89 of *Progr. Math.*, pages 177–192. Birkhäuser Boston, Boston, MA, 1991.

- [13] P. Frediani, A. Ghigi and M. Penegini. Shimura varieties in the Torelli locus via Galois coverings. *Int. Math. Res. Not.* 2015, no. 20, 10595-10623.
- [14] P. Frediani, A. Ghigi and I. Spelta. Infinitely many Shimura varieties in the Jacobian locus for $g \leq 4$. arXiv:1910.13245. To appear in *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*
- [15] P. Frediani, G.P. Grosselli. Shimura curves in the Prym loci of ramified double covers. arXiv:2007.09646.
- [16] P. Frediani, G.P. Grosselli, A. Mohajer Higher dimensional Shimura varieties in the Prym loci of ramified double covers. arXiv:2101.09016.
- [17] P. Frediani, G.P. Pirola, On the geometry of the second fundamental form of the Torelli map. *Proceedings of the AMS.* Vol. 149, N. 3, 2021, 1011-1024.
- [18] P. Frediani, M. Penegini, P. Porru Shimura varieties in the Torelli locus via Galois coverings of elliptic curves. *Geometriae Dedicata* 181 (2016) 177-192.
- [19] P. Frediani, P. Porru, On the Bielliptic and Bihyperelliptic Loci, *Michigan Math. J.*, 69, (2020), 571-600.
- [20] Friedman, Robert; Smith, Roy, The generic Torelli theorem for the Prym map. *Invent. Math.* 67 (1982), no. 3, 473-490.
- [21] S. Grushevsky and M. Möller, Explicit formulas for infinitely many Shimura curves in genus 4. *Asian J. Math.* 22 (2018), no. 2, 381-390.
- [22] A. Ikeda, Global Prym-Torelli Theorem for double coverings of elliptic curves, *Algebr. Geom.* 7 (2020), no. 5, 544-560.
- [23] V. I. Kanev. A global Torelli theorem for Prym varieties at a general point. *Izv. Akad. Nauk SSSR Ser. Mat.*, 46(2):244-268, 431, 1982.
- [24] H. Lange and A. Ortega. Prym varieties of cyclic coverings. *Geom. Dedicata*, 150:391-403, 2011.
- [25] V.O. Marcucci, G.P. Pirola. Generic Torelli theorem for Prym varieties of ramified coverings. *Compos. Math.* 148 (2012), no. 4, 1147-1170.
- [26] Marcucci, V, Naranjo, J.C., Prym varieties of double coverings of elliptic curves, *Int. Math. Res. Notices* 6 (2014), 1689-1698.
- [27] Mohajer, A, Shimura varieties and abelian covers of the line. arXiv:2001.08536v1. To appear in *Houston Journal of Math.*
- [28] Mohajer, A, On Shimura subvarieties of the Prym locus. arXiv:1804.10131v2. To appear in *Communications in Algebra*. DOI: 10.1080/00927872.2021.1878205.
- [29] A. Mohajer, K. Zuo, *On Shimura subvarieties generated by families of abelian covers of \mathbb{P}^1* . *Journal of Pure and Applied Algebra*, 222 (4), 2018, 931-949.
- [30] B. Moonen. Linearity properties of Shimura varieties. I. *J. Algebraic Geom.*, 7(3):539-567, 1998.
- [31] B. Moonen. Special subvarieties arising from families of cyclic covers of the projective line. *Doc. Math.*, 15:793-819, 2010.
- [32] B. Moonen and F. Oort. The Torelli locus and special subvarieties. In *Handbook of Moduli: Volume II*, pages 549-94. International Press, Boston, MA, 2013.
- [33] D. Mumford. A note of Shimura's paper "Discontinuous groups and abelian varieties". *Math. Ann.*, 181:345-351, 1969.
- [34] D. S., Nagaraj; S. Ramanan, S. Polarizations of type $(1,2,\dots,2)$ on abelian varieties. *Duke Math. J.* 80 (1995), no. 1, 157-194.
- [35] Naranjo, J.C., Ortega, A., Verra, A., *Trans. Amer. Math. Soc.* 371 (2019), 3627-3646.
- [36] J.C. Naranjo, A. Ortega, Global Prym-Torelli for double coverings ramified in at least 6 points, arXiv:2005.11108. To appear in *Journal of Algebraic Geometry*.
- [37] F. Oort and J. Steenbrink, The local Torelli problem for algebraic curves, *Journées de Géométrie Algébrique d'Angers* 1979 (1979), 157-204.
- [38] R. Pardini, *On the period map for abelian covers of projective varieties*. *Annali della Scuola Normale Superiore di Pisa - Classe di Scienze* 26.4 (1998): 719-735.
- [39] M. Penegini. Surfaces isogenous to a product of curves, braid groups and mapping class groups. In *Beauville surfaces and groups*, volume 123 of *Springer Proc. Math. Stat.*, pages 129-148. Springer, Cham, 2015.
- [40] J. C. Rohde. *Cyclic coverings, Calabi-Yau manifolds and complex multiplication*, volume 1975 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2009.
- [41] A. Wright, *Shwarz triangle mappings and Teichmüller curves: abelian square-tiled surfaces*, *J. Mod. Dyn.* 6 (2012) 405- 426.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PAVIA, VIA FERRATA 5, I-27100 PAVIA, ITALY
 Email address: paola.frediani@unipv.it