

# An analytic model for gravitational collapse of spherical matter under mixed pressure

Shuichi Yokoyama,\*

*Center for Gravitational Physics,  
Yukawa Institute for Theoretical Physics, Kyoto University,  
Kitashirakawa Oiwake-cho, Sakyo-Ku, Kyoto 606-8502, Japan*

## Abstract

We investigate spherically symmetric gravitational collapse of thick matter shell without radiation in the Einstein gravity with cosmological constant. The orbit of the infalling thick matter is determined by imposing an equation of state for the matter near interface, where pressure constituted of the transverse component and the longitudinal one is proportional to energy density. We present analytic solutions for the equation of state and discuss parameter region free from any shell crossing singularity. We finally show that adopting the definition presented in arXiv:2005.13233 the total energy in this time-dependent system is invariant under the given time evolution.

---

\*shuichi.yokoyama[at]yukawa.kyoto-u.ac.jp

# 1 Introduction

The study of dynamical process for matter to collapse into a black hole is important not only in observational viewpoints but also in theoretical ones of general relativity. Oppenheimer and Snyder initiated to study a classical gravitational implosion of a compact star analytically [1], where homogeneous spherical pressureless gas falls into the center of the system without any radiation. (See also [2].) Subsequently gravitational collapses in various situations including the one with radiation were studied [3–8], which may incubate conceptually important ideas such as the cosmic censorship [9] and the singularity theorem [4]. (See also [10–13].)

A traditional approach to construct an analytic solution to describe gravitational collapse is to prepare two discontinuous space-times, one of which entails an event horizon and the other does not, and glue them smoothly at their boundaries. Indeed the Oppenheimer-Snyder solution can be obtained [14] by sewing two Tolman-Bondi space-times [15, 16], whose metric has the trivial time-component but can describe the Schwarzschild space-time in Lemaitre coordinates [17]. Such a sewing condition is also referred to as the junction condition, which is formulated in several styles [18–21]. (See also [22].) This technique was used to construct analytic solutions to describe spherically symmetric gravitational collapse of dust matter [23] including cosmological constant [24–27]. (See also [28] for references therein.)

Another approach is to take one good coordinate system for the entire space-time to describe gravitational collapse as a part of it [29]. This approach has an advantage to make it easy to get the whole picture while it may be heuristic and does not always make the calculation simplified [30, 31]. However an adequate coordinate system also makes the calculation considerably simple, which will pay off enough for an effort to find.

A main purpose of this paper is to take the second approach and extend results of spherically symmetric gravitational collapse of shell of light in [29] to that of general fluid including cosmological constant. (See also [32–34] for earlier study of matter collapse.) Upon this extension the absence of a shell crossing singularity [14] becomes a non-trivial obstruction, which constrains the parameter space of consistent solutions. (See also [35].) A characteristic point of our extension is to consider gravitational collapse of general fluid under mixed pressure of both the transverse component and the longitudinal one in its equation of state. We can still solve such a gravitational collapsing system with fluid under such a general pressure analytically.

Furthermore we compute the total energy of the system adopting the definition of charges presented in [36]. (See also [37].) The proposed definition of charges is done so as to extend the one in the flat space-time to a general curved space-time with manifest invariance under general coordinate transformation. This definition provides a simple way to test whether a charge is conserved or not by checking the associated vector field for the charge to satisfy an equation referred to as the conservation equation [38]. We show that the time evolution vector field in the matter collapsing system studied in this paper satisfies the conservation condition, and confirm that the total energy is indeed invariant under the given time evolution by explicit computation.

The rest of this paper is organized as follows. In section 2 we analyze a model of gravitational collapse of spherical thick matter shell, which outermost shell obeys an equation of state with mixed pressure of the transverse component and the longitudinal one, and solve it avoiding a shell crossing singularity. In section 3 we compute the total energy flow of this system, and show that the time evolution vector field satisfies the conservation condition and that the total energy is invariant under the given time evolution. Section 4 is devoted to summary and discussion.

## 2 Model of gravitational collapse of thick matter shell

In this section we fix the setup to study the formation of a spherically symmetric black hole in a  $d$ -dimensional spacetime with a general value of cosmological constant  $\Lambda$ , where  $d$  is greater than two.<sup>1</sup> For this purpose, following [29], we consider the original form of the Eddington-Finkelstein coordinate system<sup>2</sup>

$$ds^2 = -(1+u)dt^2 - 2udt dr + (1-u)dr^2 + r^2\tilde{g}_{ij}dx^i dx^j. \quad (2.1)$$

Here  $\tilde{g}_{ij}$  is the metric for a  $d-2$  dimensional Einstein manifold whose Ricci tensor is  $\tilde{R}_{ij} = (d-3)\tilde{g}_{ij}$ , and  $u$  is a function of  $t$  and  $r$  such that

$$u(t, r) = -\frac{2\Lambda r^2}{(d-2)(d-1)} - \frac{m(t, r)}{r^{d-3}}, \quad (2.2)$$

where  $m(t, r)$  is a matter profile function which could describe the dynamics of formation or deformation of a spherically symmetric black hole. The metric (2.1) with vanishing  $m(t, r)$  describes the AdS or dS space-time for  $\Lambda$  negative or positive, respectively, while for positive constant  $m(t, r)$  it describes a static spherically symmetric black hole. This can be easily seen by transforming (2.1) into the usual Schwarzschild metric when  $u$  is a function only of  $r$ :

$$ds^2 = -(1+u(r))(dx^0)^2 + \frac{1}{1+u(r)}dr^2 + r^2\tilde{g}_{ij}dx^i dx^j, \quad (2.3)$$

where  $x^0 = t + \int dr \frac{u(r)}{1+u(r)}$ . Thus the event horizon emerges if  $m(t, r) > 0$ , while another event horizon called the de-Sitter or cosmological event horizon will also appear when  $\Lambda > 0$  [39, 40]. Note that in the original form of the Eddington-Finkelstein coordinates the event horizon and the infinite redshift surface are identical.

---

<sup>1</sup>Our primary interest is in four dimensional system, but performing analysis in general dimensions we can also extract some interesting information at  $d=3$  as seen below.

<sup>2</sup>The Eddington-Finkelstein coordinate system may not be fully general to describe a time-dependent spherically symmetric system. For example, it does not contain the Tolman-Bondi spacetime, which has two unknown functions:  $ds_{\text{TB}}^2 = -dt^2 + X(t, r)dr^2 + Y(t, r)\tilde{g}_{ij}dx^i dx^j$ . However, the original form of the Eddington-Finkelstein coordinates suits to describe spherical matter collapse without any radiation, while other coordinate systems such as the Vaidya one may not.

The time direction in the metric (2.1) can be freely chosen. Here we determine it so that the infalling null radial geodesic does not depend on the function  $u$ , so that it becomes the same as in the flat spacetime,  $t = -r + \text{const}$ . Then the outgoing radial light ray is  $dr/dt = (1+u)/(1-u)$ . This suggests that there is an infinite redshift at  $u = -1$ . Note that any light does not come out to the spacial infinity if the cosmological constant is nonzero.

We wish to study whether there exists any matter which collapses gravitationally in the manner described by (2.1). To this end, we first determine the form of the matter energy momentum tensor so as to satisfy the Einstein equation,  $R^\mu_\nu - \delta^\mu_\nu \frac{R}{2} + \delta^\mu_\nu \Lambda = 8\pi G_N T^\mu_\nu$ , by computing the curvature tensors for the metric (2.1). The result is as follows.

$$\begin{aligned} T^t_t &= \frac{d-2}{16\pi G_N} \frac{\partial_r(r^{d-3}\delta u)}{r^{d-2}}, \\ T^t_r &= \frac{d-2}{16\pi G_N} \frac{\partial_t u}{r} = -T^r_t, \\ T^r_r &= \frac{d-2}{16\pi G_N} \left( \frac{\partial_r(r^{d-3}\delta u)}{r^{d-2}} - 2\frac{\partial_t u}{r} \right), \\ T^i_j &= \frac{\delta^i_j}{16\pi G_N} \frac{(\partial_r - \partial_t)^2(r^{d-3}\delta u)}{r^{d-3}}, \end{aligned} \tag{2.4}$$

where  $G_N$  is the Newton constant, and  $\delta u(t, r) = -\frac{m(t, r)}{r^{d-3}}$ .<sup>3</sup> Note that the energy momentum tensor of this model is not perfect fluid, so this model describes a non-adiabatic process of spherical gravitational collapse or blast. Remark that the  $\Lambda$  dependence cancels due to the fine-tuned coefficient of  $\Lambda$  in (2.2).<sup>4</sup> Then the density,  $\rho = -T^t_t$ , the radial component of pressure,  $p_\perp = T^r_r$ , and the angular component denoted by  $p_\parallel$ , where  $T^i_j = p_\parallel \delta^i_j$ , read

$$\begin{aligned} \rho &= \frac{d-2}{16\pi G_N} \frac{\partial_r m(t, r)}{r^{d-2}}, \\ p_\perp &= \frac{d-2}{16\pi G_N} \left( \frac{-\partial_r m(t, r)}{r^{d-2}} + 2\frac{\partial_t m}{r^{d-2}} \right), \\ p_\parallel &= \frac{1}{16\pi G_N} \frac{(\partial_r - \partial_t)^2(-m(t, r))}{r^{d-3}}. \end{aligned} \tag{2.5}$$

Now we assume the matter profile to form shell structure, so that it is a function of the one dimensional subspace such that

$$m(t, r) = m\theta(r)F\left(\frac{t+h(r)}{\Delta}\right). \tag{2.6}$$

---

<sup>3</sup>One might wonder whether the strategy here to determine the energy momentum tensor from a given metric configuration is valid or meaningful. In fact, the Einstein equation can only determine the form of the energy momentum tensor and is not sufficient to specify the matter. Taking advantage of this we perform a general analysis of gravitational collapse of unspecified matter or fluid as seen below. A goal is to elucidate some important property of gravitational collapse of fluid from this general analysis so as to extract a general lesson for further analysis.

<sup>4</sup>We have checked that (2.4) satisfies the covariant conservation  $\nabla_\mu T^\mu_\nu = 0$  in four dimensions.

Here  $m$  is a positive constant related to the black hole mass at the end,  $h(r)$  is a function to describe an orbit of a piece of matter consisting of the shell. We insert the step function  $\theta(r)$ , which plays no role to construct the solution in this section but a technically important role to compute the contribution of the energy of emergent curvature singularity in the next section.  $F(x)$  is an “upslope” function: it vanishes from  $x = -\infty$  to the origin, at which it starts to increase monotonically from 0 to 1 up to  $x = 1$  and keeps the same value afterwards from  $x = 1$  to  $x = \infty$ .<sup>5</sup> In this normalization of the upslope function  $\Delta$  represents the thickness of the shell. The upslope function plays a role of the continuous label of the matter shell such that the concentric slice of matter shell labeled by  $F = \lambda$  ( $0 \leq \lambda \leq 1$ ) has the world-line  $t = -h(r) + F^{-1}(\lambda)\Delta$ , where  $F^{-1}$  is the inverse function of  $F$  only in the region  $[0, 1]$  so that  $F^{-1}(0) = 0, F^{-1}(1) = 1$ . In particular,  $F(0) = 0$  labels the innermost shell while  $F(1) = 1$  labels the outermost one. Note that a discrete sequence of thin matter shells can be realized by choosing the upslope function as a multi-step function. This upslope function is used as a physical shell attribute in this system as we shall see in section 3. In what follows we assume  $F'(0) = F'(1) = 0$  for the continuity of the density and the radial component of pressure, and set  $h(0) = 0$  using the time shift.

Our goal is to determine the form of the function  $h(r)$  for given parameters  $\Delta, m$  and  $F(x)$ . We do this for the matter near the interface specified by  $F = 1 - \epsilon$  with  $\epsilon$  a small parameter to satisfy an equation of state such that

$$P = w\rho, \tag{2.7}$$

where  $w$  is a constant,  $\rho$  is the density, and  $P$  is the pressure the matter at the interface receives. As seen from (2.5) the pressure in the equation of state can be generally composed of the radial component and the angular one. In the following subsections, we determine  $h(r)$  in a couple of cases for  $P$  to be composed only of the angular component, only of the radial component, and of their mixture in order.

Since we impose a condition only at the junction, it is not guaranteed that a solution obtained in this way is consistent in the whole space-time region. We require a solution to be free from a shell crossing singularity [14]. The absence of such a shell crossing singularity restricts the parameter region. In order to avoid a shell crushing singularity at the junction we impose a condition for  $h(r)$  to increase monotonically. If this is not satisfied, then the shell ansatz (2.6) breaks down in some space-time region. In addition we also impose a condition that the emergent infinite redshift surface monotonically increases. Note that for positive cosmological constant the smaller event horizon is the black hole one, whose radius is increasing, while the bigger one is the cosmological event horizon, whose radius is decreasing during the matter collapsing into a black hole [41]. The infinite redshift surface is given by an equation  $u = -1$ , where  $u$  is given by (2.2), since the outgoing radial null geodesic is determined from  $dr/dt = (1+u)/(1-u)$  as shown above. Combining with (2.6)

---

<sup>5</sup>On the other hand, in the case to study the matter explosion, a “downslope” function is suitable, which is described by  $F(1-x)$ . Such a solution to describe the process of black hole diffusion will be obtained by flipping the time direction for the solutions obtained below.

the emergent infinite redshift surface can be explicitly written as

$$t = -h(r) + \Delta F^{-1} \left( \frac{r^{d-3}}{m} \left( 1 - \frac{\Lambda}{(d-2)(d-1)} r^2 \right) \right) \quad (2.8)$$

up to when the surface reach the final black hole horizon radius  $r = r_H$ , which is determined from  $\frac{r_H^{d-3}}{m} \left( 1 - \frac{\Lambda}{(d-2)(d-1)} r_H^2 \right) = 1$ . The absence of a shell crossing singularity requires the right hand side to increase monotonically with respect to  $r$ .

The two conditions can be summarized as

$$0 < h'(r) < \frac{\Delta}{m} r^{d-4} \left( d - 3 - \frac{\Lambda}{d-2} r^2 \right) (F^{-1})' \left( \frac{r^{d-3}}{m} \left( 1 - \frac{\Lambda}{(d-2)(d-1)} r^2 \right) \right) \quad (2.9)$$

for  $r < r_H$ . We have comments on the second inequality in (2.9), which comes from the second condition above. The second inequality is sensitive to the signature of the cosmological constant. If the cosmological constant is not positive, then the right-hand side can be arbitrary large by taking  $\Delta$  to be large for any upslope function  $F$ . This means that the second inequality in (2.9) is satisfied if  $\Delta$  is sufficiently large when  $\Lambda \leq 0$ . On the other hand, a careful analysis is necessary for the case of a positive cosmological constant. It is interesting to comment at three dimensions. At  $d = 3$ , (2.9) reduces to

$$0 < h'(r) < \frac{\Delta}{m} (-\Lambda r) (F^{-1})' \left( \frac{1}{m} \left( 1 - \frac{\Lambda}{2} r^2 \right) \right). \quad (2.10)$$

This inequality cannot be satisfied unless the cosmological constant is negative. This shows that a (spherically symmetric) black hole in three dimensions cannot form unless there is no negative cosmological constant in the Einstein gravity [42, 43].

Below we determine the form of  $h(r)$ , and subsequently study a qualitative parameter region to satisfy (2.9) implied by numerical calculation.

## 2.1 Transverse pressure

We first consider a case where the pressure in the equation of state is composed only of the angular component, so that the equation of state is  $p_{\parallel} = w_{\parallel}^0 \rho$ . Plugging this and (2.5) into (2.7) we find

$$y(h'(r) - 1)^2 + h''(r) = -\frac{w_{\parallel}}{r} h'(r), \quad (2.11)$$

where  $w_{\parallel} = (d-2)w_{\parallel}^0$ ,  $y = \frac{F''(1-\epsilon)}{F'(1-\epsilon)} \Delta$ . We rewrite (2.11) in more useful form later as

$$y w_{\parallel} H(r)^2 + H'(r) = -\frac{w_{\parallel}}{r} H(r) - \frac{1}{r}, \quad (2.12)$$

where

$$H(r) = \frac{h'(r) - 1}{w_{\parallel}}. \quad (2.13)$$

For  $y = 0$  this differential equation can be easily solved as  $h(r) = \frac{c_0}{1-w_{\parallel}} r^{1-w_{\parallel}}$ , where we used  $h(0) = 0$  and  $c_0$  is an integration constant, unless  $w_{\parallel} = 1$ . A solution with  $w_{\parallel} = 1$  is obtained by formally taking the limit of this solution and ignoring a constant term:  $h(r) = c_0 \log r$ . We fix the integration constant  $c_0$  by requesting that the shell of the pressureless matter of  $w_{\parallel} = 0$  is the light-like shell or null-dust. This is satisfied if  $c_0 = 1$  at  $w_{\parallel} = 0$ . For simple presentation we set  $c_0 = 1$  in what follows:

$$h(r) = \frac{r^{1-w_{\parallel}}}{1-w_{\parallel}}. \quad (2.14)$$

This implies that  $w_{\parallel}$  has to be smaller than unity for a reasonable solution.

For  $y > 0$ , it is difficult to solve (2.11) analytically in terms of  $h(r)$ , while it is possible in terms of  $h'(r)$  or to solve (2.12) as follows.

$$H(r) = \frac{\Gamma(2-w_{\parallel})J_{-w_{\parallel}}(2\sqrt{ryw_{\parallel}}) - c_y\Gamma(w_{\parallel})J_{w_{\parallel}}(2\sqrt{ryw_{\parallel}})}{\sqrt{ryw_{\parallel}}\left(\Gamma(2-w_{\parallel})J_{1-w_{\parallel}}(2\sqrt{ryw_{\parallel}}) + c_y\Gamma(w_{\parallel})J_{w_{\parallel}-1}(2\sqrt{ryw_{\parallel}})\right)}, \quad (2.15)$$

where  $c_y$  is an integration constant and  $J_{\nu}(z)$  is the Bessel function of the first kind. Note that in order to reach this form we used the following recurrence equation for the Bessel function

$$J_{\pm(w-2)}(z) = \frac{\pm 2(w-1)}{z} J_{\pm(w-1)}(z) - J_{\pm w}(z). \quad (2.16)$$

We fix the integration constant  $c_y$  for this solution to reduce to the one (2.14) by taking the limit  $y \rightarrow 0$ . To this end we expand the solution (2.19) at  $y \sim 0$  keeping  $|w_{\parallel}| \ll 1$ . Employing the asymptotic expansion of the Bessel function  $J_{\alpha}(z) \sim \frac{1}{\Gamma(\alpha+1)}\left(\frac{z}{2}\right)^{\alpha}$  around  $z \sim 0$ , we find

$$H'(r) = \frac{1}{w_{\parallel}} - \frac{(1-w_{\parallel})(ryw_{\parallel})^{-w_{\parallel}}}{c_y} + \dots, \quad (2.17)$$

where the ellipsis contains higher ordered terms. Therefore it is sufficient to choose as

$$c_y = \frac{1-w_{\parallel}}{y^{w_{\parallel}}w_{\parallel}^{w_{\parallel}-1}} \quad (2.18)$$

for the reduction to (2.14) with  $y \rightarrow 0$ . As a result we obtain

$$H(r) = \frac{1}{\sqrt{ryw_{\parallel}}} \frac{(yw_{\parallel})^{w_{\parallel}}\Gamma(2-w_{\parallel})J_{-w_{\parallel}}(2\sqrt{ryw_{\parallel}}) + (w_{\parallel}-1)w_{\parallel}\Gamma(w_{\parallel})J_{w_{\parallel}}(2\sqrt{ryw_{\parallel}})}{(yw_{\parallel})^{w_{\parallel}}\Gamma(2-w_{\parallel})J_{1-w_{\parallel}}(2\sqrt{ryw_{\parallel}}) - (w_{\parallel}-1)w_{\parallel}\Gamma(w_{\parallel})J_{w_{\parallel}-1}(2\sqrt{ryw_{\parallel}})}. \quad (2.19)$$

Note that we could not further perform the integration analytically to obtain  $h(r)$ .

Let us study a region where  $h(r)$  is a monotonically increasing function or  $h'(r)$  is positive. It can be confirmed that the condition is always satisfied if both  $y$  and  $w_{\parallel}$  are sufficiently small by using the asymptotic formula of the Bessel function of the first kind




Figure 1: The blue and yellow curves depict  $h'(r)$  and  $h(r)$ , respectively, with parameters chosen as shown.

$J_\nu(z) \sim \sqrt{\frac{2}{\pi|z|}} \cos(z - \frac{\pi}{4} - \frac{\nu\pi}{2})$  for  $z \sim \infty$ . However, numerical results suggest that the solution does not meet the condition and hit a crushing singularity from a certain  $r$  if  $y$  or  $w_{\parallel}$  is close to one as seen from Figure 1. In fact the numerical solution for  $h(r)$  starts oscillating at large  $r$  and is not reliable in this regime.

We shall give a comment on the second inequality in (2.9) in the next section drawing a graph with parameters suitably chosen.

## 2.2 Longitudinal pressure

Next we consider a case where the pressure in (2.7) is given by the radial component, so that  $p_{\perp} = w_{\perp}\rho$ . It seems that this case was not studied before, but it is theoretically possible and becomes a basis for the study of mixed pressure in the next subsection. Substituting (2.5) into (2.7) we obtain

$$h'(r) = \frac{2}{1+w_{\perp}} \rightsquigarrow h(r) = \frac{2}{1+w_{\perp}}r. \quad (2.20)$$

As a result the collapse of matter shell to satisfy  $p_{\perp} = w_{\perp}\rho$  outermost is equivalent to the one to satisfy  $p_{\parallel} = 0$  with  $F''(1) = 0$  and the integration constant  $c_0 = \frac{2}{1+w_{\perp}}$ .

## 2.3 Mixed pressure

At the end of this section we consider the collapse of matter under mixed pressure such that  $P = p_{\parallel} + vp_{\perp}$ , where  $v$  is a non-negative parameter. Employing (2.5) the equation of state (2.7) can be computed as

$$y(h'(r) - 1)^2 + h''(r) - \frac{\bar{v}}{r} = -\frac{\bar{w}}{r}h'(r), \quad (2.21)$$

where  $\bar{v} = 2(d-2)v$ ,  $\bar{w} = (d-2)(v+w)$ . This can be further rewritten as

$$y(\bar{w} - \bar{v})H(r)^2 + H'(r) = -\frac{\bar{w}}{r}H(r) - \frac{1}{r}, \quad (2.22)$$

where

$$H(r) = \frac{h'(r) - 1}{\bar{w} - \bar{v}}. \quad (2.23)$$

This differential equation (2.22) in this form can be obtained from the previous one (2.12) by replacing  $yw_{\parallel} \rightarrow y(\bar{w} - \bar{v})$  and  $w_{\parallel} \rightarrow \bar{w}$ . Therefore the solution for (2.22) can be also obtained from that for (2.12), which is given by (2.19), by the same replacement:

$$\frac{h'(r) - 1}{\bar{w} - \bar{v}} = \frac{\Gamma(2 - \bar{w})J_{-\bar{w}}(2x) - c\Gamma(\bar{w})J_{\bar{w}}(2x)}{x(\Gamma(2 - \bar{w})J_{1-\bar{w}}(2x) + c\Gamma(\bar{w})J_{\bar{w}-1}(2x))}, \quad (2.24)$$

where  $c$  is an integration constant,  $x = \sqrt{ry(\bar{w} - \bar{v})}$ . We fix the integration constant  $c$  by requesting this solution to reduce to (2.19) by taking the limit  $\bar{v} \rightarrow +0$ . This can be easily fixed by the above replacement of parameters for (2.18) taking into account  $\bar{v} \sim 0$ :

$$c = \frac{\bar{w}(1 - \bar{w})}{(y\bar{w})^{\bar{w}}} + \alpha\bar{v}, \quad (2.25)$$

where  $\alpha$  is an arbitrary number which can be dependent on  $\bar{v}$  and  $\bar{w}$  in a non-singular manner. We fix  $\alpha$  so that this solution reduces to the one with only radial pressure, (2.20), in the limit  $\bar{v}, \bar{w} \rightarrow \infty$  holding their ratio fixed. For simplicity we take the limit  $\bar{v}, \bar{w} \rightarrow \infty$  as well as  $y \rightarrow 0$  with  $y(\bar{w} - \bar{v})$  fixed. Then  $x$  becomes constant in this limit. If we choose  $\alpha \neq 0$ , the leading term of the above solution (2.24) can be computed as

$$h'(r) - 1 \sim \frac{(\bar{w} - \bar{v})(-J_{\bar{w}}(2x))}{xJ_{\bar{w}-1}(2x)} \sim -1 + \frac{\bar{v}}{\bar{w}}, \quad (2.26)$$

where we used the asymptotic formula  $J_{\nu}(z) \sim \frac{1}{\sqrt{2\pi\nu}}\left(\frac{ez}{2\nu}\right)^{\nu}$  with  $\nu \sim \infty$ . Thus any nonzero  $\alpha$  gives a desired solution. For  $\alpha = 1$ , the final solution is expressed as

$$h'(r) = 1 + \frac{(\bar{w} - \bar{v})[(y\bar{w})^{\bar{w}}\Gamma(2 - \bar{w})J_{-\bar{w}}(2x) - \{\bar{w}(1 - \bar{w}) + (y\bar{w})^{\bar{w}\bar{v}}\}\Gamma(\bar{w})J_{\bar{w}}(2x)]}{x[(y\bar{w})^{\bar{w}}\Gamma(2 - \bar{w})J_{1-\bar{w}}(2x) + \{\bar{w}(1 - \bar{w}) + (y\bar{w})^{\bar{w}\bar{v}}\}\Gamma(\bar{w})J_{\bar{w}-1}(2x)]}. \quad (2.27)$$

Let us comment on the valid parameter region where  $h(r)$  is a monotonically increasing function or  $h'(r)$  is positive including the parameter  $\bar{v}$ . Numerical results suggest that  $h'(r) > 0$  for sufficiently small  $\bar{v}$  but not with  $\bar{v}$  closed to one keeping both  $y$  and  $\bar{w}$  sufficiently small. If one of  $y, \bar{w}, \bar{v}$  becomes close to one, we observe that the solution hits a shell crossing singularity at some  $r$ . To summarize, the valid parameter regime is (i)  $\bar{v}, |\bar{w}|, y \ll 1$  with  $\bar{v}y > 0$ , or (ii)  $|\bar{v}|, |\bar{w}| \gg 1$  with  $\bar{v}y > 0, y \ll 1, |\bar{v}/\bar{w}| \sim 1$ .

### 3 Conservation of energy

In this section we investigate the flow of energy and energy density in the system studied in the previous section.

Since all the matter collapses into a black hole in the process, the total energy is expected to be conserved. It was shown in [44] that the Misner-Sharp mass [3] in similar gravitational collapsing systems is invariant under the co-moving time differentiation [16].

On the other hand, there has been a proposal of a manifestly covariant definition of the total energy in field theory on a general curved space-time [36]

$$E = \int_{\mathbf{R}^{d-1}} d^{d-1}x \sqrt{|g|} T^t{}_\mu n^\mu, \quad (3.1)$$

where  $n^\mu$  is the time evolution vector field given by  $n^\mu = -\delta_t^\mu$ . In what follows we confirm that the energy defined by (3.1) is invariant under the usual time evolution.

An advantage in the definition (3.1) is that whether the charge of the form (3.1) conserves or not can be easily determined by whether the vector field to define the charge satisfies the conservation condition [38]

$$T^\nu{}_\mu \nabla_\nu n^\mu = 0. \quad (3.2)$$

In the current situation one can easily check that this equation is satisfied using (2.4). Thus we conclude that the charge  $E$  defined by (3.1) is conserved for all the cases studied in section 2 as long as the solution is meaningful without any shell crossing singularity.

More explicitly we compute the total energy separating the time periods:  $\{t < 0\}$ , when the thick matter shell is on-going falling before the black hole forms,  $\{0 \leq t < \Delta\}$ , the black hole is forming, and  $\{\Delta \leq t\}$ , the thick matter shell completes infalling. Plugging (2.4) into (3.1) we obtain

$$E = \frac{d-2}{16\pi G_N} V_{d-2} \int_0^\infty dr \partial_r (m(t, r)), \quad (3.3)$$

where  $V_{d-2} = \int d^{d-2}x \sqrt{\bar{g}}$  is the volume of the internal Einstein manifold.

When  $t < 0$ , the innermost shell specified by  $F = 0$  does not reach the origin:  $0 < h^{-1}(-t)$ . Therefore we can compute the total energy as

$$E = \frac{d-2}{16\pi G_N} V_{d-2} m(F(1) - F(0)) = M, \quad (3.4)$$

where  $M = \frac{d-2}{16\pi G_N} V_{d-2} m$ .

At  $t = 0$  the black hole starts to form, and it becomes growing up to  $t = \Delta$ . In this period both the infalling matter shell and the black hole exist. We denote the energy of the former by  $M_1$  and that of the latter by  $M_2$ . Then the total energy (3.3) is given by the sum of two:  $E = M_1 + M_2$ .  $M_1$  is obtained by acting the differentiation with respect to  $r$  on the upslope function  $F$  in  $m(t, r)$ . Thus it is computed as

$$M_1 = \frac{d-2}{16\pi G_N} V_{d-2} \int_0^\infty dr \partial_r F\left(\frac{t+h(r)}{\Delta}\right) = M\left(1 - F\left(\frac{t}{\Delta}\right)\right), \quad (3.5)$$

where we used  $h(0) = 0$  and the assumption that  $h(r)$  increases monotonically to infinity. On the other hand,  $M_2$  is obtained by acting the differentiation with respect to  $r$  on the

| Time                        | $t$   | $t < 0$ | $0 \leq t < \Delta$          | $\Delta \leq t$ |
|-----------------------------|-------|---------|------------------------------|-----------------|
| Energy of collapsing matter | $M_1$ | $M$     | $M(1 - F(\frac{t}{\Delta}))$ | 0               |
| Mass of black hole          | $M_2$ | 0       | $MF(\frac{t}{\Delta})$       | $M$             |
| Total energy                | $E$   | $M$     | $M$                          | $M$             |

Table 1: The energy flow is shown. The total energy is conserved.

step function  $\theta$  in  $m(t, r)$ , which converts to the delta function. Therefore it is computed as<sup>6</sup>

$$M_2 = \frac{d-2}{16\pi G_N} V_{d-2} m \int_0^\infty dr (\delta(r) F(\frac{t+h(r)}{\Delta})) = MF(\frac{t}{\Delta}). \quad (3.6)$$

The total energy is  $E = M$  as expected.

The matter shell ends falling at  $t = \Delta$ . Afterwards the system becomes static, so is the energy flow. The energy configuration is obtained by taking the limit  $t \rightarrow \Delta - 0$ .

We summarize the result of energy flow in Table 2. For instruction, we draw a picture of the Carter-Penrose diagram for the spherical collapse of thick light shell in Figure 2.

We can also compute the energy inside the radius  $R$  in the system denoted by  $E(t, R)$ :

$$\begin{aligned} E(t, R) &= - \int_{\sqrt{x^2} \leq R} d^{d-1} x \sqrt{|g|} T_t^t = \frac{d-2}{16\pi G_N} V_{d-2} \int_0^R dr \partial_r (m(t, r)) \\ &= MF(\frac{t+h(R)}{\Delta}). \end{aligned} \quad (3.7)$$

This is a monotonically increasing function inside the fluid shells as long as  $h(r)$  is. This can be used as an observable shell attribute.

For concreteness we draw a picture of gravitational collapse of thick matter shell under pressure from the transverse direction without any shell crossing singularity in four dimensions. To this end we first determine the location of the final event horizons after the thick matter shell ends infalling. The determining equation of the event horizons is

$$r^3 + \frac{6}{\Lambda}(-r + m) = 0. \quad (3.8)$$

This is solved by using the Cardano's formula as  $r = r_k$ , where

$$r_k = \omega^k \sqrt[3]{-\frac{3m}{\Lambda} + \sqrt{(\frac{3m}{\Lambda})^2 - (\frac{2}{\Lambda})^3}} + \omega^{3-k} \sqrt[3]{-\frac{3m}{\Lambda} - \sqrt{(\frac{3m}{\Lambda})^2 - (\frac{2}{\Lambda})^3}}, \quad (3.9)$$

---

<sup>6</sup>Here we present the computation of the delta function in the radial coordinate in a naive expression. This part can be computed in a more formally rigorous fashion by rewriting  $\delta(r)$  in terms of the  $d$ -dimensional delta function  $\delta^d(x)$  employing an equation obtained by evaluating the equation  $\int d^d x \delta^d(x) = 1$  with the polar coordinates. However such a formal description with extra complication might make physics obscure. Here we adopt a simple description to elucidate physics at the cost of the rigorous expression.




Figure 2: The Carter-Penrose diagram of the spherical collapse of the thick light shell is depicted. We almost use conventional notation.  $O$  is the origin,  $O^+$  is the location of the curvature singularity, and the band in the middle describes the thick infalling light shell. The gradation of the color schematically describes the density of the energy. In addition three Eddington-Finkelstein time slices are schematically drawn, where  $t_- < 0 < t_0 < \Delta < t_+$ . Any intersecting point of the thick light shell with the time slice and that of the curvature singularity contributes to the integral of the total energy at the time. In particular the contribution from the curvature singularity is the mass of the black hole.

with  $k = 0, 1, 2$ ,  $\omega = \frac{-1+\sqrt{3}}{2}$  the cube root of unity. The discriminant is computed as  $3(\frac{6}{\Lambda})^2(-9m^2 + \frac{8}{\Lambda})$ . Therefore for negative cosmological constant the discriminant is negative and only one positive solution exists as  $r = r_0$ . On the other hand, for positive cosmological constant smaller than  $\Lambda_0 = \frac{8}{9m^2}$  all three solutions for (3.8) are real, two of which are positive,  $r_1, r_2 > 0$ , while for  $\Lambda > \Lambda_0$  there is no positive real solution, so that the horizons do not exist [45]. For positive cosmological constant, as the final mass of the black hole

| Cosmological constant | $\Lambda$       | $\Lambda < 0$ | 0   | $0 < \Lambda \leq \Lambda_0$ | $\Lambda_0 < \Lambda$ |
|-----------------------|-----------------|---------------|-----|------------------------------|-----------------------|
| Black hole horizon    | $r_{\text{BH}}$ | $r_0$         | $m$ | $r_1$                        | -                     |
| Cosmological horizon  | $r_{\text{dS}}$ | -             | -   | $r_2$                        | -                     |

Table 2: The locations of the final black hole event horizon and the cosmological one after the end of the matter infalling are shown with the final mass of the black hole fixed. “-” means the absence of the corresponding horizon.

increases, the black hole event horizon radius  $r_1$  increases while the cosmological one  $r_2$  decreases [41].

For concreteness we choose an upslope function as  $F(x) = \sin^2 \frac{\pi x}{2}$ , whose inverse function is  $F^{-1}(y) = \frac{2}{\pi} \arcsin \sqrt{y}$ . Then the growing infinite redshift surface is

$$t = -h(r) + \Delta F^{-1}\left(\frac{r}{m}\left(1 - \frac{\Lambda}{6}r^2\right)\right), \quad (3.10)$$

up to when the surface intersects the interface or the outermost thick matter shell. Afterwards it matches the final black hole horizon  $r = r_{\text{BH}}$ . The time for the surface to reach




Figure 3: A solution of the gravitational collapse of the thick matter shell for positive cosmological constant without any shell crushing singularity is depicted. The parameters are chosen such that  $\bar{v} = 0, \bar{w} = 0.1, \Delta = 10, m = 4, \Lambda = 0.05, \epsilon = 0.05$ , and an upslope function  $F(x) = \sin^2 \frac{\pi x}{2}$ . For negative cosmological constant the corresponding picture is qualitatively the same except the disappearance of the cosmological event horizon.

$r = r_{BH}$  is  $t_{BH} = \Delta - h(r_{BH})$ . The light-ray which passes through the intersecting point of the infinite redshift surface and the interface,  $(r_{BH}, t_{BH})$ , does not go out, and any light-ray emitted after the last light-ray is trapped inside the horizon. Thus the orbit of the last light-ray describes the radius of the growing black hole event horizon, which is larger than the infinite redshift radius at an equal time until it reaches the final black hole horizon radius. The orbit of the last light-ray satisfies an integral equation  $t = \int_{r_{BH}}^r \frac{1-u}{1+u} dr + t_{BH}$ . In particular, before an outgoing light-ray reaches the thick matter shell, the orbit can be computed as  $t = \frac{1+\Lambda r^2}{1-\Lambda r^2}(r - r_{BH}) + t_{BH}$ . We draw a figure for a consistent solution in this setup in Fig. 3.

## 4 Discussion

We have investigated an analytic model of spherical gravitational collapse of thick matter shell without any radiation, where the metric is described by the Eddington-Finkelstein coordinate system with the mass parameter replaced with a function of the radial coordinate and the time. We have argued a condition which makes a solution free from a shell crossing singularity, which was determined by requesting the monotonic infalling matter and the monotonic grow of infinite redshift surface in general dimensions.

For general analysis we have imposed an equation of state for the outermost matter consisting of the shell where pressure as a mixture of the longitudinal component and the transverse one is proportional to the energy density. Solving this equation of state at interface we have determined the form of the function describing the orbit of the matter shell or its derivative analytically, and discussed a parameter region where the solution is free from a shell crossing singularity. We have finally confirmed that employing the

definition of the energy presented in [36] the total energy of the system is invariant under the ordinary time translation.

In order to avoid a shell crushing singularity for the infalling matter, we have imposed the function to describe the infalling matter as well as the emergent infinite redshift surface to monotonically increase. This condition is summarized as (2.9). At three dimensions this simple inequality leads to a well-known result that negative cosmological constant is required for a black hole to exist. It would be interesting to investigate this inequality at four dimensions and extract any non-trivial condition for a black hole to be present.

This paper has clarified that a gravitational collapse of matter shell generally suffers from shell crossing singularity and that even so it is possible to construct analytic solutions by choosing parameters suitably. This is an encouraging result to find such a solution describing gravitational collapse by specifying matter. The construction of such a solution with matter specified may give a hint to clarify the meaning of the off-diagonal component of the energy momentum tensor. As seen from the construction, the anisotropic form of fluid is important to construct the analytic solutions collapsing into a black hole. It would be interesting to understand the physical importance of the heat flux in this process.

The system studied in this paper describes a rather simple gravitational collapse, but will be useful as a toy model. It is also possible to investigate this system by adding other structures keeping the spherical symmetry. One is to make the radius part of the metric dependent on time. In such a situation, the relative velocity of adjacent fluid element inside the same shell becomes non-trivial and will play an important role to construct a solution [3]. Another interesting extension is to take into account the radiation and compute the energy of emitted gravitational wave. (See also [32, 46, 47].) Although the definition of the energy used in this paper is based on the matter energy-momentum tensor, it could be possible to also compute the energy of the gravitational wave by computing the difference between the energy of the matter at the initial state and that of the final one. It would be of interest to compare the result to the one obtained by any other known approach.

We hope to come back to these issues in near future.

## Acknowledgement

The author would like to thank Sinya Aoki, Tetsuya Onogi for participation at an early stage of the project. He is also grateful to Tetsuya Shiromizu, Shigeki Sugimoto and especially Sinya Aoki for useful discussion and valuable comments on the draft. He appreciates an anonymous referee for giving accurate comments shedding light on physical aspects underlying the paper. This work is supported in part by the Grant-in-Aid of the Japanese Ministry of Education, Sciences and Technology, Sports and Culture (MEXT) for Scientific Research (No. JP19K03847).

## References

- [1] J. R. Oppenheimer and H. Snyder, “On Continued Gravitational Contraction,” *Physical Review* **56** (Sept., 1939) 455–459.
- [2] B. Datt, “Über eine Klasse von Lösungen der Gravitationsgleichungen der Relativität,” *Zeitschrift für Physik* **108** (1938) 314–321.
- [3] C. W. Misner and D. H. Sharp, “Relativistic equations for adiabatic, spherically symmetric gravitational collapse,” *Phys. Rev.* **136** (1964) B571–B576.
- [4] R. Penrose, “Gravitational collapse and space-time singularities,” *Phys. Rev. Lett.* **14** (1965) 57–59.
- [5] B. K. Harrison, K. S. Thorne, M. Wakano, and J. A. Wheeler, *Gravitation Theory and Gravitational Collapse*. 1965.
- [6] P. C. Vaidya, “An Analytical Solution for Gravitational Collapse with Radiation,” *Astrophys. J.* **144** (June, 1966) 943.
- [7] G. C. McVittie, “An Example of Gravitational Collapse in General Relativity,” *Astrophys. J.* **143** (Mar., 1966) 682.
- [8] H. Nariai, “A Simple Model for Gravitational Collapse with Pressure Gradient,” *Progress of Theoretical Physics* **38** (07, 1967) 92–106,  
<https://academic.oup.com/ptp/article-pdf/38/1/92/5366896/38-1-92.pdf>.
- [9] R. Penrose, “Gravitational collapse: The role of general relativity,” *Riv. Nuovo Cim.* **1** (1969) 252–276.
- [10] S. W. Hawking and R. Penrose, “The Singularities of gravitational collapse and cosmology,” *Proc. Roy. Soc. Lond.* **A314** (1970) 529–548.
- [11] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2, 2011.
- [12] R. M. Wald, “Gravitational collapse and cosmic censorship,” 10, 1997.  
[gr-qc/9710068](https://arxiv.org/abs/gr-qc/9710068).
- [13] J. M. M. Senovilla and D. Garfinkle, “The 1965 penrose singularity theorem,” *Classical and Quantum Gravity* **32** (jun, 2015) 124008.
- [14] D. M. Eardley and L. Smarr, “Time function in numerical relativity. Marginally bound dust collapse,” *Phys. Rev. D* **19** (1979) 2239–2259.
- [15] R. C. Tolman, “Effect of Inhomogeneity on Cosmological Models,” *Proceedings of the National Academy of Science* **20** (Mar., 1934) 169–176.

- [16] H. Bondi, “Spherically symmetrical models in general relativity,” *Mon. Not. Roy. Astron. Soc.* **107** (1947) 410–425.
- [17] G. Lemaître, “L’Univers en expansion,” *Annales de la Société Scientifique de Bruxelles* **53** (Jan., 1933) 51.
- [18] G. Darrois, *Les équations de la gravitation einsteinienne*. No. 25 in Mémorial des sciences mathématiques. Gauthier-Villars, 1927.
- [19] S. O’Brien and J. L. Synge, “Jump conditions at discontinuities in general relativity.” *Commun. Dublin Inst. Advanced Stud.*, Ser. A 9, 20 p. (1952)., 1952.
- [20] A. Lichnerowicz, *Theories relativistes de la gravitation et de l’électromagnétisme. Relativité générale et théories unitaires*. 1955.
- [21] W. Israel, “Discontinuities in spherically symmetric gravitational fields and shells of radiation,” *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences* **248** (1958), no. 1254, 404–414.
- [22] W. Bonner and P. Vickers, “Junction conditions in general relativity,” *General Relativity and Gravitation* **13(1)** (1981) 29–36.
- [23] T. P. Singh and L. Witten, “Cosmic censorship and spherical gravitational collapse with tangential pressure,” *Class. Quant. Grav.* **14** (1997) 3489–3499, [gr-qc/9701002](#).
- [24] D. Garfinkle and C. Vuille, “Gravitational collapse with a cosmological constant,” *General Relativity and Gravitation* **23** (Apr., 1991) 471–475.
- [25] M. Cissoko, J. C. Fabris, J. Gariel, G. Le Denmat, and N. O. Santos, “Gravitational dust collapse with cosmological constant,” [gr-qc/9809057](#).
- [26] D. Markovic and S. Shapiro, “Gravitational collapse with a cosmological constant,” *Phys. Rev. D* **61** (2000) 084029, [gr-qc/9912066](#).
- [27] K. Lake, “Gravitational collapse of dust with a cosmological constant,” *Phys. Rev. D* **62** (2000) 027301, [gr-qc/0002044](#).
- [28] P. S. Joshi and D. Malafarina, “Recent developments in gravitational collapse and spacetime singularities,” *Int. J. Mod. Phys. D* **20** (2011) 2641–2729, [1201.3660](#).
- [29] R. J. Adler, J. D. Bjorken, P. Chen, and J. S. Liu, “Simple analytic models of gravitational collapse,” *Am. J. Phys.* **73** (2005) 1148–1159, [gr-qc/0502040](#).
- [30] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*. W. H. Freeman, San Francisco, 1973.

- [31] L. D. Landau and E. M. Lifschits, *The Classical Theory of Fields*, vol. Volume 2 of *Course of Theoretical Physics*. Pergamon Press, Oxford, 1975.
- [32] G. L. Alberghi, R. Casadio, G. P. Vacca, and G. Venturi, “Gravitational collapse of a shell of quantized matter,” *Class. Quant. Grav.* **16** (1999) 131–147, [gr-qc/9808026](#).
- [33] G. L. Alberghi, R. Casadio, and D. Fazi, “Classical dynamics and stability of collapsing thick shells of matter,” *Class. Quant. Grav.* **23** (2006) 1493–1506, [gr-qc/0601062](#).
- [34] B. K. Tippett and V. Husain, “Gravitational collapse of quantum matter,” *Phys. Rev. D* **84** (2011) 104031, [1106.1118](#).
- [35] P. S. Joshi and R. V. Saraykar, “Shell-crossings in Gravitational Collapse,” *Int. J. Mod. Phys. D* **22** (2013) 1350027, [1205.3263](#).
- [36] S. Aoki, T. Onogi, and S. Yokoyama, “Conserved charges in general relativity,” *Int. J. Mod. Phys. A* **36** (2021), no. 10, [2150098](#), [2005.13233](#).
- [37] V. Fock, *Theory of Space, Time, and Gravitation*. Pergamon Press, New York, USA, 1959.
- [38] S. Aoki, T. Onogi, and S. Yokoyama, “Charge Conservation, Entropy Current, and Gravitation,” [2010.07660](#).
- [39] B. Carter, “The commutation property of a stationary, axisymmetric system,” *Commun. Math. Phys.* **17** (1970) 233–238.
- [40] B. Carter, “Black holes equilibrium states,” in *Les Houches Summer School of Theoretical Physics: Black Holes*. 1973.
- [41] G. W. Gibbons and S. W. Hawking, “Cosmological Event Horizons, Thermodynamics, and Particle Creation,” *Phys. Rev. D* **15** (1977) 2738–2751.
- [42] M. Banados, C. Teitelboim, and J. Zanelli, “The Black hole in three-dimensional space-time,” *Phys. Rev. Lett.* **69** (1992) 1849–1851, [hep-th/9204099](#).
- [43] D. Ida, “No black hole theorem in three-dimensional gravity,” *Phys. Rev. Lett.* **85** (2000) 3758–3760, [gr-qc/0005129](#).
- [44] L. Herrera and N. O. Santos, “Local anisotropy in self-gravitating systems,” *Phys. Rept.* **286** (1997) 53–130.
- [45] S. A. Hayward, T. Shiromizu, and K.-i. Nakao, “A Cosmological constant limits the size of black holes,” *Phys. Rev. D* **49** (1994) 5080–5085, [gr-qc/9309004](#).
- [46] G. L. Alberghi, R. Casadio, G. P. Vacca, and G. Venturi, “Gravitational collapse of a radiating shell,” *Phys. Rev. D* **64** (2001) 104012, [gr-qc/0102014](#).

- [47] V. Baccetti, R. B. Mann, and D. R. Terno, “Role of evaporation in gravitational collapse,” *Class. Quant. Grav.* **35** (2018), no. 18, 185005, 1610.07839.