

Four-Loop Gauge and Three-Loop Yukawa Beta Functions in a General Renormalizable Theory

Alexander BEDNYAKOV^{1,2,*} and Andrey PIKELNER^{1,†}

¹*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna 141980, Russia*

²*P.N. Lebedev Physical Institute of the Russian Academy of Sciences, Moscow 119991, Russia*

We present the beta functions of gauge and Yukawa couplings in general four-dimensional quantum field theory, at four and three loops, respectively. The essence of our approach is fixing unknown coefficients in the most general ansatz for beta functions by direct calculation in several simplified models. We apply our results to the Standard Model and its extension with an arbitrary number of Higgs doublets and provide expressions for all four-loop gauge couplings beta functions with matrix Yukawa interactions.

INTRODUCTION

Renormalization group equations (RGE) find their numerous applications in many physical problems formulated in the language of quantum field theory (QFT). Being a convenient tool to improve perturbation theory results, RGE allow one to obtain high-precision predictions for various quantities ranging from critical exponents in the theory of critical phenomena to observables in the Standard Model (SM) and its extensions.

In the early 1980s, Machacek and Vaughn [1–3] presented their two-loop RG functions in $\overline{\text{MS}}$ scheme for all dimensionless couplings in general four-dimensional renormalizable QFT. Several misprints have been corrected during the subsequent 35 years, and the classical result has been extended to include RG functions for dimensionful parameters [4–7]. The two-loop general expressions became highly demanded in studies of the SM extensions. After the appearance of computer codes aimed to generate RGE for user-specified models, two-loop RG studies have become *de facto* standard in a New Physics analysis.

After discovering the Higgs boson at the LHC [8, 9] and a burst of activity on the vacuum stability problem (see, e.g., [10]), it became clear that three-loop RG functions [11–16] can play an essential role in precision studies of the SM and its extensions. Partial four-loop results in the SM [17–19] and three-loop beta functions in the Two-Higgs-Doublet Model (THDM) [20] became available during the past few years.

Recently, two major steps toward general high-order results have been made. First of all, RG functions in general theories were represented by a linear combination of independent tensor structures (TS) corresponding to contractions of various indices. One can match these “template” expressions to known results in specific models and extract model-independent coefficients [21, 22]. Second, new ideas based on the so-called Weyl Consistency Conditions (WCC) [23–25] allow one to find relations between known and unknown TS coefficients, thus, putting constraints on to-be-computed numbers. In particular, WCC relate gauge, Yukawa, and self-coupling beta functions computed at four, three, and two loops, respectively.

Let us also mention that WCC allows to resolve [26] a well-known issue with γ_5 ambiguity (see, e.g., [27]) in $\overline{\text{MS}}$ RG functions. Poole and Thomsen in Ref. [26] use WCC to relate the ambiguous four-loop terms in strong coupling beta function [17, 18] to unambiguous three-loop contributions to Yukawa beta functions [14, 28] and confirm the prescription advocated in Ref. [17]. The relation holds for any four-loop gauge beta function in a general QFT, and it was immediately used for the gauge sector of the SM with diagonal Yukawa matrices [29].

There is also an (unphysical) ambiguity in RG functions due to possible unitary contributions [20, 30, 31] to field renormalization in models with flavor symmetries. In what follows, we assume a natural choice of Hermitian anomalous dimension and refer to recent work [32] for more details.

In the current study, we present for the first time complete results for all TS coefficients in gauge and Yukawa beta functions at four and three loops, respectively. Partial results are already available in the literature [21, 22, 25], and we want to emphasize the difference between our approach and methods used in previous studies. In the latter, given WCC, the unknown beta-function coefficients are fixed from *known* results for specific (usually physical) models. One of the most powerful constraints of this type is provided by the three-loop calculation in THDM [20].

On the contrary, we, for the first time, *design simple toy models* with specific gauge group structure, each giving constraints on different unknown coefficients. We use diagrams for TS provided in Supplementary Materials of Ref. [25] as a guide, but it is fair to say that we found the required models by trial and error. We use TSs implemented in the prominent code `RGBeta` [33] to speed up our investigation.

General four-dimensional QFT that covers most of possible phenomenological applications can be written in the

Here g , y_s , and y_u are gauge and Yukawa couplings of our interest, and λ_s , λ_{sh} and λ_h are the required scalar self-couplings.

We also study a gauge theory with $\mathcal{G} = SU(n_1) \times SU(n_2) \times SU(n_3)$ describing interactions of a Dirac fermion Ψ in fundamental representation of each factor in \mathcal{G} and three adjoint scalars ϕ_i , each charged only under one $SU(n_i)$:

$$\begin{aligned} \mathcal{L}_{\mathbf{M3}} = & -\frac{1}{2g_i^2} \text{Tr}(F_{\mu\nu}^i F_{\mu\nu}^i) + \text{Tr}[(D_\mu \phi_i)(D_\mu \phi_i)] + i\bar{\Psi} \gamma_\mu (D_\mu \Psi) \\ & - y_i [\bar{\Psi} \phi_i \Psi + \text{h.c.}] - \frac{\lambda_{ij}}{8} \text{Tr}(\phi_i \phi_i) \text{Tr}(\phi_j \phi_j) - \frac{\lambda_i}{24} \text{Tr}(\phi_i \phi_i \phi_i \phi_i), \end{aligned} \quad (6)$$

where summation over $i, j = 1, 3$ is assumed. The gauge and Yukawa couplings are denoted by g_i and y_i , respectively, and we have nine independent self-couplings in the model, λ_i and symmetric λ_{ij} .

Finally, we consider a $U(1)$ model with three Dirac fermions arranged as $\Psi = (\psi_1, \psi_2)$ and ψ . They interact with charged (h) and neutral (s) Higgs bosons via matrix $(y_1)_{ij}$, vector $(y_2)_i$, $(y_3)_i$, and scalar y_4 Yukawa couplings:

$$\begin{aligned} \mathcal{L}_{\mathbf{M4}} = & -\frac{1}{2g^2} (F_{\mu\nu} F_{\mu\nu}) + |D_\mu h|^2 + \frac{1}{2} (\partial_\mu s)^2 + i\bar{\Psi}_i \gamma_\mu (D_\mu \Psi_i) + i\bar{\psi} \gamma_\mu (D_\mu \psi) \\ & - [(y_1)_{ij} s \bar{\Psi}_i P_R \Psi_j + (y_2)_i h \bar{\Psi}_i P_R \psi + (y_3)_i h^* \bar{\psi} P_R \Psi_i + y_4 s \bar{\psi} P_R \psi + \text{h.c.}] - \frac{\lambda_s s^4}{24} - \frac{\lambda_{sh}}{2} s^2 (h^\dagger h) - \frac{\lambda_h}{2} (h^\dagger h)^2. \end{aligned} \quad (7)$$

The $U(1)$ charges satisfy $Q_h + Q_\psi = Q_\Psi$, and the sums run over $i = 1, 2$.

This choice of models is also motivated by the fact that we can easily implement them both in `RGBeta` [33] and `DIANA` [35]. We use the former to obtain the beta functions in terms of unknown coefficients, while the latter allows us to utilize our standard setup [13, 17] and compute required two- and three-point functions with `FORCER` [36].

To extract RG functions for gauge and Yukawa couplings, we need one-loop renormalization of the self couplings. We again use `RGBeta` to generate the necessary Z -factors.

FIXING COEFFICIENTS

With explicit results of calculation in models **M1** (4), **M2** (5), **M3** (6) and **M4** (7) at hand, we are in position to apply all the collected constraints and fix all beta-function coefficients. We summarize our procedure in the Table I, where we show how the number of unknowns $u_{\mathfrak{g}}$ and $u_{\mathfrak{h}}$ reduces after sequential application of available constraints. We start with WCC connecting $\mathfrak{g}^{(4)}$ and $\mathfrak{h}^{(3)}$ and interpret them as constraints on gauge beta-function coefficients. Applying further constraints, we obtain new relations (we denote the corresponding number by \mathbf{n}), and also a set of identities (the corresponding number is given by c) for cross-checking.

Type of the beta function	Gauge		Yukawa	
Number of equations and unknowns	$r = \mathbf{n} + c$	$u_{\mathfrak{g}}$	$r = \mathbf{n} + c$	$u_{\mathfrak{h}}$
Initial number of unknown coefficients	-	202	-	308
Weyl Consistency Conditions	128 +0	74	133 +0	175
Four-loop SM gauge beta functions	63 +84	11	-	-
Three-loop matrix Yukawa beta functions in the SM	-	-	128 +17	47
Three-loop matrix Yukawa beta functions in THDM	-	-	33 +213	14
Four-loop QCD beta function for general group	2 +11	9	-	-
$SU(n_1) \times SU(n_2)$ gauge theory (4) (M1)	5 +25	4	-	-
$SU(n)$ gauge theory (5) (M2)	2 +55	2	4 +76	10
$SU(n_1) \times SU(n_2) \times SU(n_3)$ gauge theory (6) (M3)	-	-	9 +89	1
$U(1)$ gauge theory (7) (M4)	-	-	1 +199	0
Constraints from symmetric T_{IJ}	2 +8	0	-	-
Final number of unknowns	-	0	-	0

TABLE I. Reduction of the number of unknown coefficients $u_{\mathfrak{g}}$ and $u_{\mathfrak{h}}$ after sequential application of constraints. Here $r = \mathbf{n} + c$ is the rank of the system without any previous constraints included, \mathbf{n} corresponds to new independent constraints, and c relations are automatically satisfied due to previous steps.

After matching template expressions with our toy models we are left with two unknowns in the gauge sector. In Ref. [25] the authors conjectured that T_{IJ} tensor entering WCC can be symmetric. This provides additional 10 constraints. We use 2 of them to constrain the remaining coefficients. The other 8 equations become identities and, thus, verify the assumption on T_{IJ} . It is worth noting that we use models **M1-M3** to constrain all TSs but the difference $\eta_{172}^{(3)} - \eta_{173}^{(3)} \equiv 2\delta$ in Yukawa beta function. To deal with δ , we developed a special model **M4**, which explicitly confirms our initial guess that $\delta = 0$.

With this procedure we fix all coefficients in gauge and Yukawa beta functions at four and three loops, respectively.

RESULTS AND DISCUSSION

We combine the constraints from WCC [23] and our explicit computations in toy models to fix all the coefficients in the ansatz for the 4-3-2 ordering given in Ref. [25]. As an application of our general expressions, we derive all four-loop gauge beta functions in the SM extension with the arbitrary number n_d of Higgs doublets (NHDM). We keep matrix Yukawa couplings, and by setting $n_d = 1$ or $n_d = 2$, we obtain the SM and THDM result. Direct computation in such a scenario would be very complicated.

Let us return to the ambiguities in theories possessing a flavor symmetry. In Refs. [25, 30, 32], for the coupling y_{aij} a flavor-improved, i.e unambiguous, version B_{aij} of the Yukawa beta function β_{aij} is introduced

$$B_{aij}^{(3)} = \beta_{aij}^{(3)} - S_{ik}^{(3)} y_{akj} - S_{jk}^{(3)} y_{aik} - S_{ab}^{(3)} y_{bij}. \quad (8)$$

Here $S_{ij}^{(3)}$ and $S_{ab}^{(3)}$ are three-loop quantities that can be represented as linear combinations of antisymmetric two-point TS for fermions and scalars, respectively [25, 30]. There are six coefficients $f_{1-6}^{(3)}$ entering $S_{ij}^{(3)}$ and three coefficients $s_{1-3}^{(3)}$ entering $S_{ab}^{(3)}$. All but one ($f_3^{(3)}$) numerical coefficients are predicted from WCC² in terms of $\mathbf{g}_i^{(4)}$ and $\eta_i^{(3)}$ computed in this Letter. The authors of Ref. [32] calculated $f_4^{(3)} = -3/8$ and $f_5^{(3)} = -5/16$. Given our results, we provide full set of three-loop corrections to S_{ij} and S_{ab} :

$$f_1^{(3)} = 0, \quad f_2^{(3)} = \frac{29}{8} - 3\zeta_3, \quad f_3^{(3)} = \frac{21}{8} - 3\zeta_3, \quad f_4^{(3)} = -\frac{3}{8}, \quad f_5^{(3)} = -\frac{5}{16}, \quad f_6^{(3)} = -\frac{7}{16}. \quad (9)$$

$$s_1^{(3)} = \frac{7}{2} - 6\zeta_3, \quad s_2^{(3)} = \frac{5}{8}, \quad s_3^{(3)} = -\frac{3}{4}. \quad (10)$$

These coefficients can be tested by direct calculations along the lines of Refs. [30, 32].

CONCLUSION

The calculation of four-loop Gauge and three-loop Yukawa beta functions performed in this Letter complement recent six-loop results in general pure scalar theory [37], and represents the most advanced achievement in this field. The obtained TS coefficients can be incorporated into modern computer codes, giving access to a new precision level for model building. The dummy-field method (see, e.g., Ref. [38]) applied to our results provides us with scale dependence of such important quantities as fermion mass matrices.

We make all our results, including the TS coefficients and four-loop gauge-coupling beta functions in the SM, THDM, and NHDM, available as Supplementary Material. We also provide a modified version of the RGBeta package with all our findings included [39].

We thank A.E. Thomsen for correspondence regarding RGBeta and Ref. [25]. We are grateful to the Joint Institute for Nuclear Research for letting us use their supercomputer ‘‘Govorun.’’ The work of A.P. is supported by the Foundation for the Advancement of Theoretical Physics and Mathematics ‘‘BASIS.’’ The work of A.B. is supported by the Grant of the Russian Federation Government, Agreement No. 14.W03.31.0026.

* bednya@theor.jinr.ru

² See Supplementary Material of Ref. [25].

† pikelner@theor.jinr.ru

- [1] Marie E. Machacek and Michael T. Vaughn, “Two Loop Renormalization Group Equations in a General Quantum Field Theory. 2. Yukawa Couplings,” *Nucl.Phys.* **B236**, 221 (1984).
- [2] Marie E. Machacek and Michael T. Vaughn, “Two Loop Renormalization Group Equations in a General Quantum Field Theory. 1. Wave Function Renormalization,” *Nucl.Phys.* **B222**, 83 (1983).
- [3] Marie E. Machacek and Michael T. Vaughn, “Two Loop Renormalization Group Equations in a General Quantum Field Theory. 3. Scalar Quartic Couplings,” *Nucl.Phys.* **B249**, 70 (1985).
- [4] Ming-xing Luo, Hua-wen Wang, and Yong Xiao, “Two loop renormalization group equations in general gauge field theories,” *Phys. Rev.* **D67**, 065019 (2003), [arXiv:hep-ph/0211440](https://arxiv.org/abs/hep-ph/0211440) [hep-ph].
- [5] A. V. Bednyakov, “On three-loop RGE for the Higgs sector of 2HDM,” *JHEP* **11**, 154 (2018), [arXiv:1809.04527](https://arxiv.org/abs/1809.04527) [hep-ph].
- [6] Ingo Schienbein, Florian Staub, Tom Steudtner, and Kseniia Svirina, “Revisiting RGEs for general gauge theories,” *Nucl. Phys.* **B939**, 1–48 (2019), [arXiv:1809.06797](https://arxiv.org/abs/1809.06797) [hep-ph].
- [7] Lohan Sartore, “General RGEs for dimensionful couplings in the $\overline{\text{MS}}$ scheme,” *Phys. Rev. D* **102**, 076002 (2020), [arXiv:2006.12307](https://arxiv.org/abs/2006.12307) [hep-ph].
- [8] Georges Aad *et al.* (ATLAS), “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,” *Phys. Lett.* **B716**, 1–29 (2012), [arXiv:1207.7214](https://arxiv.org/abs/1207.7214) [hep-ex].
- [9] Serguei Chatrchyan *et al.* (CMS), “Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC,” *Phys. Lett.* **B716**, 30–61 (2012), [arXiv:1207.7235](https://arxiv.org/abs/1207.7235) [hep-ex].
- [10] A. V. Bednyakov, B. A. Kniehl, A. F. Pikelner, and O. L. Veretin, “Stability of the Electroweak Vacuum: Gauge Independence and Advanced Precision,” *Phys. Rev. Lett.* **115**, 201802 (2015), [arXiv:1507.08833](https://arxiv.org/abs/1507.08833) [hep-ph].
- [11] A. G. M. Pickering, J. A. Gracey, and D. R. T. Jones, “Three loop gauge beta function for the most general single gauge coupling theory,” *Phys. Lett.* **B510**, 347–354 (2001), [Erratum: *Phys. Lett.* B535,377(2002)], [arXiv:hep-ph/0104247](https://arxiv.org/abs/hep-ph/0104247) [hep-ph].
- [12] Luminita N. Mihaila, Jens Salomon, and Matthias Steinhauser, “Gauge Coupling Beta Functions in the Standard Model to Three Loops,” *Phys.Rev.Lett.* **108**, 151602 (2012), [arXiv:1201.5868](https://arxiv.org/abs/1201.5868) [hep-ph].
- [13] A.V. Bednyakov, A.F. Pikelner, and V.N. Velizhanin, “Anomalous dimensions of gauge fields and gauge coupling beta-functions in the Standard Model at three loops,” *JHEP* **1301**, 017 (2013), [arXiv:1210.6873](https://arxiv.org/abs/1210.6873) [hep-ph].
- [14] A.V. Bednyakov, A.F. Pikelner, and V.N. Velizhanin, “Yukawa coupling beta-functions in the Standard Model at three loops,” *Phys.Lett.* **B722**, 336–340 (2013), [arXiv:1212.6829](https://arxiv.org/abs/1212.6829).
- [15] K.G. Chetyrkin and M.F. Zoller, “ β -function for the Higgs self-interaction in the Standard Model at three-loop level,” *JHEP* **1304**, 091 (2013), [arXiv:1303.2890](https://arxiv.org/abs/1303.2890) [hep-ph].
- [16] A.V. Bednyakov, A.F. Pikelner, and V.N. Velizhanin, “Higgs self-coupling beta-function in the Standard Model at three loops,” *Nucl.Phys.* **B875**, 552–565 (2013), [arXiv:1303.4364](https://arxiv.org/abs/1303.4364).
- [17] A. V. Bednyakov and A. F. Pikelner, “Four-loop strong coupling beta-function in the Standard Model,” *Phys. Lett.* **B762**, 151–156 (2016), [arXiv:1508.02680](https://arxiv.org/abs/1508.02680) [hep-ph].
- [18] M. F. Zoller, “Top-Yukawa effects on the β -function of the strong coupling in the SM at four-loop level,” *JHEP* **02**, 095 (2016), [arXiv:1508.03624](https://arxiv.org/abs/1508.03624) [hep-ph].
- [19] K. G. Chetyrkin and M. F. Zoller, “Leading QCD-induced four-loop contributions to the β -function of the Higgs self-coupling in the SM and vacuum stability,” *JHEP* **06**, 175 (2016), [arXiv:1604.00853](https://arxiv.org/abs/1604.00853) [hep-ph].
- [20] Florian Herren, Luminita Mihaila, and Matthias Steinhauser, “Gauge and Yukawa coupling beta functions of two-Higgs-doublet models to three-loop order,” *Phys. Rev.* **D97**, 015016 (2018), [arXiv:1712.06614](https://arxiv.org/abs/1712.06614) [hep-ph].
- [21] Tom Steudtner, “General scalar renormalisation group equations at three-loop order,” *JHEP* **12**, 012 (2020), [arXiv:2007.06591](https://arxiv.org/abs/2007.06591) [hep-th].
- [22] Tom Steudtner, “Towards general scalar-Yukawa renormalisation group equations at three-loop order,” *JHEP* **05**, 060 (2021), [arXiv:2101.05823](https://arxiv.org/abs/2101.05823) [hep-th].
- [23] I. Jack and H. Osborn, “Constraints on RG Flow for Four Dimensional Quantum Field Theories,” *Nucl. Phys. B* **883**, 425–500 (2014), [arXiv:1312.0428](https://arxiv.org/abs/1312.0428) [hep-th].
- [24] Oleg Antipin, Marc Gillioz, Jens Krog, Esben Mølgaard, and Francesco Sannino, “Standard Model Vacuum Stability and Weyl Consistency Conditions,” *JHEP* **08**, 034 (2013), [arXiv:1306.3234](https://arxiv.org/abs/1306.3234) [hep-ph].
- [25] Colin Poole and Anders Eller Thomsen, “Constraints on 3- and 4-loop β -functions in a general four-dimensional Quantum Field Theory,” *JHEP* **09**, 055 (2019), [arXiv:1906.04625](https://arxiv.org/abs/1906.04625) [hep-th].
- [26] C. Poole and A. E. Thomsen, “Weyl Consistency Conditions and γ_5 ,” *Phys. Rev. Lett.* **123**, 041602 (2019), [arXiv:1901.02749](https://arxiv.org/abs/1901.02749) [hep-th].
- [27] F. Jegerlehner, “Facts of life with γ_5 ,” *Eur. Phys. J.* **C18**, 673–679 (2001), [arXiv:hep-th/0005255](https://arxiv.org/abs/hep-th/0005255) [hep-th].
- [28] K. G. Chetyrkin and M. F. Zoller, “Three-loop β -functions for top-Yukawa and the Higgs self-interaction in the Standard Model,” *JHEP* **06**, 033 (2012), [arXiv:1205.2892](https://arxiv.org/abs/1205.2892) [hep-ph].
- [29] Joshua Davies, Florian Herren, Colin Poole, Matthias Steinhauser, and Anders Eller Thomsen, “Gauge Coupling β Functions to Four-Loop Order in the Standard Model,” *Phys. Rev. Lett.* **124**, 071803 (2020), [arXiv:1912.07624](https://arxiv.org/abs/1912.07624) [hep-ph].
- [30] Jean-Francois Fortin, Benjamin Grinstein, and Andreas Stergiou, “Limit Cycles and Conformal Invariance,” *JHEP* **01**, 184 (2013), [arXiv:1208.3674](https://arxiv.org/abs/1208.3674) [hep-th].
- [31] A. V. Bednyakov, A. F. Pikelner, and V. N. Velizhanin, “Three-loop SM beta-functions for matrix Yukawa couplings,” *Phys. Lett.* **B737**, 129–134 (2014), [arXiv:1406.7171](https://arxiv.org/abs/1406.7171) [hep-ph].
- [32] Florian Herren and Anders Eller Thomsen, “On ambiguities and divergences in perturbative renormalization group functions,” *JHEP* **06**, 116 (2021), [arXiv:2104.07037](https://arxiv.org/abs/2104.07037) [hep-th].
- [33] Anders Eller Thomsen, “Introducing RGBeta: a Mathematica package for the evaluation of renormalization group β -

- functions,” *Eur. Phys. J. C* **81**, 408 (2021), [arXiv:2101.08265 \[hep-ph\]](#).
- [34] Predrag Cvitanovic, “Group theory for Feynman diagrams in non-Abelian gauge theories,” *Phys. Rev.* **D14**, 1536–1553 (1976).
- [35] M. Tentyukov and J. Fleischer, “A Feynman diagram analyzer DIANA,” *Comput. Phys. Commun.* **132**, 124–141 (2000), [arXiv:hep-ph/9904258 \[hep-ph\]](#).
- [36] B. Ruijl, T. Ueda, and J. A. M. Vermaseren, “Forcer, a FORM program for the parametric reduction of four-loop massless propagator diagrams,” *Comput. Phys. Commun.* **253**, 107198 (2020), [arXiv:1704.06650 \[hep-ph\]](#).
- [37] A. Bednyakov and A. Pikelner, “Six-loop beta functions in general scalar theory,” *JHEP* **04**, 233 (2021), [arXiv:2102.12832 \[hep-ph\]](#).
- [38] Florian Staub, “SARAH 4 : A tool for (not only SUSY) model builders,” *Comput. Phys. Commun.* **185**, 1773–1790 (2014), [arXiv:1309.7223 \[hep-ph\]](#).
- [39] <https://github.com/apik/RGBeta>.