

ON DIMENSION THEORY OF SUPERMODULES, SUPER-RINGS AND SUPERSCHEMES

A. N. ZUBKOV AND P.S.KOLESNIKOV

ABSTRACT. We introduce the notion of Krull super-dimension of supermodules over certain super-commutative Noetherian super-rings. We investigate how this notion relates to the notion of odd regular sequence from [7] and how it behaves with respect to the transition to the graded and bigraded supermodules and super-rings associated with the original ones. We also apply these results to the super-dimension theory of superschemes of finite type and their morphisms.

INTRODUCTION

The notion of Krull dimension plays crucial role in the theory of commutative rings and in the algebraic geometry. Recall that the Krull dimension $\text{Kdim}(R)$ of a commutative ring R is the supremum of heights of the prime ideals of R . If $\text{Kdim}(R) < \infty$, then $\text{Kdim}(R)$ is just the largest nonnegative integer n such that there is a chain $\mathfrak{p}_0 \subseteq \dots \subseteq \mathfrak{p}_n$ of pairwise different prime ideals. Geometrically, $\text{Kdim}(R)$ is nothing else but the dimension of the affine scheme $\text{Spec}(R)$ (cf. [2], Chapter II, Example 3.2.7).

In what follows all super-rings are supposed to be super-commutative, unless stated otherwise. If R is a super-ring, then each prime superideal \mathfrak{P} of R has a form $\mathfrak{p} \oplus R_1$, where $\mathfrak{p} = \mathfrak{P}_0$ is a prime ideal of the ring R_0 . In other words, the prime spectrum of R coincides with the prime spectrum of R_0 , and it does not detect the *odd dimension* of the affine superscheme $\text{SSpec}(R)$. Therefore, a relevant notion of Krull super-dimension of a super-ring has to include an *odd component* also. Such a notion has been recently suggested in [4]. Let R be a super-ring with $\text{Kdim}(R_0)$ to be finite. Then a collection of odd elements y_1, \dots, y_s is called a *system of odd parameters* of R if

$$\text{Kdim}(R_0/\text{Ann}_{R_0}(y_1 \dots y_s)) = \text{Kdim}(R_0).$$

In other words, the elements y_1, \dots, y_s form a system of odd parameters of R if there is a longest prime chain $\mathfrak{p}_0 \subseteq \dots \subseteq \mathfrak{p}_n$ in R such that $\text{Ann}_{R_0}(y_1 \dots y_s) \subseteq \mathfrak{p}_0$. Finally, a *Krull super-dimension* $\text{Ksdim}(R)$ of R is a couple $n | l$, where $n = \text{Kdim}(R_0)$ and l is the supremum of lengths of all systems of odd parameters of R . We also use the notations $\text{Ksdim}_0(R)$ and $\text{Ksdim}_1(R)$ for n and l respectively.

In general, the odd super-dimension l can be infinite. But if R is Noetherian, then l is always finite.

The purpose of this work is to define the notion of super-dimension of a supermodule over a super-ring, based on the above notion of Krull super-dimension, and investigate its properties and several applications.

The article is organized as follows. In the first section we describe Artinian super-rings. In the second section the notion of super-dimension of supermodules over

certain Noetherian super-rings is introduced and some its elementary properties are studied. We also discuss how the odd parameters relate to odd regular elements in the sense of [7]. Indeed, one can easily see that any system of odd M -regular elements of a supermodule M form a system of odd parameters of M . It was incorrectly stated in Lemma 4.2(3), [4], that any system of odd regular elements of a super-ring R can be included into a longest system of odd parameters. In the second and third sections we construct two counterexamples for this statement in the categories of supermodules and super-rings respectively. Moreover, in the category of super-rings we develop a fragment of a *superized Hochschild cohomology* theory, with the help of which the second counterexample is constructed.

Let R be a (not necessary Noetherian) super-ring and I be a superideal of R . We associate with R a graded super-ring

$$\mathbf{gr}_I(R) = \bigoplus_{k \geq 0} I^k / I^{k+1}$$

and a bigraded super-ring

$$\mathbf{bgr}_I(R) = \bigoplus_{k, l \geq 0} I_0^k I_1^l R / (I_0^{k+1} I_1^l R + I_0^k I_1^{l+1} R).$$

These super-rings are Noetherian, provided R is. If $\text{Ksdim}_0(R) < \infty$, then both $\mathbf{gr}_I(R)$ and $\mathbf{bgr}_I(R)$ satisfy this condition as well.

Similarly, with any R -supermodule M we associate a graded $\mathbf{gr}_I(R)$ -supermodule

$$\mathbf{gr}_I(M) = \bigoplus_{k \geq 0} I^k M / I^{k+1} M$$

and a bigraded $\mathbf{bgr}_I(R)$ -supermodule

$$\mathbf{bgr}_I(M) = \bigoplus_{k, l \geq 0} I_0^k I_1^l M / (I_0^{k+1} I_1^l M + I_0^k I_1^{l+1} M).$$

Let R be a Noetherian super-ring with $\text{Ksdim}_0(R) < \infty$ and I be a superideal of R that is contained in the radical of R . In the fifth section we prove that for any finitely generated R -supermodule M there hold $\text{Ksdim}_0(M) = \text{Ksdim}_0(\mathbf{gr}_I(M))$ and $\text{Ksdim}_1(M) \geq \text{Ksdim}_1(\mathbf{gr}_I(M))$.

The latter inequality can be strict in general. However, if $I = I_R = RR_1$, then $\text{Ksdim}(M) = \text{Ksdim}(\mathbf{gr}_I(M))$. In particular, if the elements

$$z_1, \dots, z_t \in I_R / I_R^2 \simeq R_1 / R_1^3$$

form a longest system of odd parameters of the supermodule $\mathbf{gr}_{I_R}(M)$, then their representatives $y_1, \dots, y_t \in R$ form a longest system of odd parameters of M .

In the six section we develop the dimension theory of certain local super-rings. More precisely, let R be a local super-ring with the maximal superideal \mathfrak{M} . Assume that R contains a coefficient field $K \simeq R_0 / \mathfrak{m}$. Let \mathfrak{N} be a \mathfrak{M} -primary superideal of R such that $(\mathfrak{n} + R_1^2) / R_1^2$ is generated by the parameters of $\overline{R} = R / I_R \simeq R_0 / R_1^2$. Then $\text{Ksdim}_0(\mathbf{bgr}_{\mathfrak{N}}(R)) = \text{Ksdim}_0(R)$ but $\text{Ksdim}_1(\mathbf{bgr}_{\mathfrak{N}}(R)) \leq \text{Ksdim}_1(R)$. However, if \mathfrak{N} has a form $\mathfrak{n} \oplus R_1$, then $\text{Ksdim}(\mathbf{bgr}_{\mathfrak{N}}(R)) = \text{Ksdim}(R)$.

Let B denote $\mathbf{bgr}_{\mathfrak{N}}(R)$ and for each couple of nonnegative integers k, l let $B(k, l)$ denote the corresponding homogeneous component of the bigraded super-ring B . We can associate with B a *Hilbert polynomial* $g_{\mathfrak{N}}(x, y) = \sum_{l \geq 0} g_l(x) y^l$ such that

$$\sum_{0 \leq t \leq k} \dim B(t, l) = g_l(k)$$

for all sufficiently large k and for any $l \geq 0$. Then the degree of each polynomial $g_l(x)$ is at most $d = \text{Ksdim}_0(R)$ and if $\mathfrak{R}_1 = R_1$, then

$$\text{Ksdim}_1(R) = \max\{l \mid \text{the degree of } g_l(x) \text{ is equal to } d\}.$$

In the fifth and seventh sections we discuss some applications of the above results for the dimension theory of superschemes. For example, we prove that for any flat morphism $f : X \rightarrow Y$ of irreducible superschemes of finite type and for any nonsingular point $y \in Y^e$, the odd dimension of each irreducible component of the fiber X_y is at most $\text{sdim}_1(X) - \text{sdim}_1(Y)$, contrary to the well known fact that its even dimension is equal to $\text{sdim}_0(X) - \text{sdim}_0(Y)$. Moreover, this inequality can be strict in general.

1. ARTINIAN SUPER-RINGS

As in the classical setting, a super-ring R is called *Artinian* if R satisfies the descending chain condition for super-ideals (DCC).

Lemma 1.1. *A super-ring R is Artinian if and only if R_0 is an Artinian ring and R_0 -module R_1 is Artinian as well.*

Proof. Use the same arguments as in Lemma 1.4, [4]. □

The following proposition superizes Proposition (2.C), [3].

Proposition 1.2. *A super-ring R is Artinian if and only if it has finite length as R -supermodule.*

Proof. The part "if" is obvious.

Assume that R is Artinian. By Lemma 1.1 R_0 is Artinian, hence R_0 (and R as well) has only finite number of maximal ideals, say $\mathfrak{m}_1, \dots, \mathfrak{m}_t$ (correspondingly, $\mathfrak{M}_1 = \mathfrak{m}_1 \oplus R_1, \dots, \mathfrak{M}_t = \mathfrak{m}_t \oplus R_1$). Moreover, $\text{rad}(R_0) = \bigcap_{1 \leq i \leq t} \mathfrak{m}_i = \mathfrak{m}_1 \dots \mathfrak{m}_t$ is nilpotent (cf. Proposition (2.C), [3]). On the other hand, the ideal R_1^2 is locally nilpotent, hence $R_1^2 \subseteq \text{rad}(R_0)$ and R_1^2 is also nilpotent, say $R_1^{2k} = 0$ for sufficiently large k .

Set $I = \mathfrak{M}_1 \dots \mathfrak{M}_t$. Since $I \subseteq \text{rad}(R_0) + I_R$ and $I_R^{2k} = 0$, we have $I^{2k+l} = 0$, provided $\text{rad}(R_0)^l = 0$. Set $N = 2k + l$. Now, one can construct a chain of super-ideals

$$\begin{aligned} R \supseteq \mathfrak{M}_1 &\supseteq \mathfrak{M}_1 \mathfrak{M}_2 \supseteq \dots \supseteq \mathfrak{M}_1 \dots \mathfrak{M}_{t-1} \supseteq I \supseteq \\ I \mathfrak{M}_1 &\supseteq I \mathfrak{M}_1 \mathfrak{M}_2 \supseteq \dots \supseteq I \mathfrak{M}_1 \dots \mathfrak{M}_{t-1} \supseteq I^2 \supseteq \dots \supseteq I^N = 0, \end{aligned}$$

where each quotient of the chain is a vector (super)space over a field R/\mathfrak{M}_i for some i , hence a finite dimensional vector (super)space. Proposition is proven. □

Corollary 1.3. *A super-ring R is Artinian if and only if R is a Noetherian super-ring of Krull super-dimension $0|n$ for some non-negative integer n .*

Proof. If R is Artinian, then Proposition 1.2 obviously implies that R is Noetherian. Moreover, by Corollary (2.C), [3], $\text{Ksdim}_0(R) = \text{Kdim}(R_0) = 0$. Conversely, the same corollary and Lemma 1.4, [4], infer that if R is Noetherian and $\text{Kdim}(R_0) = 0$, then R_0 is Artinian and I_R is nilpotent. Repeating the arguments of Proposition 1.2, we complete the proof. □

2. SUPER-DIMENSION OF SUPERMODULES

Let B be a commutative ring and let N be a B -module. Recall that the *dimension* of N is defined by $\dim(N) = \text{Kdim}(B/\text{Ann}_B(N))$.

From now on we assume that R is a Noetherian super-ring with $\text{Kdim}(R_0) < \infty$, unless stated otherwise. For any set of odd elements $y_1, \dots, y_s \in R_1$ and any subset $L \subseteq \underline{s} = \{1, 2, \dots, s\}$, let y^L denote the product $\prod_{i \in L} y_i$.

If M is an R -supermodule, then $\dim(M)$ is always regarded as the dimension of M as the R_0 -module. Further, the *super-dimension* of M is defined as

$$\text{sdim}(M) = \text{Ksdim}(R/\text{Ann}_R(M)).$$

More precisely, $\text{sdim}(M)$ is a vector with two coordinates $\text{sdim}_0(M)$ and $\text{sdim}_1(M)$, where $\text{sdim}_0(M) = \dim(M)$. In its turn, $\text{sdim}_1(M)$ is equal to the largest positive integer l such that there are the elements $y_1, \dots, y_l \in R_1$, which satisfy

$$(\star) \quad \dim(y^l M) = \dim(M),$$

otherwise $\text{sdim}_1(M) = 0$.

In what follows, any collection of odd elements y_1, \dots, y_l , which satisfies the condition (\star) , is called a *system of odd parameters* of M . In other words, the elements $y_1, \dots, y_l \in R_1$ form a system of odd parameters of M if and only if there is a prime chain $\mathfrak{p}_0 \subseteq \dots \subseteq \mathfrak{p}_l$ in R_0 , such that $\text{Ann}_{R_0}(y^l M) \subseteq \mathfrak{p}_0$ and

$$\mathfrak{p}_0/\text{Ann}_{R_0}(M) \subseteq \dots \subseteq \mathfrak{p}_l/\text{Ann}_{R_0}(M)$$

is a longest prime chain in $R_0/\text{Ann}_{R_0}(M)$. We say that the above system of odd parameters subordinates the prime ideal \mathfrak{p}_0 .

For example, it is obvious that the Krull super-dimension of a super-ring R coincides with the super-dimension of R , regarded as a left R -supermodule. Moreover, any system of odd parameters of the super-ring R is the system of odd parameters of the left R -supermodule R and vice versa.

Proposition 2.1. *Choose a system of generators of R_0 -module R_1 , say y_1, \dots, y_d . Then the following conditions are equivalent :*

- (1) *There is a system of odd parameters of M of cardinality $l \geq 1$;*
- (2) *For some $1 \leq i_1 < \dots < i_l \leq d$ the elements y_{i_1}, \dots, y_{i_l} form a system of odd parameters of M ;*
- (3) $\dim(R_1^l M) = \dim(M)$.

In particular, we have

$$\text{sdim}_1(M) = \max\{l \mid \dim(R_1^l M) = \dim(M)\}.$$

Proof. It is obvious modification of the proof of Proposition 4.1, [4]. \square

Lemma 2.2. *If M is a finitely generated R -supermodule, then the following conditions are equivalent :*

- (1) *M is a supermodule of finite length.*
- (2) *$R/\text{Ann}_R(M)$ is an Artinian super-ring.*
- (3) $\text{sdim}(M) = 0 \mid n$ *for some nonnegative integer n .*
- (4) $\dim(M) = 0$.

Proof. The equivalences $(2) \leftrightarrow (3)$ and $(2) \leftrightarrow (4)$ immediately follow by Corollary 1.3. If M is an R -supermodule of finite length, then any its super-submodule is of finite length. Without loss of a generality, one can assume that M is generated

by homogeneous elements m_1, \dots, m_t . Then $\text{Ann}_R(M) = \bigcap_{1 \leq i \leq t} \text{Ann}_R(m_i)$ and $R/\text{Ann}_R(M)$ is a subdirect product of Artinian super-rings $R/\text{Ann}_R(m_i)$, hence Artinian as well. Thus (1) implies (2). Conversely, there is a natural surjective supermodule morphism $(R/\text{Ann}_R(M))^{\oplus t} \rightarrow M$. Therefore, M is an R -supermodule of finite length, provided $R/\text{Ann}_R(M)$ is an Artinian super-ring. Lemma is proven. \square

Lemma 2.3. *Let R be an Artinian super-ring. Then for any non zero R -supermodule M there is*

$$\text{sdim}_1(M) = \max\{l \mid R_1^l M \neq 0\}.$$

Proof. By Corollary (2.C), [3], R_0 (respectively, R) has finitely many prime ideals (respectively, super-ideals) all of which are maximal. Since $\dim(M) = 0$, a nonempty collection of odd elements z_1, \dots, z_l form a system of odd parameters of M if and only if $\text{Ann}_{R_0}(z^l M)$ is a proper ideal of R_0 if and only if $z^l M \neq 0$. Proposition 2.1 infers the statement. \square

Lemma 2.4. *We have the following :*

- (1) *If I is an nilpotent superideal of R , then $\text{sdim}_0(M/IM) = \text{sdim}_0(M)$.*
- (2) *If a super-submodule N of M contains yM , where y belongs to a (not necessary longest) system of odd parameters, then $\text{sdim}_0(N) = \text{sdim}_0(M)$.*

Proof. (1) Assume that $I^l = 0$ for sufficiently large nonnegative integer l . If an even element r satisfies $rM \subseteq IM$, then $r^l M = 0$, that is $\text{Ann}_{R_0}(M/IM)$ is contained in the radical of $\text{Ann}_{R_0}(M)$.

- (2) If $y = y_1$ and y_1, \dots, y_s is a system of odd parameters, then

$$\text{Ann}_{R_0}(M) \subseteq \text{Ann}_{R_0}(N) \subseteq \text{Ann}_{R_0}(yM) \subseteq \text{Ann}_{R_0}(y^s M).$$

\square

Lemma 2.5. *Let N be a super-submodule of M , such that $\text{sdim}_0(N) = \text{sdim}_0(M)$. Then $\text{sdim}_1(N) \leq \text{sdim}_1(M)$.*

Proof. Assume that y_1, \dots, y_s is a system of odd parameters of N . Then

$$\text{Ann}_{R_0}(M) \subseteq \text{Ann}_{R_0}(y^s M) \subseteq \text{Ann}_{R_0}(y^s N)$$

implies that y_1, \dots, y_s is a system of odd parameters of M . \square

Recall that an odd element $r \in R_1$ is M -regular if the kernel of the odd R -linear map $m \mapsto rm, m \in M$, coincides with rM (cf. [7], (2.1.1)). A sequence of odd elements r_1, \dots, r_t is called M -regular if each r_i is M -regular modulo $(Rr_1 + \dots + Rr_{i-1})M$.

Proposition 2.6. *The following statements hold :*

- (1) *Let y_1, \dots, y_t be a fragment of a longest system of odd parameters of M . Set $I = Ry_1 + \dots + Ry_t$. Then*

$$\text{sdim}_0(M) = \text{sdim}_0(IM) = \text{sdim}_0(M/IM)$$

and

$$\text{sdim}_1(M) \geq \text{sdim}_1(IM) \geq \text{sdim}_1(M) - 1, \text{sdim}_1(M/IM) \geq \text{sdim}_1(M) - t.$$

(2) If y_1, \dots, y_t is an odd M -regular sequence, then $\text{sdim}_1(M) \geq t$ and

$$\text{sdim}_1(M/IM) = \text{sdim}_1(y^{\underline{t}}M) \leq \text{sdim}_1(M) - t.$$

Moreover, $\text{sdim}_1(M/IM)$ is equal to $\text{sdim}_1(M) - t$ if and only if y_1, \dots, y_t can be extended to a longest system of odd parameters of M .

Proof. (1) Lemma 2.4 implies the first two equations. If y_1, \dots, y_l is a longest system of odd parameters of M , that extends y_1, \dots, y_t , then y_2, \dots, y_l is obviously a system of odd parameters of y_1M . By Lemma 2.5 we have

$$\text{sdim}_1(M) \geq \text{sdim}_1(IM) \geq \text{sdim}_1(y_1M) \geq \text{sdim}_1(M) - 1.$$

Further, we obviously have

$$\text{Ann}_{R_0}(M) \subseteq \text{Ann}_{R_0}((y^{\underline{t}}M + IM)/IM) \subseteq \text{Ann}_{R_0}(y^{\underline{t}}M).$$

In other words, y_{t+1}, \dots, y_l form a system of odd parameters for M/IM .

(2) The map $m \mapsto y^{\underline{t}}m, m \in M$, induces an isomorphism $M/IM \simeq y^{\underline{t}}M$ of R -supermodules, which has parity $|y^{\underline{t}}|$ (see [7], Corollary 3.1.2). In particular, $\text{sdim}_1(M/IM) = \text{sdim}_1(y^{\underline{t}}M)$. Moreover, if $ry^{\underline{t}}M = 0$, then $rM \subseteq IM$ and $r^{t+1}M = 0$. Therefore, y_1, \dots, y_t form a system of odd parameters of M and $\text{sdim}_1(M) \geq t$.

If y_{t+1}, \dots, y_l is a longest system of odd parameters for M/IM , then y_1, \dots, y_l is a system of odd parameters for M , hence $l \leq s$. Indeed, one easily sees that

$$\text{Ann}_{R_0}(y^{\underline{l}}M) = \text{Ann}_{R_0}((y^{\underline{l}}M + IM)/IM).$$

Combining with (1), we obtain that $l = s$ if and only if y_1, \dots, y_t can be extended to a longest system of odd parameters of M . \square

Remark 2.7. In [4], Lemma 4.2 (3) erroneously states that any odd regular sequence can be included into the longest system of parameters. In the above proposition we give the correct formulation of this statement.

Let A be a commutative Artinian ring. In the rest part of this section we construct a supermodule M over the polynomial super-ring $R = A[Z_1, Z_2, Z_3, Y]$ with free odd generators Z_1, Z_2, Z_3, Y , such that Y is an odd M -regular element, but $\text{sdim}_1(M/YM) < \text{sdim}_1(M) - 1$. In particular, Y can not be included in any system of odd parameters of M .

Let V be an A -supermodule of finite length. Set $M = V \oplus \Pi V$. To define a structure of R -supermodule on M one needs to define an action of the free generators of R on M by some odd A -linear endomorphisms of M , which supercommute to each other.

Set $Y(v) = \Pi v, Y(\Pi v) = 0, v \in V$. Thus it is obvious that Y is M -regular.

Lemma 2.8. Assume that an action of R on M is defined so that

$$Z_i Z_j M \subseteq YM, 1 \leq i, j \leq 3 \text{ and } Z_1 Z_2 Z_3 M \neq 0.$$

Then $\text{sdim}_1(M) = 3$, but $\text{sdim}_1(M/YM) \leq 1$.

Proof. Since R is Artinian, Lemma 2.3 implies that $\text{sdim}_1(M) = 3$. In fact, $R_1^3 M \neq 0$ but $R_1^4 M = Y Z_1 Z_2 Z_3 M \subseteq Y Z_1 (YM) = 0$. Finally, Y can not be included in any system of odd parameters of length 3, since the product of the elements of such system belongs to the superideal $\sum_{1 \leq i < j \leq 3} R Y Z_i Z_j$, which acts trivially on M . Proposition 2.6 (2) concludes the proof. \square

The action of each Z_i can be expressed as $Z_i(v) = \phi_i(v) + \Pi\psi_i(v)$, where ϕ_i, ψ_i are odd and even endomorphisms of A -supermodule V respectively. Then $Z_i(\Pi v) = (Z_i Y)(v) = -Y(Z_i(v)) = -Y(\phi_i(v)) = -\Pi\phi_i(v)$.

Lemma 2.9. *The odd endomorphisms of M , induced by the free generators of R , supercommute to each other and satisfy the conditions of Lemma 2.8 if and only if the following equations and one inequality hold :*

- (1) $\phi_i\psi_i - \psi_i\phi_i = 0, \phi_i\phi_j = 0, 1 \leq i, j \leq 3$;
- (2) $\psi_i\phi_j - \phi_i\psi_j - \phi_j\psi_i + \psi_j\phi_i = 0, 1 \leq i \neq j \leq 3$;
- (3) $\phi_1\psi_2\phi_3 \neq 0$.

Proof. All these equations can be easily derived from the formulas

$$(Z_i Z_j)(v) = (\phi_i\phi_j)(v) + \Pi((\psi_i\phi_j - \phi_i\psi_j)(v)), 1 \leq i, j \leq 3,$$

and

$$(Z_1 Z_2 Z_3)(v) = \Pi((\phi_1\psi_2\phi_3)(v)).$$

□

Consider a (not necessary super-commutative) A -superalgebra B , generated by the "formal" elements ϕ'_i, ψ'_i , subject to the relations (1) and (2) from Lemma 2.9, and the relations $p = 0$, where p is any monomial in ϕ'_i, ψ'_i of degree at least 4, $1 \leq i \leq 3$. It is clear that B is a graded superalgebra, provided all ϕ'_i and ψ'_i are odd and even elements of degree 1 respectively. Moreover, B has finite length as an A -supermodule.

Lemma 2.10. *The product $\phi'_1\psi'_2\phi'_3$ is not zero in B .*

Proof. One has to show that $\phi'_1\psi'_2\phi'_3$ can not be expressed as any A -linear combination of the polynomials xp or px , where p is one of the left hand side polynomials in the relations (1), (2) and x is one of the free generators of the absolutely free A -superalgebra $A \langle \phi'_i, \psi'_i \mid 1 \leq i \leq 3 \rangle$. It is obvious that if $p = 0$ is a relation (1), then both xp and px do not contain the monomial $\phi'_1\psi'_2\phi'_3$ at all. Therefore, $\phi'_1\psi'_2\phi'_3$ should be an A -linear combination of xp and px , where $p = 0$ is a relation (2). There are only two such polynomials :

$$\phi'_1(\psi'_2\phi'_3 - \phi'_2\psi'_3 - \phi'_3\psi'_2 + \psi'_3\phi'_2) \text{ and } (\psi'_2\phi'_1 - \phi'_2\psi'_1 - \phi'_1\psi'_2 + \psi'_1\phi'_2)\phi'_3.$$

Modulo the relations (1) they congruent to

$$\phi'_1\psi'_2\phi'_3 + \phi'_1\psi'_3\phi'_2 \text{ and } -\phi'_2\psi'_1\phi'_3 - \phi'_1\psi'_2\phi'_3.$$

It is clear now that no linear combination of the last polynomials is equal to $\phi'_1\psi'_2\phi'_3$. Lemma is proven. □

We set $V = B$ and define ϕ_i and ψ_i as the operators $b \mapsto \phi'_i b$ and $b \mapsto \psi'_i b, b \in B, 1 \leq i \leq 3$. It is evident that they satisfy (1), (2), (3) of Lemma 2.9.

3. ODD REGULAR ELEMENTS AND A SUPERIZED HOCHSCHILD COCHAIN COMPLEX

Let A be a super-ring. In this section we describe an homological approach to the following question. How to construct a super-ring R with an odd regular element $y \in R$ such that $R/Ry \simeq A$? By the definition, $yR \simeq \Pi A$ as a \mathbb{Z} -supermodule, on which A acts as $a\Pi b = (-1)^{|a|}\Pi(ab), a, b \in A$.

For the sake of simplicity we assume that the sequence of \mathbb{Z} -supermodules

$$0 \rightarrow Ry \rightarrow R \rightarrow R/Ry \rightarrow 0$$

is split. For example, this is the case if R is a superalgebra over a field. Then R is isomorphic to $A \oplus \Pi A$ as a \mathbb{Z} -supermodule, and the multiplication on R can be defined as

$$a \circ b = ab + \Pi \pi(a, b), a \circ \Pi b = (-1)^{|a|} \Pi(ab), (\Pi b) \circ a = \Pi(ba), (\Pi a) \circ (\Pi b) = 0, a, b \in A,$$

where $\pi : A \otimes A \rightarrow A$ is an odd bilinear super-skew symmetric map. The super-skew symmetry means that $\pi(a, a) = 0, \pi(b, c) = (-1)^{|b||c|} \pi(c, b), a \in A_1, b, c \in A$. If 2 is invertible in A , then the first condition is redundant. Note also that the regular element y can be identified with $\Pi 1$.

Lemma 3.1. *The map π defines a super-ring structure on $A \oplus \Pi A$ if and only if for any $a, b, c \in A$ the following conditions hold :*

- (1) $\pi(1, a) = 0$;
- (2) $\pi(ab, c) - \pi(a, bc) + \pi(a, b)c - (-1)^{|a|} a\pi(b, c) = 0$.

Proof. Since $1 \circ a = a$, the first condition obviously follows. For the second condition just compare

$$(a \circ b) \circ c = (ab)c + \Pi(\pi(ab, c) + \pi(a, b)c)$$

and

$$a \circ (b \circ c) = a(bc) + \Pi(\pi(a, bc) + (-1)^{|a|} a\pi(b, c)).$$

□

Assume that A is arbitrary (not necessary super-commutative) super-ring. Let M be an A -superbimodule.

We define a *superized Hochschild cochain complex* (compare with [8], p.301)

$$\tilde{C}^n(A, M) = \text{Hom}_{\mathbb{Z}}(A^{\otimes(n+1)}, M), n \geq 0,$$

with coboundary maps

$$\delta_n(f)(a_0, \dots, a_{n+1}) = \sum_{0 \leq i \leq n} (-1)^i f(a_0, \dots, a_i a_{i+1}, \dots, a_{n+1})$$

$$-(-1)^{|f||a_0|} a_0 f(a_1, \dots, a_{n+1}) + (-1)^{n+1} f(a_0, \dots, a_n) a_{n+1}, f \in \tilde{C}^m(A, M), n \geq 0.$$

Note that each δ_n is graded with respect to the natural \mathbb{Z}_2 -grading of \mathbb{Z} -supermodule $\tilde{C}^n(A, M)$. Thus the cohomology groups of this complex, denoted by $SHH^n(A, M)$, are naturally \mathbb{Z}_2 -graded.

Lemma 3.2. *For any $n \geq 0$ we have $\delta_{n+1} \delta_n = 0$.*

Proof. There is

$$\delta_{n+1}(\delta_n(f))(a_0, \dots, a_{n+2}) = \sum_{0 \leq i \leq n+1} (-1)^i \delta_n(f)(a_0, \dots, a_i a_{i+1}, \dots, a_{n+2})$$

$$-(-1)^{|f||a_0|} a_0 \delta_n(f)(a_1, \dots, a_{n+2}) + (-1)^{n+2} \delta_n(f)(a_0, \dots, a_{n+1}) a_{n+2}.$$

In the sum

$$\sum_{0 \leq i \leq n+1} (-1)^i \delta_n(f)(a_0, \dots, a_i a_{i+1}, \dots, a_{n+2})$$

each summand $f(a_0, \dots, a_i a_{i+1}, \dots, a_j a_{j+1}, \dots, a_{n+2})$ appears two times, once with the sign $(-1)^{i+j-1}$ and once with the sign $(-1)^{i+j}$, hence they are cancelled. Similarly, a summand $f(a_0, \dots, a_i a_{i+1} a_{i+2}, \dots, a_{n+2})$ appears two times, once with the sign $(-1)^{2i}$ and once with the sign $(-1)^{2i+1}$, hence cancelled as well. Thus

$$\begin{aligned}
 & \sum_{0 \leq i \leq n+1} (-1)^i \delta_n(f)(a_0, \dots, a_i a_{i+1}, \dots, a_{n+2}) = \\
 & -(-1)^{|a_0 a_1| |f|} a_0 a_1 f(a_2, \dots, a_{n+2}) + (-1)^{n+1} f(a_0 a_1, \dots, a_{n+1}) a_{n+2} \\
 & - \sum_{0 < i < n+1} (-1)^{i+|a_0| |f|} a_0 f(a_1, \dots, a_i a_{i+1}, \dots, a_{n+2}) \\
 & + (-1)^{n+1} \sum_{0 < i < n+1} (-1)^i f(a_0, \dots, a_i a_{i+1}, \dots, a_{n+1}) a_{n+2} \\
 & - (-1)^{n+1+|a_0| |f|} a_0 f(a_1, \dots, a_{n+1} a_{n+2}) + f(a_0, \dots, a_n) a_{n+1} a_{n+2} = \\
 & (-1)^{|a_0| |f|} a_0 [(-1)^{|a_1| |f|} a_1 f(a_2, \dots, a_{n+2}) + \sum_{0 < i \leq n+1} (-1)^{i+1} f(a_1, \dots, a_i a_{i+1}, \dots, a_{n+2})] \\
 & + (-1)^{n+1} [\sum_{0 \leq i < n+1} (-1)^i f(a_0, \dots, a_i a_{i+1}, \dots, a_{n+1}) + (-1)^{n+1} f(a_0, \dots, a_n) a_{n+1}] a_{n+2} = \\
 & (-1)^{|a_0| |f|} a_0 [\delta_n(f)(a_1, \dots, a_{n+2}) - (-1)^{n+1} f(a_1, \dots, a_{n+1}) a_{n+2}] \\
 & + (-1)^{n+1} [\delta_n(f)(a_0, \dots, a_{n+1}) + (-1)^{|a_0| |f|} f(a_1, \dots, a_{n+1}) a_{n+2}] = \\
 & (-1)^{|a_0| |f|} \delta_n(f)(a_1, \dots, a_{n+2}) + (-1)^{n+1} \delta_n(f)(a_0, \dots, a_{n+1}) a_{n+2}.
 \end{aligned}$$

Lemma is proven. \square

From now on A is again a super-commutative super-ring and M is a left A -supermodule, regarded as a right A -supermodule via the right action

$$ma = (-1)^{|a||m|} am, a \in A, m \in M.$$

Let $C^n(A, M)$ denote the \mathbb{Z} -supersubmodule of $\tilde{C}^n(A, M)$ consisting of all $f \in \tilde{C}^n(A, M)$, such that for any $a_0, \dots, a_n \in A$ there is

- (1) $f(1, a_1, \dots, a_n) = 0$;
- (2) $f(a_n, a_{n-1}, \dots, a_0) = (-1)^{\frac{n(n-1)}{2} + \sum_{0 \leq i < j \leq n} |a_i| |a_j|} f(a_0, \dots, a_n)$;
- (3) if 2 is not invertible in A , then $f(a, \dots, a) = 0$, provided $n \geq 1$ and a is odd.

Lemma 3.3. *If A is a super-commutative super-ring, then for any $n \geq 0$ we have $\delta_n(C^n(A, M)) \subseteq C^{n+1}(A, M)$, that is $\{C^n(A, M)\}_{n \geq 0}$ is a subcomplex of the complex $\{\tilde{C}^m(A, M)\}_{m \geq 0}$.*

Proof. Let d_n and ϵ_n denote $\sum_{0 \leq i < j \leq n} |a_i| |a_j|$ and $\frac{n(n-1)}{2}$ respectively. For arbitrary $f \in C^n(A, M)$ we have

$$\begin{aligned}
 \delta_n(f)(a_{n+1}, \dots, a_0) &= \sum_{0 \leq i \leq n} (-1)^i f(a_{n+1}, \dots, a_{n+1-i} a_{n-i}, \dots, a_0) \\
 &- (-1)^{|f| |a_{n+1}|} a_{n+1} f(a_n, \dots, a_0) + (-1)^{n+1} f(a_{n+1}, \dots, a_1) a_0 = \\
 &(-1)^{\epsilon_n + d_{n+1} + n} \delta_n(f)(a_0, \dots, a_{n+1}).
 \end{aligned}$$

Similarly, there is

$$\delta_n(f)(1, a_1, \dots, a_{n+1}) = f(a_1, \dots, a_{n+1}) - f(a_1, \dots, a_{n+1}) = 0.$$

It is clear that $\delta_n(f)$ satisfies the condition (3), whenever $n \geq 1$. If $n = 0$, then

$$\delta_0(f)(a, a) = f(a^2) - (-1)^{|f|}af(a) - f(a)a = 0$$

for $a \in A_1$. Lemma is proven. \square

The cohomology groups of the complex $\{C^n(A, M)\}_{n \geq 0}$ are denoted by $SH^n(A, M)$.

The statement of Lemma 3.1 can be formulated so that $\pi(a, b)$ defines a super-ring structure on $A \oplus \Pi A$ if and only if π is an odd cocycle from $C^1(A, A)$. The resulting super-ring is denoted by A_π .

Let R and R' are super-rings with regular elements y and y' respectively. We say that an isomorphism of super-rings $\phi : R \rightarrow R'$ is *adapted* to y and y' , whenever $\phi(y) = y'$ and the induced super-ring morphism $R/Ry \rightarrow R'/R'y'$ is an isomorphism. If it the case, then R and R' are said to be *adaptively* isomorphic. If we denote $R/Ry \simeq R'/R'y'$ by A , then there are cocycles $\pi, \pi' \in C^1(A, A)$ such that R and R' can be identified with A_π and $A_{\pi'}$ respectively.

Proposition 3.4. *In the above notations, R is adaptively isomorphic to R' if and only if the cosets of π and π' in $SH^1(A, A)$ are the same. Moreover, the exact sequence*

$$0 \rightarrow Ry \rightarrow R \rightarrow A \rightarrow 0$$

is split if and only if the coset of π is zero.

Proof. Recall that R and R' are isomorphic to $A \oplus \Pi A$ as \mathbb{Z} -supermodules. Let $\phi : R \rightarrow R'$ be an adapted isomorphism. It can be defined by

$$a \mapsto a + \Pi f(a), \Pi a \mapsto \Pi a, a \in A,$$

where $f \in C^0(A)_1$. Then ϕ is a super-ring morphism if and only if

$$\begin{aligned} \phi(a \circ b) &= \phi(ab + \Pi\pi(a, b)) = ab + \Pi(f(ab) + \pi(a, b)) = \phi(a) \circ \phi(b) = \\ &ab + \Pi(\pi'(a, b) + f(a)b + (-1)^{|f||a|}af(b)) \end{aligned}$$

for any $a, b \in A$, hence if and only if $\pi' - \pi = \delta_0(f)$. The proof of the second statement is similar and we leave it for the reader. Proposition is proven. \square

In the rest part of this section we construct an Artinian super-ring R with regular element $y \in R$ such that $\text{sdim}_1(R/Ry) < \text{sdim}_1(R) - 1$.

Let B be an Artinian ring. Consider the super-ring $A' = B[Y_1, Y_2, Y_3, Y_4]/I'$, where the super-ideal I' is generated by the elements $Y_{i_1}Y_{i_2}Y_{i_3}$, $1 \leq i_1 < i_2 < i_3 \leq 4$. It is clear that A' is an Artinian B -superalgebra and $\text{sdim}_1(A') = 2$.

Let A be an A' -superalgebra. Choose the elements $t_{i_1 i_2 i_3} \in A_0$, $1 \leq i_1, i_2, i_3 \leq 4$, such that $t_{i_{\sigma(1)} i_{\sigma(2)} i_{\sigma(3)}} = (-1)^\sigma t_{i_1 i_2 i_3}$ for any substitution σ and $t_{i_1 i_2 i_3} = 0$ if one of the indices i_1, i_2, i_3 coincides with the other. Define a B -bilinear map $\pi' : A' \otimes A' \rightarrow A$ by

- (a) $\pi'(1, A') = \pi'(A', 1) = 0$;
- (b) $\pi'(Y_i, Y_j) = 0$, $1 \leq i, j \leq 4$;
- (c) $\pi'(Y_i Y_j, Y_k) = \pi'(Y_k, Y_i Y_j) = t_{ijk}$, $1 \leq i, j, k \leq 4$;
- (d) $\pi'(Y_i Y_j, Y_s Y_k) = -t_{skj} Y_i$, $1 \leq i \neq j, k \neq s \leq 4$.

Lemma 3.5. *The map π' is a cocycle from $C^1(A', A)$ if and only if*

- (\star) $t_{skj} Y_i + t_{ski} Y_j = 0$, $1 \leq i, j, k, s \leq 4$;
- ($\star\star$) $t_{ski} Y_i = 0$, $1 \leq i, s, k \leq 4$.

Proof. The condition (\star) and $(\star\star)$ are derived from (d) and the eccential equations

$$\pi'(Y_i Y_j, Y_s Y_k) = -\pi'(Y_j Y_i, Y_s Y_k), \quad \pi'(Y_i^2, Y_s Y_k) = 0.$$

The super-skew symmetricity of π' is equivalent to the condition

$$t_{ijk} Y_s = t_{skj} Y_i.$$

Using (\star) and the definition of t -s we have

$$t_{ijk} Y_s = t_{jki} Y_s = -t_{jks} Y_i = -t_{ksj} Y_i = t_{skj} Y_i.$$

It sufficies to check the condition (2) from Lemma 3.1 for the free generators of free B -supermodule A' only, that is one can assume that a, b, c belong to

$$\{1, Y_i, Y_j Y_k \mid 1 \leq i \leq 4, 1 \leq j < k \leq 4\}.$$

If one of the elements a, b, c is equal to 1, then (2) holds automatically. Therefore, one can assume that

$$a, b, c \in \{Y_i, Y_j Y_k \mid 1 \leq i \leq 4, 1 \leq j < k \leq 4\}.$$

Note also that if $a, b, c \in \{Y_j Y_k \mid 1 \leq j < k \leq 4\}$, then all summands in (2) are equal to zero.

Below we list the rest cases.

If $a = Y_i, b = Y_j, c = Y_k$, then (2) follows from (b) and (c).

If $a = Y_i, b = Y_j, c = Y_s Y_k$, then (2) is equivalent to (d).

If $a = Y_i, b = Y_s Y_k, c = Y_j$, then (2) is equivalent to (\star) .

If $a = Y_s Y_k, b = Y_i, c = Y_j$, then (2) is derived from (\star) and the definition of t -s.

If $a = Y_i Y_j, b = Y_s Y_k, c = Y_t$, then (2) is equivalent to the equation

$$(t_{skj} Y_t + t_{skt} Y_j) Y_i = 0,$$

which follows from (\star) .

If $a = Y_i Y_j, b = Y_t, c = Y_s Y_k$, then (2) is equivalent to the equation

$$t_{ijt} Y_s Y_k = t_{skt} Y_i Y_j,$$

that can be derived from (\star) and the definition of t -s as

$$\begin{aligned} t_{ijt} Y_s Y_k &= t_{jti} Y_s Y_k = -t_{jts} Y_i Y_k = t_{jts} Y_k Y_i = t_{tsj} Y_k Y_i = \\ &= -t_{tsk} Y_j Y_i = t_{tsk} Y_i Y_j = t_{skt} Y_i Y_j. \end{aligned}$$

Finally, if $a = Y_t, b = Y_i Y_j, c = Y_s Y_k$, then (2) is equivalent to an equation, which is in turn equivalent to the equation from the previous case modulo (\star) . Lemma is proven. \square

Set $B = C[t_{ijk} \mid 1 \leq i < j < k \leq 4]/J$, where C is a (commutative) Artinian ring and the ideal J is generated by all products $t_{ijk} t_{skl}$. Then B is also Artinian.

For arbitrary triple of indices i, j, k we define t_{ijk} to be zero if $i = j$ or $k \in \{i, j\}$. Otherwise, i, j, k is a permutation of some ordered triple $i' < j' < k'$ and we set $t_{ijk} = (-1)^\sigma t_{i'j'k'}$, where $\sigma : i' \mapsto i, j' \mapsto j, k' \mapsto k$.

Define A as a quotient of the super-ring A' modulo the superideal I generated by the left-hand sides of (\star) and $(\star\star)$. The following lemma is obvious.

Lemma 3.6. *We have $\pi'(I, A') = 0$, hence π' naturally induces a cocylce $\pi \in C^1(A, A)$.*

Lemma 3.7. *The element $t_{123} Y_4$ does not belong to I .*

Proof. If this element does, then it should be a C -linear combination of the left-hand sides of (\star) and $(\star\star)$. The C -module A'_1 is freely generated by the elements

$$t_{ijk}Y_s, 1 \leq i < j < k \leq 4, s \in \{i, j, k\},$$

and

$$v_1 = t_{234}Y_1, v_2 = t_{134}Y_2, v_3 = t_{234}Y_3, v_4 = t_{123}Y_4.$$

The first elements are just left-hand sides of (\star) . Let M and N denote the C -span of the left-hand sides of $(\star\star)$ and the C -span of v_1, v_2, v_3, v_4 respectively. It is clear that $A'_1 = M \oplus N$.

If an element $z = t_{skj}Y_i + t_{ski}Y_j$ does not belong to M , then the indices i, j, s, k are pairwise different and $z \in N$. Without loss of a generality one can assume that $s < k$ and $i < j$. There are six such elements :

$$\begin{aligned} z_1 &= t_{143}Y_2 + t_{142}Y_3 = -t_{134}Y_2 - t_{124}Y_3 = -v_2 - v_3, \\ z_2 &= t_{134}Y_2 + t_{132}Y_4 = t_{134}Y_2 - t_{123}Y_4 = v_2 - v_4, \\ z_3 &= t_{124}Y_3 + t_{123}Y_4 = v_3 + v_4, \\ z_4 &= t_{234}Y_1 + t_{231}Y_4 = t_{234}Y_1 + t_{123}Y_4 = v_1 + v_4, \\ z_5 &= t_{243}Y_1 + t_{241}Y_3 = -t_{234}Y_1 + t_{124}Y_3 = -v_1 + v_3, \\ z_6 &= t_{342}Y_1 + t_{341}Y_2 = t_{234}Y_1 + t_{134}Y_2 = v_1 + v_2. \end{aligned}$$

Straightforward calculations show that

$$z_1 = -z_5 - z_6, z_2 = -z_4 + z_6, z_3 = z_4 + z_5$$

and the C -submodule $N' = \sum_{1 \leq i \leq 6} Cz_i$ is freely generated by the elements z_4, z_5, z_6 . Moreover, v_4 does not belong to N' . Lemma is proven. \square

Set $R = A_\pi$. Recall that R has an regular element $y = \Pi 1$. Then the odd super-dimension of $A \simeq R/Ry$ is at most 2. On the other hand, $Y_1 \circ Y_2 \circ Y_3 \circ Y_4 = \Pi(\pi(Y_1Y_2, Y_3)Y_4) = \Pi(t_{12,3}Y_4) \neq 0$, that is $\text{sdim}_1(R) = 4$.

4. GRADED AND BIGRADED SUPER-RINGS AND SUPERMODULES ASSOCIATED WITH NOETHERIAN SUPER-RINGS AND THEIR SUPERMODULES

Recall that a (not necessary Noetherian) super-ring R is called *graded* if $R = \bigoplus_{n \geq 0} R(n)$, where $R(n)$ is an $R(0)$ -super-submodule of R and $R(n)R(n') \subseteq R(n+n')$ for any nonnegative integers n, n' . Respectively, an R -supermodule M is called *graded* if $M = \bigoplus_{n \geq 0} M(n)$, where $M(n)$ is an $R(0)$ -super-submodule of M and $R(n)M(n') \subseteq M(n+n')$ for any nonnegative integers n, n' .

A super-ring R is called *bigraded* if $R = \bigoplus_{k, l \geq 0} R(k, l)$, and for any $k, k', l, l' \geq 0$ the following conditions hold :

- (1) $R(k, l)$ is a $R(0, 0)$ -super-submodule of R .
- (2) $R(k, l)R(k', l') \subseteq R(k+k', l+l')$.

Respectively, an R -supermodule M is called *bigraded* if $M = \bigoplus_{k, l \geq 0} M(k, l)$, and for any $k, k', l, l' \geq 0$ the following conditions hold :

- (1) $M(k, l)$ is a $R(0, 0)$ -super-submodule of M .
- (2) $R(k, l)M(k', l') \subseteq M(k+k', l+l')$.

It is clear that a bigraded super-ring R is also a graded super-ring with respect to the grading $R(n) = \bigoplus_{k+l=n} R(k, l), n \geq 0$. Moreover, if M is a bigraded R -supermodule, then it is also graded R -supermodule with respect to the similar grading $M(n) = \bigoplus_{k+l=n} M(k, l), n \geq 0$.

The polynomial superalgebra $A[X_1, \dots, X_r \mid Y_1, \dots, Y_s]$ is a bigraded superalgebra, where each "scalar" $a \in A$ has degree $(0, 0)$, X_i and Y_j have degrees $(1, 0)$ and $(0, 1)$ respectively, $1 \leq i \leq r, 1 \leq j \leq s$.

Let I be a super-ideal of R . Then we associate with R a graded super-ring $\text{gr}_I(R)$ such that $\text{gr}_I(R)(n) = I^n/I^{n+1}, n \geq 0$. Moreover, if M is a left R -supermodule, then one can define a graded $\text{gr}_I(R)$ -supermodule $\text{gr}_I(M)$ such that $\text{gr}_I(M)(n) = I^n M/I^{n+1} M$.

Let $I_{k,l}$ denote the super-ideal $I_0^k I_1^l R$. It is obvious that $R = I_{0,0}$ and $I = I_{1,0} + I_{0,1}$. Furthermore, if M is an R -supermodule, then we set $M_{k,l} = I_{k,l} M$. Since $I_{k,l} I_{k',l'} \subseteq I_{k+k',l+l'}$, the superspaces $\bigoplus_{k,l \geq 0} I_{k,l}/II_{k,l}$ and $\bigoplus_{k,l \geq 0} M_{k,l}/IM_{k,l}$ are a bigraded superalgebra and a bigraded supermodule over it, respectively. We denote them by $\text{bgr}_I(R)$ and $\text{bgr}_I(M)$ correspondingly.

Assume again that R is a Noetherian super-ring with $\text{Ksdim}_0(R) = \text{Kdim}(R_0) < \infty$.

Lemma 4.1. *The super-rings $\text{gr}_I(R)$ and $\text{bgr}_I(R)$ are Noetherian for any super-ideal I of R . Moreover, $\text{Ksdim}_0(\text{gr}_I(R)) < \infty$ and $\text{Ksdim}_0(\text{bgr}_I(R)) < \infty$ as well.*

Proof. There is a natural superalgebra epimorphism $\text{bgr}_I(R) \rightarrow \text{gr}_I(R)$ that is induced by the embeddings $I_{k,l} \rightarrow I^{k+l}$. Thus all we need is to prove that $\text{bgr}_I(R)$ is a Noetherian super-ring and its even part has finite Krull dimension.

Assume that I is generated by r even and l odd elements x_1, \dots, x_r and y_1, \dots, y_l respectively. There is a natural epimorphism

$$(R/I)[X_1, \dots, X_r \mid Y_1, \dots, Y_s] \rightarrow \text{bgr}_I(R),$$

of bigraded superalgebras, induced by the map

$$X_i \mapsto x_i + II_{1,0}, Y_j \mapsto y_j + II_{0,1}, 1 \leq i \leq r, 1 \leq j \leq s.$$

Therefore, one has to show that $(R/I)[X_1, \dots, X_r \mid Y_1, \dots, Y_s]$ is a Noetherian super-ring and its even part has finite Krull dimension.

Note that $(R/I)[X_1, \dots, X_r \mid Y_1, \dots, Y_s]$ is a finitely generated $(R/I)_0[X_1, \dots, X_r]$ -module. Since $(R/I)_0[X_1, \dots, X_r]$ is a Noetherian ring, the first statement follows.

Finally, $((R/I)[X_1, \dots, X_r \mid Y_1, \dots, Y_s])_0$ is integral over $(R/I)_0[X_1, \dots, X_r]$, and Theorem 20 and Theorem 22 from [3] imply the second statement. \square

5. GRADING AND SUPER-DIMENSION OF SUPEMODULES

Let R be a Noetherian super-ring with $\text{Kdim}(R_0) < \infty$. Let I be a superideal of R such that $I \subseteq \text{rad}(R)$. The graded super-ring $\text{gr}_I(R)$ is denoted by B . Then $B = \bigoplus_{n \geq 0} B(n)$, where $B(n) = I^n/I^{n+1}, n \geq 0$. Observe that for any positive integer n , there holds $B(n) = B(1)^n$.

Let M be a finitely generated R -supermodule. Let C denote $\text{gr}_I(M)$. Then $C = \bigoplus_{n \geq 0} C(n)$, where $C(n) = I^n M/I^{n+1} M, n \geq 0$. As above, for any positive integer n there is $C(n) = B(1)^n C(0)$. In particular, a homogeneous element $b \in B(k)$ belongs to $\text{Ann}_B(C)$ if and only if $bC(0) = 0$. Furthermore, since the super-ideal $\bigoplus_{n \geq 1} B(n)$ is nilpotent, we have

$$\text{sdim}_0(C) = \text{Kdim}(B(0)_0/\text{Ann}_{B(0)_0}(C(0))).$$

The prime (super)spectrum of R can be naturally identified with the prime (super)spectrum of B via

$$\mathfrak{P} \mapsto \tilde{\mathfrak{P}} = \mathfrak{P}/I \oplus I/I^2 \oplus \dots,$$

so that

$$\mathrm{gr}_{I_{\mathfrak{P}}}(M_{\mathfrak{P}}) \simeq \mathrm{gr}_I(M)_{\tilde{\mathfrak{P}}} = B_{\tilde{\mathfrak{P}}}.$$

Lemma 5.1. *The following hold*

$$\mathrm{sdim}_0(M) = \mathrm{sdim}_0(\mathrm{gr}_I(M)) \text{ and } \mathrm{sdim}_1(M) \geq \mathrm{sdim}_1(\mathrm{gr}_I(M)).$$

Proof. By the above remark $\mathrm{sdim}_0(\mathrm{gr}_I(M)) = \mathrm{sdim}_0(M/IM)$ and Lemma 2.4 infers the first equality. Further, if the elements z_1, \dots, z_t generate $B(0)_1 \oplus B(1)_1$ as a $B(0)_0$ -module, then they generate B_1 as a B_0 -module. Moreover, their representatives $y_1, \dots, y_t \in R_1$ generate R_1 as an R_0 -module (cf. [3], Corollary (1.M)(b)). By Proposition 2.1, for some $1 \leq s_1 < \dots < s_l \leq t$ the elements z_{s_1}, \dots, z_{s_l} form a longest system of odd parameters of $C = \mathrm{gr}_I(M)$. In other words, there is a prime ideal \mathfrak{q} of R_0 such that the system z_{s_1}, \dots, z_{s_l} subordinates the prime ideal $\mathfrak{q}/I_0 \oplus (\oplus_{n \geq 1} B(n)_0)$, that is $\mathrm{Ann}_{B(0)_0}(z^S C) \subseteq \mathfrak{q}/I_0$, where $S = \{s_1, \dots, s_l\}$ and \mathfrak{q}/I_0 is congruent to the first member of a longest prime chain in $B(0)_0/\mathrm{Ann}_{B(0)_0}(C(0))$. We have

$$(\mathrm{Ann}_{R_0}(M) + I_0)/I_0 \subseteq \mathrm{Ann}_{B(0)_0}(C(0)) \subseteq \mathrm{Ann}_{B(0)_0}(z^S C).$$

Moreover, Lemma 2.4 implies that $\mathrm{Ann}_{B(0)_0}(C(0))$ is contained in the radical of the ideal $(\mathrm{Ann}_{R_0}(M) + I_0)/I_0$, hence $\mathfrak{q}/\mathrm{Ann}_{R_0}(M)$ is the first member of a longest prime chain in $R_0/\mathrm{Ann}_{R_0}(M)$. Since

$$(\mathrm{Ann}_{R_0}(y^S M) + I_0)/I_0 \subseteq \mathrm{Ann}_{B(0)_0}(z^S C),$$

we conclude that $\mathrm{Ann}_{R_0}(y^S M) \subseteq \mathfrak{q}$. Lemma is proven. \square

Proposition 5.2. *Let \mathfrak{p} be a prime ideal of R_0 , such that $\mathfrak{p}/\mathrm{Ann}_{R_0}(M)$ is the first member of a longest prime chain of $R_0/\mathrm{Ann}_{R_0}(M)$. Set $\mathfrak{P} = \mathfrak{p} \oplus R_1$. Then*

$$\mathrm{sdim}_1(M_{\mathfrak{P}}) = \max\{k \mid (R_1^k M)_{\mathfrak{P}} \neq 0\} \leq \mathrm{sdim}_1(M).$$

If there is a longest system of odd parameters in M , which subordinates \mathfrak{p} , then the above inequality turns into equation.

Proof. We use the following elementary observations. For any $z_1, \dots, z_t \in R_1$ and $s_1, \dots, s_t \in S = R_0 \setminus \mathfrak{p}$, there holds

$$\mathrm{Ann}_{S^{-1}R_0}\left(\frac{z_1}{s_1} \dots \frac{z_t}{s_t} M\right) = S^{-1}\mathrm{Ann}_{R_0}(z^{\underline{t}} M).$$

Besides, $\mathrm{Ann}_{R_{\mathfrak{P}}}(M_{\mathfrak{P}}) = S^{-1}\mathrm{Ann}_R(M)$ and $R_{\mathfrak{P}}/\mathrm{Ann}_{R_{\mathfrak{P}}}(M_{\mathfrak{P}})$ is an Artinian super-ring with the unique prime superideal $\mathfrak{P}_{\mathfrak{P}}/\mathrm{Ann}_{R_{\mathfrak{P}}}(M_{\mathfrak{P}}) = S^{-1}(\mathfrak{P}/\mathrm{Ann}_R(M))$. Since the super-dimension of $M_{\mathfrak{P}}$ as an $R_{\mathfrak{P}}$ -supermodule is the same as the super-dimension of $M_{\mathfrak{P}}$ as an $R_{\mathfrak{P}}/\mathrm{Ann}_{R_{\mathfrak{P}}}(M_{\mathfrak{P}})$ -supermodule, Lemma 2.3 implies the first equality.

Further, the elements $\frac{z_1}{s_1}, \dots, \frac{z_t}{s_t}$ form a system of odd parameters if and only if $\mathrm{Ann}_{S^{-1}R_0}\left(\frac{z_1}{s_1} \dots \frac{z_t}{s_t} M\right) \subseteq S^{-1}\mathfrak{p}$ if and only if $\mathrm{Ann}_{R_0}(z^{\underline{t}} M) \subseteq \mathfrak{p}$. Proposition is proven. \square

Let $P(M)$ denote the subset of the prime spectrum of R consisting of all \mathfrak{P} such that $\text{Ann}_R(M) \subseteq \mathfrak{P}$ and $\mathfrak{p}/\text{Ann}_{R_0}(M)$ is the first member of a longest prime chain in $R_0/\text{Ann}_{R_0}(M)$. The following corollary is now evident.

Corollary 5.3. *There is*

$$\text{sdim}_1(M) = \max_{\mathfrak{P} \in P(M)} \max\{k \mid (R_1^k M)_{\mathfrak{P}} \neq 0\}.$$

In general, the inequality in Lemma 5.1 is not necessary equation. But in the following important case it is.

Theorem 5.4. *If $I = I_R$, then*

$$\text{sdim}(M) = \text{sdim}(\text{gr}_I(M)).$$

Proof. Let y_1, \dots, y_t be the same elements as in Lemma 5.1, and assume that for some $1 \leq i_1 < \dots < i_l \leq t$ the elements y_{i_1}, \dots, y_{i_l} form a longest system of odd parameters of M . Let \mathfrak{q} be a prime ideal of R_0 to which this system subordinates. Set $\Omega = \mathfrak{q} \oplus R_1$.

All we need is to show that R_Ω -supermodule M_Ω satisfies the conclusion of our theorem. In fact, if it is the case, then

$$\begin{aligned} \text{sdim}_1(M) &= \text{sdim}_1(M_\Omega) = \text{sdim}_1(\text{gr}_{I_\Omega}(M_\Omega)) = \text{sdim}_1(\text{gr}_I(M)_{\tilde{\Omega}}) \\ &\leq \text{sdim}_1(\text{gr}_I(M)), \end{aligned}$$

and by Lemma 5.1 M satisfies the conclusion of our theorem as well.

Therefore, without loss of a generality, one can assume that R is a local super-ring with the maximal superideal Ω such that $\Omega/\text{Ann}_R(M)$ is the unique prime superideal of the Artinian super-ring $R/\text{Ann}_R(M)$. Arguing as in Proposition 5.2, we obtain $\text{sdim}_1(M) = \max\{l \mid R_1^l M \neq 0\}$. On the other hand, $B = \text{gr}_I(R)$ is a local super-ring with the maximal superideal $\tilde{\Omega}$ and the arguments of Lemma 5.1 show that $\tilde{\Omega}/\text{Ann}_B(C)$ is the unique prime superideal of the Artinian super-ring $B/\text{Ann}_B(C)$ as well. In particular, $\text{sdim}_1(C) = \max\{l \mid B_1^l C \neq 0\}$. Then we have

$$\max\{l \mid (B_1)^l C \neq 0\} = \max\{l \mid B_1^l C \neq 0\} = \max\{l \mid C(l) \neq 0\}.$$

On the other hand, $(R_1)^l M \neq 0$ if and only if $(R_1)^l M \neq (R_1)^{l+1} M$ if and only if $C(l) \neq 0$, hence our theorem follows. \square

Corollary 5.5. *For any super-ring R there holds $\text{Ksdim}(R) = \text{Ksdim}(\text{gr}_{I_R}(R))$.*

Corollary 5.6. *If the elements $z_1, \dots, z_t \in I_R/I_R^2 \simeq R_1/R_1^3$ form a longest system of odd parameters of the supermodule $\text{gr}_{I_R}(M)$, then their representatives y_1, \dots, y_t form a longest system of odd parameters of M .*

Example 5.7. *Let M be the $A[Z_1, Z_2, Z_3, Y]$ -supermodule from the final part of the third section. Set $I = RY$. Then $R \simeq \text{gr}_I(R)$ and $\text{sdim}_1(\text{gr}_I(M)) = 2 < \text{sdim}_1(M)$. Indeed, $M/YM \simeq V$ and $YM \simeq \text{IV}$ as A -supermodules. Moreover, each Z_i acts on them by the rule $Z_i(v) = \phi_i(v)$, $Z_i(\Pi v) = -\Pi(\phi_i(v))$, $v \in V$. In particular, any product $Z_i Z_j$ acts trivially on $\text{gr}_I(M)$, hence Z_1, Y is one of the longest systems of odd parameters of $\text{gr}_I(M)$.*

Let X be an irreducible superscheme of finite type over a field K . Recall that the *super-dimension* of X is defined as $\text{Ksdim}(\mathcal{O}_X(U))$, where U is arbitrary (nonempty) open affine super-subscheme of X (cf. [4]).

Let \mathcal{J} be a locally nilpotent quasi-coherent superideal sheaf on X . Then one can associate with X an irreducible superscheme $\mathrm{gr}_{\mathcal{J}}(X)$ with the same *underlying topological space* X^e and with the structural superalgebra sheaf to be the sheafification of the presheaf $\bigoplus_{n \geq 0} \mathcal{J}^n / \mathcal{J}^{n+1}$. The functor $X \rightarrow \mathrm{gr}_{\mathcal{J}}(X)$ is an endofunctor of the category of superschemes of finite type that takes immersions to immersions. Moreover, if $\mathcal{J} = \mathcal{I}_X = (\mathcal{O}_X(\mathcal{O}_X)_1)^+$, a superscheme morphism $f : X \rightarrow Y$ is an isomorphism if and only if $\mathrm{gr}_{\mathcal{I}_X}(f)$ is (see [6], Proposition 9.3).

Proposition 5.8. *For any locally nilpotent quasi-coherent superideal sheaf \mathcal{J} on X there holds*

$$\mathrm{sdim}_0(X) = \mathrm{sdim}_0(\mathrm{gr}_{\mathcal{J}}(X)), \quad \mathrm{sdim}_1(X) \geq \mathrm{sdim}_1(\mathrm{gr}_{\mathcal{J}}(X)),$$

and

$$\mathrm{sdim}(X) = \mathrm{sdim}(\mathrm{gr}_{\mathcal{I}_X}(X)).$$

Proof. Arguing as in the proof of Proposition 9.3 (1), [6], one can assume that X is affine, say $X \simeq \mathrm{SSpec}(A)$. Then Corollary 5.5 concludes the proof. \square

6. DIMENSION THEORY OF CERTAIN LOCAL SUPERALGEBRAS

Let A be a Noetherian super-ring, I be a superideal of A and M be an A -supermodule. The I -adic completions $\varprojlim_n A/I^n$ and $\varprojlim_n M/I^n M$ of A and M are denoted by \widehat{A} and \widehat{M} respectively. It is evident that \widehat{M} has a natural structure of \widehat{A} -supermodule.

Lemma 6.1. *The following statements hold :*

- (1) *If M is a finitely generated A -supermodule, then \widehat{M} is canonically isomorphic to $M \otimes_A \widehat{A}$.*
- (2) *\widehat{A} is a flat A -supermodule.*

Proof. Use Proposition 1.9, [4], and copy the proof of Theorem 55 from [3]. \square

Let R be a local Noetherian super-ring with the maximal superideal $\mathfrak{M} = \mathfrak{m} \oplus R_1$. Let \mathfrak{N} be a (not necessary prime) \mathfrak{M} -primary superideal, that is $\mathfrak{N}_0 = \mathfrak{n}$ is a \mathfrak{m} -primary ideal.

Lemma 6.2. *A superideal \mathfrak{N} is \mathfrak{M} -primary if and only if R/\mathfrak{N} is Artinian if and only if $\mathfrak{M}^t \subseteq \mathfrak{N}$ for some positive integer t .*

Proof. Use Corollary 1.3 and note that by Lemma 1.5, [4], the \mathfrak{M} -adic topology on R coincides with the \mathfrak{m} -adic topology as well as the \mathfrak{N} -adic topology on R coincides with the \mathfrak{n} -adic topology. \square

From now on we assume that R contains a field, unless stated otherwise.

Lemma 6.3. *Let A be a Noetherian ring and I be a nilpotent ideal of A . If B is a subring of A such that A/I is a finitely generated $(B+I)/I$ -module, then A is a finitely generated B -module also.*

Proof. Just note that each quotient I^k/I^{k+1} is a finitely generated B/I -module, hence a finitely generated B -module as well. \square

Lemma 6.4. *Assume that the residue classes of the elements x_1, \dots, x_d form a system of (even) parameters of $\overline{R} = R_0/R_1^2$. If \mathfrak{n} is generated by these elements modulo R_1^2 , then $d = \mathrm{Ksdim}_0(R) = \mathrm{Ksdim}_0(\mathrm{bgr}_{\mathfrak{N}}(R))$.*

Proof. Let B denote $\text{bgr}_{\mathfrak{N}}(R)$. Then

$$B = \bigoplus_{k,l \geq 0} B(k, l), B(k, l) = \mathfrak{n}^k \mathfrak{N}_1^l R / (\mathfrak{n}^{k+1} \mathfrak{N}_1^l R + \mathfrak{n}^k \mathfrak{N}_1^{l+1} R), k, l \geq 0.$$

Observe that $B(0, 0)_0 \simeq R_0/\mathfrak{n}$ is a local algebra with the nilpotent maximal ideal $\mathfrak{m}/\mathfrak{n}$, hence complete. By Cohen structure theorem B contains a *coefficient field* $K \simeq R/\mathfrak{M}$ and $B(0, 0) = R/\mathfrak{N}$ is a finite dimensional K -superalgebra.

Since the superideal $\bigoplus_{k \geq 0, l > 0} B(k, l)$ is nilpotent, we have $\text{Ksdim}_0(B) = \text{Ksdim}_0(C)$, where C is the (graded) super-subring $\bigoplus_{k \geq 0} B(k, 0)$. Moreover, the ring

$$C_0 = \bigoplus_{k \geq 0} \mathfrak{n}^k / (\mathfrak{n}^{k+1} + \mathfrak{n}^k \mathfrak{N}_1 R_1)$$

is a quotient of $\text{gr}_{\mathfrak{n}}(R_0)$ modulo the nilpotent ideal

$$\bigoplus_{k \geq 0} (\mathfrak{n}^k \mathfrak{N}_1 R_1 + \mathfrak{n}^{k+1}) / \mathfrak{n}^{k+1},$$

hence $\text{Ksdim}_0(B) = \text{Kdim}(\text{gr}_{\mathfrak{n}}(R_0))$.

Let $\bar{\mathfrak{n}}$ denote $(\mathfrak{n} + R_1^2)/R_1^2$. The ring $\text{gr}_{\bar{\mathfrak{n}}}(\bar{R})$ is a quotient of $\text{gr}_{\mathfrak{n}}(R_0)$ modulo the nilpotent ideal

$$\bigoplus_{k \geq 0} (\mathfrak{n}^k \cap R_1^2 + \mathfrak{n}^{k+1}) / \mathfrak{n}^{k+1},$$

that implies $\text{Ksdim}_0(B) = \text{Kdim}_0(\text{gr}_{\bar{\mathfrak{n}}}(\bar{R}))$. Arguing as in Proposition 11.20 and Corollary 11.21, [1], one can show that $\text{gr}_{\bar{\mathfrak{n}}}(\bar{R})$ is a finitely generated module over the polynomial subalgebra $K[t_1, \dots, t_d]$, where t_i is $x_i + R_1^2 \pmod{\bar{\mathfrak{n}}^2}$, $1 \leq i \leq d$. Theorem 20 from [3] concludes the proof. \square

Assume that R is complete with respect to the \mathfrak{M} -adic or, equivalently, to the $\mathfrak{m}R$ -adic topology. The ring R_0 is also complete and therefore, it contains a coefficient field $K \simeq R/\mathfrak{M}$. Then the map $X_i \mapsto z_i = x_i + R_1^2$, $1 \leq i \leq d$, is extended for a continuous monomorphism $K[[X_1, \dots, X_d]] \rightarrow \bar{R}$ of complete local algebras. Besides, \bar{R} is a finitely generated module over $K[[z_1, \dots, z_d]]$ (see [1], Chapter 11, Exercise 2). Lemma 6.3 implies that R_0 is a finitely generated $K[[x_1, \dots, x_d]]$ -module. Moreover, $K[[x_1, \dots, x_d]]$ is a power series algebra, freely generated by the elements x_1, \dots, x_d as a *topological* algebra. Let D denote $K[[x_1, \dots, x_d]]$.

Lemma 6.5. *The following statements are equivalent :*

- (1) *The odd elements y_1, \dots, y_l form a system of odd parameters of R ;*
- (2) $\text{Ann}_D(y^l) = 0$;
- (3) *The map $Y_k \mapsto y_k$, $1 \leq k \leq l$, is extended for the natural isomorphism*

$$K[[X_1, \dots, X_d \mid Y_1, \dots, Y_l]] \simeq D[[y_1, \dots, y_l]].$$

Proof. The proof can be copied from Proposition 4.3, [4]. \square

Let y_1, \dots, y_s generate R_1 as a D -module. The proof of the following lemma is an obvious modification of the proof of Lemma 4.4, [4].

Lemma 6.6. *There holds*

$$\text{Ksdim}_1(R) = \max\{|I| \mid I \subseteq \underline{s} \text{ and } \text{Ann}_D(y^I) = 0\} = \max\{l \mid \text{Ann}_D(R_1^l) = 0\}.$$

Proposition 6.7. *Under the conditions of Lemma 6.4 we have*

$$\text{Ksdim}_1(\text{bgr}_{\mathfrak{N}}(R)) \leq \text{Ksdim}_1(R).$$

Proof. Let \widehat{R} denote the completion of R in the \mathfrak{N} -adic topology. Since this topology coincides with the \mathfrak{M} -adic topology, Theorem 5.12 from [4] implies $\text{Ksdim}(R) = \text{Ksdim}(\widehat{R})$. Moreover, Lemma 6.1 implies that $\text{bgr}_{\mathfrak{N}}(R)$ is naturally isomorphic to $\text{bgr}_{\mathfrak{N}}(\widehat{R})$. Therefore, without loss of a generality one can assume that R is complete.

Using Lemma 6.3 and the elementary observation that some elements are algebraically independent (over the ground field K) whenever they are algebraically independent modulo an ideal, we obtain that the elements $\overline{x_j} = x_j \pmod{\mathfrak{N}_{1,0}\mathfrak{N}}$ freely generate a polynomial subalgebra $K[\overline{x_1}, \dots, \overline{x_d}]$ of B , and B is a finitely generated $K[\overline{x_1}, \dots, \overline{x_d}]$ -supermodule.

Let \overline{D} denote $K[\overline{x_1}, \dots, \overline{x_d}]$. Since \overline{D} is a graded subalgebra of B , there is a generating set of \overline{D} -supermodule B consisting of homogeneous elements, say $u_1 \in B(k_1, l_1), \dots, u_t \in B(k_t, l_t)$. Choose also their representatives $v_1 \in \mathfrak{N}_{k_1, l_1}, \dots, v_t \in \mathfrak{N}_{k_t, l_t}$.

If $\text{Ksdim}_1(B) = l$, then there are odd elements u_{i_1}, \dots, u_{i_l} such that

$$\text{Ann}_{\overline{D}}(u_{i_1} \dots u_{i_l}) = 0.$$

It remains to show that the elements v_{i_1}, \dots, v_{i_l} form a system of odd parameters in R . If they do not, then by Lemma 6.5 (2) there is a nonzero power series $f \in D$ such that $f v_{i_1} \dots v_{i_l} = 0$. Thus, if g is the first nonzero homogeneous component of f , say of degree m , then $\overline{g} = g \pmod{\mathfrak{N}_{m+1,0}\mathfrak{N}} \neq 0$ and $\overline{g} u_{i_1} \dots u_{i_l} = 0$, a contradiction. Proposition is proven. \square

Later we will show that the above inequality is strict in general. But in some particular cases it turns to be equality.

Assume that \mathfrak{N} is a \mathfrak{M} -primary superideal from Proposition 6.7 such that $\mathfrak{N}_1 = R_1$. We call such (\mathfrak{M} -primary) superideal *special*.

Let A be either a polynomial K -algebra or a power series K -algebra, freely generated by the elements v_1, \dots, v_d . Set $\mathfrak{v} = Av_1 + \dots + Av_d$.

Lemma 6.8. *If M is a finitely generated A -module, then the function*

$$g_M : n \mapsto \dim M/\mathfrak{v}^{n+1}M$$

is a polynomial of degree at most d for all sufficiently large n .

Proof. If A is a polynomial algebra, then its completion in \mathfrak{v} -adic topology is isomorphic to $K[[v_1, \dots, v_d]]$. Moreover, $\widehat{\mathfrak{v}} = \widehat{A}\mathfrak{v} = \widehat{A}v_1 + \dots + \widehat{A}v_d$ and Lemma 6.1 implies that \widehat{M} is a finitely generated \widehat{A} -module. Moreover, we have

$$\widehat{M}/\widehat{\mathfrak{v}}^{n+1}\widehat{M} \simeq M/\mathfrak{v}^{n+1}M \otimes_A \widehat{A} \simeq M/\mathfrak{v}^{n+1}M$$

for any $n \geq 0$. Therefore, without loss of a generality one can assume that A is a power series algebra. Then Corollary 11.5 from [1], or [3], (12.C), conclude the proof. \square

Lemma 6.9. *$\text{Ann}_A(M) = 0$ if and only if the degree of g_M is exactly d .*

Proof. As it has been already observed, one can assume that A is a power series algebra. If $\text{Ann}_A(M) \neq 0$, then $I_m = \text{Ann}_A(m) \neq 0$ for any $m \in M$. Moreover, $\text{Kdim}(A/I_m) = t \leq d - 1$ and there is a \mathfrak{v}/I_m -primary ideal \mathfrak{t} , generated by t elements. Thus the degree of $g_{Am} \leq d - 1$ (see Corollary 11.5 and Proposition 11.6, [1], or [3], (12.C)). Note that M has a finite filtration whose quotients are monic

A -modules. The induction on the length of this filtration and Proposition (12.D), [3], imply that the degree of $g_M \leq d - 1$, hence the part "if".

Conversely, if the degree of $g_M \leq d - 1$, then by Proposition (12.D), [3], the degree of $g_N \leq d - 1$ for each monic A -submodule N . Thus $\text{Ann}_A(N) \neq 0$ and since M is finitely generated, $\text{Ann}_A(M) \neq 0$. Lemma is proven. \square

Theorem 6.10. *If \mathfrak{N} is a special \mathfrak{M} -primary superideal, then $\text{Ksdim}(\text{bgr}_{\mathfrak{N}}(R)) = \text{Ksdim}(R)$.*

Proof. For arbitrary nonnegative integers k, l we have

$$B(k, l) = \frac{\mathfrak{n}^k R_1^l + \mathfrak{n}^k R_1^{l+1}}{\mathfrak{n}^{k+1} R_1^l + \mathfrak{n}^k R_1^{l+2} + \mathfrak{n}^k R_1^{l+1}} \simeq \frac{\mathfrak{n}^k R_1^l}{\mathfrak{n}^{k+1} R_1^l + \mathfrak{n}^k R_1^{l+2}}.$$

As above, one can assume that R is complete in the \mathfrak{n} -adic topology. Then $C = \text{gr}_{I_R}(R)$ is complete in the $\bar{\mathfrak{n}}$ -adic topology, where $\bar{\mathfrak{n}} = (\mathfrak{n} + R_1^2)/R_1^2$ is generated by the even parameters $z_i = x_i + R_1^2, 1 \leq i \leq d$. Besides, these elements freely generate a power series subalgebra $K[[z_1, \dots, z_d]]$ of $C(0) = \bar{R}$ also, and C is a finitely generated $K[[z_1, \dots, z_d]]$ -supermodule. Since $C(l) = C(1)^l$ for any positive integer l , Lemma 6.5 infers $\text{Ksdim}_1(C) = \max\{l \mid \text{Ann}_D(C(l)) = 0\}$.

Consider a bigraded superalgebra $T = \text{gr}_{\bar{\mathfrak{n}}}(C)$, whose (bigraded) components are

$$T(k, l) = \bar{\mathfrak{n}}^k C(l) / \bar{\mathfrak{n}}^{k+1} C(l) \simeq \frac{\mathfrak{n}^k R_1^l + R_1^{l+2}}{\mathfrak{n}^{k+1} R_1^l + R_1^{l+2}} \simeq \frac{\mathfrak{n}^k R_1^l}{\mathfrak{n}^{k+1} R_1^l + (\mathfrak{n}^k R_1^l \cap R_1^{l+2})}.$$

We have $\mathfrak{n}^k R_1^{l+2} \subseteq \mathfrak{n}^k R_1^l \cap R_1^{l+2}$ for any $k, l \geq 0$, thus a canonical epimorphism $B \rightarrow T$ of bigraded superalgebras. As it has been already observed, the map $\bar{x}_i \mapsto t_i$ induces an isomorphism $\bar{D} \rightarrow K[t_1, \dots, t_d]$ of polynomial rings. Since both B and T are finitely generated supermodules over \bar{D} and $K[t_1, \dots, t_d]$ respectively, Lemma 4.4, [4], implies $\text{Ksdim}_1(B) \geq \text{Ksdim}_1(T)$. Therefore, if we will prove that $\text{Ksdim}_1(T) = \text{Ksdim}_1(C)$, then Proposition 6.7 and Theorem 5.4 imply the statement.

Combining Lemma 6.6 with Lemma 6.9, one sees that

$$\text{Ksdim}_1(C) = \max\{l \mid g_{C(l)} \text{ has the degree } d\},$$

where C is regarded as a $K[[z_1, \dots, z_d]]$ -supermodule. Recall that T is a graded superalgebra with respect to the grading $T(l) = \bigoplus_{k \geq 0} T(k, l), l \geq 0$. Moreover, if \mathfrak{v} denotes the ideal of $K[t_1, \dots, t_d]$ generated by t_1, \dots, t_d , then

$$\mathfrak{v}^k T(l) = \bigoplus_{s \geq k} T(s, l) \text{ and } \mathfrak{v}^k T(l) / \mathfrak{v}^{k+1} T(l) \simeq T(k, l), k, l \geq 0.$$

Furthermore, for any $l \geq 1$ there is $T(l) = T(1)^l$, and as above we have

$$\text{Ksdim}_1(T) = \max\{l \mid g_{T(l)} \text{ has the degree } d.\}$$

It remains to note that

$$g_{C(l)}(n) = \sum_{0 \leq k \leq n} \dim T(k, l) = g_{T(l)}(n)$$

for any nonnegative integer n , hence our theorem follows. \square

Let \mathfrak{w} denote the ideal of \bar{D} generated by the elements $\bar{x}_1, \dots, \bar{x}_d$. Set $B(l) = \bigoplus_{k \geq 0} B(k, l), l \geq 0$. Then we have

$$\mathfrak{w}^k B(l) = \bigoplus_{s \geq k} B(s, l) \text{ and } \mathfrak{w}^k B(l) / \mathfrak{w}^{k+1} B(l) \simeq B(k, l), k, l \geq 0,$$

similarly to the above.

Corollary 6.11. *We have $\text{Ksdim}_1(R) = \max\{l \mid g_{B(l)} \text{ has the degree } d\}$.*

Proof. Since $B \rightarrow T$ is a surjective morphism of bigraded superalgebras, for any $k, l \geq 0$ we have $g_{B(l)}(k) \geq g_{T(l)}(k)$. Moreover, B is a finitely generated \overline{D} -supermodule, hence the degree of each $g_{B(l)}$ is at most d . In particular, if the degree of some $g_{T(l)}$ is equal to d , then the degree of $g_{B(l)}$ is equal to d as well. As above, we have $B(l) = B(1)^l$ for any $l \geq 1$. Thus

$$\text{sdim}_1(B) = \max\{l \mid g_{B(l)} \text{ has the degree } d\} \leq \text{sdim}_1(T) \leq \text{sdim}_1(B).$$

□

This corollary allows us to define a *Hilbert polynomial* of R associated with a special \mathfrak{M} -primary superideal \mathfrak{N} as

$$g_{\mathfrak{N}}(x, y) = \sum_{l \geq 0} g_{B(l)}(x) y^l.$$

Then Corollary 6.11 can be reformulated as follows. Let $g_{\mathfrak{N}}(x, y)$ be a Hilbert polynomial of R associated with a special \mathfrak{M} -primary superideal \mathfrak{N} , expressed as

$$g_{\mathfrak{N}}(x, y) = \sum_{l \geq 0} g_l(x) y^l.$$

Then the odd superdimension of R is equal to the largest exponent l such that the degree of $g_l(x)$ is equal to $\text{sdim}_0(R)$.

7. FIBRES OF MORPHISMS OF SUPERSCHEMES

Lemma 7.1. *Let A be a local regular super-ring. Let z be a member of a minimal set of generators of A_0 -module A_1 . Then $\text{Ksdim}_1(A/Az) = \text{Ksdim}_1(A) - 1$.*

Proof. Let $z = z_1, \dots, z_s$ be a minimal set of generators of A_0 -module A_1 . Then z_1, \dots, z_s form an odd regular sequence in A , which is also the longest system of odd parameters of A (see Proposition 5.2, [4], or Corollary 3.3, [7]). Moreover, A/Az is again regular, hence

$$\text{Ksdim}_1(A/Az) = \dim_K((\overline{\mathfrak{P}}/\overline{\mathfrak{P}}^2)_1) = \dim_K((\mathfrak{P}/\mathfrak{P}^2)_1) - 1 = \text{Ksdim}_1(A),$$

where \mathfrak{P} is the maximal superideal of A and $\overline{\mathfrak{P}} = (\mathfrak{P} + Az)/Az$. □

From now on a local super-ring contains a field, unless stated otherwise.

Theorem 7.2. *Let $A \rightarrow B$ be a flat local morphism of local superalgebras and A is regular. Then*

$$\text{Ksdim}_1(B) \geq \text{Ksdim}_1(A) + \text{Ksdim}_1(B/\mathfrak{P}B),$$

where \mathfrak{P} is the maximal superideal of A .

Proof. Set $\text{Ksdim}(A) = r|s$. As it has been already observed, any minimal system of generators of A_0 -module A_1 has the cardinality s , and it is the longest system of odd parameters of A as well as the (longest) odd regular sequence of A . Moreover, we have the exact sequence

$$0 \rightarrow \sum_{1 \leq i \leq s} Az_i = \text{Ann}_A(z^s) \rightarrow A \rightarrow Az^s \rightarrow 0,$$

where $A \rightarrow Az^{\mathbb{z}}$ is defined as $a \mapsto az^{\mathbb{z}}, a \in A$. Since B is flat over A , we have the exact sequence

$$0 \rightarrow B \otimes_A \left(\sum_{1 \leq i \leq s} Az_i \right) \rightarrow B \otimes_A A \rightarrow B \otimes_A Az^{\mathbb{z}} \rightarrow 0,$$

which is naturally isomorphic to

$$0 \rightarrow \sum_{1 \leq i \leq s} Bz_i \rightarrow B \rightarrow Bz^{\mathbb{z}} \rightarrow 0,$$

hence z_1, \dots, z_s form an odd regular sequence in B as well.

Further, the superalgebra $B/Bz^{\mathbb{z}} \simeq A/Az^{\mathbb{z}} \otimes_A B$ is a flat $A/Az^{\mathbb{z}}$ -supermodule. By Lemma 2.6 (2) and Lemma 7.1, we have

$$\text{Ksdim}_1(B/Bz^{\mathbb{z}}) \geq \text{Ksdim}_1(B) - s, \quad \text{Ksdim}_1(A/Az^{\mathbb{z}}) = \text{Ksdim}_1(A) - s = 0.$$

Since $Bz^{\mathbb{z}} \subseteq B\mathfrak{P}$ and $A/Az^{\mathbb{z}} \simeq \overline{A}$, it remains to prove our theorem in the case when A is a regular ring. In other words, one has to show that

$$\text{Ksdim}_1(B) \geq \text{Ksdim}_1(B/\mathfrak{p}B),$$

where \mathfrak{p} is the maximal ideal of A . The ideal \mathfrak{p} is generated by a regular sequence a_1, \dots, a_r , where $r = \text{Kdim}(A)$ (see Theorem 11.22 and Lemma 11.23, [1]).

Again, since B is flat over A , the element a_1 is regular in B and B/Ba_1 is flat over A/Aa_1 . If we will show that $\text{Ksdim}_1(B) \geq \text{Ksdim}_1(B/Ba_1)$, then Corollary 11.18, [1], combined with the induction on r , concludes the proof.

Theorem 19 (2), [3], implies

$$\text{Kdim}(B_0) = \text{Kdim}(\overline{B}) = \text{Kdim}(A) + \text{Kdim}(B_0/B_0\mathfrak{p}) = \text{Kdim}(A) + \text{Kdim}(\overline{B}/\overline{B}\mathfrak{p}).$$

Let b_1, \dots, b_l form a system of parameters of $\overline{B}/\overline{B}\mathfrak{p}$. Then $b_1, \dots, b_l, a_1, \dots, a_r$ form a system of parameters of \overline{B} . Using Theorem 5.12, [4], one can replace B by \widehat{B} and B/Ba_1 by $\widehat{B}/\widehat{B}a_1 \simeq \widehat{B}/\widehat{B}a_1$. Moreover, the elements $b_1, \dots, b_l, a_1, \dots, a_r$ remain parameters of $\widehat{B} \simeq \overline{B}$. Besides, by Lemma 6.1 (2) \widehat{B} is flat over A , whence a_1 is regular in \widehat{B} .

If B is complete, then Lemma 6.6 infers

$$\text{Ksdim}_1(B) = \max\{l \mid \text{Ann}_D(B_1^l) = 0\},$$

where $D = K[[b_1, \dots, b_l, a_1, \dots, a_r]]$ and K is a coefficient field of B . Similarly, Corollary 11.18, [1], implies

$$\text{Kdim}(B_0/B_0a_1) = \text{Kdim}(B_0) - 1,$$

and thus

$$\text{Ksdim}_1(B/Ba_1) = \max\{l \mid \text{Ann}_{D'}(B_1^l/(B_1^l \cap Ba_1)) = 0\},$$

where $D' = K[[b_1, \dots, b_l, a_2, \dots, a_r]]$.

Assume that an element $x \in B_1^l$ is not annihilated by any nonzero element from D' modulo $B_1^l \cap Ba_1$. Each element $f \in D \setminus 0$ can be represented as $a_1^m(f_1 + a_1^n f_2)$, $f_1 \in D' \setminus 0$, $f_2 \in D$, $m \geq 0$, $n > 0$. The congruence

$$(f_1 + a_1^n f_2)x \equiv f_1 x \pmod{B_1^l \cap Ba_1}$$

implies $(f_1 + a_1^n f_2)x \neq 0$, hence $f_1 x \neq 0$, since a_1 is not zero divisor in B . Thus $\text{Ann}_D(x) = 0$ and our theorem follows. \square

Let $f : X \rightarrow Y$ be a morphism of superschemes. If $y \in Y^e$ is a point in the underlying topological space of Y , then similarly to [2], Chapter II, §3, one can define the *fiber* of the morphism f over the point y to be the superscheme

$$X_y = X \times_Y \text{SSpec}(K(y)),$$

where $K(y) = \mathcal{O}_{y,Y}/\mathfrak{m}_y$ and the closed embedding $\text{SSpec}(K(y)) \rightarrow Y$ is defined by the natural super-ring morphism $\mathcal{O}(Y) \rightarrow \mathcal{O}_{y,Y}$. In general, the canonical projection $X_y \rightarrow X$ is not necessary immersion (cf. [6], section 1.4).

Lemma 7.3. *The topological space X_y^e is naturally isomorphic to $f^{-1}(y)$.*

Proof. The topological space X_y^e is identified with the underlying topological space of $(X_y)_{ev}$. By Lemma 2.3, [6], $(X_y)_{ev} \simeq (X_{ev})_y$. It remains to refer to [2], Chapter II, Exercise 3.10. \square

Recall that a superscheme Z is a *closed super-subscheme* of X , if there is a closed embedding $\iota : Z^e \rightarrow X^e$ such that the sheaf $\iota^*\mathcal{O}_Z$ is an epimorphic image of the sheaf \mathcal{O}_X . Equivalently, for any $z \in Z^e$ the induced local morphism $\mathcal{O}_{\iota(z),X} \rightarrow \mathcal{O}_{z,Z}$ is surjective. Note that a nonempty closed subset Z of X^e may have many possible closed super-subscheme structures. In particular, the odd dimension of Z can vary with respect to the choice of such a structure.

Let X be a superscheme of finite type over a field K . Let Z be an irreducible component of X^e equipped with a closed super-subscheme structure.

Lemma 7.4. *There is a point $z \in Z$ such that $\text{sdim}_1(Z) \leq \text{Ksdim}_1(\mathcal{O}_{z,X})$.*

Proof. Let $U \simeq \text{SSpec}(A)$ be an open affine super-subscheme of X such that $U \cap Z \neq \emptyset$. Then $U \cap Z \simeq \text{SSpec}(A/I)$ is an irreducible component of U , where $\sqrt{I_0} = \mathfrak{p}$ is a minimal prime ideal of A_0 . The required point z corresponds to the prime superideal $\mathfrak{P} = \mathfrak{p} \oplus A_1$. In fact, $\text{sdim}_1(Z) = \text{Ksdim}_1(A/I)$ and if the cosets of the elements y_1, \dots, y_k form a system of odd parameters of A/I , then

$$(\text{Ann}_{A_0}(y^{\underline{k}}) + I_0)/I_0 \subseteq \text{Ann}_{A_0/I_0}(\bar{y}^{\underline{k}}) \subseteq \mathfrak{p}/I_0$$

implies $\text{Ann}_{A_0}(y^{\underline{k}}) \subseteq \mathfrak{p}$. In particular, $S^{-1}A_1^{\underline{k}} \neq 0$, where $S = A_0 \setminus \mathfrak{p}$. Since the superalgebra $\mathcal{O}_{z,X} \simeq S^{-1}A$ is Artinian, Lemma 2.3 concludes the proof. \square

Recall that a superscheme morphism $f : X \rightarrow Y$ is called *flat* whenever for any $x \in X^e$ the induced local morphism $\mathcal{O}_{f(x),Y} \rightarrow \mathcal{O}_{x,X}$ is flat.

It is known that for arbitrary flat morphism of irreducible schemes $f : X \rightarrow Y$ each irreducible component of X_y , where $f^{-1}(y) \neq \emptyset$, has the dimension $\dim(X) - \dim(Y)$ (cf. [2], Chapter III, Corollary 9.6). In particular, $\dim(X_y) = \dim(X) - \dim(Y)$.

Theorem 7.5. *Let $f : X \rightarrow Y$ be a flat morphism of irreducible superschemes. Let $y \in Y^e$ be a nonsingular point such that $f^{-1}(y) \neq \emptyset$ and let Z be an irreducible component of X_y^e equipped with a closed (irreducible) super-subscheme structure. Then*

$$\text{sdim}_1(X) - \text{sdim}_1(Y) \geq \text{sdim}_1(Z).$$

Proof. Without loss of generality one can assume that X and Y are affine, say $X = \text{SSpec}(B)$ and $Y = \text{SSpec}(A)$ (argue as in Lemma 2.1, [6]). Then f is dual to a (flat) superalgebra morphism $A \rightarrow B$, the point y corresponds to a prime superideal \mathfrak{P} of A and each point x from X_y^e corresponds to some prime superideal

Ω of B such that $A \cap \Omega = \mathfrak{P}$. Moreover, the fiber $X_y \simeq \text{SSpec}(B_{\mathfrak{P}}/(B\mathfrak{P})_{\mathfrak{P}})$ is also affine.

Note that Theorem 7.2 can be interpreted as follows. For any $x \in X_y^e$ there holds

$$\text{sdim}_1(\mathcal{O}_{x,X}) \geq \text{sdim}_1(\mathcal{O}_{y,Y}) + \text{sdim}_1(\mathcal{O}_{x,X_y}).$$

In fact, we have

$$\mathcal{O}_{x,X} \simeq B_{\Omega}, \mathcal{O}_{y,Y} \simeq A_{\mathfrak{P}}$$

and

$$\mathcal{O}_{x,X_y} \simeq ((B/B\mathfrak{P})_{\mathfrak{P}})_{(\Omega/B\mathfrak{P})_{\mathfrak{P}}} \simeq B_{\Omega}/\mathfrak{P}B_{\Omega}.$$

Thus, if x is as in Lemma 7.4, then the equalities $\text{sdim}_1(\mathcal{O}_{x,X}) = \text{sdim}_1(X)$ and $\text{sdim}_1(\mathcal{O}_{y,Y}) = \text{sdim}_1(Y)$ imply our statement (see [4], Remark 6.2). \square

The following example shows that the inequality in Theorem 7.2 can be strict.

Example 7.6. *In the example from the third section set $C = K$. Then A is obviously a local Artinian superalgebra. Moreover, $R = A_{\pi}$ is a local Artinian superalgebra and a free $K[y]$ -supermodule of the rank $\dim A$ as well, where $y = \Pi 1$. The superalgebra $K[y]$ is obviously regular and local with the maximal superideal $\mathfrak{P} = Ky$. Thus the natural embedding $K[y] \rightarrow R$ is flat and local, and we have*

$$\text{Ksdim}_1(R) > \text{Ksdim}_1(K[y]) + \text{Ksdim}_1(R/R\mathfrak{P}).$$

REFERENCES

- [1] V.F.Atiyah, I.G.Macdonald, *Introduction to Commutative Algebra*, Addison-Wesley, Reading, Mass., 1969.
- [2] R.Hartshorne, *Algebraic Geometry*, Springer, 1977.
- [3] H. Matsumura, *Commutative algebra*, 2nd ed., Math. Lec. Notes Series **56**, Benjamin/Cunning Publishing Company, 1980.
- [4] A.Masuoka and A.N.Zubkov, On the notion of Krull super-dimension, *Journal of Pure and Applied Algebra*, 224 (2020), no. 5, 106245.
- [5] A. Masuoka and A. N. Zubkov, *Quotient sheaves of algebraic supergroups are superschemes*. *J. Algebra*, 348 (2011), 135–170.
- [6] A. Masuoka and A. N. Zubkov, *Group superschemes*, arXiv : 2011.14207 v1.
- [7] T. Schmitt, *Regular sequences in \mathbb{Z}_2 -graded commutative algebra*, *J.Algebra*, 124 (1989), 60–118.
- [8] Weibel, Charles A., *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, 38. Cambridge University Press, Cambridge, 1994. xiv+450 pp.
- [9] Zubkov, A. N., *On some properties of Noether superschemes*, (Russian) *Algebra Logika* 57 (2018), no. 2, 197–213; translation in *Algebra Logic* 57 (2018), no. 2, 130–140.

DEPARTMENT OF MATHEMATICAL SCIENCE, UAEU, AL-AIN, UNITED ARAB EMIRATES; SOBOLEV INSTITUTE OF MATHEMATICS, OMSK BRANCH, PEVTZOVA 13, 644043 OMSK, RUSSIA

Email address: a.zubkov@yahoo.com

SOBOLEV INSTITUTE OF MATHEMATICS, AKAD. KOPTYUG PROSP., 4, 630090 NOVOSIBIRSK, RUSSIA

Email address: pavelisk@math.nsc.ru