

ON HOMOGENEOUS SPACES WITH FINITE ANTI-SOLVABLE STABILIZERS

GIANCARLO LUCCHINI ARTECHE

ABSTRACT. We say that a group is anti-solvable if all of its composition factors are non-abelian. We consider a particular family of anti-solvable finite groups containing the simple alternating groups for $n \neq 6$ and all 26 sporadic simple groups. We prove that, if K is a perfect field and X is a homogeneous space of a smooth algebraic K -group G with finite geometric stabilizers lying in this family, then X is dominated by a G -torsor. In particular, if $G = \mathrm{SL}_n$, all such homogeneous spaces have rational points.

MSC codes: 14M17, 12G05, 14G05.

We say that a finite group F is *aut-split* if the exact sequence

$$1 \rightarrow \mathrm{Inn}(F) \rightarrow \mathrm{Aut}(F) \rightarrow \mathrm{Out}(F) \rightarrow 1,$$

splits, where $\mathrm{Inn}(F)$ denotes the group of inner automorphisms of F and $\mathrm{Out}(F)$ is simply defined as the quotient $\mathrm{Aut}(F)/\mathrm{Inn}(F)$. A full characterization of aut-split finite simple groups was given in [LMM03]. These include the alternating group A_n for $n = 5$ or $n \geq 7$ and all 26 sporadic groups. As for groups of Lie type, some notation is needed. If F is a finite simple group of Lie type over \mathbb{F}_q with $q = p^m$ and p prime, denote by d the order of the group of diagonal automorphisms (i.e. those induced by conjugation by diagonal matrices) modulo inner automorphisms. Then F is aut-split if and only if one of the following holds:

- F is a Chevalley group, not of type $D_\ell(q)$, and $(\frac{q-1}{d}, d, m) = 1$;
- $F = D_\ell(q)$ and $(\frac{q^\ell-1}{d}, d, m) = 1$;
- F is a twisted group, not of type ${}^2D_\ell(q)$, and $(\frac{q+1}{d}, d, m) = 1$;
- $F = {}^2D_\ell(q)$ and either ℓ is odd or $p = 2$.

We say that F is *anti-solvable* if all of its composition factors (which are simple groups) are non-abelian. In this note we are interested in anti-solvable groups whose composition factors are aut-split (i.e. they belong to the list above). This includes of course the groups in the list themselves. The main result of this note is the following.

Theorem 1. *Let K be a perfect field and let G be a smooth algebraic K -group. Let X be a homogeneous space of G with geometric stabilizer \bar{F} . Assume that \bar{F} is finite, anti-solvable, and that all of its composition factors are aut-split. Then X is dominated by a G -torsor. In particular, if $G = \mathrm{SL}_n$, then X has a rational point.*

Of course, the particular case of $G = \mathrm{SL}_n$ is a natural consequence of the triviality of torsors under SL_n , which is valid over any field K . Now, recall that the *Springer class*,

Key words and phrases. homogeneous spaces, rational points, non-abelian cohomology, finite simple groups.

The author was partially supported by Fondecyt Grant 1210010.

which is class in a non-abelian 2-cohomology set associated to every homogeneous space X , corresponds to the obstruction to being dominated by a G -torsor (cf. [DLA17, §2] or [FSS98, §1], or [Spr66] for the original article by Springer). Using this tool we immediately see that Theorem 1 is implied by the following result on non-abelian Galois cohomology.

Theorem 2. *Let K be a perfect field and let L be a finite K -lien (or K -kernel) whose underlying group is anti-solvable and all of its composition factors are aut-split. Then $H^2(K, L)$ has exactly one class and it is neutral.*

The rest of this note is devoted to proving Theorem 2. We start with the particular case where the group is itself aut-split (for instance, if there is only one composition factor).

Proposition 3. *Let K be a perfect field and let L be a finite K -lien (or K -kernel) whose underlying group is aut-split. Then $H^2(K, L)$ has a neutral class (in particular, the K -lien L is representable). If moreover L has a trivial center, then this is the only class in $H^2(K, L)$.*

Proof. Let F be the underlying finite group of L and let Γ_K be the absolute Galois group of K . By the definition of a K -lien we have a homomorphism $\kappa : \Gamma_K \rightarrow \text{Out}(F)$. Since F is aut-split, we can compose this with a section $\text{Out}(F) \rightarrow \text{Aut}(F)$ and get a group action of Γ_K on F . Since F is finite, this defines a K -form of F and, since this action lifts κ , it corresponds to a neutral class $\eta \in H^2(K, L)$. The second assertion follows immediately from [Spr66, Prop. 1.17] (see also [Mac95, IV, Thm. 8.8]). \square

Proof of Theorem 2. Let L be a K -lien and let F be its underlying group as in Theorem 2. Since F is anti-solvable, it has trivial center and hence the K -lien L is representable (cf. [Mac95, IV, Thm. 8.7]). We fix then a class $\eta \in H^2(K, L)$, which is unique again by the triviality of the center and [Spr66, Prop. 1.17]. We need to prove that it is neutral.

If F has no proper nontrivial characteristic subgroups (i.e. it is *characteristically simple*), then it is a direct product of copies of a single simple group (cf. [Gor80, Ch. 2, Thm. 1.4]), which has to be aut-split then by hypothesis. Since every automorphism of such a product permutes its direct factors, it is easy to see that F is aut-split as well. We are done then by Proposition 3.

Assume now that F has a proper nontrivial characteristic subgroup N . It is clear that both N and F/N also satisfy the hypotheses of Theorem 2, so we argue by induction. Since N is characteristic, the arrow $\text{Aut}(F) \rightarrow \text{Aut}(F/N)$ is well-defined and thus the homomorphism $\kappa : \Gamma_K \rightarrow \text{Out}(F)$ gives a homomorphism $\Gamma_K \rightarrow \text{Out}(F/N)$, which gives us a K -lien L' whose underlying group is F/N . By induction hypothesis, we know that $H^2(K, L')$ has exactly one class η' and that it is neutral. We interpret this in the language of group extensions (cf. again [DLA17, §2.2.2] or [FSS98, 1.18] or [Spr66, 1.13–14]). The class η' corresponds to a split extension which, since the arrow $H^2(K, L) \rightarrow H^2(K, L')$ clearly sends η to η' , fits into a commutative diagram

$$\begin{array}{ccccccc}
 1 & \longrightarrow & F & \longrightarrow & E & \longrightarrow & \Gamma_K \longrightarrow 1 \\
 & & \downarrow & & \downarrow \pi & \longleftarrow s & \parallel \\
 1 & \longrightarrow & F/N & \longrightarrow & E' & \longrightarrow & \Gamma_K \longrightarrow 1.
 \end{array}$$

Consider the preimage $\pi^{-1}(s(\Gamma_K)) \subset E$, which fits into the following commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & N & \longrightarrow & \pi^{-1}(s(\Gamma_K)) & \longrightarrow & \Gamma_K \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \parallel \\ 1 & \longrightarrow & F & \longrightarrow & E & \longrightarrow & \Gamma_K \longrightarrow 1. \end{array}$$

The top row corresponds to a class in $H^2(K, L'')$ for some K -lien L'' with underlying group N . By induction hypothesis again we know that this class is neutral and hence the extension is split, giving a section $\Gamma_K \rightarrow \pi^{-1}(s(\Gamma_K))$ and hence a section $\Gamma_K \rightarrow E$ by composition, proving that the class η is neutral. \square

Remark 4. In [Ber67], Bercov characterizes anti-solvable finite groups whose composition factors are aut-split as iterated twisted wreath products of simple groups. Note however that we did not need this result in the proof above.

Remark 5. Theorems 1 and 2 are false if we take away the aut-split hypothesis on the composition factors, already for simple groups. Indeed, consider the group A_6 and the order 2 element $\phi \in \text{Out}(A_6)$ coming from the nontrivial class of outer automorphisms of S_6 . Taking $K = \mathbb{R}$, we can define a K -lien L via the homomorphism $\kappa : \Gamma_{\mathbb{R}} \rightarrow \text{Out}(A_6)$ that sends the nontrivial element in $\Gamma_{\mathbb{R}}$ to ϕ . By the same arguments used above we see that $H^2(K, L)$ has only one class η . But this class is not neutral, since this would amount to the existence of a lift of ϕ to an automorphism $\varphi \in \text{Aut}(A_6)$ of order 2, which does not exist. Using [DLA17, Cor. 3.3] we may construct a homogeneous space X of SL_n over \mathbb{R} whose Springer class is η . This implies that X is not dominated by a torsor.

REFERENCES

- [Ber67] **R. Bercov.** On groups without abelian composition factors. *J. Algebra* 5, 106–109, 1967.
- [DLA17] **C. Demarche, G. Lucchini Arteche.** Le principe de Hasse pour les espaces homogènes : réduction au cas des stabilisateurs finis. *Compos. Math.* 155, 1568–1593, 2019.
- [FSS98] **Y. Z. Flicker, C. Scheiderer, R. Sujatha.** Grothendieck’s theorem on non-abelian H^2 and local-global principles. *J. Amer. Math. Soc.* 11(3), 731–750, 1998.
- [Gor80] **D. Gorenstein.** *Finite Groups*. Second edition. Chelsea Publishing Co., New York, 1980.
- [LMM03] **A. Lucchini, F. Menegazzo, M. Morigi.** On the existence of a complement for a finite simple group in its automorphism group. *Illinois J. Math.* 47, 395–418, 2003.
- [Mac95] **S. Mac Lane** *Homology*. Reprint of the 1975 edition. Classics in Mathematics. Springer-Verlag, Berlin, 1995.
- [Spr66] **T. A. Springer.** Nonabelian H^2 in Galois cohomology. *Algebraic Groups and Discontinuous Subgroups (Proc. Sympos. Pure Math., Boulder, Colo., 1965)*, 164–182, Amer. Math. Soc., Providence, RI, 1966.

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD DE CHILE, LAS PALMERAS
3425, ÑUÑO A, SANTIAGO, CHILE
Email address: luco@uchile.cl