

Tautological rings of Hilbert modular varieties

Simon Cooper

November 16, 2021

Abstract

We compute the tautological ring for a Hilbert modular variety at an unramified prime. The method generalises that of van der Geer from the Siegel case.

1 Introduction

In this note we shall compute the tautological ring of Hilbert modular varieties in characteristic p .

For a Shimura variety S_K , its tautological ring $T^\bullet(S_K)$ is the subring of its Chow ring with rational coefficients generated by the Chern classes of automorphic vector bundles. This ring was first defined and studied by van der Geer [vdG99] in the Siegel case and later by Wedhorn-Ziegler [Zie18] for Shimura varieties admitting an embedding into a Siegel Shimura variety (i.e. Hodge-type). Given an integral canonical model of a Shimura variety, conjectured to exist at hyperspecial level by Milne [Mil92] (and proven to exist for Hodge- and abelian-type by Kisin [Kis10] and Vasiu [Vas99]), one can also define the tautological ring for the reduction of a Shimura variety to characteristic p . When suitable toroidal compactifications S_K^{tor} exist there is the folk-lore conjecture that

$$T^\bullet(S_K^{tor}) \cong H^{2\bullet}(\mathbf{X}^\vee(\mathbb{C}), \mathbb{Q})$$

where \mathbf{X}^\vee is the compact dual. Van der Geer proved this in the Siegel case in all characteristics and Wedhorn-Ziegler proved that it holds for Shimura varieties of Hodge-type in characteristic p (where suitable toroidal compactifications of integral models exist by work of Madapusi-Pera [MP19]). We are interested in the related question:

Question 1.1. *Is there a similar description of $T^\bullet(S_K)$?*

In the Siegel case the answer is yes. Indeed, van der Geer proved that $T^\bullet(A_g) \cong H^{2\bullet}(\mathbf{X}_{g-1}^\vee)$, where \mathbf{X}_{g-1} is the double Siegel half-space. In this note we compute $T^\bullet(S_K)$ for a Hilbert modular variety in characteristic p (see the beginning of section 5 for a precise statement).

Theorem 1. *Let X_K be a Hilbert modular variety at an unramified prime.*

$$T^\bullet(X_K) \cong H^{2\bullet}(X^\vee)/(top\ degree\ part)$$

In particular, the only extra relation is the vanishing of the top Chern class of the Hodge vector bundle pulled back from the Siegel case. It is not clear what a general description for Hodge-type Shimura varieties could be. For example, S_K could itself be compact.

The proof of Theorem 1 generalises the method of van der Geer from the Siegel case, using properness of subvarieties to obtain positivity of certain classes. Pullback gives a surjective map of graded rings

$$H^{2\bullet}(X^\vee) \cong T^\bullet(S_K^{tor}) \longrightarrow T^\bullet(S_K)$$

Our aim is to determine the kernel of this morphism of graded rings. Section 2 recalls the Hilbert modular datum and in section 3 we review Wedhorn-Ziegler's definition of $T^\bullet(S_K)$ and computation of $T^\bullet(S_K^{tor})$ for the Hilbert modular datum. In section 4 we choose an explicit Siegel embedding and explain that the pullback of the Hodge vector bundle splits into line bundles whose Chern classes generate $T^\bullet(S_K^{tor})$. This automatically gives a new relation coming from the vanishing of the top Chern class of the Hodge vector bundle in the Siegel case [vdG99, 1.2]. The final section tackles the problem of showing that there are no further relations. This reduces to an analysis of the highest degree part, which is at most d -dimensional. There are d elements given by multiples of $d-1$ generators and we show that they are linearly independent. This requires the properness of certain strata closures with a description of their cycle classes and is the novel difficulty in the Hilbert modular case: in the Siegel case the highest degree part is 1-dimensional so it sufficed to find a non-zero class.

Acknowledgements. *The author would like to thank W. Goldring for his support and guidance and D. Petersen for helpful discussions. The author is supported by grant KAW 2018.0356.*

2 Hilbert modular datum

We begin by recalling the Shimura datum which gives a Hilbert modular variety and describing the cocharacter datum one obtains from this in characteristic p .

Let F be a totally real field of degree $d > 1$ over \mathbb{Q} . Define \mathbf{G} as the algebraic group

$$\begin{array}{ccc} \mathbf{G} & \longrightarrow & \mathbb{G}_{m,\mathbb{Q}} \\ \downarrow & & \downarrow \\ Res_{F/\mathbb{Q}}GL_2 & \xrightarrow{\det} & Res_{F/\mathbb{Q}}\mathbb{G}_m \end{array}$$

Define \mathbf{X} as the $\mathbf{G}(\mathbb{R})$ -conjugacy class of the cocharacter $h: \mathbb{S} \rightarrow G_{\mathbb{R}}$ given on \mathbb{R} -points $\mathbb{C}^{\times} \rightarrow GL_2(\mathbb{R})^d$ by

$$x + iy \mapsto \left(\begin{pmatrix} x & y \\ y & -x \end{pmatrix} \right)_{i \leq d}$$

(\mathbf{G}, \mathbf{X}) is a Shimura datum, henceforth referred to as the Hilbert modular datum given by F . The associated Shimura variety is the Hilbert modular variety and is of PEL-type. Indeed, Raynaud constructed a model over \mathbb{Z} as a moduli space of principally-polarised abelian varieties with real multiplication by \mathcal{O}_F .

Definition 2.1. *Let p be a rational prime unramified in F . Fix an algebraic closure k of \mathbb{F}_p and a hyperspecial level $K = K^p K_p$. Denote X_K the special fibre of the integral model of the Hilbert modular variety at level K and $X_{K,k}$ its base change to k . Fix a smooth proper toroidal compactification X_K^{tor} , which exists by [MP19].*

Let G be the special fibre of a smooth reductive \mathbb{Z}_p -model of \mathbf{G} . We obtain a cocharacter datum (G, μ) as in [GK19, 1.2.2]. If p is inert then G is the fibre product

$$\begin{array}{ccc} G & \longrightarrow & \mathbb{G}_{m, \mathbb{F}_p} \\ \downarrow & & \downarrow \\ \text{Res}_{\mathbb{F}_{p^d}/\mathbb{F}_p} GL_2 & \xrightarrow{\det} & \text{Res}_{\mathbb{F}_{p^d}/\mathbb{F}_p} \mathbb{G}_m \end{array}$$

and if p splits then G is the fibre product

$$\begin{array}{ccc} G & \longrightarrow & \mathbb{G}_{m, \mathbb{F}_p} \\ \downarrow & & \downarrow \\ (GL_{2, \mathbb{F}_p})^d & \xrightarrow{\det} & (\mathbb{G}_{m, \mathbb{F}_p})^d \end{array}$$

Fix an embedding $\mathbb{F}_p \hookrightarrow k$ so we can identify $\{\tau: \mathbb{F}_p \hookrightarrow k\} \cong \mathbb{Z}/d\mathbb{Z}$. For any unramified prime, the group $G_{\mathbb{F}_{p^d}}$ can be described as the d -tuples in GL_2^d which have the same determinant in each factor. Any field extension $\mathbb{F}_{p^d} \subset K$ contains the field of definition $\kappa = \mathbb{F}_p$ of μ and

$$\mu_K: \mathbb{G}_{m, K} \longrightarrow G_K$$

is given by $t \mapsto \left(\left(\begin{pmatrix} t & 0 \\ 0 & 1 \end{pmatrix} \right)_{i \leq d}, t \right)$.

Let P be the parabolic subgroup associated with μ given on k -valued points $P(k) = \left\{ \left(\begin{pmatrix} a_i & b_i \\ 0 & d_i \end{pmatrix}, a_i d_i \right) \right\} \subset G(k)$ (this is called $P_-(\mu)$ in [Zie18], \mathbf{P}^- in [GK19]). This has a Levi subgroup $L = \text{Cent}(\mu) = \left\{ \left(\begin{pmatrix} a_i & 0 \\ 0 & d_i \end{pmatrix}, a_i d_i \right) \right\}$ which is a maximal torus T .

3 Tautological ring of a toroidal compactification

In this section we review the computation of $T^\bullet(X_K^{tor})$.

For a Hodge-type Shimura variety S_K in characteristic p , Wedhorn-Ziegler [Zie18] proved that $T^\bullet(S_K^{tor}) \cong A^\bullet(G\text{-Zip}^\mu)$, where $G\text{-Zip}^\mu$ is the stack introduced in [PWZ15] defined from the cocharacter datum (G, μ) and classifying certain semi-linear algebraic data. In characteristic p the morphism $\sigma: S_K \rightarrow [P \backslash *]$ given by the P -torsor on S_K appearing in the G -zip H_{dR}^1 factors as:

$$\begin{array}{ccc}
 S_K^{tor} & \xrightarrow{\zeta^{tor}} & G\text{-Zip}^\mu \xrightarrow{\beta} [P \backslash *] \\
 \uparrow j & \searrow \zeta & \\
 S_K & &
 \end{array}$$

Definition 3.1. Let S_K be a Hodge-type Shimura variety in characteristic p . Its tautological ring is defined $T^\bullet(S_K) := \text{Im}(\sigma^*) = \text{Im}(\zeta^* \circ \beta^*) \subset A^\bullet(S_K)$. Similarly, for a toroidal compactification $T^\bullet(S_K^{tor}) := \text{Im}(\zeta^{tor,*} \circ \beta^*)$.

In general, ζ is smooth ([Zha18, 3.1.2]) while smoothness of ζ^{tor} is expected to follow from [LS18] (and is known in the PEL case). Wedhorn-Ziegler proved that β^* is surjective and $\zeta^{tor,*}$ is injective, giving $T^\bullet(S_K^{tor}) := \text{Im}(\zeta^{tor,*} \circ \beta^*) \cong A^\bullet(G\text{-Zip}^\mu)$.

Now let (G, μ) be the cocharacter datum associated with X_K introduced in section 2. Then $\pi: G\text{-ZipFlag}^\mu \rightarrow G\text{-Zip}^\mu$ is an isomorphism, where $G\text{-ZipFlag}^\mu$ is the stack of zip flags defined in [GK19, 2.1]. Thus, on Chow rings

$$A^\bullet(G\text{-Zip}^\mu_k) \cong A^\bullet(G\text{-ZipFlag}^\mu_k)$$

The Weyl group $W = W_{G,T} = \{\pm 1\}^{\mathbb{Z}/d\mathbb{Z}}$ acts on the abelian group $X^*(T_k)$ via its action on T_k . This extends to an action on the symmetric algebra $S := \text{Sym}(X^*(T_k) \otimes_{\mathbb{Z}} \mathbb{Q})$ and we denote $I = S_+^W$. Then [Zie18, 8.2] gives

$$A^\bullet(G\text{-ZipFlag}^\mu_k) = S/I \cong \mathbb{Q}[z_1, \dots, z_d]/(z_1^2, \dots, z_d^2)$$

These generators $z_i \in S$ are given by characters

$$z_i: \begin{pmatrix} a_j & 0 \\ 0 & d_j \end{pmatrix} \mapsto d_i$$

in $X^*(T_k)$. Thus,

$$T^\bullet(X_{K,k}^{tor}) \cong \frac{\mathbb{Q}[z_1, \dots, z_d]}{(z_1^2, \dots, z_d^2)}$$

Remark 3.1. *This description is essentially independent of the prime. However, one has an action of Frobenius σ on S and this depends on the prime. For example, if p is inert then $\sigma \cdot z_i = z_{i+1}$ for all i , whereas if p is split then the action is trivial. Use σ to also refer to the corresponding permutation of $\{1, \dots, d\}$.*

Remark 3.2. *It is possible to interpret the relation z_i^2 geometrically. In the notation of section 5, there is a now-where vanishing section in $H^0(\overline{Z_{w_i}}, L_i \otimes L_{\sigma(i)}^p)$ giving $0 = c_1(L_i \otimes L_{\sigma(i)}^p |_{\overline{Z_{w_i}}}) = (z_i + pz_{\sigma(i)})(z_i - pz_{\sigma(i)})$ for each i .*

4 A Siegel embedding

Notation. *Denote $x_i := j^*(z_i) \in T^\bullet(X_{K,k})$. Denote $V(\rho)$ the vector bundle on $[G \backslash *]$ given by a representation ρ of G .*

In this section we prove that the relation $x_1 \dots x_d = 0$ holds in $T^\bullet(X_{K,k})$. Fixing a Siegel embedding gives a description of z_i in terms of the Hodge vector bundle. Namely, the pullback of the Hodge vector bundle \mathbb{E} splits on $X_{K,k}$ into line bundles ω_i with $x_i = c_1(\omega_i)$. The relation follows because $c_d(\mathbb{E}) = 0$ [vdG99, 1.2]. Note that we have made a judicious choice of embedding to give precisely $x_i = c_1(\omega_i)$; this is simply to ease computation. We will work with the following explicit description of the general symplectic group involved in the Siegel Shimura datum.

Example 4.1. *Consider the symplectic vector space $V = \mathbb{F}_p^{2d}$ with symplectic form given by $J := \begin{pmatrix} 0 & -I_d \\ I_d & 0 \end{pmatrix}$. Then $GS\!p(V) = \{M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \mid M^T J M = \lambda J \text{ for some } \lambda \in k\}$. Let $\mathcal{A}_{d,K'}$ the moduli of principally-polarised abelian varieties with level K' hyperspecial at p . Then as in section 3 there is a morphism $\sigma_d: \mathcal{A}_{d,K'} \rightarrow GS\!p(V)\text{-Zip} \rightarrow [P_d \backslash *]$.*

The Siegel parabolic $P_d = \left\{ \begin{pmatrix} A & B \\ 0 & D \end{pmatrix} \right\}$ has a Levi quotient isomorphic to $GL_d \times \mathbb{G}_m$. The Hodge vector bundle on $\mathcal{A}_{d,K'}$ is $\mathbb{E} := \sigma_d^* V(\rho_d)$ given by the representation $\rho_d: P_d \twoheadrightarrow GL_d \times \mathbb{G}_m \xrightarrow{P_1} GL_d$. Explicitly

$$\begin{pmatrix} A & B \\ 0 & D \end{pmatrix} \mapsto D$$

Now consider the standard embedding $\iota: G \hookrightarrow GS\!p(V) = GS\!p_{2d}$. This can be described by "embedded squares" over k :

$$\left(\begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix}, \lambda \right) \mapsto \begin{pmatrix} A & B \\ C & D \end{pmatrix}$$

where $A = \text{diag}(a_1, \dots, a_d)$, $B = \text{diag}(b_1, \dots, b_d)$ etc. The parabolic $P \subset G$ maps into the Siegel parabolic $P_d \subset GS\!p(V)$.

We can explicitly study the pullback of the Hodge vector bundle to $X_{K,k}$. The following diagram commutes, giving $\iota^*\mathbb{E} = \sigma^*V(\rho)$.

$$\begin{array}{ccc} \mathcal{A}_{d,K'} & \longrightarrow & [P_d \setminus *] \\ \uparrow & & \uparrow \\ X_{K,k} & \longrightarrow & [P \setminus *] \end{array}$$

Explicitly, $\rho: P \rightarrow P_d \rightarrow GL_d$ is given by

$$\begin{pmatrix} a_i & b_i \\ 0 & d_i \end{pmatrix} \mapsto D$$

where $D = \text{diag}(d_1, \dots, d_d)$. Consider the characters of P given by

$$\rho_j: \begin{pmatrix} a_i & b_i \\ 0 & d_i \end{pmatrix} \mapsto d_j$$

Then $\rho = \rho_1 \oplus \dots \oplus \rho_d$ and so $V(\rho) \simeq V(\rho_1) \oplus \dots \oplus V(\rho_d)$ as vector bundles on $[P \setminus *]$. Considering the commutative diagram

$$\begin{array}{ccccccc} X^*(P) & \longrightarrow & X^*(T) & & & & \\ \downarrow & & \downarrow & & & & \\ A^\bullet([P \setminus *]) & \longrightarrow & S & \longrightarrow & S/IS & \longrightarrow & T^\bullet(X_{K,k}) \end{array}$$

we see that $x_i = \sigma^*c_1(V(\rho_i))$.

Proposition 4.1. $x_1 \cdot \dots \cdot x_d = 0$ in $T^\bullet(X_{K,k})$.

Proof. On $X_{K,k}$ we have $\iota^*\mathbb{E} = \sigma^*V(\rho)$ so

$$x_1 \dots x_d = c_d(\sigma^*V(\rho)) = c_d(\iota^*\mathbb{E}) = \iota^*c_d(\mathbb{E}) = 0$$

with the last equality given in [vdG99, 1.2]. □

5 Positivity of classes: proof of Theorem 1

In this section we prove that there are no further relations.

Theorem 1. *Let F be a totally real field, p an unramified prime, k an algebraic closure of \mathbb{F}_p , K a hyperspecial level and $X_{K,k}$ the Hilbert modular variety at level K in characteristic p .*

$$T^\bullet(X_{K,k}) = \frac{\mathbb{Q}[x_1, \dots, x_d]}{(x_1^2, \dots, x_d^2, x_1 \cdot \dots \cdot x_d)}$$

The proof of Theorem 1 follows from:

Proposition 5.1. *Let p be a prime unramified in F . The d classes*

$$\gamma_i := x_1 \dots \widehat{x}_i \dots x_d$$

are linearly independent in $A^{d-1}(X_{K,k})$.

The closures of Ekedahl-Oort strata $\overline{X_{K,w}} = \zeta^{-1}(\overline{Z_w})$ give cycles in $T^\bullet(X_K)$ [Zie18, 6.1]. The codimension 1 strata are those indexed by $w_i = (\epsilon_j) \in \{\pm 1\}^d$ with $\epsilon_j = (-1)^{\delta_{ij}+1}$ (length $d-1$). There are sections s_i of $L_i \otimes L_{\sigma(i)}^{-p}$, where $L_i := \beta^*V(\rho_i)$ on $G\text{-Zig}^\mu$, such that $Z(s_i) = \overline{Z_{w_i}}$. To be precise, there is a separating system of partial Hasse invariants for the ordinary locus Z_{ord} as defined in [GK18, 3.4.2]. The divisors $D_i := Z(\zeta^*s_i) = \overline{X_{K,w_i}}$ are the Goren-Oort strata closures defined and shown to be smooth and proper in [GO00] and [Gor01]. Indeed, one can see by the modular description of X_K^{tor} that $\overline{X_{K,w_i}}$ doesn't intersect the toroidal boundary: a semi-abelian variety with real multiplication is either abelian or ordinary.

Proof of Proposition 5.1. Denote $\gamma_i^{tor} := z_1 \dots \widehat{z}_i \dots z_d$ in $A^{d-1}(X_{K,k}^{tor})$ so that $\gamma_i = j^* \gamma_i^{tor}$. We obtain the following commutative diagram.

$$\begin{array}{ccc}
 & X_{K,k}^{tor} & \\
 \nearrow & \uparrow & \searrow \zeta^{tor} \\
 D_i & j & G\text{-Zig}_k^\mu \\
 \searrow f_i & \downarrow & \nearrow \zeta \\
 & X_{K,k} &
 \end{array}$$

Assume that $\sum_{i=1}^d a_i \gamma_i = 0$ in $A^{d-1}(X_{K,k})$. For each i the projection formula [Ful98, 2.6(b)] for the morphism $f_i \circ j$ applied to the class $\sum_{i=1}^d a_i \gamma_i^{tor}$ in $A^{d-1}(X_{K,k}^{tor})$ gives

$$(f_i \circ j)_*(f_i \circ j)^* \sum_{i=1}^d a_i \gamma_i^{tor} = [D_i] \cdot \sum_{i=1}^d a_i \gamma_i^{tor}$$

$(f_i \circ j)^* \sum_{i=1}^d a_i \gamma_i^{tor} = f_i^! \sum_{i=1}^d a_i \gamma_i = 0$ by assumption. Since $[D_i] = \zeta^{tor,*}[\overline{Z_i}] = z_i - pz_{\sigma(i)}$ this implies that for all $i \in \mathbb{Z}/d\mathbb{Z}$

$$\begin{aligned}
 0 &= [D_i] \cdot \sum_{i=1}^d a_i \gamma_i^{tor} \\
 &= (z_i - pz_{\sigma(i)}) \cdot \sum_{i=1}^d a_i \gamma_i^{tor} \\
 &= (a_i - pa_{\sigma(i)})(z_1 \dots z_d)
 \end{aligned}$$

Since $\deg(z_1 \dots z_d) = d > 0$ we have that $a_i - pa_{\sigma(i)} = 0$ for all i . It remains to show that the only solution is $a_i = 0$ for all i . For p inert, $\sigma(i) = i + 1$ (modulo d) and this system of linear equations is given by the circulant matrix

$$A := \begin{pmatrix} 1 & -p & 0 & \dots & 0 \\ 0 & 1 & -p & & \\ \vdots & & \ddots & \ddots & \\ & & & 1 & -p \\ -p & & & & 1 \end{pmatrix}$$

A finite sequence of row operations gives an upper triangular matrix with diagonal entries $1, \dots, 1, 1 \pm p^n$ for some $n \neq 0$. The determinant of this matrix is non-zero, so the only solution to the system of linear equations is

$$a_1 = \dots = a_d = 0$$

A similar argument yields the other cases. Hence, $\gamma_1, \dots, \gamma_d$ are linearly independent. \square

The proof of Theorem 1 can now be completed by showing that there are no relations in lower degrees (below d).

Proof of Theorem 1. $T^\bullet(X_{K,k})$ is a quotient of $\frac{\mathbb{Q}[x_1, \dots, x_d]}{(x_1^2, \dots, x_d^2, x_1 \dots x_d)}$. The degree of the generators x_i is 1 so the maximum degree is degree $d - 1$. Suppose that there is a relation

$$\sum_{I \subset \{1, \dots, d\}, |I|=l} a_I z_I = 0$$

in $T^l(X_{K,k})$, indexed over multisets, with $l < d$. For any I_0 there exists some $i \notin I_0$. Let $J = \{1, \dots, i-1, i+1, \dots, d\} \setminus I_0$. Multiplying by z_J gives $\sum_I a_I z_{I \sqcup J} = 0$ in $T^{d-1}(X_{K,k})$. This can be rewritten uniquely as $\sum_I a_I z_{I \sqcup J} = \sum b_j \gamma_j$ by Proposition 5.1. If i is not in $I \sqcup J$ then either $I = I_0$ or there is a repeated index so $z_{I \sqcup J} = 0$. Hence, a_{I_0} is the only potentially non-zero coefficient of γ_i and by linear independence in $T^{d-1}(X_{K,k})$ we have $a_{I_0} = 0$. Since this holds for each I_0 we have that the relation is trivial.

Thus, there are no relations in lower degree and $T^\bullet(X_{K,k}) \cong \frac{\mathbb{Q}[x_1, \dots, x_d]}{(x_1^2, \dots, x_d^2, x_1 \dots x_d)}$ \square

References

- [Ful98] William Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 1998.

- [GK18] Wushi Goldring and Jean-Stefan Koskivirta. Automorphic vector bundles with global sections on G -zip ^{\mathbb{Z}} -schemes. *Compos. Math.*, 154(12):2586–2605, 2018.
- [GK19] Wushi Goldring and Jean-Stefan Koskivirta. Strata Hasse invariants, Hecke algebras and Galois representations. *Invent. Math.*, 217:887–984, 2019.
- [GO00] E. Z. Goren and F. Oort. Stratifications of Hilbert modular varieties. *J. Algebraic Geom.*, 9:111–154, 2000.
- [Gor01] Eyal Z. Goren. Hasse invariants for Hilbert modular varieties. *Israel J. Math.*, 122:157–174, 2001.
- [Kis10] Mark Kisin. Integral models for Shimura varieties of abelian type. *J. Amer. Math. Soc.*, 23(4):967–1012, 2010.
- [LS18] Kai-Wen Lan and Benoît Stroh. Compactifications of subschemes of integral models of Shimura varieties. *Forum Math. Sigma*, 6:Paper No. e18, 105, 2018.
- [Mil92] James S. Milne. The points on a Shimura variety modulo a prime of good reduction. In *The zeta functions of Picard modular surfaces*, pages 151–253. Univ. Montréal, Montreal, QC, 1992.
- [MP19] Keerthi Madapusi Pera. Toroidal compactifications of integral models of Shimura varieties of Hodge type. *Ann. Sci. Éc. Norm. Supér. (4)*, 52(2):393–514, 2019.
- [PWZ15] Richard Pink, Torsten Wedhorn, and Paul Ziegler. F -zips with additional structure. *Pacific J. Math.*, 274:183–236, 2015.
- [Vas99] Adrian Vasiu. Integral canonical models of Shimura varieties of pre-abelian type. *Asian J. Math.*, 3(2):401–518, 1999.
- [vdG99] Gerard van der Geer. Cycles on the moduli space of abelian varieties. In *Moduli of curves and abelian varieties*, pages 65–89. Friedr. Vieweg, Braunschweig, 1999.
- [Zha18] Chao Zhang. Ekedahl-Oort strata for good reductions of Shimura varieties of Hodge type. *Canad. J. Math.*, 70(2):451–480, 2018.
- [Zie18] Torsten Wedhorn; Paul Ziegler. Tautological rings of Shimura varieties and cycle classes of Ekedahl-Oort strata. *preprint*, 2018. <http://arxiv.org/abs/1811.04843>.