

# Involution factorizations of Ewens random permutations

Charles Burnette  
 Department of Mathematics  
 Xavier University of Louisiana  
 New Orleans, LA 70125  
 cburnet2@xula.edu

June 23, 2021

## Abstract

An involution is a bijection that is its own inverse. Given a permutation  $\sigma$  of  $[n]$ , let  $\text{invol}(\sigma)$  denote the number of ways to express  $\sigma$  as a composition of two involutions of  $[n]$ . The statistic  $\text{invol}$  is asymptotically lognormal when the symmetric groups  $\mathfrak{S}_n$  are each equipped with Ewens Sampling Formula probability measures of some fixed positive parameter  $\theta$ . This paper strengthens and generalizes previously determined results on the limiting distribution of  $\log(\text{invol})$  for uniform random permutations, i.e. the specific case of  $\theta = 1$ . We also investigate the first two moments of  $\text{invol}$  itself.

**Keywords:** asymptotic lognormality, Erdős-Turán law, Ewens Sampling Formula, Feller coupling, involutions, Mellin transform, saddle-point method.

**2020 Mathematics Subject Classification:** 05A05, 05A16, 60C05

## 1 Introduction

An involution is a bijection that is its own inverse. The involutions in the symmetric group  $\mathfrak{S}_n$  are precisely those permutations whose cycles are all of length 1 or 2. This paper is concerned with decomposing random permutations into two involutive factors and the probabilistic analysis thereof. Recent interest in involution factorizations in  $\mathfrak{S}_n$  can be traced to Roberts and Vivaldi [31] who proposed that the dynamical behavior of reversible rational maps over a finite field can be modeled by compositions of two random involutions with a prescribed number of fixed points. Unrestricted products of two involutions, however, are of singular interest as well. For instance, Petersen and Tenner [30] outline hypothetical links between enumerating involution factorizations, rim-hook tableaux, and the irreducible characters of  $\mathfrak{S}_n$  that are still in need of further clarification. Other applications have surfaced in comparative genomics [6], [11] and the analysis of algorithms [21], [35].

If  $\sigma$  is a permutation of  $[n]$ , let  $\text{invol}(\sigma)$  denote the number of function compositional factorizations of  $\sigma$  into the product of two involutions of  $[n]$ , that is, the number of ordered pairs  $(\tau_1, \tau_2)$  of involutions in  $\mathfrak{S}_n$  such that  $\sigma = \tau_2 \circ \tau_1$ . It is known that

$$\text{invol}(\sigma) = \mathbf{B}(\sigma) \prod_{k=1}^n \sum_{j=0}^{\lfloor c_k/2 \rfloor} \frac{c_k!}{(2k)^j j! (c_k - 2j)!}, \quad (1.1)$$

where  $c_k = c_k(\sigma)$  denotes the number of cycles of length  $k$  (or, more succinctly,  $k$ -cycles) that  $\sigma$  possesses and  $\mathbf{B}(\sigma) = \prod k^{c_k}$  is the product of its cycle lengths. To the author's knowledge, the earliest complete proofs of (1.1) are independently attributed to Lugo [26] and Petersen and Tenner [30]. Both of their derivations rely on identifying involutions with partial matchings and, consequently, the composition of two involutions with 2-edge-colored graphs in which every vertex is incident to exactly two edges, one edge per color. A derivation of (1.1) that merely exploits the canonical group action of  $\mathfrak{S}_n$  on  $[n]$  appears in [10].

From (1.1), one notices that  $\text{invol}(\sigma)$  is a class function; its value depends only on the cycle structure of  $\sigma$ . It is therefore natural to wonder what the distribution of  $\text{invol}$  is for permutations drawn from conjugation-invariant probability measures on  $\mathfrak{S}_n$  where cycles are multiplicatively weighted. The archetypal example of such a measure is the Ewens Sampling Formula with parameter  $\theta > 0$ , hereafter abbreviated  $\text{ESF}(\theta)$ . (Read Crane's survey article [12] for a lucid and concise synopsis of the literature on  $\text{ESF}(\theta)$ .)

For each positive integer  $n$ , the Ewens probability measure  $\mathbb{P}_{\theta,n}$  is a deformation of the uniform measure  $\nu_n$  on  $\mathfrak{S}_n$  obtained by incorporating the following sampling bias:

$$\mathbb{P}_{\theta,n}(\sigma = \sigma_0) = \frac{n! \theta^K}{\theta^{(n)}} \nu_n(\sigma = \sigma_0), \quad (1.2)$$

where  $K = \sum c_k$  is the total number of cycles that  $\sigma$  has and  $\theta^{(n)}$  is the rising factorial  $\theta(\theta+1)\cdots(\theta+n-1)$ . (Note, in particular, that  $\mathbb{P}_{1,n} = \nu_n$ .) More specifically, the probability that a permutation of  $[n]$  chosen according to  $\text{ESF}(\theta)$  has cycle index  $(a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n$  is

$$\mathbb{P}_{\theta,n}(c_k = a_k \text{ for } k = 1, \dots, n) = \mathbb{1} \left\{ \sum_{\ell=1}^n \ell a_\ell = n \right\} \frac{n!}{\theta^{(n)}} \prod_{k=1}^n \left( \frac{\theta}{k} \right)^{a_k} \frac{1}{a_k!}, \quad (1.3)$$

where  $\mathbb{1}\{\cdot\}$  denotes the indicator function. We shall also refer to random permutations chosen from  $\text{ESF}(\theta)$  as  $\theta$ -weighted permutations, just as Lugo does in [27].

A pivotal feature of  $\text{ESF}(\theta)$  is that the joint distribution of the cycle counts  $c_k$  converges to that of independent Poisson random variables of respective intensity  $\theta/k$ . Now consider a sequence  $\alpha = (\alpha_1, \alpha_2, \dots)$  of mutually independent random variables with  $\alpha_k \sim \text{Poisson}(\theta/k)$ , for  $k = 1, 2, \dots$ , and define

$$\text{invol}_n(\alpha) = \mathbf{B}_n(\alpha) \prod_{k=1}^n \sum_{j=0}^{\lfloor \alpha_k/2 \rfloor} \frac{\alpha_k!}{(2k)^j j! (\alpha_k - 2j)!}, \quad \text{where} \quad \mathbf{B}_n(\alpha) = \prod_{k=1}^n k^{\alpha_k}. \quad (1.4)$$

Note that

$$\mathbb{E}|\log \text{invol}_n(\alpha) - \log \mathbf{B}_n(\alpha)| = \sum_{k=1}^n \mathbb{E} \log \left( \sum_{j=0}^{\lfloor \alpha_k/2 \rfloor} \frac{\alpha_k!}{(2k)^j j! (\alpha_k - 2j)!} \right) \quad (1.5)$$

$$\leq \sum_{k=1}^n \frac{1}{2k} \mathbb{E} \alpha_k^2 = O(1), \quad (1.6)$$

whereas

$$\text{Var}(\log \mathbf{B}_n(\alpha)) = \sum_{k=1}^{\infty} \frac{\theta \log^2 k}{k} = \frac{\theta}{3} \log^3 n (1 + o(1)). \quad (1.7)$$

Markov's inequality thus dictates that

$$\frac{|\log \text{invol}_n(\alpha) - \log \mathbf{B}_n(\alpha)|}{\sqrt{\text{Var}(\log \mathbf{B}_n(\alpha))}} \xrightarrow{\mathbb{P}} 0 \quad (1.8)$$

One would then suspect that something similar holds for  $\log \text{invol}$  and  $\log \mathbf{B}$  in  $\text{ESF}(\theta)$ , assuming both are plausibly well-approximated by their Poissonized counterparts.

In this paper, we will establish both central limit and functional limit theorems for  $\log(\text{invol})$  over Ewens random permutations. This shall be accomplished by generalizing the key results of [10]. It was particularly shown in [10] that  $\text{invol}$  is asymptotically lognormal when each  $\mathfrak{S}_n$  is endowed with the uniform probability measure. The proof of this hinged on the observation that, for asymptotically almost all permutations  $\sigma \in \mathfrak{S}_n$ , the statistic  $\text{invol}(\sigma)$  is readily controlled by the simpler statistic  $\mathbf{B}(\sigma)$  together with the fact that  $\mathbf{B}$  is itself an asymptotically lognormal random variable over uniform random permutations. As we shall soon witness, these facts remain true in the setting of  $\text{ESF}(\theta)$ .

On the other hand, the distribution of  $\text{invol}$  itself, when left untouched by any rescaling or standardization, is heavily right-skewed. Essentially all of its range is concentrated far below its expected value. To demonstrate this, we will derive the leading-term asymptotics of the first and second moments of  $\text{invol}$  for general  $\theta$ -weighted permutations. The latter will highlight certain discrepancies in the variability of  $\text{invol}$  for  $0 < \theta \leq 1$  versus  $\theta > 1$ .

Some classical results and techniques from asymptotic analysis are cited in a piecemeal fashion throughout this paper. As is common for random permutation and integer partition statistics, we will encounter generating functions that demand a careful examination by way of the saddle-point method and Mellin transforms. Technical preliminaries are briefly discussed in an appendix at the end.

## Acknowledgments

This paper refines various elements of the author's doctoral thesis. The author wishes to thank his advisor, Eric Schmutz, for his extensive guidance. The author also expresses his

gratitude to Dirk Zeindler for suggesting that the results of [10] can be generalized to the Ewens Sampling Formula, and Hsien-Kuei Hwang for technical discussions on how to tackle the variance of invol.

## 2 The Expected and Typical Values of invol

Let  $\mathbb{P}_n$  denote the probability measure on  $\mathfrak{S}_n$  corresponding to  $\text{ESF}(\theta)$ . (For the sake of notational brevity, we will henceforth omit the  $\theta$  in  $\mathbb{P}_{\theta,n}$  and associated operators.)

**Theorem 2.1.** *The average number of involution factorizations for a permutation chosen according to the Ewens measure  $\mathbb{P}_n$  is*

$$\mathbb{E}_n \text{invol} = \frac{\theta^{\frac{1-\theta^2}{4}} \Gamma(\theta)}{2^{\frac{\theta^2}{2}+1} e^{\frac{\theta}{2}} \sqrt{\pi}} e^{2\sqrt{\theta n}} n^{\frac{\theta^2}{4}-\theta+\frac{1}{4}} \left(1 + O(n^{-\frac{1}{2}})\right).$$

*Proof.* For each  $\lambda \in \mathbb{Z}_{\geq 0}^n$ , set  $\mathcal{A}_\lambda = \{\sigma \in \mathfrak{S}_n : \sigma \text{ has cycle type } 1^{\lambda_1} \cdots n^{\lambda_n}\}$ . Consider the power series

$$F(z) = 1 + \sum_{n=1}^{\infty} (\mathbb{E}_n \text{invol}) \frac{\theta^{(n)}}{n!} z^n. \quad (2.1)$$

By the law of total expectation,

$$F(z) = 1 + \sum_{n=1}^{\infty} \left( \sum_{\lambda \vdash n} \mathbb{P}_n(\sigma \in \mathcal{A}_\lambda) \mathbb{E}_n(\text{invol}(\sigma) \mid \sigma \in \mathcal{A}_\lambda) \right) \frac{\theta^{(n)}}{n!} z^n, \quad (2.2)$$

where  $\lambda \vdash n$  means that  $\lambda \in \mathbb{Z}_{\geq 0}^n$  and  $\sum k \lambda_k = n$ . As pointed out in Theorem 2.5 of [10],

$$\text{invol}(\sigma) = \prod_{k=1}^n k^{c_k(\sigma)} \frac{He_{c_k(\sigma)}(i\sqrt{k})}{(i\sqrt{k})^{c_k(\sigma)}}, \quad (2.3)$$

where  $He_m$  is the probabilists' Hermite polynomial  $He_m(x) = m! \sum_{r=0}^{\lfloor m/2 \rfloor} \frac{(-1)^r}{r!(m-2r)!} \frac{x^{m-2r}}{2^r}$ . Thus

$$F(z) = 1 + \sum_{n=1}^{\infty} \left( \sum_{\lambda \vdash n} \left( \frac{n!}{\theta^{(n)}} \prod_{k=1}^n \left( \frac{\theta}{k} \right)^{\lambda_k} \frac{1}{\lambda_k!} \right) \left( \prod_{k=1}^n k^{\lambda_k} \frac{He_{\lambda_k}(i\sqrt{k})}{(i\sqrt{k})^{\lambda_k}} \right) \right) \frac{\theta^{(n)}}{n!} z^n \quad (2.4)$$

$$= \prod_{k=1}^{\infty} \left( \sum_{\lambda_k=0}^{\infty} \frac{He_{\lambda_k}(i\sqrt{k})}{\lambda_k!} \left( \frac{\theta z^k}{i\sqrt{k}} \right)^{\lambda_k} \right) \quad (2.5)$$

Invoking the exponential generating function  $\exp(xt - t^2/2) = \sum_{n=0}^{\infty} He_n(x) \frac{t^n}{n!}$  (an assortment of derivations are available in [23], [24], [25]), we see that

$$F(z) = \prod_{k=1}^{\infty} \exp\left(\theta z^k + \frac{\theta^2 z^{2k}}{2k}\right) = \frac{\exp(\theta z/(1-z))}{(1-z^2)^{\frac{\theta^2}{2}}}. \quad (2.6)$$

Hence

$$\mathbb{E}_n \text{invol} = \frac{n!}{\theta^{(n)}} [z^n] \frac{\exp(\theta z / (1 - z))}{(1 - z^2)^{\frac{\theta^2}{2}}}. \quad (2.7)$$

Applying Corollary 6.4 with  $\beta = -\theta^2/2$ ,  $\phi(z) = e^{-\theta}(1+z)^{-\frac{\theta^2}{2}}$ , and  $\alpha = \theta$ , together with the Stirling's approximations  $n! = \sqrt{2\pi n} e^{-n} n^n (1 + O(n^{-1}))$  and  $\theta^{(n)} = \frac{\Gamma(\theta+n)}{\Gamma(\theta)} = \frac{\sqrt{2\pi}}{\Gamma(\theta)} e^{-n} n^{n+\theta-\frac{1}{2}} (1 + O(n^{-1}))$  completes the proof.  $\square$

Another consequence of the proof of Theorem 2.1 is that

$$F(u, z) = \frac{\exp(uz/(1-z))}{(1-z^2)^{\frac{u^2}{2}}} \quad (2.8)$$

is the bivariate generating function for the expected value of  $\mathbb{1}\{K(\sigma) = m\} \text{invol}(\sigma)$  over uniform random permutations, counted according to the size of the ground set (indicated by the variable  $z$ ) and the total number of cycles (indicated by the variable  $u$ ). The diagonal of  $F(u, z)$  is given by

$$\lim_{z \rightarrow 0} F(x/z, z) = e^{x + \frac{x^2}{2}}, \quad (2.9)$$

which is the exponential generating function for the class of involutions. This makes sense, as the only permutation of  $[n]$  with  $n$  cycles is the identity permutation, which is precisely the square of all involutions in  $\mathfrak{S}_n$ . We can also retrieve the conditional expectation of  $\text{invol}$  for permutations with  $m$  cycles from the ‘‘vertical’’ generating function

$$[u^m] F(u, z) = \sum_{k=\lceil m/2 \rceil}^m \frac{\binom{k}{m-k}}{k! 2^{m-k}} \left( \frac{z}{1-z} \right)^{2k-m} \left( \log \frac{1}{1-z^2} \right)^{m-k}. \quad (2.10)$$

For example,  $[u] F(u, z) = z/(1-z)$ , and so  $[uz^n] F(u, z) = 1$  for  $n \geq 1$ . This too makes sense as the probability of picking an  $n$ -cycle is  $1/n$ , and there are precisely  $n$  ways to write an  $n$ -cycle as a product of two involutions.

**Theorem 2.2.** *For each positive integer  $m$ ,*

$$\mathbb{E}_n(\text{invol}(\sigma) \mid K(\sigma) = m) = \frac{n^m}{m! (\log n)^{m-1}} \left( 1 + O\left( \frac{1}{\log n} \right) \right)$$

as  $n \rightarrow \infty$ .

*Proof.* Because  $K$  is a sufficient statistic for the underlying parameter  $\theta$ , the conditional distribution of  $\text{invol}$  does not depend on  $\theta$ , and so

$$\mathbb{E}_n(\text{invol}(\sigma) \mid K(\sigma) = m) = \frac{[u^m z^n] F(u, z)}{[m] / n!}, \quad (2.11)$$

where  $\begin{bmatrix} n \\ m \end{bmatrix}$  are the unsigned Stirling numbers of the first kind. Based on the Flajolet-Odlyzko transfer theorem [18]

$$[z^n](1-z)^\alpha \left( \log \frac{1}{1-z} \right)^k = \frac{1}{\Gamma(-\alpha)} n^{-\alpha-1} (\log n)^k + O(n^{-\alpha-2} (\log n)^k) \text{ for } \alpha \notin \mathbb{Z}_{\geq 0}, \quad (2.12)$$

we find that

$$[u^m z^n] F(u, z) = \frac{1}{m!} [z^n] \left( \frac{z}{1-z} \right)^m + O(n^{m-2}) = \frac{n^{m-1}}{m!(m-1)!} + O(n^{m-2}) \quad (2.13)$$

Use the following crude simplification of Wilf's asymptotic [32]

$$\frac{\begin{bmatrix} n \\ m \end{bmatrix}}{n!} = \frac{(\log n)^{m-1}}{(m-1)!n} + O\left(\frac{(\log n)^{m-2}}{n}\right) \quad (2.14)$$

to finish the calculation. □

Most of the contribution to the mean comes from a small subset of permutations that admit an exceptionally large number of involution factorizations. Scrupulous glances at (1.1) should convince the reader that giving a permutation  $\sigma$  numerous occurrences of one very long cycle length or an exorbitant number of cycles of one distinct length exaggerates the magnitude of  $\text{invol}(\sigma)$ . But it is rare for a permutation to have such a cycle profile. The *typical* permutation has much fewer involution factorizations. Direct combinatorial arguments can verify this for uniform random permutations, which have roughly  $e^{2\sqrt{n}}/\sqrt{8\pi en}$  involution factorizations on average.

**Theorem 2.3.** *Let  $\xi_1$  and  $\xi_2$  be integers such that  $20 \leq \xi_1, \xi_2 \leq n$  and  $\xi_1 T_{\xi_2} < n$ , where  $T_r$  denotes the  $r^{\text{th}}$  triangular number. Then at least  $n!(2 - \exp(15/(\xi_1+1)! + 3/\xi_2))$  permutations  $\sigma \in \mathfrak{S}_n$  satisfy the inequality*

$$\text{invol}(\sigma) < (\xi_2!)^{\frac{\xi_1}{2}-1} \lfloor (2(n - (\xi_1 - 1)T_{\xi_2}))^{\frac{1}{2}} \rfloor \frac{(\xi_1!)^{\xi_2} \exp\left(\frac{\xi_1 \xi_2}{2} + \frac{2}{3} \xi_2^{\frac{3}{2}} \sqrt{\xi_1 + 1} + \sqrt{\xi_2(\xi_1 + 1)} + \frac{\xi_2}{2}\right)}{\xi_1^{\xi_1 \xi_2 / 2}}.$$

*Proof.* Let  $\mathcal{P}_{\xi_1, \xi_2}$  be the set of all permutations  $\sigma \in \mathfrak{S}_n$  for which  $c_k(\sigma) \leq \xi_1$  for all  $k \leq \xi_2$  and  $c_k(\sigma) \leq 1$  for all  $k > \xi_2$ . Erdős and Turán proved in Lemma III of [15] that

$$1 - \frac{|\mathcal{P}_{\xi_1, \xi_2}|}{n!} \leq e^{\frac{15}{(\xi_1+1)!} + \frac{3}{\xi_2}} - 1. \quad (2.15)$$

The maximum value of  $\text{invol}(\sigma)$  over  $\mathcal{P}_{\xi_1, \xi_2}$  is attained by a permutation with the most cycles possible (cf Proposition 3.3.1 in [9]). To wit, the multiplicities of the shortest cycles must be optimal. So if we set

$$\xi_3 = \min\{r \in \mathbb{Z} : T_r + (\xi_1 - 1)T_{\xi_2} \geq n\} = \lfloor 2(n - (\xi_1 - 1)T_{\xi_2})^{\frac{1}{2}} \rfloor \quad (2.16)$$

and apply (6.1) and (6.2), then for every  $\sigma \in \mathcal{P}_{\xi_1, \xi_2}$ ,

$$\text{invol}(\sigma) = \mathbf{B}(\sigma) \prod_{k=1}^{\xi_2} \sum_{j=0}^{\lfloor c_k(\sigma)/2 \rfloor} \frac{c_k(\sigma)!}{(2k)^j j! (c_k(\sigma) - 2j)!} \quad (2.17)$$

$$< \left( \prod_{k=1}^{\xi_2} k^{\xi_1} \right) \cdot \left( \prod_{k=\xi_2+1}^{\xi_3} k \right) \cdot \left( \prod_{k=1}^{\xi_2} \frac{\xi_1! \exp\left(\frac{\xi_1}{2} + \sqrt{k(\xi_1+1)} + \frac{1}{2}\right)}{(k\xi_1)^{\frac{\xi_1}{2}}} \right) \quad (2.18)$$

$$= (\xi_2!)^{\frac{1}{2}\xi_1-1} \xi_3! \frac{(\xi_1!)^{\xi_2} \exp\left(\frac{\xi_1\xi_2}{2} + \frac{2}{3}\xi_2^{\frac{3}{2}}\sqrt{\xi_1+1} + \sqrt{\xi_2(\xi_1+1)} + \frac{\xi_2}{2}\right)}{\xi_1^{\frac{\xi_1\xi_2}{2}}}. \quad (2.19)$$

□

Setting  $\xi_1 = \xi_2$  in Theorem 2.3 and employing (6.3) along with Stirling's approximation begets the following corollary.

**Corollary 2.4.** *If  $\xi = \xi(n) \rightarrow \infty$ , but slowly enough so that  $\xi < \sqrt[3]{2n}$ , then all but  $o(n!)$  permutations  $\sigma \in \mathfrak{S}_n$  satisfy the inequality*

$$\text{invol}(\sigma) < (2\pi)^{\frac{3}{4}\xi-1} \xi^{\xi^2-\frac{1}{4}\xi-1} (2n - \xi^3)^{\frac{1}{2}} \sqrt{2n - \xi^3} e^{2\xi - \frac{1}{3}\xi^2}.$$

In particular, taking  $\xi = o(n^{1/4})$  produces an a.a.s. upper bound which is itself  $o(\mathbb{E}_n \text{invol})$ . The upcoming Lemma 3.2 shows that this holds for all  $\theta$ -weighted permutations, regardless of what  $\theta$  is. However, we can conclude something even stronger. Based on Theorem 3.4, if  $\sigma$  is a  $\theta$ -weighted permutation of  $[n]$  and  $\epsilon > 0$ , then w.h.p.,

$$e^{(\frac{\theta}{2}-\epsilon)\log^2 n} < \text{invol}(\sigma) < e^{(\frac{\theta}{2}+\epsilon)\log^2 n}.$$

### 3 A Central Limit Theorem for $\log \text{invol}$

In this section, we emulate the proof of Theorem 4.3 in [10], except now each  $\mathfrak{S}_n$  is equipped with the ESF( $\theta$ ) probability measure  $\mathbb{P}_n$  with  $\theta > 0$  fixed. We proceed in three steps. First,  $\text{invol}(\sigma)$  is well-approximated by  $\mathbf{B}(\sigma)$  for permutations  $\sigma$  in which only  $k$ -cycles with  $k$  under a certain threshold  $\xi$ , to be stipulated *a posteriori*, can occur more than once but still not more than  $\xi$  times. Call the set of all such permutations  $\mathcal{P}_\xi$ . The second step is to show that  $\mathbb{P}_n(\sigma \in \mathcal{P}_\xi) \rightarrow 1$  as  $n \rightarrow \infty$ . Lastly, the previous two steps are combined to compare the distribution of  $\log \text{invol}$  to the distribution of  $\log \mathbf{B}$ .

There are two notable improvements here over the presentation of [9] and [10]. Our present proof of the aforementioned second step is cleaner and considerably streamlined. Furthermore, a more concrete error term for the approximate Gaussian law is added. Perhaps unsurprisingly, the error term matches the convergence rate of the Erdős-Turán limit law for the order of a random permutation that was first determined in [5].

**Lemma 3.1.** (Burnette, Schmutz) *Suppose  $\sigma \in \mathcal{P}_\xi$ . Then there is a constant  $c > 0$ , not dependent on  $\sigma$  nor  $\xi$ , such that  $\mathbf{B}(\sigma) \leq \text{invol}(\sigma) \leq \mathbf{B}(\sigma) \cdot (c\xi^\xi)^\xi$ .*

Lemma 3.1 is entirely deterministic and thus needs not be revisited. The next lemma generalizes Lemma III in [15].

**Lemma 3.2.** *If  $\xi = \xi(n) \rightarrow \infty$ , then  $\mathbb{P}_n(\sigma \notin \mathcal{P}_\xi) = O(\frac{1}{\xi})$ .*

*Proof.* Due to (1.3) and the exponential formula for labeled combinatorial structures,

$$\mathbb{P}_n(\sigma \in \mathcal{P}_\xi) = \frac{n!}{\theta^{(n)}} [z^n] \left[ \left( \prod_{k=1}^{\lfloor \xi \rfloor} \sum_{j=1}^{\lfloor \xi \rfloor} \left( \frac{\theta z^k}{k} \right)^j \frac{1}{j!} \right) \cdot \left( \prod_{k=\lfloor \xi \rfloor+1}^{\infty} \left( 1 + \frac{\theta z^k}{k} \right) \right) \right] \quad (3.1)$$

The product in (3.1) can be written as  $(1-z)^{-\theta} \Omega(z)$ , where

$$\Omega(z) = \left( \prod_{k=1}^{\lfloor \xi \rfloor} \left( 1 - e^{-\frac{\theta z^k}{k}} \sum_{j=\lfloor \xi \rfloor+1}^{\infty} \left( \frac{\theta z^k}{k} \right)^j \frac{1}{j!} \right) \right) \cdot \left( \prod_{k=\lfloor \xi \rfloor+1}^{\infty} \left( 1 - e^{-\frac{\theta z^k}{k}} \sum_{j=2}^{\infty} \left( \frac{\theta z^k}{k} \right)^j \frac{1}{j!} \right) \right) \quad (3.2)$$

Equating the corresponding coefficients reveals that

$$\mathbb{P}_n(\sigma \in \mathcal{P}_\xi) - 1 = \frac{n!}{\theta^{(n)}} \sum_{\ell=1}^n \binom{\theta + n - \ell - 1}{n - \ell} [z^\ell] \Omega(z) \leq \sum_{\ell=1}^n [z^\ell] \Omega(z) \quad (3.3)$$

If we replace every instance of the term  $-e^{-\frac{\theta z^k}{k}}$  by  $e^{\frac{\theta z^k}{k}}$  in each factor of  $\Omega(z)$ , we obtain an entire function  $\Omega^*(z)$  whose coefficients are all positive and majorize the absolute values of the corresponding coefficients of  $\Omega(z)$ . Hence

$$\mathbb{P}_n(\sigma \notin \mathcal{P}_\xi) = 1 - \mathbb{P}_n(\sigma \in \mathcal{P}_\xi) \quad (3.4)$$

$$\leq \sum_{\ell=1}^n [z^\ell] \Omega^*(z) \quad (3.5)$$

$$< \Omega^*(1) - \Omega^*(0) \quad (3.6)$$

$$= \left( \prod_{k=1}^{\lfloor \xi \rfloor} \left( 1 + e^{\frac{\theta}{k}} \sum_{j=\lfloor \xi \rfloor+1}^{\infty} \left( \frac{\theta}{k} \right)^j \frac{1}{j!} \right) \right) \cdot \left( \prod_{k=\lfloor \xi \rfloor+1}^{\infty} \left( 1 + e^{\frac{\theta}{k}} \sum_{j=2}^{\infty} \left( \frac{\theta}{k} \right)^j \frac{1}{j!} \right) \right) - 1 \quad (3.7)$$

For sufficiently large  $n$ , we have

$$\prod_{k=1}^{\lfloor \xi \rfloor} \left( 1 + e^{\frac{\theta}{k}} \sum_{j=\lfloor \xi \rfloor+1}^{\infty} \left( \frac{\theta}{k} \right)^j \frac{1}{j!} \right) < \exp \left( \sum_{k=1}^{\lfloor \xi \rfloor} e^{\frac{\theta}{k}} \sum_{j=\lfloor \xi \rfloor+1}^{\infty} \left( \frac{\theta}{k} \right)^j \frac{1}{j!} \right) \quad (3.8)$$



$$< \exp\left(\frac{e^{2\theta}\theta^{[\xi]+1}}{([\xi]+1)!} \sum_{k=1}^{[\xi]} \frac{1}{k^{[\xi]+1}}\right) \quad (3.9)$$

$$< \exp\left(\frac{K_1\theta^{[\xi]+1}}{([\xi]+1)!}\right) \quad (3.10)$$

for some constant  $K_1$ , and

$$\prod_{k=[\xi]+1}^{\infty} \left(1 + e^{\frac{\theta}{k}} \sum_{j=2}^{\infty} \left(\frac{\theta}{k}\right)^j \frac{1}{j!}\right) < \exp\left(\sum_{k=[\xi]}^{\infty} e^{\frac{\theta}{k}} \sum_{j=2}^{\infty} \left(\frac{\theta}{k}\right)^j \frac{1}{j!}\right) \quad (3.11)$$

$$< \exp\left(\frac{e^{\theta+2\theta^2}}{2} \sum_{k=[\xi]}^{\infty} \frac{1}{k^2}\right) \quad (3.12)$$

$$< \exp\left(\frac{K_2}{\xi}\right) \quad (3.13)$$

for some constant  $K_2$ . Therefore, as  $n \rightarrow \infty$ ,

$$\mathbb{P}_n(\sigma \notin \mathcal{P}_\xi) < \exp\left(\frac{K_1\theta^{[\xi]+1}}{([\xi]+1)!} + \frac{K_2}{\xi}\right) - 1 = O\left(\frac{1}{\xi}\right). \quad (3.14)$$

□

We now cite the asymptotic lognormality of  $\mathbf{B}$  for  $\theta$ -weighted permutations.

**Lemma 3.3.** (Arratia, Barbour, Tavaré)

$$\sup_{x \in \mathbb{R}} \left| \mathbb{P}_n\left(\frac{\log \mathbf{B}(\sigma) - \mu_n}{\sigma_n} \leq x\right) - \Phi(x) \right| = O\left(\frac{1}{\sqrt{\log n}}\right),$$

where  $\mu_n := \sum_{k=1}^n \frac{\theta \log k}{k} \sim \frac{\theta}{2} \log^2 n$ ,  $\sigma_n^2 := \sum_{k=1}^n \frac{\theta \log^2 k}{k} \sim \frac{\theta}{3} \log^3 n$ , and  $\Phi(x) := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt$ .

Erdős and Turán [16] were the first to confirm the asymptotic lognormality of  $\mathbf{B}$  for uniform random permutations. Progressively sharper versions of the Erdős-Turán limit law have emerged on multiple occasions, e.g. [8], [13], [28], [29]. Proofs of Lemma 3.3 itself appear in the aggregate works of Arratia, Barbour, and Tavaré [2], [3], [4], [5].

**Theorem 3.4.**

$$\sup_{x \in \mathbb{R}} \left| \mathbb{P}_n\left(\frac{\log \text{invol}(\sigma) - \mu_n}{\sigma_n} \leq x\right) - \Phi(x) \right| = O\left(\frac{1}{\sqrt{\log n}}\right)$$

*Proof.* Because  $\text{invol}(\sigma) \geq \mathbf{B}(\sigma)$  for all  $\sigma \in \mathfrak{S}_n$ , the upper bound is immediate:

$$\mathbb{P}_n\left(\frac{\log \text{invol}(\sigma) - \mu_n}{\sigma_n} \leq x\right) \leq \mathbb{P}_n\left(\frac{\log \mathbf{B}(\sigma) - \mu_n}{\sigma_n} \leq x\right) = \Phi(x) + O\left(\frac{1}{\sqrt{\log n}}\right). \quad (3.15)$$

For the lower bound, we capitalize on the uniform continuity of  $\Phi$ . We also invoke the conditional bound from Lemma 3.1, where  $\xi$  will now be chosen to our liking. Due to Lemma 3.2, this bound holds with probability  $1 - O(\frac{1}{\xi})$ .

The remainder of the proof is virtually identical to that of Theorem 4.3 in [10], which is included here for the sake of completeness. Let  $\epsilon > 0$  be a fixed but arbitrarily small positive number. We can choose  $\delta > 0$  so that  $|\Phi(x) - \Phi(a)| < \epsilon$  whenever  $|x - a| < \delta$ . If we choose  $\xi = \sqrt{\log n}$ , then we have  $\log((c\xi^\xi)^\xi) = o(\sigma_n)$ . Therefore we can choose  $N_\epsilon$  so that, for all  $n \geq N_\epsilon$ , we have  $\log((c\xi^\xi)^\xi) < \frac{\delta\sigma_n}{2}$ . It follows that, uniformly in  $x$ ,

$$\mathbb{P}_n\left(\frac{\log \text{invol}(\sigma) - \mu_n}{\sigma_n} \leq x\right) \geq \mathbb{P}_n\left(\frac{\log \mathbf{B}(\sigma) + \log((c\xi^\xi)^\xi) - \mu_n}{\sigma_n} \leq x\right) + O\left(\frac{1}{\sqrt{\log n}}\right) \quad (3.16)$$

$$\geq \mathbb{P}_n\left(\frac{\log \mathbf{B}(\sigma) + \delta\sigma_n/2 - \mu_n}{\sigma_n} \leq x\right) + O\left(\frac{1}{\sqrt{\log n}}\right) \quad (3.17)$$

$$= \mathbb{P}_n\left(\frac{\log \mathbf{B}(\sigma) - \mu_n}{\sigma_n} \leq x - \frac{\delta}{2}\right) + O\left(\frac{1}{\sqrt{\log n}}\right) \quad (3.18)$$

$$= \Phi\left(x - \frac{\delta}{2}\right) + O\left(\frac{1}{\sqrt{\log n}}\right) > \Phi(x) - \epsilon + O\left(\frac{1}{\sqrt{\log n}}\right). \quad (3.19)$$

Yet  $\epsilon > 0$  was arbitrary, and so  $\mathbb{P}_n\left(\frac{\log \text{invol}(\sigma) - \mu_n}{\sigma_n} \leq x\right) \geq \Phi(x) + O(\frac{1}{\sqrt{\log n}})$ .  $\square$

## 4 A Functional Limit Theorem for $\log \text{invol}$

A stochastic process variant of Theorem 3.4 can also be formulated. Let  $C^{(n)} = (c_1^{(n)}, c_2^{(n)}, \dots)$  be distributed according to  $\text{ESF}(\theta)$  and  $W_n$  be the random element of the Skorokhod space  $\mathcal{D}[0, 1]$  of càdlàg functions on  $[0, 1]$  defined by

$$W_n(t) = \frac{\log \text{invol}_{\lfloor nt \rfloor}(C^{(n)}) - \frac{\theta t^2}{2} \log^2 n}{\sqrt{\frac{\theta}{3} \log^3 n}}. \quad (4.1)$$

**Theorem 4.1.** *It is possible to construct  $C^{(n)}$  and a standard Brownian motion  $W$  on the same probability space in such a way that*

$$\mathbb{E}\left\{\sup_{0 \leq t \leq 1} |W_n(t) - W(t^3)|\right\} = O\left(\frac{\log \log n}{\sqrt{\log n}}\right).$$

The Feller coupling, which expresses the Ewens Sampling Formula in terms of the spacings between successes in a sequence of independent Bernoulli trials, offers the ideal terrain for this task. Arratia, Barbour, and Tavaré [1] recently wrote a brief expository note about the Feller coupling to act as a companion to [12]. To summarize, consider a sequence  $\beta = (\beta_1, \beta_2, \dots)$  of independent Bernoulli random variables in which  $\Pr(\beta_j = 1) = \theta/(\theta + j - 1)$ ,  $j = 1, 2, \dots$ . If for all positive integers  $k$  and  $n$  we define

$$c_k^{(n)} = \#k - \text{spacings in } 1\beta_2\beta_3 \cdots \beta_n 1, \quad (4.2)$$

it can be shown that the distribution of  $C^{(n)} = (c_1^{(n)}, c_2^{(n)}, \dots)$  is exactly the same as  $\text{ESF}(\theta)$ . If we further define

$$Z_{k,m} = \#k - \text{spacings in } \beta_{m+1}\beta_{m+2} \cdots, \quad k, m \geq 1, \quad (4.3)$$

then  $Z_{k,m} \xrightarrow{\text{a.s.}} 0$  as  $m \rightarrow \infty$  due to the Borel-Cantelli lemma, and so  $Z := (Z_{1,0}, Z_{2,0}, \dots)$  adheres to the Poisson point process that  $C^{(n)}$  converges to.

We now assemble the pieces needed to prove Theorem 4.1. Our plan is to show that within the Feller coupling,  $\log \text{invol}_n(C^{(n)})$  is “close enough” to  $\log \text{invol}_n(Z)$ , which in turn is “close enough” to  $\log \mathbf{B}_n(Z)$ , which is known to conform to a Wiener process. First, we need the following lemma, which is a trivial consequence of the preceding definitions.

**Lemma 4.2.** *For all positive integers  $k$  and  $n$ ,*

$$Z_{k,0} - Z_{k,n} - \mathbb{1}\{L_n + R_n = k + 1\} \leq c_k^{(n)} \leq Z_{k,0} + \mathbb{1}\{L_n = k\},$$

where  $L_j = \min\{j \geq 1 : \beta_{n-j+1} = 1\}$  and  $R_j = \min\{j \geq 1 : \beta_{n+j} = 1\}$ .

**Lemma 4.3.**

$$\mathbb{E} \left\{ \sup_{0 \leq t \leq 1} \left| \log \text{invol}_{\lfloor n^t \rfloor}(C^{(n)}) - \log \mathbf{B}_{\lfloor n^t \rfloor}(Z) \right| \right\} = O(\log n).$$

*Proof.* If  $a$  and  $b$  are two vectors such that  $a_k \leq b_k$  for  $k = 1, 2, \dots$ , then

$$\text{invol}_n(b) = \text{invol}_n(a) \left( \prod_{k=1}^n k^{b_k - a_k} \right) \cdot \left( \prod_{k=1}^n \frac{\sum_{j=0}^{\lfloor b_k/2 \rfloor} \frac{b_k!}{(2k)^j j! (b_k - 2j)!}}{\sum_{j=0}^{\lfloor a_k/2 \rfloor} \frac{a_k!}{(2k)^j j! (a_k - 2j)!}} \right) \quad (4.4)$$

$$\leq \text{invol}_n(b) n^{\|b-a\|_1} \prod_{k=1}^n \exp\left(\frac{b_k^2}{2k}\right) \quad (4.5)$$

where  $\|b-a\|_1$  denotes the usual 1-norm. It follows from Theorem 4.2 that for all  $j \leq n$ ,

$$\log \text{invol}_{\lfloor j \rfloor}(Z) - (Y_n + 1) \log n - D_n \leq \log \text{invol}_{\lfloor j \rfloor}(C^{(n)}) \leq \log \text{invol}_{\lfloor j \rfloor}(Z) + \log n + D_n^* \quad (4.6)$$

where  $Y_n = \sum_{k=1}^n Z_{k,n}$ ,  $D_n = \sum_{k=1}^n \frac{(c_k^{(n)})^2}{2k}$ , and  $D_n^* = \sum_{k=1}^n \frac{(Z_{k,0}+1)^2}{2k} \geq D_n$ . Hence

$$|\log \text{invol}_{\lfloor n^t \rfloor}(C^{(n)}) - \log \text{invol}_{\lfloor n^t \rfloor}(Z)| \leq (Y_n + 1) \log n + D_n^* \quad (4.7)$$

for  $0 \leq t \leq 1$ . Barbour and Tavaré proved in [5] that  $\mathbb{E}Y_n \leq \theta^2$ . Additionally,

$$\mathbb{E}D_n^* = \sum_{k=1}^n \frac{1}{2k} \left( \frac{\theta^2}{k^2} + \frac{3\theta}{k} + 1 \right) = O(\log n). \quad (4.8)$$

Hence,

$$\mathbb{E} \left\{ \sup_{0 \leq t \leq 1} |\log \text{invol}_{\lfloor n^t \rfloor}(C^{(n)}) - \log \text{invol}_{\lfloor n^t \rfloor}(Z)| \right\} = O(\log n). \quad (4.9)$$

Due to (1.5) and (1.6),

$$\mathbb{E} \left\{ \sup_{0 \leq t \leq 1} |\log \text{invol}_{\lfloor n^t \rfloor}(Z) - \log \mathbf{B}_{\lfloor n^t \rfloor}(Z)| \right\} = O(1). \quad (4.10)$$

Triangle inequality implies the desired result.  $\square$

The remainder of the proof of Theorem 4.1 is immediate in light of the following.

**Theorem 4.4.** (Barbour, Tavaré [5]) *Suppose for all  $t \in [0, 1]$  we define*

$$B_n(t) = \frac{\log \mathbf{B}_{\lfloor n^t \rfloor}(Z) - \frac{\theta t^2}{2} \log^2 n}{\sqrt{\frac{\theta}{3} \log^3 n}}.$$

*Then  $B_n$  converges weakly in  $\mathcal{D}[0, 1]$  to a standard Brownian motion  $B$ , in the sense that*

$$\mathbb{E} \left\{ \sup_{0 \leq t \leq 1} |B_n(t) - B(t^3)| \right\} = O\left(\frac{\log \log n}{\sqrt{\log n}}\right).$$

Although we will decline from doing so, we remark that it is straightforward to compute explicit bounds for the constants implied by the error terms in Theorems 3.4 and 4.1. Most of the heavy lifting necessary to quantify how close the distributions of  $\log \text{invol}$  and  $\log \mathbf{B}$  are to each other was already done in the proof of Lemma 3.2. (You just need to approximate  $K_2$ .) The estimates needed to measure the distance between the distributions of  $\log \mathbf{B}_n(C^{(n)})$  and  $\log \mathbf{B}_n(Z)$  already appear in [5]. From there, a generic Berry-Esséen inequality will disclose how quickly  $\log \mathbf{B}_n(Z)$  tends to its Gaussian limit law. The constants implied by (4.8) and (4.9) require more delicate asymptotic evaluation, but can be safely ignored since the dominant error term for Theorem 4.1 resides in Theorem 4.4.

## 5 A Close Look at the Second Moment of invol

Ultimately, the asymptotic lognormality of  $\text{invol}$  for  $\theta$ -weighted permutations prevents, say, the method of moments from succeeding. (See, for instance, Example 30.2 on pg. 389 of Billingsley [7].) Ignoring that, we can still observe that the second moment of  $\text{invol}$  is quite obstreperous; its growth rate vastly outpaces the first moment, and the extent to which it does depends on whether  $\theta > 1$  or not.

Lemma 3.2 hints strongly at the presence of this disparity. One easy consequence of its proof is that

$$\mathbb{P}_n(c_k(\sigma) \leq 1 \text{ for every } k \geq \xi) = \mathbb{P}_n(\sigma \in \mathcal{P}_\xi)(1 + o(1)).$$

However, the Ewens Sampling Formula favors permutations with fewer cycles whenever  $\theta < 1$ . This inflates the likelihood of drawing a permutation with several instances of one very long cycle length, which is one of the events we wish to avoid in order to efficiently bound  $\text{invol}(\sigma)$ .

Meanwhile, if  $\theta > 1$ , then we are more likely to draw a permutation  $\sigma$  with a lot of short cycles. As was noted earlier, this is actually when  $\text{invol}(\sigma)$  is at its biggest. So  $\mathbb{E}_n(\text{invol})^2$  will grow even faster than it does in the  $\theta < 1$  case and falls under an entirely different asymptotic regime.

The ensuing analysis is included here not only to contextualize the sheer futility of the method of moments in this situation but also to help cast light on the statistical spread of  $\text{invol}$  for  $\theta$ -weighted permutations and how  $\theta = 1$  acts as a “phase transition point” for the variance. Reprising the notation presented in the proof of Theorem 2.1, we consider the generating function

$$G(z) = 1 + \sum_{n=1}^{\infty} \mathbb{E}_n(\text{invol})^2 \frac{\theta^{(n)}}{n!} z^n \quad (5.1)$$

$$= 1 + \sum_{n=1}^{\infty} \left( \sum_{\lambda \vdash n} \mathbb{P}_n(\sigma \in \mathcal{A}_\lambda) \mathbb{E}_n((\text{invol})^2 \mid \sigma \in \mathcal{A}_\lambda) \right) \frac{\theta^{(n)}}{n!} z^n \quad (5.2)$$

$$= 1 + \sum_{n=1}^{\infty} \left( \sum_{\lambda \vdash n} \left( \frac{n!}{\theta^{(n)}} \prod_{k=1}^n \left( \frac{\theta}{k} \right)^{\lambda_k} \frac{1}{\lambda_k!} \right) \left( \prod_{k=1}^n k^{2\lambda_k} \frac{(He_{\lambda_k}(i\sqrt{k}))^2}{(i\sqrt{k})^{2\lambda_k}} \right) \right) \frac{\theta^{(n)}}{n!} z^n \quad (5.3)$$

$$= \prod_{k=1}^{\infty} \left( \sum_{\lambda_k=0}^{\infty} \frac{(He_{\lambda_k}(i\sqrt{k}))^2}{\lambda_k!} \cdot (-\theta z^k)^{\lambda_k} \right) \quad (5.4)$$

We now recall the probabilists’ version of the Mehler kernel (see Kibble [22])

$$\sum_{n=0}^{\infty} He_n(x) He_n(y) \frac{t^n}{n!} = \frac{1}{\sqrt{1-t^2}} \exp \left( -\frac{t^2(x^2 + y^2) - 2txy}{2(1-t^2)} \right)$$

to observe that

$$G(z) = \prod_{k=1}^{\infty} \frac{1}{\sqrt{1 - \theta^2 z^{2k}}} \exp\left(\frac{\theta k z^k}{1 - \theta z^k}\right). \quad (5.5)$$

If  $\theta > 1$ , then  $G$  has a dominant singularity at  $z = 1/\theta$  and the function  $\phi$  given by

$$\phi(z) = \frac{1}{\sqrt{1 + \theta z}} \prod_{k=2}^{\infty} \frac{1}{\sqrt{1 - \theta^2 z^{2k}}} \exp\left(\frac{\theta k z^k}{1 - \theta z^k}\right) \quad (5.6)$$

is regular in the disc  $|z| < 1/\theta$ . Corollary 6.4 can thus be applied to get

$$[z^n]G(z) = \frac{\theta^{n-\frac{1}{4}} K_{\theta} \exp(2\sqrt{\theta n} + \theta/2)}{n^{\frac{1}{4}} 2\sqrt{2\pi}} \left(1 + O(n^{-\frac{1}{2}})\right) \quad (5.7)$$

where

$$K_{\theta} = \prod_{k=2}^{\infty} \frac{1}{\sqrt{1 - \theta^{2-2k}}} \exp\left(\frac{k}{\theta^{k-1} - 1}\right). \quad (5.8)$$

Hence

$$\mathbb{E}_n(\text{invol})^2 = \frac{n!}{\theta^{(n)}} [z^n]G(z) = \frac{K_{\theta} \Gamma(\theta)}{2\sqrt{2\pi}} n^{\frac{3}{4}-\theta} e^{2\sqrt{\theta n} + \frac{\theta}{2}} \theta^{n-\frac{1}{4}} \left(1 + O(n^{-\frac{1}{2}})\right), \quad (5.9)$$

from which it follows that  $\mathbb{E}_n(\text{invol})^2 = \omega(\mathbb{E}_n \text{invol})$ . However, if  $0 < \theta \leq 1$ , then  $G$  has a radius of convergence of 1, but behaves too chaotically around the unit circle to be analytically continued beyond it. In this case, we appeal to the exp-log reorganization of (5.5):

$$G(z) = \exp\left(\sum_{k=1}^{\infty} \left[-\frac{1}{2} \log(1 - \theta^2 z^{2k}) + \frac{\theta k z^k}{1 - \theta z^k}\right]\right) \quad (5.10)$$

$$= \exp\left(\sum_{k=1}^{\infty} \sum_{\ell=1}^{\infty} \left[\frac{\theta^{2\ell} z^{2k\ell}}{2\ell} + k\theta^{\ell} z^{k\ell}\right]\right) \quad (5.11)$$

$$= \exp\left(\sum_{\ell=1}^{\infty} \left[\frac{\theta^{2\ell}}{2\ell} \frac{z^{2\ell}}{(1 - z^{2\ell})} + \frac{\theta^{\ell} z^{\ell}}{(1 - z^{\ell})^2}\right]\right). \quad (5.12)$$

From this, we can attain the following upper bound on  $|\log G(z)|$  for when  $0 < |z| < 1$ :

$$|\log G(z)| \leq \frac{\theta|z|}{|1 - z|^2} + \sum_{\ell=1}^{\infty} \frac{\theta^{2\ell}}{2\ell} \frac{|z|^{2\ell}}{|1 - z^{2\ell}|} + \sum_{\ell=2}^{\infty} \frac{\theta^{\ell}|z|^{\ell}}{|1 - z^{\ell}|^2} \quad (5.13)$$

$$< \frac{\theta}{|1 - z|^2} + \sum_{\ell=1}^{\infty} \frac{\theta^{2\ell}}{(2\ell)^2} \frac{2\ell|z|^{2\ell}}{(1 - |z|^{2\ell})} + \sum_{\ell=2}^{\infty} \frac{\theta^{\ell}|z|^{\ell}}{(1 - |z|^{\ell})^2} \quad (5.14)$$

$$< \frac{\theta}{|1 - z|^2} + \frac{1}{1 - |z|} \sum_{\ell=1}^{\infty} \frac{\theta^{2\ell}}{(2\ell)^2} \frac{2\ell}{(|z|^{-1} + |z|^{-2} + \dots + |z|^{-2\ell})}$$

$$+ \frac{1}{(1-|z|)^2} \sum_{\ell=2}^{\infty} \frac{\theta^\ell}{(1+|z|+\dots+|z|^{\ell-1})(1+|z|^{-1}+\dots+|z|^{-(\ell-1)})} \quad (5.15)$$

$$< \frac{\theta}{|1-z|^2} + \frac{1}{1-|z|} \sum_{\ell=1}^{\infty} \frac{\theta^{2\ell}}{4\ell^2} + \frac{1}{(1-|z|)^2} \sum_{\ell=2}^{\infty} \frac{\theta^\ell}{\ell^2} \quad (5.16)$$

$$= \frac{\theta}{|1-z|^2} + \frac{\text{Li}_2(\theta^2)}{4(1-|z|)} + \frac{\text{Li}_2(\theta) - \theta}{(1-|z|)^2}. \quad (5.17)$$

(Apply the arithmetic mean-harmonic mean inequality on the terms of the second summation to transition from (5.15) to (5.16).)

Now in setting  $z = e^{-u}$ , the function

$$L(u) = \log G(e^{-u}) = \sum_{\ell=1}^{\infty} \left[ \frac{\theta^{2\ell}}{2\ell} \frac{e^{-2\ell u}}{(1-e^{-2\ell u})} + \frac{\theta^\ell e^{-\ell u}}{(1-e^{-\ell u})^2} \right]$$

is a harmonic sum well-suited for a Mellin transform analysis. Consult chapter 6 of [14] to deduce that the Mellin transform of  $L(u)$  is

$$L^*(s) = (2^{-s-1}\text{Li}_{s+1}(\theta^2)\zeta(s) + \text{Li}_s(\theta)\zeta(s-1))\Gamma(s), \quad \Re(s) > 1,$$

where  $\text{Li}_s(z) = \sum_{n \geq 1} \frac{z^n}{n^s}$  is the polylogarithm function of order  $s$ .

Note that  $\text{Li}_s(z)$ , when treated as a function of  $s$  and with  $z$  held constant, is analytic for  $z \in [0, 1)$ , because  $\text{Li}_s(z)$  converges absolutely for all  $s \in \mathbb{C}$  when  $|z| < 1$ . If  $z = 1$ , then the polylogarithm is simply the Riemann zeta function which has a simple pole at  $s = 1$ . We therefore consider the cases  $0 < \theta < 1$  and  $\theta = 1$  separately.

**Case 1:**  $\theta \in (0, 1)$ . The Mellin transform  $L^*(s)$  has simple poles at  $s = 2, 1, 0, -1, \dots$ . Using the standard transfer rule for the transform, we ascertain that

$$L(t) \underset{t \rightarrow 0^+}{=} \frac{\text{Li}_2(\theta)}{t^2} + \frac{\text{Li}_2(\theta^2)}{4t} + \frac{1}{4} \log(1-\theta^2) - \frac{\theta}{12(1-\theta)} + O(t) \quad (5.18)$$

which, because of the Laurent series expansions of the powers of  $t = -\log(z)$  at  $z = 1$  and the fact that (5.18) extends to  $\mathbb{C} - \mathbb{R}_{<0}$ , corresponds to

$$\log G(z) \underset{z \rightarrow 1}{=} \frac{\text{Li}_2(\theta)}{(1-z)^2} + \frac{\text{Li}_2(\theta^2) - 4\text{Li}_2(\theta)}{4(1-z)} + \frac{1}{12}\text{Li}_2(\theta) - \frac{1}{8}\text{Li}_2(\theta^2) + \frac{1}{4} \log(1-\theta^2) - \frac{\theta}{12(1-\theta)} + O(1-z) \quad (5.19)$$

from within the unit disc. We can now conclude that

$$G(z) \underset{z \rightarrow 1}{=} \tau(z)[1 + O(1-z)], \quad (5.20)$$

provided  $|z| < 1$  and

$$\tau(z) = (1-\theta^2)^{1/4} \exp\left(\frac{\text{Li}_2(\theta)}{(1-z)^2} + \frac{\text{Li}_2(\theta^2) - 4\text{Li}_2(\theta)}{4(1-z)} + \frac{1}{12}\text{Li}_2(\theta) - \frac{1}{8}\text{Li}_2(\theta^2) - \frac{\theta}{12(1-\theta)}\right) \quad (5.21)$$

We claim next that  $[z^n]G(z) = [z^n]\tau(z)(1 + o(1))$ . To see this, consider  $\gamma$ , the positively oriented circle  $|z| = 1 - (2\text{Li}_2(\theta)/n)^{1/3}$ , and split  $\gamma$  into the two contours

$$\gamma_1 = \left\{ z \in \gamma : |1 - z| < \left( \frac{3\text{Li}_2(\theta)}{n} \right)^{\frac{1}{3}} \right\}$$

and  $\gamma_2 = \gamma - \gamma_1$ . By Cauchy's integral formula,

$$[z^n](G(z) - \tau(z)) = \frac{1}{2\pi i} \int_{\gamma_1} \frac{G(z) - \tau(z)}{z^{n+1}} dz + \frac{1}{2\pi i} \int_{\gamma_2} \frac{G(z) - \tau(z)}{z^{n+1}} dz \quad (5.22)$$

The length of  $\gamma_1$  is  $O(n^{-1/3})$ . This together with (5.20) shows that the first integral is

$$= O\left( \int_{\gamma_1} |z|^{-n} |1 - z| \exp\left( \frac{\text{Li}_2(\theta)}{(1 - |z|)^2} + \frac{\text{Li}_2(\theta^2) - 4\text{Li}_2(\theta)}{4(1 - |z|)} \right) |dz| \right) \quad (5.23)$$

$$= O\left( \exp\left( \frac{2\text{Li}_2(\theta)n^{\frac{2}{3}}}{(2\text{Li}_2(\theta))^{\frac{2}{3}}} \right) \cdot n^{-\frac{1}{3}} \cdot \exp\left( \frac{\text{Li}_2(\theta)n^{\frac{2}{3}}}{(2\text{Li}_2(\theta))^{\frac{2}{3}}} + \frac{(\text{Li}_2(\theta^2) - 4\text{Li}_2(\theta))n^{\frac{1}{3}}}{4(2\text{Li}_2(\theta))^{\frac{2}{3}}} \right) \right) \cdot O(n^{-\frac{1}{3}}) \quad (5.24)$$

$$= O\left( n^{-\frac{2}{3}} \exp\left( \frac{3\text{Li}_2(\theta)n^{\frac{2}{3}}}{(2\text{Li}_2(\theta))^{\frac{2}{3}}} + \frac{(\text{Li}_2(\theta^2) - 4\text{Li}_2(\theta))n^{\frac{1}{3}}}{4(2\text{Li}_2(\theta))^{\frac{1}{3}}} \right) \right) \quad (5.25)$$

On the other hand, estimate (5.17) applies over  $\gamma_2$ , and so the second integral in (5.22) is

$$= O\left( \int_{\gamma_2} |z|^{-n} \left[ \exp\left( \frac{\theta}{|1 - z|^2} + \frac{\text{Li}_2(\theta^2)}{4(1 - |z|)} + \frac{\text{Li}_2(\theta) - \theta}{(1 - |z|)^2} \right) + \exp\left( \frac{\text{Li}_2(\theta)}{|1 - z|^2} + \frac{\text{Li}_2(\theta^2) - 4\text{Li}_2(\theta)}{4|1 - z|} \right) \right] |dz| \right) \quad (5.26)$$

$$= O\left( \exp\left( \frac{2\text{Li}_2(\theta)n^{\frac{2}{3}}}{(2\text{Li}_2(\theta))^{\frac{2}{3}}} \right) \cdot \exp\left( \frac{\theta n^{\frac{2}{3}}}{(3\text{Li}_2(\theta))^{\frac{2}{3}}} + \frac{\text{Li}_2(\theta^2)n^{\frac{1}{3}}}{4(2\text{Li}_2(\theta))^{\frac{1}{3}}} + \frac{(\text{Li}_2(\theta) - \theta)n^{\frac{2}{3}}}{(2\text{Li}_2(\theta))^{\frac{2}{3}}} \right) \right. \\ \left. + \exp\left( \frac{2\text{Li}_2(\theta)n^{\frac{2}{3}}}{(2\text{Li}_2(\theta))^{\frac{2}{3}}} \right) \cdot \exp\left( \frac{\text{Li}_2(\theta)n^{2/3}}{(3\text{Li}_2(\theta))^{\frac{2}{3}}} + \frac{(\text{Li}_2(\theta^2) - 4\text{Li}_2(\theta))n^{\frac{1}{3}}}{4(3\text{Li}_2(\theta))^{\frac{1}{3}}} \right) \right) \quad (5.27)$$

$$= O\left( \exp\left( An^{\frac{2}{3}} \right) \right), \quad (5.28)$$

for some  $A < 3\text{Li}_2(\theta)/(2\text{Li}_2(\theta))^{2/3}$ . Therefore

$$[z^n]G(z) = [z^n]\tau(z) + O\left( n^{-\frac{2}{3}} \exp\left( \frac{3\text{Li}_2(\theta)n^{\frac{2}{3}}}{(2\text{Li}_2(\theta))^{\frac{2}{3}}} + \frac{(\text{Li}_2(\theta^2) - 4\text{Li}_2(\theta))n^{\frac{1}{3}}}{4(2\text{Li}_2(\theta))^{\frac{1}{3}}} \right) \right) \quad (5.29)$$

It is now a sophomore's homework exercise to use Theorem 6.3 on  $\tau(z)$ , especially considering that the required singular expansion is already accounted for in (5.18). The error term



in (5.29) is smaller than the leading term asymptotic of  $[z^n]\tau(z)$  by a decaying factor of  $O(\exp(-\text{Li}_2(\theta)n^{1/3}/(2\text{Li}_2(\theta))^{1/3}))$ . In the end, we find that

$$[z^n]G(z) \sim \frac{(1-\theta^2)^{\frac{1}{4}}(2\text{Li}_2(\theta))^{\frac{1}{6}}}{\sqrt{6\pi n^{\frac{2}{3}}}} \exp\left(\frac{3\text{Li}_2(\theta)n^{\frac{2}{3}}}{(2\text{Li}_2(\theta))^{\frac{2}{3}}} + \frac{\text{Li}_2(\theta^2)n^{\frac{1}{3}}}{(2\text{Li}_2(\theta))^{\frac{1}{3}}} - \frac{\theta}{12(1-\theta)}\right), \quad (5.30)$$

and so

$$\mathbb{E}_n(\text{invol})^2 \sim \frac{(1-\theta^2)^{\frac{1}{4}}(2\text{Li}_2(\theta))^{\frac{1}{6}}\Gamma(\theta)n^{\frac{1}{3}-\theta}}{\sqrt{6\pi}} \exp\left(\frac{3\text{Li}_2(\theta)n^{\frac{2}{3}}}{(2\text{Li}_2(\theta))^{\frac{2}{3}}} + \frac{\text{Li}_2(\theta^2)n^{\frac{1}{3}}}{(2\text{Li}_2(\theta))^{\frac{1}{3}}} - \frac{\theta}{12(1-\theta)}\right). \quad (5.31)$$

**Case 2:**  $\theta = 1$ . Then  $G(z)$  is precisely the generating function of  $\mathbb{E}_n(\text{invol})^2$ , and the Mellin transform becomes

$$L^*(s) = (2^{-s-1}\zeta(s+1) + \zeta(s-1))\zeta(s)\Gamma(s), \quad (5.32)$$

which has simple poles at  $s = 2, 1, 0, -1, \dots$ , and a double pole at  $s = 0$ . Hence

$$L(t) \underset{t \rightarrow 0^+}{=} \frac{\pi^2}{6t^2} + \frac{\pi^2 - 12}{24t} + \frac{1}{4} \log t - \log \sqrt[4]{2\pi} + \frac{1}{24} + O(t), \quad (5.33)$$

which corresponds to

$$\log G(z) \underset{z \rightarrow 1}{=} \frac{\pi^2}{6(1-z)^2} - \frac{\pi^2 - 4}{8(1-z)} + \frac{1}{4} \log(1-z) + \frac{7}{24} - \frac{\pi^2}{144} - \log \sqrt[4]{2\pi} + O(1-z) \quad (5.34)$$

from within the unit disc. Therefore

$$G(z) = \tau(z)[1 + O(1-z)], \quad (5.35)$$

provided  $|z| < 1$  and

$$\tau(z) = (2\pi(1-z))^{\frac{1}{4}} \exp\left(\frac{\pi^2}{6(1-z)^2} - \frac{\pi^2 - 4}{8(1-z)} + \frac{7}{24} - \frac{\pi^2}{144}\right). \quad (5.36)$$

At this point, the outstanding justifications are practically the same as they were for Case 1. It is tedious, but not too difficult to figure out that

$$[z^n]G(z) \sim \frac{1}{\sqrt[4]{2} \sqrt[12]{\pi} (3n)^{\frac{7}{12}}} \exp\left(\frac{1}{2}(3\pi n)^{\frac{2}{3}} - \frac{\pi^2 - 4}{8} \left(\frac{3n}{\pi^2}\right)^{\frac{1}{3}} + \frac{7}{24} - \frac{\pi^2}{144}\right). \quad (5.37)$$

## Concluding Remarks

The nonuniformity of `invol` makes it evident that the composition of two uniformly random involutions is not a uniformly random permutation. Furthermore, if we let  $t_n$  denote the total number of involutions in  $\mathfrak{S}_n$ , then

$$t_n^2 \sim n! \cdot \frac{e^{2\sqrt{n}}}{\sqrt{8\pi en}} \sim |\mathfrak{S}_n| \cdot \mathbb{E}_{1,n} \text{invol}. \quad (5.38)$$

(Refer to pg. 583, Example VIII.9, [19].) Therefore, using pairs of random involutions to generate random permutations introduces a significant sampling bias.

Like  $\text{ESF}(\theta)$  for  $\theta > 1$ , a composition of two random involutions is likelier to have a lot of cycles. However, the distribution of  $c_k$  looks quite different in this model. The factors of the exponential generating function in (2.6) suggests that  $c_k$  converges in law to the distribution of  $X + 2Y_k$ , where  $X$  and  $Y_k$  are independent Poisson random variables of mean 1 and  $1/(2k)$ , respectively. Lugo proved as much in an unpublished manuscript [26]. The author intends to expand on these and other ideas in a followup paper.

## References

- [1] Arratia, Richard and Barbour, A. D. and Tavaré, Simon, Exploiting the Feller coupling for the Ewens sampling formula, *Statist. Sci.*, **31**, (2016), no. 1, 27–29.
- [2] Arratia, R. and Barbour, A. D. and Tavaré, S., Limits of logarithmic combinatorial structures, *Ann. Probab.*, **28**, (2000), no. 4, 1620–1644.
- [3] Arratia, Richard and Barbour, A. D. and Tavaré, Simon, Logarithmic combinatorial structures: a probabilistic approach, *EMS Monographs in Mathematics*, (2003) ISBN 3-03719-000-0.
- [4] Arratia, Richard and Tavaré, Simon, Limit theorems for combinatorial structures via discrete process approximations, *Random Structures Algorithms*, **3**, (1992), no. 3, 321–345.
- [5] Barbour, A. D. and Tavaré, Simon, A rate for the Erdős-Turán law, *Combin. Probab. Comput.*, **3**, (1994), no. 2, 167–176.
- [6] Bhatia, Sangeeta and Egri-Nagy, Attila and Francis, Andrew R., Algebraic double cut and join: a group-theoretic approach to the operator on multichromosomal genomes, *J. Math. Biol.*, **71**, (2015), no. 5, 1149–1178.
- [7] Billingsley, Patrick, Probability and measure, *Wiley Series in Probability and Mathematical Statistics*, John Wiley & Sons, Inc., New York, (1995) ISBN 0-471-00710-2.

- [8] Bovey, J.D. An approximate probability distribution for the order of elements of the symmetric group, *Bull. London Math. Soc.*, **12**, (1980), no. 1, 41–46.
- [9] Burnette, Charles, Factoring permutations into the product of two involutions: a probabilistic, combinatorial, and analytic approach, Thesis (Ph.D.)– Drexel University, 2017.
- [10] Burnette, Charles and Schmutz, Eric, Representing random permutations as the product of two involutions, *Online J. Anal. Comb.*, **11**, (2016), Paper No. 6.
- [11] Chindelevitch, Leonid and Meidanis, João, On the rank-distance median of 3 permutations, *BMC Bioinformatics*, **19**, (2018), no. 8, 256–176.
- [12] Crane, Harry, The ubiquitous Ewens sampling formula, *Statist. Sci.*, **31**, (2016), no. 1, 1–19.
- [13] DeLaurentis, J. M. and Pittel, B. G., Random permutations and Brownian motion, *Pacific J. Math.*, **119**, (1985), no. 2, 287–301,
- [14] Erdélyi, A. and Magnus, W. and Oberhettinger, F. and Tricomi, F. G., Tables of integral transforms. Vol. I, McGraw-Hill Book Company, Inc., New York-Toronto-London, (1954), ISBN 0-070-19549-3.
- [15] Erdős, P. and Turán, P., On some problems of a statistical group-theory. I, *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, **4**, (1965), 175–186.
- [16] Erdős, P. and Turán, P., On some problems of a statistical group-theory. III, *Acta Math. Acad. Sci. Hungar.*, **18**, (1967), 309–320.
- [17] Flajolet, Philippe and Gourdon, Xavier and Dumas, Philippe, Mellin transforms and asymptotics: harmonic sums, *Theoret. Comput. Sci.*, **144**, (1995), nos. 1–2, 3–58.
- [18] Flajolet, Philippe and Odlyzko, Andrew, Singularity analysis of generating functions, *SIAM J. Discrete Math.*, **3**, (1990), no. 2, 216–240.
- [19] Flajolet, Philippe and Sedgewick, Robert, Analytic combinatorics, Cambridge University Press, Cambridge, (2009), ISBN 978-0-521-89806-5.
- [20] Hayman, W. K., A generalisation of Stirling’s formula, *J. Reine Angew. Math.*, **196**, (1956), 67–95.
- [21] Kawahara, Jun and Saitoh, Toshiki and Yoshinaka, Ryo, The time complexity of permutation routing via matching, token swapping and a variant, *J. Graph Algorithms Appl.*, **23**, (2019), 29–70.
- [22] Kibble, W. F., An extension of a theorem of Mehler’s on Hermite polynomials, *Proc. Cambridge Philos. Soc.*, **41**, (1941), 12–15.

- [23] Kim, D. S. and Kim, T. and Kim, Y.-H. and Dolgy, D. V., A note on Eulerian polynomials associated with Bernoulli and Euler numbers and polynomials, *Adv. Stud. Contemp. Math. (Kyungshang)*, **22**, (2012), 379–389.
- [24] Kim, Taekyun, Identities involving Frobenius-Euler polynomials arising from non-linear differential equations, *J. Number Theory*, **132**, (2012), 2854–2865.
- [25] Kim, T. and Mansour, T., Umbral calculus associated with Frobenius-type Eulerian polynomials, *Russ. J. Math. Phys.*, **21**, (2014), 484–493.
- [26] Lugo, Michael T., The cycle structure of compositions of random involutions, (2009), arXiv:0911.3604 [math.CO].
- [27] Lugo, Michael, Profiles of permutations, *Electron. J. Combin.*, **16**, (2009), no. 1, Research Paper 99, 20.
- [28] Manstavičius, E., The Berry-Esseen bound in the theory of random permutations, *Ramanujan J.*, **2**, (1998), nos. 1-2, 185–199.
- [29] Nicolas, J.-L., Distribution statistique de l'ordre d'un élément du groupe symétrique, *Acta Math. Hungar.*, **45**, (1985), nos. 1-2, 69–84.
- [30] Petersen, T. Kyle and Tenner, Bridget Eileen, How to write a permutation as a product of involutions (and why you might care), *Integers*, **13**, (2013), Paper No. A63, 20.
- [31] Roberts, John A. G. and Vivaldi, Franco, A combinatorial model for reversible rational maps over finite fields, *Nonlinearity*, **22**, (2009), no. 8, 1965–1982.
- [32] Wilf, Herbert S., The asymptotic behavior of the Stirling numbers of the first kind, *J. Combin. Theory Ser. A*, **64**, (1993), no. 2, 344–349.
- [33] Wright, E. Maitland, The Coefficients of a Certain Power Series, *J. London Math. Soc.*, **S1-7**, (1932), no. 4, 256–262.
- [34] Wright, E. M., On the coefficients of power series having exponential singularities. II, *J. London Math. Soc.*, **24**, (1949), 304–309.
- [35] Yang, Qingxuan and Ellis, John and Mamakani, Khalegh and Ruskey, Frank, In-place permuting and perfect shuffling using involutions, *Inform. Process. Lett.*, **113**, (2013), 386–391.

## 6 Appendix

This appendix compiles some technical prerequisites which are used sporadically. The first two subsections center around special cases of the saddle-point method.

## 6.1 Saddle-Point Bounds

The following proposition (pg. 549, Corollary VIII.1, [19]) can be used to bound the Hermite polynomial factors of  $\text{invol}(\sigma)$ .

**Theorem 6.1.** *Let  $f(z)$ , not a polynomial, be analytic at 0 with non-negative coefficients and radius of convergence  $R \leq +\infty$ . Assume that  $\lim_{x \rightarrow R^-} f(x) = +\infty$ . Then*

$$[z^n]f(z) \leq \frac{f(r_n)}{r_n^n}$$

where  $r_n \in (0, R)$  is the unique solution of the equation  $xf'(x)/f(x) = n + 1$ .

Using the exponential generating function for the probabilists' Hermite polynomials yields

$$\frac{He_{c_k}(i\sqrt{k})}{(i\sqrt{k})^{c_k}} = c_k! [z^{c_k}] \exp\left(z + \frac{z^2}{2k}\right) \leq \frac{c_k! \exp\left(r_{c_k} + \frac{r_{c_k}^2}{2k}\right)}{r_{c_k}^{c_k}}, \quad (6.1)$$

where  $r_{c_k} = \frac{1}{2}(\sqrt{4kc_k + k^2} + k)$ . A tidier bound can be acquired by noticing that

$$\sqrt{kc_k} \leq r \leq \sqrt{k(c_k + 1)}. \quad (6.2)$$

Of course, a full appeal to the saddle-point method produces a more accurate estimate.

**Theorem 6.2.** (Hayman [20]) *If  $P(z)$  is a polynomial of degree  $m$ , then as  $n \rightarrow \infty$ ,*

$$[z^n]e^{P(z)} \sim \frac{e^{P(r_n)}}{\sqrt{2\pi m n r_n^n}},$$

where  $r_n$  is the unique positive root of the equation  $xP'(x) = n$ .

In this case, the saddle point shifts to  $\frac{1}{2}(\sqrt{4kc_k + k^2} + k)$ . So for sufficiently large  $c_k$ , we observe that

$$[z^{c_k}] \exp\left(z + \frac{z^2}{2k}\right) \leq \frac{\exp\left(\frac{c_k}{2} + \sqrt{kc_k}\right)}{2\sqrt{\pi c_k} (kc_k)^{c_k/2}} (1 + o(1)). \quad (6.3)$$

## 6.2 Wright's Expansions

The following consequence of the saddle-point method applied to functions with exponential singularities is often helpful in unraveling the asymptotic development of various random permutation statistics.

**Theorem 6.3.** (Wright [34]) *Suppose*

$$f(z) = (1 - z)^{-\beta} \phi(z) e^{P(z)},$$

where  $\phi$  is regular in the unit disc,  $\phi(1) \neq 0$ ,

$$P(z) = \sum_{m=1}^M \frac{\alpha_m}{(1-z)^{\rho_m}},$$

$\rho_1, \dots, \rho_M$  are real,  $\beta, \alpha_1, \dots, \alpha_M$  are complex,  $\rho_1 > 0$ , and  $\alpha_1 \neq 0$ . If in addition

$$P(e^{-u}) = Au^{-\rho} \left( 1 + \sum_{\ell=1}^L A_\ell u^{\sigma_\ell} \right) + O(u^K), \quad u \rightarrow 0^+,$$

where  $A \in \mathbb{C} - \mathbb{R}_{\leq 0}$ ,  $0 < \sigma_1 < \sigma_2 < \dots < \sigma_L \leq \rho$ , and  $L, A, \rho$ , the  $\sigma_\ell$ , and the  $A_\ell$  are all calculable in terms of the  $\rho_m$  and  $\alpha_m$ , then

$$[z^n]f(z) = \frac{N^{\beta-\frac{1}{2}} e^{\Omega(N)} \phi(1)}{\sqrt{2\pi n(\rho+1)}} (1 + O(n^{-K})), \quad n \rightarrow \infty,$$

where

$$N = \left( \frac{n}{A\rho} \right)^{\frac{1}{\rho+1}},$$

$$\Omega(N) = A(1+\rho)N^\rho + A \sum_{\ell=1}^L A_\ell N^{\rho-\sigma_\ell} + \sum_{s=2}^{\infty} \frac{(-1)^{s-1} (A\rho)^{1-\frac{\rho}{\rho+1}}}{s!(\rho+1)} \left[ \frac{d^{s-2}}{dy^{s-2}} (y^{\frac{\rho}{\rho+1}-2} (\vartheta(y))^s) \right] \Big|_{y=N}, \text{ and}$$

$$\vartheta(y) = \frac{1}{\rho} \sum_{\ell=1}^L (\rho - \sigma_\ell) A_\ell \left( \frac{y}{A\rho} \right)^{-\frac{\sigma_\ell}{\rho+1}}.$$

With one exception in section 5, we solely need this simpler instance of Theorem 6.3.

**Corollary 6.4.** (Wright [33]) *The leading-term asymptotic for*

$$s_n = [z^n](1-z)^\beta \phi(z) \exp\left(\frac{\alpha}{1-z}\right),$$

where  $\beta$  is a complex number,  $\phi$  is regular in the unit disc,  $\phi(1) \neq 0$ , and  $\alpha$  is a nonzero real number, is given by

$$s_n = \frac{1}{n^{\frac{\beta}{2}+\frac{3}{4}}} \left[ \frac{\exp(2\sqrt{\alpha n})}{2\sqrt{\pi}} \phi(1) e^{\frac{\alpha}{2}} \alpha^{\frac{\beta}{2}+\frac{1}{4}} \right] \left( 1 + O(n^{-\frac{1}{2}}) \right).$$

### 6.3 The Mellin Transform

To conclude this section, we review some pertinent facts about the Mellin transform as a means of extracting asymptotic estimates. Given a locally integrable function  $f$  defined over  $\mathbb{R}_{>0}$ , the Mellin transform of  $f$  is the complex-variable function

$$f^*(s) := \int_0^\infty f(x) x^{s-1} dx.$$

The transform is also occasionally denoted by  $\mathcal{M}[f]$  or  $\mathcal{M}[f(x); s]$ . If  $f(x) = O(x^{-a})$  as  $x \rightarrow 0^+$  and  $f(x) = O(x^{-b})$  as  $x \rightarrow +\infty$ , then  $f^*$  is an analytic function on the “fundamental strip”  $a < \Re(z) < b$  and is often continuable to a meromorphic function on the whole of  $\mathbb{C}$ .

Notably, the Mellin transform is a linear operator that satisfies the rescaling rule

$$\mathcal{M}[f(\mu x); s] = \mu^{-s} f^*(s),$$

for any  $\mu > 0$ . This together with the linearity of  $\mathcal{M}$  yields the “harmonic sum rule”

$$\mathcal{M}\left[\sum_k \lambda_k f(\mu_k x); s\right] = \left(\sum_k \lambda_k \mu_k^{-s}\right) f^*(s),$$

provided that the integral and summation are interchangeable. Sums of the form  $\sum_k \lambda_k f(\mu_k x)$  are known as harmonic sums with base function  $f$ , amplitudes  $\lambda_k$ , and frequencies  $\mu_k$ . So in other words, the Mellin transform of a harmonic sum factorizes as the product of the Mellin transform of the base function and a generalized Dirichlet series associated to the corresponding sequence of amplitudes and frequencies.

Akin to many other valuable integral transforms, the Mellin transform is essentially involutive. If  $f$  is continuous in an interval containing  $x$ , then  $f(x)$  can be recovered from  $f^*(s)$  via the inversion formula

$$f(x) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} f^*(s) x^{-s} ds,$$

where the abscissa  $c$  must be chosen from the fundamental strip of  $f$ . If the continuation of  $f^*(s)$  decays quickly enough as  $\Im(c) \rightarrow +\infty$ , then the inverse Mellin integral and Cauchy’s residue theorem can be combined to derive an asymptotic series for  $f(x)$  as  $x \rightarrow 0^+$ . We have the following transfer rule in particular: a pole of order  $k + 1$  at  $s = s_0$  with

$$f^*(s) \underset{s \rightarrow s_0}{\sim} a_{s_0} \frac{(-1)^k k!}{(s - s_0)^{k+1}}$$

translates to the term  $a_{s_0} x^{-s_0} (\log x)^k$  in the singular expansion of  $f(x)$  near  $x = 0$ . (The specific requirements for this transfer rule are all fulfilled by the special functions appearing in this manuscript, as the gamma function decays rapidly along vertical lines in the complex plane whereas polylogarithms undergo only moderate growth.) Detailed explanations along with further uses of the Mellin transform towards solving asymptotic enumeration problems can be found in [17] and [19].