

Maximally Chaotic Dynamical System
of
Infinite Dimensionality

George Savvidy

Institute of Nuclear and Particle Physics
Demokritos National Research Center
Ag. Paraskevi, Athens, Greece

Abstract

We analyse the infinite-dimensional limit of the maximally chaotic dynamical systems that are defined on N -dimensional tori. These hyperbolic systems found successful application in computer algorithms that generate high-quality pseudorandom numbers for advanced Monte Carlo simulations. The chaotic properties of these systems are increasing with N because the corresponding Kolmogorov-Sinai entropy grows linearly with N . We calculated the spectrum and the entropy of the system that appears in the infinite dimensional limit. We demonstrated that the limiting system has exponentially expanding and contracting foliations and therefore belongs to the Anosov C -systems of infinite dimensionality. The limiting system defines the hyperbolic evolution of the continuous functions very similar to the evolution of a velocity function describing the hydrodynamic flow of fluids. We compare the chaotic properties of the limiting system with those of the hydrodynamic flow of incompressible ideal fluid on a torus investigated by Arnold. This maximally chaotic system can find application in Monte Carlo method, statistical physics and digital signal processing.

PACS: 45.30.+s; 05.45.Jn; 89.70.Cf; 05.45.Pq

1 *Introduction*

The maximally chaotic dynamical systems continue to attract great attention of the researchers due to their rich physical properties and extended areas of application. In recent years the classical and quantum-mechanical concepts of maximally chaotic systems were developed in series of publications with application to the physics of black holes [1, 2, 3, 4, 5, 6], of non-Abelian gauge fields [7, 8] and string theory [9], to the fluid dynamics [13, 14, 15] and astrophysics [16, 17, 18, 19], foundation of statistical physics [20, 21, 22, 23, 24] and computer science [25, 26, 27]. There is a growing evidence that intrinsic properties of the Hawking black hole radiation can be understood in terms of classical and quantum theory of chaos [28, 29, 30, 31, 32, 33, 34]. It was conjectured that the resolution of the black hole information paradox of black holes radiation, which behaves as a black body radiation with finite temperature and is similar to the thermodynamic system characterised by entropy and other thermodynamic quantities, can be formulated in terms of maximally chaotic dynamical systems [1, 2, 3, 4, 5, 6, 8]. We are interested to analyse deterministic dynamical systems, which show up physical properties intrinsically related to the thermodynamical behaviour [20, 21, 22, 23, 24, 35, 36, 37].

There is an intuitive understanding about how strong chaotic behaviour of a classical system can be, but it seems natural to define a maximally chaotic system as a system that has a nonzero Kolmogorov-Sinai entropy and therefore belongs to the so called K-systems [38, 39, 40, 41, 8]. The examples of maximally chaotic systems were investigated and constructed in the earlier investigations of Artin, Hadamard, Hedlund, Hopf, Birkhoff, von Neumann and many others researchers working in ergodic theory [42, 43, 44, 45, 46, 47]. A large class of maximally chaotic dynamical systems was constructed by Anosov [41, 48, 49, 51, 52, 53, 54]. These are the systems that fulfil the C-condition that leads to the exponential instability of the trajectories [55], to the mixing of all orders and to a positive Kolmogorov-Sinai entropy. As a result it is natural to call them C-K systems [8].

The geodesic flows on closed Riemannian manifolds of variable negative sectional curvatures fulfil the C-condition and therefore represent a rich class of maximally chaotic dynamical systems [41, 35]. A progress in understanding of the chaotic behaviour of the non-Abelian gauge fields, of the N-body systems in Newtonian gravity and of some cosmological models in general relativity were achieved by the application of the methods and results of the ergodic theory and the theory of geodesic C-K flows [7, 8, 56, 57, 58, 59, 60, 61, 16, 17, 18, 19].




Figure 1: The left figure shows the distribution of the eigenvalues of the operators $A(N)$, where $N = 256$. On the right figure is the distribution of the eigenvalues of the infinite-dimensional operator A in λ plane. The unit circles are shown in both figures.

A class of hyperbolic C-K systems defined on a torus found application in computer algorithms that generate high-quality pseudorandom numbers for the advanced Monte Carlo simulations [25, 26, 27]. The chaotic properties of C-K systems on a torus are increasing with the dimensionality N of the systems because the entropy increases linearly with N as $h(A) = \frac{2}{\pi} N$. It seems important to investigate the infinite-dimensional limit of these systems that found successful application in Monte Carlo method [62, 63, 64, 65, 67] and to understand the exceptional properties of the limiting system.

2 Chaotic Dynamical Systems on N -dimensional Tori

The C-K system that realises automorphisms of a torus with coordinates (u_1, \dots, u_N) is defined as an integer matrix transformation [41, 25, 26, 27, 40, 46, 47, 54]:

$$u_i^{(k+1)} = \sum_{j=1}^N A_{ij}(N) u_j^{(k)} \pmod{1}, \quad k = 0, 1, 2, \dots \quad (2.1)$$

where the components of the vector $u^{(k)}$ are $u^{(k)} = (u_1^{(k)}, \dots, u_N^{(k)})$. The phase space of the system is the N -dimensional torus \mathcal{T}^N appearing in factorisation of the Euclidean space E^N with coordinates $u = (u_1, \dots, u_N)$ over an integer lattice \mathcal{Z}^N endowed with the invariant Liouville's measure $d\mu = du_1 \dots du_N$ [41]. The automorphisms (2.1) fulfils the C-condition if and only if the integer matrix $A(N)$ has no eigenvalues on the unit circle and has the determinant equal to one, that is, the eigenvalues $\{\Lambda = \lambda_1, \dots, \lambda_N\}$ fulfil the following conditions:

$$1) \text{Det}A(N) = \lambda_1 \lambda_2 \dots \lambda_N = 1, \quad 2) |\lambda_i| \neq 1, \quad \forall i. \quad (2.2)$$

The eigenvalues are divided into sets with modulus smaller and larger than one:

$$0 < |\lambda_\alpha| < 1 \text{ for } \alpha = 1 \dots d, \quad 1 < |\lambda_\beta| < \infty \text{ for } \beta = d+1 \dots N. \quad (2.3)$$

The C-K system (2.1) has nonzero entropy and its value $h(A)$ can be calculated in terms of its eigenvalues [41, 25]:

$$h(A) = \sum_{\beta} \ln |\lambda_\beta| = - \sum_{\alpha} \ln |\lambda_\alpha|. \quad (2.4)$$

The eigenvalues λ_β larger than one are nothing else but the Lyapunov exponents and characterise the chaotic properties of the system as it follows from the above definition of entropy (2.4).

We are interested to consider the infinite-dimensional limit $N \rightarrow \infty$ of the system (2.1) when the operator $A(N)$ is given by the $N \times N$ matrix with all integer entries $A_{ij} \in \mathbb{Z}$ and has the following form [25, 26, 27]:

$$A(N) = \begin{pmatrix} 1 & 1 & 1 & 1 & \dots & 1 & 1 \\ 1 & 2 & 1 & 1 & \dots & 1 & 1 \\ & & & \dots & & & \\ 1 & N & N-1 & N-2 & \dots & 3 & 2 \end{pmatrix}. \quad (2.5)$$

It has the determinant equal to one and the spectrum of inverse matrix has the form [27]

$$\lambda_j = 1 - 2 \exp(i \pi j/N) + \exp(2i \pi j/N), \quad j = -N, -N+2, \dots, N-2, N \quad (2.6)$$

shown in Fig.1. The kernel $A(N)u_0 = 0$ of the operator $A(N)$ consists of only one vector with zero components $u_0 = (0, \dots, 0)$. The eigenvalues fulfil the C-condition (2.2) and the entropy of the system can be calculated for the large values of N as a sum over eigenvalues:

$$h(A) = \sum_{\beta} \ln |\lambda_\beta| = \sum_{-2\pi/3 < \phi_j < 2\pi/3} \ln(4 \cos^2(\phi_j/2)) \sim \frac{2}{\pi} N, \quad (2.7)$$

where the eigenvalues are given in (2.6) and $\phi_j = \pi \frac{j}{N}$. The entropy increases linearly with the dimension N of the operator $A(N)$. We note that the special form of the matrix $A(N)$ in (2.5) has highly desirable property of having a widely spread, nearly continuum spectrum of eigenvalues (2.6) shown in Fig.1 and indicating that the exponential mixing takes place in many scales [25].

3 Infinite-dimensional Limit of $A(N)$ System

We are interested to define and investigate the system that appears in the infinite-dimensional limit of $A(N)$ when $N \rightarrow \infty$. The size of state vector $u = (u_1, \dots, u_N)$ tends to infinity and it

seems natural to expect that it can be represented by a continuous function ψ defined in an infinite-dimensional space $\mathcal{H} = \{\psi(u)\}$, possibly a Hilbert space, and the operator $A(N)$ will reduce to a differential operator A acting in \mathcal{H} . The existence of such a limit would mean that A defines a "measure preserving" transformation of continuous functions $A^n\psi = \psi^{(n)}$ in \mathcal{H} that will have maximally strong chaotic properties. Our intension is to define this limiting system, to explore its properties and possible applications in Monte Carlo method. It seems that the investigation of infinite-dimensional "fully chaotic" transformation of continuous functions may also help to understand better a chaotic/turbulent motion of fluids. As it was demonstrated by Arnold [13, 14], the solutions of the partial differential equation describing the evolution of the hydrodynamic flow of incompressible ideal fluid can be considered as a continuous measure preserving transformation of fluid velocity and the evolution is partially chaotic, that is, the flow is exponentially unstable in some directions and is stable in other directions (the details will be discussed in the last paragraph).

In many areas of mathematics the consideration of infinite-dimensional limits is an ambiguous procedure, and a priori there is no guarantee that a sensible limit of a given finite-dimensional structure exists. In our case the dimension of the N -dimensional torus $S^1 \otimes \dots \otimes S^1$ tends to infinity and it is unclear what type of phase space should be taken in the limit. The hint that a sensible limit may exist comes from the fact that as $N \rightarrow \infty$ the eigenvalues (2.6) fill out the cardioid curve more and more dense without producing any deformation of the cardioid curve that can be seen in figure Fig.1 [27]. The other important hint is that the inverse matrix $A(N)^{-1}$ is reminiscent to the matrix that represents the discrete version of the second-order differential operator [27]. If one supposes that the state vector becomes a function $\psi(x)$ with its argument on a real line $x \in R^1$, then it seems natural to look for a differential operator of the second order and the one that will reproduce the eigenvalue spectrum distributed on cardioid curve. Having in mind the above consideration let us consider the differential operator

$$A_x = 1 - 2 \exp\left(\frac{d}{dx}\right) + \exp\left(2\frac{d}{dx}\right) = \frac{d^2}{dx^2} + \frac{d^3}{dx^3} + \frac{7}{12} \frac{d^4}{dx^4} + \dots \quad (3.8)$$

acting in the Hilbert space of functions $\mathcal{H} = \{\psi(x)\}$ defined on the interval $x \in [-\infty, +\infty]$. The series expansion of the operator has indeed a second-order differential operator and also infinitely many high derivative terms. Its spectral characteristics are defined by the eigenvalue equation $A_x\psi(x) = \lambda\psi(x)$. Searching the eigenfunctions in the form of plane

waves

$$\psi_a(x) = e^{iax} \quad (3.9)$$

one can find that the spectrum represents a continuous cardioid curve on the complex plane

$$\lambda(a) = 1 - 2e^{ia} + e^{2ia} = 4e^{i(a+\pi)} \sin^2\left(\frac{a}{2}\right). \quad (3.10)$$

It is similar to the discrete spectrum (2.6) and has the periodic structure

$$\lambda(a + 2\pi k) = \lambda(a), \quad k = 0, \pm 1, \pm 2, \dots \quad (3.11)$$

The spectrum is continuous and the eigenvalues are distributed in the complex plane representing the cardioid curve shown in Fig.1. As the real momentum parameter a varies on the real line interval the eigenvalues run around the cardioid infinitely many time (3.11). The eigenvalues of the operator A_x can be divided into two sets $\{\lambda_\alpha\}$ and $\{\lambda_\beta\}$ with modulus smaller and larger than one: $0 < |\lambda_\alpha| < 1$, $1 < |\lambda_\beta|$. The eigenvalues λ_α and λ_β can be found using (3.10):

$$\begin{aligned} \lambda_\alpha &= 4e^{i(a+\pi)} \sin^2\left(\frac{a}{2}\right), \quad \text{when } -\frac{\pi}{3} + 2\pi k < a < +\frac{\pi}{3} + 2\pi k, \\ \lambda_\beta &= 4e^{i(a+\pi)} \sin^2\left(\frac{a}{2}\right), \quad \text{when } +\frac{\pi}{3} + 2\pi k < a < +\frac{5\pi}{3} + 2\pi k, \end{aligned} \quad (3.12)$$

where $k = 0, \pm 1, \pm 2, \dots$. This structure of the spectrum repeats itself with the period 2π . There are eigenvalues $|\lambda| = 1$ corresponding to $a = \pm\frac{\pi}{3} + 2\pi k$ where the cardioid intersects a unit circle shown in Fig.1.

4 The Kolmogorov-Sinai Entropy of Limiting System

In order to establish the fact that the operator A_x is defining a measure-preserving transformation one should calculate the determinant of the operator A_x . The measure-preserving transformations of the phase spaces is a characteristic property of Hamiltonian systems that is expressed in terms of the Liouville's theorem and represent a large class of dynamical systems that are considered in ergodic theory [20, 21, 22, 23, 24]. Using the fact that $\ln \text{Det} A_x = \text{Tr} \ln A_x$ we will have

$$\ln \text{Det} A_x = \sum_{k=-\infty}^{\infty} \int_{-\pi+2\pi k}^{+\pi+2\pi k} \ln[e^{i(a+\pi)} 4 \sin^2\left(\frac{a}{2}\right)] \frac{da}{2\pi} = 0, \quad (4.13)$$

that is, the determinant is equal to one $\text{Det} A_x = 1$ and the operator A_x is defining a measure-preserving transformation (the kernel subspace \mathcal{K} of the operator A_x will be defined in (4.23)).

One can define now the homomorphism of the Hilbert space $\mathcal{H} = \{\psi(x)\}$ in terms of the operator A_x as

$$\psi^{(1)}(x) = A_x \psi(x) \quad (4.14)$$

and the dynamical system on the infinite dimensional phase space \mathcal{H} as:

$$\psi^{(n)}(x) = A_x^n \psi(x) \quad \text{mod } 1, \quad n = 0, 1, 2, \dots \quad (4.15)$$

The homomorphism (4.15) is defined by mod 1 operation meaning that the functions $\psi^{(n)}(x)$ are wrapping around an infinitely long cylinder [68] $R^1 \otimes S^1$ ($\psi : R^1 \rightarrow S^1$).

The determinant of the operator A is equal to one and the eigenvalues are distributed inside and outside of the unit circle, and we have an example of infinite-dimensional hyperbolic C-K system of the type (2.2). To get convinced that the operator A_x represents a hyperbolic C-K system one should establish the existence of exponentially expanding and contracting foliations [41]. Let us consider the evolution of the infinitesimal perturbation $\psi \rightarrow \psi + \delta\psi$ under the action of A operator in analogy with the geodesic deviation equation:

$$\delta\psi^{(n)}(x) = A_x^n \delta\psi(x). \quad (4.16)$$

The deviation δL can be evaluated by using the standard inner product in Hilbert space and the mean value theorem [69]:

$$\begin{aligned} \delta L_n &= \langle \delta\psi^+ | A_x^n \delta\psi \rangle = \int_{-\infty}^{+\infty} \delta\psi^+(x) A_x^n \delta\psi(x) dx \\ &= \int_{-\infty}^{+\infty} \int_{\Delta a} \frac{da'}{2\pi} \int_{\Delta a} \frac{da}{2\pi} e^{-ia'x} \delta\phi^+(a') \left[4 \sin^2 \left(\frac{a}{2} \right) \right]^n e^{in(a+\pi)} e^{iax} \delta\phi(a) dx \\ &= \frac{1}{2\pi} \int_{\Delta a} da e^{in(a+\pi)} \left[4 \sin^2 \left(\frac{a}{2} \right) \right]^n |\delta\phi(a)|^2 = e^{in(\bar{a}+\pi)} e^{n \ln [4 \sin^2 (\frac{\bar{a}}{2})]} |\overline{\delta\phi^2}|, \end{aligned} \quad (4.17)$$

where \bar{a} is a number in the interval $\Delta a \subset (\pi/3, 5\pi/3)$ and $|\overline{\delta\phi^2}| = \frac{1}{2\pi} \int_{\Delta a} |\delta\phi|^2 da$. The absolute value of the deviation is growing exponentially with the iteration time n as

$$|\delta L_n| \sim |\overline{\delta\phi^2}| e^{n \ln [4 \sin^2 (\frac{\bar{a}}{2})]}. \quad (4.18)$$

In the above perturbation the Fourier spectrum of the function $\delta\psi(x)$ is localised in the region of the spectrum Δa , where the eigenvalues λ_β are larger than one, $\Delta a \subset (\pi/3, 5\pi/3)$, and are defined in (3.12). The integration over a and a' was specified to be in that region Δa and the perturbation function had the following form:

$$\delta\psi(x) = \int_{\Delta a} \delta\phi(a) e^{iax} \frac{da}{2\pi}. \quad (4.19)$$

In a similar way one can get convinced that the exponential contraction takes place when the Fourier spectrum of the perturbation function is localised in the part of the spectrum λ_α , where the eigenvalues are less than one $\Delta a \subset (-\pi/3, \pi/3)$ in (3.12). The existence of exponentially expanding and contracting foliations is a sufficient condition for a dynamical system to expose strong statistical/chaotic properties and, in particular, to have nonzero Kolmogorov-Sinai entropy and to be classified as a hyperbolic C-K system. In physical terms this means that the Fourier amplitudes $\phi(a)$ of the initial state vector $\psi(x)$ are stretched and compressed depending on whether the value of a is in the interval $a \in (\frac{\pi}{3} + 2\pi k, \frac{5\pi}{3} + 2\pi k)$ or in the interval $a \in (-\frac{\pi}{3} + 2\pi k, \frac{\pi}{3} + 2\pi k)$, where $k = 0, \pm 1, \pm 2, \dots$.

The above consideration allows to calculate the Kolmogorov entropy of the system per unit of the iteration time n as it was defined by Kolmogorov [39]. The new aspect that appears in this infinite-dimensional system case is that the spectrum (3.10) is continuous and repeats itself infinitely many times. For that reason the standard Kolmogorov definition [39] of the entropy per unit iteration time is equal to infinity. That can be observed also from the equation (2.7) when $N \rightarrow \infty$. In this circumstances one can propose to calculate the entropy per unit period (3.11) of the spectrum (3.10):

$$h(A_x) = \sum_{\alpha} \ln \frac{1}{|\lambda_{\alpha}|} = - \int_{-\pi/3}^{\pi/3} \ln[4 \sin^2(\frac{a}{2})] \frac{da}{2\pi} = 2i[Li_2(e^{i5\pi/3}) - Li_2(e^{i\pi/3})] \sim \frac{2}{\pi}, \quad (4.20)$$

where we used the fact that $\prod_{\alpha} \lambda_{\alpha} \prod_{\beta} \lambda_{\beta} = 1$ and $Li_n(z)$ is the polylogarithm function. This result is understandable in the sense that the finite-dimensional matrix system (2.5) considered above had the entropy $\sim \frac{2}{\pi} N$, where N is the dimension of the matrix operator. As far as the operator A can be considered as the infinite dimensional limit $N \rightarrow \infty$ of (2.5), the standard Kolmogorov entropy of the system (4.15) tends to infinity but its entropy "per spectral period" is finite. The Fig.(2) demonstrates an example of the iteration (4.15) of a smooth function.

The inverse operator G is defined by the equation

$$A_x G(x - y) = \sum_{k=-\infty}^{\infty} \delta(x - y - k) \quad (4.21)$$

and has the following solution:

$$G(x - y) = \frac{1}{2}(x - y)^2 \sum_{k=-\infty}^{\infty} e^{2\pi i k(x-y)}. \quad (4.22)$$

The general solution of (4.21) can be expressed as a sum the fixed solution (4.22) and an arbitrary element of the kernel \mathcal{K} . The kernel subspace \mathcal{K} of the operator A_x is defined by




Figure 2: The figure demonstrates the result of a triple iteration (4.15) $\psi^{(3)}(x) = A_x^3\psi(x)$ of the smooth function $\psi(x) = \sin(x) + \sin(2x) - \cos(3x) + \sin(4x)$.

the equation $A\psi_0(x) = 0$ and has the following solution:

$$\psi_0(x) = (c_1x + c_2) \sum_{k=-\infty}^{\infty} a_k e^{2\pi kix}, \quad (4.23)$$

where c_1, c_2, a_k are arbitrary constants. The spectrum belongs to the discrete values of spectral parameter $a = 2\pi k$ in (3.9) and have a zero measure in the continuous spectrum of the limiting system (3.8). The inverse transformation $\psi^{(-n)} = G^n\psi$, $n = 0, 1, 2, \dots$ is therefore defined modulo kernel (4.23). We will defined it in its most simple form (4.22):

$$\psi^{(-1)} = \int G(x-y)\psi(y)dy = \sum_{k=-\infty}^{\infty} \frac{1}{2}k^2 \psi(x-k) \quad \text{mod } 1. \quad (4.24)$$

Thus the evolution of the system is given in both "time directions" by (4.24) and (4.14).

Because the determinant of the operator A_x is equal to one on a quotient space \mathcal{H}/\mathcal{K} [70] of functions on a cylinder $R^1 \otimes S^1$, where \mathcal{K} is a kernel (4.23), it is natural to think that the operator A_x defines a measure preserving transformation. In this circumstances one can try to define a measure that is invariant with respect to the transformations generated by A_x . The volume element \mathcal{V} in the quotient space \mathcal{H}/\mathcal{K} can be defined by using the Wiener-Feynman functional integral $\mathcal{V}_C = \int_C F[\psi] \mathcal{D}\psi(x)$, where C is a subset in \mathcal{H}/\mathcal{K} and the functional $F[A_x\psi] = F[\psi]$ should be invariant under the action of the transformations generated by the operator A_x . The measure $\mathcal{D}\psi(x)$ is invariant because the determinant of the corresponding Jacobian operator is equal to one (4.13). The invariant functional $F[\psi]$ can be constructed by projecting the state vectors on the eigenfunctions $\phi(\pm\frac{\pi}{3} + 2\pi k) = \int_{-\infty}^{\infty} e^{i(\pm\frac{\pi}{3} + 2\pi k)x} \psi(x) dx$ (3.9) that correspond to the eigenvalues of modulus one $|\lambda(\pm\frac{\pi}{3} + 2\pi k)| = 1$, $k = 0, \pm 1, \pm 2, \dots$. The invariant functional $F[\psi]$ can be defined as:

$$F[\psi] = \exp \left[- \sum_{k=\pm 1, \pm 2, \dots} \frac{|\phi(\frac{\pi}{3} + 2\pi k)|^2 + |\phi(-\frac{\pi}{3} + 2\pi k)|^2}{2} \right]. \quad (4.25)$$

5 *Flow of Incompressible Ideal Fluid*

Were similar systems investigated and successfully used in the past? The solutions of the partial differential equation describing the evolution $t \rightarrow g_t(x)$ of the hydrodynamic flow of incompressible ideal fluid filled in a two dimensional torus $x \in \mathcal{T}^2$ can be considered as a continuous area preserving diffeomorphisms $SDiff(\mathcal{T}^2)$ of a torus $\mathcal{T}^2 \rightarrow \mathcal{T}^2$. In Arnold approach [13] the ideal fluid flow is described by the geodesics $g_t(x) \in G$ on the diffeomorphism group $G = SDiff(\mathcal{T}^2)$ with the velocity $v_t(x) = \dot{g}_t(x)g_t^{-1}(x)$ belonging to the corresponding algebra $\mathfrak{g} = \text{sdiff}(\mathcal{T}^2)$ of divergence free vector fields. The Riemannian metric on the group G is induced from the metric on a torus [13] and the stability of the geodesic flows on the group G can be analysed by investigating the behaviour of the corresponding sectional curvatures $K(v, \delta v)$ [13, 41, 71]. It was found that the flows that are defined by a parallel velocity field on \mathcal{T}^2 are unstable because the sectional curvatures are negative and the flow is exponentially unstable. In other directions the sectional curvatures are positive and the flows are stable [13]. A similar stability analysis was performed for the hydrodynamic flow on two-dimensional sphere \mathcal{S}^2 in [14, 73] and on high-dimensional torus \mathcal{T}^N in [72, 73]. In all these cases the flow is exponentially unstable in some directions and is stable in some other directions, resulting in the limitation of predictability of the hydrodynamic flow and leading to the principal difficulties of a long-term "weather forecasting" [13]. Comparing these systems with the system considered above one can observe that here we have discrete in time transformations of the phase space and, secondly, the system (3.8), (4.14), (4.24) shows up exponential instability of its geodesics in the full quotient phase space \mathcal{H}/\mathcal{K} . This full phase space chaotic behaviour can find application in Monte Carlo method, statistical physics and in digital signal processing in communication systems, the subjects of future investigation.

6 *Conclusion*

We were able to define the infinite-dimensional limit of maximally chaotic dynamical systems on N -dimensional tori when the dimension N tends to infinity. The limiting system A_x is represented by a nonlocal differential operator acting on the infinite-dimensional Hilbert space of continuous functions. We calculated the eigenvalue spectrum of the limiting operator, as well as its determinant and the corresponding Kolmogorov-Sinai entropy. We investigated the exponentially expanding and contracting foliations and demonstrated that the limiting

system belongs to the class of hyperbolic Anosov C-systems. The limiting system represents a unique example of a C-system of infinite dimensionality, which has a quite simple form and is chaotic in its full phase space.

7 Acknowledgement

The author would like to thank Konstantin Savvidy for stimulating discussions. This work was supported by the Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Grant Agreement No 644121.

8 Appendix

The motion of an abstract "rigid body" rotating in high-dimensional Euclidean space $\vec{r} \in E$ that is invariant under the isometry group $g(\xi) \in G$ can be described in terms of geodesic flow on a corresponding group manifold [13]. The stationary frame coordinates of the "rigid body" are defined as $\vec{r} = g_t \vec{Q}$, where $g_t = g(\xi(t))$ is a time-dependent element of the matrix group G and \vec{Q} are the frame coordinates rigidly fixed to the rotating "body" $\dot{\vec{Q}} = 0$. Thus

$$\dot{\vec{r}} = \dot{g}_t \vec{Q} = \dot{g}_t g_t^{-1} g_t \vec{Q} = \hat{\omega}_s \vec{r}. \quad (8.26)$$

The matrix of angular velocity in stationary frame

$$\hat{\omega}_s = \dot{g}_t g_t^{-1} \quad (8.27)$$

is a right-invariant one-form ($d(gg_0)(gg_0)^{-1} = dgg^{-1}$, where g_0 is a fixed element of the group G). The matrix of angular velocity in rotating frame

$$\hat{\Omega}_c = g_t^{-1} \dot{g}_t \quad (8.28)$$

is left invariant because $(g_0 g)^{-1} d(g_0 g) = g^{-1} dg$. It follows that*

$$\hat{\omega}_s = \dot{g}_t g_t^{-1} = g_t g_t^{-1} \dot{g}_t g_t^{-1} = g_t \hat{\Omega}_c g_t^{-1}. \quad (8.29)$$

The kinetic energy is defined as a sum of the kinetic energies of all "parts" of the rotating "body" through the velocities (8.26):

$$T = \frac{1}{2} \sum_a m_a \dot{\vec{r}}_a \dot{\vec{r}}_a = \frac{1}{2} Tr(\hat{I}_s \hat{\omega}_s \hat{\omega}_s^+) = -\frac{1}{2} Tr(\hat{I}_s \hat{\omega}_s \hat{\omega}_s), \quad (8.30)$$

*The general relation between operators in stationary and rotating frames is $A_s = g_t A_c g_t^{-1}$.

where one should use the relation $\vec{r} = g_t \vec{Q}$, and therefore

$$\hat{I}_s = \sum_a m_a r_a r_a^+ = g_t \hat{I} g_t^{-1}, \quad \hat{I} = \sum_a m_a Q_a Q_a^+. \quad (8.31)$$

The matrix \hat{I} is a symmetric positive definite constant matrix that determines the "moment of inertia" in the frame rigidly fixed to the rotating "body". The matrix of angular momentum in stationary frame is

$$\hat{m}_s = \hat{I}_s \hat{\omega}_s \quad (8.32)$$

and the corresponding angular momentum in rotating frame can be defined by projection of \hat{m}_s into the rotating frame:

$$\hat{m}_s = g_t \hat{M}_c g_t^{-1}, \quad (8.33)$$

thus[†]

$$\hat{M}_c = \hat{I} \hat{\Omega}_c, \quad (8.34)$$

where one should use the relations (8.32), (8.31) and (8.27), (8.28). In terms of rotating frame coordinates the kinetic energy (8.30) will take the form

$$T = -\frac{1}{2} \text{Tr}(\hat{I}_s \hat{\omega}_s \hat{\omega}_s) = -\frac{1}{2} \text{Tr}(\hat{m}_s \hat{\omega}_s) = -\frac{1}{2} \text{Tr}(\hat{M}_c \hat{\Omega}_c) = -\frac{1}{2} \text{Tr}(\hat{I} \hat{\Omega}_c \hat{\Omega}_c), \quad (8.35)$$

where we used the relations (8.33) and (8.29). As it follows from (8.35) and (8.40), in mathematical terms the matrix \hat{I} defines the alternative Euclidean structure on the group algebra $\langle a, b \rangle_I = \text{Tr}(T_a \hat{I} T_b)$, where T^a are the generators of the algebra \mathfrak{g} and $\hat{\Omega}_c = \sum_a T_a \Omega^a$.

Using the relation (8.33) and the conservation of the angular momentum in stationary frame $\dot{\hat{m}}_s = 0$ one can get the generalised Euler equation

$$\dot{g}_t \hat{M}_c g_t^{-1} + g_t \dot{\hat{M}}_c g_t^{-1} - g_t \hat{M}_c g_t^{-1} \dot{g}_t g_t^{-1} = 0,$$

which can be represented in the following standard form:

$$\dot{\hat{M}}_c + \hat{\Omega}_c \hat{M}_c - \hat{M}_c \hat{\Omega}_c = 0,$$

or equivalently as

$$\frac{d\hat{M}_c}{dt} = [\hat{M}_c; \hat{\Omega}_c]. \quad (8.36)$$

In terms of angular velocity it takes the following form:

$$\hat{I} \frac{d\hat{\Omega}_c}{dt} = [\hat{I} \hat{\Omega}_c; \hat{\Omega}_c], \quad \frac{d\hat{\Omega}_c}{dt} = \hat{I}^{-1} [\hat{I} \hat{\Omega}_c; \hat{\Omega}_c] = \Gamma(\hat{\Omega}_c, \hat{\Omega}_c). \quad (8.37)$$

[†]The square of angular momentum is conserved: $\text{Tr}(\hat{m}_s^2) = \text{Tr}[(g_t \hat{M}_c g_t^{-1})^2] = \text{Tr}(\hat{M}_c^2)$.

The last Euler equation can be represented in the form of geodesic equation

$$\frac{d\Omega^a}{dt} + \Gamma_{bd}^a \Omega^b \Omega^d = 0, \quad (8.38)$$

where Γ_{bd}^a are the Christoper symbols of the metric (8.40). The kinetic energy defines the left invariant metric on the group:

$$ds^2 = Tr(\hat{I}\hat{\Omega}_c \hat{\Omega}_c)dt^2 = Tr(\hat{I} g^{-1}dg g^{-1}dg) = Tr(\hat{I} g^{-1} \frac{\partial g}{\partial \xi^a} g^{-1} \frac{\partial g}{\partial \xi^b}) d\xi^a d\xi^b, \quad (8.39)$$

where the ξ^a are parameters of the Lie group G and

$$ds^2 = g_{ab} d\xi^a d\xi^b, \quad g_{ab} = Tr(g^{-1} \frac{\partial g}{\partial \xi^b} \hat{I} g^{-1} \frac{\partial g}{\partial \xi^a}). \quad (8.40)$$

The calculation of the components of the Riemann tensor and of the sectional curvatures

$$K(\xi, \eta) = \frac{R_{abcd}\xi^a\eta^b\xi^c\eta^d}{|\xi \wedge \eta|^2} \quad (8.41)$$

allows to investigate the stability of the geodesic flows even in the cases when G is the infinite-dimensional group of diffeomorphisms $SDiff(\mathcal{M})$ that describes the flow of incompressible ideal fluid filled in a manifold \mathcal{M} [13, 14, 72, 73].

9 Data Availability Statement.

The data that support the findings of this study are available within the article and also in the references [25, 26, 27, 62, 63, 64, 65, 67].

References

- [1] S. H. Shenker and D. Stanford, *Black holes and the butterfly effect*, JHEP **1403** (2014) 067 doi:10.1007/JHEP03(2014)067 [arXiv:1306.0622 [hep-th]].
- [2] J. Maldacena, S. H. Shenker and D. Stanford, *A bound on chaos*, JHEP **1608** (2016) 106 doi:10.1007/JHEP08(2016)106 [arXiv:1503.01409 [hep-th]].
- [3] G. Gur-Ari, M. Hanada and S. H. Shenker, *Chaos in Classical D0-Brane Mechanics*, JHEP **1602** (2016) 091 doi:10.1007/JHEP02(2016)091 [arXiv:1512.00019 [hep-th]].
- [4] J. S. Cotler *et al.*, *Black Holes and Random Matrices*, JHEP **1705** (2017) 118 Erratum: [JHEP **1809** (2018) 002] doi:10.1007/JHEP09(2018)002, 10.1007/JHEP05(2017)118.

- [5] I. Y. Aref'eva, P. B. Medvedev, O. A. Rytchkov and I. V. Volovich, *Chaos in M(matrix) theory*, Chaos Solitons Fractals **10** (1999) 213, [hep-th/9710032].
- [6] Y. Sekino and L. Susskind, *Fast Scramblers*, JHEP **10** (2008), 065 doi:10.1088/1126-6708/2008/10/065 [arXiv:0808.2096 [hep-th]].
- [7] G. K. Savvidy, *Classical and Quantum Mechanics of non-Abelian Gauge Fields*, Nucl. Phys. B **246** (1984) 302. doi:10.1016/0550-3213(84)90298-0
- [8] G. Savvidy, *Maximally chaotic dynamical systems*, Annals Phys. **421** (2020), 168274 doi:10.1016/j.aop.2020.168274
- [9] D. J. Gross and V. Rosenhaus, *Chaotic scattering of highly excited strings*, [arXiv:2103.15301 [hep-th]].
- [10] G. 't Hooft, *Black holes, Hawking radiation, and the information paradox*, Nucl. Phys. B Proc. Suppl. **43** (1995), 1-11 doi:10.1016/0920-5632(95)00444-E
- [11] G. 't Hooft, *The Scattering matrix approach for the quantum black hole: An Overview*, Int. J. Mod. Phys. A **11** (1996), 4623-4688 doi:10.1142/S0217751X96002145 [arXiv:gr-qc/9607022 [gr-qc]].
- [12] P. Betzios, N. Gaddam and O. Papadoulaki, *Black Hole S-matrix for a scalar field*, [arXiv:2012.09834 [hep-th]].
- [13] V. Arnold, *Sur la géométrie des groupes de Lie de dimension infinie et ses applications en hydrodynamique des fluides parfaits*, Ann.Inst. Fourier **16**, No 1 (1966) 319-361
- [14] A. Arakelian and G. K. Savvidy, *Geometry of a Group of Area Preserving Diffeomorphisms*, Phys. Lett. B **223** (1989), 41-46 doi:10.1016/0370-2693(89)90916-7
- [15] S. Endlich, A. Nicolis, R. Rattazzi and J. Wang, *The Quantum mechanics of perfect fluids*, JHEP **04** (2011), 102 doi:10.1007/JHEP04(2011)102 [arXiv:1011.6396 [hep-th]].
- [16] S. Chandrasekhar, *Principles of Stellar Dynamics*, Chicago: University of Chicago Press; London: Cambridge University Press, 1942.
- [17] V. Gurzadyan and G. Savvidy, *Collective relaxation of stellar systems*, Astron. Astrophys. **160** (1986) 203

- [18] G. W. Gibbons, *The Jacobi-metric for timelike geodesics in static spacetimes*, Class. Quant. Grav. **33** (2016) no.2, 025004 doi:10.1088/0264-9381/33/2/025004.
- [19] V. A. Belinsky, I. M. Khalatnikov and E. M. Lifshitz, *A General Solution of the Einstein Equations with a Time Singularity*, Adv. Phys. **31** (1982), 639-667 doi:10.1080/00018738200101428
- [20] J.W.Gibbs, *Elementary principles in statistical mechanics*, Charles Scribner's Sons, New York 1902
- [21] E. Hopf, *Proof of Gibbs Hypothesis on the Tendency Toward Statistical Equilibrium*, Mathematics **18** (1932) 333.
- [22] N.S.Krylov, *Works on the foundation of statistical physics*, M.- L. Izdatelstvo Acad.Nauk. SSSR, 1950; (Princeton University Press, 1979)
- [23] V.Arnold and A.Avez, *Ergodic Problems of Classical Mechanics*, (The Mathematical physics monograph series) Benjamin (July 5, 1968), 286pp.
- [24] I. P. Kornfeld, S. V. Fomin, Y. G. Sinai, *Ergodic Theory*, Springer, 1982
- [25] G. Savvidy and N. Ter-Arutyunyan-Savvidy, *On the Monte Carlo simulation of physical systems*, J.Comput.Phys. **97** (1991) 566; Preprint EFI-865-16-86-YEREVAN, Jan. 1986.
- [26] K.Savvidy, *The MIXMAX random number generator*, Comput.Phys.Commun. 196 (2015) 161-165. (<http://dx.doi.org/10.1016/j.cpc.2015.06.003>); arXiv:1404.5355
- [27] K. Savvidy and G. Savvidy, *Spectrum and Entropy of C-systems. MIXMAX random number generator*, Chaos Solitons Fractals **91** (2016) 33 doi:10.1016/j.chaos.2016.05.003.
- [28] S. W. Hawking, *Breakdown of Predictability in Gravitational Collapse*, Phys. Rev. D **14** (1976), 2460-2473 doi:10.1103/PhysRevD.14.2460
- [29] S. W. Hawking, *The Information Paradox for Black Holes*, [arXiv:1509.01147 [hep-th]].
- [30] D. N. Page, *Information in black hole radiation*, Phys. Rev. Lett. **71** (1993), 3743-3746 doi:10.1103/PhysRevLett.71.3743 [arXiv:hep-th/9306083 [hep-th]].

- [31] L. Susskind, L. Thorlacius and J. Uglum, *The Stretched horizon and black hole complementarity*, Phys. Rev. D **48** (1993), 3743-3761 doi:10.1103/PhysRevD.48.3743 [arXiv:hep-th/9306069 [hep-th]].
- [32] S. W. Hawking, M. J. Perry and A. Strominger, *Soft Hair on Black Holes*, Phys. Rev. Lett. **116** (2016) no.23, 231301 doi:10.1103/PhysRevLett.116.231301 [arXiv:1601.00921 [hep-th]].
- [33] A. Strominger and A. Zhiboedov, *Gravitational Memory, BMS Supertranslations and Soft Theorems*, JHEP **01** (2016), 086 doi:10.1007/JHEP01(2016)086 [arXiv:1411.5745 [hep-th]].
- [34] G. Penington, S. H. Shenker, D. Stanford and Z. Yang, *Replica wormholes and the black hole interior*, [arXiv:1911.11977 [hep-th]].
- [35] D. V. Anosov and Ya. G. Sinai, *Certain smooth ergodic systems*, Uspehi Mat. Nauk **22** (1967), no. 5 (137), 107-172; Russian Math. Surveys **22** (1967), 103-167.
- [36] V. P. Leonov, *On the central limit theorem for ergodic endomorphisms of the compact commutative groups*, Dokl. Acad. Nauk SSSR, **124** No: 5 (1969) 980-983
- [37] M. C. Gutzwiller, *Stochastic Behaviour in Quantum Scattering*, Physica **7D** (1983) 341-355.
- [38] A.N. Kolmogorov, *New metrical invariant of transitive dynamical systems and automorphisms of Lebesgue spaces*, Dokl. Acad. Nauk SSSR, **119** (1958) 861-865
- [39] A.N. Kolmogorov, *On the entropy per unit time as a metrical invariant of automorphism*, Dokl. Acad. Nauk SSSR, **124** (1959) 754-755
- [40] Ya.G. Sinai, *On the Notion of Entropy of a Dynamical System*, Doklady of Russian Academy of Sciences, **124** (1959) 768-771.
- [41] D. V. Anosov, *Geodesic flows on closed Riemannian manifolds with negative curvature*, Trudy Mat. Inst. Steklov., Vol. **90** (1967) 3 - 210
- [42] Emil Artin, *Ein mechanisches system mit quasi-ergodischen bahnen*, E. Abh. Math. Semin. Univ. Hambg. (1924) 3: 170.

- [43] J. Hadamard, *Sur le billiard non Euclidean*, Soc. Sci. Bordeaux, Proc. Verbaux 1898, 147 (1898); J. Math Pure Appl. 4 (1898) 27.
- [44] G. Hedlund, *The dynamics of geodesic flow*, Bull.Am.Math.Soc. **45** (1939) 241-246
- [45] E. Hopf, *Ergodic theory and the geodesic flow on surfaces of constant negative curvature*, Bull. Amer. Math. Soc, **77** (1971) 863- 877.
- [46] V. A. Rokhlin, *Metric properties of endomorphisms of compact commutative groups*, Izv. Akad. Nauk SSSR Ser. Mat., Volume **28**, Issue 4 (1964) 867- 874
- [47] V.A. Rokhlin, *On the entropy of automorphisms of compact commutative groups*, Teor. Ver. i Pril., vol. **3**, issue 3 (1961) p. 351
- [48] S. Smale, *Differentiable dynamical systems*. Bull. Am. Math. Soc. **73** (1967) 747-817
- [49] Ya. G. Sinai, *Markov partitions and C-diffeomorphisms*, Funkcional. Anal, i Prilozen. **2** (1968),64-89; Functional Anal. Appl. **2** (1968) 61-82.
- [50] Ya. G. Sinai, *Proceedings of the International Congress of Mathematicians*, Uppsala (1963) 540-559.
- [51] G. A. Margulis, *Certain measures that are connected with C-flows on compact manifolds*, Funkcional. Anal, i Prilozen. **4** (1970) 62-76; Functional Anal. Appli. **4** (1970) 55-67.
- [52] R.Bowen, *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms*. (Lecture Notes in Math., 470: A. Dold and B. Eckmann, editors). Springer-Verlag 1975.
- [53] R.Bowen, *Periodic orbits for hyperbolic flows*, Amer. J. Math.,**94** (1972), 1-30.
- [54] A. L. Gines, *Metrical properties of the endomorphisms on m-dimensional torus*, Dokl. Acad. Nauk SSSR, **138** (1961) 991-993
- [55] *The property to which many authors refer as to a "butterfly effect".*
- [56] G. Basesyan, S. Matinyan and G. Savvidy, *Nonlinear plane waves in the massless Yang-Mills theory*, Pisma Zh. Eksp. Teor. Fiz. **29** (1979) 641-644
- [57] S. Matinyan, G. Savvidy and N. Ter-Arutyunyan-Savvidi, *Classical Yang-Mills mechanics. Nonlinear colour oscillations*, Zh. Eksp. Teor. Fiz. **80** (1980) 830-838

- [58] B.V. Chirikov, D.L. Shepelyansky, *Stochastic oscillation of classical Yang-Mills fields*, JETP Lett. **34** (1981) 163.
- [59] G. M. Asatrian and G. K. Savvidy, *Configuration Manifold of Yang-Mills Classical Mechanics*, Phys. Lett. A **99** (1983) 290. doi:10.1016/0375-9601(83)90887-3
- [60] G. Savvidy, *The Yang-Mills classical mechanics as a Kolmogorov system*, Phys. Lett. **130B** (1983) 303-307
- [61] G. Savvidy, *Yang-Mills Quantum Mechanics*, Phys. Lett. **159B** (1985) 325. doi:10.1016/0370-2693(85)90260-6
- [62] K. Savvidy, *MIXMAX software in HEPFORGE.ORG*, <http://mixmax.hepforge.org>
- [63] K. Savvidy, *MIXMAX software in Boost C++ Libraries*, https://www.boost.org/doc/libs/1_76_0/boost/random/mixmax.hpp
- [64] CERN, *CLHEP- A Class Library for High Energy Physics*, <http://proj-clhep.web.cern.ch/proj-clhep/>; <https://gitlab.cern.ch/CLHEP/CLHEP/blob/develop/Random/src/MixMaxRng.cc>
- [65] CERN, *ROOT, Data Analysis Framework, An open-source analysis framework used by high energy physics and others*, https://root.cern.ch/doc/master/classROOT_1_1Math_1_1MixMaxEngine.html
- [66] At CERN the CMS experiment, in its full simulation software for Run-2, includes MIXMAX as default engine. https://indico.cern.ch/event/587955/contributions/2937635/attachments/1679273/2706817/PosterCMS_SIM_v4.pdf
- [67] CERN, *MIXMAX workshop*, <https://indico.cern.ch/event/404547>
- [68] *One can imagine this cylinder as appearing in the "decompactification" of a 2-torus $S^1 \otimes S^1$, when cutting a 2-torus with a non-contractible circle S^1 and stretching the open boundary circles to the infinities.*
- [69] *If $f(x)$ is a continuous function and $g(x)$ is a nonnegative integrable function on $[a, b]$, then there exists some number $\bar{x} \in (a, b)$, such that $\int_a^b f(x)g(x)dx = f(\bar{x}) \int_a^b g(x)dx$.*
- [70] *These are the functions orthogonal to the $\psi_0(x)$ (4.23).*

[71] See Appendix for details.

[72] A. M. Lukatzki, *On curvature of the group of measure preserving diffeomorphisms of n -dimensional torus*, Uspekhi Mat. Nauk **36** (1981) 187

[73] N. K. Smolentsev, *Diffeomorphism Groups of Compact Manifolds*, J. Math. Sciences **146** (2007) 6213