

Gravitational radiation from binary systems in massive graviton theories

Tanmay Kumar Poddar,^{1,*} Subhendra Mohanty,^{1,†} and Soumya Jana^{2,‡}

¹*Theoretical Physics Division, Physical Research Laboratory, Ahmedabad - 380009, India*

²*Department of Physics, Sitananda College, Nandigram, 721631, India*

Abstract

Theories with massive gravitons have peculiarity called the van Dam-Veltman-Zakharov discontinuity in that the massive theory propagator does not go to the massless graviton propagator in the zero graviton mass limit. This results in large deviation in Newton's law for massive graviton theories even when the graviton mass vanishes. We test the vDVZ in massive graviton theories for single graviton vertex process namely the gravitational radiation from a classical source. We calculate the gravitational radiation from compact binaries using the perturbative Feynman diagram method. We perform this calculation for Einstein's gravity with massless gravitons and verify that the Feynman diagram calculation reproduces the quadrupole formula. Using the same procedure we calculate the gravitational radiation for three massive graviton theories: (1) the Fierz-Pauli theory (2) the modified Fierz-Pauli theory without the vDVZ discontinuity and (3) the Dvali-Gabadadze-Porrati theory with a momentum dependent graviton mass. We put limits on the graviton mass in each of these theories from observations of binary pulsar timings.

* Email Address: tanmay@prl.res.in

† Email Address: mohanty@prl.res.in

‡ Email Address: soumyajana.physics@gmail.com

I. INTRODUCTION

Einstein's general relativity (GR), since its inception in 1916, has passed all experimental tests [1]. In order to move towards the correct quantum theory of gravity it is important to test which variations of classical GR fail the experimental tests or have some theoretical inconsistencies. One such variation of GR which has been widely studied is the Fierz-Pauli (FP) theory of massive gravity [2]. In a scalar or vector field theory a massive particle exchange gives rise to a $(1/r)e^{-m_g r}$ Yukawa potential which goes to the $1/r$ potential in the $m_g \rightarrow 0$ limit. The FP theory of massive graviton has the peculiarity that in the zero graviton mass limit the Lagrangian goes smoothly to Einstein-Hilbert (EH) linearized gravity theory, while the graviton propagator has additional contributions from the scalar modes of the metric which do not decouple in the zero graviton mass limit. As a result the Newtonian potential in the zero-mass limit of FP theory is a factor $(4/3)$ larger than the prediction from the EH theory (which of-course agrees with the Newtonian potential). This peculiarity of the FP theory where the action goes to the EH theory in the zero mass limit but the graviton propagator does not was first pointed out by van Dam and Veltman [3] and independently by Zakharov [4] and this feature which arises in most massive gravity theories [5–8] is called the van Dam-Veltman-Zakharov (vDVZ) discontinuity. Experimental constraints on the graviton mass are listed in [9].

It is of interest to ask if instead of a graviton exchange diagram we consider a one graviton vertex process like gravitational radiation then, whether there is a difference in the result between the predictions of GR and the predictions of massive gravity theories in the $m_g \rightarrow 0$ limit and whether a manifestation of the vDVZ discontinuity can be seen in this phenomenon.

GR in the weak field limit can be treated as a quantum field theory of spin-2 fields in the Minkowski space [10–13]. Any classical gravity interaction like Newtonian potential between massive bodies or bending of light by a massive body can be described by a tree level graviton exchange diagram. The result of the tree level diagrams should match the weak field classical GR results. The derivation of gravitational radiation from binary stars as single vertex Feynman diagram of graviton emission from a classical source has been performed in [14, 15] and the results match with the result of Peter and Mathews [16] who

used the quadrupole formula of classical GR.

The first evidence of Gravitational Wave (GW) radiation was obtained from precision measurements of the Hulse -Taylor binary system [17–19]. The orbital period loss of the compact binary system confirms Einstein’s GR [16] to $\sim 0.1\%$ accuracy [20]. Following the Hulse-Taylor binary there have been other precession measurements from compact binary systems [21–23].

Binary stars can also radiate other ultra-light fields like axions or gauge bosons. The angular frequency of pulsar binaries is $\Omega \sim 10^{-19}$ eV and particles with mass lower than Ω can be radiated in a manner similar to the radiation of gravitational waves. The Feynman diagram method is pedagogically simpler to generalise the calculation of scalars and gauge bosons. A calculation of radiation of ultra-light scalars [14], axions [24] and gauge bosons [25] has been performed with this method and compared with experimental observations of binary pulsars (or pulsar-white dwarf binaries). This enables us to probe the couplings of ultra-light dark matter [26, 27] which are predicted to be in the mass range $\sim 10^{-21} - 10^{-22}$ eV to be probed with binary pulsar timing measurements.

In this paper we study massive graviton theories with a single vertex process namely graviton radiation from binary stars and we consider three models (1) the Fierz-Pauli ghost free theory which has a vDVZ discontinuity in the propagator, (2) a modification of Fierz-Pauli theory where there is a cancellation between the ghost and the scalar degrees so that there is no vDVZ discontinuity [28–31] and (3) the Dvali-Gabadadze-Porrati (DGP) theory [32–34] which is ghost-free but the extra scalar degree of freedom gives rise to the vDVZ discontinuity. The mass term in DGP gravity is momentum dependent which serves the purpose of suppressing the long range interactions in a virtual graviton exchange process. For real gravitons the graviton mass is tachyonic. We compare our results with observations and put limits on the graviton mass allowed in each of these theories.

We also compare our results with the earlier classical field calculations in massive gravity theories [29, 35–39].

This paper is organised as follows. In Section II we discuss the Fierz-Pauli theory and derive the formula for energy loss by graviton radiation using the Feynman diagram method. In Section III we do the same study for the modified FP theory without the vDVZ discon-




FIG. 1: Emission of graviton from a classical source.

tinuity and in Section IV we study the DGP theory. In Section V we compare the results with observations from the Hulse-Taylor binary (PSR B1913+16) and pulsar white dwarf binary (PSR J1738+0333) and put limits on the graviton mass for each of the massive gravity theories discussed. We also discuss the limits of applicability of the perturbation theory from the Vainshtein criterion and the corresponding limits on the range of graviton mass established from binary stars. In the Section VI we summarise the results and discuss future directions. In the Appendix A we give the detailed derivation of the Feynman diagram method of calculating gravitational radiation from compact binaries in GR for comparison with massive gravity theory results discussed in this paper.

II. FIERZ-PAULI MASSIVE GRAVITY THEORY

The Fierz-Pauli theory [2] is described by the action

$$\begin{aligned}
 S &= \int d^4x \left[-\frac{1}{2}(\partial_\mu h_{\nu\rho})^2 + \frac{1}{2}(\partial_\mu h)^2 - (\partial_\mu h)(\partial^\nu h_\nu^\mu) + (\partial_\mu h_{\nu\rho})(\partial^\nu h^{\mu\rho}) \right. \\
 &\quad \left. + \frac{1}{2}m_g^2(h_{\mu\nu}h^{\mu\nu} - h^2) + \frac{\kappa}{2}h_{\mu\nu}T^{\mu\nu} \right] \\
 &= \int d^4x \left[\frac{1}{2}h_{\mu\nu}\mathcal{E}^{\mu\nu\alpha\beta}h_{\alpha\beta} + \frac{1}{2}m_g^2h_{\mu\nu}(\eta^{\mu(\alpha}\eta^{\beta)\nu} - \eta^{\mu\nu}\eta^{\alpha\beta})h_{\alpha\beta} + \frac{\kappa}{2}h_{\mu\nu}T^{\mu\nu} \right], \quad (2.1)
 \end{aligned}$$

where the operator $\mathcal{E}^{\mu\nu\alpha\beta}$ is given in Eq.A4. The mass term breaks the gauge symmetry $h_{\mu\nu} \rightarrow h_{\mu\nu} - \partial_\mu \xi_\nu - \partial_\nu \xi_\mu$. We will assume that the energy-momentum is conserved, $\partial_\mu T^{\mu\nu} = 0$.

The e.o.m from Eq.2.1 is

$$(\square + m_g^2) h_{\mu\nu} - \eta_{\mu\nu} (\square + m_g^2) h - \partial_\mu \partial^\alpha h_{\alpha\nu} - \partial_\nu \partial^\alpha h_{\alpha\mu} + \eta_{\mu\nu} \partial^\alpha \partial^\beta h_{\alpha\beta} + \partial_\mu \partial_\nu h = -\kappa T_{\mu\nu}. \quad (2.2)$$

Taking the divergence of Eq.2.2 we have

$$m_g^2 (\partial^\mu h_{\mu\nu} - \partial_\nu h) = 0. \quad (2.3)$$

These are 4 constraint equations which reduce the independent degrees of freedom of the graviton from 10 to 6.

Using Eq.2.3 in Eq.2.2 we obtain

$$\square h_{\mu\nu} - \partial_\mu \partial_\nu h + m_g^2 (h_{\mu\nu} - \eta_{\mu\nu} h) = -\kappa T_{\mu\nu}. \quad (2.4)$$

Taking the trace of this equation we obtain the relation

$$h = \frac{\kappa}{3m_g^2} T. \quad (2.5)$$

Therefore trace h is not a propagating mode but is determined algebraically from the trace of the stress tensor. This is the ghost mode as the kinetic term for h in e.o.m Eq.2.2 appears with the wrong sign. Therefore in the Fierz-Pauli theory the ghost mode does not propagate. The number of independent propagating degrees of freedom of the Fierz Pauli theory is therefore 5. These are 2 tensor modes, 2 three-vector d.o.f which do not couple to the energy-momentum tensor and 1 scalar which couples to the trace of the energy-momentum tensor.

The propagator in the FP theory is given formally by

$$[\mathcal{E}^{\mu\nu\alpha\beta} + m_g^2 (\eta^{\mu(\alpha} \eta^{\beta)\nu} - \eta^{\mu\nu} \eta^{\alpha\beta})] D_{\alpha\beta\rho\sigma}^{(m)}(x-y) = \delta_{(\rho}^{\mu} \delta_{\sigma)}^{\nu} \delta^4(x-y). \quad (2.6)$$

Going to momentum space ($\partial_\mu \rightarrow ik_\mu$) we can find $D_{\alpha\beta\rho\sigma}^{(m)}(k)$ from Eq.2.6. The propagator for the Pauli-Fierz massive graviton turns out to be

$$D_{\alpha\beta\rho\sigma}^{(m)}(k) = \frac{1}{-k^2 + m_g^2} \left(\frac{1}{2} (P_{\alpha\rho} P_{\beta\sigma} + P_{\alpha\sigma} P_{\beta\rho}) - \frac{1}{3} P_{\alpha\beta} P_{\rho\sigma} \right), \quad (2.7)$$

where

$$P_{\alpha\beta} \equiv \eta_{\alpha\beta} - \frac{k_\alpha k_\beta}{m_g^2}. \quad (2.8)$$

In tree level processes where there is a graviton exchange between conserved currents, the amplitude is of the form

$$\mathcal{A}_{FP} = \frac{\kappa^2}{4} T^{\alpha\beta} D_{\alpha\beta\mu\nu}^{(m)} T'^{\mu\nu}. \quad (2.9)$$

The momentum dependent terms will vanish due to conservation of the stress tensor $k_\mu T^{\mu\nu} = k_\nu T^{\mu\nu} = 0$. Hence, for tree level calculations one may drop the momentum dependent terms in Eq.2.7 and the propagator for the FP theory may be written as

$$D_{\mu\nu\alpha\beta}^{(m)}(p) = \frac{1}{-k^2 + m_g^2} \left(\frac{1}{2}(\eta_{\alpha\mu}\eta_{\beta\nu} + \eta_{\alpha\nu}\eta_{\beta\mu}) - \frac{1}{3}\eta_{\alpha\beta}\eta_{\mu\nu} + (k\text{-dependent terms}) \right). \quad (2.10)$$

When the graviton is treated as a quantum field, the Feynman propagator is defined as in the massless theory Eq.A16,

$$\begin{aligned} D_{\mu\nu\alpha\beta}^{(m)}(x-y) &= \langle 0|T(\hat{h}_{\mu\nu}(x)\hat{h}_{\alpha\beta}(y))|0\rangle \\ &= \int \frac{d^4k}{(2\pi)^4} \frac{1}{-k^2 + m_g^2 + i\epsilon} e^{ik(x-y)} \sum_\lambda \epsilon_{\mu\nu}^\lambda(k) \epsilon_{\alpha\beta}^{*\lambda}(k). \end{aligned} \quad (2.11)$$

Comparing Eq.2.10 and Eq.2.11 we see that the polarisation sum for the FP massive gravity theory can be written as

$$\sum_\lambda \epsilon_{\mu\nu}^\lambda(k) \epsilon_{\alpha\beta}^{*\lambda}(k) = \frac{1}{2}(\eta_{\mu\alpha}\eta_{\nu\beta} + \eta_{\nu\alpha}\eta_{\mu\beta}) - \frac{1}{3}\eta_{\alpha\beta}\eta_{\mu\nu} + (k\text{-dependent terms}). \quad (2.12)$$

In processes where there is graviton emission from an external leg as in the case of gravitational wave radiation from an classical current, the amplitude square will have the form

$$|\mathcal{M}|^2 = \left(\frac{\kappa^2}{4}\right) \sum_\lambda |\epsilon_{\mu\nu}^\lambda(k) T^{\mu\nu}(k')|^2 = \left(\frac{\kappa^2}{4}\right) \sum_\lambda \epsilon_{\mu\nu}^\lambda(k) \epsilon_{\alpha\beta}^{*\lambda}(k) T^{\mu\nu}(k') T^{*\alpha\beta}(k'). \quad (2.13)$$

Since $T^{\mu\nu}$ is a conserved current, the momentum dependent pieces in the polarisation sum will give zero and we can drop them from Eq.2.12 for the calculations of diagrams with graviton emission from external legs as we will do in this paper.

We see that when the propagator Eq.A11 and polarisation sum Eq.A17 of the massless graviton theory is compared with the corresponding quantities Eq.2.6 and Eq.2.12, the massive theory differs from the massless theory even in the $m_g \rightarrow 0$ limit. There is an extra contribution of $(1/6)T^*T'$ to the amplitude Eq.2.9 in the FP theory. This is the contribution of the scalar degree of freedom of $g_{\mu\nu}$ which does not decouple in the $m_g \rightarrow 0$ limit.

Consider the Newtonian potential between two massive bodies. The amplitude for the diagram with one graviton exchange is in GR is

$$\mathcal{A}_{GR} = \frac{\kappa^2}{4} T^{\mu\nu} D_{\mu\nu\alpha\beta}^{(0)}(k) T'^{\alpha\beta}. \quad (2.14)$$

The stress tensor for massive bodies at rest in a given reference frame is of the form $T^{\mu\nu} = (M_1, 0, 0, 0)$ and $T'^{\alpha\beta} = (M_2, 0, 0, 0)$ and the massless graviton propagator in GR is Eq.A11. The potential derived from Eq.2.14 is the usual Newtonian form

$$\begin{aligned} V_{GR} &= \frac{\kappa^2}{4} \int \frac{d^3k}{(2\pi)^3} e^{ik \cdot r} \frac{1}{-k^2} \left(T_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} T^\alpha_\alpha \right) T'^{\mu\nu} \\ &= \frac{GM_1 M_2}{r}. \end{aligned} \quad (2.15)$$

On the other hand in the Fierz-Pauli theory the one graviton exchange amplitude Eq.2.9 is

$$\mathcal{A}_{FP} = \frac{\kappa^2}{4} \frac{1}{-k^2 + m_g^2} \left(T_{\mu\nu} - \frac{1}{3} \eta_{\mu\nu} T^\alpha_\alpha \right) T'^{\mu\nu}, \quad (2.16)$$

and the gravitational potential between two massive bodies in the FP theory is

$$\begin{aligned} V_{FP} &= \frac{\kappa^2}{4} \int \frac{d^3k}{(2\pi)^3} e^{ik \cdot r} \frac{1}{-k^2 + m_g^2} \left(T_{\mu\nu} - \frac{1}{3} \eta_{\mu\nu} T^\alpha_\alpha \right) T'^{\mu\nu} \\ &= \left(\frac{4}{3} \right) \frac{GM_1 M_2}{r} e^{-m_g r}. \end{aligned} \quad (2.17)$$

The FP theory gives a Yukawa potential as expected however in the $m_g \rightarrow 0$ limit the gravitational potential between massive bodies in the FP theory is a factor $4/3$ larger than the Newtonian potential arising from GR. This is ruled out from solar system tests of gravity [40] even in the $m_g \rightarrow 0$ limit. We note here that the bending of light by massive bodies is unaffected (in $m_g \rightarrow 0$ limit) as the stress tensor for photons $T^\mu_\nu = (\omega, 0, 0, -\omega)$ is traceless and the scattering amplitudes $\mathcal{A}_{FP}(m_g \rightarrow 0) = \mathcal{A}_{GR}$. Experimental observations [41] of the bending of radio waves by the Sun matches GR to 1%. The two observations together imply

that the extra factor of $(4/3)$ in the Newtonian potential of FP theory cannot be absorbed by redefining G .

The fact that the action of the FP theory 2.1 goes to the Einstein-Hilbert action Eq.A3 in the $m_g \rightarrow 0$ limit while the propagator Eq.2.10 does not go to the massless form Eq.A11, is what is called the vDVZ discontinuity pointed out by van Dam and Veltman [3] and Zakharov [4].

It has been pointed out by Vainshtein [42, 43] that the linear FP theory breaks down at distances much larger than the Schwarzschild radius $R_s = 2GM$ below which the linearised GR is no longer valid as $(\kappa h_{\mu\nu} \sim$ below this distance). The scalar mode in FP theory becomes strongly coupled with decreasing m_g and the minimum radius from a massive body at which the linearised FP theory is valid is called the Vainshtein radius and is given by $r_V = (R_s/m_g^4)^{1/5}$. We will discuss the Vainshtein radius of different theories of gravity discussed in this paper and how this consideration limits the bounds on m_g from binary systems derived in this paper in Section V A.

A. Graviton radiation from binaries in Fierz-Pauli theory

We consider the graviton radiation from the compact binary systems classically. The pictorial representation of graviton emission from a classical source is shown in Fig.1. The classical graviton current $T^{\mu\nu}$ is determined from the Kepler's orbit and the interaction vertex is $\frac{1}{2}\kappa h_{\mu\nu}T^{\mu\nu}$, where $h_{\mu\nu}$ is the graviton field and $\kappa = \sqrt{32\pi G}$. G is the Newton's universal gravitational constant. Here we use linearized gravity formulation with an extension of non zero graviton mass term Eq.2.1 to calculate the energy loss of a compact binary system due to graviton emission.

The emission rate of graviton from the interaction Lagrangian between the gravity and source $(\frac{1}{2}\kappa h_{\mu\nu}T^{\mu\nu})$ is given by

$$d\Gamma = \frac{\kappa^2}{4} \sum_{\lambda} |T_{\mu\nu}(k')\epsilon^{\mu\nu}(k)|^2 2\pi\delta(\omega - \omega') \frac{d^3k}{(2\pi)^3} \frac{1}{2\omega}, \quad (2.18)$$

where $T_{\mu\nu}(k')$ is the classical graviton current in the momentum space. Expanding the

modulus squared in Eq.2.18, we can write

$$d\Gamma = \frac{\kappa^2}{8(2\pi)^2} \sum_{\lambda} \left(T_{\mu\nu}(k') T_{\alpha\beta}^*(k') \epsilon^{\mu\nu}(k) \epsilon^{*\alpha\beta}(k) \right) \frac{d^3k}{\omega} \delta(\omega - \omega'). \quad (2.19)$$

Using the polarization sum of the Fierz-Pauli theory 2.12 this expression becomes

$$d\Gamma = \frac{\kappa^2}{8(2\pi)^2} \int \left[T_{\mu\nu}(k') T_{\alpha\beta}^*(k') \right] \left[\frac{1}{2} (\eta^{\mu\alpha} \eta^{\nu\beta} + \eta^{\mu\beta} \eta^{\nu\alpha} - \eta^{\mu\nu} \eta^{\alpha\beta}) + \frac{1}{6} \eta^{\mu\nu} \eta^{\alpha\beta} \right] \frac{d^3k}{\omega} \eta(\omega - \omega'). \quad (2.20)$$

The extra $(1/6)\eta^{\mu\nu}\eta^{\alpha\beta}$ term compared to the massless graviton case is the contribution of the scalar mode in FP theory. Simplifying we obtain

$$d\Gamma = \frac{\kappa^2}{8(2\pi)^2} \int \left[|T_{\mu\nu}(k')|^2 - \frac{1}{3} |T^{\mu}_{\mu}(k')|^2 \right] \delta(\omega - \omega') \omega \left(1 - \frac{m_g^2}{\omega^2} \right)^{\frac{1}{2}} d\omega d\Omega_k, \quad (2.21)$$

where we have used $d^3k = k^2 dk d\Omega$ and the dispersion relation $k^2 = (\omega^2 - m_g^2)$. From the emission rate we can calculate the rate of energy loss due to massive graviton emission which is

$$\frac{dE}{dt} = \frac{\kappa^2}{8(2\pi)^2} \int \left[|T_{\mu\nu}(k')|^2 - \frac{1}{3} |T^{\mu}_{\mu}(k')|^2 \right] \delta(\omega - \omega') \omega^2 \left(1 - \frac{m_g^2}{\omega^2} \right)^{\frac{1}{2}} d\omega d\Omega_k. \quad (2.22)$$

For the massive graviton, the dispersion relation is

$$|\mathbf{k}|^2 = \omega^2 \left(1 - \frac{m_g^2}{\omega^2} \right). \quad (2.23)$$

Hence, the unit vector along the momentum direction of graviton is $\hat{k}^i = \frac{k^i}{\omega \sqrt{1 - \frac{m_g^2}{\omega^2}}}$. Using the relation $k_{\mu} T^{\mu\nu} = 0$ and Eq.2.23, we can write the T_{00} and T_{i0} components of the stress tensor in terms of T_{ij} as follows

$$T_{0j} = -\sqrt{1 - \frac{m_g^2}{\omega^2}} \hat{k}^i T_{ij}, \quad T_{00} = \left(1 - \frac{m_g^2}{\omega^2} \right) \hat{k}^i \hat{k}^j T_{ij}. \quad (2.24)$$

Hence, we can write

$$\left[|T_{\mu\nu}(k')|^2 - \frac{1}{3} |T^{\mu}_{\mu}(k')|^2 \right] \equiv \Lambda_{ij,lm} T^{ij*} T^{lm}, \quad (2.25)$$

where,

$$\Lambda_{ij,lm} = \left[\delta_{il} \delta_{jm} - 2 \left(1 - \frac{m_g^2}{\omega^2} \right) \hat{k}_j \hat{k}_m \delta_{il} + \frac{2}{3} \left(1 - \frac{m_g^2}{\omega^2} \right)^2 \hat{k}_i \hat{k}_j \hat{k}_l \hat{k}_m - \frac{1}{3} \delta_{ij} \delta_{lm} + \frac{1}{3} \left(1 - \frac{m_g^2}{\omega^2} \right) \left(\delta_{ij} \hat{k}_l \hat{k}_m + \delta_{lm} \hat{k}_i \hat{k}_j \right) \right]. \quad (2.26)$$

Therefore, we can write Eq.2.22 as

$$\frac{dE}{dt} = \frac{\kappa^2}{8(2\pi)^2} \int \Lambda_{ij,lm} T^{ij*} T^{lm} \delta(\omega - \omega') \omega^2 \left(1 - \frac{m_g^2}{\omega^2}\right)^{\frac{1}{2}} d\omega d\Omega_k. \quad (2.27)$$

We can do the angular integrals using the relations A26 and obtain,

$$\begin{aligned} \int d\Omega_k \Lambda_{ij,lm} T^{ij*}(\omega') T^{lm}(\omega') &= \frac{8\pi}{5} \left(\left[\frac{5}{2} - \frac{5}{3} \left(1 - \frac{m_g^2}{\omega'^2}\right) + \frac{2}{9} \left(1 - \frac{m_g^2}{\omega'^2}\right)^2 \right] T^{ij} T^{*ij} \right. \\ &\quad \left. + \left[-\frac{5}{6} + \frac{5}{9} \left(1 - \frac{m_g^2}{\omega'^2}\right) + \frac{1}{9} \left(1 - \frac{m_g^2}{\omega'^2}\right)^2 \right] |T^i_i|^2 \right), \end{aligned} \quad (2.28)$$

Hence, the rate of energy loss becomes

$$\begin{aligned} \frac{dE}{dt} &= \frac{8G}{5} \int \left[\left\{ \frac{5}{2} - \frac{5}{3} \left(1 - \frac{m_g^2}{\omega'^2}\right) + \frac{2}{9} \left(1 - \frac{m_g^2}{\omega'^2}\right)^2 \right\} T^{ij} T^{*ij} \right. \\ &\quad \left. + \left\{ -\frac{5}{6} + \frac{5}{9} \left(1 - \frac{m_g^2}{\omega'^2}\right) + \frac{1}{9} \left(1 - \frac{m_g^2}{\omega'^2}\right)^2 \right\} |T^i_i|^2 \right] \delta(\omega - \omega') \omega^2 \left(1 - \frac{m_g^2}{\omega^2}\right)^{\frac{1}{2}} d\omega. \end{aligned} \quad (2.29)$$

In the massless gravity theory the prefactors of $T^{ij} T^{*ij}$ and $|T^i_i|^2$ are 1 and $-1/3$ respectively. Note that the $m_g \rightarrow 0$ limit of Eq.2.29 gives different prefactors. In the massive graviton limit, all the five polarization components contribute to the energy loss instead of two as in the massless limit. Therefore, from Eq.2.29, we will not obtain the energy loss for massless limit by simply putting $m_g \rightarrow 0$. In AppendixA we obtain the energy loss due to massless graviton radiation from compact binary systems.

From Eq.A44 we get

$$\left[T_{ij}(\omega') T_{ji}^*(\omega') - \frac{1}{3} |T^i_i(\omega')|^2 \right] = 4\mu^2 \omega'^4 a^4 f(n, e). \quad (2.30)$$

where $n_0 = \frac{m_g}{\Omega}$, and

$$\begin{aligned} f(n, e) &= \frac{1}{32n^2} \left\{ [J_{n-2}(ne) - 2eJ_{n-1}(ne) + 2eJ_{n+1}(ne) + \frac{2}{n} J_n(ne) - J_{n+2}(ne)]^2 + \right. \\ &\quad \left. (1 - e^2) [J_{n-2}(ne) - 2J_n(ne) + J_{n+2}(ne)]^2 + \frac{4}{3n^2} J_n^2(ne) \right\}. \end{aligned} \quad (2.31)$$

The final expression of dE/dt for massive theory can be written in the compact form as

$$\frac{dE}{dt} = \frac{32G}{5}\mu^2 a^4 \Omega^6 \sum_{n=1}^{\infty} n^6 \sqrt{1 - \frac{n_0^2}{n^2}} \left[f(n, e) \left(\frac{19}{18} + \frac{11}{9} \frac{n_0^2}{n^2} + \frac{2}{9} \frac{n_0^4}{n^4} \right) + \frac{5J_n^2(ne)}{108n^4} \left(1 - \frac{n_0^2}{n^2} \right)^2 \right]. \quad (2.32)$$

We can split Eq.2.32 as

$$\begin{aligned} \frac{dE}{dt} = & \frac{32G}{5}\mu^2 a^4 \Omega^6 \sum_{n=1}^{\infty} n^6 \sqrt{1 - \frac{n_0^2}{n^2}} \left[f(n, e) \left(1 + \frac{4}{3} \frac{n_0^2}{n^2} + \frac{1}{6} \frac{n_0^4}{n^4} \right) - \frac{5J_n^2(ne)}{36n^4} \frac{n_0^2}{n^2} \left(1 - \frac{n_0^2}{4n^2} \right) \right] + \\ & \frac{32G}{5}\mu^2 a^4 \Omega^6 \sum_{n=1}^{\infty} n^6 \sqrt{1 - \frac{n_0^2}{n^2}} \left[\frac{1}{18} f(n, e) \left(1 - \frac{n_0^2}{n^2} \right)^2 + \frac{5J_n^2(ne)}{108n^4} \left(1 + \frac{n_0^2}{2n^2} \right)^2 \right], \end{aligned} \quad (2.33)$$

where the first term in Eq.2.33 denotes the energy loss in the massive gravity theory without vDVZ discontinuity (Eq.3.16) and the second term denotes the contribution due to the scalar mode associated with $\frac{1}{6}\eta_{\mu\nu}\eta_{\alpha\beta}$.

The rate of energy loss in the Keplerian orbit leads to the decrease in orbital period decay at a rate

$$\dot{P}_b = -6\pi G^{-\frac{3}{2}}(m_1 m_2)^{-1}(m_1 + m_2)^{-\frac{1}{2}} a^{\frac{5}{2}} \left(\frac{dE}{dt} \right). \quad (2.34)$$

The energy loss or the power radiated from the binary system increases with increasing the eccentricity as it is clear from Fig.2, since the energy loss in the first term is proportional to $n^6 f(n, e)$. The radiation is dominated by the higher harmonics for $e \approx 1$. The radiation has a peak at some particular value of n for a given eccentric orbit.

III. MASSIVE GRAVITY WITHOUT VDVZ DISCONTINUITY

In the Fierz-Pauli theory Eq.2.1 there is no ghost owing to the fact that the relative coefficients of the h^2 and $h_{\mu\nu}h^{\mu\nu}$ terms is chosen as -1 . Generalising the theory beyond this point will lead to the appearance of ghosts. There is a special choice of coefficient where the ghost term cancels the extra scalar contribution to the propagator. In this theory therefore there is no vDVZ discontinuity and there are no ghosts [30, 31]. Phenomenologically this theory has the simple generalisation of the spin-2 graviton with 2 polarizations which




FIG. 2: Variation of $n^6 f(n, e)$ with n .

obey the dispersion relation $k_0^2 = |\vec{k}|^2 + m_g^2$. Consider the one parameter generalisation of the Fierz-Pauli theory

$$S = \int d^4x \left[\frac{1}{2} h_{\mu\nu} \mathcal{E}^{\mu\nu\alpha\beta} h_{\alpha\beta} + \frac{1}{2} m_g^2 h_{\mu\nu} (\eta^{\mu(\alpha} \eta^{\beta)\nu} - (1-a) \eta^{\mu\nu} \eta^{\alpha\beta}) h_{\alpha\beta} + \frac{\kappa}{2} h_{\mu\nu} T^{\mu\nu} \right], \quad (3.1)$$

where $a = 0$ corresponds to the Fierz-Pauli theory Eq.2.1. We will derive the equations assuming $a \neq 0$ and see which values of a can solve the problem of vDVZ discontinuity which is generic in massive gravity theories.

The e.o.m from Eq.3.1 are

$$(\square + m_g^2) h_{\mu\nu} - \eta_{\mu\nu} (\square + m_g^2(1-a)) h - \partial_\mu \partial^\alpha h_{\alpha\nu} - \partial_\nu \partial^\alpha h_{\alpha\mu} + \eta_{\mu\nu} \partial^\alpha \partial^\beta h_{\alpha\beta} + \partial_\mu \partial_\nu h = -\kappa T_{\mu\nu}. \quad (3.2)$$

Taking the divergence of Eq.3.2 we have

$$m_g^2 (\partial^\mu h_{\mu\nu} - (1-a) \partial_\nu h) = 0. \quad (3.3)$$

These are 4 constraint equations which reduce the independent degrees of freedom of the graviton from 10 to 6.

Using Eq.3.3 in Eq.3.2 we obtain

$$(\square + m_g^2)h_{\mu\nu} - a\eta_{\mu\nu}\square h - (1 - 2a)\partial_\mu\partial_\nu h - m_g^2\eta_{\mu\nu}(1 - a)h = -\kappa T_{\mu\nu}. \quad (3.4)$$

Taking the trace of this equation we obtain

$$-2a\square h - (3m_g^2 - 4m_g^2a)h = -\kappa T. \quad (3.5)$$

We see that the h is now a propagating field if $a \neq 0$. The kinetic term for h appears with a minus sign so h is a ghost field. The homogenous equation for h can be written as

$$\square h - m_h^2 h = 0 \quad (3.6)$$

with the ghost mass given by

$$m_h^2 = \frac{m_g^2}{2} \left(1 + 3 \left(1 - \frac{1}{a} \right) \right). \quad (3.7)$$

The propagator of the deformed Fierz-Pauli theory Eq.3.1 is given

$$[\mathcal{E}^{\mu\nu\alpha\beta} + m_g^2 (\eta^{\mu(\alpha}\eta^{\beta)\nu} - \eta^{\mu\nu}\eta^{\alpha\beta}(1 - a))] D_{\alpha\beta\rho\sigma}^{(a)}(x - y) = \delta_{(\rho}^{\mu} \delta_{\sigma)}^{\nu} \delta^4(x - y). \quad (3.8)$$

This equation can be inverted to give the $D_{\alpha\beta\rho\sigma}^{(a)}$ which in momentum space turns out of the form

$$D_{\alpha\beta\mu\nu}^{(a)}(k) = \frac{1}{-k^2 + m_g^2} \left(\frac{1}{2}(\eta_{\alpha\mu}\eta_{\beta\nu} + \eta_{\alpha\nu}\eta_{\beta\mu}) - \frac{1}{3}\eta_{\alpha\beta}\eta_{\mu\nu} \right) + \frac{i}{k^2 + m_h^2} \left(\frac{1}{6}\eta_{\alpha\beta}\eta_{\mu\nu} \right) + (k\text{-dependent terms}). \quad (3.9)$$

This shows that there are two types of contributions to the propagator, 2 helicity states of spin-2 massive gravitons and a massive scalar with mass m_h . This part is identical to the propagator of the Fierz-Pauli theory. In 3.9 there is an additional contribution from the ghost mode with the kinetic operator k^2 with the wrong sign and mass m_h given in Eq.3.7. The remaining 3 vector degrees of freedom don't couple to the energy momentum tensor and we ignore their contribution here. Now if we choose the parameter $a = 1/2$, the mass of the ghost mode Eq.3.7 becomes $m_h^2 = -m_g^2$. The ghost mode for $a = 1/2$ becomes tachyonic. Substituting $m_h^2 = -m_g^2$ in Eq.3.9 we see that the propagator simplifies to the form

$$D_{\alpha\beta\mu\nu}^{(1/2)}(k) = \frac{1}{-k^2 + m_g^2} \left(\frac{1}{2}(\eta_{\alpha\mu}\eta_{\beta\nu} + \eta_{\alpha\nu}\eta_{\beta\mu}) - \frac{1}{2}\eta_{\alpha\beta}\eta_{\mu\nu} \right) + (k\text{-dependent terms}). \quad (3.10)$$

The ghost term with tachyonic mass cancels the extra scalar contribution to the propagator and we are left with the tensor structure of the propagator which is the same as for the massless gravitons Eq.A11 but which have the dispersion relations of massive gravitons, $k_0^2 = |\vec{k}|^2 + m_g^2$. In the limit $m_g \rightarrow 0$ the propagator goes to the massless propagator form Eq.A11 and thus there is no vDVZ discontinuity. From the tensor structure of Eq.3.10 it is clear that for even for $m_g \neq 0$ the polarisation sum takes the same form Eq.A17 as that of the massless theory.

The gravitational potential in this theory takes the Yukawa form

$$V^{(1/2)}(r) = \frac{GM_1M_2}{r} e^{-m_g r}, \quad (3.11)$$

and the extra factor of $(4/3)$ which was there in the FP theory Eq.2.17 is absent due to cancellation of the scalar graviton mode with the ghost contribution in the propagator. The Yukawa corrections to the $1/r$ potential will give rise to a perihelion precession in planetary orbits [44]. Constraints on the Yukawa potential between planets and the sun which give bounds in the mass of the exchanged particle have been obtained in [44]. The long range Yukawa potential caused by axions can also affect the gravitational light bending and Shapiro time delay which is discussed in [45].

This theory which avoids contributions from the extra scalars mode is phenomenologically the most acceptable. The classical calculation of energy loss from binaries in this spin-2 massive gravity theory was done by Finn and Sutton [29]. Our calculation which we present now is the QFT version of this calculation. We find that the result of our tree level QFT calculation agrees in the leading order with the result of [29].

From the direct detection of gravitational waves by Virgo and Ligo [46], mass of the spin-2 graviton is $m_g < 1.2 \times 10^{-22}$ eV which is derived from the experimental upper bound on the dispersion of the gravitational wave event GW150914.

A. Graviton radiation in massive gravity without vDVZ discontinuity

Following the steps described in the Appendix A, we compute the rate of energy loss due to the massive graviton radiation as

$$\frac{dE}{dt} = \frac{\kappa^2}{8(2\pi)^2} \int \left[|T_{\mu\nu}(k')|^2 - \frac{1}{2} |T^\mu{}_\mu(k')|^2 \right] \delta(\omega - \omega') \omega^2 \left(1 - \frac{m_g^2}{\omega^2}\right)^{\frac{1}{2}} d\omega d\Omega_k \quad (3.12)$$

$$= \frac{\kappa^2}{8(2\pi)^2} \int \tilde{\Lambda}_{ij,lm} T^{ij*} T^{lm} \delta(\omega - \omega') \omega^2 \left(1 - \frac{m_g^2}{\omega^2}\right)^{\frac{1}{2}} d\omega d\Omega_k, \quad (3.13)$$

where

$$\begin{aligned} \tilde{\Lambda}_{ij,lm} = & \left[\delta_{il} \delta_{jm} - 2 \left(1 - \frac{m_g^2}{\omega^2}\right) \hat{k}_j \hat{k}_m \delta_{il} + \frac{1}{2} \left(1 - \frac{m_g^2}{\omega^2}\right)^2 \hat{k}_i \hat{k}_j \hat{k}_l \hat{k}_m - \frac{1}{2} \delta_{ij} \delta_{lm} \right. \\ & \left. + \frac{1}{2} \left(1 - \frac{m_g^2}{\omega^2}\right) (\delta_{ij} \hat{k}_l \hat{k}_m + \delta_{lm} \hat{k}_i \hat{k}_j) \right]. \end{aligned} \quad (3.14)$$

After computation of the angular integration we obtain

$$\begin{aligned} \frac{dE}{dt} = & \frac{8G}{5} \int \left[\left\{ \frac{5}{2} - \frac{5}{3} \left(1 - \frac{m_g^2}{\omega'^2}\right) + \frac{1}{6} \left(1 - \frac{m_g^2}{\omega'^2}\right)^2 \right\} T^{ij} T_{ij}^* \right. \\ & \left. + \left\{ -\frac{5}{4} + \frac{5}{6} \left(1 - \frac{m_g^2}{\omega'^2}\right) + \frac{1}{12} \left(1 - \frac{m_g^2}{\omega'^2}\right)^2 \right\} |T^i{}_i|^2 \right] \delta(\omega - \omega') \omega^2 \left(1 - \frac{m_g^2}{\omega^2}\right)^{\frac{1}{2}} d\omega. \end{aligned} \quad (3.15)$$

Finally we get the expression for the rate of energy loss as

$$\frac{dE}{dt} = \frac{32G}{5} \mu^2 a^4 \Omega^6 \sum_{n=1}^{\infty} n^6 \sqrt{1 - \frac{n_0^2}{n^2}} \left[f(n, e) \left(1 + \frac{4n_0^2}{3n^2} + \frac{1n_0^4}{6n^4}\right) - \frac{5J_n^2(ne)}{36n^4} \frac{n_0^2}{n^2} \left(1 - \frac{n_0^2}{4n^2}\right) \right]. \quad (3.16)$$

We note that the expression reduces to that of GR in the limit $n_0 = 0$. Thus there is no vDVZ discontinuity. To the leading order in n_0^2 this agrees with the result of the classical calculation of Finn and Sutton [29].

IV. DVALI-GABADADZE-PORRATI (DGP) THEORY

The GR theory is a non linear theory which obeys diffeomorphism invariance. However this symmetry is broken in theories with a massive graviton. In FP theory, if the graviton

is expanded around curved spacetime a ghost degree of freedom appears [47]. To obtain a consistent massive gravity theory free from any ghost, one can go to higher dimension. One such massive gravity theory in higher dimension using a braneworld model framework is the DGP theory [32–34, 48]. In the higher dimensions the massless gravity theory has a general covariance symmetry. The number of polarisation states of the spin-2 massless graviton in 5-dimensions is (5). When the extra dimension compactifies the number of massive graviton degrees of freedom in 4-d remains 5 and there is no (Boulware Deser) BD ghost. The DGP theory in a cosmological background can account for the cosmological constant [49]. The mass of gravitons is momentum dependent so that one can modify the infrared theory (at cosmological scales) while retaining Newtonian theory at solar system scales. The scalar degree of freedom of the graviton however still contributes to the vDVZ discontinuity which remains a problem for phenomenology of the DGP theory of massive gravity [48].

In the five dimensional DGP theory, the matter field is localized in a four dimensional brane world which leads to an induced curvature term on the brane. The Planck scales of the five dimensional DGP theory with the four dimensional brane world are denoted by M_5 and M_{pl} respectively.

The action of five dimensional DGP model [32–34] with the matter field localized in four dimensional brane world at $y = 0$ is

$$\mathcal{S} \supset \int d^4x dy \left(\frac{M_5^3}{4} \sqrt{-^{(5)}g^{(5)}} R + \delta(y) \left[\sqrt{-g} \frac{M_{pl}^2}{2} R[g] + \mathcal{L}_m(g, \psi_i) \right] \right), \quad (4.1)$$

where ψ_i denotes the matter field with the energy stress tensor $T_{\mu\nu}$ in the brane world.

The resulting modified linearized Einstein equation on the $y = 0$ brane is [6]

$$(\square h_{\mu\nu} - \partial_\mu \partial_\nu h) - m_0 \sqrt{-\square} (h_{\mu\nu} - h \eta_{\mu\nu}) = -\frac{\kappa}{2} T_{\mu\nu}(x), \quad (4.2)$$

where $m_0 = \frac{M_5^3}{M_{pl}^2}$, $M_{pl}^2 = 1/8\pi G = 4/\kappa^2$. Here, the Fierz-Pauli mass term $(h_{\mu\nu} - h \eta_{\mu\nu})$ appears naturally from the higher dimensional DGP theory. This corresponds to the linearized massive gravity with a scale-dependent effective mass $m_g^2(\square) = m_0 \sqrt{-\square}$. The propagator is

$$D_{\alpha\beta\mu\nu}^{(5)}(k) = \frac{i}{(-\omega^2 + |\mathbf{k}|^2) + m_0(\omega^2 - |\mathbf{k}|^2)^{1/2}} \left(\frac{1}{2}(\eta_{\alpha\mu}\eta_{\beta\nu} + \eta_{\alpha\nu}\eta_{\beta\mu}) - \frac{1}{3}\eta_{\alpha\beta}\eta_{\mu\nu} \right). \quad (4.3)$$

The terms in the brackets represent the polarization sum which is identical to that of the FP theory Eq.2.12. In the $m_0 \rightarrow 0$ limit the DGP propagator does not go to the massless form Eq.A11 and the DGP theory also has the vDVZ discontinuity.

The dispersion relation corresponding to real gravitons in the DGP model is given by the pole of the propagator Eq.4.3,

$$\omega^2 = |\mathbf{k}|^2 - m_0^2, \quad (4.4)$$

where $|\mathbf{k}|$ is the magnitude of the propagation vector. We note that in the DGP theory the graviton has a tachyonic mass.

Following the same steps of FP theory that we have done in the previous section, we write down energy loss due to massive graviton radiation in DGP theory. All the relevant expressions in DGP theory differ from those of the FP theory by replacing $m_g^2 \rightarrow -m_0^2$ and $\tilde{n}_0^2 = m_0^2/\Omega^2 = -n_0^2$, i.e.

$$\frac{dE}{dt} = \frac{\kappa^2}{8(2\pi)^2} \int \left[|T_{\mu\nu}(k')|^2 - \frac{1}{3} |T^\mu{}_\mu(k')|^2 \right] \delta(\omega - \omega') \omega^2 \left(1 + \frac{m_0^2}{\omega^2}\right)^{\frac{1}{2}} d\omega d\Omega_k. \quad (4.5)$$

The components of the stress tensor in $x - y$ plane is given in Eq.A44. The dispersion relation gives $\hat{k}^i = \frac{k^i}{\omega\sqrt{1+\frac{m_0^2}{\omega^2}}}$. The other components of $T_{\mu\nu}$ can be obtained by using the current conservation relation $k_\mu T^{\mu\nu} = 0$ which yields,

$$T_{0j} = -\sqrt{1 + \frac{m_0^2}{\omega^2}} \hat{k}^i T_{ij}, \quad T_{00} = \left(1 + \frac{m_0^2}{\omega^2}\right) \hat{k}^i \hat{k}^j T_{ij}. \quad (4.6)$$

Hence, the term in the third bracket of Eq.4.5 can be written in terms of the projection operator $\tilde{\Lambda}_{ij,lm}$ as

$$\left[|T_{\mu\nu}(k')|^2 - \frac{1}{3} |T^\mu{}_\mu(k')|^2 \right] = \tilde{\Lambda}_{ij,lm} T^{ij*} T^{lm}, \quad (4.7)$$

where

$$\tilde{\Lambda}_{ij,lm} = \left[\delta_{il}\delta_{jm} - 2\left(1 + \frac{m_0^2}{\omega^2}\right) \hat{k}_j \hat{k}_m \delta_{il} + \frac{2}{3}\left(1 + \frac{m_0^2}{\omega^2}\right)^2 \hat{k}_i \hat{k}_j \hat{k}_l \hat{k}_m - \frac{1}{3} \delta_{ij} \delta_{lm} + \frac{1}{3}\left(1 + \frac{m_0^2}{\omega^2}\right) (\delta_{ij} \hat{k}_l \hat{k}_m + \delta_{lm} \hat{k}_i \hat{k}_j) \right]. \quad (4.8)$$

The final expression of dE/dt for massive DGP theory can be written in the compact form as

$$\frac{dE}{dt} = \frac{32G}{5} \mu^2 a^4 \Omega^6 \sum_{n=1}^{\infty} n^6 \sqrt{1 + \frac{\tilde{n}_0^2}{n^2}} \left[f(n, e) \left(\frac{19}{18} - \frac{11}{9} \frac{\tilde{n}_0^2}{n^2} + \frac{2}{9} \frac{\tilde{n}_0^4}{n^4} \right) + \frac{5J_n^2(ne)}{108n^4} \left(1 + \frac{\tilde{n}_0^2}{n^2} \right)^2 \right]. \quad (4.9)$$

V. CONSTRAINTS FROM OBSERVATIONS

TABLE I: Summary of the measured orbital parameters and the orbital period derivative values from observation and GR for PSR B1913+16 [20] and PSR J1738+0333 [23]. The uncertainties in the last digits are quoted in the parenthesis.

Parameters	PSR B1913+16	PSR J1738+0333
Pulsar mass m_1 (solar masses)	1.438 ± 0.001	$1.46^{+0.06}_{-0.05}$
Companion mass m_2 (solar masses)	1.390 ± 0.001	$0.181^{+0.008}_{-0.007}$
Eccentricity e	$0.6171340(4)$	$(3.4 \pm 1.1) \times 10^{-7}$
Orbital period P_b (d)	$0.322997448918(3)$	$0.3547907398724(13)$
Intrinsic $\dot{P}_b(10^{-12}\text{ss}^{-1})$	-2.398 ± 0.004	$(-25.9 \pm 3.2) \times 10^{-3}$
GR $\dot{P}_b(10^{-12}\text{ss}^{-1})$	-2.40263 ± 0.00005	$-27.7^{+1.5}_{-1.9} \times 10^{-3}$

In this section we calculate the graviton mass from the observation of orbital period decay of the Hulse -Taylor binary system (PSR B1913+16) and a pulsar white-dwarf binary system (PSR J1738+0333). The orbital parameters of the two compact binary systems and the orbital period derivative values from observation and GR are given in Table I. Massless graviton has two states of polarization and the rate of energy loss of the compact binary system due to the emission of massless graviton radiation is given by Eq. (A48) and it agrees with the Peters Mathews formula [16]. The variation of $F(e)$ with the eccentricity is shown in FIG.3 which implies that the energy loss due to the GR value is largely enhanced by the eccentricity enhancement factor $F(e)$. Its value is always greater than one for non zero eccentric orbit. Large eccentric binary orbit has strong speed variation as it moves from periastron to apiastron which leads to produce a large amount of radiation in higher harmonics of orbital frequency. In the following, we compare three massive theories of gravity and find limits on the graviton mass for PSR B1913+16 and PSR J1738+0333.




FIG. 3: Variation of $F(e)$ with the eccentricity.

A. Vainshtein radius and limits of linear theory

We have used the leading order perturbation of the metric for calculating the graviton emission. In linearised Einstein's gravity the perturbation theory holds as long as $\kappa h_{\mu\nu} \ll 1$. This implies that perturbation theory breaks down at radius smaller than $R_s = 2GM$ of the source. In the Fierz-Pauli theory with non-zero graviton mass the scalar mode becomes strongly coupled as $m_g \rightarrow 0$. The smallest radius until which the perturbation theory can be applied is Vainshtein radius [42] given by

$$r_V = \left(\frac{R_s}{m_g^4} \right)^{1/5}. \quad (5.1)$$

The Vaishtein radius is much larger than the R_s and perturbative calculations of the Fierz-Pauli theory are valid in regions with $r > r_V$ away from the source. In our application of binary pulsar radiation, classically the gravitational field is evaluated at the radiation zone such that $r_V < \lambda$ (where $\lambda \sim \pi/\Omega$ is the wavelength of the gravitational waves radiated). In the FP theory this implies that we must have

$$\lambda \sim \pi\Omega^{-1} > r_V = \left(\frac{R_s}{m_g^4} \right)^{1/5}. \quad (5.2)$$

We therefore have a lower bound on the graviton mass above which the perturbative calculations is valid given by

$$m_g > \frac{\Omega^{5/4}}{\pi^{5/4}} (2GM)^{1/4} \quad (5.3)$$

Using the numbers as shown in Table.I for PSR B1913+16, we find that the region of m_g of the Fierz-Pauli theory where the perturbative calculation is valid is $m_g > 3.06 \times 10^{-22}$ eV. For PSR J1738+0333, we use the Vainshtein limit and obtain the region of graviton mass for the validity of the perturbative calculation $m_g > 2.456 \times 10^{-22}$ eV for FP theory

For the case of $a = 1/2$ massive gravity theory, the strongly coupled scalar is cancelled by the ghost mode and there is no Vainshtein limiting radius.

For the DGP theory the Vainshtein radius is given by [32, 43]

$$r_V = \left(\frac{R_g}{m_g^2} \right)^{1/3}. \quad (5.4)$$

Again we must have $\lambda \sim \pi\Omega^{-1} > r_V$ which gives a lower bound on the graviton mass in the DGP theory above which the perturbative calculation is valid, given by

$$m_g > \frac{\Omega^{3/2}}{\pi^{3/2}} (2GM)^{1/2}. \quad (5.5)$$

This number is 7.84×10^{-24} eV for PSR B1913+16 and 1.406×10^{-24} eV for PSR J1738+0333.

B. Constraints from observation for FP Theory

The massive graviton has five states of polarization and of these the scalar and the tensor modes couple to the energy momentum tensor and contribute to the energy loss for the compact binary systems. In the massless limit $m_g \rightarrow 0$ of the FP theory, the extra scalar mode does not decouple and one encounters vDVZ discontinuity. In FIG.4(a) and FIG.4(b), we show the variation of the orbital period loss with the graviton mass for PSR B1913+16 and In FIG.4(c) and FIG.4(d) we obtain the same variation for PSR J1738+0333. The dotted lines denote the corresponding Vainshtein limit for the two binary system.

In the region $m_g \sim \Omega$ the energy loss falls with increasing m_g as the phase of graviton momentum shrinks. There is a region where the theoretical curve goes through the observational band as shown FIG.4(b) and FIG.4(d) where the variation of orbital period derivative is shown with the observational uncertainty for the two compact binary systems.

This range value which corresponds to the graviton mass $m_g \in (6.88 - 6.96) \times 10^{-19}$ eV (FIG.4(b)) for PSR B1913+16 and $m_g \in (2.31 - 2.48) \times 10^{-19}$ eV for PSR J1738+0333. There




FIG. 4: In the upper panel, we have plotted the orbital period decay as a function of mass of the graviton for (a) FP theory and (b) FP theory from observations for PSR B1913+16. In the lower panel of (c) and (d) we have shown the same variation for PSR J1738+0333

is no common mass range in the overlap region for the two binary systems for any value of m_g .

For the FP theory therefore, the limit on graviton mass from observations PSR B1913+16 together with PSR J1738+0333 comes from the Vainstein limit $m_g < 3.06 \times 10^{-22}$ eV.

C. Constraints from observation for DGP Theory

In DGP theory, the massless limit $m_0 \rightarrow 0$ of the DGP theory does not simply give the massless result and here also one encounters vDVZ discontinuity due to the extra contribution of the scalar gravitons. In FIG.5(a) and FIG.5(b), we show the variation of the orbital period




FIG. 5: In the upper panel, we have plotted the orbital period decay as a function of mass parameter of the graviton for (a) DGP theory and (b) DGP theory from observation for PSR B1913+16. In the lower panel of (c) and (d) we have shown the same variation for PSR J1738+0333

loss with m_0 for PSR B1913+16 and in FIG.5(c) and FIG.5(d) we obtain the same for PSR J1738+0333. The dotted lines denote the corresponding Vainshtein limit for the two binary system which are $m_g > 7.84 \times 10^{-24}$ eV for PSR B1913+16 and $m_g > 1.406 \times 10^{-24}$ eV for PSR J1738+0333.

As in FP theory, in the DGP theory also there is some region where the theoretical prediction crosses the observed band value which corresponds to the graviton mass $m_0 \in (2.45 - 2.47) \times 10^{-19}$ eV (FIG.4(b)) for PSR B1913+16 and $m_0 \in (0.31 - 1.41) \times 10^{-19}$ eV for PSR J1738+0333. Since, for DGP theory also, there is no common mass range in the overlap region for the two binary systems, it implies any graviton mass value in the region

above the Vainshtein limit is ruled out and the experimental bound from observations is $m_0 < 7.84 \times 10^{-24} \text{eV}$.

D. No vDVZ discontinuity theory

In massive gravity theory without vDVZ discontinuity, where the scalar mode is cancelled by the ghost mode and there is no Vainshtein limiting radius which applies to our calculation.

In Fig.6 we have shown the variation of orbital period loss with the graviton mass for the two compact binary systems and in the low graviton mass limit, the orbital period loss for this theory and massless theory become degenerate.



There exist two regions where the theoretical prediction agrees with the observational band. For PSR B1913+16 this corresponds to the graviton mass $m_g \in (6.32 - 6.50) \times 10^{-19} \text{eV}$ and $m_g < 1.81 \times 10^{-20} \text{eV}$ (FIG.6(b)). For PSR J1738+0333, the corresponding graviton mass range are $m_g \in (2.18 - 2.34) \times 10^{-19} \text{eV}$ and $m_g < 5.29 \times 10^{-20} \text{eV}$. Here for the two binary systems we find common graviton mass region where there is agreement with both observations and the bound on graviton mass is $m_g < 1.81 \times 10^{-20} \text{eV}$.

All the bounds derived in the paper are at 68% C.L.



VI. CONCLUSIONS

In this paper we put constraints on three massive gravity theories from binary pulsar observations. We show that the bounds on gravitational mass from binary observations are highly model dependent as the predictions for the gravitational luminosity for different graviton mass models have significant differences.


In massive gravity theories like FP and DGP with an extra propagating scalar, the contribution of the extra scalar to the energy loss is of the same order as that of the tensor gravitational waves and the region $m_g < \Omega$ is ruled out from binary pulsar observations. As the graviton mass approaches and becomes larger than Ω the energy radiated drops with increasing mass. For each binary system there is therefore a range of graviton mass where the theoretical predictions are within observational limits. We found that the allowed



(a) Massive gravity without vDVZ discontinuity (b) massive gravity without vDVZ discontinuity from observation



(c) Massive gravity without vDVZ discontinuity from observation (d) Massive gravity without vDVZ discontinuity from observation



(e) massive gravity without vDVZ discontinuity from observation

FIG. 6: We have plotted the orbital period decay as a function of mass parameter of the graviton for (a) massive gravity theory without vDVZ discontinuity and (b) massive gravity theory without vDVZ discontinuity from observation for PSR B1913+16 for higher graviton mass and (c) for smaller graviton mass . In (d) and (e) we have shown the same variation for PSR J1738+0333

ranges of graviton mass from PSR B1913+16 and PSR PSR J1738+0333 do not have any overlap. Therefore combining observations from the two pulsars we see that no range of graviton mass is consistent with both pulsar observations. In these theories the linear order calculation breaks down below a Vainshtein radius.

To summarise, observations from PSR B1913+16 and PSR J1738+0333 rule out all value of graviton mass and from the Vainshtein limit we can put the upper bounds $m_g < 3.06 \times 10^{-22}$ eV for the FP theory and $m_0 < 1.406 \times 10^{-24}$ eV for the DGP theory. For the No-vDVZ discontinuity theory the upper bound from combined PSR B1913+16 and PSR J1738+0333 data is $m_g < 1.81 \times 10^{-20}$ eV. All bounds quoted in the paper are with 68% C.L.

There are other massive gravity theories like Lorentz violating gravitational mass [50–52] and more general Lorentz violating graviton bilinear terms [53, 54] which we have not covered in the Lorentz covariant calculation in this paper. We will address these theories in a separate publication.

The diagrammatic method can also be used for computing the wave-form of gravitational waves observed in direct detection experiments like LIGO and VIRGO [55, 56]. It will be interesting to test massive gravity theory predictions [36, 37] with direct observations and in particular to constrain the scalar and vector modes of gravity from direct detection [46].

Appendix A: ENERGY LOSS BY MASSLESS GRAVITON RADIATION FROM BINARIES

The action for the graviton field $h_{\mu\nu}$ is obtained by starting with the Einstein-Hilbert action for gravity and matter fields

$$S_{EH} = \int d^4x \sqrt{-g} \left[-\frac{1}{16\pi G} R + \mathcal{L}_m \right], \quad (\text{A1})$$

and expanding the metric $g_{\mu\nu} = \eta_{\mu\nu} + \kappa h_{\mu\nu}$ to the linear order in $h_{\mu\nu}$, where $\kappa = \sqrt{32\pi G}$ is the gravitational coupling. For consistency the inverse metric $g^{\mu\nu}$ and square root of

determinant $\sqrt{-g}$ should be expanded to quadratic order

$$\begin{aligned}
g_{\mu\nu} &= \eta_{\mu\nu} + \kappa h_{\mu\nu}, \\
g^{\mu\nu} &= \eta^{\mu\nu} - \kappa h^{\mu\nu} + \kappa^2 h^{\mu\lambda} h_{\lambda}^{\nu} + \mathcal{O}(\kappa^3), \\
\sqrt{-g} &= 1 + \frac{\kappa}{2} h + \frac{\kappa^2}{8} h^2 - \frac{\kappa^2}{4} h^{\mu\nu} h_{\mu\nu} + \mathcal{O}(\kappa^3),
\end{aligned} \tag{A2}$$

where $\eta_{\mu\nu} = \text{diagonal}(1, -1, -1, -1)$ is the background Minkowski metric and $h = h_{\mu}^{\mu}$. Indices are raised and lowered by $\eta^{\mu\nu}$ and $\eta_{\mu\nu}$.

The linearised Einstein-Hilbert action for the graviton field $h_{\mu\nu}$ is then given by

$$\begin{aligned}
S_{EH} &= \int d^4x \left[-\frac{1}{2} (\partial_{\mu} h_{\nu\rho})^2 + \frac{1}{2} (\partial_{\mu} h)^2 - (\partial_{\mu} h) (\partial^{\nu} h_{\nu}^{\mu}) + (\partial_{\mu} h_{\nu\rho}) (\partial^{\nu} h^{\mu\rho}) + \frac{\kappa}{2} h_{\mu\nu} T^{\mu\nu} \right] \\
&\equiv \int d^4x \left[\frac{1}{2} h_{\mu\nu} \mathcal{E}^{\mu\nu\alpha\beta} h_{\alpha\beta} + \frac{\kappa}{2} h_{\mu\nu} T^{\mu\nu} \right],
\end{aligned} \tag{A3}$$

where the kinetic operator $\mathcal{E}^{\mu\nu\alpha\beta}$ has the form

$$\mathcal{E}^{\mu\nu\alpha\beta} = (\eta^{\mu(\alpha} \eta^{\beta)\nu} - \eta^{\mu\nu} \eta^{\alpha\beta}) \square - \eta^{\mu(\alpha} \partial^{\beta)} \partial^{\nu} - \eta^{\nu(\alpha} \partial^{\beta)} \partial^{\mu} + \eta^{\alpha\beta} \partial^{\mu} \partial^{\nu} + \eta^{\mu\nu} \partial^{\alpha} \partial^{\beta}, \tag{A4}$$

and indices enclosed by brackets denote symmetrisation, $A^{(\mu} B^{\nu)} = \frac{1}{2}(A^{\mu} B^{\nu} + A^{\nu} B^{\mu})$. The massless graviton propagator $D_{\mu\nu\alpha\beta}^{(0)}$ is the inverse of the kinetic operator $\mathcal{E}^{\mu\nu\alpha\beta}$

$$\mathcal{E}^{\mu\nu\alpha\beta} D_{\alpha\beta\rho\sigma}^{(0)}(x-y) = \delta_{(\rho}^{\mu} \delta_{\sigma)}^{\nu} \delta^4(x-y). \tag{A5}$$

The massless graviton action Eq.A3 has the gauge symmetry $h_{\mu\nu} \rightarrow h_{\mu\nu} - \partial_{\mu} \xi_{\nu} - \partial_{\nu} \xi_{\mu}$ due to which the operator $\mathcal{E}^{\mu\nu\alpha\beta}$ cannot be inverted so the propagator cannot be determined from the relation Eq.A11. To invert the kinetic operator we need to choose a gauge. The gauge choice for which the propagator has the simplest form is the de-Dhonder gauge choice in which,

$$\partial^{\mu} h_{\mu\nu} - \frac{1}{2} \partial_{\nu} h = 0, \tag{A6}$$

where $h = h^{\alpha}_{\alpha}$. We can incorporate this gauge choice by adding the following gauge fixing term to the lagrangian Eq.A3,

$$S_{gf} = - \int d^4x \left(\partial^{\mu} h_{\mu\nu} - \frac{1}{2} \partial_{\nu} h \right)^2. \tag{A7}$$

The total action with the gauge fixing term turns out to be of the form

$$\begin{aligned} S_{EH} + S_{gf} &= \int d^4x \left(\frac{1}{2} h_{\mu\nu} \square h^{\mu\nu} - \frac{1}{4} h \square h + \frac{\kappa}{2} h_{\mu\nu} T^{\mu\nu} \right) \\ &= \int d^4x \left(\frac{1}{2} h_{\mu\nu} \mathcal{K}^{\mu\nu\alpha\beta} h_{\alpha\beta} + \frac{\kappa}{2} h_{\mu\nu} T^{\mu\nu} \right), \end{aligned} \quad (\text{A8})$$

where $\mathcal{K}^{\mu\nu\alpha\beta}$ is the kinetic operator in the de Donder gauge given by

$$\mathcal{K}^{\mu\nu\alpha\beta} = \left(\frac{1}{2} (\eta^{\mu\alpha} \eta^{\nu\beta} + \eta^{\mu\beta} \eta^{\nu\alpha}) - \frac{1}{2} \eta^{\mu\nu} \eta^{\alpha\beta} \right) \square. \quad (\text{A9})$$

The propagator in the de Donder gauge is the inverse of the kinetic operator Eq.A9 and is given by

$$\mathcal{K}^{\mu\nu\alpha\beta} D_{\alpha\beta\rho\sigma}^{(0)}(x-y) = \delta_{(\rho}^{\mu} \delta_{\sigma)}^{\nu} \delta^4(x-y). \quad (\text{A10})$$

This relation can be used to solve for $D_{\alpha\beta\rho\sigma}^{(0)}$ which in the momentum space ($\partial_\mu = ik_\mu$) is then given

$$D_{\mu\nu\alpha\beta}^{(0)}(k) = \frac{1}{-k^2} \left(\frac{1}{2} (\eta_{\mu\alpha} \eta_{\nu\beta} + \eta_{\mu\beta} \eta_{\nu\alpha}) - \frac{1}{2} \eta_{\mu\nu} \eta_{\alpha\beta} \right). \quad (\text{A11})$$

We treat the graviton as a quantum field by expanding it in terms of creation and annihilation operators,

$$\hat{h}_{\mu\nu}(x) = \sum_{\lambda} \int \frac{d^3k}{(2\pi)^3} \frac{1}{\sqrt{2\omega_k}} \left[\epsilon_{\mu\nu}^{\lambda}(k) a_{\lambda}(k) e^{-ik \cdot x} + \epsilon_{\mu\nu}^{*\lambda}(k) a_{\lambda}^{\dagger}(k) e^{ik \cdot x} \right]. \quad (\text{A12})$$

Here $\epsilon_{\mu\nu}^{\lambda}(k)$ are the polarization tensors which obey the orthogonality relation

$$\epsilon_{\mu\nu}^{\lambda}(k) \epsilon^{*\lambda'\mu\nu}(k) = \delta_{\lambda\lambda'}, \quad (\text{A13})$$

while $a_{\lambda}(k)$ and $a_{\lambda}^{\dagger}(k)$ are graviton annihilation and creation operator which obey the canonical commutation relations

$$\left[a_{\lambda}(k), a_{\lambda'}^{\dagger}(k') \right] = \delta^4(k-k') \delta_{\lambda\lambda'}. \quad (\text{A14})$$

The Feynman propagator of gravitons is defined as the time ordered two point function

$$D_{\mu\nu\alpha\beta}^{(0)}(x-y) \equiv \langle 0 | T(\hat{h}_{\mu\nu}(x) \hat{h}_{\alpha\beta}(y)) | 0 \rangle, \quad (\text{A15})$$

which may be evaluated using Eq.A12 to give

$$D_{\mu\nu\alpha\beta}^{(0)}(x-y) = \int \frac{d^4k}{(2\pi)^4} \frac{1}{-k^2 + i\epsilon} e^{ik(x-y)} \sum_{\lambda} \epsilon_{\mu\nu}^{\lambda}(k) \epsilon_{\alpha\beta}^{*\lambda}(k). \quad (\text{A16})$$

Comparing Eq.A11 and Eq.A16 we have the expression for the polarization sum of massless spin-2 gravitons

$$\sum_{\lambda=1}^2 \epsilon_{\mu\nu}^{\lambda}(k) \epsilon_{\alpha\beta}^{*\lambda}(k) = \frac{1}{2}(\eta_{\mu\alpha}\eta_{\nu\beta} + \eta_{\mu\beta}\eta_{\nu\alpha}) - \frac{1}{2}\eta_{\mu\nu}\eta_{\alpha\beta}. \quad (\text{A17})$$

This will be used in the computation of massless graviton radiation from classical sources.

We now calculate the energy loss due to the radiation of massless graviton from compact binary systems [14, 15] by evaluating the Feynman diagram shown in FIG.1. We treat the current $T_{\mu\nu}$ of the binary stars as classical source and the gravitons as quantum fields. From the interaction Lagrangian Eq.A8 we see that the interaction vertex is $\frac{1}{2}\kappa h^{\mu\nu}T_{\mu\nu}$, therefore we can write the emission rate of massless gravitons with polarisation tensor $\epsilon_{\lambda}^{\mu\nu}(k')$ from the classical source $T_{\mu\nu}(k)$ as

$$d\Gamma = \frac{\kappa^2}{4} \sum_{\lambda=1}^2 |T_{\mu\nu}(k') \epsilon_{\lambda}^{\mu\nu}(k)|^2 2\pi \delta(\omega - \omega') \frac{d^3k}{(2\pi)^3} \frac{1}{2\omega}. \quad (\text{A18})$$

Expanding the modulus squared in Eq.A18, we can write

$$d\Gamma = \frac{\kappa^2}{8(2\pi)^2} \sum_{\lambda=1}^2 \left(T_{\mu\nu}(k') T_{\alpha\beta}^*(k') \epsilon_{\lambda}^{\mu\nu}(k) \epsilon_{\lambda}^{*\alpha\beta}(k) \right) \frac{d^3k}{\omega} \delta(\omega - \omega'). \quad (\text{A19})$$

Using the polarization sum of massless spin-2 gravitons from Eq.A17, we can write the emission rate as

$$\begin{aligned} d\Gamma &= \frac{\kappa^2}{8(2\pi)^2} \int \left[T_{\mu\nu}(k') T_{\alpha\beta}^*(k') \right] \left[\frac{1}{2}(\eta^{\mu\alpha}\eta^{\nu\beta} + \eta^{\mu\beta}\eta^{\nu\alpha} - \eta^{\mu\nu}\eta^{\alpha\beta}) \right] \frac{d^3k}{\omega} \delta(\omega - \omega'). \\ &= \frac{\kappa^2}{8(2\pi)^2} \int \left[|T_{\mu\nu}(k')|^2 - \frac{1}{2}|T^{\mu}{}_{\mu}(k')|^2 \right] \delta(\omega - \omega') \omega d\omega d\Omega_k, \end{aligned} \quad (\text{A20})$$

where we use $d^3k = k^2 dk d\Omega$. Thus, the rate of energy loss due to massless graviton radiation becomes

$$\frac{dE}{dt} = \frac{\kappa^2}{8(2\pi)^2} \int \left[|T_{\mu\nu}(k')|^2 - \frac{1}{2}|T^{\mu}{}_{\mu}(k')|^2 \right] \delta(\omega - \omega') \omega^2 d\omega d\Omega_k. \quad (\text{A21})$$

Using the conserved current relation $k_{\mu}T^{\mu\nu} = 0$, we can write the T^{00} and T^{i0} components of the stress tensor in terms of the T^{ij} components,

$$T_{0j} = -\hat{k}^i T_{ij}, \quad T_{00} = \hat{k}^i \hat{k}^j T_{ij}. \quad (\text{A22})$$

Therefore, we can write

$$\left[|T_{\mu\nu}(k')|^2 - \frac{1}{2}|T^\mu{}_\mu(k')|^2\right] = \Lambda_{ij,lm}^0 T^{ij*} T^{lm}, \quad (\text{A23})$$

where,

$$\Lambda_{ij,lm}^0 = \left[\delta_{il}\delta_{jm} - 2\hat{k}_j\hat{k}_m\delta_{il} + \frac{1}{2}\hat{k}_i\hat{k}_j\hat{k}_l\hat{k}_m - \frac{1}{2}\delta_{ij}\delta_{lm} + \frac{1}{2}\left(\delta_{ij}\hat{k}_l\hat{k}_m + \delta_{lm}\hat{k}_i\hat{k}_j\right) \right]. \quad (\text{A24})$$

We do the angular integrals

$$\int d\Omega_k \Lambda_{ij,lm}^0 T^{ij*}(\omega') T^{lm}(\omega') = \frac{8\pi}{5} \left(T_{ij}(\omega') T_{ji}^*(\omega') - \frac{1}{3} |T^i{}_i(\omega')|^2 \right), \quad (\text{A25})$$

using the relations

$$\int d\Omega \hat{k}^i \hat{k}^j = \frac{4\pi}{3} \delta_{ij}, \quad \int d\Omega \hat{k}^i \hat{k}^j \hat{k}^l \hat{k}^m = \frac{4\pi}{15} (\delta_{ij}\delta_{lm} + \delta_{il}\delta_{jm} + \delta_{im}\delta_{jl}). \quad (\text{A26})$$

The stress tensor or the current density for this compact binary system is

$$T_{\mu\nu}(x') = \mu \delta^3(\mathbf{x}' - \mathbf{x}(t)) U_\mu U_\nu, \quad (\text{A27})$$

where $\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass of the binary system and m_1 and m_2 are the masses of the two stars in the binary system. $U_\mu = (1, \dot{x}, \dot{y}, 0)$ is the non relativistic four velocity of the reduced mass of the binary system in the $x - y$ plane of the Keplerian orbit. We can write the Keplerian orbit in the parametric form as

$$x = a(\cos \xi - e), \quad y = a\sqrt{(1 - e^2)} \sin \xi, \quad \Omega t = \xi - e \sin \xi, \quad (\text{A28})$$

where a and e are the semi-major axis and eccentricity of the elliptic orbit respectively. Since the angular velocity of an eccentric orbit is not constant, we can write the fourier transform of the current density in terms of the n harmonics of the fundamental frequency $\Omega = \left[G \frac{(m_1 + m_2)}{a^3} \right]^{\frac{1}{2}}$. Using Eq.A28, we can write the Fourier transforms of the velocity components in the Kepler orbit as

$$\dot{x}_n = \frac{1}{T} \int_0^T e^{in\Omega t} \dot{x} dt = -ia\Omega J'_n(ne), \quad (\text{A29})$$

and

$$\dot{y}_n = \frac{1}{T} \int_0^T e^{in\Omega t} \dot{y} dt = \frac{a\sqrt{(1 - e^2)}}{e} \Omega J_n(ne), \quad (\text{A30})$$

where we have used $T = 2\pi/\Omega$ and the Bessel function identity $J_n(z) = \frac{1}{2\pi} \int_0^{2\pi} e^{i(n\xi - z \sin \xi)} d\xi$. The prime over the Bessel function denotes the derivative with respect to the argument. Hence the Fourier transforms of the orbital coordinates become

$$x_n = \frac{\dot{x}_n}{-i\Omega n} = \frac{a}{n} J'_n(ne), \quad y_n = \frac{\dot{y}_n}{-i\Omega n} = \frac{ia\sqrt{1-e^2}}{ne} J_n(ne). \quad (\text{A31})$$

Now we will calculate the Fourier transforms of different components of the stress tensor with $\omega' = n\Omega$ as below. Thus,

$$\begin{aligned} T_{ij}(\mathbf{k}', \omega') &= \frac{1}{T} \int_0^T \int T_{ij}(\mathbf{x}, t) e^{-i(\mathbf{k}' \cdot \mathbf{x} - \omega' t)} d^3x dt \\ &= \int T_{ij}(\mathbf{x}, \omega') e^{-i\mathbf{k}' \cdot \mathbf{x}} d^3x. \end{aligned} \quad (\text{A32})$$

Expanding $e^{-i\mathbf{k}' \cdot \mathbf{x}} \approx 1 - i\mathbf{k}' \cdot \mathbf{x} - \dots$ and retaining the leading order term $\mathbf{k}' \cdot \mathbf{x} \sim \Omega a \ll 1$ for binary orbit, we can write Eq.A32 as

$$T_{ij}(\mathbf{k}', \omega') \simeq T_{ij}(\omega') = \int T_{ij}(\mathbf{x}, \omega') d^3x. \quad (\text{A33})$$

From conservation of the stress-energy tensor, i.e. $\partial_\mu T^{\mu\nu}(\mathbf{x}, t) = 0$, we get

$$\partial^i \partial^j T_{ij}(\mathbf{x}, \omega') = -\omega'^2 T_{00}(\mathbf{x}, \omega'). \quad (\text{A34})$$

Multiplying both side of the Eq.A34 by $x_k x_l$ and integrating over all \mathbf{x} we get

$$T_{kl}(\omega') = -\frac{\omega'^2}{2} \int T_{00}(\mathbf{x}, \omega') x_k x_l d^3x \quad (\text{A35})$$

$$= -\frac{\mu\omega'^2}{2T} \int_0^T \int \delta^3(\mathbf{x}' - \mathbf{x}(t)) e^{i\omega' t} x_k x_l d^3x dt \quad (\text{A36})$$

$$= -\frac{\mu\omega'^2}{2T} \int_0^T x'_k(t) x'_l(t) e^{i\omega' t} dt, \quad (\text{A37})$$

where in the Eq.A36 we used the Eq.A27. Doing integration by parts of Eq.A37 and using the Bessel function identities¹, we can write the different components of stress tensor in the $x - y$ plane. The xx -component of stress tensor in the Fourier space is

$$\begin{aligned} T_{xx}(\omega') &= -\frac{\mu\omega'^2}{2T} \int_0^T x^2(t) e^{i\omega' t} dt \\ &= -\frac{\mu\omega'^2}{4\pi} \int_0^{2\pi} a^2 (\cos \xi - e)^2 e^{in\beta} d\beta, \end{aligned} \quad (\text{A38})$$

¹ $J_{n-1}(z) - J_{n+1}(z) = 2J'_n(z)$, $J_{n-1}(z) + J_{n+1}(z) = \frac{2n}{z} J_n(z)$

where we used Eq.A28 and $\omega' = n\Omega$, $\beta = \Omega t$, and $T = 2\pi/\Omega$. Doing integration by parts of Eq.A38 we get

$$\begin{aligned}
T_{xx}(\omega') &= \frac{\mu\omega'^2}{4\pi in} \int_0^{2\pi} e^{in\beta} \frac{d}{d\beta} (\cos \xi - e)^2 d\beta \\
&= \frac{\mu\omega'^2}{2\pi in} \int_0^{2\pi} \sin \xi (\cos \xi - e) e^{in\beta} d\xi \\
&= -\frac{\mu\omega'^2}{8\pi n} \int_0^{2\pi} [(e^{2i\xi} - e^{-2i\xi}) - 2e(e^{i\xi} - e^{-i\xi})] e^{in\beta} d\xi \\
&= -\frac{\mu\omega'^2}{4n} [J_{n-2}(ne) - 2eJ_{n-1}(ne) + 2eJ_{n+1}(ne) - J_{n+2}(ne)], \tag{A39}
\end{aligned}$$

where in the last step we used the definition of the Bessel function and $\beta = \Omega t = \xi - e \sin \xi$.

Similarly,

$$\begin{aligned}
T_{yy}(\omega') &= -\frac{\mu\omega'^2}{2T} \int_0^T y^2(t) e^{i\omega't} dt \\
&= -\frac{\mu\omega'^2(1-e^2)}{4\pi} \int_0^{2\pi} a^2 \sin^2 \xi e^{in\beta} d\beta. \tag{A40}
\end{aligned}$$

Adding Eq.A38 and Eq.A40 we get

$$\begin{aligned}
T_{yy}(\omega') + T_{xx}(\omega') &= -\frac{\mu\omega'^2 a^2}{4\pi} \int_0^{2\pi} (1 - e \cos \xi)^2 e^{in\beta} d\beta \\
&= \frac{\mu\omega'^2 a^2 e}{2\pi in} \int_0^{2\pi} \sin \xi e^{in\beta} d\beta, \\
&= \frac{\mu\omega'^2 a^2}{2\pi n^2} \int_0^{2\pi} e \cos \xi e^{in\beta} d\xi, \\
&= \frac{\mu\omega'^2 a^2}{2\pi n^2} \int_0^{2\pi} e^{in\beta} d\xi = \frac{\mu\omega'^2 a^2}{n^2} J_n^2(ne). \tag{A41}
\end{aligned}$$

Therefore

$$\begin{aligned}
T_{yy}(\omega') &= -T_{xx}(\omega') + \frac{\mu\omega'^2 a^2}{n^2} J_n^2(ne) \\
&= \frac{\mu\omega'^2 a^2}{4n} \left[J_{n-2}(ne) - 2eJ_{n-1}(ne) + \frac{4}{n} J_n(ne) + 2eJ_{n+1}(ne) - J_{n+2}(ne) \right]. \tag{A42}
\end{aligned}$$

The xy -component of the Stress Tensor in the Fourier space is

$$\begin{aligned}
T_{xy}(\omega') &= -\frac{\mu\omega'^2}{2T} \int_0^T x(t)y(t)e^{i\omega't} dt \\
&= -\frac{\mu\omega'^2\sqrt{1-e^2}}{4\pi} \int_0^{2\pi} a^2(\cos\xi - e) \sin\xi e^{in\beta} d\beta \\
&= \frac{\mu\omega'^2 a^2 \sqrt{1-e^2}}{4\pi in} \int_0^{2\pi} (\cos(2\xi) - e \cos\xi) e^{in\beta} d\xi, \\
&= -i \frac{\mu\omega'^2 a^2 \sqrt{1-e^2}}{4\pi n} \int_0^{2\pi} (\cos(2\xi) - 1) e^{in\beta} d\xi \\
&= -i \frac{\mu\omega'^2 a^2 \sqrt{1-e^2}}{4n} [J_{n+2}(ne) - 2J_n(ne) + J_{n-2}(ne)] \tag{A43}
\end{aligned}$$

For convenience we summarize the final expressions of $T_{ij}(\omega')$ as

$$\begin{aligned}
T_{xx}(\omega') &= -\frac{\mu\omega'^2 a^2}{4n} [J_{n-2}(ne) - 2eJ_{n-1}(ne) + 2eJ_{n+1}(ne) - J_{n+2}(ne)], \\
T_{yy}(\omega') &= \frac{\mu\omega'^2 a^2}{4n} [J_{n-2}(ne) - 2eJ_{n-1}(ne) + \frac{4}{n}J_n(ne) + 2eJ_{n+1}(ne) - J_{n+2}(ne)], \\
T_{xy}(\omega') &= \frac{-i\mu\omega'^2 a^2}{4n} (1-e^2)^{\frac{1}{2}} [J_{n-2}(ne) - 2J_n(ne) + J_{n+2}(ne)]. \tag{A44}
\end{aligned}$$

Using Eq.A44, we get two useful results

$$\begin{aligned}
T_{ij}(\omega')T^{ij*}(\omega') &= \frac{\mu^2\omega'^4 a^4}{8n^2} \left\{ [J_{n-2}(ne) - 2eJ_{n-1}(ne) + 2eJ_{n+1}(ne) + \frac{2}{n}J_n(ne) - J_{n+2}(ne)]^2 \right. \\
&\quad \left. + (1-e^2)[J_{n-2}(ne) - 2J_n(ne) + J_{n+2}(ne)]^2 + \frac{4}{n^2}J_n^2(ne) \right\} \\
&= 4\mu^2\omega'^4 a^4 \left(f(n, e) + \frac{J_n^2(ne)}{12n^4} \right), \tag{A45}
\end{aligned}$$

where

$$\begin{aligned}
f(n, e) &= \frac{1}{32n^2} \left\{ [J_{n-2}(ne) - 2eJ_{n-1}(ne) + 2eJ_{n+1}(ne) + \frac{2}{n}J_n(ne) - J_{n+2}(ne)]^2 + \right. \\
&\quad \left. (1-e^2)[J_{n-2}(ne) - 2J_n(ne) + J_{n+2}(ne)]^2 + \frac{4}{3n^2}J_n^2(ne) \right\} \tag{A46}
\end{aligned}$$

and

$$|T^i_i|^2 = \frac{\mu^2\omega'^4 a^4}{n^4} J_n^2(ne). \tag{A47}$$

Thus the energy loss due to massless graviton radiation becomes

$$\begin{aligned}
\frac{dE}{dt} &= \frac{\kappa^2}{8(2\pi)^2} \int \frac{8\pi}{5} \left[T_{ij}(\omega') T_{ji}^*(\omega') - \frac{1}{3} |T^i{}_i(\omega')|^2 \right] \delta(\omega - \omega') \omega^2 d\omega, \\
&= \frac{32G}{5} \sum_{n=1}^{\infty} (n\Omega)^2 \mu^2 a^4 (n\Omega)^4 f(n, e) \\
&= \frac{32G}{5} \Omega^6 \left(\frac{m_1 m_2}{m_1 + m_2} \right)^2 a^4 (1 - e^2)^{-7/2} \left(1 + \frac{73}{24} e^2 + \frac{37}{96} e^4 \right). \tag{A48}
\end{aligned}$$

This expression is called Einstein’s quadrupole gravitational radiation which matches with the Peters-Mathews result [16]. From the energy loss formula we can calculate the change in time period ($P_b = 2\pi/\Omega$). From Kepler’s law $\Omega^2 a^3 = G(M_1 + M_2)$ we have $\dot{a}/a = (2/3)(\dot{P}_b/P_b)$. The gravitational energy is $E = -GM_1 M_2/2a$ which implies $\dot{a}/a = -(\dot{E}/E)$. Using these two relations we get $\dot{P}_b/P_b = -(3/2)(\dot{E}/E)$.

-
- [1] S. Weinberg, Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity. John Wiley and Sons, New York, 1972.
- [2] M. Fierz and W. Pauli, “On relativistic wave equations for particles of arbitrary spin in an electromagnetic field,” Proc. Roy. Soc. Lond. A **173** (1939) 211–232.
- [3] H. van Dam and M. J. G. Veltman, “Massive and massless Yang-Mills and gravitational fields,” Nucl. Phys. B **22** (1970) 397–411.
- [4] V. I. Zakharov, “Linearized gravitation theory and the graviton mass,” JETP Lett. **12** (1970) 312.
- [5] K. Hinterbichler, “Theoretical Aspects of Massive Gravity,” Rev. Mod. Phys. **84** (2012) 671–710, [arXiv:1105.3735 \[hep-th\]](#).
- [6] C. de Rham, “Massive Gravity,” Living Rev. Rel. **17** (2014) 7, [arXiv:1401.4173 \[hep-th\]](#).
- [7] E. Mitsou, Aspects of Infrared Non-local Modifications of General Relativity. PhD thesis, Geneva U., 2015. [arXiv:1504.04050 \[gr-qc\]](#).
- [8] A. Joyce, B. Jain, J. Khoury, and M. Trodden, “Beyond the Cosmological Standard Model,” Phys. Rept. **568** (2015) 1–98, [arXiv:1407.0059 \[astro-ph.CO\]](#).

- [9] C. de Rham, J. T. Deskins, A. J. Tolley, and S.-Y. Zhou, “Graviton Mass Bounds,” Rev. Mod. Phys. **89** no. 2, (2017) 025004, [arXiv:1606.08462](#) [[astro-ph.CO](#)].
- [10] R. P. Feynman, Feynman lectures on gravitation. 1996.
- [11] S. Weinberg, “Photons and Gravitons in *S*-Matrix Theory: Derivation of Charge Conservation and Equality of Gravitational and Inertial Mass,” Phys. Rev. **135** (1964) B1049–B1056.
- [12] M. J. G. Veltman, “Quantum Theory of Gravitation,” Conf. Proc. C **7507281** (1975) 265–327.
- [13] J. F. Donoghue, M. M. Ivanov, and A. Shkerin, “EPFL Lectures on General Relativity as a Quantum Field Theory,” [arXiv:1702.00319](#) [[hep-th](#)].
- [14] S. Mohanty and P. Kumar Panda, “Particle physics bounds from the Hulse-Taylor binary,” Phys. Rev. D **53** (1996) 5723–5726, [arXiv:hep-ph/9403205](#).
- [15] S. Mohanty, Astroparticle Physics and Cosmology: Perspectives in the Multimessenger Era, vol. 975. 2020.
- [16] P. C. Peters and J. Mathews, “Gravitational radiation from point masses in a Keplerian orbit,” Phys. Rev. **131** (1963) 435–439.
- [17] R. A. Hulse and J. H. Taylor, “Discovery of a pulsar in a binary system,” Astrophys. J. Lett. **195** (1975) L51–L53.
- [18] J. H. Taylor and J. M. Weisberg, “A new test of general relativity: Gravitational radiation and the binary pulsar PS R 1913+16,” Astrophys. J. **253** (1982) 908–920.
- [19] J. M. Weisberg and J. H. Taylor, “Observations of Post-Newtonian Timing Effects in the Binary Pulsar PSR 1913+16,” Phys. Rev. Lett. **52** (1984) 1348–1350.
- [20] J. M. Weisberg and Y. Huang, “Relativistic Measurements from Timing the Binary Pulsar PSR B1913+16,” Astrophys. J. **829** no. 1, (2016) 55, [arXiv:1606.02744](#) [[astro-ph.HE](#)].
- [21] M. Kramer et al., “Tests of general relativity from timing the double pulsar,” Science **314** (2006) 97–102, [arXiv:astro-ph/0609417](#).
- [22] J. Antoniadis et al., “A Massive Pulsar in a Compact Relativistic Binary,” Science **340** (2013) 6131, [arXiv:1304.6875](#) [[astro-ph.HE](#)].
- [23] P. C. C. Freire, N. Wex, G. Esposito-Farese, J. P. W. Verbiest, M. Bailes, B. A. Jacoby,

- M. Kramer, I. H. Stairs, J. Antoniadis, and G. H. Janssen, “The relativistic pulsar-white dwarf binary PSR J1738+0333 II. The most stringent test of scalar-tensor gravity,” Mon. Not. Roy. Astron. Soc. **423** (2012) 3328, [arXiv:1205.1450 \[astro-ph.GA\]](#).
- [24] T. Kumar Poddar, S. Mohanty, and S. Jana, “Constraints on ultralight axions from compact binary systems,” Phys. Rev. D **101** no. 8, (2020) 083007, [arXiv:1906.00666 \[hep-ph\]](#).
- [25] T. Kumar Poddar, S. Mohanty, and S. Jana, “Vector gauge boson radiation from compact binary systems in a gauged $L_\mu - L_\tau$ scenario,” Phys. Rev. D **100** no. 12, (2019) 123023, [arXiv:1908.09732 \[hep-ph\]](#).
- [26] W. Hu, R. Barkana, and A. Gruzinov, “Cold and fuzzy dark matter,” Phys. Rev. Lett. **85** (2000) 1158–1161, [arXiv:astro-ph/0003365](#).
- [27] L. Hui, J. P. Ostriker, S. Tremaine, and E. Witten, “Ultralight scalars as cosmological dark matter,” Phys. Rev. D **95** no. 4, (2017) 043541, [arXiv:1610.08297 \[astro-ph.CO\]](#).
- [28] M. Visser, “Mass for the graviton,” Gen. Rel. Grav. **30** (1998) 1717–1728, [arXiv:gr-qc/9705051](#).
- [29] L. S. Finn and P. J. Sutton, “Bounding the mass of the graviton using binary pulsar observations,” Phys. Rev. D **65** (2002) 044022, [arXiv:gr-qc/0109049](#).
- [30] G. Gambuti and N. Maggiore, “A note on harmonic gauge(s) in massive gravity,” Phys. Lett. B **807** (2020) 135530, [arXiv:2006.04360 \[gr-qc\]](#).
- [31] G. Gambuti and N. Maggiore, “Fierz–Pauli theory reloaded: from a theory of a symmetric tensor field to linearized massive gravity,” Eur. Phys. J. C **81** no. 2, (2021) 171, [arXiv:2102.10813 \[gr-qc\]](#).
- [32] G. R. Dvali, G. Gabadadze, and M. Porrati, “4-D gravity on a brane in 5-D Minkowski space,” Phys. Lett. B **485** (2000) 208–214, [arXiv:hep-th/0005016](#).
- [33] G. R. Dvali, G. Gabadadze, and M. Porrati, “Metastable gravitons and infinite volume extra dimensions,” Phys. Lett. B **484** (2000) 112–118, [arXiv:hep-th/0002190](#).
- [34] G. R. Dvali and G. Gabadadze, “Gravity on a brane in infinite volume extra space,” Phys. Rev. D **63** (2001) 065007, [arXiv:hep-th/0008054](#).
- [35] P. Van Nieuwenhuizen, “Radiation of massive gravitation,” Phys. Rev. D **7** (1973) 2300–2308.
- [36] C. M. Will, “Bounding the mass of the graviton using gravitational wave observations of

- inspiralling compact binaries,” Phys. Rev. D **57** (1998) 2061–2068, [arXiv:gr-qc/9709011](#).
- [37] S. L. Larson and W. A. Hiscock, “Using binary stars to bound the mass of the graviton,” Phys. Rev. D **61** (2000) 104008, [arXiv:gr-qc/9912102](#).
- [38] C. de Rham, A. J. Tolley, and D. H. Wesley, “Vainshtein Mechanism in Binary Pulsars,” Phys. Rev. D **87** no. 4, (2013) 044025, [arXiv:1208.0580 \[gr-qc\]](#).
- [39] L. Shao, N. Wex, and S.-Y. Zhou, “New Graviton Mass Bound from Binary Pulsars,” Phys. Rev. D **102** no. 2, (2020) 024069, [arXiv:2007.04531 \[gr-qc\]](#).
- [40] C. Talmadge, J. P. Berthias, R. W. Hellings, and E. M. Standish, “Model Independent Constraints on Possible Modifications of Newtonian Gravity,” Phys. Rev. Lett. **61** (1988) 1159–1162.
- [41] E. Fomalont, S. Kopeikin, G. Lanyi, and J. Benson, “Progress in Measurements of the Gravitational Bending of Radio Waves Using the VLBA,” Astrophys. J. **699** (2009) 1395–1402, [arXiv:0904.3992 \[astro-ph.CO\]](#).
- [42] A. I. Vainshtein, “To the problem of nonvanishing gravitation mass,” Phys. Lett. B **39** (1972) 393–394.
- [43] E. Babichev and C. Deffayet, “An introduction to the Vainshtein mechanism,” Class. Quant. Grav. **30** (2013) 184001, [arXiv:1304.7240 \[gr-qc\]](#).
- [44] T. Kumar Poddar, S. Mohanty, and S. Jana, “Constraints on long range force from perihelion precession of planets in a gauged $L_e - L_{\mu,\tau}$ scenario,” Eur. Phys. J. C **81** no. 4, (2021) 286, [arXiv:2002.02935 \[hep-ph\]](#).
- [45] T. Kumar Poddar, “Constraints on axionic fuzzy dark matter from light bending and Shapiro time delay,” [arXiv:2104.09772 \[hep-ph\]](#).
- [46] **LIGO Scientific, Virgo** Collaboration, B. P. Abbott *et al.*, “Tests of general relativity with GW150914,” Phys. Rev. Lett. **116** no. 22, (2016) 221101, [arXiv:1602.03841 \[gr-qc\]](#). [Erratum: Phys.Rev.Lett. 121, 129902 (2018)].
- [47] D. G. Boulware and S. Deser, “Can gravitation have a finite range?,” Phys. Rev. D **6** (1972) 3368–3382.
- [48] G. Dvali, “Predictive Power of Strong Coupling in Theories with Large Distance Modified Gravity,” New J. Phys. **8** (2006) 326, [arXiv:hep-th/0610013](#).

- [49] C. Deffayet, G. R. Dvali, and G. Gabadadze, “Accelerated universe from gravity leaking to extra dimensions,” Phys. Rev. D **65** (2002) 044023, [arXiv:astro-ph/0105068](#).
- [50] V. A. Rubakov, “Lorentz-violating graviton masses: Getting around ghosts, low strong coupling scale and VDVZ discontinuity,” [arXiv:hep-th/0407104](#).
- [51] S. L. Dubovsky, “Phases of massive gravity,” JHEP **10** (2004) 076, [arXiv:hep-th/0409124](#).
- [52] V. A. Rubakov and P. G. Tinyakov, “Infrared-modified gravities and massive gravitons,” Phys. Usp. **51** (2008) 759–792, [arXiv:0802.4379 \[hep-th\]](#).
- [53] V. A. Kostelecký and M. Mewes, “Testing local Lorentz invariance with gravitational waves,” Phys. Lett. B **757** (2016) 510–514, [arXiv:1602.04782 \[gr-qc\]](#).
- [54] V. A. Kostelecký and M. Mewes, “Testing local Lorentz invariance with short-range gravity,” Phys. Lett. B **766** (2017) 137–143, [arXiv:1611.10313 \[gr-qc\]](#).
- [55] **LIGO Scientific** Collaboration, J. Aasi et al., “Advanced LIGO,” Class. Quant. Grav. **32** (2015) 074001, [arXiv:1411.4547 \[gr-qc\]](#).
- [56] **VIRGO** Collaboration, F. Acernese et al., “Advanced Virgo: a second-generation interferometric gravitational wave detector,” Class. Quant. Grav. **32** no. 2, (2015) 024001, [arXiv:1408.3978 \[gr-qc\]](#).