

REDUCTION FOR BRANCHING MULTIPLICITIES

PIERRE-EMMANUEL CHAPUT AND NICOLAS RESSAYRE

ABSTRACT. A reduction formula for the branching coefficients of the restrictions of representations of a semisimple group to a semisimple subgroup is proved in [KT03, DW11, Res11b]. This formula holds when the highest weights of the representations belong to a codimension 1 face of the Horn cone, which by [Res10] corresponds to some Schubert coefficient equal to 1. We prove a similar reduction formula when this Schubert coefficient is equal to 2, and show some properties of the class of the branch divisor corresponding to a generically finite morphism naturally defined in this context.

1. INTRODUCTION

Fix an inclusion $G \subset \hat{G}$ of complex connected reductive groups. We are interested in the branching problem for the decomposition of irreducible \hat{G} -modules as representations of G . The appearing multiplicities are nonnegative integers parametrized by the pairs of dominant weights for G and \hat{G} respectively. The support of this multiplicity function is known to be a finitely generated semigroup. It generates a convex polyhedral cone $\Gamma_{\mathbb{Q}}(G \subset \hat{G})$, that we call the *Horn cone*. The set of codimension one faces of this cone is in bijection with the set of Levi-movable pairs (see Section 6.1) of Schubert classes in some projective homogeneous spaces $G/P \subset \hat{G}/\hat{P}$ such that the Schubert coefficient c of the first Schubert class with the pullback of the second Schubert class in G/P is equal to one. See [Res10].

Moreover, the multiplicities on those faces satisfy reduction rules: if L, \hat{L} denote the Levi subgroups of P, \hat{P} , the branching multiplicities for the inclusion $G \subset \hat{G}$ on such a face are equal to branching multiplicities for the inclusion $L \subset \hat{L}$. See [Res11b].

Actually, to any pair of Schubert classes such that the Schubert coefficient c is positive corresponds an inequality satisfied by $\Gamma_{\mathbb{Q}}(G \subset \hat{G})$. Our aim is to study the associated reduction rules for the multiplicities on the associated face when $c = 2$. Although most of our results are general we consider in this introduction the case of the tensor product decomposition for the linear group.

Fix an n -dimensional vector space V . Let $\Lambda_n^+ = \{(\lambda_1 \geq \dots \geq \lambda_n \geq 0 : \lambda_i \in \mathbb{N})\}$ denote the set of partitions. For $\lambda \in \Lambda_n^+$, let $S^\lambda V$ be the corresponding Schur module, that is the irreducible $\mathrm{GL}(V)$ -module of highest weight $\sum \lambda_i \epsilon_i$ (notation as in [Bou02]). The Littlewood-Richardson coefficients (or LR coefficients for short) $c_{\lambda, \mu}^\nu$ are defined by

$$(1) \quad S^\lambda V \otimes S^\mu V \simeq \bigoplus_{\nu \in \Lambda_n^+} \mathbb{C}^{c_{\lambda, \mu}^\nu} \otimes S^\nu V,$$

where $\mathbb{C}^{c_{\lambda, \mu}^\nu}$ is a multiplicity space.

Let $1 \leq r \leq n - 1$ and $\mathrm{Gr}(r, n)$ be the Grassmannian of r -dimensional linear subspaces of V . Recall that the Schubert basis (σ^I) of the Chow ring $A^*(\mathrm{Gr}(r, n), \mathbb{Z})$ is parametrized by the subsets I of $\{1, \dots, n\}$ with r elements. The Schubert coefficients $c_{I, J}^K$ are defined by

$$(2) \quad \sigma^I \sigma^J = \sum_K c_{I, J}^K \sigma^K.$$

Actually, $c_{I, J}^K$ is also a LR coefficient by [Les47], but this coincidence is specific to the type A .

Given a partition λ and a subset I , let λ_I be the partition whose parts are λ_i with $i \in I$. Let also \bar{I} denote the complementary subset $\{1, \dots, n\} \setminus I$.

Theorem 1. [KT03, DW11]

This work was supported by the ANR GeoLie project, of the French Agence Nationale de la Recherche.

(1) Let $1 \leq r \leq n-1$ and $I, J, K \subset \{1, \dots, n\}$ be subsets with r elements such that $c_{I,J}^K \neq 0$. For any $(\lambda, \mu, \nu) \in (\Lambda_n^+)^3$, if $c_{\lambda,\mu}^\nu \neq 0$ then

$$(3) \quad |\lambda_I| + |\mu_J| \geq |\nu_K|.$$

(2) Conversely, fix $(\lambda, \mu, \nu) \in (\Lambda_n^+)^3$ such that $|\lambda| + |\mu| = |\nu|$. If for any $1 \leq r \leq n-1$ and I, J, K of cardinal r such that $c_{I,J}^K = 1$, inequality (3) holds, then $c_{\lambda,\mu}^\nu \neq 0$.

The reduction result that we mentioned above, corresponding to the semisimple part $\mathrm{SL}_r \times \mathrm{SL}_{n-r} \subset \mathrm{SL}_n$ of the Levi subgroup is the following:

Theorem 2. [KTT09, Rot11, DW11, Res11b] Assume that $c_{I,J}^K = 1$. Let λ, μ, ν be partitions such that

$$(4) \quad |\lambda_I| + |\mu_J| = |\nu_K|.$$

Then

$$(5) \quad c_{\lambda,\mu}^\nu = c_{\lambda_I, \mu_J}^{\nu_K} \cdot c_{\lambda_{\overline{T}}, \mu_{\overline{T}}}^{\nu_{\overline{K}}}.$$

Formula (5) is a multiplicativity property. Let us first report on a similar property for Belkale-Kumar [BK06] coefficients (BK coefficients for short). See Section 6.1 for a short description of these numbers which are all Schubert coefficients or zero. Consider an inclusion $P \subset Q$ of parabolic subgroups of a reductive algebraic group G , and the corresponding fibration $G/P \rightarrow G/Q$. Richmond [Ric12] proved that any BK coefficient d of G/P is the product of two BK coefficients for G/Q and Q/P . In type A, this implies that a non-zero BK coefficient for any two steps flag manifold is a product of two LR coefficients: $d = c_1 c_2$.

If moreover $c_1 = 1$, Theorem 2 implies that c_2 itself is the product of two LR coefficients: $c_2 = c'_2 c''_2$. Thus $d = c_1 c'_2 c''_2$ is the product of *three* LR coefficients. This is the content of [KP11, Theorem 3], which even more generally states that on a k -step flag variety, any BK coefficient can be factorized as a product of $\frac{k(k-1)}{2}$ LR coefficients. Unfortunately, this assertion needs $c_1 = 1$ and is not correct in general, as we show in Remark 8. Our original motivation was to correct this result. We get such a correction if $c_1 = 2$.

In [Res11b], the variety $Y \subset \mathrm{Gr}(r, n) \times (G/B)^3$ is defined, where a quadruple (V, X_1, X_2, X_3) belongs to Y if and only if $V \in \mathrm{Gr}(r, n)$ belongs to the intersection of the three Schubert varieties defined by I, J, K and the three flags X_1, X_2, X_3 .

The hypothesis $c_{I,J}^K = 1$ implies that the projection $\Pi : Y \rightarrow (G/B)^3$ is birational. By the projection formula, since Y has rational singularities, it follows that the sections of line bundles on $(G/B)^3$ identify to those on Y . Let $C \subset (G/B)^3$ be a product of flag varieties under L . In [Res11b], it is shown that taking G -invariants and restricting to C is the same as restricting to C and then taking L -invariants, which leads to (5).

Fix now r and $I, J, K \subset \{1, \dots, n\}$ of cardinal r such that $c_{I,J}^K = 2$. We can still define $\Pi : Y \rightarrow (G/B)^3$ in a similar way, but it is now generically finite of degree 2. We show in Subsection 3.4 that we can associate to this generically finite morphism a branch divisor in $(G/B)^3$ and that the pushforward of the structure sheaf \mathcal{O}_Y is expressed in terms of this branch divisor. In Theorem 7, we determine the triple (α, β, γ) of dominant weights defining the branch divisor class. This is done using the morphism $Y \rightarrow \mathrm{Gr}(r, n)$ which describes Y as a relative product of Schubert varieties over the Grassmannian. This shows that Y is normal and allows computing the ramification divisor as the relative canonical sheaf, adapting in this relative setting previous computations of the canonical sheaf of Schubert varieties ([Ram87, Theorem 4.2] and [Per07, Proposition 4.4]), see Section 4.5.

Under an assumption of Levi-movability (see Sections 6.1 and 6.2), which is automatically satisfied in type A, we get an alternative characterization of this triple in Proposition 36. Namely, (α, β, γ) is the unique minimal element of the set of triples $(\lambda, \mu, \nu) \in (\Lambda_n^+)^3$ such that

$$|\lambda_I| + |\mu_J| = |\nu_K| \quad \text{and} \quad 0 \neq c_{\lambda,\mu}^\nu < c_{\lambda_I, \mu_J}^{\nu_K} \cdot c_{\lambda_{\overline{T}}, \mu_{\overline{T}}}^{\nu_{\overline{K}}}.$$

The interested reader can find at [CR21] a program allowing to compute (α, β, γ) and its generalisations to any classical group (and under the more general assumption $c_{I,J}^K \neq 0$).

Arguing as in [Res11b], we deduce from our description of $\Pi_*\mathcal{O}_Y$ our general result, Theorem 6. It holds for any inclusion $G \subset \hat{G}$. For $\hat{G} = G \times G$ and G/P a Grassmannian, it states the following modification of (5):

Theorem 3. *Assume $c_{I,J}^K = 2$. Let λ, μ, ν be partitions such that $|\lambda_I| + |\mu_J| = |\nu_K|$. Then*

$$(6) \quad c_{\lambda,\mu}^\nu + c_{\lambda-\alpha,\mu-\beta}^{\nu-\gamma} = c_{\lambda_I,\mu_J}^{\nu_K} \cdot c_{\lambda_{\mathcal{T}},\mu_{\mathcal{T}}}^{\nu_{\mathcal{K}}}.$$

Observe that although the Horn cone is defined by inequations of the form (4) for I, J, K such that $c_{I,J}^K = 1$ inside $(\Lambda_n^+)^3$, the inequalities coming from the fact $\lambda, \mu, \nu \in \Lambda_n^+$ make it possible that an element (λ, μ, ν) in the Horn cone satisfies an equality $|\lambda_I| + |\mu_J| = |\nu_K|$ for some triple (I, J, K) such that $c_{I,J}^K = 2$ but for no triple such that $c_{I,J}^K = 1$: see Remark 3. For such an element, Theorem 1 cannot be applied but Theorem 3 can.

Under the assumption of Levi-movability, the mentioned characterization of (α, β, γ) implies that it satisfies (4). Then, an immediate induction expresses the multiplicity as an alternating sum of similar coefficients for the reduction $L \subset \hat{L}$. This is Corollary 37 below. In the setting of Theorem 3, we get:

Theorem 4. *Assume $c_{I,J}^K = 2$. Let λ, μ, ν be partitions such that $|\lambda_I| + |\mu_J| = |\nu_K|$. Then,*

$$(7) \quad c_{\lambda,\mu}^\nu = \sum_{k \geq 0} (-1)^k c_{\lambda_I - k\alpha_I, \mu_J - k\beta_J}^{\nu_K - k\gamma_K} \cdot c_{\lambda_{\mathcal{T}} - k\alpha_{\mathcal{T}}, \mu_{\mathcal{T}} - k\beta_{\mathcal{T}}}^{\nu_{\mathcal{K}} - k\gamma_{\mathcal{K}}}.$$

Given three subsets I, J, K as above, it is a difficult task to describe the face it defines in the Horn cone: to the best of our knowledge, even the dimension of such a face is not known in general, and our experience is that a computer will only give very limited information related to this problem. From a theoretical point of view as well as a computational point of view, both finding linear equations defining this face and points on it is challenging. In Section 7.3, we describe completely the faces corresponding to some relevant examples.

We also believe that the study of the ramification or branch divisor of $Y \rightarrow (G/B)^3$ might be interesting in itself. In general, we have a quite poor understanding of the geometry of these divisors, although we know their classes as Weil divisors, by Theorem 7. In the example treated in Section 7.2, we have a very explicit and nice description of the ramification and branch divisors in terms of projective geometry in \mathbb{P}^2 . Moreover, we observe that they are irreducible.

Finally, we give in Section 7.4 examples in type B that show that the Levi-movability assumption in Theorem 4 is necessary.

Our methods do not extend easily to the case when the multiplicity $c_{I,J}^K$ is more than 2. Indeed, a key point in our arguments is that any finite degree 2 morphism is cyclic, allowing to compute the pushforward of the structural sheaf. Example 4 shows that the situation is deeply different when $c_{I,J}^K > 2$. Maybe in the case of multiplicity 3, [Mir85] could be of some help. It might also be helpful considering a deformation of Π which gives a cyclic covering. Another natural possibility would be to study directly the combinatorics as done in [KP11] with the new understanding we have of the geometry involved.

Acknowledgement. The authors are partially supported by the French National Agency (Project GeoLie ANR-15-CE40-0012).

CONTENTS

1.	Introduction	1
2.	Main results	4
2.1.	Notation	4
2.2.	Our reduction result for the branching problem	6
2.3.	A formula for the branch divisor class	7
3.	Preliminaries in Algebraic Geometry	9
3.1.	Recollections on Intersection theory	9
3.2.	Canonical, ramification and branch divisors	10

3.3.	Pushforward of the structure sheaf under a finite morphism of degree 2	10
3.4.	Generically finite morphisms of degree 2	11
4.	Preliminaries on Schubert varieties	12
4.1.	Reminder on Schubert varieties	12
4.2.	Inversion sets	12
4.3.	Relative Schubert varieties and their Chow classes	14
4.4.	Relative Bott-Samelson varieties	15
4.5.	Canonical divisor of a relative Schubert variety	16
5.	Proof of the main theorems	17
5.1.	Chow class of $Y(\mathfrak{v})$ for any \mathfrak{v}	17
5.2.	Canonical class of $Y(\mathfrak{v})$	19
5.3.	Computation of the class of the branch divisor	21
5.4.	Proof of our reduction formula	23
6.	Properties of the branch divisor class	24
6.1.	The Belkale-Kumar product	24
6.2.	The branch divisor class in the Levi-movable case	25
7.	Examples	26
7.1.	Type A	26
7.2.	A detailed example in $\text{Gr}(3, 6)$	27
7.3.	A bigger example	30
7.4.	Examples in type B	32
Appendix A. Appendix: puzzles		33
A.1.	Some computations	33
A.2.	The puzzles for $c_{\text{Gr}(3,6)}(J, J, J) = 2$ with $J = \{2, 4, 6\}$	34
A.3.	The puzzles for $c_{\text{Gr}(6,9)}(I, I, I) = 3$ with $I = \{2, 3, 5, 6, 8, 9\}$	34
A.4.	The puzzles for $c_{\mathcal{F}(3,6;9)}(\omega, \omega, \omega) = 6$ with $\omega = (\{3, 6, 9\} \subset \{2, 3, 5, 6, 8, 9\})$	34
References		36

2. MAIN RESULTS

2.1. **Notation.** In this section, we introduce most of our needed notation.

2.1.1. *Algebraic groups.* Come back to a general inclusion $G \subset \hat{G}$, or even more generally a finite morphism $G \rightarrow \hat{G}$ of connected reductive algebraic groups. We also assume that G and \hat{G} are simply connected.

Fix a maximal torus T and a Borel subgroup B such that $T \subset B$. Let \hat{B} be a Borel subgroup of \hat{G} containing B and let \hat{T} be a maximal torus of \hat{B} containing T . We denote by $X(T)$ and $X(\hat{T})$ the groups of characters of T and \hat{T} respectively, and by $X(T)^+$ and $X(\hat{T})^+$ the subsets of dominant weights.

Given two dominant weights ζ and $\hat{\zeta}$ for G and \hat{G} respectively, we denote by V_ζ and $V_{\hat{\zeta}}$ the highest weight modules of G and \hat{G} respectively. In this paper, the following integers are what we are mostly interested in:

$$m_{G \subset \hat{G}}(\zeta, \hat{\zeta}) = \dim(V_\zeta \otimes V_{\hat{\zeta}})^G,$$

where the exponent G means subset of G -invariant vectors. Indeed, they encode the branching law since

$$V_{\hat{\zeta}} = \bigoplus_{\zeta \in X(T)^+} \mathbb{C}^{m_{G \subset \hat{G}}(\zeta, \hat{\zeta})} \otimes V_\zeta^*,$$

as G -modules, where V_ζ^* denotes the dual G -module of V_ζ .

We choose a one-parameter subgroup τ of T , which is also a one-parameter subgroup of \hat{T} , and thus defines parabolic subgroups $P = P(\tau) \subset G$ and $\hat{P} = \hat{P}(\tau) \subset \hat{G}$. We thus have the following inclusions:

$$(8) \quad \begin{array}{ccccccc} T & \hookrightarrow & B & \hookrightarrow & P & \hookrightarrow & G \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \hat{T} & \hookrightarrow & \hat{B} & \hookrightarrow & \hat{P} & \hookrightarrow & \hat{G} \end{array}$$

Denote by $\iota : G/P \rightarrow \hat{G}/\hat{P}$ the inclusion morphism. We denote by L the Levi factor of P containing T , and similarly for \hat{L} . We denote by S the neutral component of the center of L . Given a character γ of T or \hat{T} , we denote by $\gamma|_S$ its restriction to the torus S .

Denote by W and W_P the Weyl groups of G and P respectively, and by W^P the set of minimal length representatives of the cosets in W/W_P , and similarly \widehat{W} , $W^{\hat{P}}$ and $W_{\hat{P}}$.

Consider the group $\mathbb{G} = G \times \hat{G}$ and use bolded letters to refer to this group. Namely $\mathbb{T} = T \times \hat{T}$, $\mathbb{B} = B \times \hat{B}$, $\mathbb{P} = P \times \hat{P}$, $\mathbb{L} = L \times \hat{L}$ and $\mathbb{W} = W \times \widehat{W}$. The set Φ of roots of \mathbb{G} is

$$\Phi = (\Phi \times \{0\}) \cup (\{0\} \times \hat{\Phi}),$$

where Φ and $\hat{\Phi}$ denote the sets of roots for G and \hat{G} respectively.

Let Δ , $\hat{\Delta}$ and $\mathbb{\Delta}$ be the sets of simple roots for G , \hat{G} and \mathbb{G} respectively. For $\alpha \in \mathbb{\Delta}$, we denote by ϖ_α the associated fundamental weight. Let $\rho_{\mathbb{G}}$ (resp. $\rho_{\mathbb{L}}$) denote the half sum of positive roots of \mathbb{G} (resp. \mathbb{L}). Set also $\rho^{\mathbb{L}} = \rho_{\mathbb{G}} - \rho_{\mathbb{L}}$. Similarly, we use ω_α , $\rho_{\hat{G}}$...

2.1.2. *Schubert classes.* Let $p : G/P \rightarrow \text{Spec}(\mathbb{C})$ be the structure morphism, and denote, for ξ in the Chow group $A^*(G/P)$,

$$(9) \quad \chi_{G/P}(\xi) = p_* \xi \in A^0(\text{Spec}(\mathbb{C})) \simeq \mathbb{Z}.$$

Since G/P is rationally connected, $A^{\dim(G/P)} = \mathbb{Z}[pt]$ and $\chi_{G/P}(\xi)$ is the coefficient of $[pt]$ in the homogeneous part of degree $\dim(G/P) - \ell(\xi)$ of ξ .

For $v \in W^P$, τ_v denotes the class of the Schubert variety $\overline{BvP/P}$ in $A^{\dim(G/P) - \ell(v)}(G/P)$. Let w_0 be the longest element of W and $w_{0,P}$ be the longest element of W_P . Poincaré duality takes the nice form

$$(10) \quad \chi_{G/P}(\tau_v \cdot \tau_w) = \delta_{v^\vee, w} \text{ with } v^\vee := w_0 v w_{0,P}.$$

Similarly, $(\tau_{\hat{v}})_{\hat{v} \in W^{\hat{P}}}$ denotes the Schubert basis of $A^*(\hat{G}/\hat{P})$. When $P = B$ and $u \in W$, we denote by $\sigma_u \in A^*(G/B)$ the Chow class of $\overline{BuB/B} \subset G/B$. Moreover, we define $\sigma^u := \sigma_{w_0 u}$ and $\tau^v := \tau_{w_0 v w_{0,P}}$.

Let

$$\begin{aligned} \delta : G/P &\longrightarrow \mathbb{G}/\mathbb{P} \\ gP/P &\longmapsto (gP/P, g\hat{P}/\hat{P}) \end{aligned}$$

denote the small diagonal map and consider the pullback $\delta^* : A^*(\mathbb{G}/\mathbb{P}) \rightarrow A^*(G/P)$. For $\mathfrak{v} = (v, \hat{v}) \in W^{\mathbb{P}}$, observe that

$$(11) \quad \delta^*(\tau_{\mathfrak{v}}) = \tau_v \cdot \iota^*(\tau_{\hat{v}}).$$

For $\mathfrak{v} = (v, \hat{v}) \in W^{\mathbb{P}} = W^P \times W^{\hat{P}}$ we set

$$(12) \quad c(\mathfrak{v}) = c(v, \hat{v}) := \chi_{G/P}(\delta^*(\tau_{\mathfrak{v}})), \text{ so that } \iota^*(\tau_{\hat{v}}) = \sum_{v \in W^P} c(v, \hat{v}) \tau^v.$$

Since $\tau_{\mathfrak{v}}$ has degree $\dim G/P + \dim \hat{G}/\hat{P} - \ell(\mathfrak{v})$, we have $c(\mathfrak{v}) = 0$ unless $\ell(\mathfrak{v}) = \dim(\hat{G}/\hat{P}) (= \ell(v) + \ell(\hat{v}))$.

2.1.3. *Bruhat orders.* A covering relation in the left weak Bruhat order is a pair (v, w) such that there exists

a simple root α with $v = s_\alpha w$ and $\ell(v) = \ell(w) + 1$. We use the graph $\begin{array}{c} v \\ \alpha \Big| \\ w \end{array}$ to depict such a relation.

A covering relation in the strong Bruhat order is a pair (v, w) such that there exists a positive root β with $v = s_\beta w$ and $\ell(v) = \ell(w) + 1$. We use the graph $\begin{array}{c} v \\ \beta \Big| \\ w \end{array}$ to depict such a relation. For example the graph on the left on Figure 1 is the Bruhat graph of SL_3 .

We also define the (weak) twisted Bruhat graph by labelling the previous edges by $w^{-1}\alpha$ and $w^{-1}\beta$ in place of α and β . We use green color for these twisted labels to avoid any confusion. Hence $\begin{array}{c} v \\ \beta \Big| \\ w \end{array}$ means




FIGURE 1. Bruhat graphs for SL_3

$v = s_\beta w$ and $\begin{array}{c} v \\ \vdots \gamma \\ w \end{array}$ means $v = ws_\gamma$, with in both cases $\ell(v) = \ell(w) + 1$. For $SL_3(\mathbb{C})$ we get the right graph of Figure 1.

We denote by $v \leq w$ the strong Bruhat order and by $v \leq_w w$ the left weak Bruhat order.

2.1.4. *Chevalley formula.* Given $\zeta \in X(T)$, we denote by $\mathcal{L}_{G/B}(\zeta)$ the G -linearized line bundle on G/B such that B acts with weight $-\zeta$ on the fiber over B/B . Similarly, define $\mathcal{L}_{\hat{G}/\hat{B}}(\hat{\zeta})$. For any v in W^P and any character ζ of P , we have the Chevalley formula, proved for example in [Pra05]:

$$(13) \quad c_1(\mathcal{L}_{G/P}(\zeta)) \cup \tau_v = \sum \langle \zeta, \gamma^\vee \rangle \tau_{v'},$$

where the sum runs over the covering relations $\begin{array}{c} v \\ \vdots \gamma \\ v' \end{array}$ in the Bruhat graph of G/P . Here, c_1 denotes the first Chern class.

2.2. **Our reduction result for the branching problem.** In [Res11b], the following result is proved:

Theorem 5. *Let $\mathfrak{v} = (v, \hat{v}) \in W^{\mathbb{P}}$ be such that $c(\mathfrak{v}) = 1$. Let $\zeta \in X(T)^+$ and $\hat{\zeta} \in X(\hat{T})^+$ be such that $v^{-1}(\zeta)|_S + \hat{v}^{-1}(\hat{\zeta})|_S$ is trivial. Then $v^{-1}(\zeta)$ and $\hat{v}^{-1}(\hat{\zeta})$ are dominant weights for L and \hat{L} respectively. Moreover*

$$m_{G\subset\hat{G}}(\zeta, \hat{\zeta}) = m_{L\subset\hat{L}}(v^{-1}(\zeta), \hat{v}^{-1}(\hat{\zeta})).$$

The conclusion of Theorem 5 does not hold when $c(\mathfrak{v}) = 2$. Our reduction result is a modification of this conclusion which holds when $c(\mathfrak{v}) = 2$.

Consider the incidence variety

$$(14) \quad Y(\mathfrak{v}) = \{(x, gB/B, \hat{g}\hat{B}/\hat{B}) : x \in \overline{gBvP/P} \text{ and } \iota(x) \in \overline{\hat{g}\hat{B}\hat{v}\hat{P}/\hat{P}}\} \subset G/P \times \mathbb{G}/\mathbb{B},$$

endowed with its projection $\Pi : Y(\mathfrak{v}) \rightarrow \mathbb{G}/\mathbb{B}$ (in the sequel ι will be omitted and G/P will be considered as a subvariety of \hat{G}/\hat{P}). Consider the ramification divisor class K_Π of Π (see Section 3.2 for details) and the branch divisor class $[B_\Pi] = \Pi_*(K_\Pi)$ in $Cl(\mathbb{G}/\mathbb{B})$ (note that Π is proper since Y is projective). Since \mathbb{G}/\mathbb{B} is smooth, $[B_\Pi]$ is an element of the Picard group and has an expression as

$$[B_\Pi] = \mathcal{L}_{G/B}(\sum_{\alpha \in \Delta} n_\alpha \varpi_\alpha) \otimes \mathcal{L}_{\hat{G}/\hat{B}}(\sum_{\hat{\alpha} \in \hat{\Delta}} n_{\hat{\alpha}} \varpi_{\hat{\alpha}})$$

for some well defined integers n_α and $n_{\hat{\alpha}}$.

Assume now that $c(\mathfrak{v}) = 2$. Then, we prove in Proposition 12 that the integers n_α and $n_{\hat{\alpha}}$ are even. Define θ and $\hat{\theta}$ in $X(T)$ and $X(\hat{T})$ by setting

$$(15) \quad \theta = \sum_{\alpha} \frac{n_{\alpha}}{2} \varpi_{\alpha} \quad \hat{\theta} = \sum_{\hat{\alpha}} \frac{n_{\hat{\alpha}}}{2} \varpi_{\hat{\alpha}}.$$

Theorem 6. *Let $\mathfrak{v} = (v, \hat{v}) \in W^{\mathbb{P}}$ be such that $c(\mathfrak{v}) = 2$. Let $\zeta \in X(T)^+$ and $\hat{\zeta} \in X(\hat{T})^+$ be such that $v^{-1}(\zeta)|_S + \hat{v}^{-1}(\hat{\zeta})|_S$ is trivial. Then*

$$m_{G \subset \hat{G}}(\zeta, \hat{\zeta}) + m_{G \subset \hat{G}}(\zeta - \theta, \hat{\zeta} - \hat{\theta}) = m_{L \subset \hat{L}}(v^{-1}\zeta, \hat{v}^{-1}\hat{\zeta}).$$

Theorem 6 is proved in Section 5.4, using the same idea as in [Res11b] and a formula for the sheaf $\Pi_* \mathcal{O}_{Y(\mathfrak{v})}$ which is proved in Section 3.3 for any generically finite morphism of degree 2.

Consider now the case when $\hat{G} = G \times G$ corresponding to the tensor product decomposition. Then $\hat{P} = P \times P$. Assume moreover that P is cominuscle. Then Theorem 6 can be improved by expressing $m_{G \subset G \times G}(\zeta, \hat{\zeta})$ as an alternating sum of multiplicities for the tensor product for L as in Theorem 4 (see Corollary 37).

2.3. A formula for the branch divisor class. In this section, P and \hat{P} are any parabolic subgroups of G and \hat{G} respectively as in Diagram (8), but not necessarily associated to a given one-parameter subgroup of T . We let $K_{\Pi} := K_{Y(\mathfrak{v})} - \Pi^* K_{\mathbb{G}/\mathbb{B}}$. If $c(\mathfrak{v}) \neq 0$, it is the ramification divisor class of Π , and we still call it in this way when $c(\mathfrak{v}) = 0$ although Π is contractant in this case. Under the assumption $c(\mathfrak{v}) = 2$, $\Pi_* K_{\Pi}$ is twice the class of the branch divisor. More details are provided in Sections 3.2 and 3.3. For $\beta \in \Phi$, we set:

$$(16) \quad \mathfrak{h}(\beta) = \langle \rho_{\mathbb{G}}, \beta^{\vee} \rangle$$

Theorem 7. *Let $\mathfrak{v} \in W^{\mathbb{P}}$ such that $\ell(\mathfrak{v}) = \dim(\hat{G}/\hat{P})$. Consider the map*

$$\Pi : Y(\mathfrak{v}) \longrightarrow \mathbb{G}/\mathbb{B},$$

and its ramification divisor class $K_{\Pi} \in \text{Cl}(Y)$. Then, in $\text{Cl}(\mathbb{G}/\mathbb{B})$, we have

$$\Pi_* K_{\Pi} = \sum_{\mathfrak{v}'} \left(c_1(\mathcal{L}(2\rho_{|T}^{\hat{L}})) \cdot \delta^*(\tau_{\mathfrak{v}'}) + 2c(\mathfrak{v}) - \sum_{\mathfrak{v}''} (\mathfrak{h}(\mathfrak{d}) + 1)c(\mathfrak{v}'') \right) \mathcal{L}_{\mathbb{G}/\mathbb{B}}(\varpi_{\mathfrak{a}}),$$

where the sums over \mathfrak{v}' and \mathfrak{v}'' respectively run over the covering relations $\begin{array}{c} \mathfrak{v}' \\ \alpha \downarrow \\ \mathfrak{v} \end{array}$ and $\begin{array}{c} \mathfrak{v}' \\ \mathfrak{d} \downarrow \\ \mathfrak{v}'' \end{array}$ in the Bruhat

graph of \mathbb{G}/\mathbb{P} , with $\mathfrak{a} \in \Delta$ and $\mathfrak{d} \in \Phi^+(\mathbb{G})$ is the twisted label. Here c_1 denotes the first Chern class.

The first term $c_1(\mathcal{L}(2\rho_{|T}^{\hat{L}})) \cdot \delta^*(\tau_{\mathfrak{v}'})$ is an integer as an element of $A^{\dim(G/P)}(G/P) \simeq \mathbb{Z}$. We emphasise that it is an intersection number: see (11).

Let's outline the proof of this result, which is spread over several Sections of the paper. The term $\Pi_* \Pi^* K_{\mathbb{G}/\mathbb{B}}$ is computed via the projection formula. To compute $\Pi_* K_{Y(\mathfrak{v})}$, we observe that projection on G/P expresses $Y(\mathfrak{v})$ as a fibered product of two relative Schubert varieties. Our first task is to express the canonical class of a relative Schubert variety, extending previous works on the absolute case (see e.g. [Ram87, Theorem 4.2] and [Per07, Proposition 4.4]). This is done using the fact that (relative) Schubert varieties are normal and Cohen-Macaulay and using the Bott-Samelson resolution (Proposition 27).

Then, using the fibered product structure of $Y(\mathfrak{v})$, Proposition 32 expresses the canonical class of $Y(\mathfrak{v})$. We get the expression (47) of K_{Π} as a sum of line bundles coming from \mathbb{G}/\mathbb{B} and G/P , and Weil divisors of the form $[Y(\mathfrak{v}')] for elements \mathfrak{v}' covered by \mathfrak{v} in the Bruhat order. To push forward to \mathbb{G}/\mathbb{B} those Weil divisors as well as the line bundle coming from G/P , we express the class of $Y(\mathfrak{v})$ in the Chow ring of $\mathbb{G}/\mathbb{B} \times G/P$, thanks to a result by Brion valid for any orbit closure of a spherical subgroup, see Propositions 20 and 29.$

Notation 1. In the case when $G \subset \hat{G} = G^{n-1}$ and $\mathbb{G}/\mathbb{P} = (G/P)^n$ for some integer $n \geq 2$, write c_1^α for the coefficient of $\Pi_*(K_\Pi)$ at $\mathcal{L}_{\mathbb{G}/\mathbb{B}}(\varpi_\alpha, 0, \dots, 0)$. We let $(\zeta, \hat{\zeta}) = (\zeta_1, \zeta_2, \dots, \zeta_n)$, $\mathfrak{v} = (v_1, v_2, \dots, v_n)$, $(\theta, \hat{\theta}) = (\theta^1, \theta^2, \dots, \theta^n)$ and $m_{G \subset \hat{G}}(\zeta, \hat{\zeta}) = m_G(\zeta_1, \zeta_2, \dots, \zeta_n)$.

Even more specifically, we call “tensor product case” the case when $n = 3$, since then $m_G(\zeta_1, \zeta_2, \zeta_3)$ is the multiplicity of $V_{\zeta_1}^*$ in $V_{\zeta_2} \otimes V_{\zeta_3}$. In this case, we set $\Gamma(G) = \Gamma(G \subset \hat{G})$.

Corollary 8. Let $n \geq 2$ be an integer. Assume that $\hat{G} = G^{n-1}$ and $\hat{P} = P^{n-1}$. Let $\mathfrak{v} = (v_1, \dots, v_n) \in (W^P)^n$ such that $\ell(\mathfrak{v}) = (n-1) \dim(G/P)$ and $c(\mathfrak{v}) \neq 0$. Consider the map

$$\Pi : Y(\mathfrak{v}) \rightarrow \mathbb{G}/\mathbb{B},$$

and its ramification divisor class $K_\Pi \in \text{Cl}(Y)$. Fix $\alpha \in \Delta$ and denote by c_1^α the coefficient of $\Pi_* K_\Pi$ along

$\mathcal{L}((\varpi_\alpha, 0, \dots, 0))$. If $\begin{array}{c} v'_1 \\ \alpha \\ v_1 \end{array}$ is a covering relation in the Bruhat graph of G/P then

$$c_1^\alpha = (n-1)c_1(\mathcal{L}(2\rho^L))\tau_{v'_1}\tau_{v_2}\dots\tau_{v_n} + 2c(\mathfrak{v}) - \sum_{\mathfrak{v}''} (\hat{h}(\mathfrak{v}) + 1)c(\mathfrak{v}''),$$

where $\mathfrak{v}' = (v'_1, v_2, \dots, v_n)$ and the sum runs over the covering relations $\begin{array}{c} \mathfrak{v}' \\ \vdots \\ \mathfrak{v}'' \end{array}$. Otherwise $c_1^\alpha = 0$.

Remark 1. In the case when moreover G/P is cominusculc, strong and weak Bruhat order coincide, so the

covering relations $\begin{array}{c} \mathfrak{v}' \\ \vdots \\ \mathfrak{v}'' \end{array}$ may be replaced by the covering relations $\begin{array}{c} \mathfrak{v}' \\ | \\ \mathfrak{v}'' \end{array}$.

Proof of Corollary 8: We only need to remark that since $\hat{G}/\hat{P} = (G/P)^{n-1}$, we have $\mathcal{L}(2\rho_{|\hat{T}}^{\hat{L}}) = (n-1)\mathcal{L}_{G/P}(2\rho^L)$ and $\delta^*(\tau_{\mathfrak{v}'}) = \tau_{v'_1}\tau_{v_2}\dots\tau_{v_n}$. \square

Theorem 7 is proved in Section 5.3. Here we make one comment. Let $\mathfrak{v}' = (v', \hat{v}') \in W^{\mathbb{P}}$ such that $\ell(\mathfrak{v}') = \dim(\hat{G}/\hat{P}) + 1$ like in the first sum of Theorem 7. Set

$$C(\mathfrak{v}') = g \cdot \overline{Bv'P/P} \cap \iota^{-1}(\hat{g} \cdot \overline{B\hat{v}'P/P}),$$

for general elements $g \in G, \hat{g} \in \hat{G}$. Then $C(\mathfrak{v}') \subset G/P$ is a curve. The degree of the pullback of the anticanonical bundle $\mathcal{L}(2\rho_{|\hat{T}}^{\hat{L}})$ on $C(\mathfrak{v}')$ is the first term appearing in the expression of $\Pi_* K_\Pi$. By Chevalley’s formula applied to $c_1(\mathcal{L}(2\rho_{|\hat{T}}^{\hat{L}})) \cdot \tau_{v'_1}$, we have an explicit formula

$$(17) \quad c_1(\mathcal{L}(2\rho_{|\hat{T}}^{\hat{L}})) \cdot \delta^*(\tau_{\mathfrak{v}'}) = \sum_{\mathfrak{v}''} \langle 2\rho_{|\hat{T}}^{\hat{L}}, \gamma^\vee \rangle c(\mathfrak{v}''),$$

where the sum runs over the covering relations $\begin{array}{c} \mathfrak{v}' \\ \vdots \\ \mathfrak{v}'' \end{array}$ with $\gamma \in \Phi^+(G)$.

If $c(\mathfrak{v}) = 1$ then Π is birational so $\Pi_*(K_\Pi) = 0$. Theorem 7 gives a funny but combinatorially involved necessary condition for the equality $c(\mathfrak{v}) = 1$. In Example 3, we check a family of examples.

At the opposite, if $c(\mathfrak{v}) \geq 2$ then $\Pi_*(K_\Pi) \neq 0$. Indeed, if $\Pi_*(K_\Pi) = 0$ then by removing the singular locus of $Y(\mathfrak{v})$ and the ramification divisor R_Π , we get an unramified cover onto the complement in \mathbb{G}/\mathbb{B} of a codimension two subvariety. But, this complement is simply connected as \mathbb{G}/\mathbb{B} is by the Bruhat decomposition. This is a contradiction.

Assume now that $c(\mathfrak{v}) = 0$. In this case, Π is contractant, thus $\Pi_* \Pi^* K_{\mathbb{G}/\mathbb{B}} = 0$. However, we have no information on $\Pi_* K_{Y(\mathfrak{v})}$, and this can be non zero: see Example 2.

Example 1. Let $G = \mathrm{SL}_3$, $\hat{G} = G \times G$, $P = B$, $\hat{P} = B \times B$, and $\mathfrak{v} = (s_2s_1, s_2s_1, s_1s_2)$. Then $\Pi_*K_\Pi = 0$.

Proof. We use Corollary 8 to check this. Recall that the Bruhat graph of SL_3 is depicted in Figure 1, and that the Poincaré dual of τ_{s_i} is $\tau_{w_0s_i} = \tau_{s_i s_{3-i}}$. Fix first $\mathfrak{w} = (\alpha_1, 0, 0)$, which implies $\mathfrak{v}' = (s_1s_2s_1, s_2s_1, s_1s_2)$. Then $2c_1(\mathcal{L}(2\rho))\tau_{v'_1} \dots \tau_{v'_3} + 2c(\mathfrak{v}) = 2c_1(\mathcal{L}(2\rho))(\tau_{s_1} + \tau_{s_2}) + 2 \times 1 = 2\langle 2\rho, \alpha_1^\vee + \alpha_2^\vee \rangle + 2 = 10$. The possibilities for \mathfrak{v} with $c(\mathfrak{v}'') \neq 0$ are $(0, \alpha_1 + \alpha_2, 0)$, $(0, 0, \alpha_1 + \alpha_2)$, $(\alpha_1, 0, 0)$ and $(\alpha_2, 0, 0)$. In these cases $c(\mathfrak{v}'') = 1$ and $\hat{h}(\mathfrak{v}) = 2$ twice and $\hat{h}(\mathfrak{v}) = 1$ twice. We get $c_1^{\alpha_1} = 0$ as expected.

Fix now $\mathfrak{w} = (0, 0, \alpha_2)$ with $\mathfrak{v}' = (s_2s_1, s_2s_1, s_2s_1s_2)$. Then $2c_1(\mathcal{L}(2\rho))\tau_{v'_1} \dots \tau_{v'_3} + 2c(\mathfrak{v}) = 2c_1(\mathcal{L}(2\rho))(\tau_{s_1}) + 2 \times 1 = 6$. The possibilities for \mathfrak{v} with $c(\mathfrak{v}'') \neq 0$ are $(\alpha_1, 0, 0)$, $(0, \alpha_1, 0)$ and $(0, 0, \alpha_1)$ and give $3 \times 2 = 6$ for the sum.

The remaining cases are either trivial or obtained by exchanging v_1 and v_2 . \square

Example 2. Let $G = \mathrm{SL}_3$, $\hat{G} = G \times G$, $P = B$, $\hat{P} = B \times B$, and $\mathfrak{v} = (s_2s_1, s_1s_2s_1, s_1)$. We have $c_1^{\alpha_1} = -2$.

Proof. For $c_1^{\alpha_1}$, with the notation of Corollary 8, we have $\mathfrak{v}' = (s_1s_2s_1, s_1s_2s_1, s_1)$ and the elements \mathfrak{v}'' such that $c(\mathfrak{v}'') \neq 0$ are the three triples $(s_1s_2s_1, s_1s_2s_1, e)$, $(s_1s_2s_1, s_1s_2, s_1)$, $(s_1s_2, s_1s_2s_1, s_1)$. The right hand-side of the formula given in Corollary 8 is thus $2c_1(\mathcal{L}(2\rho))\tau_{s_1} + 2c(\mathfrak{v}) - 3 \times (1+1) = 4+0-6 = -2$. \square

We now consider a family of examples again in the birational case:

Example 3. Let $\hat{G} = G \times G$ and $\hat{P} = P \times P$. Let $v \in W^P$ be arbitrary and let $\mathfrak{v} = (v, v^\vee, w^P)$. Then we have $\Pi_*R_\Pi = 0$.

Proof. We check that Corollary 8 gives $\Pi_*R_\Pi = 0$. Up to exchanging v and v^\vee , the only case to consider is when $\mathfrak{w} = (\alpha, 0, 0)$ with $\alpha \in \Delta$, $s_\alpha v \in W^P$ and $\ell(s_\alpha v) = \ell(v) + 1$. Set $\gamma_0 = -v^{-1}\alpha$ and $\mathfrak{v}' = (s_\alpha v, v^\vee, w^P)$.

Consider the Chevalley formula (13) for $c_1(\mathcal{L}(2\rho^L)) \cdot \tau_{\mathfrak{v}'}$. The only summand surviving after multiplication by $\tau_{s_\alpha v}$ is $\langle 2\rho^L, \gamma \rangle \tau_{(s_\alpha v)}$. Moreover, $c(\mathfrak{v}'') = 1$ for this term and $\gamma = w_{0,P}(\gamma_0)$. Finally $2c_1(\mathcal{L}(2\rho))\tau_{v'_1} \dots \tau_{v'_3} + 2c(\mathfrak{v}) = \langle 4\rho^L, w_{0,P}(\gamma_0^\vee) \rangle + 2 = \langle 4\rho^L, \gamma_0^\vee \rangle + 2$.

The only case with $c(\mathfrak{v}'') \neq 0$ and \mathfrak{v} in the first factor is $\mathfrak{v} = (\gamma_0, 0, 0)$. Its contribution in the sum is $\langle \rho, \gamma_0^\vee \rangle + 1$. Similarly, with \mathfrak{v} in the second factor, we get $\langle w_{0,P}(\rho), \gamma_0^\vee \rangle + 1$.

Assume now that γ is in the third copy. It is a descent of w^P , hence an element of $\Delta - \Delta(L)$. We have $\tau_{w^P s_\gamma} = c_1(\mathcal{L}(\varpi_\gamma))$, so the associated coefficient $c(\mathfrak{v}'')$ is $\langle \varpi_\gamma, \gamma_0^\vee \rangle$ by the Chevalley formula. The term $\hat{h}(\gamma) + 1$ equals $\langle \rho, \gamma^\vee \rangle + 1 = 2$. The contribution of these terms is $\sum_{\gamma \in \Delta - \Delta(L)} 2\langle \varpi_\gamma, \gamma_0^\vee \rangle = 2\langle \rho - \rho_L, \gamma_0^\vee \rangle$.

Finally,

$$c_1^\alpha = 4\langle \rho^L, \gamma_0^\vee \rangle + 2 - (\langle \rho, \gamma_0^\vee \rangle + 1) - (\langle w_{0,P}(\rho), \gamma_0^\vee \rangle + 1) - 2\langle \rho - \rho_L, \gamma_0^\vee \rangle$$

which vanishes, since $\rho^L = \rho - \rho_L$ and $w_{0,P}(\rho) = \rho^L - \rho_L$. \square

In Section 7, we give several examples of Theorem 7 in the case $c(\mathfrak{v}) = 2$, which is the case we are mainly interested in. Let us now give an example with $c(\mathfrak{v}) = 3$:

Example 4. In $\mathrm{Gr}(4, 8)$, let $u = [35681247] = (s_2s_3s_5s_4)^\vee$, $v = [24681357] = (s_2s_4s_6s_3s_5s_4)^\vee$ and $w = [24681357] = (s_2s_4s_3s_6s_5s_4)^\vee$. Then $c(u, v, w) = c_{211,321}^{4321} = 3$. We computed with Theorem 7 that

$$[B_\Pi] = \mathcal{L}_{(G/B)^3}(4(\varpi_2 + \varpi_4 + \varpi_6), 4(\varpi_2 + \varpi_4 + \varpi_6), 4\varpi_3 + 6\varpi_6).$$

In particular, $[B_\Pi]$ is not divisible by 3, which shows that the following Lemma 10 and Proposition 12 don't have obvious analogues for morphisms of degree 3.

3. PRELIMINARIES IN ALGEBRAIC GEOMETRY

3.1. Recollections on Intersection theory. We now recall useful notions in Intersection Theory and give some details on degree two morphisms and the associated ramification and branch divisors.

Let $f : Y \rightarrow X$ be a dominant morphism between irreducible varieties of the same dimension. We say that f is *generically finite*. The *degree of f* is defined to be $\deg(f) = [\mathbb{C}(Y) : \mathbb{C}(X)]$.

Let X be a quasi-projective irreducible variety. We denote by $\mathrm{Cl}(X)$ the Weil divisor class group, by $\mathrm{CaCl}(X)$ the Cartier divisor class group and by $\mathrm{Pic}(X)$ the Picard group. By [Har77, Prop. II.6.15], we have a canonical isomorphism $\mathrm{CaCl}(X) \simeq \mathrm{Pic}(X)$, mapping D on $\mathcal{O}(D)$. There is also a morphism $c_1 : \mathrm{Pic}(X) \rightarrow \mathrm{Cl}(X)$ (see [Ful84, p. 30]) that is neither injective nor surjective. Nevertheless, if X is

normal, c_1 is injective and $\text{CaCl}(X) = \text{Pic}(X)$ can be seen as a subgroup of $\text{Cl}(X)$. Moreover if X is locally factorial (e.g. is smooth) then this morphism is an isomorphism.

We also consider the Chow ring $A^*(X)$ and identify $\text{Cl}(X)$ with $A^1(X)$. Given two irreducible subvarieties Z_1 and Z_2 that intersect transversally in X , we have

$$(18) \quad [Z_1 \cap Z_2] = [Z_1] \cdot [Z_2] \text{ in } A^*(X).$$

Here $[Z_1 \cap Z_2]$ denotes the sum of the classes of the irreducible components of $Z_1 \cap Z_2$.

3.2. Canonical, ramification and branch divisors. Set $n = \dim(X)$. If X is smooth, the canonical divisor class $K_X \in \text{Pic}(X)$ on X is defined to be the line bundle $\wedge^n T^*X$. Assume that X is normal. Let X_{reg} denote the regular locus of X and TX_{reg} denote its tangent bundle. Then $\wedge^n T^*X_{\text{reg}}$ is a line bundle \mathcal{L} on X_{reg} . The first Chern class $c_1(\mathcal{L}) \in \text{Cl}(X_{\text{reg}}) = \text{Cl}(X)$ is called the canonical divisor class of X and denoted by K_X . The canonical sheaf \mathcal{K}_X is defined to be the pushforward of the sheaf $U \mapsto \mathcal{L}(U)$ by the inclusion $X_{\text{reg}} \subset X$.

Let $f : Y \rightarrow X$ be a generically finite morphism between irreducible varieties. Assume moreover that Y is normal and X is smooth. The determinant of the restriction of f to the smooth locus Y_{reg} of Y define a Cartier divisor R_f in Y_{reg} . By construction, the associated line bundle $\mathcal{O}(R_f) \in \text{Pic}(Y_{\text{reg}})$ is $K_f := K_Y - f^*K_X$. Its closure, still denoted by R_f has a class $[R_f]$ in $\text{Cl}(Y)$ called ramification divisor class of f . We have

$$(19) \quad [R_f] = K_Y - f^*K_X \in \text{Cl}(Y),$$

where $f^* : \text{Pic}(X) \rightarrow \text{Pic}(Y)$ is the pullback of line bundles. Observe that R_f is an effective Weil divisor. We write R_f as $R_f = \sum_{i \in I} a_i E_i + \sum_{j \in J} c_j R_j$, with prime Weil divisors E_i, R_j such that the divisors E_i are contracted and the divisors R_j are not, and positive integers a_i, c_j . We define the Weil divisors

$$(20) \quad E_f = \sum_{i \in I} a_i E_i, \quad R_f^1 = \sum_{j \in J} c_j R_j.$$

We define the branch divisor $B_f := f_* R_f^1 = \sum_{j \in J} c_j \deg(f|_{R_j}) f(R_j)$, and its class $[B_f] = f_*[R_f] = \sum_{j \in J} c_j \deg(f|_{R_j}) [f(R_j)] \in \text{Cl}(X)$ (since f is proper, $f_* : \text{Cl}(Y) \rightarrow \text{Cl}(X)$ is well defined).

3.3. Pushforward of the structure sheaf under a finite morphism of degree 2. We consider a finite morphism $f : S \rightarrow X$ and study $f_*\mathcal{O}_S$. The next lemmas are well-known to specialists, see for example [Laz80, Lemma 1.1] or [Par91]. We provide details for the singular case we are interested in.

As it is explained in [BK05, p.9], we have:

Lemma 9. *Let $f : S \rightarrow X$ be a finite surjective morphism of degree d and assume that X is normal. Then there exists a splitting $f_*\mathcal{O}_S = \mathcal{O}_X \oplus L$ with L a coherent sheaf on X .*

We now consider the degree 2 case and exploit the fact that f is a cyclic cover.

Lemma 10. *Let $f : S \rightarrow X$ be a finite surjective morphism of degree 2 and assume that S is normal and X is normal and locally factorial. Then, in the splitting $f_*\mathcal{O}_S = \mathcal{O}_X \oplus L$ as above, L is a line bundle, and $[L^{\otimes 2}] = -[B_f] \in \text{Pic}(X) \simeq \text{Cl}(X)$. Moreover, B_f and R_f are reduced Weil divisors. As Cartier divisors, we have $f^*\mathcal{O}_X(B_f) = \mathcal{O}_Y(2R_f)$.*

Proof. Let us restrict ourselves over an affine open subset $U \subset X$ and set $A = \mathbb{C}[U]$ and $B = \mathbb{C}[f^{-1}(U)]$.

Let $b \in B \setminus \text{Frac}(A)$. The ring B being integral over A , there exists a monic polynomial P in $A[T]$ of minimal degree such that $P(b) = 0$. Then P is irreducible in $A[T]$. By assumption, A is unique factorization domain. Then, P is also irreducible as an element of $\mathbb{C}(X)[T]$, by [Eis95, Exercise 3.4.c]. It follows that P is the minimal polynomial of $b \in \mathbb{C}(S)$ over $\mathbb{C}(X)$. But $[\mathbb{C}(S) : \mathbb{C}(X)] = 2$, hence P has degree 2.

Write $P = T^2 + a_1T + a_2$ with a_1 and a_2 in A . Up to changing b by $b - \frac{a_1}{2}$, one may assume that $a_1 = 0$. Assume that $-a_2 = t^2c$ in A . Set $b = tb'$. Then $b'^2 - c = 0$, and $b' \in \mathbb{C}(X)$ is integral over A . By normality of S , this implies that $b' \in B$. In particular, up to changing b , one may assume that the minimal polynomial of b is $T^2 - a$ with some square free element $a \in A$.

We claim that $B = A \oplus Ab$. Let $y = \alpha + \beta b \in B$ with $\alpha, \beta \in \text{Frac}(A)$. The matrix of $x \mapsto xy$ is $\begin{pmatrix} \alpha & a\beta \\ \beta & \alpha \end{pmatrix}$. Then $\text{Tr}(y) = 2\alpha$ and $N(y) = \alpha^2 - a\beta^2$ are integral over A . Hence $\alpha \in A$ and $a\beta^2 \in A$. Since A is factorial and a is square free, this implies that $\beta \in A$. This proves the claim.

It follows that $f^{-1}(U) = \text{Spec}(A[t]/(t^2 - a))$ and f corresponds to the inclusion $A \subset A[t]/(t^2 - a)$. Then L is the kernel of Tr , namely $tA \subset A[t]/(t^2 - a)$. This is a free A -module. This proves that L is a line bundle.

A local computation as in [BPVdV84, Lemma 16.1] shows that R_f is locally defined by t , so it is reduced. We get $f^*\mathcal{O}_X(B_f) = \mathcal{O}_S(2R_f)$, since these locally free sheaves are generated by $a = t^2$. Moreover, a being square-free, is an equation of $f(R_f) \cap U$ in $\mathbb{C}[U]$. It follows that $L^{\otimes 2}(U)$ identifies with $aA = \mathcal{O}(-B_f)(U) \subset \mathbb{C}(X)$. This proves that $L^{\otimes 2} \simeq \mathcal{O}(-B_f)$. Finally, f restricts to an isomorphism from the hypersurface defined by the equation $t = 0$ to $B_f \cap U$, proving that the Weil divisor B_f is reduced. \square

Remark 2. *The degree 2 is a special case since any finite morphism of degree 2 is a cyclic covering. Already in degree 3, this is no longer true, and the above Lemma fails: see Example 4 below.*

3.4. Generically finite morphisms of degree 2. We now consider the case of a degree 2 map, but not necessarily finite: we let $f : Y \rightarrow X$ be a generically finite morphism of degree 2, with X and Y projective,

Y irreducible normal, and X normal and locally factorial. We let $Y \begin{array}{c} \xrightarrow{p} S \xrightarrow{h} X \\ \searrow f \end{array}$ be the Stein

factorization of f .

Lemma 11. *With the above notation, assume we have an irreducible divisor $C_S \subset S$. Then there is a unique irreducible component C_Y of $p^{-1}(C_S)$ which maps onto C_S , the restriction $C_Y \xrightarrow{p} C_S$ is birational, and we have the equivalence*

$$C_Y \subset R_f \iff C_S \subset R_h.$$

Proof. Since p is surjective, there is an irreducible component C_Y of $p^{-1}(C_S)$ mapping onto C_S . This component is unique and $C_Y \xrightarrow{p} C_S$ is birational since p has connected fibers. Since Y, S and X are normal, for a generic point $y \in C_Y$, Y, S, X are normal and locally factorial at $y, p(y), f(y)$, respectively. Since $C_Y \xrightarrow{p} C_S$ is dominant, $T_y p$ is invertible, by Zariski's Main Theorem again. Therefore $T_y f$ will be degenerate if and only if $T_{p(y)} h$ is degenerate. In other words, $C_Y \subset R_f$ if and only if $C_S \subset R_h$. \square

In the next proposition, we use the prime divisors E_i and the divisor R_f^1 introduced in Section 3.2.

Proposition 12. *Let $f : Y \rightarrow X$ be a generically finite morphism of degree 2, with X and Y projective, Y irreducible normal, and X normal and locally factorial. Then there is a line bundle L on X such that $f_*\mathcal{O}_Y = \mathcal{O}_X \oplus L$. Moreover, in $\text{Pic}(X) = \text{Cl}(X)$, we have $2L + f_*(K_Y) - f_*f^*(K_X) = 0$ and $B_f = B_h$ is reduced. As Weil divisors on Y , we have*

$$(21) \quad f^*\mathcal{O}_X(B_f) = 2R_f^1 + \sum_{i \in I} b_i E_i$$

for some non-negative integers b_i .

Proof. We keep the notation of the beginning of the section, and we want to prove that $p_*R_f = R_h$. By Lemma 10, R_h is reduced: $R_h = \sum_{C_S \subset R_h} C_S$. By Lemma 11, for a component C_S of R_h , there is a unique component C_Y of R_f such that $p(C_Y) = C_S$. Since $C_Y \xrightarrow{p} C_S$ is birational, we get $p_*C_Y = C_S$. Moreover, p is an isomorphism at a generic point of C_Y , so C_Y is a reduced component of R_f . For a contracted component C'_Y of R_f , we have $p_*[C'_Y] = 0$. Thus $p_*R_f = \sum_{C_Y \subset R_f} p_*C_Y = \sum_{C_S \subset R_h} C_S = R_h$. Applying h_* and using Lemma 10, we get that $B_f = f_*R_f = h_*R_h = B_h$ is reduced.

By Lemma 9, let L be such that $h_*\mathcal{O}_S = \mathcal{O}_X \oplus L$. By Lemma 10, $h_*[R_h] = -2L$, so $f_*[R_f] = [B_f] = -2[L]$. It remains to prove that $f_*\mathcal{O}_Y = h_*\mathcal{O}_S$. This follows from Zariski's Main Theorem that asserts that $p_*\mathcal{O}_Y = \mathcal{O}_S$.

Let us write the Weil divisor $f^*\mathcal{O}_X(B_f)$ as $\sum_{i \in I} b_i E_i + \sum_{j \in J} d_j R_j$. Then (21) is equivalent to $b_i \geq 0$ for $i \in I$ and $d_j = 2$ for $j \in J$. We write the Cartier divisor $\mathcal{O}_X(B_f)$ as (U_i, ξ_i) , with ξ_i a local equation of B_f on U_i . Then the Cartier divisor $f^*\mathcal{O}_X(B_f)$ is defined by $(f^{-1}(U_i), \xi_i \circ f)$. Since $\xi_i \circ f$ is regular, we have $b_i \geq 0$ for all i . Moreover, if $j \in J$, then restricting to a neighborhood of a point of R_j where p is a local isomorphism and using Lemma 10, we get $d_j = 2$. \square

We now describe the branch locus as a set.

Proposition 13. *Let $f : Y \rightarrow X$ be a generically finite morphism of degree 2, with X and Y projective, Y irreducible normal, and X normal and locally factorial. Then*

$$\text{Supp}(B_f) = \{x \in X : f^{-1}(x) \text{ is connected}\}$$

Proof. We keep our notation. First, we claim that $B_h = B_f$. This is equivalent to saying that $h(R_h) = f(R_f^1)$. These two varieties are closed in X and of pure codimension one. Lemma 11 yields a bijection between the irreducible components of R_h and those of R_f^1 , implying that $h(R_h)$ and $f(R_f^1)$ have the same irreducible components, proving the claim.

Observe now that B_h is the set of $x \in X$ such that $h^{-1}(x)$ is a point (see the proof of Lemma 10). The fibers of p being connected by Zariski's Main Theorem, the proposition follows. \square

4. PRELIMINARIES ON SCHUBERT VARIETIES

In this section, we fix a semisimple group G with maximal torus T and Borel subgroup B containing T . Let $P \supset B$ be a standard parabolic subgroup. We are interested in the Schubert varieties in G/P , their relative versions in $G/B \times G/P$ and their Bott-Samelson resolutions. We collect both combinatorial and geometric material.

4.1. Reminder on Schubert varieties. For $w \in W$ (resp. $w \in W^P$), denote by $\mathcal{X}_w^B = \overline{BwB/B}$ (resp. $\mathcal{X}_w^P = \overline{BwP/P}$) the associated Schubert subvariety of G/B (resp. G/P). We use notation for the Bruhat orders introduced in Section 2.1.

Lemma 14. *Let $v \in W^P$. Then*

- (1) *The variety \mathcal{X}_v^P is normal. In particular, $\text{Pic}(\mathcal{X}_v^P) = \text{CaCl}(\mathcal{X}_v^P)$ embeds in $\text{Cl}(\mathcal{X}_v^P)$.*
- (2) *We have*

$$\text{Cl}(\mathcal{X}_v^P) = \bigoplus \mathbb{Z}[\mathcal{X}_{v'}^P],$$

where the direct sum runs over the covering relations $\begin{array}{c} v \\ \vdots \\ v' \end{array}$ in the Bruhat graph of G/P .

Proof. Good references for assertions 1 and 2 are [Mat88, XII, Lemme 75] and [BK05, §3.2]. \square

4.2. Inversion sets.

4.2.1. Inversion sets of Weyl group elements. As before, we use notation $W, W^P, W_P, \Delta, \Phi$ and Φ^+ . The inversion set $\Phi(w)$ of an element w of W is defined to be $\Phi(w) = \Phi^+ \cap w^{-1}(\Phi^-)$ and it satisfies $\ell(w) = \#\Phi(w)$. One gets a map

$$\begin{array}{ccc} W & \longrightarrow & \mathcal{S}(\Phi^+) \\ w & \longmapsto & \Phi(w) \end{array}$$

($\mathcal{S}(\Phi^+)$ denotes the power set of Φ^+) that is injective.

The strong and left weak Bruhat orders are characterized by

$$(22) \quad \begin{array}{ccc} v \leq w & \iff & \mathcal{X}_v^B \subset \mathcal{X}_w^B; \\ v \leq_w w & \iff & \Phi(v) \subset \Phi(w). \end{array}$$

The second point is [BB05, Proposition 1.3.1]. The direct implication is a consequence of the easy fact that $w = s_\alpha v$ and $\ell(w) = \ell(v) + 1$ imply that

$$(23) \quad \Phi(w) = \Phi(v) \cup \{v^{-1}(\alpha)\}.$$

A reduced expression $w = s_{\alpha_1} \dots s_{\alpha_\ell}$ corresponds to a sequence $e \leq_w s_{\alpha_\ell} \leq_w s_{\alpha_{\ell-1}} s_{\alpha_\ell} \leq_w \dots \leq_w w$ of weak covering relations and hence to a sequence $\emptyset \subset \mathcal{I}_1 \subset \mathcal{I}_2 \subset \dots \subset \Phi(w)$ of inversion sets of Φ^+ with cardinality increasing by one at each step. In particular, given such a reduced expression we get a numbering

$$(24) \quad \Phi(w) = \{\gamma_1, \dots, \gamma_\ell\} \text{ with } \gamma_i = s_{\alpha_\ell} \dots s_{\alpha_{\ell-i+2}}(\alpha_{\ell-i+1}).$$

Note that $w^{-1} = s_{\alpha_\ell} \dots s_{\alpha_1}$ is also a reduced expression, thus

$$(25) \quad \Phi(w^{-1}) = \{\beta_1, \dots, \beta_\ell\} \text{ with } \beta_i = s_{\alpha_1} \dots s_{\alpha_{i-1}}(\alpha_i).$$

Lemma 15. *Let $v' \leq v$ in W with $\ell(v) = \ell(v') + 1$ be a covering relation in the Bruhat order. Fix a reduced expression $\mathbf{v} = s_{\alpha_1} \dots s_{\alpha_\ell}$ of v . Then there exists a unique k such that a reduced expression of v' is obtained from \mathbf{v} by deleting s_{α_k} .*

Proof. This is the content of [BB05, Lemma 1.3.1]. A geometric interpretation of this result can be obtained using the Bott-Samelson resolution. Indeed, if the statement would be false, one would get nontrivial finite fibers for the Bott-Samelson resolution. By Zariski's Main Theorem, this contradicts the normality of the Schubert varieties. \square

A useful result on the inversion sets is:

Lemma 16. *See e.g. [Kum02, Corollary 1.3.22.3]. Let $w \in W$. We have*

$$\sum_{\beta \in \Phi(w^{-1})} \beta = \rho - w(\rho).$$

We also use the following variation, see [Per07, Lemma 4.18]. The roots β_i are those defined by (25) and the function \hat{h} by (16).

Lemma 17. *Let $v' \leq v$ in W with $\ell(v) = \ell(v') + 1$. Let $\mathbf{v} = s_{\alpha_1} \dots s_{\alpha_\ell}$ be a reduced expression of v and $1 \leq i \leq \ell$ such that v' is obtained by deleting s_{α_i} . Write $\Phi(v^{-1}) = \{\beta_1, \dots, \beta_\ell\}$ as in (25).*

Then, $v' = s_{\beta_i} v$ and $\Phi(v'^{-1}) = \{\beta_1, \dots, \beta_{i-1}, s_{\beta_i}(\beta_{i+1}), \dots, s_{\beta_i}(\beta_\ell)\}$. Moreover,

$$\left\langle \sum_{k=i}^{\ell} \beta_k, \beta_i^\vee \right\rangle = \hat{h}(-v^{-1}\beta_i) + 1.$$

Proof. One can easily check the first assertion. We provide an argument for the second assertion simpler than that in [Per07]. By Lemma 16, we have

$$\begin{aligned} v'(\rho) - v(\rho) &= \sum_{\theta \in \Phi(v^{-1})} \theta - \sum_{\theta \in \Phi(v'^{-1})} \theta \\ &= \beta_i + \sum_{k=i+1}^{\ell} (\beta_k - s_{\beta_i}(\beta_k)) \\ &= \beta_i + \sum_{k=i+1}^{\ell} \langle \beta_k, \beta_i^\vee \rangle \beta_i. \end{aligned}$$

Set $S = \langle \sum_{k=i}^{\ell} \beta_k, \beta_i^\vee \rangle = 2 + \langle \sum_{k=i+1}^{\ell} \beta_k, \beta_i^\vee \rangle$. We get

$$\langle v'(\rho) - v(\rho), \beta_i^\vee \rangle = 2(S - 1).$$


On the other hand,

$$\langle v'(\rho) - v(\rho), \beta_i^\vee \rangle = \langle s_{\beta_i} v(\rho) - v(\rho), \beta_i^\vee \rangle = -2 \langle v(\rho), \beta_i^\vee \rangle,$$

since $s_{\beta_i}(\zeta) = \zeta - \langle \zeta, \beta_i^\vee \rangle \beta_i$ for any weight ζ . The lemma follows. \square


4.2.2. *Diamond lemmas.* We first state the following diamond lemma in the weak Bruhat graph.

Lemma 18. *Let $\alpha, \beta, \gamma, \delta \in \Delta$ such that*



is a subgraph of the weak Bruhat graph. Then $\alpha = \delta$, $\beta = \gamma$ and $s_\alpha s_\beta = s_\beta s_\alpha$.


Proof. Let v denote the Weyl group element of the bottom vertex. We have $s_\gamma s_\alpha v = s_\delta s_\beta v$ whose length is $\ell(v) + 2$. In particular $s_\gamma s_\alpha = s_\delta s_\beta$ has length 2: one may assume that v is trivial. Then, the four inversion sets of the vertices are



Since $\alpha \neq \beta$, we deduce that $\alpha = s_\beta(\delta) = \delta - \langle \delta, \beta^\vee \rangle \beta$. Hence α, β and δ are linearly dependent simple roots. Since $\alpha \neq \beta$ we deduce that $\alpha = \delta$. Now, the linear relation implies $\langle \delta, \beta^\vee \rangle = 0$, so s_δ and s_β commute. By symmetry this ends the proof. \square

For $\alpha \in \Delta$, let $P_\alpha \subset G$ be the corresponding minimal standard parabolic subgroup. We prove a diamond lemma mixing strong and weak Bruhat orders:

Lemma 19. *Let $w \in W, \alpha \in \Delta, \gamma \in \Phi^+$ with $s_\alpha w \neq ws_\gamma$. Let P be a standard parabolic subgroup of G .*



Proof. For (i), since $\mathcal{X}_{ws_\gamma}^B$ contains \mathcal{X}_w^B and not $\mathcal{X}_{s_\alpha w}^B$, it cannot be P_α -stable. Hence $\ell(s_\alpha ws_\gamma) = \ell(w) + 2$ and the first assertion follows.

In the second assertion, assume for a contradiction that $\ell(s_\alpha ws_\gamma) = \ell(ws_\gamma) + 1$. Since \mathcal{X}_w^B is P_α -stable and contains $\mathcal{X}_{ws_\gamma}^B$, one must have $s_\alpha ws_\gamma = w$, a contradiction. Thus $\ell(s_\alpha ws_\gamma) = \ell(ws_\gamma) - 1$.

To prove (iii), let $s_{\alpha'} \in W_P$. Since $s_\alpha w \in W^P$, $\ell(s_\alpha ws_{\alpha'}) = \ell(w) + 2$, so $\ell(ws_{\alpha'}) = \ell(w) + 1$. Thus, $w \in W^P$. If $s_\alpha ws_\gamma \notin W^P$ then $\dim(\mathcal{X}_{s_\alpha ws_\gamma}^P) \leq \dim(\mathcal{X}_{s_\alpha w}^P) = \dim(\mathcal{X}_{ws_\gamma}^P)$. Then $\mathcal{X}_{s_\alpha ws_\gamma}^P = \mathcal{X}_{s_\alpha w}^P = \mathcal{X}_{ws_\gamma}^P$ and $s_\alpha w = ws_\gamma$, a contradiction. \square

4.3. Relative Schubert varieties and their Chow classes. Fix $v \in W^P$. The usual Schubert varieties $\mathcal{X}_v^B = \overline{BvB/B} \subset G/B$ and $\mathcal{X}_v^P = \overline{BvP/P} \subset G/P$ are called *absolute* Schubert varieties. Consider the *relative* Schubert variety $\mathcal{Y}_v = G \times^B \overline{BvP/P}$. It is a closed subvariety of $G/B \times G/P$ and, more precisely, it is the G -orbit closure of $(B/B, vP/P)$ for the diagonal G -action.

The variety \mathcal{Y}_v can be seen from different points of view:

- (1) $\mathcal{Y}_v = \overline{G \times^B BvP/P}$ and $\mathcal{Y}_v = \overline{G \times^P Pv^{-1}B/B}$;
- (2) $\mathcal{Y}_v = \overline{G \cdot (B/B, vP/P)} = \overline{G \cdot (v^{-1}B/B, P/P)} \subset G/B \times G/P$;
- (3) $\mathcal{Y}_v = \{(gB/B, hP/P) : g^{-1}hP/P \in \overline{BvP/P}\} \subset G/B \times G/P$;
- (4) $\mathcal{Y}_v = \{(gB/B, hP/P) : h^{-1}gB/B \in \overline{Pv^{-1}B/B}\} \subset G/B \times G/P$.

Recall that the Schubert classes σ^w, τ_u were defined in Section 2.1.2. Think about $G/B \times G/P$ as a G^2 -flag variety endowed with the action of G diagonally embedded in G^2 . Then \mathcal{Y}_v is an orbit closure of a spherical subgroup in a flag variety. By [Bri98, Theorem 1.5], we have in particular

Proposition 20. *The Chow class of \mathcal{Y}_v is given by:*

$$[\mathcal{Y}_v] = \sum \sigma^{v'v^{-1}} \otimes \tau_{v'} \in A^*(G/B \times G/P),$$

where the sum runs over the set of $v' \in W^P$ such that $v \leq_w v'$.

4.4. Relative Bott-Samelson varieties. Resolutions of Schubert varieties can be produced as Bott-Samelson varieties. We recall this construction in a relative context.

Definition 1. *Let $\mathbf{v} = (\alpha_1, \dots, \alpha_\ell)$ be a sequence of elements in Δ . We call (relative) Bott-Samelson variety the variety $\mathcal{Y}_{\mathbf{v}}$ above G/B defined by*

$$\mathcal{Y}_{\mathbf{v}} := (G \times P_{\alpha_1} \times \dots \times P_{\alpha_\ell}) / B^{\ell+1}.$$

Here we consider the action of $B^{\ell+1}$ defined by

$$(b_0, b_1, \dots, b_\ell) \cdot (g, p_1, \dots, p_\ell) = (gb_0^{-1}, b_0p_1b_1^{-1}, b_1p_2b_2^{-1}, \dots, b_{\ell-1}p_\ell b_\ell^{-1}).$$

Let $v = s_{\alpha_1} \dots s_{\alpha_\ell}$ and assume that $v \in W^P$ and $\ell(v) = \ell$. As in [BK05, §2.2(6)], we may define a morphism ω from $\mathcal{Y}_{\mathbf{v}}$ to the relative Schubert variety \mathcal{Y}_v , by the following diagram:

$$(26) \quad \begin{array}{ccc} & \mathcal{Y}_{\mathbf{v}} := (G \times P_{\alpha_1} \times \dots \times P_{\alpha_\ell}) / B^{\ell+1} & \\ & \downarrow \omega & \\ & \mathcal{Y}_v := G \times^B \overline{BvP/P} & \\ \pi \swarrow & & \searrow \gamma \\ G/B & & G/P \end{array}$$

Forgetting the last term gives a chain of morphisms

$$\mathcal{Y}_{(\alpha_1, \dots, \alpha_\ell)} \xrightarrow{\rho_\ell} \mathcal{Y}_{(\alpha_1, \dots, \alpha_{\ell-1})} \xrightarrow{\rho_{\ell-1}} \dots \longrightarrow \mathcal{Y}_{(\alpha_1)} \xrightarrow{\rho_0} G/B,$$

which are known to be \mathbb{P}^1 -bundles (see [BK05, p.66]). In particular, $\mathcal{Y}_{\mathbf{v}}$ is smooth, and more precisely, we have:

Lemma 21. [BK05, Theorem 3.4.3] *The morphism ω is a rational resolution. Namely, $\omega_* \mathcal{O}_{\mathcal{Y}} = \mathcal{O}_{\mathcal{Y}}$ and for $i > 0$, $R^i \omega_* \mathcal{O}_{\mathcal{Y}} = 0$.*

Since \mathcal{X}_v^P , and thus \mathcal{Y}_v , are smooth in codimension 1, their canonical classes are well-defined as Weil divisors, and since they have rational singularities, we may compute them using the following lemma:

Lemma 22. [BK05, Lemma 3.4.2] *Let $\omega : \mathcal{Y} \rightarrow \mathcal{Y}$ be a rational resolution. Then \mathcal{Y} is Cohen-Macaulay with dualizing sheaf $\omega_* \mathcal{K}_{\mathcal{Y}}$.*

Lemma 21 thus gives:

Lemma 23. *With the notation of Definition 1, we have*

$$\mathcal{K}_{\mathcal{Y}_v} = \omega_* \mathcal{K}_{\mathcal{Y}_{\mathbf{v}}} \text{ as sheaves and } K_{\mathcal{Y}_v} = \omega_* K_{\mathcal{Y}_{\mathbf{v}}} \text{ in } \text{Cl}(\mathcal{Y}_v).$$

Standard Weil divisors and line bundles are defined on those Bott-Samelson varieties as follows [BK05, p.67]:

Definition 2. *Let k be such that $1 \leq k \leq \ell$ and let ζ be a character of B . Then:*

- \mathcal{D}_k is the quotient by $B^{\ell+1}$ of $G \times P_{\alpha_1} \times \dots \times B \times \dots \times P_{\alpha_\ell}$, where we replaced P_{α_k} by B . This is a divisor in $\mathcal{Y}_{\mathbf{v}}$, and there is an isomorphism $\sigma : \mathcal{Y}_{(\alpha_1, \dots, \widehat{\alpha}_k, \dots, \alpha_\ell)} \rightarrow \mathcal{D}_k \subset \mathcal{Y}_{\mathbf{v}}$.

- ζ and k define a character ζ of $B^{\ell+1}$ by $\zeta(b_0, \dots, b_\ell) = \zeta(b_k)$. We define the line bundle $\mathcal{L}_k(\zeta)$ by its total space $(G \times P_{\alpha_1} \times \dots \times P_{\alpha_\ell} \times \mathbb{C}_{-\zeta}) / B^{\ell+1}$ and natural morphism to \mathcal{Y}_v .

In the following lemma, we use heavier notation than the notation in the previous definition: we denote by ${}^v\mathcal{L}_k(\zeta)$ the bundle on \mathcal{Y}_v denoted by $\mathcal{L}_k(\zeta)$ above.

Given $v = (\alpha_1, \dots, \alpha_\ell)$ as above and $1 \leq k \leq \ell$, set $v(\leq k) = (\alpha_1, \dots, \alpha_k)$. The map $(g, p_1, \dots, p_\ell) \mapsto (g, p_1, \dots, p_k)$ induces a proper surjection

$$\rho_{\leq k} : \mathcal{Y}_v \longrightarrow \mathcal{Y}_{v(\leq k)}.$$

Note that $\rho_\ell = \rho_{\leq \ell-1}$.

Lemma 24. *Let v and k be as above. Let $1 \leq i \leq k$ and $\zeta \in X(T)$. Then*

$$\rho_{\leq k}^*({}^{v(\leq k)}\mathcal{L}_k(\zeta)) = {}^v\mathcal{L}_k(\zeta) \quad \text{and} \quad \rho_{\leq k}^*(\mathcal{O}(\mathcal{D}_i)) = \mathcal{O}(\mathcal{D}_i).$$

This Lemma shows that there is no risk of confusion using the lighter notation $\mathcal{L}_k(\zeta)$.

Lemma 25. *Let v , k and ζ be as above. Then, in $\text{Pic}(\mathcal{Y}_v)$, we have*

$$\mathcal{L}_k(\zeta) = \mathcal{L}_{k-1}(s_{\alpha_k}(\zeta)) + \langle \zeta, \alpha_k^\vee \rangle \mathcal{O}(\mathcal{D}_k).$$

Proof. Using Lemma 24, it is sufficient to prove the case $k = \ell$. This lemma is proved in [Dem74, Proposition 1] with some slightly different notation. Therefore we give here a quick argument. Let, as above, $v = (\alpha_1, \dots, \alpha_\ell)$ and $v' = (\alpha_1, \dots, \alpha_{\ell-1})$. A fiber of ρ_ℓ is $P_{\alpha_\ell}/B \simeq \mathbb{P}^1$ and the bundle $\mathcal{L}_\ell(\zeta)$ restricts to $\mathcal{O}_{\mathbb{P}^1}(\langle \zeta, \alpha_\ell^\vee \rangle)$. Thus

$$(27) \quad \mathcal{L}_\ell(\zeta) = \rho_\ell^* L \otimes \mathcal{O}(\langle \zeta, \alpha_\ell^\vee \rangle \mathcal{D}_\ell)$$

for some line bundle L on \mathcal{Y}_v to be determined.

To determine L , we apply σ^* to equation (27), where $\sigma : \mathcal{Y}_{v(\leq \ell-1)} \rightarrow \mathcal{D}_\ell \subset \mathcal{Y}_v$ is as in Definition 2. We have $\sigma^* \rho_\ell^* L = L$, and $\sigma^* \mathcal{O}(\mathcal{D}_\ell) = \mathcal{O}_{\mathcal{Y}_v}(\mathcal{D}_\ell)|_{\mathcal{D}_\ell} = \mathcal{L}_{\ell-1}(\alpha_\ell)$ since the normal bundle of \mathcal{D}_ℓ is $\mathcal{L}_{\ell-1}(\alpha_\ell)$. Thus we get

$$L = \mathcal{L}_{\ell-1}(\zeta - \langle \zeta, \alpha_\ell^\vee \rangle \alpha_\ell) = \mathcal{L}_{\ell-1}(s_{\alpha_\ell}(\zeta)).$$

□

Since ω is proper, the pushforward $\omega_* : A^1(\mathcal{Y}_v) \rightarrow A^1(\mathcal{Y}_v)$ is well defined.

Lemma 26. *Let k be such that $1 \leq k \leq \ell$. Then, in $A^1(\mathcal{Y}_v)$, we have*

$$\omega_*[\mathcal{D}_k] = \begin{cases} [\mathcal{Y}_{s_{\alpha_1} \cdots \widehat{s_{\alpha_k}} \cdots s_{\alpha_\ell}}] & \text{if } s_{\alpha_1} \cdots \widehat{s_{\alpha_k}} \cdots s_{\alpha_\ell} \text{ is reduced;} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. This follows from the fact that the restriction of ω to \mathcal{D}_k is birational onto $\mathcal{Y}_{s_{\alpha_1} \cdots \widehat{s_{\alpha_k}} \cdots s_{\alpha_\ell}}$ if the product $s_{\alpha_1} \cdots \widehat{s_{\alpha_k}} \cdots s_{\alpha_\ell}$ is reduced and contracts \mathcal{D}_k otherwise. □

4.5. Canonical divisor of a relative Schubert variety. We keep the notation of the previous subsection, in particular those of Definition 1. We express the canonical classes of \mathcal{Y}_v and \mathcal{Y}_v :

Proposition 27. *Assume that $v = (\alpha_1, \dots, \alpha_\ell)$ is a reduced expression of $v \in W^P$ and let $\gamma_1 = \alpha_\ell$, $\gamma_i = s_{\alpha_\ell} \cdots s_{\alpha_{\ell-i+2}}(\alpha_{\ell-i+1})$. We have:*

$$-K_{\mathcal{Y}_v} = \pi^* \mathcal{L}_{G/B}(\rho + v(\rho)) + \sum_{i=1}^{\ell} (\ell(\gamma_i) + 1) [\mathcal{D}_i] \text{ in } \text{Cl}(\mathcal{Y}_v) = \text{Pic}(\mathcal{Y}_v).$$

Moreover, we have:

$$-K_{\mathcal{Y}_v} = \pi^* \mathcal{L}_{G/B}(\rho + v(\rho)) + \sum_{v'} (\ell(\gamma) + 1) [\mathcal{Y}_{v'}] \text{ in } \text{Cl}(\mathcal{Y}_v),$$

where the sum runs over the covering relations $\begin{array}{c} v \\ \vdots \\ \gamma \\ \vdots \\ v' \end{array}$ in the twisted Bruhat graph of W^P .

Proof. An immediate induction using Lemma 25 shows that

$$\mathcal{L}_k(\alpha_k) = \mathcal{L}_{G/B}(s_{\alpha_1} \cdots s_{\alpha_k}(\alpha_k)) + \sum_{i=1}^k \langle s_{\alpha_{i+1}} \cdots s_{\alpha_k}(\alpha_k), \alpha_i^\vee \rangle \mathcal{O}(\mathcal{D}_i).$$

Applying $s_{\alpha_1} \cdots s_{\alpha_i}$, we get the equality $\langle s_{\alpha_{i+1}} \cdots s_{\alpha_k}(\alpha_k), \alpha_i^\vee \rangle = \langle \beta_k, \beta_i^\vee \rangle$, since we have the relation $s_{\alpha_1} \cdots s_{\alpha_k}(\alpha_k) = -\beta_k$. Finally

$$(28) \quad \mathcal{L}_k(\alpha_k) = \mathcal{L}_{G/B}(-\beta_k) + \sum_{i=1}^k \langle \beta_k, \beta_i^\vee \rangle \mathcal{O}(\mathcal{D}_i).$$

The fibers of ρ_ℓ are isomorphic to \mathbb{P}^1 and thus the relative tangent bundle of ρ_ℓ is $\mathcal{L}_\ell(\alpha_\ell)$. Therefore, the sequence $0 \rightarrow T\rho_\ell \rightarrow T\mathcal{Y}_v \rightarrow \rho_\ell^*(T\mathcal{Y}_{v(\ell)}) \rightarrow 0$ reads:

$$(29) \quad 0 \rightarrow \mathcal{L}_\ell(\alpha_\ell) \rightarrow T\mathcal{Y}_v \rightarrow \rho_\ell^*(T\mathcal{Y}_{v(\ell)}) \rightarrow 0.$$

Taking account that $\mathcal{Y}_\emptyset = G/B$ and hence $K_{\mathcal{Y}_\emptyset} = \mathcal{L}_{G/B}(-2\rho)$, an immediate induction gives

$$K_{\mathcal{Y}_v} = \mathcal{L}_\ell(-\alpha_\ell) + \rho_\ell^*(K_{\mathcal{Y}_{v(\ell)}}) = \mathcal{L}_{G/B}(-2\rho) + \sum_{k=1}^{\ell} \mathcal{L}_k(-\alpha_k).$$

Injecting (28), we get

$$(30) \quad \begin{aligned} K_{\mathcal{Y}_v} &= \mathcal{L}_{G/B}(-2\rho + \sum_{k=1}^{\ell} \beta_k) + \sum_{k=1}^{\ell} \sum_{i=1}^k \langle \beta_k, \beta_i^\vee \rangle \mathcal{O}(-\mathcal{D}_i) \\ &= \mathcal{L}_{G/B}(-2\rho + \sum_{k=1}^{\ell} \beta_k) + \sum_{i=1}^{\ell} \langle \sum_{k=i}^{\ell} \beta_k, \beta_i^\vee \rangle \mathcal{O}(-\mathcal{D}_i). \end{aligned}$$

We already observed that $\{\beta_1, \dots, \beta_\ell\} = \Phi(v^{-1})$. Then Lemma 16 shows that $\sum_{k=1}^{\ell} \beta_k = \rho - v(\rho)$. Moreover, Lemma 17 shows that $\langle \sum_{k=i}^{\ell} \beta_k, \beta_i^\vee \rangle = \mathfrak{h}(\gamma_i) + 1$, and we get the given formula for $K_{\mathcal{Y}_v}$.

The formula for $K_{\mathcal{Y}_v}$ then follows by applying ω_* , by Lemma 23. Since $\mathcal{L}_{G/B}(\zeta)$ is defined by a pullback, $\omega_*(\mathcal{L}_{G/B}(\zeta)) = \pi^*(\mathcal{L}_{G/B}(\zeta))$. Lemma 26 allows to handle the terms $\mathcal{O}(\mathcal{D}_i)$. We can then conclude using Lemma 15 that realizes a bijection between the covering relations $v' \leq v$ in the weak Bruhat order and

$$\{1 \leq i \leq \ell : \omega \text{ does not contract } \mathcal{D}_i\}.$$

The given formula for $K_{\mathcal{Y}_v}$ follows. □

5. PROOF OF THE MAIN THEOREMS

5.1. Chow class of $Y(\mathfrak{v})$ for any \mathfrak{v} . In this subsection, $\mathfrak{v} = (v, \hat{v})$ denotes any element of $W^{\mathbb{P}}$ and we consider $Y(\mathfrak{v})$ defined by (14). Observe that the projection $\mathcal{Y}_v \rightarrow \hat{G}/\hat{P}$ is a locally trivial fibration with fiber $\hat{P}\hat{v}^{-1}\hat{B}/\hat{B}$. Recall that G/P is embedded in \hat{G}/\hat{P} and denote by \mathcal{Y}_v^G the preimage of G/P in \mathcal{Y}_v . Then $\mathcal{Y}_v^G \simeq G \times^P \hat{P}\hat{v}^{-1}\hat{B}/\hat{B}$.

One can obtain $Y(\mathfrak{v})$ as a transverse intersection:

Lemma 28. *Consider $\hat{G}/\hat{B} \times \mathcal{Y}_v$ and $G/B \times \mathcal{Y}_v$ as subvarieties of $G/B \times \hat{G}/\hat{B} \times \hat{G}/\hat{P}$. Then $Y(\mathfrak{v})$ is the transverse intersection*

$$Y(\mathfrak{v}) = (\hat{G}/\hat{B} \times \mathcal{Y}_v) \cap (G/B \times \mathcal{Y}_v).$$

Similarly, $Y(\mathfrak{v})$ is also the transverse intersection $(\hat{G}/\hat{B} \times \mathcal{Y}_v) \cap (G/B \times \mathcal{Y}_v^G)$ in $G/B \times \hat{G}/\hat{B} \times G/P$.

Proof. It is plain that $Y(\mathfrak{v})$ is equal to the two intersections of the lemma; we prove that these intersections are transverse. Let $x \in G/B$, $\hat{x} \in \hat{G}/\hat{B}$ and $\hat{z} \in \hat{G}/\hat{P}$ such that $(x, \hat{x}, \hat{z}) \in (\hat{G}/\hat{B} \times \mathcal{Y}_v) \cap (G/B \times \mathcal{Y}_v)$. Let $(T_{(x, \hat{x}, \hat{z})}(\mathcal{Y}_v \times \hat{G}/\hat{B}))^\perp$ denote the (conormal) space of linear forms on $T_{(x, \hat{x}, \hat{z})}(G/B \times \hat{G}/\hat{B} \times \hat{G}/\hat{P})$ which vanish on the tangent space $T_{(x, \hat{x}, \hat{z})}(\mathcal{Y}_v \times \hat{G}/\hat{B})$. We have

$$\begin{aligned} (T_{(x, \hat{x}, \hat{z})}(\hat{G}/\hat{B} \times \mathcal{Y}_v))^\perp &\subset T_x^*G/B \times \{0\} \times T_{\hat{z}}^*\hat{G}/\hat{P}, \text{ and} \\ (T_{(x, \hat{x}, \hat{z})}(G/B \times \mathcal{Y}_v))^\perp &\subset \{0\} \times T_{\hat{x}}^*\hat{G}/\hat{B} \times T_{\hat{z}}^*\hat{G}/\hat{P}. \end{aligned}$$

It follows that the intersection $\left(T_{(x,\hat{x},\hat{z})}(\mathcal{Y}_v \times \hat{G}/\hat{B})\right)^\perp \cap \left(T_{(x,\hat{x},\hat{z})}(\mathcal{Y}_{\hat{v}} \times G/B)\right)^\perp$ of these conormal spaces is included in $\{0\} \times \{0\} \times T_{\hat{z}}^* \hat{G}/\hat{P}$, and thus reduced to $\{0\}$ since the projection $G/B \times \mathcal{Y}_{\hat{v}} \rightarrow \hat{G}/\hat{P}$ is surjective. This means that the intersection is transverse. \square

We now compute $[Y(\mathfrak{v})]$ in $A^*(\mathbb{G}/\mathbb{B} \times G/P)$ in the Schubert basis.

Proposition 29. *With the above notation, in $A^*(\mathbb{G}/\mathbb{B} \times G/P)$, we have*

$$(31) \quad [Y(\mathfrak{v})] = \sum \sigma^{\mathfrak{v}'\mathfrak{v}^{-1}} \otimes \delta^*(\tau_{\mathfrak{v}'}),$$

where the sum runs over the set of $\mathfrak{v}' \in \mathbb{W}^{\mathbb{P}}$ such that $\mathfrak{v} \leq_{\mathbb{W}} \mathfrak{v}'$.

Proof. By Lemma 28 and formula (18), we have

$$(32) \quad [Y(\mathfrak{v})] = [\hat{G}/\hat{B} \times \mathcal{Y}_v] \cdot [G/B \times \mathcal{Y}_{\hat{v}}^G] \in A^*(\mathbb{G}/\mathbb{B} \times G/P).$$

Proposition 20 gives a formula for $[\mathcal{Y}_v] \in A^*(G/B \times G/P)$:

$$(33) \quad [\mathcal{Y}_v \times \hat{G}/\hat{B}] = \sum \sigma^{v'v^{-1}} \otimes 1 \otimes \tau_{v'},$$

where the sum runs over the set of $v' \in W^P$ such that $v \leq_{\mathbb{W}} v'$.

Consider now the class $[\mathcal{Y}_{\hat{v}}^G] \in A^*(\hat{G}/\hat{B} \times G/P)$ and the regular embedding $i : \hat{G}/\hat{B} \times G/P \rightarrow \hat{G}/\hat{B} \times \hat{G}/\hat{P}$. Let i^* denote the associated Gysin map. Since $\mathcal{Y}_{\hat{v}}^G$ is the transverse intersection of $\mathcal{Y}_{\hat{v}}$ and $\hat{G}/\hat{B} \times G/P$ in $\hat{G}/\hat{B} \times \hat{G}/\hat{P}$, we have

$$(34) \quad [\mathcal{Y}_{\hat{v}}^G] = i^*[\mathcal{Y}_{\hat{v}}] \in A^*(\hat{G}/\hat{B} \times G/P).$$

But $[\mathcal{Y}_{\hat{v}}]$ has an expression given by Proposition 20 as a linear combination of terms $\sigma^{\hat{v}'\hat{v}^{-1}} \otimes \tau_{\hat{v}'}$. Using the relation $i^*(\sigma^{\hat{v}'\hat{v}^{-1}} \otimes \tau_{\hat{v}'}) = \sigma^{\hat{v}'\hat{v}^{-1}} \otimes i^*(\tau_{\hat{v}'})$, we get in $A^*(\hat{G}/\hat{B} \times G/P)$

$$(35) \quad [\mathcal{Y}_{\hat{v}}^G] = \sum \sigma^{\hat{v}'\hat{v}^{-1}} \otimes i^*(\tau_{\hat{v}'}),$$

and hence in $A^*(\mathbb{G}/\mathbb{B} \times G/P)$

$$(36) \quad [\mathcal{Y}_{\hat{v}}^G \times G/B] = \sum 1 \otimes \sigma^{\hat{v}'\hat{v}^{-1}} \otimes i^*(\tau_{\hat{v}'}),$$

where the two last sums run over the set of $\hat{v}' \in \widehat{W}^{\hat{P}}$ such that $\hat{v} \leq_{\mathbb{W}} \hat{v}'$. Multiplying (33) with (36) and using (11), one gets the formula of the proposition. \square

We now compute its pushforward by Π :

Corollary 30. *Assume that $\ell(\mathfrak{v}) = \dim(\hat{G}/\hat{P}) - 1$ and consider the projection $\tilde{\Pi} : \mathbb{G}/\mathbb{B} \times G/P \rightarrow \mathbb{G}/\mathbb{B}$. Then, in $A^1(\mathbb{G}/\mathbb{B})$, we have*

$$\tilde{\Pi}_*([Y(\mathfrak{v})]) = \sum c(\mathfrak{v}') \sigma^{s_{\alpha}},$$

where the sum runs over the covering relations $\begin{array}{c} \mathfrak{v}' \\ \alpha \downarrow \\ \mathfrak{v} \end{array}$ in the Bruhat graph of \mathbb{G}/\mathbb{P} .


Proof. In general, for a class $\sum a_{\mathfrak{u},v} \sigma^{\mathfrak{u}} \otimes \tau_v$ in $A^*(\mathbb{G}/\mathbb{B} \times G/P)$, we have

$$(37) \quad \tilde{\Pi}_* \left(\sum_{\mathfrak{u},v} a_{\mathfrak{u},v} \sigma^{\mathfrak{u}} \otimes \tau_v \right) = \sum_{\mathfrak{u}} a_{\mathfrak{u},e} \sigma^{\mathfrak{u}}.$$

But in (31), the degree of $\tau_{\mathfrak{v}'}$ equals $\dim(G/P)$ if and only if $\ell(\mathfrak{v}') = \dim(\hat{G}/\hat{P})$ if and only if $\ell(\mathfrak{v}'\mathfrak{v}^{-1}) = 1$. Hence the terms in the sum (31) that do not vanish after applying $\tilde{\Pi}_*$ are indexed by the $\mathfrak{v}' = s_{\alpha}\mathfrak{v}$ for some simple root $\alpha \in \Delta$ such that $\ell(s_{\alpha}\mathfrak{v}) = \ell(\mathfrak{v})$. The corresponding term is $\sigma^{s_{\alpha}} \otimes \delta^*(\tau_{s_{\alpha}\mathfrak{v}}) = c(s_{\alpha}\mathfrak{v}) \sigma^{s_{\alpha}} \otimes [pt]$. This implies the given formula. \square

5.2. **Canonical class of $Y(\mathfrak{v})$.** In the previous subsection, we computed the canonical class of a relative Schubert variety, and we now express $Y(\mathfrak{v})$ as a fibered product of two relative Schubert varieties and explain how to deduce the canonical class of Π .

Fix reduced expressions \mathfrak{v} and $\hat{\mathfrak{v}}$ of v and \hat{v} respectively, and let us use Definition 1 for v and \hat{v} . We define $\mathcal{Y}_{\hat{\mathfrak{v}}}^G$ from $\mathcal{Y}_{\hat{\mathfrak{v}}}$ in a way similar to what we did for $\mathcal{Y}_{\mathfrak{v}}^G$ from $\mathcal{Y}_{\mathfrak{v}}$. We have morphisms:



Observe that $Y(\mathfrak{v})$ as in (14) is nothing else than $\mathcal{Y}_{\mathfrak{v}} \times_{G/P} \mathcal{Y}_{\hat{\mathfrak{v}}}^G$ and that Π is the restriction of $\tilde{\Pi} = (\pi, \hat{\pi})$ to $Y(\mathfrak{v})$. We also let $\mathfrak{v} = (\mathfrak{v}, \hat{\mathfrak{v}})$ and $Y(\mathfrak{v}) = \mathcal{Y}_{\mathfrak{v}} \times_{G/P} \mathcal{Y}_{\hat{\mathfrak{v}}}^G$. We have the following natural morphisms:

$$(38) \quad \begin{array}{ccccc} \mathcal{Y}_{\mathfrak{v}} & \xleftarrow{\epsilon} & Y(\mathfrak{v}) & \xrightarrow{\hat{\epsilon}} & \mathcal{Y}_{\hat{\mathfrak{v}}}^G \\ \omega \downarrow & & \downarrow \Omega & & \downarrow \hat{\omega} \\ G/B \times G/P & \xleftarrow{\epsilon} & \mathcal{Y}_{\mathfrak{v}} & \xrightarrow{\hat{\epsilon}} & \mathcal{Y}_{\hat{\mathfrak{v}}}^G \hookrightarrow \hat{G}/\hat{B} \times G/P \\ & & \Gamma \swarrow & & \searrow \Pi \\ & & G/P & & G/B \times \hat{G}/\hat{B} \end{array}$$

The first properties of the morphisms appearing in this diagram we need are:

Lemma 31. *With the above notation,*

- (1) ϵ is a locally trivial fibration with fiber $\overline{\hat{P}\hat{v}^{-1}\hat{B}/\hat{B}} = \mathcal{X}_{w_0, \hat{P}\hat{v}}^{\hat{B}}$.
- (2) $\hat{\epsilon}$ is a locally trivial fibration with fiber $\overline{Pv^{-1}B/B} = \mathcal{X}_{w_0, Pv}^B$.
- (3) ϵ and $\hat{\epsilon}$ are locally trivial fibrations.

In particular, these four morphisms are flat and induce pullbacks between Chow rings.

Proof. We already observed when we defined $\mathcal{Y}_{\hat{\mathfrak{v}}}^G$ at the beginning of Section 5.1 that $\mathcal{Y}_{\hat{\mathfrak{v}}}^G$ is a locally trivial fibration over G/P with fiber $\overline{\hat{P}\hat{v}^{-1}\hat{B}/\hat{B}}$. Since $Y(\mathfrak{v}) = \mathcal{Y}_{\mathfrak{v}} \times_{G/P} \mathcal{Y}_{\hat{\mathfrak{v}}}^G$, the first assertion follows. The second one works similarly.

By the Bruhat decomposition, the morphism $G \rightarrow G/P$ is locally trivial in Zariski topology. Hence the maps $\mathcal{Y}_{\hat{\mathfrak{v}}}^G \rightarrow G/P$ and $\mathcal{Y}_{\mathfrak{v}} \rightarrow G/P$ are locally trivial fibrations. The last assertion follows. \square

We can now express the canonical bundle of $Y(\mathfrak{v})$ in terms of the canonical bundles of $\mathcal{Y}_{\mathfrak{v}}$ and $\mathcal{Y}_{\hat{\mathfrak{v}}}$:

Proposition 32. *We have $K_{Y(\mathfrak{v})} = \epsilon^* K_{\mathcal{Y}_{\mathfrak{v}}} + \hat{\epsilon}^* K_{\mathcal{Y}_{\hat{\mathfrak{v}}}^G} - \Gamma^* K_{G/P}$ in $\text{Cl}(Y(\mathfrak{v}))$.*

Proof. Let us first prove this result at the level of Bott-Samelson varieties. We have by definition of $Y(\mathfrak{v})$ a fibered square of smooth varieties:

$$\begin{array}{ccc}
\mathbf{Y}(\mathbf{v}) & \xrightarrow{\epsilon \times \hat{\epsilon}} & \mathcal{Y}_{\mathbf{v}} \times \mathcal{Y}_{\hat{\mathbf{v}}}^G \\
\Gamma \circ \Omega \downarrow & \square & \downarrow \gamma \times \hat{\gamma} \\
G/P & \xrightarrow{\Delta} & G/P \times G/P
\end{array}$$

Let $\mathbf{y} = (\mathbf{y}, \hat{\mathbf{y}}) \in \mathbf{Y}(\mathbf{v})$ and let $u = \Gamma \circ \Omega(\mathbf{y}) \in G/P$. We thus have a fibered square of tangent spaces:

$$\begin{array}{ccc}
T_{\mathbf{y}}\mathbf{Y}(\mathbf{v}) & \longrightarrow & T_{\mathbf{y}}\mathcal{Y}_{\mathbf{v}} \oplus T_{\hat{\mathbf{y}}}\mathcal{Y}_{\hat{\mathbf{v}}}^G \\
\downarrow & \square & \downarrow \\
T_u G/P & \xrightarrow{d\Delta} & T_u G/P \oplus T_u G/P
\end{array}$$

The bottom horizontal map being injective, so is the top horizontal one. The right vertical map being surjective, $(T_{\mathbf{y}}\mathcal{Y} \oplus T_{\hat{\mathbf{y}}}\mathcal{Y}_{\hat{\mathbf{v}}}^G)/T_{\mathbf{y}}\mathbf{Y}(\mathbf{v})$ identifies with $(T_u G/P \oplus T_u G/P)/T_u G/P$. Moreover, on $\mathbf{Y}(\mathbf{v})$, the bundle $(TG/P \oplus TG/P)/TG/P$ is isomorphic to $\Omega^* \Gamma^* TG/P$. We get the short exact sequence on $\mathbf{Y}(\mathbf{v})$

$$0 \longrightarrow T_{\mathbf{Y}(\mathbf{v})} \longrightarrow T_{\mathcal{Y}_{\mathbf{v}}} \oplus T_{\mathcal{Y}_{\hat{\mathbf{v}}}^G} \longrightarrow \Omega^* \Gamma^* T_{G/P} \longrightarrow 0.$$

From this we deduce that

$$\begin{aligned}
(39) \quad K_{\mathbf{Y}(\mathbf{v})} &= (K_{\mathcal{Y}_{\mathbf{v}}} + K_{\mathcal{Y}_{\hat{\mathbf{v}}}^G})|_{\mathbf{Y}(\mathbf{v})} - \Omega^* \Gamma^* K_{G/P} \\
&= \epsilon^* K_{\mathcal{Y}_{\mathbf{v}}} + \hat{\epsilon}^* K_{\mathcal{Y}_{\hat{\mathbf{v}}}^G} - \Omega^* \Gamma^* K_{G/P}.
\end{aligned}$$

We now apply Ω_* . Lemma 21 yields $\Omega_* \mathcal{O}_{\mathbf{Y}(\mathbf{v})} = \mathcal{O}_{Y(\mathbf{v})}$ and $\Omega_* K_{\mathbf{Y}(\mathbf{v})} = K_{Y(\mathbf{v})}$. From the first point and projection formula, we deduce in $\text{Cl}(Y(\mathbf{v}))$

$$(40) \quad \Omega_* \Omega^* \Gamma^* K_{G/P} = \Gamma^* K_{G/P}.$$

$$\begin{array}{ccc}
\mathbf{Y}(\mathbf{v}) & \xrightarrow{(\epsilon, \hat{\epsilon})} & \mathcal{Y}_{\mathbf{v}} \times \mathcal{Y}_{\hat{\mathbf{v}}}^G \\
\Omega \downarrow & \square & \downarrow \omega \\
Y(\mathbf{v}) & \xrightarrow{(\epsilon, \hat{\epsilon})} & \mathcal{Y}_{\mathbf{v}} \times \mathcal{Y}_{\hat{\mathbf{v}}}^G
\end{array}$$

Moreover, we have a fibered product . By the flat base change $(\epsilon, \hat{\epsilon})$ and [Har77,

Proposition III.9.3], we deduce that

$$(41) \quad \Omega_*(\epsilon^*, \hat{\epsilon}^*)(\mathcal{K}_{\mathcal{Y}_{\mathbf{v}}} \otimes \mathcal{K}_{\mathcal{Y}_{\hat{\mathbf{v}}}^G}) = (\epsilon^*, \hat{\epsilon}^*)\omega_*(\mathcal{K}_{\mathcal{Y}_{\mathbf{v}}} \otimes \mathcal{K}_{\mathcal{Y}_{\hat{\mathbf{v}}}^G}).$$

But, by [KM98, Theorem 5.10], $\omega_* \mathcal{K}_{\mathcal{Y}_{\mathbf{v}}} = \mathcal{K}_{\mathcal{Y}_{\mathbf{v}}}$ and $\omega_* \mathcal{K}_{\mathcal{Y}_{\hat{\mathbf{v}}}^G} = \mathcal{K}_{\mathcal{Y}_{\hat{\mathbf{v}}}^G}$. We deduce that

$$(42) \quad \Omega_*(\epsilon^* \mathcal{K}_{\mathcal{Y}_{\mathbf{v}}} + \hat{\epsilon}^* \mathcal{K}_{\mathcal{Y}_{\hat{\mathbf{v}}}^G}) = \epsilon^* \mathcal{K}_{\mathcal{Y}_{\mathbf{v}}} + \hat{\epsilon}^* \mathcal{K}_{\mathcal{Y}_{\hat{\mathbf{v}}}^G}.$$

By the equalities (39), (40), (41), and (42), we get the result of the proposition. \square

For later use, we also compute $\Pi_* \circ \Gamma^*$ (recall from (9) that χ is the coefficient of the point class):

Lemma 33. *The map $\Pi_* \circ \Gamma^* : A^*(G/P) \longrightarrow A^*(\mathbb{G}/\mathbb{B})$ is determined by*

$$\Pi_* \circ \Gamma^*(\xi) = \sum \chi_{G/P}(\xi \cdot \delta^*(\tau_{\mathbf{v}'})) \sigma^{\mathbf{v}' \vee^{-1}} \quad \forall \xi \in A^*(G/P),$$

where the sum runs over the set of $\mathbf{v}' \in \mathbb{W}^{\mathbb{P}}$ such that $\mathbf{v} \leq_{\mathbb{W}} \mathbf{v}'$. If $\xi \in A^d(G/P)$, then all the non-vanishing terms in this summand satisfy $\ell(\mathbf{u}) = d$.

Proof. Let $\xi \in A^*(G/P)$ and consider the following commutative diagram:

$$(43) \quad \begin{array}{ccc}
Y(\mathbf{v}) & \xrightarrow{i} & \mathbb{G}/\mathbb{B} \times G/P \\
& \searrow \Gamma & \downarrow p \\
& & G/P.
\end{array}$$

By the projection formula, we have

$$i_* \Gamma^* \xi = i_* i^* p^* \xi = p^* \xi \cup i_* [Y(\mathbf{v})].$$

Proposition 29 gives $i_*[Y(\mathfrak{v})]$, and since $p^*\xi = 1 \otimes \xi$, one gets

$$(44) \quad i_*\Gamma^*\xi = \sum_{\substack{\mathfrak{v}' \in \mathbb{W}^{\mathbb{P}} \\ \mathfrak{u}\mathfrak{v} \xrightarrow{\mathbb{P}} \mathfrak{u}\mathfrak{v}}} \sigma^{\mathfrak{v}'\mathfrak{v}^{-1}} \otimes (\xi \cdot \delta^*(\tau_{\mathfrak{v}'})),$$

where the sum runs over the set of $\mathfrak{v}' \in \mathbb{W}^{\mathbb{P}}$ such that $\mathfrak{v} \leq_{\mathbb{W}} \mathfrak{v}'$. Since $\Pi_*\Gamma^*\xi = \tilde{\Pi}_*i_*\Gamma^*\xi$, we deduce the formula of the proposition.

Assume now that ξ is homogeneous of degree d . As in the proof of Corollary 30, the terms to keep when applying $\tilde{\Pi}_*$ to (44) satisfy $\ell(\mathfrak{u}) = d$. \square

5.3. Computation of the class of the branch divisor. We compute $[B_{\Pi}] = \Pi_*K_{\Pi}$ using the already done computation of $K_{Y(\mathfrak{v})}$.

Proof of Theorem 7: recall that $Y(\mathfrak{v}) \subset \mathbb{G}/\mathbb{B} \times G/P$ and that Π resp. Γ is the projection of $Y(\mathfrak{v})$ on \mathbb{G}/\mathbb{B} resp. G/P . Consider also as before

$$\tilde{\Pi} : \mathbb{G}/\mathbb{B} \times G/P \longrightarrow \mathbb{G}/\mathbb{B},$$

the restriction to $Y(\mathfrak{v})$ of which is Π .

By the definition of K_{Π} in Section 3.2 and Proposition 32, we have

$$(45) \quad K_{\Pi} = \epsilon^*K_{\mathcal{Y}_v} + \hat{\epsilon}^*K_{\mathcal{Y}_v^G} - \Gamma^*K_{G/P} - \Pi^*(K_{G/B} + K_{\hat{G}/\hat{B}}).$$

Consider first $K_{\mathcal{Y}_v^G}$ in $A^1(\mathcal{Y}_v^G)$. Let $j : \mathcal{Y}_v^G \longrightarrow \mathcal{Y}_v$ denote the inclusion. As the pullback of the inclusion $\iota : G/P \longrightarrow \hat{G}/\hat{P}$, it is a regular embedding and we can consider the associated Gysin map j^* . At any point y of \mathcal{Y}_v^G with projection $\bar{y} \in G/P$ we have the exact sequence:

$$0 \longrightarrow T_y\mathcal{Y}_v^G \longrightarrow T_y\mathcal{Y}_v \longrightarrow T_{\bar{y}}\hat{G}/\hat{P}/T_{\bar{y}}G/P \longrightarrow 0.$$

We deduce that in $\text{Cl}(\mathcal{Y}_v^G)$, we have

$$(46) \quad K_{\mathcal{Y}_v^G} = j^*K_{\mathcal{Y}_v} - \hat{\gamma}^*((K_{\hat{G}/\hat{P}})|_{G/P}) + \hat{\gamma}^*(K_{G/P}).$$

Using the formulas $-K_{\hat{G}/\hat{P}} = \mathcal{L}_{\hat{G}/\hat{P}}(2\rho^{\hat{L}})$, $-K_{G/B} = \mathcal{L}_{G/B}(2\rho_G)$, and Proposition 27, injecting (46) in (45), one gets

$$(47) \quad \begin{aligned} &K_{\Pi} = A + B - C - D \quad \text{with} \\ &\begin{cases} A = \Pi^*c_1(\mathcal{L}_{\mathbb{G}/\mathbb{B}}(\rho_{\mathbb{G}} - \mathfrak{v}(\rho_{\mathbb{G}}))) \\ B = \Gamma^*c_1(\mathcal{L}_{G/P}(2\rho_{|T}^{\hat{L}})) \\ C = \sum_{\gamma} (\hat{h}(\gamma) + 1)\epsilon^*([\mathcal{Y}_{v'}]) \\ D = \sum_{\hat{\gamma}} (\hat{h}(\hat{\gamma}) + 1)\hat{\epsilon}^* \circ j^*([\mathcal{Y}_{v'}]), \end{cases} \end{aligned}$$

where the sums C and D run respectively over the pictures $\begin{array}{c} v \\ \vdots \\ \gamma \\ \vdots \\ v' \end{array}$ and $\begin{array}{c} \hat{v} \\ \vdots \\ \hat{\gamma} \\ \vdots \\ \hat{v}' \end{array}$ in the twisted Bruhat graphs of W^P and $W^{\hat{P}}$ respectively.

We now pushforward by Π each summand of (47). First of all, Lemma 33 gives

$$\Pi_*B = \sum_{\substack{\alpha \\ \mathfrak{v}}} \left(c_1(\mathcal{L}_{G/P}(2\rho_{|T}^{\hat{L}})) \cdot \delta^*(\tau_{s_{\alpha}\mathfrak{v}}) \right) \sigma^{s_{\alpha}},$$

where the sum runs over the covering relations $\begin{array}{c} s_{\alpha}\mathfrak{v} \\ \alpha \\ \mathfrak{v} \end{array}$ in the Bruhat graph of \mathbb{G}/\mathbb{P} . This is the first term of the formula of Theorem 7.

For $\zeta \in X(\mathbb{T})$, the projection formula gives $\Pi_* \circ \Pi^* c_1(\mathcal{L}_{\mathbb{G}/\mathbb{B}}(\zeta)) = \deg(\Pi) c_1(\mathcal{L}_{\mathbb{G}/\mathbb{B}}(\zeta))$, and the Chevalley formula gives $c_1(\mathcal{L}_{\mathbb{G}/\mathbb{B}}(\zeta)) = \sum_{\alpha \in \Delta} \langle \zeta, \alpha^\vee \rangle \sigma^{s_\alpha}$. Thus

$$(48) \quad \Pi_* A = c(\mathfrak{v}) \sum_{\alpha \in \Delta} \langle \rho_{\mathbb{G}} - \mathfrak{v}(\rho_{\mathbb{G}}), \alpha^\vee \rangle \sigma^{s_\alpha}.$$

Consider first a term $\alpha \in \Delta$ such that $s_\alpha \mathfrak{v} \in \mathfrak{v}W_{\mathbb{P}}$. Then $s_\alpha \mathfrak{v} = \mathfrak{v}s_{\mathfrak{v}}$ with $\mathfrak{v} = \mathfrak{v}^{-1}(\alpha) \in \Phi(\mathbb{L})$. Since $\mathfrak{v} \in W^{\mathbb{P}}$, $\ell(\mathfrak{v}) + 1 \geq \ell(s_\alpha \mathfrak{v}) = \ell(\mathfrak{v}s_{\mathfrak{v}}) = \ell(\mathfrak{v}) + \ell(s_{\mathfrak{v}})$. Then \mathfrak{v} is a simple root of \mathbb{L} . In particular, $\langle \mathfrak{v}(\rho_{\mathbb{G}}), \alpha^\vee \rangle = 1$ and the corresponding term in the sum (48) vanishes.

Assume now that $\ell(s_\alpha \mathfrak{v}) = \ell(\mathfrak{v}) + 1$ and $s_\alpha \mathfrak{v} \in W^{\mathbb{P}}$. Then $\langle \mathfrak{v}(\rho_{\mathbb{G}}) - \rho_{\mathbb{G}}, \alpha^\vee \rangle = \mathfrak{h}(\mathfrak{v}) - 1$ for $\mathfrak{v} = \mathfrak{v}^{-1}(\alpha)$. Assume finally that $\ell(s_\alpha \mathfrak{v}) = \ell(\mathfrak{v}) - 1$ and $s_\alpha \mathfrak{v} \in W^{\mathbb{P}}$. Then, similarly, we have $\langle \rho_{\mathbb{G}} - \mathfrak{v}(\rho_{\mathbb{G}}), \alpha^\vee \rangle = \mathfrak{h}(\mathfrak{v}) + 1$, for $\mathfrak{v} = -\mathfrak{v}^{-1}\alpha$. Summing up, we get

$$(49) \quad \Pi_* A = \sum -c(\mathfrak{v})(\mathfrak{h}(\mathfrak{v}) - 1)\sigma^{s_\alpha} + \sum c(\mathfrak{v})(\mathfrak{h}(\mathfrak{v}) + 1)\sigma^{s_\alpha},$$

where the two sums run respectively over the pictures $\begin{array}{c} s_\alpha \mathfrak{v} \\ \vdots \\ \mathfrak{v} \end{array}$ and $\begin{array}{c} \mathfrak{v} \\ \vdots \\ s_\alpha \mathfrak{v} \end{array}$ in the twisted Bruhat graphs of $W^{\mathbb{P}}$.

We now compute $\Pi_* \circ \epsilon^*([\mathcal{Y}_{v'}])$, where $v' = s_\alpha v \in W^{\mathbb{P}}$, $\alpha \in \Delta$ and $\ell(v) = \ell(v') + 1$. Set $\mathfrak{v}' = (v', \hat{v})$. Set $Y' = Y(\mathfrak{v}') = \mathcal{Y}_{v'} \times_{G/P} \mathcal{Y}_{\hat{v}}^G$, so that we have

$$(50) \quad \epsilon^*[\mathcal{Y}_{v'}] = [Y'] \in \text{Cl}(Y(\mathfrak{v})).$$

Consider the class $i_*[Y']$ of Y' in $A^*(\mathbb{G}/\mathbb{B} \times G/P)$ (recall the diagram (43)). Then

$$(51) \quad \Pi_*(\epsilon^*[\mathcal{Y}_{v'}]) = \tilde{\Pi}_* i_*[Y'].$$

We apply Corollary 30 to \mathfrak{v}' to get

$$(52) \quad \tilde{\Pi}_* i_*[Y'] = \sum c(s_\alpha \mathfrak{v}') \sigma^{s_\alpha},$$

where the sum runs over the covering relation $\begin{array}{c} s_\alpha \mathfrak{v}' \\ \alpha \\ \mathfrak{v}' \end{array}$. Then

$$(53) \quad \Pi_* C = \sum (\mathfrak{h}(\gamma) + 1) c(\mathfrak{v}'') \sigma^{s_\alpha},$$

where the sum runs over the subgraphs $\begin{array}{c} \mathfrak{v} \\ \gamma \setminus / \\ \mathfrak{v}' \end{array}$ in the Bruhat graph of \mathbb{G}/\mathbb{P} with $\alpha \in \Delta$ and $\gamma \in \Phi^+(G)$.

We now compute a term $\Pi_* \circ \hat{\epsilon}^* \circ j^*(\mathcal{Y}_{\hat{v}'})$ where $\hat{v}' = s_{\hat{\alpha}} \hat{v} \in \widehat{W}^{\hat{P}}$, $\hat{\alpha} \in \hat{\Delta}$ and $\ell(\hat{v}) = \ell(\hat{v}') + 1$. Since the intersection $\mathcal{Y}_{\hat{v}'} \cap \mathcal{Y}_{\hat{v}}^{\hat{G}} = \mathcal{Y}_{\hat{v}'}^{\hat{G}}$ is transverse, we have $j^*([\mathcal{Y}_{\hat{v}'}]) = [\mathcal{Y}_{\hat{v}'}^{\hat{G}}]$. Then

$$\Pi_* \circ \hat{\epsilon}^* \circ j^*([\mathcal{Y}_{\hat{v}'}]) = \Pi_*([\mathcal{Y}_{\hat{v}} \times_{G/P} \mathcal{Y}_{\hat{v}'}])$$

is given by Corollary 30. One gets that

$$(54) \quad \Pi_* D = \sum (\mathfrak{h}(\hat{\gamma}) + 1) c(\mathfrak{v}'') \sigma^{s_\alpha},$$

where the sum runs over the subgraphs $\begin{array}{c} \mathfrak{v} \\ \hat{\gamma} \setminus / \\ \mathfrak{v}' \end{array}$ in the Bruhat graph of \mathbb{G}/\mathbb{P} with $\alpha \in \Delta$ and $\hat{\gamma} \in \Phi^+(\hat{G})$.

Putting together (53) and (54), we get

$$(55) \quad \Pi_*(C + D) = \sum_{22} (\mathfrak{h}(\mathfrak{v}) + 1) c(\mathfrak{v}'') \sigma^{s_\alpha},$$

where the sum runs over the subgraphs $\begin{array}{c} \mathfrak{v} \\ \delta \searrow \\ \mathfrak{v}' \end{array} \begin{array}{c} \mathfrak{v}'' \\ \nearrow \\ \alpha \end{array}$ in the Bruhat graph of \mathbb{G}/\mathbb{P} .

By Lemma 19, the terms in (55) with $\mathfrak{v} \neq \mathfrak{v}''$ are exactly the terms in the second sum of Theorem 7. Let us now consider a term in (55) with $\mathfrak{v} = \mathfrak{v}''$, meaning that $s_\alpha \mathfrak{v} = \mathfrak{v} s_\delta$. If $\ell(s_\alpha \mathfrak{v}) < \ell(\mathfrak{v})$, then the coefficient of σ^{s_α} in $\Pi_* A$ and in $\Pi_*(C+D)$ is $c(\mathfrak{v})(h(\mathfrak{v})+1)$, by (48) and (55). So these terms cancel, and accordingly Theorem 7 states that this coefficient vanishes. If $\ell(s_\alpha \mathfrak{v}) > \ell(\mathfrak{v})$, then the coefficient of σ^{s_α} in $\Pi_* A$ is $-c(\mathfrak{v})(h(\mathfrak{v})-1)$ and in $\Pi_*(C+D)$ it is 0. This is also compatible with Theorem 7. \square

5.4. Proof of our reduction formula. In this section, we prove Theorem 6.

5.4.1. The birational case. We recall from [Res11b] the main construction in the proof of Theorem 5, because this will allow introducing useful notation. Recall that $\mathbb{G} = G \times \hat{G}$, $\mathfrak{v} = (v, \hat{v})$ and set $\mathbb{L} = L \times \hat{L}$ and $\zeta = (\zeta, \hat{\zeta})$. Consider the variety $X = \mathbb{G}/\mathbb{B}$ endowed with the diagonal G -action. Let $\mathcal{L} \rightarrow X$ be the line bundle defined by ζ . By Borel-Weil theorem, we have

$$(56) \quad m_{G \times \hat{G}}(\zeta, \hat{\zeta}) = \dim(\mathrm{H}^0(X, \mathcal{L})^G).$$

Let $C \subset X$ be the irreducible component of the fixed point set X^τ of τ in X containing the point $x_0 := \mathfrak{v}^{-1}\mathbb{B}/\mathbb{B}$. Let $B_{\mathbb{L}} = \mathbb{B} \cap \mathbb{L}$. Observe that the stabilizer of x_0 in \mathbb{L} is $B_{\mathbb{L}}$, so that the variety C is isomorphic to $\mathbb{L}/B_{\mathbb{L}}$. Moreover, $\mathcal{L}|_C$ is isomorphic to the line bundle defined by $\mathfrak{v}^{-1}\zeta$. In particular

$$(57) \quad m_{LC\hat{L}}(v^{-1}\zeta, \hat{v}^{-1}\hat{\zeta}) = \dim(\mathrm{H}^0(C, \mathcal{L})^L).$$

The Białynicki-Birula cell C^+ is

$$C^+ = \{x \in X \mid \lim_{t \rightarrow 0} \tau(t)x \in C\} = \mathbb{P}_{\mathfrak{v}^{-1}\mathbb{B}}/\mathbb{B}.$$

Consider

$$(58) \quad \begin{array}{ccc} G \times^P \overline{C^+} & \xrightarrow{\Pi} & X \\ \downarrow & & \\ G/P & & \end{array}$$

Observe that $Y(\mathfrak{v}) = G \times^P \overline{C^+}$ endowed with $\Pi : Y(\mathfrak{v}) \rightarrow X$ is nothing but the incidence variety defined in (14).

The proof in [Res11b] goes on proving the following, which does not need the hypothesis $c(\mathfrak{v}) = 1$:

Theorem 34. *Assume that $v^{-1}(\zeta)|_S + \hat{v}^{-1}(\hat{\zeta})|_S$ is trivial. With the notation of (58), there is a natural isomorphism*

$$\mathrm{H}^0(Y(\mathfrak{v}), \Pi^* \mathcal{L})^G \simeq \mathrm{H}^0(C, \mathcal{L}|_C)^L.$$

In [Res11b], to prove Theorem 34, we first prove that $\mathrm{H}^0(C, \mathcal{L}|_C)^L = \mathrm{H}^0(C^+, \mathcal{L}|_{C^+})^P$ using the action of τ . Since $\mathrm{H}^0(C^+, \mathcal{L}|_{C^+})^P = \mathrm{H}^0(G \times^P C^+, \mathcal{L})^G$, it remains to prove that each rational G -invariant on $Y(\mathfrak{v})$ that is regular on the open set $G \times^P C^+$ extends to $Y(\mathfrak{v})$. By normality of the Schubert varieties, it is sufficient to prove that σ has no pole along the boundary. This last step is made by quite explicit local computation.

The fiber of Π over any general point in x is isomorphic to the intersection of two subvarieties of G/P of Chow class τ_v and $L^* \tau_{\hat{v}}$: therefore the assumption $c(\mathfrak{v}) = 1$ implies that Π is birational and gives further $\mathrm{H}^0(Y(\mathfrak{v}), \Pi^* \mathcal{L}) \simeq \mathrm{H}^0(X, \mathcal{L})$, from which Theorem 5 follows.

5.4.2. *The degree 2 case.* We now deal with the case $c(\mathfrak{v}) = 2$. Then Π is no longer birational, but it is generically finite of degree 2, and we apply the projection formula as in Section 3.4. We denote by $B_\Pi \subset X$ the branch divisor. Then, in the Schubert basis of $X = \mathbb{G}/\mathbb{B}$, $[B_\Pi]$ expands as

$$(59) \quad [B_\Pi] = \sum_{\alpha} n_{\alpha} \sigma^{s_{\alpha}} \otimes 1 + \sum_{\hat{\alpha}} n_{\hat{\alpha}} 1 \otimes \sigma^{s_{\hat{\alpha}}} \in A^*(\mathbb{G}/\mathbb{B}),$$

for some well defined integers n_{α} and $n_{\hat{\alpha}}$. These coefficients are described by Theorem 7.

We proved in Lemma 10 that $\frac{1}{2}[B_\Pi] \in \text{Pic}(X)$. Recall that θ and $\hat{\theta}$ in $X(T)$ and $X(\hat{T})$ are defined by

$$(60) \quad \theta = \sum_{\alpha} \frac{n_{\alpha}}{2} \varpi_{\alpha} \quad \hat{\theta} = \sum_{\hat{\alpha}} \frac{n_{\hat{\alpha}}}{2} \varpi_{\hat{\alpha}},$$

in such a way that $\frac{1}{2}[B_\Pi]$ is the line bundle on $X = G/B \times \hat{G}/\hat{B}$ associated to the character $(\theta, \hat{\theta})$ of $B \times \hat{B}$.

Proof of Theorem 6. Observe that C can be seen as a subvariety of $Y(\mathfrak{v}) = G \times^P \overline{C^+}$ via the closed immersion $x \mapsto [e, x]$, and that $\Pi^*(\mathcal{L})|_C$ is equal to $\mathcal{L}|_C$. In particular equality (57) can be rewritten as

$$m_{LC\hat{L}}(v^{-1}\zeta, \hat{v}^{-1}\hat{\zeta}) = \dim H^0(C, \Pi^*(\mathcal{L})|_C)^L.$$

Now, by Theorem 34, there is an isomorphism

$$H^0(C, \Pi^*(\mathcal{L})|_C)^L \simeq H^0(Y(\mathfrak{v}), \Pi^*\mathcal{L})^G.$$

By the projection formula and Proposition 12,

$$H^0(Y(\mathfrak{v}), \Pi^*\mathcal{L}) = H^0(X, \mathcal{L} \otimes \Pi_* \mathcal{O}_{Y(\mathfrak{v})}) = H^0(X, \mathcal{L}) \oplus H^0\left(X, \mathcal{L}\left(-\frac{1}{2}[B_\Pi]\right)\right).$$

But by the Borel-Weil Theorem

$$H^0\left(X, \mathcal{L}\left(-\frac{1}{2}[B_\Pi]\right)\right) = V_{\zeta-\theta} \otimes V_{\hat{\zeta}-\hat{\theta}}.$$

Considering the dimensions of G -invariant subspaces, we get the theorem. \square

6. PROPERTIES OF THE BRANCH DIVISOR CLASS

6.1. The Belkale-Kumar product. Fix a one-parameter subgroup τ of T and set $P = P(\tau)$. For $w \in W^P$, recall that $\tau^w = \tau_{w^\vee}$ is a class of degree $\ell(w)$, see Section 2.1. For $w \in W^P$, define the BK degree of $\tau^w \in A^*(G/P)$ to be

$$\text{BK-deg}(\tau^w) := \langle w^{-1}(\rho) - \rho, \tau \rangle.$$

Let w_1, w_2 and w_3 in W^P . By [BK06, Proposition 17], if $c(w_1^\vee, w_2^\vee, w_3) \neq 0$ that is if τ^{w_3} appears in the product $\tau^{w_1} \cdot \tau^{w_2}$ then

$$(61) \quad \text{BK-deg}(\tau^{w_3}) \leq \text{BK-deg}(\tau^{w_1}) + \text{BK-deg}(\tau^{w_2}).$$

In other words the BK degree filters the Chow ring. Let \odot denotes the associated graded product on $A^*(G/P)$. For later use, note that \odot coincides with the usual product when G/P is cominuscule (See [BK06]).

Assume now that G is embedded in \hat{G} and set $\hat{P} = \hat{P}(\tau)$. Recall that $\iota : G/P \rightarrow \hat{G}/\hat{P}$ denotes the inclusion. In [RR11] a graded ring morphism is defined

$$\iota^\odot : (A^*(\hat{G}/\hat{P}), \odot) \rightarrow (A^*(G/P), \odot).$$

As in (12), define the integers $c^\odot(v, \hat{v})$ by

$$(62) \quad \forall \hat{v} \in W^{\hat{P}}, \quad \iota^\odot(\tau_{\hat{v}}) = \sum_{v \in W^P} c^\odot(v, \hat{v}) \tau^v.$$

Fix $\mathfrak{v} = (v, \hat{v}) \in W^{\mathbb{P}}$ such that $\ell(\mathfrak{v}) = \dim(\hat{G}/\hat{P})$. Consider now the incidence variety $\Pi : Y(\mathfrak{v}) = G \times^P \overline{C^+} \rightarrow \mathbb{G}/\mathbb{B}$ as above. Note that $c(\mathfrak{v})$ is not zero if and only if Π is dominant. By equivariance this means that there exists a point $x \in C^+$ such the tangent map of Π at $[e : x]$ is invertible.

By [RR11, Proposition 2.3] and the definition of ι° , the coefficient $c^\circ(\mathfrak{v})$ is not zero if and only if there exists $y_0 \in C$ such that the tangent map of Π at $[e : y_0]$ is invertible. Moreover, if $c^\circ(\mathfrak{v}) \neq 0$, then $\tau(\mathbb{C}^*)$ acts trivially on the line bundle $(K_\Pi)_{|C}$, so that for $y \in C^+$ with $y_0 = \lim_{t \rightarrow 0} \tau(t)y$, the tangent map of Π at $[e : y]$ is invertible if and only if the tangent map of Π at $[e : y_0]$ is. Writing $y_0 = (l, \hat{l}) \cdot \mathfrak{v}^{-1} \mathbb{B}/\mathbb{B}$, the tangent map of Π at $[e : y_0]$ is invertible if and only if the intersection

$$\iota(lv^{-1}BvP/P) \cap (\hat{l}\hat{v}^{-1}\hat{B}\hat{v}\hat{P}/\hat{P})$$

is transverse at P/P in \hat{G}/\hat{P} . In [BK06], this condition is called Levi-movability. To sum up, we obtain:

Lemma 35. *In the setting of Theorem 7, assume that*

$$\tau_v \circ \iota^\circ(\tau_{\hat{v}}) = k[pt],$$

with $k > 0$. Then, for general (l, \hat{l}) in \mathbb{L} , the intersection $\iota(lv^{-1}BvP/P) \cap (\hat{l}\hat{v}^{-1}\hat{B}\hat{v}\hat{P}/\hat{P})$ is transverse at P/P and the tangent map of Π at $[e : y_0]$ is invertible, with $y_0 = (l, \hat{l}) \cdot \mathfrak{v}^{-1} \mathbb{B}/\mathbb{B}$.

6.2. The branch divisor class in the Levi-movable case. Consider the support

$$\Gamma(G, \hat{G}) = \{(\zeta, \hat{\zeta}) \in X(T)^+ \times X(\hat{T})^+ : m_{G \subset \hat{G}}(\zeta, \hat{\zeta}) \neq 0\}$$

of the branching multiplicities. It is known (see [Éla92]) to be a finitely generated semigroup and hence it generates a convex polyhedral cone $\Gamma_{\mathbb{Q}}(G, \hat{G}) \subset (X(T) \oplus X(\hat{T})) \otimes \mathbb{Q}$, that we call Horn cone since the introduction. Let S be the center of the Levi subgroup L of G . Given $\mathfrak{v} = (v, \hat{v}) \in W^{\mathbb{P}}$ such that $c(\mathfrak{v}) \neq 0$, the points $(\zeta, \hat{\zeta})$ in $\Gamma_{\mathbb{Q}}(G, \hat{G})$ satisfy

$$\langle v^{-1}(\zeta), \tau \rangle + \langle \hat{v}^{-1}(\hat{\zeta}), \tau \rangle \leq 0$$

for any dominant (for G) one-parameter subgroup τ of S . In particular, the set of pairs $(\zeta, \hat{\zeta}) \in (X(T) \times X(\hat{T})) \cap \Gamma_{\mathbb{Q}}(G, \hat{G})$ such that $v^{-1}(\zeta)|_S + \hat{v}^{-1}(\hat{\zeta})|_S$ is trivial is a face $\mathcal{F}(\mathfrak{v})$ of the Horn cone.

Under the Levi-movability assumption, the branch divisor class $\frac{1}{2}[B_\Pi]$ is the minimal element of $\mathcal{F}(\mathfrak{v})$ that does not satisfy the conclusion of Theorem 5:

Proposition 36. *In the setting of Theorem 7, assume that $c^\circ(\mathfrak{v}) > 0$. Then $\frac{1}{2}[B_\Pi] \in \mathcal{F}(\mathfrak{v})$, and we have, for any $(\zeta, \hat{\zeta}) \in \mathcal{F}(\mathfrak{v})$,*

$$m_{G \subset \hat{G}}(\zeta, \hat{\zeta}) \leq m_{L \subset \hat{L}}(v^{-1}(\zeta), \hat{v}^{-1}(\hat{\zeta})).$$

Assume more specifically that $c^\circ(\mathfrak{v}) = 2$. Then the set of pairs $(\zeta, \hat{\zeta}) \in \mathcal{F}(\mathfrak{v})$ where this inequality is strict is exactly the set

$$\frac{1}{2}[B_\Pi] + (\mathcal{F}(\mathfrak{v}) \cap \Gamma(G, \hat{G})).$$

Proof. First, let us prove that $[B_\Pi]$ belongs to the face $\mathcal{F}(\mathfrak{v})$. If $c^\circ(\mathfrak{v}) = 1$, then $\frac{1}{2}[B_\Pi] = 0$ and there is nothing to prove. We assume $c^\circ(\mathfrak{v}) > 1$. Recall that $\mathcal{O}(B_\Pi) = \mathcal{L}(\theta, \hat{\theta}) \in \text{Pic}(\mathbb{G}/\mathbb{B})$. We have $(\theta, \hat{\theta}) \in \mathcal{F}(\mathfrak{v})$ if and only if $v^{-1}(\theta)|_S + \hat{v}^{-1}(\hat{\theta})|_S$ is trivial that is if S acts trivially on $\mathcal{O}(B_\Pi)_{|C}$.

We pick $(l, \hat{l}) \in \mathbb{L}$ as provided by Lemma 35 and we set $y_0 = (l, \hat{l}) \cdot \mathfrak{v}^{-1} \mathbb{B}/\mathbb{B}$. In particular $[e : y_0] \in Y(\mathfrak{v}) = G \times^P \overline{C^+}$ is an isolated point in the fiber $\Pi^{-1}(y_0)$. By transversality and degree assumption, the fiber $\Pi^{-1}(y_0)$ cannot be reduced to this point. In particular, this fiber is not connected. By Proposition 13, y_0 does not belong to the support $\text{Supp}(B_\Pi)$ of B_Π .

Since $\mathcal{O}(B_\Pi)$ is the line bundle on the smooth variety X associated to some divisor with $\text{Supp}(B_\Pi)$ as support, $\mathcal{O}(B_\Pi)$ has a canonical section σ that does not vanish at y_0 . By unicity up to scalar multiplication of this section and G -invariance of $\text{Supp}(B_\Pi)$, it has to be an eigenvector for the action of G . Since G is semi-simple this section is a G -invariant element of $H^0(X, \mathcal{O}(B_\Pi))$, meaning that σ is a G -equivariant section of $\mathcal{O}(B_\Pi)$.

But S fixes y_0 and hence acts on the fiber $\mathcal{O}(B_\Pi)_{y_0}$. This action has to fix $\sigma(y_0)$ and hence is trivial.

Let $f \in H^0(\mathbb{G}/\mathbb{B}, \mathcal{L})^G$ that restricts to 0 on C . Then f vanishes on C^+ and thus $\Pi^* f$ vanishes on $Y(\mathfrak{v})$. Thus, $H^0(\mathbb{G}/\mathbb{B}, \mathcal{L})^G$ injects into $H^0(C, \mathcal{L}_{|C})^L$, and the inequality of the proposition follows.

Assuming that $c^\odot(\mathfrak{v}) = 2$, Theorem 6 shows that the inequality will be strict if and only if $m_{G\subset\hat{G}}(\zeta - \theta, \hat{\zeta} - \hat{\theta}) \neq 0$, which means that $(\zeta, \hat{\zeta})$ belongs to $(\theta, \hat{\theta}) + (\mathcal{F}(\mathfrak{v}) \cap \Gamma(G, \hat{G}))$. \square

Corollary 37. *In the setting of Theorem 6, assume that $c^\odot(v, \hat{v}) = 2$. Then*

$$m_{G\subset\hat{G}}(\zeta, \hat{\zeta}) = \sum_{k \geq 0} (-1)^k m_{L\subset\hat{L}}(v^{-1}(\zeta - k\theta), \hat{v}^{-1}(\hat{\zeta} - k\hat{\theta})).$$

Proof. Direct induction from Theorem 6. Indeed Proposition 36 allows to apply the theorem to each weight $(\zeta - k\theta, \hat{\zeta} - k\hat{\theta})$. \square

7. EXAMPLES

7.1. Type A. In this section, we assume that G has type A . In type A_{n-1} , it is more convenient to work with representations of $\mathrm{GL}_n(\mathbb{C})$ instead of $\mathrm{SL}_n(\mathbb{C})$. For $\mathrm{GL}_n(\mathbb{C})$, we have

$$\Lambda_{\mathrm{GL}_n}^+ := X(T)^+ = \left\{ \sum_{i=1}^n \lambda_i \varepsilon_i : \lambda_1 \geq \dots \geq \lambda_n \in \mathbb{Z} \right\},$$

with notation as in [Bou02]. Given $\lambda \in X(T)^+$, set $\bar{\lambda} = \sum_{i=1}^{n-1} (\lambda_i - \lambda_n) \varepsilon_i$. Then $\bar{\lambda}$ is a partition and the $\mathrm{GL}_n(\mathbb{C})$ -representation V_λ of highest weight λ is isomorphic to the $\mathrm{SL}_n(\mathbb{C})$ -representation of highest weight $\bar{\lambda}$ as an $\mathrm{SL}_n(\mathbb{C})$ -representation. For $\lambda, \mu, \nu \in \Lambda_{\mathrm{GL}_n}^+$, set $m_{\mathrm{GL}_n}(\lambda, \mu, \nu) = \dim(V_\lambda \otimes V_\mu \otimes V_\nu)^{\mathrm{GL}_n(\mathbb{C})}$. Because of the action of the center of $\mathrm{GL}_n(\mathbb{C})$, we have

$$m_{\mathrm{GL}_n}(\lambda, \mu, \nu) \neq 0 \implies |\lambda| + |\mu| + |\nu| = 0,$$

where $|\lambda| = \sum_i \lambda_i$ and $|\mu|$ and $|\nu|$ are defined similarly. Set $\nu^* = \sum_i -\nu_{n+1-i} \varepsilon_i$ so that V_{ν^*} is the dual representation of V_ν . If λ, μ and ν^* are partitions (that is $\lambda_n, \mu_n, \nu_n^* \geq 0$) then

$$(63) \quad m_{\mathrm{GL}_n}(\lambda, \mu, \nu) = c_{\lambda, \mu}^{\nu^*}$$

is the LR coefficient. The Horn semigroup of $\mathrm{GL}_n(\mathbb{C})$ is

$$\Gamma(\mathrm{GL}_n(\mathbb{C})) = \{(\lambda, \mu, \nu) \in (\Lambda_{\mathrm{GL}_n}^+)^3 : m_{\mathrm{GL}_n}(\lambda, \mu, \nu) \neq 0\}.$$

Let $1 \leq r \leq n-1$ and consider $\mathrm{Gr}(r, n) = G/P$. Given a subset $I \subset \{1, \dots, n\}$ with r elements, define $v_I \in W^P$ by $v_I(\{1, \dots, r\}) = I$ and set $I^\vee = \{n+1-i \mid i \in I\}$. Recall from (2) in the introduction that $c_{I, J}^K$ denotes the structure constants of the Chow ring of the Grassmannian $\mathrm{Gr}(r, n)$ equipped with its Schubert basis. For any subsets $I, J, K \subset \{1, \dots, n\}$ with r elements, we have

$$(64) \quad c(v_I, v_J, v_K) = c_{I, J}^K.$$

Recall also that given a partition λ and a subset I , λ_I is the partition whose parts are λ_i with $i \in I$. Assume that $c(v_I, v_J, v_K) \neq 0$. The associated Horn inequality (3) expressed in terms of $\Gamma(\mathrm{GL}_n(\mathbb{C}))$ becomes

$$(65) \quad (\lambda, \mu, \nu) \in \Gamma(\mathrm{GL}_n(\mathbb{C})) \implies |\lambda_I| + |\mu_{J^\vee}| + |\nu_{K^\vee}| \leq 0.$$

Two interpretations. Fix $0 < r < n$. Let $I = \{i_1 < \dots < i_r\}$ be a subset of $\{1, \dots, n\}$ with r elements. Let

$$\lambda(I) = i_r - r \geq \dots \geq i_1 - 1$$

be the associated partition.

An amazing property of the LR coefficients is that they are also the multiplicities of the tensor product decomposition for the linear groups. Namely, $c_{I, J}^K$ is also the multiplicity $c_{\lambda(I), \lambda(J)}^{\lambda(K)}$ of $V_{\mathrm{SL}_r}(\lambda(K))$ in $V_{\mathrm{SL}_r}(\lambda(I)) \otimes V_{\mathrm{SL}_r}(\lambda(J))$. Hence

$$(66) \quad c_{I, J}^K = c_{\lambda(I), \lambda(J)}^{\lambda(K)}.$$

In this section, we only consider the case $G \subset \hat{G} = G \times G$, for a given classical group G of type A or B . In particular, we use Notation 1. In each case, we have a vector space V endowed with a basis (e_1, \dots, e_n)

and eventually a symmetric bilinear form. In type B , we choose the bilinear form β such that $\beta(e_i, e_j) = 1$ if and only if $j = n + 1 - i$, and equals 0 otherwise. With this convention, the set of diagonal resp. triangular matrices in G is a maximal torus T resp. Borel subgroup B .


The classical Grassmannians $\text{Gr} = G/P$ occur. The Schubert varieties in Gr are indexed by subsets I of $\{1, \dots, n\}$ in the following way:

$$X_I = \overline{B \cdot V_I} \text{ with } V_I := \text{Span}(e_i : i \in I), \quad \text{and} \quad \tau_I = [X_I] \in A^{\text{codim}(X_I)}(\text{Gr}).$$

This also gives a bijection between a set of subsets of $\{1, \dots, n\}$ and W^P . Moreover, set $\bar{i} = n + 1 - i$. and $I^\vee = \{\bar{i} \mid i \in I\}$. This operation is the Poincaré duality: $\tau^I = \tau_{I^\vee}$.

In type A , we also consider the two steps flag manifolds $\text{Fl}(p, q; n) = \{F_1 \subset F_2 \subset V : \dim(F_1) = p, \dim(F_2) = q\}$. The Schubert varieties are then indexed by “flags” $(I_p \subset I_q)$ of subsets of $\{1, \dots, n\}$: $X_{(I_p \subset I_q)} = \overline{B \cdot (V_{I_p}, V_{I_q})}$.

7.2. A detailed example in $\text{Gr}(3, 6)$. In this subsection, $G = \text{SL}_6$, $G/P = \text{Gr}(3, 6)$, and $\hat{G} = G^2$. We let $I = J = K = \{2, 4, 6\} \in W^P$. Set $\mathfrak{v} = (v_I, v_J, v_K)$. The inversion set $\Phi(v_I)$ of v_I is depicted with black nodes:



Since $\lambda(I^\vee) = \lambda(J^\vee) = 21$ and $\lambda(K) = 321$, we have $c(\mathfrak{v}) = c_{21, 21}^{321} = 2$, by (64) and (66). We describe geometrically the divisors R_Π and B_Π in this example, but we start with Lemma 38 about configurations of triangles in the plane.

Definition 3. Given a 3-dimensional vector space E , a sextuple $(A_1, A_2, A_3, B_1, B_2, B_3)$ of points in the projective plane $\mathbb{P}E$ is called a bitriangle if the three sets of points $\{A_i, A_j, B_k\}$ for i, j, k distinct in $\{1, 2, 3\}$ are colinear. In the generic situation, A_1, A_2, A_3 define a triangle and the three points B_1, B_2, B_3 are on the sides of this triangle.

The variety of all bitriangles in $\mathbb{P}E$ will be denoted by $\mathcal{T}(E)$. Given two vector spaces E and E' of dimension 3, and two bitriangles

$$(67) \quad T = (A_1, A_2, A_3, B_1, B_2, B_3) \in \mathcal{T}(E), \quad T' = (A'_1, A'_2, A'_3, B'_1, B'_2, B'_3) \in \mathcal{T}(E'),$$

a morphism $u : T \rightarrow T'$ is a linear map $u : E \rightarrow E'$ such that for all i , $u(A_i) \subset A'_i$ and $u(B_i) \subset B'_i$ (here A_i, B_i resp. A'_i, B'_i are considered as 1-dimensional subspaces of E resp. E'). Note that the set of morphisms of bitriangles is a subspace of the vector space of linear maps $E \rightarrow E'$.

Observe that $\mathcal{T}(E)$ is irreducible of dimension 9 and that the modality of the action of $\text{PSL}(E)$ is one.

Lemma 38. There is exactly one divisor \mathcal{D} in $\mathcal{T}(E) \times \mathcal{T}(E')$ such that for all pairs (T, T') in \mathcal{D} , the space of morphisms from T to T' is not reduced to $\{0\}$. It may be described as the divisor of pairs of isomorphic bitriangles.

Proof. First of all, the variety \mathcal{D} of pairs of isomorphic bitriangles is a divisor, since the modality of the action of $\text{PSL}(E)$ on the space of bitriangles is 1. Moreover, tautologically, if T and T' are isomorphic, then the space of morphisms $T \rightarrow T'$ is not reduced to $\{0\}$.

On the other hand, let $u : T \rightarrow T'$ be a morphism of bitriangles, with T and T' as in (67). If u has rank 3, then T and T' are isomorphic and $(T, T') \in \mathcal{D}$.

Assume first that T is generic. If u has rank 2, then at least 5 of the six points $A_1, A_2, A_3, B_1, B_2, B_3$ have a well defined image in $\mathbb{P}E'$ and all these images belong to the line in $\mathbb{P}E'$ defined by the image of u . It follows that 5 of the 6 points $A'_1, A'_2, A'_3, B'_1, B'_2, B'_3$ are on a line, and the set of such bitriangles has codimension 2 in $\mathcal{T}(E')$. If u has rank 1, similarly, we don't find a divisor in $\mathcal{T}(E')$ since 3 of the six points $A'_1, A'_2, A'_3, B'_1, B'_2, B'_3$ must be equal in this case. The case where T is degenerate is similar. \square

Proposition 39. *The hypersurface B_Π is irreducible, it is equal to the variety Δ defined below in (70), and*

$$\mathcal{O}_{(G/B)^3}(B_\Pi) = \mathcal{L}_{(G/B)^3}(2\varpi_2 + 2\varpi_4, 2\varpi_2 + 2\varpi_4, 2\varpi_2 + 2\varpi_4).$$

The hypersurface R_Π is also irreducible, it is the preimage of B_Π by Π , and it also has the description given below in (71).

Proof. Let $U \subset (G/B)^3$ be the open subset of triples $({}_1X_\bullet, {}_2X_\bullet, {}_3X_\bullet)$ such that $\mathbb{C}^6 = {}_1X_2 \oplus {}_2X_2 \oplus {}_3X_2$, ${}_1X_4 \cap {}_2X_4 \cap {}_3X_4 = \{0\}$, and ${}_iX_4 \cap {}_jX_2 = \{0\}$ for $i \neq j$. We first investigate the intersection $B_\Pi \cap U$. In fact, let $({}_1X_\bullet, {}_2X_\bullet, {}_3X_\bullet) \in U$: the fiber $\Pi^{-1}({}_1X_\bullet, {}_2X_\bullet, {}_3X_\bullet)$ can be explicitly described as follows.

Let $V_3 \in \text{Gr}(3, 6)$. Then $(V_3; {}_1X_\bullet, {}_2X_\bullet, {}_3X_\bullet)$ is an element of $Y(\mathfrak{v})$ if and only if

$$(68) \quad \forall i \in \{1, 2, 3\}, \dim V_3 \cap {}_iX_2 \geq 1 \text{ and } \dim V_3 \cap {}_iX_4 \geq 2.$$

From $\dim V_3 \cap {}_iX_2 \geq 1$ and the genericity assumption, we deduce that V_3 can be written as

$$(69) \quad V_3 = L_1 \oplus L_2 \oplus L_3,$$

with $L_i \subset {}_iX_2$ a subspace of dimension 1. Moreover, under our genericity assumption, we have an isomorphism ${}_iX_2 \simeq \mathbb{C}^6 / {}_kX_4$ if $i \neq k$. If i, j, k are distinct, we get isomorphisms ${}_iX_2 \simeq \mathbb{C}^6 / {}_kX_4$ and $\mathbb{C}^6 / {}_kX_4 \simeq {}_jX_2$, and we denote $\varphi_{i,j} : {}_iX_2 \rightarrow {}_jX_2$ the isomorphism obtained by composition.

Observe that $L_1 \oplus L_2 \oplus L_3$ will meet ${}_kX_4$ in dimension 2 if and only if the subspace of \mathbb{C}^6 generated by L_i, L_j and ${}_kX_4$ has dimension 5, which is equivalent to the equality $L_j = \varphi_{i,j}(L_i)$.

Thus, $V_3 = L_1 \oplus L_2 \oplus L_3$ satisfies (68) if and only if for $i \neq j$, $L_j = \varphi_{i,j}(L_i)$. In other words, $L_2 = \varphi_{1,2}(L_1)$, $L_3 = \varphi_{2,3}(L_2)$, and $L_1 \subset {}_1X_2$ is an eigenline of $\varphi_{3,1} \circ \varphi_{2,3} \circ \varphi_{1,2}$.

By Proposition 13, $B_\Pi \cap U$ is the set of triples $({}_1X_\bullet, {}_2X_\bullet, {}_3X_\bullet)$ such that $\varphi_{3,1} \circ \varphi_{2,3} \circ \varphi_{1,2}$ has only one eigenvalue or is the identity.

We denote by Δ this divisor:

$$(70) \quad \Delta = \{({}_1X_\bullet, {}_2X_\bullet, {}_3X_\bullet) \mid \varphi_{3,1} \circ \varphi_{2,3} \circ \varphi_{1,2} \text{ has only one eigenvalue}\}$$

We now show that there is only one divisor in the ramification of Π . To show this, we observe that for a general element $(V_3, {}_1X_\bullet, {}_2X_\bullet, {}_3X_\bullet)$ of a divisor of $Y(\mathfrak{v})$, $\dim(V_3 \cap {}_iX_2) = 1$ and $\dim(V_3 \cap {}_iX_4) = 2$ for $i \in \{1, 2, 3\}$. Therefore, this defines a bitriangle T in $\mathbb{P}V_3$, with vertices $A_i = V_3 \cap {}_iX_2$ and edges $E_i = V_3 \cap {}_iX_4$, and this defines also a bitriangle T' in \mathbb{C}^6 / V_3 , with vertices $A'_i = p(V_3 \cap {}_iX_2)$ and edges $E'_i = p({}_iX_4)$, where $p : \mathbb{C}^6 \rightarrow \mathbb{C}^6 / V_3$ denotes the projection.

In this way, we factorize the morphism $Y(\mathfrak{v}) \rightarrow \text{Gr}(3, 6)$ as a composition of rational maps $Y(\mathfrak{v}) \xrightarrow{f} \mathcal{T}(E) \times_{\text{Gr}(3,6)} \mathcal{T}(Q) \rightarrow \text{Gr}(3, 6)$, where E and Q denote the tautological bundles on $\text{Gr}(3, 6)$, and $\mathcal{T}(E)$ and $\mathcal{T}(Q)$ denote the relative varieties of bitriangles therein. The rational map f is defined in codimension 1 and equidimensional. It follows that $f(R_\Pi)$ is defined and at least a divisor in $\mathcal{T}(E) \times \mathcal{T}(Q)$.

On the other hand, an element $u \in T_{V_3} \text{Gr}(3, 6) \simeq \text{Hom}(V_3, \mathbb{C}^6 / V_3)$ belongs to the tangent space to the Schubert variety defined by ${}_iX_\bullet$ if and only if $u(A_i) \subset A'_i$ and $u(E_i) \subset E'_i$. Since this must hold for all $i \in \{1, 2, 3\}$, u defines a morphism of bitriangles (recall Definition 3). It thus follows from Lemma 38 that the image of R_Π by f is the divisor of isomorphic bitriangles in $\mathcal{T}(E) \times \mathcal{T}(Q)$:

$$(71) \quad R_\Pi = \{y \in Y : f(y) \in \mathcal{T}(E) \times \mathcal{T}(Q) \text{ is a pair of isomorphic bitriangles.}\}$$

It follows that R_Π is irreducible, equal to $\Pi^{-1}(B_\Pi)$, and that $B_\Pi = \Delta$. Moreover, the given formula for $\mathcal{O}_{(G/B)^3}(B_\Pi)$ can be proved using Corollary 8 and can also be seen more geometrically as a consequence of (70). \square

We now check directly Theorem 6 in this case. First we give a full description of the face in the Horn cone defined by \mathfrak{v} . Recall that $I = J = K = \{2, 4, 6\}$.

Lemma 40. *The span of the face \mathcal{F}_{IJK} of $\Gamma(\mathrm{GL}_6)$ defined by the equation $|\lambda_I| + |\mu_J| + |\nu_K| = 0$ has equations*

$$(72) \quad \begin{cases} |\lambda| + |\mu| + |\nu| = 0; \\ \lambda_1 = \lambda_2, \lambda_3 = \lambda_4, \text{ and } \lambda_5 = \lambda_6; \\ \mu_1 = \mu_2, \mu_3 = \mu_4, \text{ and } \mu_5 = \mu_6; \\ \nu_1 = \nu_2, \nu_3 = \nu_4, \text{ and } \nu_5 = \nu_6. \end{cases}$$

Proof. The equalities (72) clearly imply $|\lambda_I| + |\mu_J| + |\nu_K| = 0$. On the other hand, if $|\lambda_I| + |\mu_J| + |\nu_K| = 0$, we get

$$0 = 2|\lambda_I| + 2|\mu_J| + 2|\nu_K| \leq |\lambda| + |\mu| + |\nu| = 0.$$

This proves that the middle inequality is an equality. So (72) hold. \square

Remark 3. *The face \mathcal{F}_{IJK} in Lemma 40 is contained in no face corresponding to some LR coefficient equal to one.*

Proof. Let (I', J', K') of the same cardinality r such that $c(I', J', K') \neq 0$ and the elements of \mathcal{F}_{IJK} satisfy $|\lambda_{I'}| + |\mu_{J'}| + |\nu_{K'}| = 0$. The description of the span of \mathcal{F}_{IJK} in Lemma 40 implies that the linear form $(\lambda, \mu, \nu) \mapsto |\lambda_{I'}| + |\mu_{J'}| + |\nu_{K'}|$ belongs to the span of the 10 linear forms vanished in equations (72): there are rational numbers $n, d, a_1, a_3, a_5, b_1, b_3, b_5, c_1, c_3, c_5$ such that

$$\begin{aligned} n(|\lambda_{I'}| + |\mu_{J'}| + |\nu_{K'}|) &= d(|\lambda| + |\mu| + |\nu|) \\ &\quad + a_1(\lambda_1 - \lambda_2) + a_3(\lambda_3 - \lambda_4) + a_5(\lambda_5 - \lambda_6) \\ &\quad + b_1(\mu_1 - \mu_2) + b_3(\mu_3 - \mu_4) + b_5(\mu_5 - \mu_6) \\ &\quad + c_1(\nu_1 - \nu_2) + c_3(\nu_3 - \nu_4) + c_5(\nu_5 - \nu_6) \end{aligned}$$

for any λ, μ, ν . We may assume $n = 6$. Choosing $\lambda = \mu = \nu = 1^6$, one gets $d = r$. Since $0 < r < 6$, the linear form $6(|\lambda_{I'}| + |\mu_{J'}| + |\nu_{K'}|) - r(|\lambda| + |\mu| + |\nu|)$ has positive coefficient on λ_i if and only if $i \in I'$, and negative coefficient otherwise. It follows that I', J' , and K' all intersect $\{1, 2\}$, $\{3, 4\}$ and $\{5, 6\}$ in exactly one element. In particular $r = 3$. Now, the degree condition implied by the non-vanishing of the Schubert coefficient $c(I', J', K')$ implies that $I' = J' = K' = \{2, 4, 6\}$. In particular, $c_{I', J'}^{K'} = 2$. \square

By the remark, Theorem 2 cannot be applied to the points in this face. However, applying our reduction formula, we compute explicitly those coefficients:

Lemma 41. *Let $\lambda, \mu, \nu \in \Lambda_{\mathrm{GL}_3}^+$ such that $|\lambda| + |\mu| + |\nu| = 0$. Then $m_{\mathrm{GL}_3}(\lambda, \mu, \nu)$ is equal to the number of integers in the interval*

$$[\max(\mu_1 - \lambda_2, \mu_2, -\nu_3 - \lambda_1, \mu_1 + \nu_1, -\nu_2 - \lambda_2, \mu_1 + \mu_2 + \nu_2), \min(\mu_1, -\nu_3 - \lambda_2, \mu_1 + \mu_2 + \nu_1)].$$

Moreover, set $\lambda^2 = (\lambda_1, \lambda_1, \lambda_2, \lambda_2, \lambda_3, \lambda_3)$, and define similarly μ^2 and ν^2 . Then

$$m_{\mathrm{GL}_6}(\lambda^2, \mu^2, \nu^2) = \frac{m_{\mathrm{GL}_3}(\lambda, \mu, \nu)(m_{\mathrm{GL}_3}(\lambda, \mu, \nu) + 1)}{2}.$$

Proof. The first equality follows from [KT99]. In fact, it is explained in [PN20, Proposition 9] that $c_{\lambda, \mu}^{\nu}$ is the number of integers in the interval

$$[\max(\mu_1 - \lambda_2, \mu_2, \nu_1 - \lambda_1, \mu_1 - \nu_3, \nu_2 - \lambda_2, \mu_1 + \mu_2 - \nu_2), \min(\mu_1, \nu_1 - \lambda_2, \mu_1 + \mu_2 - \nu_3)].$$

Using the fact that $V_{(\nu_1, \nu_2, \nu_3)}^* \simeq V_{(-\nu_3, -\nu_2, -\nu_1)}$, we get our formula. We want to compute $m_{\mathrm{GL}_6}(\lambda^2, \mu^2, \nu^2)$ using Corollary 37 with $\mathfrak{v} = (v_I, v_J, v_K)$ and $I = J = K = \{2, 4, 6\}$. By Proposition 39, one half of the branch divisor corresponds to the triple of partitions $((2, 2, 1, 1, 0, 0), (2, 2, 1, 1, 0, 0), (-1, -1, -2, -2, -3, -3))$. Thus, a term $m_L(v^{-1}(\gamma - k\theta), \hat{v}^{-1}(\hat{\gamma} - k\hat{\theta}))$ in Corollary 37 is equal in our context to

$$\left(m_{\mathrm{GL}_3}(\lambda - k(210), \mu - k(210), \nu + k(123)) \right)^2.$$

Observe that by the first point, $m_{\mathrm{GL}_3}(\lambda - k(210), \mu - k(210), \nu + k(123))$ is $m_{\mathrm{GL}_3}(\lambda, \mu, \nu) - k$ or 0. Indeed, when λ, μ, ν are replaced respectively by $\lambda - (210), \mu - (210), \nu + (123)$, all the integers that appear in the max decrease by 1 and all the integers that appear in the min decrease by 2. Corollary 37 therefore gives

$$m_{\mathrm{GL}_6}(\lambda^2, \mu^2, \nu^2) = \sum_{k=0}^{m_{\mathrm{GL}_3}(\lambda, \mu, \nu)} (-1)^{m_{\mathrm{GL}_3}(\lambda, \mu, \nu) + k} k^2.$$

This is equal to $\frac{m_{\text{GL}_3}(\lambda, \mu, \nu)(m_{\text{GL}_3}(\lambda, \mu, \nu)+1)}{2}$, as stated. \square

Remark 4. As we can observe here, the Littlewood-Richardson coefficients are close to being polynomials in the coefficients of the weights. This is a general phenomenon. Kostant [Kos58, Theorem 6.2] proves that the multiplicity of a weight in an irreducible representation is a partition function. Steinberg [Ste61] deduces that the multiplicity of an irreducible submodule in the tensor product of two representations is again a partition function. Rassart [Ras04, Theorem 4.1] deduces that the Horn cone can be subdivided into subcones where the Littlewood-Richardson coefficients are polynomial in the three weights (this holds in type A; in general the coefficients are only quasi-polynomial).

7.3. A bigger example. In the following example, the face of $\Gamma(G)$ associated to the LR-coefficient equal to two is contained in a face associated to some LR-coefficient equal to one. Hence, one can apply both Theorems 2 and 3. The conclusion is that, even if Theorem 2 apply, our Theorem 3 is useful.

Here $n = 10$, $r = 6$ and $G/P = \text{Gr}(6, 10)$. Set $I = \{2, 4, 5, 6, 9, 10\}$, $J = \{3, 4, 6, 7, 9, 10\}$ and $K = \{2, 4, 5, 7, 8, 10\}$. Set $\mathfrak{v} = (v_I, v_J, v_K)$ in such a way that $c(\mathfrak{v}) = c_{\lambda(I^V), \lambda(J^V)}^{\lambda(K)} = c_{3222, 2211}^{433221} = 2$. Then, applying Theorem 7, one gets:

$$(73) \quad \frac{1}{2}[B_{\Pi}] = \mathcal{L}_{(G/B)^3}(\varpi_2 + 2\varpi_6, \varpi_4 + \varpi_7, \varpi_2 + \varpi_5 + \varpi_8).$$

First, we prove a statement analogous to Lemma 40:

Lemma 42. *The span of the face \mathcal{F}_{IJK} defined by the equation $|\lambda_I| + |\mu_J| + |\nu_K| = 0$ in $\Gamma(\text{GL}_{10})$ has equations*

$$(74) \quad \begin{cases} |\lambda| + |\mu| + |\nu| = 0; \\ (\lambda_5 + \lambda_6) + (\mu_7 + \mu_{10}) + (\nu_5 + \nu_8) = 0; \\ \lambda_1 = \lambda_2, \lambda_3 = \lambda_4, \text{ and } \lambda_7 = \lambda_8 = \lambda_9 = \lambda_{10}; \\ \mu_1 = \mu_2 = \mu_3 = \mu_4, \mu_5 = \mu_6, \text{ and } \mu_8 = \mu_9; \\ \nu_1 = \nu_2, \nu_3 = \nu_4, \nu_6 = \nu_7, \text{ and } \nu_9 = \nu_{10}. \end{cases}$$

Proof. Let F be the linear space defined by the equations (74). We observe that the following triples (λ, μ, ν) span F and satisfy $m_{\text{GL}_{10}}(\lambda, \mu, \nu) = 1$:

$$\begin{array}{cccc} (1^{10}, 0, (-1)^{10}) & (1^{10}, 0, (-1)^{10}) & (1^{10}, 0, (-1)^{10}) & (1^6, (-1)^4(-2)^6, 1^{10}) \\ (1^6, (-1)^{10}, 1^4) & (1^5, (-1)^{10}, 1^5) & (1^2, (-1)^{10}, 1^8) & (1^5, (-1)^7(-2)^3, 1^8) \\ (1^6, (-1)^9 - 2, 1^5) & (1^6, (-1)^7(-2)^3, 1^7) & (1^6, (-1)^6(-2)^4, 1^8) & (1^6, (-1)^4(-2)^3(-3)^3, 2^5 1^3) \\ (1^6, (-1)^7(-2)^3, 2^2 1^3) & (1^6, (-1)^4(-2)^6, 2^2 1^6) & (1^6, (-1)^4(-2)^3(-3)^3, 2^5 1^3) & (2^4 1^2, (-2)^7(-3)^3, 2^5 1^3) \end{array}$$

Conversely, let (λ, μ, ν) satisfying $|\lambda_I| + |\mu_J| + |\nu_K| = 0$ and $m_{\text{GL}_{10}}(\lambda, \mu, \nu) \neq 0$. To have nicer formulas, we write a instead of 10. We have

$$(75) \quad \begin{aligned} 0 = 2|\lambda_I| + 2|\mu_J| + 2|\nu_K| &\leq |\lambda_{\{12\}}| + |\lambda_{\{34\}}| + 2\lambda_5 + 2\lambda_6 + |\lambda_{\{789a\}}| \\ &+ |\mu_{\{1234\}}| + |\mu_{\{56\}}| + 2\mu_7 + |\mu_{\{89\}}| + 2\mu_a \\ &+ |\nu_{\{12\}}| + |\nu_{\{34\}}| + 2\nu_5 + |\nu_{\{67\}}| + 2\nu_8 + |\nu_{\{9a\}}| \\ &= |\lambda_{\{56\}}| + |\mu_{\{7a\}}| + |\nu_{\{58\}}| \\ &\leq 0 \end{aligned}$$

The last inequality is the Horn inequality associated to the LR coefficient $c(v_{\{56\}}, v_{\{7a\}}, v_{\{58\}}) = c_{44,2}^{64} = 1$. It follows that all inequalities in (75) must be equalities, which implies that (λ, μ, ν) belongs to F . \square

Remark 5. We do not know about a general procedure to compute the face given by a triple I, J, K with $c(v_I, v_J, v_K) > 1$. When $c(v_I, v_J, v_K) = 1$, it is known that the corresponding face has codimension 1, but when $c(v_I, v_J, v_K) > 1$ it seems an interesting problem to determine the linear span of the corresponding face, or at least its dimension.

Remark 6. In this case, Lemma 42 shows that the face \mathcal{F}_{IJK} is contained in some regular face associated to some LR coefficient equal to one (namely $c(v_{\{56\}}, v_{\{7a\}}, v_{\{58\}}) = 1$). Then both Theorems 2 and 4 can be applied to the points of the face. These two statements are not concurrent but complementary. Indeed, by

applying these results consecutively, one gets an expression of each LR coefficient on the face of Lemma 42 as an alternating sum of products of three LR coefficients. The use of a two step flag variety allows to recover this result from Corollary 37 more conceptually. Indeed, set $I_1 = \{5, 6\}$, $J_1 = \{7, 10\}$ and $K_1 = \{5, 8\}$. Set also $I_2 = I$, $J_2 = J$, and $K_2 = K$. The pairs $(I_1 \subset I_2)$, $(J_1 \subset J_2)$ and $(K_1 \subset K_2)$ define three Schubert classes in $\text{Fl}(2, 6; 10)$.

Consider the fibration $\text{Fl}(2, 6; 10) \rightarrow \text{Gr}(6, 10)$ with fiber $\text{Gr}(2, 6)$. The three given Schubert varieties in $\text{Fl}(2, 6; 10)$ map onto X_{I_2} , X_{J_2} and X_{K_2} respectively with fibers isomorphic to $X_{\{34\}}$, $X_{\{46\}}$ and $X_{\{35\}}$ in $\text{Gr}(2, 6)$. The associated Schubert coefficient in $\text{Gr}(6, 10)$ is 2 and in $\text{Gr}(2, 6)$ it is $c_{22,1}^2 = 1$. Then (see e.g. [Ric12]), the associated Schubert coefficient for $\text{Fl}(2, 6; 10)$ is 2.

Similarly, consider the fibration $\text{Fl}(2, 6; 10) \rightarrow \text{Gr}(2, 10)$ with fiber $\text{Gr}(4, 8)$. Here we find a Schubert coefficient 1 in the base and 2 in the fiber.

Recalling the BK coefficients c° from (62), these two assertions imply that

$$(76) \quad c^\circ((I_1 \subset I_2), (J_1 \subset J_2), (K_1 \subset K_2)) = 2$$

and we are in position to apply Corollary 37. Since the incidence variety \tilde{Y} in $\text{Fl}(2, 6; 10) \times (G/B)^3$ maps birationally on that corresponding to (I_2, J_2, K_2) in $\text{Gr}(6, 10)$ these two varieties have the same branch divisor class given by (73):

$$\frac{1}{2}[B_\Pi] = \mathcal{L}(3^2 2^4, 2^4 1^3, 3^2 2^3 1^3).$$

Let $(\lambda, \mu, \nu) \in (\Lambda_{\text{GL}_{10}}^+)^3$ on the span of the considered face (see Lemma 42):

$$(77) \quad \begin{aligned} \lambda &= \lambda_1^2 \lambda_3^2 \lambda_5 \lambda_6 \lambda_7^4 \\ \mu &= \mu_1^4 \mu_5^2 \mu_7 \mu_8^2 \mu_{10} \\ \nu &= \nu_1^2 \nu_3^2 \nu_5 \nu_6^2 \nu_8 \nu_9^2 \end{aligned}$$

The restriction of $\mathcal{L}(\lambda) \otimes \mathcal{L}(\mu) \otimes \mathcal{L}(\nu)$ to C is

$$\begin{aligned} \lambda_{\{56\}} &= \lambda_5 \lambda_6 & \lambda_{\{249a\}} &= \lambda_1 \lambda_3 \lambda_7^2 & \lambda_{\{1378\}} &= \lambda_1 \lambda_3 \lambda_7^2 \\ \mu_{\{7a\}} &= \mu_7 \mu_{10} & \mu_{\{3469\}} &= \mu_1^2 \mu_5 \mu_8 & \mu_{\{1258\}} &= \mu_1^2 \mu_5 \mu_8 \\ \nu_{\{58\}} &= \nu_5 \nu_8 & \nu_{\{247a\}} &= \nu_1 \nu_3 \nu_6 \nu_9 & \nu_{\{1369\}} &= \nu_1 \nu_3 \nu_6 \nu_9. \end{aligned}$$

Similarly, the restriction of $\frac{1}{2}[B_\Pi]$ to C is as an $(\text{SL}_2 \times \text{SL}_4 \times \text{SL}_4)$ -linearized line bundle

$$\begin{array}{cccc} 2 & 2 & 3 & 2 & 0 & 0 & 3 & 2 & 0 & 0 \\ 1 & 0 & 2 & 2 & 1 & 0 & 2 & 2 & 1 & 0 \\ 2 & 1 & 3 & 2 & 1 & 0 & 3 & 2 & 1 & 0. \end{array}$$

It has the same invariant sections as the $(\text{GL}_2 \times \text{GL}_4 \times \text{GL}_4)$ -linearized line bundle

$$\begin{array}{cccc} 0 & 0 & 3 & 2 & 0 & 0 & 3 & 2 & 0 & 0 \\ 1 & 0 & 2 & 2 & 1 & 0 & 2 & 2 & 1 & 0 \\ 0 & -1 & -1 & -2 & -3 & -4 & -1 & -2 & -3 & -4. \end{array}$$

Hence Corollary 37 gives:

$$\begin{aligned} m_{\text{GL}_{10}}(\lambda, \mu, \nu) &= \sum_k (-1)^k m_{\text{GL}_2}(\lambda_5 \lambda_6, \mu_{7-k} \mu_{10}, \nu_5 \nu_{8+k}) \\ &\quad \times m_{\text{GL}_4}(\lambda_{1-3k} \lambda_{3-2k} \lambda_7^2, (\mu_{1-2k})^2 \mu_{5-k} \mu_8, \nu_{1+k} \nu_{3+2k} \nu_{6+3k} \nu_{9+4k})^2. \end{aligned}$$

For $\lambda = [41, 41, 36, 36, 35, 24, 0, 0, 0, 0]$, $\mu = [-41, -41, -41, -41, -48, -48, -49, -65, -65, -72]$ and $\nu = [49, 49, 42, 42, 40, 25, 25, 22, 2, 2]$, this formula gives

$$6 = 1 \times 3^2 - 1 \times 2^2 + 1 \times 1^2.$$

Given a point as in (77), one can also apply Theorem 2 to the triple (I_1, J_1, K_1) to get

$$m_{\text{GL}_{10}}(\lambda, \mu, \nu) = m_{\text{GL}_2}(\lambda_5 \lambda_6, \mu_7 \mu_{10}, \nu_5 \nu_8) m_{\text{GL}_8}(\lambda_1^2 \lambda_3^2 \lambda_7^4, \mu_1^4 \mu_5^2 \mu_8^2, \nu_1^2 \nu_3^2 \nu_6^2 \nu_9^2).$$

But, for $\tilde{I} = \{2, 4, 7, 8\}$, $\tilde{J} = \{3, 4, 6, 8\}$ and $\tilde{K} = \{2, 4, 6, 8\}$ we have $c(\tilde{I}, \tilde{J}, \tilde{K}) = c_{32,221}^{4321} = 2$ and one can apply Theorem 4 to the second factor. Since $\frac{1}{2}[B_\Pi] = (\varpi_2 + \varpi_4, \varpi_4 + \varpi_6, \varpi_2 + \varpi_6) = (2^2 1^2, 2^4 1^2, 2^2 1^4)$, one

gets that $m_{\text{GL}_{10}}(\lambda, \mu, \nu)$ is equal to

$$m_{\text{GL}_2}(\lambda_5 \lambda_6, \mu_7 \mu_{10}, \nu_5 \nu_8) \sum_{k \geq 0} (-1)^k m_{\text{GL}_4}(\lambda_1 - 2k, \lambda_3 - k, \lambda_7^2, \mu_1^2 + k, \mu_5 + 2k, \mu_8 + 3k, \nu_1 - 2k, \nu_3 - k, \nu_6 - k, \nu_9)^2.$$

Applied to the explicit example above, one gets

$$6 = 1 \times (3^2 - 2^2 + 1^2).$$

7.4. Examples in type B. In this section, we give some examples with $G = \text{Spin}(2n+1)$. New phenomena occur compared to the type A. First, the Grassmannians are not always cominuscule and these examples show that the Levi-movability assumption is needed in Proposition 36 and Corollary 37. Second, the Horn semigroup is not saturated.

Many results in this section were obtained with the help of a computer, see [CR21]. Given a weight ζ , we associate a partition λ to it with parts λ_i such that $\zeta = \sum \lambda_i \epsilon_i$, with the notation of [Bou02], (note that, given ζ all the coefficients λ_i are in \mathbb{Z} or they are all in $\frac{1}{2} + \mathbb{Z}$).

Number the simple roots as follows: $\circ_1 - \circ_2 - \dots - \circ_{n-2} - \circ_{n-1} \text{---} \circ_n$. Note that the Horn inequality associated to a triple (I, J, K) of subsets of $\{1, \dots, n\} \cup \{n+2, \dots, 2n+1\}$ such that $c(I, J, K) \neq 0$ is

$$(78) \quad |\lambda_{I \cap [1;n]}| - |\lambda_{I \cap [n+2;2n+1]}| + |\mu_{J \cap [1;n]}| - |\mu_{J \cap [n+2;2n+1]}| + |\nu_{K \cap [1;n]}| - |\nu_{K \cap [n+2;2n+1]}| \leq 0.$$

Let $n = 4$. In the Chow ring of the quadric Q^7 , for $I = J = \{7\}$ and $K = \{6\}$, we have $\sigma_{\{7\}} \odot \sigma_{\{7\}} \odot \sigma_{\{6\}} = 2[pt]$ and (78) means

$$(79) \quad \lambda_3 + \mu_3 + \nu_4 \geq 0.$$

Using [CR21], we get $\theta = (\varpi_2, \varpi_2, \varpi_3) = (1^2, 1^2, 1^3)$ and (79) is an equality, as predicted by Proposition 36. Here $L^{\text{ss}} = \text{Spin}(7)$. We put superscript ϖ^G and ϖ^L to make the difference between weights of G and L . We have $v_I^{-1}(\varpi_2^G) = v_I^{-1}(\epsilon_1^G + \epsilon_2^G) = \epsilon_2^G + \epsilon_3^G$, and its restriction to L is $\epsilon_1^L + \epsilon_2^L$, namely ϖ_2^L . Similarly, $(v_K^{-1}(\varpi_3^G))^L = 2\varpi_3^L$. Using the equality $m_L(\varpi_2, \varpi_2, 2\varpi_3) = 1$ and the following Remark 7, we get:

$$m_L(kv_I^{-1}\theta^1, kv_J^{-1}\theta^2, kv_K^{-1}\theta^3) = m_L(k\varpi_2, k\varpi_2, 2k\varpi_3) = 1.$$

Corollary 37 gives $m_G(k\varpi_2, k\varpi_2, k\varpi_3) = 1$ if k is even and 0 if k is odd, consistently with the non-saturation of the tensor semigroup $\Gamma(G) := \{(\zeta_1, \zeta_2, \zeta_3) \in (\Lambda_G^+)^3 : m_G(\zeta_1, \zeta_2, \zeta_3) \neq 0\}$.

Remark 7. *The group Spin_7 has a dense orbit in $\text{OG}(2, 7) \times \text{OG}(2, 7) \times \text{OG}(3, 7)$.*

Proof. We consider the isotropic space \mathbb{C}_a^2 generated by e_1 and e_2 , the isotropic space \mathbb{C}_b^2 generated by e_6 and e_7 , and the non-isotropic space \mathbb{C}^3 generated by e_3, e_4, e_5 . Let V_3 be the isotropic space generated by $v_1 = e_1 + e_3, v_2 = e_5 - e_7$ and $v_3 = e_2 + e_4 - e_6$. We claim that the isotropy of the triple $(\mathbb{C}_a^2, \mathbb{C}_b^2, V_3)$ in Spin_7 is 1-dimensional, which proves the statement since $\dim(\text{OG}(2, 7) \times \text{OG}(2, 7) \times \text{OG}(3, 7)) = 20 = \dim(\text{Spin}_7) - 1$.

Let indeed g be in the isotropy. Since g stabilizes \mathbb{C}_a^2 and \mathbb{C}_b^2 , it stabilizes \mathbb{C}^3 as it is the subspace orthogonal to $\mathbb{C}_a^2 \oplus \mathbb{C}_b^2$. The action of g on $\mathbb{C}_b^2 \simeq (\mathbb{C}_a^2)^*$ is the dual of that of g on \mathbb{C}_a^2 .

Observe that g must stabilize the line generated by v_1 , since it is the intersection of V_3 and $\mathbb{C}_a^2 \oplus \mathbb{C}^3$. Similarly it stabilizes the line generated by v_2 . It follows that there exists a scalar λ such that $g \cdot e_1 = \lambda e_1, g \cdot e_3 = \lambda e_3, g \cdot e_7 = \lambda^{-1} e_7$ and $g \cdot e_5 = \lambda^{-1} e_5$. Then g preserves $\mathbb{C}_a^2 \cap e_7^\perp$, which is the line generated by e_2 . It follows that there exists a scalar μ such that $g \cdot e_2 = \mu e_2, g \cdot e_6 = \mu^{-1} e_6$ and $g \cdot e_4 = e_4$. Since g preserves the line generated by v_3 , we have $\mu = 1$. Thus g is uniquely defined by λ and the isotropy has dimension 1. \square

In the Chow ring of $\text{OG}(4, 9)$, for $I = J = \{3, 6, 8, 9\}$ and $K = \{1, 3, 6, 8\}$, we have $\sigma_I \odot \sigma_J \odot \sigma_K = 2[pt]$. Inequality (78) means

$$(80) \quad \lambda_1 + \lambda_2 + \lambda_4 + \mu_1 + \mu_2 + \mu_4 + \nu_2 + \nu_4 \geq \lambda_3 + \mu_3 + \nu_1 + \nu_3.$$

Using [CR21], we get $\theta = (\varpi_3, \varpi_3, \varpi_1 + \varpi_3) = (1^3 0, 1^3 0, 2 1^2 0)$ and once again, (80) is an equality.

Here $L^{\text{ss}} = \text{SL}(4)$. We have $v_I^{-1}(\theta^1) = v_J^{-1}(\theta^2) = v_I^{-1}(\epsilon_1^G + \epsilon_2^G + \epsilon_3^G) = -\epsilon_4^G - \epsilon_3^G + \epsilon_1^G$ and this restricts to $2\epsilon_1^L + \epsilon_2^L = \varpi_1^L + \varpi_2^L$. We also have $v_K^{-1}(\theta^3)^L = \varpi_1^L + \varpi_2^L + \varpi_3^L$. We get $m_L(kv_I^{-1}\theta^1, kv_J^{-1}\theta^2, kv_K^{-1}\theta^3) = c_{2k, 2k, 2k}^{3k, 2k, k} = k + 1$. Corollary 37 gives $m_G(k\varpi_3, k\varpi_3, k\varpi_1 + \varpi_3) = \lceil \frac{k}{2} \rceil$.

In the Chow ring of $\text{OG}(4, 9)$, for $I = \{3, 6, 8, 9\}$ and $J = K = \{2, 4, 7, 9\}$, we have $\sigma_I \sigma_J \sigma_K = 2[pt]$ and $\sigma_I \odot \sigma_J \odot \sigma_K = 0$. Inequality (78) means

$$(81) \quad \lambda_1 + \lambda_2 + \lambda_4 + \mu_1 + \mu_3 + \nu_1 + \nu_3 \geq \lambda_3 + \mu_2 + \mu_4 + \nu_2 + \nu_4.$$

Using [CR21], we get $\theta = (\varpi_3, \varpi_2 + \varpi_4, \varpi_2 + \varpi_4) = (1^3 0, \frac{1}{2}(3^2 1^2), \frac{1}{2}(3^2 1^2))$, and this does not belong to the face. This shows that the Levi-movability assumption in Proposition 36 is necessary. The restriction of $\frac{1}{2}[B_{\Pi}]$ to C is $(21, 321, 321)$. Corollary 37 does not apply.

APPENDIX A. APPENDIX: PUZZLES

A.1. Some computations. In this appendix, we give an example for recent results [KP11, Res11a, Ric12] about the Schubert coefficients when G/P is a partial flag variety. It is a counterexample to [KP11, Theorem 3]. Since [KP11, Theorem 3] is “proved” by constructing a bijection between hives, we picture the concerned hives.

Even more precisely, we consider the flag variety $\text{Fl}(3, 6; 9)$ parametrizing flags $(V_3 \subset V_6)$ in \mathbb{C}^9 with $\dim(V_3) = 3$ and $\dim(V_6) = 6$. We have $\text{Fl}(3, 6; 9) = G/P$ and $\text{Gr}(6, 9) = \text{SL}_9/Q$ with $P \subset Q \subset G = \text{SL}_9$ the corresponding parabolic subgroups.

Consider the Schubert class $\omega = (\{3, 6, 9\} \subset \{2, 3, 5, 6, 8, 9\})$ for $\text{Fl}(3, 6; 9)$. The projection of the Schubert variety X_ω on $\text{Gr}(6, 9)$ is X_I with $I = \{2, 3, 5, 6, 8, 9\}$. Its intersection with the fiber $\text{Gr}(3, 6)$ is X_J with $J = \{2, 4, 6\}$.

Example 5. *We have the equalities*

$$c_{\text{Gr}(3,6)}(J, J, J) = 2, \quad c_{\text{Gr}(6,9)}(I, I, I) = 3, \quad \text{and} \quad c_{\text{Fl}(3,6,9)}(\omega, \omega, \omega) = 6.$$


Remark 8. *Since $6 = 3 * 2$, this example is compatible with [DW11, Theorem 7.14], [Ric12, Theorem 1.1], and [Res11a, Theorem A]. However, it shows that [KP11, Theorem 3] cannot be correct, since the right hand side of [KP11, Theorem 3] is equal to $2 * 2 * 2 = 8$.*

Actually, the map in [KP11, Theorem 2] is not surjective. The "inverse map" they construct should certainly be a left-inverse, but it is never verified that it is a right-inverse.


In fact, let us denote by R the puzzle displayed on the left in Appendix A.2 (with a Rhombus over the edge drawn with a thick line) and by T the puzzle on the right (with a Triangle). Performing the map (D_{23}, D_{13}, D_{12}) of [KP11, Theorem 2] on the six puzzles of Appendix A.4, we get the six triples (R, R, R) , (T, R, R) , (T, T, R) , (T, T, T) , (R, R, T) , (R, T, T) . So the map in [KP11, Theorem 2] is injective, but its image misses the two triples (R, T, R) and (T, R, T) .

As a conclusion, Theorems 2 and 3 in [KP11] are not correct, whereas we have no objection for the other statements in that paper.







A.2. The puzzles for $c_{\text{Gr}(3,6)}(J, J, J) = 2$ with $J = \{2, 4, 6\}$.



A.3. The puzzles for $c_{\text{Gr}(6,9)}(I, I, I) = 3$ with $I = \{2, 3, 5, 6, 8, 9\}$.



A.4. The puzzles for $c_{\mathcal{F}(3,6;9)}(\omega, \omega, \omega) = 6$ with $\omega = (\{3, 6, 9\} \subset \{2, 3, 5, 6, 8, 9\})$.



UNIVERSITÉ DE LORRAINE, CNRS, INSTITUT ÉLIE CARTAN DE LORRAINE, UMR 7502, VANDOEUVRE-LÈS-NANCY, F-54506, FRANCE

Email address: pierre-emmanuel.chaput@univ-lorraine.fr

UNIVERSITÉ CLAUDE BERNARD LYON I, INSTITUT CAMILLE JORDAN (ICJ), UMR CNRS 5208, 43 BOULEVARD DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX

Email address: ressayre@math.univ-lyon1.fr

REFERENCES

- [BB05] Anders Björner and Francesco Brenti, *Combinatorics of Coxeter groups*, Graduate Texts in Mathematics, vol. 231, Springer, New York, 2005.
- [BK05] Michel Brion and Shrawan Kumar, *Frobenius splitting methods in geometry and representation theory*, Progress in Mathematics, vol. 231, Birkhäuser Boston, Inc., Boston, MA, 2005.
- [BK06] Prakash Belkale and Shrawan Kumar, *Eigenvalue problem and a new product in cohomology of flag varieties*, Invent. Math. **166** (2006), no. 1, 185–228.
- [Bou02] Nicolas Bourbaki, *Elements of Mathematics* (Berlin), Springer-Verlag, Berlin, 2002, Translated from the 1968 French original by Andrew Pressley.
- [BPVdV84] W. Barth, C. Peters, and A. Van de Ven, *Compact complex surfaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 4, Springer-Verlag, Berlin, 1984.
- [Bri98] M. Brion, *The behaviour at infinity of the Bruhat decomposition*, Comment. Math. Helv. **73** (1998), no. 1, 137–174.
- [CR21] P.-E. Chaput and N. Ressayre, *Sage program for computing branching divisors*, <http://math.univ-lyon1.fr/~ressayre/Info/branchDivClassGrass.tar>, 2021, Based on Anders Buch qcalc package.
- [Dem74] Michel Demazure, *Désingularisation des variétés de Schubert généralisées*, Ann. Sci. École Norm. Sup. (4) **7** (1974), 53–88.
- [DW11] Harm Derksen and Jerzy Weyman, *The combinatorics of quiver representations*, Ann. Inst. Fourier (Grenoble) **61** (2011), no. 3, 1061–1131.
- [Eis95] David Eisenbud, *Commutative algebra*, Graduate Texts in Mathematics, vol. 150, Springer-Verlag, New York, 1995, With a view toward algebraic geometry.
- [Éla92] Alexander G. Élashvili, *Invariant algebras*, Lie groups, their discrete subgroups, and invariant theory (Providence, RI), Adv. Soviet Math., vol. 8, Amer. Math. Soc., Providence, RI, 1992, pp. 57–64.
- [Ful84] William Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 2, Springer-Verlag, Berlin, 1984.
- [Har77] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York-Heidelberg, 1977, Graduate Texts in Mathematics, No. 52.
- [KM98] János Kollár and Shigefumi Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998, With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [Kos58] Bertram Kostant, *A formula for the multiplicity of a weight*, Proc. Nat. Acad. Sci. U.S.A. **44** (1958), 588–589.
- [KP11] Allen Knutson and Kevin Purbhoo, *Product and puzzle formulae for GL_n Belkale-Kumar coefficients*, Electron. J. Combin. **18** (2011), no. 1, Paper 76, 20.
- [KT99] Allen Knutson and Terence Tao, *The honeycomb model of $GL_n(\mathbb{C})$ tensor products. I. Proof of the saturation conjecture*, J. Amer. Math. Soc. **12** (1999), no. 4, 1055–1090.
- [KT03] Allen Knutson and Terence Tao, *Puzzles and (equivariant) cohomology of Grassmannians*, Duke Math. J. **119** (2003), no. 2, 221–260.
- [KTT09] Ronald C. King, Christophe Tollu, and Frédéric Toumazet, *Factorisation of Littlewood-Richardson coefficients*, J. Combin. Theory Ser. A **116** (2009), no. 2, 314–333.
- [Kum02] Shrawan Kumar, *Kac-Moody groups, their flag varieties and representation theory*, Progress in Mathematics, vol. 204, Birkhäuser Boston, Inc., Boston, MA, 2002.
- [Laz80] Robert Lazarsfeld, *A Barth-type theorem for branched coverings of projective space*, Math. Ann. **249** (1980), no. 2, 153–162.
- [Les47] Léonce Lesieur, *Les problèmes d’intersection sur une variété de Grassmann*, C. R. Acad. Sci. Paris **225** (1947), 916–917.
- [Mat88] Olivier Mathieu, *Formules de caractères pour les algèbres de Kac-Moody générales*, Astérisque (1988), no. 159-160, 267.
- [Mir85] Rick Miranda, *Triple covers in algebraic geometry*, Amer. J. Math. **107** (1985), no. 5, 1123–1158.
- [Par91] Rita Pardini, *Abelian covers of algebraic varieties*, J. Reine Angew. Math. **417** (1991), 191–213.
- [Per07] Nicolas Perrin, *Small resolutions of minuscule Schubert varieties*, Compos. Math. **143** (2007), no. 5, 1255–1312.
- [PN20] Maxime Pelletier and Ressayre Nicolas, *Some unexpected properties of littlewood-richardson coefficients*, 2020.
- [Pra05] Piotr Pragacz, *Multiplying Schubert classes*, Trends Math., Birkhauser, Basel, 2005, pp. 163–174.
- [Ram87] A. Ramanathan, *Equations defining Schubert varieties and Frobenius splitting of diagonals*, Inst. Hautes Études Sci. Publ. Math. (1987), no. 65, 61–90.

- [Ras04] Etienne Rassart, *A polynomiality property for Littlewood-Richardson coefficients*, J. Combin. Theory Ser. A **107** (2004), no. 2, 161–179.
- [Res10] Nicolas Ressayre, *Geometric invariant theory and generalized eigenvalue problem*, Invent. Math. **180** (2010), 389–441.
- [Res11a] N. Ressayre, *Multiplicative formulas in Schubert calculus and quiver representation*, Indag. Math. (N.S.) **22** (2011), no. 1-2, 87–102.
- [Res11b] Nicolas Ressayre, *Reductions for branching coefficients*, J. of Lie Theory (to appear) (2011), arXiv:1102.0196.
- [Ric12] Edward Richmond, *A multiplicative formula for structure constants in the cohomology of flag varieties*, Michigan Math. J. **61** (2012), no. 1, 3–17.
- [Rot11] Mike Roth, *Reduction rules for Littlewood-Richardson coefficients*, Int. Math. Res. Not. IMRN (2011), no. 18, 4105–4134.
- [RR11] Nicolas Ressayre and Edward Richmond, *Branching Schubert calculus and the Belkale-Kumar product on cohomology*, Proc. Amer. Math. Soc. **139** (2011), no. 3, 835–848.
- [Ste61] Robert Steinberg, *A general Clebsch-Gordan theorem*, Bull. Amer. Math. Soc. **67** (1961), 406–407.