

The quantum switch is uniquely defined by its action on unitary operations

Qingxiuxiong Dong¹, Marco Túlio Quintino^{2,3,1}, Akihito Soeda¹, and Mio Murao^{1,4}

¹Department of Physics, Graduate School of Science, The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

²Institute for Quantum Optics and Quantum Information (IQOQI), Austrian Academy of Sciences, Boltzmanngasse 3, A-1090 Vienna, Austria

³Vienna Center for Quantum Science and Technology (VCQ), Faculty of Physics, University of Vienna, Boltzmanngasse 5, A-1090 Vienna, Austria

⁴Trans-scale Quantum Science Institute, The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

The quantum switch is a physical process that creates a coherent control between different unitary operations which is often described as a process which transforms a pair of unitary operations (U_1, U_2) into a controlled unitary operation that coherently applies them in different orders as $|0\rangle\langle 0| \otimes U_1 U_2 + |1\rangle\langle 1| \otimes U_2 U_1$. This description, however, does not directly define its action on non-unitary operations. The action of quantum switch on non-unitary operations is then chosen to be a “natural” extension of its action on unitary operation. Since, in general, the action of a process on non-unitary operations is not uniquely determined by its action on only unitary operations, in principle, there could be a set of inequivalent extensions of quantum switch for non-unitary operations. In this paper, we prove that there is a unique way to extend the actions of quantum switch to non-unitary operations. In other words, contrary to the general case, the action of quantum switch on non-unitary operations is completely determined by its action on unitary operations. We also discuss the general problem of when the complete description of a quantum process is uniquely determined by its action on unitary operations and identify a set of single-slot processes which are completely defined by their action on unitary operations.

1 Introduction

The quantum switch [1] is a higher-order quantum operation [2–5] that transforms a pair of input unitary operations into a “coherent superposition of different orders” of the input operations. It is an example of a higher-order quantum operation that cannot

be implemented within the standard quantum circuit model, but there is no fundamental principle that forbids its implementation in quantum mechanics. For that, quantum switch is a well-studied higher-order quantum operation in the context of indefinite causal order [1, 6, 7], for revealing how the causal structure may affect the quantum information processing in quantum mechanics. It is also proposed as a resource for discriminating non-signaling channels [8], quantum computation [9], and quantum communication [10–15]. Also, although we only consider quantum switch that acts on a pair of input operations, the concept of quantum switch can be generalized to multiple input operations as studied in Ref. [9].

The action of quantum switch is to produce a coherent control of causal orders. In Ref. [1], the action of quantum switch is originally defined on unitary operations, and then generalized to quantum operations beyond unitary operations. That is, given two unitary operations \tilde{U}_1 and \tilde{U}_2 characterized by the unitary operators U_1 and U_2 as input operations¹, quantum switch transforms them into the coherent control of different orders, namely, the unitary operation given by

$$(U_1, U_2) \mapsto |0\rangle\langle 0|_C \otimes U_2 U_1 + |1\rangle\langle 1|_C \otimes U_1 U_2, \quad (1)$$

by using each of U_1 and U_2 only once. If the control qubit is in $|0\rangle_C$, U_1 is applied on the target system first, and followed by U_2 , and if the control qubit is in $|1\rangle_C$, U_2 is applied first followed by U_1 . If the control qubit is in a superposition of $|0\rangle$ and $|1\rangle$, then the two different orders should be “superposed” in a coherent way.

The action of quantum switch for unitary operations is defined in a natural way, but its action for general quantum operations is not trivial from the definition for unitary operations. In the definition of quan-

¹In this work, we denote quantum operations and maps with a tilde, and supermaps with a double tilde. In this way a unitary operation \tilde{U} described by the operator U is defined via $\tilde{U}(\rho) = U\rho U^\dagger$.

Qingxiuxiong Dong: qingxiuxiong.dong@gmail.com

Marco Túlio Quintino: mtcq.mm@gmail.com

Akihito Soeda: soeda@phys.s.u-tokyo.ac.jp

Mio Murao: murao@phys.s.u-tokyo.ac.jp

tum switch [1], the action on general quantum operations is chosen to be the most “natural” one based on its Kraus representation [16] as follows: given two quantum operations represented by the Kraus operators $\{K_i\}_i$ and $\{L_j\}_j$, respectively, the output operation of quantum switch is given by the Kraus operators $\{W_{ij}\}_{ij}$ defined by

$$W_{ij} := |0\rangle\langle 0|_C \otimes L_j K_i + |1\rangle\langle 1|_C \otimes K_i L_j. \quad (2)$$

Following this definition, different results have been established. One interesting result is that quantum switch can help in enhancing certain communication tasks [11–13]. In particular, if two input operations are the depolarizing channels, which do not transfer any information, the output operation is not depolarizing channel anymore, and can be used for transferring information. The action of quantum switch on the depolarizing channels looks different from that on unitary operations, that is, coherently controlled two differently causally ordered operations, since the composition of two depolarizing channels is a depolarizing channel. The definition of the quantum switch for general operations presented in Eq.(2) is widely used in various studies, but before this paper, it was not clear if the definition in Eq.(2) follows uniquely by the behavior of the quantum switch on unitary operators described in Eq.(1).

Before proceeding we present a simple example to show that, in general, the behavior of a quantum process for unitary operations does not uniquely define its action on general operations. Our example is illustrated in terms of quantum circuits shown in Fig. 1 and consists in presenting two processes which despite being different, have identical action on unitary operations. This example follows from the fact that since unitary operations are unital, that is, $\tilde{U}(I) = I$, if $\tilde{D}(\rho) := I/d \text{Tr } \rho$ is the completely depolarizing channel and $id(\rho) = \rho$ is the identity channel, we have that

$$\tilde{D} \circ \tilde{U} \circ \tilde{D} = id \circ \tilde{U} \circ \tilde{D} = \tilde{D}. \quad (3)$$

But, when considering non-unital channels $\tilde{\Lambda}$, that is $\tilde{\Lambda}(I) \neq I$, the identity in Eq. (3) does not hold. In particular, if we take the quantum operation $\tilde{\Lambda}(\rho) = |0\rangle\langle 0| \text{Tr } \rho$ which replaces the input state ρ by the quantum state $|0\rangle\langle 0|$, we obtain

$$\tilde{D} \circ \tilde{\Lambda} \circ \tilde{D} = \tilde{D} \quad (4)$$

$$id \circ \tilde{\Lambda} \circ \tilde{D} = \tilde{\Lambda}. \quad (5)$$

Although simple, this example can be used to show that *almost all* higher-order quantum operations share this property, namely, the action on unitary operations does not uniquely define the action on non-unitary operations. In particular, for any fixed circuit operation, we can construct two new circuits by

\forall all unitary operations \tilde{U} :

$$\boxed{\tilde{D}} \text{---} \boxed{\tilde{U}} \text{---} \boxed{\tilde{D}} = \boxed{\tilde{D}} \text{---} \boxed{\tilde{U}} \text{---} \boxed{id}$$

but \exists non-unitary operations such as $\tilde{\Lambda}$:

$$\boxed{\tilde{D}} \text{---} \boxed{\tilde{\Lambda}} \text{---} \boxed{\tilde{D}} \neq \boxed{\tilde{D}} \text{---} \boxed{\tilde{\Lambda}} \text{---} \boxed{id}$$

Figure 1: Two quantum circuits implementing higher-order quantum operations that transform an arbitrary unitary operation into the depolarizing channel. Here \tilde{U} denotes the input unitary operation, \tilde{D} denotes the depolarizing channel, and id denotes the identity channel. While the actions of both of them are equivalent on unitary operations, their actions on some quantum operations, e.g., the quantum operation $\tilde{\Lambda}(\rho) = |0\rangle\langle 0| \text{Tr } \rho$ which replaces the input state ρ by the quantum state $|0\rangle\langle 0|$, are different.

considering probabilistic mixtures of the fixed operations with the identity channel or the depolarizing channel. For any non-trivial probability mixture, the same argument used in our previous example shows that these two new circuits may act identically on unitary operations, but they act differently for non-unital operations. Moreover, this argument does not rely on quantum circuits and can be used for any general process (potentially with indefinite causal order [1]) which transforms quantum operations. We just need to recognize that the two different circuits presented in Fig. 1 are valid processes, and by performing any non-trivial convex combinations, one can always construct examples of two higher-order transformations which the action on unitary operations does not uniquely define the action on general.

Our main question then is, does there exist a higher-order quantum operation which respects the constraints of the quantum switch on unitary operations presented in Eq. (1), but does not respect the constraints of the quantum switch for general operations presented in Eq. (2)?

It is worth mentioning that the difficulty of the problem follows the fact that general quantum operations cannot be written as a linear combination of unitary operations. For instance, a non-unital channel can never be written as a linear combination of unitary operations. However, linearity is not the only condition in which higher-order operations should respect. It is crucial to respect a notion of complete positivity. That is, since quantum operations should be completely positive (CP), it is inevitable to impose that processes transforming quantum operations should be completely CP preserving [2–4, 6, 7, 11].

When dealing with the “lower-order case” where one considers quantum operations which transform quantum states into quantum states, linearity itself is enough to ensure that the action of quantum operations on pure quantum states uniquely defines the action of this operation on any quantum state. This holds because the linear space spanned by pure quan-

tum states forms the whole set of self-adjoint operators. An analogous problem then would be to define the action of a quantum operation for a restricted set of pure states and then analyze how the CP condition restricts the extension of this operation to the set of all quantum states in a non-trivial way, problem which is discussed in Refs. [17–20]. We also remark Ref. [21] which investigates when the marginals of a linear map uniquely determine its whole action and Ref. [22] which analyses when compatible tuples of measurements admit a unique joint measurement.

In this paper, we analyze how the action of higher-order processes on unitary quantum operations restricts its action on general quantum operations. In particular, we show that the constraints of linearity and completely CP preserving, required by such quantum processes, ensure that the action of quantum switch on general quantum operations uniquely defines their action on unitary operations. We then consider the general problem of understanding the restrictions imposed by the completely CP preserving property and apply our proof techniques to problems which are not directly related to the quantum switch. In particular, we identify a class of single slot processes which, similarly to the switch, their action on unitary operations uniquely defines their action on general operations. This class of single input-operation processes includes identity supermaps, unitary transposition, and qubit unitary complex conjugation. This result strengthens the theoretical background for the studies on quantum switch, and provides a physical meaning to the experimental realization of quantum switch because the assumptions we pose are necessary to promise the action of quantum switch on general quantum operations.

This paper is organized as follows. In Sec. 2, we provide the statement of our main theorem. We also introduce the Choi representation, and state the main theorem in terms of the Choi operators. In Sec. 3, we show the consequences that can be obtained only from the linearity, which are also used in later proofs. In Sec. 4, we provide a similar result on the supermaps with a single input operation instead of quantum switch. The conclusion is given in Sec. 5. In Appendix A, we provide the proofs for the results from the linearity stated in Sec. 3. In Appendix B, we provide the proof of the result on the supermaps with a single input operation stated in Sec. 4, and in Appendix C, we provide the proof of the result on quantum switch stated in Sec. 2. We provide the proof for quantum switch in the latter Appendix because while the technique for both proofs is similar, the former one is much simpler. In Appendix D and E, we provide the detailed calculations that are used in Appendix C.




Figure 2: Conceptual figure of a two-slot supermap which may not respect a definite causal order such as the quantum switch. Two-slot supermaps transform a pair of quantum operations as input and output a quantum operation from \mathcal{P} to \mathcal{F} . In general, the action of a supermap on unitary operations does not uniquely define its action on non-unitary operations.

2 The Uniqueness of Quantum Switch

In this paper, we derive the suitable action of quantum switch on arbitrary quantum operations from its action on unitary operations. In particular, we derive the action of quantum switch from the following three conditions:

1. The action of quantum switch on unitary operations is given by Eq. (1).
2. Quantum switch is a linear supermap.
3. Quantum switch is a completely CP preserving supermap.

The precise conditions will be presented in the following of this section. The second condition is equivalent to that quantum switch uses each input operation once, and the third condition is a necessary condition for a supermap to be (probabilistically) implementable [3, 4]. Note that since we do not assume any other linear constraint known as the causal condition for quantum switch, these conditions do not exclude the possibility of a probabilistic action of quantum switch on certain input operations.

Theorem 1. *If the action of a 2-slot supermap on unitary operations is given by Eq. (1) for arbitrary input unitary operation U_1 and U_2 , and is implementable by using each input operation once, then its action on arbitrary quantum operations is uniquely determined by the Kraus operators Eq. (2).*

In order to prove Theorem 1, we use the Choi representation of supermaps [2, 3, 23, 24]. That is, since the Choi-Jamiołkowski isomorphism is a bijection, we prove the uniqueness of the extension of quantum switch by proving that the corresponding Choi operator W is uniquely determined.

Here we first introduce the Choi representation of linear maps [23, 24], which will be useful to represent quantum operations. Then we discuss how to represent supermaps that describe higher-order quantum

operations as linear operators via the Choi representation [2, 4]. For a linear map $\tilde{\Lambda} : \mathcal{L}(\mathcal{I}) \rightarrow \mathcal{L}(\mathcal{O})$, the corresponding Choi operator is given by

$$J_{\mathcal{A}} := \sum_{ij} |i\rangle\langle j| \otimes \tilde{\Lambda}(|i\rangle\langle j|) \in \mathcal{L}(\mathcal{I} \otimes \mathcal{O}), \quad (6)$$

where $\{|i\rangle\}$ is an orthonormal basis for \mathcal{I} . The condition that a map $\tilde{\Lambda}$ is completely positive (CP) corresponds to the positivity of $J_{\mathcal{A}}$ as $J_{\mathcal{A}} \geq 0$. In this paper, since unitary operations play an important role, we also denote a unitary operation \tilde{U} with the corresponding unitary operator U , and its Choi operator as $J_U := \sum_{ij} |i\rangle\langle j| \otimes \tilde{U}(|i\rangle\langle j|)$.

A linear supermap, also simply referred as a supermap, is linear transformations on maps and are useful to represent higher-order transformations. The quantum switch has a particular structure of a two-slot process hence its supermap may have the form of

$$\tilde{\mathcal{W}} : [\mathcal{L}(\mathcal{I}_1) \rightarrow \mathcal{L}(\mathcal{O}_1) \otimes \mathcal{L}(\mathcal{I}_2) \rightarrow \mathcal{L}(\mathcal{O}_2)] \rightarrow \mathcal{L}(\mathcal{P}) \rightarrow \mathcal{L}(\mathcal{F}), \quad (7)$$

where

$$\begin{aligned} & \tilde{\mathcal{W}}(\alpha \tilde{\mathcal{A}}_1 \otimes \tilde{\mathcal{A}}_2 + \beta \tilde{\mathcal{B}}_1 \otimes \tilde{\mathcal{B}}_2) \\ &= \alpha \tilde{\mathcal{W}}(\tilde{\mathcal{A}}_1 \otimes \tilde{\mathcal{A}}_2) + \beta \tilde{\mathcal{W}}(\tilde{\mathcal{B}}_1 \otimes \tilde{\mathcal{B}}_2) \end{aligned} \quad (8)$$

for arbitrary input maps $\tilde{\mathcal{A}}_1, \tilde{\mathcal{A}}_2, \tilde{\mathcal{B}}_1, \tilde{\mathcal{B}}_2$ and arbitrary $\alpha, \beta \in \mathbb{C}$. In Eq. (7), $\mathcal{I}_k, \mathcal{O}_k \cong \mathbb{C}^d$ for $k = 1, 2$ are the Hilbert spaces of the two input operations, $\mathcal{P}, \mathcal{F} \cong \mathbb{C}^{2d}$ denote the Hilbert spaces of the output operation as Fig. 2 shows. Since linear maps can be represented by operators with the Choi isomorphism, supermaps may be viewed as maps action on Choi operators. Hence, by taking the Choi representation of this induced map, we can also represent a supermap as in Eq. (7) by an operator $W \in \mathcal{L}(\mathcal{P} \otimes \mathcal{I}_1 \otimes \mathcal{O}_1 \otimes \mathcal{I}_2 \otimes \mathcal{I}_2 \otimes \mathcal{F})$, also if $A \in \mathcal{L}(\mathcal{I}_1 \otimes \mathcal{O}_1)$ and $B \in \mathcal{L}(\mathcal{I}_2 \otimes \mathcal{O}_2)$ are the Choi operators of quantum operations, the Choi operator of the output operation is given by [1, 2, 4]

$$\text{Tr}_{\mathcal{I}_1 \mathcal{O}_1 \mathcal{I}_2 \mathcal{O}_2} [W (I_{\mathcal{P}} \otimes A_{\mathcal{I}_1 \mathcal{O}_1} \otimes B_{\mathcal{I}_2 \mathcal{O}_2} \otimes I_{\mathcal{F}})^t], \quad (9)$$

where t stands for transposition in the computational basis.

A supermap $\tilde{\mathcal{W}} : [\mathcal{L}(\mathcal{I}) \rightarrow \mathcal{L}(\mathcal{O})] \rightarrow \mathcal{L}(\mathcal{P}) \rightarrow \mathcal{L}(\mathcal{F})$, is CP preserving if it transforms CP maps into CP maps. Also, a supermap is completely CP preserving if all of its trivial extensions transforms CP maps into CP maps, that is, for every CP linear map which may make use of an auxiliary space $\tilde{\mathcal{C}} : \mathcal{L}(\mathcal{I} \otimes \mathcal{H}_{\text{aux}}) \rightarrow \mathcal{L}(\mathcal{O} \otimes \mathcal{H}_{\text{aux}})$, the output-map $\tilde{\mathcal{W}} \otimes \tilde{id}(\tilde{\mathcal{C}})$ is also CP, where $\tilde{id} : \mathcal{L}(\mathcal{H}_{\text{aux}}) \rightarrow \mathcal{L}(\mathcal{H}_{\text{aux}})$ is the identity supermap on the auxiliary space defined by $\tilde{id}(\tilde{\mathcal{A}}) = \tilde{\mathcal{A}}$ for every map $\tilde{\mathcal{A}}$. In Choi operator representation completely CP preserving condition of $\tilde{\mathcal{W}}$ is simply

given by the positivity constraint $W \geq 0$ [2, 4], similar to the CP condition for quantum operation case in Choi representation. If, in addition we impose that a supermap corresponds to a superchannel, that is, a deterministic quantum supermap, extra linear constraints should be imposed. For the case of general two-slot superchannels which may not have a definite causal order these constraints constitute in imposing that the supermap transforms pair of independent quantum channels into a quantum channel [1, 6, 7]. We note however that such linear constraints are not used in the proof of the uniqueness of the extension of quantum switch. For an introduction on deterministic quantum transformations we suggest the introductory sections of Ref. [25] and [26] and references therein.

We denote the Choi operator of quantum switch defined by the Kraus operators given in Ref. [1] as $W_0 \in \mathcal{L}(\mathcal{P} \otimes \mathcal{I}_1 \otimes \mathcal{O}_1 \otimes \mathcal{I}_2 \otimes \mathcal{I}_2 \otimes \mathcal{F})$. In particular, this operator can be calculated as $W_0 = |W_0\rangle\langle W_0|$ with

$$\begin{aligned} |W_0\rangle &= |00\rangle_{\mathcal{P}_C \mathcal{F}_C} |I\rangle_{\mathcal{P}_T \mathcal{I}_1} |I\rangle_{\mathcal{O}_1 \mathcal{I}_2} |I\rangle_{\mathcal{O}_2 \mathcal{F}_T} \\ &+ |11\rangle_{\mathcal{P}_C \mathcal{F}_C} |I\rangle_{\mathcal{P}_T \mathcal{I}_2} |I\rangle_{\mathcal{O}_2 \mathcal{I}_1} |I\rangle_{\mathcal{O}_1 \mathcal{F}_T} \\ &= \sum_{i,j,k=0}^{d-1} (|ijjkik00\rangle + |jkikik11\rangle), \end{aligned} \quad (10)$$

where $|I\rangle := \sum_i |ii\rangle$ and the Hilbert spaces for the vector in the last line is in the order of $\mathcal{I}_1 \otimes \mathcal{O}_1 \otimes \mathcal{I}_2 \otimes \mathcal{O}_2 \otimes \mathcal{P}_T \otimes \mathcal{F}_T \otimes \mathcal{P}_C \otimes \mathcal{F}_C$. Note that the normalization of the Choi operators is given by $\text{Tr} W = \text{Tr} W_0 = 2d^3$, and each elements of the matrix W_0 is 0 or 1.

In this paper, we show that the only possible W satisfies $W = W_0$ under the following assumptions: when input operations are unitary operations denoted by $J = J_{U_1} \otimes J_{U_2}$, the action of quantum switch is given by W_0 as² $\text{Tr}_{in} W_0 J^t$, where $\mathcal{H}_{in} := \mathcal{I}_1 \otimes \mathcal{O}_1 \otimes \mathcal{I}_2 \otimes \mathcal{I}_2$ stands for the total input-channel space of the switch; the linearity of the supermap, which is automatically satisfied as the linearity is required when the Choi operator is defined; and the positivity $W \geq 0$. In terms of Choi representation, Theorem 1 is equivalent to the following Lemma.

Lemma 1. *Let $W_0 = |W_0\rangle\langle W_0|$ where $|W_0\rangle$ is defined by Eq. (10). If there exists $W \geq 0$ such that $\text{Tr}_{in} [W (J_{U_1} \otimes J_{U_2})^t] = \text{Tr}_{in} [W_0 (J_{U_1} \otimes J_{U_2})^t]$ holds for all U_1 and U_2 , then $W = W_0$ holds.*

The main proof of Lemma 1 is given in Appendix C. Here we provide a sketch of the proof. We first consider the linear span of the Choi operator of unitary operations, as the action of a supermap on this linear span is uniquely determined due to the linearity. In

²When we write $\text{Tr}_{in} W_0 J^t$, we are suppressing the identity operators, that is, in the full notation we have $\text{Tr}_{in} [W_0 (I_{\mathcal{P}} \otimes J^t \otimes I_{\mathcal{F}})]$. To avoid a heavy notation, whenever clear from context, we will suppress the identity operator in these trace relations.

Sec. 3 and Appendix A, we provide a set of operators that are in the linear span. In Appendix C, we consider the consequences from the positivity of the Choi operator $W \geq 0$, which also relies on the results on the linear span. Remark that the Choi operator of quantum switch given by Eq. (10) has a simple form, namely, it is a matrix with a submatrix filled with only 1 and the remaining being 0. The proof can be separated into two parts. In the first part, we determine all of the diagonal elements which are 1, where the main calculations are separated into Appendix D. After determining all of the diagonal elements, the remaining is to show that the submatrix with the diagonal elements being 1 is filled with 1. In the second part of the proof, we evaluate the element-wise 1-norm, i.e., the sum of absolute values of all elements in the computational basis, and show that it is equal to the number of elements of the submatrix, where the main calculations are separated into Appendix E.

3 The Linear Span of the Choi Operators of Unitary Operations

While the positivity (of the Choi operator) is a key feature for seeking the unique extension of quantum switch, we first focus on the linearity in this section. From the linearity, the action on any input operations that is a linear combination of unitary operations is given in the same way, that is, for any $J \in \text{span}\{J_{U_1} \otimes J_{U_2}\}$ where

$$\begin{aligned} & \text{span}\{J_{U_1} \otimes J_{U_2}\} \\ & := \{O \mid O = \sum_{i_1, i_2} c_{i_1 i_2} (J_{U_{i_1}} \otimes J_{U_{i_2}}), c_{i_1 i_2} \in \mathbb{C}\}, \end{aligned} \quad (11)$$

the action of quantum switch on J is given by $\text{Tr}_{in} W_0 J^t$. For that, we consider the operators in $\text{span}\{J_{U_1} \otimes J_{U_2}\}$. Since $\text{span}\{J_{U_1} \otimes J_{U_2}\} = \text{span}\{J_U\} \otimes \text{span}\{J_U\}$ holds, it is enough to consider the operators in $\text{span}\{J_U\}$.

We first summarize the results in this section: the following operators are in $\text{span}\{J_U\}$.

1. $|ii\rangle\langle jj|, |ij\rangle\langle ji| \in \text{span}\{J_U\}$ for $i \neq j$.
2. $|ij\rangle\langle i'j'| \in \text{span}\{J_U\}$ when i, j, i', j' are pairwise different.
3. $|ij\rangle\langle i'j'| \in \text{span}\{J_U\}$ if only two of i, j, i', j' are the same and others are different.
4. $\sum_{j=i+k} |ij\rangle\langle ij| \in \text{span}\{J_U\}$ for $k = 0, \dots, d-1$.
5. $(|ij\rangle\langle ii| - |jj\rangle\langle ji|), (|ji\rangle\langle ii| - |jj\rangle\langle ij|) \in \text{span}\{J_U\}$ for $i \neq j$.

The proof is given in Lemmas 3-7.

We remark that the set of the operators listed here does not span the whole space of $\text{span}\{J_U\}$ unless

$d = 2$. There are only $2d(d-1) + d(d-1)(d-2)(d-3) + 6d(d-1)(d-2) + d + 2d(d-1) = d(d^3 - 3d + 3)$ elements here, but the dimension of $\text{span}\{J_U\}$ is $(d^2 - 1)^2 + 1$ (Lemma 3 of Ref. [27]), which is strictly greater than $d(d^3 - 3d + 3)$ for $d > 2$. As we show in Appendix C that Lemma 1 can be proved by only considering the action of quantum switch on these operators, requiring the action of quantum switch on all unitary operations is not necessary in general to uniquely determine the action of quantum switch on general quantum operations. Instead, requiring the action of quantum switch on a restricted set of unitary operations is enough for uniquely determining the action on the rest of unitary operations and general quantum operations.

Another important fact is that the action of quantum switch on the depolarizing channels is determined only by the linearity, because

$$\frac{I}{d} \otimes \frac{I}{d} \in \text{span}\{J_{U_1} \otimes J_{U_2}\} \quad (12)$$

holds, where $\frac{I}{d}$ is the Choi operator of the depolarizing channel. This is also obvious because the depolarizing channel is a probabilistic mixture of uniformly random unitary operations. Thus, the results of Refs. [11–13] can be obtained with only the assumption of the linearity. In contrast, if we apply quantum switch to non-unital quantum operations, the action may not be determined from only the linearity.

4 Uniqueness of single input supermaps

In this section, we provide a similar theorem on the supermaps with a single input operation instead of quantum switch.

Consider the operations $\tilde{\mathcal{C}}$ which acts on as the identity supermap on unitary operations, that is, for every $\tilde{\mathcal{U}}$ we have $\tilde{\mathcal{C}}(\tilde{\mathcal{U}}) = \tilde{\mathcal{U}}$. We show that by imposing that $\tilde{\mathcal{C}}$ is completely CP preserving, $\tilde{\mathcal{C}}$ also behaves as the identity supermap for any arbitrary input operation $\tilde{\Lambda}$, that is, $\tilde{\mathcal{C}}(\tilde{\Lambda}) = \tilde{\Lambda}$. In other words, the Choi operator of $\tilde{\mathcal{C}}$ is uniquely determined by

$$C_0 = \sum_{i,j,k,l=1}^d |ij\rangle\langle kl| \otimes |ij\rangle\langle kl| \in \mathcal{L}(\mathcal{I} \otimes \mathcal{O} \otimes \mathcal{P} \otimes \mathcal{F}). \quad (13)$$

Theorem 2. *If the action of a 1-slot supermap on unitary operations is given by $\tilde{\mathcal{U}} \mapsto \tilde{\mathcal{U}}$ for an arbitrary input unitary operation $\tilde{\mathcal{U}}$, and is implementable by using the input operation once, then its action on arbitrary quantum operations is uniquely given by $\tilde{\Lambda} \mapsto \tilde{\Lambda}$ for an arbitrary input quantum operation $\tilde{\Lambda}$.*

Similarly, Theorem 2 is equivalent to the following Lemma.

Lemma 2. Let C_0 be the Choi operator of the identity supermap defined by Eq. (13). If there exists $C \geq 0$ such that $\text{Tr}_{\mathcal{IO}}(C J_U^t) = \text{Tr}_{\mathcal{IO}}(C_0 J_U^t)$ holds for all U , then $C = C_0$ holds.

The proof of Lemma 2 is given in Appendix B.

While Theorem 2 only holds supermaps which act as the identity on unitary operations, this result can be easily extended to cases to supermaps which act on unitary operations as $\tilde{U} \mapsto \tilde{B} \circ \tilde{U} \circ \tilde{A}$, where \tilde{A} and \tilde{B} are fixed unitary operations. This class of supermaps is precisely the set of reversibility preserving one-slot superchannels (also known as one-partite pure processes [7]) which do not modify the dimension of the input-operation [25].

Corollary 1. If the action of a 1-slot supermap on unitary operations is given by $\tilde{U} \mapsto \tilde{B} \circ \tilde{U} \circ \tilde{A}$ for an arbitrary input unitary operation \tilde{U} and two fixed unitary operations \tilde{A} and \tilde{B} , and is implementable by using the input operation once, then its action on arbitrary quantum operations is uniquely given by $\tilde{\Lambda} \mapsto \tilde{B} \circ \tilde{\Lambda} \circ \tilde{A}$ for an arbitrary input quantum operation $\tilde{\Lambda}$.

Proof. Corollary 1 can be reduced to Theorem 2 as follows. Consider the condition for unitary operations

$$\text{Tr}_{\mathcal{IO}}(C' J_U^t) = \text{Tr}_{\mathcal{IO}}(C_0 J_U^t). \quad (14)$$

By introducing $C = (A_I \otimes B_F)^\dagger C' (A_I \otimes B_F)$, the l.h.s. can be written as

$$\begin{aligned} \text{Tr}_{\mathcal{IO}}(C' J_U^t) &= \text{Tr}_{\mathcal{IO}}[(A_I \otimes B_F) C (A_I \otimes B_F)^\dagger J_U^t] \\ &= B_F \{ \text{Tr}_{\mathcal{IO}}[C (A_I^\dagger J_U^t A_I)] \} B_F^\dagger \\ &= B_F \{ \text{Tr}_{\mathcal{IO}}[C (J_{UA^*})^t] \} B_F^\dagger, \end{aligned} \quad (15)$$

whereas the r.h.s. can be written as

$$\text{Tr}_{\mathcal{IO}}(C_0 J_U^t) = B_F \{ \text{Tr}_{\mathcal{IO}}[C_0 (J_{UA^*})^t] \} B_F^\dagger. \quad (16)$$

Since UA^* is also a unitary operator and spans the whole linear span of $\text{span}\{J_U\}$, the uniqueness of C' is equivalent to the uniqueness of the identity supermap C . \square

Corollary 1 can be used to show that the process leading to qubit unitary complex conjugation analyzed in [28] is uniquely defined by its action on unitary operations. This holds true because $\tilde{U}^* = \tilde{Y} \circ \tilde{U} \circ \tilde{Y}$, where \tilde{Y} denotes the Pauli Y unitary operation.

Another important supermap that can be proven to be unique is the unitary transposition [26, 29]. Here it is interesting to remark that unitary transposition cannot be implemented deterministic and its probability of success is limited by $1/d^2$. Note however that our proofs only rely on linearity and the fact that quantum supermaps should be completely CP preserving and do not make use of normalization or

causal constraints. Due to that, we can also tackle probabilistic quantum transformations such as unitary transposition.

Corollary 2 (Unitary transposition). Consider the 1-slot supermap on unitary operations that transforms a unitary operator U into its transpose operator U^T with with some fixed success probability p . More formally, let $C \geq 0$ be the Choi operator of a supermap which respects $\text{Tr}_{\mathcal{IO}}(C J_U^t) = p J_{U^T}$ for all unitary operators U . The action of C on a general operation with Choi operator $J_\Lambda \in \mathcal{L}(\mathcal{I} \otimes \mathcal{O})$ is necessarily given by $\text{Tr}_{\mathcal{IO}}(C J_\Lambda^t) = p F J_\Lambda F$ where F denotes the flip (swap) operator defined by $F|\phi\psi\rangle = |\psi\phi\rangle$ for all $|\phi\rangle$ and $|\psi\rangle$.

Proof. A possible Choi operator for unitary transposition is given by $C'_0 = F_{\mathcal{IO}} C_0 F_{\mathcal{IO}}^\dagger$. Now, consider the condition for unitary operations

$$\text{Tr}_{\mathcal{IO}}(C' J_U^t) = \text{Tr}_{\mathcal{IO}}(C'_0 J_U^t). \quad (17)$$

In this case, by introducing $C = (F_{\mathcal{IO}})^\dagger C' (F_{\mathcal{IO}})$, the l.h.s. can be written as

$$\begin{aligned} \text{Tr}_{\mathcal{IO}}(C' J_U^t) &= \text{Tr}_{\mathcal{IO}}[(F_{\mathcal{IO}}) C (F_{\mathcal{IO}})^\dagger J_U^t] \\ &= \text{Tr}_{\mathcal{IO}}[C (F_{\mathcal{IO}})^\dagger J_U^t (F_{\mathcal{IO}})] \\ &= \text{Tr}_{\mathcal{IO}}[C (J_{U^t})^t] \end{aligned} \quad (18)$$

whereas the r.h.s. can be written as

$$\text{Tr}_{\mathcal{IO}}(C'_0 J_U^t) = \text{Tr}_{\mathcal{IO}}[C (J_{U^t})^t]. \quad (19)$$

Since U^t is also a unitary operator and spans the whole linear span of $\text{span}\{J_U\}$, the uniqueness of C' is equivalent to the uniqueness of the identity supermap C . \square

Remark 1. In general, for a given action of a supermap on unitary operations, an extension of the domain of the supermap is not necessarily unique under the three assumptions: the action on unitary operation; the linearity; and the positivity. Thus, it is natural to consider what extra condition may lead to the uniqueness. The results of Corollary 1 shows the Choi operator of the supermap $\tilde{U} \mapsto \tilde{B} \circ \tilde{U} \circ \tilde{A}$ is a rank-1 operator. One could wonder that, if there exists a rank-1 positive semidefinite operator extension, then there is a unique extension. Unfortunately, this hypothesis is not enough for the uniqueness of the extension as we will now show.

Consider a supermap transforming all two-dimensional unitary operations to the two-dimensional unitary operation I , the identity channel, with some probability. That is, there exists a Choi operator C_0 such that $\text{Tr}_{\mathcal{IO}}(C_0 J_U^t) \propto J_I$. For $0 \leq p \leq 1$, we define the Choi operator $C_p \in \mathcal{L}((\mathcal{I} \otimes \mathcal{O}) \otimes (\mathcal{P} \otimes \mathcal{F}))$

$$C_p := \begin{pmatrix} 1 & p & p & 1 \\ p & 1 & 1 & p \\ p & 1 & 1 & p \\ 1 & p & p & 1 \end{pmatrix} \otimes \phi^+, \quad (20)$$

where ϕ^+ denotes the maximally entangled state which is proportional to the Choi operator of the identity channel. We can check that for any $0 \leq p \leq 1$, the positivity constraint $C_p \geq 0$ is respected and $\text{Tr}_{\mathcal{I}\mathcal{O}}(C_p J_U^t) \propto J_I$. But note however that when $p = 1$, C_p is a rank-1 operator. Hence, the existence of a rank-1 is not sufficient for proving the uniqueness of the corresponding supermap on unitary operations.

5 Conclusion and Future Scope

In this paper we have addressed the question of whether the description of a higher-order process is uniquely determined by its action on unitary operations. We proved that, differently than general processes, the action of the quantum switch on unitary operations uniquely defines its action on general quantum operations. Our proof follows from linearity and the fact that every higher-order quantum process transforming quantum operations must respect the completely CP preserving property. We have also shown this uniqueness property hold for different classes of higher-order quantum processes which include unitary identity supermap, unitary transposition [26], and qubit unitary complex conjugate [28].

In order to tackle more general processes, it would be useful to find a simple criterium to identify which processes are uniquely defined by its action on unitary operations. Here particular cases of interest would be to analyze the n -slot quantum switch [9], and processes which may be viewed as generalizations of the switch, such as the ones which can be described by coherent control of causal order [25, 30, 31].

We remark that when multiple uses of an input operation are allowed, the action on unitary operation cannot be uniquely determined for general quantum operations in general. In particular, if a higher-order quantum operation can be implemented with k uses of an input operation, then if more than k uses is allowed, the uniqueness does not hold because it is always possible to discard a certain number of input operations. Even though, the completely CP preserving condition may be used to restrict the form of the supermap.

Another interesting future direction would be to understand how the constraints of normalization would limit the possible extensions of quantum processes. In this work we have obtained our results simply by arguments based on positivity, but the normalization constraints for deterministic transformations or linear constraints arising from causal relations [4] may considerably restrict the possibilities of extending the action of processes to non-unitary operations. We note that, when extending lower-order operations from a restricted set of quantum states, example 5 of Ref. [19] shows that requiring the transformation to be deterministic may be a severe constraint. We then expect that stronger claims may be proven by imposing that

a higher-order process should be deterministic.

As we show here, it is possible to strongly restrict the form of a supermap given only its action on unitary operations, and such a technique may reduce the difficulty on the analysis of the performance of higher-order quantum operation, since it is enough to perform an optimization on a restricted set of supermaps. We hope that questions, answers, and the methods covered in this paper contribute to our understanding on how positivity conditions restrict higher-order quantum operations and that this work paves the way for understanding the extension of general quantum processes.

Acknowledgments

We acknowledge Simon Milz for comments on the first version of this manuscript. This work was supported by MEXT Quantum Leap Flagship Program (MEXT QLEAP) Grant No. JPMXS0118069605, JPMXS0120351339, Japan Society for the Promotion of Science (JSPS) KAKENHI Grant No. 17H01694, 18H04286, 18K13467, 21H03394 and Advanced Leading Graduate Course for Photon Science (ALPS). M. T. Quintino acknowledges the Austrian Science Fund (FWF) through the SFB project ‘‘BeyondC’’ (sub-project F7103), a grant from the Foundational Questions Institute (FQXi) Fund and a grant from the John Templeton Foundation (sub-project F7103) as part of the The Quantum Information Structure of Spacetime (QISS) Project (qiss.fr). The opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the John Templeton Foundation. This project has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 801110 and the Austrian Federal Ministry of Education, Science and Research (BMBWF). It reflects only the authors’ view, the EU Agency is not responsible for any use that may be made of the information it contains.

References

- [1] G. Chiribella, G. M. D’Ariano, P. Perinotti, and B. Valiron, Quantum computations without definite causal structure, *Phys. Rev. A* **88**, 022318 (2013), [arXiv:0912.0195 \[quant-ph\]](#).
- [2] G. Chiribella, G. M. D’Ariano, and P. Perinotti, Quantum Circuit Architecture, *Phys. Rev. Lett.* **101**, 060401 (2008), [arXiv:0712.1325 \[quant-ph\]](#).
- [3] G. Chiribella, G. M. D’Ariano, and P. Perinotti, Transforming quantum operations: Quantum supermaps, *EPL (Europhysics Letters)* **83**, 30004 (2008), [arXiv:0804.0180 \[quant-ph\]](#).
- [4] G. Chiribella, G. M. D’Ariano, and P. Perinotti, Theoretical framework for quantum networks,

- Phys. Rev. A* **80**, 022339 (2009), [arXiv:0904.4483 \[quant-ph\]](#).
- [5] A. Bisio and P. Perinotti, Theoretical framework for higher-order quantum theory, *Proceedings of the Royal Society of London Series A* **475**, 20180706 (2019), [arXiv:1806.09554 \[quant-ph\]](#).
- [6] O. Oreshkov, F. Costa, and Č. Brukner, Quantum correlations with no causal order, *Nature Communications* **3**, 1092 (2012), [arXiv:1105.4464 \[quant-ph\]](#).
- [7] M. Araújo, A. Feix, M. Navascués, and Č. Brukner, A purification postulate for quantum mechanics with indefinite causal order, *Quantum* **1**, 10 (2017), [arXiv:1611.08535 \[quant-ph\]](#).
- [8] G. Chiribella, Perfect discrimination of no-signalling channels via quantum superposition of causal structures, *Phys. Rev. A* **86**, 040301 (2012), [arXiv:1109.5154 \[quant-ph\]](#).
- [9] M. Araújo, F. Costa, and Č. Brukner, Computational advantage from quantum-controlled ordering of gates, *Phys. Rev. Lett.* **113**, 250402 (2014), [arXiv:1401.8127 \[quant-ph\]](#).
- [10] P. A. Guérin, A. Feix, M. Araújo, and Č. Brukner, Exponential Communication Complexity Advantage from Quantum Superposition of the Direction of Communication, *Phys. Rev. Lett.* **117**, 100502 (2016), [arXiv:1605.07372 \[quant-ph\]](#).
- [11] D. Ebler, S. Salek, and G. Chiribella, Enhanced Communication with the Assistance of Indefinite Causal Order, *Phys. Rev. Lett.* **120**, 120502 (2018), [arXiv:1711.10165 \[quant-ph\]](#).
- [12] S. Salek, D. Ebler, and G. Chiribella, Quantum communication in a superposition of causal orders, (2018), [arXiv:1809.06655 \[quant-ph\]](#).
- [13] G. Chiribella, M. Banik, S. Sankar Bhattacharya, T. Guha, M. Alimuddin, A. Roy, S. Saha, S. Agrawal, and G. Kar, Indefinite causal order enables perfect quantum communication with zero capacity channel, (2018), [arXiv:1810.10457 \[quant-ph\]](#).
- [14] K. Goswami, Y. Cao, G. A. Paz-Silva, J. Romero, and A. G. White, Increasing communication capacity via superposition of order, *Phys. Rev. Research* **2**, 033292 (2020), [arXiv:1807.07383 \[quant-ph\]](#).
- [15] Y. Guo, X.-M. Hu, Z.-B. Hou, H. Cao, J.-M. Cui, B.-H. Liu, Y.-F. Huang, C.-F. Li, G.-C. Guo, and G. Chiribella, Experimental Transmission of Quantum Information Using a Superposition of Causal Orders, *Phys. Rev. Lett.* **124**, 030502 (2020), [arXiv:1811.07526 \[quant-ph\]](#).
- [16] K. Kraus, *States, effects and operations: fundamental notions of quantum theory* (Springer, 1983).
- [17] A. Chefles, Deterministic quantum state transformations, *Physics Letters A* **270**, 14–19 (2000), [arXiv:quant-ph/9911086 \[quant-ph\]](#).
- [18] A. Chefles, R. Jozsa, and A. Winter, On the existence of physical transformations between sets of quantum states, *International Journal of Quantum Information* **02**, 11–21 (2004), [quant-ph/0307227](#).
- [19] T. Heinosaari, M. A. Jivulescu, D. Reeb, and M. M. Wolf, Extending quantum operations, *Journal of Mathematical Physics* **53**, 102208–102208 (2012), [arXiv:1205.0641 \[math-ph\]](#).
- [20] Q. Dong, M. T. Quintino, A. Soeda, and M. Murao, Implementing positive maps with multiple copies of an input state, *Phys. Rev. A* **99**, 052352 (2019), [arXiv:1808.05788 \[quant-ph\]](#).
- [21] E. Haapasalo, T. Heinosaari, and J. P. Pellonpää, When do pieces determine the whole? Extreme marginals of a completely positive map, *Reviews in Mathematical Physics* **26**, 1450002 (2014), [arXiv:1209.5933 \[quant-ph\]](#).
- [22] L. Guerini and M. Terra Cunha, Uniqueness of the joint measurement and the structure of the set of compatible quantum measurements, *Journal of Mathematical Physics* **59**, 042106 (2018), [arXiv:1711.04804 \[quant-ph\]](#).
- [23] M.-D. Choi, Completely positive linear maps on complex matrices, *Linear Algebra and its Applications* **10**, 285–290 (1975).
- [24] A. Jamiołkowski, Linear transformations which preserve trace and positive semidefiniteness of operators, *Reports on Mathematical Physics* **3**, 275–278 (1972).
- [25] W. Yokojima, M. T. Quintino, A. Soeda, and M. Murao, Consequences of preserving reversibility in quantum superchannels, *Quantum* **5**, 441 (2021), [2003.05682](#).
- [26] M. T. Quintino, Q. Dong, A. Shimbo, A. Soeda, and M. Murao, Probabilistic exact universal quantum circuits for transforming unitary operations, *Phys. Rev. A* **100**, 062339 (2019), [arXiv:1909.01366 \[quant-ph\]](#).
- [27] Q. Dong, M. T. Quintino, A. Soeda, and M. Murao, Success-or-Draw: A Strategy Allowing Repeat-Until-Success in Quantum Computation, *Phys. Rev. Lett.* **126**, 150504 (2021), [arXiv:2011.01055 \[quant-ph\]](#).
- [28] J. Miyazaki, A. Soeda, and M. Murao, Complex conjugation supermap of unitary quantum maps and its universal implementation protocol, *Phys. Rev. Research* **1**, [arXiv:1706.03481 \(2019\)](#), [arXiv:1706.03481 \[quant-ph\]](#).
- [29] M. T. Quintino, Q. Dong, A. Shimbo, A. Soeda, and M. Murao, Reversing unknown quantum transformations: A universal quantum circuit for inverting general unitary operations, *Phys. Rev. Lett.* **123**, 210502 (2019), [arXiv:1810.06944 \[quant-ph\]](#).
- [30] J. Barrett, R. Lorenz, and O. Oreshkov, Cyclic quantum causal models, *Nature Communications* **12**, 885 (2021), [arXiv:2002.12157 \[quant-ph\]](#).

- [31] J. Wechs, H. Dourdent, A. A. Abbott, and C. Branciard, Quantum circuits with classical versus quantum control of causal order, (2021), [arXiv:2101.08796 \[quant-ph\]](#).

A The Elements in the Span of the Choi Operator of Unitary Operations

Lemma 3. $|ii\rangle\langle jj|, |ij\rangle\langle ji| \in \text{span}\{J_U\}$ holds for $i \neq j$.

Proof. Since only the linear span is considered, we use unnormalized quantum states for convenience. Consider the following maximally entangled state

$$|\psi_{\theta,\phi}\rangle = |ii\rangle + e^{i\theta}|jj\rangle + e^{i\phi}|KK\rangle, \quad (21)$$

where $|KK\rangle := \sum_{k \neq i,j} |kk\rangle$, then $|\psi_{\theta,\phi}\rangle\langle\psi_{\theta,\phi}| = J_U$ for some U and $|\psi_{\theta,\phi}\rangle\langle\psi_{\theta,\phi}| \in \text{span}\{J_U\}$. Since

$$|\psi_{\theta,\phi}\rangle\langle\psi_{\theta,\phi}| = |ii\rangle\langle ii| + |jj\rangle\langle jj| + |KK\rangle\langle KK| \quad (22)$$

$$+ e^{-i\theta}|ii\rangle\langle jj| + e^{-i\phi}|ii\rangle\langle KK| + e^{i(\theta-\phi)}|jj\rangle\langle KK| \quad (23)$$

$$+ e^{i\theta}|jj\rangle\langle ii| + e^{i\phi}|KK\rangle\langle ii| + e^{-i(\theta-\phi)}|KK\rangle\langle jj|, \quad (24)$$

we obtain

$$\int_0^{2\pi} d\theta \int_0^{2\pi} d\phi e^{i\theta} |\psi_{\theta,\phi}\rangle\langle\psi_{\theta,\phi}| = |ii\rangle\langle jj| \in \text{span}\{J_U\} \quad (25)$$

for all $i \neq j$. The same calculation with the maximally entangled state

$$|\psi'_{\theta,\phi}\rangle = |ij\rangle + e^{i\theta}|ji\rangle + e^{i\phi}|KK\rangle, \quad (26)$$

leads to $|ij\rangle\langle ji| \in \text{span}\{J_U\}$ for all $i \neq j$. \square

Lemma 4. $|ij\rangle\langle i'j'| \in \text{span}\{J_U\}$ holds for $i \neq j \neq i' \neq j'$.

Proof. Consider the maximally entangled state

$$|\psi_{\theta,\phi}\rangle = |ij\rangle + e^{i\theta}|i'j'\rangle + e^{i\phi}|KK\rangle \quad (27)$$

where $|KK\rangle := |ji\rangle + |j'i'\rangle + \sum_{k \neq i,j,i',j'} |kk\rangle$. Then the same calculation of Lemma 3 proves $|ij\rangle\langle i'j'| \in \text{span}\{J_U\}$. \square

Lemma 5. $|ij\rangle\langle i'j'| \in \text{span}\{J_U\}$ holds if only two of i, j, i', j' are the same and others are different.

Proof. There are 6 cases that only two of i, j, i', j' are the same and others are different. If $|ij\rangle\langle i'j'| \in \text{span}\{J_U\}$, then the conjugate transpose also satisfies $|i'j'\rangle\langle ij| \in \text{span}\{J_U\}$. Thus, it is enough to consider the following four cases with $i \neq j \neq k$.

$$|ii\rangle\langle jk| \in \text{span}\{J_U\} \quad (28)$$

$$|ij\rangle\langle jk| \in \text{span}\{J_U\} \quad (29)$$

$$|ij\rangle\langle ik| \in \text{span}\{J_U\} \quad (30)$$

$$|ij\rangle\langle kj| \in \text{span}\{J_U\} \quad (31)$$

For each of them, consider the following maximally entangled states, and the same calculation of Lemma 3 proves that they are in $\text{span}\{J_U\}$, respectively.

$$1. |\psi_{\theta,\phi}\rangle = |ii\rangle + e^{i\theta}|jk\rangle + e^{i\phi}|LL\rangle \text{ with } |LL\rangle := |kj\rangle + \sum_{l \neq i,j,k} |ll\rangle.$$

$$2. |\psi_{\theta,\phi}\rangle = |ij\rangle + e^{i\theta}|jk\rangle + e^{i\phi}|LL\rangle \text{ with } |LL\rangle := |ki\rangle + \sum_{l \neq i,j,k} |ll\rangle.$$

$$3. |\psi_{\theta,\phi}\rangle = |i\rangle(|j\rangle + e^{i\theta}|k\rangle) + e^{i\phi}|LL\rangle \text{ with } |LL\rangle := |j\rangle(|j\rangle - e^{i\theta}|k\rangle) + |ki\rangle + \sum_{l \neq i,j,k} |ll\rangle.$$

$$4. |\psi_{\theta,\phi}\rangle = (|i\rangle + e^{i\theta}|k\rangle)|j\rangle + e^{i\phi}|LL\rangle \text{ with } |LL\rangle := (|i\rangle - e^{i\theta}|k\rangle)|i\rangle + |jk\rangle + \sum_{l \neq i,j,k} |ll\rangle.$$

\square

Lemma 6. $\sum_{j=i+k} |ij\rangle\langle ij| \in \text{span}\{J_U\}$ holds for $k = 0, \dots, d-1$.

Proof. Consider the maximally entangled state

$$|\psi_{\theta_i}\rangle = \sum_i e^{i\theta_i} |i, i+k\rangle. \quad (32)$$

Note that we consider the d -dimensional system and $|i+k\rangle$ indicates $|i+k \pmod{d}\rangle$. Then we obtain

$$\int_0^{2\pi} \prod_i d\theta_i |\psi_{\theta_i}\rangle\langle\psi_{\theta_i}| = \sum_i |i, i+k\rangle\langle i, i+k| = \sum_{j=i+k} |ij\rangle\langle ij| \in \text{span}\{J_U\}. \quad (33)$$

□

Lemma 7. $(|ij\rangle\langle ii| - |jj\rangle\langle ji|), (|ji\rangle\langle ii| - |jj\rangle\langle ij|) \in \text{span}\{J_U\}$ holds for $i \neq j$.

Proof. Consider the maximally entangled state

$$|\psi_{\theta, \phi}^1\rangle = (|i\rangle + e^{i\theta_2}|j\rangle)(|i\rangle + e^{i\theta_2}|j\rangle) + e^{i\theta_1}(|i\rangle - e^{i\theta_2}|j\rangle)(|i\rangle - e^{i\theta_2}|j\rangle) + e^{i\phi}|KK\rangle \quad (34)$$

$$|\psi_{\theta, \phi}^2\rangle = (|i\rangle + e^{i\theta_2}|j\rangle)(|i\rangle - e^{i\theta_2}|j\rangle) + e^{i\theta_1}(|i\rangle - e^{i\theta_2}|j\rangle)(|i\rangle + e^{i\theta_2}|j\rangle) + e^{i\phi}|KK\rangle \quad (35)$$

where $|KK\rangle := \sum_{k \neq i, j} |kk\rangle$. Here

$$\int_0^{2\pi} d\theta_1 d\theta_2 d\phi e^{i\theta_1} e^{-i\theta_2} |\psi_{\theta, \phi}^1\rangle\langle\psi_{\theta, \phi}^1| = |ij\rangle\langle ii| + |ji\rangle\langle ii| - |jj\rangle\langle ij| - |jj\rangle\langle ji| \quad (36)$$

$$\int_0^{2\pi} d\theta_1 d\theta_2 d\phi e^{i\theta_1} e^{-i\theta_2} |\psi_{\theta, \phi}^2\rangle\langle\psi_{\theta, \phi}^2| = -|ij\rangle\langle ii| + |ji\rangle\langle ii| - |jj\rangle\langle ij| + |jj\rangle\langle ji|, \quad (37)$$

holds, and by considering the sum and difference, we obtain

$$(|ij\rangle\langle ii| - |jj\rangle\langle ji|), (|ji\rangle\langle ii| - |jj\rangle\langle ij|) \in \text{span}\{J_U\}. \quad (38)$$

□

B The proof of Lemma 2

Consider the identity supermap $\tilde{\mathcal{C}}$ that transforms an input unitary operation \tilde{U} on $\mathcal{I} \otimes \mathcal{O} \cong \mathbb{C}^d \otimes \mathbb{C}^d$ to \tilde{U} on $\mathcal{P} \otimes \mathcal{F} \cong \mathbb{C}^d \otimes \mathbb{C}^d$. We show that this supermap transforms an arbitrary input operation $\tilde{\Lambda}$ on $\mathcal{I} \otimes \mathcal{O}$ to itself on $\mathcal{P} \otimes \mathcal{F}$ by assuming the completely CP preserving property. That is, the Choi operator of the identity supermap C satisfies

$$C = \sum_{i, j, k, l=1}^d |ij\rangle\langle kl| \otimes \tilde{\mathcal{C}}(|ij\rangle\langle kl|) \in \mathcal{L}(\mathcal{I} \otimes \mathcal{O} \otimes \mathcal{P} \otimes \mathcal{F}), \quad (39)$$

where $\tilde{\mathcal{C}}$ denotes the corresponding map of C , i.e., $\tilde{\mathcal{C}}(J) = \text{Tr}_{\mathcal{IO}}(CJ^t)$ for $J \in \mathcal{L}(\mathcal{I} \otimes \mathcal{O})$, and we show that this Choi operator is uniquely determined as

$$C_0 = \sum_{i, j, k, l=1}^d |ij\rangle\langle kl| \otimes |ij\rangle\langle kl| \in \mathcal{L}(\mathcal{I} \otimes \mathcal{O} \otimes \mathcal{P} \otimes \mathcal{F}) \quad (40)$$

under the following assumptions: its action on unitary operations is determined as $\text{Tr}_{\mathcal{IO}}(CJ_U^t) = \text{Tr}_{\mathcal{IO}}(C_0J_U^t)$ for an arbitrary unitary operation U ; and it is a positive operator $C \geq 0$. Note that the action of $\tilde{\mathcal{C}}$ is given by $\tilde{\mathcal{C}}(J_U) = \text{Tr}_{\mathcal{IO}}(CJ_U^t) = \text{Tr}_{\mathcal{IO}}(C_0J_U^t) = J_U$ by assumption. We first consider the diagonal elements of C , then consider the off-diagonal elements of C .

(First part: diagonal elements of C) Since $\sum_{i, j=1}^d |ij\rangle\langle ij| = I \in \text{span}\{J_U\}$, the action of $\tilde{\mathcal{C}}$ is determined as $\tilde{\mathcal{C}}(I) = I$, and we obtain

$$\sum_{i, j=1}^d |ij\rangle\langle ij| = \tilde{\mathcal{C}}\left(\sum_{i, j=1}^d |ij\rangle\langle ij|\right) = \sum_{i, j=1}^d \tilde{\mathcal{C}}(|ij\rangle\langle ij|). \quad (41)$$

Thus, for $|ijkl\rangle \in \mathcal{I} \otimes \mathcal{O} \otimes \mathcal{P} \otimes \mathcal{F}$

$$\begin{aligned} \sum_{i,j=1}^d \langle ijkl|C|ijkl\rangle &= \sum_{i,j=1}^d \langle kl|\text{Tr}(C|i_j\rangle\langle i_j|^t)|kl\rangle = \sum_{i,j=1}^d \langle kl|\tilde{\mathcal{C}}(|i_j\rangle\langle i_j|)|kl\rangle \\ &= \langle kl|\sum_{i,j=1}^d \tilde{\mathcal{C}}(|i_j\rangle\langle i_j|)|kl\rangle = \langle kl|(\sum_{i,j=1}^d |i_j\rangle\langle i_j|)|kl\rangle = 1, \end{aligned} \quad (42)$$

holds for all k, l , and we obtain

$$\text{Tr } C = \sum_{i,j,k,l=1}^d \langle ijkl|C|ijkl\rangle = d^2. \quad (43)$$

Since every diagonal element of a positive semidefinite operator is non-negative, Eq. (42) shows that every diagonal element of C , say a , satisfies $0 \leq a \leq 1$, and Eq. (43) shows that the sum of all diagonal elements of C is equal to d^2 .

Since $|i_j\rangle\langle j_i|, |i_i\rangle\langle j_j| \in \text{span}\{J_U\}$ holds for all $i \neq j$ from Lemma 3, the action of $\tilde{\mathcal{C}}$ is determined as

$$\tilde{\mathcal{C}}(|i_j\rangle\langle j_i|) = |i_j\rangle\langle j_i|, \quad \tilde{\mathcal{C}}(|i_i\rangle\langle j_j|) = |i_i\rangle\langle j_j|, \quad (44)$$

and the Choi operator C satisfies

$$\langle ijij|C|jiji\rangle = 1, \quad \langle iiii|C|jjjj\rangle = 1. \quad (45)$$

Since $C \geq 0$ all principal minors of C are non-negative. Consider the following two-dimensional submatrices of C

$$\begin{pmatrix} \langle ijij|C|ijij\rangle & \langle ijij|C|jiji\rangle \\ \langle jiji|C|ijij\rangle & \langle jiji|C|jiji\rangle \end{pmatrix}, \quad \begin{pmatrix} \langle iiii|C|iiii\rangle & \langle iiii|C|jjjj\rangle \\ \langle jjjj|C|iiii\rangle & \langle jjjj|C|jjjj\rangle \end{pmatrix}, \quad (46)$$

with $i \neq j$. Eq. (45) indicates that all the off-diagonal elements of these matrices are 1, and the diagonal elements are between 0 and 1 due to Eq. (42). Then these matrices are positive semi-definite only if

$$\langle ijij|C|ijij\rangle = 1, \quad \langle iiii|C|iiii\rangle = 1 \quad (47)$$

for all $i \neq j$.

Since the sum of all diagonal elements is equal to d^2 due to Eq. (43), and Eq. (47) provides the $d(d-1)+d = d^2$ elements that are equal to 1, the other diagonal elements of C are equal to 0. For positive semi-definite matrices, if a diagonal element is 0, all elements in that row or column are also 0. Thus, we have

$$\langle i_1 j_1 k_1 l_1|C|i_2 j_2 k_2 l_2\rangle \neq 0 \quad (48)$$

only if $(i_1, j_1, k_1, l_1) = (i, j, i, j)$ and $(i_2, j_2, k_2, l_2) = (k, l, k, l)$ for some i, j, k, l .

(Second part: off-diagonal elements of W) The second part of the proof is to show that all the elements which are non-zero are equal to 1. Consider a Choi operator $J \in \text{span}\{J_U\}$ with $\langle ij|J|kl\rangle \neq 0$ for all $i \neq j, k \neq l$, e.g., the Choi operator of the Fourier transform. Since $J \in \text{span}\{J_U\}$, the equality

$$\langle ij|\tilde{\mathcal{C}}(J)|kl\rangle = \langle ij|J|kl\rangle \quad (49)$$

holds for all $i \neq j, k \neq l$. On the other hand, the action determined by the Choi operator C is given by

$$\langle ij|\tilde{\mathcal{C}}(J)|kl\rangle = \langle ij|\text{Tr}_{\mathcal{I}\mathcal{O}}[C(J_{\mathcal{I}\mathcal{O}}^t \otimes I_{\mathcal{P}\mathcal{F}})]|kl\rangle \quad (50)$$

$$= \text{Tr}(\langle ij|C|kl\rangle J^t) \quad (51)$$

$$= \sum_{i',j',k',l'} \langle i'j'|C|k'l'kl\rangle \langle k'l'|J^t|i'j'\rangle \quad (52)$$

$$= \langle ijij|C|klkl\rangle \langle kl|J^t|ij\rangle \quad (53)$$

$$= \langle ijij|C|klkl\rangle \langle ij|J|kl\rangle, \quad (54)$$

where the fourth equality holds due to Eq. (47). Therefore, by considering the condition that both Eq. (49) and Eq. (54) hold, we obtain $\langle ijij|C|klkl\rangle = 1$ for all $i \neq j, k \neq l$. Similarly, consider a Choi operator $J \in \text{span}\{J_U\}$ with $\langle ii|J|kl\rangle \neq 0$ for all i and $k \neq l$, e.g., the Choi operator of the Fourier transform, we obtain $\langle iiii|C|klkl\rangle = 1$ for all i and $k \neq l$. Similar calculations show that $\langle ijji|C|kkkk\rangle = 1$ for $i \neq j$ and all k , and $\langle iiii|C|kkkk\rangle = 1$ for all i and k . Thus, we uniquely determine all the elements of C and obtain $C = C_0$.

C The Proof of Lemma 1

The Choi operator W_0 of quantum switch that we prove is given by Eq. (10). For $J = |ijkl\rangle\langle i'j'k'l'| \in \mathcal{L}(\mathcal{I}_1 \otimes \mathcal{O}_1 \otimes \mathcal{I}_2 \otimes \mathcal{O}_2)$, the action of W_0 can be evaluated as

$$\begin{aligned} \text{Tr}_{in}(W_0 J^t) &= \langle ijkl|W_0\rangle\langle W_0||i'j'k'l'\rangle = \delta_{jk}\delta_{j'k'}|il00\rangle\langle i'l'00| + \delta_{il}\delta_{i'l'}|kj11\rangle\langle k'j'11| \\ &\quad + \delta_{jk}\delta_{i'l'}|il00\rangle\langle k'j'11| + \delta_{il}\delta_{j'k'}|kj11\rangle\langle i'l'00|. \end{aligned} \quad (55)$$

Note that for $J \in \text{span}\{J_{U_1} \otimes J_{U_2}\}$, $\text{Tr}_{in}(W J^t) = \text{Tr}_{in}(W_0 J^t)$ holds by assumption.

Here each element of W_0 is 0 or 1. In order to prove Theorem 1, it is enough to prove that each element of also W satisfies $|(W)_{ij}| = 0$ or 1. This is because if an arbitrary W satisfies $|(W)_{ij}| = 0$ or 1 but $W \neq W_0$, then some elements of $\epsilon W + (1-\epsilon)W_0$, which is also a candidate for W , must have a value satisfying $0 < |(W)_{ij}| < 1$, which contradicts with the assumption that W satisfies $|(W)_{ij}| = 0$ or 1.

The rest of the proof is divided into two parts: the first part considers the diagonal elements of W , and the second part considers the off-diagonal elements of W .

(First part: diagonal elements of W) Consider $J = |ijkl\rangle\langle i'j'k'l'|$. For all i, j, k, l , there exist i', j', k', l' such that $J \in \text{span}\{J_{U_1} \otimes J_{U_2}\}$. That is, if $i = j$ then let $i' = j' (\neq i)$ so that $|ij\rangle\langle i'j'| \in \text{span}\{J_U\}$, and in this case, we relabel the variables i', j' as j . Similarly, if $i \neq j$ then let $i' = j, j' = i$ so that $|ij\rangle\langle i'j'| \in \text{span}\{J_U\}$. (See Sec. 3 or Lemma 3) The same arguments hold for k, l . We then obtain the followings for $i \neq j$ and $k \neq l$:

$$\widetilde{\mathcal{W}}(|ijkl\rangle\langle jilk|) = (\delta_{jk}|il00\rangle + \delta_{il}|kj11\rangle)(\delta_{il}\langle jk00| + \delta_{jk}\langle li11|) \quad (56)$$

$$\widetilde{\mathcal{W}}(|ijkk\rangle\langle jill|) = (\delta_{jk}|ik00\rangle + \delta_{ik}|kj11\rangle)(\delta_{il}\langle jl00| + \delta_{jl}\langle li11|) \quad (57)$$

$$\widetilde{\mathcal{W}}(|iikl\rangle\langle jllk|) = (\delta_{ik}|il00\rangle + \delta_{il}|ki11\rangle)(\delta_{jl}\langle jk00| + \delta_{jk}\langle lj11|) \quad (58)$$

$$\widetilde{\mathcal{W}}(|iikk\rangle\langle jlll|) = (\delta_{ik}|ik00\rangle + \delta_{ik}|ki11\rangle)(\delta_{jl}\langle jl00| + \delta_{jl}\langle lj11|), \quad (59)$$

where $\widetilde{\mathcal{W}}$ denotes the corresponding map of W , i.e., $\widetilde{\mathcal{W}}(J) = \text{Tr}_{in} W J^t$. Also, consider that $I \otimes I \in \text{span}\{J_{U_1} \otimes J_{U_2}\}$, we obtain

$$\text{Tr} \widetilde{\mathcal{W}}(I \otimes I) = \text{Tr}(\text{Tr}_{in} W) = \text{Tr} W = 2d^3. \quad (60)$$

By considering Eqs. (56) – (60) and the positivity $W \geq 0$, we obtain the following elements of W with $i \neq j, k \neq l$

$$\langle ijklil00|W|ijklil00\rangle = 1 \text{ for } j = k \quad (61)$$

$$\langle ijklkj11|W|ijklkj11\rangle = 1 \text{ for } i = l \quad (62)$$

$$\langle ijkkik00|W|ijkkik00\rangle = 1 \text{ for } j = k \quad (63)$$

$$\langle ijkkkj11|W|ijkkkj11\rangle = 1 \text{ for } i = k \quad (64)$$

$$\langle iiklil00|W|iiklil00\rangle = 1 \text{ for } i = k \quad (65)$$

$$\langle iiklki11|W|iiklki11\rangle = 1 \text{ for } i = l \quad (66)$$

$$\langle iikkik00|W|iikkik00\rangle = 1 \text{ for } i = k \quad (67)$$

$$\langle iikkki11|W|iikkki11\rangle = 1 \text{ for } i = k, \quad (68)$$

or equivalently,

$$\langle ijjlil00|W|ijjlil00\rangle = 1 \text{ for } i \neq j, j \neq l \quad (69)$$

$$\langle ijkkkj11|W|ijkkkj11\rangle = 1 \text{ for } i \neq j, k \neq i \quad (70)$$

$$\langle ijjjij00|W|ijjjij00\rangle = 1 \text{ for } i \neq j \quad (71)$$

$$\langle ijiiij11|W|ijiiij11\rangle = 1 \text{ for } i \neq j \quad (72)$$

$$\langle iiiiil00|W|iiiiil00\rangle = 1 \text{ for } i \neq l \quad (73)$$

$$\langle iikiki11|W|iikiki11\rangle = 1 \text{ for } i \neq k \quad (74)$$

$$\langle iiiiii00|W|iiiiii00\rangle = 1 \quad (75)$$

$$\langle iiiiii11|W|iiiiii11\rangle = 1. \quad (76)$$

The proof is given in Appendix D. The number of terms in 1st, 3rd, 5th, and 7th lines (or 2nd, 4th, 6th, 8th lines) is given by $d(d-1)^2, d(d-1), d(d-1), d$ respectively, and the total number of terms in these 8 lines is $2d^3$. Since $\text{Tr} W = 2d^3$, all the other diagonal terms of W are 0, that is, $\langle \psi|W|\psi'\rangle = 0$ for all $|\psi\rangle, |\psi'\rangle$ not of these forms.

Remark 2. Here we illustrate the key idea of the proof in Appendix D with a simpler example. Consider a linear map $\widetilde{W} : \mathcal{L}(\mathbb{C}^2) \rightarrow \mathcal{L}(\mathbb{C}^2)$ and the corresponding Choi operator $W \in \mathcal{L}(\mathbb{C}^4)$ satisfying

$$\widetilde{W}(|i\rangle\langle j|) = |i\rangle\langle j| \quad \text{for } (i, j) = (0, 1) \text{ and } (i, j) = (1, 0) \quad (77)$$

$$\text{Tr} W = 2 \quad (78)$$

$$W \geq 0, \quad (79)$$

then the matrix W has the form of

$$W = \begin{pmatrix} a & b & 0 & 1 \\ c & d & 0 & 0 \\ 0 & 0 & e & f \\ 1 & 0 & g & h \end{pmatrix} \quad (80)$$

with $a + d + e + h = 2$. From the positivity $W \geq 0$, the determinant of all its minors has to be non-negative, hence we obtain $ah - 1 \geq 0$ and $a, d, e, h \geq 0$. By combining these constraints we can verify that

$$2 = a + d + e + h \quad (81)$$

$$\geq 2\sqrt{ah} + e + d, \quad \left[\text{since } \frac{ah}{2} \geq \sqrt{ah} \right] \quad (82)$$

$$\geq 2 + e + d, \quad [\text{since } ah \geq 1] \quad (83)$$

$$\geq 2. \quad [\text{since } e \geq 0, d \geq 0] \quad (84)$$

Hence, we obtain $a + h = 2\sqrt{ah}$ as a necessary condition. Thus, we obtain $a = d = 1$ and the diagonal terms are uniquely determined as $(a, d, e, h) = (1, 0, 0, 1)$.

(Second part: off-diagonal elements of W) In this part, we show that every element of W satisfies $|(W)_{ij}| = 0$ or 1. From the positivity, if a diagonal element is 0, then all the elements of that row or column is 0. Therefore, we can assume that W is given by

$$\begin{aligned} \text{Tr}_{in} W J^t &= \langle ijkl | W | i'j'k'l' \rangle = a_{ijkli'j'k'l'} \delta_{jk} \delta_{j'k'} |il00\rangle\langle i'l'00| \\ &\quad + b_{ijkli'j'k'l'} \delta_{il} \delta_{i'l'} |kj11\rangle\langle k'j'11| \\ &\quad + c_{ijkli'j'k'l'} \delta_{jk} \delta_{i'l'} |il00\rangle\langle k'j'11| \\ &\quad + d_{ijkli'j'k'l'} \delta_{il} \delta_{j'k'} |kj11\rangle\langle i'l'00| \end{aligned} \quad (85)$$

with $|a_{ijkli'j'k'l'}|, |b_{ijkli'j'k'l'}|, |c_{ijkli'j'k'l'}|, |d_{ijkli'j'k'l'}| \leq 1$. (cf. Eq. (55)) As we mentioned, it is enough to show that all of the absolute values of them (with the non-vanishing Kronecker deltas) are 1. Considering that there are at most $(2d^3)^2$ non-zero elements, this is also equivalent to show that $\|W\|_1 = (2d^3)^2$, where $\|W\|_1$ denotes the element-wise 1-norm (i.e., the sum of absolute values of all elements in the computational basis). Since $\|W\|_1 \leq (2d^3)^2$ is trivial, we prove $\|W\|_1 \geq (2d^3)^2$ in the following.

Remark 3. Before proceeding to the proof, we briefly explain the key idea of the proof of this part by adding an extra assumption which makes the proof much simpler. Let $J = \sum_{ijkli'j'k'l'} |ijkl\rangle\langle i'j'k'l'|$, we have

$$\|W\|_1 = \sum_{ijkli'j'k'l'} \|\langle ijkl | W | i'j'k'l' \rangle\|_1 \geq \left\| \sum_{ijkli'j'k'l'} \langle ijkl | W | i'j'k'l' \rangle \right\|_1 = \|\text{Tr}_{in} W J^t\|_1. \quad (86)$$

Assume that $J \in \text{span}\{J_{U_1} \otimes J_{U_2}\}$ (which is the extra assumption and is not true for the case of the proof), then the r.h.s. can be evaluated using $\text{Tr}_{in} W J^t = \text{Tr}_{in} W_0 J^t$ as

$$\|W\|_1 \geq \|\text{Tr}_{in} W J^t\|_1 = \|\text{Tr}_{in} W_0 J^t\|_1 = (2d^3)^2, \quad (87)$$

which completes the proof.

In order to evaluate $\|W\|_1$, we first provide a grouping of $|ijkl\rangle\langle i'j'k'l'| = |ij\rangle\langle i'j'| \otimes |kl\rangle\langle k'l'|$, which covers all possible elements and also satisfies that each group is in $\text{span}\{J_{U_1} \otimes J_{U_2}\}$. See Sec. 3 for the proof that they are in $\text{span}\{J_{U_1} \otimes J_{U_2}\}$. We first classify $|ij\rangle\langle i'j'|$ into the following groups. (i, j, i', j' are assumed to be different by default)

1. If all i, j, i', j' are different, or if only two of them are the same, or $i = j, i' = j'$ or $i = j', j = i'$, then we use $|ij\rangle\langle i'j'| \in \text{span}\{J_{U_1} \otimes J_{U_2}\}$.

2. If $i = i', j = j'$ or $i = i' = j = j'$, then we use $\sum_{j=i+k} |ij\rangle\langle ij| \in \text{span}\{J_{U_1} \otimes J_{U_2}\}$.

3. If $i = j = j'$ or $i = j = j'$ or $i = i' = j'$ or $j = i' = j$, then we use $(|ij\rangle\langle ii| - |jj\rangle\langle ji|), (|ji\rangle\langle ii| - |jj\rangle\langle ij|) \in \text{span}\{J_{U_1} \otimes J_{U_2}\}$.

We denote the three groups as G_1 , G_2 and G_3 respectively. To make the statement clear, we provide the definitions explicitly as follows:

$$\begin{aligned} G_1 = \{ & |ij\rangle\langle i'j'| \mid i \neq j \neq i' \neq j', \\ & \text{or } i = j, i \neq i' \neq j' \quad (\text{and the remaining 5 patterns}), \\ & \text{or } i = j, i' = j', i \neq i', \\ & \text{or } i = j', j = i', i \neq j\}, \end{aligned} \quad (88)$$

$$G_2 = \left\{ \sum_{j=i+k} |ij\rangle\langle ij| \mid k = 0, \dots, d-1 \right\}, \quad (89)$$

$$G_3 = \{|ij\rangle\langle ii| - |jj\rangle\langle ji|, |ji\rangle\langle ii| - |jj\rangle\langle ij| \mid i \neq j\}. \quad (90)$$

The number of elements of each group is given by

$$\begin{aligned} |G_1| &= d(d-1)(d-2)(d-3) + 6d(d-1)(d-2) + 2d(d-1) \\ &= d(d-1)(d^2 + d - 4), \end{aligned} \quad (91)$$

$$|G_2| = d, \quad (92)$$

$$|G_3| = 2d(d-1). \quad (93)$$

Note that $|G_1| + d|G_2| + 2|G_3| = d^4$ holds. The same grouping is chosen for $|kl\rangle\langle k'l'|$. Since the three groups have no overlap between each other, we obtain

$$\|W\|_1 \geq \sum_{a,b=1}^3 \sum_{J \in G_a \otimes G_b} \|\text{Tr}_{in} W J^t\|_1. \quad (94)$$

Moreover, since all J considered here satisfies $J \in \text{span}\{J_{U_1} \otimes J_{U_2}\}$, we can evaluate the r.h.s. with $\text{Tr}_{in} W J^t = \text{Tr}_{in} W_0 J^t$.

The summation $\sum_{J \in G_a \otimes G_b} \|\text{Tr}_{in} W J^t\|_1$ for each a, b is evaluated as follows. See Appendix E for calculations. Note that it is enough to consider only 6 cases that satisfy $a \leq b$ because of the symmetry between the two systems.

$$\sum_{J \in G_1 \otimes G_1} \|\text{Tr}_{in} W J^t\|_1 = 2d(d-1)(2d^4 + 2d^3 - 18d^2 + 11d + 8) \quad (95)$$

$$\sum_{J \in G_1 \otimes G_2} \|\text{Tr}_{in} W J^t\|_1 = 2d(d-1)(2d^2 - d - 4) \quad (96)$$

$$\sum_{J \in G_2 \otimes G_2} \|\text{Tr}_{in} W J^t\|_1 = 2d^2(d+1) \quad (97)$$

$$\sum_{J \in G_1 \otimes G_3} \|\text{Tr}_{in} W J^t\|_1 = 4d(d-1)(4d^2 - 5d - 2) \quad (98)$$

$$\sum_{J \in G_2 \otimes G_3} \|\text{Tr}_{in} W J^t\|_1 = 4d(d-1)(d+2) \quad (99)$$

$$\sum_{J \in G_3 \otimes G_3} \|\text{Tr}_{in} W J^t\|_1 = 16d^2(d-1). \quad (100)$$

By using these equations, $\|W\|_1$ is evaluated as

$$\|W\|_1 \geq (2d^3)^2, \quad (101)$$

which completes the proof of Lemma 1.

D Evaluation of the Diagonal Terms in the Proof of Lemma 1

Here we provide the proof of Eqs. (61) – (68) from Eqs. (56) – (60) and the positivity $W \geq 0$. Let

$$S_1 = \{|ijklil00\rangle \mid i \neq j, k \neq l, j = k\} \cup \{|ijklkj11\rangle \mid i \neq j, k \neq l, i = l\} \quad (102)$$

$$S_2 = \{|jkkik00\rangle \mid i \neq j, j = k\} \quad (103)$$

$$S_3 = \{|jkkkj11\rangle \mid i \neq j, i = k\} \quad (104)$$

$$S_4 = \{|iklil00\rangle \mid k \neq l, i = k\} \quad (105)$$

$$S_5 = \{|iklki11\rangle \mid k \neq l, i = l\} \quad (106)$$

$$S_6 = \{|ikkik00\rangle \mid i = k\} \quad (107)$$

$$S_7 = \{|ikkki11\rangle \mid i = k\}, \quad (108)$$

and it is enough to prove that

$$\prod_{|\psi\rangle \in S_a} \langle \psi | W | \psi \rangle \geq 1 \quad (109)$$

holds for $a = 1, \dots, 7$. This is because if Eq. (109) holds, the inequality

$$2d^3 \geq \sum_{a=1}^7 \sum_{|\psi\rangle \in S_a} \langle \psi | W | \psi \rangle \geq \sum_{a=1}^7 |S_a| \left(\prod_{|\psi\rangle \in S_a} \langle \psi | W | \psi \rangle \right)^{\frac{1}{|S_a|}} \geq \sum_{a=1}^7 |S_a| = 2d^3 \quad (110)$$

also holds, and by considering the condition for equality, we obtain

$$\langle \psi | W | \psi \rangle = 1 \quad (111)$$

for all $|\psi\rangle \in S_a$ for all a . Here we only prove Eq. (109) for $a = 1, 2, 6$. The case of $a = 4$ can be proved in the same way as the case of $a = 2$ by considering the symmetry between the two input operations. The rest cases $a = 3, 5, 7$ can be proved in the same way as $a = 2, 4, 6$, respectively, by considering the symmetry between the action of the control qubit being 0 and 1.

(Case 1: S_1) In this case, we show the following inequality holds

$$\begin{aligned} \prod_{|\psi\rangle \in S_1} \langle \psi | W | \psi \rangle &= \left(\prod_{i \neq j, k \neq l, j = k} \langle ijklil00 | W | ijklil00 \rangle \right) \left(\prod_{i \neq j, k \neq l, i = l} \langle ijklkj11 | W | ijklkj11 \rangle \right) \\ &= \left(\prod_{i \neq k, k \neq l} \langle ikklil00 | W | ikklil00 \rangle \right) \left(\prod_{i \neq k, k \neq l} \langle kilkli11 | W | kilkli11 \rangle \right) \\ &= \prod_{i \neq k, k \neq l} \langle ikklil00 | W | ikklil00 \rangle \langle kilkli11 | W | kilkli11 \rangle \geq 1. \end{aligned} \quad (112)$$

Let $|\psi_{ikl}^0\rangle := |ikklil00\rangle$ and $|\psi_{ikl}^1\rangle := |kilkli11\rangle$, then $\langle \psi_{ikl}^0 | W | \psi_{ikl}^1 \rangle = 1$ holds for arbitrary i, k, l satisfying $i \neq k, k \neq l$ because of Eq. (56). By considering the positivity of the submatrix

$$\begin{pmatrix} \langle \psi_{ikl}^0 | W | \psi_{ikl}^0 \rangle & \langle \psi_{ikl}^0 | W | \psi_{ikl}^1 \rangle \\ \langle \psi_{ikl}^1 | W | \psi_{ikl}^0 \rangle & \langle \psi_{ikl}^1 | W | \psi_{ikl}^1 \rangle \end{pmatrix} = \begin{pmatrix} \langle \psi_{ikl}^0 | W | \psi_{ikl}^0 \rangle & 1 \\ 1 & \langle \psi_{ikl}^1 | W | \psi_{ikl}^1 \rangle \end{pmatrix} \geq 0, \quad (113)$$

we obtain the inequality

$$\langle \psi_{ikl}^0 | W | \psi_{ikl}^0 \rangle \langle \psi_{ikl}^1 | W | \psi_{ikl}^1 \rangle \geq 1, \quad (114)$$

which completes the proof of Eq. (112).

(Case 2: S_2) In this case, we show the following inequality holds

$$\begin{aligned} \prod_{|\psi\rangle \in S_2} \langle \psi | W | \psi \rangle &= \prod_{i \neq j, j = k} \langle ijkkik00 | W | ijkkik00 \rangle \\ &= \prod_{i \neq k} \langle \psi_{ik} | W | \psi_{ik} \rangle = \prod_{i < k} \langle \psi_{ik} | W | \psi_{ik} \rangle \langle \psi_{ki} | W | \psi_{ki} \rangle \geq 1, \end{aligned} \quad (115)$$

where we define $|\psi_{ik}\rangle := |ikkkik00\rangle$ for $i, k = 0, \dots, d-1$. Since $\langle\psi_{ik}|W|\psi_{ki}\rangle = 1$ holds arbitrary i, k satisfying $i \neq k$ because of Eq. (57), we consider the positivity of the submatrix

$$\begin{pmatrix} \langle\psi_{ik}|W|\psi_{ik}\rangle & \langle\psi_{ik}|W|\psi_{ki}\rangle \\ \langle\psi_{ki}|W|\psi_{ik}\rangle & \langle\psi_{ki}|W|\psi_{ki}\rangle \end{pmatrix} = \begin{pmatrix} \langle\psi_{ik}|W|\psi_{ik}\rangle & 1 \\ 1 & \langle\psi_{ki}|W|\psi_{ki}\rangle \end{pmatrix} \geq 0, \quad (116)$$

and obtain

$$\langle\psi_{ik}|W|\psi_{ik}\rangle\langle\psi_{ki}|W|\psi_{ki}\rangle \geq 1 \quad (117)$$

for any $i \neq k$, which completes the proof of Eq. (115).

(Case 3: S_6) In this case, we show the following inequality holds

$$\prod_{|\psi\rangle \in S_7} \langle\psi|W|\psi\rangle = \prod_{i=k} \langle iikkkik00|W|iikkkik00\rangle = \prod_k \langle kkkkkkk00|W|kkkkkkk00\rangle \geq 1. \quad (118)$$

Let $|\psi_k\rangle := |kkkkkkk00\rangle$ for $k = 0, \dots, d-1$, then $\langle\psi_k|W|\psi_{k'}\rangle = 1$ holds for $k \neq k'$ because of Eq. (59). By considering the positivity of the submatrix

$$\begin{pmatrix} \langle\psi_0|W|\psi_0\rangle & \langle\psi_0|W|\psi_1\rangle & \cdots & \langle\psi_0|W|\psi_{d-1}\rangle \\ \langle\psi_1|W|\psi_0\rangle & \langle\psi_1|W|\psi_1\rangle & \ddots & \langle\psi_1|W|\psi_{d-1}\rangle \\ \vdots & \ddots & \ddots & \vdots \\ \langle\psi_{d-1}|W|\psi_0\rangle & \langle\psi_{d-1}|W|\psi_1\rangle & \cdots & \langle\psi_{d-1}|W|\psi_{d-1}\rangle \end{pmatrix} \geq 0, \quad (119)$$

where all of the off-diagonal elements are 1, we obtain Eq. (118).

In total, we uniquely determined the diagonal elements of W .

E Evaluation of the Summation in the Proof of Lemma 1

We evaluate the summation $\sum_{J \in G_a \otimes G_b} \|\text{Tr}_{in} W J^t\|_1$ for each a, b in the following. Note that it is enough to consider only 6 cases that satisfy $a \leq b$ because of the symmetry between the two systems. In the following, when we take summation on J , it also indicates the summation on the corresponding variables in the form $|ijkl\rangle\langle i'j'k'l'|$.

(Case 1: $G_1 \otimes G_1$) In this case, the equality

$$\sum_{J \in G_1 \otimes G_1} \|\text{Tr}_{in} W J^t\|_1 = \sum_{J \in G_1 \otimes G_1} \|\langle ijkl|W\rangle\langle W|i'j'k'l'\rangle\|_1 \quad (120)$$

$$= \sum_{J \in G_1 \otimes G_1} (\|\delta_{jk}\delta_{j'k'}|il00\rangle\langle i'l'00|\|_1 + \|\delta_{il}\delta_{i'l'}|kj11\rangle\langle k'j'11|\|_1 + \|\delta_{jk}\delta_{i'l'}|il00\rangle\langle k'j'11|\|_1 + \|\delta_{il}\delta_{j'k'}|kj11\rangle\langle i'l'00|\|_1), \quad (121)$$

holds. Moreover, the summation over the 1st and 2nd terms or the 3rd and 4th terms are equal because of the symmetry, and thus we only evaluate the 1st and 3rd terms here.

For the 1st term of Eq. (121), the equality

$$\sum_{J \in G_1 \otimes G_1} \|\delta_{jk}\delta_{j'k'}|il00\rangle\langle i'l'00|\|_1 = \sum_{J \in G_1 \otimes G_1} \delta_{jk}\delta_{j'k'} \quad (122)$$

holds. In order to evaluate the summation, we consider the following two cases. If $j \neq j'$, there are $d(d-1)(d-2)(d-3) + 5d(d-1)(d-2) + 2d(d-1)$ possible choices on (i, j, i', j') and for each choice of (i, j, i', j') , there are only $(d-2)(d-3) + 5(d-2) + 2$ possible choices on (k, l, k', l') because $k = j, k' = j'$ is necessary for the non-vanishing Kronecker delta. Similarly, if $j = j'$, there are $d(d-1)(d-2)$ possible choices on (i, j, i', j') and for each choice of (i, j, i', j') , there are $(d-1)(d-2)$ possible choices on (k, l, k', l') . For the 3rd term of Eq. (121), the equality

$$\sum_{J \in G_1 \otimes G_1} \|\delta_{jk}\delta_{i'l'}|il00\rangle\langle k'j'11|\|_1 = \sum_{J \in G_1 \otimes G_1} \delta_{jk}\delta_{i'l'} \quad (123)$$

holds. Similarly, if $j \neq i'$, there are $d(d-1)(d-2)(d-3) + 5d(d-1)(d-2) + d(d-1)$ possible choices on (i, j, i', j') and for each choice of (i, j, i', j') , there are $(d-2)(d-3) + 5(d-2) + 1$ possible choices on (k, l, k', l') .

If $j = i'$, there are $d(d-1)(d-2) + d(d-1)$ possible choices on (i, j, i', j') and for each choice of (i, j, i', j') , there are $(d-1)(d-2) + (d-1)$ possible choices on (k, l, k', l') .

In total, we obtain

$$\sum_{J \in G_1 \otimes G_1} \|\text{Tr}_{in} W J^t\|_1 = 2 \sum_{J \in G_1 \otimes G_1} \delta_{jk} \delta_{j'k'} + 2 \sum_{J \in G_1 \otimes G_1} \delta_{jk} \delta_{i'l'} \quad (124)$$

$$= 2d(d-1)(2d^4 + 2d^3 - 18d^2 + 11d + 8). \quad (125)$$

(Case 2: $G_1 \otimes G_2$) In this case, the equality

$$\sum_{J \in G_1 \otimes G_2} \|\text{Tr}_{in} W J^t\|_1 = \sum_{J \in G_1} \sum_{m=0}^{d-1} \sum_{k=0}^{d-1} \|\sum_{i,j,k} \langle ijk, k+m | W \rangle \langle W | i'j'k, k+m \rangle\|_1 \quad (126)$$

$$= \sum_{J \in G_1} \sum_{m=0}^{d-1} \sum_{k=0}^{d-1} \|\sum_{i,j,k} (\delta_{jk} \delta_{j'k} |i, k+m, 00\rangle \langle i', k+m, 00| + \delta_{i,k+m} \delta_{i',k+m} |kj11\rangle \langle kj'11| + \delta_{jk} \delta_{i',k+m} |i, k+m, 00\rangle \langle kj'11| + \delta_{i,k+m} \delta_{j'k} |kj11\rangle \langle i', k+m, 00|)\|_1 \quad (127)$$

$$= \sum_{J \in G_1} \sum_{m,k} (\delta_{jk} \delta_{j'k} + \delta_{i,k+m} \delta_{i',k+m} + \delta_{jk} \delta_{i',k+m} + \delta_{i,k+m} \delta_{j'k}) \quad (128)$$

holds. The summation over the 1st and 2nd terms or the 3rd and 4th terms are equal, and thus we only evaluate the 1st and 3rd terms here.

The summation over the 1st term of Eq. (128) can be evaluated as $d^2(d-1)(d-2)$, because for each $k, m = 0, \dots, d-1$, there are $(d-1)(d-2)$ possible choices on (i, j, i', j') due to $j = j' = k$. The summation over the 3rd term of Eq. (128) can be evaluated as follows: for each $k = 0, \dots, d-1$, if $m = 0$, there are $(d-1)(d-2) + (d-1)$ possible choices on (i, j, i', j') , and if $m \neq 0$, there are $(d-2)(d-3) + 5(d-2) + 1$ possible choices on (i, j, i', j') .

In total, we obtain

$$\sum_{J \in G_1 \otimes G_2} \|\text{Tr}_{in} W J^t\|_1 = 2\{d^2(d-1)(d-2) + d[(d-1)(d-2) + (d-1)] + d(d-1)[(d-2)(d-3) + 5(d-2) + 1]\} \quad (129)$$

$$= 2d(d-1)(2d^2 - d - 4). \quad (130)$$

(Case 3: $G_2 \otimes G_2$) In this case, the summation is evaluated as

$$\sum_{J \in G_2 \otimes G_2} \|\text{Tr}_{in} W J^t\|_1 = \sum_{n,m=0}^{d-1} \sum_{i,k=0}^{d-1} (\delta_{i+n,k} \delta_{i+n,k} + \delta_{i,k+m} \delta_{i,k+m} + \delta_{i+n,k} \delta_{i,k+m} + \delta_{i,k+m} \delta_{i+n,k}) \quad (131)$$

$$= 2 \sum_{n,m=0}^{d-1} \sum_{i,k=0}^{d-1} (\delta_{i+n,k} \delta_{i+n,k} + \delta_{i+n,k} \delta_{i,k+m}) \quad (132)$$

$$= 2(d^3 + d^2) = 2d^2(d+1). \quad (133)$$

(Case 4: $G_1 \otimes G_3$) Here we consider G_3 with only the elements of the form $|kl\rangle\langle kk| - |ll\rangle\langle lk|$. The elements of the other form can be evaluated in the same way since the two input systems are symmetric. We denote this group as G'_3 , that is,

$$G'_3 = \{|kl\rangle\langle kk| - |ll\rangle\langle lk| \mid k \neq l\} \quad (134)$$

and the following equality holds

$$\sum_{J \in G_1 \otimes G_3} \|\text{Tr}_{in} W J^t\|_1 = 2 \sum_{J \in G_1 \otimes G'_3} \|\text{Tr}_{in} W J^t\|_1. \quad (135)$$

In this case, the summation can be evaluated as

$$\sum_{J \in G_1 \otimes G'_3} \|\text{Tr}_{in} W J^t\|_1 = \sum_{J \in G_1} \sum_{k \neq l} \|\langle ijk|W\rangle\langle W|i'j'kk\rangle - \langle ijll|W\rangle\langle W|i'j'lk\rangle\|_1 \quad (136)$$

$$\begin{aligned} &= \sum_{J \in G_1} \sum_{k \neq l} \|\langle \delta_{jk}\delta_{j'k}|il00\rangle\langle i'k00| - \delta_{jl}\delta_{j'l}|il00\rangle\langle i'k00| \\ &\quad + (\delta_{il}\delta_{i'k}|kj11\rangle\langle kj'11| - \delta_{il}\delta_{i'k}|lj11\rangle\langle lj'11|) \\ &\quad + (\delta_{jk}\delta_{i'k}|il00\rangle\langle kj'11| - \delta_{jl}\delta_{i'k}|il00\rangle\langle lj'11|) \\ &\quad + (\delta_{il}\delta_{j'k}|kj11\rangle\langle i'k00| - \delta_{il}\delta_{j'l}|lj11\rangle\langle i'k00|)\|_1 \end{aligned} \quad (137)$$

$$\begin{aligned} &= \sum_{J \in G_1} \sum_{k \neq l} (\|\delta_{jk}\delta_{j'k}|il00\rangle\langle i'k00| - \delta_{jl}\delta_{j'l}|il00\rangle\langle i'k00|\|_1 \\ &\quad + \|\delta_{il}\delta_{i'k}|kj11\rangle\langle kj'11| - \delta_{il}\delta_{i'k}|lj11\rangle\langle lj'11|\|_1 \\ &\quad + \|\delta_{jk}\delta_{i'k}|il00\rangle\langle kj'11| - \delta_{jl}\delta_{i'k}|il00\rangle\langle lj'11|\|_1 \\ &\quad + \|\delta_{il}\delta_{j'k}|kj11\rangle\langle i'k00| - \delta_{il}\delta_{j'l}|lj11\rangle\langle i'k00|\|_1) \end{aligned} \quad (138)$$

$$= \sum_{J \in G_1} \sum_{k \neq l} (\delta_{jk}\delta_{j'k} + \delta_{jl}\delta_{j'l} + 2\delta_{il}\delta_{i'k} + \delta_{jk}\delta_{i'k} + \delta_{jl}\delta_{i'k} + \delta_{il}\delta_{j'k} + \delta_{il}\delta_{j'l}) \quad (139)$$

$$= 2 \sum_{J \in G_1} \sum_{k \neq l} (\delta_{jk}\delta_{j'k} + \delta_{il}\delta_{i'k} + \delta_{jk}\delta_{i'k} + \delta_{jl}\delta_{i'k}), \quad (140)$$

where the third equality holds because the states of control qubits are different, the fourth equality holds because $k \neq l$, and the last equality holds because the summation over the 1st and 2nd, 4th and 7th, 5th and 6th terms are the same, respectively. For each (k, l) , by counting the number of possible choices of (i, j, i', j') , we obtain

$$\sum_{J \in G_1 \otimes G'_3} \|\text{Tr}_{in} W J^t\|_1 = 2d(d-1) \sum_{J \in G_1} (\delta_{jk}\delta_{j'k} + \delta_{il}\delta_{i'k} + \delta_{jk}\delta_{i'k} + \delta_{jl}\delta_{i'k}), \quad (141)$$

$$\begin{aligned} &= 2d(d-1) \times \{[(d-1)(d-2)] + [(d-2)(d-3) + 5(d-2) + 2] \\ &\quad + [(d-1)(d-2) + (d-1)] + [(d-2)(d-3) + 5(d-2) + 1]\} \end{aligned} \quad (142)$$

$$= 2d(d-1)(4d^2 - 5d - 2), \quad (143)$$

and thus, we obtain

$$\sum_{J \in G_1 \otimes G_3} \|\text{Tr}_{in} W J^t\|_1 = 4d(d-1)(4d^2 - 5d - 2). \quad (144)$$

(Case 5: $G_2 \otimes G_3$) As in the case 4, we consider G'_3 instead, and using the following equality:

$$\sum_{J \in G_2 \otimes G_3} \|\text{Tr}_{in} W J^t\|_1 = 2 \sum_{J \in G_2 \otimes G'_3} \|\text{Tr}_{in} W J^t\|_1. \quad (145)$$

In this case, the summation can be evaluated as

$$\begin{aligned} & \sum_{J \in G_2 \otimes G'_3} \|\text{Tr}_{in} W J^t\|_1 \\ &= \sum_{k \neq l} \sum_{m=0}^{d-1} \left\| \sum_{i=0}^{d-1} (\langle i, i+m, k | W \rangle \langle W | i, i+m, k \rangle - \langle i, i+m, l | W \rangle \langle W | i, i+m, l \rangle) \right\|_1 \end{aligned} \quad (146)$$

$$\begin{aligned} &= \sum_{k \neq l} \sum_{m=0}^{d-1} \left\| \sum_{i=0}^{d-1} \delta_{i+m, k} \delta_{i+m, k} |il00\rangle \langle ik00| - \delta_{i+m, l} \delta_{i+m, l} |il00\rangle \langle ik00| \right\|_1 \\ &+ \left\| \sum_i \delta_{il} \delta_{ik} |k, i+m, 11\rangle \langle k, i+m, 11| - \delta_{il} \delta_{ik} |l, i+m, 11\rangle \langle l, i+m, 11| \right\|_1 \\ &+ \left\| \sum_i \delta_{i+m, k} \delta_{ik} |il00\rangle \langle k, i+m, 11| - \delta_{i+m, l} \delta_{ik} |il00\rangle \langle l, i+m, 11| \right\|_1 \\ &+ \left\| \sum_i \delta_{il} \delta_{i+m, k} |k, i+m, 11\rangle \langle ik00| - \delta_{il} \delta_{i+m, l} |l, i+m, 11\rangle \langle ik00| \right\|_1 \end{aligned} \quad (147)$$

$$\begin{aligned} &= \sum_{k \neq l} \sum_{m, i=0}^{d-1} (\delta_{i+m, k} \delta_{i+m, k} + \delta_{i+m, l} \delta_{i+m, l} + \delta_{il} \delta_{ik} + \delta_{il} \delta_{ik} \\ &\quad + \delta_{i+m, k} \delta_{ik} + \delta_{i+m, l} \delta_{ik} + \delta_{il} \delta_{i+m, k} + \delta_{il} \delta_{i+m, l}) \end{aligned} \quad (148)$$

$$= d(d-1) \times (d + d + 0 + 0 + 1 + 1 + 1 + 1) \quad (149)$$

$$= 2d(d-1)(d+2). \quad (150)$$

Here, the second equality holds because the states of control qubits are different, the third equality holds because $k \neq l$, and the fourth equality holds by counting the number of possible choices of (i, j, i', j') for each (k, l) . In total, we obtain

$$\sum_{J \in G_2 \otimes G_3} \|\text{Tr}_{in} W J^t\|_1 = 4d(d-1)(d+2). \quad (151)$$

(Case 6: $G_3 \otimes G_3$) In this case, we divide the first G_3 into two groups, i.e., G'_3 and G''_3 , where G'_3 defined by Eq. (134) is the one used in cases 4 and 5, and G''_3 is defined as the remaining of G_3 , that is,

$$G''_3 = \{|ji\rangle \langle ii| - |jj\rangle \langle ij| \mid i \neq j\}. \quad (152)$$

Here we evaluate both $G'_3 \otimes G'_3$ and $G''_3 \otimes G'_3$, and the result can be obtained by

$$\sum_{J \in G_3 \otimes G_3} \|\text{Tr}_{in} W J^t\|_1 = 2 \left(\sum_{J \in G'_3 \otimes G'_3} \|\text{Tr}_{in} W J^t\|_1 + \sum_{J \in G''_3 \otimes G'_3} \|\text{Tr}_{in} W J^t\|_1 \right). \quad (153)$$

We first consider the summation over $G'_3 \otimes G'_3$, which is evaluated as

$$\begin{aligned} & \sum_{J \in G'_3 \otimes G'_3} \|\text{Tr}_{in} W J^t\|_1 \\ &= \sum_{i \neq j} \sum_{k \neq l} \left\| \langle ijk | W \rangle \langle W | iikk \rangle - \langle ijll | W \rangle \langle W | iilk \rangle \right. \\ &\quad \left. - \langle jjkl | W \rangle \langle W | jikk \rangle + \langle jjll | W \rangle \langle W | jilk \rangle \right\|_1 \end{aligned} \quad (154)$$

$$\begin{aligned} &= \sum_{i \neq j, k \neq l} \left\| (\delta_{jk} \delta_{ik} - \delta_{jl} \delta_{il}) |il00\rangle \langle ik00| - (\delta_{jk} \delta_{ik} - \delta_{jl} \delta_{il}) |jl00\rangle \langle jk00| \right\|_1 \\ &\quad + \left\| (\delta_{il} \delta_{ik} - \delta_{jl} \delta_{jk}) |kj11\rangle \langle ki11| - (\delta_{il} \delta_{ik} - \delta_{jl} \delta_{jk}) |lj11\rangle \langle li11| \right\|_1 \\ &\quad + \left\| (\delta_{jk} \delta_{ik} |il00\rangle \langle ki11| - \delta_{jl} \delta_{ik} |il00\rangle \langle li11| - \delta_{jk} \delta_{jk} |jl00\rangle \langle ki11| + \delta_{jl} \delta_{jk} |jl00\rangle \langle li11|) \right\|_1 \\ &\quad + \left\| (\delta_{il} \delta_{ik} |kj11\rangle \langle ik00| - \delta_{il} \delta_{il} |lj11\rangle \langle ik00| - \delta_{jl} \delta_{ik} |kj11\rangle \langle jk00| + \delta_{jl} \delta_{il} |lj11\rangle \langle jk00|) \right\|_1 \end{aligned} \quad (155)$$

$$= \sum_{i \neq j, k \neq l} 3\delta_{jk} \delta_{ik} + 3\delta_{jl} \delta_{il} + 3\delta_{il} \delta_{ik} + 3\delta_{jl} \delta_{jk} + 2\delta_{jl} \delta_{ik} + \delta_{jk} \delta_{jk} + \delta_{il} \delta_{il} \quad (156)$$

$$= \sum_{i \neq j, k \neq l} 2\delta_{jl} \delta_{ik} + \delta_{jk} \delta_{jk} + \delta_{il} \delta_{il} \quad (157)$$

$$= 2d(d-1) + 2d(d-1)^2 = 2d^2(d-1), \quad (158)$$

where the second equality holds because the states of control qubits are different, and the third equality holds because $i \neq j$ and $k \neq l$. Similarly, the summation over $G_3'' \otimes G_3'$ can be evaluated as

$$\begin{aligned}
& \sum_{J \in G_3'' \otimes G_3'} \|\text{Tr}_{in} W J^t\|_1 \\
&= \sum_{i \neq j, k \neq l} \|\langle jikl|W\rangle\langle W||iikk\rangle - \langle jill|W\rangle\langle W||iilk\rangle \\
&\quad - \langle jjkl|W\rangle\langle W||ijkk\rangle + \langle jjll|W\rangle\langle W||ijlk\rangle\|_1 \tag{159}
\end{aligned}$$

$$\begin{aligned}
&= \sum_{i \neq j, k \neq l} \|(\delta_{ik}\delta_{ik} - \delta_{il}\delta_{il} - \delta_{jk}\delta_{jk} + \delta_{jl}\delta_{jl})|jl00\rangle\langle ik00|\|_1 \\
&\quad + \|(\delta_{jl}\delta_{ik}|ki11\rangle\langle ki11| - \delta_{jl}\delta_{ik}|li11\rangle\langle li11| - \delta_{jl}\delta_{ik}|kj11\rangle\langle kj11| + \delta_{jl}\delta_{ik}|lj11\rangle\langle lj11|)\|_1 \\
&\quad + \|(\delta_{ik}\delta_{ik}|jl00\rangle\langle ki11| - \delta_{il}\delta_{ik}|jl00\rangle\langle li11| - \delta_{jk}\delta_{ik}|jl00\rangle\langle kj11| + \delta_{jl}\delta_{ik}|jl00\rangle\langle lj11|)\|_1 \\
&\quad + \|(\delta_{jl}\delta_{ik}|ki11\rangle\langle ik00| - \delta_{jl}\delta_{il}|li11\rangle\langle ik00| - \delta_{jl}\delta_{jk}|kj11\rangle\langle ik00| + \delta_{jl}\delta_{jl}|lj11\rangle\langle ik00|)\|_1 \tag{160}
\end{aligned}$$

$$= \sum_{i \neq j, k \neq l} 2\delta_{ik}\delta_{ik} + \delta_{il}\delta_{il} + \delta_{jk}\delta_{jk} + 2\delta_{jl}\delta_{jl} + 4\delta_{jl}\delta_{ik} + \delta_{il}\delta_{ik} + \delta_{jk}\delta_{ik} + 2\delta_{jl}\delta_{ik} + \delta_{jl}\delta_{il} + \delta_{jl}\delta_{jk} \tag{161}$$

$$= \sum_{i \neq j, k \neq l} 4\delta_{jl}\delta_{ik} + 2\delta_{ik}\delta_{ik} + 2\delta_{jl}\delta_{jl} + 2\delta_{jl}\delta_{ik} + \delta_{il}\delta_{il} + \delta_{jk}\delta_{jk} \tag{162}$$

$$= d(d-1)[4 + 2(d-1) + 2(d-1) + 2 + (d-1) + (d-1)] \tag{163}$$

$$= 6d^2(d-1) \tag{164}$$

where the second equality holds because the states of control qubits are different. For the third equality, the expansion can be done for all but the first term using $i \neq j$ and $k \neq l$. For the first term, there exist some patterns where more than one terms of the Kronecker deltas are non-zero, for example, $\delta_{ik}\delta_{ik}$ and $\delta_{jl}\delta_{jl}$ can be non-zero simultaneously. However, since they have same sign, the expansion is possible and the third equality holds.

In total, we obtain

$$\sum_{J \in G_3 \otimes G_3} \|\text{Tr}_{in} W J^t\|_1 = 16d^2(d-1). \tag{165}$$

Finally, we have evaluated all terms in Eq. (94), and we obtain

$$\|W\|_1 \geq (2d^3)^2, \tag{166}$$

which completes the proof of Lemma 1.