

# Lower Bound Estimates of the Order of Meromorphic Solutions to Non-Homogeneous Linear Differential-Difference Equations

Rachid BELLAAMA and Benharrat BELAÏDI<sup>1</sup>

Department of Mathematics  
 Laboratory of Pure and Applied Mathematics  
 University of Mostaganem (UMAB)  
 B. P. 227 Mostaganem-(Algeria)  
 rachidbellaama10@gmail.com  
 benharrat.belaidi@univ-mosta.dz

**Abstract.** In this article, we deal with the order of growth of solutions of non-homogeneous linear differential-difference equation

$$\sum_{i=0}^n \sum_{j=0}^m A_{ij} f^{(j)}(z + c_i) = F(z),$$

where  $A_{ij}$ ,  $F(z)$  are entire or meromorphic functions and  $c_i$  ( $0, 1, \dots, n$ ) are non-zero distinct complex numbers. Under the sufficient condition that there exists one coefficient having the maximal lower order or having the maximal lower type strictly greater than the order or the type of other coefficients, we obtain estimates of the lower bound of the order of meromorphic solutions of the above equation.

2010 *Mathematics Subject Classification*: 30D35, 39B32, 39A10.

*Key words*: Linear difference equation, linear differential-difference equations, Meromorphic solution, Order, Type, Lower order, Lower type.

## 1 Introduction and statement of main results

Throughout this article, a meromorphic function means a function that is meromorphic in the whole complex plane  $\mathbb{C}$ . We use the basic notations such as  $m(r, f)$ ,  $N(r, f)$ ,  $T(r, f)$  and fundamental results of Nevanlinna's

---

<sup>1</sup>Corresponding author

value distribution theory ([8, 12, 24]). Further, we denote respectively by  $\rho(f), \mu(f), \tau(f), \underline{\tau}(f)$ , the order, the lower order, the type and the lower type of a meromorphic function  $f$ , also when  $f$  is an entire function, we use the notations  $\tau_M(f), \underline{\tau}_M(f)$  respectively for the type and lower type of  $f$  (see e.g. [8, 12, 14, 24]).

Recently, a lot of results were rapidly obtained for complex difference and complex difference equations ([1 – 3, 5, 7, 10, 11, 15 – 17, 19, 25, 26]). The back-ground for these studies lies in the recent difference counterparts of Nevanlinna theory. The key result here is the difference analogue of the lemma on the logarithmic derivative obtained by Halburd-Korhonen [10, 11] and Chiang-Feng [7], independently. Several authors have investigated the properties of meromorphic solutions of complex linear difference equation

$$A_n f(z + c_n) + A_{n-1} f(z + c_{n-1}) + \cdots + A_1 f(z + c_1) + A_0 f(z) = A_{n+1}, \quad (1.1)$$

where  $A_j(z)$  ( $j = 0, \dots, n + 1$ ) are entire or meromorphic functions and  $c_i$  ( $1, \dots, n$ ) are non-zero distinct complex numbers, when one coefficient has maximal order (lower order) or among coefficients having the maximal order (lower order), exactly one has its type (lower type) strictly greater than others and achieved some important results (see e.g. [2, 7, 15, 19, 25]). Very recently in [2], the authors have studied the growth of meromorphic solutions of (1.1) when one coefficient having maximal lower order or having the maximal lower type strictly greater than the order or the type of other coefficients, we have obtained the following theorems.

**Theorem A** ([2]) *Let  $A_j(z)$  ( $j = 0, \dots, n + 1$ ) be entire functions, and let  $k, l \in \{0, 1, \dots, n + 1\}$ . If the following three assumptions hold simultaneously:*

- (1)  $\max\{\mu(A_k), \rho(A_j), j \neq k, l\} = \rho \leq \mu(A_l) < \infty, \mu(A_l) > 0$ ;
- (2)  $\underline{\tau}_M(A_l) > \underline{\tau}_M(A_k)$ , when  $\mu(A_l) = \mu(A_k)$ ;
- (3)  $\max\{\tau_M(A_j) : \rho(A_j) = \mu(A_l), j \neq k, l\} = \tau_1 < \underline{\tau}_M(A_l)$ , when  $\mu(A_l) = \max\{\rho(A_j), j \neq k, l\}$ .

*Then every meromorphic solution  $f$  of (1.1) satisfies  $\rho(f) \geq \mu(A_l)$  if  $A_{n+1} \not\equiv 0$ . Furthermore, if  $A_{n+1}(z) \equiv 0$ , then every meromorphic solution  $f \not\equiv 0$  of (1.1) satisfies  $\rho(f) \geq \mu(A_l) + 1$ .*

**Theorem B** ([2]) *Let  $A_j(z)$  ( $j = 0, \dots, n+1$ ) be meromorphic functions, and let  $k, l \in \{0, 1, \dots, n+1\}$ . If the following five assumptions hold simultaneously:*

(1)  $\max\{\mu(A_k), \rho(A_j), j \neq k, l\} = \rho \leq \mu(A_l) < \infty;$

(2)  $\tau(A_l) > \tau(A_k)$ , when  $\mu(A_l) = \mu(A_k);$

(3)

$$\tau_1 = \sum_{\rho(A_j)=\mu(A_l), j \neq l, k} \tau(A_j) < \tau(A_l) < +\infty$$

when  $\mu(A_l) = \max\{\rho(A_j), j \neq l, k\};$

(4)

$$\tau_1 + \tau(A_k) < \tau(A_l) < +\infty$$

when  $\mu(A_l) = \mu(A_k) = \max\{\rho(A_j), j \neq k, l\};$

(5)  $\lambda\left(\frac{1}{A_l}\right) < \mu(A_l) < \infty.$

*Then every meromorphic solution  $f$  of (1.1) satisfies  $\rho(f) \geq \mu(A_l)$  if  $A_{n+1} \not\equiv 0$ . Furthermore, if  $A_{n+1}(z) \equiv 0$ , then every meromorphic solution  $f \not\equiv 0$  of (1.1) satisfies  $\rho(f) \geq \mu(A_l) + 1$ .*

Historically, the study of complex differential-difference equations can be traced back to Naftalevich's research. By making use of operator theory and iteration method, Naftalevich [20] considered the meromorphic solutions on complex differential-difference equations. However, there are few investigations on complex differential-difference field using Nevanlinna theory. Recently, some papers (see, [6, 18, 21, 22, 26]) focusing on complex differential-difference emerged. In [6], Chen and Zheng have investigated the growth of solutions of the homogeneous differential-difference equation

$$\sum_{i=0}^n \sum_{j=0}^m A_{ij} f^{(j)}(z + c_i) = 0 \tag{1.2}$$

and obtained the following results.

**Theorem C** ([6]) *Let  $A_{ij}(z)$  ( $i = 0, \dots, n; j = 0, \dots, m$ ) be entire functions, and  $a, l \in \{0, 1, \dots, n\}, b \in \{0, 1, \dots, m\}$  such that  $(a, b) \neq (l, 0)$ . If the following three assumptions hold simultaneously :*

- (1)  $\max\{\mu(A_{ab}), \rho(A_{ij}), (i, j) \neq (a, b), (l, 0)\} = \rho \leq \mu(A_{l0}) < \infty, \mu(A_{l0}) > 0$ ;
- (2)  $\underline{\tau}_M(A_{l0}) > \underline{\tau}_M(A_{ab})$ , when  $\mu(A_{l0}) = \mu(A_{ab})$ ;
- (3)  $\underline{\tau}_M(A_{l0}) > \max\{\tau_M(A_{ij}) : \rho(A_{ij}) = \mu(A_{l0}), (i, j) \neq (a, b), (l, 0)\}$ , when  $\mu(A_{l0}) = \max\{\rho(A_{ij}) : (i, j) \neq (a, b), (l, 0)\}$ . Then any non zero meromorphic solution  $f$  of (1.2) satisfies  $\rho(f) \geq \mu(A_{l0}) + 1$ .

**Theorem D** ([6]) *Let  $A_{ij}(z)$  ( $i = 0, \dots, n; j = 0, \dots, m$ ) be meromorphic functions, and  $a, l \in \{0, 1, \dots, n\}, b \in \{0, 1, \dots, m\}$  such that  $(a, b) \neq (l, 0)$ . If the following four assumptions hold simultaneously:*

- (1)  $\delta(\infty, A_{l0}) = \liminf_{r \rightarrow +\infty} \frac{m(r, A_{l0})}{T(r, A_{l0})} = \delta > 0$ ;
- (2)  $\max\{\mu(A_{ab}), \rho(A_{ij}), (i, j) \neq (a, b), (l, 0)\} = \rho \leq \mu(A_{l0}) < \infty, \mu(A_{l0}) > 0$ ;
- (3)  $\delta \underline{\tau}(A_{l0}) > \underline{\tau}(A_{ab})$ , when  $\mu(A_{l0}) = \mu(A_{ab})$ ;
- (4)  $\delta \underline{\tau}(A_{l0}) > \max\{\tau(A_{ij}) : \rho(A_{ij}) = \mu(A_{l0}), (i, j) \neq (a, b), (l, 0)\}$  when  $\mu(A_{l0}) = \max\{\rho(A_{ij}) : (i, j) \neq (a, b), (l, 0)\}$ . Then any non zero meromorphic solution  $f$  of (1.2) satisfies  $\rho(f) \geq \mu(A_{l0}) + 1$ .

In this paper, by combining complex differential and difference equations, we extend the results of Theorems C and D for the complex non-homogeneous linear differential-difference equation

$$\sum_{i=0}^n \sum_{j=0}^m A_{ij} f^{(j)}(z + c_i) = F. \quad (1.3)$$

The main results of this paper state as follows.

**Theorem 1.1** *Let  $A_{ij}(z)$  ( $i = 0, \dots, n; j = 0, \dots, m$ ),  $F(z)$  be entire functions, and let  $k, p$  ( $0 \leq k \leq n, 0 \leq p \leq m$ ) be integers. If there exists an integer  $l$  ( $0 \leq l \leq n$ ), such that  $A_{l0}$  satisfies the following three assumptions simultaneously:*

- (1)  $\max\{\mu(A_{kp}), \rho(F), \rho(A_{ij}), (i, j) \neq (l, 0), (k, p)\} = \rho \leq \mu(A_{l0}) < \infty, \mu(A_{l0}) > 0$ ;
- (2)  $\underline{\tau}_M(A_{l0}) > \underline{\tau}_M(A_{kp})$ , when  $\mu(A_{l0}) = \mu(A_{kp})$ ;

$$(3) \max\{\tau_M(S) : \rho(S) = \mu(A_{l0}), S \in \{F, A_{ij} : (i, j) \neq (l, 0), (k, p)\}\} = \tau_1 < \underline{\tau}_M(A_{l0}), \text{ when } \mu(A_{l0}) = \max\{\rho(F), \rho(A_{ij}), (i, j) \neq (l, 0), (k, p)\}.$$

Then every meromorphic solution  $f$  of (1.3) satisfies  $\rho(f) \geq \mu(A_{l0})$  if  $F(z) \not\equiv 0$ . Further, if  $F(z) \equiv 0$ , then every meromorphic solution  $f \not\equiv 0$  of (1.2) satisfies  $\rho(f) \geq \mu(A_{l0}) + 1$ .

**Theorem 1.2** Let  $A_{ij}(z) (i = 0, \dots, n; j = 0, \dots, m), F(z)$  be meromorphic functions, and let  $k, p$  ( $0 \leq k \leq n, 0 \leq p \leq m$ ) be integers. If there exists an integer  $l$  ( $0 \leq l \leq n$ ), such that  $A_{l0}$  satisfies the following five assumptions simultaneously:

$$(1) \max\{\mu(A_{kp}), \rho(F), \rho(A_{ij}), (i, j) \neq (l, 0), (k, p)\} = \rho \leq \mu(A_{l0}) < \infty;$$

$$(2) \underline{\tau}(A_{l0}) > \underline{\tau}(A_{kp}), \text{ when } \mu(A_{l0}) = \mu(A_{kp});$$

$$(3) \tau_1 = \sum_{\rho(A_{ij})=\mu(A_{l0}), (i,j) \neq (l,0), (k,p)} \tau(A_{ij}) + \tau(F) < \underline{\tau}(A_{l0}) < +\infty$$

when  $\mu(A_{l0}) = \max\{\rho(F), \rho(A_{ij}), (i, j) \neq (l, 0), (k, p)\}$ ;

$$(4) \tau_1 + \underline{\tau}(A_{kp}) < \underline{\tau}(A_{l0}) < +\infty$$

when  $\mu(A_{l0}) = \mu(A_{kp}) = \max\{\rho(F), \rho(A_{ij}), (i, j) \neq (l, 0), (k, p)\}$ ;

$$(5) \lambda\left(\frac{1}{A_{l0}}\right) < \mu(A_{l0}) < \infty.$$

Then every meromorphic solution  $f$  of (1.3) satisfies  $\rho(f) \geq \mu(A_{l0})$  if  $F(z) \not\equiv 0$ . Further, if  $F(z) \equiv 0$ , then every meromorphic solution  $f \not\equiv 0$  of (1.2) satisfies  $\rho(f) \geq \mu(A_{l0}) + 1$ .

## 2 Some Preliminary Lemmas

**Lemma 2.1** ([9]) Let  $f$  be a transcendental meromorphic function of finite order  $\rho(f)$ , and let  $k$  and  $j$  be integers satisfying  $k > j \geq 0$ . Then for any

given  $\varepsilon(> 0)$ , there exists a subset  $E_1 \subset (1, +\infty)$  which has finite logarithmic measure such that for all  $z$  satisfying  $|z| = r \notin [0, 1] \cup E_1$ , we have

$$\left| \frac{f^{(k)}(z)}{f^{(j)}(z)} \right| \leq |z|^{(k-j)(\rho(f)-1+\varepsilon)}.$$

**Lemma 2.2** ([7]) *Let  $f$  be a meromorphic function of finite order  $\rho$ , and let  $c_1, c_2 (c_1 \neq c_2)$  be two arbitrary complex numbers. Let  $\varepsilon > 0$  be given, then there exists a subset  $E_2 \subset (1, +\infty)$  with finite logarithmic measure such that for all  $z$  satisfying  $|z| = r \notin [0, 1] \cup E_2$ , we have*

$$\exp\{-r^{\rho-1+\varepsilon}\} \leq \left| \frac{f(z+c_1)}{f(z+c_2)} \right| \leq \exp\{r^{\rho-1+\varepsilon}\}.$$

**Lemma 2.3** ([8]) *Let  $f$  be a meromorphic function,  $c$  be a non-zero complex constant. Then we have that for  $r \rightarrow +\infty$*

$$(1+o(1))T(r-|c|, f(z)) \leq T(r, f(z+c)) \leq (1+o(1))T(r+|c|, f(z)).$$

Consequently

$$\rho(f(z+c)) = \rho(f), \quad \mu(f(z+c)) = \mu(f).$$

**Lemma 2.4** ([4]) *Let  $f$  be a meromorphic function of finite order  $\rho$ . Then for any given  $\varepsilon > 0$ , there exists a set  $E_3 \subset (1, +\infty)$  having finite linear measure and finite logarithmic measure such that for all  $z$  satisfying  $|z| = r \notin [0, 1] \cup E_3$  and sufficiently large  $r$ , we have*

$$\exp\{-r^{\rho+\varepsilon}\} \leq |f(z)| \leq \exp\{r^{\rho+\varepsilon}\}.$$

**Lemma 2.5** ([13]) *Let  $f$  be an entire function with  $\mu(f) < \infty$ . Then for any given  $\varepsilon(> 0)$ , there exists a subset  $E_4 \subset (1, +\infty)$  with infinite logarithmic measure such that for all  $r \in E_4$ , we have*

$$\mu(f) = \lim_{\substack{r \rightarrow +\infty \\ r \in E_4}} \frac{\log \log M(r, f)}{\log r}$$

and

$$M(r, f) < \exp\{r^{\mu(f)+\varepsilon}\}.$$

**Lemma 2.6** ([23]) *Let  $f$  be an entire function with  $0 < \mu(f) < \infty$ . Then for any given  $\varepsilon(> 0)$ , there exists a subset  $E_5 \subset (1, +\infty)$  with infinite logarithmic measure such that for all  $r \in E_5$ , we have*

$$\underline{\mathcal{I}}_M(f) = \lim_{\substack{r \rightarrow +\infty \\ r \in E_5}} \frac{\log M(r, f)}{\log r}$$

and

$$M(r, f) < \exp\{(\underline{\mathcal{I}}_M(f) + \varepsilon)r^{\mu(f)}\}.$$

**Lemma 2.7** ([7]) *Let  $f$  be a meromorphic function of finite order  $\rho(f) < \infty$ , and let  $c_1, c_2$  be two distinct complex numbers. Then for each  $\varepsilon > 0$ , we have*

$$m\left(r, \frac{f(z + c_1)}{f(z + c_2)}\right) = O(r^{\rho(f)-1+\varepsilon}).$$

**Lemma 2.8** ([12]) (*Logarithmic Derivative Lemma*) *Let  $f$  be a meromorphic function and  $k \geq 1$  be an integer. Then, we have*

$$m\left(r, \frac{f^{(k)}}{f}\right) = O(\log T(r, f) + \log r)$$

possibly outside a set  $E_6 \subset [0, +\infty)$  of a finite linear measure. If  $\rho(f) < \infty$ , then

$$m\left(r, \frac{f^{(k)}}{f}\right) = O(\log r).$$

**Lemma 2.9** ([25]) *Let  $f$  be a meromorphic function with  $\mu(f) < \infty$ . Then for any given  $\varepsilon(> 0)$ , there exists a subset  $E_7 \subset (1, +\infty)$  with infinite logarithmic measure such that for all  $r \in E_7$ , we have*

$$T(r, f) < r^{\mu(f)+\varepsilon}.$$

**Lemma 2.10** ([19]) *Let  $f$  be a meromorphic function with  $0 < \mu(f) < \infty$ . Then for any given  $\varepsilon(> 0)$ , there exists a subset  $E_8 \subset (1, +\infty)$  with infinite logarithmic measure such that for all  $r \in E_8$ , we have*

$$T(r, f) < (\underline{\mathcal{I}}(f) + \varepsilon)r^{\mu(f)}.$$

### 3 Proof of main results

**Proof of Theorem 1.1** If  $f$  has infinite order, then the result holds. Now, we suppose that  $\rho(f) < \infty$ . We divide (1.3) by  $f(z + c_l)$  to get

$$\begin{aligned} -A_{l0}(z) &= \sum_{i=0, i \neq l, k}^n \sum_{j=0}^m A_{ij} \frac{f^{(j)}(z + c_i) f(z + c_i)}{f(z + c_i) f(z + c_l)} + \sum_{j=0, j \neq p}^m A_{kj} \frac{f^{(j)}(z + c_k) f(z + c_k)}{f(z + c_k) f(z + c_l)} \\ &+ A_{kp} \frac{f^{(p)}(z + c_k) f(z + c_k)}{f(z + c_k) f(z + c_l)} + \sum_{j=1}^m A_{lj} \frac{f^{(j)}(z + c_l)}{f(z + c_l)} - \frac{F(z)}{f(z + c_l)}. \end{aligned} \quad (3.1)$$

Therefore

$$\begin{aligned} |A_{l0}(z)| &\leq \sum_{i=0, i \neq l, k}^n \sum_{j=0}^m |A_{ij}| \left| \frac{f^{(j)}(z + c_i)}{f(z + c_i)} \right| \left| \frac{f(z + c_i)}{f(z + c_l)} \right| \\ &+ \sum_{j=0, j \neq p}^m |A_{kj}| \left| \frac{f^{(j)}(z + c_k)}{f(z + c_k)} \right| \left| \frac{f(z + c_k)}{f(z + c_l)} \right| \\ &+ |A_{kp}| \left| \frac{f^{(p)}(z + c_k)}{f(z + c_k)} \right| \left| \frac{f(z + c_k)}{f(z + c_l)} \right| + \sum_{j=1}^m |A_{lj}| \left| \frac{f^{(j)}(z + c_l)}{f(z + c_l)} \right| + \left| \frac{F(z)}{f(z + c_l)} \right|. \end{aligned} \quad (3.2)$$

From Lemma 2.1 and Lemma 2.3, for any given  $\varepsilon (> 0)$ , there exists a subset  $E_1 \subset (1, +\infty)$  which has finite logarithmic measure such that for all  $z$  satisfying  $|z| = r \notin [0, 1] \cup E_1$ , we have

$$\left| \frac{f^{(j)}(z + c_i)}{f(z + c_i)} \right| \leq |z|^{j(\rho(f+c_i)-1+\varepsilon)} = |z|^{j(\rho(f)-1+\varepsilon)}, \quad (i, j) \neq (l, 0). \quad (3.3)$$

It follows by Lemma 2.2 that for any  $\varepsilon (> 0)$ , there exists a subset  $E_2 \subset (1, +\infty)$  with finite logarithmic measure such that for all  $z$  satisfying  $|z| = r \notin [0, 1] \cup E_2$ , we have

$$\left| \frac{f(z + c_i)}{f(z + c_l)} \right| \leq \exp\{r^{\rho(f)-1+\varepsilon}\}, \quad i \neq l. \quad (3.4)$$

From Lemma 2.3, we get

$$\rho(f(z + c_l)) = \rho\left(\frac{1}{f(z + c_l)}\right) = \rho(f).$$



So, by Lemma 2.4, for any given  $\varepsilon > 0$ , there exists a subset  $E_3 \subset (1, +\infty)$  having finite linear measure and finite logarithmic measure such that for all  $z$  satisfying  $|z| = r \notin [0, 1] \cup E_3$  sufficiently large, we have

$$\left| \frac{1}{f(z + c_l)} \right| \leq \exp\{r^{\rho(f)+\varepsilon}\}. \quad (3.5)$$

In the following, we divide the proof into four cases:

**Case (i).** We suppose that  $\rho < \mu(A_{l_0})$ .

For  $S \in \{F, A_{ij} : (i, j) \neq (l, 0), (k, p)\}$ , by the definition of  $\rho(S)$ , for any given  $\varepsilon > 0$  and sufficiently large  $r$ , we have

$$|S(z)| \leq \exp\{r^{\rho(S)+\varepsilon}\} \leq \exp\{r^{\rho+\varepsilon}\}. \quad (3.6)$$

It follows by the definition of  $\mu(A_{l_0})$ , for sufficiently small  $\varepsilon > 0$  and sufficiently large  $r$ , we have

$$|A_{l_0}(z)| \geq \exp\{r^{\mu(A_{l_0})-\varepsilon}\}. \quad (3.7)$$

It also follows by the definition of  $\mu(A_{kp})$  and Lemma 2.5, for any given  $\varepsilon (> 0)$ , there exists a subset  $E_4 \subset (1, +\infty)$  with infinite logarithmic measure such that for all  $r \in E_4$ , we have

$$|A_{kp}(z)| \leq \exp\{r^{\mu(A_{kp})+\varepsilon}\}. \quad (3.8)$$

By substituting (3.3) – (3.8) into (3.2), for all  $z$  satisfying  $|z| = r \in E_4 \setminus ([0, 1] \cup E_1 \cup E_2 \cup E_3)$ , we obtain

$$\begin{aligned} \exp\{r^{\mu(A_{l_0})-\varepsilon}\} &\leq \sum_{i=0, i \neq l, k}^n \sum_{j=0}^m \exp\{r^{\rho+\varepsilon}\} |z|^{j(\rho(f)-1+\varepsilon)} \exp\{r^{\rho(f)-1+\varepsilon}\} \\ &\quad + \sum_{j=0, j \neq p}^m \exp\{r^{\rho+\varepsilon}\} |z|^{j(\rho(f)-1+\varepsilon)} \exp\{r^{\rho(f)-1+\varepsilon}\} \\ &\quad + |z|^{p(\rho(f)-1+\varepsilon)} \exp\{r^{\mu(A_{kp})+\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\} \\ &\quad + \sum_{j=1}^m \exp\{r^{\rho+\varepsilon}\} |z|^{j(\rho(f)-1+\varepsilon)} + \exp\{r^{\rho+\varepsilon}\} \exp\{r^{\rho(f)+\varepsilon}\}. \end{aligned}$$

$$\begin{aligned} &\leq ((n-1)(m+1) + 2m) r^{m(\rho(f)-1+\varepsilon)} \exp\{r^{\rho+\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\} \\ &+ r^{p(\rho(f)-1+\varepsilon)} \exp\{r^{\mu(A_{kp})+\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\} + \exp\{r^{\rho+\varepsilon}\} \exp\{r^{\rho(f)+\varepsilon}\}. \end{aligned} \quad (3.9)$$

Now, we may choose sufficiently small  $\varepsilon$  satisfying  $0 < 3\varepsilon < \mu(A_{l_0}) - \rho$ , we deduce from (3.9) that for  $|z| = r \in E_4 \setminus ([0, 1] \cup E_1 \cup E_2 \cup E_3)$ ,  $r \rightarrow +\infty$

$$\exp\{r^{\mu(A_{l_0})-2\varepsilon}\} \leq \exp\{r^{\rho(f)+\varepsilon}\}.$$

Therefore,  $\mu(A_{l_0}) \leq \rho(f) + 3\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0})$ .

Further, if  $F \equiv 0$ , then by substituting (3.3), (3.4) and (3.6) – (3.8) into (3.2), for all  $z$  satisfying  $|z| = r \in E_4 \setminus ([0, 1] \cup E_1 \cup E_2)$ , we obtain

$$\begin{aligned} \exp\{r^{\mu(A_{l_0})-\varepsilon}\} &\leq (nm + n + m - 1) r^{m(\rho(f)-1+\varepsilon)} \exp\{r^{\rho+\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\} \\ &+ r^{p(\rho(f)-1+\varepsilon)} \exp\{r^{\mu(A_{kp})+\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\}. \end{aligned} \quad (3.10)$$

By choosing sufficiently small  $\varepsilon$  satisfying  $0 < 3\varepsilon < \mu(A_{l_0}) - \rho$ , we deduce from (3.10) that for  $|z| = r \in E_4 \setminus ([0, 1] \cup E_1 \cup E_2)$ ,  $r \rightarrow +\infty$

$$\exp\{r^{\mu(A_{l_0})-2\varepsilon}\} \leq \exp\{r^{\rho(f)-1+\varepsilon}\},$$

that is,  $\mu(A_{l_0}) \leq \rho(f) - 1 + 3\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0}) + 1$ .

**Case (ii)** We suppose that  $\beta = \max\{\rho(F), \rho(A_{ij}), (i, j) \neq (l, 0), (k, p)\} < \mu(A_{l_0}) = \mu(A_{kp})$ ,  $\underline{\tau}_M(A_{l_0}) > \underline{\tau}_M(A_{kp})$ .

For  $S \in \{F, A_{ij} : (i, j) \neq (l, 0), (k, p)\}$ , by the definition of  $\rho(S)$ , for any given  $\varepsilon (> 0)$ , and sufficiently large  $r$ , we have

$$|S(z)| \leq \exp\{r^{\rho(S)+\varepsilon}\} \leq \exp\{r^{\beta+\varepsilon}\}. \quad (3.11)$$

From the definition of  $\underline{\tau}_M(A_{l_0})$ , for sufficiently small  $\varepsilon > 0$  and sufficiently large  $r$ , we have

$$|A_{l_0}(z)| \geq \exp\{(\underline{\tau}_M(A_{l_0}) - \varepsilon) r^{\mu(A_{l_0})}\}. \quad (3.12)$$

Also, from the definition of  $\underline{\tau}_M(A_{kp})$  and Lemma 2.6, for any given  $\varepsilon (> 0)$ , there exists a subset  $E_5 \subset (1, +\infty)$  with infinite logarithmic measure such that for all  $r \in E_5$ , we have

$$|A_{kp}(z)| \leq \exp\{(\underline{\tau}_M(A_{kp}) + \varepsilon) r^{\mu(A_{kp})}\} = \exp\{(\underline{\tau}_M(A_{kp}) + \varepsilon) r^{\mu(A_{l_0})}\}. \quad (3.13)$$

By substituting (3.3) – (3.5) and (3.11) – (3.13) into (3.2), for all  $z$  satisfying  $|z| = r \in E_5 \setminus ([0, 1] \cup E_1 \cup E_2 \cup E_3)$ , we get

$$\begin{aligned} & \exp\{(\mathcal{I}_M(A_{l_0}) - \varepsilon)r^{\mu(A_{l_0})}\} \\ & \leq (nm + n + m - 1) r^{m(\rho(f)-1+\varepsilon)} \exp\{r^{\beta+\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\} \\ & + r^{p(\rho(f)-1+\varepsilon)} \exp\{(\mathcal{I}_M(A_{kp}) + \varepsilon)r^{\mu(A_{l_0})}\} \exp\{r^{\rho(f)-1+\varepsilon}\} + \exp\{r^{\beta+\varepsilon}\} \exp\{r^{\rho(f)+\varepsilon}\}. \end{aligned} \quad (3.14)$$

Therefore, we may choose sufficiently small  $\varepsilon$ ,  $0 < 2\varepsilon < \min\{\mu(A_{l_0}) - \beta, \mathcal{I}_M(A_{l_0}) - \mathcal{I}_M(A_{kp})\}$ , then from (3.14) for  $r \in E_5 \setminus ([0, 1] \cup E_1 \cup E_2 \cup E_3)$  sufficiently large, we obtain

$$\exp\{(\mathcal{I}_M(A_{l_0}) - \mathcal{I}_M(A_{kp}) - 2\varepsilon)r^{\mu(A_{l_0})-\varepsilon}\} \leq \exp\{r^{\rho(f)+\varepsilon}\}.$$

Then,  $\mu(A_{l_0}) \leq \rho(f) + 2\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, so  $\rho(f) \geq \mu(A_{l_0})$ .

Further, if  $F \equiv 0$ , then by substituting (3.3), (3.4) and (3.11) – (3.13) into (3.2), for all  $z$  satisfying  $|z| = r \in E_5 \setminus ([0, 1] \cup E_1 \cup E_2)$ , we have

$$\begin{aligned} & \exp\{(\mathcal{I}_M(A_{l_0}) - \varepsilon)r^{\mu(A_{l_0})}\} \\ & \leq (nm + n + m - 1) r^{m(\rho(f)-1+\varepsilon)} \exp\{r^{\beta+\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\} \\ & + r^{p(\rho(f)-1+\varepsilon)} \exp\{(\mathcal{I}_M(A_{kp}) + \varepsilon)r^{\mu(A_{l_0})}\} \exp\{r^{\rho(f)-1+\varepsilon}\}. \end{aligned} \quad (3.15)$$

Now, we may choose sufficiently small  $\varepsilon$ ,  $0 < 2\varepsilon < \min\{\mu(A_{l_0}) - \beta, \mathcal{I}_M(A_{l_0}) - \mathcal{I}_M(A_{kp})\}$ , then from (3.15) for  $r \in E_5 \setminus ([0, 1] \cup E_1 \cup E_2)$  sufficiently large, we get

$$\exp\{(\mathcal{I}_M(A_{l_0}) - \mathcal{I}_M(A_{kp}) - 2\varepsilon)r^{\mu(A_{l_0})-\varepsilon}\} \leq \exp\{r^{\rho(f)-1+\varepsilon}\},$$

that is,  $\mu(A_{l_0}) \leq \rho(f) - 1 + 2\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0}) + 1$ .

**Case (iii)** When  $\mu(A_{l_0}) = \max\{\rho(F), \rho(A_{ij}), (i, j) \neq (l, 0), (k, p)\} > \mu(A_{kp})$ , and  $\max\{\tau_M(S) : \rho(S) = \mu(A_{l_0}), S \in \{F, A_{ij} : (i, j) \neq (l, 0), (k, p)\}\} = \tau_1 < \mathcal{I}_M(A_{l_0})$ .

For  $S \in \{F, A_{ij} : (i, j) \neq (l, 0), (k, p)\}$ , by the definitions of  $\rho(S)$  and  $\tau_M(S)$ , for any given  $\varepsilon > 0$  and sufficiently large  $r$ , we have

$$|S(z)| \leq \begin{cases} \exp\{r^{\rho(S)+\varepsilon}\} \leq \exp\{r^{\mu(A_{l_0})-\varepsilon}\}, & \text{if } \rho(S) < \mu(A_{l_0}), \\ \exp\{(\tau_1 + \varepsilon)r^{\mu(A_{l_0})}\}, & \text{if } \rho(S) = \mu(A_{l_0}). \end{cases} \quad (3.16)$$

Then, by substituting (3.3) – (3.5), (3.8), (3.12) and (3.16) into (3.2), for all  $z$  satisfying  $|z| = r \in E_4 \setminus ([0, 1] \cup E_1 \cup E_2 \cup E_3)$  sufficiently large, we obtain

$$\begin{aligned}
& \exp\{(\underline{\tau}_M(A_{l_0}) - \varepsilon) r^{\mu(A_{l_0})}\} \\
& \leq O(r^{m(\rho(f))-1+\varepsilon}) \exp\{(\tau_1 + \varepsilon) r^{\mu(A_{l_0})}\} \exp\{r^{\rho(f)-1+\varepsilon}\} \\
& \quad + O(r^{m(\rho(f))-1+\varepsilon}) \exp\{r^{\mu(A_{l_0})-\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\} \\
& \quad \quad + r^{p(\rho(f))-1+\varepsilon}) \exp\{r^{\mu(A_{k_p})+\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\} \\
& \quad \quad \quad + O(r^{m(\rho(f))-1+\varepsilon}) \exp\{r^{\mu(A_{l_0})-\varepsilon}\} \\
& + O(r^{m(\rho(f))-1+\varepsilon}) \exp\{(\tau_1 + \varepsilon) r^{\mu(A_{l_0})}\} + \exp\{(\tau_1 + \varepsilon) r^{\mu(A_{l_0})}\} \exp\{r^{\rho(f)+\varepsilon}\}.
\end{aligned} \tag{3.17}$$

Now, we may choose sufficiently small  $\varepsilon$  satisfying

$$0 < 2\varepsilon < \min\{\mu(A_{l_0}) - \mu(A_{k_p}), \underline{\tau}_M(A_{l_0}) - \tau_1\},$$

then from (3.17) for sufficiently large  $r \in E_4 \setminus ([0, 1] \cup E_1 \cup E_2 \cup E_3)$ , we get

$$\exp\{(\underline{\tau}_M(A_{l_0}) - \tau_1 - 2\varepsilon) r^{\mu(A_{l_0})-\varepsilon}\} \leq \exp\{r^{\rho(f)+\varepsilon}\}.$$

That means,  $\mu(A_{l_0}) \leq \rho(f) + 2\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0})$ .

Further, if  $F \equiv 0$ , then by substituting (3.3), (3.4), (3.8), (3.12) and (3.16) into (3.2), for all  $z$  satisfying  $|z| = r \in E_4 \setminus ([0, 1] \cup E_1 \cup E_2)$  sufficiently large, we have

$$\begin{aligned}
& \exp\{(\underline{\tau}_M(A_{l_0}) - \varepsilon) r^{\mu(A_{l_0})}\} \\
& \leq O(r^{m(\rho(f))-1+\varepsilon}) \exp\{(\tau_1 + \varepsilon) r^{\mu(A_{l_0})}\} \exp\{r^{\rho(f)-1+\varepsilon}\} \\
& \quad + O(r^{m(\rho(f))-1+\varepsilon}) \exp\{r^{\mu(A_{l_0})-\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\} \\
& \quad \quad + r^{p(\rho(f))-1+\varepsilon}) \exp\{r^{\mu(A_{k_p})+\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\} \\
& + O(r^{m(\rho(f))-1+\varepsilon}) \exp\{r^{\mu(A_{l_0})-\varepsilon}\} + O(r^{m(\rho(f))-1+\varepsilon}) \exp\{(\tau_1 + \varepsilon) r^{\mu(A_{l_0})}\}.
\end{aligned} \tag{3.18}$$

Now, we may choose sufficiently small  $\varepsilon$  satisfying

$$0 < 2\varepsilon < \min\{\mu(A_{l_0}) - \mu(A_{k_p}), \underline{\tau}_M(A_{l_0}) - \tau_1\},$$

then from (3.18) for sufficiently large  $r \in E_4 \setminus ([0, 1] \cup E_1 \cup E_2)$ , we get

$$\exp\{(\underline{\tau}_M(A_{l_0}) - \tau_1 - 2\varepsilon) r^{\mu(A_{l_0})-\varepsilon}\} \leq \exp\{r^{\rho(f)-1+\varepsilon}\}.$$

That means,  $\mu(A_{l_0}) \leq \rho(f) - 1 + 2\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0}) + 1$ .

**Case (iv)** We suppose that  $\max\{\rho(A_{ij}), \rho(F), (i, j) \neq (l, 0), (k, p)\} = \mu(A_{kp}) = \mu(A_{l_0})$  and  $\max\{\underline{\tau}_M(A_{kp}), \tau_M(S) : \rho(S) = \mu(A_{l_0}), S \in \{F, A_{ij} : (i, j) \neq (l, 0), (k, p)\}\} = \tau_2 < \underline{\tau}_M(A_{l_0})$ .

It follows by substituting (3.3) – (3.5), (3.12), (3.13) and (3.16) into (3.2), for all  $z$  satisfying  $|z| = r \in E_5 \setminus ([0, 1] \cup E_1 \cup E_2 \cup E_3)$  sufficiently large, we have

$$\begin{aligned}
& \exp\{(\underline{\tau}_M(A_{l_0}) - \varepsilon) r^{\mu(A_{l_0})}\} \\
& \leq O(r^{m(\rho(f)-1+\varepsilon)} \exp\{(\tau_2 + \varepsilon) r^{\mu(A_{l_0})}\} \exp\{r^{\rho(f)-1+\varepsilon}\}) \\
& \quad + O(r^{m(\rho(f)-1+\varepsilon)} \exp\{r^{\mu(A_{l_0})-\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\}) \\
& \quad + r^{p(\rho(f)-1+\varepsilon)} \exp\{(\underline{\tau}_M(A_{kp}) + \varepsilon) r^{\mu(A_{l_0})}\} \exp\{r^{\rho(f)-1+\varepsilon}\} \\
& \quad + O(r^{m(\rho(f)-1+\varepsilon)} \exp\{r^{\mu(A_{l_0})-\varepsilon}\}) + O(r^{m(\rho(f)-1+\varepsilon)} \exp\{(\tau_2 + \varepsilon) r^{\mu(A_{l_0})}\}) \\
& \quad + \exp\{(\tau_2 + \varepsilon) r^{\mu(A_{l_0})}\} \exp\{r^{\rho(f)+\varepsilon}\}. \tag{3.19}
\end{aligned}$$

Now, we may choose sufficiently small  $\varepsilon$  satisfying

$$0 < 2\varepsilon < \underline{\tau}_M(A_{l_0}) - \tau_2,$$

from (3.19) for sufficiently large  $r \in E_5 \setminus ([0, 1] \cup E_1 \cup E_2 \cup E_3)$ , we get

$$\exp\{(\underline{\tau}_M(A_{l_0}) - \tau_2 - 2\varepsilon) r^{\mu(A_{l_0})-\varepsilon}\} \leq \exp\{r^{\rho(f)+\varepsilon}\},$$

that means,  $\mu(A_{l_0}) \leq \rho(f) + 2\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0})$ .

Further, if  $F \equiv 0$ , by substituting (3.3), (3.4), (3.12), (3.13) and (3.16) into (3.2), for all  $z$  satisfying  $|z| = r \in E_5 \setminus ([0, 1] \cup E_1 \cup E_2)$  sufficiently large, we have

$$\begin{aligned}
\exp\{(\underline{\tau}_M(A_{l_0}) - \varepsilon) r^{\mu(A_{l_0})}\} & \leq O(r^{m(\rho(f)-1+\varepsilon)} \exp\{(\tau_2 + \varepsilon) r^{\mu(A_{l_0})}\} \exp\{r^{\rho(f)-1+\varepsilon}\}) \\
& \quad + O(r^{m(\rho(f)-1+\varepsilon)} \exp\{r^{\mu(A_{l_0})-\varepsilon}\} \exp\{r^{\rho(f)-1+\varepsilon}\}) \\
& \quad + r^{p(\rho(f)-1+\varepsilon)} \exp\{(\underline{\tau}_M(A_{kp}) + \varepsilon) r^{\mu(A_{l_0})}\} \exp\{r^{\rho(f)-1+\varepsilon}\}
\end{aligned}$$

$$+ O\left(r^{m(\rho(f)-1+\varepsilon)} \exp\{r^{\mu(A_{l_0})-\varepsilon}\}\right) + O\left(r^{m(\rho(f)-1+\varepsilon)} \exp\{(\tau_2 + \varepsilon)r^{\mu(A_{l_0})}\}\right). \quad (3.20)$$

Now, we may choose sufficiently small  $\varepsilon$  satisfying

$$0 < 2\varepsilon < \underline{\tau}_M(A_{l_0}) - \tau_2,$$

from (3.20) for sufficiently large  $r \in E_5 \setminus ([0, 1] \cup E_1 \cup E_2)$ , we get

$$\exp\{(\underline{\tau}_M(A_{l_0}) - \tau_2 - 2\varepsilon)r^{\mu(A_{l_0})-\varepsilon}\} \leq \exp\{r^{\rho(f)-1+\varepsilon}\}.$$

That means,  $\mu(A_{l_0}) \leq \rho(f) - 1 + 2\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0}) + 1$ . The proof of Theorem 1.1 is complete.

**Proof of Theorem 1.2** If  $f$  has infinite order, then the result holds. Now, we suppose that  $\rho(f) < \infty$ . By (3.1), we have

$$\begin{aligned} T(r, A_{l_0}(z)) &= m(r, A_{l_0}(z)) + N(r, A_{l_0}(z)) \\ &\leq \sum_{i=0, i \neq l, k}^n \sum_{j=0}^m m(r, A_{ij}(z)) + m(r, A_{kp}(z)) \\ &+ \sum_{j=0, j \neq p}^m m(r, A_{kj}(z)) + \sum_{j=1}^m m(r, A_{lj}(z)) + \sum_{i=0, i \neq l, k}^n \sum_{j=0}^m m\left(r, \frac{f^{(j)}(z + c_i)}{f(z + c_i)}\right) \\ &+ \sum_{i=0, i \neq l, k}^n m\left(r, \frac{f(z + c_i)}{f(z + c_l)}\right) + \sum_{j=1}^m m\left(r, \frac{f^{(j)}(z + c_k)}{f(z + c_k)}\right) + 2m\left(r, \frac{f(z + c_k)}{f(z + c_l)}\right) \\ &+ \sum_{j=1}^m m\left(r, \frac{f^{(j)}(z + c_l)}{f(z + c_l)}\right) + m(r, F(z)) + m\left(r, \frac{1}{f(z + c_l)}\right) \\ &+ N(r, A_{l_0}(z)) + O(1) \leq \sum_{i=0, i \neq l, k}^n \sum_{j=0}^m T(r, A_{ij}(z)) + T(r, A_{kp}(z)) \\ &+ \sum_{j=0, j \neq p}^m T(r, A_{kj}(z)) + \sum_{j=1}^m T(r, A_{lj}(z)) + \sum_{i=0, i \neq l, k}^n \sum_{j=1}^m m\left(r, \frac{f^{(j)}(z + c_i)}{f(z + c_i)}\right) \\ &+ \sum_{i=0, i \neq l, k}^n m\left(r, \frac{f(z + c_i)}{f(z + c_l)}\right) + \sum_{j=1}^m m\left(r, \frac{f^{(j)}(z + c_k)}{f(z + c_k)}\right) + 2m\left(r, \frac{f(z + c_k)}{f(z + c_l)}\right) \end{aligned}$$

$$+ \sum_{j=1}^m m \left( r, \frac{f^{(j)}(z + c_l)}{f(z + c_l)} \right) + T(r, F(z)) + T \left( r, \frac{1}{f(z + c_l)} \right) + N(r, A_{l_0}(z)) + O(1).$$

By Lemma 2.3 and the first main theorem of Nevanlinna, when  $r$  sufficiently large, we have

$$T \left( r, \frac{1}{f(z + c_l)} \right) = T(r, f(z + c_l)) + O(1) \leq (1 + o(1))T(r + |c_l|, f) \leq 2T(2r, f).$$

So, for  $r$  sufficiently large we obtain

$$\begin{aligned} T(r, A_{l_0}(z)) &\leq \sum_{i=0, i \neq l, k}^n \sum_{j=0}^m T(r, A_{ij}(z)) + T(r, A_{kp}(z)) \\ &\quad + \sum_{j=0, j \neq p}^m T(r, A_{kj}(z)) + \sum_{j=1}^m T(r, A_{lj}(z)) \\ &+ \sum_{i=0, i \neq l, k}^n \sum_{j=1}^m m \left( r, \frac{f^{(j)}(z + c_i)}{f(z + c_i)} \right) + \sum_{i=0, i \neq l, k}^n m \left( r, \frac{f(z + c_i)}{f(z + c_l)} \right) + T(r, F(z)) \\ &\quad + 2T(2r, f) + \sum_{j=1}^m m \left( r, \frac{f^{(j)}(z + c_l)}{f(z + c_l)} \right) \\ &+ \sum_{j=1}^m m \left( r, \frac{f^{(j)}(z + c_k)}{f(z + c_k)} \right) + 2m \left( r, \frac{f(z + c_k)}{f(z + c_l)} \right) + N(r, A_{l_0}(z)) + O(1). \end{aligned} \quad (3.21)$$

By Lemma 2.7, for any given  $\varepsilon (> 0)$ , we have

$$m \left( r, \frac{f(z)}{f(z + c_l)} \right) = O(r^{\rho(f)-1+\varepsilon}), \quad m \left( r, \frac{f(z + c_j)}{f(z + c_l)} \right) = O(r^{\rho(f)-1+\varepsilon}), \quad j \neq l. \quad (3.22)$$

It follows by Lemma 2.8, there exists a subset  $E_6 \subset [0, +\infty[$  of a finite linear measure such that for all  $r \notin E_6$  sufficiently large, we have

$$m \left( r, \frac{f^{(j)}(z + c_i)}{f(z + c_i)} \right) = O(\log r) \quad (i = 0, \dots, n; j = 1, \dots, m). \quad (3.23)$$

From the definition of  $\lambda \left( \frac{1}{A_{l_0}} \right)$ , for any given  $\varepsilon > 0$  and sufficiently large  $r$ , we have

$$N(r, A_{l_0}) \leq r^{\lambda \left( \frac{1}{A_{l_0}} \right) + \varepsilon}. \quad (3.24)$$

In the following, we divide the proof into four cases:

**Case (i)** We suppose that  $\rho < \mu(A_{l0})$ .

For  $S \in \{F, A_{ij} : (i, j) \neq (l, 0), (k, p)\}$ , from the definition of  $\rho(S)$  and  $\rho(f)$  for any given  $\varepsilon > 0$  and sufficiently large  $r$ , we have

$$T(r, S) \leq r^{\rho(S)+\varepsilon} \leq r^{\rho+\varepsilon}, \quad (3.25)$$

$$T(r, f) \leq r^{\rho(f)+\varepsilon}. \quad (3.26)$$

It follows from the definition of  $\mu(A_{l0})$ , for sufficiently small  $\varepsilon > 0$  and sufficiently large  $r$ , we have

$$T(r, A_{l0}) \geq r^{\mu(A_{l0})-\varepsilon}. \quad (3.27)$$

It follows from the definition of  $\mu(A_{kp})$  and Lemma 2.9, for any given  $\varepsilon (> 0)$ , there exists a subset  $E_7 \subset (1, +\infty)$  with infinite logarithmic measure such that for all  $r \in E_7$ , we have

$$T(r, A_{kp}) \leq r^{\mu(A_{kp})+\varepsilon}. \quad (3.28)$$

By substituting (3.22) – (3.28) into (3.21) for sufficiently large  $r \in E_7 \setminus E_6$ , we obtain

$$\begin{aligned} r^{\mu(A_{l0})-\varepsilon} &\leq ((n-1)(m+1) + 2m) r^{\rho+\varepsilon} + r^{\mu(A_{kp})+\varepsilon} + O(r^{\rho(f)-1+\varepsilon}) \\ &\quad + 2(2r)^{\rho(f)+\varepsilon} + r^{\rho+\varepsilon} + r^{\lambda\left(\frac{1}{A_{l0}}\right)+\varepsilon} + O(\log r). \end{aligned} \quad (3.29)$$

We may choose sufficiently small  $\varepsilon$  satisfying

$$0 < 3\varepsilon < \min \left\{ \mu(A_{l0}) - \rho, \mu(A_{l0}) - \lambda \left( \frac{1}{A_{l0}} \right) \right\},$$

it follows from (3.29) that for  $r \in E_7 \setminus E_6$ ,  $r \rightarrow +\infty$

$$r^{\mu(A_{l0})-2\varepsilon} \leq r^{\rho(f)+\varepsilon},$$

that means,  $\mu(A_{l0}) \leq \rho(f) + 3\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l0})$ . Further, if  $F \equiv 0$ , then by substituting (3.22) – (3.25), (3.27) and (3.28) into (3.21) for sufficiently large  $r \in E_7 \setminus E_6$ , we obtain

$$r^{\mu(A_{l0})-\varepsilon} \leq ((n-1)(m+1) + 2m) r^{\rho+\varepsilon} + r^{\mu(A_{kp})+\varepsilon} + O(r^{\rho(f)-1+\varepsilon})$$



$$+ r^{\lambda\left(\frac{1}{A_{l0}}\right)+\varepsilon} + O(\log r). \quad (3.30)$$

We may choose sufficiently small  $\varepsilon$  satisfying

$$0 < 3\varepsilon < \min \left\{ \mu(A_{l0}) - \rho, \mu(A_{l0}) - \lambda \left( \frac{1}{A_{l0}} \right) \right\},$$

from (3.30) that for  $r \in E_7 \setminus E_6$ ,  $r \rightarrow +\infty$

$$r^{\mu(A_{l0})-2\varepsilon} \leq r^{\rho(f)-1+\varepsilon},$$

that means,  $\mu(A_{l0}) \leq \rho(f) - 1 + 3\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l0}) + 1$ .

**Case (ii)** We suppose that  $\max\{\rho(A_{ij}), \rho(F), (i, j) \neq (l, 0), (k, p)\} = \beta < \mu(A_{l0}) = \mu(A_{kp})$ ,  $\underline{\tau}(A_{l0}) > \underline{\tau}(A_{kp})$ .

For  $S \in \{F, A_{ij} : (i, j) \neq (l, 0), (k, p)\}$ , by the definition of  $\rho(S)$ , for any given  $\varepsilon (> 0)$  and sufficiently large  $r$ , we obtain

$$T(r, S) \leq r^{\rho(S)+\varepsilon} \leq r^{\beta+\varepsilon}. \quad (3.31)$$

From the definition of  $\underline{\tau}(A_{l0})$ , for sufficiently small  $\varepsilon > 0$  and sufficiently large  $r$ , we have

$$T(r, A_{l0}) \geq (\underline{\tau}(A_{l0}) - \varepsilon) r^{\mu(A_{l0})}. \quad (3.32)$$

It follows from the definition of  $\underline{\tau}(A_{kp})$  and Lemma 2.10, for any given  $\varepsilon (> 0)$ , there exists a subset  $E_8 \subset (1, +\infty)$  with infinite logarithmic measure such that for all  $r \in E_8$ , we have

$$T(r, A_{kp}) \leq (\underline{\tau}(A_{kp}) + \varepsilon) r^{\mu(A_{kp})} \leq (\underline{\tau}(A_{kp}) + \varepsilon) r^{\mu(A_{l0})}. \quad (3.33)$$

By substituting (3.22) – (3.24), (3.26) and (3.31) – (3.33) into (3.21), for sufficiently large  $r \in E_8 \setminus E_6$ , we obtain

$$\begin{aligned} (\underline{\tau}(A_{l0}) - \varepsilon) r^{\mu(A_{l0})} &\leq ((n-1)(m+1) + 2m) r^{\beta+\varepsilon} + (\underline{\tau}(A_{kp}) + \varepsilon) r^{\mu(A_{l0})} \\ &+ O(r^{\rho(f)-1+\varepsilon}) + 2(2r)^{\rho(f)+\varepsilon} + r^{\beta+\varepsilon} + r^{\lambda\left(\frac{1}{A_{l0}}\right)+\varepsilon} + O(\log r). \end{aligned} \quad (3.34)$$

Now, we may choose sufficiently small  $\varepsilon$  satisfying

$$0 < 2\varepsilon < \min \left\{ \mu(A_{l0}) - \beta, \underline{\tau}(A_{l0}) - \underline{\tau}(A_{kp}), \mu(A_{l0}) - \lambda \left( \frac{1}{A_{l0}} \right) \right\},$$

so from (3.34) for sufficiently large  $r \in E_8 \setminus E_6$ , we have

$$(\underline{\tau}(A_{l_0}) - \underline{\tau}(A_{kp}) - 2\varepsilon)r^{\mu(A_{l_0})-\varepsilon} \leq r^{\rho(f)+\varepsilon},$$

that means,  $\mu(A_{l_0}) \leq \rho(f) + 2\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0})$ . Further, if  $F \equiv 0$ , then by substituting (3.22) – (3.24) and (3.31) – (3.33) into (3.21), for sufficiently large  $r \in E_8 \setminus E_6$ , we obtain

$$\begin{aligned} (\underline{\tau}(A_{l_0}) - \varepsilon) r^{\mu(A_{l_0})} &\leq ((n-1)(m+1) + 2m) r^{\beta+\varepsilon} + (\underline{\tau}(A_{kp}) + \varepsilon) r^{\mu(A_{l_0})} \\ &+ O(r^{\rho(f)-1+\varepsilon}) + r^{\lambda\left(\frac{1}{A_{l_0}}\right)+\varepsilon} + O(\log r). \end{aligned} \quad (3.35)$$

Now, we may choose sufficiently small  $\varepsilon$  satisfying

$$0 < 2\varepsilon < \min \left\{ \mu(A_{l_0}) - \beta, \underline{\tau}(A_{l_0}) - \underline{\tau}(A_{kp}), \mu(A_{l_0}) - \lambda \left( \frac{1}{A_{l_0}} \right) \right\}.$$

From (3.35) for sufficiently large  $r \in E_8 \setminus E_6$ , we get

$$(\underline{\tau}(A_{l_0}) - \underline{\tau}(A_{kp}) - 2\varepsilon)r^{\mu(A_{l_0})-\varepsilon} \leq r^{\rho(f)-1+\varepsilon},$$

that means,  $\mu(A_{l_0}) \leq \rho(f) - 1 + 2\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0}) + 1$ .

**Case (iii)** We suppose that  $\mu(A_{l_0}) = \max\{\rho(F), \rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\} > \mu(A_{kp})$ , and  $\tau_1 = \sum_{\rho(A_{ij})=\mu(A_{l_0}), (i,j) \neq (l,0), (k,p)} \tau(A_{ij}) + \tau(F) < \underline{\tau}(A_{l_0})$ . Then, there exists a subset  $J \subseteq \{0, 1, \dots, n\} \times \{0, 1, \dots, m\} \setminus \{(l, 0), (k, p)\}$  such that for all  $(i, j) \in J$ , when  $\rho(A_{ij}) = \mu(A_{l_0})$ , we have  $\sum_{(i,j) \in J} \tau(A_{ij}) < \underline{\tau}(A_{l_0}) - \tau(F)$ , and for  $(i, j) \in \Pi = \{0, 1, \dots, n\} \times \{0, 1, \dots, m\} \setminus (J \cup \{(l, 0), (k, p)\})$  we have  $\rho(A_{ij}) < \mu(A_{l_0})$ . Hence, for any given  $\varepsilon > 0$  and sufficiently large  $r$ , we get

$$T(r, A_{ij}) \leq \begin{cases} (\tau(A_{ij}) + \varepsilon) r^{\mu(A_{l_0})}, & \text{if } (i, j) \in J, \\ r^{\rho(A_{ij})+\varepsilon} \leq r^{\mu(A_{l_0})-\varepsilon}, & \text{if } (i, j) \in \Pi \end{cases} \quad (3.36)$$

and

$$T(r, F) \leq \begin{cases} (\tau(F) + \varepsilon) r^{\mu(A_{l_0})}, & \text{if } \rho(F) = \mu(A_{l_0}), \\ r^{\rho(F)+\varepsilon} \leq r^{\mu(A_{l_0})-\varepsilon}, & \text{if } \rho(F) < \mu(A_{l_0}). \end{cases} \quad (3.37)$$

Then, by substituting (3.22)–(3.24), (3.26), (3.28), (3.32), (3.36) and (3.37) into (3.21), for all  $z$  satisfying  $|z| = r \in E_7 \setminus E_6$  sufficiently large  $r$ , we obtain

$$(\underline{\tau}(A_{l_0}) - \varepsilon) r^{\mu(A_{l_0})} \leq \sum_{(i,j) \in J} (\tau(A_{ij}) + \varepsilon) r^{\mu(A_{l_0})} + \sum_{(i,j) \in \Pi} r^{\mu(A_{l_0})-\varepsilon} + r^{\mu(A_{kp})+\varepsilon}$$

$$\begin{aligned}
& + (\tau(F) + \varepsilon) r^{\mu(A_{l_0})} + r^{\lambda\left(\frac{1}{A_{l_0}}\right)+\varepsilon} + 2(2r)^{\rho(f)+\varepsilon} + O(r^{\rho(f)-1+\varepsilon}) + O(\ln r) \\
& \leq (\tau_1 + (nm + n + m)\varepsilon) r^{\mu(A_{l_0})} + O(r^{\mu(A_{l_0})-\varepsilon}) + r^{\mu(A_{kp})+\varepsilon} \\
& \quad + r^{\lambda\left(\frac{1}{A_{l_0}}\right)+\varepsilon} + 2(2r)^{\rho(f)+\varepsilon} + O(r^{\rho(f)-1+\varepsilon}) + O(\log r). \tag{3.38}
\end{aligned}$$

Now, we choose sufficiently small  $\varepsilon$  satisfying

$$0 < \varepsilon < \min \left\{ \frac{\mu(A_{l_0}) - \mu(A_{kp})}{2}, \frac{\underline{\tau}(A_{l_0}) - \tau_1}{nm + n + m + 1}, \frac{\mu(A_{l_0}) - \lambda\left(\frac{1}{A_{l_0}}\right)}{2} \right\},$$

then from (3.38) for sufficiently large  $r \in E_7 \setminus E_6$ , we get

$$(\underline{\tau}(A_{l_0}) - \tau_1 - (nm + n + m + 1)\varepsilon) r^{\mu(A_{l_0})-\varepsilon} \leq r^{\rho(f)+\varepsilon},$$

that means,  $\mu(A_{l_0}) \leq \rho(f) + 2\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0})$ .

Further, if  $F \equiv 0$ , then by substituting (3.22) – (3.24), (3.28), (3.32) and (3.36) into (3.21), for all  $z$  satisfying  $|z| = r \in E_7 \setminus E_6$  sufficiently large  $r$ , we obtain

$$\begin{aligned}
(\underline{\tau}(A_{l_0}) - \varepsilon) r^{\mu(A_{l_0})} & \leq (\tau_1 + (nm + n + m - 1)\varepsilon) r^{\mu(A_{l_0})} + O(r^{\mu(A_{l_0})-\varepsilon}) \\
& \quad + r^{\mu(A_{kp})+\varepsilon} + r^{\lambda\left(\frac{1}{A_{l_0}}\right)+\varepsilon} + O(r^{\rho(f)-1+\varepsilon}) + O(\log r). \tag{3.39}
\end{aligned}$$

Now, we choose sufficiently small  $\varepsilon$  satisfying

$$0 < \varepsilon < \min \left\{ \frac{\mu(A_{l_0}) - \mu(A_{kp})}{2}, \frac{\underline{\tau}(A_{l_0}) - \tau_1}{nm + n + m}, \frac{\mu(A_{l_0}) - \lambda\left(\frac{1}{A_{l_0}}\right)}{2} \right\},$$

then from (3.39) for sufficiently large  $r \in E_7 \setminus E_6$ , we get

$$(\underline{\tau}(A_{l_0}) - \tau_1 - (nm + n + m)\varepsilon) r^{\mu(A_{l_0})-\varepsilon} \leq r^{\rho(f)-1+\varepsilon},$$

that means,  $\mu(A_{l_0}) \leq \rho(f) - 1 + 2\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0}) + 1$ .

**Case (iv)** We suppose that  $\max\{\rho(A_j), \rho(F), (i, j) \neq (l, 0), (k, p)\} = \mu(A_{l_0}) = \mu(A_{kp})$  with  $\tau_1 + \underline{\tau}(A_{kp}) < \underline{\tau}(A_{l_0})$ .

It follows by substituting (3.22) – (3.24), (3.26), (3.32), (3.33), (3.36) and (3.37) into (3.21), for all sufficiently large  $r \in E_8 \setminus E_6$ , we have

$$\begin{aligned} & (\underline{\tau}(A_{l_0}) - \varepsilon) r^{\mu(A_{l_0})} \leq (\tau_1 + (nm + n + m)\varepsilon) r^{\mu(A_{l_0})} + O(r^{\mu(A_{l_0}) - \varepsilon}) \\ & + (\underline{\tau}(A_{kp}) + \varepsilon) r^{\mu(A_{l_0})} + r^{\lambda\left(\frac{1}{A_{l_0}}\right) + \varepsilon} + 2(2r)^{\rho(f) + \varepsilon} + O(r^{\rho(f) - 1 + \varepsilon}) + O(\log r). \end{aligned} \quad (3.40)$$

Now, we choose sufficiently small  $\varepsilon$  satisfying

$$0 < \varepsilon < \min \left\{ \frac{\underline{\tau}(A_{l_0}) - \tau_1 - \underline{\tau}(A_{kp})}{nm + n + m + 2}, \frac{\mu(A_{l_0}) - \lambda\left(\frac{1}{A_{l_0}}\right)}{2} \right\},$$

then from (3.40) for sufficiently large  $r \in E_8 \setminus E_6$ , we get

$$(\underline{\tau}(A_{l_0}) - \tau_1 - \underline{\tau}(A_{kp}) - (nm + n + m + 2)\varepsilon) r^{\mu(A_{l_0}) - \varepsilon} \leq r^{\rho(f) + \varepsilon},$$

that means,  $\mu(A_{l_0}) \leq \rho(f) + 2\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0})$ .

Further, if  $F \equiv 0$ , then by It follows by substituting (3.22) – (3.24), (3.32), (3.33) and (3.36), for all sufficiently large  $r \in E_8 \setminus E_6$ , we have

$$\begin{aligned} & (\underline{\tau}(A_{l_0}) - \varepsilon) r^{\mu(A_{l_0})} \leq (\tau_1 + (nm + n + m - 1)\varepsilon) r^{\mu(A_{l_0})} + O(r^{\mu(A_{l_0}) - \varepsilon}) \\ & + (\underline{\tau}(A_{kp}) + \varepsilon) r^{\mu(A_{l_0})} + r^{\lambda\left(\frac{1}{A_{l_0}}\right) + \varepsilon} + O(r^{\rho(f) - 1 + \varepsilon}) + O(\log r). \end{aligned} \quad (3.41)$$

Now, we choose sufficiently small  $\varepsilon$  satisfying

$$0 < \varepsilon < \min \left\{ \frac{\underline{\tau}(A_{l_0}) - \tau_1 - \underline{\tau}(A_{kp})}{nm + n + m + 1}, \frac{\mu(A_{l_0}) - \lambda\left(\frac{1}{A_{l_0}}\right)}{2} \right\},$$

then from (3.41) for sufficiently large  $r \in E_8 \setminus E_6$ , we get

$$(\underline{\tau}(A_{l_0}) - \tau_1 - \underline{\tau}(A_{kp}) - (nm + n + m + 1)\varepsilon) r^{\mu(A_{l_0}) - \varepsilon} \leq r^{\rho(f) - 1 + \varepsilon},$$

so that means,  $\mu(A_{l_0}) \leq \rho(f) - 1 + 2\varepsilon$ , since  $\varepsilon > 0$  is arbitrary, then  $\rho(f) \geq \mu(A_{l_0}) + 1$ . The proof of Theorem 1.2 is complete.

## 4 Examples

**Example 4.1** Consider the non-homogeneous differential-difference equation with entire coefficients

$$A_{02}(z)f''(z) + A_{11}(z)f'(z+1) + A_{01}(z)f'(z) + A_{10}(z)f(z+1) + A_{00}(z)f(z) = F(z). \quad (4.1)$$

**Case 1.**  $\max\{\rho(F), \mu(A_{kp}), \rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\} < \mu(A_{l0})$ . In (4.1) for

$$A_{00}(z) = \pi^2 + 2\pi^4 z^2, \quad A_{10}(z) = e^{-\pi^2 z^2 - \pi^2}, \quad A_{01}(z) = 2\pi^2(z+1)e^{2\pi^2 z + \pi^2},$$

$$A_{11}(z) = -2\pi^2 z, \quad A_{02}(z) = -\frac{1}{2}, \quad F(z) = e^{2\pi^2 z},$$

we have

$$\max\{\rho(F), \mu(A_{11}), \rho(A_{ij}) : (i, j) \neq (1, 0), (1, 1)\} = 1 < \mu(A_{10}) = 2.$$

We see that the conditions of Theorem 1.1 are verified. The function

$$f(z) = e^{\pi^2 z^2}$$

is a solution of equation (4.1) which satisfies  $\rho(f) = 2 \geq \mu(A_{10}) = 2$ .

**Case 2.**  $\max\{\rho(F), \rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\} < \mu(A_{l0}) = \mu(A_{kp})$  with  $\underline{\tau}_M(A_{l0}) > \underline{\tau}_M(A_{kp})$ . In (4.1) for

$$A_{00}(z) = 2\pi^2, \quad A_{10}(z) = 2\pi^2(z+1)e^{z^2} + e^{-\pi^2 z^2 - \pi^2}, \quad A_{01}(z) = 2\pi^2 z,$$

$$A_{11}(z) = -e^{z^2}, \quad A_{02}(z) = -1, \quad F(z) = e^{2\pi^2 z},$$

we get  $\max\{\rho(F), \rho(A_{ij}) : (i, j) \neq (1, 0), (1, 1)\} = 1 < \mu(A_{10}) = \mu(A_{11}) = 2$  and  $\underline{\tau}_M(A_{10}) = \pi^2 > \underline{\tau}_M(A_{11}) = 1$ . Hence, the conditions of Theorem 1.1 are satisfied. The function

$$f(z) = e^{\pi^2 z^2}$$

is a solution of equation (4.1) and  $f$  satisfies  $\rho(f) = 2 \geq \mu(A_{10}) = 2$ .

**Case 3.**  $\mu(A_{l0}) = \max\{\rho(F), \rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\} > \mu(A_{kp})$  with  $\underline{\tau}_M(A_{l0}) > \tau_1 = \max\{\tau_M(S) : \rho(S) = \mu(A_{l0}), S \in \{F, A_{ij} : (i, j) \neq (l, 0), (k, p)\}\}$ . In (4.1) for

$$A_{00}(z) = \pi^2 + 2\pi^4 z^2, \quad A_{10}(z) = e^{-\frac{4}{5}\pi^2 z^2 - \pi^2}, \quad A_{01}(z) = 2\pi^2(z+1)e^{2\pi^2 z + \pi^2},$$

$$A_{11}(z) = -2\pi^2 z, \quad A_{02}(z) = -\frac{1}{2}, \quad F(z) = e^{\frac{1}{5}\pi^2 z^2 + 2\pi^2 z},$$

we have  $\mu(A_{10}) = \max\{\rho(F), \rho(A_{ij}) : (i, j) \neq (1, 0), (0, 1)\} = 2 > \mu(A_{01}) = 1$  and  $\underline{\tau}_M(A_{10}) = \frac{4\pi^2}{5} > \tau_1 = \tau_M(F) = \frac{\pi^2}{5}$ . Obviously, the conditions of Theorem 1.1 are verified. The function

$$f(z) = e^{\pi^2 z^2}$$

is a solution of equation (4.1) which satisfies  $\rho(f) = 2 \geq \mu(A_{10}) = 2$ .

**Case 4.**  $\mu(A_{l0}) = \mu(A_{kp}) = \max\{\rho(F), \rho(A_{ij}) : (i, j) \neq (l, 0), (k, 0)\}$ , and  $\underline{\tau}_M(A_{l0}) > \max\{\tau_1, \underline{\tau}_M(A_{kp})\}$ . In (4.1) for

$$A_{00}(z) = 2\pi^2, \quad A_{10}(z) = 2\pi^2(z+1)e^{z^2} + e^{-\frac{4}{5}\pi^2 z^2 - \pi^2}, \quad A_{01}(z) = 2\pi^2 z,$$

$$A_{11}(z) = -e^{z^2}, \quad A_{02}(z) = -1, \quad F(z) = e^{\frac{1}{5}\pi^2 z^2 + 2\pi^2 z},$$

we get  $\mu(A_{10}) = \mu(A_{11}) = \max\{\rho(F), \rho(A_{ij}) : (i, j) \neq (1, 0), (1, 1)\} = 2$  and  $\underline{\tau}_M(A_{10}) = \frac{4\pi^2}{5} > \max\{\tau_1, \underline{\tau}_M(A_{11})\} = \max\{\tau_M(F), \underline{\tau}_M(A_{11})\} = \frac{\pi^2}{5}$ . We see that the conditions of Theorem 1.1 are satisfied. The function

$$f(z) = e^{\pi^2 z^2}$$

is a solution of equation (4.1) which satisfies  $\rho(f) = 2 \geq \mu(A_{10}) = 2$ .

**Example 4.2** Consider the homogeneous differential-difference equation with entire coefficients

$$A_{11}(z)g'(z-1) + A_{20}(z)g(z+3) + A_{00}(z)g(z) = 0. \quad (4.2)$$

**Case 1.**  $\max\{\mu(A_{kp}), \rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\} < \mu(A_{l0})$ . In (4.2), for

$$A_{00}(z) = 1, \quad A_{20}(z) = (4\pi i(1-z) - e^{4\pi iz})e^{-16\pi iz}, \quad A_{11}(z) = 1,$$

we have

$$\max\{\mu(A_{11}), \rho(A_{ij}) : (i, j) \neq (2, 0), (1, 1)\} = 0 < \mu(A_{20}) = 1.$$

So, the conditions of Theorem 1.1 are satisfied. The meromorphic function

$$g(z) = e^{2\pi iz^2}$$

is a solution of equation (4.2) and  $g$  satisfies  $\rho(g) = 2 \geq \mu(A_{20}) + 1 = 2$ .

**Case 2.**  $\max\{\rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\} < \mu(A_{l0}) = \mu(A_{kp})$ , with  $\underline{\tau}_M(A_{l0}) > \underline{\tau}_M(A_{kp})$ . In (4.2), for

$$A_{00}(z) = 1, \quad A_{20}(z) = (4\pi i(1-z) - e^{2\pi iz}) e^{-14\pi iz}, \quad A_{11}(z) = e^{2\pi iz},$$

we get  $\mu(A_{20}) = \mu(A_{11}) = 1 > \max\{\rho(A_{ij}) : (i, j) \neq (2, 0), (1, 1)\} = 0$  and  $\underline{\tau}_M(A_{20}) = 14\pi > \underline{\tau}_M(A_{11}) = 2\pi$ . Obviously, the conditions of Theorem 1.1 are verified. The meromorphic function

$$g(z) = e^{2\pi iz^2}$$

is a solution of equation (4.2) and  $g$  satisfies  $\rho(g) = 2 \geq \mu(A_{20}) + 1 = 2$ .

**Case 3.**  $\mu(A_{l0}) = \mu(A_{kp}) = \max\{\rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\}$  with  $\underline{\tau}_M(A_{l0}) > \tau_1 = \max\{\tau_M(A_{ij}) : \rho(A_{ij}) = \mu(A_{l0}), (i, j) \neq (l, 0), (k, p)\}$ . In (4.2), for

$$A_{00}(z) = e^{-2\pi iz}, \quad A_{20}(z) = (4\pi i(1-z) - 1) e^{-14\pi iz}, \quad A_{11}(z) = e^{2\pi iz},$$

we have  $\mu(A_{20}) = \mu(A_{11}) = \max\{\rho(A_{ij}) : (i, j) \neq (2, 0), (1, 1)\} = 1$  and  $\underline{\tau}_M(A_{20}) = 14\pi > \max\{\tau_M(A_{00}), \underline{\tau}_M(A_{11})\} = 2\pi$ . It is clear that the conditions of Theorem 1.1 are satisfied. The meromorphic function

$$g(z) = e^{2\pi iz^2}$$

is a solution of equation (4.2) and  $g$  satisfies  $\rho(g) = 2 \geq \mu(A_{20}) + 1 = 2$ .

**Example 4.3** Consider the non-homogeneous differential-difference equation with meromorphic coefficients

$$A_{11}(z)f'(z-1) + A_{01}(z)f'(z) + A_{20}(z)f(z+1) + A_{10}(z)f(z-1) = F(z). \quad (4.3)$$

**Case 1.**  $\max\{\rho(F), \mu(A_{kp}), \rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\} < \mu(A_{l0})$ . In (4.3) for

$$A_{10}(z) = e^{-\pi^3 z^3 + 3\pi^3 z^2 - 3\pi^3 z + \pi^3}, \quad A_{20}(z) = 3\pi^3 (2z - 1) e^{-3\pi^3 z^2 - 3\pi^3 z - \pi^3},$$

$$A_{01}(z) = -1, \quad A_{11}(z) = e^{3\pi^3 z^2 - 3\pi^3 z + \pi^3}, \quad F(z) = \tan(\pi z),$$

we have

$$\max\{\rho(F), \mu(A_{11}), \rho(A_{ij}) : (i, j) \neq (1, 0), (1, 1)\} = 2 < \mu(A_{10}) = 3,$$

$$\lambda\left(\frac{1}{A_{10}}\right) = 0 < \mu(A_{10}) = 3.$$

It is easy to see that the conditions of Theorem 1.2 are verified. The function

$$f(z) = e^{\pi^3 z^3} \tan(\pi z)$$

is a solution of equation (4.3) which satisfies  $\rho(f) = 3 \geq \mu(A_{10}) = 3$ .

**Case 2.**  $\max\{\rho(F), \rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\} < \mu(A_{10}) = \mu(A_{kp})$  with  $\underline{\tau}(A_{10}) > \underline{\tau}(A_{kp})$ . In (4.3) for

$$\begin{aligned} A_{10}(z) &= e^{-\pi^3 z^3 + 3\pi^3 z^2 - 3\pi^3 z + \pi^3} \\ &\quad + \left(3\pi^3(2z - 1) - 3z^2\pi^3 - \pi \tan(\pi z) + \frac{\pi}{\tan(\pi z)}\right) e^{-z^3}, \\ A_{20}(z) &= 3\pi^3 z^2 + \pi \tan(\pi z) + \frac{\pi}{\tan(\pi z)}, \quad A_{01}(z) = -e^{3\pi^3 z^2 + 3\pi^3 z + \pi^3}, \\ A_{11}(z) &= e^{-z^3}, \quad F(z) = \tan(\pi z), \end{aligned}$$

we get

$$\max\{\rho(F), \rho(A_{ij}) : (i, j) \neq (1, 0), (1, 1)\} = 2 < \mu(A_{10}) = \mu(A_{11}) = 3,$$

$$\lambda\left(\frac{1}{A_{10}}\right) = 1 < \mu(A_{10}) = 3$$

and

$$\underline{\tau}(A_{10}) = \pi^2 > \underline{\tau}(A_{11}) = \frac{1}{\pi}.$$

Hence, the conditions of Theorem 1.2 are satisfied. The function

$$f(z) = e^{\pi^3 z^3}$$

is a solution of equation (4.3) and  $f$  satisfies  $\rho(f) = 3 \geq \mu(A_{10}) = 3$ .



**Case 3.**  $\mu(A_{l0}) = \max\{\rho(F), \rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\} > \mu(A_{kp})$  with  $\underline{\tau}(A_{l0}) > \tau_1 = \sum_{\rho(A_{ij})=\mu(A_{l0}), (i,j)\neq(l,0),(k,p)} \tau(A_{ij}) + \tau(F)$ . In (4.3) for

$$A_{10}(z) = e^{-2\pi^3 z^3 + 3\pi^3 z^2 - 3\pi^3 z + \pi^3}, \quad A_{20}(z) = 3\pi^3 (2z - 1) e^{-3\pi^3 z^2 - 3\pi^3 z - \pi^3},$$

$$A_{01}(z) = -1, \quad A_{11}(z) = e^{3\pi^3 z^2 - 3\pi^3 z + \pi^3}, \quad F(z) = \frac{\tan(\pi z)}{e^{\pi^3 z^3}},$$

we have

$$\mu(A_{10}) = \max\{\rho(F), \rho(A_{ij}) : (i, j) \neq (1, 0), (1, 1)\} = 3 > \mu(A_{11}) = 2,$$

$$\lambda\left(\frac{1}{A_{10}}\right) = 0 < \mu(A_{10}) = 3$$

and  $\underline{\tau}(A_{10}) = 2\pi^2 > \tau_1 = \tau(F) = \pi^2$ . We can see that the conditions of Theorem 1.2 are verified. The function

$$f(z) = e^{\pi^3 z^3} \tan(\pi z)$$

is a solution of equation (4.3) which satisfies  $\rho(f) = 3 \geq \mu(A_{10}) = 3$ .

**Case 4.**  $\mu(A_{l0}) = \mu(A_{k0}) = \max\{\rho(F), \rho(A_{ij}), : (i, j) \neq (l, 0), (k, 0)\}$ , and  $\underline{\tau}(A_{l0}) > \tau_1 + \underline{\tau}(A_{kp})$ . In (4.3) for

$$A_{10}(z) = e^{-2\pi^3 z^3 + 3\pi^3 z^2 - 3\pi^3 z + \pi^3}, \quad A_{20}(z) = 3\pi^3 (2z - 1) e^{\left(\frac{\pi}{4}z\right)^3 - 3\pi^3 z^2 - 3\pi^3 z - \pi^3},$$

$$A_{01}(z) = -e^{\left(\frac{\pi}{4}z\right)^3}, \quad A_{11}(z) = e^{\left(\frac{\pi}{4}z\right)^3 + 3\pi^3 z^2 - 3\pi^3 z + \pi^3}, \quad F(z) = \frac{\tan(\pi z)}{e^{\pi^3 z^3}},$$

we get

$$\mu(A_{10}) = \mu(A_{11}) = \max\{\rho(F), \rho(A_{ij}) : (i, j) \neq (1, 0), (1, 1)\} = 3,$$

$$\lambda\left(\frac{1}{A_{10}}\right) = 0 < \mu(A_{10}) = 3$$

and  $\tau_1 + \underline{\tau}(A_{11}) = \tau(A_{01}) + \tau(A_{20}) + \tau(F) + \underline{\tau}(A_{11}) = \left(\frac{2}{43} + 1\right)\pi^2 + \frac{\pi^2}{43} = \frac{67}{64}\pi^2 < \underline{\tau}(A_{10}) = 2\pi^2$ . Obviously, the conditions of Theorem 1.2 are satisfied. The function

$$f(z) = e^{\pi^3 z^3} \tan(\pi z)$$

is a solution of equation (4.3) which satisfies  $\rho(f) = 3 \geq \mu(A_{10}) = 3$ .

**Example 4.4** Consider the homogeneous (differential)-difference equation with meromorphic coefficients

$$A_{11}(z)h'(z + i\pi) + A_{20}(z)h(z + 2i\pi) + A_{00}(z)h(z) = 0. \quad (4.4)$$

**Case 1.**  $\max\{\mu(A_{kp}), \rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\} < \mu(A_{l0})$ . In (4.4), for

$$A_{00}(z) = -1, \quad A_{20}(z) = e^{12\pi z^2 + 24\pi^2 iz - 16\pi^3} - e^{6\pi z^2 + 18\pi^2 iz - 14\pi^3},$$

$$A_{11}(z) = \frac{\cos(2iz)}{6i(z + i\pi)^2 \cos(2iz) + 2i \sin(2iz)},$$

we have  $\max\{\mu(A_{11}), \rho(A_{ij}) : (i, j) \neq (2, 0), (1, 1)\} = 1 < \mu(A_{20}) = 2$  and

$$\lambda\left(\frac{1}{A_{20}}\right) = 0 < \mu(A_{20}) = 2.$$

Obviously, the conditions of Theorem 1.2 are verified. The meromorphic function

$$h(z) = \frac{e^{2iz^3}}{\cos(2iz)}$$

is a solution of equation (4.4) and  $h$  satisfies  $\rho(h) = 3 \geq \mu(A_{20}) + 1 = 3$ .

**Case 2.**  $\max\{\rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\} < \mu(A_{l0}) = \mu(A_{kp})$  with  $\underline{\tau}(A_{l0}) > \underline{\tau}(A_{kp})$ . In (4.4), for

$$A_{00}(z) = 1, \quad A_{20}(z) = -2e^{12\pi z^2 + 24\pi^2 iz - 16\pi^3},$$

$$A_{11}(z) = \frac{e^{6\pi z^2 + 6\pi^2 iz - 2\pi^3} \cos(2iz)}{6i(z + i\pi)^2 \cos(2iz) + 2i \sin(2iz)},$$

we get

$$\mu(A_{20}) = \mu(A_{11}) = 2 > \max\{\rho(A_{ij}) : (i, j) \neq (2, 0), (1, 1)\} = \rho(A_{00}) = 0,$$

$$\lambda\left(\frac{1}{A_{20}}\right) = 0 < \mu(A_{20}) = 2$$

and

$$\underline{\tau}(A_{20}) = 12 > \underline{\tau}(A_{11}) = 6.$$

It is clear that the conditions of Theorem 1.2 are satisfied. The meromorphic function

$$h(z) = \frac{e^{2iz^3}}{\cos(2iz)}$$

is a solution of equation (4.4) and  $h$  satisfies  $\rho(h) = 3 \geq \mu(A_{20}) + 1 = 3$ .

**Case 3.**  $\mu(A_{l0}) = \mu(A_{kp}) = \max\{\rho(A_{ij}) : (i, j) \neq (l, 0), (k, p)\}$ , with  $\tau(A_{l0}) > \sum_{\rho(A_{ij})=\mu(A_{l0}), (i,j)\neq(l,0),(k,p)} \tau(A_{ij}) + \tau(A_{kp})$ . In (4.4), for

$$A_{00}(z) = e^{\pi z^2}, \quad A_{20}(z) = -2e^{13\pi z^2 + 24\pi^2 iz - 16\pi^3},$$

$$A_{11}(z) = \frac{e^{7\pi z^2 + 6\pi^2 iz - 2\pi^3} \cos(2iz)}{6i(z + i\pi)^2 \cos(2iz) + 2i \sin(2iz)},$$

we have

$$\mu(A_{20}) = \mu(A_{11}) = \max\{\rho(A_{ij}) : (i, j) \neq (2, 0), (1, 1)\} = \rho(A_{00}) = 2,$$

$$\lambda\left(\frac{1}{A_{20}}\right) = 0 < \mu(A_{20}) = 2$$

and

$$\tau(A_{20}) = 13 > \tau(A_{00}) + \tau(A_{11}) = 1 + 7 = 8.$$

It is easy to see that the conditions of Theorem 1.2 are verified. The meromorphic function

$$h(z) = \frac{e^{2iz^3}}{\cos(2iz)}$$

is a solution of equation (4.4) and  $h$  satisfies  $\rho(h) = 3 \geq \mu(A_{20}) + 1 = 3$ .

**Acknowledgements.** This paper was supported by the Directorate-General for Scientific Research and Technological Development (DGRSDT).

## References

- [1] B. Belaïdi and Y. Benkarouba, *Some properties of meromorphic solutions of higher order linear difference equations*. Ser. A: Appl. Math. Inform. and Mech. vol. 11, 2 (2019), 75–95.
- [2] B. Belaïdi and R. Bellaama, *Meromorphic solutions of higher order non-homogeneous linear difference equations*. Bulletin of the Transilvania University of Brasov, Series III: Mathematics, Informatics, Physics. Vol. 13, Issue 2 (2020), 433–450.

- [3] B. Belaïdi and R. Bellaama, *Study of the growth properties of meromorphic solutions of higher-order linear difference equations*. Arab. J. Math. (2021). <https://doi.org/10.1007/s40065-021-00324-2>
- [4] Z. X. Chen, *The zero, pole and orders of meromorphic solutions of differential equations with meromorphic coefficients*. Kodai Math. J. 19 (1996), no. 3, 341–354.
- [5] Z. X. Chen, *Complex differences and difference equations*, Mathematics Monograph Series 29. Science Press, Beijing (2014).
- [6] Z. Chen and X. M. Zheng, *Growth of meromorphic solutions of general complex linear differential-difference equation*. Acta Univ. Apulensis Math. Inform. No. 56 (2018), 1–12.
- [7] Y. M. Chiang and S. J. Feng, *On the Nevanlinna characteristic of  $f(z + \eta)$  and difference equations in the complex plane*. Ramanujan J. 16 (2008), no. 1, 105–129.
- [8] A. Goldberg and I. Ostrovskii, *Value distribution of meromorphic functions*. Transl. Math. Monogr., vol. 236, Amer. Math. Soc., Providence RI, 2008.
- [9] G. G. Gundersen, *Estimates for the logarithmic derivative of a meromorphic function, plus similar estimates*. J. London Math. Soc. (2) 37 (1988), no. 1, 88–104.
- [10] R. G. Halburd and R. J. Korhonen, *Difference analogue of the lemma on the logarithmic derivative with applications to difference equations*. J. Math. Anal. Appl. 314 (2006), no. 2, 477–487.
- [11] R. G. Halburd and R. J. Korhonen, *Nevanlinna theory for the difference operator*. Ann. Acad. Sci. Fenn. Math. 31 (2006), no. 2, 463–478.
- [12] W. K. Hayman, *Meromorphic functions*. Oxford Mathematical Monographs Clarendon Press, Oxford 1964.
- [13] H. Hu and X. M. Zheng, *Growth of solutions to linear differential equations with entire coefficients*. Electron. J. Differential Equations 2012, No. 226, 15 pp.
- [14] I. Laine, *Nevanlinna theory and complex differential equations*. de Gruyter Studies in Mathematics, 15. Walter de Gruyter & Co., Berlin, 1993.
- [15] I. Laine and C. C. Yang, *Clunie theorems for difference and  $q$ -difference polynomials*. J. Lond. Math. Soc. (2) 76 (2007), no. 3, 556–566.
- [16] Z. Latreuch and B. Belaïdi, *Growth and oscillation of meromorphic solutions of linear difference equations*. Mat. Vesnik 66 (2014), no. 2, 213–222.
- [17] H. F. Liu and Z. Q. Mao, *On the meromorphic solutions of some linear difference equations*. Adv. Difference Equ. 2013, 2013:133, 1–12.

- [18] K. Liu and C. J. Song, *Meromorphic solutions of complex differential-difference equations*. Results Math. 72 (2017), no. 4, 1759–1771.
- [19] I. Q. Luo and X. M. Zheng, *Growth of meromorphic solutions of some kind of complex linear difference equation with entire or meromorphic coefficients*. Math. Appl. (Wuhan) 29 (2016), no. 4, 723–730.
- [20] A. G. Naftalevič, *Meromorphic solutions of a differential-difference equation*. (Russian) Uspehi Mat. Nauk 16 1961 no. 3 (99), 191–196.
- [21] X. G. Qi and L. Z. Yang, *A note on meromorphic solutions of complex differential-difference equations*. Bull. Korean Math. Soc. 56 (2019), no. 3, 597–607.
- [22] S. Z. Wu and X. M. Zheng, *Growth of meromorphic solutions of complex linear differential-difference equations with coefficients having the same order*. J. Math. Res. Appl. 34 (2014), no. 6, 683–695.
- [23] S. Z. Wu and X. M. Zheng, *Growth of solutions to some higher-order linear differential equations in  $\mathbb{C}$  and in unit disc  $\Delta$* . (Chinese) Math. Appl. (Wuhan) 29 (2016), no. 1, 20–30.
- [24] C. C. Yang and H. X. Yi, *Uniqueness theory of meromorphic functions*. Mathematics and its Applications, 557. Kluwer Academic Publishers Group, Dordrecht, 2003.
- [25] X. M. Zheng and J. Tu, *Growth of meromorphic solutions of linear difference equations*. J. Math. Anal. Appl. 384 (2011), no. 2, 349–356.
- [26] Y. P. Zhou and X. M. Zheng, *Growth of meromorphic solutions to homogeneous and non-homogeneous linear (differential-)difference equations with meromorphic coefficients*. Electron. J. Differential Equations 2017, Paper No. 34, 15 pp.