

Variational principles for asymptotic variance of general Markov processes

Lu-Jing Huang Yong-Hua Mao Tao Wang*

Abstract

A variational formula for the asymptotic variance of general Markov processes is obtained. As application, we get a upper bound of the mean exit time of reversible Markov processes, and some comparison theorems between the reversible and non-reversible diffusion processes.

Keywords: Markov process, asymptotic variance, variational formula, the mean exit time, comparison theorem, semi-Dirichlet form

Mathematics subject classification(2020): 60J25, 60J46, 60J60

1 Introduction and main results

Asymptotic variance is a popular criterion to evaluate the performance of Markov processes, and widely used in Markov chain Monte Carlo(see e.g. [1, 5, 19, 20, 24]).

There are numerous studies of the asymptotic variance in the literature. For reversible Markov processes, the asymptotic variance can be presented by a spectral calculation, which brings a lot of applications (see [7, 14, 23] etc.). The comparisons on efficiency of reversible Markov processes, in terms of the asymptotic variance, has been extensively researched(see e.g. [1, 11, 17, 20, 27]). Recently, there are also some comparison results between reversible and non-reversible Markov processes, see e.g. [2, 6, 12, 26] for discrete-time Markov chains, and [8, 13, 22] for diffusions. However, the study of the asymptotic variance of non-reversible Markov processes is still a challenge since the lack of spectral theory of non-symmetric operators. Very recently, [11] gives some variational formulas for the asymptotic variance of general discrete-time Markov chains by solving Poisson equation, and obtains some estimates and comparison results of the asymptotic variance.

In this paper we extend the results in [11] to the general Markov process by constructing the weak solution of Poisson equation with the help of the semi-Dirichlet form.

Let $X = \{X_t\}_{t \geq 0}$ be a positive recurrent (or ergodic) Markov process on a Polish space (S, \mathcal{S}) , with strongly continuous contraction transition semigroup $\{P_t\}_{t \geq 0}$ and stationary distribution π . Denote $L^2(\pi)$ by the space of square integrable functions with scalar product $(u, v) := \int_S u(x)v(x)\pi(dx)$ and norm $\|u\| = (u, u)^{1/2}$. Let $L_0^2(\pi)$ be the subspace of functions in $L^2(\pi)$ with mean-zero, i.e.

$$L_0^2(\pi) = \{u \in L^2(\pi) : \pi(u) := \int_S u(x)\pi(dx) = 0\}.$$

*Corresponding author: wang_tao@mail.bnu.edu.cn

Denote $(L, \mathcal{D}(L))$ by the infinitesimal generator induced by $\{P_t\}_{t \geq 0}$ in $L^2(\pi)$.

Define a bilinear form associated with L as

$$\mathcal{E}(u, v) = (-Lu, v), \quad u, v \in \mathcal{D}(L),$$

and for $\alpha \geq 0$,

$$\mathcal{E}_\alpha(u, v) := \mathcal{E}(u, v) + \alpha(u, v), \quad u, v \in \mathcal{D}(L).$$

Since π is the stationary distribution and P_t is L^2 -contractive, for any $u \in \mathcal{D}(L)$,

$$\mathcal{E}(u, u) = -\pi(uLu) = \lim_{t \rightarrow 0} \frac{\pi(u^2) - \pi((P_t u)^2)}{t} \geq 0,$$

that is, \mathcal{E} is non-negative definite.

$(L, \mathcal{D}(L))$ is said to satisfy the sector condition if there exists a constant $K > 0$ such that

$$|\mathcal{E}(u, v)| \leq K \mathcal{E}(u, u)^{1/2} \mathcal{E}(v, v)^{1/2}, \quad \text{for } u, v \in \mathcal{D}(L). \quad (1.1)$$

Remarkably, if process X is reversible:

$$\pi(dx)P_t(x, dy) = \pi(dy)P_t(y, dx), \quad \text{for all } t \geq 0, \pi\text{-a.s. } x, y \in S,$$

then the sector condition is always true with $K = 1$ by Cauchy-Schwartz inequality.

Under the sector condition, we can obtain a unique semi-Dirichlet form $(\mathcal{E}, \mathcal{F})$, where \mathcal{F} is the completion of $\mathcal{D}(L)$ with respect to $\overline{\mathcal{E}}_1^{1/2}$ ($\overline{\mathcal{E}}_1$ is the symmetric part of \mathcal{E}_1), see [18, Chapter 1, Theorem 2.15]) for more details.

We say that the semigroup $\{P_t\}_{t \geq 0}$ is L^2 -exponentially ergodic, if there exist constants $C, \lambda_1 > 0$ such that for $u \in L_0^2(\pi)$,

$$\|P_t u\| \leq C \|u\| e^{-\lambda_1 t}.$$

It is well known that when process X is reversible, C can be chosen as 1 and (the optimal) λ_1 is nothing but the spectral gap:

$$\lambda_1 = \inf\{\mathcal{E}(u, u) : u \in \mathcal{F}, \pi(u) = 0 \text{ and } \pi(u^2) = 1\}. \quad (1.2)$$

Now for $f \in L_0^2(\pi)$, we consider the following asymptotic variance for X and f :

$$\sigma^2(X, f) = \limsup_{t \rightarrow \infty} \mathbb{E}_\pi \left[\left(\frac{1}{\sqrt{t}} \int_0^t f(X_s) ds \right)^2 \right]. \quad (1.3)$$

Under the L^2 -exponential ergodicity and the sector condition, our first main result presents a variational formula for the asymptotic variance as follows.

Theorem 1.1. *Suppose that the semigroup $\{P_t\}_{t \geq 0}$ associated with process X is L^2 -exponentially ergodic. Then the limit in (1.3) exists and is finite for $f \in L_0^2(\pi)$. If in addition the associated semi-Dirichlet form $(\mathcal{E}, \mathcal{F})$ satisfies the sector condition, then for $f \in L_0^2(\pi)$,*

$$2/\sigma^2(X, f) = \inf_{u \in \mathcal{M}_{f,1}} \sup_{v \in \mathcal{M}_{f,0}} \mathcal{E}(u + v, u - v), \quad (1.4)$$

where $\mathcal{M}_{f,\delta} = \{u \in \mathcal{F} : (u, f) = \delta\}$, $\delta = 0, 1$.

Particularly, if process X is reversible, then (1.4) is reduced to

$$2/\sigma^2(X, f) = \inf_{u \in \mathcal{M}_{f,1}} \mathcal{E}(u, u). \quad (1.5)$$

Remark 1.2. (1) For fixed $f \in L_0^2(\pi)$, from the proof below we will see that functions $Gf := \int_0^\infty P_t f dt$ and $G^*f := \int_0^\infty P_t^* f dt$ are both in \mathcal{F} , here P_t^* is the dual operator of P_t in $L^2(\pi)$. This is a main reason that we need the semi-Dirichlet form $(\mathcal{E}, \mathcal{F})$ in (1.4). However, if the generator L is bounded in $L^2(\pi)$, then $\mathcal{D}(L) = L^2(\pi)$, so that $Gf, G^*f \in \mathcal{D}(L)$. In this case,

$$2/\sigma^2(X, f) = \inf_{\substack{u \in L^2(\pi), \\ \pi(uf)=1}} \sup_{\substack{v \in L^2(\pi), \\ \pi(vf)=0}} ((-L)(u+v), u-v).$$

The proof of this result can be obtained immediately by replacing $P - I$ by L in the proof of [11, Theorem 1.1].

- (2) The assumption of the L^2 -exponential ergodicity of $\{P_t\}_{t \geq 0}$ is not too strong for non-reversible Markov processes, since [9] gives a geometrically ergodic Markov chain such that the asymptotic variance is infinite for some $f \in L_0^2(\pi)$.
- (3) Variational formula for the asymptotic variance has been studied in [15, Chapter 4]. It is based on a variational formula for positive definite operators in analysis and resolvent equations. Here we obtain a new variational formula.

As a direct application of Theorem 1.1, bound of the mean exit time of the process is obtained. For that, let $\Omega \subset S$ be an open set, denote by $\tau_\Omega = \inf\{t \geq 0 : X_t \notin \Omega\}$ the first exit time from Ω of process X .

Corollary 1.3. *Suppose that process X is reversible with L^2 -exponentially ergodic semigroup $\{P_t\}_{t \geq 0}$ and stationary distribution π . Let $\Omega \subset S$ be an open set with $\pi(\Omega) \in (0, 1)$, then*

$$\mathbb{E}_\pi \tau_\Omega \leq \frac{\pi(\Omega)}{2\lambda_1 \pi(\Omega^c)},$$

where λ_1 is the spectral gap defined in (1.2).

Note that in [10, Remark 3.6(1)], we gave another upper bound for the mean exit time. Explicitly, $\mathbb{E}_\pi \tau_\Omega \leq 1/(\lambda_1 \pi(\Omega^c))$ for open set $\Omega \subset S$ satisfying $\pi(\Omega^c) > 0$. It is obvious that the upper bound in Corollary 1.3 is more precise than that.

For the reversible case, similar to [11, Theorem 1.3], we could derive variational formula (1.5) without the assumption of the L^2 -exponential ergodicity. Since the proof is quite similar, we omit it in this paper.

Theorem 1.4. *Suppose that X is a reversible ergodic Markov process with stationary distribution π . Then for fixed $f \in L_0^2(\pi)$,*

$$2/\sigma^2(X, f) = \inf_{u \in \mathcal{M}_{f,1}} \mathcal{E}(u, u).$$

Note that in Theorem 1.4, maybe $\sigma^2(X, f) = \infty$ for some $f \in L_0^2(\pi)$.

The remaining part of this paper is organized as follows. In Section 2 we apply our main result in two situations. The first application is extending the comparison result for the asymptotic variance of one dimensional diffusions in [25, Theorem 1] to multi-dimensional reversible diffusions. We note that [25, Theorem 1] is proved by discrete approximation which is different from our idea, and the less assumptions are requested

in our proof. Another application is a comparison result between reversible and non-reversible diffusions on Riemannian Manifolds, which shows the asymptotic variance of a non-reversible diffusion is smaller. The similar result can be found in [8, 13] (for example, [13] proves a similar result on compact manifolds by using a spectral theorem), we provide a complete different proof by the new variational formula. Finally, the proofs of Theorem 1.1 and Corollary 1.3 are given in Section 3.

2 Applications

2.1 Reversible diffusions

First, we recall the comparison theorem proved in [25, Theorem 1]. Fix a C^1 probability density function $\mu : [I_1, I_2] \rightarrow (0, \infty)$, where $-\infty \leq I_1 < I_2 \leq \infty$. Given a C^1 positive function η on $[I_1, I_2]$ and consider a one-dimensional Langevin diffusion:

$$dX_t^\eta = \eta(X_t^\eta) dB_t + \left(\frac{1}{2} \eta^2(X_t^\eta) \log \mu'(X_t^\eta) + \eta(X_t^\eta) \eta'(X_t^\eta) \right) dt.$$

Under some additional conditions (see [25, Page 133]), [25] proves that for any $f \in L_0^2(\mu)$, and two C^1 positive functions η, η_1 on $[I_1, I_2]$ such that $\eta_1(x) \leq \eta(x)$ for all $x \in [I_1, I_2]$,

$$\sigma^2(X^m, f) \geq \sigma^2(X^\eta, f).$$

Note that in [25], the above conclusion is proved by discrete approximation. In fact, we can obtain the above result by a direct calculation as follows. For convenience, we only consider the case on half-line.

Fix a C^1 probability density function $\pi : [0, \infty) \rightarrow (0, \infty)$. Given a C^1 positive function a on $[0, \infty)$ and consider a one-dimensional diffusion X^a with reflecting boundary 0 and generator:

$$L_a = a(x) \frac{d^2}{dx^2} + b(x) \frac{d}{dx}, \quad (2.1)$$

where $b(x) = a(x)(\pi'(x)/\pi(x)) + a'(x)$. Let $\pi(dx) = \pi(x)dx$. It is easy to see that L_a is symmetric on $L^2(\pi)$. Choose a point $x_0 > 0$ and set

$$c(x) = \int_{x_0}^x \frac{b(y)}{a(y)} dy \quad \text{and} \quad \varphi(x) = \int_0^x e^{-c(y)} dy.$$

So we have

$$\pi(x) = e^{c(x)} \pi(x_0) a(x_0) / a(x). \quad (2.2)$$

Assume that X^a is non-explosive, that is,

$$\int_0^\infty \varphi'(y) \pi([0, y]) dy = \infty,$$

then X is ergodic with stationary distribution $\pi(dx)$ (see e.g. [4, Table 5.1]).

For fixed function $f \in L_0^2(\pi)$, consider Poisson equation $-L_a u = f$. By some direct calculations and (2.2), the equation has strong solution

$$u(x) = \int_0^x e^{-c(y)} \left(\int_y^\infty f(z) \frac{e^{c(z)}}{a(z)} dz \right) dy = \frac{1}{\pi(x_0) a(x_0)} \int_0^\infty f(z) \varphi(x \wedge z) \pi(dz).$$

Since $\sigma^2(X, f) = 2(u, f)$ by Lemma 3.1 and (3.3) below, from $\pi(f) = 0$ and the integration by parts we have that

$$\begin{aligned}
\frac{1}{2}\sigma^2(X^a, f) &= \frac{1}{a(x_0)\pi(x_0)} \int_0^\infty \int_0^\infty f(x)f(y)\varphi(x \wedge y)\pi(dy)\pi(dx) \\
&= \frac{2}{a(x_0)\pi(x_0)} \int_0^\infty \varphi(x)f(x) \int_x^\infty f(y)\pi(dy)\pi(dx) \\
&= -\frac{2}{a(x_0)\pi(x_0)} \int_0^\infty \varphi(x)f(x) \int_0^x f(y)\pi(dy)\pi(dx) \\
&= -\frac{1}{a(x_0)\pi(x_0)} \int_0^\infty \varphi(x) \left[\left(\int_0^x f(y)\pi(dy) \right)^2 \right]' dx \\
&= \int_0^\infty \left(\int_0^x f(y)\pi(dy) \right)^2 \frac{1}{a(x)\pi(x)} dx.
\end{aligned} \tag{2.3}$$

Using the above representation, we obtain the following comparison theorem directly.

Theorem 2.1. *Let a, a_1 be two C^1 positive function on $[0, \infty)$. Then Langevin diffusions X^a and X^{a_1} , with generators of form (2.1), possess the same stationary distribution π . Moreover, if $a \geq a_1$, then for any $f \in L_0^2(\pi)$,*

$$\sigma^2(X^a, f) \leq \sigma^2(X^{a_1}, f).$$

In particular, for fixed $f \in L_0^2(\pi)$, $\sigma^2(X^{ka}, f)$ is non-increasing for $k \in (0, \infty)$.

For multi-dimensional reversible diffusion processes, explicit representation (2.3) for the asymptotic variance is difficult to obtain. However, we could use Theorem 1.4 to get the similar comparison result as follows.

Let $V \in C^2(\mathbb{R}^d)$ with $\int_{\mathbb{R}^d} e^{V(x)} dx < \infty$. Consider the reversible diffusion process X^A generated by elliptic operator

$$L_A = \sum_{i,j} a_{ij}(x) \frac{\partial^2}{\partial x_i \partial x_j} + \sum_i b_i(x) \frac{\partial}{\partial x_i},$$

where $A(x) = (a_{ij}(x))_{1 \leq i,j \leq d}$, $x \in \mathbb{R}^d$ are positive definite matrices with $a_{ij} \in C^2(\mathbb{R}^d)$ and

$$b_i(x) = \sum_j a_{ij}(x) \frac{\partial}{\partial x_j} V(x) + \sum_j \frac{\partial}{\partial x_j} a_{ij}(x).$$

Assume that X^A is non-explosive. By [21, Theorem 4.2.1], we see that process X^A is ergodic with stationary distribution

$$\pi(dx) := \frac{e^{V(x)}}{\int_{\mathbb{R}^d} e^{V(y)} dy} dx.$$

Denote by $(\mathcal{E}_A(\cdot, \cdot), \mathcal{F}_A)$ the Dirichlet form associated with the process X^A . Explicitly, we see that

$$\mathcal{E}_A(u, v) = \int_{\mathbb{R}^d} \nabla u \cdot A \nabla v d\pi, \quad \text{for } u, v \in \mathcal{F}_A := \{u \in L^2(\pi) : \mathcal{E}_A(u, u) < \infty\}. \tag{2.4}$$

Theorem 2.2. Let $V \in C^2(\mathbb{R}^d)$ with $\int_{\mathbb{R}^d} e^{V(x)} dx < \infty$, $A(x) = (a_{ij}(x))_{1 \leq i, j \leq d}$ and $A_1(x) = (a_{ij}^1(x))_{1 \leq i, j \leq d}$, $x \in \mathbb{R}^d$ be positive definite matrices satisfying $a_{ij}, a_{ij}^1 \in C^2(\mathbb{R}^d)$ for $1 \leq i, j \leq d$. Suppose that $A_1 \leq A$ in the sense that $A(x) - A_1(x)$ is non-negative definite for all $x \in \mathbb{R}^d$. Then for any $f \in L_0^2(\pi)$,

$$\sigma^2(X^{A_1}, f) \geq \sigma^2(X^A, f). \quad (2.5)$$

In particular, for fixed $f \in L_0^2(\pi)$, $\sigma^2(X^{kA}, f)$ is non-increasing for $k \in (0, \infty)$.

Proof. Since $A_1 \leq A$, by (2.4) it is easy to check that $\mathcal{F}_{A_1} \supseteq \mathcal{F}_A$ and

$$\mathcal{E}_{A_1}(u, u) \leq \mathcal{E}_A(u, u) \quad \text{for all } u \in \mathcal{F}_A.$$

Fix $f \in L_0^2(\pi)$. The inequality (2.5) is trivial when $\sigma^2(X^{A_1}, f) = \infty$. Now assume that $\sigma^2(X^{A_1}, f) < \infty$. It follows from Theorem 1.4 that

$$2/\sigma^2(X^{A_1}, f) = \inf_{\substack{u \in \mathcal{F}_{A_1}, \\ \pi(fu)=1}} \mathcal{E}_{A_1}(u, u) \leq \inf_{\substack{u \in \mathcal{F}_A, \\ \pi(fu)=1}} \mathcal{E}_A(u, u) = 2/\sigma^2(X^A, f).$$

Hence, the proof is completed. \square

2.2 Non-reversible diffusions on Riemannian Manifolds

In this section, we turn to non-reversible case. Let M be a connected, complete Riemannian manifold with empty boundary or convex boundary, and $\langle \cdot, \cdot \rangle$ be the inner product under the Riemannian metric. Denote dx and Δ by the Riemannian volume and Laplace operator on M , respectively.

Let $\pi(dx) := e^{-U(x)} dx$ be a probability measure on M with potential function $U \in C^2(M)$. We consider the following diffusion operator:

$$\mathfrak{L}\varphi = \Delta\varphi - \langle \nabla U - Z, \nabla\varphi \rangle, \quad (2.6)$$

where Z is a C^1 vector field on M . Denote by \mathfrak{L}^* the dual operator of \mathfrak{L} on $L^2(\pi)$:

$$\mathfrak{L}^*\varphi = \Delta\varphi - \langle \nabla U + Z, \nabla\varphi \rangle - (\operatorname{div} Z - \langle \nabla U, Z \rangle)\varphi,$$

where div is the divergence operator. It is well known that π is the invariant measure of \mathfrak{L} if and only if $(\mathfrak{L}^*1, \varphi) = 0$ for $\varphi \in C_0^\infty(M)$, i.e.,

$$\int_M (\operatorname{div} Z - \langle \nabla U, Z \rangle)\varphi d\pi = \int_M \operatorname{div}(Ze^{-U})\varphi dx = 0.$$

From now on we assume that

$$\operatorname{div}(Ze^{-U}) \equiv 0. \quad (2.7)$$

Then by [3, Corollary 3.6], the diffusion X with generator \mathfrak{L} is ergodic with stationary distribution π .

Denote the symmetric part of \mathfrak{L} with respect to π by $\overline{\mathfrak{L}} := \Delta - \langle \nabla U, \nabla \cdot \rangle$. and let \overline{X} be the diffusion generated by $\overline{\mathfrak{L}}$.

Define $(\mathcal{E}, \mathcal{F})$ as the semi-Dirichlet form generated by \mathfrak{L} , and denote its symmetric part and antisymmetric part by $\overline{\mathcal{E}}$, $\widehat{\mathcal{E}}$ respectively. So from the integration by parts and (2.7), we have

$$\overline{\mathcal{E}}(\varphi, \phi) = \int_M \langle \nabla \varphi, \nabla \phi \rangle d\pi \quad \text{and} \quad \widehat{\mathcal{E}}(\varphi, \phi) = \int_M \phi \langle Z, \nabla \varphi \rangle d\pi \quad \varphi, \phi \in C_0^\infty(M).$$

Indeed, it is easy to check that $(\overline{\mathcal{E}}, \mathcal{F})$ is the Dirichlet form generated by $\overline{\mathfrak{L}}$.

We suppose that the following **Assumption A** holds:

- (A1) $|\Delta U| \leq \epsilon_* |\nabla U|^2 + C_U$ for some $\epsilon_* < 1$ and $C_U \geq 0$;
- (A2) there is a constant K such that $|Z| \leq K(|\nabla U| + 1)$;
- (A3) the symmetric Dirichlet form $(\overline{\mathcal{E}}, \mathcal{F})$ satisfies the Poincaré inequality, i.e., there exists a constant $\lambda_1 > 0$ such that

$$\|\varphi\|^2 \leq \lambda_1^{-1} \overline{\mathcal{E}}(\varphi, \varphi) \quad \text{for all } \varphi \in \mathcal{F},$$

where $\|\cdot\|$ is $L^2(\pi)$ -norm.

We note that (A3) is equivalent to the L^2 -exponential ergodicity of semigroup of diffusion \overline{X} .

Lemma 2.3. *If Assumption A and (2.7) hold, then $(\mathcal{E}, \mathcal{F})$ satisfies the sector condition (1.1). Therefore, Theorem 1.1 holds for the diffusion X .*

Proof. Since $(\overline{\mathcal{E}}, \mathcal{F})$ is symmetric, it satisfies the sector condition, we only need to check the sector condition for the antisymmetric part $\widehat{\mathcal{E}}$.

Fix $\phi, \varphi \in C_0^\infty(M)$. By Cauchy-Schwarz inequality and (A2) we have

$$\begin{aligned} \int_M \langle \phi Z, \nabla \varphi \rangle d\pi &\leq K \int_M (|\nabla U| + 1) |\phi \nabla \varphi| d\pi \\ &\leq K \int_M |\phi \nabla \varphi| d\pi + K \overline{\mathcal{E}}(\varphi, \varphi)^{1/2} \|\nabla U \phi\|. \end{aligned} \tag{2.8}$$

For the last term above, the integration by parts on manifold, Cauchy-Schwarz inequality and (A1) yield that

$$\begin{aligned} \|\nabla U \phi\|^2 &= - \int_M \langle \phi^2 \nabla U, \nabla e^{-U} \rangle dx = \int_M \operatorname{div}(\phi^2 \nabla U) e^{-U} dx \\ &= \int_M \langle 2\phi \nabla \phi, \nabla U \rangle d\pi + \int_M \Delta U \phi^2 d\pi \\ &= \int_M \langle 2\phi \nabla \phi, \nabla U \rangle d\pi + \int_M (\epsilon_* |\nabla U|^2 + C_U) \phi^2 d\pi. \end{aligned} \tag{2.9}$$

Now fix $\varepsilon > 0$ such that $\epsilon_* + \varepsilon < 1$. Combining inequality $|xy| \leq (x^2/\varepsilon + \varepsilon y^2)/2$ with (2.9) and (A3) we have

$$\begin{aligned} \|\nabla U \phi\|^2 &\leq 2 \int |\nabla \phi| |\phi \nabla U| d\pi + \int (\epsilon_* |\nabla U|^2 + C_U) \phi^2 d\pi \\ &\leq \frac{1}{\varepsilon} \overline{\mathcal{E}}(\phi, \phi) + (\epsilon_* + \varepsilon) \|\phi \nabla U\|^2 + C_U \|\phi\|^2 \\ &\leq \frac{1}{\varepsilon} \overline{\mathcal{E}}(\phi, \phi) + (\epsilon_* + \varepsilon) \|\phi \nabla U\|^2 + C_U \lambda_1^{-1} \overline{\mathcal{E}}(\phi, \phi), \end{aligned}$$

which implies that

$$\|\nabla U|\phi\|^2 \leq \frac{\lambda_1 + C_U \varepsilon}{(1 - \epsilon_* - \varepsilon)\varepsilon\lambda_1} \overline{\mathcal{E}}(\phi, \phi).$$

Combining this with (2.8) and (A3), we obtain that $\widehat{\mathcal{E}}$ satisfies the sector condition on \mathcal{F} . \square

From Lemma 2.3 and Theorem 1.1, we obtain the following comparison result.

Theorem 2.4. *Suppose that Assumption A holds. Then for any $f \in L_0^2(\pi)$,*

$$\sigma^2(X, f) \leq \sigma^2(\overline{X}, f).$$

Proof. Since the conditions in Theorem 1.1 are satisfied by Lemma 2.3, we obtain by taking $v = 0$ that

$$\begin{aligned} 2/\sigma^2(X, f) &= \inf_{u \in \mathcal{M}_{f,1}} \sup_{v \in \mathcal{M}_{f,0}} \mathcal{E}(u + v, u - v) \\ &\geq \inf_{u \in \mathcal{M}_{f,1}} \mathcal{E}(u, u) = \inf_{u \in \mathcal{M}_{f,1}} \overline{\mathcal{E}}(u, u) = 2/\sigma^2(\overline{X}, f). \end{aligned}$$

\square

Remark 2.5. Similar comparison result in Theorem 2.4 can be found in [8, 13]. For example, [13] proves the comparison theorem by using a spectral theorem (see [13, Section 3.4.3]). Here we provide a completely different proof by the new variational formula.

Example 2.6. ([16, Example 5.2]) *Let $M = \mathbb{R}^2$, potential function $U(x) = (1/2\pi)e^{-|x|^2/2}$ and vector field*

$$Z = -cx_2 \frac{\partial}{\partial x_1} + cx_1 \frac{\partial}{\partial x_2},$$

where c is a positive constant. Consider the 2-dimensional Ornstein-Uhlenbeck diffusion with rotation:

$$\mathfrak{L} := \frac{1}{2} \left(\frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} \right) - (x_1 + cx_2) \frac{\partial}{\partial x_1} - (x_2 - cx_1) \frac{\partial}{\partial x_2}.$$

Its invariant probability measure is $\pi(dx) = (1/2\pi)e^{-|x|^2/2}dx$. The symmetric part of \mathfrak{L} with respect to π is

$$\overline{\mathfrak{L}} := \left(\frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} \right) - x_1 \frac{\partial}{\partial x_1} - x_2 \frac{\partial}{\partial x_2}.$$

Since the symmetric Ornstein-Uhlenbeck diffusion generated by $\overline{\mathfrak{L}}$ is exponentially ergodic, (A3) is satisfied. A direct calculation shows that $\operatorname{div}(Ze^{-U}) = 0$ and (A1), (A2) are satisfied. Hence, Theorems 1.1 and 2.4 are valid.

3 Proofs of Theorem 1.1 and Corollary 1.3

Recall that $X = \{X_t\}_{t \geq 0}$ is a positive recurrent (or ergodic) Markov process on a Polish space (S, \mathcal{S}) , with strongly continuous contraction transition semigroup $\{P_t\}_{t \geq 0}$ and stationary distribution π . $(L, \mathcal{D}(L))$, $(\mathcal{E}, \mathcal{F})$ are its associated infinitesimal generator in $L^2(\pi)$ and semi-Dirichlet form, respectively. For fixed $f \in L^2_0(\pi)$, we want to study the asymptotic variance of X and f defined in (1.3). Indeed, from [15, Section 2.5], we see that the asymptotic variance can be represented by P_t as follows:

$$\sigma^2(X, f) = 2 \lim_{t \rightarrow \infty} \int_0^t (1 - \frac{s}{t})(P_s f, f) ds. \quad (3.1)$$

To prove Theorem 1.1, first we do some preparations. For any $\alpha > 0$, set $G_\alpha f = \int_0^\infty e^{-\alpha s} P_s f ds$ for $f \in L^2(\pi)$. From [18, Chapter 1, Proposition 1.10] we see that $(G_\alpha)_{\alpha > 0}$ is the strong continuous contraction resolvent associated to L and $G_\alpha f \in \mathcal{D}(L)$ for all $f \in L^2(\pi)$. If the semigroup $\{P_t\}_{t \geq 0}$ is L^2 -exponentially ergodic, then it is known that $Gf := \int_0^\infty P_s f ds \in L^2(\pi)$ for $f \in L^2_0(\pi)$.

Lemma 3.1. *Suppose that the semigroup $\{P_t\}_{t \geq 0}$ is L^2 -exponentially ergodic and its corresponding semi-Dirichlet form $(\mathcal{E}, \mathcal{F})$ satisfies the sector condition (1.1). Then for all $f \in L^2_0(\pi)$, we have $Gf \in \mathcal{D}(L)$ and*

$$\mathcal{E}(Gf, u) = (f, u), \quad u \in \mathcal{F}.$$

Proof. We first prove that $Gf \in \mathcal{D}(L)$ for all $f \in L^2_0(\pi)$. Note that the generator L is closed and densely defined, that is, $\mathcal{D}(L)$ is complete with respect to the graph norm $\|Lu\| + \|u\|, u \in \mathcal{D}(L)$ (see e.g. [18, Chapter 1, Proposition 1.10]). Thus for fixed $f \in L^2_0(\pi)$, we only need to prove that $\|G_{1/n}f - Gf\| \rightarrow 0$ as $n \rightarrow \infty$ and $\{G_{1/n}f\}_{n \geq 1}$ is a Cauchy sequence under $\|L \cdot\|$ by $G_{1/n}f \in \mathcal{D}(L), n \geq 1$. Indeed, it follows from L^2 -exponential ergodicity and Hölder inequality that

$$\begin{aligned} \|G_{1/n}f - Gf\| &= \left\| \int_0^\infty (1 - e^{-s/n}) P_s f ds \right\| \leq \int_0^\infty (1 - e^{-s/n}) \|P_s f\| ds \\ &\leq C \|f\| \int_0^\infty (1 - e^{-s/n}) e^{-\lambda_1 s} ds \\ &= C \|f\| \frac{1/n}{\lambda_1(\lambda_1 + 1/n)} \rightarrow 0, \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (3.2)$$

On the other hand, since $Lf = (\alpha - G_\alpha^{-1})f$ for all $\alpha > 0$ and $f \in L^2_0(\pi)$, we have

$$\begin{aligned} \|L(G_{1/n}f - G_{1/m}f)\| &= \|(1/n - G_{1/n}^{-1})G_{1/n}f - (1/m - G_{1/m}^{-1})G_{1/m}f\| \\ &= \left\| \frac{1}{n}G_{1/n}f - \frac{1}{m}G_{1/m}f \right\| \\ &\leq \frac{1}{n} \|G_{1/n}f - G_{1/m}f\| + \left| \frac{1}{n} - \frac{1}{m} \right| \|G_{1/m}f\| \\ &\rightarrow 0, \quad \text{as } n, m \rightarrow \infty. \end{aligned}$$

Therefore $Gf \in \mathcal{D}(L)$.

Next we prove that $-LGf = f$ for $f \in L_0^2(\pi)$. Arguing similarly as we did in (3.2), $\lim_{\beta \rightarrow 0} \|G_\beta f - Gf\| = 0$ for $f \in L_0^2(\pi)$. Combining this fact with the property $G_\beta - G_\alpha = (\alpha - \beta)G_\alpha G_\beta$, we obtain that

$$Gf - G_\alpha f = \alpha G_\alpha Gf, \quad \text{for } \alpha > 0, f \in L_0^2(\pi).$$

Using this equality and the fact $Gf \in \mathcal{D}(L)$ shows that for any $\alpha > 0$ and $f \in L_0^2(\pi)$,

$$G_\alpha(-LGf) = G_\alpha(G_\alpha^{-1} - \alpha)Gf = Gf - \alpha G_\alpha Gf = G_\alpha f.$$

That is, $-LGf = f$ for all $f \in L_0^2(\pi)$.

From above analysis and [18, Chapter 1, Corollary 2.10] we could obtain that for any $f \in L_0^2(\pi), u \in \mathcal{F}$,

$$\mathcal{E}(Gf, u) = ((-L)Gf, u) = (f, u).$$

□

We now proceed to prove Theorem 1.1.

Proof of Theorem 1.1. For fixed $f \in L_0^2(\pi)$, we first claim that the limit in (1.3), i.e. (3.1), exists and $\sigma^2(X, f) = 2(Gf, f) < \infty$. Indeed, for $t > 0$,

$$2 \int_0^t (1 - \frac{s}{t})(P_s f, f) ds = 2 \int_0^t (P_s f, f) ds - \frac{2}{t} \int_0^t s(P_s f, f) ds.$$

Since $\{P_t\}_{t \geq 0}$ is L^2 -exponentially ergodic, we arrive at

$$\begin{aligned} \frac{1}{t} \left| \int_0^t s(P_s f, f) ds \right| &\leq \frac{1}{t} \int_0^t s \|P_s f\| \|f\| ds \leq \frac{C \|f\|^2}{t} \int_0^t s e^{-\lambda_1 s} ds \\ &\leq \frac{1 - (1 + \lambda_1 t) e^{-\lambda_1 t}}{t} C \|f\|^2 \rightarrow 0, \quad \text{as } t \rightarrow \infty, \end{aligned}$$

and

$$\left| \int_t^\infty (P_s f, f) ds \right| \leq C \|f\|^2 \int_t^\infty e^{-\lambda_1 s} ds \rightarrow 0, \quad \text{as } t \rightarrow \infty.$$

Therefore, by combining above analysis, we obtain that the limit in (3.1) exists and

$$\sigma^2(X, f) = 2 \int_0^\infty (P_s f, f) ds < \infty.$$

By the Fubini-Tonelli's theorem and L^2 -exponential ergodicity again we get

$$\int_0^\infty (P_s f, f) ds = \int_0^\infty \int_S f P_s f d\pi ds = \int_S \int_0^\infty f P_s f ds d\pi = (Gf, f).$$

Thus

$$\sigma^2(X, f) = 2(Gf, f) < \infty. \quad (3.3)$$

To prove (1.4), we set $w = Gf/(Gf, f)$, $w^* = G^*f/(Gf, f)$ and $u_0 = (w + w^*)/2$, $v_0 = (w - w^*)/2$. Then $u_0 \in \mathcal{M}_{f,1}$ and $v_0 \in \mathcal{M}_{f,0}$ by noting

$$(Gf, f) = \int_0^\infty (P_s f, f) ds = \int_0^\infty (f, P_s^* f) ds = (G^* f, f).$$

Now let $v_1 = v - v_0$ for any $v \in \mathcal{M}_{f,0}$. By the definition of w, w^*, v_0 and Lemma 3.1, we have $\pi(v_1 f) = 0$ and

$$\mathcal{E}(v_1, w^*) = \mathcal{E}(w, v_1) = \frac{1}{(Gf, f)} \mathcal{E}(Gf, v_1) = \frac{1}{(Gf, f)} (f, v_1) = 0.$$

Therefore, using this fact with $\mathcal{E}(w, w^*) = 1/(Gf, f)$ and $\mathcal{E}(u, u) \geq 0$ for all $u \in \mathcal{F}$ gives that

$$\mathcal{E}(u_0 + v, u_0 - v) = \mathcal{E}(w - v_1, w^* + v_1) = \mathcal{E}(w, w^*) - \mathcal{E}(v_1, v_1) \leq 1/(Gf, f),$$

which implies that

$$1/(Gf, f) \geq \inf_{u \in \mathcal{M}_{f,1}} \sup_{v \in \mathcal{M}_{f,0}} \mathcal{E}(u + v, u - v). \quad (3.4)$$

For the converse inequality, let $u_1 = u - u_0$ for any $u \in \mathcal{M}_{f,1}$. Since $u_0 \in \mathcal{M}_{f,1}$, we also have $\pi(u_1 f) = 0$. Similar argument shows that

$$\mathcal{E}(u + v_0, u - v_0) = \mathcal{E}(w + u_1, w^* + u_1) = \mathcal{E}(w, w^*) + \mathcal{E}(u_1, u_1) \geq 1/(Gf, f).$$

Therefore,

$$1/(Gf, f) \leq \inf_{u \in \mathcal{M}_{f,1}} \sup_{v \in \mathcal{M}_{f,0}} \mathcal{E}(u + v, u - v). \quad (3.5)$$

So we obtain (1.4) by combining (3.4), (3.5) and the fact $\sigma^2(X, f) = 2(Gf, f)$.

When process X is reversible, $\mathcal{E}(\cdot, \cdot)$ is symmetric, i.e.,

$$\mathcal{E}(u, v) = \mathcal{E}(v, u), \quad \text{for } u, v \in \mathcal{F}.$$

Thus

$$\mathcal{E}(u + v, u - v) = \mathcal{E}(u, u) - \mathcal{E}(v, v) \leq \mathcal{E}(u, u).$$

That is, the supremum in (1.4) is attained by $v = 0$ for any fixed $u \in \mathcal{M}_{f,1}$. Hence, we obtain (1.5). \square

By using Theorem 1.1, we prove Corollary 1.3 as follows.

Proof of Corollary 1.3. Fix an open set $\Omega \subset S$ with $\pi(\Omega) \in (0, 1)$. It follows from [10, Theorem 3.3] that

$$1/\mathbb{E}_\pi \tau_\Omega = \inf_{u \in \mathcal{N}_{\Omega,1}} \mathcal{E}(u, u), \quad (3.6)$$

where $\mathcal{N}_{\Omega,1} := \{u \in \mathcal{F} : u|_{\Omega^c} = 0 \text{ and } \pi(u) = 1\}$. Take

$$f = \frac{\mathbf{1}_\Omega - \pi(\Omega)}{1 - \pi(\Omega)}.$$

It is easy to check that $\pi(f) = 0$ and $\|f\|^2 = \pi(\Omega)/\pi(\Omega^c)$. Notice that for any $u \in \mathcal{N}_{\Omega,1}$, by simple calculation we have $\pi(uf) = 1$, thus $u \in \mathcal{M}_{f,1}$. So we see that $\mathcal{N}_{\Omega,1} \subset \mathcal{M}_{f,1}$. Combining this fact with (1.5) and (3.6), we obtain that

$$2/\sigma^2(X, f) = \inf_{u \in \mathcal{M}_{f,1}} \mathcal{E}(u, u) \leq \inf_{u \in \mathcal{N}_{\Omega,1}} \mathcal{E}(u, u) = 1/\mathbb{E}_\pi \tau_\Omega.$$

That is, $\mathbb{E}_\pi \tau_\Omega \leq \sigma^2(X, f)/2$. Moreover, from the reversibility and L^2 -exponential ergodicity we have

$$\sigma^2(X, f)/2 = \int_0^\infty (P_s f, f) ds \leq \int_0^\infty \|P_s f\| \|f\| ds \leq \|f\|^2 / \lambda_1.$$

Hence,

$$\mathbb{E}_\pi \tau_\Omega \leq \frac{\|f\|^2}{2\lambda_1} = \frac{\pi(\Omega)}{2\lambda_1 \pi(\Omega^c)}.$$

□

Acknowledgement Lu-Jing Huang acknowledges support from NSFC (No. 11901096), NSF-Fujian(No. 2020J05036), the Program for Probability and Statistics: Theory and Application (No. IRTL1704), and the Program for Innovative Research Team in Science and Technology in Fujian Province University (IRTSTFJ). Yong-Hua Mao and Tao Wang acknowledge support by the National Key R&D Program of China (2020YFA0712900) and the National Natural Science Foundation of China (Grant No.11771047).

References

- [1] C. Andrieu and S. Livingstone. Peskun-Tierney ordering for Markov chain and process Monte Carlo: beyond the reversible scenario. *arXiv:1906.06197*, 2019.
- [2] J. Bierkens. Non-reversible Metropolis-Hastings. *Stat. Comput.*, 26(6):1213–1228, 2016.
- [3] V. I. Bogachev, M. Rockner, and F.-Y. Wang. Elliptic equations for invariant measures on finite and infinite dimensional manifolds. *J. Math. Pure. Appl.*, 80(2):177–221, 2001.
- [4] M.-F. Chen. *Eigenvalues, inequalities, and ergodic theory*. Springer-Verlag London, Ltd., London, 2004.
- [5] T.-L. Chen, W.-K. Chen, C.-R. Hwang, and H.-M. Pai. On the optimal transition matrix for Markov chain Monte Carlo sampling. *SIAM J. Control Optim.*, 50(5):2743–2762, 2012.
- [6] T.-L. Chen and C.-R. Hwang. Accelerating reversible Markov chains. *Statist. Probab. Lett.*, 83(9):1956–1962, 2013.
- [7] Y. Derriennic and M. Lin. The central limit theorem for Markov chains with normal transition operators, started at a point. *Probab. Theory Related Fields*, 119:508–528, 2001.
- [8] A.-B. Duncan, T. Lelièvre, and G.-A. Pavliotis. Variance reduction using nonreversible Langevin samplers. *J. Stat. Phys.*, 163:457–491, 2016.
- [9] O. Häggström. On the central limit theorem for geometrically ergodic Markov chains. *Probab. Theory Related Fields*, 132:74–82, 2005.

- [10] L.-J. Huang, K.-Y. Kim, Y.-H. Mao, and T. Wang. Variational principles for the exit time of Hunt processes generated by semi-Dirichlet forms. *ArXiv 2011.04334*, 2020.
- [11] L.-J. Huang and Y.-H. Mao. Variational formulas of asymptotic variance for general discrete-time Markov chains. *arXiv:2012.13895*, 2020.
- [12] C.-R. Hwang. Accelerating Monte Carlo Markov processes. *Cosmos*, 1(1):87–94, 2005.
- [13] C.-R. Hwang, R. Normanda, and S.-J. Wu. Variance reduction for diffusions. *Stoch. Proc. Appl.*, 125(9):3522–3540, 2015.
- [14] C. Kipnis and S.-R.-S. Varadhan. Central limit theorem for additive functionals of reversible Markov processes and applications to simple exclusions. *Comm. Math. Phys.*, 104(1):1–19, 1986.
- [15] T. Komorowski, C. Landim, and S. Olla. *Fluctuations in Markov processes: time symmetry and martingale approximation*. Springer-Verlag, Berlin, 2012.
- [16] S. Kusuoka and I. Shigekawa. Exponential convergence of Markovian semigroups and their spectra on L^p -spaces. *Kyoto J. Math.*, 54(2):367—399, 2014.
- [17] F. Leisen and A. Mira. An extension of Peskun and Tiernery orderings to continuous time Markov chains. *Statist. Sinica*, 18:1641–1651, 2008.
- [18] Z.-M. Ma and M. Röckner. *Introduction to the Theory of (Non-Symmetric) Dirichlet forms*. Springer-Verlag Berlin Heidelberg, 1992.
- [19] F. Maire, R. Douc, and J. Olsson. Comparison of asymptotic variances of inhomogeneous Markov chains with applications to Markov chain Monte Carlo methods. *Ann. Statist.*, 42(4):1483–1510, 2014.
- [20] P.-H. Peskun. Optimum Monte-Carlo sampling using Markov chains. *Biometrika*, 60(3):607–612, 1973.
- [21] G. Da Prato and J. Zabczyk. *Ergodicity for Infinite Dimensional Systems*. Cambridge University Press, 1996.
- [22] L. Rey-Bellet and K. Spiliopoulos. Irreversible Langevin samplers and variance reduction: a large deviations approach. *Nonlinearity*, 7:2081–2103, 2015.
- [23] G.-O. Roberts and J.-S. Rosenthal. Geometric ergodicity and hybrid Markov chains. *Electron. Comm. Probab.*, 2:13–25, 1997.
- [24] G.-O. Roberts and J.-S. Rosenthal. Variance bounding Markov chains. *Ann. Appl. Probab.*, 18(3):1201–1214, 2008.
- [25] G. O. Roberts and J. S. Rosenthal. Minimising MCMC variance via diffusion limits, with an application to simulated tempering. *Ann. Appl. Probab.*, 24:131–149, 2014.
- [26] Y. Sun, F. Gomez, and J. Schmidhuber. Improving the asymptotic performance of Markov chain Monte-Carlo by inserting vortices. *Neural Information Processing Systems*, pages 2235–2243, 2010.

- [27] L. Tierney. A note on Metropolis-Hastings kernels for general state spaces. *Ann. Appl. Probab.*, 8(1):1–9, 1998.

Lu-Jing Huang: College of Mathematics and Informatics, Fujian Normal University, Fuzhou, 350007, P.R. China. E-mail: huanglj@fjnu.edu.cn

Yong-Hua Mao: Laboratory of Mathematics and Complex Systems(Ministry of Education), School of Mathematical Sciences, Beijing Normal University, Beijing 100875, P.R. China. E-mail: maoyh@bnu.edu.cn

Tao Wang: Laboratory of Mathematics and Complex Systems(Ministry of Education), School of Mathematical Sciences, Beijing Normal University, Beijing 100875, P.R. China. E-mail: wang_tao@mail.bnu.edu.cn