

Local and uniform moduli of continuity of chi-square processes

Michael B. Marcus Jay Rosen *

March 19, 2022

1 Introduction

Let $\eta = \{\eta(t); t \in [0, 1]\}$ be a mean zero continuous Gaussian process with covariance $U = \{U(s, t), s, t \in [0, 1]\}$, with $U(0, 0) > 0$. Let $\{\eta_i; i = 1, \dots, k\}$ be independent copies of η and set,

$$Y_k(t) = \sum_{i=1}^k \eta_i^2(t), \quad t \in [0, 1]. \quad (1.1)$$

The stochastic process $Y_k = \{Y_k(t), t \in [0, 1]\}$ is referred to as a chi-square process of order k with kernel U .

Theorem 1.1 *Let $\phi(t)$ be a positive function on $[0, \delta]$ for some $\delta > 0$. If*

$$\limsup_{t \rightarrow 0} \frac{\eta(t) - \eta(0)}{\phi(t)} = 1 \quad a.s., \quad (1.2)$$

then for all integers $k \geq 1$,

$$\limsup_{t \rightarrow 0} \frac{Y_k(t) - Y_k(0)}{\phi(t)} = 2Y_k^{1/2}(0) \quad a.s. \quad (1.3)$$

When $k = 1$ this is particularly simple. Since η is symmetric it follows from (1.2) that,

$$\liminf_{t \rightarrow 0} \frac{\eta(t) - \eta(0)}{\phi(t)} = -1 \quad a.s. \quad (1.4)$$

*Research of Jay Rosen was partially supported by grants from the Simons Foundation.

⁰Key words and phrases: chi-square processes, moduli of continuity

⁰AMS 2020 subject classification: 60F15, 60G15, 60G17

Therefore, writing $Y_1(t) - Y_1(0) = (\eta(t) - \eta(0))(\eta(t) + \eta(0))$ and using the continuity of η , we see that

$$\limsup_{t \rightarrow 0} \frac{Y_1(t) - Y_1(0)}{\phi(t)} = 2(\eta(0) \vee -\eta(0)) \quad a.s., \quad (1.5)$$

which is (1.3).

A result similar to Theorem 1.1 for the limiting behavior of chi-square sequences at infinity is given in [2, Lemma 6.5].

Set

$$\sigma^2(u, v) = E(\eta(u) - \eta(v))^2 \quad \text{and} \quad \tilde{\sigma}^2(x) = \sup_{|u-v| \leq x} \sigma^2(u, v). \quad (1.6)$$

Theorem 1.2 *Assume that $\inf_{t \in [0,1]} U(t, t) > 0$ and,*

$$\lim_{x \rightarrow 0} \tilde{\sigma}^2(x) \log 1/x = 0. \quad (1.7)$$

Let $\varphi(t)$ be a positive function on $[0, 1]$. Then if

$$\lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{\eta(u) - \eta(v)}{\varphi(|u-v|)} = 1 \quad a.s., \quad (1.8)$$

for all intervals $\Delta \subset [0, 1]$, it follows that for all intervals $\Delta \subset [0, 1]$ and all integers $k \geq 1$,

$$\lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{Y_k(u) - Y_k(v)}{\varphi(|u-v|)} = 2 \sup_{u \in \Delta} Y_k^{1/2}(u), \quad a.s. \quad (1.9)$$

When η is a continuous Gaussian process with stationary increments, $\sigma^2(u, v)$ in (1.6) can be written as $\sigma^2(u - v, 0)$. In this case if $\tilde{\sigma}^2(x)$ is asymptotic to an increasing function at 0, then (1.8) implies (1.7). We discuss this further in Remark 2.1.

An extensive treatment of Gaussian processes satisfying (1.2) and (1.8) is given in [1, Chapter 7].

2 Proofs

Proof of Theorem 1.1 Let $\eta_i(t)$, $i = 1, \dots, k$, be independent copies of $\eta(t)$. We write,

$$\begin{aligned} \eta_i^2(t) - \eta_i^2(0) &= (\eta_i(t) - \eta_i(0))(\eta_i(t) + \eta_i(0)) \\ &= (\eta_i(t) - \eta_i(0))(2\eta_i(0) + (\eta_i(t) - \eta_i(0))) \\ &= 2(\eta_i(t) - \eta_i(0))\eta_i(0) + (\eta_i(t) - \eta_i(0))^2. \end{aligned} \quad (2.1)$$

By (1.2)

$$\begin{aligned} \limsup_{t \rightarrow 0} \frac{\sum_{i=1}^k (\eta_i(t) - \eta_i(0))^2}{\phi(t)} & \quad (2.2) \\ & \leq \sum_{i=1}^k \limsup_{t \rightarrow 0} \frac{|\eta_i(t) - \eta_i(0)|}{\phi(t)} \lim_{t \rightarrow 0} |\eta_i(t) - \eta_i(0)| = 0. \end{aligned}$$

Consequently, using (1.1) we see that,

$$\limsup_{t \rightarrow 0} \frac{Y_k(t) - Y_k(0)}{\phi(t)} = \limsup_{t \rightarrow 0} \frac{2 \sum_{i=1}^k (\eta_i(t) - \eta_i(0)) \eta_i(0)}{\phi(t)}. \quad (2.3)$$

Write,

$$\begin{aligned} & (\eta_i(t) - \eta_i(0)) \eta_i(0) \quad (2.4) \\ & = \left(\eta_i(t) - \frac{U(0,t)}{U(0,0)} \eta_i(0) \right) \eta_i(0) - \left(\frac{U(0,0) - U(0,t)}{U(0,0)} \right) \eta_i^2(0). \end{aligned}$$

We show below that

$$\limsup_{t \rightarrow 0} \frac{|U(0,0) - U(0,t)|}{U(0,0)\phi(t)} = 0. \quad (2.5)$$

Consequently,

$$\limsup_{t \rightarrow 0} \frac{Y_k(t) - Y_k(0)}{\phi(t)} = \limsup_{t \rightarrow 0} \frac{2 \sum_{i=1}^k \left(\eta_i(t) - \frac{U(0,t)}{U(0,0)} \eta_i(0) \right) \eta_i(0)}{\phi(t)}. \quad (2.6)$$

Let $\{\xi_i(t), t \in [0, 1]\}$, $i = 1, \dots, k$, be independent copies of a mean zero Gaussian process $\{\xi(t), t \in [0, 1]\}$, and set $\vec{\xi}(t) = (\xi_1(t), \dots, \xi_k(t))$. Let $\vec{v} \in R^k$ with $\|\vec{v}\|_2 = 1$. Computing the covariances we see that,

$$\{(\vec{v} \cdot \vec{\xi}(t)), t \in [0, 1]\} \stackrel{law}{=} \{\xi(t), t \in [0, 1]\}. \quad (2.7)$$

(This relationship is used by P. Revesz in [3, Theorem 18.1] to obtain LILs for Brownian motion in R^k .)

Therefore, since $(\eta_i(t) - (U(0,t)/U(0,0))\eta_i(0))$ and $\eta_i(0)$ are independent for $i = 1, \dots, k$, we see that,

$$\begin{aligned} & \left\{ \left(\vec{\eta}(t) - \frac{U(0,t)}{U(0,0)} \vec{\eta}(0) \right) \cdot \frac{\vec{\eta}(0)}{\|\vec{\eta}(0)\|_2}, t \in [0, 1] \right\} \quad (2.8) \\ & \stackrel{law}{=} \left\{ \left(\eta(t) - \frac{U(0,t)}{U(0,0)} \eta(0) \right), t \in [0, 1] \right\}, \end{aligned}$$

where $\vec{\eta}(t) = (\eta_1(t), \dots, \eta_k(t))$. Consequently, (2.6) implies that

$$\limsup_{t \rightarrow 0} \frac{Y_k(t) - Y_k(0)}{\phi(t) \|\vec{\eta}(0)\|_2} \stackrel{law}{=} \limsup_{t \rightarrow 0} \frac{2 \left(\eta(t) - \frac{U(0,t)}{U(0,0)} \eta(0) \right)}{\phi(t)}. \quad (2.9)$$

Using (2.4) again and (2.5) we see that this implies that,

$$\limsup_{t \rightarrow 0} \frac{Y_k(t) - Y_k(0)}{\phi(t) \|\vec{\eta}(0)\|_2} \stackrel{law}{=} \limsup_{t \rightarrow 0} \frac{2(\eta(t) - \eta(0))}{\phi(t)} = 2, \quad (2.10)$$

where the last equality uses (1.2). Since,

$$\|\vec{\eta}(0)\|_2 = \left(\sum_{i=1}^k \eta_i^2(0) \right)^{1/2} = Y_k^{1/2}(0), \quad (2.11)$$

we obtain (1.3).

To obtain (2.5) we first note that it follows from (1.2) that,

$$\phi(t) = (E(\eta(t) - \eta(0))^2)^{1/2} h(t), \quad (2.12)$$

for some function h such that $\lim_{t \downarrow 0} h(t) = \infty$. To see this, suppose that it is false. Then there exists a sequence $\{t_k\}$, with $\lim_{k \rightarrow \infty} t_k = 0$, such that $\sup_k h(t_k) \leq M$. Therefore, if (1.2) holds, we would have,

$$\sup_k \frac{\eta(t_k) - \eta(0)}{(E(\eta(t_k) - \eta(0))^2)^{1/2}} \leq M \quad a.s. \quad (2.13)$$

This is not possible because $\{\eta(t_k) - \eta(0)/(E(\eta(t_k) - \eta(0))^2)^{1/2}\}$ is a sequence of standard normal random variables.

Since,

$$U(0,0) - U(0,t) = E((\eta(t) - \eta(0)) \eta(0)) \leq E \left((\eta(t) - \eta(0))^2 \right)^{1/2} U^{1/2}(0,0), \quad (2.14)$$

we have,

$$\frac{U(0,0) - U(0,t)}{\phi(t)} \leq \frac{U^{1/2}(0,0)}{h(t)}. \quad (2.15)$$

Using the fact that $\lim_{t \downarrow 0} h(t) = \infty$ we get (2.5). \square

Proof of Theorem 1.2 Note that (1.8) implies that $\{\eta(t), t \in [0, 1]\}$ and therefore $\{\eta^2(t), t \in [0, 1]\}$ are uniformly continuous on $[0, 1]$, which in turn implies that for all $k \geq 1$, $\{Y_k(t); t \in [0, 1]\}$ is uniformly continuous on $[0, 1]$.

To show,

$$\lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{Y_k(u) - Y_k(v)}{\varphi(|u-v|)} \geq 2 \sup_{t \in \Delta} Y_k^{1/2}(t), \quad a.s. \quad (2.16)$$

it suffices to show that for any $d \in \Delta$,

$$\lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{Y_k(u) - Y_k(v)}{\varphi(|u-v|)} \geq 2Y_k^{1/2}(d), \quad a.s. \quad (2.17)$$

This is because, (2.17) holding almost surely implies that for any countable dense set $\Delta' \subset \Delta$,

$$\lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{Y_k(u) - Y_k(v)}{\varphi(|u-v|)} \geq 2 \sup_{d \in \Delta'} Y_k^{1/2}(d), \quad a.s. \quad (2.18)$$

which implies (2.16).

Let $u, v, d \in \Delta$. We write,

$$\begin{aligned} \eta_i^2(u) - \eta_i^2(v) &= (\eta_i(u) - \eta_i(v))(\eta_i(u) + \eta_i(v)) \\ &= (\eta_i(u) - \eta_i(v))(2\eta_i(d) + (\eta_i(u) - \eta_i(d)) + (\eta_i(v) - \eta_i(d))). \end{aligned} \quad (2.19)$$

It follows from (1.8) that,

$$\begin{aligned} &\lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{\sum_{i=1}^k (\eta_i(u) - \eta_i(v))(\eta_i(u) - \eta_i(d))}{\varphi(|u-v|)} \\ &\leq \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{\sum_{i=1}^k |\eta_i(u) - \eta_i(v)| \sup_{u \in \Delta} |\eta_i(u) - \eta_i(d)|}{\varphi(|u-v|)} \\ &\leq \sum_{i=1}^k \sup_{u \in \Delta} |\eta_i(u) - \eta_i(d)| := \Delta^*. \end{aligned} \quad (2.20)$$

Therefore, using (2.16)–(2.20) and (2.11) with 0 replaced by d we see that,

$$\begin{aligned} &\lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{Y_k(u) - Y_k(v)}{\varphi(|u-v|) Y_k^{1/2}(d)} \\ &\geq \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{2 \sum_{i=1}^k (\eta_i(u) - \eta_i(v)) \eta_i(d)}{\|\vec{\eta}(d)\|_2 \varphi(|u-v|)} - \frac{2\Delta^*}{\|\vec{\eta}(d)\|_2}. \end{aligned} \quad (2.21)$$

To simplify the notation we take $V(u, v) = U(u, v)/U(d, d)$. Write

$$\eta_i(u) - \eta_i(v) = V(v, d)\eta_i(u) - V(u, d)\eta_i(v) + G_i(u, v), \quad (2.22)$$

where,

$$G_i(u, v) = (1 - V(v, d))\eta_i(u) - (1 - V(u, d))\eta_i(v). \quad (2.23)$$

In this notation,

$$\begin{aligned} & \frac{\sum_{i=1}^k (\eta_i(u) - \eta_i(v))\eta_i(d)}{\|\vec{\eta}(d)\|_2 \varphi(|u - v|)} - \frac{\sum_{i=1}^k G_i(u, v)\eta_i(d)}{\|\vec{\eta}(d)\|_2 \varphi(|u - v|)} \\ &= \frac{\sum_{i=1}^k (V(v, d)\eta_i(u) - V(u, d)\eta_i(v)) \eta_i(d)}{\|\vec{\eta}(d)\|_2 \varphi(|u - v|)}. \end{aligned} \quad (2.24)$$

Note that for all $u, v \in [0, 1]$,

$$E((V(v, d)\eta_i(u) - V(u, d)\eta_i(v))\eta_i(d)) = 0.$$

This shows that $\eta_i(d)$ is independent of $\{V(v, d)\eta_i(u) - V(u, d)\eta_i(v); u, v \in [0, 1]\}$. Therefore by (2.7),

$$\begin{aligned} & \left\{ \sum_{i=1}^k (V(v, d)\eta_i(u) - V(u, d)\eta_i(v)) \frac{\eta_i(d)}{\|\vec{\eta}(d)\|_2}; \quad u, v \in [0, 1] \right\} \\ & \stackrel{law}{=} \left\{ V(v, d)\eta(u) - V(u, d)\eta(v); \quad u, v \in [0, 1] \right\}. \end{aligned} \quad (2.25)$$

It follows that,

$$\begin{aligned} & \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{\sum_{i=1}^k (\eta_i(u) - \eta_i(v))\eta_i(d)}{\|\vec{\eta}(d)\|_2 \varphi(|u - v|)} \\ & \quad + \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{\sum_{i=1}^k |G_i(u, v)|\eta_i(d)}{\|\vec{\eta}(d)\|_2 \varphi(|u - v|)} \\ & \geq \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{V(v, d)\eta(u) - V(u, d)\eta(v)}{\varphi(|u - v|)}. \end{aligned} \quad (2.26)$$

Using (2.22) we can write,

$$\begin{aligned}
& \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{V(v, d)\eta(u) - V(u, d)\eta(v)}{\varphi(|u-v|)} & (2.27) \\
& \geq \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{(\eta(u) - \eta(v))}{\varphi(|u-v|)} - \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{|G(u, v)|}{\varphi(|u-v|)} \\
& = 1 - \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{|G(u, v)|}{\varphi(|u-v|)},
\end{aligned}$$

where

$$G(u, v) = (1 - V(v, d))\eta(u) - (1 - V(u, d))\eta(v) \quad (2.28)$$

and we use (1.8) for the last line in (2.27).

It follows from (2.26) and (2.27) that,

$$\lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{\sum_{i=1}^k (\eta_i(u) - \eta_i(v))\eta_i(d)}{\|\vec{\eta}(d)\|_2 \varphi(|u-v|)} \geq 1 - \mathcal{H}, \quad (2.29)$$

where,

$$\mathcal{H} = \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{\sum_{i=1}^k |G_i(u, v)| |\eta_i(d)|}{\|\vec{\eta}(d)\|_2 \varphi(|u-v|)} + \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{|G(u, v)|}{\varphi(|u-v|)}$$

Using the Schwartz inequality followed by the triangle inequality we note that,

$$\frac{\sum_{i=1}^k |G_i(u, v)| |\eta_i(d)|}{\|\vec{\eta}(d)\|_2 \varphi(|u-v|)} \leq \frac{\sum_{i=1}^k |G_i(u, v)|}{\varphi(|u-v|)}.$$

Therefore,

$$\mathcal{H} \leq \sum_{i=0}^k \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{|G_i(u, v)|}{\varphi(|u-v|)}, \quad a.s., \quad (2.30)$$

where for notational convenience we have set $G_0 = G$.

Set,

$$\sigma^2(u, v) = E(\eta(u) - \eta(v))^2, \quad (2.31)$$

and

$$\tilde{\sigma}^2(x) = \sup_{|u-v| \leq x} \sigma^2(u, v). \quad (2.32)$$

Then it follows from (1.8) that we can write,

$$\varphi(h) = \tilde{\sigma}(h)g(h), \text{ where necessarily, } \lim_{h \rightarrow 0} g(h) = \infty. \quad (2.33)$$

This follows by a minor modification of the argument used to prove (2.12). Note that for any sequence $h_k \rightarrow 0$ we can find sequences $\{u_k\}, \{v_k\}$ in Δ , with $|u_k - v_k| \leq h_k$ such that $\tilde{\sigma}(h_k) \leq 2\sigma(u_k, v_k)$. Now, suppose that $\limsup_{h \rightarrow 0} g(h) = M$. Then by (1.8) we would have,

$$\sup_{k \rightarrow \infty} \frac{\eta(u_k) - \eta(v_k)}{\sigma(u_k, v_k)} \leq 4M \quad a.s.$$

This is not possible because each term $(\eta(u_k) - \eta(v_k))/\sigma(u_k, v_k)$ is a standard normal random variable.

We show in Lemma 2.1 below that,

$$|G(u, v)| \leq \frac{\sigma(d, v)}{U^{1/2}(d, d)} |\eta(u) - \eta(v)| + \frac{\sigma(u, v)}{U^{1/2}(d, d)} |\eta(v)|. \quad (2.34)$$

Therefore, using (1.8) and (2.33) we have,

$$\begin{aligned} \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{|G(u, v)|}{\varphi(|u-v|)} &\leq \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{|\eta(u) - \eta(v)| \sigma(d, v)}{U^{1/2}(d, d) \varphi(|u-v|)} \\ &\quad + \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{\sigma(u, v)}{U^{1/2}(d, d) \varphi(|u-v|)} \eta(v) \\ &\leq \sup_{d, v \in \Delta} \frac{\sigma(d, v)}{U^{1/2}(d, d)} + \lim_{h \rightarrow 0} \frac{1}{g(h)} \sup_{v \in \Delta} \frac{|\eta(v)|}{U^{1/2}(d, d)} = \frac{\tilde{\sigma}(|\Delta|)}{U^{1/2}(d, d)}. \end{aligned} \quad (2.35)$$

where $\tilde{\sigma}(|\Delta|)$ is defined in (2.32). We now use (2.21), (2.29) and (2.35) to see that,

$$\begin{aligned} \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{Y_k(u) - Y_k(v)}{\varphi(|u-v|)} &\quad (2.36) \\ &\geq 2Y_k^{1/2}(d) \left(1 - (k+1) \frac{\tilde{\sigma}(|\Delta|)}{U^{1/2}(d, d)} \right) - 2\Delta^*, \quad a.s., \end{aligned}$$

where $d \in \Delta$ and Δ^* is defined in (2.20).

Suppose that $|\Delta| = 1/n$. Then it follows from Lemma 2.2 that,

$$P \left(\Delta^* \geq k((1+2C)\tilde{\sigma}^2(1/n) \log n)^{1/2} \right) \leq \frac{2k}{nC}. \quad (2.37)$$

Now let $\Delta(d, n) \subseteq \Delta$, be an interval of size $1/n$ that contains d . It follows from (2.36) and (2.37) applied to $\Delta(d, n)$ and $\Delta^*(d, n)$ that ,

$$\begin{aligned}
& \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{Y_k(u) - Y_k(v)}{\varphi(|u-v|)} & (2.38) \\
& \geq \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta(d, n)}} \frac{Y_k(u) - Y_k(v)}{\varphi(|u-v|)} \\
& \geq 2Y_k^{1/2}(d) \left(1 - (k+1) \frac{\tilde{\sigma}(|\Delta|)}{U^{1/2}(d, d)} \right) - 2k((1+2C)\tilde{\sigma}^2(1/n) \log n)^{1/2},
\end{aligned}$$

except, possibly, on a set of measure $2k/n^C$. Taking $n \rightarrow \infty$, and using (1.7), gives (2.17) and consequently (2.16), which is the lower bound in (1.9).

We now obtain the upper bound in (1.9). Let $\Delta_{m,n} = \Delta \cap [\frac{m-1}{n}, \frac{m+1}{n}]$. Analogous to (2.38) we have,

$$\begin{aligned}
& \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta_{m,n}}} \frac{Y_k(u) - Y_k(v)}{\varphi(|u-v|)} & (2.39) \\
& \leq 2Y_k^{1/2}(m/n) \left(1 + (k+1) \frac{\tilde{\sigma}(2/n)}{U^{1/2}(\frac{m}{n}, \frac{m}{n})} \right) + k((1+2C)\tilde{\sigma}^2(2/n) \log n)^{1/2} \\
& := 2Y_k^{1/2}(m/n) \mathcal{A}(\tilde{\sigma}(2/n), k, n, C),
\end{aligned}$$

except, possibly, on a set of measure $2k/n^C$. The proof of (2.39) proceeds in essentially the same way as the proof of (2.38). In the proof of the lower bound in (2.38) we subtract several terms. In proving the upper bound in (2.39) we add these terms. It then follows that,

$$\begin{aligned}
& \lim_{h \rightarrow 0} \sup_{\substack{|u-v| \leq h \\ u, v \in \Delta}} \frac{Y_k(u) - Y_k(v)}{\varphi(|u-v|)} & (2.40) \\
& \leq 2 \sup_{m=1, \dots, n-1} Y_k^{1/2}(m/n) \mathcal{A}(\tilde{\sigma}(2/n), k, n, C) \\
& \leq 2 \sup_{v \in \Delta} Y_k^{1/2}(v) \mathcal{A}(\tilde{\sigma}(2/n), k, n, C),
\end{aligned}$$

except possibly on a set of measure $2k/n^{C-1}$. Taking the limit as $n \rightarrow \infty$ gives the upper bound in (1.9). \square

Lemma 2.1

$$|G(u, v)| \leq \frac{\sigma(d, v)}{U^{1/2}(d, d)} |\eta(u) - \eta(v)| + \frac{\sigma(u, v)}{U^{1/2}(d, d)} |\eta(v)|. \quad (2.41)$$

Proof: We write,

$$G(u, v) = (1 - V(v, d))(\eta(u) - \eta(v)) + (V(u, d) - V(v, d))\eta(v).$$

Note that,

$$\begin{aligned} |(V(u, d) - V(v, d))| &= \frac{|(E(\eta(u) - \eta(v))\eta(d))|}{U(d, d)} \\ &\leq \frac{(E(\eta(u) - \eta(v))^2 E\eta^2(d))^{1/2}}{U(d, d)} \\ &\leq \frac{\sigma(u, v)}{U^{1/2}(d, d)}, \end{aligned}$$

and,

$$\begin{aligned} (1 - V(v, d)) &= \frac{(E(\eta(d) - \eta(v))\eta(d))}{U(d, d)} \\ &\leq \frac{(E(\eta(d) - \eta(v))^2 E\eta^2(d))^{1/2}}{U(d, d)} \\ &\leq \frac{\sigma(d, v)}{U^{1/2}(d, d)}. \end{aligned}$$

□

Lemma 2.2

$$P \left(\sum_{i=1}^k \sup_{u \in \Delta} |\eta_i(u) - \eta_i(d)| \geq k((1 + 2C)\tilde{\sigma}^2(|\Delta|) \log 1/|\Delta|)^{1/2} \right) \leq 2k|\Delta|^C. \quad (2.42)$$

Proof: Let \mathbf{a} be the median of $\sup_{u \in \Delta} (\eta_1(u) - \eta_1(d))$. It follows from [1, Lemma 5.4.3] that,

$$P \left(\sup_{u \in \Delta} |\eta_1(u) - \eta_1(d)| \geq \mathbf{a} + \tilde{\sigma}(|\Delta|)t \right) \leq 2e^{-t^2/2}. \quad (2.43)$$

Since by [1, (7.113)],

$$\mathbf{a} = o(\tilde{\sigma}^2(|\Delta|) \log 1/|\Delta|^{1/2}), \quad (2.44)$$

we see that

$$P \left(\sup_{u \in \Delta} |\eta_1(u) - \eta_1(d)| \geq ((1 + 2C)\tilde{\sigma}^2(|\Delta|) \log 1/|\Delta|^{1/2}) \right) \leq 2|\Delta|^C, \quad (2.45)$$

which gives (2.42). \square

Remark 2.1 When η has stationary increments, $\sigma^2(u, v)$ in (1.6) can be written as $\sigma^2(u - v)$. For $x \in [0, \delta]$ define,

$$\bar{\sigma}^2(x) = \mu (s : \sigma^2(x) \leq x),$$

where μ is Lebesgue measure. Clearly, $\bar{\sigma}^2(x)$ is an increasing function. It is called the increasing rearrangement of $\sigma^2(x)$. (See [1, Section 6,4] for more details.) We show in [1, (6.138)] that when η is continuous a.s.,

$$\lim_{x \rightarrow 0} \bar{\sigma}^2(x) \log(1/x) = 0.$$

This shows that if $\tilde{\sigma}^2(x)$ is asymptotic to an increasing function at 0, then (1.8), which implies that η is continuous a.s., implies (1.7).

References

- [1] M. B. Marcus and J. Rosen, *Markov Processes, Gaussian Processes and Local Times*, Cambridge University Press, New York, (2006).
- [2] M. B. Marcus and J. Rosen, *Asymptotic Properties of Permenental Sequences*, Springer Briefs in Prob. and Math. Stat., Switzerland, (2021).
- [3] P. Revesz, *Random Walk in Random and Non-random Environments*, Cambridge University Press, New York, (1990).

Michael B. Marcus
 253 West 73rd. St., Apt. 2E
 New York, NY 10023, USA
 mbmarcus@optonline.net

Jay Rosen
 Department of Mathematics
 College of Staten Island, CUNY
 Staten Island, NY 10314, USA
 jrosen30@optimum.net