

A generalization of van der Corput's difference theorem with applications to recurrence and multiple ergodic averages

SOHAIL FARHANGI

*Department of mathematics and computer science, University of Adam Mickiewicz,
Poznań, 61-712, Greater Poland Voivodeship, Poland
(e-mail: sohail.farhangi@gmail.com)*

Abstract. We prove a generalization of van der Corput's difference theorem for sequences of vectors in a Hilbert space. We use this generalization to obtain applications regarding recurrence and multiple ergodic averages of noncommutative measure preserving group actions when certain rigidity and ergodicity assumptions are satisfied.

Key words: van der Corput, recurrence, noncommutative ergodic theory

2020 Mathematics Subject Classification: 47A35 (Primary), 37B20 (Secondary)

1. Introduction

In [22] van der Corput proved Theorem 1.1 which is now known as van der Corput's Difference Theorem (henceforth abbreviated as vdCDT).

THEOREM 1.1 Theorem 1.3.1 in [19]. *If $(x_n)_{n=1}^{\infty} \subseteq [0, 1]$ is a sequence for which $(x_{n+h} - x_n)_{n=1}^{\infty} \pmod{1}$ is uniformly distributed for all $h \in \mathbb{N}$, then $(x_n)_{n=1}^{\infty}$ is uniformly distributed.*

In Ergodic Theory, the following Hilbertian analogues of Theorem 1.1 were introduced by Bergelson in [2] and are of great use.

THEOREM 1.2. *Let \mathcal{H} be a Hilbert space and $(x_n)_{n=1}^{\infty} \subseteq \mathcal{H}$ a bounded sequence of vectors.*

(i) *If for every $h \in \mathbb{N}$ we have*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \langle x_{n+h}, x_n \rangle = 0, \text{ then } \lim_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=1}^N x_n \right\| = 0. \quad (1.1)$$

(ii) If

$$\lim_{h \rightarrow \infty} \limsup_{N \rightarrow \infty} \left| \frac{1}{N} \sum_{n=1}^N \langle x_{n+h}, x_n \rangle \right| = 0, \text{ then } \lim_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=1}^N x_n \right\| = 0. \quad (1.2)$$

(iii) If

$$\lim_{H \rightarrow \infty} \frac{1}{H} \sum_{h=1}^H \limsup_{N \rightarrow \infty} \left| \frac{1}{N} \sum_{n=1}^N \langle x_{n+h}, x_n \rangle \right| = 0, \text{ then } \lim_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=1}^N x_n \right\| = 0. \quad (1.3)$$

See [3], [7], [17], [12], and [18] for some examples of applications of parts (i), (ii), and (iii) of Theorem 1.2 in Ergodic Theory, and see [9] for a survey of modern developments regarding vdCDT and its many variations. One of the main results of this paper is to prove a generalization of Theorem 1.2(ii) as Theorem 2.7, which we do not state here since we have not yet discussed the prerequisite terminology. We then deduce Corollary 2.8, which is a generalization of Theorem 1.2(i) that we will use to obtain applications in the theory of recurrence as Theorems 1.5 and 1.7. We will also use Corollary 2.8 to obtain applications in ergodic theory by showing that certain multiple ergodic averages converge to their expected value in Theorems 1.8 and 1.9.

We will assume throughout this paper that if (X, \mathcal{B}, μ) is a probability space and $R : X \rightarrow X$ is a measure preserving transformation, then R is invertible. We let \mathcal{I}_R be the σ -algebra of R -invariant sets, and $\mathcal{K}_{\text{rat}}(R)$ the σ -algebra of the rational Kronecker factor of R . Let us now recall a result of Frantzikinakis which is related to our applications.

THEOREM 1.3 (Corollary 1.7 in [14]). *Let $a : \mathbb{R}_+ \rightarrow \mathbb{R}$ be a Hardy field function for which there exist some $\epsilon > 0$ and $d \in \mathbb{Z}_+$ satisfying*

$$\lim_{n \rightarrow \infty} \frac{a(n)}{t^{d+\epsilon}} = \lim_{n \rightarrow \infty} \frac{t^{d+1}}{a(n)} = \infty. \quad (1.4)$$

Furthermore, let (X, \mathcal{B}, μ) be a probability space and $T, S : X \rightarrow X$ be measure preserving transformations. Suppose that the system (X, \mathcal{B}, μ, T) has zero entropy. Then

(i) *For every $f, g \in L^\infty(X, \mu)$ we have*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f \cdot S^{\lfloor a(n) \rfloor} g = \mathbb{E}[f | \mathcal{I}_T] \cdot \mathbb{E}[g | \mathcal{I}_S], \quad (1.5)$$

where the limit is taken in $L^2(X, \mu)$.

(ii) *For every $A \in \mathcal{B}$ we have*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mu(A \cap T^{-n} A \cap S^{-\lfloor a(n) \rfloor} A) \geq \mu(A)^3. \quad (1.6)$$

To understand the significance of Theorem 1.3, we observe that in Theorem 1.1. of [8] it is shown that for any finitely generated solvable group G of exponential growth*, there exists a measure preserving system $(X, \mathcal{B}, \mu, (T_g)_{g \in G})$, $A \in \mathcal{B}$ with $\mu(A) > 0$, and $g_1, g_2 \in G$ for which $\mu(T_{g_1}^n A \cap T_{g_2}^n A) = 0$ for all $n \in \mathbb{N}$. While the previous result shows that general recurrence results do not extend to solvable groups, Theorem 1.3 shows that some recurrence results can still be obtained when commutativity assumptions are replaced with structural assumptions. It is worth noting that the assumptions of Theorem 1.3 require $a(n)$ to be sufficiently far from polynomial sequence. An analogous result for polynomial sequences was conjectured by Frantzikinakis in [14], then obtained by Frantzikinakis and Host in [15].

THEOREM 1.4. *Let (X, \mathcal{B}, μ) be a probability space and let $T, S : X \rightarrow X$ be measure preserving transformations. Suppose that the m.p.s. (X, \mathcal{B}, μ, T) has zero entropy and $f, g \in L^\infty(X, \mu)$.*

(i) *If $p \in \mathbb{Z}[x]$ is a polynomial of degree at least 2, then*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f \cdot S^{p(n)} g \tag{1.7}$$

converges in $L^2(X, \mu)$, and is equal to 0 if either $\mathbb{E}[f | \mathcal{K}_{rat}(T)] = 0$ or $\mathbb{E}[g | \mathcal{K}_{rat}(S)] = 0$.

(ii) *For every $A \in \mathcal{B}$ with $\mu(A) > 0$ and $\epsilon > 0$, the set*

$$\left\{ n \in \mathbb{N} \mid \mu(A \cap T^{-n} A \cap S^{-p(n)} A) > \mu(A)^3 - \epsilon \right\} \tag{1.8}$$

is syndetic.

In this paper we will also obtain many results regarding ergodic averages with non-commuting transformations. On the one hand, our results are weaker than those of Frantzikinakis and Host since we have to restrict ourselves to the class of rigid systems, a proper subclass of zero entropy systems[†], but on the other hand, our methods are flexible enough to apply to a wider class of sequences and to situations in which there are more than two transformations.

Definition 1.1. A measure preserving system (X, \mathcal{B}, μ, T) is **rigid** if there exists an increasing sequence $(k_m)_{m=1}^\infty \subseteq \mathbb{N}$ for which $\lim_{m \rightarrow \infty} \|U_T^{k_m} f - f\| = 0$ for all $f \in L^2(X, \mu)$.

The class of rigid measure preserving systems is rather large. In Proposition 2.9 of [5] it is shown that for a standard probability space (X, \mathcal{B}, μ) , the set \mathcal{S} of measure preserving transformations $S : X \rightarrow X$ for which (X, \mathcal{B}, μ, S) is rigid and weakly mixing simultaneously, is a generic set within the set of all measure preserving transformations of (X, \mathcal{B}, μ) .

*A finitely generated solvable group has exponential growth if and only if it contains no nilpotent subgroups of finite index, so the assumption of exponential growth ensures that the group G is not virtually nilpotent.

[†]See Remark 3.6 for a brief discussion about the limitations of our methods.

THEOREM 1.5 (cf. Theorem 3.2). *Let (X, \mathcal{B}, μ) be a probability space and let $T, S : X \rightarrow X$ be measure preserving transformations. Suppose that the m.p.s. (X, \mathcal{B}, μ, T) is rigid, and that the m.p.s. (X, \mathcal{B}, μ, S) is totally ergodic. Let $(k_n)_{n=1}^{\infty} \subseteq \mathbb{N}$ be a sequence for which $((k_{n+h} - k_n)\alpha \pmod{1})_{n=1}^{\infty}$ is uniformly distributed for all $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $h \in \mathbb{N}$.*

(i) *For any $f, g \in L^{\infty}(X, \mu)$ we have*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f \cdot S^{k_n} g = \mathbb{E}[f | \mathcal{I}_T] \int_X g d\mu, \quad (1.9)$$

with convergence taking place in $L^2(X, \mu)$.

(ii) *If $A \in \mathcal{B}$ then*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mu(A \cap T^{-n} A \cap S^{-k_n} A) \geq \mu(A)^3. \quad (1.10)$$

(iii) *If $((k_{n+h} - k_n)\alpha)_{n=1}^{\infty}$ is uniformly distributed in its orbit closure for all $\alpha \in \mathbb{R}$ then (i) and (ii) hold even when (X, \mathcal{B}, μ, S) is not ergodic.*

We see that Theorem 1.5 applies when $k_n = p(n)$ and $p : \mathbb{Z} \rightarrow \mathbb{Z}$ is a polynomial of degree at least 2, which gives a positive answer to Question 1.4 when S is totally ergodic and T is rigid. The assumption of total ergodicity on S is mainly used to simplify the proof and to obtain the expected value of the averages in consideration. We also outline the arguments needed to show that the limits in Question 1.4 exist without the assumption of total ergodicity on S in Remark 3.3. We may take $k_n = \lfloor a(n) \rfloor$ with $a(n)$ as in Theorem 1.3 or $k_n = \lfloor n^2 \log^2(n) \rfloor$ as examples satisfying Theorem 1.5(iii). We note that Theorem 1.3 does not apply when $a(n) = n^2 \log^2(n)$, but the remark after Corollary 1.7 of [14] shows that a variation of Theorem 1.3(ii) applies in this case.

Let us now recall a result of Frantzikinakis, Lesigne, and Wierdl that is related to our next main result, Theorem 1.7.

THEOREM 1.6 (Theorem 1.4 and Corollary 4.4 of [17]). *Let $k \geq 2$ be an integer and $\alpha \in \mathbb{R}$ be irrational. Let $R = \{n \in \mathbb{N} \mid n^k \alpha \in [\frac{1}{4}, \frac{3}{4}]\}$.*

(i) *If (X, \mathcal{B}, μ) is a probability space and $T_1, T_2, \dots, T_{k-1} : X \rightarrow X$ are commuting measure preserving transformations, then for any $A \in \mathcal{B}$ with $\mu(A) > 0$, there exists $n \in R$ for which*

$$\mu(A \cap T_1^{-n} A \cap T_2^{-n} A \cap \dots \cap T_{k-1}^{-n} A) > 0. \quad (1.11)$$

(ii) *There exists a m.p.s. (X, \mathcal{B}, μ, T) and a set $A \in \mathcal{B}$ satisfying $\mu(A) > 0$ such that for all $n \in R$ we have*

$$\mu(A \cap T^{-n} A \cap T^{-2n} A \cap \dots \cap T^{-kn} A) > 0. \quad (1.12)$$

Our next theorem strengthens the conclusion of Theorem 1.6(i).

THEOREM 1.7 (cf. Theorem 3.4). *Let $k \geq 2$ be an integer and $\alpha \in \mathbb{R}$ be irrational. Let $R = \{n \in \mathbb{N} \mid n^k \alpha \in [\frac{1}{4}, \frac{3}{4}]\}$. Let (X, \mathcal{B}, μ) be a probability space and $T_1, T_2, \dots, T_{k-1} : X \rightarrow X$ commuting measure preserving transformations. Let $S : X \rightarrow X$ be an measure preserving transformation for which (X, \mathcal{B}, μ, S) is rigid, and $\{S, T_1, T_2, \dots, T_{k-1}\}$ generate a nilpotent group. For any $A \in \mathcal{B}$ with $\mu(A) > 0$, there exists $n \in R$ for which*

$$\mu(A \cap S^{-n} A \cap T_1^{-n} A \cap T_2^{-n} A \cap \dots \cap T_{k-1}^{-n} A) > 0. \quad (1.13)$$

It is worth noting that the example of a m.p.s. (X, \mathcal{B}, μ, T) given in [17] satisfying Theorem 1.6(ii) is also a uniquely ergodic minimal topological dynamical system with zero topological entropy, and A can be taken to be an open set. Consequently, the analogue of Theorem 1.7 in which (X, \mathcal{B}, μ, S) has zero entropy is not true. However, our assumption that $\{S, T_1, T_2, \dots, T_{k-1}\}$ generate a nilpotent group is an artifact of our method of proof, and it is currently not clear as to whether or not this assumption can be relaxed.

Before stating our applications regarding multiple ergodic averages, we must define some terminology. A polynomial $p \in \mathbb{Q}[x]$ is an *integer polynomial* if for every integer x , $p(x)$ is also an integer. A collection of polynomials $p_1, \dots, p_K \in \mathbb{Q}[x]$ is *independent* if any nontrivial rational linear combination of them is not constant, and it is *strongly independent* if any nontrivial rational linear combination of them is not linear.

THEOREM 1.8 (cf. Theorem 3.7). *Let (X, \mathcal{B}, μ) be a probability space and $T, S : X \rightarrow X$ be measure preserving transformations. Suppose that (X, \mathcal{B}, μ, T) is rigid and (X, \mathcal{B}, μ, S) is totally ergodic. Let $p_1, \dots, p_K \in \mathbb{Q}[x]$ be a collection of integer polynomials such that $\{p_1(n+h) - p_1(n), p_2(n+h) - p_1(n) \dots, p_K(n+h) - p_1(n), p_2(n) - p_1(n), \dots, p_K(n) - p_1(n)\}$ is independent for all $h \in \mathbb{N}$. For any $f_0, f_1, \dots, f_K \in L^\infty(X, \mu)$ and $c \in \mathbb{Z} \setminus \{0\}$, we have*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f_0 \prod_{i=1}^K S^{p_i(n)} f_i = \mathbb{E}[f_0 | \mathcal{I}_T] \prod_{i=1}^K \int_X f_i d\mu, \quad (1.14)$$

with convergence taking place in $L^2(X, \mu)$.

THEOREM 1.9 (cf. Theorem 3.9). *Let (X, \mathcal{B}, μ) be a probability space and $T, S_1, S_2 : X \rightarrow X$ be measure preserving transformations. Suppose that (X, \mathcal{B}, μ, T) is rigid and $(X, \mathcal{B}, \mu, S_2)$ is weakly mixing. Let $p : \mathbb{R} \rightarrow \mathbb{R}$ either be a polynomial of degree at least 2, or of the form $p(n) = n^\alpha \log(n)^\beta$ with $\alpha > 1$ and $\beta \leq 0$.[‡] For any $f_0, f_1, f_2 \in L^\infty(X, \mu)$ satisfying $\int_X f_2 d\mu = 0$, we have*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f_0 S_1^n f_1 S_2^{\lfloor p(n) \rfloor} f_2 = 0, \quad (1.15)$$

with convergence taking place in $L^2(X, \mu)$.

[‡]The astute reader will observe that p could be any function for which $(g_1(n), g_2(n), g_3(n), g_4(n)) = (p(n+\ell+h), p(n+\ell), p(n+h), p(n))$ satisfies the hypothesis of Theorem A of [6], and that we have picked two aesthetic examples of such functions for the sake of simplicity.

REMARK 1.10. Firstly, we will see as a consequence of our method of proof that Theorems 1.5, 1.7, 1.8, and 1.9 do not need (X, \mathcal{B}, μ, T) to be rigid, but that they only need T to act rigidly on the function f ($f = \mathbb{1}_A$ when dealing with recurrence) at hand, i.e., that $\|T^{n_k} f - f\|_2 \xrightarrow[k \rightarrow \infty]{} 0$ for some sequence $(n_k)_{k=1}^\infty \subseteq \mathbb{N}$, which is equivalent to f coming from a rigid factor of (X, \mathcal{B}, μ, T) . Furthermore, it is clear that the aforementioned results are linear with respect to the first term, so we may further assume without loss of generality that f is in the L^2 -closure of linear combinations of functions coming from (potentially varying) rigid factors of (X, \mathcal{B}, μ, T) . Secondly, we average over the Følner sequence $\{[1, N]\}_{N=1}^\infty$ solely for the sake of convenience, and our results still hold with the same method of proof when averaging over any other Følner sequence $\{[a_n, b_n]\}_{n=1}^\infty$.

2. Van der Corput's difference theorem and the ergodic hierarchy of mixing properties

The contents of this section are a subset of the contents of Chapters 2.2 and 2.3 of [13] that we need to prove Corollary 2.8, which we will use in the proofs of Theorems 1.5, 1.7, 1.8, and 1.9.

Let \mathcal{H} be a Hilbert space. In this section we will discuss how to construct a Hilbert space \mathcal{H} out of sequences of vectors coming from \mathcal{H} . We will then use \mathcal{H} to prove our generalization of van der Corput's difference theorem for sequences of vectors coming from \mathcal{H} .

Let $\|\cdot\|$ and $\langle \cdot, \cdot \rangle$ denote the norm and inner product on \mathcal{H} and let $\|\cdot\|_{\mathcal{H}}$ and $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ denote the norm and inner product on \mathcal{H} . We denote the collection of square averageable sequences by

$$SA(\mathcal{H}) := \{(f_n)_{n=1}^\infty \subseteq \mathcal{H} \mid \limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \|f_n\|^2 < \infty\}. \quad (2.1)$$

Let $(f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty \in SA(\mathcal{H})$ and observe that

$$\begin{aligned} \limsup_{N \rightarrow \infty} \frac{1}{N} \left| \sum_{n=1}^N \langle f_n, g_n \rangle \right| &\leq \limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \|f_n\| \cdot \|g_n\| \\ &\leq \left(\limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \|f_n\|^2 \right)^{\frac{1}{2}} \left(\limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \|g_n\|^2 \right)^{\frac{1}{2}} < \infty. \end{aligned} \quad (2.2)$$

It follows that we may use diagonalization to construct an increasing sequence of positive integers $(N_q)_{q=1}^\infty$ for which

$$\lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \langle x_{n+h}, y_n \rangle \quad (2.3)$$

exists whenever $(x_n)_{n=1}^\infty, (y_n)_{n=1}^\infty \in \{(f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty\}$ and $h \in \mathbb{N}$. We now construct a new Hilbert space $\mathcal{H} = \mathcal{H}((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$ from $(f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty$ and $(N_q)_{q=1}^\infty$ as follows. For all $(x_n)_{n=1}^\infty, (y_n)_{n=1}^\infty \in \{(f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty\}$ and $h \in \mathbb{N}$, we define

$$\langle (x_{n+h})_{n=1}^\infty, (y_n)_{n=1}^\infty \rangle_{\mathcal{H}} = \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \langle x_{n+h}, y_n \rangle. \quad (2.4)$$

We see that $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ is a sesquilinear form on $\mathcal{H}' = \text{Span}_{\mathbb{C}}(\{(f_{n+h})_{n=1}^\infty\}_{h=0}^\infty \cup \{(g_{n+h})_{n=1}^\infty\}_{h=0}^\infty)$ with scalar multiplication and addition occurring pointwise. Letting

$$\mathcal{H}'' = \{(e_n)_{n=1}^\infty \in SA(\mathcal{H}) \mid \forall \epsilon > 0 \exists (h_n(\epsilon))_{n=1}^\infty \in \mathcal{H}' \text{ s.t.} \quad (2.5)$$

$$\limsup_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|e_n - h_n(\epsilon)\|^2 < \epsilon\}, \text{ and}$$

$$S = \{(x_n)_{n=1}^\infty \in \mathcal{H}'' \mid \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|x_n\|^2 = 0\}, \quad (2.6)$$

we see that \mathcal{H}''/S is a pre-Hilbert space. We will soon see that \mathcal{H}'' is sequentially closed under the topology induced by $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ (cf. Theorem 2.1), so we define $\mathcal{H}((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty) = \mathcal{H}''/S$. We call $\mathcal{H}((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$ the Hilbert space induced by $((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$, and we may write \mathcal{H} in place of $\mathcal{H}((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$ if $((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$ is understood from the context.

For $(f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty \in SA(\mathcal{H})$ and $(N_q)_{q=1}^\infty \subseteq \mathbb{N}$ we say that $((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$ is a **permissible triple** if $\mathcal{H}((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$ is well defined. We say that $((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$ is a **weakly permissible triple** if

$$\lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1+|h|}^{N_q} \langle f_{n+h}, g_n \rangle \quad (2.7)$$

exists for every $h \in \mathbb{Z}$. Given $(x_n)_{n=1}^\infty \subseteq \mathcal{H}$ for which $(x_n)_{n=1}^\infty \in \mathcal{H}''$, we may view $(x_n)_{n=1}^\infty$ as an element of \mathcal{H} by identifying $(x_n)_{n=1}^\infty$ with its equivalence class in \mathcal{H}''/S . We will now show that \mathcal{H} is a Hilbert space by verifying that it is complete.

THEOREM 2.1. *Let \mathcal{H} be a Hilbert space and $(f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty \in SA(\mathcal{H})$. Let $((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$ be a permissible triple and $\mathcal{H} = \mathcal{H}((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$. If $\{(\xi_{n,m})_{n=1}^\infty\}_{m=1}^\infty \subseteq \mathcal{H}''$ is a Cauchy sequence with respect to the metric induced by $\|\cdot\|_{\mathcal{H}}$, then there exists $(\xi_n)_{n=1}^\infty \in \mathcal{H}''$ for which*

$$\lim_{m \rightarrow \infty} \left(\lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|\xi_{n,m} - \xi_n\|^2 \right) = 0. \quad (2.8)$$

In particular, \mathcal{H} is a Hilbert space.

Proof. We proceed by modifying the proof of the main result in section §2 of chapter II of [10]. Let $(\epsilon_m)_{m=1}^\infty$ be a sequence of real numbers tending to 0 for which

$$\lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|\xi_{n,m} - \xi_{n,k}\|^2 < \epsilon_m \quad (2.9)$$

whenever $k \geq m$. By induction, let $T_0 = N_0 = 0$ and let $(T_m)_{m=1}^\infty \subseteq \mathbb{N}$ be such that conditions (i)-(iii) below hold.

(i) For every $m \geq 1$, every $k \geq m$, and every $T \geq T_k$

$$\frac{1}{N_T} \sum_{n=1}^{N_T} \|\xi_{n,k} - \xi_{n,m}\|^2 < \epsilon_m. \quad (2.10)$$

(ii) For every $m \geq 1$ and every $k \geq m$

$$\frac{1}{N_{T_k} - N_{T_{k-1}}} \sum_{n=N_{T_{k-1}}+1}^{N_{T_k}} \|\xi_{n,k} - \xi_{n,m}\|^2 < \epsilon_m. \quad (2.11)$$

(iii) For every $m \geq 1$

$$\frac{1}{N_{T_m}} \sum_{j=1}^{m-1} \sum_{n=N_{T_{j-1}}+1}^{N_{T_j}} \|\xi_{n,j} - \xi_{n,m}\|^2 < \epsilon_m. \quad (2.12)$$

Now let us define $(\xi_n)_{n=1}^\infty$ by $\xi_n = \xi_{n,m}$ where m is such that $N_{T_{m-1}} < n \leq N_{T_m}$. To conclude the proof, we note that for $m \geq 1$, $k > m$, and $T_{k-1} < T \leq T_k$ we have

$$\begin{aligned} & \sum_{n=1}^{N_T} \|\xi_{n,m} - \xi_n\|^2 \quad (2.13) \\ &= \sum_{j=1}^{m-1} \sum_{n=N_{T_{j-1}}+1}^{N_{T_j}} \|\xi_{n,j} - \xi_{n,m}\|^2 + \sum_{j=m}^{k-1} \sum_{n=N_{T_{j-1}}+1}^{N_{T_j}} \|\xi_{n,m} - \xi_n\|^2 + \sum_{n=N_{T_{k-1}}+1}^{N_T} \|\xi_{n,m} - \xi_n\|^2 \\ &\leq N_{T_m} \epsilon_m + \sum_{j=m}^{k-1} (N_{T_j} - N_{T_{j-1}}) \epsilon_m + \sum_{n=1}^{N_T} \|\xi_{n,k} - \xi_{n,m}\|^2 \\ &\leq N_{T_{k-1}} \epsilon_m + N_T \epsilon_m \leq 2N_T \epsilon_m. \end{aligned}$$

□

Definition 2.1. Let \mathcal{H} be a Hilbert space, $U : \mathcal{H} \rightarrow \mathcal{H}$ a unitary operator, $\xi \in \mathcal{H}$, and $(f_n)_{n=1}^\infty \in \text{SA}(\mathcal{H})$.

- (i) ξ is a **strongly mixing element** of (\mathcal{H}, U) if for any $g \in \mathcal{H}$ we have $\lim_{n \rightarrow \infty} \langle U^n \xi, g \rangle = 0$.
- (ii) ξ is a **rigid element** of (\mathcal{H}, U) if there exists an increasing sequence $(k_m)_{m=1}^\infty \subseteq \mathbb{N}$ for which $\lim_{m \rightarrow \infty} \|U^{k_m} \xi - \xi\| = 0$.

- (iii) $(f_n)_{n=1}^\infty$ is a **nearly strongly mixing sequence** if for any permissible triple of the form $((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$ we have

$$\lim_{h \rightarrow \infty} \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \langle f_{n+h}, g_n \rangle = 0. \quad (2.14)$$

- (iv) $(f_n)_{n=1}^\infty$ is a **rigid sequence** if for any permissible triple of the form $((f_n)_{n=1}^\infty, (f_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$ there exists an increasing sequence $(k_m)_{m=1}^\infty \subseteq \mathbb{N}$ for which

$$\lim_{m \rightarrow \infty} \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|f_{n+k_m} - f_n\|^2 = 0. \quad (2.15)$$

REMARK 2.2. Given a Hilbert space \mathcal{H} and a sequence $(f_n)_{n=1}^\infty \in SA(\mathcal{H})$ we may define a left shift operator S given by $S(f_n)_{n=1}^\infty = (f_{n+1})_{n=1}^\infty$, and we note that S is naturally identified with a unitary operator on any Hilbert space \mathcal{H} induced by a permissible triple from \mathcal{H} . We now see that $(f_n)_{n=1}^\infty$ is a nearly strongly mixing sequence if and only if for any permissible triple of the form $((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$, $(f_n)_{n=1}^\infty$ is a strongly mixing element of (\mathcal{H}, S) , where $\mathcal{H} = \mathcal{H}((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$. Similarly, we see that $(f_n)_{n=1}^\infty$ is a rigid sequence if and only if for any permissible triple of the form $((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$, $(f_n)_{n=1}^\infty$ is a rigid element of (\mathcal{H}, S) . It is also worth mentioning that we choose to use the terminology of “nearly strongly mixing sequences” instead of “strongly mixing sequences” because the latter term was already defined in [1]. A comparison between the notions of nearly strongly mixing sequences and strongly mixing sequences is given in Chapter 2.6 of [13].

LEMMA 2.3 (cf. Lemma 2.3.1 of [13]). *Let \mathcal{H} be a Hilbert space.*

- (i) *If $(f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty \in SA(\mathcal{H})$ are nearly strongly mixing sequences, then all*

$$(x_n)_{n=1}^\infty \in \text{Span}_{\mathbb{C}}(\{(f_{n+h})_{n=1}^\infty\}_{h=0}^\infty \cup \{(g_{n+h})_{n=1}^\infty\}_{h=0}^\infty) \quad (2.16)$$

are also nearly strongly mixing sequences.

- (ii) *Let $(f_n)_{n=1}^\infty \in SA(\mathcal{H})$ and let $\{(g_{n,m})_{n=1}^\infty\}_{m=1}^\infty \subseteq SA(\mathcal{H})$ be a family of sequences for which*

$$\lim_{m \rightarrow \infty} \left(\limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \|f_n - g_{n,m}\|^2 \right) = 0. \quad (2.17)$$

If $(g_{n,m})_{n=1}^\infty$ is a nearly strongly mixing sequence for all $m \in \mathbb{N}$, then $(f_n)_{n=1}^\infty$ is also a nearly strongly mixing sequence.

LEMMA 2.4. Let \mathcal{H} be a Hilbert space and $(f_n)_{n=1}^\infty \in SA(\mathcal{H})$ a nearly strongly mixing sequence. If $(g_n)_{n=1}^\infty \in SA(\mathcal{H})$ is a rigid sequence, then

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \langle f_n, g_n \rangle = 0. \quad (2.18)$$

Proof. Let $(M_q)_{q=1}^\infty \subseteq \mathbb{N}$ be any increasing sequence for which

$$\lim_{q \rightarrow \infty} \frac{1}{M_q} \sum_{n=1}^{M_q} \langle f_n, g_n \rangle \quad (2.19)$$

exists. Let $(N_q)_{q=1}^\infty$ be a subsequence of $(M_q)_{q=1}^\infty$ for which $((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$ is a permissible triple and let $\mathcal{H} = \mathcal{H}((f_n)_{n=1}^\infty, (g_n)_{n=1}^\infty, (N_q)_{q=1}^\infty)$. Let $\epsilon > 0$ be arbitrary, and let $k \in \mathbb{N}$ be such that

$$\lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|g_{n+k} - g_n\|^2 < \epsilon, \text{ and} \quad (2.20)$$

$$\lim_{q \rightarrow \infty} \left| \frac{1}{N_q} \sum_{n=1}^{N_q} \langle f_{n+k}, g_n \rangle \right| < \epsilon. \quad (2.21)$$

We now see that

$$\begin{aligned} & \lim_{q \rightarrow \infty} \left| \frac{1}{M_q} \sum_{n=1}^{M_q} \langle f_n, g_n \rangle \right| = \lim_{q \rightarrow \infty} \left| \frac{1}{N_q} \sum_{n=1}^{N_q} \langle f_n, g_n \rangle \right| \quad (2.22) \\ & \leq \lim_{q \rightarrow \infty} \left| \frac{1}{N_q} \sum_{n=k+1}^{N_q} \langle f_n, g_n - g_{n-k} \rangle \right| + \lim_{q \rightarrow \infty} \left| \frac{1}{N_q} \sum_{n=k+1}^{N_q} \langle f_n, g_{n-k} \rangle \right| \\ & \leq \left(\lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=k+1}^{N_q} \|f_n\|^2 \right)^{\frac{1}{2}} \left(\lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|g_{n+k} - g_n\|^2 \right)^{\frac{1}{2}} + \lim_{q \rightarrow \infty} \left| \frac{1}{N_q} \sum_{n=1}^{N_q} \langle f_{n+k}, g_n \rangle \right| \\ & \leq \sqrt{\epsilon} \left(\limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \|f_n\|^2 \right)^{\frac{1}{2}} + \epsilon \end{aligned}$$

□

LEMMA 2.5. Let (X, \mathcal{B}, μ) be a probability space, $\mathcal{H} = L^2(X, \mu)$, and $(f_n)_{n=1}^\infty \in SA(\mathcal{H})$ a nearly strongly mixing sequence. If $(g_n)_{n=1}^\infty \in SA(\mathcal{H})$ is a bounded rigid sequence, then

$$\lim_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=1}^N f_n g_n \right\| = 0. \quad (2.23)$$

Proof. Let us assume for the sake of contradiction that for some $\epsilon > 0$ and $(N_q)_{q=1}^\infty \subseteq \mathbb{N}$ we have

$$\lim_{q \rightarrow \infty} \left\| \frac{1}{N_q} \sum_{n=1}^{N_q} f_n g_n \right\| \geq \epsilon. \quad (2.24)$$

By passing to a subsequence of $(N_q)_{q=1}^\infty$ if necessary, we may assume without loss of generality that

$$\lim_{q \rightarrow \infty} \left(\left\| \frac{1}{N_q} \sum_{n=1}^{N_{q-1}} f_n g_n \right\| + \frac{N_{q-1}}{N_q} \right) = 0. \quad (2.25)$$

For $q \in \mathbb{N}$ let

$$\xi'_q = \frac{1}{N_{q+1}} \sum_{n=N_{q+1}}^{N_{q+1}} f_n g_n \text{ and } \xi_q = \frac{\xi'_q}{\|\xi'_q\|}. \quad (2.26)$$

Now consider the sequence $(G_n)_{n=1}^\infty \in \text{SA}(\mathcal{H})$ given by $G_n = \xi_q$ for $N_q < n \leq N_{q+1}$. Since

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \|G_{n+1} - G_n\| = 0, \quad (2.27)$$

we see that $(G_n)_{n=1}^\infty$ is a rigid sequence. Next, we will verify that $(\overline{g_n} G_n)_{n=1}^\infty$ is a rigid sequence. Let $(k_m)_{m=1}^\infty \subseteq \mathbb{N}$ be such that

$$\lim_{m \rightarrow \infty} \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|g_{n+k_m} - g_n\|^2 = 0. \quad (2.28)$$

Letting $C = \sup_{n \in \mathbb{N}} (\|\overline{g_n}\|^2 + \|G_n\|^2)$, we see that

$$\begin{aligned} & \frac{1}{2} \lim_{m \rightarrow \infty} \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|\overline{g_{n+k_m}} G_{n+k_m} - \overline{g_n} G_n\|^2 \\ & \leq \lim_{m \rightarrow \infty} \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|\overline{g_{n+k_m}} G_{n+k_m} - \overline{g_{n+k_m}} G_n\|^2 + \lim_{m \rightarrow \infty} \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|\overline{g_{n+k_m}} G_n - \overline{g_n} G_n\|^2 \\ & \leq \lim_{m \rightarrow \infty} \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|\overline{g_{n+k_m}}\|^2 \|G_{n+k_m} - G_n\|^2 + \lim_{m \rightarrow \infty} \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|\overline{g_{n+k_m}} - \overline{g_n}\|^2 \|G_n\|^2 \\ & \leq C \lim_{m \rightarrow \infty} \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|G_{n+k_m} - G_n\|^2 + C \lim_{m \rightarrow \infty} \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \|\overline{g_{n+k_m}} - \overline{g_n}\|^2 = 0. \end{aligned} \quad (2.29)$$

Lastly, it suffices to observe that

$$\begin{aligned}
\lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \langle f_n, \overline{g_n} G_n \rangle &= \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \langle f_n g_n, G_n \rangle \\
&= \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=N_{q-1}+1}^{N_q} \langle f_n g_n, \xi_{q-1} \rangle = \lim_{q \rightarrow \infty} \|\xi'_{q-1}\| \geq \epsilon,
\end{aligned} \tag{2.30}$$

which contradicts Lemma 2.4. \square

It is worth noting that a similar method of proof can be used to show Lemma 2.6, which verifies that Theorem 2.7 and Corollary 2.8 are indeed generalizations of Theorems 1.2(ii) and 1.2(i) respectively. We only used that fact that $\mathcal{H} = L^2(X, \mu)$ so that multiplication would be well defined, which is necessary for our applications in Section 3.

LEMMA 2.6 (cf. Lemma 2.3.1(iv) or Lemma 3.2.7 of [13]). *Let \mathcal{H} be a Hilbert space. If $(f_n)_{n=1}^{\infty} \in SA(\mathcal{H})$ is a nearly strongly mixing sequence, then*

$$\lim_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=1}^N f_n \right\| = 0. \tag{2.31}$$

THEOREM 2.7. *Let \mathcal{H} be a Hilbert space and $(f_n)_{n=1}^{\infty} \in SA(\mathcal{H})$. If*

$$\lim_{h \rightarrow \infty} \limsup_{N \rightarrow \infty} \left| \sum_{n=1}^N \langle f_{n+h}, f_n \rangle \right| = 0, \tag{2.32}$$

then $(f_n)_{n=1}^{\infty}$ is a nearly strongly mixing sequence.

Proof. Let $(g_n)_{n=1}^{\infty} \subseteq \mathcal{H}$ and $(N_q)_{q=1}^{\infty} \subseteq \mathbb{N}$ be such that $((f_n)_{n=1}^{\infty}, (g_n)_{n=1}^{\infty}, (N_q)_{q=1}^{\infty})$ is a weakly permissible triple. By passing to a subsequence of $(N_q)_{q=1}^{\infty}$ if necessary, we may assume without loss of generality $((f_n)_{n=1}^{\infty}, (g_n)_{n=1}^{\infty}, (N_q)_{q=1}^{\infty})$ is a permissible triple, so may define $\mathcal{H} = \mathcal{H}((f_n)_{n=1}^{\infty}, (g_n)_{n=1}^{\infty}, (N_q)_{q=1}^{\infty})$. Let $S : \mathcal{H} \rightarrow \mathcal{H}$ denote the unitary operator induced by the left shift. From equation (2.32) we see that

$$\lim_{h \rightarrow \infty} \langle S^h((f_n)_{n=1}^{\infty}), (f_n)_{n=1}^{\infty} \rangle_{\mathcal{H}} = 0, \tag{2.33}$$

so $(f_n)_{n=1}^{\infty}$ is a strongly mixing element of (\mathcal{H}, S) by Lemma 1 of [21]. It follows that

$$0 = \lim_{h \rightarrow \infty} \langle S^h((f_n)_{n=1}^{\infty}), (g_n)_{n=1}^{\infty} \rangle_{\mathcal{H}} = \lim_{h \rightarrow \infty} \lim_{q \rightarrow \infty} \frac{1}{N_q} \sum_{n=1}^{N_q} \langle f_{n+h}, g_n \rangle. \tag{2.34}$$

Since $(g_n)_{n=1}^{\infty}$ and $(N_q)_{q=1}^{\infty}$ were both arbitrary, we see that $(f_n)_{n=1}^{\infty}$ is a nearly strongly mixing sequence. \square

COROLLARY 2.8. Let \mathcal{H} be a Hilbert space and $(f_n)_{n=1}^\infty \in SA(\mathcal{H})$. If

$$\lim_{N \rightarrow \infty} \sum_{n=1}^N \langle f_{n+h}, f_n \rangle = 0 \quad (2.35)$$

for all $h \in \mathbb{N}$, then $(f_n)_{n=1}^\infty$ is a nearly strongly mixing sequence.

Results that generalize Theorem 1.1 in the same fashion that Theorem 2.7 generalizes Theorem 1.2(ii) are discussed in Chapter 2.4 of [13].

3. Applications to recurrence in measure preserving systems

LEMMA 3.1. If (X, \mathcal{B}, μ, T) is a rigid measure preserving system and $f \in L^2(X, \mu)$, then $(U_T^n f)_{n=1}^\infty$ is a rigid sequence.

Proof. Let $(k_m)_{m=1}^\infty \subseteq \mathbb{N}$ be a sequence along which T is rigid. We now see that

$$\lim_{m \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \|U_T^{n+k_m} f - U_T^n f\|^2 = \lim_{m \rightarrow \infty} \|U_T^{k_m} f - f\|^2 = 0. \quad (3.1)$$

□

THEOREM 3.2. Let (X, \mathcal{B}, μ) be a probability space and let $T, S : X \rightarrow X$ be measure preserving transformations. Suppose that the m.p.s. (X, \mathcal{B}, μ, T) is rigid, and that the m.p.s. (X, \mathcal{B}, μ, S) is totally ergodic. Let $(k_n)_{n=1}^\infty \subseteq \mathbb{N}$ be a sequence for which $((k_{n+h} - k_n)\alpha \pmod{1})_{n=1}^\infty$ is uniformly distributed for all $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $h \in \mathbb{N}$.

(i) For any $f, g \in L^\infty(X, \mu)$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f \cdot S^{k_n} g = \mathbb{E}[f | \mathcal{I}_T] \cdot \mathbb{E}[g | \mathcal{I}_S], \quad (3.2)$$

where $\mathcal{I}_R = \{A \in \mathcal{B} \mid R^{-1}A = A\}$ is the σ -algebra of R -invariant sets and with convergence taking place in $L^2(X, \mu)$.

(ii) If $A \in \mathcal{B}$ then

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mu(A \cap T^{-n}A \cap S^{-k_n}A) \geq \mu(A)^3. \quad (3.3)$$

(iii) If $((k_{n+h} - k_n)\alpha)_{n=1}^\infty$ is uniformly distributed in its orbit closure for all $\alpha \in \mathbb{R}$ then (i) and (ii) hold even when (X, \mathcal{B}, μ, S) is not ergodic.

For the proofs of (i)-(iii) all limits of sequences of vectors in $L^2(X, \mu)$ will be with respect to norm convergence in $L^2(X, \mu)$.

Proof of (i). Let $g' = g - \mathbb{E}[g | \mathcal{I}_S]$. We will use Corollary 2.8 to show that $(S^{k_n} g')_{n=1}^\infty$ is a nearly strongly mixing sequence in \mathcal{H} . Since $(\langle S^n g', g' \rangle)_{n=1}^\infty$ is a positive definite sequence we may apply the Bochner-Herglotz Theorem and pick some positive finite measure ν on

$[0, 1]$ for which $\hat{\nu}(n) = \langle S^n g', g' \rangle$ for all $n \in \mathbb{N}$. Since (X, \mathcal{B}, μ, S) is totally ergodic and $\mathbb{E}[g' | \mathcal{I}_S] = 0$ we see that $\nu(\mathbb{Q} \cap [0, 1]) = 0$. We note that for all $h \in \mathbb{N}$ we have

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \langle S^{k_{n+h}} g', S^{k_n} g' \rangle &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \langle S^{k_{n+h}-k_n} g', g' \rangle \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \hat{\nu}(k_{n+h} - k_n) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \int_0^1 e^{2\pi i(k_{n+h}-k_n)x} d\nu(x) \\ &= \int_0^1 \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N e^{2\pi i(k_{n+h}-k_n)x} d\nu(x) = 0, \end{aligned} \quad (3.4)$$

where the last equality follows from the fact that $((k_{n+h} - k_n) \alpha)_{n=1}^{\infty} \pmod{1}$ is uniformly distributed for all $\alpha \in [0, 1] \setminus \mathbb{Q}$ and $\nu(\mathbb{Q} \cap [0, 1]) = 0$.

Since (X, \mathcal{B}, μ, T) is rigid, $(T^n f)_{n=1}^{\infty}$ is a bounded rigid sequence in $\text{SA}(L^2(X, \mu))$ by Lemma 3.1. We may now use Lemma 2.5 to see that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f \cdot S^{k_n} g' = 0. \quad (3.5)$$

Furthermore, we may use the mean ergodic theorem to see that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f \cdot \mathbb{E}[g | \mathcal{I}_S] = \mathbb{E}[f | \mathcal{I}_T] \cdot \mathbb{E}[g | \mathcal{I}_S], \quad (3.6)$$

from which the desired result follows. \square

Proof of (ii). Lemma 1.6 of [11] tells us that for any bounded nonnegative $h \in L^2(X, \mu)$ we have

$$\int_X h \cdot \mathbb{E}[h | \mathcal{I}_T] \cdot \mathbb{E}[h | \mathcal{I}_S] \geq \left(\int_X h d\mu \right)^3, \quad (3.7)$$

so the desired result follows from part (i) after setting $h = \mathbb{1}_A$, \square

Proof of (iii). Since items (ii) was proven as a result of item (i), it suffices to only show that item (i) in this new situation. To this end, it suffices to repeat the proof of (i) and observe that the measure ν now satisfies $\nu(\{0, 1\}) = 0$ instead of $\nu([0, 1] \cap \mathbb{Q}) = 0$. Nonetheless, we see that the last equation of (3.4) will still hold since for all $x \in (0, 1)$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N e^{2\pi i(k_{n+h}-k_n)x} = 0. \quad (3.8)$$

\square

REMARK 3.3. We will now outline a proof of how to show that the limit in Theorem 3.2(i) exists without the assumption of total ergodicity on (X, \mathcal{B}, μ, S) when $k_n = n^2$. Let

$\mathcal{H}_{S, \mathbb{Q}}$ denote the closure of the span of the rational eigen functions of S in $L^2(X, \mu)$. Let $g = g_1 + g_2$ with $g_1 \in \mathcal{H}_{S, \mathbb{Q}}$ and $g_2 \in \mathcal{H}_{S, \mathbb{Q}}^\perp$. We argue we before to show that $(S^{n^2} g_2)_{n=1}^\infty$ is a nearly strongly mixing sequence, so $(T^n f \cdot S^{n^2} g_2)_{n=1}^\infty$ has a Cesàro average that converges to 0 in norm. Now let $\epsilon > 0$ be arbitrary and let g'_2 be a finite sum of rational eigen functions of S for which $\|g_2 - g'_2\| < \epsilon$, and let $W \in \mathbb{N}$ be such that $T^W g'_2 = g'_2$. Since $n^2 \equiv (n + W)^2 \pmod{W}$, we see that

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f \cdot S^{n^2} g_2 &\approx \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f \cdot S^{n^2} g'_2 \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{w=1}^W \sum_{n=1}^{\lfloor \frac{N}{w} \rfloor} T^{Wn+w} f \cdot S^{w^2} g'_2 = \sum_{w=1}^W \mathbb{E}[T^w f | \mathcal{I}_{T^w}] S^{w^2} g'_2, \end{aligned} \tag{3.9}$$

where the final equality is a consequence of the mean ergodic theorem.

We will now outline how to show that there is an n for which $\mu(A \cap T^{-n} A \cap S^{-kn} A) > 0$ in Theorem 3.2(ii) without the additional assumption of total ergodicity on (X, \mathcal{B}, μ, S) . Let $0 < \epsilon < \frac{1}{2} \mu(A)^2$, and note that $\{n \mid \mu(A \cap S^{-n^2} A) > \mu(A)^2 - \epsilon\}$ is an IP* set (cf. page 43 of [4]). Since T is rigid, there is an IP set B such that $\mu(A \cap T^{-n} A) > \mu(A) - \epsilon$ for all $n \in B$. We now see that for $n \in B$ we have $\mu(A \cap T^{-n} A \cap S^{-n^2} A) > 0$. It is worth remarking here that sets of the form $\{n \in \mathbb{N} \mid n^k \alpha \in [\frac{1}{4}, \frac{3}{4}]\}$ are complements of IP* sets, so Theorem 3.4 is not a result of a similar argument.

THEOREM 3.4. *Let $k \geq 2$ be an integer and $\alpha \in \mathbb{R}$ be irrational. Let $R = \{n \in \mathbb{N} \mid n^k \alpha \in [\frac{1}{4}, \frac{3}{4}]\}$. Let (X, \mathcal{B}, μ) be a probability space and $T_1, T_2, \dots, T_{k-1} : X \rightarrow X$ commuting invertible measure preserving transformations. Let $S : X \rightarrow X$ be an invertible measure preserving transformation for which (X, \mathcal{B}, μ, S) is rigid, and $\{S, T_1, T_2, \dots, T_{k-1}\}$ generate a nilpotent group. For any $A \in \mathcal{B}$ with $\mu(A) > 0$, there exists $n \in R$ for which*

$$\mu(A \cap S^{-n} A \cap T_1^{-n} A \cap T_2^{-n} A \cap \dots \cap T_{k-1}^{-n} A) > 0. \tag{3.10}$$

Proof. We remark that our proof is essentially the same as that in [17] other than the fact that we use Corollary 2.8 in place of Theorem 1.2(i), and that we use nilpotent ergodic theorems of Walsh and Leibman in place of the Furstenberg-Katznelson multiple recurrence theorem. We begin with a useful proposition.

PROPOSITION 3.5. *Let $k \in \mathbb{N}$ be at least 2, (X, \mathcal{B}, μ) be a probability space, and $T_1, \dots, T_{k-1} : X \rightarrow X$ be commuting measure preserving transformations. Let $p(x) \in \mathbb{R}_{\geq 0}[x]$ have degree at least k and an irrational leading coefficient, and let $g : \mathbb{T} \rightarrow \mathbb{C}$ be a Riemann integrable function satisfying $\int_{\mathbb{T}} g(x) dx = 0$. For any $f_1, \dots, f_{k-1} \in L^\infty(X, \mu)$ the sequence*

$$(T_1^n f_1 T_2^n f_2 \dots T_{k-1}^n f_{k-1} g(p(n)))_{n=1}^\infty \tag{3.11}$$

is a nearly strongly mixing sequence. □

Proof of Proposition 3.5. Using Lemma 2.3 we see that it suffices to prove the desired result for $g(x) = e^{2\pi imx}$ with $m \in \mathbb{N}$. We now proceed by induction on k . Let $a_n = T_1^n f_1 T_2^n f_2 \cdots T_{k-1}^n f_k e^{2\pi im p(n)}$, $g_{i,h} = T_i^h f_i \tilde{f}_i$, $\tilde{T}_i = T_i T_1^{-1}$, and observe that for $h \in \mathbb{N}$

$$\begin{aligned} & \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \langle a_{n+h}, a_n \rangle \tag{3.12} \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \int_X T_1^n g_{1,h} T_2^n g_{2,h} \cdots T_{k-1}^n g_{k-1,h} e^{2\pi im(p(n+h) - p(n))} d\mu \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \int_X g_{1,h} \tilde{T}_2^n g_{2,h} \cdots \tilde{T}_{k-1}^n g_{k-1,h} e^{2\pi im(p(n+h) - p(n))} d\mu = 0, \end{aligned}$$

where the last equality follows from the uniform distribution of $(p(n+h) - p(n))_{n=1}^\infty \pmod{1}$ when $k = 2$, and from the inductive hypothesis and Lemma 2.5 when $k > 2$. The fact that $(a_n)_{n=1}^\infty$ is a nearly strongly mixing sequence now follows from Corollary 2.8. \square

Returning to the proof of Theorem 3.4, we observe that $g(x) = \mathbb{1}_{[\frac{1}{4}, \frac{3}{4}]}(x) - \frac{1}{2}$ is a Riemann integrable function satisfying $\int_{\mathbb{T}} g(x) dx = 0$. Lemma 3.1 shows us that that $(\mathbb{1}_{S^{-n}A})_{n=1}^\infty \in \text{SA}(L^2(X, \mu))$ is a rigid sequence, from which we deduce that $(\mathbb{1}_{A \cap S^{-n}A})_{n=1}^\infty$ is also a rigid sequence. Since the sequence in equation (3.11) with $f_i = \mathbb{1}_A$ for all i is a nearly strongly mixing sequence, it follows from Lemma 2.5 that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mu(A \cap S^{-n}A \cap T_1^{-n}A \cap T_2^{-n}A \cap \cdots \cap T_{k-1}^n A) e^{2\pi imn^k \alpha} = 0, \tag{3.13}$$

for any $m \in \mathbb{N}$. We now see that

$$\begin{aligned} & \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mathbb{1}_{[\frac{1}{4}, \frac{3}{4}]}(n^k \alpha) \mu(A \cap S^{-n}A \cap T_1^{-n}A \cap T_2^{-n}A \cap \cdots \cap T_{k-1}^n A) \tag{3.14} \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \frac{1}{2} \mu(A \cap S^{-n}A \cap T_1^{-n}A \cap T_2^{-n}A \cap \cdots \cap T_{k-1}^n A) > 0, \end{aligned}$$

where the existence of the last limit follows from Theorem 1.1 of [23], and the last inequality follows from Theorem NM of [20].

REMARK 3.6. In [17], for each $k \geq 2$ a uniquely ergodic system (X_k, T_k) with unique invariant measure μ is constructed such that $\mu(A \cap T^{-n}A \cap T^{-2n}A \cap \cdots \cap T^{-kn}A) = 0$ for all $n \in R_k$ and some open set A . When $k = 2$, we may take $(X_2, T_2) = ([0, 1]^2, T)$, where $T(x, y) = (x + \alpha, y + x) \pmod{1}$ with α irrational. Since $([0, 1]^2, T)$ has zero topological (and hence measurable) entropy, we see that Theorem 3.4 cannot be extended to the situation in which S has zero entropy. Moreover, a careful examination of the proof of Theorem 3.4 shows that Lemma 2.5 cannot be extended to the case in which $(g_n)_{n=1}^\infty$ is a bounded sequence of zero entropy. In fact, the sequence in equation (3.11) is not just a nearly strongly

mixing sequence, it is a nearly orthogonal sequence (cf. Definition 2.2.5(v) in [13]) since it satisfies the hypothesis of Corollary 2.8, and even so the correlation of this sequence with a sequence of zero entropy need not be zero. In particular, this shows that the generalizations of vdCDT from this paper and Chapter 2 of [13] are not strong enough on their own to prove Theorems 1.3 or 1.4.

THEOREM 3.7. *Let (X, \mathcal{B}, μ) be a probability space and $T, S : X \rightarrow X$ be measure preserving transformations. Suppose that (X, \mathcal{B}, μ, T) is rigid and (X, \mathcal{B}, μ, S) is totally ergodic. Let $p_1, \dots, p_K \in \mathbb{Q}[x]$ be a collection of integer polynomials such that $\{p_1(n+h) - p_1(n), p_2(n+h) - p_1(n), \dots, p_K(n+h) - p_1(n), p_2(n) - p_1(n), \dots, p_K(n) - p_1(n)\}$ is independent for all $h \in \mathbb{N}$. For any $f_0, f_1, \dots, f_K \in L^\infty(X, \mu)$ and $c \in \mathbb{Z} \setminus \{0\}$, we have*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f_0 \prod_{i=1}^K S^{p_i(n)} f_i = \mathbb{E}[f_0 | \mathcal{I}_T] \prod_{i=1}^K \int_X f_i d\mu, \quad (3.15)$$

with convergence taking place in $L^2(X, \mu)$.

Before proving Theorem 3.7, we recall a theorem of Frantzikinakis and Kra from [16].

THEOREM 3.8. *Let (X, \mathcal{B}, μ, S) be a totally ergodic system, and let $p_1, \dots, p_K \in \mathbb{Q}[x]$ be a collection of independent integer polynomials. Then for $f_1, \dots, f_K \in L^\infty(X, \mu)$, we have*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \prod_{i=1}^K T^{p_i(n)} f_i = \prod_{i=1}^K \int_X f_i d\mu, \quad (3.16)$$

with convergence taking place in $L^2(X, \mu)$.

Proof of Theorem 3.7. We see from Lemma 3.1 that $(T^n f_0)_{n=1}^\infty$ is a rigid sequence, so using Lemma 2.5 and standard linearity arguments, it suffices to show that $(\prod_{i=1}^K S^{p_i(n)} f_i)_{n=1}^\infty$ is a nearly strongly mixing sequence if $\int_X f_i d\mu = 0$ for some $1 \leq i \leq K$. To this end, we first observe that for $h \in \mathbb{N}$ we have

$$\begin{aligned} & \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \int_X \left(\prod_{i=1}^K S^{p_i(n+h)} f_i \right) \left(\overline{f_{K+1}} \prod_{i=1}^K S^{p_i(n)} \overline{f_i} \right) d\mu \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \int_X \left(\prod_{i=1}^K S^{p_i(n+h) - p_i(n)} f_i \right) \left(\prod_{i=2}^K S^{p_i(n) - p_i(n)} \overline{f_i} \right) \overline{f_1} d\mu \\ &= \left(\prod_{i=1}^K \int_X f_i d\mu \right) \left(\prod_{i=2}^K \int_X \overline{f_i} d\mu \right) \int_X \overline{f_1} d\mu = 0, \end{aligned} \quad (3.17)$$

where the penultimate equality follows from Theorem 3.8, so the desired result now follows from Corollary 2.8. \square

THEOREM 3.9. *Let (X, \mathcal{B}, μ) be a probability space and $T, S_1, S_2 : X \rightarrow X$ be measure preserving transformations. Suppose that (X, \mathcal{B}, μ, T) is rigid and $(X, \mathcal{B}, \mu, S_2)$ is weakly mixing. Let $p : \mathbb{R} \rightarrow \mathbb{R}$ either be a polynomial of degree at least 2, or of the form $p(n) =$*

$n^\alpha \log(n)^\beta$ with $\alpha > 1$ and $\beta \leq 0$.[§] For any $f_0, f_1, f_2 \in L^\infty(X, \mu)$ satisfying $\int_X f_2 d\mu = 0$, we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f_0 S_1^n f_1 S_2^{\lfloor p(n) \rfloor} f_2 = 0, \quad (3.18)$$

with convergence taking place in $L^2(X, \mu)$.

Proof. We see from Lemma 3.1 that $(T^n f_0)_{n=1}^\infty$ is a rigid sequence, so using Lemma 2.5 and standard linearity arguments, it suffices to show that $(S_1^n f_1 S_2^{\lfloor p(n) \rfloor} f_2)_{n=1}^\infty$ is a nearly strongly mixing sequence if $\mathbb{E}[f_i | \mathcal{I}_{S_i}] = 0$ for some $i \in \{1, 2\}$. To this end we attempt to use Corollary 2.8, we observe that for all $h \in \mathbb{N}$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \sum_{m=1}^N \int_X S_1^{n+h} f_1 S_2^{\lfloor p(n+h) \rfloor} f_2 S_1^n \bar{f}_1 S_2^{\lfloor p(n) \rfloor} \bar{f}_2 d\mu \quad (3.19)$$

$$= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \sum_{m=1}^N \int_X S_2^{\lfloor p(n+h) \rfloor} S_1^{-n} f_2 S_2^{\lfloor p(n) \rfloor} S_1^{-n} \bar{f}_2 S_1^h f_1 \bar{f}_1 d\mu. \quad (3.20)$$

We observe that the constant sequence $(S_1^h f_1 \bar{f}_1)_{n=1}^\infty$ is a rigid sequence. If we show that $(S_2^{\lfloor p(n+h) \rfloor} S_1^{-n} f_2 S_2^{\lfloor p(n) \rfloor} S_1^{-n} \bar{f}_2)_{n=1}^\infty$ is a nearly strongly mixing sequence, then Lemma 2.5 will imply that the quantity in Equation (3.20) is 0, which will yield the desired result. We now see that for all $\ell \in \mathbb{N}$ we have

$$\begin{aligned} & \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \int_X S_2^{\lfloor p(n+\ell+h) \rfloor} S_1^{-n-\ell} f_2 S_2^{\lfloor p(n+\ell) \rfloor} S_1^{-n-\ell} \bar{f}_2 S_2^{\lfloor p(n+h) \rfloor} S_1^{-n} \bar{f}_2 S_2^{\lfloor p(n) \rfloor} S_1^{-n} f_2 d\mu \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \int_X S_2^{\lfloor p(n+\ell+h) \rfloor} S_1^{-\ell} f_2 S_2^{\lfloor p(n+\ell) \rfloor} S_1^{-\ell} \bar{f}_2 S_2^{\lfloor p(n+h) \rfloor} \bar{f}_2 S_2^{\lfloor p(n) \rfloor} f_2 d\mu \quad (3.21) \\ &= \int_X S_1^{-\ell} f_2 d\mu \int_X S_1^{-\ell} \bar{f}_2 d\mu \int_X f_2 d\mu \int_X \bar{f}_2 d\mu = 0, \end{aligned}$$

where the penultimate equality follows from Theorem A of [6], so we may use Corollary 2.8 once again to obtain the desired result. \square

REFERENCES

- [1] D. Berend and V. Bergelson. Mixing sequences in Hilbert spaces. *Proc. Amer. Math. Soc.*, 98(2):239–246, 1986.
- [2] V. Bergelson. Weakly mixing PET. *Ergodic Theory Dynam. Systems*, 7(3):337–349, 1987.
- [3] V. Bergelson. Ergodic Ramsey theory—an update. In *Ergodic theory of \mathbb{Z}^d actions (Warwick, 1993–1994)*, volume 228 of *London Math. Soc. Lecture Note Ser.*, pages 1–61. Cambridge Univ. Press, Cambridge, 1996.
- [4] V. Bergelson. Ultrafilters, IP sets, dynamics, and combinatorial number theory. In *Ultrafilters across mathematics*, volume 530 of *Contemp. Math.*, pages 23–47. Amer. Math. Soc., Providence, RI, 2010.

[§]The astute reader will observe that p could be any function for which $(g_1(n), g_2(n), g_3(n), g_4(n)) = (p(n+\ell+h), p(n+\ell), p(n+h), p(n))$ satisfies the hypothesis of Theorem A of [6], and that we have picked two aesthetic examples of such functions for the sake of simplicity.

- [5] V. Bergelson, A. del Junco, M. Lemańczyk, and J. Rosenblatt. Rigidity and non-recurrence along sequences. *Ergodic Theory Dynam. Systems*, 34(5):1464–1502, 2014.
- [6] V. Bergelson and I. J. Håland Knutson. Weak mixing implies weak mixing of higher orders along tempered functions. *Ergodic Theory Dynam. Systems*, 29(5):1375–1416, 2009.
- [7] V. Bergelson and A. Leibman. Polynomial extensions of van der Waerden's and Szemerédi's theorems. *J. Amer. Math. Soc.*, 9(3):725–753, 1996.
- [8] V. Bergelson and A. Leibman. Failure of the Roth theorem for solvable groups of exponential growth. *Ergod. Th. and Dynam. Sys.*, 24:45–53, 2004.
- [9] V. Bergelson and J. Moreira. Van der Corput's difference theorem: some modern developments. *Indag. Math. (N.S.)*, 27(2):437–479, 2016.
- [10] H. Bohr and E. Følner. On some types of functional spaces. A contribution to the theory of almost periodic functions. *Acta Math.*, 76:31–155, 1945.
- [11] Q. Chu. Multiple recurrence for two commuting transformations. *Ergodic Theory Dynam. Systems*, 31(3):771–792, 2011.
- [12] M. Einsiedler and T. Ward. *Ergodic theory with a view towards number theory*, volume 259 of *Graduate Texts in Mathematics*. Springer-Verlag London, Ltd., London, 2011.
- [13] S. Farhangi. *Topics in ergodic theory and ramsey theory*. PhD dissertation, the Ohio State University, 2022.
- [14] N. Frantzikinakis. Furstenberg systems of hardy field sequences and applications. *J. D'Analyse Math.*, 2022 (To Appear).
- [15] N. Frantzikinakis and B. Host. Multiple recurrence and convergence without commutativity. *arXiv:2111.01518v2 [math.DS]*, 2021.
- [16] N. Frantzikinakis and B. Kra. Polynomial averages converge to the product of integrals. volume 148, pages 267–276. 2005. *Probability in mathematics*.
- [17] N. Frantzikinakis, E. Lesigne, and M. Wierdl. Sets of k -recurrence but not $(k + 1)$ -recurrence. *Ann. Inst. Fourier (Grenoble)*, 56(4):839–849, 2006.
- [18] H. Furstenberg. *Recurrence in ergodic theory and combinatorial number theory*. Princeton University Press, Princeton, N.J., 1981. M. B. Porter Lectures.
- [19] L. Kuipers and H. Niederreiter. *Uniform distribution of sequences*. Pure and Applied Mathematics. Wiley-Interscience [John Wiley & Sons], New York-London-Sydney, 1974.
- [20] A. Leibman. Multiple recurrence theorem for measure preserving actions of a nilpotent group. *Geom. Funct. Anal.*, 147(5):853–931, 1998.
- [21] A. Rényi. On mixing sequences of sets. *Acta Math. Acad. Sci. Hungar.*, 9:215–228, 1958.
- [22] J. G. van der Corput. Diophantische Ungleichungen. I. Zur Gleichverteilung Modulo Eins. *Acta Math.*, 56(1):373–456, 1931.
- [23] M. Walsh. Norm convergence of nilpotent ergodic averages. *Ann. of Math.*, 175(2):1667–1688, 2012.