

On extinction time distribution of a 2-type linear-fractional branching process in a varying environment with asymptotically constant mean matrices

Hua-Ming WANG*

Abstract

In this paper we study a 2-type linear-fractional branching process in varying environment with asymptotically constant mean matrices. Let ν be the extinction time and for $k \geq 1$ let M_k be the mean matrix of offspring distribution of individuals of the $(k - 1)$ -th generation. Under certain conditions, we show that $P(\nu = n)$ and $P(\nu > n)$ are asymptotically equivalent to some functions of products of spectral radii of the mean matrices. This paper complements a former result [*arXiv*: 2007.07840] which requires in addition a condition $\forall k \geq 1, \det(M_k) < -\varepsilon$ for some $\varepsilon > 0$. Such a condition excludes a large class of mean matrices. As byproducts, we also get some results on asymptotics of products of nonhomogeneous matrices which have their own interests.

Keywords: Branching process in varying environment, extinction time, product of nonhomogeneous matrices, spectral radius, continued fraction.

MSC 2020: 60J80, 60J10, 15B48

1 Introduction

1.1 Background and motivation. It is known that many new phenomena, which homogeneous Galton-Watson processes do not possess, arise when considering the branching processes in varying environments (BPVE hereafter). Thus BPVE has been extensively studied by many authors. For details of the single-type case, we refer the reader to [1], [6], [7], [10], [11], [12], [15] and references therein.

Compared with the single-type case, the situation of the multitype setting is less satisfying. The convergence of the normalized population size and criteria for almost sure extinction can be found in [2], [3], [9] and a recent article [4]. We notice also that in [4], Dolgopyat et al. gave also the asymptotics of the survival probabilities. Their proof relies on a generalization of the Perron-Frobenius theorem suitable for studying the product of nonhomogeneous nonnegative matrices. Let ν be the extinction time and M_k be the mean matrix of offspring distribution of individuals of the $(k - 1)$ -th generation. They showed that

$$(1) \quad \frac{1}{C} \left(\sum_{k=1}^n \lambda_1^{-1} \cdots \lambda_n^{-1} \right)^{-1} < P(\nu > n) < C \left(\sum_{k=1}^n \lambda_1^{-1} \cdots \lambda_n^{-1} \right)^{-1},$$

*Email:hmking@ahnu.edu.cn; School of Mathematics and Statistics, Anhui Normal University, Wuhu, 241003, China

where $0 < C < \infty$ is a certain constant and for $k \geq 1$, λ_k is a number associated to the generalized Perron-Frobenius theorem and depends on the mean matrices $M_n, n \geq k$ of the branching process, see [4, Proposition 2.1 and Lemma 2.2]. Although (1) provides a lower and an upper bound for the survival probability, it does not give the asymptotical equivalence of the survival probability and furthermore, those number $\lambda_k, k \geq 1$ are hard to compute explicitly. For this consideration, Wang and Yao [18] considered a 2-type linear-fractional branching process with asymptotically constant mean matrices. Instead of those number λ_k , they used the spectral radii of the mean matrices of the offspring distributions, which can be explicitly computed. For this special setting, they give not only the asymptotical equivalence of $P(\nu > n)$ but also that of $P(\nu = n)$. But, they need a condition

$$(2) \quad \forall k \geq 1, M_k(12)M_k(21) > M_k(11)M_k(22) + \varepsilon$$

for some $\varepsilon > 0$, which excludes a large class of mean matrices. They added such a restriction because their proof depends on some result on the asymptotical equivalence between the elements of products of *positive* matrices and the products of spectral radii of those matrices, which was studied in [17], and some delicate analysis of the tails and the critical tails of *positive* continued fractions. Without (2), one has to deal with product of matrices with some *negative elements* and also continued fractions with *negative coefficients*.

In this paper, by generalizing the result of product of matrices in [17] and analyzing the tails and approximants of continued fractions with negative coefficients, we remove the assumption (2) to characterize asymptotics of the tail probability and the probability that the extinction time equals n as $n \rightarrow \infty$.

1.2 Model and main results. Suppose that $M_k, k \geq 1$ is a sequence of nonnegative 2-by-2 matrices and $\gamma_k = (\gamma_k^{(1)}, \gamma_k^{(2)})$, $k \geq 1$ is a sequence of nonnegative row vectors. To avoid the degenerate case, we require that $\forall k \geq 1$, all elements of $M_k M_{k+1}$ are strictly positive and $\gamma_k \neq \mathbf{0}$. For $\mathbf{s} = (s_1, s_2)^t \in [0, 1]^2$ and $k \geq 1$, let

$$\mathbf{f}_k(\mathbf{s}) = (f_k^{(1)}(\mathbf{s}), f_k^{(2)}(\mathbf{s}))^t = \mathbf{1} - \frac{M_k(\mathbf{1} - \mathbf{s})}{\mathbf{1} + \gamma_k(\mathbf{1} - \mathbf{s})}$$

which is known as the probability generating function of a linear-fractional distribution. Here and in what follows, \mathbf{v}^t denotes the transpose of a vector \mathbf{v} and $\mathbf{1} = (\mathbf{e}_1 + \mathbf{e}_2)^t = (1, 1)^t$, with $\mathbf{e}_1 = (1, 0)$, $\mathbf{e}_2 = (0, 1)$.

Suppose that $Z_n = (Z_{n,1}, Z_{n,2}), n \geq 0$ is a stochastic process such that

$$E(\mathbf{s}^{Z_n} | Z_0, \dots, Z_{n-1}) = [\mathbf{f}_n(\mathbf{s})]^{Z_{n-1}}, n \geq 1,$$

where $[\mathbf{f}_n(\mathbf{s})]^{Z_{n-1}} := [f_n^{(1)}(\mathbf{s})]^{Z_{n-1,1}} [f_n^{(2)}(\mathbf{s})]^{Z_{n-1,2}}$. We call the process $Z_n, n \geq 0$ a two-type linear-fractional branching process in a varying environment. Matrices $M_k, k \geq 1$ are usually referred to as the mean matrices of the branching process. Denote by

$$\nu = \min\{n : Z_n = \mathbf{0}\}$$

the extinction time of $\{Z_n\}$ which we concern.

Throughout, we assume $b_k, d_k > 0$, $a_k, \theta_k \geq 0$, $a_k + \theta_k > 0$, $k \geq 1$ and put

$$(3) \quad M_k := \begin{pmatrix} a_k & b_k \\ d_k & \theta_k \end{pmatrix}, \gamma_k := \mathbf{e}_1 M_k, \forall k \geq 1.$$

We introduce the following conditions on the number $a_k, b_k, d_k, \theta_k, k \geq 1$.

(B1) Suppose that $b, d > 0, a, \theta \geq 0$ are some numbers such that $a + \theta > 0, a_k \rightarrow a, b_k \rightarrow b, d_k \rightarrow d, \theta_k \rightarrow \theta$ as $k \rightarrow \infty$ and assume further that

$$\sum_{k=2}^{\infty} |a_k - a_{k-1}| + |b_k - b_{k-1}| + |d_k - d_{k-1}| + |\theta_k - \theta_{k-1}| < \infty.$$

Suppose now condition (B1) holds and for $k \geq 1$, set

$$(4) \quad A_k := \begin{pmatrix} \tilde{a}_k & \tilde{b}_k \\ \tilde{d}_k & 0 \end{pmatrix} \text{ with } \tilde{a}_k = a_k + \frac{b_k \theta_{k+1}}{b_{k+1}}, \tilde{b}_k = b_k, \tilde{d}_k = d_k - \frac{a_k \theta_k}{b_k}.$$

Letting $\Lambda_k = \begin{pmatrix} 1 & 0 \\ \theta_k/b_k & 1 \end{pmatrix}, k \geq 1$, then for $n \geq k \geq 1$, we have

$$(5) \quad A_k = \Lambda_k^{-1} M_k \Lambda_{k+1} \text{ and } \mathbf{e}_1 \prod_{i=k}^n M_i \mathbf{1} = \mathbf{e}_1 \prod_{i=k}^n A_i (1, 1 - \theta_{n+1}/b_{n+1})^t.$$

For linear-fractional setting, the distribution of ν can be formulated explicitly by M_k , see (70) and (71) below. However, the elements of $\prod_{i=k}^n M_i$ are hard to compute and evaluate whereas those of $\prod_{i=k}^n A_i$ are workable because they have some correspondence with continued fractions due to the special structure of the matrices $A_i, i \geq 1$. Therefore, instead of M_k , we will work with A_k below.

We need in addition the following conditions which are mutually exclusive.

(B2)_a $\exists k_0 > 0$, such that $\frac{\tilde{a}_k}{b_k} = \frac{\tilde{a}_{k+1}}{b_{k+1}}, \frac{\tilde{d}_k}{b_k} \neq \frac{\tilde{d}_{k+1}}{b_{k+1}}, \forall k \geq k_0$ and

$$\lim_{k \rightarrow \infty} \frac{\tilde{d}_{k+2}/\tilde{b}_{k+2} - \tilde{d}_{k+1}/\tilde{b}_{k+1}}{\tilde{d}_{k+1}/\tilde{b}_{k+1} - \tilde{d}_k/\tilde{b}_k} \text{ exists.}$$

(B2)_b $\exists k_0 > 0$, such that $\frac{\tilde{a}_k}{b_k} \neq \frac{\tilde{a}_{k+1}}{b_{k+1}}, \frac{\tilde{d}_k}{b_k} = \frac{\tilde{d}_{k+1}}{b_{k+1}}, \forall k \geq k_0$ and

$$\lim_{k \rightarrow \infty} \frac{\tilde{a}_{k+2}/\tilde{b}_{k+2} - \tilde{a}_{k+1}/\tilde{b}_{k+1}}{\tilde{a}_{k+1}/\tilde{b}_{k+1} - \tilde{a}_k/\tilde{b}_k} \text{ exists.}$$

(B2)_c $\exists k_0 > 0$, such that $\frac{\tilde{a}_k}{b_k} \neq \frac{\tilde{a}_{k+1}}{b_{k+1}}, \frac{\tilde{d}_k}{b_k} \neq \frac{\tilde{d}_{k+1}}{b_{k+1}}, \forall k \geq k_0$ and

$$\tau := \lim_{k \rightarrow \infty} \frac{\tilde{d}_{k+1}/\tilde{b}_{k+1} - \tilde{d}_k/\tilde{b}_k}{\tilde{a}_{k+1}/\tilde{b}_{k+1} - \tilde{a}_k/\tilde{b}_k} \neq \frac{-(a + \theta) \pm \sqrt{(a + \theta)^2 + 4(bd - a\theta)}}{2b}$$

exists as a finite or infinite number. In addition, if τ is finite, assume further that $\lim_{k \rightarrow \infty} \frac{\tilde{a}_{k+2}/\tilde{b}_{k+2} - \tilde{a}_{k+1}/\tilde{b}_{k+1}}{\tilde{a}_{k+1}/\tilde{b}_{k+1} - \tilde{a}_k/\tilde{b}_k}$ exists. Otherwise, if $\tau = \infty$, assume further that $\lim_{k \rightarrow \infty} \frac{\tilde{d}_{k+2}/\tilde{b}_{k+2} - \tilde{d}_{k+1}/\tilde{b}_{k+1}}{d_{k+1}/b_{k+1} - d_k/b_k}$ exists.

The conditions **(B2)_a**, **(B2)_b** and **(B2)_c** look a bit complicated. But they are not so difficult to fulfilled. We refer the reader to [18, Lemma 1] for examples for which **(B1)** and one of **(B2)_a**, **(B2)_b** and **(B2)_c** hold.

Note that under condition **(B1)**, we have

$$\lim_{k \rightarrow \infty} M_k = M := \begin{pmatrix} a & b \\ d & \theta \end{pmatrix}, \lim_{k \rightarrow \infty} A_k = A := \begin{pmatrix} a + \theta & b \\ d - a\theta/b & 0 \end{pmatrix}$$

whose eigenvalues are

$$(6) \quad \varrho := \varrho(M) = \varrho(A) = \frac{a + \theta + \sqrt{(a + \theta)^2 + 4(bd - a\theta)}}{2},$$

$$(7) \quad \varrho_1 := \varrho_1(M) = \varrho_1(A) = \frac{a + \theta - \sqrt{(a + \theta)^2 + 4(bd - a\theta)}}{2}.$$

It is clear that $|\varrho_1| < \varrho$. In the literature, $\varrho(B) := \sup\{|\lambda| : \lambda \text{ is an eigenvalue of } B\}$ is usually referred to as the spectral radius of a matrix B . In what follows, we always denote by $\varrho(B)$ the spectral radius of a matrix B and when we simply write ϱ and ϱ_1 , their values will be always those defined in (6) and (7) respectively.

In the rest of the paper, $f(n) \sim g(n)$ means $\lim_{n \rightarrow \infty} f(n)/g(n) = 1$, $f(n) = o(g(n))$ means $\lim_{n \rightarrow \infty} f(n)/g(n) = 0$ and unless otherwise specified, c and C are some universal strictly positive constants, which may change from line to line. We always assume that empty product equals identity and empty sum equals 0. The convention $\sqrt{-1} = i$ will also be adopted.

Now we are ready to state the main result.

Theorem 1. *Suppose that condition (B1) and one of the conditions (B2)_a, (B2)_b and (B2)_c hold. Assume further that $|\varrho_1| < 1$, $bd \neq a\theta$ and $\kappa_i > 0, \varkappa_i > 0, i = 1, 2$ are proper constants. Then for $i \in \{1, 2\}$*

$$(8) \quad P(\nu > n | Z_0 = \mathbf{e}_i) \sim \frac{\kappa_i}{\sum_{k=1}^{n+1} \varrho(M_1)^{-1} \cdots \varrho(M_{k-1})^{-1}} \text{ as } n \rightarrow \infty.$$

Furthermore, if $\varrho_1 \neq \frac{1}{2} \left(a + b + 1 - \sqrt{(a + b + 1)^2 + 4\frac{bd - a\theta}{\theta - b}} \right)$, then for $i \in \{1, 2\}$

$$(9) \quad P(\nu = n | Z_0 = \mathbf{e}_i) \sim \frac{\varkappa_i \varrho(M_1)^{-1} \cdots \varrho(M_n)^{-1}}{\left(\sum_{k=1}^{n+1} \varrho(M_1)^{-1} \cdots \varrho(M_{k-1})^{-1} \right)^2}, \text{ as } n \rightarrow \infty;$$

otherwise, if $\varrho_1 = \frac{1}{2} \left(a + b + 1 - \sqrt{(a + b + 1)^2 + 4\frac{bd - a\theta}{\theta - b}} \right)$, then for $i \in \{1, 2\}$,

$$(10) \quad P(\nu = n | Z_0 = \mathbf{e}_i) = o \left(\frac{\varrho(M_1)^{-1} \cdots \varrho(M_n)^{-1}}{\left(\sum_{k=1}^{n+1} \varrho(M_1)^{-1} \cdots \varrho(M_{k-1})^{-1} \right)^2} \right), \text{ as } n \rightarrow \infty.$$

Remark 1. By some elementary computation, one can show that if $bd > a\theta$, then $\varrho_1 = \frac{1}{2} \left(a + b + 1 - \sqrt{(a + b + 1)^2 + 4\frac{bd - a\theta}{\theta - b}} \right)$ if and only if $\theta = b + 1$, see [18] for details.

The proof of Theorem 1 relies on the asymptotics of the product of nonhomogeneous matrices $A_k, k \geq 1$, which have their own interests. See the two theorems below.

Theorem 2. *Suppose that condition (B1) and one of the conditions (B2)_a, (B2)_b and (B2)_c hold. Then for $i, j \in \{1, 2\}$ and $k \geq 1$, there exists a number $c(k, i, j) \neq 0$ such that*

$$(11) \quad \lim_{m \rightarrow \infty} \frac{\mathbf{e}_i A_k \cdots A_m \mathbf{e}_j^t}{\varrho(A_k) \cdots \varrho(A_m)} = c(k, i, j).$$

Remark 2. In [17], similarly result is proved if for each $k \geq 1$, all elements of A_k are nonnegative, see Theorem 1 therein. But now, under condition (B1), \tilde{a}_k might be negative. So, though the basic idea is somewhat the same, the proof of Theorem 2 differs on a large extent from the one of Theorem 1 in [17].

Theorem 3. *Suppose that condition (B1) and one of the conditions (B2)_a, (B2)_b and (B2)_c hold. If $bd \neq a\theta$, then*

$$(12) \quad \sum_{k=1}^{n+1} \mathbf{e}_1 A_k \cdots A_n \mathbf{e}_1^t \sim c \sum_{k=1}^{n+1} \varrho(A_k) \cdots \varrho(A_n), \text{ as } n \rightarrow \infty.$$

Remark 3. Although we have shown in Theorem 2 that for each $k \geq 1$, $\exists c(k) \neq 0$ such that $\mathbf{e}_1 A_k \cdots A_n \mathbf{e}_1^t \sim c(k) \varrho(A_k) \cdots \varrho(A_n)$ as $n \rightarrow \infty$, it is not an easy task to prove Theorem 3, because that the constants $c(k), k \geq 1$ involved there are mutually different and moreover every summand in $\sum_{k=1}^{n+1} \mathbf{e}_1 A_k \cdots A_n \mathbf{e}_1^t$ depends on n .

1.3 Outline of the paper. The remainder of the paper is organized as follows. In Section 2, we give some preliminary results of continued fractions and also some facts on products of the matrices M_k and A_k . Section 3 and Section 4 are devoted to proving Theorem 2 and Theorem 3 respectively. Finally, in Section 5, based on Theorem 2 and Theorem 3, we finish the proof of Theorem 1.

2 Preliminary results

Products of 2-by-2 matrices are closely related to continued fractions and therefore continued fractions are important tools to prove Theorem 1. To begin with, we introduce some basics of continued fractions.

2.1 Continued fractions and their tails

Let $\beta_k, \alpha_k, k \geq 1$ be certain real numbers. For $1 \leq k \leq n$, We denote by

$$(13) \quad \xi_{k,n} \equiv \frac{\beta_k}{\alpha_k} + \frac{\beta_{k+1}}{\alpha_{k+1}} + \cdots + \frac{\beta_n}{\alpha_n} := \frac{\beta_k}{\alpha_k + \frac{\beta_{k+1}}{\alpha_{k+1} + \frac{\beta_n}{\alpha_n}}}$$

the $(n - k + 1)$ -th approximant of a continued fraction

$$(14) \quad \xi_k := \frac{\beta_k}{\alpha_k} + \frac{\beta_{k+1}}{\alpha_{k+1}} + \frac{\beta_{k+2}}{\alpha_{k+2}} + \cdots.$$

We call $\xi_k, k \geq 1$ in (14) the tails and $h_k := \frac{\beta_k}{\alpha_{k-1} + \frac{\beta_{k-1}}{\alpha_{k-2} + \cdots + \frac{\beta_2}{\alpha_1}}, k \geq 2$ the critical tails of the continued fraction $\frac{\beta_1}{\alpha_1} + \frac{\beta_2}{\alpha_2} + \cdots$ respectively. We remark that in the literature, the n -th tail of a continued fraction $\frac{\beta_1}{\alpha_1} + \frac{\beta_2}{\alpha_2} + \cdots$ is usually denoted by $f^{(n)} = \frac{\beta_{n+1}}{\alpha_{n+1}} + \frac{\beta_{n+2}}{\alpha_{n+2}} + \cdots$ and the critical tails are also slightly different from $h_n, n \geq 1$ above.

If $\lim_{n \rightarrow \infty} \xi_{k,n}$ exists, we say that the continued fraction ξ_k is convergent and its value is defined as $\lim_{n \rightarrow \infty} \xi_{k,n}$. The lemma below will be use times and again.

Lemma 1. *If $\lim_{n \rightarrow \infty} \alpha_n = \alpha \neq 0, \lim_{n \rightarrow \infty} \beta_n = \beta$, and $\alpha^2 + 4\beta \geq 0$, then for any $k \geq 1, \lim_{n \rightarrow \infty} \xi_{k,n}$ exists and furthermore*

$$\lim_{k \rightarrow \infty} h_k = \lim_{k \rightarrow \infty} \xi_k = \frac{\alpha}{2} \left(\sqrt{1 + 4\beta/\alpha^2} - 1 \right).$$

The proof Lemma 1 can be found in many references, we refer the reader to [13], see discussion between (4.1) and (4.2) on page 81 therein.

2.2 Some facts on matrices A_k and M_k

By assumptions on the numbers a_k, b_k, d_k and θ_k , we have

$$\begin{aligned}
(15) \quad & A_k \cdots A_n = \Lambda_k^{-1} M_k \cdots M_n \Lambda_{n+1}, n \geq k \geq 1, \\
& \mathbf{e}_1 A_k \cdots A_n \mathbf{e}_1^t = \mathbf{e}_1 M_k \cdots M_n (1, \theta_{n+1}/b_{n+1})^t > 0, n > k \geq 1, \\
(16) \quad & \mathbf{e}_1 A_k \cdots A_n \mathbf{e}_2^t = \mathbf{e}_1 M_k \cdots M_n \mathbf{e}_2^t > 0, n > k \geq 1, \\
(17) \quad & \mathbf{e}_2 A_k \cdots A_n \mathbf{e}_1^t = (-\theta_k/b_k, 1) M_k \cdots M_n (1, \theta_{n+1}/b_{n+1})^t, n \geq k \geq 1.
\end{aligned}$$

Note that the conditions of Theorem 1 can not ensure $\Delta_k := \left(a_k + \frac{b_k \theta_{k+1}}{b_{k+1}}\right)^2 + 4(b_k d_k - a_k \theta_k) > 0$ for each $k \geq 1$. Therefore, the matrix A_k may possess complex eigenvalues. However, since $\lim_{k \rightarrow \infty} \Delta_k = (a - \theta)^2 + 4bd > 0$, there exists a number $N_1 > 0$ such that $\Delta_k > 0, \forall k \geq N_1$, and thus

$$\begin{aligned}
(18) \quad \varrho(A_k) &= \frac{\sqrt{\tilde{a}_k^2 + 4\tilde{b}_k \tilde{d}_k} + \tilde{a}_k}{2} \\
&= \frac{1}{2} \left(a_k + \frac{b_k \theta_{k+1}}{b_{k+1}} + \sqrt{\left(a_k + \frac{b_k \theta_{k+1}}{b_{k+1}}\right)^2 + 4(b_k d_k - a_k \theta_k)} \right), \forall k \geq N_1.
\end{aligned}$$

Furthermore, if $bd \neq a\theta$, then $\lim_{n \rightarrow \infty} \tilde{d}_n = d - a\theta/b \neq 0$. Thus, if $bd < a\theta$, then

$$\begin{aligned}
& \exists \varepsilon > 0 \text{ and } N_2 > 0, \text{ such that } \tilde{d}_n < -\varepsilon, \forall n \geq N_2 \text{ and consequently} \\
& \mathbf{e}_2 A_k \cdots A_n \mathbf{e}_1^t < 0, n \geq k \geq N_2 \text{ and } \mathbf{e}_2 A_k \cdots A_n \mathbf{e}_2^t < 0, n > k \geq N_2;
\end{aligned}$$

otherwise, if $bd > a\theta$, then

$$\begin{aligned}
& \exists \varepsilon > 0 \text{ and } N_2 > 0, \text{ such that } \tilde{d}_n > \varepsilon, \forall n \geq N_2 \text{ and consequently} \\
& \mathbf{e}_2 A_k \cdots A_n \mathbf{e}_1^t > 0, n \geq k \geq N_2 \text{ and } \mathbf{e}_2 A_k \cdots A_n \mathbf{e}_2^t > 0, n > k \geq N_2.
\end{aligned}$$

3 Asymptotics of elements of $\prod_{j=k}^m A_j$

The main task of this section is to prove Theorem 2. The proof is based on the spectral radius estimation derived in [8] and some analysis of the fluctuations of the tails and critical tails of continued fractions.

3.1 Lower and upper bounds for $\frac{\mathbf{e}_1 A_k \cdots A_m \mathbf{e}_1^t}{\varrho(A_k) \cdots \varrho(A_m)}$

The main result of this subsection is the following lemma.

Lemma 2. *Suppose that condition (B1) holds. Then $\exists N_0 > 0$ such that for each $k \geq N_0$, $C_k^{-1} < \frac{\mathbf{e}_1 A_k \cdots A_m \mathbf{e}_1^t}{\varrho(A_k) \cdots \varrho(A_m)} < C_k, \forall m \geq k$, where $C_k > 0$ is a proper number.*

Lemma 2 is a direct consequence of the following four auxiliary lemmas.

Lemma 3. *Suppose the condition (B1) is satisfied. Then with N_1 the one in (18), for each $k \geq N_1$, $\exists C_k > 0$ and $\mathcal{B}_k > 0$ such that*

$$\prod_{j=k}^n \varrho(A_j) \sim C_k \prod_{j=k}^n \varrho(M_j) \text{ and } \sum_{j=k}^{n+1} \prod_{i=k}^{j-1} \varrho(A_i)^{-1} \sim \mathcal{B}_k \sum_{j=k}^{n+1} \prod_{i=k}^{j-1} \varrho(M_i)^{-1}, \text{ as } n \rightarrow \infty.$$

For the proof of Lemma 3, we refer the reader to [18].

Lemma 4. *Suppose that condition (B1) holds. Then there exist constants $0 < \zeta < \gamma < \infty$ such that for $m \geq k \geq 1$, $\zeta \leq \frac{\varrho(M_k \cdots M_m)}{\varrho(M_k) \cdots \varrho(M_m)} \leq \gamma$.*

Proof. For vectors $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$ and $\mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}$, set

$$\frac{\mathbf{v}}{\mathbf{u}} := \begin{pmatrix} v_1/u_1 \\ v_2/u_2 \end{pmatrix}, \quad \left(\frac{\mathbf{v}}{\mathbf{u}}\right)_{\min} := \min \left\{ \frac{v_1}{u_1}, \frac{v_2}{u_2} \right\}, \quad \left(\frac{\mathbf{v}}{\mathbf{u}}\right)_{\max} := \max \left\{ \frac{v_1}{u_1}, \frac{v_2}{u_2} \right\}.$$

Let \mathbf{v}_n be a right eigenvector of A_n corresponding to the largest eigenvalue $\varrho(A_n)$. Then we can choose \mathbf{v}_n to be $\mathbf{v}_n = (\varrho(A_n) - \theta_n, d_n)^t$. For $m \geq k \geq 1$, write

$$\begin{aligned} \gamma_{k,m} &:= \left(\frac{\mathbf{v}_k}{\mathbf{v}_{k-1}}\right)_{\max} \cdots \left(\frac{\mathbf{v}_{m+1}}{\mathbf{v}_m}\right)_{\max} \left(\frac{\mathbf{v}_m}{\mathbf{v}_k}\right)_{\max}, \\ \zeta_{k,m} &:= \left(\frac{\mathbf{v}_k}{\mathbf{v}_{k-1}}\right)_{\min} \cdots \left(\frac{\mathbf{v}_{m+1}}{\mathbf{v}_m}\right)_{\min} \left(\frac{\mathbf{v}_m}{\mathbf{v}_k}\right)_{\min}. \end{aligned}$$

Applying [8, Theorem 1, page 228], for $k \geq 1$, we have

$$\zeta_{k,m} \leq \frac{\varrho(M_k \cdots M_m)}{\varrho(M_k) \cdots \varrho(M_m)} \leq \gamma_{k,m}.$$

It remains to show that both $\zeta_{k,m}^{-1}$ and $\gamma_{k,m}$, $m \geq k \geq 1$ are uniformly bounded away from ∞ . To this end, set $\epsilon_n = (\mathbf{v}_n/\mathbf{v}_{n-1})_{\max} - 1$, $n \geq 2$. Then by condition (B1), for $n \geq 2$, we have

$$\begin{aligned} |\epsilon_n| &\leq \max \left\{ \left| \frac{\varrho(M_n) - \theta_n}{\varrho(M_{n-1}) - \theta_{n-1}} - 1 \right|, \left| \frac{d_n}{d_{n-1}} - 1 \right| \right\} \\ &\leq \left| \frac{\varrho(M_n) - \theta_n}{\varrho(M_{n-1}) - \theta_{n-1}} - 1 \right| + \left| \frac{d_n}{d_{n-1}} - 1 \right| \\ &\leq c(|\varrho(M_n) - \varrho(M_{n-1})| + |d_n - d_{n-1}| + |\theta_n - \theta_{n-1}|) \\ &\leq c(|a_n - a_{n-1}| + |b_n - b_{n-1}| + |d_n - d_{n-1}| + |\theta_n - \theta_{n-1}|) < \infty, \end{aligned}$$

which implies $\sum_{n=2}^{\infty} \log(1 + |\epsilon_n|) < \infty$. As a consequence,

$$\gamma_{k,m} \leq \max \left\{ \frac{\varrho(M_m) - \theta_m}{\varrho(M_k) - \theta_k}, \frac{d_m}{d} \right\} \prod_{n=2}^{\infty} (1 + |\epsilon_n|) < \gamma$$

for some number $\gamma < \infty$ independent of k and m .

Since $\zeta_{k,m}^{-1} = \left(\frac{\mathbf{v}_{k-1}}{\mathbf{v}_k}\right)_{\max} \cdots \left(\frac{\mathbf{v}_m}{\mathbf{v}_{m+1}}\right)_{\max} \left(\frac{\mathbf{v}_k}{\mathbf{v}_m}\right)_{\max}$, a similar argument also yields that $\zeta_{k,m}^{-1}$, $m \geq k \geq 1$ are uniformly bounded away from ∞ . Consequently, Lemma 4 is proved. \square

Lemma 5. *Suppose that condition (B1) holds and for $k \geq 1$, let ξ_k be the one in (14) with $\alpha_n = \frac{\tilde{a}_n}{\tilde{b}_n}$, $\beta_n = \frac{\tilde{d}_n}{\tilde{b}_n}$, $n \geq 1$. Then $\exists N_3 > 0$ such that*

$$(19) \quad \lim_{m \rightarrow \infty} \frac{\varrho(A_k \cdots A_m)}{\mathbf{e}_1 A_k \cdots A_m \mathbf{e}_1^t} = 1 + b\varrho^{-1}\xi_k > c > 0, k \geq N_3.$$

Proof. For $m \geq k \geq 1$, write $A_{k,m} := A_k \cdots A_m = \begin{pmatrix} A_{k,m}(11) & A_{k,m}(12) \\ A_{k,m}(21) & A_{k,m}(22) \end{pmatrix}$.

Noticing that

$$A_{k,m} = \begin{pmatrix} \tilde{a}_k & \tilde{b}_k \\ \tilde{d}_k & 0 \end{pmatrix} \cdots \begin{pmatrix} \tilde{a}_m & \tilde{b}_m \\ \tilde{d}_m & 0 \end{pmatrix} = \tilde{b}_k \cdots \tilde{b}_m \begin{pmatrix} \alpha_k & 1 \\ \beta_k & 0 \end{pmatrix} \cdots \begin{pmatrix} \alpha_m & 1 \\ \beta_m & 0 \end{pmatrix},$$

thus we get

$$\frac{A_{k,m+1}(22)}{A_{k,m+1}(12)} = \frac{A_{k,m}(21)}{A_{k,m}(11)},$$

and by forward and backward induction we have

$$(20) \quad \frac{A_{k,m}(21)}{A_{k,m}(11)} = \frac{\beta_k}{\alpha_k + \alpha_{k+1} + \cdots + \alpha_{m-1} + \alpha_m} =: \xi_{k,m},$$

$$(21) \quad \frac{A_{k,m}(12)}{A_{k,m}(11)} = \frac{1}{\alpha_m + \alpha_{m-1} + \alpha_{m-2} + \cdots + \alpha_{k+1} + \alpha_k} =: h_{m,k},$$

$$\frac{A_{k,m}(22)}{A_{k,m}(21)} = \frac{1}{\alpha_m + \alpha_{m-1} + \alpha_{m-2} + \cdots + \alpha_{k+2} + \alpha_{k+1}}.$$

Since $\alpha_k \rightarrow (a + \theta)/b$ and $\beta_k \rightarrow (bd - a\theta)/b^2$ as $n \rightarrow \infty$, applying Lemma 1 we get

$$(22) \quad \lim_{m \rightarrow \infty} \frac{A_{k,m}(21)}{A_{k,m}(11)} = \lim_{k \rightarrow \infty} \frac{A_{k,m}(22)}{A_{k,m}(12)} = \xi_k,$$

$$(23) \quad \lim_{m \rightarrow \infty} \frac{A_{k,m}(12)}{A_{k,m}(11)} = \lim_{k \rightarrow \infty} \frac{A_{k,m}(22)}{A_{k,m}(21)} = b\varrho^{-1},$$

$$(24) \quad \lim_{m \rightarrow \infty} \xi_k = -\frac{\varrho_1}{b}.$$

We are ready to prove (19). Write

$$(25) \quad P_{k,m} = A_{k,m}(12)A_{k,m}(21) - A_{k,m}(11)A_{k,m}(22).$$

If $(A_{k,m}(11) + A_{k,m}(22))^2 + 4P_{k,m} \geq 0$, then

$$(26) \quad \varrho(A_{k,m}) = \frac{A_{k,m}(11) + A_{k,m}(22)}{2} + \frac{\sqrt{(A_{k,m}(11) + A_{k,m}(22))^2 + 4P_{k,m}}}{2},$$

otherwise, if $(A_{k,m}(11) + A_{k,m}(22))^2 + 4P_{k,m} < 0$, then

$$\varrho(A_{k,m}) = \frac{1}{2} \sqrt{(A_{k,m}(11) + A_{k,m}(22))^2 + |(A_{k,m}(11) + A_{k,m}(22))^2 + 4P_{k,m}|}.$$

Note that $\forall m \geq k \geq 1$, $A_{k,m}(11) > 0$ and by (22)-(24), we have

$$(27) \quad \lim_{k \rightarrow \infty} (1 + b\varrho^{-1}\xi_k) = 1 - \frac{\varrho_1}{\varrho} > 0,$$

$$\lim_{m \rightarrow \infty} \frac{(A_{k,m}(11) + A_{k,m}(22))^2 + 4P_{k,m}}{(A_{k,m}(11))^2} = (1 + b\varrho^{-1}\xi_k)^2.$$

Therefore there exists $N_3 > 0$, such that for each $k > N_3$, we have $1 + b\varrho^{-1}\xi_k > 0$, and for m large enough the eigenvalues of $A_{k,m}$ are real so that (26) holds. Thus, taking (22)-(24) into account, for $k \geq N_3$, we get

$$\lim_{m \rightarrow \infty} \frac{\varrho(A_k \cdots A_m)}{\mathbf{e}_1 A_k \cdots A_m \mathbf{e}_1^t} = \lim_{m \rightarrow \infty} \frac{\varrho(A_{k,m})}{A_{k,m}(11)}$$

$$= 1/2(1 + b\varrho^{-1}\xi_k + |1 + b\varrho^{-1}\xi_k|) = 1 + b\varrho^{-1}\xi_k > c > 0,$$

which proves (19). The lemma is proved. \square

Lemma 6. For $k \geq 1$, set $\varphi_k =: \frac{1+b\varrho^{-1}\xi_k}{1+b\varrho^{-1}\xi_k+b\varrho^{-1}\left(\frac{\theta_k}{b_k}-\frac{\theta}{b}\right)}$. Then, $\exists N_4 > 0$ such that

$$(28) \quad \lim_{n \rightarrow \infty} \frac{\varrho(A_k \cdots A_n)}{\varrho(M_k \cdots M_n)} = \varphi_k > c > 0, k \geq N_4.$$

Proof. Write $A_{k,n} = A_k \cdots A_n$, let $P_{k,n}$ be the one in (25) and set

$$Q_{k,n} = A_{k,n}(11) + A_{k,n}(22) + A_{k,n}(12) \left(\frac{\theta_k}{b_k} - \frac{\theta_{n+1}}{b_{n+1}} \right).$$

Since $M_k \cdots M_n = \Lambda_k A_{k,n} \Lambda_{n+1}^{-1}$, then

$$\varrho(M_k \cdots M_n) = \frac{1}{2} \left(Q_{k,n} + \sqrt{Q_{k,n}^2 + 4P_{k,n}} \right).$$

Using (22) and (23), we get

$$(29) \quad \lim_{n \rightarrow \infty} \frac{\varrho(M_k \cdots M_n)}{A_{k,n}(11)} = 1 + b\varrho^{-1}\xi_k + b\varrho^{-1} \left(\frac{\theta_k}{b_k} - \frac{\theta}{b} \right).$$

Let N_3 be the one in Lemma 5. Using (27) and the fact $\lim_{k \rightarrow \infty} \theta_k/b_k \rightarrow \theta/b$, we have $\lim_{k \rightarrow \infty} \varphi_k = 1$. Hence, $\exists N_4 > N_3$ such that for $k > N_4$, $\varphi_k > 1/2$. As a result, (29) together with (19) implies that (28) is true. \square

3.2 Fluctuations of tails and critical tails of continued fractions

For $k \geq 1$, let

$$(30) \quad f_k = \frac{\tilde{b}_k \tilde{d}_k^{-1}}{\tilde{a}_k \tilde{d}_k^{-1}} + \frac{\tilde{b}_{k-1} \tilde{d}_{k-1}^{-1}}{\tilde{a}_{k-1} \tilde{d}_{k-1}^{-1}} + \cdots + \frac{\tilde{b}_1 \tilde{d}_1^{-1}}{\tilde{a}_1 \tilde{d}_1^{-1}},$$

$$\xi_k = \frac{\tilde{b}_k^{-1} \tilde{d}_{k+1}^{-1}}{\tilde{a}_k \tilde{b}_k^{-1} \tilde{d}_{k+1}^{-1}} + \frac{\tilde{b}_{k+1}^{-1} \tilde{d}_{k+2}^{-1}}{\tilde{a}_{k+1} \tilde{b}_{k+1}^{-1} \tilde{d}_{k+2}^{-1}} + \cdots.$$

Set also

$$(31) \quad \varepsilon_k^f = f_k - \tilde{b}_{k+1} \varrho(A_{k+1})^{-1}, \quad \varepsilon_k^\xi = \xi_k - \varrho(A_k)^{-1}, k \geq 1,$$

$$\delta_k^f = \tilde{b}_k \tilde{d}_k^{-1} - \tilde{b}_{k+1} \varrho(A_{k+1})^{-1} (\tilde{a}_k \tilde{d}_k^{-1} + \tilde{b}_k \varrho(A_k)^{-1}), k \geq 2,$$

$$\delta_k^\xi = \tilde{b}_k^{-1} \tilde{d}_{k+1}^{-1} - \varrho(A_k)^{-1} (\tilde{a}_k \tilde{b}_k^{-1} \tilde{d}_{k+1}^{-1} + \varrho(A_{k+1})^{-1}), k \geq 1.$$

Suppose that condition (B1) holds. Then applying Lemma 1, we have

$$f_k \rightarrow b\varrho^{-1}, \quad \xi_k \rightarrow \varrho^{-1} \text{ and hence } \varepsilon_k^f \rightarrow 0, \quad \varepsilon_k^\xi \rightarrow 0 \text{ as } k \rightarrow \infty.$$

The following lemma, which can be proved by some arguments similar to the proofs of [17, Lemma 4] and [18, Lemma 12], studies the fluctuations of both ε_k^f and ε_k^ξ , $k \geq 1$.

Lemma 7. Suppose that condition (B1) and one of the conditions (B2)_a, (B2)_b and (B2)_c hold. Then there exists some number q with $|q| \leq 1$ such that

$$\lim_{k \rightarrow \infty} \delta_{k+1}^f / \delta_k^f = \lim_{k \rightarrow \infty} \delta_{k+1}^\xi / \delta_k^\xi = q,$$

$$\lim_{k \rightarrow \infty} \frac{\varepsilon_{k+1}^\xi}{\varepsilon_k^\xi} = q, \quad \lim_{k \rightarrow \infty} \frac{\varepsilon_{k+1}^f}{\varepsilon_k^f} = q \text{ or } \frac{a\theta - bd}{\varrho^2}.$$

3.3 Proof of Theorem 2.

For $m \geq k \geq 1$ we write $x_{k,m} := \frac{\mathbf{e}_1 A_k \cdots A_m \mathbf{e}_1^t}{\varrho(A_k) \cdots \varrho(A_m)}$ for simplicity. In view of (22) and (23), to prove (11), it suffices to show that $x_{k,m} \rightarrow c(k)$ as $m \rightarrow \infty$ for some $0 < c(k) < \infty$.

To begin with, we show first that for fixed $j \geq 1$, $x_{j,m}$, $m \geq j$ are uniformly bounded away from 0 and infinity. Indeed, using again the notation $A_{m,n} = A_m \cdots A_n$, in view of (20) and (21), for $m \geq n \geq j \geq 1$,

$$\begin{aligned} x_{j,m} &= \frac{\mathbf{e}_1 A_j \cdots A_m \mathbf{e}_1^t}{\varrho(A_j) \cdots \varrho(A_m)} \\ &= \frac{A_{j,n}(11)}{\prod_{i=j}^n \varrho(A_i)} \frac{A_{n+1,m}(11)}{\prod_{i=n+1}^m \varrho(A_i)} + \frac{A_{j,n}(12)}{\prod_{i=j}^n \varrho(A_i)} \frac{A_{n+1,m}(21)}{\prod_{i=n+1}^m \varrho(A_i)} \\ &= \frac{A_{j,n}(11)}{\prod_{i=j}^n \varrho(A_i)} \frac{A_{n+1,m}(11)}{\prod_{i=n+1}^m \varrho(A_i)} \left(1 + \frac{A_{j,n}(12)}{A_{j,n}(11)} \frac{A_{n+1,m}(21)}{A_{n+1,m}(11)} \right) \\ &= x_{j,n} x_{n+1,m} (1 + h_{n,j} \xi_{n+1,m}). \end{aligned}$$

Let $\xi_i, i \geq 1$ be those in Lemma 5. Then by (22)-(24) we get

$$\begin{aligned} \lim_{m \rightarrow \infty} (1 + h_{n,j} \xi_{n+1,m}) &= 1 + h_{n,j} \xi_{n+1}, \\ \lim_{n \rightarrow \infty} (1 + h_{n,j} \xi_{n+1}) &= 1 - \frac{\varrho_1}{\varrho} \in (0, 2). \end{aligned}$$

Thus, we can find a number $N > N_0$ such that $0 < c < 1 + h_{n,j} \xi_{n+1} < 2 - c, \forall n \geq N$, where N_0 is the one in Lemma 2.

Now fix $k > N$. There is a number $N_5 > k$ such that for $m > N_5$,

$$C^{-1} < 1/2(1 + h_{k,j} \xi_{k+1}) < 1 + h_{k,j} \xi_{k+1,m} < 2(1 + h_{k,j} \xi_{k+1}) < C.$$

Then

$$(32) \quad C^{-1} x_{j,k} x_{k+1,m} < x_{j,m} < C x_{j,k} x_{k+1,m}, \forall m > N_5, j \geq 1.$$

But by Lemma 2, we have $C_{k+1}^{-1} < x_{k+1,m} < C_{k+1}, \forall m \geq k+1$, with $C_{k+1} > 0$ certain constant. Let $\zeta_{k+1} = \max\{x_{k+1,m}, k+1 \leq m \leq N_5\}$ and set $C_{j,k} = (C x_{j,k} (\zeta_{k+1} \vee C_{k+1})) \vee \max_{j \leq m \leq N_5} x_{j,m}$, where C is the constant in (32). Then we have

$$(33) \quad C_{j,k}^{-1} < x_{j,m} < C_{j,k}, \forall m \geq j \geq 1.$$

With (33) and Lemma 7 in hands, the rest of the proof of Theorem 2 is more or less similar to the one of Theorem 1 in [17], but there are some differences in details. For convenience of the reader, we provide a complete proof here.

Next, we prove only $x_{1,m} \rightarrow c$ as $m \rightarrow \infty$ for some $0 < c < \infty$, since for $k \geq 2$ the convergence of $x_{k,m}$ as $m \rightarrow \infty$ can be proved similarly. For simplicity, we write $x_{1,m}$ as x_m and taking (33) into account, we keep always in mind that

$$(34) \quad x_m = \frac{\mathbf{e}_1 A_1 \cdots A_m \mathbf{e}_1^t}{\varrho(A_1) \cdots \varrho(A_m)} \text{ and } \forall m \geq 1, c_3 \leq x_m \leq c_4,$$

where $0 < c_3 \leq c_4 < \infty$ are proper constants.

For $n \geq 1$, writing $f_n = \frac{\mathbf{e}_1 A_1 \cdots A_n \mathbf{e}_2^t}{\mathbf{e}_1 A_1 \cdots A_n \mathbf{e}_1^t}$, then

$$(35) \quad f_n = \frac{\tilde{b}_n \mathbf{e}_1 A_1 \cdots A_{n-1} \mathbf{e}_1^t}{\mathbf{e}_1 A_1 \cdots A_{n-1} (\tilde{a}_n \mathbf{e}_1^t + \tilde{d}_n \mathbf{e}_2^t)} = \frac{\tilde{b}_n}{\tilde{a}_n + \tilde{d}_n f_{n-1}}.$$

Iterating (35), we find that $f_n, n \geq 1$ coincide with the ones defined in (30). Therefore, letting $\varepsilon_n^f, n \geq 1$ be the ones in (31), an application of Lemma 7 yields that for some number q with $|q| \leq 1$,

$$(36) \quad \lim_{m \rightarrow \infty} \frac{\varepsilon_{m+1}^f}{\varepsilon_m^f} = q \text{ or } \frac{a\theta - bd}{\varrho^2}.$$

It is easy to check that $0 < \frac{|a\theta - bd|}{\varrho^2} < 1$ since $a\theta - bd \neq 0$.

Case 1: Suppose $|q| < 1$. Then $\lambda_0 := \max\{|q|, \frac{|a\theta - bd|}{\varrho^2}\} < 1$. Fix $0 < \lambda < \lambda_0$. By (36), there exists some $m_0 > 0$ such that $\left| \frac{\varepsilon_{m+1}^f}{\varepsilon_m^f} \right| \leq \lambda$ for all $m > m_0$. Hence

$$(37) \quad \sum_{m=2}^{\infty} |\varepsilon_m^f| = \sum_{m=2}^{\infty} |f_m - \tilde{b}_{m+1} \varrho(A_{m+1})^{-1}| < \infty.$$

Taking (18) into account, for $m \geq 1$ we have

$$(38) \quad \begin{aligned} x_{m+1} - x_m &= \frac{\tilde{a}_{m+1} - \varrho(A_{m+1}) + \tilde{d}_{m+1} \frac{\mathbf{e}_1 A_1 \cdots A_m \mathbf{e}_2^t}{\mathbf{e}_1 A_1 \cdots A_m \mathbf{e}_1^t}}{\frac{\varrho(A_1) \cdots \varrho(A_{m+1})}{\mathbf{e}_1 A_1 \cdots A_m \mathbf{e}_1^t}} \\ &= \varrho(A_{m+1})^{-1} x_m (\tilde{a}_{m+1} - \varrho(A_{m+1}) + \tilde{d}_{m+1} f_m) \\ &= \varrho(A_{m+1})^{-1} x_m \tilde{d}_{m+1} (f_m - \tilde{b}_{m+1} \varrho(A_{m+1})^{-1}) \\ &= \varrho(A_{m+1})^{-1} x_m \tilde{d}_{m+1} \varepsilon_m^f. \end{aligned}$$

Since $\varrho(A_m), m \geq 1$ are uniformly bounded away from 0 and ∞ , then by (34), we have for some constant $0 < c_5 < \infty$,

$$(39) \quad |x_{m+1} - x_m| \leq c_5 |\varepsilon_m^f|, \forall m \geq 1.$$

Taking (34), (37) and (39) together, we conclude that for some constant $0 < c < \infty$, $\lim_{m \rightarrow \infty} x_m = c$.

Case 2: Suppose $q = 1$ and $\lim_{m \rightarrow \infty} \frac{\varepsilon_{m+1}^f}{\varepsilon_m^f} = \frac{a\theta - bd}{\varrho^2}$. Since $\frac{|a\theta - bd|}{\varrho^2} < 1$, the proof goes exactly the same as Case 1.

Case 3: Suppose $q = 1$ and $\lim_{m \rightarrow \infty} \frac{\varepsilon_{m+1}^f}{\varepsilon_m^f} = q$. Then there exists some number $m_1 > 0$ such that $\varepsilon_m^f = f_m - \tilde{b}_{m+1} \varrho(A_{m+1})^{-1}, m \geq m_1$ are all strictly positive or strictly negative, and consequently

$$(40) \quad \frac{\tilde{a}_{m+1} + \tilde{d}_{m+1} f_m}{\tilde{a}_{m+1} + \tilde{d}_{m+1} \tilde{b}_{m+1} \varrho(A_{m+1})^{-1}} < 1 \text{ (or } > 1), \text{ for all } m \geq m_1.$$

But

$$(41) \quad \begin{aligned} \frac{x_{m+1}}{x_m} &= \frac{1}{\varrho(A_{m+1})} \left(\tilde{a}_{m+1} + \tilde{d}_{m+1} \frac{\mathbf{e}_1 A_1 \cdots A_m \mathbf{e}_2^t}{\mathbf{e}_1 A_1 \cdots A_m \mathbf{e}_1^t} \right) \\ &= \frac{1}{\varrho(A_{m+1})} \left(\tilde{a}_{m+1} + \tilde{d}_{m+1} f_m \right) = \frac{\tilde{a}_{m+1} + \tilde{d}_{m+1} f_m}{\tilde{a}_{m+1} + \tilde{d}_{m+1} \tilde{b}_{m+1} \varrho(A_{m+1})^{-1}}. \end{aligned}$$

Thus, by (40), $\frac{x_{m+1}}{x_m} < 1$ (or > 1) for all $m \geq m_1$, that is, $x_m, m \geq m_1$ is monotone in m . As a consequence, it follows from (34) that for some constant $0 < c < \infty$ $\lim_{m \rightarrow \infty} x_m = c$.

Case 4. Suppose that $q = -1$ and $\lim_{m \rightarrow \infty} \frac{\varepsilon_{m+1}^f}{\varepsilon_m^f} = \frac{a\theta - bd}{\varrho^2}$. In this case, the proof is the same as Case 2.

Case 5. Suppose that $q = -1$ and $\lim_{m \rightarrow \infty} \frac{\varepsilon_{m+1}^f}{\varepsilon_m^f} = -1$. Combining (38) with (41), we have

$$\frac{x_{m+1} - x_m}{x_m - x_{m-1}} = \frac{\varrho(A_m)}{\varrho(A_{m+1})} \frac{\tilde{a}_m + \tilde{d}_m f_{m-1}}{\tilde{a}_m + \tilde{d}_m \tilde{b}_m \varrho(A_m)^{-1}} \frac{\tilde{d}_{m+1}}{\tilde{d}_m} \frac{\varepsilon_m^f}{\varepsilon_{m-1}^f} \rightarrow -1,$$

as $m \rightarrow \infty$. So there exists some number $m_2 > 0$ such that

$$\frac{x_{m+1} - x_m}{x_m - x_{m-1}} < 0 \quad \text{for all } m > m_2.$$

Since $\varepsilon_m^f = f_m - \tilde{b}_{m+1} \varrho(A_{m+1})^{-1} \rightarrow 0$ as $m \rightarrow \infty$, then by (38), we have $x_{m+1} - x_m \rightarrow 0$ as $m \rightarrow \infty$. We thus come to the conclusion that $x_{m+1} - x_m$ converges to 0 in an alternating manner as $m \rightarrow \infty$. Therefore,

$$c := \lim_{m \rightarrow \infty} x_m = x_1 + \sum_{m=1}^{\infty} (x_{m+1} - x_m)$$

exists and by (34), we must have $0 < c < \infty$. Theorem 2 is proved. \square

4 Product of matrices and continued fractions

Based on Theorem 2, the purpose of this section is to finish the proof of Theorem 3. Our method is some delicate analysis on continued fractions and their approximants.

4.1 Product of matrices expressed in terms of the approximants of continued fractions

Our approach to prove (12) is to express $\mathbf{e}_1 A_k \cdots A_n \mathbf{e}_1^t$ in terms of the approximants of some continued fractions. For $1 \leq k \leq n$, set

$$(42) \quad y_{k,n} := \mathbf{e}_1 A_k \cdots A_n \mathbf{e}_1^t \quad \text{and} \quad \xi_{k,n} := \frac{y_{k+1,n}}{y_{k,n}}.$$

Noting that the empty product equals identity, thus $y_{n+1,n} = 1$. Therefore,

$$(43) \quad \xi_{k,n}^{-1} \cdots \xi_{n,n}^{-1} = y_{k,n} = \mathbf{e}_1 A_k \cdots A_n \mathbf{e}_1^t,$$

$$(44) \quad \sum_{k=1}^{n+1} \mathbf{e}_1 A_k \cdots A_n \mathbf{e}_1^t = \sum_{k=1}^{n+1} \xi_{k,n}^{-1} \cdots \xi_{n,n}^{-1} = \frac{\sum_{k=1}^{n+1} \xi_{1,n} \cdots \xi_{k-1,n}}{\xi_{1,n} \cdots \xi_{n,n}}.$$

Lemma 8. For $1 \leq k \leq n$, $\xi_{k,n}$ defined in (42) coincides with the one in (13) with $\beta_k = \tilde{b}_k^{-1} \tilde{d}_{k+1}^{-1}$ and $\alpha_k = \tilde{a}_k \tilde{b}_k^{-1} \tilde{d}_{k+1}^{-1}$.

Proof. Clearly, $\xi_{n,n} = \frac{1}{y_{n,n}} = \frac{1}{\tilde{a}_n} = \frac{\tilde{b}_n^{-1} \tilde{d}_{n+1}^{-1}}{\tilde{a}_n \tilde{b}_n^{-1} \tilde{d}_{n+1}^{-1}}$. For $1 \leq k < n$, note that

$$(45) \quad \begin{aligned} \xi_{k,n} &= \frac{y_{k+1,n}}{y_{k,n}} = \frac{\mathbf{e}_1 A_{k+1} \cdots A_n \mathbf{e}_1^t}{\mathbf{e}_1 A_k \cdots A_n \mathbf{e}_1^t} = \frac{\mathbf{e}_1 A_{k+1} \cdots A_n \mathbf{e}_1^t}{(\tilde{a}_k \mathbf{e}_1 + \tilde{b}_k \mathbf{e}_2) A_{k+1} \cdots A_n \mathbf{e}_1^t} \\ &= \frac{1}{\tilde{a}_k + \tilde{b}_k \frac{\mathbf{e}_2 A_{k+1} \cdots A_n \mathbf{e}_1^t}{\mathbf{e}_1 A_{k+1} \cdots A_n \mathbf{e}_1^t}} = \frac{1}{\tilde{a}_k + \tilde{b}_k \tilde{d}_{k+1} \frac{\mathbf{e}_1 A_{k+2} \cdots A_n \mathbf{e}_1^t}{\mathbf{e}_1 A_{k+1} \cdots A_n \mathbf{e}_1^t}} \end{aligned}$$

$$= \frac{\tilde{b}_k^{-1} \tilde{d}_{k+1}^{-1}}{\tilde{a}_k \tilde{b}_k^{-1} \tilde{d}_{k+1}^{-1} + \xi_{k+1,n}}.$$

Thus, the lemma can be proved by iterating (45). \square

In the remainder of this section, we always assume that all conditions of Theorem 3 hold and $\xi_k, \xi_{k,n}, n \geq k \geq 1$ are those defined in (13) and (14) with $\beta_k = \tilde{b}_k^{-1} \tilde{d}_{k+1}^{-1}$ and $\alpha_k = \tilde{a}_k \tilde{b}_k^{-1} \tilde{d}_{k+1}^{-1}$. Then we have

$$\lim_{k \rightarrow \infty} \beta_k =: \beta = (bd - a\theta)^{-1} \neq 0 \text{ and } \lim_{k \rightarrow \infty} \alpha_k =: \alpha = \frac{a + \theta}{bd - a\theta} \neq 0.$$

Consequently, it follows from Lemma 1 that

$$(46) \quad \lim_{n \rightarrow \infty} \xi_{k,n} = \xi_k, \lim_{k \rightarrow \infty} \xi_k =: \xi = \frac{\alpha}{2} \left(\sqrt{1 + 4\beta/\alpha^2} - 1 \right) = \varrho^{-1} > 0.$$

Moreover, by (15) and (42) we have

$$(47) \quad \xi_k > 0, \xi_{k,n} > 0, \forall n \geq k \geq 1.$$

In view of (43) and (44), the following proposition is crucial to prove Theorem 3.

Proposition 1. *As $n \rightarrow \infty$, we have*

$$(48) \quad \xi_{1,n} \cdots \xi_{n,n} \sim c \varrho(A_1)^{-1} \cdots \varrho(A_n)^{-1},$$

$$(49) \quad \xi_{1,n} \cdots \xi_{n,n} \sim c \xi_1 \cdots \xi_n,$$

$$(50) \quad \sum_{k=1}^{n+1} \xi_{1,n} \cdots \xi_{k-1,n} \sim c \sum_{k=1}^{n+1} \xi_1 \cdots \xi_{k-1}.$$

Proof. To begin with, taking (15) and (43) into consideration, applying Theorem 2, we obtain (48). Since $bd \neq a\theta$ and $\lim_{k \rightarrow \infty} \tilde{d}_k = b^{-1}(bd - a\theta) \neq 0$, there exist $k_0 > 0$ and $\varepsilon > 0$ such that

$$(51) \quad \text{either } \tilde{d}_k \leq -\varepsilon, \forall k \geq k_0 \text{ or } \tilde{d}_k \geq \varepsilon, \forall k \geq k_0.$$

In view of (46) and (47), if we can show that for some $k_0 > 0$, $\xi_{k_0,n} \cdots \xi_{n,n} \sim c \xi_{k_0} \cdots \xi_n$ and $\sum_{k=k_0}^{n+1} \xi_{k_0,n} \cdots \xi_{k-1,n} \sim c \sum_{k=k_0}^{n+1} \xi_{k_0} \cdots \xi_{k-1}$ as $n \rightarrow \infty$, then (49) and (50) are also true. Therefore, instead of (51), we assume $\exists \varepsilon > 0$ such that either $\tilde{d}_k \leq -\varepsilon, \forall k \geq 1$ or $\tilde{d}_k \geq \varepsilon, \forall k \geq 1$. If $\tilde{d}_k \geq \varepsilon, \forall k \geq 1$, then Proposition 1 has been proved in [18], see Lemma 10 therein. Thus we need only to deal with the case

$$(52) \quad \exists \varepsilon > 0 \text{ such that } \tilde{d}_k \leq -\varepsilon, \forall k \geq 1.$$

To prove (49) and (50), we need the following two lemmas, whose proof will be postponed to the end of this subsection.

Lemma 9. *Suppose that (52) holds and fix $k \geq 1$. Then $\xi_{k,n}, n \geq k$ is monotone increasing in n and thus $\xi_{k,n} < \xi_k, n \geq k$.*

Lemma 10. *Suppose that (52) holds. Then $\exists 0 < r < 1$ and $k_1 > 0$ such that $\frac{\xi_{k,n} - \xi_k}{\xi_{k+1,n} - \xi_{k+1}} \leq r, \forall n > k > k_1$, and consequently*

$$\xi_k - \xi_{k,n} \leq r^{n-k} (\xi_n - \xi_{n,n}), \forall n \geq k > k_1.$$

With Lemma 9 and Lemma 10 in hands, we show next that

$$(53) \quad \exists c > 0 \text{ such that } c < \frac{\xi_{1,n} \cdots \xi_{n,n}}{\xi_1 \cdots \xi_n} < 1, n \geq 1.$$

Indeed, applying Lemma 9, we get

$$(54) \quad \xi_{1,n} \xi_{2,n} \cdots \xi_{n,n} < \xi_1 \xi_2 \cdots \xi_n, n \geq 1.$$

For a lower bound, notice that $\xi_{k,n} > 0, \forall n \geq k > 1$ and $\xi_{n,n} = \tilde{a}_n^{-1} \rightarrow (a + \theta)^{-1}$ as $n \rightarrow \infty$. Then by (46), (47) and Lemma 9, we have

$$(55) \quad c^{-1} < \xi_n < c \text{ and } c^{-1} < \xi_n - \xi_{n,n} < c, \forall n \geq 1,$$

for some number $c > 0$. Therefore, letting r be the one in Lemma 10, there is a number $k_2 > 0$ such that

$$1 - \frac{(\xi_n - \xi_{n,n})r^{k_2}}{\inf_{k \geq 1} \xi_k} > 0, \forall n \geq 1.$$

With k_1 the number in Lemma 10, applying Lemma 10, we have for n large enough,

$$\begin{aligned} \frac{\xi_{k_1+1,n} \cdots \xi_{n,n}}{\xi_{k_1+1} \cdots \xi_n} &= \frac{(\xi_{k_1+1,n} - \xi_{k_1+1})}{\xi_{k_1+1}} \frac{\xi_{k_1+2,n} \cdots \xi_{n,n}}{\xi_{k_1+2} \cdots \xi_n} + \frac{\xi_{k_1+2,n} \cdots \xi_{n,n}}{\xi_{k_1+2} \cdots \xi_n} \\ &\geq \left(1 - \frac{(\xi_n - \xi_{n,n})r^{n-k_1-1}}{\xi_{k_1+1}}\right) \frac{\xi_{k_1+2,n} \cdots \xi_{n,n}}{\xi_{k_1+2} \cdots \xi_n}, \end{aligned}$$

which implies $\liminf_{n \rightarrow \infty} \frac{\xi_{k_1+1,n} \cdots \xi_{n,n}}{\xi_{k_1+1} \cdots \xi_n} \geq \liminf_{n \rightarrow \infty} \frac{\xi_{k_1+2,n} \cdots \xi_{n,n}}{\xi_{k_1+2} \cdots \xi_n}$. Using the same tricky times and again, we get

$$\liminf_{n \rightarrow \infty} \frac{\xi_{k_1+1,n} \cdots \xi_{n,n}}{\xi_{k_1+1} \cdots \xi_n} > \liminf_{n \rightarrow \infty} \frac{\xi_{n-k_2,n} \cdots \xi_{n,n}}{\xi_{n-k_2} \cdots \xi_n}.$$

Consequently, since $\lim_{n \rightarrow \infty} \xi_{k,n} = \xi_k$ and $\lim_{n \rightarrow \infty} \frac{\xi_{n-k_2,n} \cdots \xi_{n,n}}{\xi_{n-k_2} \cdots \xi_n} = c$ for some number $c > 0$, we have

$$(56) \quad \liminf_{n \rightarrow \infty} \frac{\xi_{1,n} \cdots \xi_{n,n}}{\xi_1 \cdots \xi_n} = \liminf_{n \rightarrow \infty} \frac{\xi_{k_1+1,n} \cdots \xi_{n,n}}{\xi_{k_1+1} \cdots \xi_n} \geq \liminf_{n \rightarrow \infty} \frac{\xi_{n-k_2,n} \cdots \xi_{n,n}}{\xi_{n-k_2} \cdots \xi_n} = c > 0.$$

Taking (54) and (56) together, we finish the proof of (53).

Now, we are ready to prove (49). In view of (48), it suffices to show

$$(57) \quad \xi_1 \cdots \xi_n \sim \varrho(A_1)^{-1} \cdots \varrho(A_n)^{-1}, \text{ as } n \rightarrow \infty.$$

To this end, write $x_n = \frac{\xi_1 \cdots \xi_n}{\varrho(A_1)^{-1} \cdots \varrho(A_n)^{-1}}, n \geq 1$. Then taking (48) and (53) together, we get

$$(58) \quad C^{-1} < x_n < C, \forall n \geq 1.$$

With (58) in hand, using Lemma 7 and mimicking the proof of the convergence of x_k in the proof of Theorem 2, we can show that $\lim_{n \rightarrow \infty} x_n = c$ for some number $c > 0$. Consequently, (57) is proved and so is (49).

Finally, we turn to prove (50). For the case $\varrho < 1$, the proof is the same as the one in [18, Lemma 10]. Thus we need only to prove the case $\varrho \geq 1$. Suppose now $\varrho \geq 1$. In

order to prove (50), it suffices to show $\lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n \xi_{1,n} \cdots \xi_{k,n} - \sum_{k=1}^n \xi_1 \cdots \xi_k}{\sum_{k=1}^n \xi_1 \cdots \xi_k} = 0$ which is equivalent to

$$(59) \quad \lim_{n \rightarrow \infty} \frac{\sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_{k,n}}{\sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_k} = 1$$

since $\lim_{k \rightarrow \infty} \xi_{k,n} = \xi_k > 0$, where k_1 is the number in Lemma 10. To prove (59), on one hand, note that by Lemma 9 we have $\xi_{k,n} < \xi_k, \forall n \geq k \geq 1$ which implies

$$(60) \quad \sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_{k,n} < \sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_k, \forall n \geq k_1.$$

On the other hand, taking the lemmas 9 and 10 into account, we have for $n \geq k_1$

$$\begin{aligned} \sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_{k,n} &= \xi_{k_1+1} \sum_{j=1}^{n-k_1} \prod_{k=k_1+2}^{k_1+j} \xi_{k,n} + (\xi_{k_1+1,n} - \xi_{k_1+1}) \sum_{j=1}^{n-k_1} \prod_{k=k_1+2}^{k_1+j} \xi_{k,n} \\ &\geq \xi_{k_1+1} + \xi_{k_1+1} \sum_{j=2}^{n-k_1} \prod_{k=k_1+2}^{k_1+j} \xi_{k,n} + (\xi_{n,n} - \xi_n) r^{n-(k_1+1)} \sum_{j=2}^{n-k_1} \prod_{k=k_1+2}^{k_1+j} \xi_k. \end{aligned}$$

Iterating the above inequality, we get

$$(61) \quad \begin{aligned} \sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_{k,n} &> \sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_k \\ &+ (\xi_{n,n} - \xi_n) \sum_{i=0}^{n-(k_1+1)} \left(\prod_{k=k_1+1}^{k_1+i} \xi_k \right) r^{n-(k_1+i+1)} \sum_{j=i+1}^{n-k_1} \prod_{k=k_1+i+2}^{k_1+j} \xi_k \\ &=: \text{(I)} + \text{(II)}. \end{aligned}$$

It follows from (55) that the absolute value of the second term (II) equals to

$$(62) \quad \begin{aligned} (\xi_n - \xi_{n,n}) \sum_{i=0}^{n-(k_1+1)} \frac{r^{n-(k_1+i+1)}}{\xi_{k_1+i+1}} \left(\prod_{k=k_1+1}^{k_1+i+1} \xi_k \right) \sum_{j=i+1}^{n-k_1} \prod_{k=k_1+i+2}^{k_1+j} \xi_k \\ \leq c \sum_{i=0}^{n-(k_1+1)} r^{n-(k_1+i+1)} \left(\prod_{k=k_1+1}^{k_1+i+1} \xi_k \right) \sum_{j=i+1}^{n-k_1} \prod_{k=k_1+i+2}^{k_1+j} \xi_k. \end{aligned}$$

Clearly, for any $0 \leq i \leq n - (k_1 + 1)$,

$$(63) \quad \frac{\left(\prod_{k=k_1+1}^{k_1+i+1} \xi_k \right) \sum_{j=i+1}^{n-k_1} \prod_{k=k_1+i+2}^{k_1+j} \xi_k}{\sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_k} \leq 1$$

and an application of [18, Lemma 8] yields that

$$(64) \quad \lim_{n \rightarrow \infty} \frac{\prod_{k=k_1+1}^{n-i} \xi_k}{\sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_k} = 0, \forall i \geq 1,$$

since $\xi_k > 0, \forall k \geq 1$ and $\lim_{k \rightarrow \infty} \xi_k = \varrho^{-1} \leq 1$.

Let $\varepsilon > 0$ be an arbitrary number and fix k_3 such that $r^{k_3} < \varepsilon$. Then using (63) and (64), we have from (62) that

$$\begin{aligned} \overline{\lim}_{n \rightarrow \infty} \frac{|(\text{II})|}{\sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_k} &\leq c \overline{\lim}_{n \rightarrow \infty} \sum_{i=0}^{n-(k_1+1)-k_3} r^{n-(k_1+i+1)} \\ &+ c \overline{\lim}_{n \rightarrow \infty} \sum_{i=n-k_1-k_3}^{n-(k_1+1)} r^{n-(k_1+i+1)} \frac{\left(\prod_{k=k_1+1}^{k_1+i+1} \xi_k \right) \sum_{j=i+1}^{n-k_1} \prod_{k=k_1+i+2}^{k_1+j} \xi_k}{\sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_k} \\ &= c \overline{\lim}_{n \rightarrow \infty} \sum_{i=0}^{n-(k_1+1)-k_3} r^{n-(k_1+i+1)} = cr^{k_3}/(1-r) \\ &\leq c\varepsilon/(1-r). \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, $\overline{\lim}_{n \rightarrow \infty} \frac{|(\text{II})|}{\sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_k} = 0$, which together with (61) implies that

$$(65) \quad \underline{\lim}_{n \rightarrow \infty} \frac{\sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_{k,n}}{\sum_{j=1}^{n-k_1} \prod_{k=k_1+1}^{k_1+j} \xi_k} = 1.$$

Putting (60) and (65) together, we can infer that (59) is true and thus Proposition 1 is proved. \square

To end this subsection, we give the proof of Lemma 9 and Lemma 10.

Proof of Lemma 9. We see from (15) and (42) that $\xi_{k,n} > 0$ for all $n \geq k \geq 1$. Since $\forall k \geq 1$, $\tilde{a}_k > 0, \tilde{b}_k > 0$ and $\tilde{d}_k < -\varepsilon < 0$ for some $\varepsilon > 0$, we always have $\alpha_k < 0, \beta_k < 0, \forall k \geq 1$. Now fix $k \geq 1$ and for $n \geq k$, write

$$(66) \quad C_{k,n} = \alpha_n C_{k,n-1} + \beta_n C_{k,n-2}, \quad D_{k,n} = \alpha_n D_{k,n-1} + \beta_n D_{k,n-2}$$

with initial values

$$(67) \quad C_{k-2} = D_{k-1} = 1, C_{k-1} = D_{k-2} = 0.$$

Then by Euler-Minding formula (see [14], (1.2.14) on page 7) we obtain

$$\xi_{k,n} = \frac{C_{k,n}}{D_{k,n}} = - \sum_{j=k}^n \frac{\prod_{i=k}^j (-\beta_i)}{D_{k,j} D_{k,j-1}}$$

which leads to

$$(68) \quad \xi_{k,n+1} - \xi_{k,n} = - \frac{\prod_{i=k}^{n+1} (-\beta_j)}{D_{k,n+1} D_{k,n}}, n \geq k.$$

By induction, we have from (66) and (67) that

$$\begin{aligned} \begin{pmatrix} D_{k,n} & D_{k,n-1} \\ C_{k,n} & C_{k,n-1} \end{pmatrix} &= \begin{pmatrix} \alpha_k & 1 \\ \beta_k & 0 \end{pmatrix} \cdots \begin{pmatrix} \alpha_n & 1 \\ \beta_n & 0 \end{pmatrix} \\ &= \left(\prod_{i=k}^n \tilde{b}_i \tilde{d}_{i+1} \right)^{-1} \begin{pmatrix} \tilde{a}_k & \tilde{b}_k \tilde{d}_{k+1} \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} \tilde{a}_n & \tilde{b}_n \tilde{d}_{n+1} \\ 1 & 0 \end{pmatrix} \end{aligned}$$

$$= \left(\prod_{i=k}^n \tilde{b}_i \tilde{d}_{i+1} \right)^{-1} \begin{pmatrix} 1 & 0 \\ 0 & \tilde{d}_k^{-1} \end{pmatrix} A_k \cdots A_n \begin{pmatrix} 1 & 0 \\ 0 & \tilde{d}_{n+1} \end{pmatrix}$$

where for the third equality we use the facts

$$\begin{pmatrix} \tilde{a}_i & \tilde{b}_i \tilde{d}_{i+1} \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} \tilde{a}_i & \tilde{b}_i \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & \tilde{d}_{i+1} \end{pmatrix},$$

$$A_i = \begin{pmatrix} \tilde{a}_i & \tilde{b}_i \\ \tilde{d}_i & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & \tilde{d}_i \end{pmatrix} \begin{pmatrix} \tilde{a}_i & \tilde{b}_i \\ 1 & 0 \end{pmatrix}, i \geq 1.$$

As a consequence, taking (15) and (16) into account we get

$$(69) \quad \frac{D_{k,n-1}}{D_{k,n}} = \tilde{d}_{n+1} \frac{\mathbf{e}_1 A_k \cdots A_n \mathbf{e}_2}{\mathbf{e}_1 A_k \cdots A_n \mathbf{e}_1} < 0, \forall n \geq k.$$

Putting (68) and (69) together, we conclude that

$$\xi_{k,n+1} - \xi_{k,n} > 0, \forall n \geq k \geq 1,$$

that is to say, $\xi_{k,n}, n \geq k$ is monotone increasing in n . Lemma 9 is proved. \square

Proof of Lemma 10. Taking Lemma 9 into account, since $\xi_{k,n} > 0, \forall n \geq k \geq 1$, then by some easy computation, we have from (13) and (14) that

$$0 < \frac{\xi_{k,n} - \xi_k}{\xi_{k+1,n} - \xi_{k+1}} = -\frac{\xi_{k,n}}{\alpha_k + \xi_{k+1}} < -\frac{\xi_k}{\alpha_k + \xi_{k+1}}.$$

But $-\frac{\xi_k}{\alpha_k + \xi_{k+1}} \rightarrow -\frac{\xi}{\alpha + \xi} = \frac{\alpha + \sqrt{\alpha^2 + 4\beta}}{\alpha - \sqrt{\alpha^2 + 4\beta}} < 1$ as $k \rightarrow \infty$. As a result, for some proper number $0 < r < 1, \exists k_1 > 0$ such that $-\frac{\xi_k}{\alpha_k + \xi_{k+1}} < r, \forall k > k_1$, which finishes the proof of the lemma. \square

4.2 Proof of Theorem 3

Theorem 3 is a direct consequence of Proposition 1. In fact, it follows from (44) and Proposition 1 that

$$\sum_{k=1}^{n+1} \mathbf{e}_1 A_k \cdots A_n \mathbf{e}_1^t = \frac{\sum_{k=1}^{n+1} \xi_{1,n} \cdots \xi_{k-1,n}}{\xi_{1,n} \cdots \xi_{n,n}} \sim c \frac{\sum_{k=1}^{n+1} \xi_1 \cdots \xi_{k-1}}{\varrho(A_1)^{-1} \cdots \varrho(A_n)^{-1}}, \text{ as } n \rightarrow \infty.$$

But by (48) and (49), we obtain

$$\xi_1 \cdots \xi_n \sim c \varrho(A_1)^{-1} \cdots \varrho(A_n)^{-1}, \text{ as } n \rightarrow \infty$$

which implies that

$$\sum_{k=1}^{n+1} \xi_1 \cdots \xi_{k-1} \sim c \sum_{k=1}^{n+1} \varrho(A_1)^{-1} \cdots \varrho(A_{k-1})^{-1}, \text{ as } n \rightarrow \infty.$$

As a result, we have

$$\sum_{k=1}^{n+1} \mathbf{e}_1 A_k \cdots A_n \mathbf{e}_1^t \sim c \frac{\sum_{k=1}^{n+1} \varrho(A_1)^{-1} \cdots \varrho(A_{k-1})^{-1}}{\varrho(A_1)^{-1} \cdots \varrho(A_n)^{-1}} = c \sum_{k=1}^{n+1} \varrho(A_k) \cdots \varrho(A_n)$$

as $n \rightarrow \infty$. Thus Theorem 3 is proved. \square

5 Proof of Theorem 1

For $n \geq 0$ and $\mathbf{s} = (s_1, s_2)^t \in [0, 1]^2$, let

$$F_n^{(i)}(\mathbf{s}) \equiv E(\mathbf{s}^{Z_n} | Z_0 = \mathbf{e}_i) := E(s_1^{Z_n,1} s_2^{Z_n,2} | Z_0 = \mathbf{e}_i), i = 1, 2$$

and set $\mathbf{F}_n(\mathbf{s}) = (F_n^{(1)}(\mathbf{s}), F_n^{(2)}(\mathbf{s}))^t$. It follows by induction (see Dyakonova [5, Lemma 1]) that for $n \geq 0$,

$$\mathbf{F}_n(\mathbf{s}) = \mathbf{f}_1(\mathbf{f}_2(\cdots \mathbf{f}_n(\mathbf{s}) \cdots)) = \mathbf{1} - \frac{\prod_{k=1}^n M_k(\mathbf{1} - \mathbf{s})}{1 + \sum_{k=1}^n \gamma_k \prod_{i=k+1}^n M_i(\mathbf{1} - \mathbf{s})}$$

which leads to

$$\mathbf{F}_n(\mathbf{0}) = \mathbf{1} - \frac{\prod_{k=1}^n M_k \mathbf{1}}{1 + \sum_{k=1}^n \gamma_k \prod_{i=k+1}^n M_i \mathbf{1}}.$$

For the asymptotics of $P(\nu > n | Z_0 = \mathbf{e}_1)$ and $P(\nu = n | Z_0 = \mathbf{e}_1)$, we refer the reader to [18]. Here we treat only $P(\nu > n | Z_0 = \mathbf{e}_2)$ and $P(\nu = n | Z_0 = \mathbf{e}_2)$. Taking (3) into account, for $n \geq 1$ we have

$$(70) \quad P(\nu > n | Z_0 = \mathbf{e}_2) = 1 - F_n^{(2)}(\mathbf{0}) = \frac{\mathbf{e}_2 \prod_{k=1}^n M_k \mathbf{1}}{\sum_{k=1}^{n+1} \mathbf{e}_1 \prod_{i=k}^n M_i \mathbf{1}},$$

and consequently

$$(71) \quad P(\nu = n | Z_0 = \mathbf{e}_2) = \frac{\mathbf{e}_2 \prod_{k=1}^{n-1} M_k \mathbf{1}}{\sum_{k=1}^n \mathbf{e}_1 \prod_{i=k}^{n-1} M_i \mathbf{1}} - \frac{\mathbf{e}_2 \prod_{k=1}^n M_k \mathbf{1}}{\sum_{k=1}^{n+1} \mathbf{e}_1 \prod_{i=k}^n M_i \mathbf{1}}.$$

With matrices $A_i, i \geq 1$ the ones defined in (4), using (5), by some very careful computation, we have from (70) and (71) that

$$(72) \quad P(\nu > n | Z_0 = \mathbf{e}_2) = \frac{(\theta_1/b_1, 1) \prod_{k=1}^n A_k(1, \lambda_{n+1})^t}{\sum_{k=1}^{n+1} \mathbf{e}_1 \prod_{i=k}^n A_i(1, \lambda_{n+1})^t},$$

$$(73) \quad P(\nu = n | Z_0 = \mathbf{e}_2) = \frac{1}{\sum_{k=1}^{n+1} \mathbf{e}_1 \prod_{i=k}^n A_i(1, \lambda_{n+1})^t} \frac{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t}{\sum_{k=1}^n \mathbf{e}_1 \prod_{i=k}^{n-1} A_i(1, \lambda_n)^t} \\ \times \left(\frac{\theta_1}{b_1} G_{n-1,1} + \frac{\mathbf{e}_2 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t} G_{n-1,2} \right)$$

where for $n \geq 1$, $\lambda_n \equiv 1 - \frac{\theta_n}{b_n}$,

$$(74) \quad G_{n-1,1} \equiv \frac{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k(1, \lambda_n)^t \sum_{k=1}^{n+1} \mathbf{e}_1 \prod_{i=k}^n A_i(1, \lambda_{n+1})^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t} \\ - \frac{\mathbf{e}_1 \prod_{k=1}^n A_k(1, \lambda_{n+1})^t \sum_{k=1}^n \mathbf{e}_1 \prod_{i=k}^{n-1} A_i(1, \lambda_n)^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t}$$

and

$$(75) \quad G_{n-1,2} \equiv \frac{\mathbf{e}_2 \prod_{k=1}^{n-1} A_k(1, \lambda_n)^t \sum_{k=1}^{n+1} \mathbf{e}_1 \prod_{i=k}^n A_i(1, \lambda_{n+1})^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t} \\ - \frac{\mathbf{e}_2 \prod_{k=1}^n A_k(1, \lambda_{n+1})^t \sum_{k=1}^n \mathbf{e}_1 \prod_{i=k}^{n-1} A_i(1, \lambda_n)^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t}.$$

$G_{n,1}$ and $G_{n,2}$ defined in (74) and (75) look very complicated. Next lemma shows that they converge to the same limit.

Lemma 11. *Suppose that condition (B1) holds and $|\varrho_1| < 1$. Then, there exists a number $G \geq 0$ such that*

$$(76) \quad \lim_{n \rightarrow \infty} G_{n,2} = G = \lim_{n \rightarrow \infty} G_{n,1}.$$

Moreover, $G \neq 0$ if and only if $\varrho_1 \neq \frac{1}{2} \left(a + b + 1 - \sqrt{(a + b + 1)^2 + 4 \frac{bd - a\theta}{\theta - b}} \right)$.

Proof. The proof of the second equality in (76) can be found in [18], so we check here only the first one. To this end, set

$$f_n \equiv \frac{\mathbf{e}_2 \prod_{k=1}^n A_k \mathbf{e}_2^t}{\mathbf{e}_2 \prod_{k=1}^n A_k \mathbf{e}_1^t} \text{ and } H_n \equiv \sum_{k=1}^n \mathbf{e}_1 \prod_{i=k}^n A_i (f_n \mathbf{e}_1^t - \mathbf{e}_2^t), n \geq 1.$$

Clearly, we have $f_1 = 0, H_1 = -\tilde{b}_1 = -b_1$ and by some subtle computation,

$$(77) \quad \begin{aligned} G_{n-1,2} &= 1 + (\tilde{b}_n \lambda_n \lambda_{n+1} + \tilde{a}_n \lambda_n - \tilde{d}_n) H_{n-1} \\ &\quad + (\tilde{b}_n \lambda_n \lambda_{n+1} + \tilde{a}_n \lambda_n - \tilde{d}_n + \lambda_n) f_{n-1}. \end{aligned}$$

Note that

$$(78) \quad f_n = \frac{\mathbf{e}_2 A_1 \cdots A_n \mathbf{e}_2^t}{\mathbf{e}_2 A_1 \cdots A_n \mathbf{e}_1^t} = \frac{\tilde{b}_n \mathbf{e}_2 A_1 \cdots A_{n-1} \mathbf{e}_1^t}{\mathbf{e}_2 A_1 \cdots A_{n-1} (\tilde{a}_n \mathbf{e}_1^t + \tilde{d}_n \mathbf{e}_2^t)} = \frac{\tilde{b}_n}{\tilde{a}_n + \tilde{d}_n f_{n-1}},$$

which leads to

$$\tilde{d}_n f_n f_{n-1} = \tilde{b}_n - \tilde{a}_n f_n, n \geq 2.$$

Consequently, for $n \geq 2$,

$$(79) \quad \begin{aligned} H_n &= \sum_{k=1}^n \mathbf{e}_1 \prod_{i=k}^n A_i (f_n \mathbf{e}_1^t - \mathbf{e}_2^t) \\ &= \tilde{a}_n f_n - \tilde{b}_n + \sum_{k=1}^{n-1} \mathbf{e}_1 \prod_{i=k}^{n-1} A_i ((\tilde{a}_n f_n - \tilde{b}_n) \mathbf{e}_1^t + \tilde{d}_n f_n \mathbf{e}_2^t) \\ &= -\tilde{d}_n f_n f_{n-1} - \tilde{d}_n f_n \sum_{k=1}^{n-1} \mathbf{e}_1 \prod_{i=k}^{n-1} A_i (f_{n-1} \mathbf{e}_1^t - \mathbf{e}_2^t) \\ &= -\tilde{d}_n f_n f_{n-1} - \tilde{d}_n f_n H_{n-1}. \end{aligned}$$

Since $f_1 = 0$ and $H_1 = -b_1$, iterating (79), we get

$$(80) \quad H_n = (-1)^n b_1 \tilde{d}_2 f_2 \cdots \tilde{d}_n f_n + \sum_{k=1}^{n-1} (-1)^{n-k} f_k \tilde{d}_{k+1} f_{k+1} \cdots \tilde{d}_n f_n, n \geq 2$$

and iterating (78), we get

$$f_n = \frac{\tilde{b}_n \tilde{d}_n^{-1}}{\tilde{a}_n \tilde{d}_n^{-1}} + \frac{\tilde{b}_{n-1} \tilde{d}_{n-1}^{-1}}{\tilde{a}_{n-1} \tilde{d}_{n-1}^{-1}} + \cdots + \frac{\tilde{b}_2 \tilde{d}_2^{-1}}{\tilde{a}_2 \tilde{d}_2^{-1}}, n \geq 2.$$

Noticing that $\lim_{n \rightarrow \infty} \tilde{b}_n \tilde{d}_n^{-1} = b^2 (bd - a\theta)^{-1} \neq 0$ and $\lim_{n \rightarrow \infty} \tilde{a}_n \tilde{d}_n^{-1} = b(a + \theta)(bd - a\theta)^{-1}$, thus applying Lemma 1, we get

$$(81) \quad \lim_{n \rightarrow \infty} f_n = -\frac{b\varrho_1}{bd - a\theta}$$

and thus $\lim_{n \rightarrow \infty} \tilde{d}_n f_n = -\varrho_1$. Since $|\varrho_1| < 1$, we conclude from (80) that

$$(82) \quad \lim_{n \rightarrow \infty} H_n = -\frac{b\varrho_1^2}{(bd - a\theta)(1 - \varrho_1)}.$$

Letting $n \rightarrow \infty$ in (77), owing to (81) and (82), we get

$$\lim_{n \rightarrow \infty} G_{n-1,2} = G := \frac{(b - \theta)\varrho_1^2 - (b - \theta)(a + b + 1)\varrho_1 + bd - a\theta}{(bd - a\theta)(1 - \varrho_1)}.$$

Consequently, the first equality in (76) is true. Moreover, since $|\varrho_1| < 1$, then by the definition of G , it is easy to see that

$$G \neq 0 \text{ if and only if } \varrho_1 \neq \frac{1}{2} \left(a + b + 1 - \sqrt{(a + b + 1)^2 + 4\frac{bd - a\theta}{\theta - b}} \right).$$

What's left for us to do is to check the nonnegativity of the number G . For this purpose, applying (76) and taking (15)-(17) and Theorem 2 into account, we have

$$(83) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \left(\frac{\theta_1}{b_1} G_{n-1,1} + \frac{\mathbf{e}_2 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t} G_{n-1,2} \right) \\ &= G \lim_{n \rightarrow \infty} \left(\frac{\theta_1}{b_1} + \frac{\mathbf{e}_2 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t} \right) = GL \end{aligned}$$

exists, where using again (15)-(17) and Theorem 2, we see that

$$\begin{aligned} L &:= \lim_{n \rightarrow \infty} \left(\frac{\theta_1}{b_1} + \frac{\mathbf{e}_2 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t} \right) = \lim_{n \rightarrow \infty} \frac{\mathbf{e}_2 \prod_{k=1}^{n-1} M_k(1, \theta_n/b_n)^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} M_k(1, \theta_n/b_n)^t} \\ &= \lim_{n \rightarrow \infty} \frac{\mathbf{e}_2 \prod_{k=1}^{n-1} M_k(1, \theta_n/b_n)^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t} \geq \lim_{n \rightarrow \infty} \frac{d_1 d_{n-1} \mathbf{e}_1 \prod_{k=2}^{n-2} M_k \mathbf{e}_2^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t} \\ &= \lim_{n \rightarrow \infty} \frac{d_1 d_{n-1} \mathbf{e}_1 \prod_{k=2}^{n-2} A_k \mathbf{e}_2^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t} > 0. \end{aligned}$$

But in view of (73), since $\sum_{k=1}^n \mathbf{e}_1 \prod_{i=k}^{n-1} A_i(1, \lambda_n)^t = \sum_{k=1}^n \mathbf{e}_1 \prod_{i=k}^{n-1} M_i \mathbf{1} > 0$ and $\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t = \mathbf{e}_1 \prod_{k=1}^{n-1} M_k(1, \theta_{k+1}/b_{k+1})^t > 0$, we must have $GL \geq 0$ and consequently, $G \geq 0$. The lemma is proved. \square

Next we continue with the proof of Theorem 1. For $n \geq 1$, write

$$S_n := \frac{\sum_{k=1}^{n+1} \varrho(A_1)^{-1} \cdots \varrho(A_{k-1})^{-1}}{\varrho(A_1)^{-1} \cdots \varrho(A_n)^{-1}}.$$

We need in addition the following lemma.

Lemma 12. *Under the conditions of Theorem 1, we have*

$$(84) \quad \mathbf{e}_1 \prod_{i=1}^n A_n \mathbf{e}_1^t \sim c\varrho(A_1) \cdots \varrho(A_n),$$

$$(85) \quad \mathbf{e}_1 \prod_{i=1}^n A_n \mathbf{e}_2^t \sim c\varrho(A_1) \cdots \varrho(A_n),$$

$$(86) \quad (\theta_1/b_1, 1) \prod_{k=1}^n A_k(1, \lambda_{n+1})^t \sim c \varrho(A_1) \cdots \varrho(A_n),$$

as $n \rightarrow \infty$ and for some number $0 < \phi_1, \phi_2 < \infty$,

$$(87) \quad \lim_{n \rightarrow \infty} \frac{\sum_{k=1}^{n+1} \mathbf{e}_1 \prod_{i=k}^n A_i(1, \lambda_{n+1})^t}{S_n} = \phi_1 \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{S_{n-1}}{S_n} = \phi_2.$$

Proof. Applying Theorem 2, since $\mathbf{e}_1 \prod_{i=1}^n A_n \mathbf{e}_i^t > 0, i \in \{1, 2\}$ by (15) and (16), we get (84) and (85). Also, we infer from Theorem 2 that

$$\lim_{n \rightarrow \infty} \frac{(\theta_1/b_1, 1) \prod_{k=1}^n A_k(1, \lambda_{n+1})^t}{\varrho(A_1) \cdots \varrho(A_n)}$$

exists. But using (85), we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{(\theta_1/b_1, 1) \prod_{k=1}^n A_k(1, \lambda_{n+1})^t}{\varrho(A_1) \cdots \varrho(A_n)} &= \lim_{n \rightarrow \infty} \frac{\mathbf{e}_2 \prod_{k=1}^n M_k(1, 1)^t}{\varrho(A_1) \cdots \varrho(A_n)} \\ &\geq \lim_{n \rightarrow \infty} \frac{d_1 \mathbf{e}_1 \prod_{k=2}^n M_k \mathbf{e}_2^t}{\varrho(A_1) \cdots \varrho(A_n)} = \lim_{n \rightarrow \infty} \frac{d_1 \mathbf{e}_1 \prod_{k=2}^n A_k \mathbf{e}_2^t}{\varrho(A_1) \cdots \varrho(A_n)} > 0. \end{aligned}$$

So (86) is proved true. Finally, with Theorem 3 in hand, the proof of (87) is similar to the one of [18, Lemma 4] and we will not repeat it here. \square

Now we are ready to finish the proof of Theorem 1. Note that from (73) and the lemmas 3 and 12, we have

$$\begin{aligned} P(\nu = n | Z_0 = \mathbf{e}_2) &\sim c \frac{\varrho(A_1) \cdots \varrho(A_n)}{S_n^2} \frac{S_n}{S_{n-1}} \left(\frac{\theta_1}{b_1} G_{n-1,1} + \frac{\mathbf{e}_2 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t} G_{n-1,2} \right) \\ &\sim c \frac{\varrho(M_1)^{-1} \cdots \varrho(M_n)^{-1}}{\left(\sum_{k=1}^{n+1} \varrho(M_1)^{-1} \cdots \varrho(M_{k-1})^{-1} \right)^2} \left(\frac{\theta_1}{b_1} G_{n-1,1} + \frac{\mathbf{e}_2 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t}{\mathbf{e}_1 \prod_{k=1}^{n-1} A_k \mathbf{e}_1^t} G_{n-1,2} \right). \end{aligned}$$

Then it follows from (83) that

$$\lim_{n \rightarrow \infty} \frac{P(\nu = n | Z_0 = \mathbf{e}_2) \left(\sum_{k=1}^{n+1} \varrho(M_1)^{-1} \cdots \varrho(M_{k-1})^{-1} \right)^2}{\varrho(M_1)^{-1} \cdots \varrho(M_n)^{-1}} = GL \geq 0,$$

where $L > 0$ is a proper constant. But by Lemma 11, $G > 0$ if and only if $\varrho_1 \neq \frac{1}{2} \left(a + b + 1 - \sqrt{(a + b + 1)^2 + 4 \frac{bd - a\theta}{\theta - b}} \right)$. We thus conclude that (9) and (10) hold.

Finally, we turn to prove (8). Note that by (72), (86) and (87), we obtain

$$P(\nu > n | Z_0 = \mathbf{e}_2) \sim c \frac{\varrho(A_1) \cdots \varrho(A_n)}{S_n} = \frac{c}{\sum_{k=1}^{n+1} \varrho(A_1)^{-1} \cdots \varrho(A_{k-1})^{-1}},$$

as $n \rightarrow \infty$. As a result, applying Lemma 3, we get (8). Theorem 1 is proved. \square

Acknowledgements: This project is partially supported by National Natural Science Foundation of China (Grant No. 12071003).

References

- [1] BHATTACHARYA, N. AND PERLMAN, M. (2017). Time inhomogeneous branching processes conditioned on non-extinction. *arXiv: 1703.00337*.
- [2] BIGGINS, J. D., COHN, H. AND NERMAN, O. (1999). Multi-type branching in varying environment. *Stochastic Process. Appl.* **83**, 357-400.
- [3] COHN, H. AND WANG, Q. (2003). Multitype branching limit behavior. *Ann. Appl. Probab.* **13**, 490-500.
- [4] DOLGOPYAT, D., HEBBAR, P., KORALOV, L. AND PERLMAN, M. (2018). Multi-type branching processes with time-dependent branching rates. *J. Appl. Probab.* **55**, 701-727.
- [5] DYAKONOVA, E. E. (1999). Asymptotic behaviour of the probability of non-extinction for a multi-type branching process in a random environment. *Discrete Math. Appl.* **9**, 119-136.
- [6] FUJIMAGARI, T. (1980). On the extinction time distribution of a branching process in varying environments. *Adv. in Appl. Probab.* **12**, 350-366.
- [7] JAGERS, P. (1974). Galton-Watson processes in varying environment. *J. Appl. Probab.* **11**, 174-178.
- [8] JOHNSON, C. R. AND BRU, R. (1990). The spectral radius of a product of nonnegative matrices. *Linear Algebra Appl.* **141**, 227-240.
- [9] JONES, O. D. (1997). On the convergence of multitype branching processes with varying environments. *Ann. Appl. Probab.* **7**, 772-801.
- [10] KERSTING, G. (2020). A unifying approach to branching processes in a varying environment. *J. Appl. Probab.* **57**, 196-220.
- [11] KERSTING, G. AND VATUTIN, V. (2017). *Discrete time branching processes in random environment*. John Wiley & Sons, Inc., USA.
- [12] LINDVALL, T. (1974). Almost sure convergence of branching processes in varying and random environments. *Ann. Probab.* **2**, 344-346.
- [13] LORENTZEN, L. (1995). Computation of limit periodic continued fractions. A survey. *Numer. Algorithms* **10**, 69-111.
- [14] LORENTZEN, L. AND WAADELAND, H. (2008). *Continued fractions. 2nd. Ed., Volume 1: convergence theory*. Atlantis Press, Paris.
- [15] MACPHEE, I. M. AND SCHUH, H. J. (1983). A Galton-Watson branching process in varying environments with essentially constant means and two rates of growth. *Aust. N. Z. J. Stat.* **25**, 329-338.
- [16] SUN, H. Y. AND WANG, H. M. (2020). On a maximum of nearest-neighbor random walk with asymptotically zero drift on lattice of positive half line. *arXiv: 2004.12422*.
- [17] SUN, H. Y. AND WANG, H. M. (2020). Asymptotics of product of nonnegative 2-by-2 matrices with applications to random walks with asymptotically zero drifts. *arXiv: 2004.13440*.
- [18] WANG, H. M. AND YAO, H. (2021). Two-type linear fractional branching processes in varying environments with asymptotically constant mean matrices. *arXiv: 2007.07840*.