

General Bayesian Loss Function Selection and the use of Improper Models

Jack Jewson^{1,2} and David Rossell^{1,2}

¹*Department of Business and Economics, Universitat Pompeu Fabra, Barcelona, Spain*

²*Data Science Center, Barcelona Graduate School of Economics, Spain*

jack.jewson@upf.edu, david.rossell@upf.edu

June 2021

Abstract

Statisticians often face the choice between using probability models or a paradigm defined by minimising a loss function. Both approaches are useful and, if the loss can be re-cast into a proper probability model, there are many tools to decide which model or loss is more appropriate for the observed data, in the sense of explaining the data's nature. However, when the loss leads to an improper model, there are no principled ways to guide this choice. We address this task by combining the Hyvärinen score, which naturally targets infinitesimal relative probabilities, and general Bayesian updating, which provides a unifying framework for inference on losses and models. Specifically we propose the \mathcal{H} -score, a general Bayesian selection criterion and prove that it consistently selects the (possibly improper) model closest to the data-generating truth in Fisher's divergence. We also prove that an associated \mathcal{H} -posterior consistently learns optimal hyper-parameters featuring in loss functions, including a challenging tempering parameter in generalised Bayesian inference. As salient examples, we consider robust regression and non-parametric density estimation where popular loss functions define improper models for the data and hence cannot be dealt with using standard model selection tools. These examples illustrate advantages in robustness-efficiency trade-offs and provide a Bayesian implementation for kernel density estimation, opening a new avenue for Bayesian non-parametrics.

Keywords: Loss functions; Improper models; General Bayes; Hyvärinen score; Robust regression; Kernel density estimation.

1 Introduction

A common task in Statistics is selecting which amongst a set of models is most appropriate for an observed data set y . Tools to address this problem include a variety of penalised likelihood, shrinkage prior, and Bayesian model selection methods. Under suitable conditions, these approaches possess the important property of consistently selecting the model closest to the data-generating truth in Kullback-Leibler divergence (for example, see [Rossell \(2021\)](#) and references therein for a recent discussion). However, many data analysis methods are not defined in terms of probability models but as minimising a given loss function, for example to gain robustness or flexibility. It is then no longer clear how to use the data to guide the choice of the most appropriate loss function, or any associated hyper-parameters. A key observation is that while the likelihood $f_k(y; \theta_k)$ of a model k with parameters θ_k always defines a loss function $\ell_k(y; \theta_k) = -\log f_k(y; \theta_k)$ ([Good, 1952](#)), the converse is not true. The exponential of an arbitrary loss $f_k(y; \theta_k) = \exp\{-\ell_k(y; \theta_k)\}$ may not integrate to a finite constant and therefore, defines an improper model on y . For example, this occurs in robust regression with Tukey’s loss ([Figure 1](#)) and in kernel density estimation. In these scenarios traditional *model* selection tools are not applicable to choose the more appropriate loss. Neither are methods to evaluate predictive performance such as cross-validation, since they require specifying a loss or criterion to evaluate performance in the first place, and do not attain consistent model selection even in simple settings ([Shao, 1997](#)). Methods to tackle intractable but finite normalisation constants, such as approximate Bayesian computation ([Beaumont et al., 2002](#); [Robert, 2016](#)), also do not apply since they require simulating from a proper model.

We propose methodology to evaluate how well each given model and loss captures the unknown data-generating distribution $g(y)$. The main idea is viewing each loss k as defining a (possibly improper) model $\exp\{-\ell_k(y, \theta_k)\}$, and then measuring how well it approximates $g(y)$ via Fisher’s divergence. As we shall see, Fisher’s divergence and its related Hyvärinen score ([Hyvärinen, 2005](#)) do not depend on normalising constants, and hence give a strategy to compare improper models. Note, that one could conceivably define the likelihood in ways other than $\exp\{-\ell_k(y, \theta_k)\}$. Conversely, defining losses as negative log-likelihoods provides the only smooth, local, proper scoring rule ([Bernardo, 1979](#)), and also the only transformation that leads to consistent parameter estimation for certain general class of likelihoods ([Bissiri and Walker, 2012](#)). Further, it seems reasonable that the loss should be additive over independent pieces of information, and that the likelihood of an improper model should factorize under such independence, and for both properties to hold one must take the exponent of the negative

loss. Our framework consistently selects the best model in Fisher’s divergence, and in particular the (proper) data-generating model if it is under consideration. We also show how, after a model is chosen, one can learn important hyper-parameters such as the likelihood tempering in generalised Bayes and PAC-Bayes, robustness-efficiency trade-offs in regression and the level of smoothing in kernel density estimation.

The use of probability models versus algorithms is one of the most fundamental, long-standing debates in Statistics. In an influential piece, [Breiman et al. \(2001\)](#) argued that models are not realistic enough to represent reality in any useful manner, nor flexible enough to predict accurately complex real-world phenomena. Despite advances in flexible and non-parametric models, this view remains in the current era where predictive machine learning proliferates, and shows ample potential to tackle large and/or complex data. However, their limitations notwithstanding, probability models remain a fundamental tool for research. Paraphrasing [Efron \(2020\)](#): “Abandoning mathematical models comes close to abandoning the historic scientific goal of understanding nature.” We agree with the view that there are many situations where models facilitate describing the phenomenon under study. We seek to enrich this paradigm by noting that loss functions define improper models that also lead to natural interpretations, in terms of relative probabilities, and proposing a strategy to learn which loss gives a better description of the process underlying the data.

Our strategy is to view f_k as expressing relative (as opposed to absolute) probabilities, for example $f_k(y_0, \theta_k)/f_k(y_1, \theta_k)$ describes how much more likely is it to observe $y = y_0$ than $y = y_1$. A convenient manner to describe such ratios is by comparing the gradient of $\log f_k(y, \theta_k)$ to the gradient of the log data-generating density $\log g(y)$. This can be achieved by minimising Fisher’s divergence

$$D_F(g||f_k) := \frac{1}{2} \int \|\nabla_y \log g(y) - \nabla_y \log f_k(y; \theta_k)\|_2^2 g(y) dy, \quad (1)$$

where ∇_y is the gradient operator. Minimising Fisher’s divergence is equivalent to minimising the Hyvärinen score ([Hyvärinen, 2005](#)). The Hyvärinen score has been used for models with intractable, but finite normalising constants ([Hyvärinen, 2005](#)) and more recently to conduct Bayesian model selection using improper priors ([Dawid and Musio, 2015](#); [Shao et al., 2019](#)). We consider for the first time its use to select between possibly improper models, and associated hyper-parameters.

The paper proceeds as follows. Section 2 reviews recent developments in Bayesian updating with loss functions, discusses our motivating examples and some failures of standard methodology. Section 3 explains how we interpret the inference provided by an improper model in terms of relative probabilities, and their relation to Fisher’s divergence and the \mathcal{H} -score. It also outlines our methodology: the

definition of an \mathcal{H} -posterior, a \sqrt{n} -consistency result to learn parameters and hyper-parameters, and the definition of the integrated \mathcal{H} -score and \mathcal{H} -Bayes factors as a criterion to choose among possibly improper models. Section 4 gives consistency rates for \mathcal{H} -Bayes factors, including important non-standard cases where optimal hyper-parameters lie at the boundary, as can happen when considering nested models. Section 5 applies our procedure to robust regression. Section 6 produces a Bayesian implementation of kernel density estimation, which cannot be tackled by standard Bayesian methods, since kernel densities define an improper model for the observed data. All proofs and some additional technical results are in the supplementary material. Code to reproduce the examples of Sections 5 and 6 can be found in the repository <https://github.com/jejewson/HyvarinenImproperModels>.

2 Problem formulation

We define the problem and notation and then provide the necessary foundations by reviewing general Bayesian updating, providing two motivating examples, identifying the shortcomings of standard approaches, and finally introducing Fisher’s divergence and the Hyvärinen score.

Denote by $y = (y_1, \dots, y_n)$ an observed continuous outcome, where $y_i \in \mathbb{R}$ are independent draws from an unknown data-generating distribution with density $g(\cdot)$. One is given a set of M probability models and L loss functions which, in general, may or may not include g . As usual each model $k = 1, \dots, M$ is associated to a density $f_k(y; \theta_k, \kappa_k)$, where θ_k are parameters of interest and κ_k are hyperparameters, $(\theta_k, \kappa_k) \in \Theta_k \times \Phi_k \subseteq \mathbb{R}^{d_k}$. Any such density defines a loss $\ell_k(y; \theta_k, \kappa_k) = -\log f_k(y; \theta_k, \kappa_k)$. Similarly, denote by $\ell_k(y; \theta_k, \kappa_k)$ for $k = M + 1, \dots, M + L$ the given loss functions. For $k > M$, we refer to $f_k(y; \theta_k, \kappa_k) = \exp\{-\ell_k(y; \theta_k, \kappa_k)\}$ as the (possibly improper) density associated to ℓ_k . In general such f_k need not integrate to a finite number with respect to y , i.e. f_k may define an improper model on y . Our goal is to choose which among f_1, \dots, f_{L+M} provides a better representation of g , in a sense made precise below.

2.1 General Bayesian updating

In the frequentist paradigm it is natural to infer parameters by minimising loss functions, a classical example being M -estimation (Huber and Ronchetti, 1981). Loss functions are also used in the PAC-Bayes paradigm, where one considers the posterior distribution on the parameters

$$\pi_k(\theta_k | y, \kappa_k) \propto \pi_k(\theta_k | \kappa_k) \exp\{-\ell_k(y; \theta_k, \kappa_k)\} \tag{2}$$

where $\pi_k(\theta_k \mid \kappa_k)$ is a given prior distribution and \propto denotes "proportional to". See [Guedj \(2019\)](#) for a review on PAC-Bayes, and [Grünwald \(2012\)](#) for the safe-Bayes paradigm, which can be seen as a particular case where ℓ_k is a tempered negative log-likelihood. At this stage we consider inference for θ_k for a given hyper-parameter κ_k , we discuss learning κ_k later. As a key result supporting the interpretation of (2) as conditional probabilities akin to Bayes rule, [Bissiri et al. \(2016\)](#) showed that (2) leads to a principled updating of beliefs, and referred to the framework as general Bayesian updating.

These results allow Bayesian inference on parameters θ_k based on loss functions. The properties of the general Bayesian posterior have been well-studied, for example under suitable regularity conditions [Chernozhukov and Hong \(2003\)](#) and [Lyddon et al. \(2019\)](#) showed that it is asymptotically normal. However, the emphasis of prior work is on inference for θ_k . To our knowledge viewing $\exp\{-\ell_k(y; \theta_k, \kappa_k)\}$ as an improper density has not been considered, which is critical for interpretation and posterior predictive inference, nor has the problem of choosing which loss best represents the data.

2.2 Motivating applications

We introduce two problems which, despite being classical, cannot be tackled with standard inference. We first consider robust regression where one contemplates a parametric model and a robust loss, and wishes to assess which represents the data best. To our knowledge there are no solutions for this problem. We next consider learning the bandwidth in kernel density estimation, where the goal is predictive inference on future data. While there are many frequentist solutions, Bayesian methods are hampered by the associated loss defining an unnormalisable model for the observed data.

2.2.1 Robust regression with Tukey's loss

Consider the linear regression of y_i on an p -dimensional vector x_i ,

$$y_i = x_i^T \beta + \epsilon_i, \text{ with } E(\epsilon_i) = 0, V(\epsilon_i) = \sigma^2 \text{ for } i = 1, \dots, n.$$

Consider first a Gaussian model, denoted $k = 1$, so that $f_1(y_i; \theta_1) = \mathcal{N}(y_i; x_i^T \beta, \sigma^2)$, where $\theta_1 = \{\beta, \sigma^2\} \in \mathbb{R}^p \times \mathbb{R}_+$, and there are no hyper-parameters ($\kappa_1 = \emptyset$). The negative log-likelihood gives the least-squares loss $\ell_1(y; \theta_1) = \sum_{i=1}^n \ell_1(y_i; \theta_1)$, where

$$\ell_1(y_i; \theta_1) = -\log f_1(y_i; \beta, \sigma^2) = \frac{1}{2} \log(2\pi\sigma^2) + \frac{(y_i - x_i^T \beta)^2}{2\sigma^2}. \quad (3)$$




Figure 1: Squared-error loss and Tukey’s loss (**left**) and corresponding (improper) densities (**right**). The improper densities for Tukey’s loss are scaled to match the mode of the Gaussian density.

Since the least-squares loss is non-robust to outliers, one may consider alternatives. A classical choice is Tukey’s loss (Beaton and Tukey, 1974), which we denote $k = 2$, given by

$$\ell_2(y_i; \theta_2, \kappa_2) = \begin{cases} \frac{1}{2} \log(2\pi\sigma^2) + \frac{(y_i - x_i^T \beta)^2}{2\sigma^2} - \frac{(y_i - x_i^T \beta)^4}{2\sigma^4 \kappa_2^2} + \frac{(y_i - x_i^T \beta)^6}{6\sigma^2 \kappa_2^4}, & \text{if } |y_i - x_i \beta| \leq \kappa_2 \sigma \\ \frac{1}{2} \log(2\pi\sigma^2) + \frac{\kappa_2^2}{6}, & \text{otherwise} \end{cases} \quad (4)$$

where $\theta_2 = \{\beta, \sigma^2\} \in \mathbb{R}^p \times \mathbb{R}_+$ and $\kappa_2 \in \mathbb{R}_+$ is a cut-off hyper-parameter. Note that (4) is parametrised such that the units of the cut-off parameter κ_2 are standard deviations away from the mean.

The density $f_2(y; \theta_2, \kappa_2) = \exp\{-\sum_{i=1}^n \ell_2(y_i; \theta_2, \kappa_2)\}$ integrates to infinity, defining an improper model. Figure 1 plots Tukey’s loss for several κ_2 and their corresponding densities. (4) is similar to (3) when $|y_i - x_i^T \beta|$ is close to 0, while for large $|y_i - x_i \beta|$ it becomes flat, bounding the influence of outliers. The Gaussian model is recovered from Tukey’s loss when $\kappa_2 = \infty$. As we shall see, such nested comparisons pose methodological challenges that motivated our developments. As a technical remark, in robust statistics σ^2 is typically estimated separately from β , either as part of a two-stage procedure (see e.g. Chang et al., 2018) or using S-estimation (Rousseeuw and Yohai, 1984). Instead, our framework allows to jointly estimate (β, σ^2) .

We note that one can add more losses into our framework, for example those in Black and Rangarajan (1996); Basu et al. (1998) or Wang et al. (2020). Also, our framework is not limited to linear regression. One may replace $x_i^T \theta$ by a non-linear function, for example from a deep learning or Gaussian process regression.

Setting κ_2 in Tukey’s loss is related to the so-called robustness-efficiency trade-off, an at least

60-year old unsettled issue; see [Box \(1953\)](#) and [Tukey \(1960\)](#). While $\kappa_2 = \infty$ gives the most efficient parameter estimates if the data are Gaussian, they are least robust outliers. Decreasing κ_2 increases robustness, but can significantly reduce estimation efficiency if κ_2 is set too small. We address this issue by learning from the data whether or not to estimate parameters in a robust fashion (i.e. selecting between ℓ_1 and ℓ_2) and when appropriate, exactly how robust to be (i.e. estimating κ_2).

Despite the importance of these questions, we are aware of limited work setting κ_2 in a principled, data-driven manner. Rule-of-thumb methods are popular, e.g. setting “ $\kappa_2 = 4.6851$, gives approximately 95% asymptotic efficiency of L2 minimization on the standard normal distribution of residuals” ([Belagiannis et al., 2015](#)), setting $\kappa_2 = 1.547$ to obtain a breakdown of 1/2 ([Rousseeuw and Yohai, 1984](#)), or a balance of breakdown and efficiency (e.g. [Riani et al., 2014](#)). Other methods, rely on estimating quantiles of the data (e.g. [Sinova and Van Aelst \(2016\)](#)) minimising an estimate of parameter mean squared error ([Li et al., 2021](#)) or minimising the maximum change in parameter estimates from perturbing one observation ([Li et al., 2021](#)).

2.2.2 Non-parametric Kernel Density Estimation

Suppose that $y_i \sim g$ independently for $i = 1, \dots, n$ and one wishes to estimate g . The kernel density estimate at a given value x is given by

$$\hat{g}_h(x) = \frac{1}{nh} \sum_{i=1}^n K\left(\frac{x - y_i}{h}\right) \tag{5}$$

where the kernel $K(\cdot)$ is a symmetric, finite variance probability density, and h is the bandwidth parameter. For simplicity we focus on the Gaussian kernel $K(x) = N(x; 0, 1)$.

The bandwidth h is an important parameter controlling the smoothness and accuracy of \hat{g}_h . Popular strategies to set the bandwidth are rule-of-thumb and plug-in methods (e.g. [Silverman, 1986](#)), cross-validation ([Habbema et al., 1974](#); [Robert, 1976](#)) and minimising integrated square error ([Rudemo, 1982](#); [Bowman, 1984](#)).

Unfortunately, standard Bayesian inference cannot be used to learn h from data. The reason is that although (5) defines a proper probability distribution for a future observation x , a Bayesian framework requires a model for the observed data given the parameter. In our notation, the model likelihood $f(y; \theta, \kappa) \propto \prod_{i=1}^n \hat{g}_h(y_i)$ with $\theta = \emptyset$ and $\kappa = \{h\}$ has an infinite normalising constant. To

see this, note that

$$\hat{g}_h(y_i) = \frac{1}{nh} \sum_{j=1}^n K\left(\frac{y_i - y_j}{h}\right) = \frac{1}{nh\sqrt{2\pi}} + \frac{1}{nh\sqrt{2\pi}} \sum_{j \neq i}^n \exp\left\{-\frac{(y_i - y_j)^2}{2h^2}\right\}, \quad (6)$$

where the first term integrates to infinity with respect to y_i . Hence, (5) illustrates a situation where one has an ‘algorithm’ for producing a density estimate for future observations that defines an improper probability model for the observed data.

2.3 The failure of standard technology

As we discussed a main challenge is that standard tools are not, in general, applicable to compare improper models. Another challenge occurs when one wishes to estimate a hyperparameter κ_k of a given improper model k , e.g. Tukey’s cut-off or the kernel bandwidth. For example, the general Bayesian might consider mimicking standard Bayes or marginal likelihood estimation by defining

$$\hat{\kappa}_k := \arg \max_{\kappa_k \in \mathcal{K}} \int \pi_k(\theta_k) \exp\{-\ell_k(y; \theta_k, \kappa_k)\} d\theta_k. \quad (7)$$

Unfortunately, such procedure often produces degenerate estimates. For example, from (4), it is clear that for fixed $\theta_2 = \{\beta, \sigma^2\}$ Tukey’s loss is increasing in κ_2 and therefore (7) selects $\hat{\kappa}_2 = 0$ independently of the data. Similarly, in the kernel density estimation example (7) selects $h = 0$ (Habbema et al., 1974; Robert, 1976).

2.4 Fisher’s divergence and the Hyvärinen score

Hyvärinen (2005) proposed a score matching approach that is computationally convenient when a model’s normalising constant is intractable. Score matching estimates parameters by minimising Fisher’s divergence to the data generating density $g(\cdot)$ in (1), that is

$$\theta_k^* := \arg \min_{\theta_k \in \Theta_k} D_F(g(\cdot) || f_k(\cdot; \theta_k)) = \arg \min_{\theta_k \in \Theta_k} \mathbb{E}_{z \sim g} [H(z; f_k(\cdot; \theta_k))], \quad (8)$$

where the right-hand side is obtained using integration by parts (Hyvärinen, 2005) and

$$H(z; f_k(\cdot; \theta_k)) := 2 \frac{\partial^2}{\partial z^2} \log f_k(z; \theta_k) + \left(\frac{\partial}{\partial z} \log f_k(z; \theta_k) \right)^2, \quad (9)$$

is the Hyvärinen score (\mathcal{H} -score). Given $y \sim g$ one can estimate θ_k^* by minimising

$$\hat{\theta}_k := \arg \min_{\theta_k \in \Theta_k} \frac{1}{n} \sum_{i=1}^n H(y_i; f_k(\cdot; \theta_k)) \quad (10)$$

A critical feature for our purposes is that the \mathcal{H} -score depends only on the first and second derivatives of $\log f_k$ and hence the normalising constant does not play a role, independently of whether it is finite or not. The \mathcal{H} -score enjoys desirable properties. For example, Dawid et al. (2016) proved that $\hat{\theta}_k$ is consistent and asymptotically normal, and Dawid and Musio (2015); Shao et al. (2019) that its prequential application leads to consistent Bayesian model selection under improper priors.

Closest to our work, Matsuda et al. (2019) proposed the score matching information criteria to select between models with intractable, but finite, normalising constants. This criterion estimates Fisher’s divergence by correcting the in-sample Hyvärinen score by an estimate of its asymptotic bias. We emphasise two main distinctions with our work. The first is the extension to improper models. Second, these authors used cross-validations and predictive criteria similar to the AIC, which do not lead to consistent model selection, whereas we focus on structural learning where one seeks guarantees on recovering the loss that best approximates the data-generating g .

3 Inference for improper models

We now present our framework. Section 3.1 interprets an improper model in terms relative probabilities and motivates Fisher’s divergence as a criterion to fit such a model. Section 3.2 proposes using the \mathcal{H} -score to define a general Bayesian posterior to learn hyper-parameters and choose among a collection of models, some or all of which may be improper. Section 3.3 proposes a Laplace approximation to the \mathcal{H} -Bayes factors, which we use both in our theoretical treatment and examples. Finally, Section 3.4 argues for using the \mathcal{H} -score within a two-step procedure, first selecting a model and estimating hyper-parameters using the \mathcal{H} -score, then reverting to standard general Bayes to learn the parameters of interest.

3.1 Inference through relative probabilities

Our goal is to select which model f_k describes the data best, in terms of helping interpret the data-generating g . The main difficulty is that, since f_k may be improper, it is unclear how to define “best”. Our strategy is to view f_k as expressing relative probabilities, in contrast to the usual absolute probabilities. For example $f_k(y_0, \theta_k)/f_k(y_1, \theta_k)$ describes how much more likely one is to observe $y = y_0$ over $y = y_1$. As an illustration, consider Tukey’s loss in (4). For any pair (y_0, y_1) such that

$|y_0 - x^T \beta|, |y_1 - x^T \beta| < \kappa_2 \sigma$ are small, Tukey’s loss is approximately equal to the squared loss, hence

$$\frac{f_2(y_0; \theta_2, \kappa_2)}{f_2(y_1; \theta_2, \kappa_2)} \approx \frac{\mathcal{N}(y_0; x^T \beta, \sigma^2)}{\mathcal{N}(y_1; x^T \beta, \sigma^2)}, \quad (11)$$

In contrast, for any pair such that $|y_0 - x^T \beta|, |y_1 - x^T \beta| > \kappa_2 \sigma$ we have $f_2(y_0, \theta_2, \kappa_2)/f_2(y_1, \theta_2, \kappa_2) = 1$. That is, Tukey’s loss induces relative beliefs that observations near the mode behave like Gaussian variables, while all faraway observations are equally likely. This encodes the notion that one does not know much about the tails beyond their being thick, which is difficult to express using a proper probability distribution.

We argue that Fisher’s divergence is well-suited to evaluate how closely the relative probabilities of any such f_k approximate those from g . Assuming that the gradients of $g(y)$ and $f_k(y)$ are finite for all y , Fisher’s divergence in (1) can be expressed as

$$D_F(g||f_\theta) = \int \left\| \lim_{\epsilon \rightarrow 0} \frac{\log \frac{g(y+\epsilon)}{g(y)} - \log \frac{f_k(y+\epsilon; \theta)}{f_k(y; \theta)}}{\epsilon} \right\|_2^2 g(y) dy. \quad (12)$$

Therefore, minimising Fisher’s divergence (equivalently, the Hyvärinen score) targets a f_k that approximates the relative probabilities of g in an infinitesimal neighbourhood around y , in the quadratic error sense, on the average with respect to $g(y)$. This observation extends the usual motivation for the Hyvärinen score as a replacement of likelihood inference when the normalising constant is intractable to being a justifiable criteria to score improper models.

3.2 The \mathcal{H} -score

We consider a general Bayesian framework where the loss is defined by applying the \mathcal{H} -score to the density $f_k(y; \theta_k, \kappa_k) = \exp\{-\ell_k(y; \theta_k, \kappa_k)\}$, which gives the general posterior

$$\pi^H(\theta_k, \kappa_k | y) \propto \pi_k(\theta_k, \kappa_k) \exp \left\{ - \sum_{i=1}^n H(y_i; f_k(\cdot; \theta_k, \kappa_k)) \right\}. \quad (13)$$

We refer to (13) as the \mathcal{H} -posterior. Note that (13) is different from the general Bayesian posterior directly associated to ℓ_k in (2). An important property of (13) is that it provides a consistent estimator for parameters θ_k and hyper-parameters κ_k . Specifically, Proposition 1 shows that, under regularity conditions, $\tilde{\eta}_k = (\tilde{\theta}_k, \tilde{\kappa}_k)$ maximising (13) recovers the optimal $\eta_k^* = (\theta_k^*, \kappa_k^*)$ according to Fisher’s divergence. This extends the consistency results of Hyvärinen (2005) (Corollary 3), who considered the well-specified case where $g(y) = f_k(y; \theta_k^*, \kappa_k^*)$, which in particular requires f_k to be a proper model.

Proposition 1 also extends Dawid et al. (2016) (Theorem 2), who proved asymptotic normality for \mathcal{H} -score based estimators. Said asymptotic normality does in general not hold in interesting cases where η_k^* lies on the boundary of the parameter space. For example, for Tukey’s loss if the data are truly Gaussian then $\kappa_2^* = \infty$. By Proposition 1, even if normality does not hold, one still attains \sqrt{n} consistency to estimate κ_2^* . Further, Proposition 1 extends previous results by explicitly considering improper models and the learning of their hyperparameters.

Proposition 1 requires some mild regularity conditions A1-A3, stated and discussed in Section A.1.2. Briefly, A1 requires continuous second derivatives of the Hyvärinen score, that it has a unique minimiser, and that its first derivative has finite variance. A2 requires that the Hyvärinen score is dominated by an integrable function, which can be easily seen to hold for Tukey’s loss, for example. Finally, A3 requires that the Hessian of the \mathcal{H} -score is positive and finite around η_j^* .

Proposition 1. Let $y = (y_1, \dots, y_n) \sim g$, $\tilde{\eta}_k = (\tilde{\theta}_k, \tilde{\kappa}_k)$ maximise (13), and $\eta_k^* = (\theta_k^*, \kappa_k^*)$ minimise Fisher’s divergence from $f_k(\theta_k, \kappa_k)$ to g . Assume Conditions A1-A2 in Section A.1.2. Then, as $n \rightarrow \infty$,

$$\|\tilde{\eta}_k - \eta_k^*\|_2 = o_p(1),$$

where $\|\cdot\|_2$ is the L_2 -norm. Further, if Condition A3 also holds, then

$$\|\tilde{\eta}_k - \eta_k^*\|_2 = O_p(1/\sqrt{n}).$$

A consequence of Proposition 1 is that one can use (13) to learn tempering hyper-parameters. Specifically, suppose that one considers a family of losses $w_k \ell_k(\cdot)$, where $w_k > 0$ is a tempering parameter. While w_k does not affect the point estimate of θ_k given by (2), it plays an important role in driving the posterior uncertainty on θ_k . Within our framework, one may define

$$f_k(y; \theta_k, \kappa'_k) = \exp\{-\ell'_k(y; \theta_k, \kappa'_k)\} = \exp\{-w_k \ell_k(y; \theta_k, \kappa_k)\}$$

where $\kappa'_k = (\kappa_k, w_k)$ and $\ell'_k(\cdot) = w_k \ell_k(\cdot)$. By Proposition 1, one can consistently learn the Fisher-divergence optimal κ'_k , and in particular w_k . In contrast, in the general Bayes posterior (2) it is challenging to estimate such w_k . Current strategies to set w_k are optimising an upper-bound on generalisation error in PAC-Bayes (Catoni, 2007), estimating w_k via marginalisation similar to the “Safe Bayesian” fractional likelihood approach of Grünwald (2012), or using information theoretic

arguments to calibrate w_k to match certain limiting sampling distributions (Holmes and Walker, 2017; Lyddon et al., 2019). These strategies essentially view w_k as a tuning parameter. In contrast, in our framework w_k is viewed as a parameter of interest that controls the dispersion of the improper model and affects its interpretation. See Section 6 for an illustration in kernel density estimation.

Recall that our main goal is model comparison. To this end, in analogy to the marginal likelihood in standard Bayesian model selection, we define the integrated \mathcal{H} -score

$$\mathcal{H}_k(y) = \int \pi_k(\theta_k, \kappa_k) \exp \left\{ - \sum_{i=1}^n H(y_i; f_k(\cdot; \theta_k, \kappa_k)) \right\} d\theta_k d\kappa_k. \quad (14)$$

Also, analogously to Bayes factors and posterior model probabilities, we define the \mathcal{H} -Bayes factor as $B_{kl}^{(\mathcal{H})} := \mathcal{H}_k(y) / \mathcal{H}_l(y)$ and

$$\pi(k | y) = \frac{\mathcal{H}_k(y) \pi(k)}{\sum_l \mathcal{H}_l(y) \pi(l)} = \left(1 + \sum_{l \neq k} B_{lk}^{(\mathcal{H})} \frac{\pi(l)}{\pi(k)} \right)^{-1}, \quad (15)$$

where $\pi(l)$ are given model prior probabilities. In our examples we use uniform $\pi(k)$, since we focus on the comparison of a few models, but in high-dimensional settings it may be desirable to set $\pi(k)$ to favour simpler models.

We note that an interesting alternative strategy for model comparison, also based on the \mathcal{H} -score, is to extend the prequential framework of Dawid and Musio (2015) and Shao et al. (2019) designed for improper priors. Therein one could adopt a general Bayesian framework, replacing the likelihood by $f_k(x_i; \theta_k, \kappa_k) = \exp\{\ell_k(x_i; \theta_k, \kappa_k)\}$. Prequential approaches enjoy desirable properties, such as consistency and leading to joint coherent inference on the model and parameter values. Unfortunately, prequential inference is computationally hard, particularly when considering several models. First, inference needs to be updated n times to calculate the one-step-ahead predictive distribution. Second, said updates are not permutation invariant, so one should in principle consider the $n!$ orderings of the data. Thus, while interesting, we leave such line of research for future work.

3.3 Laplace approximation and BIC-type criterion

While there are many strategies for computing integrals such as $\mathcal{H}_k(y)$ in (14) exactly (see [Llorente et al. \(2020\)](#) for a review), it is convenient to have faster options. We consider a Laplace approximation

$$\tilde{\mathcal{H}}_k(y) := (2\pi)^{\frac{d_k}{2}} \pi_k(\tilde{\eta}_k) \exp \left\{ - \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) \right\} |A_k(\tilde{\eta}_k)|^{-\frac{1}{2}}, \quad (16)$$

$$\text{with } \tilde{\eta}_k := \arg \min_{\eta_k} \sum_{i=1}^n H(y_i; f_k(\cdot; \eta_k)) - \log \pi_k(\eta_k), \quad (17)$$

being the mode of the log \mathcal{H} -posterior, $A_k(\eta_k)$ its hessian at η_k , and $\eta_k = (\theta_k, \kappa_k)$.

Computational tractability is important when one considers many models or the integrand is expensive to evaluate, e.g. in our kernel density examples it requires $\mathcal{O}(n^2)$ operations. Further, the availability of a closed-form expression facilitates its theoretical study (Section 4). See [Kass et al. \(1990\)](#) for results on the validity of Laplace approximations. We do not undertake such a study, instead we prove our results directly for the approximation (16) that we actually use for inference.

Although we motivate our methodology from a Bayesian standpoint, we note that $\tilde{\mathcal{H}}_k(y)$ can be viewed as Bayesian-inspired information criteria analogous to the BIC ([Schwarz, 1978](#)). Specifically, in regular settings where A_k is of order $1/n$, one could take the leading terms in $\log \tilde{\mathcal{H}}_k(y)$ to obtain

$$- \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) - \frac{d_k}{2} \log(n) + \log \pi_k(\tilde{\eta}_k)$$

as a model selection criterion. This expression is analogous to the BIC, except for the log prior density term, which converges to a constant for any π_k bounded away from 0 and infinity. The log prior term can play a relevant role however when considering non-local priors where $\pi_k(\eta_k)$ can be equal to 0 for certain η_k , see Section 4.

3.4 Two-step inference and model averaging

As discussed the \mathcal{H} -posterior (13) asymptotically recovers the parameters (θ_k^*, κ_k^*) minimising Fisher's divergence, whereas the general Bayesian posterior (2) recovers the parameters minimising the expected loss $\tilde{\theta}_k^* = \arg \min_{\theta_k \in \Theta_k} \int \ell_k(y; \theta_k, \kappa_k) g(y) dy$, for a given κ_k .

We adopt the pragmatic view that, while one may consider the \mathcal{H} -posterior to choose a model k and learn the associated hyper-parameter κ_k , after said choice one may want to obtain standard inference under the selected model. That is, one desires to learn

$$\arg \min_{\theta_k \in \Theta_k} \int \ell_k(y; \theta_k, \kappa_k^*) g(y) dy. \quad (18)$$

For example, suppose that the \mathcal{H} -score selects a proper probability model (e.g. the Gaussian model). One may then wish that inference collapses to standard Bayesian inference under that model. This is easily achieved with a two-step procedure. First, one uses (15) to select \hat{k} and (13) to estimate $\hat{\kappa}_{\hat{k}}$. Second, given $(\hat{k}, \hat{\kappa}_{\hat{k}})$ one uses the general Bayesian posterior (2) for $\theta_{\hat{k}}$. A further alternative to selecting a single model is to mimic Bayesian model averaging (Hoeting et al., 1999), where the estimates under each model are weighted according to the posterior probabilities in (15).

4 Consistency of \mathcal{H} -score model selection

We now state Theorem 1, our main result that (16) consistently selects the model closest in Fisher's divergence to the data-generating g . When several models attain the same minimum, as may happen when considering nested models, then (16) selects that of smallest dimension. The proof does not require that the Hyvärinen score is asymptotically normal, which holds under the conditions in Theorem 2 of Dawid et al. (2016), but simply the \sqrt{n} -consistency proven in Proposition 1.

Theorem 1 mirrors standard results for Bayes factors (Theorem 1 in Dawid (1999)), the main difference being that it involves Fisher's rather than Kullback-Leibler divergence. As discussed after the theorem, when the optimal hyper-parameter occurs at the boundary the model selection consistency provided by Theorem 1 may not hold, unless one uses a suitable adjustment. Before stating the theorem, we interpret its implications. Part (i) considers a situation where one compares two models l and k such that the former is closer to g in Fisher's divergence. Then $\tilde{B}_{kl}^{(\mathcal{H})}$ converges to 0 at an exponential rate in n . Part (ii) considers that both models attain the same Fisher's divergence, for example for Tukey's loss and the Gaussian model when the data are truly Normal. Then, $\tilde{B}_{kl}^{(\mathcal{H})}$ favours the smaller model at a polynomial rate in n .

Theorem 1. Assume Conditions A1-A4, and let $\eta_k^* = (\theta_k^*, \kappa_k^*)$ be as in Proposition 1.

(i) Suppose that $\mathbb{E}_g[H(y; f_l(\cdot; \eta_l^*))] < \mathbb{E}_g[H(y; f_k(\cdot; \eta_k^*))]$. Then

$$\frac{1}{n} \log \tilde{B}_{kl}^{(\mathcal{H})} = \mathbb{E}_g[H(y; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(y; f_k(\cdot; \eta_k^*))] + o_p(1).$$

(ii) Suppose that $\mathbb{E}_g[H(y; f_l(\cdot; \eta_l^*))] = \mathbb{E}_g[H(y; f_k(\cdot; \eta_k^*))]$. Then

$$\log \tilde{B}_{kl}^{(\mathcal{H})} = \frac{d_l - d_k}{2} \log(n) + O_p(1).$$

The result requires Conditions A1-A4 given and discussed in Section A.1.2. Conditions A1-A3 are mild and also summarised before Proposition 1, whereas A4 imposes a Lipschitz condition on the \mathcal{H} -score and its Hessian. In fact, under extended Conditions A1-A7 it is possible to prove that in the nested case the criterion of (Matsuda et al., 2019) does not guarantee consistent model selection, see Corollary A.1. This result is analogous to predictive criteria such as cross-validation or Akaike’s information criterion not leading to consistent model selection, see Shao (1997).

As an important remark, the slower rate in Part (ii) is due to Condition A1 that the prior density $\pi_l(\eta_l^*) > 0$ at the optimal η_l^* , where l is the larger model. This defines a so-called local prior, in contrast to non-local priors (Johnson and Rossell, 2012; Rossell and Telesca, 2017) which place zero density at the value $\kappa_{l \setminus k}^*$ where the more complicated model l recovers the simpler model k . Since non-local priors violate A1, Corollary A.1 extends Theorem 1 to show that non-local priors attain faster rates in Part (ii), while maintaining the exponential rates in Part (i). We will demonstrate that this improvement can have non-negligible practical implications in Section 5.2.

Another practically-relevant remark is that A3 requires a finite expected Hessian near the optimal (θ_k^*, κ_k^*) , which can be problematic in certain settings. For example, in Tukey’s loss if data are truly Gaussian then $\kappa_2^* = \infty$, which leads to an infinite Hessian. For Tukey’s loss the problem can be avoided by reparameterising $\nu_2 = 1/\kappa_2^2$, for which the Hessian is finite, but more generally such a reparameterisation may not be obvious or not exist. These cases provide a further use for non-local priors. By Corollary A.1, one may set a non-local prior that vanishes sufficiently fast at the boundary (basically, a faster rate than that at which the Hessian diverges) to attain model selection consistency.

5 Robust Regression with Tukey’s loss

We revisit the robust regression in Section 2.2, where one considers a Gaussian model and the improper model defined by Tukey’s loss. Section 5.1 illustrates that when the data contain outliers, the \mathcal{H} -score chooses Tukey’s model and learns its cut-off hyper-parameter in a manner that leads to robust estimation. Section 5.2 shows the opposite situation, where data are truly Gaussian, and the benefits of setting a non-local prior on Tukey’s cut-off hyper-parameter to improve the model selection consistency rate. Finally, Section 5.3 shows two gene expression datasets, one exhibiting Gaussian behavior and the other thicker tails. We compare our results to the SMIC (Matsuda et al., 2019) which, despite not being designed to compare improper models, to our knowledge is the only existing criterion that can

be used for this task.

The \mathcal{H} -scores for the squared loss (ℓ_1) and Tukey's loss (ℓ_2) (see Section A.2.1) are

$$H_1(y; f(\cdot; x, \theta_1)) = \sum_{i=1}^n -\frac{2}{\sigma^2} + \frac{(y_i - x_i^T \beta)^2}{\sigma^4} \quad (19)$$

$$H_2(y; f(\cdot; x, \theta_2, \kappa_2)) = \sum_{|y_i - x_i^T \beta| \leq \kappa_2 \sigma} \left\{ \left(\frac{(y_i - x_i^T \beta)}{\sigma^2} - \frac{2(y_i - x_i^T \beta)^3}{\kappa_2^2 \sigma^4} + \frac{(y_i - x_i^T \beta)^5}{\kappa_2^4 \sigma^6} \right)^2 - 2 \left(\frac{1}{\sigma^2} - \frac{6(y_i - x_i^T \beta)^2}{\kappa_2^2 \sigma^4} + \frac{5(y_i - x_i^T \beta)^4}{\kappa_2^4 \sigma^6} \right) \right\}. \quad (20)$$

Note that minimising H_2 in (20) has a trivial degenerate solution when $(\beta, \sigma, \kappa_2)$ are such that only one observation satisfies $y_i - x_i^T \beta \leq \kappa_2 \sigma$, and $y_i = x_i^T \beta$ for that observation. To avoid such solutions, we define a constraint on the parameters related to the notion of the breakdown point of an estimator, i.e. the number of observations that can be perturbed without causing arbitrary changes to the estimator (Rousseeuw and Yohai, 1984). The constraint is to focus attention on $(\beta, \sigma, \kappa_2)$ such that

$$\frac{1}{\rho(\kappa_2, \kappa_2)} \sum_{i=1}^n \rho \left(\frac{y_i - x_i^T \beta}{\sigma}, \kappa_2 \right) \leq \frac{n}{2} - p, \quad (21)$$

where $\rho \left(\frac{y_i - x_i^T \beta}{\sigma}, \kappa_2 \right) = \ell_2(y_i; x_i, \theta_2, \kappa_2) - \frac{1}{2} \log(2\pi\sigma^2)$, ℓ_2 is as in (4), and $p = \dim(x_i)$. See Section A.3.2 for the derivation and further discussion. Further, to satisfy the differentiability conditions of Theorem 1 and enable the use of standard second order optimisation software, we implemented a differentiable approximation to the indicator function in Tukey's loss (see Section A.3.3).

5.1 The marginal \mathcal{H} -score in κ

Our first example illustrates the properties of the \mathcal{H} -score for calibrating the robustness-efficiency trade-off, in a setting where the data contain outliers. We simulated $n = 500$ observations from the data-generating $g(y) = 0.9\mathcal{N}(y; 0, 1) + 0.1\mathcal{N}(y; 5, 3)$. Related to our breakdown point discussion above, in such a scenario one wishes to estimate the parameters of the larger component (uncontaminated data), in a manner that is robust to the presence of data from the smaller component (outliers). We compare the estimation from the Gaussian model f_1 , which is correctly specified for 90% of the data, with that of the robust improper model arising from Tukey's loss f_2 .

A first question of interest is studying the ability of the \mathcal{H} -score to learn the cutoff hyper-parameter κ_2 . To this end, we measured the evidence for different κ_2 provided by the marginal \mathcal{H} -score

$$\mathcal{H}_2(y; \kappa_2) = \int \pi_2(\theta_2) \exp \left(- \sum_{i=1}^n H(y_i; f_2(\cdot; \theta_2, \kappa_2)) \right) d\theta_2. \quad (22)$$

We set the priors $\sigma^2 \sim \mathcal{IG}(0.1, 0.1)$ and $\mu|\sigma^2 \sim \mathcal{N}(0, 5\sigma^2)$ and consider a grid $\kappa_2 \in \{1, 1.5, 2, \dots, 10\}$. The prior on (μ, σ^2) is further constrained by the breakdown condition (21).




Figure 2: **Left:** The integrated \mathcal{H} -score $\mathcal{H}_2(y; \kappa_2)$ for varying values of κ_2 (black) and box plots of $\hat{\mu}_2(\kappa_2)$ across $B = 1000$ samples from $g(y)$. The grey line indicates the truly zero mean of the uncontaminated component. **Right:** Histogram of $n = 500$ observations from $g(y) = 0.9\mathcal{N}(y; 0, 1) + 0.1\mathcal{N}(y; 5, 3)$ and fitted densities of Tukey's loss with $\hat{\kappa}_2 = 5$ and the Gaussian model (the height of Tukey's loss was set to match the mode of $g(y)$).

The left panel in Figure 2 shows $\mathcal{H}_2(y; \kappa_2)$ (solid line), which is highest for the estimate $\hat{\kappa}_2 = 5$, along with box-plots for $\hat{\mu}(\kappa_2)$, the posterior mean of μ for each fixed κ_2 (obtained from 1,000 repeat samples from $g(y)$). For $\hat{\kappa}_2 = 5$ this estimator is on average close to the uncontaminated data mean, and exhibits small variance. For small κ_2 the average estimate is similar, i.e. is robust to the outliers, but there is larger variance due to a greater proportion of the data being above Tukey's loss threshold. For large κ_2 , $\hat{\mu}(\kappa_2)$ deviates from zero, indicating lack of robustness to outliers, and in the limit as κ_2 grows, $\hat{\mu}(\kappa_2)$ recovers the sample mean.

The right panel of Figure 2 is a histogram of the observed data, the data-generating $g(y)$ and the estimated Gaussian and improper Tukey-based model at $\hat{\kappa}_2 = 5$. The latter provides a better description of the central component of the data, in the sense of capturing the log-gradient of $g(y)$ around the mode, and excludes the outliers.

5.2 Non-local priors and model selection consistency

We demonstrate the selection consistency (Theorem 1) when data are truly Gaussian, and that setting a non-local prior on the cutoff hyper-parameter κ_2 speeds up this selection (Corollary A.1). We

simulated 100 independent data sets of sizes $n = 100, 1,000, 10,000$ and $100,000$ from the data-generating $g(y_i) = \mathcal{N}(y_i; x_i^T \beta, \sigma^2)$, where the first entry in $x_i \in \mathbb{R}^6$ corresponds to the intercept and the remaining entries are Gaussian with unit variances and 0.5 pairwise covariances, $\beta = (0, 0.5, 1, 1.5, 0, 0)$ and $\sigma^2 = 1$. We set priors $\sigma^2 \sim \mathcal{IG}(2.01, 0.5)$ and $\beta | \sigma^2 \sim \mathcal{N}(0, 5\sigma^2 I)$ for both models.

Recall that Tukey’s loss collapses to the Gaussian model for $\kappa_2 = \infty$, and otherwise adds certain flexibility by allowing one to consider an improper model. If this extra flexibility is not needed, following Occam’s razor one wants to choose the Gaussian model. While Theorem 1 guarantees this to occur asymptotically, our experiments show that setting a local prior on κ_2 leads to poor performance, even for $n = 100,000$. Specifically, we compare a (local) half-Gaussian prior $\pi_2^{\text{LP}}(\nu_2) \propto 1_{\nu_2 \geq 0} \mathcal{N}(\nu_2; 0, 1)$ where $\nu_2 = 1/\kappa_2^2$ and a (non-local) inverse-gamma prior $\pi_2^{\text{NLP}}(\nu_2) = \mathcal{IG}(\nu_2; a_0, b_0)$. We set default $(a_0, b_0) = (4.35, 1.56)$ chosen to assign prior probability $P(\kappa_2 \in (1, 3)) = 0.95$, i.e. the range between $1\text{-}\sigma$ and $3\text{-}\sigma$ that cuts off 0.3% of Gaussian data outside the $3\text{-}\sigma$ region, and includes the 68.3% of Gaussian data within $1\text{-}\sigma$. See Section A.3.4 for further discussion of these priors. By Corollary A.2, the log- \mathcal{H} -Bayes factor under the non-local prior should favor the Gaussian model at least at a \sqrt{n} rate, in contrast to the local prior’s $\log(n)$ rate.

Figure 3 compares the SMIC with our integrated \mathcal{H} -score under the local and non-local priors. Score differences are plotted such that negative values indicate correctly selecting the Gaussian model. Firstly, there is no evidence of SMIC being consistent as n grows, even for $n = 100,000$ the wrong model was selected 11% of the time. The \mathcal{H} -score under the local prior has a decreasing median in n , but exhibits heavy tails and even for $n = 100,000$ it also failed to select the Gaussian model 11% of the time. Under the non-local prior, already for $n = 1,000$ the correct decision was made 99% of the time. These experiments illustrate the benefits of non-local priors to penalise parameter values near the boundary ($1/\kappa_2^2 = 0$, in this example). Recall also that, as discussed in Section 4, in general in such situations a local prior need not even attain consistency.

5.3 Real datasets

We considered two gene expression data sets from Rossell and Rubio (2018). In the first, the data are well-approximated by a Gaussian distribution, whereas the second exhibits thicker tails. In both examples we used the \mathcal{H} -score to compare the Gaussian model (ℓ_1) and Tukey’s loss (ℓ_2), using the priors discussed in Section 5.2: $\sigma^2 \sim \mathcal{IG}(2.01, 0.5)$ and $\beta | \sigma^2 \sim \mathcal{N}(0, 5\sigma^2 I)$ for both models, and the non-local inverse-gamma prior $\pi_2^{\text{NLP}}(\nu_2) = \mathcal{IG}(\nu_2; 4.35, 1.56)$.




Figure 3: Selecting between the Gaussian model and Tukey’s loss across $N = 100$ data sets generated from a Gaussian model. SMIC ($\text{SMIC}_1(y) - \text{SMIC}_2(y)$) vs integrated \mathcal{H} -score with local and non-local priors ($\log \tilde{\mathcal{H}}_2(y) - \log \tilde{\mathcal{H}}_1(y)$) with negative values correctly selecting the Gaussian model.

5.3.1 TGF- β data

The dataset from [Calon et al. \(2012\)](#) concerns gene expression data for $n = 262$ colon cancer patients. Previous work ([Rossell and Telesca, 2017](#); [Rossell and Rubio, 2018](#)) focused on selecting genes that have an effect on the expression levels of TGF- β , a gene known to play an important role in colon cancer progression. Instead, we study the relation between TGF- β and the 7 genes (listed in Section [A.4.1](#)) that appear in the ‘TGF- β 1 pathway’ according to the *KEGGREST* package in *R* ([Tenenbaum, 2016](#)), so that $p = 8$ after including the intercept.

The top panels in Figure 4 summarises the results. The integrated \mathcal{H} -score for the Gaussian model was $\tilde{\mathcal{H}}_1(y) = 272.88$ and that for Tukey’s loss $\tilde{\mathcal{H}}_2(y) = 233.90$, providing strong evidence for the Gaussian model. This is in agreement with the SMIC ($\text{SMIC}_1(y) = -283.93$ and $\text{SMIC}_2(y) = -277.55$, where minimisation is desired) and results in [Rossell and Rubio \(2018\)](#), who found evidence for Gaussian over (thicker) Laplace tails. The left panel shows the fitted densities which, in conjunction with the Q - Q Normal plots in Figure [A.2](#), show that the residual distribution is well-approximated by a Gaussian. The right panel shows the squared difference between the \mathcal{H} -posterior mean parameter estimates of each β_j under Tukey’s model minus that under the Gaussian, and the differences between their sampling variances (estimated via bootstrap, dashed gray line). Both models returned very




Figure 4: **Top:** TGF- β data, where the \mathcal{H} -score selected the Gaussian model. **Bottom:** DLD data, where the \mathcal{H} -score selected Tukey’s loss. **Left:** fitted Tukey-based density to the residuals. **Right:** Squared difference between the \mathcal{H} -posterior mean estimates of each β_j under Tukey’s model minus that under the Gaussian (solid black line), and difference between their variances (estimated with $B = 500$ bootstrap re-samples)

similar parameter estimates, but the Gaussian had smaller variance for all parameters. Altogether, these results strongly support that the Gaussian model should be selected over Tukey’s.

5.3.2 DLD dataset

We consider an RNA-sequencing data set from Yuan et al. (2016) measuring gene expression for $n = 192$ patients with different types of cancer. Rossell and Rubio (2018) studied the impact of 57 predictors on the expression of DLD, a gene that can perform several functions such as metabolism regulation. To illustrate our methodology, we selected the 15 variables with the 5 highest loadings in the first 3 principal components, and used the integrated \mathcal{H} -score to choose between the Gaussian and

Tukey’s loss. Section [A.4.1](#) lists the selected variables.

The bottom panels of Figure [4](#) summarise the results. The \mathcal{H} -score strongly supported Tukey’s loss ($\tilde{\mathcal{H}}_1(y) = 155.57$ for the Gaussian model, $\tilde{\mathcal{H}}_2(y) = 783.94$ for Tukey’s), with \mathcal{H} -posterior mean estimate $\hat{\kappa}_2 = 4.12$. Indeed, the bottom left panel indicates that the residuals have thicker-than-Gaussian tails, see also the Q - Q Normal residual plot in Figure [A.2](#). The bottom right panel illustrates two things. Firstly, the estimated coefficients of 6 of the 16 predictors differ quite considerably between the Gaussian model and Tukey’s loss (solid line). Second, the latter often have smaller variance (estimated via bootstrap). Both observations align with the presence of thicker-than-normal tails, which can cause parameter estimation biases and inflated variance. Notably, the \mathcal{H} -score agrees with [Rossell and Rubio \(2018\)](#), who selected Laplace over Gaussian tails, but disagrees with the SMIC ($\text{SMIC}_1(y_{1:192}) = -145.72$ and $\text{SMIC}_2(y_{1:192}) = -141.67$, where minimisation is desired). We speculate that the SMIC results may have been affected by outliers, which could cause instability in the SMIC asymptotic bias estimation.

6 Kernel density estimation

We revisit the kernel density estimate $\hat{g}_h(\cdot)$ from Section [2.2.2](#). The associated loss is

$$\ell(y_i; y, h, w) = -w \log \hat{g}_h(y_i). \tag{23}$$

This loss has two hyper-parameters $\kappa = (h, w)$, where $w > 0$. As discussed in Section [3.2](#), standard Bayesian inference on the bandwidth h is not possible, since the loss does not define a proper probability model for the observed data y . We also included a tempering hyper-parameter w to illustrate how it can provide added flexibility and improve performance. A downside of including w however is that, when $w \neq 1$, the estimated density $\hat{g}_{h,w}(x) = \exp\{-\ell(x; y, h, w)\}$ for a future observation x is no longer a proper model, as it was when $w = 1$ (see Section [2.2.2](#)), and its normalising constant is intractable. However, such one-dimensional integrals are easily approximated by numerical integration. We note that while kernel density estimation is a well-studied area, to the best of our knowledge the incorporation of a tempering parameter w has not been considered, and is hence a further innovation enabled by the Hyvärinen score.

The \mathcal{H} -score associated to [\(23\)](#) and its derivation are in Section [A.2.2](#). Analogously to Tukey’s loss, we must take care to avoid certain cases with degenerate solutions where the in-sample Hyvärinen score attains negative infinity by setting $h = 0$. Such a mode can be avoided by lower-bounding h

away from 0, and for illustration in our examples we set a lower bound of $n^{-1/5}/100$. Such a lower bound is justified as popular bandwidth plug-in methods are $O(n^{-1/5})$ and dividing by a further 100 provides further flexibility. Our method therefore refines previous methods by learning h from the data, but at the same time ensures one does not drastically under-smooth. The estimated \hat{h} 's in the next section are all away from this lower bound.

6.1 Gaussian mixture implementations

We simulated data from four Gaussian mixture models considered by [Marron and Wand \(1992\)](#) (with two further examples provided in [Section A.4.2](#)). All scenarios consider a data-generating

$$g(y) = \sum_{j=1}^J m_j N(y; \mu_j, \sigma_j)$$

with parameters chosen as follows.

- *Bimodal*: $J = 2$ components, $\mu_1 = -1.5$, $\mu_2 = 1.5$, $\sigma_1 = \sigma_2 = 1/2$ and $m_1 = m_2 = 0.5$.
- *Trimodal*: $J = 3$ components, $\mu_1 = -1.2$, $\mu_2 = 0$, $\mu_3 = 1.2$, $\sigma_1 = \sigma_3 = 3/5$, $\sigma_2 = 1/4$, $m_1 = m_3 = 0.45$, and $m_2 = 0.1$.
- *Claw*: $J = 6$ components, $\mu_1 = 0$, $\sigma_1 = 1$ and $m_1 = 0.5$ then for $j = 2, \dots, 6$: $\mu_j = -2 + j/2$, $\sigma_j = 0.1$ and $m_j = 0.1$.
- *Skewed*: $J = 8$ components, for $j = 1, \dots, 8$: $\sigma_j = (2/3)^{j-1}$, $\mu_j = 3(\sigma_j - 1)$ and $m_j = 1/8$.

We compared the \mathcal{H} -score estimated density with fixed $w = 1$ to that when learning w from the data. We further compare against the default kernel estimate in *R*, which sets the bandwidth using Silverman's rule-of-thumb ([Silverman, 1986](#)), and a Bayesian non-parametric density estimate using a Dirichlet Process mixture of Gaussians (DPMM), implemented in *R*'s *dirichletprocess* package ([Ross and Markwick, 2018](#)) using their default parameters. We sampled $n = 1,000$ observations from each simulation setting above and standardised the data to zero mean and unit variance, as recommended for the *dirichletprocess* package. For the \mathcal{H} -score kernel density estimate (KDE), we set the prior $\pi(h) \propto \mathcal{IG}(1, 1)\mathbb{I}(h \geq n^{-1/5}/100)$ for the bandwidth parameter, reparameterised $w = h + 1 + \phi$ to reduce the posterior dependence between (h, w) , and set the prior $\pi(\phi | h) \propto \mathcal{N}(\phi; 0, 1)\mathbb{I}(h + 1 + \phi > 0)$.

Figure 5 provides histograms of the standardised data and the estimated densities, while Table 1 shows Fisher's divergence of the fitted models to the data-generating $g(y)$. The \mathcal{H} -score density




Figure 5: Density estimation for Gaussian mixture data. Histograms of observed data, standardised to 0 mean and unit variance, and estimated density by DPMM, R 's density function with the bandwidth rule of [Silverman \(1986\)](#), the \mathcal{H} -posterior estimate with no tempering ($w = 1$), and with tempering ($w \neq 1$).

estimate generally captured the modes better than the competing methods. This was particularly evident in the *Claw* data where the DPMM and R 's kernel estimate missed the 5 modes, and to a lesser extent in the *Trimodal* and *Skewed* data sets. Lastly, in most examples the \mathcal{H} -score estimate improved by learning the hyper-parameter w , relative to $w = 1$, suggesting that the tempering parameter can add useful flexibility.

7 Discussion

We considered a novel problem, that of selecting between probability models and possibly improper models defined via loss functions. Despite being non-standard, the latter can be naturally interpreted in terms of relative probabilities, and open an avenue to increase the flexibility at the disposal

Table 1: Fisher’s divergence between the density estimates and the data-generating Gaussian mixtures in four simulation scenarios. The best performing method in each is highlighted in bold face

	KDE	DPMM	Hyvärinen KDE ($w = 1$)	Hyvärinen KDE ($w \neq 1$)
<i>Bimodal</i>	1.03	0.09	0.26	0.09
<i>Claw</i>	13.77	15.35	3.39	2.51
<i>Trimodal</i>	0.26	0.47	0.19	0.19
<i>Skewed</i>	21.34	17.21	10.55	9.52

of the data analyst. These relative probabilities motivated our use of Fisher’s divergence and the Hyvärinen score, which led to defining an associated \mathcal{H} -posterior and \mathcal{H} -Bayes factors to estimate hyper-parameters and model selection, respectively. We used Laplace approximations for computational convenience, and proved that one can recover the model that is closest to the data-generating truth in Fisher’s divergence. If a proper model is chosen, one can use a two-step procedure to ensure that inference collapses to standard Bayesian methods. We illustrated how the method provides promising results in robust regression and kernel density estimation, providing a new strategy to address the robustness-efficiency trade-off for the former and enabling Bayesian inference in the latter.

Our asymptotic results mimic those for Bayesian model selection in proper models. There is however an important issue when considering hyper-parameters whose optimal value lies at the boundary, where consistency may fail for standard priors and it may be necessary to use non-local priors. As future work, one could attempt to improve the finite sample performance of the \mathcal{H} -score using weighting procedures to calibrate its sampling distribution, e.g. as proposed by Dawid et al. (2016). Another interesting direction is to extend the robust regression examples to variable selection. Rossell and Rubio (2018) showed the importance of specifying a well-fitting error distribution to avoid non-negligible drops in statistical power, which is particularly important in high-dimensional cases where $p > n$. Considering robust improper models such as that stemming from Tukey’s loss may improve the selection performance here.

We also provided a novel strategy to learn likelihood tempering hyper-parameters from the data, which could be combined with various tempered likelihood approaches (e.g. Grünwald, 2012; Holmes and Walker, 2017; Miller and Dunson, 2018) to estimate this value for standard likelihoods. Yet another interesting future extension is to consider multivariate responses, for example in graphical or factor models where outliers can be hard to detect (e.g. Hall et al., 2005; Filzmoser et al., 2008)

and result in significantly worse inference. In summary, we believe there are many settings where the consideration of improper models can lead to fruitful research avenues.

Acknowledgements

JJ and DR were partially funded by the Ayudas Fundación BBVA a Equipos de Investigación Científica 2017 and Government of Spain’s Plan Nacional PGC2018-101643-B-I00 grants. DR was also partially funded by the Europa Excelencia grant EUR2020-112096 and Ramón y Cajal Fellowship RYC-2015-18544.

References

- Basu, A., Harris, I. R., Hjort, N. L. and Jones, M. (1998), ‘Robust and efficient estimation by minimising a density power divergence’, *Biometrika* **85**(3), 549–559.
- Beaton, A. E. and Tukey, J. W. (1974), ‘The fitting of power series, meaning polynomials, illustrated on band-spectroscopic data’, *Technometrics* **16**(2), 147–185.
- Beaumont, M. A., Zhang, W. and Balding, D. J. (2002), ‘Approximate Bayesian computation in population genetics’, *Genetics* **162**(4), 2025–2035.
- Belagiannis, V., Rupprecht, C., Carneiro, G. and Navab, N. (2015), Robust optimization for deep regression, in ‘Proceedings of the IEEE international conference on computer vision’, pp. 2830–2838.
- Bernardo, J. M. (1979), ‘Expected information as expected utility’, *the Annals of Statistics* pp. 686–690.
- Bissiri, P. G. and Walker, S. (2012), ‘On Bayesian learning via loss functions’, *Journal of statistical planning and inference* **142**(12), 3167–3173.
- Bissiri, P., Holmes, C. and Walker, S. G. (2016), ‘A general framework for updating belief distributions’, *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **78**(5), 1103–1130.
- Black, M. J. and Rangarajan, A. (1996), ‘On the unification of line processes, outlier rejection, and robust statistics with applications in early vision’, *International journal of computer vision* **19**(1), 57–91.

- Bowman, A. W. (1984), ‘An alternative method of cross-validation for the smoothing of density estimates’, *Biometrika* **71**(2), 353–360.
- Box, G. E. (1953), ‘Non-normality and tests on variances’, *Biometrika* **40**(3/4), 318–335.
- Breiman, L. et al. (2001), ‘Statistical modeling: The two cultures (with comments and a rejoinder by the author)’, *Statistical science* **16**(3), 199–231.
- Calon, A., Espinet, E., Palomo-Ponce, S., Tauriello, D. V., Iglesias, M., Céspedes, M. V., Sevillano, M., Nadal, C., Jung, P., Zhang, X. H.-F. et al. (2012), ‘Dependency of colorectal cancer on a TGF- β -driven program in stromal cells for metastasis initiation’, *Cancer cell* **22**(5), 571–584.
- Carpenter, B., Gelman, A., Hoffman, M., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M. A., Guo, J., Li, P. and Riddell, A. (2016), ‘Stan: A probabilistic programming language’, *Journal of Statistical Software* **20**.
- Catoni, O. (2007), Pac-Bayesian supervised classification: the thermodynamics of statistical learning, IMS.
- Chang, L., Roberts, S. and Welsh, A. (2018), ‘Robust lasso regression using Tukey’s biweight criterion’, *Technometrics* **60**(1), 36–47.
- Chernozhukov, V. and Hong, H. (2003), ‘An MCMC approach to classical estimation’, *Journal of Econometrics* **115**(2), 293–346.
- Dawid, A. (1999), The trouble with Bayes factors, Technical report, University College London.
- Dawid, A. P. and Musio, M. (2015), ‘Bayesian model selection based on proper scoring rules’, *Bayesian analysis* **10**(2), 479–499.
- Dawid, A. P., Musio, M. and Ventura, L. (2016), ‘Minimum scoring rule inference’, *Scandinavian Journal of Statistics* **43**(1), 123–138.
- Efron, B. (2020), ‘Prediction, estimation, and attribution’, *Journal of the American Statistical Association* **115**(530), 636–655.
- Ferentios, K. (1982), ‘On tcebycheff’s type inequalities’, *Trabajos de Estadística y de Investigación Operativa* **33**(1), 125.
- Filzmoser, P., Maronna, R. and Werner, M. (2008), ‘Outlier identification in high dimensions’, *Computational Statistics & Data Analysis* **52**(3), 1694–1711.
- Good, I. J. (1952), ‘Rational decisions’, *Journal of the Royal Statistical Society, Series B* **14**, 107–114.

- Grünwald, P. (2012), The safe Bayesian, *in* ‘International Conference on Algorithmic Learning Theory’, Springer, pp. 169–183.
- Guedj, B. (2019), ‘A primer on PAC-Bayesian learning’, *arXiv preprint arXiv:1901.05353*.
- Habbema, J., JDF, H., Van den Broek, K. et al. (1974), ‘A stepwise discriminant analysis program using density estimation.’.
- Hall, P., Marron, J. and Neeman, A. (2005), ‘Geometric representation of high dimension, low sample size data’, *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **67**(3), 427–444.
- Hoeting, J. A., Madigan, D., Raftery, A. E. and Volinsky, C. T. (1999), ‘Bayesian model averaging: a tutorial’, *Statistical science* pp. 382–401.
- Hoffman, M. D. and Gelman, A. (2014), ‘The No-U-Turn sampler: adaptively setting path lengths in Hamiltonian Monte Carlo.’, *Journal of Machine Learning Research* **15**(1), 1593–1623.
- Holmes, C. and Walker, S. (2017), ‘Assigning a value to a power likelihood in a general Bayesian model’, *Biometrika* **104**(2), 497–503.
- Huber, P. J. and Ronchetti, E. (1981), ‘Robust statistics, series in probability and mathematical statistics’.
- Hyvärinen, A. (2005), ‘Estimation of non-normalized statistical models by score matching’, *Journal of Machine Learning Research* **6**(Apr), 695–709.
- Johnson, V. E. and Rossell, D. (2012), ‘Bayesian model selection in high-dimensional settings’, *Journal of the American Statistical Association* **107**(498), 649–660.
- Kass, R. E., Tierney, L. and Kadane, J. B. (1990), ‘The validity of posterior expansions based on Laplace’s method’, *Essays in Honor of George Barnard*, eds. S. Geisser, J. S. Hodges, S. J. Press and A. Zellner, Amsterdam: North-Holland pp. 473–488.
- Li, Q., Chen, H. and Zhu, F. (2021), ‘Robust estimation for Poisson integer-valued GARCH models using a new hybrid loss’, *Journal of Systems Science and Complexity* pp. 1–19.
- Llorente, F., Martino, L., Delgado, D. and Lopez-Santiago, J. (2020), ‘Marginal likelihood computation for model selection and hypothesis testing: an extensive review’, *arXiv* **2005.08334**, 1–91.
- Lyddon, S., Holmes, C. and Walker, S. (2019), ‘General Bayesian updating and the loss-likelihood bootstrap’, *Biometrika* **106**(2), 465–478.

- Marron, J. S. and Wand, M. P. (1992), ‘Exact mean integrated squared error’, *The Annals of Statistics* pp. 712–736.
- Matsuda, T., Uehara, M. and Hyvarinen, A. (2019), ‘Information criteria for non-normalized models’, *arXiv preprint arXiv:1905.05976* .
- Miller, J. W. and Dunson, D. B. (2018), ‘Robust Bayesian inference via coarsening’, *Journal of the American Statistical Association* pp. 1–13.
- Molenberghs, G. and Verbeke, G. (2006), *Models for discrete longitudinal data*, Springer Science & Business Media.
- Riani, M., Cerioli, A. and Torti, F. (2014), ‘On consistency factors and efficiency of robust S-estimators’, *Test* **23**(2), 356–387.
- Robert, C. P. (2016), Approximate Bayesian computation: A survey on recent results, *in* ‘Monte Carlo and Quasi-Monte Carlo Methods’, Springer, pp. 185–205.
- Robert, P. (1976), ‘On the choice of smoothing parameters for Parzen estimators of probability density functions’, *IEEE Transactions on Computers* **25**(11), 1175–1179.
- Ross, G. J. and Markwick, D. (2018), ‘dirichletprocess: An R package for fitting complex Bayesian nonparametric models’.
- Rossell, D. (2021), ‘Concentration of posterior model probabilities and normalized L0 criteria’, *Bayesian Analysis (in press)*, 1–27.
- Rossell, D. and Rubio, F. J. (2018), ‘Tractable Bayesian variable selection: beyond normality’, *Journal of the American Statistical Association* **113**(524), 1742–1758.
- Rossell, D. and Telesca, D. (2017), ‘Nonlocal priors for high-dimensional estimation’, *Journal of the American Statistical Association* **112**(517), 254–265.
- Rousseeuw, P. J. and Leroy, A. M. (2005), *Robust regression and outlier detection*, Vol. 589, John Wiley & sons.
- Rousseeuw, P. and Yohai, V. (1984), Robust regression by means of S-estimators, *in* ‘Robust and nonlinear time series analysis’, Springer, pp. 256–272.
- Rudemo, M. (1982), ‘Empirical choice of histograms and kernel density estimators’, *Scandinavian Journal of Statistics* pp. 65–78.
- Schwarz, G. (1978), ‘Estimating the dimension of a model’, *The Annals of Statistics* **6**(2), 461–464.

- Shao, J. (1997), ‘An asymptotic theory for linear model selection’, *Statistica sinica* pp. 221–242.
- Shao, S., Jacob, P. E., Ding, J. and Tarokh, V. (2019), ‘Bayesian model comparison with the Hyvärinen score: computation and consistency’, *Journal of the American Statistical Association* pp. 1–24.
- Silverman, B. W. (1986), *Density estimation for statistics and data analysis*, Vol. 26, CRC press.
- Sinova, B. and Van Aelst, S. (2016), Tukey’s biweight loss function for fuzzy set-valued M-estimators of location, in ‘International Conference on Soft Methods in Probability and Statistics’, Springer, pp. 447–454.
- Tenenbaum, D. (2016), ‘Kegcrest: Client-side rest access to kegg’, *R package version 1(1)*.
- Tukey, J. W. (1960), ‘A survey of sampling from contaminated distributions’, *Contributions to probability and statistics* pp. 448–485.
- Van der Vaart, A. W. (2000), *Asymptotic statistics*, Vol. 3, Cambridge university press.
- Wang, L., Peng, B., Bradic, J., Li, R. and Wu, Y. (2020), ‘A tuning-free robust and efficient approach to high-dimensional regression’, *Journal of the American Statistical Association* pp. 1–44.
- Yuan, T., Huang, X., Woodcock, M., Du, M., Dittmar, R., Wang, Y., Tsai, S., Kohli, M., Boardman, L., Patel, T. et al. (2016), ‘Plasma extracellular rna profiles in healthy and cancer patients’, *Scientific reports* **6**(1), 1–11.

A Supplementary Material

Section A.1 contains the proofs of the theoretical results associated with Sections 3 and 4 of the main paper. Section A.2 provides the \mathcal{H} -score for the squared, Tukey’s and the kernel density estimation losses. Section A.3 describes implementation details for Tukey’s loss and kernel density examples, including smooth approximations that enable the use of second order optimisation and sampling methods, and how to set a non-local prior on Tukey’s cut-off hyper-parameter κ_2 . Finally, Section A.4 provides additional empirical results that complement those presented in the main paper.

A.1 Proofs

The proofs are structured as follows. First we establish our notation (Section A.1.1) in full, before stating and discussing the conditions for the theorems in Section A.1.2. Sections A.1.3 and A.1.4 provide the proof of Proposition 1, for ease of presentation we split the proof between two separate results. Section A.1.3 provides Lemma A.1 which proves consistency, $o_p(1)$, of the maximum \mathcal{H} -posterior estimates, while Section A.1.4 provides Proposition A.1 extending this to $O_p(1/\sqrt{n})$ convergence. This is then used in Section A.1.5 to prove our main result that the Laplace approximation to the \mathcal{H} -Bayes factor provides consistent selection amongst models and losses to that which minimises Fisher’s divergence to the data generating distribution $g(y)$. Further, Section A.1.6 contains Corollary A.1 which proves that the SMIC of Matsuda et al. (2019) does not provide consistent model/loss selection between nested models, Corollary A.1 in Section A.1.7 extends the \mathcal{H} -score consistency results to situations where non-local priors are placed on the additional parameters of nested models, and lastly Corollary A.2 extends this to the special case of the inverse-gamma non-local prior applied when selecting between a Gaussian model and Tukey’s loss in Section 5.

A.1.1 Notation

Consider models k and l with possibly improper densities $f_j(y; \theta_j, \kappa_j) \propto \exp\{-\ell_j(y; \theta_j, \kappa_j)\}$, for $j \in \{k, l\}$, where κ_j already includes w_j where applicable. To ease notation let $\eta_j = (\theta_j, \kappa_j)$ and call its dimension d_j . Without loss of generality assume l is at least as complicated as k , i.e. that $d_l \geq d_k$. We obtain \mathcal{H} -Bayes Factor rates under the assumption that $y \sim g$. In so doing

$$\eta_j^* := \arg \min_{\eta_j} \mathbb{E}_g [H(y; f_j(\cdot; \eta_j))].$$

are the parameter minimising Fisher’s divergence from model f_j to $g(y)$.

Given sample $y \sim g$, define the in-sample \mathcal{H} -score minimising estimate, $\hat{\eta}_j$, and given and parameter prior $\pi(\eta_j)$ the penalised maximum \mathcal{H} -posterior estimate, $\tilde{\eta}_j$, as

$$\hat{\eta}_j := \arg \min_{\eta_j} \sum_{i=1}^n H(y_i; f_j(\cdot; \eta_j)), \quad \tilde{\eta}_j := \arg \min_{\eta_j} \sum_{i=1}^n H(y_i; f_j(\cdot; \eta_j)) - \log \pi_j(\eta_j),$$

where throughout we suppress the dependence on n to avoid cumbersome notation. The empirical and expected Hessian matrix for model j are $A_j(\eta_j)$ and $A_j^*(\eta_j)$ respectively defined as

$$\begin{aligned} A_j(\eta_j) &:= \nabla_{\eta_j}^2 \left(\sum_{i=1}^n H(y_i; f_j(\cdot; \eta_j)) - \log \pi_j(\eta_j) \right), \\ A_j^*(\eta_j) &:= \nabla_{\eta_j}^2 \mathbb{E}_g [H(y_i; f_j(\cdot; \eta_j))], \end{aligned} \quad (24)$$

The Laplace approximation to the \mathcal{H} -Bayes factor is then given by

$$\tilde{B}_{kl}^{(\mathcal{H})} := \frac{\tilde{\mathcal{H}}_k(y)}{\tilde{\mathcal{H}}_l(y)} = (2\pi)^{\frac{d_k - d_l}{2}} \frac{|A_l(\tilde{\eta}_l)|^{1/2} \pi_k(\tilde{\eta}_k)}{|A_k(\tilde{\eta}_k)|^{1/2} \pi_l(\tilde{\eta}_l)} e^{\sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) - \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k))} \quad (25)$$

For some of the results below it is sometimes easier to consider the estimating equations associated with minimising the Hyvärinen score. Define the empirical and expected vectors of first derivatives, $\Psi_j(\eta_j)$ and $\Psi_j^*(\eta_j)$ respectively, as

$$\begin{aligned} \Psi_j(\eta_j) &:= \nabla_{\eta_j} \sum_{i=1}^n H(y_i; f_j(\cdot; \eta_j)), \quad \phi_j(\eta_j) := \nabla_{\eta_j} \log \pi_j(\eta_j) \\ \tilde{\Psi}_j(\eta_j) &:= \Psi_j(\eta_j) + \phi_j(\eta_j), \quad \Psi_j^*(\eta_j) := \nabla_{\eta_j} \mathbb{E}_g [H(y; f_j(\cdot; \eta_j))]. \end{aligned}$$

A.1.2 Technical Conditions

Next, we list the technical conditions required to prove the consistency of the \mathcal{H} -Bayes factor to the model minimising Fisher's divergence, Theorem 1.

A1. Standard regularity assumptions on the models $j = l, k$ and their priors

- The parameter and hyperparameter spaces $\Theta_j \times \Phi_j$ are compact.
- The priors are continuous in η_j , twice continuously differentiable and $\pi_j(\eta_j) > 0 \forall \eta_j$.
- Model densities $f_j(y; \eta_j) \propto \exp\{-\ell_j(y; \eta_j)\}$, are twice continuously differentiable in y , ensuring that their Hyvärinen score, $H(y; f_j(\cdot; \eta_j))$, is continuous.
- The variance of the first derivatives of the Hyvärinen score $\mathbb{V}_G[\Psi_j(\eta_j)] = \mathbb{V}_G[\nabla_{\eta_j} H(y; f_j(\cdot; \eta_j))] < \infty$.

- The Hyvärinen scores, $H(y; f_j(\cdot; \eta_j))$, are themselves twice continuously differentiable in parameter η_j .
- These exists unique η_j^* and $\tilde{\eta}_j$ for all n such that the first derivatives of the empirical and expected Hyvärinen score, $\tilde{\Psi}_j(\tilde{\eta}_j) = \Psi_j^*(\eta_j^*) = 0$.

A2. For model $j = l, k$, there exists a function $b_j(\cdot; z) : \mathbb{R}^{d_j} \mapsto \mathbb{R}$ such that $|H(z; f_j(\cdot; \eta_j))| < b_j(\eta_j; z)$ for all z where $\int b_j(\eta_j; z) d\eta_j < \infty$.

A3. For model $j = l, k$, the expected Hessian of the Hyvärinen scores (24) is such that

$$0 < |A_j^*(\eta_j^*)| < \infty,$$

and there exist constant, M_A and $\delta_A > 0$

$$0 < M_A \leq \left| \frac{1}{n} A_j(\eta_j) \right|, \text{ for all } \eta_j \text{ such that } \|\eta_j - \eta_j^*\|_2 < \delta,$$

with probability tending to 1 as $n \rightarrow \infty$.

A4. For model $j = l, k$ there exist functions $m_H : \mathbb{R} \mapsto \mathbb{R}_{\geq 0}$ and $m_A(\cdot) : \mathbb{R} \mapsto \mathbb{R}_{\geq 0}$ with $\mathbb{E}[m_H(\cdot)] < \infty$ and $\mathbb{E}[m_A(\cdot)] < \infty$ so that the following Lipschitz conditions hold

$$\begin{aligned} \left| H(z; f_j(\cdot; \eta_j^{(a)})) - H(z; f_j(\cdot; \eta_j^{(b)})) \right| &\leq m_H(z) \left\| \eta_j^{(a)} - \eta_j^{(b)} \right\|_2, \quad \forall \eta_j^{(a)}, \eta_j^{(b)}, \\ \left\| A_j^{(1)}(\eta_j^{(a)}) - A_j^{(1)}(\eta_j^{(b)}) \right\|_2 &\leq m_A(z) \left\| \eta_j^{(a)} - \eta_j^{(b)} \right\|_2, \quad \forall \eta_j^{(a)}, \eta_j^{(b)}, \end{aligned}$$

for all $z \in \mathbb{R}$, where the notation $A_j^{(1)}(\cdot)$ emphasises that only one observation, z , is involved.

These conditions are weak and standard. A1 constitute standard assumptions required to ensure the Hyvärinen score applied to model/loss f_j/ℓ_j , $j = l, k$ is itself continuous and has continuous first and second derivatives (in parameters, e.g. condition S1 [Matsuda et al. \(2019\)](#)), as well as a ensuring the existence and uniqueness of a Hyvärinen score minimising parameters and that its first derivative has finite variance. A2 requires the existence of an function dominating the Hyvärinen score applied to loss $j = k, l$. Note that e.g. for Tukey's loss the improper-model itself is not dominated by an integrable function, but both the first and second log-derivatives vanish above the threshold and therefore the Hyvärinen score applied to Tukey's loss is dominated by a integrable function. A3 ensures the expected Hessian is positive and finite at the limiting parameter η_j^* ([Molenberghs and Verbeke, 2006](#), e.g.), and that within a neighborhood of η_j^* the finite sample average Hessian is also

positive and finite with probability tending to 1. This allows us to deploy a reverse Lipschitz type inequality to bound differences between parameters using differences in the Hyvärinen score’s gradient evaluated at those parameters. A4 then explicitly imposes Lipschitz bounds on the Hyvärinen score applied to ℓ_j and its Hessian in order to use the parameter convergence of Proposition 1 to prove the required functions convergence to prove Theorem 1 (similarly to the assumption of (Van der Vaart, 2000, Theorem 5.21, 5.23)). In fact these bounds are looser than are required as we only need the Lipschitz to hold in the region of η^* . One could further use first order Taylor expansions to provide such Lipschitz bounds which would require the bounding of the first and third derivatives of the Hyvärinen score applied to ℓ_j which are standard assumptions (e.g. condition S3 Matsuda et al. (2019), A6 of Molenberghs and Verbeke (2006)). These however, result in slightly more specific conditions and therefore we considered the more general conditions for brevity.

A.1.3 Lemma A.1 (Hyvärinen score parameter consistency)

Before proving Theorem 1 it is useful to first prove Proposition 1, providing the consistency of the Hyvärinen score minimising parameters estimates to the Fisher’s divergence minimising parameters. For clarity of presentation here we split Proposition 1 into Lemma A.1 proving $o_p(1)$ and Proposition A.1 proving $O_p(1/\sqrt{n})$. Here we first prove Lemma A.1. We note Hyvärinen (2005) (Corollary 3) provided a similar result except this further required that $g = f_j(\cdot; \theta_j^*)$ which we do not require.

Lemma A.1 (Hyvärinen score parameter consistency). Assume Conditions A1-A2. Given a sample $y \sim g$, then as $n \rightarrow \infty$ the maximum \mathcal{H} -posterior parameters of model j have the following asymptotic behaviour

$$\|\tilde{\eta}_j - \eta_j^*\|_2 = o_p(1), \tag{26}$$

where $\|\cdot\|_2$ is the L_2 -norm.

Proof. Firstly, $\|\tilde{\eta}_j - \hat{\eta}_j\|_2 = o_p(1)$ as by A1 $\pi_j(\eta_j) > 0$ for all η_j .

Then, Theorem 5.7 of Van der Vaart (2000) provides general conditions for $\|\hat{\eta}_j - \eta_j^*\| = o_p(1)$ of any M -estimator, of which the the in-sample Hyvärinen score is an example. As is discussed, (Van der Vaart, 2000, p. 46) sufficient conditions for $o_p(1)$ convergence are, compactness of the set $\Theta_j \times \Phi_j$ (A1), uniqueness of η_j^* (A1), continuity of $\eta_j \mapsto H(z; f_j(\cdot; \eta_j))$ for all z (A1), and the dominance of $H(z; f_j(\cdot; \eta_j))$ by an integrable function (A2). As a result, under our conditions $\|\tilde{\eta}_j - \eta_j^*\|_2 = o_p(1)$. \square

A.1.4 Proposition 1 (Hyvärinen score parameter consistency at rate $O_p(1/\sqrt{n})$)

To complete the proof of Proposition 1 we extend Lemma A.1 to establish that under Condition A3 also, the convergence of the Hyvärinen score minimising parameters happens at rate $O_p(1/\sqrt{n})$.

Proposition A.1 (Hyvärinen score parameter consistency at rate $O_p(1/\sqrt{n})$). Assume Conditions A1-A3. Given a sample $y \sim g$, then as $n \rightarrow \infty$ the maximum \mathcal{H} -posterior parameters of model j have the following asymptotic behaviour

$$\|\tilde{\eta}_j - \eta_j^*\|_2 = O_p(1/\sqrt{n}), \quad (27)$$

where $\|\cdot\|_2$ is the L_2 -norm.

Proof. We achieve this by first proving that the gradient of the penalised empirical Hyvärinen loss function $\frac{1}{n}\tilde{\Psi}_j(\eta_j)$ converges to the gradient of the expected Hyvärinen loss function $\Psi_j^*(\eta_j)$ at rate $O_p(1/\sqrt{n})$ (Step 1). We then use this and A3 on the Hessian matrix of the Hyvärinen score to extend the $O_p(1/\sqrt{n})$ convergence of the gradients to that of the parameters (Step 2).

Step 1: Proof that $\left\|\frac{1}{n}\tilde{\Psi}_j(\eta_j) - \Psi_j^*(\eta_j)\right\|_2 = O_p(1/\sqrt{n})$

To do so firstly note that by the triangle inequality of norm $\|\cdot\|_2$

$$\begin{aligned} & P\left(\left\|\sqrt{n}\left(\frac{1}{n}\Psi_j(\eta_j) - \frac{1}{n}\phi_j(\eta_j) - \Psi_j^*(\eta_j)\right)\right\|_2 > M\right) \\ & \leq P\left(\left\|\sqrt{n}\left(\frac{1}{n}\Psi_j(\eta_j) - \Psi_j^*(\eta_j)\right)\right\|_2 + \left\|\sqrt{n}\left(\frac{1}{n}\phi_j(\eta_j)\right)\right\|_2 > M\right) \end{aligned}$$

for any $M > 0$ and under the assumption that $\pi_j(\theta_j) > 0$ (A1). Now for $O_p(1/\sqrt{n})$ convergence we are interested in the behaviour of this probability for sufficiently large $n > N$ and therefore,

$$\begin{aligned} & P\left(\left\|\sqrt{n}\left(\frac{1}{n}\Psi_j(\eta_j) - \Psi_j^*(\eta_j)\right)\right\|_2 + \left\|\sqrt{n}\left(\frac{1}{n}\phi_j(\eta_j)\right)\right\|_2 > M\right) \\ & = P\left(\left\|\sqrt{n}\left(\frac{1}{n}\Psi_j(\eta_j) - \Psi_j^*(\eta_j)\right)\right\|_2 > M - \left\|\frac{1}{\sqrt{n}}\phi_j(\eta_j)\right\|_2\right) \\ & \leq P\left(\left\|\sqrt{n}\left(\frac{1}{n}\Psi_j(\eta_j) - \Psi_j^*(\eta_j)\right)\right\|_2 > M'(\eta_j)\right) \quad \forall n \geq N, \end{aligned}$$

where $M'(\eta_j) := M - \left|\frac{1}{\sqrt{N}}\phi_j(\eta_j)\right|$. Next note that vector valued,

$$\begin{aligned} \mathbb{E}_g \left[\sqrt{n} \left(\frac{1}{n} \Psi_j(\eta_j) - \Psi_j^*(\eta_j) \right) \right] &= 0 \\ \mathbb{V}_G \left[\sqrt{n} \left(\frac{1}{n} \Psi_j(\eta_j) - \Psi_j^*(\eta_j) \right) \right] &= \mathbb{V}_G [\Psi_j(\eta_j)], \end{aligned}$$

where $\sum \mathbb{V}_G[\Psi_j(\eta_j)] < \infty$ (A1). Then we can apply the L_2 -norm vector extension of Chebyshev's inequality (Ferentios, 1982) to vector valued $X_n = \sqrt{n} \left(\frac{1}{n} \Psi_j(\eta_j) - \Psi_j^*(\eta_j) \right)$ which provides that

$$P \left(\left\| \sqrt{n} \left(\frac{1}{n} \Psi_j(\eta_j) - \Psi_j^*(\eta_j) \right) \right\|_2 > M'(\eta_j) \right) < \frac{\sum \mathbb{V}_G[\Psi_j(\eta_j)]}{(M'(\eta_j))^2}$$

which proves that

$$P \left(\left\| \sqrt{n} \left(\frac{1}{n} \tilde{\Psi}_j(\eta_j) - \Psi_j^*(\eta_j) \right) \right\|_2 > M \right) < \frac{\sum \mathbb{V}_G[\Psi_j(\eta_j)]}{(M'(\eta_j))^2}, \quad \forall n > N.$$

Lastly, we note that $\frac{\sum \mathbb{V}_G[\Psi_j(\eta_j)]}{(M'(\eta_j))^2}$ can be made arbitrarily small by making M arbitrarily big and under the assumption that $\pi_j(\theta_j) > 0$ (A1) and therefore $\left\| \frac{1}{n} \tilde{\Psi}_j(\eta_j) - \Psi_j^*(\eta_j) \right\|_2 = O_p(1/\sqrt{n})$ for all η_j .

Step 2: Proof that $\|\tilde{\eta}_j - \eta_j^*\|_2 = O_p(1/\sqrt{n})$

Next, we use A3 on the behaviour of the empirical Hessians around η_j^* to extend the fact that $\left\| \frac{1}{n} \tilde{\Psi}_j(\eta_j) - \Psi_j^*(\eta_j) \right\|_2 = O_p(1/\sqrt{n})$ to prove that $\|\tilde{\eta}_j - \eta_j^*\|_2 = O_p(1/\sqrt{n})$.

Plugging in η_j^* for η_j and using $\tilde{\Psi}_j(\tilde{\eta}_j) = \Psi_j^*(\eta_j^*) = 0$ (by A1) gives,

$$\left\| \frac{1}{n} \tilde{\Psi}_j(\eta_j^*) - \Psi_j^*(\eta_j^*) \right\|_2 = \left\| \frac{1}{n} \tilde{\Psi}_j(\eta_j^*) - \frac{1}{n} \tilde{\Psi}_j(\tilde{\eta}_j) \right\|_2 = O_p(1/\sqrt{n})$$

Next, consider a first order Taylor expansion of $\frac{1}{n} \tilde{\Psi}_j(\cdot)$ about η_j^* evaluated at $\tilde{\eta}_j$. Note that $\frac{1}{n} \tilde{\Psi}_j$ is a vector valued function and as a result we consider a Taylor expansion of each entry of the vector.

$$\begin{aligned} \frac{1}{n} \tilde{\Psi}_j(\tilde{\eta}_j) &= \frac{1}{n} \tilde{\Psi}_j(\eta_j^*) + \nabla_{\eta_j} \frac{1}{n} \tilde{\Psi}_j(\tilde{\eta}_j^\dagger) (\tilde{\eta}_j - \eta_j^*) \\ \Rightarrow \left\| \frac{1}{n} \tilde{\Psi}_j(\tilde{\eta}_j) - \frac{1}{n} \tilde{\Psi}_j(\eta_j^*) \right\|_2 &= \left\| \frac{1}{n} A_j(\tilde{\eta}_j^\dagger) (\tilde{\eta}_j - \eta_j^*) \right\|_2 \end{aligned}$$

for some $\tilde{\eta}_j^\dagger = \alpha \tilde{\eta}_j + (1 - \alpha) \eta_j^*$ for $\alpha \in [0, 1]^{d_j}$. The fact that $\tilde{\Psi}_j(\cdot)$ is already the vector valued first derivative of the Hyvärinen score meant that $\nabla_{\eta_j} \frac{1}{n} \tilde{\Psi}_j(\tilde{\eta}_j^\dagger) = \frac{1}{n} A_j(\tilde{\eta}_j^\dagger)$. Now the right hand side can be bounded as follows

$$\begin{aligned} \left\| \frac{1}{n} A_j(\tilde{\eta}_j^\dagger) (\tilde{\eta}_j - \eta_j^*) \right\|_2 &= \sqrt{(\tilde{\eta}_j - \eta_j^*)^T \left(\frac{1}{n} A_j(\tilde{\eta}_j^\dagger) \right)^T \frac{1}{n} A_j(\tilde{\eta}_j^\dagger) (\tilde{\eta}_j - \eta_j^*)} \\ &\in \left[\lambda_{\min}(\tilde{\eta}_j^\dagger) \|\tilde{\eta}_j - \eta_j^*\|_2, \lambda_{\max}(\tilde{\eta}_j^\dagger) \|\tilde{\eta}_j - \eta_j^*\|_2 \right] \end{aligned}$$

where $\lambda_{\min}(\tilde{\eta}_j^\dagger)$ and $\lambda_{\max}(\tilde{\eta}_j^\dagger)$ are the smallest and largest Eigen-values of $\frac{1}{n} A_j(\tilde{\eta}_j^\dagger)$ respectively. As a result we can bound

$$\left\| \frac{1}{n} \tilde{\Psi}_j(\tilde{\eta}_j) - \frac{1}{n} \tilde{\Psi}_j(\eta_j^*) \right\|_2 \geq \lambda_{\min}(\tilde{\eta}_j^\dagger) \|\tilde{\eta}_j - \eta_j^*\|_2.$$

Further, $\|\tilde{\eta}_j - \eta_j^*\|_2 = o_p(1)$ from Lemma A.1 and by A3ii) there exists $\delta_A > 0$ such that $\lambda_{\min}(\eta_j) > M_A$ with probability tending to 1 for $\|\eta_j - \eta_j^*\|_2 \leq \delta_A$. Therefore, with probability tending to 1

$$\left\| \frac{1}{n} \tilde{\Psi}_j(\tilde{\eta}_j) - \frac{1}{n} \tilde{\Psi}_j(\eta_j^*) \right\|_2 \geq M_A \|\tilde{\eta}_j - \eta_j^*\|_2.$$

M_A is a constant and as a result $\|\tilde{\eta}_j - \eta_j^*\|_2 = O_p(1/\sqrt{n})$, as required. □

Combining Lemma A.1 and Proposition A.1 proves Proposition 1.

A.1.5 Theorem 1 (\mathcal{H} -Bayes Factor Consistency)

Given Proposition 1 and the conditions stated in Section A.1.2 we are now ready to prove the paper's main theorem. First we restate Theorem 1.

Theorem 1 (\mathcal{H} -Bayes Factor Consistency). Assume Conditions A1-A4 given in Section A.1.2. Given a sample $y \sim g$ the Laplace approximation to the \mathcal{H} -Bayes factor (given by $\tilde{B}_{kl}^{(\mathcal{H})} := \tilde{\mathcal{H}}_k(y)/\tilde{\mathcal{H}}_l(y)$ for $\tilde{\mathcal{H}}_j(y)$ defined in (16)) of loss k over loss l has the following asymptotic behaviour as $n \rightarrow \infty$

(i) When $\mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))] < 0$ then

$$\frac{1}{n} \log \tilde{B}_{kl}^{(\mathcal{H})} = \mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))] + o_p(1) \quad (28)$$

That is that when the more complex model l decreases Fisher's divergence relative to model k , the \mathcal{H} -Bayes factor accrues evidence in favour of model l , $\tilde{B}_{kl}^{(\mathcal{H})} \rightarrow 0$, at an exponential rate.

(ii) When $\mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] = \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))]$, with k being the simpler model

$$\log \tilde{B}_{kl}^{(\mathcal{H})} = \frac{d_l - d_k}{2} \log(n) + O_p(1) \quad (29)$$

That is that when the models are equally preferable according to Fisher's divergence then $\tilde{B}_{kl}^{(\mathcal{H})} \rightarrow \infty$ at a polynomial rate in n .

Proof. The proof of Theorem 1 can be broken down into 3 stages. Step 1 deals with how the prior densities of models l and k contribute to $\tilde{B}_{kl}^{(\mathcal{H})}$. Step 2 does the same for the Hessians of the Hyvärinen score applied to models l and k . Lastly, Step 3 deals with the in-sample Hyvärinen score contribution, considering the two cases of the theorem.

Step 1: Priors

From Proposition 1, given that $\|\tilde{\eta}_j - \eta_j^*\|_2 = O_p(1/\sqrt{n})$ and under the assumption that $\pi(\eta_j)$ is continuous in η_j (A1), the Continuous Mapping Theorem provides $\pi_j(\tilde{\eta}_j) \xrightarrow{P} \pi_j(\eta_j^*)$, $j \in \{l, k\}$. Further, assuming that $\pi_l(\eta_l^*) > 0$ (A1), it follows that

$$\frac{\pi_k(\tilde{\eta}_k)}{\pi_l(\tilde{\eta}_l)} \xrightarrow{P} \frac{\pi_k(\eta_k^*)}{\pi_l(\eta_l^*)},$$

a strictly positive finite constant. Note: that for non-local priors either $\pi_j(\eta_j^*)$ may be zero, in that case the limit may be 0 or ∞ . We address this point again later Theorem A.1.

Step 2: Hessians

Now for the Hessians firstly we rewrite $|A_j(\tilde{\eta}_j)| = n^{d_j/2} |\frac{1}{n}A_j(\tilde{\eta}_j)|$. Then we want to prove that $\frac{1}{n}A_j(\tilde{\eta}_j) \xrightarrow{P} A_j^*(\eta_j^*)$. Firstly,

$$\frac{1}{n}A_j(\tilde{\eta}_j) - A_j^*(\eta_j^*) = \frac{1}{n}A_j(\tilde{\eta}_j) - \frac{1}{n}A_j(\eta_j^*) + \frac{1}{n}A_j(\eta_j^*) - A_j^*(\eta_j^*).$$

Then, we use the weak law of large numbers and the Lipschitz condition (A4) to show that

$$\begin{aligned} \left\| \frac{1}{n}A_j(\tilde{\eta}_j) - A_j^*(\eta_j^*) \right\|_2 &= \left\| \frac{1}{n}A_j(\tilde{\eta}_j) - \frac{1}{n}A_j(\eta_j^*) + \frac{1}{n}A_j(\eta_j^*) - A_j^*(\eta_j^*) \right\|_2 \\ &\leq \left\| \frac{1}{n}A_j(\tilde{\eta}_j) - \frac{1}{n}A_j(\eta_j^*) \right\|_2 + \left\| \frac{1}{n}A_j(\eta_j^*) - A_j^*(\eta_j^*) \right\|_2 \quad (\text{tri. in}) \\ &\leq \frac{1}{n} \sum_{i=1}^n \|A_j(\tilde{\eta}_j) - A_j(\eta_j^*)\|_2 + o_p(1) \quad (\text{tri. in \& WLLN.}) \\ &\leq \frac{1}{n} \|\tilde{\eta}_j - \eta_j^*\|_2 \sum_{i=1}^n m_A(y_i) + o_p(1) = o_p(1), \end{aligned}$$

where $\frac{1}{n} \sum_{i=1}^n m_A(y_i) \xrightarrow{P} \mathbb{E}[m_A(y)] < \infty$ by the weak law of large numbers and $\|\tilde{\eta}_j - \eta_j^*\|_2 = O_p(1/\sqrt{n})$ as proved in Proposition 1. As a result we have that

$$\frac{|A_l(\tilde{\eta}_l)|^{1/2}}{|A_k(\tilde{\eta}_k)|^{1/2}} \xrightarrow{P} n^{\frac{(d_l - d_k)}{2}} \frac{|A_l^*(\eta_l^*)|^{1/2}}{|A_k^*(\eta_k^*)|^{1/2}}.$$

Step 3: In-sample Hyvärinen score difference

Combining Steps 1-2, the right hand terms of (25) is asymptotically equivalent to

$$\frac{|A_l^*(\eta_l^*)|^{1/2}}{|A_k^*(\eta_k^*)|^{1/2}} \frac{\pi_k(\eta_k^*)}{\pi_l(\eta_l^*)} \left(\frac{2\pi}{n} \right)^{\frac{d_k - d_l}{2}} e^{\sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) - \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k))}, \quad (30)$$

where the first two terms are constants (A1 and A3). The only step left is to characterize the last term, the in-sample \mathcal{H} -scoreratio. To do this it is necessary to distinguish two cases:

1. Model l decreases Fisher's divergence relative to model k , $\mathbb{E}_g [H(y; f_l(\cdot; \eta_l^*))] < \mathbb{E}_g [H(y; f_k(\cdot; \eta_k^*))]$.
This encompasses the cases where the models are non-nested or nested models where the bigger of the two models is true.
2. Model l does not decrease Fisher's divergence relative to model k , $\mathbb{E}_g [H(y; f_l(\cdot; \eta_l^*))] = \mathbb{E}_g [H(y; f_k(\cdot; \eta_k^*))]$.
This is the case when the models are nested and the simplest model k is sufficient for minimising Fisher's divergence.

Case 1)

First note that the logarithm of (30) is equivalent to

$$\begin{aligned} \frac{1}{n}c - \frac{d_k - d_l}{2n} \log\left(\frac{2\pi}{n}\right) + \sum_{i=1}^n \frac{1}{n} H(y_i; f_l(\cdot; \tilde{\eta}_l)) - \sum_{i=1}^n \frac{1}{n} H(y_i; f_k(\cdot; \tilde{\eta}_k)) = \\ \sum_{i=1}^n \frac{1}{n} H(y_i; f_l(\cdot; \tilde{\eta}_l)) - \sum_{i=1}^n \frac{1}{n} H(y_i; f_k(\cdot; \tilde{\eta}_k)) + O(1/n), \end{aligned}$$

for a constant $c \in \mathbb{R}$ (A1 and A3) given by the first two terms in (30). Now

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) - \frac{1}{n} \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) = \mathbb{E}_g[H(y; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(y; f_k(\cdot; \eta_k^*))] \\ + \frac{1}{n} \sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) - \mathbb{E}_g[H(y; f_l(\cdot; \eta_l^*))] - \left\{ \frac{1}{n} \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) - \mathbb{E}_g[H(y; f_k(\cdot; \eta_k^*))] \right\}, \end{aligned}$$

and for each of the terms on the bottom line ($j \in \{k, l\}$)

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n H(y_i; f_j(\cdot; \tilde{\eta}_j)) - \mathbb{E}_g[H(y; f_j(\cdot; \eta_j^*))] \\ = \frac{1}{n} \sum_{i=1}^n H(y_i; f_j(\cdot; \tilde{\eta}_j)) - \frac{1}{n} \sum_{i=1}^n H(y_i; f_j(\cdot; \eta_j^*)) + \frac{1}{n} \sum_{i=1}^n H(y_i; f_j(\cdot; \eta_j^*)) - \mathbb{E}_g[H(y; f_j(\cdot; \eta_j^*))]. \end{aligned}$$

We then use the weak law of large numbers and the Lipschitz condition (A4) to show that

$$\begin{aligned}
& \left| \frac{1}{n} \sum_{i=1}^n H(y_i; f_j(\cdot; \tilde{\eta}_j)) - \mathbb{E}_g[H(y; f_j(\cdot; \eta_j^*))] \right| \\
&= \left| \frac{1}{n} \sum_{i=1}^n H(y_i; f_j(\cdot; \tilde{\eta}_j)) - \frac{1}{n} \sum_{i=1}^n H(y_i; f_j(\cdot; \eta_j^*)) + \frac{1}{n} \sum_{i=1}^n H(y_i; f_j(\cdot; \eta_j^*)) - \mathbb{E}_g[H(y; f_j(\cdot; \eta_j^*))] \right| \\
&\leq \left| \frac{1}{n} \sum_{i=1}^n H(y_i; f_j(\cdot; \tilde{\eta}_j)) - \frac{1}{n} \sum_{i=1}^n H(y_i; f_j(\cdot; \eta_j^*)) \right| + \left| \frac{1}{n} \sum_{i=1}^n H(y_i; f_j(\cdot; \eta_j^*)) - \mathbb{E}_g[H(y; f_j(\cdot; \eta_j^*))] \right| \quad (\text{tri. in}) \\
&\leq \frac{1}{n} \sum_{i=1}^n |H(y_i; f_j(\cdot; \tilde{\eta}_j)) - H(y_i; f_j(\cdot; \eta_j^*))| + o_p(1) \quad (\text{tri. in \& WLLN}) \\
&\leq \frac{1}{n} \sum_{i=1}^n m_H(y_i) \|\tilde{\eta}_j - \eta_j^*\|_2 + o_p(1) = o_p(1)
\end{aligned}$$

where $\frac{1}{n} \sum_{i=1}^n m_H(y_i) \xrightarrow{P} \mathbb{E}[m_H(y)] < \infty$ by the weak law of large numbers and $\|\tilde{\eta}_j - \eta_j^*\|_2 = O_p(1/\sqrt{n})$ by Proposition 1. As a result

$$\frac{1}{n} \log \frac{\mathcal{H}_k(y)}{\mathcal{H}_l(y)} = \mathbb{E}_g[H(y; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(y; f_k(\cdot; \eta_k^*))] + o_p(1),$$

where $\mathbb{E}_g[H(y; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(y; f_k(\cdot; \eta_k^*))] < 0$ by assumption. That is, the \mathcal{H} -score ratio accrues evidence in favour of model l at an exponential rate.

Case 2)

Now $\mathbb{E}_g[H(y; f_l(\cdot; \eta_l^*))] = \mathbb{E}_g[H(y; f_k(\cdot; \eta_k^*))]$ and we want to prove that the \mathcal{H} -Bayes factor convergence towards the simpler model k . The key is to prove that

$$\sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) - \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) = O_p(1).$$

Note that as model k is nested in model l , there exists $\eta_{l \setminus k}^*$ such that

$$H(y_i; f_k(\cdot; \eta_k)) = H(y_i; f_l(\cdot; (\eta_k, \eta_{l \setminus k}^*))) \quad \forall \eta_k,$$

where $\eta_{l \setminus k}^*$ is the value of the hyperparameters of model l that recover model k . Henceforth, denote $\tilde{\eta}'_k = (\tilde{\eta}_k, \eta_{l \setminus k}^*)$. We can rewrite our objective of interest as

$$\sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) - \log \pi_l(\tilde{\eta}'_k) - \left\{ \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) - \log \pi_l(\tilde{\eta}'_k) \right\},$$

provided $\pi_l(\eta_l) > 0, \forall \eta_l$ (A1). This facilitates the Taylor expansion of $\tilde{H}_l(\tilde{\eta}'_k) := \sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}'_k)) - \log \pi_l(\tilde{\eta}'_k)$ about $\tilde{\eta}_l$, assuming that its first and second derivatives are finite.

$$\tilde{H}_l(\tilde{\eta}'_k) = \tilde{H}_l(\tilde{\eta}_l) + (\tilde{\eta}_k - \tilde{\eta}_l) \tilde{\Psi}_l(\tilde{\eta}_l) + \frac{1}{2}(\tilde{\eta}_k - \tilde{\eta}_l)^T \nabla_{\tilde{\eta}_l}^2 A_l(\tilde{\eta}_l^\dagger) (\tilde{\eta}_k - \tilde{\eta}_l),$$

for some $\tilde{\eta}_l^\dagger = \alpha \tilde{\eta}_k + (1 - \alpha) \tilde{\eta}_l$ with $\alpha \in [0, 1]^{d_l}$.

Now by definition we have that $\tilde{\Psi}_l(\tilde{\eta}_l) = 0$ for all n and therefore plugging this back into our target equation we get that

$$\begin{aligned} & \sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) - \log \pi_l(\tilde{\eta}'_k) - \left\{ \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) - \log \pi_l(\tilde{\eta}_l) \right\} \\ &= \log \pi_l(\tilde{\eta}_l) - \log \pi_l(\tilde{\eta}'_k) + \frac{1}{2}(\tilde{\eta}_k - \tilde{\eta}_l)^T A_l(\tilde{\eta}_l^\dagger) (\tilde{\eta}_k - \tilde{\eta}_l) \\ &= \log \pi_l(\tilde{\eta}_l) - \log \pi_l(\tilde{\eta}'_k) + \frac{1}{2} \sqrt{n} (\tilde{\eta}_k - \tilde{\eta}_l)^T \frac{1}{n} A_l(\tilde{\eta}_l^\dagger) \sqrt{n} (\tilde{\eta}_k - \tilde{\eta}_l). \end{aligned}$$

Now, $\|\tilde{\eta}_j - \eta_j^*\|_2 = O_p(1/\sqrt{n})$, $j = \{k, l\}$ from Proposition 1, where in this nested case $\eta_k^* = \eta_l^* = \eta^*$, which further provides that $\|\tilde{\eta}_l^\dagger - \eta^*\|_2 = O_p(1/\sqrt{n})$ also. From Step 1 we know that $\log \pi_j(\tilde{\eta}_j) \xrightarrow{P} \log \pi_j(\eta_j^*)$, $j = k, l$, and therefore $\log \pi_l(\tilde{\eta}_l) - \log \pi_l(\tilde{\eta}'_k) \xrightarrow{P} 0$. Lastly, Step 2 proved that $\left\| \frac{1}{n} A_j(\tilde{\eta}_j) - A_j^*(\eta_j^*) \right\|_2 = o_p(1)$ and therefore

$$\begin{aligned} \sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) - \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) &= \frac{1}{2} \sqrt{n} (O_p(1/\sqrt{n}) (A_l^*(\eta_l^*) + o_p(1)) \sqrt{n} (O_p(1/\sqrt{n})) \\ &= O_p(1) + O_p(1) o_p(1) = O_p(1) \end{aligned}$$

As a result, that $\log\mathcal{H}$ -score accrues evidence in favour of simpler model k at a $\log(n)$ rate. That is, the \mathcal{H} -score accrues evidence at polynomial rate. □

A.1.6 Corollary A.1 (SMIC inconsistency for nested models)

Here we investigate whether the SMIC model selection criteria provided by Matsuda et al. (2019) can provide the same model/loss selection consistency as Theorem 1, in particular focusing on nested models. We do so under the same conditions as Theorem 1, where the Laplace approximation of the \mathcal{H} -score were shown to be consistent, but with minor further conditions. Firstly, define

$$\begin{aligned} I_j(\eta_j) &:= \sum_{i=1}^n (\nabla_{\eta_j} H(y_i; f_j(\cdot; \eta_j))) (\nabla_{\eta_j} H(y_i; f_j(\cdot; \eta_j)))^T \\ I_j^*(\eta_j) &:= \mathbb{E}_g \left[(\nabla_{\eta_j} H(z; f_j(\cdot; \eta_j))) (\nabla_{\eta_j} H(z; f_j(\cdot; \eta_j)))^T \right]. \end{aligned}$$

We then consider conditions

A5. For model $j = l, k$ there exist functions $m_I(\cdot)$ with $\mathbb{E}[m_I(\cdot)] < \infty$ so that the following Lipschitz condition holds

$$\left\| I_j^{(1)}(\eta_j^{(a)}) - I_j^{(1)}(\eta_j^{(b)}) \right\|_2 \leq m_I(z) \left\| \eta_j^{(a)} - \eta_j^{(b)} \right\|_2, \quad \forall \eta_j^{(a)}, \eta_j^{(b)},$$

where the notation $I_j^{(1)}(\cdot)$ emphasises that only one observation, z , is involved.

A6 Models k and l are such that $\|\tilde{\eta}_j - \eta_j^*\|_2 > o_p(1/\sqrt{n})$.

A5 provides a standard Lipschitz condition for the matrices $I_j(\eta_j)$, while A6 says that the convergence of $\tilde{\eta}_j$ to η_j^* is not faster than $1/\sqrt{n}$. While Proposition 1 proved this convergence of $O_p(1/\sqrt{n})$, it is rare to obtain parametric convergence of $o_p(1/\sqrt{n})$, for simplicity here we simply assume this to be the case. Further, note that while we desired maximisation of our \mathcal{H} -score, the SMIC is to be minimised.

Corollary A.1 (SMIC inconsistency for nested models). Assume Conditions A1-A4, A5 and parametric convergence according to A6. Given a sample $y \sim g$, the difference in the SMIC proposed by Matsuda et al. (2019) of loss k over loss l has the following asymptotic behaviour as $n \rightarrow \infty$.

(i) When $\mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] = \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))]$, with k being the simpler model

$$\text{SMIC}_k - \text{SMIC}_l > o_p(1) \tag{31}$$

That is that when the models are equally preferable according to Fisher's divergence the SMIC will not consistently select the simpler model.

Proof. To prove this we first establish that the trace terms, $\text{tr} \left(\frac{1}{n} I_j(\tilde{\eta}_j) \left(\frac{1}{n} A_j(\tilde{\eta}_j) \right)^{-1} \right)$, tend to constants. Then we show that under the conditions the difference $\left\{ \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) - \sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) \right\}$ is not $o_p(1)$ and therefore for any n there is non-zero probability of $\text{SMIC}_k - \text{SMIC}_l < 0$ when the model k was sufficient according to Fisher's divergence.

From Matsuda et al. (2019) (34)

$$\begin{aligned} \text{SMIC}_k - \text{SMIC}_l &= \left\{ \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) - \sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) \right\} \\ &\quad + \text{tr} \left(\frac{1}{n} I_k(\tilde{\eta}_k) \left(\frac{1}{n} A_k(\tilde{\eta}_k) \right)^{-1} \right) - \text{tr} \left(\frac{1}{n} I_l(\tilde{\eta}_l) \left(\frac{1}{n} A_l(\tilde{\eta}_l) \right)^{-1} \right). \end{aligned}$$

Firstly, **Step 2** of Theorem 1 proved that

$$\left\| \frac{1}{n} A_j(\tilde{\eta}_j) - A_j^*(\eta_j^*) \right\|_2 = o_p(1), \quad j = l, k.$$

Next, we use similar arguments, and A5 to prove that $\left\| \frac{1}{n} I_j(\tilde{\eta}_j) - I_j^*(\eta_j^*) \right\|_2 = o_p(1)$.

$$\begin{aligned} \left\| \frac{1}{n} I_j(\tilde{\eta}_j) - I_j^*(\eta_j^*) \right\|_2 &= \left\| \frac{1}{n} I_j(\tilde{\eta}_j) - \frac{1}{n} I_j(\eta_j^*) + \frac{1}{n} I_j(\eta_j^*) - I_j^*(\eta_j^*) \right\|_2 \\ &\leq \left\| \frac{1}{n} I_j(\tilde{\eta}_j) - \frac{1}{n} I_j(\eta_j^*) \right\|_2 + \left\| \frac{1}{n} I_j(\eta_j^*) - I_j^*(\eta_j^*) \right\|_2 \quad (\text{tri. in}) \\ &\leq \frac{1}{n} \sum_{i=1}^n \left\| I_j(\tilde{\eta}_j) - I_j(\eta_j^*) \right\|_2 + o_p(1) \quad (\text{tri. in \& WLLN.}) \\ &\leq \frac{1}{n} \left\| \tilde{\eta}_j - \eta_j^* \right\|_2 \sum_{i=1}^n m_I(y_i) + o_p(1) = o_p(1), \end{aligned}$$

where $\frac{1}{n} \sum_{i=1}^n m_I(y_i) \xrightarrow{P} \mathbb{E}[m_I(y)] < \infty$ by the weak law of large numbers and $\left\| \tilde{\eta}_j - \eta_j^* \right\|_2 = O_p(1/\sqrt{n})$ as proved in Proposition 1. Therefore, for constant $C^* = \text{tr} \left(\frac{1}{n} I_k^*(\eta_k^*) \left(\frac{1}{n} A_k^*(\eta_k^*) \right)^{-1} \right) - \text{tr} \left(\frac{1}{n} I_l^*(\eta_l^*) \left(\frac{1}{n} A_l^*(\eta_l^*) \right)^{-1} \right)$ we have that

$$\text{SMIC}_k - \text{SMIC}_l = \left\{ \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) - \sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) \right\} + C^* + o_p(1).$$

Lastly, **Step 3, Case 2)** of Theorem 1 proved that in the nested case

$$\left\{ \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) - \sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) \right\} = o_p(1) + \frac{1}{2} \sqrt{n} (\tilde{\eta}_k - \tilde{\eta}_l)^T \left(\frac{1}{n} A_l^*(\eta_l^*) + o_p(1) \right) \sqrt{n} (\tilde{\eta}_k - \tilde{\eta}_l).$$

As a result unless $\left\| \tilde{\eta}_k - \tilde{\eta}_l \right\|_2 = o_p(1/\sqrt{n})$, which when $\mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] = \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))]$ will not happen by assumption (A6), then

$$\left\{ \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) - \sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) \right\} > o_p(1)$$

as a result there exists $\epsilon^* > 0$ and $M^* > 0$ such that for all n

$$P \left(\left\{ \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) - \sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) \right\} > M^* \right) > \epsilon.$$

The consequences of this are as follows, in the nested modelling scenario $\left\{ \sum_{i=1}^n H(y_i; f_k(\cdot; \tilde{\eta}_k)) - \sum_{i=1}^n H(y_i; f_l(\cdot; \tilde{\eta}_l)) \right\} \geq 0$, therefore if $C^* > 0$, then model k will never be preferred to model l . For nested models however it is often the case that C^* is less than 0. In this case it is possible that M^*

is such that $M^* + C^* > 0$ and therefore there is still non-zero probability that model l is chosen over model k even as $n \rightarrow \infty$. Knowing whether this is the case is impossible without knowledge of the data generating density g and therefore a practitioner can not be sure the SMIC will provide consistent model selection. Therefore, we describe it as inconsistent for this task. □

Note Corollary A.1 considered the SMIC evaluated at the penalised Hyvärinen score minimiser $\tilde{\eta}_k$ in order to simplify the extension from Theorem 1, however it is straightforward to see how this also holds for the unpenalised $\hat{\eta}_j$ also. A further consequence of this result is that cases where $A_j^*(\eta_j^*)$ is not a finite matrix, as discussed in Section 4, will also affect the performance of the SMIC. For the SMIC there is no natural method to ameliorate this.

A.1.7 Theorem A.1 (\mathcal{H} -Bayes Factor Consistency under non-local priors)

The conditions of Theorem 1 required that $\pi_j(\theta_j) > 0$ for all θ_j and therefore these conditions will be violated if a non-local prior (Johnson and Rossell, 2012; Rossell and Telesca, 2017) is placed on any model parameters. Here, we extend Theorem 1 to allow for these, particularly focusing on nested models. In order to do so we consider the following extended conditions defining the form of the nested models and the non-local prior.

A7 Models k and l are such that $\eta_k = \{\theta_k\}$ with $\theta_k \in \Theta_k = \Theta$ (i.e. no hyperparameters $\kappa_k = \emptyset$) and $\eta_l = \{\theta_l, \kappa_l\}$ with $\kappa_l \neq \emptyset$ and $\theta_l \in \Theta_l = \Theta$. Define $\kappa_{l \setminus k}$ as the hyperparameters of model l that recovers model k ,

$$H(y_i; f_k(\cdot; \eta_k)) = H(y_i; f_l(\cdot; \{\eta_k, \kappa_{l \setminus k}\})), \quad \text{for all } y_i.$$

A8 Models l and k satisfying A7 have priors

- i) The prior $\pi_k(\eta_k) = \pi_k(\theta_k)$ for model k satisfies A1 and the prior for model l is $\pi_l(\eta_l) = \pi_l(\theta_l, \kappa_l) = \pi_l^{\text{NLP}}(\kappa_l)\pi_l(\theta_l|\kappa_l)$ where $\pi_l(\theta_l|\kappa_l)$ satisfy A1 also.
- ii) The non-local prior on κ_l is $\pi_l^{\text{NLP}}(\kappa_l) = d_l(\|\kappa_l - \kappa_{l \setminus k}\|_2)\pi_l^{\text{LP}}(\kappa_l)$, where $\pi_l^{\text{LP}}(\kappa_l)$ satisfies A1, $d_l(z) > 0$ for all $z \neq 0$, and there is a monotonically decreasing function $c(z) : \mathbb{R} \mapsto \mathbb{R}$ such that $\lim_{z \rightarrow 0} c(z) = \infty$, $c(z/a) \leq c(z)/c(a)$, and $\frac{c(z)}{-\log d_l(z)} = O(1)$ as $z \rightarrow 0$.

A7 defines the form of the nested models we consider. Both models share the same parameter space Θ , but the simpler model k has no hyperparameters. An example of A7 is where k is the Gaussian model and l Tukey's loss and $\kappa_{l \setminus k} = \infty$. A8 specifies that a non-local prior is placed on the additional hyperparameter of model l which penalises their being too close to the value that recovered the simpler model k at a rate as least as fast as $c(\cdot)$. The parameters shared by both models are given local priors.

We define the non-local prior adjusted Laplace approximate Bayes factor as

$$\tilde{B}_{kl}^{\mathcal{H}-\text{NLP}} := \tilde{\mathcal{H}}_k(y) / \tilde{\mathcal{H}}_l^{\text{LP}}(y) \times 1/d_l(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2), \quad (32)$$

where $\tilde{\mathcal{H}}_k(y)$ and $\tilde{\mathcal{H}}_l^{\text{LP}}(y)$ are Laplace approximations to the marginal likelihoods (16) whose respective priors $\pi_k(\eta_k) = \pi_k(\theta_k)$ and $\pi_l(\eta_l) = \pi_l(\theta_l | \kappa_l) \pi_l^{\text{LP}}(\kappa_l)$ satisfy the conditions of A1 according to A8. This allows us to invoke Theorem 1 for the asymptotic behaviour of $\tilde{\mathcal{H}}_k(y) / \tilde{\mathcal{H}}_l^{\text{LP}}(y)$. Some justification for this objective is provided in Remark A.1 at the end of this section.

The following Theorem proves that (32) maintains consistent model selection of the \mathcal{H} -Bayes factor as proved in Theorem 1, and can improve the rate at which the simpler of two nested models is selected when it is sufficient for minimising Fisher's divergence.

Theorem A.1 (\mathcal{H} -Bayes Factor Consistency under Non-Local Priors). Assume Conditions A1-A4, models satisfying A7 and non-local priors satisfying A8.

- (i) When $\mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))] < 0$ then

$$\frac{1}{n} \log \tilde{B}_{kl}^{\mathcal{H}-\text{NLP}} = \mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))] + o_p(1). \quad (33)$$

That is that when the more complex model l decreases Fisher's divergence relative to model k , the non-local \mathcal{H} -Bayes factor accrues evidence in favour of model l , $\tilde{B}_{kl}^{\mathcal{H}-\text{NLP}} \rightarrow 0$, at an exponential rate.

- (ii) When $\mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] = \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))]$, with k being the simpler model, there exists a constant M_d such that with arbitrarily high probability

$$\log \tilde{B}_{kl}^{\mathcal{H}-\text{NLP}} > c(1/\sqrt{n}) \left(\frac{d_l - d_k}{2} \frac{\log(n)}{c(1/\sqrt{n})} + o_p(1) + M_d \right). \quad (34)$$

That is that when the models are equally preferable according to Fisher's divergence then $\tilde{B}_{kl}^{\mathcal{H}-\text{NLP}} \rightarrow \infty$ at a rate depending on the NLP specification via the function $c(\cdot)$.

Proof. To prove this we decompose the log non-local \mathcal{H} -Bayes factor in the local \mathcal{H} -Bayes factor, which was dealt with by Theorem 1, and the non-local penalty term. Then, we consider the two cases of the theorem. When the more complicated model is true, we show that the non-local penalty term tends to a positive constant while when the simpler model is sufficient, we show that the properties of the non-local penalty function control the rate of convergence.

Firstly, from (32)

$$\log \tilde{B}_{kl}^{\mathcal{H}-\text{NLP}} = \log \tilde{B}_{kl}^{(\mathcal{H})} - \log d_l(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2), \quad (35)$$

and by Slutsky's Theorem we can investigate both terms separately. The term $\tilde{B}_{kl}^{(\mathcal{H})}$ was constructed so that it satisfied the conditions of Theorem 1 and therefore we can invoke these results here. Further, $\tilde{\eta}_l = \{\tilde{\theta}_l, \tilde{\kappa}_l\}$ where defined as the Hyvärinen score minimisers penalised by the local prior, therefore satisfying the conditions of Proposition 1 and ensuring that $\|\tilde{\eta}_j - \eta_j^*\|_2 = O_p(1/\sqrt{n})$. Next, we consider the two cases.

Case 1)

From Theorem 1 we have that

$$\frac{1}{n} \log \tilde{B}_{kl}^{(\mathcal{H})} = \mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))] + o_p(1).$$

Further, for $\mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))] < 0$ it must be the case that $\kappa_l^* \neq \kappa_{l \setminus k}$ (by A7). Therefore, $\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 \not\rightarrow 0$ and by the continuous mapping theorem we have that $|d(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) - d_l(\|\kappa_l^* - \kappa_{l \setminus k}\|_2)| = O_p(1/\sqrt{n})$ with $d_l(\|\kappa_l^* - \kappa_{l \setminus k}\|_2) > 0$ as a strictly positive constant (A8ii). As a result $\frac{1}{n} d(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) = O(1/n)$ and

$$\frac{1}{n} \log \tilde{B}_{kl}^{\mathcal{H}-\text{NLP}} = \mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))] + o_p(1).$$

Case 2)

In this case, $\mathbb{E}_g[H(y; f_l(\cdot; \eta_l^*))] = \mathbb{E}_g[H(y; f_k(\cdot; \eta_k^*))] = 0$ which by A7 requires that $\kappa_l^* = \kappa_{l \setminus k}$. Therefore, $\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 = O_p(1/\sqrt{n})$, which by the continuous mapping theorem implies $d_l(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) \rightarrow 0$. Here we establish the asymptotic rate of this convergence.

By (35), in order to establish (34) we require that

$$\forall \epsilon, \quad \exists M_d, N_d > 0 \text{ such that } P(-\log d_l(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) > M_d c(1/\sqrt{n})) > 1 - \epsilon, \quad \forall n > N_d$$

From A8ii), $\frac{c(z)}{-\log d_l(z)} = O(1)$ as $z \rightarrow 0$ which guarantees the existence of M_{δ_κ} and $\delta_\kappa > 0$ such that

$$\frac{-\log d_l(z)}{c(z)} > M_{\delta_\kappa} \text{ for all } z \text{ such that } |z| < \delta_\kappa.$$

As a result, by the law of total probability

$$\begin{aligned}
& P(-\log d_l(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) > M_d c(1/\sqrt{n})) \\
&= P(-\log d_l(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) > M_d c(1/\sqrt{n}) \mid \|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 \leq \delta_\kappa) P(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 \leq \delta_\kappa) \\
&\quad + P(-\log d_l(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) > M_d c(1/\sqrt{n}) \mid \|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 > \delta_\kappa) P(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 > \delta_\kappa) \\
&\geq P(c(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) > \frac{M_d}{M_{\delta_\kappa}} c(1/\sqrt{n}) \mid \|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 \leq \delta_\kappa) P(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 \leq \delta_\kappa).
\end{aligned}$$

Given that $\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 = O_p(1/\sqrt{n})$ from Proposition 1 and δ_κ is a constant, the term $P(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 \leq \delta_\kappa)$ is arbitrarily close to 1 as $n \rightarrow \infty$. Regarding the other terms, by the decreasing monotonicity of $c(z)$ (A8ii) the conditioning events is such that

$$\{\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 \leq \delta_\kappa\} = \{c(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) \geq c(\delta_\kappa)\},$$

and therefore for $\frac{M_d}{M_{\delta_\kappa}} c(1/\sqrt{n}) \geq c(\delta_\kappa)$ we have that

$$P\left(c(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) > \frac{M_d}{M_{\delta_\kappa}} c(1/\sqrt{n}) \mid c(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) > c(\delta_\kappa)\right) \geq P\left(c(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) > \frac{M_d}{M_{\delta_\kappa}} c(1/\sqrt{n})\right).$$

Since $\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 = O_p(1/\sqrt{n})$, by definition for every $\epsilon' > 0$ there exists $M_{\epsilon'} > 0$ and $N_{\epsilon'} > 0$ such that

$$P(\sqrt{n}\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 < M_{\epsilon'}) > 1 - \epsilon', \quad \forall n > N_{\epsilon'}.$$

Further, since $c(z)$ is strictly decreasing and $c(z/a) \leq c(z)/c(a)$ by Assumption A8ii), we have that

$$P(c(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) / c(1/\sqrt{n}) > c(M_{\epsilon'})) \geq P(c(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2 / (1/\sqrt{n})) > c(M_{\epsilon'})) > 1 - \epsilon' \quad \forall n > N_{\epsilon'}.$$

Consider the particular choice $M_d = c(M_{\epsilon'})M_{\delta_\kappa}$ so that $\frac{M_d}{M_{\delta_\kappa}} = c(M_{\epsilon'})$ giving that

$$P(c(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) > \frac{M_d}{M_{\delta_\kappa}} c(1/\sqrt{n})) > 1 - \epsilon' \quad \forall n > N_{\epsilon'}$$

Therefore, taking $(1 - \epsilon') > (1 - \epsilon)$

$$P(-\log d_l(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) > M_d c(1/\sqrt{n})) > (1 - \epsilon') P(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) > (1 - \epsilon)$$

for large enough n . Further, invoking Theorem 1 we have that with arbitrarily high probability there

exists constant M_d such that for sufficiently large n

$$\begin{aligned}
\log \tilde{B}_{kl}^{\mathcal{H}-NLP} &= \log \tilde{B}_{kl}^{(\mathcal{H})} - \log d_l(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2) \\
&> \log \tilde{B}_{kl}^{(\mathcal{H})} + M_d \times c(1/\sqrt{n}) \\
&= \frac{d_l - d_k}{2} \log(n) + O_p(1) + M_d \times c(1/\sqrt{n}) \\
&= c(1/\sqrt{n}) \left(\frac{d_l - d_k}{2} \frac{\log(n)}{c(1/\sqrt{n})} + \frac{O_p(1)}{c(1/\sqrt{n})} + M_d \right).
\end{aligned}$$

□

We follow Theorem A.1 with the special case corollary considering the inverse-gamma non-local prior applied to $\nu_2 = \frac{1}{\kappa_2^2}$ where κ_2 was the cut-off parameter of Tukey's loss (Section 5.2). First, we generalise that prior set-up from Section 5.2 as the following condition.

A9 Under the model setup of A7

- i) Let $\kappa_l = \{\nu\} \in \mathbb{R}$ be a univariate hyperparameter with $\kappa_{l \setminus k} = \nu_0$ and $\nu \geq \nu_0$.
- ii) $\pi_l^{\text{LP}}(\nu) = 2\mathbf{1}_{\nu \geq \nu_0} \mathcal{N}(\nu - \nu_0; 0, s_0^2)$, a half-Gaussian distribution with scale s_0 .
- iii) $\pi_l^{\text{NLP}}(\nu) = \mathcal{IG}(\nu - \nu_0; a_0, b_0)$, an inverse-gamma distribution with shape a_0 and scale b_0 .

We note that the corresponding local prior to an inverse-gamma non-local prior would normally be a gamma distribution. However, this does not have finite second derivative as $\nu - \nu_0 \rightarrow 0$ and would therefore violate the conditions of Theorem 1.

Corollary A.2 (\mathcal{H} -Bayes Factor Consistency under an Inverse-Gamma Non-Local Prior). Assume Conditions A1-A4, models satisfying A7 and non-local priors satisfying A8(i) and A9.

- (i) When $\mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))] < 0$ then

$$\frac{1}{n} \log \tilde{B}_{kl}^{\mathcal{H}-NLP} = \mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] - \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))] + o_p(1). \quad (36)$$

That is that when the more complex model l decreases Fisher's divergence relative to model k , the non-local \mathcal{H} -Bayes factor accrues evidence in favour of model l , $\tilde{B}_{kl}^{\mathcal{H}-NLP} \rightarrow 0$, at an exponential rate.

- (ii) When $\mathbb{E}_g[H(z; f_l(\cdot; \eta_l^*))] = \mathbb{E}_g[H(z; f_k(\cdot; \eta_k^*))]$, with k being the simpler model, there exists a constant M_d such that with arbitrarily high probability

$$\log \tilde{B}_{kl}^{\mathcal{H}-NLP} > \sqrt{n}(o_p(1) + M_d). \quad (37)$$

That is that when the models are equally preferable according to Fisher's divergence then $\log \tilde{B}_{kl}^{\mathcal{H}-\text{NLP}} \rightarrow \infty$ at a rate at least as fast as \sqrt{n} as $n \rightarrow \infty$.

Proof. To prove this we derive the non-local penalty function associated with the prior setup of A9 and calculate its bounding function $c(\cdot)$ associated with A8ii). This allows us to invoke Theorem A.1 to provide the desired asymptotic behaviour.

Firstly, from A9, $\nu \geq \nu_0$ and therefore $d_l(|\nu - \nu_0|) = d_l(\nu - \nu_0)$. Then, the non-local penalty function $d_l(\nu - \nu_0) = \frac{\pi_l^{\text{NLP}}(\nu)}{\pi_l^{\text{LP}}(\nu)}$ given by the prior specification of A9 is

$$\begin{aligned} d_l(\nu - \nu_0) &= \frac{\pi_l^{\text{NLP}}(\nu)}{\pi_l^{\text{LP}}(\nu)} \\ &\propto \frac{b_0^{a_0} \sqrt{2\pi} \sqrt{s_0^2}}{2\Gamma(a_0)} \left(\frac{1}{\nu - \nu_0} \right)^{a_0+1} \exp \left(\frac{(\nu - \nu_0)^2}{2s_0^2} - \frac{b_0}{\nu - \nu_0} \right), \end{aligned}$$

and therefore

$$-\log d_l(\nu - \nu_0) = -\log \frac{b_0^{a_0} \sqrt{2\pi} \sqrt{s_0^2}}{2\Gamma(a_0)} + (a_0 + 1) \log(\nu - \nu_0) - \frac{(\nu - \nu_0)^2}{2s_0^2} + \frac{b_0}{(\nu - \nu_0)}.$$

Therefore, there exists $\delta_\nu > 0$ and $M_{\delta_\nu} > 0$ such that for all ν s.t. $|\nu - \nu_0| < \delta_\nu$

$$-\log d_l(\nu - \nu_0) > M_{\delta_\nu} \frac{1}{\nu - \nu_0},$$

and as a result, $c(z) = \frac{1}{z}$. Now we consider the two cases.

Case 1)

Follows directly from the proof of Theorem A.1, noting that $d_l(z) > 0$ for all $0 < z < \infty$.

Case 2)

Using Theorem A.1 with $c(z) = \frac{1}{z}$ provides that with arbitrarily high probability there exists M_d such that for sufficiently large n

$$\begin{aligned} \log \tilde{B}_{kl}^{\mathcal{H}-\text{NLP}} &> c(1/\sqrt{n}) \left(\frac{d_l - d_k}{2} \frac{\log(n)}{c(1/\sqrt{n})} + \frac{O_p(1)}{c(1/\sqrt{n})} + M_d \right) \\ &= \sqrt{n} \left(\frac{d_l - d_k}{2} \frac{\log(n)}{\sqrt{n}} + \frac{O_p(1)}{\sqrt{n}} + M_d \right) \\ &= \sqrt{n}(o_p(1) + M_d), \end{aligned}$$

as required. □

Remark A.1 (Justification for (32)). Lastly, we provide some justification for the objective function (32) to conduct model selection using non-local priors. Using the arguments of Rossell and Telesca (2017), the integrated \mathcal{H} -score (14) associated with the non-local prior specification A8 can be rewritten as

$$\begin{aligned}\mathcal{H}_l^{\text{NLP}}(y) &= \int \pi_l^{\text{NLP}}(\kappa_l) \pi_l(\theta_l | \kappa_l) \exp \left\{ - \sum_{i=1}^n H(y_i; f_l(\cdot; \theta_l, \kappa_l)) \right\} d\theta_l d\kappa_l \\ &= \mathcal{H}_l^{\text{LP}}(y) \int d_l(\|\kappa_l - \kappa_{l \setminus k}\|_2) \frac{\pi_l^{\text{LP}}(\kappa_l) \pi_l(\theta_l | \kappa_l) \exp \left\{ - \sum_{i=1}^n H(y_i; f_l(\cdot; \theta_l, \kappa_l)) \right\}}{\mathcal{H}_l^{\text{LP}}(y)} d\theta_l d\kappa_l \\ &= \mathcal{H}_l^{\text{LP}}(y) \int d_l(\|\kappa_l - \kappa_{l \setminus k}\|_2) \pi_l^{\text{LP}}(\theta_l, \kappa_l | y) d\theta_l d\kappa_l\end{aligned}$$

where

$$\begin{aligned}\pi_l^{\text{LP}}(\theta_l, \kappa_l | y) &\propto \pi_l^{\text{LP}}(\kappa_l) \pi_l(\theta_l | \kappa_l) \exp \left\{ - \sum_{i=1}^n H(y_i; f_l(\cdot; \theta_l, \kappa_l)) \right\}, \\ \mathcal{H}_l^{\text{LP}}(y) &= \int \pi_l^{\text{LP}}(\kappa_l) \pi_l(\theta_l | \kappa_l) \exp \left\{ - \sum_{i=1}^n H(y_i; f_l(\cdot; \theta_l, \kappa_l)) \right\} d\theta_l d\kappa_l,\end{aligned}$$

As a result we can write the \mathcal{H} -Bayes factor of nested models satisfying A7 and non-local priors according to A8 as

$$B_{kl}^{\mathcal{H}\text{-NLP}} = \frac{\mathcal{H}_k(y)}{\mathcal{H}_l^{\text{NLP}}(y)} = \frac{\mathcal{H}_k(y)}{\mathcal{H}_l^{\text{LP}}(y)} \frac{1}{\int d_l(\|\kappa_l - \kappa_{l \setminus k}\|_2) \pi_l^{\text{LP}}(\theta_l, \kappa_l | y) d\theta_l d\kappa_l}.$$

Now, the terms involved in the construction of $\mathcal{H}_k(y)$ and $\mathcal{H}_l^{\text{LP}}(y)$ satisfy the conditions of Theorem 1 (by A8i), and therefore the behaviour of the Laplace approximations (16) to the \mathcal{H} -Bayes factor, $\tilde{\mathcal{H}}_k(y)/\tilde{\mathcal{H}}_l^{\text{LP}}(y)$, is detailed in Theorem 1. We further consider a somewhat looser, but convenient, approximation of $\int d_l(\|\kappa_l - \kappa_{l \setminus k}\|_2) \pi_l^{\text{LP}}(\theta_l, \kappa_l | y) d\theta_l d\kappa_l$ as $d_l(\|\tilde{\kappa}_l - \kappa_{l \setminus k}\|_2)$ where $\tilde{\kappa}_l \in \tilde{\eta}_l$ are the parameters maximising the \mathcal{H} -posterior $\pi_l^{\text{LP}}(\theta_l, \kappa_l | y)$. As a result we consider (32) to approximate the \mathcal{H} -Bayes factor under non-local priors

We highlight that calculating the approximation $\tilde{\mathcal{H}}_l^{\text{NLP}}(y)$ uses the maximum a posteriori estimates and evaluates the observed Hessian matrices according to the \mathcal{H} -posterior under the local prior, before evaluating the prior density as it appears in the Laplace approximation (16) at the non-local prior, as shown in (32). This is convenient as it allows for a direct extension of Theorem 1.

A.2 Derivation of Hyvärinen scores

Section A.2.1 provides the Hyvärinen score for the Gaussian log-likelihood and for Tukey's loss.

A.2.1 The Hyvärinen score of the Gaussian Model and Tukey's Loss

Here we provide the derivations of the Hyvärinen score applied to the Gaussian model (19) and Tukey's loss (20). Firstly, for the Gaussian model $f_1(y_i; x_i, \theta_1, \kappa_1) = \mathcal{N}(y_i; x_i^T \beta, \sigma^2)$ we have that

$$\begin{aligned}\log f_1(y_i; x_i, \theta_1, \kappa_1) &= -\frac{1}{2} \log(2\pi) - \frac{1}{2} \log \sigma^2 - \frac{(y_i - x_i^T \beta)^2}{2\sigma^2} \\ \frac{\partial}{\partial y_i} \log f_1(y_i; x_i, \theta_1, \kappa_1) &= -\frac{(y_i - x_i^T \beta)}{\sigma^2} \\ \frac{\partial^2}{\partial y_i^2} \log f_1(y_i; x_i, \theta_1, \kappa_1) &= -\frac{1}{\sigma^2},\end{aligned}$$

which results in Hyvärinen score

$$H_1(y_i; f(\cdot; x_i, \theta_1)) = -\frac{2}{\sigma^2} + \frac{(y_i - x_i^T \beta)^2}{\sigma^4}.$$

For Tukey's loss, $\log f_2(y_i; x_i, \theta_2, \kappa_2) = -\ell_2(y_i; x_i, \theta_2, \kappa_2)$ given in (4) and

$$\begin{aligned}\frac{\partial}{\partial y_i} \ell_2(y_i; x_i, \theta_2, \kappa_2) &= \begin{cases} \frac{(y_i - x_i^T \beta)}{\sigma^2} - \frac{2(y_i - x_i^T \beta)^3}{\kappa_2^2 \sigma^4} + \frac{(y_i - x_i^T \beta)^5}{\kappa_2^4 \sigma^6} & \text{if } |y_i - x_i^T \beta| \leq \kappa_2 \sigma \\ 0 & \text{otherwise} \end{cases}, \\ \frac{\partial^2}{\partial y_i^2} \ell_2(y_i; x_i, \theta_2, \kappa_2) &= \begin{cases} \frac{1}{\sigma^2} - \frac{6(y_i - x_i^T \beta)^2}{\kappa_2^2 \sigma^4} + \frac{5(y_i - x_i^T \beta)^4}{\kappa_2^4 \sigma^6} & \text{if } |y_i - x_i^T \beta| \leq \kappa_2 \sigma \\ 0 & \text{otherwise} \end{cases},\end{aligned}$$

which results in Hyvärinen score

$$\begin{aligned}H_2(y_i; f(\cdot; x_i, \theta_2, \kappa_2)) &= \mathbb{I}(|y_i - x_i^T \beta| \leq \kappa_2 \sigma) \left\{ \left(\frac{(y_i - x_i^T \beta)}{\sigma^2} - \frac{2(y_i - x_i^T \beta)^3}{\kappa_2^2 \sigma^4} + \frac{(y_i - x_i^T \beta)^5}{\kappa_2^4 \sigma^6} \right)^2 \right. \\ &\quad \left. - 2 \left(\frac{1}{\sigma^2} - \frac{6(y_i - x_i^T \beta)^2}{\kappa_2^2 \sigma^4} + \frac{5(y_i - x_i^T \beta)^4}{\kappa_2^4 \sigma^6} \right) \right\}.\end{aligned}$$

A.2.2 The Hyvärinen score of kernel density estimation

We derive the Hyvärinen score for the kernel density estimation examples implemented in Section 6.

Combining the kernel density estimate improper density (6) with the power w (23) results in in-sample

$$\hat{g}_{w,h}(y_i) = \left(\frac{1}{n\sqrt{2\pi h}} \sum_{j=1}^n \exp\left(-\frac{(y_i - y_j)^2}{2h}\right) \right)^w,$$

or equivalently defining the kernel density loss function to be the log-density

$$\begin{aligned}\log \hat{g}_{w,h}(y_i) &= w \log \left(\frac{1}{n\sqrt{2\pi h}} \sum_{j=1}^n \exp \left(-\frac{(y_i - y_j)^2}{2h} \right) \right) \\ &= w \log \left(\frac{1}{n\sqrt{2\pi h}} + \frac{1}{n\sqrt{2\pi h}} \sum_{j \neq i}^n \exp \left(-\frac{(y_i - y_j)^2}{2h} \right) \right).\end{aligned}$$

The Hyvärinen score is then composed of the first and second derivatives of the log-density, given by

$$\begin{aligned}\frac{\partial}{\partial y_i} \log \hat{g}_{w,h}(y_i) &= w \frac{-\sum_{j \neq i}^n \frac{(y_i - y_j)}{n\sqrt{2\pi\sigma^3}} \exp \left(-\frac{(y_i - y_j)^2}{2h} \right)}{\left(\frac{1}{n\sqrt{2\pi h}} + \frac{1}{n\sqrt{2\pi h}} \sum_{j \neq i}^n \exp \left(-\frac{(y_i - y_j)^2}{2h} \right) \right)} \\ \frac{\partial^2}{\partial y_i^2} \log \hat{g}_{w,h}(y_i) &= w \frac{\sum_{j \neq i}^n \frac{(y_i - y_j)^2}{n\sqrt{2\pi h^{5/2}}} \exp \left(-\frac{(y_i - y_j)^2}{2h} \right) - \sum_{j \neq i}^n \frac{1}{n\sqrt{2\pi h^{3/2}}} \exp \left(-\frac{(y_i - y_j)^2}{2h} \right)}{\left(\frac{1}{n\sqrt{2\pi h}} + \frac{1}{n\sqrt{2\pi h}} \sum_{j \neq i}^n \exp \left(-\frac{(y_i - y_j)^2}{2h} \right) \right)} \\ &\quad - w \frac{\left(\sum_{j \neq i}^n \frac{(y_i - y_j)}{n\sqrt{2\pi\sigma^3}} \exp \left(-\frac{(y_i - y_j)^2}{2h} \right) \right)^2}{\left(\frac{1}{n\sqrt{2\pi h}} + \frac{1}{n\sqrt{2\pi h}} \sum_{j \neq i}^n \exp \left(-\frac{(y_i - y_j)^2}{2h} \right) \right)^2}.\end{aligned}$$

As a result, the Hyvärinen score of the kernel density estimate is

$$\begin{aligned}H(y_i, \log \hat{g}_{w,h}(\cdot)) &= 2w \frac{\sum_{j \neq i}^n \frac{(y_i - y_j)^2}{\sigma^4} \exp \left(-\frac{(y_i - y_j)^2}{2h} \right) - \sum_{j \neq i}^n \frac{1}{\sigma^2} \exp \left(-\frac{(y_i - y_j)^2}{2h} \right)}{\left(1 + \sum_{j \neq i}^n \exp \left(-\frac{(y_i - y_j)^2}{2h} \right) \right)} \\ &\quad - (2w - w^2) \frac{\left(\sum_{j \neq i}^n \frac{(y_i - y_j)}{\sigma^2} \exp \left(-\frac{(y_i - y_j)^2}{2h} \right) \right)^2}{\left(1 + \sum_{j \neq i}^n \exp \left(-\frac{(y_i - y_j)^2}{2h} \right) \right)^2}.\end{aligned}\tag{38}$$

A.3 Implementation Details

This section provides a thorough description of the experiments conducted in Section 5 and 6 of the main paper. Section A.3.1 outlines our implementation of the \mathcal{H} -score in Stan. Section A.3.2 discusses a restriction placed on Tukey's loss parameters to prevent degenerate parameter estimates based on including a single observation in the fit. Given that Tukey's loss is not twice continuously differentiable, Section A.3.3 discusses a differentiable approximation to Tukey's loss that allows using second order sampling and optimisation methods, as well as satisfying the conditions of Theorem 1. Finally, Section A.3.4 discusses how to set a non-local prior on Tukey's loss cut-off hyper-parameter κ_2 .

Code to reproduce the examples of Sections 5 and 6 can be found in the repository <https://github.com/jejewson/HyvarinenImproperModels>.

A.3.1 Computation

To calculate the Laplace approximations of the \mathcal{H} -score we used the second order LBFGS optimisation routine implemented in the probabilistic programming language Stan (Carpenter et al., 2016). Further, *stan*'s implementation of the No U-Turn Sampler (Hoffman and Gelman, 2014) can further be used to sample from the General Bayesian posteriors. The code associated with all of our experiments is available as part of the supplementary material.

A.3.2 The breakdown of Tukey's loss

As discussed in Section 5, we define a restriction of the parameter space of Tukey's loss $(\beta, \sigma, \kappa_2)$ that prevents degenerate solutions where κ_2 is set to exclude all observations but one. Intuitively, the idea is to avoid κ_2 being too small. More specifically, we motivate the restriction via a related notion of the breakdown point. The breakdown of an estimator is the number of observations that can be arbitrarily perturbed without causing arbitrary changes to the estimator. The squared loss can be seen to have breakdown 0, while clearly for Tukey's-loss any observation outside the threshold can be perturbed arbitrarily without changing the estimator. There exists abundant literature arguing that the breakdown can be no greater than $1/2$, which intuitively means that an estimator should depend on more than half of the data. See Rousseeuw and Yohai (1984) and Rousseeuw and Leroy (2005) for discussion and thorough investigation of Tukey's loss breakdown in the context of S-estimation. The authors showed that for Tukey's breakdown to be less than $1/2$ the condition in (21) must hold. Therefore, we restrict the parameter space to $(\beta, \sigma, \kappa_2)$ satisfying (21). We note that (21) applies to S-estimation rather than to our \mathcal{H} -scores however, we found that (21) provides a simple rule that showed good performance in our examples, hence we recommend it as a default.

A.3.3 A differentiable approximation to the indicator

In order to satisfy the conditions of Theorem 1 as well as facilitate second order optimisation (or sampling) methods, it is necessary to have a log-density that is twice continuously differentiable. While Tukey's loss itself is defined to ensure it that this is the case, the Hyvärinen score applied to Tukey's loss, which is a function of its first and second derivatives is not. This is because the indicator

$\mathbb{I}(|y_i - x_i^T \beta| \leq \kappa_2 \sigma)$ is not differentiable at $|y_i - x_i^T \beta| = \kappa_2 \sigma$. We ensured the required conditions by the approximating

$$|x| \approx \sqrt{x^2 + \frac{1}{k_1}} \text{ for large } k_1$$

$$\mathbb{I}(x \geq 0) \approx \sigma_{k_2}(x) = \frac{1}{1 + \exp(-k_2 x)} \text{ for large } k_2$$

in (20). For our results we set $k_1 = k_2 = 100$.

A.3.4 Non-Local Prior Specification for Tukey’s cut-off hyper-parameter

To place a non-local prior on κ_2 , it is convenient to consider the reparametrisation $\nu_2 = \frac{1}{\kappa_2}$ (as discussed in Section 4, this is also needed to ensure that the Hessian of Tukey’s loss is finite as $\kappa_2 \rightarrow \infty$ ($\nu_2 \rightarrow 0$)). Given that the Gaussian model is recovered at $\nu_2 = 0$, any prior setting positive density at $\nu_2 = 0$ satisfies the definition of being a local prior. In our examples we considered a half-Gaussian prior $\pi_2^{\text{LP}}(\nu_2) \propto \mathcal{N}(\nu_2; 0, s_0^2)I(\nu_2 \geq 0)$ where $I(\cdot)$ is the indicator function and $s_0 = 1$. A possible alternative that also assigns non-zero density at $\nu_2 = 0$ is to consider a gamma prior with shape parameter < 1 , but the associated log-prior does not have a finite Hessian as $\nu_2 \rightarrow 0$ and violates Condition A3 of Theorem 1.

As a non-local prior we considered an inverse-gamma distribution $\pi_2^{\text{NLP}}(\nu_2) = \mathcal{IG}(\nu_2; a_0, b_0)$, which has zero density at $\kappa_2 = 0$. The values of the prior hyper-parameters (a_0, b_0) can have a considerable affect on the finite sample performance of the \mathcal{H} -score model selection, and therefore setting these in a principled manner is important. Fortunately, κ_2 has a natural interpretation that allows restricting attention to a range of values that would be reasonable in most applications. Recall from Section 2.2.1 that the role of κ_2 is to exclude observations that are more than κ_2 ‘standard deviations’ σ away from the mean μ . First, if the data are Gaussian then there is > 0.99 probability of an observation being within 3σ ’s of the mean. We thus set (a_0, b_0) such that there is high prior probability that $\kappa_2 \leq 3$. Second, given that the Gaussian has a substantial amount of central data within 1σ from the mean, we set high prior probability that $\kappa_2 > 1$, to ensure that this central component is captured. Given these considerations, we set default parameter values $(a_0, b_0) = (4.35, 1.56)$, which ensures a prior probability $P(\kappa_2 \in (1, 3)) = 0.95$. Figure A.1 plots these local and non-local priors for ν_2 and the corresponding priors under the original parametrisation, κ_2 .




Figure A.1: Left: Density of the local half-standard Gaussian and non-local $\mathcal{IG}(\nu_2; 4.35, 1.56)$ priors for ν_2 . Right: The corresponding prior density for $\kappa_2 = 1/\sqrt{\nu_2}$. The non-local prior sets $P(\kappa_2 \in (1, 3)) = 0.95$.

A.4 Supplementary results

A.4.1 Variable selection for TGF- β and DLD dataset

To illustrate our methodology for robust regression in Section 5, we selected between the Gaussian model and Tukey’s loss when regressing the TGF- β and DLD gene expressions on a subset of the variables available in the full data sets. The procedures for which variables were selected was outlined in Sections 5.3.1 and 5.3.2. To ensure that our results are reproducible, below we indicate the selected covariates and the supplementary material contains code for these variable pre-screening steps. We also provide supplementary figures providing a visual inspection of the Gaussian fit to the residuals of the TGF- β and DLD examples.

TGF- β : For the TGF- β analysis we focused on 7 of the 10172 genes available in the data set that appear in the ‘TGF- β 1 pathway’ according to the KEGGREST package in R (Tenenbaum, 2016). These were the VIT, PDE4B, ATP8B1, MAGEA11, PDE6C, PDE9A and SEPTIN4 genes.

DLD: For the DLD analysis we selected the 15 genes with the 5 highest loadings in the first 3 principal components of the original 57 predictors. This procedure selected the following genes C15orf52, BRAT1, CYP26C1, SLC35B4, GRLF1, RXRA, RAB3GAP2, NOTCH2NL, SDC4, TTC22, PTCH2, ECH1, CSF2RA, TP53AIP1, and RRP1B.

Supplementary figures: We provide Figure A.2 to visually inspect the Gaussian approximation to the fitted standardised residuals of the TGF- β and DLD datasets. The left panels overlay the fitted Gaussian models to the histogram of the residuals, while the right are Q - Q Normal residual plots. The top panels show that the TGF- β data is well-approximated by a Gaussian model, whereas the bottom panels show that the DLD data set exhibits heavier tails. These graphical inspections support the \mathcal{H} -score findings from Section 5 and Figure 4.




Figure A.2: **Top:** TGF- β data, where the integrated \mathcal{H} -score selected the Gaussian model. **Bottom:** DLD data, where the integrated \mathcal{H} -score selected Tukey’s loss. **Left:** Gaussian approximations to the residuals. **Right:** Q - Q normal plot of the fitted residuals according to the Gaussian model.

A.4.2 Further Gaussian Mixture Examples

Additionally to the experiments in Section 6.1, here we provide two further Gaussian mixture data sets from Marron and Wand (1992)

- *Asymmetric:* 2-components, $\mu_1 = 0$, $\mu_2 = 1.5$, $\sigma_1 = 1$, $\sigma_2 = 1/3$, $m_1 = 0.75$ and $m_2 = 0.25$.

- *Kurtotic*: 2-components, $\mu_1 = \mu_2 = 0$, $\sigma_1 = 1$ $\sigma_2 = 0.1$, $m_1 = 2/3$ and $m_2 = 1/3$.

Figure A.3 plots the approximation to the underlying $g(y)$ of the four methods under consideration in Section 6.1, while Table A.1 estimates the Fisher’s divergence of the estimates (posterior mean where appropriate) to $g(y)$. For both examples the DPMM provides the best estimate according the Fisher’s divergence. For the *Kurtotic* data set, the \mathcal{H} -posterior KDE’s perform comparably to the DPMM and better than the default KDE implementation. However, for the *Asymmetric* data set the \mathcal{H} -posterior KDE’s under-smooth the data and perform considerably worse than the other methods.




Figure A.3: Density estimation for Gaussian mixture data. Histograms of observed data, standardised to 0 mean and unit variance, and estimated density by DPMM, R ’s density function with the bandwidth rule of Silverman (1986), the \mathcal{H} -posterior estimate with no tempering ($w = 1$), and with tempering ($w \neq 1$).

Table A.1: Fisher’s divergence between the density estimates and the data-generating Gaussian mixtures in two simulation scenarios. The best performing method in each is highlighted in bold face.

	KDE	DPMM	Hyvärinen KDE ($w = 1$)	Hyvärinen KDE ($w \neq 1$)
<i>Asymmetric</i>	0.25	0.04	2.17	2.24
<i>Kurtotic</i>	3.14	2.05	2.55	2.44