

The Third Order Scalar Induced Gravitational Waves

Jing-Zhi Zhou,^{*} Xukun Zhang,[†] Qing-Hua Zhu,[‡] and Zhe Chang[§]

Institute of High Energy Physics Chinese Academy of Sciences Beijing 100049 China and

University of Chinese Academy of Sciences Beijing 100049 China

(Dated: September 1, 2021)

Since the gravitational waves were detected by LIGO and VIRGO, it has been promising that lots of information about the primordial Universe could be learned by further observations on stochastic gravitational waves background. The studies on gravitational waves induced by primordial curvature perturbations are of great interest. The aim of this paper is to investigate the third order induced gravitational waves. Based on the theory of cosmological perturbations, the first order scalar induces the second order scalar, vector and tensor perturbations. At the next iteration, the first order scalar, the second order scalar, vector and tensor perturbations all induce the third order tensor perturbations. We present the energy density spectrum of the third order gravitational waves for a monochromatic primordial power spectrum. The shape of the energy density spectrum of the third order gravitational waves is different from that of the second order scalar induced gravitational waves. And it is found that the third order gravitational waves sourced by the second order scalar perturbations dominate the energy density spectrum.

I. INTRODUCTION

Inflationary cosmology suggests that cosmological perturbations are originated from quantum fluctuations in causal contact at the early Universe. Since the gravitational waves were detected by LIGO and VIRGO [1], it has been promising that the further observations on stochastic gravitational waves background could test inflationary models and might shed light on the quantum reality of the early Universe [2–5].

The cosmological perturbations are decomposed as scalar, vector and tensor perturbations based on helicity decomposition of metric perturbations. The cosmological perturbations generated at inflation epochs are known as primordial perturbations. So far, the measurements of cosmic microwave background and large-scale structure indicate that the scalar parts dominate the primordial perturbations [6, 7]. Therefore, it is of great interest to study gravitational waves induced by the primordial scalar perturbations [8–26], if there would be further directed observation on the gravitational wave background. Besides, the cosmic microwave background and large-scale structure constrained the primordial spectrum on the pivot scale larger than 1Mpc [6]. It did not tell the story about perturbations that re-enter the horizon on small scales, which also contain lots of information about the early Universe. In this sense, it is necessary to apply to future space-based and ground-based detections of stochastic gravitational waves background on small scales [5, 27–29], and theoretically study cosmological perturbations from the early Universe, such as the induced gravitational waves.

^{*} zhoujingzhi@ihep.ac.cn

[†] zhangxukun@ihep.ac.cn

[‡] zhuqh@ihep.ac.cn

[§] changz@ihep.ac.cn

The induced gravitational wave as a prediction of inflationary cosmology has been studied for many years [8, 9]. The energy density spectra of the induced gravitational waves in principle include the information about primordial black hole [30–63] and primordial non-Gaussianity. [64–68]. Besides, recent studies on induced gravitational wave were also extended to gauge issue [16–19, 23–25], epochs of the Universe [15, 20, 21] and modified gravity [69].

As suggested in pioneers' study [39], the higher order induced gravitational waves have non-trivial results comparing with the second order induced gravitational waves, and play important role in constraints on primordial black hole. In this paper, we also study induced gravitational waves to the third order in a radiation-dominated era. For scalar induced gravitational waves [9, 70], the source terms of the third order gravitational waves should contain not only the first order scalar perturbations, but also other three types of the second order perturbations, since the second order perturbations can also be induced by the first order scalar perturbations. It is found that the dominated parts of the power spectrum of the third order gravitational waves for a monochromatic primordial power spectrum come from the source term of the second order scalar perturbations. This is beyond previous study [39, 70] that neglected all the second order perturbations.

This paper is organized as follows. In Sec. II, we present the equation of motion of the third order induced gravitational waves. In Sec. III, we solve the equation of motion of the third order induced gravitational waves. The explicit expressions of kernel functions are obtained. In Sec. IV, the energy density spectrum of the third order induced gravitational waves is studied. Finally, the conclusions and discussions are summarized in Sec. V.

II. EQUATIONS OF MOTION OF THIRD ORDER INDUCED GRAVITATIONAL WAVES

In flat Friedmann-Robertson-Walker (FRW) spacetime, the background metric is given by

$$g_{\mu\nu}^{(0)} dx^\mu dx^\nu = a^2(\eta) (-d\eta^2 + \delta_{ij} dx^i dx^j) , \quad (1)$$

where η is the conformal time. The third order metric perturbations in Newtonian gauge take the form of

$$ds^2 = -a^2 \left[\left(1 + 2\phi^{(1)} + \phi^{(2)} \right) d\eta^2 + V_i^{(2)} d\eta dx^i + \left(\left(1 - 2\psi^{(1)} - \psi^{(2)} \right) \delta_{ij} + \frac{1}{2} h_{ij}^{(2)} + \frac{1}{6} h_{ij}^{(3)} \right) dx^i dx^j \right] , \quad (2)$$

where $\phi^{(n)}$ and $\psi^{(n)}$ ($n = 1, 2$) are the n -order scalar perturbations, $V_i^{(2)}$ is the second order vector perturbation, and $h_{ij}^{(n)}$ ($n = 2, 3$) are the n -order tensor perturbations. Based on the Einstein field equation for cosmological perturbations sourced by perfect fluid, we obtain the equation of motion of the third order induced gravitational waves,

$$h_{ij}^{(3)''}(\eta, \mathbf{x}) + 2\mathcal{H}h_{ij}^{(3)'}(\eta, \mathbf{x}) - \Delta h_{ij}^{(3)}(\eta, \mathbf{x}) = -12\Lambda_{ij}^{lm} \mathcal{S}_{lm}^{(3)}(\eta, \mathbf{x}) . \quad (3)$$

The expression of transverse and traceless operator is

$$\Lambda_{ij}^{lm} = \mathcal{T}_i^l \mathcal{T}_j^m - \frac{1}{2} \mathcal{T}_{ij} \mathcal{T}^{lm} , \quad (4)$$

where the expression of traceless operator is $\mathcal{T}_i^l = \delta_i^l - \partial^l \Delta^{-1} \partial_i$. For illustration, the source term $S_{lm}^{(3)}(\eta, \mathbf{x})$ can be divided into four parts,

$$S_{lm}^{(3)}(\eta, \mathbf{x}) = S_{lm,1}^{(3)}(\eta, \mathbf{x}) + S_{lm,2}^{(3)}(\eta, \mathbf{x}) + S_{lm,3}^{(3)}(\eta, \mathbf{x}) + S_{lm,4}^{(3)}(\eta, \mathbf{x}), \quad (5)$$

where $S_{lm,1}^{(3)}(\eta, \mathbf{x})$ is composed of the first order scalar perturbation $\phi^{(1)}$. Corresponding expression of $S_{lm,1}^{(3)}(\eta, \mathbf{x})$ is given by

$$\begin{aligned} S_{lm,1}^{(3)}(\eta, \mathbf{x}) &= 12\phi^{(1)}\partial_l\phi^{(1)}\partial_m\phi^{(1)} - \frac{4}{\mathcal{H}}\phi^{(1)'}\partial_l\phi^{(1)}\partial_m\phi^{(1)} + \frac{2}{3\mathcal{H}^2}\Delta\phi^{(1)}\partial_l\phi^{(1)}\partial_m\phi^{(1)} + \frac{2}{3\mathcal{H}^4}\Delta\phi^{(1)}\partial_l\phi^{(1)'}\partial_m\phi^{(1)'} \\ &\quad - \frac{3}{\mathcal{H}^2}\phi^{(1)'}\partial_l\phi^{(1)'}\partial_m\phi^{(1)} - \frac{3}{\mathcal{H}^2}\phi^{(1)'}\partial_m\phi^{(1)'}\partial_l\phi^{(1)} + \frac{2}{3\mathcal{H}^3}\Delta\phi^{(1)}\partial_l\phi^{(1)'}\partial_m\phi^{(1)} + \frac{2}{3\mathcal{H}^3}\Delta\phi^{(1)}\partial_m\phi^{(1)'}\partial_l\phi^{(1)} \\ &\quad - \frac{2}{\mathcal{H}^3}\phi^{(1)'}\partial_l\phi^{(1)'}\partial_m\phi^{(1)'} - \frac{4}{\mathcal{H}^2}\phi^{(1)}\partial_l\phi^{(1)'}\partial_m\phi^{(1)'} . \end{aligned} \quad (6)$$

The source term $S_{lm,2}^{(3)}(\eta, \mathbf{x})$ is composed of the first order scalar perturbation $\phi^{(1)}$ and the second order tensor perturbation $h_{lm}^{(2)}$. The expression of $S_{lm,2}^{(3)}(\eta, \mathbf{x})$ is shown to be

$$S_{lm,2}^{(3)}(\eta, \mathbf{x}) = -\frac{1}{2}\phi^{(1)}\left(h_{lm}^{(2)''} + 2\mathcal{H}h_{lm}^{(2)'} - \Delta h_{lm}^{(2)}\right) - \phi^{(1)}\Delta h_{lm}^{(2)} - \phi^{(1)'}\mathcal{H}h_{lm}^{(2)} - \frac{1}{3}\Delta\phi^{(1)}h_{lm}^{(2)} - \partial^b\phi^{(1)}\partial_b h_{lm}^{(2)}. \quad (7)$$

The source term $S_{lm,3}^{(3)}(\eta, \mathbf{x})$ is composed of the first order scalar perturbation $\phi^{(1)}$ and the second order vector perturbation $V_l^{(2)}$, namely,

$$\begin{aligned} S_{lm,3}^{(3)}(\eta, \mathbf{x}) &= \phi^{(1)}\partial_l\left(V_m^{(2)'} + 2\mathcal{H}V_m^{(2)}\right) + \phi^{(1)}\partial_m\left(V_l^{(2)'} + 2\mathcal{H}V_l^{(2)}\right) + \phi^{(1)'}\left(\partial_l V_m^{(2)} + \partial_m V_l^{(2)}\right) \\ &\quad - \frac{\phi^{(1)}}{8\mathcal{H}}\left(\partial_m\Delta V_l^{(2)} + \partial_l\Delta V_m^{(2)}\right) - \frac{\phi^{(1)'}}{8\mathcal{H}^2}\left(\partial_m\Delta V_l^{(2)} + \partial_l\Delta V_m^{(2)}\right). \end{aligned} \quad (8)$$

And the source term $S_{lm,4}^{(3)}(\eta, \mathbf{x})$ is composed of the first order scalar perturbation $\phi^{(1)}$ and the second order scalar perturbations $\phi^{(2)}$ and $\psi^{(2)}$,

$$S_{lm,4}^{(3)}(\eta, \mathbf{x}) = \frac{1}{\mathcal{H}}\left(\phi^{(1)}\partial_l\partial_m\psi^{(2)'}\right) + \frac{1}{\mathcal{H}}\left(\phi^{(1)'}\partial_l\partial_m\phi^{(2)}\right) + \frac{1}{\mathcal{H}^2}\left(\phi^{(1)'}\partial_l\partial_m\psi^{(2)'}\right) + 3\left(\phi^{(1)}\partial_l\partial_m\phi^{(2)}\right). \quad (9)$$

In this paper, we let the second order perturbations $h_{lm}^{(2)}$, $V_l^{(2)}$ and $\phi^{(2)}$ and $\psi^{(2)}$ be induced by the first order scalar perturbations $\phi^{(1)}$ ($= \psi^{(1)}$). The equations of motion of the second order perturbations have already shown in pioneers' studies [9, 71–74]. In this case, the seeds of the third gravitational waves are all attributed to the first order primordial curvature perturbations. In the following, we will solve the motion of equation for the third order gravitational waves $h_{ij}^{(3)}$ based on the expression of Eq. (3).

III. KERNEL FUNCTION OF THIRD ORDER INDUCED GRAVITATIONAL WAVES

In order to solve the equations of motion of third order induced gravitational waves, we rewrite Eq. (3) in momentum space as

$$h^{\lambda,(3)''}(\eta, \mathbf{k}) + 2\mathcal{H}h^{\lambda,(3)'}(\eta, \mathbf{k}) + k^2h^{\lambda,(3)}(\eta, \mathbf{k}) = \sum_{i=1}^4 12\mathcal{S}_i^{\lambda,(3)}(\eta, \mathbf{k}), \quad (10)$$

where $h^{\lambda,(3)}(\eta, \mathbf{k}) = \varepsilon^{\lambda,ij}(\mathbf{k})h_{ij}^{(3)}(\eta, \mathbf{k})$ and $S^{\lambda,(3)}(\eta, \mathbf{k}) = -\varepsilon^{\lambda,lm}(\mathbf{k})S_{lm}^{(3)}(\eta, \mathbf{k})$. The $\varepsilon_{ij}^{\lambda}(\mathbf{k})$ is polarization tensor, which satisfies $\varepsilon_{ij}^{\lambda}(\mathbf{k})\varepsilon^{\bar{\lambda},ij}(\mathbf{k}) = \delta^{\lambda\bar{\lambda}}$ and $\delta_{\lambda\bar{\lambda}}\varepsilon_{ij}^{\lambda}(\mathbf{k})\varepsilon^{\bar{\lambda},lm}(\mathbf{k}) = \Lambda_{ij}^{lm}(\mathbf{k})$. Unlike the second order tensor perturbation, the third order induced gravitational waves have four types of source terms $S_i^{\lambda,(3)}(\eta, \mathbf{k})$, ($i = 1, 2, 3, 4$) (Eqs. (6)–(9)).

As mentioned, we consider that the second order perturbations $h_{ij}^{(2)}$, $V_l^{(2)}$, $\phi^{(2)}$, and $\psi^{(2)}$ are induced by the first order scalar perturbations $\phi^{(1)}$ and $\psi^{(1)}$. It is convenient to rewrite the first order scalar perturbations in the form of

$$\psi(\eta, \mathbf{p}) = \phi(\eta, \mathbf{p}) = \Phi_{\mathbf{p}}T_{\phi}(p\eta), \quad (11)$$

where $\Phi_{\mathbf{p}}$ is initial value originated from primordial curvature perturbation, and transfer function $T_{\phi}(y) = \frac{9}{y^2} \left(\frac{\sqrt{3}}{y} \sin\left(\frac{y}{\sqrt{3}}\right) - \cos\left(\frac{y}{\sqrt{3}}\right) \right)$. Therefore, we can rewrite the source terms $S_i^{\lambda,(3)}(\eta, \mathbf{k})$ in terms of initial value of the first order scalar perturbations $\Phi_{\mathbf{p}}$ in momentum space, i.e.,

$$S_1^{\lambda,(3)}(\eta, \mathbf{k}) = \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} \varepsilon^{\lambda,lm}(\mathbf{k})(p_l - q_l)q_m \Phi_{\mathbf{k}-\mathbf{p}}\Phi_{\mathbf{p}-\mathbf{q}}\Phi_{\mathbf{q}}f_1^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta), \quad (12)$$

$$S_2^{\lambda,(3)}(\eta, \mathbf{k}) = \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} \varepsilon^{\lambda,lm}(\mathbf{k})\Lambda_{lm}^{rs}(\mathbf{p})q_rq_s \Phi_{\mathbf{k}-\mathbf{p}}\Phi_{\mathbf{p}-\mathbf{q}}\Phi_{\mathbf{q}}f_2^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta), \quad (13)$$

$$S_3^{\lambda,(3)}(\eta, \mathbf{k}) = \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} \varepsilon^{\lambda,lm}(\mathbf{k})(\mathcal{T}_m^s(\mathbf{p})p_l + \mathcal{T}_l^s(\mathbf{p})p_m) \frac{p^s}{p^2} \times q_rq_s \Phi_{\mathbf{k}-\mathbf{p}}\Phi_{\mathbf{p}-\mathbf{q}}\Phi_{\mathbf{q}}f_3^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta), \quad (14)$$

$$S_4^{\lambda,(3)}(\eta, \mathbf{k}) = \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} \varepsilon^{\lambda,lm}(\mathbf{k})p_l p_m \Phi_{\mathbf{k}-\mathbf{p}}\Phi_{\mathbf{p}-\mathbf{q}}\Phi_{\mathbf{q}}f_4^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta), \quad (15)$$

and the transfer functions $f_i^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta)$ are also originated from the evolution of the first order scalar perturbations. The expressions of the transfer functions are shown as follows,

$$\begin{aligned} f_1^{(3)}(u, \bar{u}, \bar{v}, x, y) &= 12T_{\phi}(ux)T_{\phi}(\bar{u}y)T_{\phi}(\bar{v}y) - 4ux \frac{d}{d(ux)}T_{\phi}(ux)T_{\phi}(\bar{u}y)T_{\phi}(\bar{v}y) - \frac{2u^2x^2}{3}T_{\phi}(ux)T_{\phi}(\bar{u}y)T_{\phi}(\bar{v}y) \\ &- 6\bar{u}uxy \frac{d}{d(ux)}T_{\phi}(ux) \frac{d}{d(\bar{u}y)}T_{\phi}(\bar{u}y)T_{\phi}(\bar{v}y) - \frac{4u^2\bar{u}x^2y}{3}T_{\phi}(ux) \frac{d}{d(\bar{u}y)}T_{\phi}(\bar{u}y)T_{\phi}(\bar{v}y) \\ &- 4\bar{u}\bar{v}y^2T_{\phi}(ux) \frac{d}{d(\bar{u}y)}T_{\phi}(\bar{u}y) \frac{d}{d(\bar{v}y)}T_{\phi}(\bar{v}y) - 2\bar{u}\bar{v}uy^2x \frac{d}{d(ux)}T_{\phi}(ux) \frac{d}{d(\bar{u}y)}T_{\phi}(\bar{u}y) \frac{d}{d(\bar{v}y)}T_{\phi}(\bar{v}y) \\ &- \frac{2u^2\bar{u}\bar{v}x^2y^2}{3}T_{\phi}(ux) \frac{d}{d(\bar{u}y)}T_{\phi}(\bar{u}y) \frac{d}{d(\bar{v}y)}T_{\phi}(\bar{v}y), \end{aligned} \quad (16)$$

$$\begin{aligned}
f_2^{(3)}(u, \bar{u}, \bar{v}, x, y) = & -6T_\phi(ux)T_\phi(\bar{u}y)T_\phi(\bar{v}y) - 4\bar{u}yT_\phi(ux)\frac{d}{d(\bar{u}y)}T_\phi(\bar{u}y)T_\phi(\bar{v}y) - T_\phi(ux)p^2I_h^{(2)}(\bar{u}, \bar{v}, y) \\
& - 2\bar{u}\bar{v}y^2T_\phi(ux)\frac{d}{d(\bar{u}y)}T_\phi(\bar{u}y)\frac{d}{d(\bar{v}y)}T_\phi(\bar{v}y) + \frac{u}{vy}\frac{d}{d(ux)}T_\phi(ux)p^2I_h^{(2)}(\bar{u}, \bar{v}, y) \\
& - \frac{u^2}{3v^2}T_\phi(ux)p^2I_h^{(2)}(\bar{u}, \bar{v}, y) - \frac{1-u^2-v^2}{2v^2}T_\phi(ux)p^2I_h^{(2)}(\bar{u}, \bar{v}, y) ,
\end{aligned} \tag{17}$$

$$\begin{aligned}
f_3^{(3)}(u, \bar{u}, \bar{v}, x, y) = & \frac{u}{v}\frac{d}{d(ux)}T_\phi(ux)pI_V^{(2)}(\bar{u}, \bar{v}, y) + \frac{y}{8}T_\phi(ux)pI_V^{(2)}(\bar{u}, \bar{v}, y) + \frac{y^2u}{8v}\frac{d}{d(ux)}T_\phi(ux)pI_V^{(2)}(\bar{u}, \bar{v}, y) \\
& - 4T_\phi(ux)T_\phi(\bar{u}y)T_\phi(\bar{v}y) + 8\bar{u}yT_\phi(ux)\frac{d}{d(\bar{u}y)}T_\phi(\bar{u}y)T_\phi(\bar{v}y) + 16T_\phi(ux)T_\phi(\bar{u}y)T_\phi(\bar{v}y) \\
& + 4\bar{u}\bar{v}y^2T_\phi(ux)\frac{d}{d(\bar{u}y)}T_\phi(\bar{u}y)\frac{d}{d(\bar{v}y)}T_\phi(\bar{v}y) ,
\end{aligned} \tag{18}$$

$$\begin{aligned}
f_4^{(3)}(u, \bar{u}, \bar{v}, x, y) = & y\left(T_\phi(ux)\frac{\partial}{\partial y}I_\psi^{(2)}(\bar{u}, \bar{v}, y)\right) + ux\left(\frac{d}{d(ux)}T_\phi(ux)(I_\psi^{(2)}(\bar{u}, \bar{v}, y) + f_\phi^{(2)}(\bar{u}, \bar{v}, y))\right) \\
& + uxy\left(\frac{d}{d(ux)}T_\phi(ux)\frac{\partial}{\partial y}I_\psi^{(2)}(\bar{u}, \bar{v}, y)\right) + 3\left(T_\phi(ux)(I_\psi^{(2)}(\bar{u}, \bar{v}, y) + f_\phi^{(2)}(\bar{u}, \bar{v}, y))\right) ,
\end{aligned} \tag{19}$$

where we have set $|k-p|=uk$, $p=vk$, $|p-q|=\bar{u}p$, $q=\bar{v}p$, $x=k\eta$, and $y=p\eta$ for illustration. Indicated by Eqs. (17)–(19), we have to calculate the kernel function of the second order scalar, vector, tensor perturbations induced by the first order scalar perturbations via reproducing pioneers' studies ($I_\psi^{(2)}(\bar{u}, \bar{v}, y)$, $I_\phi^{(2)}(\bar{u}, \bar{v}, y)$, $I_V^{(2)}(\bar{u}, \bar{v}, y)$, and $I_h^{(2)}(\bar{u}, \bar{v}, y)$) [9, 71–74].

Corresponding to different types of the source terms in Eqs. (12)–(15), the third order gravitational wave can be divided into four types of parts,

$$h^{\lambda,(3)}(\eta, \mathbf{k}) = h_1^{\lambda,(3)}(\eta, \mathbf{k}) + h_2^{\lambda,(3)}(\eta, \mathbf{k}) + h_3^{\lambda,(3)}(\eta, \mathbf{k}) + h_4^{\lambda,(3)}(\eta, \mathbf{k}) , \tag{20}$$

where $h_i^{\lambda,(3)}$, ($i=1, 2, 3, 4$) are defined with

$$h_1^{\lambda,(3)}(\eta, \mathbf{k}) = \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} \varepsilon^{\lambda,lm}(\mathbf{k})(p_l - q_l)q_m \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} I_1^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta) , \tag{21}$$

$$h_2^{\lambda,(3)}(\eta, \mathbf{k}) = \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} \varepsilon^{\lambda,lm}(\mathbf{k}) \Lambda_{lm}^{rs}(\mathbf{p}) q_r q_s \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} I_2^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta) , \tag{22}$$

$$\begin{aligned}
h_3^{\lambda,(3)}(\eta, \mathbf{k}) = & \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} \varepsilon^{\lambda,lm}(\mathbf{k}) (\mathcal{T}_m^s(\mathbf{p})p_l + \mathcal{T}_l^s(\mathbf{p})p_m) \frac{p^s}{p^2} q_r q_s \\
& \times \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} I_3^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta) ,
\end{aligned} \tag{23}$$

$$h_4^{\lambda,(3)}(\eta, \mathbf{k}) = \int \frac{d^3p}{(2\pi)^{3/2}} \int \frac{d^3q}{(2\pi)^{3/2}} \varepsilon^{\lambda,lm}(\mathbf{k}) p_l p_m \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} I_4^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta) . \tag{24}$$

The kernel functions $I_i^{(3)}(|\mathbf{k}-\mathbf{p}|, |\mathbf{p}-\mathbf{q}|, \mathbf{q}, \eta)$ include the information of the corresponding source terms. Substituting Eqs. (12)–(15) and (21)–(24) into Eq. (10), we obtain the equations of motion of kernel functions




FIG. 1. The squares of four types of the kernel functions of the third order tensor perturbations and the second order tensor perturbation. Here we have set $|\mathbf{k} - \mathbf{p}| = p = k$ and $|\mathbf{p} - \mathbf{q}| = q = p$.

$I_i^{(3)}(u, \bar{u}, \bar{v}, x)$ in the form of

$$I_i^{(3)''}(u, \bar{u}, \bar{v}, x) + 2\mathcal{H}I_i^{(3)'}(u, \bar{u}, \bar{v}, x) + k^2 I_i^{(3)}(u, \bar{u}, \bar{v}, x) = 12f_i^{(3)}(u, \bar{u}, \bar{v}, x), \quad (i = 1, 2, 3, 4). \quad (25)$$

The solutions of kernel functions can be expressed as

$$I_i^{(3)}(u, \bar{u}, \bar{v}, x) = \frac{12}{k^2} \int_0^x d\bar{x} \left(\frac{\bar{x}}{x} \sin(x - \bar{x}) f_i^{(3)}(u, \bar{u}, \bar{v}, x) \right), \quad (i = 1, 2, 3, 4). \quad (26)$$

Since the expression of power spectrum is composed of product of kernel functions $I_i^{(3)}$, we here present the $I_i(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, \mathbf{q}, \eta) I_j(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, \mathbf{q}, \eta)$, ($i, j = 1, 2, 3, 4$) as function of $k\eta$ in Fig. 1 and Fig. 2 for selected momentums. It shows that all the kernel functions are decay with $k\eta$, and the amplitude of $(I_3^{(3)})^2$ is the largest.

IV. THE ENERGY DENSITY SPECTRUM OF THIRD ORDER INDUCED GRAVITATIONAL WAVES

In this section, we will study the energy density spectrum of the third order induced gravitational waves. An explicit expression of the power spectrum $\mathcal{P}_h^{(3)}(\eta, \mathbf{k})$ is presented. First of all, the power spectrum of the third order gravitational waves can be written as the sum of sixteen terms,

$$\mathcal{P}_h^{(3)}(\eta, \mathbf{k}) = \sum_{i,j=1}^4 \mathcal{P}_h^{ij}(\eta, \mathbf{k}). \quad (27)$$




FIG. 2. The products $k^4 |I_i \times I_j|$ ($i \neq j$) of the kernel functions of the third order tensor perturbations. Here we have set $|\mathbf{k} - \mathbf{p}| = p = k$ and $|\mathbf{p} - \mathbf{q}| = q = p$.

It is known that the two-point correlation function and the power spectra are related by $\langle h_*^\lambda(\eta, \mathbf{k}) h_*^{\lambda'}(\eta, \mathbf{k}') \rangle = \delta^{\lambda\lambda'} \delta(\mathbf{k} + \mathbf{k}') \frac{2\pi^2}{k^3} \mathcal{P}_{h_*}(\eta, \mathbf{k})$. Based on the expression of $h_i^\lambda(\eta, \mathbf{k})$ in Eqs. (21)–(24), we obtain the power spectra $\mathcal{P}_h^{ij}(\eta, \mathbf{k})$ in the form of

$$\begin{aligned} \mathcal{P}_h^{ij}(\eta, k) &= \frac{k^3 (2\pi^2)^2}{2} \int \frac{d^3 p d^3 q d^3 p' d^3 q'}{(2\pi)^6} \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}') \\ &\quad \times I_i^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, q, \eta) I_j^{(3)}(|-\mathbf{k} - \mathbf{p}'|, |\mathbf{p}' - \mathbf{q}'|, q', \eta) \mathcal{C}(\mathbf{k}, \mathbf{k}', \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}'). \end{aligned} \quad (28)$$

It is easy to find that the $\mathcal{P}_h^{ij}(\eta, \mathbf{k})$ for different i, j is determined by the polynomial $\mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}')$. We obtain the expressions of $\mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}')$ by making use of the properties of polarization vectors and polarization tensors in Eq. (22) and Eq. (23). The $\mathcal{C}(\mathbf{k}, \mathbf{k}', \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}')$ is derived from simplification of six-point correlation function, the explicit expression is shown in Eq. (A1).

For illustration, we show the expression of $\mathcal{P}_h^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}')$ for the first term of $\mathcal{P}_h^{ij}(\eta, k)$ in Eq. (A1),

$$\begin{aligned} \mathcal{P}_h^{ij} &= \frac{k^3}{32\pi^2} \int \frac{d^3 p d^3 q}{|\mathbf{k} - \mathbf{p}|^3 |\mathbf{p} - \mathbf{q}|^3 q^3} \left\{ \mathbb{P}^{ij}(k, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) \right. \\ &\quad \left. \times I_i^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, q, \eta) I_j^{(3)}(|\mathbf{k} - \mathbf{p}|, |\mathbf{p} - \mathbf{q}|, q, \eta) P_\Phi(|\mathbf{k} - \mathbf{p}|) P_\Phi(|\mathbf{p} - \mathbf{q}|) P_\Phi(q) \right\} \quad (29) \\ &= \frac{1}{2\pi} \int_0^\infty dv \int_0^\infty d\bar{v} \int_{|1-\bar{v}v|}^{1+\bar{v}v} dw \int_{|1-v|}^{1+v} du \int_{\bar{u}_-}^{\bar{u}_+} d\bar{u} \left\{ \frac{w}{u^2 \bar{u}^2 v \bar{v}^2 \sqrt{Y(1-X^2)}} \mathbb{P}^{ij}(k, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) \right. \\ &\quad \left. \times I_i^{(3)}(u, \bar{u}v, \bar{v}v, x) I_j^{(3)}(u, \bar{u}v, \bar{v}v, x) P_\Phi(ku) P_\Phi(k\bar{u}v) P_\Phi(k\bar{v}v) \right\}, \quad (30) \end{aligned}$$

where we have used the substitution as follows in the second equal,

$$|\mathbf{k} - \mathbf{p}| = uk, |\mathbf{k} - \mathbf{q}| = wk, |\mathbf{p} - \mathbf{q}| = \bar{u}p = \bar{u}vk, q = \bar{v}p = \bar{v}vk, \quad (31)$$

and

$$\bar{u}_{\pm} = \left(1 + \bar{v}^2 - \frac{(1 + v^2 - u^2)(1 + (\bar{v}v)^2 - w^2)}{2v^2} \right. \\ \left. \pm 2\bar{v} \sqrt{\left(1 - \left(\frac{1 + v^2 - u^2}{2v} \right)^2 \right) \left(1 - \left(\frac{1 + (\bar{v}v)^2 - w^2}{2\bar{v}v} \right)^2 \right)} \right)^{\frac{1}{2}}, \quad (32)$$

$$X = \frac{-1 + u^2 + v^2 - 2\bar{u}^2v^2 + v^2\bar{v}^2 + u^2v^2\bar{v}^2 - v^4\bar{v}^2 + w^2 - u^2w^2 + v^2w^2}{\sqrt{(1 - 2u^2 + u^4 - 2v^2 - 2u^2v^2 + v^4)(1 - 2v^2\bar{v}^2 + v^4\bar{v}^4 - 2w^2 - 2v^2\bar{v}^2w^2 + w^4)}}, \quad (33)$$

$$Y = (1 - 2u^2 + u^4 - 2v^2 - 2u^2v^2 + v^4)(1 - 2v^2\bar{v}^2 + v^4\bar{v}^4 - 2w^2 - 2v^2\bar{v}^2w^2 + w^4). \quad (34)$$

In order to obtain a specific result of $\mathcal{P}_h^{(3)}(\eta, \mathbf{k})$, we can choose power spectrum of the first order scalar perturbation $P_{\Phi}(k)$ in the form of a monochromatic power spectrum,

$$P_{\Phi}(k) = Ak_*\delta(k - k_*). \quad (35)$$

In this case, the power spectrum \mathcal{P}_h^{ij} reduce to

$$\mathcal{P}_h^{ij} = \frac{\mathcal{A}^3 \tilde{k}^3}{2\pi} \Theta(3 - \tilde{k}) I_i^{(3)} \left(\frac{1}{\tilde{k}}, \frac{1}{\tilde{k}}, \frac{1}{\tilde{k}}, x \right) I_j^{(3)} \left(\frac{1}{\tilde{k}}, \frac{1}{\tilde{k}}, \frac{1}{\tilde{k}}, x \right) \\ \times \int_{|1 - \frac{1}{\tilde{k}}|}^{\min\{\frac{2}{\tilde{k}}, 1 + \frac{1}{\tilde{k}}\}} dv \int_{w_-}^{w_+} dw \left\{ vw \left(\frac{1}{\sqrt{Y(1 - X^2)}} \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) \right)_{u=\frac{1}{\tilde{k}}, \bar{u}=\bar{v}=\frac{1}{\tilde{k}}} \right\}, \quad (36)$$

where we have defined $\tilde{k} \equiv \frac{k}{k_*}$ and

$$w_{\pm} = \sqrt{\frac{1}{2} + \frac{3}{2\tilde{k}^2} - \frac{1}{2}v^2 \pm \frac{1}{\tilde{k}v} \sqrt{\left(v^2 - \frac{4}{\tilde{k}^2} \right) \left(v^2 - \left(1 - \frac{1}{\tilde{k}} \right)^2 \right) \left(v - \left(1 + \frac{1}{\tilde{k}} \right)^2 \right)}}. \quad (37)$$

For illustration, we presented the expression of power spectrum \mathcal{P}_h^{ij} in Eq. (36) corresponding with the first term in Eq. (A1) only. In fact, we have obtained a complete expression of power spectrum \mathcal{P}_h^{ij} in Eq. (36) via substituting

$$\mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) \rightarrow \mathbb{P}^{ij} \equiv \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, -\mathbf{q}) + \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, -\mathbf{p}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{q}) \\ + \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, \mathbf{q} - \mathbf{p}) + \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{p} - \mathbf{k} - \mathbf{q}, \mathbf{q}, \mathbf{p} - \mathbf{k}) \\ + \mathbb{P}^{ij}(\mathbf{k}, \mathbf{p}, \mathbf{q} - \mathbf{k}, \mathbf{q}, -\mathbf{k} + \mathbf{p}). \quad (38)$$

And the explicit expressions of $\mathbb{P}^{ij}|_{u=1/\tilde{k}, \bar{u}=\bar{v}=1/\tilde{k}}$ in Eq. (36) are shown in the Appendix B.




FIG. 3. Energy density fractions of induced gravitational waves to the second order (blue curve) and to the third order (orange curve) as function of k for $k\eta = 1000$. We have set $A = 0.005$ in the plot.

In the end of this section, we calculate the third order power spectrum $\mathcal{P}_h^{(3)}$, i.e.,

$$\mathcal{P}_h^{(3)} = \sum_{i,j=1}^4 \mathcal{P}_h^{ij} . \quad (39)$$

The fraction of the gravitational waves energy density per logarithmic wavelength is given by

$$\Omega_{\text{GW}}(\eta, k) = \frac{\rho_{\text{GW}}(\eta, k)}{\rho_{\text{tot}}(\eta)} = \frac{1}{24} \left(\frac{k}{a(\eta)H(\eta)} \right)^2 \mathcal{P}_h(\eta, k) , \quad (40)$$

where

$$\mathcal{P}_h(\eta, k) = \frac{1}{4} \mathcal{P}_h^{(2)}(\eta, k) + \frac{1}{36} \mathcal{P}_h^{(3)}(\eta, k) + O\left(\mathcal{P}_h^{(4)}\right) . \quad (41)$$

Evaluating power spectrum $\mathcal{P}_h(\eta, k)$ in Eq. (40) by using Eqs. (36) and (39), we obtain the energy density fraction of the third order induced gravitational waves. As it is shown in Fig. 3, the shape of density fraction is different when considering the third order gravitational waves. There would be potential issue on the shape of the power spectrum, as we did not take an oscillation average with respect to $k\eta$. In Fig. 4, we present the third order power spectra of selected sources. It shows that the third order gravitational waves sourced by the second order scalar perturbations dominate the energy density spectrum.

V. CONCLUSIONS AND DISCUSSIONS

In this paper, we studied the third order scalar induced gravitational waves. The source terms of the third order scalar induced gravitational waves were completely considered. For illustration, we presented energy density spectrum of the third order gravitational waves for a monochromatic primordial power spectrum. The




FIG. 4. The third order power spectra of the third order gravitational waves induced by selected source as function of \tilde{k} . Here, we let $A = 0.005$ and $k\eta = 1000$. The blue, orange, and green curve represent the power spectra sourced by $S_{lm,1}^{(3)}$, both $S_{lm,1}^{(3)}$ and $S_{lm,4}^{(3)}$, and $S_{lm}^{(3)}$, respectively.

explicit integral expressions given in Eqs. (30) and (36) are shown to be useful for evaluating the energy density spectrum of the third order gravitational waves. Our results are different from the pioneer's study [70] in principle, which neglected all the second cosmological perturbations. It was found that the third order gravitational waves sourced by the second order scalar perturbations dominates the energy density spectrum.

The source term of the third order induced gravitational waves $h_{ij}^{(3)}$ is completely different from the second order induced gravitational waves $h_{ij}^{(2)}$. Since the first order scalar, vector, and tensor perturbations are independent, it is nothing wrong to set the vector and tensor perturbations to be zero. The second order induced gravitational wave sourced by the first order scalar perturbations only is well-defined. However, for the third order gravitational wave $h_{ij}^{(3)}$, the sources include the first order scalar perturbations and three types of the second order perturbations. The second order perturbations induced by the first order scalar perturbations are not independent at all. One can not set any the second order perturbation to be zero. For instance, if we set the second order vector perturbation $V_j^{(2)}$ to zero, the right hand side of equation of motion will be zero. It will bring an unphysical constraint to the first order scalar perturbation $\phi^{(1)}$. This discussion can also be applied to higher order gravitational waves.

We calculated four types of kernel functions of the third order induced gravitational waves, there are sixteen combinations of the product of kernel functions $I_i^{(3)}I_j^{(3)}$. As shown in Figs (1) and (2), the $(I_3^{(3)})^2$ is much larger than others. Therefore, it is expected that the second order vector perturbations would dominate the amplitude of the third order gravitational waves. However, for a monochromatic primordial power spectrum, we found that the \mathbb{P}_h^{33} is zero. In this case, the corresponding power spectrum \mathcal{P}_h^{33} has no relevance with the power spectrum of the third order scalar induced gravitational waves. In this sense, it is necessary to consider other primordial power spectrum in further study, for example, the power-law spectrum. And non-trivial results might be obtained for a general primordial power spectrum.

We considered the third order gravitational waves induced by the first order scalar perturbations. In princi-

ple, there should be second order primordial curvature perturbation [71] that could also induce the third order gravitational waves. Since the correlation function between the first and second order scalar perturbations refers to a careful calculation in an inflation model, relevant studies might be given in the future.

It has to be clarified that this paper studied the power spectrum of the third order gravitational waves instead of the third order power spectrum of gravitational waves. Obviously, the latter is more important for observation and should include the contribution from the fourth order gravitational waves. An incomplete study of the third order power spectrum of gravitational waves was given in Refs. [39, 70], which neglected all the contributions from the higher order cosmological perturbations. Perhaps, a complete study on the third order power spectrum of induced gravitational waves might be presented in the future.

ACKNOWLEDGMENTS

We thank Prof. Sai Wang for the useful discussions. This work has been funded by the National Nature Science Foundation of China under grant No. 12075249 and 11690022, and the Key Research Program of the Chinese Academy of Sciences under Grant No. XDPB15

Appendix A: Six-Point Correlation Function

In this appendix, we use wick theorem to simplify the six-point correlation function in Sec. IV. We define $\mathcal{C}(\mathbf{k}, \mathbf{k}', \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}')$ as

$$\begin{aligned}
\mathcal{C}(\mathbf{k}, \mathbf{k}', \mathbf{p}, \mathbf{p}', \mathbf{q}, \mathbf{q}') &= \frac{1}{(2\pi^2)^3} \int d^3\mathbf{k}' \langle \Phi_{\mathbf{k}-\mathbf{p}} \Phi_{\mathbf{p}-\mathbf{q}} \Phi_{\mathbf{q}} \Phi_{\mathbf{k}'-\mathbf{p}'} \Phi_{\mathbf{p}'-\mathbf{q}'} \Phi_{\mathbf{q}'} \rangle \\
&= \left[\left(\frac{1}{(k-p)^3 q^3 (p'-q')^3} \delta(\mathbf{q}-\mathbf{k}) \delta(\mathbf{p}') P_{\Phi}(k-p) P_{\Phi}(q) P_{\Phi}(p'-q') \right) \right. \\
&+ \left(\frac{1}{(k-p)^3 q^3 (-k-p')^3} \delta(\mathbf{q}-\mathbf{k}) \delta(-\mathbf{k}-\mathbf{p}'+\mathbf{q}') P_{\Phi}(k-p) P_{\Phi}(q) P_{\Phi}(-k-p') \right) \\
&+ \left(\frac{1}{(k-p)^3 q^3 (-k-p')^3} \delta(\mathbf{q}+\mathbf{q}') \delta(-\mathbf{k}-\mathbf{q}') P_{\Phi}(k-p) P_{\Phi}(q) P_{\Phi}(-k-p') \right) \\
&+ \left(\frac{1}{(k-p)^3 (p-q)^3 (p'-q')^3} \delta(\mathbf{p}-\mathbf{q}-\mathbf{k}) \delta(\mathbf{p}') P_{\Phi}(k-p) P_{\Phi}(p-q) P_{\Phi}(p'-q') \right) \\
&+ \left(\frac{1}{(k-p)^3 (p-q)^3 (-k-p')^3} \delta(\mathbf{p}-\mathbf{q}-\mathbf{k}) \delta(-\mathbf{k}-\mathbf{p}'+\mathbf{q}') P_{\Phi}(k-p) P_{\Phi}(p-q) P_{\Phi}(-k-p') \right) \\
&+ \left(\frac{1}{(k-p)^3 (p-q)^3 (-k-p')^3} \delta(\mathbf{p}-\mathbf{q}-\mathbf{k}) \delta(-\mathbf{k}-\mathbf{q}') P_{\Phi}(k-p) P_{\Phi}(p-q) P_{\Phi}(-k-p') \right) \\
&+ \left(\frac{1}{(k-p)^3 (p-q)^3 (p'-q')^3} \delta(\mathbf{p}) \delta(\mathbf{p}') P_{\Phi}(k-p) P_{\Phi}(p-q) P_{\Phi}(p'-q') \right) \\
&+ \left(\frac{1}{(k-p)^3 (p-q)^3 q^3} \delta(\mathbf{p}+\mathbf{p}') \delta(\mathbf{q}+\mathbf{q}') P_{\Phi}(k-p) P_{\Phi}(p-q) P_{\Phi}(q) \right) \\
&+ \left(\frac{1}{(k-p)^3 (p-q)^3 q^3} \delta(\mathbf{p}+\mathbf{p}') \delta(\mathbf{q}+\mathbf{p}'-\mathbf{q}') P_{\Phi}(k-p) P_{\Phi}(p-q) P_{\Phi}(q) \right) \\
&+ \left(\frac{1}{(k-p)^3 (p-q)^3 (-k-p')^3} \delta(\mathbf{p}) \delta(-\mathbf{k}-\mathbf{p}'+\mathbf{q}') P_{\Phi}(k-p) P_{\Phi}(p-q) P_{\Phi}(-k-p') \right) \\
&+ \left(\frac{1}{(k-p)^3 (p-q)^3 q^3} \delta(\mathbf{p}-\mathbf{q}-\mathbf{k}-\mathbf{p}') \delta(\mathbf{q}+\mathbf{q}') P_{\Phi}(k-p) P_{\Phi}(p-q) P_{\Phi}(q) \right) \\
&+ \left(\frac{1}{(k-p)^3 (p-q)^3 q^3} \delta(\mathbf{p}-\mathbf{q}+\mathbf{q}') \delta(\mathbf{q}-\mathbf{k}-\mathbf{p}') P_{\Phi}(k-p) P_{\Phi}(p-q) P_{\Phi}(q) \right) \\
&+ \left(\frac{1}{(k-p)^3 (p-q)^3 (-k-p')^3} \delta(\mathbf{p}) \delta(-\mathbf{k}-\mathbf{q}') P_{\Phi}(k-p) P_{\Phi}(p-q) P_{\Phi}(-k-p') \right) \\
&+ \left(\frac{1}{(k-p)^3 (p-q)^3 q^3} \delta(\mathbf{p}-\mathbf{k}-\mathbf{q}') \delta(\mathbf{q}+\mathbf{p}'-\mathbf{q}') P_{\Phi}(k-p) P_{\Phi}(p-q) P_{\Phi}(q) \right) \\
&+ \left. \left(\frac{1}{(k-p)^3 (p-q)^3 q^3} \delta(-\mathbf{k}+\mathbf{p}-\mathbf{q}') \delta(\mathbf{q}-\mathbf{k}-\mathbf{p}') P_{\Phi}(k-p) P_{\Phi}(p-q) P_{\Phi}(q) \right) \right]. \tag{A1}
\end{aligned}$$

Appendix B: The Expressions of $\mathbb{P}^{ij}(\eta, \mathbf{k})$

Explicit expressions of $\mathbb{P}^{ij}|_{u=1/\tilde{k}, \bar{u}=\bar{v}=1/\tilde{k}}$ in right hand side of Eq. (38) are shown as follows,

$$\begin{aligned}
\mathbb{P}^{11} &= \frac{k^4}{16\tilde{k}^8} (\tilde{k}^8 (v^6 (w^2 + 1) + v^4 (2w^4 - 3w^2 - 1) + v^2 (2w^6 - 4w^4 + 3w^2 + 1) + w^2 (w^6 \\
&- 2w^4 + 2w^2 - 1)) - \tilde{k}^6 (v^6 + v^4 (7w^2 + 4) + v^2 (10w^4 - 4w^2 + 3) + 6w^6 - 3w^2 \\
&+ 4) + \tilde{k}^4 (5v^4 + v^2 (17w^2 + 11) + 14w^4 + 13w^2 + 14) - \tilde{k}^2 (9v^2 + 15w^2 + 16) + 6), \tag{B1}
\end{aligned}$$

$$\begin{aligned}
\mathbb{P}^{12} + \mathbb{P}^{21} = & -\frac{k^4}{128\tilde{k}^{10}v^2w^2(\tilde{k}^2(v^2+w^2-1)-3)}(\tilde{k}^{12}v^{12}w^2 + 2\tilde{k}^{10}v^{10}w^2(\tilde{k}^2(10w^2+1)-11) + \tilde{k}^8v^8w^2(\tilde{k}^4(56w^4 \\
& - 89w^2 - 5) + \tilde{k}^2(12 - 203w^2) + 137) + 4(\tilde{k}^2 - 1)^2w^2(\tilde{k}^6(2w^6 - 4w^4 + 3w^2 - 1) + \tilde{k}^4(-12w^4 \\
& - 2w^2 + 11) + \tilde{k}^2(23w^2 + 37) - 15) + \tilde{k}^4v^6(\tilde{k}^8w^2(73w^6 - 200w^4 + 132w^2 - 7) + \tilde{k}^6(-416w^6 \\
& + 566w^4 + 75w^2 + 4) + \tilde{k}^4(742w^4 - 161w^2 - 12) + \tilde{k}^2(12 - 355w^2) - 4) + \tilde{k}^2v^4(\tilde{k}^{10}w^2(54w^8 - 162w^6 \\
& + 176w^4 - 77w^2 + 10) - \tilde{k}^8(394w^8 - 760w^6 + 329w^4 + 66w^2 + 4) + \tilde{k}^6(1048w^6 - 1055w^4 - 254w^2 \\
& - 8) + \tilde{k}^4(-1163w^4 + 442w^2 + 48) + 4\tilde{k}^2(95w^2 - 14) + 20) + v^2(\tilde{k}^{12}w^2(18w^{10} - 54w^8 + 61w^6 \\
& - 32w^4 + 8w^2 - 1) + \tilde{k}^{10}w^2(-162w^8 + 310w^6 - 160w^4 - 10w^2 + 23) + \tilde{k}^8(557w^8 - 576w^6 - 82w^4 \\
& + 118w^2 + 8) + \tilde{k}^6(-896w^6 + 482w^4 + 382w^2) + \tilde{k}^4(626w^4 - 437w^2 - 48) + \tilde{k}^2(64 - 85w^2) - 24), \\
\end{aligned} \tag{B2}$$

$$\begin{aligned}
\mathbb{P}^{13} + \mathbb{P}^{31} = & \frac{k^4}{32\tilde{k}^8}(\tilde{k}^2(v^2+1)-1)(\tilde{k}^6(v^6+v^4(3w^2-2)+v^2(3w^4-4w^2+2)+2w^6-3w^4 \\
& + 3w^2-1) - \tilde{k}^4(6v^4+2v^2(6w^2-1)+9w^4+2w^2-1) + 3\tilde{k}^2(4v^2+5w^2+3)-9), \\
\end{aligned} \tag{B3}$$

$$\begin{aligned}
\mathbb{P}^{14} + \mathbb{P}^{41} = & -\frac{k^4}{16\tilde{k}^8}(\tilde{k}^8(v^8+v^6(3w^2-1)+v^4(5w^4-13w^2+2)+v^2(4w^6-14w^4+13w^2-1)+2w^8 \\
& - 4w^6+5w^4-3w^2+1) - \tilde{k}^6(7v^6+v^4(19w^2+6)+v^2(22w^4-26w^2+1)+12w^6+2w^4 \\
& - 5w^2+18) + \tilde{k}^4(20v^4+v^2(41w^2+29)+29w^4+31w^2+48) - \tilde{k}^2(27v^2+33w^2+46)+15), \\
\end{aligned} \tag{B4}$$

$$\begin{aligned}
\mathbb{P}^{22} = & \frac{k^4}{512\tilde{k}^{12}v^4w^2(\tilde{k}^2(v^2+w^2-1)-3)}(5\tilde{k}^{14}v^{14}w^2 + 8(\tilde{k}^2-1)^4w^2(\tilde{k}^2(w^2-1)-3) + 10\tilde{k}^{12}v^{12}w^2(\tilde{k}^2(5w^2 \\
& - 4) - 6) + \tilde{k}^8v^{10}(\tilde{k}^6w^2(130w^4 - 295w^2 + 69) + \tilde{k}^4(-465w^4 + 396w^2 + 2) + \tilde{k}^2(279w^2 - 4) + 2) \\
& + \tilde{k}^6v^8(\tilde{k}^8w^2(165w^6 - 550w^4 + 416w^2 - 57) - \tilde{k}^6(930w^6 - 1994w^4 + 349w^2 + 14) + \tilde{k}^4(1566w^4 - 1539w^2 \\
& + 10) + \tilde{k}^2(22 - 583w^2) - 18) + 2(\tilde{k}^2 - 1)^2v^2(\tilde{k}^8(18w^8 - 36w^6 + 17w^4 + w^2) + \tilde{k}^6(-108w^6 + 106w^4 \\
& + 9w^2 - 4) + \tilde{k}^4(189w^4 - 29w^2 + 12) + \tilde{k}^2(68 - 77w^2) - 12) + \tilde{k}^4v^6(\tilde{k}^{10}w^2(120w^8 - 410w^6 + 454w^4 \\
& - 187w^2 + 22) + \tilde{k}^8(-870w^8 + 2356w^6 - 1433w^4 + 20w^2 + 14) + \tilde{k}^6(2214w^6 - 4341w^4 + 532w^2 + 56) \\
& + \tilde{k}^4(-2167w^4 + 2724w^2 - 100) + 6\tilde{k}^2(69w^2 - 4) + 54) + \tilde{k}^2v^4(\tilde{k}^{12}(40w^{12} - 120w^{10} + 117w^8 - 34w^6 \\
& - 4w^4 + w^2) + \tilde{k}^{10}(-360w^{10} + 878w^8 - 586w^6 + 18w^4 + 51w^2 - 2) + \tilde{k}^8(1197w^8 - 2406w^6 + 838w^4 \\
& + 278w^2 - 30) - 2\tilde{k}^6(855w^6 - 1543w^4 + 79w^2 + 54) + \tilde{k}^4(798w^4 - 1879w^2 + 268) + \tilde{k}^2(171w^2 - 82) - 46), \\
\end{aligned} \tag{B5}$$

$$\begin{aligned}
\mathbb{P}^{23} + \mathbb{P}^{32} = & -\frac{k^4}{128\tilde{k}^{10}v^2w^2(\tilde{k}^2(v^2+w^2-1)-3)}(5\tilde{k}^{12}v^{12}w^2 + 10\tilde{k}^{10}v^{10}w^2(\tilde{k}^2(2w^2-1)-5) + 2(\tilde{k}^2 \\
& - 1)^3w^2(\tilde{k}^4(2w^4-3w^2+1) + \tilde{k}^2(6-9w^2)+9) + \tilde{k}^6v^8(\tilde{k}^6w^2(30w^4-25w^2-1) + \tilde{k}^4(-155w^4+76w^2 \\
& + 2) + \tilde{k}^2(189w^2-4)+2) + \tilde{k}^4v^6(\tilde{k}^8w^2(25w^6-20w^4-18w^2+11) + \tilde{k}^6(-180w^6+138w^4+35w^2 \\
& - 2) + \tilde{k}^4(432w^4-231w^2-10) + \tilde{k}^2(26-327w^2)-14) + \tilde{k}^2v^4(\tilde{k}^{10}w^2(10w^8-32w^4+25w^2-4) + \tilde{k}^8(- \\
& - 100w^8+52w^6+147w^4-76w^2-2) + \tilde{k}^6w^2(348w^4-273w^2-140) + \tilde{k}^4(-507w^4+364w^2+36) \\
& + 16\tilde{k}^2(15w^2-4)+30) + (\tilde{k}^2-1)v^2(\tilde{k}^{10}w^2(10w^8-25w^6+18w^4-2w^2-1) + \tilde{k}^8(-75w^8+172w^6 \\
& - 108w^4+12w^2+2) + 2\tilde{k}^6(97w^6-201w^4+65w^2+8) - 4\tilde{k}^4(48w^4-83w^2-3) + \tilde{k}^2(39w^2-48)+18), \\
\end{aligned} \tag{B6}$$

$$\begin{aligned}
\mathbb{P}^{24} + \mathbb{P}^{24} = & \frac{k^4}{32\tilde{k}^{10}v^2w^2(\tilde{k}^2(v^2 + w^2 - 1) - 3)} (3\tilde{k}^{12}v^{12}w^2 + 2\tilde{k}^{10}v^{10}w^2(\tilde{k}^2(5w^2 - 7) - 13) + \tilde{k}^6v^8(\tilde{k}^6w^2(18w^4 \\
& - 57w^2 + 22) + \tilde{k}^4(-79w^4 + 94w^2 + 1) + \tilde{k}^2(88w^2 - 2) + 1) + (\tilde{k}^2 - 1)^2w^2(\tilde{k}^6(2w^6 - 4w^4 \\
& + 5w^2 - 3) - 3\tilde{k}^4(4w^4 + 2w^2 - 3) + \tilde{k}^2(25w^2 + 47) - 21) + \tilde{k}^4v^6(\tilde{k}^8w^2(19w^6 - 90w^4 + 89w^2 \\
& - 16) - \tilde{k}^6(118w^6 - 300w^4 + 99w^2 + 3) + \tilde{k}^4(239w^4 - 218w^2 + 1) + \tilde{k}^2(7 - 147w^2) - 5) \\
& + \tilde{k}^2v^4(\tilde{k}^{10}w^2(12w^8 - 56w^6 + 90w^4 - 51w^2 + 5) + \tilde{k}^8(-92w^8 + 268w^6 - 238w^4 + 35w^2 + 3) \\
& + \tilde{k}^6(266w^6 - 391w^4 + 97w^2 - 4) + \tilde{k}^4(-328w^4 + 193w^2 + 6) + 2\tilde{k}^2(59w^2 - 6) + 7) + v^2(\tilde{k}^{12}w^4(w^2 \\
& - 1)^2(4w^4 - 4w^2 + 7) + \tilde{k}^{10}(-36w^{10} + 54w^8 - 38w^6 + 18w^4 + 5w^2 - 1) + \tilde{k}^8(127w^8 - 46w^6 \\
& - 48w^4 + 5) - 2\tilde{k}^6(109w^6 + 9w^4 - 45w^2 - 5) + \tilde{k}^4(169w^4 - 72w^2 - 34) - 23\tilde{k}^2(w^2 - 1) - 3)
\end{aligned} \tag{B7}$$

$$\mathbb{P}^{33} = 0, \tag{B8}$$

$$\begin{aligned}
\mathbb{P}^{34} + \mathbb{P}^{43} = & -\frac{k^4}{16\tilde{k}^8}(\tilde{k}^2(v^2 + 1) - 1)(\tilde{k}^6(v^6 + v^4(3w^2 - 2) + v^2(3w^4 - 4w^2 + 2) + 2w^6 - 3w^4 \\
& + 3w^2 - 1) - \tilde{k}^4(6v^4 + 2v^2(6w^2 - 1) + 9w^4 + 2w^2 - 1) + 3\tilde{k}^2(4v^2 + 5w^2 + 3) - 9),
\end{aligned} \tag{B9}$$

$$\begin{aligned}
\mathbb{P}^{44} = & \frac{k^4}{8\tilde{k}^8}(\tilde{k}^8(v^8 + v^6(w^2 - 3) + v^4(w^4 - 7w^2 + 4) + v^2(-6w^4 + 7w^2 - 3) + w^4 - w^2 + 1) - \tilde{k}^6(5v^6 + v^4(5w^2 - 2) \\
& + v^2(2w^4 - 18w^2 - 5) + 2w^4 + w^2 + 10) + \tilde{k}^4(10v^4 + 7v^2(w^2 + 1) + w^4 + 5w^2 + 20) - \tilde{k}^2(9v^2 + 3w^2 + 14) + 3).
\end{aligned} \tag{B10}$$

-
- [1] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Phys. Rev. Lett.* **116**, 061102 (2016), arXiv:1602.03837 [gr-qc].
- [2] M. Sasaki, T. Suyama, T. Tanaka, and S. Yokoyama, *Class. Quant. Grav.* **35**, 063001 (2018), arXiv:1801.05235 [astro-ph.CO].
- [3] Z. Arzoumanian *et al.* (NANOGrav), *Astrophys. J.* **821**, 13 (2016), arXiv:1508.03024 [astro-ph.GA].
- [4] Z. Arzoumanian *et al.* (NANOGrav), *Astrophys. J.* **859**, 47 (2018), arXiv:1801.02617 [astro-ph.HE].
- [5] Z. Arzoumanian *et al.* (NANOGrav), *Astrophys. J. Lett.* **905**, L34 (2020), arXiv:2009.04496 [astro-ph.HE].
- [6] Y. Akrami *et al.* (Planck), *Astron. Astrophys.* **641**, A10 (2020), arXiv:1807.06211 [astro-ph.CO].
- [7] N. Aghanim *et al.* (Planck), *Astron. Astrophys.* **641**, A6 (2020), arXiv:1807.06209 [astro-ph.CO].
- [8] K. N. Ananda, C. Clarkson, and D. Wands, *Phys. Rev. D* **75**, 123518 (2007), arXiv:gr-qc/0612013.
- [9] D. Baumann, P. J. Steinhardt, K. Takahashi, and K. Ichiki, *Phys. Rev. D* **76**, 084019 (2007), arXiv:hep-th/0703290.
- [10] H. Assadullahi and D. Wands, *Phys. Rev. D* **81**, 023527 (2010), arXiv:0907.4073 [astro-ph.CO].
- [11] L. Alabidi, K. Kohri, M. Sasaki, and Y. Sendouda, *JCAP* **09**, 017 (2012), arXiv:1203.4663 [astro-ph.CO].
- [12] M. Guzzetti, N. Bartolo, M. Liguori, and S. Matarrese, *Riv. Nuovo Cim.* **39**, 399 (2016), arXiv:1605.01615 [astro-ph.CO].
- [13] J.-C. Hwang, D. Jeong, and H. Noh, *Astrophys. J.* **842**, 46 (2017), arXiv:1704.03500 [astro-ph.CO].
- [14] C. Caprini and D. G. Figueroa, *Class. Quant. Grav.* **35**, 163001 (2018), arXiv:1801.04268 [astro-ph.CO].
- [15] K. Kohri and T. Terada, *Phys. Rev. D* **97**, 123532 (2018), arXiv:1804.08577 [gr-qc].
- [16] K. Tomikawa and T. Kobayashi, *Phys. Rev. D* **101**, 083529 (2020), arXiv:1910.01880 [gr-qc].
- [17] V. De Luca, G. Franciolini, A. Kehagias, and A. Riotto, *JCAP* **03**, 014 (2020), arXiv:1911.09689 [gr-qc].
- [18] K. Inomata and T. Terada, *Phys. Rev. D* **101**, 023523 (2020), arXiv:1912.00785 [gr-qc].

- [19] C. Yuan, Z.-C. Chen, and Q.-G. Huang, *Phys. Rev. D* **101**, 063018 (2020), arXiv:1912.00885 [astro-ph.CO].
- [20] G. Domènech, *Int. J. Mod. Phys. D* **29**, 2050028 (2020), arXiv:1912.05583 [gr-qc].
- [21] G. Domènech, S. Pi, and M. Sasaki, *JCAP* **08**, 017 (2020), arXiv:2005.12314 [gr-qc].
- [22] S. Pi and M. Sasaki, *JCAP* **09**, 037 (2020), arXiv:2005.12306 [gr-qc].
- [23] A. Ali, Y. Gong, and Y. Lu, (2020), arXiv:2009.11081 [gr-qc].
- [24] Z. Chang, S. Wang, and Q.-H. Zhu, (2020), arXiv:2009.11994 [gr-qc].
- [25] Z. Chang, S. Wang, and Q.-H. Zhu, (2020), arXiv:2010.01487 [gr-qc].
- [26] K. T. Abe, Y. Tada, and I. Ueda, (2020), arXiv:2010.06193 [astro-ph.CO].
- [27] P. Amaro-Seoane *et al.*, “Laser interferometer space antenna,” (2017), arXiv:1702.00786 [astro-ph.IM].
- [28] J. Luo *et al.* (TianQin), *Class. Quant. Grav.* **33**, 035010 (2016), arXiv:1512.02076 [astro-ph.IM].
- [29] W.-H. Ruan, Z.-K. Guo, R.-G. Cai, and Y.-Z. Zhang, *Int. J. Mod. Phys. A* **35**, 2050075 (2020), arXiv:1807.09495 [gr-qc].
- [30] R. Saito and J. Yokoyama, *Phys. Rev. Lett.* **102**, 161101 (2009), [Erratum: *Phys.Rev.Lett.* 107, 069901 (2011)], arXiv:0812.4339 [astro-ph].
- [31] E. Bugaev and P. Klimai, *JETP Lett.* **91**, 1 (2010), arXiv:0911.0611 [astro-ph.CO].
- [32] R. Saito and J. Yokoyama, *Prog. Theor. Phys.* **123**, 867 (2010), [Erratum: *Prog.Theor.Phys.* 126, 351–352 (2011)], arXiv:0912.5317 [astro-ph.CO].
- [33] E. Bugaev and P. Klimai, *Phys. Rev. D* **83**, 083521 (2011), arXiv:1012.4697 [astro-ph.CO].
- [34] C. T. Byrnes, P. S. Cole, and S. P. Patil, *JCAP* **06**, 028 (2019), arXiv:1811.11158 [astro-ph.CO].
- [35] S. Clesse, J. García-Bellido, and S. Orani, (2018), arXiv:1812.11011 [astro-ph.CO].
- [36] Y.-F. Cai, C. Chen, X. Tong, D.-G. Wang, and S.-F. Yan, *Phys. Rev. D* **100**, 043518 (2019), arXiv:1902.08187 [astro-ph.CO].
- [37] S. Wang, T. Terada, and K. Kohri, *Phys. Rev. D* **99**, 103531 (2019), [Erratum: *Phys.Rev.D* 101, 069901 (2020)], arXiv:1903.05924 [astro-ph.CO].
- [38] Y. Tada and S. Yokoyama, *Phys. Rev. D* **100**, 023537 (2019), arXiv:1904.10298 [astro-ph.CO].
- [39] C. Yuan, Z.-C. Chen, and Q.-G. Huang, *Phys. Rev. D* **100**, 081301 (2019), arXiv:1906.11549 [astro-ph.CO].
- [40] P. Carrilho, K. A. Malik, and D. J. Mulryne, *Phys. Rev. D* **100**, 103529 (2019), arXiv:1907.05237 [astro-ph.CO].
- [41] Y. Lu, Y. Gong, Z. Yi, and F. Zhang, *JCAP* **12**, 031 (2019), arXiv:1907.11896 [gr-qc].
- [42] N. Bartolo, D. Bertacca, V. De Luca, G. Franciolini, S. Matarrese, M. Peloso, A. Ricciardone, A. Riotto, and G. Tasinato, *JCAP* **02**, 028 (2020), arXiv:1909.12619 [astro-ph.CO].
- [43] S. Bhattacharya, S. Mohanty, and P. Parashari, *Phys. Rev. D* **102**, 043522 (2020), arXiv:1912.01653 [astro-ph.CO].
- [44] R.-G. Cai, Z.-K. Guo, J. Liu, L. Liu, and X.-Y. Yang, *JCAP* **06**, 013 (2020), arXiv:1912.10437 [astro-ph.CO].
- [45] J. Lin, Q. Gao, Y. Gong, Y. Lu, C. Zhang, and F. Zhang, *Phys. Rev. D* **101**, 103515 (2020), arXiv:2001.05909 [gr-qc].
- [46] G. Ballesteros, J. Rey, M. Taoso, and A. Urbano, *JCAP* **07**, 025 (2020), arXiv:2001.08220 [astro-ph.CO].
- [47] K. Inomata, M. Kawasaki, K. Mukaida, T. Terada, and T. T. Yanagida, *Phys. Rev. D* **101**, 123533 (2020), arXiv:2003.10455 [astro-ph.CO].
- [48] J. Fumagalli, S. Renaux-Petel, J. W. Ronayne, and L. T. Witkowski, (2020), arXiv:2004.08369 [hep-th].
- [49] M. Braglia, D. K. Hazra, F. Finelli, G. F. Smoot, L. Sriramkumar, and A. A. Starobinsky, *JCAP* **08**, 001 (2020), arXiv:2005.02895 [astro-ph.CO].
- [50] I. Dalianis and K. Kritos, (2020), arXiv:2007.07915 [astro-ph.CO].
- [51] Z. Yi, Y. Gong, B. Wang, and Z.-h. Zhu, (2020), arXiv:2007.09957 [gr-qc].
- [52] A. M. Green and B. J. Kavanagh, (2020), arXiv:2007.10722 [astro-ph.CO].

- [53] H. Ragavendra, P. Saha, L. Sriramkumar, and J. Silk, (2020), arXiv:2008.12202 [astro-ph.CO].
- [54] N. Bhaumik and R. K. Jain, (2020), arXiv:2009.10424 [astro-ph.CO].
- [55] K. Kohri and T. Terada, (2020), arXiv:2009.11853 [astro-ph.CO].
- [56] Z. Zhou, J. Jiang, Y.-F. Cai, M. Sasaki, and S. Pi, Phys. Rev. D **102**, 103527 (2020), arXiv:2010.03537 [astro-ph.CO].
- [57] G. Domènech and S. Pi, (2020), arXiv:2010.03976 [astro-ph.CO].
- [58] T. Papanikolaou, V. Vennin, and D. Langlois, (2020), arXiv:2010.11573 [astro-ph.CO].
- [59] K. Inomata, M. Kawasaki, K. Mukaida, and T. T. Yanagida, (2020), arXiv:2011.01270 [astro-ph.CO].
- [60] H. Ragavendra, L. Sriramkumar, and J. Silk, (2020), arXiv:2011.09938 [astro-ph.CO].
- [61] Z. Yi, Q. Gao, Y. Gong, and Z.-h. Zhu, (2020), arXiv:2011.10606 [astro-ph.CO].
- [62] Q. Gao, Y. Gong, and Z. Yi, (2020), arXiv:2012.03856 [gr-qc].
- [63] G. Domènech, C. Lin, and M. Sasaki, (2020), arXiv:2012.08151 [gr-qc].
- [64] N. Bartolo, V. Domcke, D. G. Figueroa, J. García-Bellido, M. Peloso, M. Pieroni, A. Ricciardone, M. Sakellariadou, L. Sorbo, and G. Tasinato, JCAP **11**, 034 (2018), arXiv:1806.02819 [astro-ph.CO].
- [65] R.-g. Cai, S. Pi, and M. Sasaki, Phys. Rev. Lett. **122**, 201101 (2019), arXiv:1810.11000 [astro-ph.CO].
- [66] C. Unal, Phys. Rev. D **99**, 041301 (2019), arXiv:1811.09151 [astro-ph.CO].
- [67] A. Ota, Phys. Rev. D **101**, 103511 (2020), arXiv:2001.00409 [astro-ph.CO].
- [68] C. Yuan and Q.-G. Huang, (2020), arXiv:2007.10686 [astro-ph.CO].
- [69] K. N. Ananda, S. Carloni, and P. Dunsby, Phys. Rev. D **77**, 024033 (2008), arXiv:0708.2258 [gr-qc].
- [70] C. Yuan and Q.-G. Huang, (2021), arXiv:2103.04739 [astro-ph.GA].
- [71] K. Inomata, (2020), arXiv:2008.12300 [gr-qc].
- [72] T. H.-C. Lu, K. Ananda, and C. Clarkson, Phys. Rev. D **77**, 043523 (2008), arXiv:0709.1619 [astro-ph].
- [73] T. H.-C. Lu, K. Ananda, C. Clarkson, and R. Maartens, JCAP **02**, 023 (2009), arXiv:0812.1349 [astro-ph].
- [74] D. Yamauchi, T. Namikawa, and A. Taruya, JCAP **10**, 030 (2012), arXiv:1205.2139 [astro-ph.CO].