

PARAMETRIC GEOMETRY OF NUMBERS FOR A GENERAL FLOW

OMRI NISAN SOLAN

ABSTRACT. We extend results on parametric geometry of numbers to a general diagonal flow on the space of lattices. Moreover, we compute the Hausdorff dimension of the set of trajectories with every given behavior, with respect to a nonstandard metric and obtain bounds on the standard ones.

1. INTRODUCTION

Let X_n be the space of rank- n lattices in \mathbb{R}^n with covolume 1, and let $g_t \in \mathrm{SL}_n(\mathbb{R})$ be a 1-parameter diagonal flow. There is a long history of research on the behavior of g_t -orbits in X_n , see [6], [14], [15].

In 2011, Cheung [4] computed the Hausdorff dimension of the set of divergent $g_t = \exp(t \operatorname{diag}(-2, 1, 1))$ -trajectories in X_3 , and proved that the Hausdorff codimension is $2/3$. Later on, many extensions of his result were proved, e.g.,

Date: November 25, 2021.

I thank my advisor Barak Weiss and Eilon Solan for their comments, and acknowledge the support of ISF 2919 BSF 2016256. I wish to thank Andrei Iacob who read the paper and improved the presentation.

Paper	space	flow eigenvalues	codimension	remarks
[4]	X_3	$-2, 1, 1$	$\frac{2}{3}$	
[8]	X_3	$-2, 1, 1$	$\geq \frac{2}{3}$	using entropy method
[5]	X_{n+1}	$-n, 1, \dots, 1$	$\frac{n}{n+1}$	
[12]	X_{n+m}	$-m, \dots, -m, n, \dots, n$	$\geq \frac{nm}{n+m}$	
[17]	X_3	$-1, w_1, w_2;$ $w_1 \leq w_2$ $w_1 + w_2 = 1$	$\frac{1}{1+w_1}$	computed on $\begin{pmatrix} 1 & 0 & 0 \\ * & 1 & 0 \\ * & 0 & 1 \end{pmatrix}$ -orbits
[10]	general G/Γ	General flow	> 0	
[1]	Product space	flow weighted with respect to product	exact result	

In the first four results, all eigenvalues of the flow that are greater than 0 are equal, and so are the eigenvalues smaller than 0.

An independent research direction initiated by Schmidt and Sumner [21] and continued by Roy [18], though solving a seemingly different problem, classified the possible trajectories $g_t\Lambda$ for $\Lambda \in X_n$ up to equivalence. They use the action of

$$(1.1) \quad g_t := \exp(t \operatorname{diag}(n, \dots, n, -m, \dots, -m))$$

on X_{n+m} . We say two trajectories $g_t\Lambda_1, g_t\Lambda_2$ are equivalent if the distance between them is bounded for all $t > 0$.

Das, Fishman, Simmons, and Urbański [7] merged these two research directions and proved a general result computing the Hausdorff dimensions of sets of trajectories closed under equivalence in the setting of the $g_t = \exp(t \operatorname{diag}(n, \dots, n, -m, \dots, -m))$ -action on X_{n+m} .

In this paper we extend the result by Das, Fishman, Simmons, and Urbański [7] to a set of trajectories with respect to a general one-dimensional diagonal action $g_t \curvearrowright X_n$. We compute the dimension of the set of trajectories with any specific behavior, provide exact dimension formulas with respect to a weighted metric, which depends on g_t , and derive lower and upper bounds on the standard Hausdorff dimension.

1.1. Structural Exposition. Section 2 is devoted to the preliminary material that will be needed to formally state the main results of this paper. Sections 4-10 are devoted to Theorem 2.31, the main result of this paper, which determines the Hausdorff dimension of certain

families of trajectories with respect to a certain (non-standard) metric. Section 11 proves Corollary 2.32, which applies Theorem 2.31 to the family of divergent trajectories.

Theorem 2.31 transforms a dynamical question on the space of lattices, into a combinatorial question on combinatorial objects, namely g -templates. In Section 4 we discuss the Hausdorff dimension of sets, and present our computational tool which reduces the computation of the Hausdorff dimension to the dynamical question of computing the value of the Dimension Game introduced in [7]. In Section 5 we discuss properties of g -templates. In Sections 6 and 7 we prove geometric theorems that we will apply in the following three sections. Finally, in Sections 8 and 9 we describe strategies in the dimension game of the two players, and in Section 10 we compute the game's value.

1.1.1. *Mathematical topics.* We use ideas from three different areas: Geometry, Dynamics, and Combinatorics.

- (a) **Geometry** is used to discuss properties of the action of $\mathrm{SL}_n(\mathbb{R})$ and its subgroups g_t and H on the Grassmannian.
- (b) We discuss **dynamical** properties of orbits.
- (c) We construct **combinatorial** objects and then encounter combinatorial problems.

These ideas are not independent: the combinatorial objects are designed with a view to the geometric theorems, we iteratively apply the geometric theorems to obtain orbits with dynamical properties and the construction of each orbit follows a combinatorial object.

1.1.2. *Dependencies.* Although the proofs of the main results depend on intuition acquired in most of the chapters preceding them, in a first reading one can focus only on Sections 2, 4.4, and 5 up to Subsection 5.1, 7, the introduction of Section 8, and Sections 9 – 11.

2. PRELIMINARIES

2.1. **Geometry of Trajectories.** Let $X_n = \mathrm{SL}_n(\mathbb{R})/\mathrm{SL}_n(\mathbb{Z})$ denote the topological space of rank- n lattices in \mathbb{R}^n with determinant 1. Denote the standard right-invariant Riemannian metric on $\mathrm{SL}_n(\mathbb{R})$ by $d_{\mathrm{SL}_n(\mathbb{R})}$. It induces a metric d_{X_n} on X_n .

By right-invariance,

$$(2.1) \quad \forall g \in \mathrm{SL}_n(\mathbb{R}), \quad \Lambda \in X_n. \quad d_{X_n}(g\Lambda, \Lambda) < d_{\mathrm{SL}_n(\mathbb{R})}(g, \mathrm{Id}).$$

Fix a sequence of real weights

$$\eta_1 \leq \eta_2 \leq \cdots \leq \eta_n \text{ with } \eta_1 + \eta_2 + \cdots + \eta_n = 0,$$

for the rest of the paper. Denote $g_t := \exp(t \operatorname{diag}(\eta_1, \dots, \eta_n))$ be a 1-parameter group $\{g_t : t \in \mathbb{R}\} \subseteq \operatorname{SL}_n(\mathbb{R})$. Denote by $E_{\text{all}} := \{\eta_1, \dots, \eta_n\}$ the multiset of all weights. The multiplicity of η_i is the number of indices j such that $\eta_j = \eta_i$.

We say that two trajectories $\{g_t \Lambda : t \geq 0\}$ and $\{g_t \Lambda' : t \geq 0\}$ are equivalent if the distance $d_{X_n}(g_t \Lambda, g_t \Lambda')$ is uniformly bounded for $t > 0$.

2.2. Auxiliary Group Actions. Set

$$\begin{aligned} H &= \left\{ h \in \operatorname{SL}_n(\mathbb{R}) : g_{-t} h g_t \xrightarrow{t \rightarrow \infty} \operatorname{Id} \right\} && \text{– The expanding subgroup.} \\ H^- &= \left\{ h \in \operatorname{SL}_n(\mathbb{R}) : g_t h g_{-t} \xrightarrow{t \rightarrow \infty} \operatorname{Id} \right\} && \text{– The contracting subgroup.} \\ H^0 &= \{ h \in \operatorname{SL}_n(\mathbb{R}) : g_t h g_{-t} = h \} && \text{– The stable subgroup.} \\ H^{-0} &= H^- H^0 && \text{– The nonexpanding subgroup.} \end{aligned}$$

Observation 2.1. Expressing the definitions in coordinates we observe that

$$\begin{aligned} H &= \{ h \in \operatorname{SL}_n(\mathbb{R}) : h_{ij} = \delta_{ij} \text{ if } \eta_i \leq \eta_j \}, \\ H^- &= \{ h \in \operatorname{SL}_n(\mathbb{R}) : h_{ij} = \delta_{ij} \text{ if } \eta_i \geq \eta_j \}, \\ H^0 &= \{ h \in \operatorname{SL}_n(\mathbb{R}) : h_{ij} = 0 \text{ if } \eta_i \neq \eta_j \}, \\ H^{-0} &= \{ h \in \operatorname{SL}_n(\mathbb{R}) : h_{ij} = 0 \text{ if } \eta_i > \eta_j \}. \end{aligned}$$

If $\eta_1 < \dots < \eta_n$, the group H (resp. H^-) is the group of lower (resp. upper) triangular matrices with 1 on the diagonal, and H^0 is the group of diagonal matrices. ♣

Remark 2.2. Here we see the use of ♣ to mark the end of an observation. Likewise, ♡ will mark the end of a definition, ♢ the end of an example, and □ the end of a proof.

Lemma 2.3. *The image of the multiplication map*

$$m_{H^{-0}H} : H \times H^{-0} \rightarrow \operatorname{SL}_n(\mathbb{R}), \quad (h, q^{-0}) \mapsto q^{-0} h$$

is a dense Zariski open set that contains Id.

Proof. Since H and H^{-0} are algebraic groups and their intersection of the complexifications $H_{\mathbb{C}} \cap H_{\mathbb{C}}^{-0} = \{\operatorname{Id}\}$, it follows that $m_{H^{-0}H}$ is one-to-one. Since $\dim H + \dim H^{-0} = \dim \operatorname{SL}_n(\mathbb{R})$, $m_{H^{-0}H}$ is an open map, and hence its image is open. Since $\operatorname{SL}_n(\mathbb{R})$ is irreducible as an algebraic variety, the image is dense. □

Observation 2.4. By Eq. (2.1), for every $q \in H^{-0}$, $\Lambda \in X_n$, the trajectories $g_t\Lambda$ and $g_tq\Lambda$ are equivalent. Hence, studying the behavior of trajectories of lattices in $H^{-0}H\Lambda$ is equivalent to studying the behavior of trajectories of lattices in $H\Lambda$. ♣

The restriction d_H of the metric $d_{\mathrm{SL}_n(\mathbb{R})}$ to H does not suit us, as it has bad behavior with respect to the g_t -action. We use instead the semi-metric d_φ on H defined by

$$d_\varphi(h_1, h_2) := \exp(-\min\{t \in \mathbb{R} : d_H(g_t h_1 g_{-t}, g_t h_2 g_{-t}) \geq 1\}).$$

The semi-metric d_φ satisfies the equality

$$d_\varphi(g_t h_1 g_{-t}, g_t h_2 g_{-t}) = \exp(t) d_\varphi(h_1, h_2),$$

while d_H has no such property even approximately. We will see that d_φ is not a necessarily a metric, but there exists a constant $\alpha_\varphi < 1$ such that $d_\varphi^{\alpha_\varphi}$ is a metric, see Subsections 4.1 and 4.2.

2.3. The Harder-Narasimhan filtration. The Harder-Narasimhan filtration of a lattice was defined in [11] in the context of algebraic geometry, and adapted to lattices by Grayson [9]. Its construction for standard lattices in \mathbb{R}^n goes as follows.

For every lattice $\Gamma \subset \mathbb{R}^n$, denote by $\mathrm{covol} \Gamma$ the Euclidean volume of the torus $\mathrm{span} \Gamma / \Gamma$. By convention, $\mathrm{covol} \{0\} := 1$.

Informally, the Harder-Narasimhan filtration of a lattice Λ is a lattice filtration $\{0\} = \Gamma_{l_0} \subset \Gamma_{l_1} \subset \dots \subset \Gamma_{l_k} = \Lambda$ of “minimal covolume”, where $\mathrm{rank} \Gamma_{l_i} = l_i$. It is the set of all sublattices $\Gamma \subseteq \Lambda$ with exceptionally small covolume; if $\mathrm{rank} \Gamma = l$, even if we know the minimal covolume of all lattices for every possible rank $l' \neq l$, the covolume of Γ is still particularly small.

Formally, to every lattice $\Gamma \subseteq \Lambda$ we associate the point $p_\Gamma := (\mathrm{rank} \Gamma, \log \mathrm{covol} \Gamma) \in \mathbb{R}^2$. For every lattice $\Lambda \in X_n$ define $S_\Lambda := \{p_\Gamma : \Gamma \subseteq \Lambda\}$. Denote the extreme points of the convex hull $\mathrm{conv}(S_\Lambda)$ by $\{p_l : l \in L\}$, where the index l is the x -coordinate of p_l . For each $l \in L$, let $\Gamma_l \subseteq \Lambda$ satisfy $p_{\Gamma_l} = p_l$.

By [9], the lattices $\{\Gamma_l : l \in L\}$ form a filtration of Λ . Furthermore, Γ_l is the unique subgroup




FIGURE 1. An illustration of S_Λ . (Not based on a real lattice.)

that is associated to the point p_l for $l \in L$.

By definition, for every $l \in L$ one has $\text{covol } \Gamma_l < 1$. The filtration $F_{\text{HN}}(\Lambda) := \{\Gamma_l : l \in L\}$ is called the *Harder-Narasimhan filtration*. Note that we index the elements in the filtration by their rank, and the set of ranks is $L = L(\Lambda)$.

2.4. Various Filtration Types. We will often use linear subspaces $V \subseteq \mathbb{R}^n$. The Grassmannian $\text{Gr}(l, n)$ is the space of all l -dimensional subspaces $V \subseteq \mathbb{R}^n$. Denote by $d_{\text{Gr}(l, n)}$ the path metric on $\text{Gr}(l, n)$ derived from the Riemannian metric, which is invariant under multiplication by orthogonal matrices.

The group $\{g_t\}$ acts on $\text{Gr}(l, n)$. The set of g_t -invariant spaces is denoted $\text{Gr}(l, n)^g$. For any $V \in \text{Gr}(l, n)$ the eigenvalues of the action of g_t on V are $\exp(\eta t)$ for a multiset of $\eta \in E_{\text{all}}$. We let \mathcal{I}_l denote the set of multisets of E_{all} of size l .

For every $E \in \mathcal{I}_l$ denote by $\text{Gr}(l, n)_E^g$ the space of $V \in \text{Gr}(l, n)^g$ such that the eigenvalues of $g|_V$ are $\{\exp(t\eta) : \eta \in E\}$.

Example 2.5. Let $n = 4$, $E_{\text{all}} = \{-1, 0, 0, 1\}$, and $l = 2$. Then $\mathcal{I}_2 = \{\{-1, 0\}, \{-1, 1\}, \{0, 1\}, \{0, 0\}\}$, and

$$\begin{aligned} \text{Gr}(2, 4)_{\{-1, 0\}}^g &= \{\text{span}(e_1, v) : 0 \neq v \in \text{span}(e_2, e_3)\}, \\ \text{Gr}(2, 4)_{\{-1, 1\}}^g &= \{\text{span}(e_1, e_4)\}, \\ \text{Gr}(2, 4)_{\{0, 1\}}^g &= \{\text{span}(e_4, v) : 0 \neq v \in \text{span}(e_2, e_3)\}, \\ \text{Gr}(2, 4)_{\{0, 0\}}^g &= \{\text{span}(e_2, e_3)\}. \end{aligned}$$

◇

A *flag* is an increasing sequence of linear spaces,

$$V_\bullet : 0 = V_0 \subset V_{l_1} \subset V_{l_2} \subset \cdots \subset V_{l_k} = \mathbb{R}^n; \quad \dim V_{l_i} = l_i.$$

A *direction filtration* is a sequence

$$(2.2) \quad E_\bullet : \emptyset = E_0 \subset E_{l_1} \subset E_{l_2} \subset \cdots \subset E_{l_k} = E_{\text{all}}; E_{l_i} \in \mathcal{I}_{l_i}.$$

Denote by $L(V_\bullet) = L(E_\bullet) := \{l_0, \dots, l_k\}$.

A *height sequence* $a_\bullet : 0 = a_0, a_1, \dots, a_n = 0$, is a convex sequence of numbers, that is, for every l we have

$$\partial^2 a_l := a_{l-1} - 2a_l + a_{l+1} \geq 0.$$

The *nontrivial places* $1 \leq l \leq n-1$ of a_\bullet are those for which $\partial^2 a_l > 0$. By convention, $l = 0$ and $l = n$ are also nontrivial place. Denote the

set of nontrivial places by $L(a_\bullet)$. Using convexity we can reconstruct a_\bullet from the values $(a_l)_{l \in L(a_\bullet)}$.

A direction filtration with heights is a tuple $F = (F_{E,\bullet}, F_{H,\bullet})$ of a direction filtration and a height sequence where $L(F_{E,\bullet}) = L(F_{H,\bullet})$. The Harder-Narasimhan filtration $F_{\text{HN}}(\Lambda) : \{\Gamma_l : l \in L\}$ of a lattice Λ yields a flag $F_{\text{HN}}(\Lambda)_{V,\bullet} = \text{span } \Gamma_\bullet$, a height sequence defined by $F_{\text{HN}}(\Lambda)_{H,l} = \log \text{covol } \Gamma_l$ for every $l \in L$ and extended piecewise linearly for other l . For consistency, we will write $F_{\text{HN}}(\Lambda)_{\Gamma,l} = \Gamma_l$.

2.5. Dynamics of Linear Spaces. For every $0 \leq l \leq n$ consider the standard L^2 norm $\|\cdot\|$ on $\bigwedge^l \mathbb{R}^n$. Denote by \mathcal{B}_l^n the set of l -blades in $\bigwedge^l \mathbb{R}^n$, that is, the set of vectors $0 \neq v \in \bigwedge^l \mathbb{R}^n$ that can be written as $v = v_1 \wedge v_2 \wedge \cdots \wedge v_l$ with $v_1, v_2, \dots, v_l \in \mathbb{R}^n$. Blades are useful for the following definition:

Definition 2.6. Let $\Gamma \subset \mathbb{R}^n$ be a lattice of rank l . Let v_1, v_2, \dots, v_l be a basis of Γ . Define the *blade* of Γ by

$$\text{Bl } \Gamma := v_1 \wedge v_2 \wedge \cdots \wedge v_l.$$

It is well-defined up to sign. ♡

We identify the real Grassmanian $\text{Gr}(l, n)$, the space of l -dimensional subspaces of \mathbb{R}^n , with $\mathcal{B}_l^n / \mathbb{R}^\times$, via

$$v_1 \wedge \cdots \wedge v_l \mapsto \text{span}(v_1, \dots, v_l).$$

Here \mathbb{R}^\times denotes the multiplicative group of \mathbb{R} . The group $\text{GL}_n(\mathbb{R})$ acts naturally on the spaces $\bigwedge^l \mathbb{R}^n$ and $\text{Gr}(l, n)$.

Recall that the connected components of the g_t -invariant subspace $\text{Gr}(l, n)^g$ are $\text{Gr}(l, n)_E^g$ for every $E \in \mathcal{I}_l$. Define a partial order \trianglelefteq on \mathcal{I}_l by $E_1 \trianglelefteq E_2$ if $\#\{x \in E_1 : x > r\} \leq \#\{x \in E_2 : x > r\}$ for every $r > 0$. In other words, we can obtain E_2 from E_1 by increasing elements. Here $\#$ denotes the cardinality of a set. We write $E_1 \triangleleft E_2$ if $E_1 \trianglelefteq E_2$ and $E_1 \neq E_2$. Thus we have (and will use) two partial orders on multisets: multiset inclusion \subseteq and the new partial order \trianglelefteq .

The following result describes the dynamics of the g_t -action on $\text{Gr}(l, n)$. Its proof is given in Subsection 6.1.

Theorem 2.7. *For every $\varepsilon > 0$, and $1 \leq l \leq n-1$ there is $C > 0$ such that for every $V \in \text{Gr}(l, n)$ there exists a sequence $E_1 \triangleleft \cdots \triangleleft E_r \in \mathcal{I}_l$ and a sequence of intervals $a_1 < b_1 < a_2 < b_2 < \cdots < a_r < b_r$ such that*

$$a_1 = -\infty, \quad b_r = \infty, \quad \sum_{k=1}^{r-1} (a_{k+1} - b_k) < C$$

and for every $t \in (a_k, b_k)$, the space $g_t V$ lies in an ε -neighborhood $U_\varepsilon(\text{Gr}(l, n)_{E_k}^g)$.

Remark 2.8. Note that in Theorem 2.7, if ε is large enough the theorem is vacuous. On the other hand, if

$$\varepsilon < \varepsilon_0 := \min_{1 \leq l \leq n-1} \min_{E_1 \neq E_2 \in \mathcal{I}_l} d_{\text{Gr}(l, n)}(\text{Gr}(l, n)_{E_1}^g, \text{Gr}(l, n)_{E_2}^g)/2,$$

then we have a uniqueness property for the output. That is, if $E_1 \triangleleft \dots \triangleleft E_r \in \mathcal{I}_l$, $a_1 < b_1 < a_2 < b_2 < \dots < a_r < b_r$, and $E'_1 \triangleleft \dots \triangleleft E'_{r'} \in \mathcal{I}_l$, $a'_1 < b'_1 < a'_2 < b'_2 < \dots < a'_{r'} < b'_{r'}$ are two valid outputs of Theorem 2.7, and if $(a_i, b_i) \cap (a'_j, b'_j) \neq \emptyset$ for some $1 \leq i \leq r, 1 \leq j \leq r'$, then $E_i = E'_j$. In particular, if $b_i - a_i > C$ then E_i must be one of the E'_j .

Definition 2.9 (Exterior eigenvalues). For every $E \in \mathcal{I}_l$ denote $\eta_E := \sum_{\eta \in E} \eta$. This sum, as all operations on multisets, counts the elements $\eta \in E$ with multiplicities. The eigenvalue of g_t at an l -blade $w \in \mathcal{B}_l^n$, which corresponds to $V \in \text{Gr}(l, n)_E^g$, is $\exp(t\eta_E)$. \heartsuit

Theorem 2.7 admits the following,

Corollary 2.10. For every $w \in \mathcal{B}_l^n$ there are $-\infty = t_0 < \dots < t_r = \infty$, and $E_1 \triangleleft \dots \triangleleft E_r \in \mathcal{I}_l$ such that $\log \|g_t w\| = \exp(f(t)) + O(1)$, where $f(t)$ is a piecewise linear function with $f'(t) = \eta_{E_k}$ for every $t_{k-1} < t < t_k$.

Example 2.11 (Existence of a blade given a function). Denote by k the number of different values of η in E_{all} , and denote them by $\bar{\eta}_1 < \bar{\eta}_2 < \dots < \bar{\eta}_k$. Let $-\infty = t_0 \leq t_1 \leq \dots \leq t_k = \infty$ and let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a convex piecewise-linear function, such that $df/dt(t) = \eta_r$ for $t_{r-1} < t < t_r$. We will construct a vector $v \in \mathbb{R}^n$ such that $\log \|g_t v\| = f(t) + O(1)$ for every t .

Denote by V_η the $\exp(\eta t)$ -eigenspace of g_t . Fix $v_\eta \in V_\eta$ with $\|v_\eta\| = 1$. Note that for some constants $(a_\eta)_{\eta \in E_{\text{all}}} \subset \{-\infty\} \cup \mathbb{R}$ we have $f(t) = \max_{\eta \in E_{\text{all}}} (t\eta + a_\eta)$. Therefore, the vector $\sum_{\eta \in E_{\text{all}}} \exp(a_\eta) v_\eta$ enjoys the desired property.

If $C > 0$ is large and $t_{r-1} + C \leq t \leq t_r - C$ then the only significant summand in $g_t v$ is $a_{\eta_r} g_t v_{\eta_r}$, and hence $g_t v \cdot \mathbb{R} \in U_\varepsilon(\text{Gr}(1, n)_{\{\eta_r\}}^g)$.

By wedging such vectors we can obtain similar results for spaces and blades with $l > 1$. \diamond

2.6. A g -Template. In this section we describe information that is given in a trajectory $g_t\Lambda$ up to equivalence. Though for most choices of E_{all} the information on the trajectory that we will provide will be equivalent to the equivalence relation on trajectories, sometimes this information will be stronger than the equivalence relation. We will keep track of some of the information provided by the Harder-Narasimhan filtration $F_{\text{HN}}(g_t\Lambda)$, and the direction of its linear spaces.

We wish to encode the “behavior of $F_{\text{HN}}(g_t\Lambda)$ ” for $t \geq 0$. For every $\Gamma \subseteq \Lambda$ of rank l there is a (possibly empty) set $U_\Gamma \subseteq \mathbb{R}$ such that $g_t\Gamma = F_{\text{HN}}(g_t\Lambda)_{\Gamma,l}$ for all $t \in U_\Gamma$. It will follow that U_Γ is approximately an interval.

Inspired by Theorem 2.7, we will encode the behavior of $g_t\Gamma$ for $t \in U_\Gamma$ with multisets $E \in \mathcal{I}_l$. The partially ordered structure on \mathcal{I}_l will not suffice, as $F_{\text{HN}}(g_t\Lambda)$ might not contain a rank- l element, and $F_{\text{HN}}(g_t\Lambda)_{V,l}$ can return to the same $U_\varepsilon(\text{Gr}(l, n)_E^g)$ many times.

It will be convenient to extend \mathcal{I}_l to a category, though we will use only the combinatorial structure of a category and not the whole power of the theory. See Example 2.16 for an equivalent formulation of the concepts without resorting to categories.

Definition 2.12. Let $1 \leq l \leq n - 1$. Define a category \mathcal{I}_l^* whose objects are $\mathcal{I}_l \cup \{*\}$, where $*$ is the null object, and which has two types of arrows.

- *Transition type arrows:* for $a, b \in \mathcal{I}_l$ we have a transition arrow $a \xrightarrow{\triangleleft} b$ if $a \trianglelefteq b$. The composition of transition tow arrows $(b \xrightarrow{\triangleleft} c) \circ (a \xrightarrow{\triangleleft} b)$ is the transition arrow $a \xrightarrow{\triangleleft} c$. For every $a \in \mathcal{I}_l$ the arrow $a \xrightarrow{\triangleleft} a$ is the identity arrow.
- *Null type arrows:* for every $a, b \in \mathcal{I}_l^*$ we have a null arrow $a \xrightarrow{*} b$. The composition of a null arrow with any arrow is a null arrow.

By convention, for $l \in \{0, n\}$ we denote $\mathcal{I}_l^* := \mathcal{I}_l$, the unique arrow is the identity, the single element in \mathcal{I}_l^* is the null element. \heartsuit

The objects of the category \mathcal{I}_l^* will be used to classify the different phases of the l -dimensional spaces in $F_{\text{HN}}(g_t\Lambda)$ for various t . The object $*$ indicates there is no l -dimensional space in the flag, and any multiset E indicates that $F_{\text{HN}}(g_t\Lambda)_{V,l}$ exists and is near $\text{Gr}(l, n)_E^g$. Morphisms in the category \mathcal{I}_l^* classify which phase of the l -dimensional spaces in $F_{\text{HN}}(g_t\Lambda)$ can transform to which one as t increases.

Definition 2.13. Extend each direction filtration E_\bullet by setting $E_l = *$ whenever E_l was not defined previously.

Define the category \mathcal{I}_\bullet^* whose objects are all the direction filtrations E_\bullet , and each arrow between E_\bullet and E'_\bullet is a collection of $n + 1$ arrows $(E_l \rightarrow E'_l)_{l=0, \dots, n}$. We say that the empty filtration $\emptyset \subseteq E_{\text{all}}$ is the null element of \mathcal{I}_\bullet^* , and an arrow $E_\bullet \rightarrow E'_\bullet$ in \mathcal{I}_\bullet^* is null if the induced arrow $E_l \rightarrow E'_l$ is null for every $l = 1, \dots, n - 1$. \heartsuit

Definition 2.14. Let C be a category with a null element. For every interval $I \subseteq \mathbb{R}$, a *category flow* $f : I \rightarrow C$ is a locally constant map $f : I \setminus T \rightarrow C$ for a locally finite set $T \subseteq I$ together with the choice of a non-identity arrow $f_-(t) \rightarrow f_+(t)$ for every point $t \in T$, where $f_\pm(t) = \lim_{\varepsilon \searrow 0} f(t \pm \varepsilon)$. A point $t \in I$ is called a *null point* if

- $t \in T$ and $f_-(t) \rightarrow f_+(t)$ is the null arrow, or
- $f(t)$ is the null object.

All the other points are called *nontrivial points*.

Points in T are called *irregular point*, and the other are called *regular points*. \heartsuit

Observation 2.15. Let $I \subset \mathbb{R}$ be an interval and let $0 \leq l \leq n$. We can restrict every category flow $f_{E_\bullet} : I_\bullet^*$ to its l -th entry and obtain a category flow $f_{E,l} : I \rightarrow \mathcal{I}_l^*$. \clubsuit

Example 2.16. Let $I = (A, B)$ be an open interval. A, B may be $\pm\infty$. A category flow $f : I \rightarrow \mathcal{I}_l^*$ contains the following information:

- A collection of disjoint intervals $((a_i, b_i))_{i \in Z}$ in I , where $Z \subseteq \mathbb{Z}$ is a possibly infinite collection of successive integers, with $b_i \leq a_{i+1}$ for every tuple of consecutive integers $i, i + 1 \in Z$. In addition, the set $\{a_i, b_i : i \in Z\}$ is required to be locally finite.
- A finite set $T_i = \{a_i = c_{i,0} < c_{i,1} < c_{i,2} < \dots < c_{i,r_i} = b_i\}$ for every $i \in Z$
- A sequence of multisets $E_{i,1} \triangleleft E_{i,2} \triangleleft \dots \triangleleft E_{i,r_i}$ for every $i \in Z$.

Denote

$$T := \bigcup_{i \in Z} T_i.$$

Define a category flow by

$$f(t) = \begin{cases} E_{i,j}, & \text{if } c_{i,j-1} < t < c_{i,j}, \\ *, & \text{if } t \in I \setminus \bigcup_{i \in Z} [a_i, b_i], \end{cases}$$

and the connecting morphisms are $E_{i,j-1} \xrightarrow{\triangleleft} E_{i,j}$ at $t = c_{i,j}$ for every $i \in Z, 1 \leq j < r_i$, and the null morphism for $t = a_i, b_i$ for $i \in Z$, provided that $t \in I$ (as opposed to t being an endpoint of I). This is a complete description of a category flow $f : I \rightarrow \mathcal{I}_l^*$. \diamond

We now define our encoding for g_t -trajectories, which are termed g -templates, and play a crucial role in this paper.

Definition 2.17 (g -template). A g -template f over an interval $I \subseteq \mathbb{R}$ is a pair of a category flow $f_{E,\bullet} : I \rightarrow \mathcal{I}_\bullet^*$ and a continuous height sequence-valued map $t \mapsto f_{H,\bullet}(t)$ such that

- (1) For every $t \in I$ that is regular for $f_{E,\bullet}(t)$, the tuple $(f_{E,\bullet}(t), f_{H,\bullet}(t))$ forms a direction filtration with heights.
- (2) For every $1 \leq l < n$ and every $t \in I$ that is regular and not a null point for $f_{E,l}$,

$$(2.3) \quad \eta_{f_{E,l}(t)} = \frac{d}{dt} f_{H,l}(t).$$

Note that each $f_{E,l} : I \rightarrow \mathcal{I}_l$ is a category flow. ♡

Definition 2.18 (Nontriviality intervals). Let f be a g -template on an interval I . Let $U_l(f)$ denote the set of nontrivial points of $f_{E,l}$, and set $U_0(f) = U_n(f) := I$. The set $U_l(f)$ is a disjoint union of open intervals (as every open subset of \mathbb{R}), which will be referred to as the *nontriviality intervals of f at l* .

Define the set of all nontriviality intervals by

$$\mathcal{G}_f := \bigsqcup_{l=0}^n \pi_0(U_l(f)),$$

where $\pi_0(U_l(f))$ is the set of nontriviality intervals of f at l . ♡

Remark 2.19. We have already defined the concepts of nontrivial places of a height sequence, and nontrivial points of a g -template. They match: l is a nontrivial place of $f_{H,\bullet}(t)$ if and only if t is a nontrivial point of $f_{E,l}$.

Example 2.20. In Figure 2 we present a g -template with $E_{\text{all}} = \{-1.2, 0.5, 0.7\}$. The nontriviality intervals are depicted as solid lines and the parts outside them as dashed lines.

Figure 3 shows two cross-sections of f at the three points t_0, t_1 , and t_2 . Note that the value of $f_{H,2}(t_0)$ is defined by $f_{H,1}(t_0)$ and convexity. Similarly, $f_{H,1}(t_0)$ is defined by $f_{H,1}(t_0)$ and convexity. ◇

Definition 2.21 (Equivalence of g -templates). We say that two g -templates f, f' over $I \subseteq \mathbb{R}$ are C -close if

- (1) For every $0 \leq l \leq n$ and $t \in I$ we have $|\partial^2 f_{H,l}(t) - \partial^2 f'_{H,l}(t)| < C$.

FIGURE 2. A g -template.





FIGURE 3. The cross-sections.



- (2) For every subinterval $[a, b] \subseteq I$ of length $|a - b| \geq 2C$ such that $\partial^2 f_{H,l}(t) \geq C$ and $f_{E,l}|_{[a,b]} \equiv E \in \mathcal{I}_l$, we have $f'_{E,l}|_{[a+C,b-C]} \equiv E$. We require also the symmetric claim when swapping f and f' .

The concept of C -closeness defines a metric on the set of g -templates. We say that two g -templates are equivalent if they are C -close for some C . ♥

Remark 2.22. Condition 1 is equivalent to bounding the supremum $|f_{H,l}(t) - f'_{H,l}(t)|$, with a different constant. We define it as in Condition 1 for technical reasons.

g -templates are related to g_t -trajectories by the following definition.

Definition 2.23 (*g-template matching a lattice*). Let ε_0 be as in Remark 2.8, $0 < \varepsilon < \varepsilon_0$, $C > 0$ and let $I \subseteq \mathbb{R}$ be an interval. We say that a lattice $\Lambda \in X_n$ (ε, C)-*matches* a g -template f on I if the following two conditions are satisfied.

- (1) For every $t \in I$ and $1 \leq l \leq n-1$, we have

$$|F_{\text{HN}}(g_t \Lambda)_{H,l} - f_{H,l}(t)| < C.$$

- (2) Let $1 \leq l \leq n-1$ and $[a, b] \subseteq I$ be an interval such that for every $t \in [a, b]$ we have $f_{E,l}(t) = E \in \mathcal{I}_l$ and $\partial^2 f_{H,l}(t) \geq C$. Then for every $t \in [a + C, b - C]$, we have $F_{\text{HN}}(g_t \Lambda)_{V,l} \in U_\varepsilon(\text{Gr}(l, n)_E^g)$.

Note that such a template is unique up to g -template equivalence. If such a template exists for $I = \mathbb{R}$ we denote it by f^Λ . \heartsuit

Observation 2.24. By the uniqueness property observed in Remark 2.8, if Λ (ε, C)-matches f , then for every $\varepsilon' < \varepsilon$ there is $C' > 0$ such that Λ (ε', C')-matches f . \clubsuit

Remark 2.25. In [20] Schmidt describes a weaker form of the g -template, which is similar to the definition of g -templates, but does not remember the multisets. It equivalent to the following definition. A g -system is a height sequence-valued map $f_{H,\bullet} : \mathbb{R} \rightarrow \{\text{height sequences}\}$ such

- (1) For every $0 \leq l \leq n$ the function $t \mapsto f_{H,l}(t)$ is piecewise linear.
(2) For every $1 \leq l \leq n-1, t \in \mathbb{R}$ for which $\partial^2 f_{H,l}(t) > 0$ we have that $f_{H,l}$ is locally convex near t and

$$\frac{d}{dt} f_{H,l}(t) \in \{\eta_E : E \in \mathcal{I}_l\}.$$

if $\frac{d}{dt} f_{H,l}(t)$ is defined.

Here are two examples which show that the multisets conditions of g -templates are necessary for the existence of a lattice that (ε, C)-matches a template.

Counterexample 2.26. Let $n = 4, \eta_1 = -2, \eta_2 = -1, \eta_3 = 1, \eta_4 = 2$. Define Let $f_{H,\bullet} : \mathbb{R} \rightarrow \{\text{height sequences}\}$ be defined by

$$f_{H,\bullet}(t) = \begin{cases} 0, & \text{if } t \leq 0, \\ ta_l, & \text{if } t > 0. \end{cases}$$

where $a_0 = 0, a_1 = -1, a_2 = -1, a_3 = -1/2, a_4 = 0$. Then $f_{H,l}(t)$ is indeed piecewise linear, but there is no Λ that (ε, C)-matches f for every $\varepsilon, C > 0$. Indeed, assuming the contrary there would exist $0 \subset \Gamma_1 \subset \Gamma_2 \subset \Lambda$ of ranks 1 and 2, respectively, such that for every

$t \geq 0$ we have $\log \text{covol } g_t \Gamma_l = f_{H,l}(t) + O(1)$, $l = 1, 2$. Combining Theorem 2.7 and Corollary 8.12 we get that

$$\text{span } g_t \Gamma_1 \xrightarrow{t \rightarrow \infty} \text{Gr}(1, 4)_{\{-1\}}^g \quad \text{and} \quad \text{span } g_t \Gamma_2 \xrightarrow{t \rightarrow \infty} \text{Gr}(2, 4)_{\{-2, 1\}}^g.$$

But this contradicts the fact that no space in $\text{Gr}(2, 4)_{\{-2, 1\}}^g$ contains a space in $\text{Gr}(1, 4)_{\{-1\}}^g$. \diamond

Counterexample 2.27. Let $n = 4$, $\eta_1 = -2, \eta_2 = \eta_3 = -1, \eta_4 = 4$. Let T be arbitrarily large. Let $f_{H,\bullet} : \mathbb{R} \rightarrow \{\text{height sequences}\}$ be defined by

$$f_{H,\bullet}(t) = \begin{cases} 0, & \text{if } |t| \geq T, \\ |T - t|a_l, & \text{if } |t| < T. \end{cases}$$

where $a_0 = 0, a_1 = -1, a_2 = -2, a_3 = -1, a_4 = 0$. Then $f_{H,l}(t)$ is indeed piecewise linear, but there is no Λ that (ε, C) -matches f for every $\varepsilon, C > 0$, and T large enough as a function of C, ε . Indeed, assuming the contrary, there would exist $0 \subset \Gamma_2 \subset \Lambda$ of ranks 2 such that for every t with $|t| \leq T$ we have $\log \text{covol } g_t \Gamma_2 = -2|T - t| + O(C)$. This contradicts Theorem 2.7, as the only multisets $E \in \mathcal{I}_2$ with $\eta_E = \pm 2$ are $\{-1, -1\}$ and $\{-2, 4\}$, and $\{-1, -1\} \not\supseteq \{-2, 4\}$. \diamond

We can now state the main results of this paper.

2.7. Results.

Theorem 2.28 (Existence of Templates). *For every $0 < \varepsilon < \varepsilon_0$ there is a $C > 0$ such that every $\Lambda \in X_n$ (ε, C) -matches a g -template f on \mathbb{R} .*

Remark 2.29. Theorem 2.28 extends the results of [21] from the flow described in Eq. (1.1) to a general diagonal flow. In addition, it sharpens [20, Proposition 2.2].

Definition 2.30. If $E_\bullet \in \mathcal{I}_\bullet^*$ is a direction filtration, denote

$$\delta(E_\bullet) := \sum_{i=1}^k \sum_{\substack{\eta \in E_{l_i} \\ \eta' \in E_{l_i} \setminus E_{l_{i-1}}}} (\eta - \eta')^+,$$

where $a^+ = \max(a, 0)$.

For a g -template f on $[0, \infty)$ define

$$\Delta_0(f) = \liminf_{T \rightarrow \infty} \frac{1}{T} \int_0^T \delta(f_{E,\bullet}(t)) dt,$$

and $\Delta(f) = \sup_{f' \sim f} \Delta_0(f')$, where the supremum is over all f' which are equivalent to f . \heartsuit

Let \mathcal{F} be a set of g -templates that is closed under equivalence and Λ be a lattice. Denote

$$Y_{\Lambda, \mathcal{F}} := \{h \in H : f^{h\Lambda} \in \mathcal{F}\} \subseteq H.$$

The g -template $f^{h\Lambda}$ is defined as in Definition 2.23.

Theorem 2.31. *Let \mathcal{F} be a set of g -templates that is closed under equivalence. Then for every $\Lambda \in X_n$,*

$$\dim_{\mathbf{H}}(Y_{\Lambda, \mathcal{F}}; d_\varphi) = \sup_{f \in \mathcal{F}} \Delta(f) = \sup_{f \in \mathcal{F}} \Delta_0(f),$$

where d_φ is the expansion metric on H with respect to the conjugation by g_t , see Subsection 4.2, and $\dim_{\mathbf{H}}$ is the Hausdorff dimension.

By Corollary 4.8,

$$D := \dim_{\mathbf{H}}(H; d_\varphi) = \delta(\emptyset \subseteq E_{\text{all}}) = \sum_{\eta, \eta' \in E_{\text{all}}} (\eta - \eta')^+.$$

Corollary 2.32. *Let $n \geq 3$. Denote $\Xi := \sum_{\eta \in E_{\text{all}}} \eta^+$. Then*

$$\dim_{\mathbf{H}}(\text{Sing}(H, \Lambda; g_t); d_\varphi) = D - \Xi,$$

where $\text{Sing}(H, \Lambda; g_t)$ is the set of $h \in H$ such that $g_t h \Lambda$ diverges.

Remark 2.33. Corollary 2.32 holds for $n \geq 2$ if we replace divergence with divergence on average. The proof is similar to the proof provided here.

We can pair both results with the Comparison Theorem 4.7. We will construct two functions, $\overline{F} : [0, D] \rightarrow [0, N]$ and $\underline{F} : [0, N] \rightarrow [0, D]$, where $N = \dim H$ is the standard manifold dimension of H .

Corollary 2.34. *Let \mathcal{F} be a set of g -templates that is closed under equivalence. Then for every $\Lambda \in X_n$,*

$$\underline{F}^{-1} \left(\sup_{f \in \mathcal{F}} \Delta_0(f) \right) \leq \dim_{\mathbf{H}}(Y_{\Lambda, \mathcal{F}}; d_H) \leq \overline{F} \left(\sup_{f \in \mathcal{F}} \Delta_0(f) \right),$$

where d_H is standard right-invariant metric on H . In particular, if $n \geq 3$ then

$$\underline{F}^{-1}(D - \Xi) \leq \dim_{\mathbf{H}}(\text{Sing}(H, \Lambda; g_t); d_H) \leq \overline{F}(D - \Xi).$$

The following corollary extends the result of [18] to general flows as in Eq. (1.1). Together with Theorem 2.28 it provides a complete classification of the possible functions $t, i \mapsto F_{\text{HN}}(g_t \Lambda)_{H,i}$ up to a bounded constant for $t \geq 0$ and $0 \leq i \leq n$, and completes the proof of the correction of [20, Conjectures 2.3]. The corollary follows directly from Theorem 2.31, however it does not require the Hausdorff dimension machinery the proof of Theorem 2.31 does, and we will provide a proof that does not require this machinery at Subsection 8.6.

Corollary 2.35. *For every g -template f the set*

$$Y_{\Lambda, f} := \{h \in H : f^{h\Lambda} \sim f\}$$

is non-empty.

2.8. Notations. For every real-valued expression F we will use the notation $O(F)$, $\Theta(F)$, and $\Omega(F)$ to denote expressions that satisfy $|O(F)| < CF$, $cF < \Theta(F) < CF$, $\Omega(F) > cF$ for constants $C, c > 0$ that depend only on E_{all} .

For two real-valued expressions F, G we write $G \ggg F$ if there is an implicit function $\iota : (0, \infty) \rightarrow (0, \infty)$ such that $G > \iota(F)$. This notation will allow us to make statements of the form: “Set $x \ggg y$. Then $x > y^3 + y$ ”. Hence, the function ι is determined implicitly after the expression involving \ggg is stated. Thus, the notation $x \ggg y$ can be used as an abbreviation of an argument of the form “Choose $x > \iota(y)$, where $\iota(y)$ will be determined later implicitly”.

For a metric space (X, d) , a point $x \in X$ and $r > 0$ we will denote the closed ball around x with radius r by $B_d(x; r)$. We will use this notation even if d is not a metric, but just a function $d : X \times X \rightarrow \mathbb{R}$.

3. FURTHER RESEARCH

There are many directions one can hope to further understand, as suggested in [7, Section 6]. We mention here some which arise when considering general 1-parameter flows.

3.1. Comparison to the standard Hausdorff dimension. Corollary 2.34 bounds the standard Hausdorff dimension of trajectory sets in H .

Example 3.1. [17] deals with the case $n = 3$, $\eta_1 = -1$, $\eta_2 = w_1$, $\eta_3 = w_2$, where $w_1 + w_2 \geq 0$. Then $H = \left\{ \begin{pmatrix} 1 & 0 & 0 \\ * & 1 & 0 \\ * & * & 1 \end{pmatrix} \right\}$, $H' = \left\{ \begin{pmatrix} 1 & 0 & 0 \\ * & 1 & 0 \\ * & 0 & 1 \end{pmatrix} \right\}$.

It is shown in [17, Theorem 1.5] that

$$\dim_{\mathbf{H}}(\text{Sing}(H, \Lambda; g_t) \cap H'; d_H) = 2 - \frac{1}{1 + w_1},$$

for every lattice Λ . By [16, Theorem 7.7, p. 104] which glues the result to H -orbits, it follows that

$$\dim_{\mathbf{H}}(\text{Sing}(H, \Lambda; g_t); d_H) \geq 3 - \frac{1}{1 + w_1}.$$

On the other hand, Corollary 2.34 implies that this lower bound is tight. \diamond

Inspired by Example 3.1 we make the following conjecture.

Conjecture 3.2. *Let \mathcal{F} be a set of g -templates, closed under equivalence. Then for every $\Lambda \in X_n$,*

$$\dim_{\mathbf{H}}(Y_{\Lambda, \mathcal{F}}; d_H) = \overline{F} \left(\sup_{f \in \mathcal{F}} \Delta_0(f) \right).$$

A possible heuristic for the conjecture is the computation of the weighted planar Cantor-like set C , which is constructed by the following procedure with appropriate constants. Start with a box, and recursively choose k sub-boxes to every box uniformly at random. The side-lengths of the boxes are weighted, i.e., the l -th side-length of a sub-box is $a_l < 1$ times the l -th side-length of the original box. With probability 1, the standard Hausdorff dimension of C is given by the upper bound provided by the analog of Theorem 4.7, depending on the Hausdorff dimension of C with respect to the weighted metric.

3.2. Dimension in smaller sets. It is common to consider the Hausdorff dimension of trajectory sets in the subgroups $H' \subseteq H$, as in the Dani Correspondence. A natural question is the following.

Question 3.3. *Let \mathcal{F} be a set of g -templates closed under equivalence, and let $\Lambda \in X_n$. Can one provide a combinatorial expression for*

$$\dim_{\mathbf{H}}(Y_{\Lambda, \mathcal{F}} \cap H'; d_\varphi)?$$

In general, one does not expect that the codimension of the intersection in H' will coincide with the codimension of the original set, i.e.,

$\dim_{\mathbf{H}}(H'; d_\varphi) - \dim_{\mathbf{H}}(Y_{\Lambda, \mathcal{F}} \cap H'; d_\varphi) \neq \dim_{\mathbf{H}}(H; d_\varphi) - \dim_{\mathbf{H}}(Y_{\Lambda, \mathcal{F}}; d_\varphi)$, does not hold in general, even when it does not predict a negative dimension. Instead, we expect the dimension to be an integral involving quantities of the form $\dim_{\mathbf{H}}(H' \cap H_{V_\bullet}; d_\varphi)$ for various flags V_\bullet , (H_{V_\bullet} is defined in Definition 7.1).

3.3. Different Lie groups. It seems plausible that the techniques we use here can be generalized to general 1-parameter semisimple flows in a semisimple real Lie group G , acting on G/Γ , where $\Gamma \subset G$ is a lattice. The proof of Lemma 7.16 stands out from the other parts of the paper by being mostly a Lie-theoretic result, and its generalization may require other tools.

Let G be a complex semisimple Lie group, B the Borel subgroup, and $B \subseteq P_1, \dots, P_r \subseteq G$ the maximal parabolic subgroups of G . These groups are all maximal parabolic subgroups of G up to conjugation. Let $(g_t)_{t \in \mathbb{R}}$ be a 1-parameter semisimple flow on G . Define $(G/P)^g := \{p \in G/P : \forall t \in \mathbb{R}, g_t p = p\}$ to be the set of fixed points in G/P for every parabolic subgroup P . In particular, P may be the Borel subgroup B . Let \mathcal{I}_P denote the set of connected components of $(G/P)^g$. Note that the quotient maps $G/B \rightarrow G/P_i$ are g_t -equivariant, and hence define natural maps $\pi_i : \mathcal{I}_B \rightarrow \mathcal{I}_{P_i}$. For every parabolic subgroup P and $E_1, E_2 \in \mathcal{I}_P$, we write $E_1 \trianglelefteq E_2$ if there is $p \in G/P$ such that $\lim_{t \rightarrow -\infty} g_t p \in E_1$ and $\lim_{t \rightarrow \infty} g_t p \in E_2$.

Question 3.4.

- Is \trianglelefteq a partial order of \mathcal{I}_P for every parabolic $P \subseteq G$?
- Let $E_1, E_2 \in \mathcal{I}_B$. Do we have $E_1 \trianglelefteq E_2$ if and only if $\pi_i(E_1) \trianglelefteq \pi_i(E_2)$ for every $1 \leq i \leq r$?

For the relation between parabolic subgroups of G and the cusps of G/Γ , see [2] for the theory and [22] for an application.

4. HAUSDORFF DIMENSIONS

4.1. The Expansion Metric.

Definition 4.1. Let (X, d_X) be a metric space and $\varphi_t : X \rightarrow X$ be a group action $\mathbb{R} \curvearrowright X$. We say that φ is a *boundedly expanding action* if for some $0 < \alpha'_\varphi < \alpha_\varphi$ we have

$$\forall t \in \mathbb{R}, \frac{d}{dt} \log d_X(\varphi_t x_1, \varphi_t x_2) \in [\alpha_\varphi, \alpha'_\varphi],$$

for every $x_1, x_2 \in X$.

From now on we assume that φ is a boundedly expanding action.

Define $d_\varphi : X \times X \rightarrow [0, \infty)$ on X by

$$d_\varphi(x_1, x_2) := \begin{cases} \exp(-\min\{t \in \mathbb{R} : d_X(\varphi_t x_1, \varphi_t x_2) \geq 1\}), & \text{if } x_1 \neq x_2, \\ 0, & \text{if } x_1 = x_2, \end{cases}$$

d_φ is called the *expansion semi-metric*. ♥

Observation 4.2. Note that

$$(4.1) \quad d_\varphi(\varphi_t x_1, \varphi_t x_2) = d_t(x_1, x_2) \cdot \exp(t),$$

and the unit ball with respect to d_φ coincides with the unit ball with respect to d_X . These properties characterize d_φ . \clubsuit

In this subsection we fix a boundedly expanding group action $\mathbb{R} \curvearrowright X$, denoted φ , and denote by $0 < \alpha_\varphi < \alpha'_\varphi$ the corresponding quantities in Definition 4.1.

Lemma 4.3. *If $0 < a \leq \alpha_\varphi$ then d_φ to the power of a , namely, d_φ^a , is a metric.*

Proof. It is clear that d_φ^a is symmetric and that $d_\varphi^a(x, x) = 0$ for every $x \in X$. Since the action φ is boundedly expanding, $d_\varphi^a(x_1, x_2) > 0$ whenever $x_1 \neq x_2$. It remains to show the triangle inequality for three different points $x_1, x_2, x_3 \in X$. Assume $d_0 := d_\varphi(x_1, x_3) \geq d_\varphi(x_1, x_2), d_\varphi(x_2, x_3)$. By Eq. (4.1) w.l.o.g. we may assume that $d_\varphi(x_1, x_3) = 1$. Now note that since $d_\varphi(x_1, x_2) < 1$ we have $s_{12} := -\log(d_\varphi(x_1, x_2)) \geq 0$. The bounded expansion property yields

$$\begin{aligned} -\log d_X(x_1, x_2) &= \log d_X(\varphi_{s_{12}} x_1, \varphi_{s_{12}} x_2) - \log d_X(x_1, x_2) \\ &= \int_0^{s_{12}} \frac{d}{dt} \log d_X(\varphi_t x_1, \varphi_t x_2) dt \geq s_{12} \alpha_\varphi, \end{aligned}$$

and hence $d_X(x_1, x_2) \leq d_\varphi^{\alpha_\varphi}(x_1, x_2) \leq d_\varphi^a(x_1, x_2)$. Similarly, $d_X(x_2, x_3) \leq d_\varphi^a(x_2, x_3)$. Therefore,

$$\begin{aligned} d_\varphi^a(x_1, x_3) = 1 &= d_X(x_1, x_3) \leq d_X(x_1, x_2) + d_X(x_2, x_3) \\ &\leq d_\varphi^a(x_1, x_2) + d_\varphi^a(x_2, x_3), \end{aligned}$$

as desired. \square

Definition 4.4. Recall that for a set $A \subseteq X$ the Hausdorff dimension with respect to the metric d_φ^a , is given by

$$\dim_{\mathbf{H}}(A; d_\varphi^a) := \inf\{d \geq 0 : \mathcal{H}^d(A; d_\varphi^a) = 0\},$$

where

$$\mathcal{H}^d(A; d_\varphi^a) = \inf \left\{ \sum_i r_i^d : \begin{array}{l} A \text{ admits a cover with balls of} \\ \text{radii } r_i > 0 \text{ in the metric } d_\varphi^a \end{array} \right\},$$

is the Hausdorff measure. Since a ball of radius r with respect to $d_\varphi^{a_1}$ is a ball of radius r^{a_2/a_1} with respect to $d_\varphi^{a_2}$ we see that, for every

$$0 < a_1, a_2 < \alpha_\varphi,$$

$$\begin{aligned} \mathcal{H}^d(A; d_\varphi^{a_1}) &= \inf \left\{ \sum_i r_i^d : A \text{ admits a cover with balls of} \right. \\ &\quad \left. \text{radii } r_i > 0 \text{ in the metric } d_g^{a_1} \right\} \\ &= \inf \left\{ \sum_i r_i^{da_1/a_2} : A \text{ admits a cover with balls of} \right. \\ &\quad \left. \text{radii } r_i > 0 \text{ in the metric } d_g^{a_2} \right\} \\ &= \mathcal{H}^{da_1/a_2}(A; d_g^{a_2}), \end{aligned}$$

and hence $\dim_{\mathbf{H}}(A; d_\varphi^{a_1}) = \frac{a_2}{a_1} \dim_{\mathbf{H}}(A; d_\varphi^{a_2})$. Thus, the quantity

$$\dim_{\mathbf{H}}(A; d_\varphi) := a \dim_{\mathbf{H}}(A; d_\varphi^a),$$

which is called *Hausdorff dimension of A with respect to the semi-metric d_φ* , is independent of a . \heartsuit

Example 4.5. The simplest example is provided by $X = \mathbb{R}^n$ and $\varphi_t \vec{x} = e^t \vec{x}$. Then $d_\varphi = d_X$ and the Hausdorff dimension $\dim_{\mathbf{H}}(A; d_\varphi) = \dim_{\mathbf{H}}(A)$ for every $A \subseteq X$. \diamond

4.2. A Case of Interest. Recall that g_t is a diagonal 1-parameter group, and H is the expanding group of g_t , with the right-invariant Riemannian metric d_H . Denote $\varphi_t(h) := g_t h g_{-t}$.

Lemma 4.6. *The action φ_t is boundedly expanding with respect to the metric d_H .*

Proof. The standard Riemannian metric $d_{\text{SL}_n(\mathbb{R})}$ is defined by the bilinear form $\langle A, B \rangle_{\text{Id}} = \text{tr } AB^t$ on $T_{\text{Id}} \text{SL}_n(\mathbb{R}) = \ker \text{tr}$, where $\text{tr} : M_{n \times n}(\mathbb{R}) \rightarrow \mathbb{R}$ is the trace map, and extended to every tangent space $T_p \text{SL}_n(\mathbb{R})$, $p \in \text{SL}_n(\mathbb{R})$ by right translations.

Note that the eigenspaces of the φ_t -action on $T_{\text{Id}}(H)$ are orthogonal with respect to this bilinear form, and that the different eigenvalues are $\exp(t(\eta - \eta'))$, where $\eta > \eta' \in E_{\text{all}}$.

Consequently, for every vector $\vec{v} \in T_{\text{Id}}(H)$ we have

$$\frac{d}{dt} \|(\varphi_t)_* \vec{v}\| / \|\vec{v}\| \in [\alpha_\varphi, \eta_n - \eta_1],$$

where α_φ is the minimal positive distance between elements of E_{all} .

For every point $p \in H$ define the right translation $R_{p^{-1}} : H \rightarrow H$ to be the multiplication from the right by p^{-1} . For every $\vec{v} \in T_p H$ we have

$$\begin{aligned} \|(\varphi_t)_* \vec{v}\| &= \|(R_{E_t(p)^{-1}})_* (\varphi_t)_* \vec{v}\| = \|(\varphi_t)_* (R_{p^{-1}})_* \vec{v}\| \\ &\in [\alpha_\varphi, \eta_n - \eta_1] \cdot \|(R_{p^{-1}})_* \vec{v}\| = [\alpha_\varphi, \eta_n - \eta_1] \cdot \|\vec{v}\| \end{aligned}$$

In particular, φ_t satisfies the bounded expansion property when it is tested on tangent vectors instead of distances. Now the assertion of the lemma follows by the definition of the Riemannian metric. \square

Therefore, there is a constant α_φ such that $d_\varphi^{\alpha_\varphi}$ is a metric. The notations φ , d_φ , and α_φ will be used for the rest of the paper.

4.3. Comparison with the Standard Hausdorff Dimension. Let $0 < \zeta_1 \leq \dots \leq \zeta_N$ be the differences $\{\eta_i - \eta_j > 0 : 1 \leq j \leq i \leq n\}$ in increasing order. Let $\overline{F} : [0, \sum_{i=1}^N \zeta_i] \rightarrow [0, N]$ be the piecewise linear function defined by the values

$$\overline{F} \left(\sum_{i=1}^k \zeta_i \right) = k \quad 0 \leq k \leq N,$$

and

$$\underline{F} : [0, n] \rightarrow \left[0, \sum_{i=1}^N \zeta_i \right]$$

be the piecewise linear function defined by the values

$$\underline{F}(k) = \sum_{i=N-k+1}^N \zeta_i \quad 0 \leq k \leq N.$$

The functions \overline{F} and \underline{F}^{-1} are related. Their graphs are composed of intervals of equal lengths and slopes. In \overline{F} they are ordered from the largest slope to the smallest, and in \underline{F}^{-1} the other way around.

Theorem 4.7 (Comparison Theorem). *Let φ_t the flow on H induced by conjugation with the diagonal flow g_t and d_φ be its expansion semi-metric. Then for any set $X \subseteq H$*

$$(4.2) \quad \dim_{\mathbf{H}}(X; d_H) \leq \overline{F}(\dim_{\mathbf{H}}(X; d_\varphi))$$

and

$$(4.3) \quad \dim_{\mathbf{H}}(X; d_\varphi) \leq \underline{F}(\dim_{\mathbf{H}}(X; d_H)).$$

Corollary 4.8. *We have $\dim_{\mathbf{H}}(H; d_\varphi) = \sum_{\eta, \eta' \in E_{\text{all}}} (\eta - \eta')^+$.*

The proofs of (4.3) and (4.8) are similar, we will cover a ball of one semi-metric with balls of the other.

To prove the theorem we need some preparation. Denote by μ the two sided Haar measure on H .

Lemma 4.9. *There are constants $c_1, c_2 > 1$ such that for every $0 < r_1, r_2 < 1$ and $h_1, h_2 \in H$ such that*

$$B_{d_H}(h_1; r_1) \cap B_{d_\varphi}(h_2; r_2) \neq \emptyset,$$

we have

$$\mu(B_{d_H}(h_1; c_1 r_1) \cap B_{d_\varphi}(h_1; c_2 r_2)) = \Theta \left(\prod_{i=1}^N \min(r_1, r_2^{\zeta_i}) \right).$$

Proof. We will first show the result when $h_1 = h_2 = \text{Id}$. By Observation 2.1, every $h \in H$ has $n^2 - N$ fixed entries and N entries that can vary. Denote by $\kappa : H \rightarrow \mathbb{R}^N$ the map that assigns to each $h \in H$ its N varying coordinates. Denote by $\varphi'_t : \mathbb{R}^N \rightarrow \mathbb{R}^N$ the diagonal action multiplying the i -th coordinate by $\exp(t\zeta_i)$. Note that κ is action-equivariant, that is, the following diagram commutes:

$$\begin{array}{ccc} H & \xrightarrow{\kappa} & \mathbb{R}^N \\ \downarrow \varphi_t & & \downarrow \varphi'_t \\ H & \xrightarrow{\kappa} & \mathbb{R}^N \end{array}$$

Since κ is a manifold diffeomorphism, it distorts the standard metric by a bounded factor on any compact set in H . A direct computation shows that κ maps the Haar measure μ on H into the standard Lebesgue measure λ on \mathbb{R}^N . It follows that for some $0 < \alpha < \alpha'$ we have

$$(4.4) \quad [-\alpha r_1, \alpha r_1]^N \subseteq \kappa(B_{d_H}(\text{Id}; r_1)) \subseteq [-\alpha' r_1, \alpha' r_1]^N.$$

Substituting $r_1 = 1$ and applying $\varphi'_{\log r_2}$ to Eq. (4.4) we obtain,

$$(4.5) \quad \prod_{i=1}^N [-\alpha r_2^{\zeta_i}, \alpha r_2^{\zeta_i}] \subseteq \kappa(B_{d_\varphi}(\text{Id}; r_2)) \subseteq \prod_{i=1}^N [-\alpha' r_2^{\zeta_i}, \alpha' r_2^{\zeta_i}].$$

Combining (4.4) and (4.5) we obtain,

$$(4.6) \quad \mu(B_{d_H}(\text{Id}; r_1) \cap B_{d_\varphi}(\text{Id}; r_2)) = \Theta \left(\prod_{i=1}^N \min(r_1, r_2^{\zeta_i}) \right).$$

Now We turn to the general case. Let $c_1 = 2$, and $c_2 = 2^{1/\alpha_\varphi}$, where α_φ is such that $d_\varphi^{\alpha_\varphi}$ is a metric.

Let $0 < r_1, r_2 < 1$ and $h_1, h_2 \in H$ and assume

$$B_{d_H}(h_1; r_1) \cap B_{d_\varphi}(h_2; r_2) \neq \emptyset.$$

Fix $h_3 \in B_{d_H}(h_1; r_1) \cap B_{d_\varphi}(\text{Id}; r_2)$. Note that

$$B_{d_H}(h_1; 2r_1) \cap B_{d_\varphi}(h_3; 2^{1/\alpha_\varphi} r_2) \supseteq B_{d_H}(h_3; r_1) \cap B_{d_\varphi}(h_3; r_2).$$

The lower bound follows from the right-invariance of d_H and d_φ . The upper bound follows in much the same way, using the fact that

$$B_{d_H}(h_1; 2r_1) \cap B_{d_\varphi}(h_3; 2^{1/\alpha_\varphi} r_2) \subseteq B_{d_H}(h_3; 4r_1) \cap B_{d_\varphi}(h_3; 4^{d_\varphi} r_2).$$

□

For every $d \in \mathbb{R}$ denote

$$\bar{\chi}(d) := \sum_{i=1}^N (\zeta_i - d)^+, \quad \underline{\chi}(d) := \sum_{i=1}^N (1 - d\zeta_i)^+.$$

Lemma 4.10. *Let $r < 1$ and $d \geq 0$. We can cover a d_φ -ball of radius r with $O(r^{-\bar{\chi}(d)})$ d_H -ball of radii r^d .*

Proof. Let $H_1 \subseteq B_{d_\varphi}(\text{Id}; r)$ be a maximal set of points such that $d_H(h_1, h_2) \geq r^d$ for every $h_1, h_2 \in H_1$. Then the balls $B_{d_H}(h_1; r^d/2)$ with $h_1 \in H_1$ are pairwise disjoint and $\{B_{d_H}(h_1; r^d) : h_1 \in H_1\}$ is a cover of $B_{d_\varphi}(\text{Id}; r)$.

Applying Lemma 4.9 to $h_1 \in H_1$, $h_2 = \text{Id}$, $r_1 = r^d/(2c_1)$, $r_2 = r$ we obtain

$$\mu(B_{d_H}(h_1; r^d/2) \cap B_{d_\varphi}(\text{Id}; rc_2)) = \Theta\left(\prod_{i=1}^N \min(r^d, r^{\zeta_i})\right).$$

Similarly,

$$\mu(B_{d_H}(h_1; r^d) \cap B_{d_\varphi}(\text{Id}; rc_2)) = \Theta\left(\prod_{i=1}^N \min(r^d, r^{\zeta_i})\right).$$

The claim follows from the fact that

$$\mu(B_{d_\varphi}(\text{Id}; rc_1)) = \Theta\left(\prod_{i=1}^N r^{\zeta_i}\right).$$

□

The following result and its proof are analogous to Lemma 4.10.

Lemma 4.11. *Let $r < 1$ and $d \geq 0$. We can cover a d_H ball of radius r with $O(r^{-\bar{\chi}(d)})$ balls of radii r^d with respect to the semi-metric d_φ .*

Proof of Theorem 4.7. Let $X \subseteq H$ with $d_0 = \dim_{\mathbf{H}}(X; d_\varphi)$. Let $d > d_0$ and let $(B_{d_\varphi}(h_i; r_i))_{i \in I}$ be a cover of X with d_φ -balls such that $\sum_{i \in I} r_i^d < 1$. We will cover X with d_H -balls. Fix $d' \geq 0$. By Lemma 4.10, we can cover each d_φ -ball of radius r_i with $\Theta(r^{-\bar{\chi}(d')})$ d_H -balls of radius $r^{d'}$.

Consequently, for $d'' \geq 0$, the d'' -Hausdorff measure of X is at most

$$\sum_{i \in I} \Theta(r^{-\bar{\chi}(d')}) \cdot r_i^{d'd''} = \Theta\left(\sum_{i \in I} r^{-\bar{\chi}(d') + d'd''}\right).$$

We will derive a bound on the Hausdorff measure of X only when $-\bar{\chi}(d') + d'd'' \geq d$. Hence we ask ourselves what is the minimal $d'' > 0$ such that for some $d' \geq 0$ we have $-\bar{\chi}(d') + d'd'' \geq d$. Inverting the role of d and d'' , we want to show that for every $d'' \in [0, N]$ we have

$$(4.7) \quad \sup_{d' > 0} d'd'' - \bar{\chi}(d') = \bar{F}^{-1}(d'').$$

Note that $\bar{\chi}$ vanishes on $(-\infty, \zeta_1]$ and hence the supremum in Eq. (4.7) is obtained at $d' \geq \zeta_1 > 0$ and can be taken even for all $d' \in \mathbb{R}$. Hence, it is enough to prove

$$(4.8) \quad \sup_{d' \in \mathbb{R}} d'd'' - \bar{\chi}(d') = \bar{F}^{-1}(d'').$$

In fact, \bar{F} was defined so that \bar{F}^{-1} is the convex conjugate of $\bar{\chi}$, which is stated by Eq. (4.8). Similarly, Eq. (4.3) is a consequence of the fact that \underline{F}^{-1} is the convex conjugate of $\underline{\chi}$. \square

4.4. Dimension Games. Schmidt [19] was the first to prove Hausdorff dimension theorems via (Borel) games with a victory condition.

Das et al. [7] obtained Theorem 2.31 for simpler diagonal flows g_t using a family of zero-sum games with a payoff function (and not a victory condition), which Alice tries to maximize and Bob to minimize. We here describe the game of [7] in our setting, which is termed the (T, g) -game.

Let $T > 0$ and let $X \subseteq H$ be a Borel set. Alice and Bob play a zero-sum alternating-move game, where Alice plays an initialization step, and afterwards Alice and Bob play alternately, first Alice and then Bob.

In the initialization step Alice chooses $h_0 \in H$ and $T_0 > 0$. Along the game, Alice and Bob generate a sequence $B_0 \supset B_1 \supset \dots$, where $B_m = B_{d_\varphi}(h_m; \exp(-T_m))$ and $T_m = Tm + T_0$. Recall that $d_\varphi^{\alpha_\varphi}$ is a metric.

Formally, the game is played as follows.

Initialization step: Alice chooses $h_0 \in H, T_0 > 0$. Set $B_0 := B_{d_\varphi}(h_0; \exp(-T_0))$ and $T_m := Tm + T_0$ for every $m \geq 1$.

Alice's m -th step: Denote $B_m := B_{d_\varphi}(h_m; \exp(-T_m))$. Alice chooses a finite set

$$(4.9) \quad A_{m+1} \subset B_{d_\varphi}(h_m; (1 - \exp(-\alpha_\varphi T))^{1/\alpha_\varphi} \exp(-T_m))$$

such that $d_\varphi(h_{m+1}, h'_{m+1}) > 3^{1/\alpha_\varphi} \exp(-T_m)$ for every two distinct elements $h_{m+1}, h'_{m+1} \in A_{m+1}$.

Bob's m -th step: Bob chooses $h_{m+1} \in A_{m+1}$.

Remark 4.12. The coefficient $(1 - \exp(-\alpha_\varphi T))^{1/\alpha_\varphi}$ in Eq. (4.9) ensures that $B_{m+1} \subseteq B_m$.

Denote

$$h_\infty := \lim_{m \rightarrow \infty} h_m = \bigcap_{m=0}^{\infty} B_m$$

and $A_\bullet = (A_m)_{m=1}^{\infty}$. The payoff which Alice wants to maximize and Bob wants to minimize, is

$$(4.10) \quad \Delta(A_\bullet, h_\infty) := \begin{cases} \frac{1}{T} \liminf_{m \rightarrow \infty} \frac{1}{m} \sum_{k=1}^m \log \#A_k, & \text{if } h_\infty \in X, \\ -\infty, & \text{otherwise.} \end{cases}$$

Denote the value of this game by $D_{T,g}(X)$, i.e., Alice can guarantee any payoff less than $D_{T,g}(X)$, and Bob can deny payoffs higher than $D_{T,g}(X)$.

Recall that a metric space (X, d_X) is called *doubling* if there exists $M > 0$ such that for every $R > 0$ and every point $x \in X$ there exist points $x_1, \dots, x_M \in X$ such that

$$B_{d_X}(x; R) \subseteq \bigcup_{i=1}^M B_{d_X}(x_i; R/2).$$

The following theorem is a special case of [7, Theorem 28.2]. Using [7, Remark 28.3] we may apply it to any complete doubling space. $(H, d_\varphi^{\alpha_\varphi})$ is a doubling space by the compactness of $B_{d_\varphi^{\alpha_\varphi}}(1)$, the right-invariance of d_φ and the behavior under the φ -action. Therefore, we obtain the following result.

Theorem 4.13. *For any Borel set $Y \subseteq H$ we have $\lim_{T \rightarrow \infty} D_{Y,g}(X) = \dim_{\mathbf{H}}(Y; d_\varphi)$.*

4.5. Dimension Game Applied to Trajectory Sets. Let \mathcal{F} be a set of g_t templates closed under equivalence, and $\Lambda \in X_n$ be a lattice. To prove Theorem 2.31, we will apply Theorem 4.13 to the set $Y_{\Lambda, \mathcal{F}} := Y_{\Lambda, \mathcal{F}} \subseteq H$.

Defining the strategies seems complicated, as the resulting h_∞ of a play is known only after the game is played, and hence $f^{h_\infty \Lambda}$ can be computed up to equivalence only at the end of the (T, g) -game. Nonetheless, $f^{h_\infty \Lambda}$ can be approximated.

Observation 4.14. Since $d_\varphi(h_m, h_\infty) \leq \exp(-Tm)$,

$$h_t := g_t h_m h_\infty^{-1} g_{-t}$$

is such that $d_\varphi(h_t, \text{Id}) = O(\exp(t - Tm))$ for every $t \leq Tm$. Using the equality $h_t g_t h_\infty \Lambda = g_t h_m \Lambda$ we deduce that $f^{h_\infty \Lambda}|_{(-\infty, Tm]}$ and $f^{h_m \Lambda}|_{(-\infty, Tm]}$ are uniformly equivalent, where the implicit constants are independent of m and T . In other words, after m steps of Alice and Bob we are able to approximate $f^{h_\infty \Lambda}$ up to time Tm . ♣

The strategies of Alice and Bob will be constructed based on Observation 4.14. In each step the two players will consider $f^{h_\infty \Lambda}$ at time Tm and choose their actions to ensure a certain behavior of $f^{h_\infty \Lambda}$ at time $T(m + 1)$.

5. TEMPLATES APPROXIMATION

In this section we will prove two results. The first is purely combinatorial, and approximates a template with a simpler one that will be used in Alice's strategy. The second constructs a template for every g_t lattice orbit.

Definition 5.1 (Nontrivial places at time t). Let f be a g -template on I . For $t \in I$ denote

$$L_f(t) := \{0 \leq l \leq n : t \in U_l(f)\} = L(f_{H, \bullet}(t)),$$

that is, $L_f(t)$ consists of those l for which $f_{E, l}(t)$ is not null.

♡


Definition 5.2 (Vertices of a g -template). Let f be a g -template.

We term the irregular points t of $f_{E, \bullet}$ *vertices of f* and the irregular points of $f_{E, l}$ *vertices of f at l* . A vertex t is called *null* (resp. *non-null*) if its corresponding arrow by the category flow $f_{E, l}$ is null (resp. non-null) in the category \mathcal{I}_l^* .

Note that the null vertices of f at l are the endpoints of the nontriviality intervals $J \subseteq U_l(f)$. In addition, note that there are no vertices at $l = 0, n$. \heartsuit

Example 2.20, continued. Figure 4 depicts a g -template with $E_{\text{all}} = \{-1.2, 0.5, 0.7\}$. The green vertices are non-null and the blue vertices are null. \diamond

FIGURE 4. A g -template with its vertices.



Definition 5.3 (C -separated g -templates). A g -template f is said to be C -separated for $C > 0$ if for every null vertex t of f there is no other vertex t' of f with $|t - t'| < C$. Moreover, t cannot be a vertex of f at two values of l . \heartsuit

An implication of a template being C -separated is that it cannot change too often, and hence it will be useful in the construction of the strategy of Alice.

Observation 5.4. Let $0 \leq l_{-1} < l_0 < l_1 \leq n$ be three indices and a_\bullet be a height sequence. If a_\bullet is linear on $[l_{-1}, l_0]$ and on $[l_0, l_1]$, then

$$\partial^2 a_l = \frac{1}{l_0 - l_{-1}} a_{l_{-1}} + \frac{1}{l_1 - l_0} a_{l_1} - \frac{l_1 - l_{-1}}{(l_1 - l_0)(l_0 - l_{-1})} a_{l_0}.$$



Corollary 5.5. *Let f be a g -template on an interval I , let $1 \leq l \leq n-1$ and let $t \in I$ be a point such that $\partial^2 f_{H,l}$ is not differentiable at t . Then there exists an index l_0 such that t is a vertex of f at l_0 . Moreover one of the following is true:*

- $l = l_0$.
- l is adjacent to l_0 in $L_f(t)$.

Corollary 5.5 motivates the following definition.

Definition 5.6 (Vertex impact). Let t be a vertex of a g -template f at l_0 . Denote by $l_{-1} < l_0 < l_1$ the adjacent indices of l_0 in $L_f(t)$. We say that the vertex t at l_0 *impacts* the indices l_{-1}, l_0, l_1 . The set of vertices t that impact l contains all nondifferentiability points of $\partial^2 f_{H,l}(t)$, though not necessarily all these vertices are points of nondifferentiability. \heartsuit

Example 2.20, continued. In Figure 4 the vertex t_0 at $l = 1$ impacts 0, 1, 2, while t_1 at $l = 2$ impacts 0, 2, 3. \diamond

Definition 5.7 (C -Significant g -templates). We say that a g -template f is C -significant for $C > 0$ if for every $1 \leq l \leq n-1$ that is impacted by a vertex t of f we have $\partial^2 f_{H,l}(t) > C$ or $\partial^2 f_{H,l}(t) = 0$. \heartsuit


Remark 5.8. Let t be a vertex of a g -template f at l_0 . If t is a null vertex, then $l_0 \notin L_f(t)$ even though this vertex itself impacts l_0 . In this case $\partial^2 f_{H,l_0}(t) = 0$, and hence to check whether f is C -significant or not, one can ignore this impact instance. We still regard this phenomenon as impact in view of Corollary 5.5.

Lemma 5.9. *For every $C > 0$ there exists $C' > 0$ such that for every g -template f on the interval I there exists a C -significant, C -separated g -template f' that is C' -close to f .*

The section will be divided as follows. In Subsection 5.1 we introduce the shifting operation on g -templates. It is an operation that is able to perturb a g -template. In Subsection 5.2 we introduce combinatorial tools that allow to shift a g -template to a significant one. In Subsection 5.3 we discuss some properties of significant templates and complete the proof of Lemma 5.9. In Subsection 5.4 we prove Theorem 2.28.

5.1. Shifting Templates. In this subsection we define the shift operation on templates.

Definition 5.10 (Lower convex hull). Let $X \subset \mathbb{R}$ be a finite set and let $f : X \rightarrow \mathbb{R}$ be function. The *lower convex hull* $f^{\text{conv}} :$

FIGURE 5. The graphs of a_\bullet , ρ , $a_l + \rho_l$ and on it a_l^ρ .

$[\min X, \max X] \rightarrow \mathbb{R}$ is the maximal convex function which satisfies $f^{\text{conv}}(x) \leq f(x)$ for every $x \in X$.

The *lower convex hull* of a sequence a_0, \dots, a_n is the lower convex hull of the function $l \mapsto a_l$, then restricted back to $\{0, \dots, n\}$:

$$((l \mapsto a_l)^{\text{conv}}(l))_{l=0}^n;$$

it is alternatively referred to as *convexification*. ♡

Definition 5.11 (Shift of height sequences). Let a_\bullet be a height sequence and let $\rho = (\rho_l)_{l=0}^n$ be a concave sequence of numbers with $\rho_0 = \rho_n = 0$, i.e., the numbers $0 = \rho_0, \rho_1, \dots, \rho_n = 0$ satisfy $\rho_l \geq (\rho_{l-1} + \rho_{l+1})/2$. Define the *shifted height sequence* a_l^ρ to be the lower convex hull of the sequence $(a_l + \rho_l)_{l=0}^n$; see Example 5.13. ♡

Remark 5.12. The shift operation is continuous in a_\bullet and ρ , and the concavity of ρ implies that the shift depends only on the values of ρ in $L(a_\bullet)$. This yields an explicit formula for a_l^ρ :

$$a_l^\rho = \min_{\substack{l_1 \leq l \leq l_2 \\ \alpha l_1 + (1-\alpha)l_2 = l \\ l_1, l_2 \in L(a_\bullet)}} \alpha(a_{l_1} + \rho_{l_1}) + (1-\alpha)(a_{l_2} + \rho_{l_2})$$

Example 5.13. Figure 5 shows a height sequence a_\bullet , a concave sequence ρ , and the height sequence a_l^ρ . ◇

Definition 5.14 (Shift of g -templates). Let f be a g -template on an interval I . A *shift sequence* is a sequence $\rho = (\rho_l)_{l=0}^n$ of $n+1$ locally constant functions $\rho_l : U_l(f) \rightarrow \mathbb{R}$, such that $\rho_0 \equiv \rho_n \equiv 0$, and for every $0 = l_0 < \dots < l_k = n$ and $t \in U_{l_0}(f) \cap \dots \cap U_{l_k}(f)$ the piecewise linear function defined by $l_i \mapsto \rho_{l_i}(t)$ is concave.

Let ρ be a shift sequence. For every $t \in I$, let $\bar{\rho}(t)$ be any extension of ρ at t to a concave sequence $0 = \bar{\rho}_0(t), \bar{\rho}_1(t), \dots, \bar{\rho}_n(t) = 0$.

Let us define the *shifted g -template* f^ρ . For every t , set

$$f_{H,\bullet}^\rho(t) = (f_{H,\bullet}(t))^{\bar{\rho}(t)};$$

this is independent of the choice of $\bar{\rho}$ and continuous by Remark 5.12.

Denote by $U_l(f^\rho)$ the set of times t for which $l \in \text{supp } f_H^\rho(t)$ is a nontrivial index. This will later coincide with Definition 2.18 of $U_l(f^\rho)$.

Let $f_{E,l}^\rho$ be as follows:

- On $U_l(f^\rho)$ it coincides with $f_{E,l}$.
- Outside of $U_l(f^\rho)$ it is null.
- The connecting morphisms on the boundary $\partial U_l(f^\rho)$ are the null arrows.

One can verify that f^ρ is a g -template.

For future use we note that

$$(5.1) \quad f_{H,l}^\rho := \min_{\substack{l_1 \leq l \leq l_2 \\ \alpha l_1 + (1-\alpha)l_2 = l \\ l_1, l_2 \in L_f(t)}} \alpha(f_{H,l_1}(t) + \rho_{l_1}(t)) + (1-\alpha)(f_{H,l_2}(t) + \rho_{l_2}(t)).$$

♡

Lemma 5.15. *If $(\rho_l)_{l=0}^n$ are bounded by C , then f^ρ is $3C$ -close to f .*

Proof. Fix $t \in I$. To attain $f_H^\rho(t)$, we take $f_H(t)$, increase every value by $\rho_l(t)$ and then take the lower convex hull. It follows that $f_{H,l}(t) \leq f_{H,l}^\rho(t) \leq f_{H,l}(t) + C$, and hence $|\partial^2 f_{H,l}(t) - \partial^2 f_{H,l}^\rho(t)| \leq 2C$. If $[a, b] \subseteq I$ is such that $\partial^2 f|_{[a,b]} \geq 3C$, then $[a, b] \subseteq U_l(f^\rho)$, and hence $f_E|_{[a,b]} \equiv f_E^\rho|_{[a,b]}$. \square

A useful shift sequence is the following.


Definition 5.16 (The independent shift sequence). Let $C > 0$, let f be a g -template on an interval I , and let $\nu_J \in [0, C]$ for $J \in \mathcal{G}_f$, where \mathcal{G}_f is as in Definition 2.18. By convention, the unique nontriviality intervals J of f at 0 and n have $\nu_J = 0$.

The *independent shift sequence* ρ with parameters $C, (\nu_J)_{J \in \mathcal{G}_f}$ is defined by $\rho_l|_J = l(n-l)C + \nu_J$ for every $J \in \pi_0(U_l(f))$. From now on, we will use only independent shift sequences. A useful property of this sequence is that it can change independently on each interval. \heartsuit

Example 2.20, continued. Figures 6 and 7 depict shifts of the g -template in Figure 2.20 and the original g -template in lighter colors behind. We use the independent shift sequence ρ composed of the

constant functions $0, 1, 1, 0$, and 2ρ . Notice that the nontriviality intervals, i.e., the connected components of U_l shrinks in Figure 6, and the interval (t_0, t_1) even vanishes completely in Figure 7, as the maximum of $\partial^2 f_{H,1}|_{(t_0, t_1)}$ is 1.7 , which becomes negative when we add to it $\partial^2 \rho(t) = -2$. The condition $\partial^2 a_l + \partial^2 \rho_l \leq 0$ is sufficient for l to be a trivial place of a_\bullet^ρ , though it is not necessary. We see that the null (blue) vertices moved, and the non-null (green) did not. \diamond

FIGURE 6. A shift of a g -template.




5.2. Constructing a Significant Approximation. In this subsection we will develop a technique which will enable us to solve a large system of a type of linear inequalities, and thus construct a significant approximation and a separated approximation.

The following lemma is a step in the proof of Lemma 5.9

Lemma 5.17. *For every $C > 0$ there exists $C' > 0$ such that for every g -template f on the interval I there exists a C' -significant g -template f' , which is C' -close to f .*

In Subsubsection 5.2.1 we will go over the steps of the construction. In Subsubsection 5.2.2 we will provide proofs for lemmas stated in Subsubsection 5.2.1. Finally, in Subsubsection 5.2.3 we will characterize significant g -templates in a way that will enable us to construct such approximations.

FIGURE 7. A shift of a g -template.

5.2.1. *The Plan.* Our goal is to construct a C -significant template that is equivalent to f . This will be obtained as a shift of f by an independent shift sequence ρ with parameters $C_1, (\nu_J)_{J \in \mathcal{G}_f}$, where $\nu_J \in [0, C_1]$ for every $J \in \mathcal{G}_f$. We will see (Lemma 5.23) that the condition of being C -significant can be strengthened to a condition of the following form:

Definition 5.18. Fix $C_1 > C > 0$, $\varepsilon > 0$, and a “variable set” \mathcal{G} . An *enticement on \mathcal{G}* (with parameters $C_1 > C, \varepsilon$) is an affine function

$$\varphi : \prod_{J \in \mathcal{G}} [0, C_1] \rightarrow \mathbb{R}$$

$$\varphi((\nu_J)_{J \in \mathcal{G}}) = a + \sum_{J \in \mathcal{G}'} a_J \nu_J; \quad a, a_J \in \mathbb{R},$$

where $\mathcal{G}' \subseteq \mathcal{G}$ is a finite subset and $|a_J| \geq \varepsilon$ for every $J \in \mathcal{G}'$. We call \mathcal{G}' the *support* of the enticement φ . We say that a collection of values $(\nu_J)_{J \in \mathcal{G}} \subseteq [0, C_1]$ *satisfies* the enticement φ if

$$\varphi((\nu_J)_{J \in \mathcal{G}}) \notin [0, C].$$

Let Z be an index set. A *solution* of a collection of enticement $\{\varphi_z : z \in Z\}$ is a tuple $(\nu_J)_{J \in \mathcal{G}}$ that satisfies all enticements in the collection.

A collection of enticements $\{\varphi_z : z \in Z\}$ on \mathcal{G} is (r, R) -*finitary* if

- For every $z \in Z$ the support of the enticement φ_z has at most r elements.
- For every $\mathcal{G}' \subseteq \mathcal{G}$ there are at most R enticements φ_z with support \mathcal{G}' , where $z \in Z$.

♡

We will need a collection of enticements whose index set \mathcal{G} has a graph structure. We will now introduce the desired properties of the graph.

Definition 5.19 (Enticement compatibility with a graph). Let \mathcal{G} be a graph. An enticement φ on \mathcal{G} is said to be *compatible with the graph structure on \mathcal{G}* if its support \mathcal{G}' is a clique in \mathcal{G} . From now on all enticements on a graph will be assumed to be compatible with it. ♡

Definition 5.20 (d -constructible graph). A graph \mathcal{G} is said to be *d -constructible* if there is a filtration of full subgraphs $\emptyset = \mathcal{G}_0 \subseteq \mathcal{G}_1 \subseteq \dots \subseteq \mathcal{G}$ such that $\bigcup_m \mathcal{G}_m = \mathcal{G}$ and for every vertex $v \in \mathcal{G}_m \setminus \mathcal{G}_{m-1}$ we have $\deg_{\mathcal{G}_m}(v) \leq d$. ♡

Lemma 5.21. *Let $\{\varphi_z : z \in Z\}$ be a collection of enticements with parameters $C_1 > C, \varepsilon$ on a d -constructible graph \mathcal{G} which are (r, R) -finitary for some $r, R \in \mathbb{Z}_{\geq 0}$. If*

$$C_1 > \varepsilon^{-1} R \sum_{r'=0}^{r-1} \binom{d}{r'} C,$$

then there is a solution for the collection of enticements.

We will define a collection of enticements for each g -template f . We start by defining the underlying graph \mathcal{G}_f as $\mathcal{G}_f = \bigsqcup_{l=1}^{n-1} \pi_0(U_l(f))$ as in Definition 2.18. Two vertices $J_1, J_2 \in \mathcal{G}_f$ are connected by an edge if the intervals intersect. Abusing notation, we do not distinguish between a graph and its set of vertices.

Lemma 5.22. *For every g -template f , the graph \mathcal{G}_f is $(2n-4)$ -constructible.*

Finally we produce a collection of enticements, where the existence of a solution $(\nu_J)_{J \in \mathcal{G}_f}$ guarantees that the g -template f^ρ is C -significant, where ρ is the independent shift sequence with parameters $C, (\nu_J)_{J \in \mathcal{G}_f}$.

Lemma 5.23. *There exists $\varepsilon > 0$ depending on E_{all} such that for every $C_1 > C > 0$ the following holds. For every g -template f there exists a $(4, 49 \cdot 2^{4n} \cdot n^{12})$ -finitary collection of enticements $\{\varphi_z : z \in Z\}$ on \mathcal{G}_f with parameters $C_1 > C, \varepsilon$ such that for every solution $(\nu_J)_{J \in \mathcal{G}_f}$*

of $\{\varphi_z : z \in Z\}$, the g -template f^ρ is C -significant, where ρ is the independent shift sequence with parameters $C, (\nu_J)_{J \in \mathcal{G}_f}$.

Lemma 5.17 is a direct combination of Lemmas 5.23, 5.22 and 5.21. These lemmas also show that Lemma 5.17 holds with any

$$C_1 > \varepsilon^{-1} 49 \cdot 2^{4n} \cdot n^{12} \sum_{r=0}^3 \binom{2n-4}{r} C.$$

5.2.2. Proof of Lemmas 5.21 and 5.22.

Proof of Lemma 5.21. We will choose ν_J one by one. For every m choose a well ordering \prec on $\mathcal{G}_m \setminus \mathcal{G}_{m-1}$ and merge the well orderings into a single well ordering on \mathcal{G} , still denoted \prec , such that $\mathcal{G}_1 \setminus \mathcal{G}_0 \prec \mathcal{G}_2 \setminus \mathcal{G}_1 \prec \dots$. Note that by the degree condition, for every J there are at most d neighbors which are less or equals to J according to \prec .

We will construct the sequence ν_J for every J by transfinite recursion. Fix $J \in \mathcal{G}$ and suppose that we already defined $(\nu_{J'})_{\{J' \in \mathcal{G} : J' \prec J\}}$. Since there are at most d neighbors of J which precede J under \prec , it follows that there are at most $\sum_{r'=0}^{r-1} \binom{d}{r'}$ cliques of size at most r that contain J and are contained in $\{J' \in \mathcal{G} : J' \preceq J\}$. Hence, there are at most $R \sum_{r'=0}^{r-1} \binom{d}{r'}$ enticements whose support is contained in $\{J' \in \mathcal{G} : J' \preceq J\}$ and contains J .

Each of the enticements imposes a linear condition on ν_J of the form $a_J \nu_J + a \notin [0, C]$ for $|a_J| \geq \varepsilon$, which fails to occur on an interval of length at most C/ε . Consequently, if $C_1 > \varepsilon^{-1} C R \sum_{r'=0}^{r-1} \binom{d}{r'}$, then there is a possible value of ν_J that satisfies these enticements. \square

Remark 5.24. Although we used transfinite recursion, when we use this lemma the ordinal will not exceed $n\omega$.

Proof of Lemma 5.22. We will use the fact that no n intervals in \mathcal{G}_f have a common intersection. Denote by A^1 the set of $J \in \mathcal{G}_f$ that are minimal with respect to interval inclusion. Note that $\deg_{\mathcal{G}_f}(J) \leq 2n-4$ for every $J \in A^1$, because

- by the minimality condition, every neighbor must contain one of J 's endpoints, and
- by the intersection condition, at most $n-2$ intervals can contain every endpoint.

Let $\mathcal{G}^1 := \mathcal{G}_f \setminus A^1 \subseteq \mathcal{G}^0 := \mathcal{G}$. Let $A^2 \subseteq \mathcal{G}^1$ be the set of all intervals that are minimal with respect to interval inclusion in \mathcal{G}^1 . As above, for every $J \in A^2$ we have $\deg_{\mathcal{G}^1}(J) \leq 2n-4$.

Continue this process and construct a decreasing sequence of subgraphs $\mathcal{G}^0 \supseteq \mathcal{G}^1 \supseteq \mathcal{G}^2 \supseteq \dots$ such that for every m and every $J \in$

$\mathcal{G}^m \setminus \mathcal{G}^{m+1}$ we have $\deg_{\mathcal{G}^m}(J) \leq 2n - 4$. We have $\mathcal{G}^n = \emptyset$ because the length of the maximal increasing sequence $J_1 \subset J_2 \subset \dots \subset J_k$ of elements in \mathcal{G}^m decreases by 1 as long as \mathcal{G}^m is non-empty. We have thus constructed the desired filtration. \square

5.2.3. *Construction of Enticements.* Throughout this section we consider a tuple of variables $(\nu_J)_{J \in \mathcal{G}_f}$ and the independent shift sequence ρ with parameters $C, (\nu_J)_{J \in \mathcal{G}_f}$. Recall that $\rho_l|_J = l(n-l)C + \nu_J$. For every $0 \leq l \leq n$ define a function $\tilde{f}_{H,l}^\rho : U_l(f) \rightarrow \mathbb{R}$ by

$$\forall t \in U_l(f), \tilde{f}_{H,l}^\rho(t) := f_{H,l}(t) + \rho_l(t).$$

In view of Observation 5.4, we can formulate the following claim.

Claim 5.25. *Let t_0 be a vertex of f^ρ that impacts l_0 , and assume $\partial^2 f_{H,l_0}^\rho(t_0) \neq 0$. Then there exists $l_{-1} < l_0 < l_1 \in L_f(t_0)$, not necessarily adjacent, such that*

$$(5.2) \quad \partial^2 f_{H,l_0}^\rho(t_0) = \frac{1}{l_0 - l_{-1}} \tilde{f}_{H,l_{-1}}^\rho(t_0) + \frac{1}{l_1 - l_0} \tilde{f}_{H,l_1}^\rho(t_0) \\ - \frac{l_1 - l_{-1}}{(l_1 - l_0)(l_0 - l_{-1})} \tilde{f}_{H,l_0}^\rho(t_0).$$

In addition, one of the following holds:

- (1) For some $l \in \{l_0, l_{-1}, l_1\}$ the time t_0 is a non-null vertex of f at l .
- (2) For some $l_{1/2} \in L_f(t_0), l_0 < l_{1/2} < l_1$ the time t_0 is a null vertex of f^ρ , at $l_{1/2}$, and moreover,

$$(5.3) \quad 0 = \frac{1}{l_{1/2} - l_0} \tilde{f}_{H,l_0}^\rho(t_0) + \frac{1}{l_1 - l_{1/2}} \tilde{f}_{H,l_1}^\rho(t_0) \\ - \frac{l_1 - l_0}{(l_1 - l_{1/2})(l_{1/2} - l_0)} \tilde{f}_{H,l_{1/2}}^\rho(t_0).$$

- (3) For some $l_{-1/2} \in L_f(t_0), l_{-1} < l_{-1/2} < l_0$ the time t_0 is a null vertex of f^ρ , at $l_{-1/2}$, and moreover,

$$0 = \frac{1}{l_0 - l_{-1/2}} \tilde{f}_{H,l_0}^\rho(t_0) + \frac{1}{l_{-1/2} - l_{-1}} \tilde{f}_{H,l_{-1}}^\rho(t_0) \\ - \frac{l_0 - l_{-1}}{(l_0 - l_{-1/2})(l_{-1/2} - l_{-1})} \tilde{f}_{H,l_{-1/2}}^\rho(t_0).$$

Proof. Let t_0 be a non-null vertex of f^ρ that impacts l_0 . Choose $l_{-1} < l_0 < l_1$ to be the indices in $L_{f^\rho}(t)$ near l_0 . It follows that $\tilde{f}_{H,l_i}^\rho(t) = f_{H,l_i}^\rho(t)$ for $i = 0, 1, 2$, and using Observation 5.4, Eq. (5.2) holds.

Note that either f^ρ has a vertex t_0 at l_0 , or f_{H,l_0}^ρ is linear near t_0 . If f^ρ has a vertex t_0 at l_0 , then it must be a non-null vertex, as $\partial^2 f_{H,l_1}^\rho(t_0) > 0$, and Case (1) holds.

Otherwise, f_{H,l_0}^ρ is linear near t_0 . Since the vertex t_0 impacts l_0 , it follows that t_0 is a vertex at some $l' \neq l$. Assume $l' > l$. Since $l_1 \in L_{f^\rho}(t)$, it follows that $l' \leq l_1$. We distinguish three cases.

Case $l' = l_1$: The vertex t_0 of f^ρ at l_1 must be a non-null vertex as $l_1 \in L_{t_0}(f^\rho)$, and Case (1) holds.

Case $l' = l_{1/2} < l_1$: Since $l_{1/2} \notin L_{t_0}(f^\rho)$, it follows that t_0 is a null vertex of f^ρ at $l_{1/2}$. Consequently, there is a connected component J of $U_{l_{1/2}}(f^\rho)$ such that t_0 is an endpoint of J .

Therefore, $f_{H,l_{1/2}}^\rho(t_0) = \tilde{f}_{H,l_{1/2}}^\rho(t_0)$, and hence Eq. (5.3) holds. This yields Case (2).

The symmetric alternative $l' < l$ (as opposed to the assumption above $l' > l$), yields Cases (1) and (3). \square

Proof of Lemma 5.23. We will first construct a collection of enticements to ensure each of the Cases (1), (2), (3) of Claim 5.25 fails to give a $\partial^2 f_{H,l_0}^\rho(t_0) \in (0, C]$, for every vertex t_0 that impacts l_0 . We will then verify that it is $(4, 49 \cdot 2^{4n} \cdot n^{12})$ -finitary. To avoid Case (1), note that for every $J \in U_l(f)$ there are at most $l(n-l) \leq n^2$ non-null vertices of f at l in an interval $J \in \pi_0(U_l)$. This is the length of the maximal sequence $E_0 \triangleleft E_1 \triangleleft \dots \triangleleft E_k$.

For every $l_{-1} < l_0 < l_1$, every $J_i \in U_{l_i}(f)$, and every vertex $t_0 \in J_{-1} \cap J_0 \cap J_1$ of f at l_{-1}, l_0 or l_1 , the right-hand side of Eq. (5.2) defines an enticement with support $\{J_i : i = -1, 0, 1 \text{ and } l_i \neq 0, n\}$,

$$\begin{aligned} \varphi((\nu_J)_{J \in \mathcal{G}_f}) := & \frac{f_{H,l_{-1}}(t_0) + l_{-1}(n - l_{-1})C_1 + \nu_{J_{-1}}}{l_0 - l_{-1}} \\ & + \frac{f_{H,l_1}(t_0) + l_1(n - l_1)C_1 + \nu_{J_1}}{l_1 - l_0} \\ & - \frac{l_1 - l_{-1}}{(l_1 - l_0)(l_0 - l_{-1})} (f_{H,l_0}(t_0) + l_0(n - l_0)C_1 + \nu_{J_0}). \end{aligned}$$

In each set of three nontriviality intervals there are at most $3n^2$ vertices, hence there are at most $3n^2$ such enticements. Since some of these intervals may be for $l = 0, n$, we actually obtain at most $6n^2$ such enticements with support $\mathcal{G}' \subseteq \mathcal{G}_f$ with $\#\mathcal{G}' \leq 3$.

Case (2) is more tricky. Fix $l_{-1} < l_0 < l_{1/2} < l_1$, $J_i \in U_{l_i}$ for $i = -1, 0, 1/2, 1$, and linear parts $a_i t + b_i - l(n-l)C_1$ of $f_{H,l_i}(t)$, where $J_{-1} \cap J_0 \cap J_{1/2} \cap J_1 \neq \emptyset$. Since in Case (2) t_0 is defined by Eq. (5.3).

set

$$(5.4) \quad t_0 := - \frac{\frac{1}{l_{1/2}-l_0}(b_0 + \nu_{J_0}) + \frac{1}{l_1-l_{1/2}}(b_1 + \nu_{J_1}) - \frac{l_1-l_0}{(l_1-l_{1/2})(l_{1/2}-l_0)}(b_{1/2} + \nu_{J_{1/2}})}{\frac{1}{l_{1/2}-l_0}a_0 + \frac{1}{l_1-l_{1/2}}a_1 - \frac{l_1-l_0}{(l_1-l_{1/2})(l_{1/2}-l_0)}a_{1/2}}$$

$$= - \frac{(l_1 - l_{1/2})(b_0 + \nu_{J_0}) + (l_{1/2} - l_0)(b_1 + \nu_{J_1}) - (l_1 - l_0)(b_{1/2} + \nu_{J_{1/2}})}{(l_1 - l_{1/2})a_0 + (l_{1/2} - l_0)a_1 - (l_1 - l_0)a_{1/2}},$$

which is the solution of the equation

$$\frac{a_0 t_0 + b_0 + \nu_{J_0}}{l_{1/2} - l_0} + \frac{a_1 t_0 + b_1 + \nu_{J_1}}{l_1 - l_{1/2}} - \frac{(l_1 - l_0)(a_{1/2} t_0 + b_{1/2} + \nu_{J_{1/2}})}{(l_1 - l_{1/2})(l_{1/2} - l_0)} = 0.$$

We need to consider only the case where t_0 is well defined, that is, the denominator in Eq. (5.4) does not vanish.

Substituting Eq. (5.4) in Eq. (5.2) we see that it is enough to show that $\varphi((\nu_J)_{J \in \mathcal{G}_f}) \notin (0, C]$, where

$$\begin{aligned} \varphi((\nu_J)_{J \in \mathcal{G}_f}) &:= \frac{a_{-1}t_0 + b_{-1} + \nu_{J_{-1}}}{l_0 - l_{-1}} + \frac{a_1 t_0 + b_1 + \nu_{J_1}}{l_1 - l_0} \\ &\quad - \frac{(l_1 - l_{-1})(a_0 t_0 + b_0 + \nu_{J_0})}{(l_1 - l_0)(l_0 - l_{-1})} \\ &= \sum_{i=-1,0,1/2,1} c_i(b_i + \nu_{J_i}), \end{aligned}$$

where

$$\begin{aligned} c_{-1} &:= \frac{1}{l_0 - l_{-1}}, \\ c_0 &:= \frac{a_{1/2}l_1 - a_{-1}l_1 - a_1l_{1/2} + a_{-1}l_{1/2} + a_1l_{-1} - a_{1/2}l_{-1}}{(l_1 - l_{1/2})a_0 + (l_{1/2} - l_0)a_1 - (l_1 - l_0)a_{1/2}}(l_1 - l_0), \\ c_{1/2} &:= \frac{-a_1l_0 + a_{-1}l_0 + a_0l_1 - a_{-1}l_1 - a_0l_{-1} + a_1l_{-1}}{(l_1 - l_{1/2})a_0 + (l_{1/2} - l_0)a_1 - (l_1 - l_0)a_{1/2}}(l_1 - l_0), \\ c_1 &:= \frac{\left(\begin{aligned} &-2a_1l_0^2 + a_{1/2}l_0^2 + a_{-1}l_0^2 + 2a_0l_0l_1 - a_{1/2}l_0l_1 - a_{-1}l_0l_1 - a_0l_0l_{1/2} \\ &+ 2a_1l_0l_{1/2} - a_{-1}l_0l_{1/2} - a_0l_1l_{1/2} + a_{-1}l_1l_{1/2} - a_0l_0l_{-1} + 2a_1l_0l_{-1} \\ &- a_{1/2}l_0l_{-1} - a_0l_1l_{-1} + a_{1/2}l_1l_{-1} + 2a_0l_{1/2}l_{-1} - 2a_1l_{1/2}l_{-1} \end{aligned} \right)}{(l_1 - l_{1/2})a_0 + (l_{1/2} - l_0)a_1 - (l_1 - l_0)a_{1/2}}(l_1 - l_0)(l_0 - l_{-1}). \end{aligned}$$

The tricky part is that some of the c_i -s may vanish, and hence the support of φ might not contain all $J_{-1}, J_0, J_{1/2}, J_1$. We will get that φ is an enticement with support $\{J_i : c_i \neq 0 \text{ and } l_i \neq 0, n\}$.

Note that φ is not an enticement if $\{J_i : c_i \neq 0 \text{ and } l_i \neq 0, n\} = \emptyset$. If this happens, then $c_i(b_i + \nu_{J_i}) = 0$ for every $i = -1, 0, 1/2, 1$. Hence, $\varphi = 0$ and will not yield a contradiction to f being C -significant.

Consequently, if $\{J_i : c_i \neq 0 \text{ and } l_i \notin \{0, n\}\} \neq \emptyset$ is non-empty, then φ is an enticement on it. We will see that there are only finitely many possibilities for c_i and hence we may choose ε so that $\varepsilon < |c_i|$ whenever $c_i \neq 0$.

We now count the enticements that we constructed in Case (2) given a support \mathcal{G}' .

Each a_i is the slope of f_{H,l_i} at some $t \in U_{l_i}(f)$ and hence $a_i \in \{\eta_E : E \subseteq E_{\text{all}}\}$, where $i = -1, 0, 1/2, 1$. Consequently, a_i has at most 2^n possibilities. The values of $(c_i)_{i=-1,0,1/2,1}$ depend only on $(l_i)_{i=-1,0,1/2,1}$ and $(a_i)_{i=-1,0,1/2,1}$, which have at most n^4 and $(2^n)^4$ possibilities, respectively. Consequently, the tuple $(c_i)_{i=-1,0,1/2,1}$ has at most $n^4 \cdot 2^{4n}$ possibilities.

For every $i = -1, 0, 1/2, 1$ the value b_i has at most n^2 possibilities, given J_i . For every set $\emptyset \neq \mathcal{G}' \subseteq \mathcal{G}_f$ of size r we need to choose the role of each $J \in \mathcal{G}'$ and there at most 24 possibilities. Altogether we have constructed at most $24 \cdot 2^{4n} \cdot n^{12}$ enticements with support \mathcal{G}' .

Case (3) is analogous to Case (2) and yields $24 \cdot 2^{4n} \cdot n^{12}$ enticements for every $\emptyset \neq \mathcal{G}' \subseteq \mathcal{G}_F$. We thus obtain at most $2 \cdot 24 \cdot 2^{4n} \cdot n^{12} + 2n^6 \leq 49 \cdot 2^{4n} \cdot n^{12}$ enticements.

Since we can consider only enticements on sets of intersecting intervals, the enticements are compatible with the graph. By Lemma 5.21, there is a solution $(\nu_J)_{J \in \mathcal{G}_f}$ to this collection of enticements. Claim 5.25 ensures that if $(\nu_J)_{J \in \mathcal{G}_f}$ satisfy these enticements, then f^ρ is C -significant. \square

5.3. Properties of Significant Templates. We provide useful properties of significant templates, which in particular will allow us to prove Lemma 5.9. Recall that the implicit constants in the notations $O(-)$, $\Omega(-)$, and $\Theta(-)$ depends only on E_{all} (see Subsection 2.8).

Observation 5.26. For every g -template f , and $0 \leq l \leq n$, the function $f_{H,l}$ has at most 2^n possible slopes depending on E_{all} and hence $f_{H,l}$ is $O(1)$ -Lipschitz. Consequently, so is the function $t \mapsto \partial^2 f_{H,l}$. \clubsuit

Lemma 5.27. *For every C -significant g -template f on an interval I and every $0 \leq l \leq n$, every connected component J of $U_l(f)$ that has no common endpoint with I is of length $\Omega(C)$. Moreover, there exists $J' \subset J$ such that $\partial^2 f_{H,l}(t)|_{J'} \geq C$ and the total length of $J \setminus J'$ is $O(C)$.*

Proof. The function $t \mapsto \partial^2 f_{H,l}(t)$ is $O(1)$ -Lipschitz, is positive in J , and vanishes at the boundary of J . Since f is C -significant, the maximum in J must be at least C , and consequently, J is of length $\Omega(C)$.

Moreover, the set $J' := \{t \in J : \partial^2 f_{H,l}(t) \geq C\}$ must be an interval since if it is not connected, then the minimum of $\partial^2 f_{H,l}(t)$ on $(\text{conv } J') \setminus J'$ is less than C and is attained at a vertex that impacts l , which contradicts the C -significance of f .

Since the derivative of $\partial^2 f_{H,l}(t)$ at $J \setminus J'$ is $\Theta(1)$ we deduce the desired result. \square

Lemma 5.28. *For every C -significant g -template f and every interval $[a, b] \subset \mathbb{R}$, there are most $2n^3 + O((b-a)/C)$ vertices of f in $[a, b]$.*

Proof. We will prove that for every l there are at most $2n^2 + O((a-b)/C)$ vertices of f at l in $[a, b]$. Indeed, by Lemma 5.27, the interval I intersects at most $(2 + O((a-b)/C))$ connected components of $U_l(f)$ and each of them contains at most $l \cdot (n-l) + 2 \leq n^2$ vertices. The result follows. \square

Lemma 5.29. *There exists $N = N(E_{\text{all}})$ that satisfies the following property. For every $5C_1$ -significant g -template f , there exists a $(6, N)$ -finitary set of enticements on \mathcal{G}_f such that for any solution $(\nu_J)_{J \in \mathcal{G}_f}$ of it the g -template f^ρ is C -separated, where ρ is the independent shift sequence with parameters $C_1, (\nu_J)_{J \in \mathcal{G}_f}$.*

Proof. Consider a tuple of variables $(\nu_J)_{J \in \mathcal{G}_f}$ and consider the independent shift sequence ρ with parameters $C, (\nu_J)_{J \in \mathcal{G}_f}$. Each null vertex of f^ρ at l_0 is an endpoint of an interval $J_0 \in \pi_0(U_{l_0}(f^\rho))$. As in the proof of Lemma 5.23, t_0 can be defined by an equation

$$(5.5) \quad \frac{1}{l_0 - l_{-1}} \tilde{f}_{H,l_{-1}}^\rho(t_0) + \frac{1}{l_1 - l_0} \tilde{f}_{H,l_1}^\rho(t_0) - \frac{l_1 - l_{-1}}{(l_1 - l_0)(l_0 - l_{-1})} \tilde{f}_{H,l_0}^\rho(t_0) = 0.$$

As in the proof of Lemma 5.17, the value $f_{l_i}(t_0)$ is one of at most n^2 possible linear expression of the form $a_i t_0 + b_0 - l_i(n - l_i)C_1$, and hence

$$t_0 = - \frac{\frac{1}{l_0 - l_{-1}}(b_{-1} + \nu_{-1}) + \frac{1}{l_1 - l_0}(b_1 + \nu_1) - \frac{l_1 - l_{-1}}{(l_1 - l_0)(l_0 - l_{-1})}(b_0 + \nu_0)}{\frac{1}{l_0 - l_{-1}}a_{-1} + \frac{1}{l_1 - l_0}a_1 - \frac{l_1 - l_{-1}}{(l_1 - l_0)(l_0 - l_{-1})}a_0}.$$

If $t_0 \in J_0 \cap J_{-1} \cap J_1$ is a null vertex of f^ρ at l_0 , then it must lie in the closure $\overline{J_0 \setminus J'_0}$, which consists of at most two intervals of length $O(C_1)$, and hence there are at most $O(n^3)$ non-null vertices of f that may be close to t_0 . For every such vertex t' , the condition that t_0 lies at distance less than $O(C)$ from t' is an enticement satisfaction condition, with a support consisting of at most 3 intervals (not exactly 3 as l_{-1} or l_1 may be 0 or n).

Similarly, the condition that two such null vertices lie at distance less than C is an enticement satisfaction condition, with support contained

in the set of the intervals involved. There may be at most 6 intervals, three defining each null vertex.

As in Lemma 5.23, we obtain a finitary set of enticements. \square

Proof of Lemma 5.9. Let f be a g -template. By Lemma 5.17, we may assume that f is $5C_1$ -significant. Let Z be the set of all enticements on \mathcal{G} constructed in Lemmas 5.23 and 5.29. It is $(6, N)$ -finitary with N and ε depending only on E_{all} . By Lemma 5.21 we can solve them with $(\nu_J)_{J \in \mathcal{G}_f} \subseteq [0, C_1]$, provided that C_1 is large enough.

The solution produces an independent shift sequence $\boldsymbol{\rho}$ such that f^ρ is C -significant and C -separated by Lemmas 5.23 and 5.29, and is $3C_1$ -close to f by Lemma 5.15. \square

Lemma 5.30. *Let f be a C -separated and C -significant g -template on an interval I , such that $f_{H,l}(t_0) = 0$ for every $0 \leq l \leq n$ and every endpoint t_0 of I . Then there is an α , $0 < \alpha < 1$, which depends only on E_{all} , such that there is a one-to-one correspondence between non-triviality intervals of f and nontriviality intervals of f^ρ , where $\boldsymbol{\rho}$ is the independent shift sequence with parameters αC , $(\nu_J)_{J \in \mathcal{G}_f} \in [0, \alpha C]^{\mathcal{G}_f}$.*

Moreover, f^ρ is $C/2$ -significant and $C/2$ -separated in a uniform sense in ν_J . That is, for every pair $(\nu_J)_{J \in \mathcal{G}_f}, (\nu'_J)_{J \in \mathcal{G}_f} \in [0, \alpha C]^{\mathcal{G}_f}$ and let $\boldsymbol{\rho}, \boldsymbol{\rho}'$ be the corresponding independent shift sequences. Then for every null vertex t_0 of f and every vertex t'_1 of f , the corresponding vertices of f^ρ , namely $t_0^\rho, t_1^{\rho'}$, satisfy $|t_0^\rho - t_1^{\rho'}| \geq C/2$.

Proof. The first assertion follows from the C -significance of f . The second assertion follows from the first. \square

5.4. Proof of the Template Existence Theorem 2.28. Let Λ be a lattice, and for every $t \in \mathbb{R}$ consider the filtration $F_{\text{HN}}(g_t \Lambda)$. For every $\Gamma \subseteq \Lambda$ of rank l denote by

$$U_\Gamma = \{t \in \mathbb{R} : g_t \Gamma \in F_{\text{HN}}(g_t \Lambda)\}.$$

For many Γ this set is empty, yet for some it is not.

The set U_Γ is not necessarily an interval.

Example 5.31. Let $\Lambda \in X_2$ be a random rank-2 lattice and $\Lambda' = 2\Lambda \oplus \frac{1}{4}\mathbb{Z} \subseteq \mathbb{R}^3$ be a rank-3 lattice. Let $g_t = \exp(t \operatorname{diag}(-1, 1))$ and $g'_t = \exp(t \operatorname{diag}(-1, 1, 0))$ be diagonal flows. For every t we have $\frac{1}{4}\mathbb{Z}e_3 \subseteq g'_t \Lambda' = 2(g_t \Lambda) \oplus \frac{1}{4}\mathbb{Z}$. Since g_t act ergodically on X_2 , it follows that $g_t \Lambda$ equidistributes in X_2 for $t \rightarrow \infty$ with probability one, and hence $\frac{1}{4}\mathbb{Z}e_3$ enters and leaves $F_{\text{HN}}(g'_t \Lambda')$ infinitely many times as $t \rightarrow \infty$. \diamond

For every Γ denote by $\mathfrak{V}_\Gamma := \pi_0(U_\Gamma)$ the set of connected components of U_Γ . Denote $\mathfrak{V} := \bigsqcup_{\Gamma \subseteq \Lambda} \mathfrak{V}_\Gamma$, and call the intervals in \mathfrak{V} *basic intervals*. This union is disjoint so that if a set U appears in \mathfrak{V}_{Γ_1} and \mathfrak{V}_{Γ_2} for two different lattices Γ_1 and Γ_2 , then U appears twice in \mathfrak{V} . For every $U \in \mathfrak{V}$ denote by Γ_U the lattice Γ for which $U \in \mathfrak{V}_\Gamma$.

Let $\varepsilon > 0$ and let $C_0 > 0$ be the corresponding constant provided by Theorem 2.7. We will prove Theorem 2.28 for a tuple (ε, C''') , where C''' will be defined later. It will be sufficient to prove the theorem for ε small enough. Fix $U \in \mathfrak{V}$. By Theorem 2.7, there exists a set of disjoint intervals

$$\mathfrak{U}_U = \{(a_{i,U}, b_{i,U}) : i = 1, \dots, r_U\},$$

in U which cover U except for a set of volume C_0 , and a sequence of multisets

$$E_{1,U} \triangleleft E_{2,U} \triangleleft \dots \triangleleft E_{r_U,U}$$

such that

$$a_{1,U} < b_{1,U} < a_{2,U} < b_{2,U} < \dots < a_{r_U,U} < b_{r_U,U},$$

and $g_t \Gamma_U \in U_\varepsilon \left(\text{Gr}(l, n)_{E_{i,U}}^g \right)$ for every $t \in (a_{i,U}, b_{i,U})$.

Let $U' \subseteq U$ be the interval shorter by C from the right, for some $C > 0$ to be determined later. Define

$$\mathfrak{U}'_U = \{I \cap U' : I \in \mathfrak{U}_U, I \cap U' \neq \emptyset\} := \{(a'_{i,U}, b'_{i,U}) : i = 1, \dots, r'_U\},$$

and denote the corresponding multisets in \mathcal{I}_l by

$$E'_{1,U} \triangleleft E'_{2,U} \triangleleft \dots \triangleleft E'_{r'_U,U}.$$

Denote by a'_0 and $a'_{r'_U+1}$ the endpoints of U' .

Definition 5.32. For every $U \in \mathfrak{V}$ such that $U' \neq \emptyset$, the intervals $[b'_{i,U}, a'_{i+1,U}]$ for $0 \leq i \leq r'_U$ are called *complementary intervals constructed from U* . Let

$$S := \bigcup_{U \in \mathfrak{V}} \bigcup_{i=0}^{r'_U} [b'_{i,U}, a'_{i+1,U}] \subseteq \mathbb{R}$$

be the union of the complementary intervals. ♡

Lemma 5.33. *If $C > (n-1)C_0$, then every connected component of S is of length less than C , and hence does not intersect two different U'_1, U'_2 corresponding to basic intervals $U_1, U_2 \in \mathfrak{V}$ such that $\text{rank } \Gamma_{U_1} = \text{rank } \Gamma_{U_2}$.*

Proof. Let S_0 be a connected component of S . We will show that S_0 does not contain complementary intervals of two different intervals $U_1, U_2 \in \mathfrak{V}$ with $\text{rank } \Gamma_{U_1} = \text{rank } \Gamma_{U_2}$.

Assume, by contradiction, that $I_1 = [a'_1, b'_1], I_2 = [a'_2, b'_2]$ with $b'_1 < a'_2$ are complementary intervals contained in S_0 constructed from basic intervals U_1, U_2 respectively with lattices $\Gamma_{U_1}, \Gamma_{U_2}$ of the same rank. Assume that I_1, I_2 satisfy the following minimality property: there is no such tuple of complementary intervals both intersecting (b'_1, a'_2) .

All complementary intervals that intersect (b'_1, a'_2) consist of at most $n - 1$ basic intervals, corresponding to lattices of different ranks. By Theorem 2.7, every basic interval has complementary intervals of total length at most C_0 . Hence, (b'_1, a'_2) is covered by complementary intervals of total length at most $(n - 1)C_0$. Consequently, $a'_2 - b'_1 \leq (n - 1)C_0$. On the other hand, the distance between U'_1, U'_2 is at least C , and hence $a'_2 - b'_1 > C$. This contradicts the assumption $C > (n - 1)C_0$. \square

Let us define for every $U \in \mathfrak{V}$ an interval $U'' \subseteq U'$. If the left endpoint of U' is not in S , set $U'' := U'$. Otherwise, the left endpoint of U' is in S ; let S_0 be the connected component of S that contains that point and set $U'' := U' \setminus S_0$.

We next define for every l a category flow $\tilde{f}_{E,l} : \mathbb{R} \rightarrow \mathcal{I}_l^*$.

Claim 5.34. *There exists an increasing function $\tau : \mathbb{R} \rightarrow \mathbb{R}$ such that*

- For every $t \in \mathbb{R}$ we have $\tau(t) \notin S$.
- For every $t \in \mathbb{R}$ we have $t - C \leq \tau(t) \leq t$.
- If $t \in U''$, then $\tau(t) \in U''$ for every $U \in \mathfrak{V}$.

Proof. Write S as $S = \bigsqcup_{i \in Z} [a_i, b_i]$, with $Z = [i_0, i_1] \cap \mathbb{Z} \subseteq \mathbb{Z}$ an index set of consecutive numbers such that $a_i < b_i < a_{i+1}$ for every $i_0 \leq i < i_1$ such that $i + 1 \in Z$. Note that i_0 may be $-\infty$ and i_1 may be ∞ . If $i_0 \neq \infty$ set $b_{i_0-1} = -\infty$ and if $i_1 \neq \infty$ set $a_{i_1+1} = \infty$.

Fix $i \in Z$. By Lemma 5.33, $b_i - a_i < C$. For every basic interval U , by the definition of U'' , if $U'' \cap [a_i, b_i] \neq \emptyset$, then $a_i \in U''$. Consequently, we can choose $b_{i-1} < a'_i < a_i$ such that $b_i - a'_i < C$ and $a'_i \in U''$ for every U'' which intersects $[a_i, b_i]$. If $i_1 \neq \infty$, set $a'_{i_1+1} = \infty$.

Note that

$$\bigcup_{i \in [i_0, i_1+1]} (b_{i-1}, a'_i] \cup \bigcup_{i \in Z} (a'_i, b_i] = \mathbb{R}.$$

The function

$$\tau(t) = \begin{cases} t, & \text{if } t \in (b_{i-1}, a'_i] \text{ for } i \in [i_0, i_1 + 1], \\ a'_i, & \text{if } t \in (a'_i, b_i] \text{ for } i \in Z, \end{cases}$$

enjoys the desired properties.

□

Fix a function τ as in Claim 5.34. Define the category flow $\tilde{f}_{E,l} : \mathbb{R} \rightarrow \mathcal{I}_l^*$ as follows: For every $t \in \mathbb{R}$, let U be the unique basic interval with $\text{rank } \Gamma_U = l$ and $\tau(t) \in (a'_{i,U}, b'_{i,U})$ and set $\tilde{f}_{E,l}^l(t) := E'_{i,U}$, where $1 \leq i \leq r'_U$. Define now

$$T_l = \{t \in \mathbb{R} : \tilde{f}_{E,l}^l \text{ is not locally constant at } t\},$$

and $\tilde{f}_{E,l}(t) = \tilde{f}_{E,l}^l(t)$ at every point $t \notin T_l$, and undefined for $t \in T_l$. For every $t \in T_l$ define the connecting morphisms: For every $U \in \mathfrak{V}$ with $\text{rank } \Gamma_U = l$, if $t \in U''$, denote by i_{\pm} the indices such that $\lim_{s \searrow 0} \tilde{f}_{E,l}(t \pm s) = E'_{i_{\pm},U}$. By the monotonicity of τ we get that $i_- < i_+$ and hence $E'_{i_-,U} \triangleleft E'_{i_+,U}$. Therefore, we may choose the connecting morphism of $\tilde{f}_{E,l}$ at t to be $E'_{i_-,U} \xrightarrow{\triangleleft} E'_{i_+,U}$. For

$$t \in T_l \setminus \bigcup_{\substack{U \in \mathfrak{V} \\ \text{rank } \Gamma_U = l}} U'',$$

we define the connecting morphism to be the null arrow.

By Lemma 6.3 below, if $\varepsilon > 0$ is small enough, then $\tilde{f}_{E,\bullet}(t)$ forms a direction filtration.

For every $U \in \mathfrak{V}$, fix $t_U \in U''$, and define $\tilde{f}_{H,l}|_{U''}$ as follows: set $\tilde{f}_{H,l}(t_U) := \log \text{covol } g_{t_U} \Gamma_U$, and extrapolate $\tilde{f}_{H,l}$ to U'' by Eq. (2.3).

Note that for every interval $(a'_{i,U}, b'_{i,U})$, if $a'_{i,U} - b'_{i,U} > C$, then $\tilde{f}_{E,l}|_{(a'_{i,U}+C, b'_{i,U})} \equiv E'_{i,U}$. Corollary 2.10 implies that the blade $\text{Bl } \Gamma$ satisfies

$$(5.6) \quad \left| \tilde{f}_{H,l}(t) - \log \|g_t \text{Bl } \Gamma\| \right| = O(C),$$

for every $t \in U''$.

Let $C'' > 0$, to be determined later. For every $t \in \mathbb{R}$ define $f_{H,\bullet}(t)$ to be the lower convex hull of the function

$$(5.7) \quad l \mapsto \tilde{f}_{H,l}(t) + l(n-l)C'',$$

for every $0 \leq l \leq n$, for which $f_{H,l}(t)$ is defined.

For every $U \in \mathfrak{V}$ with $l_0 := \text{rank } \Gamma_U$ and every endpoint t_0 of U'' which corresponds to an endpoint t_1 of U we have $|t_0 - t_1| \leq C$. Since

$$\partial^2 F_{\text{HN}}(g_t \Lambda)_{H,l_0}(t_1) = 0,$$

the Lipschitz property of $t \mapsto F_{\text{HN}}(g_t \Lambda)_{H,l}(t)$ implies that

$$\partial^2 F_{\text{HN}}(g_t \Lambda)_{H,l_0}(t_0) = O(C).$$

Consequently,

$$\lim_{U'' \ni t \rightarrow t_0} \partial^2 \tilde{f}_{H,l_0}(t) = O(C).$$

Hence if $C'' = O(C)$ is large enough, then

$$\lim_{U'' \ni t \rightarrow t_0} \partial^2(\tilde{f}_{H,l_0}(t) - l(n-l)C'') < 0.$$

Consequently, with this choice of C'' , near t_0 the index l does not correspond to an extreme point of the lower convex hull of the function in Eq. (5.7), hence removing it does not cause discontinuity of the lower convex hull.

From Eq. (5.6) we have $\tilde{f}_{H,l}(t) + l(n-l)C'' \geq F_{\text{HN}}(g_t\Lambda)_{H,l}(t)$ for every $t \in U_l(f)$, provided that $C'' = O(C)$ is sufficiently large compared to C . By the definition of $f_{H,\bullet}$, we get that $f_{H,l}(t) \geq F_{\text{HN}}(g_t\Lambda)_{H,l}(t)$.

By Eq. (5.6), for every $t \in U''$ we have

$$(5.8) \quad f_{H,l}(t) \leq F_{\text{HN}}(g_t\Lambda)_{H,l}(t) + O(C'').$$

Due to the Lipschitz continuity of both functions, we get that Eq. (5.8) holds also for every $t \in U$. Consequently,

$$(5.9) \quad f_{H,l}(t) \geq F_{\text{HN}}(g_t\Lambda)_{H,l}(t)^\rho,$$

where $\rho = (l(n-l)C''')_{l=0}^n$ for some $C''' = O(C'')$, and hence Eq. (5.8) holds for every $t \in \mathbb{R}$.

Define

$$U_l(f) := \{t \in \mathbb{R} : \partial^2 f_{H,l}(t) > 0\},$$

and

$$f_{E,l}(t) := \begin{cases} \tilde{f}_{E,l}(t), & \text{if } t \in U_l(f), \\ *, & \text{if } t \notin U_l(f). \end{cases}$$

The g -template f we defined satisfies the first condition of (ε, C''') -matching (Definition 2.23) for some constant, since (5.8) holds for every $t \in \mathbb{R}$.

To show that the second condition is also satisfied, let $[a, b]$ be an interval on which $f_{E,l}(t) = E$. Then $[a, b] \subseteq U''$ for some $U \in \mathfrak{U}$ with $\text{rank } \Gamma_U = l$. The definition of $\tilde{f}_{E,l}$ implies that $\tau(t) \in (a'_{i,U}, b'_{i,U})$, where $E'_{i,U} = E$. It follows that $a'_{i,U} \leq a$ and $b'_{i,U} \geq b - C'$, and hence $\text{span}_{\mathbb{R}}(\Gamma_U) \in U_\varepsilon(\text{Gr}(l, n)_E^g)$ for every $t \in [a + C', b - C']$.

6. RESULTS ON THE GRASSMANIAN

In this section we will prove Theorem 2.7, which describes the non-recurrent nature of the g_t -action on $\text{Gr}(l, n)$, and Lemma 6.3, which asserts a functorial behavior of the following correspondence with respect to inclusion: a space $V \in U_\varepsilon(\text{Gr}(l, n)_E^g)$ corresponds to $E \in \mathcal{I}_l$. We used both results in Section 5, and we will use them in the following sections as well.

6.1. Proof of Theorem 2.7. It follows from Theorem 2.7 that the g_t -action on $\text{Gr}(l, n)$ is non-recurrent. A simpler proof for the non-recurrence asserts that $g_t \curvearrowright \text{Gr}(l, n)$ is a gradient flow with respect to some function $\rho : \text{Gr}(l, n) \rightarrow \mathbb{R}$. We will not follow this direction, as it is harder to analyze how the flow connects the invariant points.

Instead, we will prove a monotonicity result in a more direct way. First, in case $l = 1$, for every $v \in \mathbb{R}^n$ we want to quantify the fact that along the trajectory $\{g_t v : t \in \mathbb{R}\}$, the coordinates with larger index increase faster.

We will exhibit quadratic forms $Q_s : \mathbb{R}^n \rightarrow \mathbb{R}$ that separate eigenspaces of smaller eigenvalue and eigenspaces of larger eigenvalue. For every fixed $v \in \mathbb{R}^n$, the sign of $Q_s(g_t v)$ will be monotone nondecreasing in t . Then for an l -dimensional subspace $V \subseteq \mathbb{R}^n$ we can compute the signature of each $Q_s|_{g_t V}$ and control its location.

Recall that the η_j -s satisfy $\eta_1 \leq \eta_2 \leq \dots \leq \eta_n$. Denote by k the number of different values of η in E_{all} , and denote these values by $\bar{\eta}_1 < \bar{\eta}_2 < \dots < \bar{\eta}_k$. Set $\bar{\eta}_0 = -\infty$. For every $0 \leq s \leq k$ denote by Q_s^+, Q_s^-, Q_s the quadratic forms on \mathbb{R}^n :

$$Q_s^+ := \sum_{\eta_j > \bar{\eta}_s} x_j^2, \quad Q_s^- := \sum_{\eta_j \leq \bar{\eta}_s} x_j^2, \quad Q_s := Q_s^+ - Q_s^-.$$

An immediate use of this definition is the following monotonicity result.

Lemma 6.1. *For every $s = 0, 1, \dots, k$, and $v \in \mathbb{R}^n \setminus \{0\}$, if $Q_s(v) \geq 0$, then $Q_s(g_t v) > 0$ for every $t > 0$.*

Proof. The result is trivial for $s = 0, k$. Otherwise, $Q_s^+(v)$ must be positive, otherwise $Q_s^+(v) \geq Q_s^-(v) \geq 0$ and then $v = 0$. This yields

$$\begin{aligned} Q_s(g_t v) &= Q_s^+(g_t v) - Q_s^-(g_t v) \\ &\geq \exp(2\bar{\eta}_{s+1}t)Q_s^+(v) - \exp(2\bar{\eta}_s t)Q_s^-(v) \\ &> \exp(2\bar{\eta}_s t)Q_s^+(v) - \exp(2\bar{\eta}_s t)Q_s^-(v) \\ &= \exp(2\bar{\eta}_s t)Q_s(v) \geq 0. \end{aligned}$$

□

We will extend the phenomenon expressed by Lemma 6.1 to l -dimensional subspaces. Let $V \in \text{Gr}(l, n)$. Fix $0 \leq s \leq k$ and denote by $(\sigma_s^0(V), \sigma_s^+(V), \sigma_s^-(V))$ the signature of $Q_s|_V$.

There is a subspace $V_0 \subseteq V$ with $\dim V_0 = \sigma_s^0(V) + \sigma_s^+(V)$ such that $Q_s(v) \geq 0$ for every $v \in V_0$. Lemma 6.1 implies that $Q_s(g_t v) > 0$ for every $t > 0$ and $v \in V_0 \setminus \{0\}$, and hence $\sigma_s^+(g_t V) \geq \sigma_s^0(V) + \sigma_s^+(V)$. It follows that $\sigma_s^+(g_t V)$ is monotone nondecreasing and $\sigma_s^-(g_t V)$ is monotone nonincreasing. Since $\sigma_s^+(g_t V) \leq l$ for every t , it follows that $\sigma_s^0(g_t V) \neq 0$ for at most l values of t .

Note that $\sigma^+(V) \geq r$ is an open condition on V , and hence $\{t \in \mathbb{R} : \sigma_s^+(g_t V) \geq r\}$ is an open set for every fixed V . Similarly, $\{t \in \mathbb{R} : \sigma_s^-(g_t V) \geq r\}$ is open.

For every $E \in \mathcal{I}_l$ define

$$U_E := \left\{ t \in \mathbb{R} : \forall s = 1, \dots, k. \begin{array}{l} \sigma_s^+(g_t V) = \#\{\eta \in E : \eta > \bar{\eta}_s\} \\ \sigma_s^-(g_t V) = \#\{\eta \in E : \eta \leq \bar{\eta}_s\} \end{array} \right\}.$$

This is an open set since we can replace the two equalities in its definition with \geq and preserve the set. The union $\bigcup_{E \in \mathcal{I}_l} U_E$ covers all points $t \in \mathbb{R}$ exactly once, except maybe lk points in which $\sigma_s^0(g_t V) \neq 0$. By the monotonicity condition, each U_E is an interval. By definition, Q_s is nondegenerate on $g_t V$ for every $t \in U_E$.

The following lemma completes the proof of Theorem 2.7.

Lemma 6.2. *For every $\varepsilon > 0$ there exists $t_0 > 0$ such that if $[t - t_0, t + t_0] \subseteq U_E$, then $g_t V$ lies in an ε -neighborhood $U_\varepsilon(\text{Gr}(l, n)_E^g)$.*

Proof. Fix $\varepsilon > 0$ and let $t_0 > 0$ to be determined later. Assume w.l.o.g. that $t = 0$. We will construct inductively a filtration

$$(6.1) \quad \{0\} = V_0^- \subseteq V_1^- \subseteq \dots \subseteq V_k^- = V$$

such that Q_s is negative definite on $g_{t_0} V_s^-$ and $\dim V_s^- = \sigma_s^-(V)$. Let $s = 0, \dots, k-1$ and assume we already defined V_s^- . Since Q_s is negative definite on $g_{t_0} V_s^-$ and $Q_{s+1} \leq Q_s$ pointwise, it follows that Q_{s+1} is negative definite on $g_{t_0} V_s^-$. We will extend the negative definiteness of Q_{s+1} from $g_{t_0} V_s^-$ to $g_{t_0} V_{s+1}^-$ of dimension $\sigma_s^-(V) = \sigma_s^-(g_{t_0} V)$. The orthogonal space W to $g_{t_0} V_s^-$ in $g_{t_0} V$ with respect to Q_{s+1} is such that $Q_{s+1}|_W$ has signature $(0, \sigma_{s+1}^- - \sigma_s^-, \sigma_{s+1}^+)$. Hence, there is $W_0 \subseteq W$ of dimension $\sigma_{s+1}^- - \sigma_s^-$ such that Q_{s+1} is negative definite on W_0 . It follows that Q_{s+1} is negative definite on $g_{t_0} V_s^- \oplus W_0$, and hence we can define $V_{s+1}^- := V_s^- \oplus g_{-t_0} W_0$.

Similarly we can define

$$0 = V_k^+ \subseteq V_{k-1}^+ \subseteq \dots \subseteq V_0^+ = V,$$

with the property that Q_s is positive definite on $g_{-t_0}V_s^-$.

It follows that $V_s^+ \cap V_s^- = 0$. Denote $W_s := V_s^- \cap V_{s-1}^+$ for every $s = 1, \dots, k$. One can show by induction that $\bigoplus_{s=1}^n W_s = V$.

Fix $\delta > 0$. Recall that V_η denotes the $\exp(t\eta)$ -eigenspace of g_t . We will show that we can choose t_0 large enough such that for every $v \in W_{s_0}$ we have $d(v, V_{\bar{\eta}_{s_0}}) \leq \delta \|v\|$. Indeed, denote by

$$v^0 \in V_{\bar{\eta}_{s_0}}, \quad v^+ \in \bigoplus_{s=s_0+1}^k V_{\bar{\eta}_s}, \quad v^- \in \bigoplus_{s=1}^{s_0-1} V_{\bar{\eta}_s},$$

the orthogonal projections of v on the corresponding spaces. Since the three spaces are orthogonal and span \mathbb{R}^n , we have $v = v^0 + v^+ + v^-$. Since $Q_{s-1}(g_{-t_0}v) \geq 0$,

$$\begin{aligned} 0 &\leq \|g_{-t_0}v^+\|^2 + \|g_{-t_0}v^0\|^2 - \|g_{-t_0}v^-\|^2 \\ &\leq \exp(-2t_0\bar{\eta}_{s_0})\|v^+ + v^0\|^2 - \exp(-2t_0\bar{\eta}_{s_0-1})\|v^-\|^2 \\ &\leq \exp(-2t_0\bar{\eta}_{s_0})\|v\|^2 - \exp(-2t_0\bar{\eta}_{s_0-1})\|v^-\|^2. \end{aligned}$$

Consequently, $\|v^-\| \leq \exp(t_0(\bar{\eta}_{s_0-1} - \bar{\eta}_{s_0}))\|v\|$. Similarly, $\|v^+\| \leq \exp(t_0(\bar{\eta}_{s_0} - \bar{\eta}_{s_0+1}))\|v\|$. In particular, if

$$t_0 = - \left(\log \frac{1}{2} \delta \right) / \min(\bar{\eta}_s - \bar{\eta}_{s+1} : s = 1, \dots, k-1),$$

then $\|v^+\|, \|v^-\| \leq \frac{1}{2}\delta\|v\|$, and so $d(v, V_{\bar{\eta}_{s_0}}) \leq \delta\|v\|$.

Finally, for every $\varepsilon > 0$ there exists $\delta > 0$ such that the following holds. For every $E \in \mathcal{I}_l$ which gives multiplicity α_η to η , if $V' \in \text{Gr}(l, n)$ has a decomposition $V' = \bigoplus_{\eta \in E_{\text{all}}} W'_\eta$ with $\dim W'_\eta = \alpha_\eta$ such that for every $v \in W'_\eta$ one has $d(v, V_\eta) \leq \delta\|v\|$, then $V' \in U_\varepsilon(\text{Gr}(l, n)_E^g)$. Indeed, the projection of W'_η to V_η is δ close to W'_η . Hence, the direct sum of the projections is close to V' .

We deduce that $V \in U_\varepsilon(\text{Gr}(l, n)_E^g)$, as desired. \square

6.2. Functoriality of Directions. We end the section by proving that directions are functorial, which was used in the proof of Theorem 2.28, and will be used in the sequel.

Lemma 6.3. *There exists $\varepsilon > 0$ such that if*

- a) $l_1 < l_2$,
- b) $E_i \in \mathcal{I}_{l_i}$ and $V_i \in U_\varepsilon(\text{Gr}(l_i, n)_{E_i}^g)$ for $i = 1, 2$, and
- c) $V_1 \subset V_2$,

then $E_1 \subset E_2$.

Proof. Assume that to the contrary, $E_1 \not\subset E_2$. Denote

$$X := \{(V_1, V_2) \in \text{Gr}(l_1, n) \times \text{Gr}(l_2, n) : V_1 \subset V_2\}.$$

Note that X is closed. By analyzing the eigenvalues of g_t , for every $V_1 \in \text{Gr}(l_1, n)_{E_1}^g$ and $V_2 \in \text{Gr}(l_2, n)_{E_2}^g$ we see that $V_1 \not\subset V_2$. That is,

$$X \cap (\text{Gr}(l_1, n)_{E_1}^g \times \text{Gr}(l_2, n)_{E_2}^g) = \emptyset.$$

Consequently, using the metric

$$d_{\Pi}((V_1, V_2), (V'_1, V'_2)) = \max(d_{\text{Gr}(l_1, n)}(V_1, V'_1), d_{\text{Gr}(l_2, n)}(V_2, V'_2)),$$

we obtain the desired result for $\varepsilon = d_{\Pi}(X, \text{Gr}(l_1, n)_{E_1}^g \times \text{Gr}(l_2, n)_{E_2}^g)$, which is positive since it is the distance between two disjoint compact sets. \square

7. FLAG PERTURBATION THEOREMS

In this section we will prove three results on perturbation of flags, namely Lemmas 7.13, 7.16, and 7.28. These results will be used to define Alice's strategy in Section 8 and the first will be used to analyze Bob's strategy in Section 10.

From a more conceptual viewpoint, Lemma 7.13 is the motivation for the definitions of $\delta(\cdot)$ and $\Delta_0(\cdot)$ (see Definition 2.30), while Lemmas 7.16 and 7.28 are the motivation for the combinatorics of category flows.

7.1. The Counting Lemma.

Definition 7.1. Let $0 \leq l \leq n$, $E \in \mathcal{I}_l$ and $V \in \text{Gr}(l, n)_E^g$. Denote $H_V := \{h \in H : hV = V\}$. Let E_{\bullet} be a direction filtration and V_{\bullet} be a flag such that $L(V_{\bullet}) = L(E_{\bullet})$ and $V_l \in \text{Gr}(l, n)_{E_l}^g$ for every $l \in L(V_{\bullet})$. Denote

$$H_{V_{\bullet}} := \bigcap_{l \in L(V_{\bullet})} H_{V_l}.$$

♡

Recall that for every direction filtration E_{\bullet} we set

$$\delta(E_{\bullet}) = \sum_{i=1}^k \sum_{\substack{\eta \in E_{l_i} \\ \eta' \in E_{l_i} \setminus E_{l_{i-1}}}} (\eta - \eta')^+,$$

and recall the expansion metric d_{φ} on H as defined in Subsection 4.2 and Definition 4.1. This subsection analyzes the subgroup $H_{V_{\bullet}} \subseteq H$ and proves a quantitative claim analogous to the equality

$$\delta(E_{\bullet}) = \dim_{\mathbf{H}}(H_{V_{\bullet}}; d_{\varphi}),$$

where $L(V_\bullet) = L(E_\bullet)$ and $V_l \in \text{Gr}(l, n)_{E_l}^g$ for every $l \in L(V_\bullet)$. Specifically, Lemma 7.2 gives an upper bound on the number of points one can choose in $B_{d_\varphi}(\text{Id}; 1) \cap H_{V_\bullet}$ that are far from one another.

Lemma 7.2. *Let V_\bullet be a flag, and let E_\bullet be a direction filtration with $L(E_\bullet) = L(V_\bullet)$ such that*

$$(7.1) \quad \forall l \in L(V_\bullet), V_l \in \text{Gr}(l, n)_{E_l}^g.$$

Let $t > 0$.

There exists a set $A \subset B_{d_\varphi}(\text{Id}; 1)$ of size $\Theta(\exp(t\delta(E_\bullet)))$ that satisfies the following conditions:

- (1) $d_\varphi(h_1, h_2) \geq \exp(-t)$ for every distinct elements $h_1, h_2 \in A$.
- (2) $hV_l \in \text{Gr}(l, n)_{E_l}^g$ for every $h \in A$, $l \in L(V_\bullet)$, i.e., $A \subseteq H_{V_\bullet}$.

Moreover, there exists no such A larger than $\Theta(\exp(t\delta(E_\bullet)))$ even if we weaken Condition (2) to

- (2') *For every $h \in A$ there exists $h' \in H$ with $d_\varphi(h, h') < \exp(-t)$ and $h'V_l \in \text{Gr}(l, n)_{E_l}^g$ for every $l \in L(V_\bullet)$.*

Fix a flag V_\bullet and a direction filtration E_\bullet . We will express Condition (7.1) in coordinates. Denote by O_g the group of orthogonal matrices that commute with g_t . Since the eigenspaces of g_t are orthogonal, it follows that

$$O_g = \prod_{V \text{ eigenspace of } g_t} O(V),$$

and hence O_g acts transitively on the set of flags which satisfy (7.1). We can thus assume that $V_l = \text{span}\{e_j : j \in F_l\}$, where $F_l \subseteq \{1, \dots, n\}$.

Denote by $(e_i)_{i=1}^n$ the standard basis of \mathbb{R}^n and by $(e_{a,b})_{a,b=1}^n$ the standard basis of $M_{n \times n}(\mathbb{R})$.

Lemma 7.3. *Let $E \in \mathcal{I}_l$ and $V = \text{span}(e_j : j \in F) \in \text{Gr}(l, n)_E^g$ for some $F \subseteq \{1, 2, \dots, n\}$ of size l for which $\{\eta_j : j \in F\} = E$ as multisets, that is, $V \in \text{Gr}(l, n)_E^g$. The group H_V satisfies*

$$\begin{aligned} H_V &= \text{Id} + \text{span}(\{e_{a,b} : \eta_a > \eta_b, b \in F \implies a \in F\}) \\ &= \{h \in H : hV \in \text{Gr}(l, n)_E^g\}. \end{aligned}$$

Proof. The first equality follows from the definition, and similarly

$$H_V \subseteq \{h \in H : hV \in \text{Gr}(l, n)_E^g\}.$$

Assume to the contrary that $hV \in \text{Gr}(l, n)_E^g$, but $hV \neq V$. Since V and hV are g_t -invariant, $g_{-t}h g_t V = hV$. On the other hand, $g_{-t}h g_t \xrightarrow{t \rightarrow \infty} \text{Id}$. The result follows. \square

By Lemma 7.3,

$$\begin{aligned} H_{V_\bullet} &= \bigcap_{l \in L(V_\bullet)} H_{V_l} \\ &= \text{Id} + \text{span}(\{e_{a,b} : \eta_a > \eta_b, \forall l \in L(V_\bullet), b \in F_l \implies a \in F_l\}) \\ &= \{h \in H : \forall l \in L(V_\bullet), hV_l \in \text{Gr}(l, n)\}. \end{aligned}$$

We want to find a large set $A \subseteq B_{d_\varphi}(\text{Id}; 1) \cap H_{V_\bullet}$ whose elements are $\exp(-t)$ far from one another with respect to the semi-metric d_φ . Since d_φ is multiplied by $\exp(t)$ after applying φ_t , we will find a large set $\varphi_t A \subseteq B_{d_\varphi}(\text{Id}; \exp(t))$ whose elements are at distance 1 from one another. We will use a volume argument. Let μ be the bi-invariant Haar measure on H_{V_\bullet} , which exists since H_{V_\bullet} is nilpotent.

Lemma 7.4. *The φ_t -action on the measure satisfies*

$$(\varphi_t)_* \mu = \exp(-t\delta(E_\bullet))\mu.$$

Proof. By the uniqueness of the Haar measure, it follows that $(\varphi_t)_* \mu = \alpha_t \mu$ for some $\alpha_t > 0$. Since H_{V_\bullet} is a Lie group, μ is the integral with respect to an invariant volume form. Hence, to compute the Radon-Nikodym derivative $d(\varphi_t)_* \mu / d\mu \equiv \alpha_t$, it suffices to compute the action of $(\varphi_t)_*$ locally on $\Omega_{\text{Id}}^N H_{V_\bullet}$, where $N = \dim H_{V_\bullet}$. The eigenvalues of $(\varphi_t)_*$ on $T_{\text{Id}} H_{V_\bullet}$ are

$$\exp(t(\eta_a - \eta_b)) \text{ such that } \eta_a > \eta_b \text{ and } \forall l \in L(V_\bullet), b \in F_l \implies a \in F_l.$$

The eigenvalues of the dual action are the reciprocals of these eigenvalues, and hence the eigenvalue of the desired exterior power is the reciprocal of the product. A direct computation shows that the product is $\exp(t\delta(E_\bullet))$, as desired. \square

Proof of Lemma 7.2. Let $\varphi_t A \subseteq B_{d_\varphi}(\text{Id}; \exp(t)) \cap H_{V_\bullet}$ be a maximal set such that the d_φ -distance between any two of its points is at least one. By maximality, every $h \in B_{d_\varphi}(\text{Id}; \exp(t)) \cap H_{V_\bullet}$ is at d_φ -distance at most 1 from A . Consequently,

$$\begin{aligned} (\#A) \cdot \mu(B_{d_\varphi}(\text{Id}; 1)) &\geq \mu(B_{d_\varphi}(\text{Id}; \exp(t))) \\ &= \exp(t\delta(E_\bullet))(\varphi_t)_* \mu(B_{d_\varphi}(\text{Id}; \exp(t))) \\ &= \exp(t\delta(E_\bullet))\mu(\varphi_{-t} B_{d_\varphi}(\text{Id}; \exp(t))) \\ &= \exp(t\delta(E_\bullet))\mu(B_{d_\varphi}(\text{Id}; 1)), \end{aligned}$$

which yields the desired bound on $\#A$.

A similar volume argument provides an upper bound on $\#A$ if A satisfies the weaker Condition (2') instead of (2). Recall that $d_\varphi^{\alpha_\varphi}$ is a

metric (see Subsection 4.2). For every point $h \in \varphi_t A$ there is a point $h' \in H_{V_\bullet} \cap B_{d_\varphi}(h; 1)$. It follows that

$$\mu(B_{d_\varphi}^{\alpha\varphi}(h; 2)) \geq \mu(B_{d_\varphi}^{\alpha\varphi}(h'; 1)) = \mu(B_{d_\varphi}(\text{Id}; 1)).$$

In addition,

$$B_{d_\varphi}^{\alpha\varphi}(h; 2) \cap H_V \subseteq B_{d_\varphi}^{\alpha\varphi}(h'; 3) \cap H_V \subseteq B_{d_\varphi}^{\alpha\varphi}(\text{Id}; \exp(\alpha_\varphi t) + 3).$$

Consequently, the integral

$$\iota := \int_{B_{d_\varphi}^{\alpha\varphi}(\text{Id}; \exp(\alpha_\varphi t) + 3)} \# \left(A \cap B_{d_\varphi}^{\alpha\varphi}(h_1; 2) \right) d\mu(h_1)$$

counts with weight of at least $\mu(B_{d_\varphi}^{\alpha\varphi}(\text{Id}; 1))$ every point in A , and hence $\iota \geq \mu(B_{d_\varphi}^{\alpha\varphi}(\text{Id}; 1)) \# A$. On the other hand, since $d_\varphi^{\alpha\varphi}$ is a doubling metric, it follows that $\# \left(A \cap B_{d_\varphi}^{\alpha\varphi}(2; h_1) \right) = O(1)$. Thus we get the desired upper bound. \square

7.2. Adjustment of Lemma 7.2 to Almost Invariant Spaces.

Definition 7.5. Let $0 \leq l \leq n$. For every $V \in \text{Gr}(l, n)$ denote $V^\rightarrow := \lim_{t \rightarrow \infty} g_t V$. This is a non-continuous mapping $\text{Gr}(l, n) \rightarrow \text{Gr}(l, n)^g$. For every $E \in \mathcal{I}_l$, denote

$$\text{Gr}(l, n)_{\rightarrow E} := \left\{ V \in \text{Gr}(l, n) : \lim_{t \rightarrow \infty} g_t V \in \text{Gr}(l, n)_E^g \right\},$$

and

$$H_{V \rightarrow E} := \{ h \in H : hV \in \text{Gr}(l, n)_{\rightarrow E} \}.$$

Let E_\bullet be a direction filtration and V_\bullet a flag with $L(V_\bullet) = L(E_\bullet)$ and $V_l \in \text{Gr}(l, n)_{\rightarrow E_l}$ for every $l \in L(V_\bullet)$. Denote $H_{V_\bullet \rightarrow E_\bullet} := \bigcap_{l \in L(V_\bullet)} H_{V_l \rightarrow E_l}$. \heartsuit

Example 2.5, continued. To describe explicitly $\text{Gr}(2, 4)_{\rightarrow E}$ for $E \in \mathcal{I}_2$, we will use a variation of the concept of Schubert cells. We will not use this representation in the rest of the paper, and instead use the more indirect description $\text{Gr}(2, 4)_{\rightarrow E} = H^- \text{Gr}(2, 4)_E^g$. For every $M \in M_{4 \times 2}(\mathbb{R})$ and a subset $F \subseteq \{1, 2, 3, 4\}$, denote by M_F the $(\#F) \times 2$

matrix composed of the F rows of M . We get

$$\begin{aligned}
\text{Gr}(2, 4)_{\rightarrow\{-1,0\}} &= \left\{ \text{Im}(M) : M = \begin{pmatrix} 1 & 0 \\ 0 & * \\ 0 & * \\ 0 & 0 \end{pmatrix}, \text{rank } M = 2 \right\} \\
&= \{ \text{span}(e_1, v) : 0 \neq v \in \text{span}(e_2, e_3) \}, \\
\text{Gr}(2, 4)_{\rightarrow\{-1,1\}} &= \left\{ \text{Im}(M) : M = \begin{pmatrix} 1 & * \\ 0 & * \\ 0 & * \\ 0 & 1 \end{pmatrix} \right\} \\
&= \{ \text{span}(e_1, v + e_4) : v \in \text{span}(e_2, e_3) \}, \\
\text{Gr}(2, 4)_{\rightarrow\{0,1\}} &= \left\{ \text{Im}(M) : M = \begin{pmatrix} * & * \\ * & * \\ * & * \\ 0 & 1 \end{pmatrix}, \text{rank } M_{\{2,3,4\}} = 2 \right\} \\
&= \left\{ \text{span}(e_4 + u, v + \alpha e_1) : \alpha \in \mathbb{R}, \text{ and } \begin{array}{l} 0 \neq v \in \text{span}(e_2, e_3), \\ u \in \text{span}(e_1, e_2, e_3) \end{array} \right\}, \\
\text{Gr}(2, 4)_{\rightarrow\{0,0\}} &= \left\{ \text{Im}(M) : M = \begin{pmatrix} * & * \\ 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \right\} \\
&= \{ \text{span}(e_2 + \alpha e_1, e_3 + \beta e_1) : \alpha, \beta \in \mathbb{R} \},
\end{aligned}$$

where $*$ represents any real number. Let $V_0 = \text{span}(e_1 + e_2 + e_3, e_4) \in \text{Gr}(2, 4)_{\rightarrow\{0,1\}}$ and $V_1 = \text{span}(e_1 + e_2, e_1 + e_3) \in \text{Gr}(2, 4)_{\rightarrow\{0,0\}}$. Then

$$\begin{aligned}
H_{V_0 \rightarrow \{0,1\}} &= \left\{ \begin{pmatrix} 1 & 0 & 0 & 0 \\ a & 1 & 0 & 0 \\ b & 0 & 1 & 0 \\ c & d & e & 1 \end{pmatrix} : a, b, c, d, e \in \mathbb{R}, (a, b) \neq (-1, -1) \right\}, \\
H_{V_1 \rightarrow \{0,0\}} &= \left\{ \begin{pmatrix} 1 & 0 & 0 & 0 \\ a & 1 & 0 & 0 \\ b & 0 & 1 & 0 \\ c & -c & -c & 1 \end{pmatrix} : a, b, c \in \mathbb{R}, a + b + 1 \neq 0 \right\}.
\end{aligned}$$

◇

Remark 7.6. The set $H_{V_\bullet \rightarrow E_\bullet}$ is not a group. It has the property that, for every $h_0 \in H_{V_\bullet \rightarrow E_\bullet}$,

$$H_{V_\bullet \rightarrow E_\bullet} = H_{h_0 V_\bullet \rightarrow E_\bullet} h_0.$$

Our next goal is to prove Lemma 7.10, which establishes a local bijection between $H_{V \rightarrow E}$ and $H_{V \rightarrow \cdot}$.

Lemma 7.7. *Let E_\bullet be a direction filtration and let V_\bullet be a flag with $L(V_\bullet) = L(E_\bullet)$ and $V_l \in \text{Gr}(l, n)_{\rightarrow E_l}$ for every $l \in L(V_\bullet)$. There exists a matrix $q_0 \in H^-$ such that $q_0 \cdot V_l^{\rightarrow} = V_l$ for all $l \in L(V_\bullet)$. Moreover, for E_\bullet fixed we can choose the mapping $V_\bullet \mapsto q_0$ so that it is differentiable as a map from a manifold of flags to H^- .*

Proof. We will first show the claim for a single space $V \in \text{Gr}(l, n)_{\rightarrow E}$. Let $W = (V^{\rightarrow})^\perp$ be the orthogonal complement of V^{\rightarrow} . Since V^{\rightarrow} decomposes as a direct sum of its intersections with eigenspaces V_η of g_t , it follows that W is the direct sum of the orthogonal complements of $V^{\rightarrow} \cap V_\eta$ in V_η , and hence is g_t -invariant.

Since $g_t V \xrightarrow{t \rightarrow \infty} V^{\rightarrow}$ and $V^{\rightarrow} \cap W = \{0\}$, it follows that $(g_t V) \cap W = \{0\}$. Since W is g_t -invariant we conclude that $V \cap W = \{0\}$. Therefore for every $v \in V^{\rightarrow}$ there exists a unique $q_0(v) \in V$ such that $q_0(v) - v \in W$. Extend q_0 to a linear map $q_0 : \mathbb{R}^n \rightarrow \mathbb{R}^n$, by $q_0(w) = w$ for every $w \in W$. As a linear map $q_0 : \mathbb{R}^n \rightarrow \mathbb{R}^n$, we think of q_0 as a matrix.

We claim that $q_t := g_t q_0 g_{-t} \xrightarrow{t \rightarrow \infty} \text{Id}$. Indeed, q_t is the identify map on W . We will show that q_t converges to the identify map on V^{\rightarrow} . For every $v \in V^{\rightarrow}$, the element $q_t v = g_t q_0 g_{-t} v$ lies in $g_t V$ and is the unique element such that $q_t v - v \in W$. Since $g_t V \xrightarrow{t \rightarrow \infty} V^{\rightarrow}$, it follows that $g_t q_0 g_{-t} v \xrightarrow{t \rightarrow \infty} v$. Therefore, $g_t q_0 g_{-t} \xrightarrow{t \rightarrow \infty} \text{Id}$, and hence $q_0 \in H^-$. Note that for fixed E , the mapping $V \mapsto q$ is differentiable as a map from $\text{Gr}(l, n)_{\rightarrow E}$ to H^- .

To prove the general case, denote $L(V_\bullet) = \{0 = l_0 < l_1 < \dots < l_k = n\}$. Let q'_0 be the matrix in H^- such that $q'_0 V_{l_{k-1}}^{\rightarrow} = V_{l_{k-1}}$. By induction on n , applying the lemma in dimension $l_{k-1} < n$ to the flow

$$g'_t = \frac{g_t|_{V_{l_{k-1}}^{\rightarrow}}}{(\det g_t|_{V_{l_{k-1}}^{\rightarrow}})^{1/l_{k-1}}},$$

the flag $\{0\} = (q'_0)^{-1} V_0 \subset V_{l_1} \cdots \subset (q'_0)^{-1} V_{l_{k-1}}$, and the direction filtration $\emptyset = E_0 \subset E_{l_1} \subset \dots \subset E_{l_{k-1}}$, there exists $q''_0 : V_{l_{k-1}}^{\rightarrow} \rightarrow V_{l_{k-1}}^{\rightarrow}$ such that $q''_0 V_{l_i}^{\rightarrow} = q'^{-1} V_{l_i}$ for every $0 \leq i \leq k-1$ and $g'_t q''_0 g'_{-t} \xrightarrow{t \rightarrow \infty} \text{Id}$. We can extend q''_0 to a map $q''_0 : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by $q''_0|_{(V_{l_{k-1}}^{\rightarrow})^\perp} = \text{Id}$. This

map satisfies $g_t q_0'' g_{-t} \xrightarrow{t \rightarrow \infty} \text{Id}$, hence $q_0 = q_0' q_0''$ is the desired matrix. The result follows. \square

Remark 7.8. For every $\varepsilon > 0$, if $d_{\text{Gr}(l,n)}(V_l, V_l^\rightarrow) < \varepsilon$ for every $l \in L(V_\bullet)$, then, by the differentiability of the construction of q_0 , we get that $d_{H^-}(q_0, \text{Id}) = O(\varepsilon)$.

Lemma 7.9. *For every $0 \leq l \leq n$, $E \in \mathcal{I}_l$, and $V \in \text{Gr}(l, n)_E^g$, we have*

$$H_{V \rightarrow E} = H_V.$$

Proof. By the definitions of the sets, $H_V \subseteq H_{V \rightarrow E}$. Let $h \in H_{V \rightarrow E}$. By the definition of $H_{V \rightarrow E}$,

$$g_t h V \xrightarrow{t \rightarrow \infty} V^\rightarrow \in \text{Gr}(l, n)_E^g.$$

Next, the definition of H implies that $g_{-t} h g_{-t} \xrightarrow{t \rightarrow \infty} \text{Id}$, and thanks to the g_t -invariance of V ,

$$(7.2) \quad g_{-t} h V = g_{-t} h g_t V \xrightarrow{t \rightarrow \infty} V \in \text{Gr}(l, n)_E^g.$$

Consequently, $d_{\text{Gr}(l,n)}(g_t h V, \text{Gr}(l, n)_E^g) \xrightarrow{t \rightarrow \pm \infty} 0$, and hence, by Theorem 2.7, we get that $g_t h V \in \text{Gr}(l, n)_E^g$ for every t and in particular $t \mapsto g_t h V$ is constant. Eq. (7.2) implies that $hV = V$, hence $h \in H_V$. \square

Recall that $H^{-0} = H^- H^0$ is the group of matrices $A \in \text{SL}_n(\mathbb{R})$ such that $A_{ij} \neq 0$ only if $\eta_i \leq \eta_j$.

Lemma 7.10. *Let E_\bullet be a direction filtration and let V_\bullet be a flag with $L(V_\bullet) = L(E_\bullet)$ and $V_l \in \text{Gr}(l, n)_{\rightarrow E_l}$ for every $l \in L(V_\bullet)$. Let $q_0 \in H^-$ be a map such that $q_0 V_l^\rightarrow = V_l$ for every $l \in L(V_\bullet)$. Then there exist open sets $U_1 \subseteq H_{V_\bullet \rightarrow E_\bullet}$ and $U_2 \subseteq H_{V_\bullet^\rightarrow}$ containing Id , a diffeomorphism $\check{h} : U_1 \rightarrow U_2$, and a map $\check{q} : U_1 \rightarrow H^{-0}$, such that*

$$h q_0 = \check{q}(h) \check{h}(h),$$

for every $h \in U_1$. Moreover, for every compact set $K \subseteq H$ and every $\varepsilon_0 > 0$ there exists $\varepsilon_K > 0$ such that if $d_{H^-}(q_0, \text{Id}) < \varepsilon_K$, then U_1 and U_2 contains $H_{V_\bullet \rightarrow E_\bullet} \cap K$ and $H_{V_\bullet^\rightarrow} \cap K$, respectively. Furthermore, for every $h \in U_1 \cap K$, we have $d_H(\check{h}(h), h) < \varepsilon_0$.

Proof. The multiplication map

$$m_{H^{-0}H} : H \times H^{-0} \rightarrow \text{SL}_n(\mathbb{R})$$

$$m_{H^{-0}H} : (h, q^{-0}) \mapsto q^{-0} h$$

is one-to-one and has an open image. Set

$$\begin{aligned} U'_1 &:= \{h \in H : hq_0 \in \text{Im}(m_{H^0H})\}, \\ U'_2 &:= \{h \in H : hq_0^{-1} \in \text{Im}(m_{H^0H})\}. \end{aligned}$$

For every $h \in U'_1$ denote

$$(7.3) \quad (\check{h}(h), \check{q}(h)) := m_{H^0H}^{-1}(hq_0)$$

and note that the equality $hq_0 = \check{q}(h)\check{h}(h)$ implies that \check{h} induces an $\check{h} : \check{U}'_1 \rightarrow \check{U}'_2$. Note that $\text{Id} \in U'_1, U'_2$.

The sets $U_1 := U'_1 \cap H_{V_\bullet \rightarrow E_\bullet}$ and $U_2 := U'_2 \cap H_{V_\bullet \rightarrow}$ have the property that, for every $h \in U'_1$, we have

$$(7.4) \quad h \in U_1 \iff \forall l \in L(V_\bullet), hV_l \in \text{Gr}(l, n)_{\rightarrow E_l}$$

$$(7.5) \quad \iff \forall l \in L(V_\bullet), hq_0V_l^{\rightarrow} \in \text{Gr}(l, n)_{\rightarrow E_l}$$

$$(7.6) \quad \iff \forall l \in L(V_\bullet), \check{q}(h)\check{h}(h)V_l^{\rightarrow} \in \text{Gr}(l, n)_{\rightarrow E_l}$$

$$(7.7) \quad \iff \forall l \in L(V_\bullet), \check{h}(h)V_l^{\rightarrow} \in \text{Gr}(l, n)_{\rightarrow E_l}$$

$$(7.8) \quad \iff \forall l \in L(V_\bullet), \check{h}(h) \in H_{V_l^{\rightarrow}},$$

where (7.4) follows from the definition of U_1 , (7.5) follows from the definition of q_0 , (7.6) follows from Eq. (7.3), (7.7) follows from the fact that H^- preserves $\text{Gr}(l, n)_{\rightarrow E}$, and (7.8) follows from Lemma 7.9.

Therefore U_1, U_2 satisfy the desired condition.

Note that the smaller q_0 is, the closer hq_0, hq_0^{-1} are to $H \subset \text{Im}(m_{H^0H})$ and hence the second condition follows. \square

Fix a direction filtration E_\bullet and a flag V_\bullet with $L(V_\bullet) = L(E_\bullet)$ and $V_l \in \text{Gr}(l, n)_{\rightarrow E_l}$ for every $l \in L(V_\bullet)$. Let $q_0 \in H^-$ be the element from Lemma 7.7 which satisfies $q_0V_\bullet^{\rightarrow} = V_\bullet$. Lemma 7.10 gives us a measure $\mu' = \check{h}_*^{-1}\mu$ on $H_{V_l \rightarrow E_l}$, where μ is the invariant measure on $H_{V_\bullet^{\rightarrow}}$.

To extend Lemma 7.2 to $H_{V_\bullet \rightarrow E_\bullet}$ we need to compute the volume $\mu'(B_{d_\varphi}(h; \exp(-t)))$ for $h \in H_{V_\bullet \rightarrow E_\bullet}$.

Lemma 7.11. *There exists $\varepsilon > 0$ such that if $d_\varphi(q_0, \text{Id}) < \varepsilon$, then every $h_0 \in H_{V_\bullet \rightarrow E_\bullet}$ with $d_\varphi(h_0, \text{Id}) \leq 1$ satisfies*

$$\mu'(B_{d_\varphi}(h_0; \exp(-t))) = \Theta(\exp(-t\delta(E_\bullet))) \text{ as } t \rightarrow \infty.$$

Proof. Let \check{h}, \check{q} be as in Lemma 7.10. If $d_\varphi(q_0, \text{Id})$ is small enough, then $B_{d_\varphi}(\text{Id}; 2^{1/\alpha_\varphi}) \cap H_{V_\bullet \rightarrow E_\bullet}$ is in the domain of \check{h}, \check{q} and similarly $B_{d_\varphi}(\text{Id}; 2^{1/\alpha_\varphi}) \cap H_{V_\bullet}$ is in the image of \check{h} .

We first show the result when $h_0 = \text{Id}$. We want to calculate

$$(7.9) \quad \begin{aligned} & \mu(\{h \in H_{V_\bullet} : \check{h}^{-1}(h) \in B_{d_\varphi}(\text{Id}; \exp(-t))\}) \\ &= \mu(\check{h}(B_{d_\varphi}(\text{Id}; \exp(-t)) \cap H_{V_\bullet \rightarrow E_\bullet})). \end{aligned}$$

Let \check{h}' , \check{q}' be the functions generated by Lemma 7.10 applied to E_\bullet , $q_1 V_\bullet$, and q_1 , where $q_1 = g_t q_0 g_{-t}$, and $q_1 V_\bullet \rightarrow$. These functions satisfy

$$(7.10) \quad \check{q}'(h) \check{h}'(h) = h q_1 = h g_t q_0 g_{-t}.$$

Substituting $h = g_t h g_{-t}$ in Eq. (7.10) we get

$$\check{q}'(g_t h g_{-t}) \check{h}'(g_t h g_{-t}) = g_t h_0 g_{-t} = (g_t \check{q}'(h) g_{-t})(g_t \check{h}'(h) g_{-t}).$$

Since the multiplication map $m_{H^{-0}H} : H^{-0} \times H \rightarrow \text{SL}_n(\mathbb{R})$ is one-to-one, it follows that

$$\check{q}'(g_t h g_{-t}) = g_t \check{q}'(h) g_{-t} \text{ and } \check{h}'(g_t h g_{-t}) = g_t \check{h}'(h) g_{-t}.$$

Conjugating by g_t we get that the quantity in Eq. (7.9) equals

$$\exp(-t\delta(E_\bullet)) \mu(\check{h}'(B_{d_\varphi}(\text{Id}; 1) \cap H_{V_\bullet \rightarrow E_\bullet})).$$

Since q_1 is close to Id , Lemma 7.10 implies that the map $\check{h}'(h)|_{B_{d_\varphi}(\text{Id}; 2^{1/\alpha_\varphi)}$ is close to the identity map, and hence

$$\mu(\check{h}'(B_{d_\varphi}(\text{Id}; 1) \cap H_{g_t V_\bullet \rightarrow E_\bullet})) = \Theta(1),$$

and hence our prediction of the μ' -measure of a small ball around Id is correct.

For $h_0 \neq \text{Id}$, write $\check{q}'(h_0) = q_2 q_2^0$, where $q_2 \in H^-$ and $q_2^0 \in H^0$. Since

$$\forall l \in L(V_\bullet), h_0 q_0 V_l \rightarrow = \check{q}'(h_0) \check{h}'(h_0) V_l \rightarrow = q_2 q_2^0 V_l \rightarrow,$$

it follows that for $V_l' := h_0 V_l$ we have $(V_l') \rightarrow = q_2^0 V_l \rightarrow$. We can now construct μ'' on $H_{V_\bullet' \rightarrow E_\bullet}$ as we constructed μ' . In much the same way, since $d_{H^-}(q_0, \text{Id})$ is arbitrarily small and h_0 lies in a compact set, we deduce from Lemma 7.10 that $d_{H^-}(\check{q}'(h_0), \text{Id}) = d_{H^-}(q_1, \text{Id})$ is arbitrarily small, and hence

$$\mu''(B_{d_\varphi}(\text{Id}; \exp(-t)) \cap H_{V_\bullet' \rightarrow E_\bullet}) = \Theta(\exp(-t\delta(E_\bullet))).$$

By Lemma 7.10, if $d_{H^-}(q_0, \text{Id})$ is small enough, then both $H_{V_\bullet \rightarrow E_\bullet}$ and $H_{V_\bullet' \rightarrow E_\bullet}$ are manifolds in a bounded set. Since μ'' , μ' are smooth measures with positive density, and since $H_{V_\bullet \rightarrow E_\bullet} h_0^{-1} = H_{V_\bullet' \rightarrow E_\bullet}$, we obtain that the Radon-Nikodym derivative $(R_{h_0^{-1}})_* \mu' / \mu''$ is bounded on $B_{d_\varphi}(\text{Id}; \exp(-t)) \cap H_{V_\bullet' \rightarrow E_\bullet}$, where $R_{h_0^{-1}}$ is the map of multiplication

from the right by h_0^{-1} , and $(R_{h_0^{-1}})_*$ is the pushforward of measure. Hence,

$$\begin{aligned} \mu'(B_{d_\varphi}(h; \exp(-t)) \cap H_{V_\bullet \rightarrow E_\bullet}) &= (R_{h_0^{-1}})_* \mu'(B_{d_\varphi}(\text{Id}; \exp(-t)) \cap H_{V'_\bullet \rightarrow E_\bullet}) \\ &= \Theta(\mu''(B_{d_\varphi}(\text{Id}; \exp(-t)) \cap H_{V'_\bullet \rightarrow E_\bullet})) \\ &= \Theta \exp(-t\delta(E_\bullet)), \end{aligned}$$

as desired. \square

Definition 7.12. For every $l = 0, \dots, n$ and $E \in \mathcal{I}_l$ we let $\text{Gr}(l, n)_{\rightarrow E}^{\varepsilon, g}$ denote the set of vector spaces $V \in \text{Gr}(l, n)_{\rightarrow E}$ such that $d_{\text{Gr}(l, n)}(V, V^{\rightarrow}) < \varepsilon$. \heartsuit

Lemma 7.4 shows that

$$(7.11) \quad \mu(B_{d_\varphi}(h; \exp(-t))) = \Theta(\exp(-t\delta(E_\bullet))),$$

for every $h \in H_{V_\bullet}$ and $t > 0$. Combined with standard covering arguments, Eq. (7.11) proves Lemma 7.2. Replacing Eq. (7.11) with Lemma 7.11 in the proof of Lemma 7.2 we can now prove the following.

Lemma 7.13. *There is $\varepsilon_0 > 0$ such that the following holds. Let V_\bullet be a flag and let E_\bullet be a direction filtration with $L(V_\bullet) = L(E_\bullet)$, $t > 0$, and $V_l \in \text{Gr}(l, n)_{\rightarrow E_l}^{\varepsilon_0, g}$ for every $l \in L(V_\bullet)$. There is a set $A \subset B_{d_\varphi}(\text{Id}; 1)$ of size $\Theta(\exp(t\delta(E_\bullet)))$ that satisfies the following conditions.*

- (1) $d_\varphi(h_1, h_2) \geq \exp(-t)$ for all pairs of distinct elements $h_1, h_2 \in A$;
- (2) We have $h \in H_{V_\bullet \rightarrow E_\bullet}$ for every $h \in A$.

Moreover, there is no such set A larger than $\Theta(\exp(t\delta(E_\bullet)))$, even if we weaken Condition (2) to

- (2') For every $h \in A$ there exists $h' \in H_{V_\bullet \rightarrow E_\bullet}$ such that $d_\varphi(h, h') < \exp(-t)$.

7.3. The Changing Step Lemma. In view of Observation 4.14, the following lemma will construct Alice's step whenever $f_{E_\bullet}^{h_\infty \Lambda}$ changes from time T_m to time T_{m+1} via a non-null vertex. In these steps $\#A_{m+1} = 1$, that is, Alice's choice determines h_{m+1} . Consequently, Lemma 7.16 below will prove the existence of an element $h \in B_{d_\varphi}(\text{Id}; 1)$ with a certain behavior.

Definition 7.14. Let E_\bullet, E'_\bullet be direction filtrations with $L(E_\bullet) = L(E'_\bullet)$. We write $E_\bullet \preceq E'_\bullet$ if $E_l \preceq E'_l$ for every $l \in L(E_\bullet)$. \heartsuit

Remark 7.15. In Definition 7.14, the conditions $E_l \trianglelefteq E'_l$ are required independently for each l . That is, for every $l \in L(E_\bullet)$ we can increase some of the elements in E_l to obtain E'_l . The increments are not required to be compatible for different values l . We will see (Definition 7.21) a precise formulation of increasing all sets in a compatible way and show (Theorem 7.22) that the definitions are equivalent.

Lemma 7.16. *Let $E_\bullet \trianglelefteq E'_\bullet$ be direction filtrations and let V_\bullet be a flag such that $L(V_\bullet) = L(E_\bullet)$ and $V_l \in \text{Gr}(l, n)_{\rightarrow E_l}$ for every $l \in L(E_\bullet)$. There exists $h \in B_{d_\varphi}(\text{Id}; 1)$ such that $hV_l \in \text{Gr}(l, n)_{\rightarrow E'_l}$ for every $l \in L(V_\bullet)$. Moreover, the convergence rate of $\lim_{t \rightarrow \infty} g_t h V_l \in \text{Gr}(l, n)_{E'_l}^g$ is uniform on compact sets of possible V_\bullet -s.*

The proof of Lemma 7.16 occupies the rest of this subsection. We will break it up into several claims.

Claim 7.17 (Transitivity of Lemma 7.16). *The lemma is transitive with respect to $E_\bullet \trianglelefteq E'_\bullet$. That is, let $E_\bullet \trianglelefteq E'_\bullet \trianglelefteq E''_\bullet$, and assume the statement of the lemma holds for the tuples $E_\bullet \trianglelefteq E'_\bullet$ and $E'_\bullet \trianglelefteq E''_\bullet$. Then the statement also holds for $E_\bullet \trianglelefteq E''_\bullet$.*

Proof. Let V_\bullet be a flag such that $L(V_\bullet) = L(E_\bullet)$, and $V_l \in \text{Gr}(l, n)_{\rightarrow E_l}$ for every $l \in L(V_\bullet)$. Fix $t_0, t_1 > 0$ to be determined later. By the statement of the lemma applied to $E_\bullet \trianglelefteq E'_\bullet$ and the flag $g_{t_0} V_\bullet$, there exists $h_0 \in B_{d_\varphi}(\text{Id}; 1)$ such that $h_0 g_{t_0} V_l \in \text{Gr}(l, n)_{\rightarrow E'_l}$ for all $l \in L(E_\bullet)$. By the statement of the lemma for $E'_\bullet \trianglelefteq E''_\bullet$ and the flag $g_{t_1} h_0 g_{t_0} V_\bullet$ there exists $h_1 \in B_{d_\varphi}(\text{Id}; 1)$ such that $h_1 g_{t_1} h_0 g_{t_0} V_l \in \text{Gr}(l, n)_{\rightarrow E''_l}$ for all $l \in L(E_\bullet)$.

And hence also $g_{-t_1-t_0} h_1 g_{t_1+t_0} g_{-t_0} h_0 g_{t_0} V_l \in \text{Gr}(l, n)_{\rightarrow E''_l}$. If t_0, t_1 are large enough, both $g_{-t_0} h_0 g_{t_0}$ and $g_{-t_1-t_0} h_1 g_{t_1+t_0}$ are close to Id , and their product lies in $B_{d_\varphi}(\text{Id}; 1)$. \square

Claim 7.18 (The meaning of uniform convergence). *Let $0 \leq l \leq n$, $E \in \mathcal{I}_l$ and $K \subseteq \text{Gr}(l, n)_{\rightarrow E}$. The convergence $g_t V \rightarrow V^\rightarrow$ is uniform for $V \in K$ if and only if the (compact) closure $\bar{K} \subseteq \text{Gr}(l, n)$ is a subset of $\text{Gr}(l, n)_{\rightarrow E}$.*

Proof. By Lemma 7.7, for every $V \in K$ there exists $q_0(V)$ such that $V = q_0(V) V^\rightarrow$ and moreover, the function $V \mapsto q_0(V)$ is continuous. Therefore, if $K \subseteq \text{Gr}(l, n)_{\rightarrow E}$ is compact, then so is $q_0(K)$, and $g_t q_0(K) g_{-t}$ contracts uniformly.

To prove the other direction, by Lemma 7.7 once again, if $Q_0 \subseteq H^-$ is a compact neighborhood of Id , then $Q_0 \text{Gr}(l, n)_E^g$ is a compact neighborhood of $\text{Gr}(l, n)_E^g$ in $\text{Gr}(l, n)_{\rightarrow E}$. Therefore, if the convergence

rate $g_t V \rightarrow V^\rightarrow$ is uniform for $V \in K$, there exists $t_0 > 0$ such that $g_{t_0} K \subseteq Q_0 \text{Gr}(l, n)_E^g$, and hence $K \subseteq g_{-t_0} Q_0 \text{Gr}(l, n)_E^g$ as desired. \square

From Claim 7.18 it easily follows that the uniform convergence satisfies transitivity as well.

Claim 7.19 (Reduction to the case where V_\bullet is g -invariant). *It is enough to prove Lemma 7.16 for flags V_\bullet satisfying $V_l \in \text{Gr}(l, n)_{E_l}^g$ for every $l \in L(E_\bullet)$.*

Proof. Assume that Lemma 7.16 holds whenever $V_l \in \text{Gr}(l, n)_{E_l}^g$ for every $l \in L(E_\bullet)$. By Claim 7.18, there exist compact sets $K_l \subseteq \text{Gr}(l, n)_{\rightarrow E'_l}$ such that for every flag V_\bullet with $L(V_\bullet) = L(E_\bullet)$ and $V_l \in \text{Gr}(l, n)_{E_l}^g$ for every $l \in L(E_\bullet)$ there is $h \in B_{d_\varphi}(\text{Id}; 1)$ such that $hV_l \in K_l$ for every $l \in L(E_\bullet)$. We will show that for every compact set $K_l^{\text{src}} \subseteq \text{Gr}(l, n)_{\rightarrow E_l}$ there exists $K_l^{\text{tar}} \subseteq \text{Gr}(l, n)_{\rightarrow E'_l}$, such that for every flag V_\bullet with $V_l \in K_l^{\text{src}}$ for every $l \in L(E_\bullet)$ there exists $h \in B_{d_\varphi}(\text{Id}; 1)$ with $hV_l \in K_l^{\text{tar}}$. This, together with Claim 7.18, establishes Lemma 7.16.

Fix $\varepsilon > 0$ and a flag V_\bullet with $L(V_\bullet) = L(E_\bullet)$ such that $V_l \in K_l^{\text{src}}$ for every $l \in L(V_\bullet)$. After applying g_t , we may assume $K_l^{\text{src}} \subseteq U_\varepsilon(\text{Gr}(l, n)_{E_l}^g)$. Hence, for some $q_0 \in B_{d_{H^-}}(\text{Id}; O(\varepsilon))$ we have $V_l = q_0 V_l^\rightarrow$. After applying $g_{\log \varepsilon}$ to K_l we may assume that if $V_l^\rightarrow \in \text{Gr}(l, n)_{E_l}^g$, then there exists $h_0 \in B_{d_\varphi}(\text{Id}; \varepsilon)$ such that $h_0 V_l^\rightarrow = h_0 q_0^{-1} V_l \in K_l$. Since h_0 and q_0 are arbitrarily small, there exist $q_1 \in B_{d_{H^-}}(\text{Id}; 1)$ and $h_1 \in B_{d_\varphi}(\text{Id}; 1)$ such that $q_1 h_1 = h_0 q_0^{-1}$. It follows that $h_1 V_l \in B_{H^-}(\text{Id}; 1) K_l =: K_l^{\text{tar}}$, as desired. \square

To employ Claim 7.17 we will investigate the poset \mathcal{I}_\bullet of direction filtrations with dimension sequence $L = \{0 = l_0 < l_1 < \dots < l_k = n\}$, where the partial order is given by $E_\bullet \trianglelefteq E'_\bullet$, as defined in Definition 7.14.

Definition 7.20 (Multiplicity of elements in multisets). Let E be a multiset. For every element x we denote by $\text{mult}(x; E)$ the multiplicity of x in E . \heartsuit

Definition 7.21 (Elementary flip). We say that $E_\bullet \triangleleft E'_\bullet$ is an *elementary flip* if there exist $\eta < \eta' \in E_{\text{all}}$ and $1 \leq a < b \leq k$ such that

$$(7.12) \quad E'_i = \begin{cases} E_{l_i} - \{\eta\} + \{\eta'\}, & \text{if } a \leq i < b, \\ E_{l_i}, & \text{otherwise.} \end{cases}$$

We use $+$ and $-$ instead of \cup and \setminus , as these are multiset operations, under which multiplicities are added or subtracted. In this case $E_\bullet \triangleleft E'_\bullet$.

is called an (η, η', a, b) -*elementary flip*. We think of an elementary flip as an operation transforming E_\bullet into E'_\bullet . Note that a direction filtration $E_\bullet \in \mathcal{I}_\bullet$ has an (η, η', a, b) -elementary flip $E_\bullet \triangleleft E'_\bullet$ if and only if the following condition of multiplicities is satisfied:

$$\text{mult}(\eta; E_{l_{a-1}}) < \text{mult}(\eta; E_{l_a}) \text{ and } \text{mult}(\eta'; E_{l_{b-1}}) < \text{mult}(\eta'; E_{l_b}).$$

Indeed, otherwise the unique E'_\bullet defined by Eq. (7.12) will fail to satisfy $E'_{l_{a-1}} \subset E'_{l_a}$ or $E'_{l_{b-1}} \subset E'_{l_b}$, respectively. \heartsuit

Theorem 7.22 (The poset \mathcal{I}_\bullet is spanned by elementary flips). *If $E_\bullet \trianglelefteq E'_\bullet$, then there exists a sequence of direction filtration $E_\bullet = E_\bullet^0 \triangleleft E_\bullet^1 \triangleleft \dots \triangleleft E_\bullet^r = E'_\bullet$, where each consecutive $E_\bullet^j \triangleleft E_\bullet^{j+1}$ is an elementary flip.*

Proof. The plan: Since \mathcal{I}_\bullet is a finite poset, it is enough to show that for every $E_\bullet \triangleleft E'_\bullet$ there is an elementary flip $E_\bullet \triangleleft E''_\bullet$ such that $E''_\bullet \trianglelefteq E'_\bullet$.

To use the fact $E_\bullet \triangleleft E'_\bullet$, define

$$(7.13) \quad \begin{aligned} \sigma : \{0, \dots, k\} \times (\text{set}(E_{\text{all}}) \cup \{-\infty\}) &\rightarrow \mathbb{N}_0, \\ \sigma(i, \eta) &:= \#(E_{l_i} \cap (-\infty, \eta]) = \sum_{\substack{\tilde{\eta} \in \text{set}(E_{\text{all}}) \\ \tilde{\eta} \leq \eta}} \text{mult}(\tilde{\eta}, E_{l_i}). \end{aligned}$$

Here $\text{set}(E_{\text{all}})$ stands for the set of distinct elements in E_{all} and $\mathbb{N}_0 = \{0, 1, 2, \dots\}$ is the set of natural numbers. Define σ' in a similar fashion with respect to E'_\bullet instead of E_\bullet . Note that $E_\bullet \trianglelefteq E'_\bullet$ if and only if $\sigma(i, \eta) \geq \sigma'(i, \eta)$ for all i, η .

If $E_\bullet \triangleleft E''_\bullet$ is an (η, η', a, b) -elementary flip, then the functions σ, σ'' satisfy that $\sigma'' = \sigma - \mathbb{1}_R$, where R is the rectangle $\{(i, \tilde{\eta}) : a \leq i < b, \eta \leq \tilde{\eta} < \eta'\}$. We need the function σ'' to remain greater or equal to σ' to ensure $E''_\bullet \trianglelefteq E'_\bullet$. To help us find a suitable rectangle, we will identify values $0 < i_0 < i_2$ such that $\sigma(i, \eta) - \sigma'(i, \eta)$ is zero for $0 \leq i < i_0$ and all $\eta \in E_{\text{all}} \cup \{-\infty\}$ and nonzero for all $i_0 \leq i < i_2$ and some $\eta \in E_{\text{all}}$. Moreover $i \mapsto \sigma(i, \eta) - \sigma'(i, \eta)$ will be weakly increasing in i for $0 \leq i < i_2$ and every $\eta \in E_{\text{all}} \cup \{\infty\}$.

Additional notations: Since we are interested in monotonicity properties of $\sigma - \sigma'$, define the function $\tau : \{1, \dots, k\} \times (\text{set}(E_{\text{all}}) \cup \{-\infty\}) \rightarrow \mathbb{Z}$ by

$$\tau(i, \eta) := \sigma(i, \eta) - \sigma'(i, \eta) - \sigma(i-1, \eta) + \sigma'(i-1, \eta).$$

By definition, for every $\eta \in E_{\text{all}} \cup \{-\infty\}$ and $0 \leq i \leq k$ we have

$$(7.14) \quad \sigma(i, \eta) - \sigma'(i, \eta) = \sum_{i'=0}^i \tau(i', \eta) = - \sum_{i'=i+1}^k \tau(i', \eta).$$

The last equality holds since $\sigma(k, \eta) = \sigma'(k, \eta)$, because $E_{l_k} = E'_{l_k} = E_{\text{all}}$. Denoting $D_i := E_{l_i} - E_{l_{i-1}}$ and $D'_i := E'_{l_i} - E'_{l_{i-1}}$, we have

$$\tau(i, \eta) = \#\{\tilde{\eta} \in D_i - D'_i : \tilde{\eta} \geq \eta\}.$$

Note that here we regard D_i and D'_i as multisets with possibly negative multiplicities.

Choosing indices: The minimal index i_0 for which the function $\eta \mapsto \tau(i_0, \eta)$ is nonzero is the minimal i_0 for which $E_{l_{i_0}} \neq E'_{l_{i_0}}$ (which exists because $E_{\bullet} \neq E'_{\bullet}$), and satisfies

$$\tau(i_0, \eta) = \sum_{i=0}^{i_0} \tau(i, \eta) = \sigma(i_0, \eta) - \sigma'(i_0, \eta) \geq 0 \quad \text{for all } \eta \in E_{\text{all}}.$$

Since $E_{i_0} \triangleleft E'_{i_0}$, it follows that $\tau(i_0, \eta) = \sigma(i_0, \eta) - \sigma'(i_0, \eta)$ is positive for some $\eta \in E_{\text{all}}$.

Similarly, looking at the right-hand side of Eq. (7.14) we see that τ attains a negative value for some i , and we let i_2 be the minimal index such that $\eta \mapsto \tau(i_2, \eta)$ attains a negative value. It follows from the equality

$$\tau(i_2, \eta_n) = \#E_{l_{i_2}} - \#E'_{l_{i_2}} - \#E_{l_{i_2-1}} + \#E'_{l_{i_2-1}} = 0,$$

that there exists an index $1 < s_2 \leq n$ such that $\tau(i_2, \eta_{s_2-1}) < 0$ but $\tau(i_2, \eta_s) \geq 0$ for all $s \geq s_2$.

Since

$$\sum_{i=1}^{i_2} \tau(i, \eta_{s_2-1}) = \sigma(i, \eta_{s_2-1}) - \sigma'(i, \eta_{s_2-1}) \geq 0,$$

it follows that for some $i_1 < i_2$ we have $\tau(i_1, \eta_{s_2-1}) > 0$. Let $s_1 \geq 1$ be the minimal index such that $\tau(i, \eta_s) > 0$ for every $\eta_{s_1} \leq \eta_s \leq \eta_{s_2-1}$.

Obtaining the desired flip: We claim that we can perform an $(\eta_{s_1}, \eta_{s_2}, i_1, i_2)$ -elementary flip to E_{\bullet} , and obtain a sequence E''_{\bullet} for which $E''_{\bullet} \triangleleft E'_{\bullet}$.

To change one occurrence of η_{s_1} to η_{s_2} between the indices l_{i_1} and l_{i_2-1} , while preserving the requirement that the resulting sequence E''_{\bullet} is a direction filtration (i.e. the multisets E''_{\bullet} are nested, see Eq. (2.2)), it is sufficient and necessary to show that

$$(7.15) \quad \text{mult}(\eta_{s_1}; E_{l_{i_1-1}}) < \text{mult}(\eta_{s_1}; E_{l_{i_1}}) \quad \text{and} \quad \text{mult}(\eta_{s_2}; E_{l_{i_2-1}}) < \text{mult}(\eta_{s_2}; E_{l_{i_2}})$$

Since

$$\tau(i_1, \eta_{s_1}) - \tau(i_1, \eta_{s_1-1}) = \text{mult}(\eta_{s_1}; E_{l_{i_1}}) - \text{mult}(\eta_{s_1}; E_{l_{i_1-1}})$$

and

$$\tau(i_2, \eta_{s_2}) - \tau(i_2, \eta_{s_2-1}) = \text{mult}(\eta_{s_2}; E_{l_{i_2}}) - \text{mult}(\eta_{s_2}; E_{l_{i_2-1}}),$$

inequalities (7.15) are equivalent to

$$(7.16) \quad \tau(i_1, \eta_{s_1}) > \tau(i_1, \eta_{s_1-1}) \text{ and } \tau(i_2, \eta_{s_2-1}) < \tau(i_2, \eta_{s_2}),$$

which follows from the choices of s_1, s_2 . Define

$$E''_{l_i} := \begin{cases} E_{l_i} - \{\eta_{s_1}\} + \{\eta_{s_2}\}, & \text{if } i_1 \leq i < i_2, \\ E_{l_i}, & \text{otherwise.} \end{cases}$$

To show that $E''_{\bullet} \trianglelefteq E'_{\bullet}$, define σ'' as in Eq. (7.13) with respect to E''_{\bullet} . By the definition of E''_{\bullet} we obtain

$$\sigma''(i, \eta) = \begin{cases} \sigma(i, \eta) - 1, & \text{if } i_1 \leq i < i_2 \text{ and } \eta_{s_1} \leq \eta < \eta_{s_2}, \\ \sigma(i, \eta), & \text{otherwise.} \end{cases}$$

We need to show that $\sigma'' \geq \sigma'$ pointwise, which is equivalent to

$$\sigma(i, \eta) > \sigma'(i, \eta) \text{ for every } i_1 \leq i < i_2, \eta_{s_1} \leq \eta < \eta_{s_2}.$$

These inequalities follow from the fact that $\sigma(i, \eta) - \sigma'(i, \eta) = \sum_{i' < i} \tau(i', \eta)$, which is a sum of nonnegative numbers by the definition of i_2 and where the summand $\tau(i_1, \eta)$ is positive by the definition of η_{s_1} .

Altogether we have shown that there is E''_{\bullet} such that $E_{\bullet} \triangleleft E''_{\bullet} \trianglelefteq E'_{\bullet}$, and the tuple $E_{\bullet} \triangleleft E''_{\bullet}$ is an elementary flip. Iterating this procedure for $E''_{\bullet} \trianglelefteq E'_{\bullet}$ yields the desired result. \square

Example 7.23. Let $n = 4$, $E_{\text{all}} = \{-2, -1, 1, 2\}$, and $l_i = i$ for $i = 0, \dots, 4$. Denote

$$E_{\bullet} = \{\emptyset \subset \{-2\} \subset \{-2, 1\}, \{-2, -1, 1\}, E_{\text{all}}\},$$

and

$$E'_{\bullet} = \{\emptyset \subset \{2\} \subset \{-1, 2\}, \{-1, 1, 2\}, E_{\text{all}}\}.$$

We will describe in detail the process elaborated in the proof of Theorem 7.22. First, compute σ, σ', τ .

$\sigma(i, \eta) :$

$i \backslash \eta$	$-\infty$	-2	-1	1	2
0	0	0	0	0	0
1	0	1	1	1	1
2	0	1	1	2	2
3	0	1	2	3	3
4	0	1	2	3	4

 $\sigma'(i, \eta) :$

$i \backslash \eta$	$-\infty$	-2	-1	1	2
0	0	0	0	0	0
1	0	0	0	0	1
2	0	0	1	1	2
3	0	0	1	2	3
4	0	1	2	3	4

 $\tau(i, \eta) :$

$i \backslash \eta$	$-\infty$	-2	-1	1	2
1	0	1	1	1	0
2	0	0	-1	0	0
3	0	0	1	0	0
4	0	-1	-1	-1	0

Hence, $i_0 = 1$, $i_2 = 2$, $s_2 = 3$, $\eta_{s_2} = 1$, $i_1 = 1$, $s_1 = 1$, $\eta_{s_1} = -2$.
 Consequently, we obtain the elementary flip $E_{\bullet} \triangleleft E_{\bullet}^{(1)}$, where

$$E_{\bullet}^{(1)} = \{\emptyset \subset \{1\} \subset \{-2, 1\}, \{-2, -1, 1\}, E_{\text{all}}\}.$$

The blue rectangle in the $\sigma(i, \eta)$ table denotes the rectangle R for which $\sigma^{(1)} = \sigma - \mathbb{1}_R$. Continuing the process for $E_{\bullet}^{(1)} \trianglelefteq E'_{\bullet}$, we have

$\sigma^{(1)}(i, \eta) :$

$i \backslash \eta$	$-\infty$	-2	-1	1	2
0	0	0	0	0	0
1	0	0	0	1	1
2	0	1	1	2	2
3	0	1	2	3	3
4	0	1	2	3	4

 $\tau(i, \eta) :$

$i \backslash \eta$	$-\infty$	-2	-1	1	2
1	0	0	0	1	0
2	0	1	0	0	0
3	0	0	1	0	0
4	0	-1	-1	-1	0

Hence, $i_0 = 1$, $i_2 = 4$, $s_2 = 4$, $\eta_{s_2} = 2$, $i_1 = 1$, $s_1 = 3$, $\eta_{s_1} = 1$.
 Consequently, we obtain the elementary flip $E_{\bullet}^{(1)} \triangleleft E_{\bullet}^{(2)}$, where

$$E_{\bullet}^{(2)} = \{\emptyset \subset \{2\} \subset \{-2, 2\}, \{-2, -1, 2\}, E_{\text{all}}\}.$$

 $\sigma^{(2)}(i, \eta) :$

$i \backslash \eta$	$-\infty$	-2	-1	1	2
0	0	0	0	0	0
1	0	0	0	0	1
2	0	1	1	1	2
3	0	1	2	2	3
4	0	1	2	3	4

 $\tau(i, \eta) :$

$i \backslash \eta$	$-\infty$	-2	-1	1	2
1	0	0	0	0	0
2	0	1	0	0	0
3	0	0	1	0	0
4	0	-1	-1	0	0

Hence, $i_0 = 2$, $i_2 = 4$, $s_2 = 3$, $\eta_{s_2} = 1$, $i_1 = 3$, $s_1 = 2$, $\eta_{s_1} = -1$.
 Consequently, we obtain the elementary flip $E_{\bullet}^{(2)} \triangleleft E_{\bullet}^{(3)}$, where

$$E_{\bullet}^{(3)} = \{\emptyset \subset \{2\} \subset \{-2, 2\}, \{-2, 1, 2\}, E_{\text{all}}\}.$$

We can continue the algorithm for one last step, or alternatively note that $E_{\bullet}^{(3)} \triangleleft E'_{\bullet}$ is an elementary flip. \diamond

Proof of Lemma 7.16. 7.17 and Theorem 7.22 shows that it is enough to prove Lemma 7.16 whenever $E_{\bullet} \triangleleft E'_{\bullet}$ is an elementary flip. By Claim 7.19, we assume V_{\bullet} is a flag with $V_l \in \text{Gr}(l, n)_{E_l}^g$. Hence, $V_{\bullet} = uV_{\bullet}^{(0)}$, where $u \in O_g$, $V_l^{(0)} = \text{span}\{e_j : j \in F_l\}$, $F_l \subseteq \{1, \dots, n\}$ and $E_l = \{\eta_j : j \in F_l\}$. Since $E_{\bullet} \triangleleft E'_{\bullet}$ is an $(\eta, \tilde{\eta}, a, b)$ -elementary flip, there exist j_0 with $\eta_{j_0} = \eta$ and $j_0 \in F_a \setminus F_{a-1}$, and j_1 with $\eta_{j_1} = \tilde{\eta}$ and $j_1 \in F_b \setminus F_{b-1}$. The matrix $h = \text{Id} + \xi e_{j_1, j_0}$ lies in $B_{d_\varphi}(\text{Id}; 1)$ provided $\xi = \Theta(1) > 0$ is small enough. It preserves the spaces $V_{l_i}^{(0)}$ for $i < a$ or $i \geq b$. If $a \leq i < b$, then the blade corresponding to $hV_{l_i}^{(0)}$ is

$$(7.17) \quad (e_{j_0} + \xi e_{j_1}) \wedge \bigwedge_{j \in F_{l_i} \setminus \{j_0\}} e_j = \pm \bigwedge_{j \in F_{l_i}} e_j \pm \xi \bigwedge_{j \in F_{l_i} \cup \{j_1\} \setminus \{j_0\}} e_j.$$

The expression in Eq. (7.17) is the sum of two g_t -eigenvectors, with respective eigenvalues $\exp\left(t \sum_{\eta \in E_i} \eta\right)$ and $\exp\left(t \sum_{\eta \in E'_i} \eta\right)$. The second eigenvalue is exponentially larger than the first, and so

$$g_t h V_{l_i}^{(0)} \xrightarrow{t \rightarrow \infty} \text{span}\{e_j : j \in F_{l_i} \cup \{j_1\} \setminus \{j_0\}\}.$$

It follows that $(hV_l^{(0)})^{\rightarrow} \in \text{Gr}(l, n)_{\rightarrow E'_l}$ for every $l \in L$, and the first claim of Lemma 7.16 holds for $V_{\bullet}^{(0)}$. As for V_{\bullet} , we have that $h' := uhu^{-1} \in B_{d_\varphi}(\text{Id}; 1)$ satisfies $h'V_{\bullet} = uhV_{\bullet}^{(0)}$ and $h'V_{l_i} \in \text{Gr}(l, n)_{\rightarrow E'_l}$ for every $l \in L$. Moreover, $h'V_l$ lies in the compact set $O_g hV_l^{(0)} \subseteq V_l \in \text{Gr}(l, n)_{\rightarrow E'_l}$ for every $l \in L$. Claim 7.18 implies the desired for V_{\bullet} . \square

7.4. Adding Spaces to the Flag. In view of Observation 4.14, the following lemma will construct Alice's step whenever $f_{E_{\bullet}}^{h_\infty \Lambda}$ changes from time T_m to time T_{m+1} via a null vertex $* \rightarrow E$ at l . In these steps $\#A_{m+1} = 1$, that is, Alice's choice determines h_{m+1} . As in Subsection 11.1, Lemma 7.28 below will prove the existence of an element $h \in B_{d_\varphi}(\text{Id}; 1)$ with a certain behavior.

Recall that a sublattice $\Gamma \subseteq \Lambda$ is called *primitive* if there is no larger sublattice $\Gamma \subset \Gamma' \subseteq \Lambda$ with $\text{span}(\Gamma') = \text{span}(\Gamma)$.

Claim 7.24. *Let Λ be a lattice, and denote by $\lambda_1(\Lambda)$ the size of the shortest nonzero vector in Λ . For every $l = 0, \dots, n$, $V \in \text{Gr}(l, n)$ and $\varepsilon > 0$ there exists a primitive sublattice $\Gamma \subseteq \Lambda$ of rank l and covolume*

covol Γ such that

$$(7.18) \quad \log \operatorname{covol} \Gamma = \frac{l}{n} \log \operatorname{covol} \Lambda + O(\log \lambda_1(\Lambda) - \log \varepsilon),$$

and $\operatorname{span} \Gamma \in U_\varepsilon(V)$.

Remark 7.25. As always, the implicit constant in the big O notation in Eq. (7.18) depends on n , see Subsection 2.8.

Proof. It is sufficient to construct a non-primitive sublattice $\Gamma \subseteq \Lambda$. Indeed, given a non-primitive sublattice $\Gamma \subseteq \Lambda$ that satisfies the other condition of the lemma, the lattice $\Gamma' = \operatorname{span} \Gamma \cap \Lambda$ is primitive with the same span and smaller covolume. Since $\lambda_1(\Gamma') \geq \lambda_1(\Lambda)$, Minkowski's Theorem ([3, Chapter III, Theorem II]) implies that $\operatorname{covol}(\Gamma') = \Omega(\lambda_1(\Lambda)^{\operatorname{rank} \Gamma})$, hence Γ' satisfies the desired condition.

Up to multiplication by a scalar, assume that $\operatorname{covol} \Lambda = 1$. It is enough to prove the claim for $l = 1$. Indeed, fixing an orthogonal basis $(v_i)_{i=1}^l$ and applying the theorem to each of the linear spaces $\mathbb{R}v_i$ for $i = 1, \dots, l$ yields l rank-1 sublattices of Λ whose sum is the sought-for sublattice. After rotating Λ and V we may assume $V = \{x \in \mathbb{R}^n : x_1 = \dots = x_{n-1} = 0\}$. By Minkowski's Theorem, there exists a vector $v = (v_i)_{i=1}^n \in \Lambda \setminus \{0\}$ satisfying $|v_1|, \dots, |v_{n-1}| \leq \varepsilon_1$ and $|v_n| \leq \varepsilon_1^{1-n}$ for every $\varepsilon_1 > 0$. Note that if $\varepsilon_1 \leq \lambda_1(\Lambda)/2\sqrt{n}$, then

$$\|v\|^2 \leq v_1^2 + \dots + v_{n-1}^2 + v_n^2 \leq (n-1)\varepsilon_1^2 + v_n^2 \leq \lambda_1(\Lambda)^2/2 + v_n^2.$$

Consequently, $|v_n| = \Omega(\lambda_1(\Lambda))$, and hence

$$d_{\operatorname{Gr}(1,n)}(\mathbb{R}v, V)O(d_{\mathbb{R}^n}(v, V)/\|v\|) = O(\varepsilon/v_n) = O(\varepsilon/\lambda_1(\Lambda)).$$

Therefore, there is a choice of $\varepsilon_1 = \Theta(\varepsilon\lambda_1(\Lambda))$ for which $d_{\operatorname{Gr}(1,n)}(\mathbb{R}v, V) \leq \varepsilon$. Consequently, for $\Gamma = \mathbb{Z}v$ we have $\operatorname{covol} \Gamma = O((\varepsilon\lambda_1(\Lambda))^{1-n})$, as desired. \square

Remark 7.26. The bound in Claim 7.24 is not optimal. Though deriving the optimal bound may be of interest, it will not improve qualitatively the results of this paper.

Definition 7.27 (Relative Covolume). Let $\Gamma_0 \subseteq \Gamma_1 \subseteq \Gamma_2$ be lattices in \mathbb{R}^n of ranks $l_0 \leq l_1 \leq l_2$, respectively. Define the *relative covolume* of $\Gamma_0, \Gamma_1, \Gamma_2$ by

$$\log \operatorname{covol}(\Gamma_2/\Gamma_1/\Gamma_0) := \log \operatorname{covol} \Gamma_1 - \frac{l_2 - l_1}{l_2 - l_0} \log \operatorname{covol} \Gamma_0 - \frac{l_1 - l_0}{l_2 - l_0} \log \operatorname{covol} \Gamma_2.$$

The relative covolume is the signed height of $p_{\Gamma_1} = (\operatorname{rank} \Gamma_1, \log \operatorname{covol} \Gamma_1)$ above the interval $[p_{\Gamma_0}, p_{\Gamma_2}]$. It is also the normalized covolume of $\pi(\Gamma_1)$ in $\pi(\Gamma_2)$ where $\pi : \mathbb{R}^n \rightarrow \operatorname{span}(\Gamma_0)^\perp$, and we normalize so that $\operatorname{covol}(\pi(\Gamma_2)) = 1$. \heartsuit

Lemma 7.28 (Adding a lattice to a flag). *Let $1 > \varepsilon > 0$, let Λ be a lattice, let E_\bullet be a direction filtration and let Γ_\bullet be a filtration such that $L(E_\bullet) = L(\Gamma_\bullet)$ and*

$$\text{span } \Gamma_l \in \text{Gr}(l, n)_{\rightarrow E_l}^{\varepsilon, g}.$$

Let $l_0 < l_1 \in L(\Gamma_\bullet)$ be consecutive elements with $l_0 + 2 \leq l_1$, let $E_{l_0} \subset E \subset E_{l_1}$, and assume that $\partial^2 F_{\text{HN}}(\Gamma_{l_1}/\Gamma_{l_0})_{H, l} \leq C$ for every $l = 1, \dots, l_1 - l_0$. Then there exists $h \in H_{\text{span } \Gamma_\bullet \rightarrow E_\bullet}$ with $d_\varphi(h, \text{Id}) \leq 1$ and $h\Gamma_{l_0} \subseteq \Gamma \subseteq h\Gamma_{l_1}$ such that

$$\text{span } \Gamma \in \text{Gr}(l, n)_{\rightarrow E}^{O(\varepsilon), g}$$

and

$$(7.19) \quad \log \text{covol}(h\Gamma_{l_1}/\Gamma/h\Gamma_{l_0}) = O(C - \log \varepsilon).$$

Proof. Denote $V_l = \text{span } \Gamma_l$ for every $l \in L(\Gamma_\bullet)$. We will first find $V_{l_0} \subset V \subset V_{l_1}$ such that $V \in \text{Gr}(l, n)_{\rightarrow E}$. By Lemma 7.7, there exists $q_0 \in H^-$ such that $V_l = q_0 V_l^\rightarrow$ for all $l \in L(\Gamma_\bullet)$. In addition, $d_{H^-}(q_0, \text{Id}) = O(\varepsilon)$. By decomposing $V_{l_0}^\rightarrow$ and $V_{l_1}^\rightarrow$ into g_t -eigenspaces, we deduce that there exists $V' \in \text{Gr}(l, n)_E^g$ such that $V_{l_0}^\rightarrow \subset V' \subset V_{l_1}^\rightarrow$. Set $V = q_0 V'$ and note that $V^\rightarrow = V'$.

We will apply Claim 7.24 to $\Gamma_{l_1}/\Gamma_{l_0}$ and V/V_{l_0} in place Γ and V . Note that $\log \lambda_1(\Gamma_{l_1}/\Gamma_{l_0}) = O(C)$ since the height $F_{\text{HN}}(\Gamma_{l_1}/\Gamma_{l_0})_{H, 1}$ depends linearly on $\partial^2 F_{\text{HN}}(\Gamma_{l_1}/\Gamma_{l_0})_{H, l}$ for $l = 1, \dots, l_1 - l_0 - 1$, which are bounded by C . Then there exists $\Gamma' \subseteq \Gamma_{l_1}/\Gamma_{l_0}$ with $d(\text{span } \Gamma', V/V_{l_0}) \leq \varepsilon_1$ and

$$\log \text{covol } \Gamma' = O(C - \log \varepsilon_1) + \frac{l_{l_1} - l}{l_{l_1} - l_{l_0}} (\log \text{covol } \Gamma_{l_1} - \log \text{covol } \Gamma_{l_0}),$$

where ε_1 will be chosen later. Let Γ denote the inverse image of Γ' in Λ ; note that $\Gamma_{l_0} \subset \Gamma \subset \Gamma_{l_1}$. Then the covolume of Γ satisfies Eq. (7.19), provided that $\varepsilon_1 = \Theta(\varepsilon)$. Since Γ is ε_1 close to V and ε_1 can be chosen sufficiently small, there exists $m \in \text{SL}_n(\mathbb{R})$ with $m \text{span } \Gamma = V$ and $mV_l = V_l$ with $d_{\text{SL}_n(\mathbb{R})}(m, \text{Id}) = O(\varepsilon_1)$. If $\varepsilon_1 > 0$ is sufficiently small, by Lemma 2.3 we can write $m = q_1 h$ for $q_1 \in H^{-0}$ and $h \in H$ with $d_{\text{SL}_n(\mathbb{R})}(q_1, \text{Id}), d_{\text{SL}_n(\mathbb{R})}(h, \text{Id}) = O(\varepsilon_1)$. It follows that $h \in B_{d_\varphi}(\text{Id}; 1)$, $hV_l = q_1^{-1}V_l \in \text{Gr}(l, n)_{\rightarrow E_l}$, and $h\Gamma = q_1^{-1}V \in \text{Gr}(l, n)_{\rightarrow E_l}$. Since $d_{\text{SL}_n(\mathbb{R})}(q_0, \text{Id}) = O(\varepsilon)$, we see that

$$\begin{aligned} d_{\text{Gr}(l, n)}(\text{span } \Gamma, \text{Gr}(l, n)_E^g) &\leq d_{\text{Gr}(l, n)}(q_1^{-1}V, V) + d_{\text{Gr}(l, n)}(V, \text{Gr}(l, n)_E^g) \\ &\leq d_{\text{Gr}(l, n)}(q_1^{-1}V, V) + d_{\text{Gr}(l, n)}(q_0 V^\rightarrow, V^\rightarrow) \\ &\leq O(d_{\text{SL}_n(\mathbb{R})}(q_1, \text{Id}) + d_{\text{SL}_n(\mathbb{R})}(q_0, \text{Id})) = O(\varepsilon + \varepsilon_1). \end{aligned}$$

Note that we can choose $\varepsilon_1 = \Theta(\varepsilon)$. The result follows. \square

Remark 7.29. In Lemma 7.28 we can require h to be in any neighborhood of Id . Such a choice will only change the implicit constant.

8. ALICE'S STRATEGY

Fix a g -template f and a lattice $\Lambda \in X_n$ of covolume 1 for the rest of the section. Let $T > 0$ be a real number. Its choice will be formalized in Subsection 8.1. Recall that

$$Y_{\Lambda, f} := Y_{\Lambda, [f]} = \{h \in H : f^{h\Lambda} \sim f\} \subseteq H,$$

where $[f]$ is the set of g -template which are equivalent to f .

In this section we will construct Alice's strategy in the (T, g) -game of $Y_{\Lambda, f}$, see Subsection 4.4, and study some of its properties as explained in Subsection 4.5.

Definition 8.1. A vertex t_0 of f at l is called a vertex of type $E_- \rightarrow E_+$ if $f_{E, l}(t)$ changes from E_- to E_+ near t_0 , i.e.,

$$E_{\pm} = \lim_{\varepsilon \searrow 0} f(t_0 \pm \varepsilon).$$

♡

Informally, and somewhat imprecisely, Alice's strategy goes as follows: The initialization step is $h_0 = \text{Id}$, $T_0 = 0$. This defines $T_m = mT$. In the m -th step Alice chooses a set A_{m+1} of possible choices of h_{m+1} to ensure that the following three conditions are satisfied:

- (1) $L_f(T_{m+1}) = L(F_{\text{HN}}(g_{T_{m+1}} h_{m+1} \Lambda))$.
- (2) $\forall l \in L_f(T_{m+1}), F_{\text{HN}}(g_{T_{m+1}} h_{m+1} \Lambda)_{V, l} \in \text{Gr}(l, n)_{\rightarrow f_{E, l}(T_{m+1})}$.
- (3) $\forall l \in L_f(T_{m+1}), F_{\text{HN}}(g_{T_{m+1}} h_{m+1} \Lambda)_{H, l} = f_{H, l}(T_{m+1}) + O(1)$.

There will be three different types of steps. In every step in which $f_{E, \bullet}(T_m) = f_{E, \bullet}(T_{m+1})$, we use Lemma 7.13 to construct a set A_{m+1} of size $\Theta(\exp(T\delta(f_E(T_m))))$. These steps are called **Standard Steps**. If f has a null vertex of type $E \rightarrow *$ in $[T_m, T_{m+1}]$, we will use a Standard Step as well. If f has a null vertex of type $* \rightarrow E$ in $[T_m, T_{m+1}]$, we will use Lemma 7.28 to construct a singleton $A_{m+1} = \{h_{m+1}\}$. These steps are called **Adding Steps**. If f has a non-null vertex in $[T_m, T_{m+1}]$, we will use Lemma 7.16 to construct a singleton $A_{m+1} = \{h_{m+1}\}$. These steps are called **Changing Steps**.

There are several problems with this approach:

- (1) We are not able to make a lattice enter the filtration $F_{\text{HN}}(g_{T_m} h_m \Lambda)$ at the exact step we need.
- (2) There may be undesired elements in $F_{\text{HN}}(g_t h_m \Lambda)_{V, l}$.
- (3) The implicit constant in Condition 3 might increase with m .

8.0.1. *Analogy to Error Correction.* To illustrate the idea behind the solution of Problems (1) and (3), we consider an analogy. Suppose Alice has to drive an infinite route in a gridlike city. A navigation app provides to Alice a list of directions, of the form “After 3 junctions turn left”. Unfortunately, Alice always makes mistakes and never turns at the right junction. If the app does not update its instructions, Alice will eventually drift far from her original route. Therefore, after every turn, the app moderately updates the instructions.

The original route of Alice is analogous to the g -template f . The route she finally takes is the g -template $f^{h_\infty \Lambda}$. The turns Alice makes are analogous to the vertices of the g -template. The updates of the app are analogous to perturbing the g -template as described below, see Subsubsections 8.1.1 and 8.2.3.

The analogous concept to C -separated g -template is a list of instructions with bounded amount of turns every C kilometers. To add Bob to the analogy, we can think of Bob as a random noise, which is weaker the more Alice focuses, and Alice wants to focus the least, while still reaching her target.

8.0.2. *Overview of the Strategy.* Formally, a strategy is a choice of a move (h_0, T_0) in the initialization step and a map

$$\sigma : (h_0, T_0; A_1, h_1, \dots, A_m, h_m) \mapsto A_{m+1},$$

which assigns for every $m \geq 0$ and history of moves up to Alice’s m -th step (excluded), the move of Alice in the current step m . To compute the value of the game it is enough to define σ_{Alice} for histories that are consistent with σ_{Alice} up to step m .

We will use auxiliary information, that is, Alice will maintain a filtration Γ_\bullet^m of Λ and a locally constant error function $\boldsymbol{\pi}^m = (\pi_l^m)_{l=0}^n$ with $\pi_l^m : U_l(f) \rightarrow \mathbb{R}$. The strategy we will define will be a map

$$\sigma_{\text{Alice}} : (h_0, T_0; A_1, h_1, \dots, A_m, h_m, \Gamma_\bullet^m, \boldsymbol{\pi}^m) \mapsto (A_{m+1}, \Gamma_\bullet^{m+1}, \boldsymbol{\pi}^{m+1}).$$

For each history, the computation of A_{m+1} , Γ_\bullet^{m+1} , and $\boldsymbol{\pi}^{m+1}$ will have one of several forms. For each step we will decide whether it is a **Standard Step**, an **Adding Step**, or a **Changing Step**. In addition, we have three ways to update Γ_\bullet^m to Γ_\bullet^{m+1} : we can **preserve** it, and then $\Gamma_\bullet^{m+1} = \Gamma_\bullet^m$; we can **add** a lattice to it, and then $\Gamma_\bullet^{m+1} \supset \Gamma_\bullet^m$; or we can remove a lattice from it, and then $\Gamma_\bullet^{m+1} \subset \Gamma_\bullet^m$. Further, there are two ways to update $\boldsymbol{\pi}^m$: we can either **anchor** it (see Subsubsection 8.2.3) or not anchor it, and then $\boldsymbol{\pi}^{m+1} = \boldsymbol{\pi}^m$.

8.1. **Preliminaries.** Fix $C_2 \ggg C_1 \ggg T$ (see Subsection 2.8 for the definition of \ggg). The strategy will depend on T, C_1, C_2 . Recall that

f is a g -template fixed for the entire section. Let f' be the g -template f shifted so that $f'_{H,l}(0) = 0$ for every l with $0 \leq l \leq n$. To shift f to f' we use the independent shift sequence with parameters C and $(0)_{J \in \mathcal{G}_f}$, for a large enough C . By Lemma 5.15, the g -templates f, f' are equivalent. By Lemma 5.9, there is a C_2 -significant and C_2 -separated g -template f'' that is equivalent to f' . By the proof of Lemma 5.9, we get that f'' is a shift of f' , hence $f''_{H,l}(0) = 0$ for every l with $0 \leq l \leq n$. Since we care about f only up to equivalence, we may assume that f is C_2 -significant, C_2 -separated, and $f_{H,l}(0) = 0$ for every l with $0 \leq l \leq n$.

8.1.1. *Adjusting f .* Define the shift sequence

$$(8.1) \quad \rho := (2l(n-l)C_1 + C_1)_{l=1}^{n-1}.$$

For every sequence of locally constant functions $\pi = (\pi_l)_{l=1}^{n-1}$, $\pi_l : U_l(f) \rightarrow [-C_1, C_1]$, we get that $\rho + \pi$ is an independent shift sequence with parameters $2C_1, C_1 + \pi$.

By Lemma 5.30, since C_2 is large enough with respect to C_1 , for every nontriviality interval J of f at l there exists a nontriviality interval J' of f^ρ at l with $J' \subseteq J$. In addition, f^ρ is also $C_2/2$ -separated and $C_2/2$ -significant.

8.1.2. *Preliminaries on Directions.*

Definition 8.2 (Final direction). Let $E_{-1} \subset E_0 \subset E_1$ be submultisets of E_{all} . We say that E_0 is *final* in $E_{-1} \subset E_1$ if it is the maximum with respect to the partial order \leq of all E of size $\#E_0$ which satisfy $E_{-1} \subset E \subset E_1$. \heartsuit

Observation 8.3. Let E_\bullet be a direction filtration and $E_l \subset E_{l'}$ be two consecutive elements in E_\bullet . Let E be a multiset between them, i.e., $E_l \subset E \subset E_{l'}$. Then $\delta(E_\bullet) \geq \delta(E_\bullet \cup \{E\})$, with equality if and only if E is final in $E_l \subset E_{l'}$. \clubsuit

Definition 8.4 (Scalar pairs). We say that a pair of multisets $E \subset E'$ is a *scalar pair* if $E' - E$ has no two distinct elements (i.e., has one element, possibly with multiplicity). \heartsuit

Observation 8.5. If $E \subset E'$ is a scalar pair, then for every l with $\#E < l < \#E'$ there is a unique $E \subset E'' \subset E'$ of size l , and it is final in $E \subset E'$. \clubsuit

We will treat scalar pairs differently because of the following property they enjoy (which is the reason for their name):

Observation 8.6 (Scalar action). Let $E \subset E'$ be a scalar pair, with $l := \#E, l' := \#E'$. Let $V \in \text{Gr}(l, n)_E^g$ and $V' \in \text{Gr}(l', n)_{E'}^g$ be two linear spaces with $V \subset V'$. Then the induced action $g_t \curvearrowright V'/V$ is multiplication by a scalar. \clubsuit

Lemma 8.7. Let $E_{-1} \subset E_0 \subset E_1$ be multisets of sizes $l_{-1} < l_0 < l_1$ respectively, such that E_0 is final in $E \subset E_1$. Then

$$\eta_{E_0} \geq \frac{l_0 - l_{-1}}{l_1 - l_{-1}} \eta_{E_1} + \frac{l_1 - l_0}{l_1 - l_{-1}} \eta_{E_{-1}},$$

and equality holds if and only if $E_{-1} \subset E_1$ is a scalar pair.

Proof. Denote the values with multiplicity of $E_1 - E_{-1}$ by

$$\eta'_1 \leq \eta'_2 \leq \dots \leq \eta'_{l_1 - l_{-1}}.$$

This sequence is constant if and only if $E_{-1} \subset E_1$ is a scalar pair. It follows that the sequence $l'_0 \mapsto \sum_{i=1}^{l'_0 - l_{-1}} \eta'_i$ for $l_{-1} \leq l'_0 \leq l_1$ is convex, and is linear if and only if $E_{-1} \subset E_1$ is a scalar pair. The result follows since $E_0 = E_{-1} + \{\eta'_i : i = l_0 - l_{-1} + 1, \dots, l_1 - l_{-1}\}$. \square

Lemma 8.8. Let f be a g -template, let J be a nontriviality interval of f at l_0 and let t_0 one of its endpoints. Let $E_\bullet = \lim_{J \ni t \rightarrow t_0} f_{E_\bullet}(t)$, and let l_{-1}, l_1 be the adjacent elements in $L(E_\bullet)$ to l_0 satisfying $l_{-1} < l_0 < l_1$. Then $E_{l_{-1}} \subset E_{l_1}$ is not a scalar pair.

Proof. Let $J_0 \subseteq J$ be an open interval in which f has no vertices and such that t_0 is an endpoint of J_0 . By the definition of E_\bullet , for every $t \in J_0$ we have $f_{E_\bullet}(t) = E_\bullet$. Consequently, $t \mapsto f_{H,l}(t)$ is linear in J_0 for every l with $0 \leq l \leq n$, and so is $t \mapsto \partial^2 f_{H,l_0}(t)$. Since $t_0 \notin U_{l_0}(f)$, it follows that $\partial^2 f_{H,l_0}(t_0) = 0$, while for every $t \in J_0$ we have $\partial^2 f_{H,l_0}(t) \neq 0$. Consequently, $\frac{d}{dt} \partial^2 f_{H,l_0}(t)$ does not vanish for $t \in J_0$.

From Observation 5.4 we get that

$$(8.2) \quad \begin{aligned} \partial^2 f_{H,l_0}(t) &= \frac{1}{l_0 - l_{-1}} f_{H,l_{-1}} + \frac{1}{l_1 - l_0} f_{H,l_1} - \frac{l_1 - l_{-1}}{(l_1 - l_0)(l_0 - l_{-1})} f_{H,l_0}. \end{aligned}$$

Differentiating Eq. (8.2) we obtain

$$0 \neq \frac{d}{dt} \partial^2 f_{H,l_0}(t) = \frac{1}{l_0 - l_{-1}} \eta_{E_{l_{-1}}} + \frac{1}{l_1 - l_0} \eta_{E_{l_1}} - \frac{l_1 - l_{-1}}{(l_1 - l_0)(l_0 - l_{-1})} \eta_{E_{l_0}},$$

which together with Lemma 8.7 implies that $E_{l_{-1}} \subset E_{l_1}$ is not a scalar pair. \square

8.1.3. *Constants Order of Magnitude.* Our computations will use several universal constants, which depend on E_{all} but not on T , f , and Λ , and should obey certain relations. Specifically, we require that

$$(8.3) \quad \gamma_{10} \gg \gamma_9 \gg \gamma_8 \gg \gamma_7 \gg \gamma_6 \gg \gamma_5 \gg \gamma_4 \gg \gamma_3, \gamma'_3 \gg \gamma_2 \gg \gamma_1, \gamma'_1 \gg 0.$$

Note that Eq. (8.3) states that both $\gamma_4 \gg \gamma_3 \gg \gamma_2$ and $\gamma_4 \gg \gamma'_3 \gg \gamma_2$ hold, and a similar interpretation applies to γ_1, γ'_1 . We will choose $\varepsilon > 0$ small enough to satisfy several conditions that will be detailed throughout the proof. In particular, ε is small enough so that the conclusions of Lemma 7.13, Theorem 2.7, and Lemma 6.3 are satisfied.

8.1.4. *Convenient Assumption.* Throughout the section we evaluate the category flow of various g -templates on $\mathbb{N}_0 T = \{T_m : m \geq 0\}$. The category flows might be undefined at T_m for some $m \geq 0$. To avoid these technicalities we will assume that all g -templates have no vertices on $\mathbb{N}_0 T$, i.e., they define direction filtrations on $\mathbb{N}_0 T$. To justify this assumption we describe two operations that guarantee that there will be no such vertices in all templates we ever use.

First, translate f by a random small number to obtain a g -template f' with no vertices on $\mathbb{N}_0 T$. We can therefore assume w.l.o.g. that f has no vertices on $\mathbb{N}_0 T$. The other g -templates we will encounter are shifts of f , namely f^ρ for some values of ρ . Since f has no non-null vertices on $\mathbb{N}_0 T$, it follows that f^ρ has no such vertices as well. To ensure f^ρ has no null vertices on $\mathbb{N}_0 T$, we perturb ρ . The perturbations affect Subsubsection 8.1.1 below and the anchoring operation that we will define in Subsubsection 8.2.3. This procedure will not affect the arguments.

8.2. Setup for the Strategy σ_{Alice} .

8.2.1. *The State.* In each step we keep a state that consists of

- (S.1) $h_m \in H$ and $T_m = Tm$.
- (S.2) An error function $\boldsymbol{\pi}^m = (\pi_l^m)_{l=1}^{n-1}$, where $\pi_l^m : U_l(f) \rightarrow [-C_1, C_1]$ is locally constant.
- (S.3) A filtration Γ_\bullet^m of Λ .

Denote $\tilde{\Lambda}^m := g_{T_m} h_m \Lambda$ and $\tilde{\Gamma}_l^m := g_{T_m} h_m \Gamma_l^m$ for every $l \in L(\Gamma_\bullet^m)$. We require the data in the state to satisfy the following conditions:

- (S.4) $L(\Gamma_\bullet^m) \subseteq L_f(T_m)$. We will see in Remark 8.13 that a stronger condition on $L(\Gamma_\bullet^m)$ actually holds.
- (S.5) For every $l \in L(\Gamma_\bullet^m)$ we have

$$(8.4) \quad \text{span } \tilde{\Gamma}_l^m \in \text{Gr}(l, n)_{\rightarrow f_{E,l}(T_m)}^{\varepsilon, g}$$

and

$$(8.5) \quad \log \operatorname{covol}(\tilde{\Gamma}_l^m) = f_{H,l}^{\rho+\pi^m}(T_m) + O(1),$$

where ρ is defined as in Eq. (8.1).

(S.6) For every two consecutive $l < l' \in L(\Gamma_\bullet^m)$, consider lattices $\tilde{\Gamma}^m$ with $\tilde{\Gamma}_l^m \subset \tilde{\Gamma}^m \subset \tilde{\Gamma}_{l'}^m$ that project to elements in $F_{\text{HN}}(\tilde{\Gamma}_{l'}^m/\tilde{\Gamma}_l^m)$. We call such $\tilde{\Gamma}^m$ a *noise lattice*. We require that the noise lattices are insignificant, in the sense that

$$(8.6) \quad \log \operatorname{covol}(\tilde{\Gamma}_{l'}^m/\tilde{\Gamma}^m/\tilde{\Gamma}_l^m) = O(T).$$

The choice of $\tilde{\Gamma}^m$ is equivalent to the choice of Γ such that $\Gamma_l^m \subset \Gamma \subset \Gamma_{l'}^m$, where $\tilde{\Gamma}^m = g_{T_m} h_m \Gamma$.

The strategy selects a finite set A_{m+1} of possible matrices $h_{m+1} \in B_{d_\varphi}(h_m; (1 - \exp(-\alpha_\varphi T))^{1/\alpha_\varphi} \exp(-T_m))$, where α_φ is a constant for which $d_\varphi^{\alpha_\varphi}$ is a metric, see Subsection 4.2. The set A_{m+1} is such that $d_\varphi(h_{m+1}, h'_{m+1}) \geq 3^{1/\alpha_\varphi} \exp(-T_{m+1})$ for every pair of distinct elements $h_{m+1}, h'_{m+1} \in A_{m+1}$.

Instead of A_{m+1} , we can consider the set

$$(8.7) \quad \tilde{A}_{m+1} := g_{T_m} A_{m+1} h_m^{-1} g_{-T_m},$$

which satisfies

- $\tilde{A}_m \subseteq B_T := B_{d_\varphi}(\text{Id}; (1 - \exp(-\alpha_\varphi T))^{1/\alpha_\varphi})$.
- $d_\varphi(h_{m+1}, h'_{m+1}) \geq 3^{1/\alpha_\varphi} \exp(-T)$ for every pair of distinct $\tilde{h}_{m+1}, \tilde{h}'_{m+1} \in \tilde{A}_{m+1}$.

This translation is possible due to the right-invariance of the semi-metric d_φ . The set B_T is chosen in view of Eq. (4.9) in the definition of the dimension game. It has the properties that, for every $h'_{m+1} \in A_m$,

$$(8.8) \quad B_{d_\varphi}(h_{m+1}; \exp(-T_{m+1})) \subseteq B_{d_\varphi}(h_m; \exp(-T_m)).$$

Sometimes the set A_{m+1} will consist of a single element. In that case, choosing A_{m+1} is equivalent to choosing the unique element h_{m+1} in it.

Definition 8.9 (Alignment of noise lattices). Let $l_0 < l_1$ be two consecutive elements in $L(\Gamma_\bullet^m)$. We say that a noise lattice Γ with $\Gamma_{l_0}^m \subset \Gamma \subset \Gamma_{l_1}^m$ is *E-aligned* at step m if $\tilde{\Gamma}^m = g_{T_m} h_m \Gamma$ satisfies $\operatorname{span} \tilde{\Gamma}^m \in U_\varepsilon(\operatorname{Gr}(l, n)_E)$, for some $f_{E, l_0}(T_m) \subset E \subset f_{E, l_1}(T_m)$, where $\operatorname{rank} \Gamma = l$. We say that Γ is *final* at step m if it is *E* aligned and *E* is final in $f_{E, l_0}(T_m) \subset f_{E, l_1}(T_m)$. \heartsuit

8.2.2. *Initialization.* Sets $h_0 = \text{Id}$, $T_0 = 0$, and $\pi^0 = 0$ everywhere. Since $f_{H,l}(0) = 0$ for every l with $0 \leq l \leq n$, it follows that $L_{f\rho+\pi^0}(t) = \{0, n\}$, and so we choose $\Gamma_\bullet^0 := (\{0\} \subseteq \Lambda)$.

8.2.3. *Updating π^m and Γ_\bullet^m .* In this subsection we describe how π^m, Γ_\bullet^m are updated at each step.

There are three different ways to update Γ_\bullet^m to Γ_\bullet^{m+1} : **Preserve**, **Add**, and **Remove**, which respectively sets $\Gamma_\bullet^{m+1} = \Gamma_\bullet^m$, adds a single element to Γ_\bullet^m to get Γ_\bullet^{m+1} , and removes a single element from Γ_\bullet^m to get Γ_\bullet^{m+1} . We will use preserve and remove at Standard Steps, preserve at Changing Steps, and add at Adding Steps.

Definition 8.10 (Anchoring). By default, $\pi^{m+1} = \pi^m$. After every adding or changing step, we will use the **Anchor** operation. For each $1 \leq l \leq n-1$, to anchor π_l^{m+1} is to fix $\pi_l^{m+1}(T_{m+1})$ so that equality is attained in the formula

$$\log \text{covol}(\tilde{\Gamma}_l^{m+1}) = f_{E,l}(T_{m+1}) + \rho_l(T_{m+1}) + \pi_l^{m+1}(T_{m+1}),$$

i.e.,

$$\pi_l^{m+1}(T_{m+1}) := \log \text{covol}(\tilde{\Gamma}_l^{m+1}) - f_{E,l}(T_{m+1}) - \rho_l(T_{m+1}).$$

Note that this operation changes π_l^m on the entire connected component of T_m on $U_l(f)$. This is the only operation that changes π^m . If we do not anchor π_l^{m+1} , then $\pi_l^{m+1} = \pi_l^m$. \heartsuit

8.3. **Geometric Picture.** We will describe here geometric features of the behavior of the (T, g) -game.

Assume the game has been played, and has generated a sequence h_0, A_1, h_1, \dots . For every $0 \leq m_0 \leq m_1 < \infty$ such that $m_0 < \infty$, denote

$$\tilde{h}^{m_0 \rightarrow m_1} := g_{T_{m_0}} h_{m_1} h_{m_0}^{-1} g_{-T_{m_0}}.$$

Eq. (8.8) implies that $d_\varphi(h_{m_0}, h_{m_1}) \leq \exp(-T_{m_0})$ and for every $m_0 \leq m_1 \leq m_2$ we have $d_\varphi(\tilde{h}^{m_0 \rightarrow m_1}, \tilde{h}^{m_0 \rightarrow m_2}) \leq \exp(-(T_{m_1} - T_{m_0}))$. Hence, the limit $\lim_{m \rightarrow \infty} h_m$ exists. Denote $h_\infty := \lim_{m \rightarrow \infty} h_m$, and for every $0 \leq m_0 < \infty$ denote

$$\tilde{h}^{m_0 \rightarrow \infty} := g_{T_{m_0}} h_\infty h_{m_0}^{-1} g_{-T_{m_0}}.$$

For every $m \geq 0$ denote $\tilde{h}^{m+1} = \tilde{h}^{m \rightarrow m+1}$. Note that

$$(8.9) \quad h_m = g_{-T_m} \tilde{h}^m g_T \tilde{h}^{m-1} g_T \cdots g_T \tilde{h}_1 g_{T_0}.$$

Remark 8.11 (Relative notation). Some objects are defined from the absolute perspective of time 0, like Λ , Γ_{\bullet}^m , h_m , and A_m . Sometimes it is more convenient to consider these objects from the perspective of time T_m . Whenever we do so, we denote the relative objects with $\tilde{}$ (tilde). For example, $\tilde{\Gamma}_{\bullet}^m = g_{T_m} h_m \Gamma_{\bullet}^m$, $\tilde{h}_{m+1} = g_{T_m} h_{m+1} h_m^{-1} g_{-T_m}$, and $\tilde{A}_{m+1} = g_{T_m} A_{m+1} h_m^{-1} g_{-T_m}$ are the relative versions of Γ_{\bullet}^m , h_m , and A_m , respectively. The advantage of the relative perspective (with tilde) is that the formulas do not involve T_m , and are time invariant. The advantage of the absolute perspective is the locally constant nature of various definitions, such as Γ_{\bullet}^m .

The following is a corollary of Lemma 7.7.

Corollary 8.12. *There exists an $\varepsilon_0 > 0$ small enough such that the following holds. For every ε with $0 < \varepsilon < \varepsilon_0$, l with $0 \leq l \leq n$, two integers $0 \leq m_0 \leq m_1$, and $V \in \text{Gr}(l, n)$ such that $\tilde{V}^m := g_{T_m} h_m V \in \text{Gr}(l, n)_{\rightarrow E}$ for all m with $m_0 \leq m \leq m_1$ and $\tilde{V}^{m_0} \in \text{Gr}(l, n)_{\rightarrow E}^{\varepsilon, g}$, we have*

$$\log d_H(\tilde{V}^{m_1}, \text{Gr}(l, n)_E^g) = -\Omega(-\log \varepsilon + (T_{m_1} - T_{m_0})),$$

i.e., $d_H(\tilde{V}^{m_1}, \text{Gr}(l, n)_E^g)$ decays exponentially in $m_1 - m_0$. In addition, for all l -blades $w \in \bigwedge^l V$ we have

$$(8.10) \quad \log \|g_{T_{m_1}} h_{m_1} w\| = \log \|g_{T_{m_0}} h_{m_0} w\| + (m_1 - m_0)T\eta_E + O(1).$$

Proof. Since $\tilde{V}^{m_0} \in \text{Gr}(l, n)_{\rightarrow E}^{\varepsilon, g}$, by Lemma 7.7 there exists $q_0 \in H^-$ such that $d_{H^-}(q_0, \text{Id}) = O(\varepsilon)$ and $\tilde{V}^{m_0} = q_0(\tilde{V}^{m_0})^{\rightarrow}$. Apply Lemma 7.10 to \tilde{V}^{m_0} , q_0 , and E . We obtain two open sets $U_1 \subseteq H_{\tilde{V}^{m_0} \rightarrow E}$, $U_2 \subseteq H((\tilde{V}^{m_0})^{\rightarrow})$ and two maps

$$\check{h} : U_1 \xrightarrow{\sim} U_2, \text{ and } \check{q} : U_2 \rightarrow H^{-0},$$

where \check{h} is a diffeomorphism such that for every $h \in U_1$ we have

$$(8.11) \quad h q_0 = \check{q}(h) \check{h}(h).$$

Moreover, since $d_{H^-}(q_0, \text{Id})$ is arbitrarily small, U_1 and U_2 are arbitrarily large. Specifically, for some $\varepsilon_0 > 0$ small enough which depends only on n , we get that $B_{d_\varphi}(\text{Id}; 1) \subseteq U_1$. We will apply these functions

to $\tilde{h}^{m_0 \rightarrow m_1}$:

$$\begin{aligned}
\tilde{V}^{m_1} &= g_{(m_1-m_0)T} \tilde{h}^{m_0 \rightarrow m_1} \tilde{V}^{m_1} = g_{(m_1-m_0)T} \tilde{h}^{m_0 \rightarrow m_1} q_0 \cdot (\tilde{V}^{m_1})^\rightarrow \\
&= g_{(m_1-m_0)T} \check{\mathfrak{q}}(\tilde{h}^{m_0 \rightarrow m_1}) \check{\mathfrak{h}}(\tilde{h}^{m_0 \rightarrow m_1}) \cdot (\tilde{V}^{m_1})^\rightarrow \\
&= g_{(m_1-m_0)T} \check{\mathfrak{q}}(\tilde{h}^{m_0 \rightarrow m_1}) \cdot (\tilde{V}^{m_1})^\rightarrow \\
&= g_{(m_1-m_0)T} \check{\mathfrak{q}}(\tilde{h}^{m_0 \rightarrow m_1}) g_{-(m_1-m_0)T} \cdot (\tilde{V}^{m_1})^\rightarrow.
\end{aligned}$$

Since

- $\check{\mathfrak{q}}$ depends differentiably on q_0 ,
- if $q_0 = \text{Id}$ then $\check{\mathfrak{q}} \equiv \text{Id}$,
- $d_{H^-}(q_0, \text{Id}) = O(\varepsilon)$,

it follows that for all $h \in B_{d_\varphi}(\text{Id}; 1)$ we have $d_{H^-0}(\check{\mathfrak{q}}(h), \text{Id}) = O(\varepsilon)$ for every $h \in B_{d_\varphi}(\text{Id}; 1)$. In particular, $d_{H^-0}(\check{\mathfrak{q}}(\tilde{h}^{m_0 \rightarrow m_1}), \text{Id}) = O(\varepsilon)$. Write $\check{\mathfrak{q}}(\tilde{h}^{m_0 \rightarrow m_1}) = q_1^- q_1^0$ where $q_1^- \in H^-$, $q_1^0 \in H^0$. We deduce that $d_{H^-1}(q_1^-, \text{Id}) = O(\varepsilon)$. Altogether, we get

$$\begin{aligned}
&d_{\text{Gr}(l,n)}(\tilde{V}^{m_1}, \text{Gr}(l, n)_E^g) \\
&\leq d_{\text{Gr}(l,n)}(g_{(m_1-m_0)T} \check{\mathfrak{q}}(\tilde{h}^{m_0 \rightarrow m_1}) g_{-(m_1-m_0)T} \cdot (\tilde{V}^{m_1})^\rightarrow, \text{Gr}(l, n)_E^g) \\
&= d_{\text{Gr}(l,n)}(g_{(m_1-m_0)T} q_1^- q_1^0 g_{-(m_1-m_0)T} \cdot (\tilde{V}^{m_1})^\rightarrow, \text{Gr}(l, n)_E^g) \\
&= d_{\text{Gr}(l,n)}(g_{(m_1-m_0)T} q_1^- g_{-(m_1-m_0)T} q_1^0 \cdot (\tilde{V}^{m_1})^\rightarrow, \text{Gr}(l, n)_E^g) \\
&\leq d_{\text{Gr}(l,n)}(g_{(m_1-m_0)T} q_1^- g_{-(m_1-m_0)T} q_1^0 \cdot (\tilde{V}^{m_1})^\rightarrow, q_1^0 \cdot (\tilde{V}^{m_1})^\rightarrow) \\
&= O(d_{H^-}(g_{(m_1-m_0)T} q_1^- g_{-(m_1-m_0)T}, \text{Id})).
\end{aligned}$$

The distance result follows from the uniform contraction rate of conjugation of g_t on H^- .

To show Eq. (8.10), note that

$$g_{T_{m_1}} h_{m_1} w = g_{(m_1-m_0)T} \tilde{h}^{m_0 \rightarrow m_1} g_{T_{m_0}} h_{m_0} w.$$

By Eq. (8.11), we can factor $\tilde{h}^{m_0 \rightarrow m_1}$ as

$$\tilde{h}^{m_0 \rightarrow m_1} = \check{\mathfrak{q}}(\tilde{h}^{m_0 \rightarrow m_1}) \check{\mathfrak{h}}(\tilde{h}^{m_0 \rightarrow m_1}) q_0^{-1}.$$

Further by the definition of q_0 ,

$$q_0^{-1} \tilde{V}^{m_0} = q_0^{-1} g_{T_{m_0}} h_{m_0} V \in \text{Gr}(l, n)_E^g,$$

and hence $q_0^{-1}g_{T_{m_0}}h_{m_0}w \in \bigwedge^l(q_0^{-1}g_{T_{m_0}}h_{m_0}V)$ is a g_t -eigenvector of $\bigwedge^l \mathbb{R}^n$, with eigenvalue $\exp(\eta_E t)$. Consequently,

$$\begin{aligned} \log \|g_{T_{m_1}}h_{m_1}w\| &= \log \|g_{(m_1-m_0)T}\tilde{h}^{m_0 \rightarrow m_1}g_{T_{m_0}}h_{m_0}w\| \\ &= \log \|g_{(m_1-m_0)T}\check{\mathfrak{q}}(\tilde{h}^{m_0 \rightarrow m_1})\check{\mathfrak{h}}(\tilde{h}^{m_0 \rightarrow m_1})q_0^{-1}g_{T_{m_0}}h_{m_0}w\| \\ &= \log \|g_{(m_1-m_0)T}\check{\mathfrak{h}}(\tilde{h}^{m_0 \rightarrow m_1})q_0^{-1}g_{T_{m_0}}h_{m_0}w\| + O(1). \end{aligned}$$

The last equality holds because

$$d_{H^{-0}}(g_{(m_1-m_0)T}\check{\mathfrak{q}}(\tilde{h}^{m_0 \rightarrow m_1})g_{-(m_1-m_0)T}, \text{Id}) = O(1).$$

Since $\check{\mathfrak{h}}(\tilde{h}^{m_0 \rightarrow m_1})$ preserves $(\tilde{V}^{m_0})^\rightarrow = q_0^{-1}g_{T_{m_0}}h_{m_0}V$, it stabilizes the vector $q_0^{-1}g_{T_{m_0}}h_{m_0}w$ and hence

$$\begin{aligned} \log \|g_{T_{m_1}}h_{m_1}w\| &= \log \|g_{(m_1-m_0)T}q_0^{-1}g_{T_{m_0}}h_{m_0}w\| + O(1) \\ &= (m_1 - m_0)T\eta_E + \log \|q_0^{-1}g_{T_{m_0}}h_{m_0}w\| + O(1) \\ &= (m_1 - m_0)T\eta_E + \log \|g_{T_{m_0}}h_{m_0}w\| + O(1), \end{aligned}$$

as desired. \square

8.4. The Different Steps. Here we describe the three types of steps of σ_{Alice} : adding, changing and standard. For each type we provide the condition that should be satisfied so that step m has this type, the way A_{m+1} is chosen, and the way π^m and Γ_\bullet^m are chosen.

8.4.1. *Adding Step.* To determine whether step m is an Adding Step we do as follows. Consider a null vertex t_0 of f at l of the form $* \rightarrow E$. By Lemma 5.30, for every independent shift sequence ρ' with parameters $2C_1, (\nu_J)_{J \in \mathcal{G}_f} \in [0, 2C_1]^{\mathcal{G}_f}$ there is a unique vertex of $f^{\rho'}$ that corresponds to t_0 , denoted $t_0^{\rho'}$. Since $t_0^{\rho'}$ depends continuously on ρ , the following set is an interval:

$$[a_{t_0}, b_{t_0}] := \left\{ t_0^{\rho'} : \begin{array}{l} \rho' \text{ is an independent shift sequence} \\ \text{with parameters } 2C_1, (\nu_J)_{J \in \mathcal{G}_f} \in [0, 2C_1]^{\mathcal{G}_f} \end{array} \right\}.$$

Since f is C_1 -separated, $b_{t_0} - a_{t_0} = \Theta(C_1)$. Denote $m_{t_0}^a := \lfloor a_{t_0}/T \rfloor$ and $m_{t_0}^b := \lfloor b_{t_0}/T \rfloor$. By Lemma 5.30, for every such ρ' the g -template $f^{\rho'}$ has exactly one vertex in $[a_{t_0}, b_{t_0}]$, namely $t_0^{\rho'}$. Denote by m_{t_0} the first $m' \in [m_{t_0}^a, m_{t_0}^b]$ for which

$$L_f^{\rho' + \pi^{m_{t_0}^a}}(T_{m'+1}) = L_f^{\rho' + \pi^{m_{t_0}^a}}(T_{m'}) \cup \{l\}.$$

As we will see below, there will be only Standard Steps at $m \in [m_{t_0}^a, m_{t_0}^b] \setminus \{m_{t_0}\}$, and Standard Steps do not change π^m . Hence, m_{t_0} is the unique

$m' \in [m_{t_0}^a, m_{t_0}^b]$ for which

$$L_f^{\rho+\pi^{m_{t_0}^a}}(T_{m'}) = L_f^{\rho+\pi^{m_{t_0}^a}}(T_{m_{t_0}^a}),$$

and

$$L_f^{\rho+\pi^{m_{t_0}^a}}(T_{m'+1}) = L_f^{\rho+\pi^{m_{t_0}^a}}(T_{m_{t_0}^a}) \sqcup \{l\} = L_f^{\rho+\pi^{m_{t_0}^a}}(T_{m_{t_0}^b+1}).$$

Condition-Adding: Step m is an **Adding Step** if $m \in [m_{t_0}^a, m_{t_0}^b]$ and $m = m_{t_0}$ for some null vertex t_0 of f of type $* \rightarrow E$. Note that there may exist a unique such null vertex t_0 with $m \in [m_{t_0}^a, m_{t_0}^b]$.

The following condition is equivalent to Condition-Adding:

Condition-Adding': Step m is an **Adding Step** if

$$L(\Gamma_{\bullet}^m) = L_{f\rho+\pi^m}(T_m) \subsetneq L_{f\rho+\pi^m}(T_{m+1}).$$

Construction of A_{m+1} : When m is an Adding Step, A_{m+1} consists of a single element h_{m+1} . Let us explain how \tilde{h}_{m+1} is calculated. Denote by $l^- < l < l^+$ the consecutive elements of l in $L_{T_{m+1}}(f^{\rho+\pi^m})$. We will use Lemma 7.28 to obtain a lattice Γ with $\tilde{\Gamma}_{l^-}^m \subset \Gamma \subset \tilde{\Gamma}_{l^+}^m$ and an element $\tilde{h}_{m+1} \in B_T \cap H_{\text{span } \Gamma_{\bullet}^m \rightarrow f_{E,l}^{\rho+\pi^m}}$ such that

$$(8.12) \quad \tilde{h}_{m+1} \text{span } \Gamma \in \text{Gr}(l, n)_{\rightarrow f_{E,l}^{\rho+\pi^m}}^{O(\epsilon),g}(T_{m+1}).$$

To use Lemma 7.28 we have to obtain some bound on the noise lattices lying between Γ_l^m and $\Gamma_{l'}^m$. In Corollary 8.22 below we will find $\gamma_4 > 0$ depending only on E_{all} such that every noise lattice Γ' at an Adding Step with $\tilde{\Gamma}_l^m \subset \Gamma' \subset \Gamma_{l'}^m$ has $\log \text{covol}(\tilde{\Gamma}_{l'}^m/\Gamma'/\tilde{\Gamma}_l^m) \leq \gamma_4 T$.

By the bound on the noise lattices and Lemma 7.28 we have

$$\left| \log \text{covol}(\tilde{\Gamma}_{l'}^m/\Gamma'/\tilde{\Gamma}_l^m) \right| \leq \gamma_5 T,$$

for some $\gamma_5 > 0$ depending only on E_{all} .

Updating the auxiliary parameters: We add $g_{-T_m} h_m^{-1} \Gamma$ to the filtration Γ_{\bullet}^m , anchor π_l^{m+1} , and get $|\pi_l^{m+1}(T_{m+1})| \leq \gamma_5 T$.

Remark 8.13. We will compare the two sets $L(\Gamma_{\bullet}^m)$ and $L_{f\rho+\pi^m}(T_m)$ for every step m . By Condition-Adding, an index is added to $L(\Gamma_{\bullet}^m)$ exactly at Adding Steps, at $m = m_{t_0}$ for some null vertex t_0 of f of type $* \rightarrow E$.

Although the behavior of $L_{f\rho+\pi^m}(T_m)$ is different from that of $L(\Gamma_{\bullet}^m)$, we know that when going from $m = m_{t_0}^a$ to $m = m_{t_0}^b$, a single index is added to $L_{f\rho+\pi^m}(T_m)$ (this index might be added to $L_{f\rho+\pi^m}(T_m)$ at a step different from m_{t_0}).

We will see in the Standard Step that near a null vertex of the form $E \rightarrow *$ we do have $L(\Gamma_{\bullet}^m) = L_{f\rho+\pi^m}(T_m)$, and hence for every $m \notin$

$\sqcup_{t_0}[m_{t_0}^a, m_{t_0}^b]$ we have $L(\Gamma_\bullet^m) = L_{f^{\rho+\pi^m}}(T_m)$, where the union is taken over the null vertices of f of type $\ast \rightarrow E$. Note that $\sqcup_{t_0}[T_{m_{t_0}^a}, T_{m_{t_0}^b}]$ is contained in a union of $\Theta(C_1)$ -neighborhoods of the null vertices of f .

8.4.2. Changing Step. Condition-Changing: Step m is a **Changing Step** if f has a non-null vertex t_0 in $[T_m, T_{m+1})$.

In that case $f^{\rho+\pi^m}$ also has a non-null vertex in $[T_m, T_{m+1})$. Since $f^{\rho+\pi^m}$ is C_2 -separated, it follows that $L_{f^{\rho+\pi^m}}(T_m) = L_{f^{\rho+\pi^m}}(T_{m+1})$, and this condition cannot hold simultaneously with Condition-Adding.

Construction of A_{m+1} : When m is a Changing Step, A_{m+1} consists of a single element h_{m+1} . Let us explain how \tilde{h}_{m+1} is calculated. By the C_2 -separatedness of f we get that m is at a distance of at least $C_2/(2T)$ from an Adding Step, and hence, by Remark 8.13, we get that $L(\Gamma_\bullet) = L_{T_m}(f^{\rho+\pi^m})$. Denote $E_\bullet = f_{E_\bullet}^{\rho+\pi^m}(T_m)$ and $E'_\bullet = f_{E'_\bullet}^{\rho+\pi^m}(T_{m+1})$.

By Lemma 7.16, there is $\tilde{h}_{m+1} \in B_T$ such that $\tilde{h}_{m+1} \text{ span } \tilde{\Gamma}_l^m \in \text{Gr}(l, n)_{\rightarrow E'_l}$ for every $l \in L_{T_{m+1}}(f^{\rho+\pi^m})$. The uniformity in Lemma 7.16 shows that for T large enough (depending only on E_{all}) we have

$$(8.13) \quad g_T \tilde{h}_{m+1} \text{ span } \tilde{\Gamma}_l^m \in U_\varepsilon \left(\text{Gr}(l, n)_{E'_l}^g \right).$$

In addition,

$$(8.14) \quad \begin{aligned} \text{covol}(g_T \tilde{h}_{m+1} \tilde{\Gamma}_l^m) &= \Theta(\text{covol}(\tilde{h}_{m+1} \tilde{\Gamma}_l^m)) \cdot \exp(O(T)) \\ &= \Theta(\text{covol}(\tilde{\Gamma}_l^m)) \cdot \exp(O(T)) \end{aligned}$$

Updating the auxiliary parameters: We anchor π_l^{m+1} for every l at which f has a non-null vertex in $[T_m, T_{m+1})$. By Eq. (8.14) and Definition 8.10, $\pi_l^{m+1}(T_{m+1}) - \pi_l^m(T_{m+1}) = O(T)$. In this step we preserve Γ_\bullet^m .

8.4.3. Standard Step. Condition-Standard: Step m is a **Standard Step** if it is neither an Adding Step nor a Changing Step.

Construction of A_{m+1} : Denote $L = L(\Gamma_\bullet^m)$ and by E_\bullet the filtration $f_{E_\bullet}(T_m)$ restricted to L . Let \tilde{A}'_{m+1} be the set of $\tilde{h} \in H_{g_{T_m} h_m \text{ span } \Gamma_\bullet^m \rightarrow E_\bullet}$ that is given by Lemma 7.13 and satisfies:

- For every two distinct elements $\tilde{h}_{m+1}, \tilde{h}'_{m+1} \in \tilde{A}'_{m+1}$ we have $d_\varphi(\tilde{h}_{m+1}, \tilde{h}'_{m+1}) > 3^{\alpha_\varphi} \exp(-T)$.
- $\#\tilde{A}'_{m+1} = \Theta(\exp(T\delta(E_\bullet)))$.

For each noise lattice $\tilde{\Gamma}^m$ with $\tilde{\Gamma}^m \subset \tilde{\Gamma}^m \subset \tilde{\Gamma}_l^m$, let E with $E_l \subset E \subset E_l'$ be a non-final multiset of size $\text{rank } \Gamma^m$.

We will mark $\tilde{h}_{m+1} \in \tilde{A}'_{m+1}$ as *E-forbidden* if

$$\tilde{h}_{m+1} \tilde{\Gamma}^m \in U_\varepsilon(\text{Gr}(\text{rank } \Gamma, n)_E^g)$$

and there is an

$$\tilde{h}'_{m+1} \in B_{d_\varphi}(h_{m+1}; 3^{\alpha_\varphi} \exp(-T)) \cap H_{g_{T_m} h_m \text{ span } \Gamma_\bullet^m \rightarrow f_E^{\rho+\pi^m}(T_m)}$$

such that $\tilde{h}'_{m+1} \tilde{\Gamma}^m \in \text{Gr}(\text{rank } \Gamma, n)_{\rightarrow E}^{\varepsilon, g}$.

Let $\tilde{A}_{m+1} \subseteq \tilde{A}'_{m+1}$ be the set of non-forbidden elements. By Lemma 7.13, the number of E forbidden elements is

$$O(\exp(T\delta(E_\bullet \cup \{E\}))).$$

By Observation 8.3, we have $\delta(E_\bullet \cup \{E\}) < \delta(E_\bullet)$. Consequently, the number of forbidden elements is negligible relative to $\#\tilde{A}'_{m+1}$, and $\#\tilde{A}_{m+1} = \Theta(\exp(T\delta(E_\bullet)))$ for every T large enough. Following Subsubsection 8.2.1 and Eq. (8.7), define $A_{m+1} := g_{-T_m} \tilde{A}_{m+1} g_{T_m} h_m$.

Updating the auxiliary parameters: In a Standard Step we do not anchor π . If there is a vertex of type $f_{E,l}^{\rho+\pi^m}(T_m) \rightarrow *$ in $[T_m, T_{m+1})$, then we remove Γ_l^m from the sequence Γ_\bullet^m . Otherwise we preserve Γ_\bullet^m .

8.5. Proof that the Assertions Hold. We have to prove four properties: First, we need to show that with all the anchoring we did, we preserved the property that $\pi_l^m(t) \in [-C_1, C_1]$ for every $t \geq 0$ and $m \geq 0$. We also need to prove the three Assertions (S.4), (S.5) and (S.6). Only the proof of Assertion (S.6) does not follow from the above discussion.

To define the behavior of Alice in step m , we need to show that these assertions hold up to step m . To establish that the assertions hold at step m , it will be convenient to assume that the whole game has been played while Alice followed the strategy σ_{Alice} . The proof will not depend on the moves after step m , hence this assumption is harmless.

8.5.1. Bound on π . The bound on π_l^m follows from the discussion. The first time we encounter a nontriviality interval J we anchored π_l^m to a value of size $O(\gamma_5 T)$, and afterwards changed by at most $O(T)$ every time there was a non-null vertex $E \rightarrow E'$ at l . Since there are at most $l(n-l)$ such vertices in J , the result follows provided C_1 is large enough.

8.5.2. Existence of Lattices - Assertion (S.4). By the conditions of the Adding Step and the removing of a lattice from Γ_m , we get that if $l \in L(\Gamma_\bullet^m)$, then $l \in L_{f^{\rho+\pi^{m'}}}(T_m)$ for some m' , and hence $l \in L_f(T_m)$, as desired.

8.5.3. *Bounds on the Flag Elements - Assertion (S.5).* Eq. (8.4) holds at Standard Steps by Corollary 8.12, at Adding Steps by Eq. (8.12) and Corollary 8.12, and at Change Steps by Eq. (8.13).

As for Eq. (8.5), for every m in which we anchored π_l^{m+1} , step m was either a Changing Step which changed l or an Adding Step which added Γ_l^{m+1} to the filtration Γ_{\bullet}^{m+1} . In these step we constructed Γ_l^{m+1} and h_{m+1} to satisfy this very assertion. For any step $m' > m$ in which $\Gamma_l^{m''} = \Gamma_l^m$ for every m'' with $m \leq m'' \leq m'$, the assertion follows from Corollary 8.12.

8.5.4. *Bounds on Noise - Assertion (S.6).* The rest of the subsection is devoted to the proof of Assertion (S.6). We will bound the noise at step m using the bound on the noise at previous steps. To make sure the noise does not amplify, we will identify regions where the noise decays.

We need to prove that the noise is bounded by $O(T)$. To this end we will show that for every lattice Γ such that $\tilde{\Gamma}_l^m \subset \Gamma \subset \tilde{\Gamma}_{l'}^m$, with l, l' consecutive elements in $L(\Gamma_{\bullet}^m)$ we have

$$(8.15) \quad \log \text{covol}(\tilde{\Gamma}_l^m / \Gamma / \tilde{\Gamma}_{l'}^m) \leq \gamma_{10} T.$$

The constant γ_{10} is larger than the bound on the noise we used in the Adding Step 8.4.1. When $E = f_{E,l}(T_m) \subset E' = f_{E,l'}(T_m)$ is a non-scalar pair for $m = m_0, \dots, m_1$ and if $m - m_0$ is large enough, we will have a better bound of $\gamma_4 T$ in Eq. (8.15) for $m = m_1$, see Eq. (8.3).

Definition 8.14 (Rectangle). Let $[m_0, m_1]$ with $0 \leq m_0 < m_1$ be a maximal interval such that $l_0 < l_1$ are consecutive elements in $L(\Gamma_{\bullet}^m)$ for all m with $m_0 \leq m \leq m_1$. Note that we may have $m_1 = \infty$. We call such a configurations of $(m_0 < m_1, l_0 < l_1)$ a *rectangle*. Every rectangle satisfies exactly one of the following conditions:

- (R.1) $m_0 = 0$. In this case we say that $(m_0 < m_1, l_0 < l_1)$ is an *initial rectangle*.
- (R.2) It was preceded by two rectangles, $(m_{-1} < m_0 - 1, l_0 < l_{1/2})$ and $(m'_{-1} < m_0 - 1, l_{1/2} < l_1)$. In this case we say that these two rectangles *merge* into $(m_0 < m_1, l_0 < l_1)$.
- (R.3) It is split from another rectangle, see Case (R.6) below.

In addition, every rectangle satisfies exactly one of the following conditions:

- (R.4) $m_1 = \infty$. In this case we say that $(m_0 < m_1, l_0 < l_1)$ is a *final rectangle*.
- (R.5) It is merged with another rectangle, see Case (R.2) above.

(R.6) It is succeeded by two rectangles, $(m_1 + 1 < m_2, l_0 < l_{1/2})$ and $(m_1 + 1 < m'_2, l_{1/2} < l_1)$. In this case we say that $(m_0 < m_1, l_0 < l_1)$ *splits* into these two rectangles.

If $(m_0 < m_1, l_0 < l_1)$ splits, then there is an Adding Step at m_1 . If the rectangle is the merger of two other rectangles, then there was a Standard Step at $m_0 - 1$, and hence $f^{\rho + \pi^{m_0 - 1}}$ has a null vertex at some $l_0 < l < l_1$ in $[T_{m_0 - 1}, T_{m_0}]$. \heartsuit

Fix a rectangle $(m_0 < m_1, l_0 < l_1)$. We will consider the following invariant, which depends on the rectangle.

Definition 8.15 (The potential invariant). For every $m \in [m_0, m_1]$ denote

$$(8.16) \quad P_m := - \min_{\tilde{\Gamma}_{l_0}^m \subset \Gamma \subset \tilde{\Gamma}_{l_1}^m} \xi(\text{rank } \Gamma) \log \text{covol}(\tilde{\Gamma}_{l_1}^m / \Gamma / \tilde{\Gamma}_{l_0}^m),$$

where $\xi : [l, l'] \rightarrow \mathbb{R}$ is given by $\xi(x) = \frac{1}{(x-l-1)(l'+1-x)}$. \heartsuit

Note that the function $x \mapsto \frac{1}{\xi(x)}$ is positive and strictly concave.

Lemma 8.16. *The minimum in Eq. (8.16) can be taken over lattices which project to elements of the Harder-Narasimhan filtration $F_{\text{HN}}(\Gamma_{l_1}^m / \Gamma_{l_0}^m)$. Moreover, if Ψ with $\tilde{\Gamma}_{l_0}^m \subset \Psi \subset \tilde{\Gamma}_{l_1}^m$ is a primitive lattice not in the filtration, then*

$$\xi(\text{rank } \Gamma) \log \text{covol}(\tilde{\Gamma}_{l_1}^m / \Gamma / \tilde{\Gamma}_{l_0}^m) \leq (1 - \gamma_1^{-1}) P_m,$$

for some $\gamma_1 > 0$ which depends only on n .

Before the proof we will quote a quantitative version of the claim of uniqueness of lattices representing vertices of $F_{\text{HN}}(\Lambda)$. It is equivalent to [9, Corollary 1.31].

Lemma 8.17. *Let $\Gamma \subseteq \Lambda$ be a primitive lattice, such that Γ is not in the Harder-Narasimhan filtration. There exist primitive lattices Γ_0, Γ_2 with $\Gamma_0 \subset \Gamma \subset \Gamma_2 \subseteq \Lambda$ such that $\log \text{covol}(\Gamma_2 / \Gamma / \Gamma_0) \geq 0$.*

Proof of Lemma 8.16. Let Ψ with $\tilde{\Gamma}_{l_0}^m \subset \Psi \subset \tilde{\Gamma}_{l_1}^m$ be a primitive lattice which is not in the filtration. By Lemma 8.17 applied to $\Lambda = \tilde{\Gamma}_{l_1}^m / \tilde{\Gamma}_{l_0}^m$, there are Ψ', Ψ'' with

$$\tilde{\Gamma}_{l_0}^m \subseteq \Psi' \subset \Psi \subset \Psi'' \subseteq \tilde{\Gamma}_{l_1}^m$$

such that $\log \text{covol}(\Psi'' / \Psi / \Psi') \geq 0$.

Denote $l_{1/2} := \text{rank } \Psi$, $l_{1/4} := \text{rank } \Psi'$, $l_{3/4} := \text{rank } \Psi''$. Then

$$\begin{aligned}
& -\xi(l_{1/2}) \log \text{covol}(\Gamma_{l_1}^m / \Psi / \Gamma_{l_0}^m) \\
& \leq -\xi(l_{1/2}) \left(\frac{l_{3/4} - l_{1/2}}{l_{3/4} - l_{1/4}} \log \text{covol}(\Gamma_{l_1}^m / \Psi' / \Gamma_{l_0}^m) \right. \\
& \quad \left. + \frac{l_{1/2} - l_{1/4}}{l_{3/4} - l_{1/4}} \log \text{covol}(\Gamma_{l_1}^m / \Psi'' / \Gamma_{l_0}^m) \right) \\
& \leq \xi(l_{1/2}) \left(\frac{l_{3/4} - l_{1/2}}{l_{3/4} - l_{1/4}} \frac{P_m}{\xi(l_{1/4})} + \frac{l_{1/2} - l_{1/4}}{l_{3/4} - l_{1/4}} \frac{P_m}{\xi(l_{3/4})} \right) \\
& = P_m \xi(l_{1/2}) \left(\frac{l_{3/4} - l_{1/2}}{l_{3/4} - l_{1/4}} \frac{1}{\xi(l_{1/4})} + \frac{l_{1/2} - l_{1/4}}{l_{3/4} - l_{1/4}} \frac{1}{\xi(l_{3/4})} \right).
\end{aligned}$$

Since $1/\xi$ is a strictly concave function, we can choose $\gamma_1 > 0$ so that

$$1 - \gamma_1^{-1} > \max_{1 \leq l_{1/2} \leq n-1} \xi(l_{1/2}) \max_{0 \leq l_{1/4} < l_{1/2} < l_{3/4} \leq n} \frac{l_{3/4} - l_{1/2}}{l_{3/4} - l_{1/4}} \frac{1}{\xi(l_{1/4})} + \frac{l_{1/2} - l_{1/4}}{l_{3/4} - l_{1/4}} \frac{1}{\xi(l_{3/4})} < 1,$$

as desired. Note that γ_1 depends only on E_{all} . \square

Observation 8.18 (P_m is Lipschitz). For every $m \in [m_0, m_1]$ and $\tilde{\Gamma}_{l_0}^m \subset \Gamma \subset \tilde{\Gamma}_{l_1}^m$ we have

$$\left| \log \text{covol}(\tilde{\Gamma}_{l_1}^{m+1} / g_T \tilde{h}_{m+1} \Gamma / \tilde{\Gamma}_{l_0}^{m+1}) - \log \text{covol}(\tilde{\Gamma}_{l_1}^m / g_T \tilde{h}_m \Gamma / \tilde{\Gamma}_{l_0}^m) \right| \leq \gamma'_1 T,$$

for some γ'_1 , which depends only on E_{all} . This implies that $|P_{m+1} - P_m| \leq \gamma'_1 T$. \clubsuit

We next prove the following lemma, which relates the functions P_m for m in certain intervals.

Lemma 8.19. *Let $[m'_0, m'_1]$ with $m_0 \leq m'_0 < m'_1 \leq m_1$ be an interval in which $f_{E, l_0}(T_m) = E$, $f_{E', l_1}(T_m) = E'$ for every $m \in [m'_0, m'_1]$. Then:*

- (1) *If $E \subseteq E'$ is a scalar pair, then $P_{m'_1} = P_{m'_0} + O(1)$.*
- (2) *There is d_0 (depending only in E_{all}) such that if $E \subseteq E'$ is not a scalar pair $m'_1 - m'_0 \geq d_0$, then*

$$P_{m'_1} \leq \max(\gamma_3 T, P_{m'_0} - (m'_1 - m'_0)/\gamma'_3),$$

where $\gamma_3, \gamma'_3 \gg \gamma_1$.

Proof. We start by proving Item (1). Let $l_{1/2}$ with $l_0 < l_{1/2} < l_1$ and Γ with $\Gamma_{l_0}^{m_0} \subseteq \Gamma \subseteq \Gamma_{l_1}^{m_0}$ be a rank- $l_{1/2}$ lattice. Since $\text{span } \tilde{\Gamma}_{l_0}^{m_0} \in \text{Gr}(l_0, n)_{E'}^{\varepsilon, g}$ and $\text{span } \tilde{\Gamma}_{l_1}^{m_0} \in \text{Gr}(l_1, n)_{E'}^{\varepsilon, g}$, Lemma 7.7 provides an element q_0 for

which $q_0^{-1} \text{span } \Gamma_{l_0}^{m_0} = \text{span}(\Gamma_{l_0}^{m_0})^{\rightarrow}$ and $q_0^{-1} \text{span } \Gamma_{l_1}^{m_0} = \text{span}(\Gamma_{l_1}^{m_0})^{\rightarrow}$, and in addition, $d_{H^-}(q_0, \text{Id}) = O(\varepsilon)$.

Decomposition into eigenvalues shows that $q_0^{-1} \text{span } \Gamma^{m_0} \in \text{Gr}(l_{1/2}, n)_{E''}^g$, where E'' with $E \subset E'' \subset E'$ is the unique such multiset with $l_{1/2}$ elements, we get that $\text{span } \Gamma^{m_0} \in \text{Gr}(l_{1/2}, n)_{E''} \in \text{Gr}(l_{1/2}, n)_{\rightarrow E''}$. Since $d_{H^-}(q_0, \text{Id}) = O(\varepsilon)$, it follows that $\text{span } \Gamma^{m_0} \in \text{Gr}(l_{1/2}, n)_{E''} \in \text{Gr}(l_{1/2}, n)_{\rightarrow E''}^{O(\varepsilon), g}$. Assume that ε is chosen to be small enough so that Corollary 8.12 is applicable to $\text{span } \Gamma^{m_0}$, which completes the proof of (1).

We turn to prove Item (2). Assume $E \subset E'$ is not a scalar pair.

Claim 8.20. *There are $\gamma_2 \gg \gamma_1, \gamma'_1$ such that if $P_{m'_0} \geq (m'_1 - m'_0)\gamma_2 T$, then only lattices in $F_{\text{HN}}(\tilde{\Gamma}_{l_1}^{m'_0}/\tilde{\Gamma}_{l_0}^{m'_0})$ should be taken into account when computing $P_{m'_1}$. That is, if $\tilde{\Gamma}_{l_0}^{m'_0} = \Psi_0 \subset \Psi_1 \subset \dots \subset \Psi_k = \tilde{\Gamma}_{l_1}^{m'_0}$ is the filtration projecting to $F_{\text{HN}}(\tilde{\Gamma}_{l_1}^{m'_0}/\tilde{\Gamma}_{l_0}^{m'_0})$, then to compute $P_{m'_1}$ it is enough to use the lattices $\tilde{\Psi}_j^{m'_1}$ for $j = 0, \dots, k$, where*

$$\tilde{\Psi}_j^m := g_{T_m} h_m h_{m'_0}^{-1} g_{-T_{m'_0}} \Psi_j.$$

Proof. Let

$$\tilde{\Psi}^{m'_1} = g_{T_{m'_1}} h_{m'_1} h_{m'_0}^{-1} g_{-T_{m'_0}} \Psi; \tilde{\Gamma}_{l_0}^{m'_1} \subseteq \tilde{\Psi}^{m'_1} \subseteq \tilde{\Gamma}_{l_1}^{m'_1}, \tilde{\Gamma}_{l_0}^{m_0} \subseteq \Psi \subseteq \tilde{\Gamma}_{l_1}^{m_0},$$

with $P_{m'_1} = -\xi(\text{rank } \tilde{\Psi}^{m'_1}) \log \text{covol}(\tilde{\Gamma}_{l_1}^{m'_1}/\tilde{\Psi}^{m'_1}/\tilde{\Gamma}_{l_0}^{m'_1})$.

The lattice $\tilde{\Psi}^{m'_1}$ is primitive in $\tilde{\Gamma}_{l_1}^{m'_1}$, as we can replace it by the primitive closure and decrease its covolume. If Ψ is not one of the Ψ_j -s, then, by Lemma 8.16,

$$-\xi(\text{rank } \Psi) \log \text{covol}(\tilde{\Gamma}_{l_1}^m/\Psi/\tilde{\Gamma}_{l_0}^m) \leq (1 - \gamma_1^{-1})P_{m'_0}.$$

It follows from Observation 8.18 that

$$P_{m'_1} = -\xi(\text{rank } \tilde{\Psi}^{m'_1}) \log \text{covol}(\tilde{\Gamma}_{l_1}^{m'_1}/\tilde{\Psi}^{m'_1}/\tilde{\Gamma}_{l_0}^{m'_1}) \leq (1 - \gamma_1^{-1})P_{m'_0} + \gamma'_1 dT.$$

On the other hand, again by Observation 8.18, $P_{m'_1} \geq P_m - \gamma'_1 dT$. Consequently, we reach a contradiction as soon as $P_{m'_0} > 2\gamma'_1 \gamma_1 dT$. Therefore, we may choose $\gamma_2 = 2\gamma'_1 \gamma_1$. This proves Claim 8.20. \square

Now assume that $P_{m'_0} \geq \gamma_2(m'_0 - m'_1)T$. Let $\Psi = \Psi_j$ be the lattice for which $\tilde{\Psi}^{m'_1} = \tilde{\Psi}_j^{m'_1}$ is the lattice that attains the minimum in the definition of $P_{m'_1}$. Denote $l := \text{rank } \Psi$. By the discussion above, for every $m'_0 \leq m \leq m'_1$ the lattice $\tilde{\Psi}^m := g_{T_m} h_m^{-1} h_{m'_0}^{-1} g_{-T_{m'_0}} \Psi$ projects to an element in the Harder-Narasimhan filtration $F_{\text{HN}}(\tilde{\Gamma}_{l_1}^m/\tilde{\Gamma}_{l_0}^m)$, and hence is a noise lattice. Denote $\bar{\Psi}^m = g_{-(m'_1-m)T} \tilde{\Psi}^{m'_1}$, and note that

$\bar{\Psi}^m = h^{m \rightarrow m'_1} \tilde{\Psi}^d$ for $h^{m \rightarrow m'_1} \in B_{d_\varphi}(\text{Id}; 1)$; that is, $\bar{\Psi}^m$ and $\tilde{\Psi}^m$ are somewhat close.

Note that if $C_2 > (m'_1 - m'_0)T$, then by Lemma 5.28 only $O(1)$ steps between m'_0, m'_1 may not be Standard Steps. Moreover, every Standard Step m , is designed to push $\tilde{\Psi}^d$ away from $\text{Gr}(l, n)_{\rightarrow E''}$, for any non-final E'' with $E \subset E'' \subset E'$. Specifically, for every $m \in [m'_0, m'_1]$ denote by $E^m \in \mathcal{I}_l$ the unique multiset for which $\bar{\Psi}^m \in U_{\varepsilon/2}(\text{Gr}(l, n)_{E^m}^g)$, if such a multiset exists, and otherwise it remains undefined. By Theorem 2.7, E^m is not defined for at most $l(n-l)$ values of m .

We claim that if m is a Standard Step and E^m is defined and non-final, then E^{m+1} does not equal to E^m . Indeed, since

$$\bar{\Psi}^m = h^{m \rightarrow m'_1} \tilde{\Psi}^m \in U_{\varepsilon/2}(\text{Gr}(l, n)_{E^m}^g)$$

and $d_\varphi(h^{m \rightarrow m'_1}, \tilde{h}_{m+1}) \leq \exp(-T)$, it follows that

$$\tilde{\Psi}^m \in U_\varepsilon(\text{Gr}(l, n)_{E^m}^g).$$

Hence, since Alice did not forbid h_{m+1} , there is no

$$\tilde{h}' \in B_{d_\varphi}(\tilde{h}_{m+1}; 3^{\alpha_\varphi} \exp(-T)) \cap H_{\text{span } \tilde{\Gamma}^m \rightarrow f_{E, \bullet}(T_m)}$$

such that $\tilde{h}' \text{span } \tilde{\Psi}^m \in \text{Gr}(l, n)_{\rightarrow E^m}^{\varepsilon, g}$. Therefore, there is no

$$(8.17) \quad \tilde{h}'' \in B_{d_\varphi}(\text{Id}; 2^{\alpha_\varphi} \exp(-T)) \cap H_{h^{m \rightarrow m'_1} \text{span } \tilde{\Gamma}^m \rightarrow f_{E, \bullet}(T_m)}$$

such that $\tilde{h}'' \text{span } \bar{\Psi}^m \in \text{Gr}(l, n)_{\rightarrow E^m}^{\varepsilon, g}$. Since $\text{span } \bar{\Psi}^m \in U_{\varepsilon/2}(\text{Gr}(l, n)_{E^m}^g)$, the ε, g part is redundant, and there is no \tilde{h}'' such that Eq. (8.17) holds and $\tilde{h}'' \text{span } \bar{\Psi}^m \in \text{Gr}(l, n)_{\rightarrow E^m}$.

Conjugating Eq. (8.17) by g_T we deduce that there is no matrix

$$\tilde{h}''' \in B_{d_\varphi}(\text{Id}; 2^{\alpha_\varphi}) \cap H_{h^{m+1 \rightarrow m'_1} \text{span } \tilde{\Gamma}^{m+1} \rightarrow f_{E, \bullet}(T_{m+1})}$$

such that $\tilde{h}''' \bar{\Psi}^{m+1} \in \text{Gr}(l, n)_{\rightarrow E^m}$. If $\bar{\Psi}^{m+1} \in U_{\varepsilon/2}(\text{Gr}(l, n)_{E^m}^g)$, that is, $E^{m+1} = E^m$, then there exists $s \in U_{O(\varepsilon)}(\text{Id}; d_{\text{SL}_n(\mathbb{R})})$ such that $s^{-1} \bar{\Psi}^{m+1} \in \text{Gr}(l, n)_{E^m}^g$ and $s^{-1} h^{m+1 \rightarrow m'_1} \text{span } \tilde{\Gamma}^{m+1} = h^{m+1 \rightarrow m'_1} \text{span } \tilde{\Gamma}^{m+1}$.

By Lemma 2.3, if ε is small enough, then there are $q \in H^{-0}$ and $\tilde{h}''' \in H$ such that $d_{H^{-0}}(q, \text{Id}), d_H(\tilde{h}''', \text{Id}) = O(\varepsilon)$ and $s = \tilde{h}''' q$. This \tilde{h}''' satisfies the above condition for \tilde{h} , which contradicts the assumption that $E^{m+1} = E^m$.

It follows that for $m \geq m_{1/2} = m'_0 + O(1)$ we have

$$\text{span } \tilde{\Psi}^m, \text{span } \bar{\Psi}^m \in U_\varepsilon \left(\text{Gr}(l, n)_{E''_{\text{final}}}^g \right),$$

where E_{final} is the unique final multiset of size l in $E \subseteq E'$. By Observation 8.18,

$$\xi(\text{rank } \tilde{\Psi}^{m_{1/2}}) \log \text{covol}(\tilde{\Gamma}_{l_1}^{m_{1/2}} / \tilde{\Psi}^{m_{1/2}} / \tilde{\Gamma}_{l_0}^{m_{1/2}}) \leq P_m + O(\gamma'_1).$$

By Corollary 2.10, for every $m \geq m'_0$ we have

$$\begin{aligned} & \log \text{covol}(\tilde{\Gamma}_{l_1}^m / \tilde{\Psi}^m / \tilde{\Gamma}_{l_1}^m) - \log \text{covol}(\tilde{\Gamma}_{l_1}^{m_{1/2}} / \tilde{\Psi}^{m_{1/2}} / \tilde{\Gamma}_{l_1}^{m_{1/2}}) \\ &= (m - m_{1/2}) \left(\frac{\text{rank } \Psi - l}{l' - l} \eta_E + \frac{l' - \text{rank } \Psi}{l' - l} \eta_{E'} - \eta_{E_{\text{final}}} \right) + O(1) \end{aligned}$$

Define

$$\Phi := \frac{\text{rank } \Psi - l}{l' - l} \eta_E + \frac{l' - \text{rank } \Psi}{l' - l} \eta_{E'} - \eta_{E_{\text{final}}}.$$

By Lemma 8.7, $\Phi < 0$. Consequently, if $P_{m'_0} \geq (m'_1 - m'_0) \gamma_2 T$, then

$$\begin{aligned} P_{m'_1} &= \xi(l) \log \text{covol}(\tilde{\Gamma}_{l_1}^{m'_1} / \tilde{\Psi}^{m'_1} / \tilde{\Gamma}_{l_0}^{m'_1}) \\ &\leq \xi(l) \log \text{covol}(\tilde{\Gamma}_{l_1}^{m_{1/2}} / \tilde{\Psi}^{m_{1/2}} / \tilde{\Gamma}_{l_0}^{m_{1/2}}) + (m'_1 - m_{1/2}) \Phi + O(1) \\ &= P_{m'_0} + (m_{1/2} - m'_0) \gamma'_1 + (m'_1 - m_{1/2}) \Phi + O(1). \end{aligned}$$

Hence, if $m_1 - m_0 \gg \gg -\gamma_1 / \Phi$, then

$$P_{m'_1} - P_{m_0} < (m'_1 - m_0) \frac{\Phi}{2}.$$

The result follows, with d_0 being the implicit lower bound on $m'_1 - m_0$. \square

We next show that P_m is bounded by $\gamma_7 T$. We will do so by induction on all rectangles $(m_0 < m_1, l_0 < l_1)$, sorted by m_0 . That is, when proving that P_m is bounded by $\gamma_7 T$ for $(m_0 < m_1, l_0 < l_1)$, we assume this claim holds for all rectangles $(m'_0 < m'_1, l'_0 < l'_1)$ with $m'_0 < m_0$. The tricky part is to show that the implicit constant does not increase at every step of the induction. What makes this possible is the decaying behavior of P_m in case 2 of Lemma 8.19, and the fact that this case occurs near every null point.

Lemma 8.21. *There exist $\gamma_9 > 0$ and bounds $\gamma_8 > \gamma_7 > \gamma_3$ that depend only on E_{all} , such that for every rectangle $(m_0 < m_1, l_0 < l_1)$ we have*

$$P_m \leq \begin{cases} \gamma_8 T, & \text{if } m_0 \leq m < m_0 + \gamma_9, \\ \gamma_7 T, & \text{if } m_0 + \gamma_9 \leq m \leq m_1. \end{cases}$$

Moreover, if the rectangle splits (see Definition R.6) then $P_{m_1} < \gamma_3 T$.

Corollary 8.22. *At every Adding Step the noise is bounded by γ_4 . Formally, if Alice attempts to add a lattice Γ_l^{m+1} at step m , and if $l^- < l < l^+$ are the consecutive elements of l in $L_{T_{m+1}}(f^{\rho+\pi^m})$, then for every $\tilde{\Gamma}_{l^-}^m \subset \Gamma \subset \tilde{\Gamma}_{l^+}^m$ we have*

$$\log \text{covol}(\tilde{\Gamma}_{l^+}^m / \Gamma / \tilde{\Gamma}_{l^-}^m) \geq -\gamma_4 T.$$

Proof of Corollary 8.22. The claim follows from the fact that in this case there is a rectangle $(m_0 < m, l^- < l^+)$ that splits, and hence $P_m < \gamma_3 T$. \square

Recall that Corollary 8.22 was used in the definition of the Adding Step. We turn to the proof of Lemma 8.21.

Proof of Lemma 8.21. We distinguish three cases, depending on whether the rectangle is a merger of two rectangles, is split from another rectangle or is an initial rectangle.

Rectangle is merged: Assume the rectangle is merged from, say, $(m_{-1} < m_0 - 1, l_0 < l_{1/2})$ and $(m'_{-1} < m_0 - 1, l_{1/2} < l_1)$. The induction hypothesis on the two rectangles bounds the corresponding potential invariants: $P'_{m_0-1}, P''_{m_0-1} \leq \gamma_7 T'$. Since $\tilde{\Gamma}_{l_{1/2}}^{m_0-1}$ was removed at step $m_0 - 1$, and by Eq. (8.5) we get that $\text{covol}(\tilde{\Gamma}_{l_1}^{m_0-1} / \tilde{\Gamma}_{l_{1/2}}^{m_0-1} / \tilde{\Gamma}_{l_0}^{m_0-1}) = O(T)$. It follows that the bound on P'_{m_0-1}, P''_{m_0-1} gives a bound on $P_m = O(T + P''_{m_0-1} + P'_{m_0-1}) = O(\gamma_7 T)$.

Since $f^{\rho+\pi^{m_0-1}}$ has a null vertex at $[T_{m_0-1}, T_{m_0})$ and $f^{\rho+\pi^{m_0-1}}$ is $C_2/2$ -separated it follows that for all $m_0 \leq m \leq m_0 + C_2/(2T)$ the multisets $f_{E,l_0}(T_m)$ and $f_{E,l_1}(T_m)$ are constant. By Lemma 8.8, these $f_{E,l_0}(T_m) \subset f_{E,l_1}(T_m)$ do not form a scalar pair.

Let a_0, \dots, a_1 be the shortest sequence of numbers with $m_0 = a_0 < a_1 < \dots < a_k = m_1$ such that $m \mapsto (f_{E,l_0}(T_m), f_{E,l_1}(T_m))$ is constant for $a_i \leq m \leq a_{i+1} - 1$ for each $0 \leq i \leq k - 1$. Since there are at most $O(n^2)$ vertices in every nontriviality interval, it follows that $k = O(n^2)$. Lemma 8.19 implies that P_m decays on $[a_0, a_1 - 1]$, and Observation 8.18 and Lemma 8.19 imply that P_m does not expand more than $O(\gamma_1 d_0 T)$ over the succeeding intervals.

Rectangle is split: Suppose that the rectangle did split from another rectangle, w.l.o.g., $(m_{-1} < m_0 - 1, l_0 < l_2)$ split into $(m_0 < m_1, l_0 < l_1)$ and $(m_0 < m'_1, l_1 < l_2)$. By the construction of the Adding Step, $\text{covol}(\Gamma_{l_2}^{m_0} / \Gamma_{l_1}^{m_0} / \Gamma_{l_0}^{m_0}) = O(\gamma_5 T)$. In addition, if P'_m is the potential invariant of the rectangle $(m_{-1} < m_0 - 1, l_0 < l_2)$, then by induction $P'_{m_0-1} \leq \gamma_3 T$. This yields the bound $P_{m_0} = O(\gamma_5 T)$. As in the previous case, we obtain a bound on P_m , for all $m_0 \leq m \leq m_1$.

Rectangle is initial: At step 0, the only noise lattices are the lattices in $F_{\text{HN}}(\Lambda)$. It follows that $P_0 = O(T)$, provided T is large enough. As in the previous cases, we obtain a bound on P_m for all $m_0 \leq m \leq m_1$.

To bound P_{m_1} in the case that the rectangle splits, note that since $f^{\rho+\pi^{m_1}}$ is separated we get that $f_{E,l_1}(T_m), f_{E,l_2}(T_m)$ are constant for $m_1 - C_2/(2T) \leq m \leq m_1$, and hence the claim follows from Lemmas 8.19 and 8.8. \square

8.6. Existence of a Lattice with a Given Template - Proof of Corollary 2.35. In this subsection we will prove Corollary 2.35 which states that every template describes at most one lattice. Let σ_{Alice} be the strategy constructed above for the (T, g) -game. Let h_0, h_1, \dots be the output of a (T, g) -game where Alice plays via σ_{Alice} and Bob plays arbitrarily. It is sufficient to prove that the lattice $h_\infty \Lambda$ satisfies $f^{h_\infty \Lambda} \sim f$.

Lemma 8.23. *Let h_0, h_1, \dots be the output of a (T, g) -game where Alice plays via σ_{Alice} and Bob plays arbitrarily. Then $f^{h_\infty \Lambda} \sim f$.*

Before the proof we will prove the following corollary of Lemma 8.17.

Corollary 8.24. *Let Λ be a lattice and let Γ_\bullet be a filtration of Λ composed of primitive sublattices. Let Γ'_\bullet be the filtration Γ_\bullet obtained by adding any two consecutive $\Gamma_l \subset \Gamma_{l'}$ the inverse image of $F_{\text{HN}}(\Gamma_{l'}/\Gamma_l)$. Consider the piecewise linear function $f : [0, n] \rightarrow \mathbb{R}$ defined by its values on $L(\Gamma'_\bullet)$ as $f(l) = \log \text{covol } \Gamma'_l$. The following statements are equivalent:*

- (1) f is convex and strictly convex at $L(\Gamma_\bullet) \setminus \{0, n\}$.
- (2) Γ_\bullet is contained in $F_{\text{HN}}(\Lambda)$.
- (3) $\Gamma'_\bullet = F_{\text{HN}}(\Lambda)$.

Proof of Lemma 8.23. The verification that (3) \iff (2) \implies (1) follows from the definition of the Harder-Narasimhan filtration and the definition of f . To show that (1) \implies (2), we use induction on n . For $n = 1$ there is nothing to prove. Assume (1) holds for Λ , but not (2), i.e., for some $l \in L(\Gamma_\bullet)$ with $l \neq 0, n$ we have $\Gamma_l \notin F_{\text{HN}}(\Lambda)$. By the induction hypothesis applied to $\Gamma_l, \Lambda/\Gamma_l$, we conclude that Γ'_\bullet is composed of $F_{\text{HN}}(\Gamma_l)$ and the inverse image of $F_{\text{HN}}(\Lambda/\Gamma_l)$ in Λ . Applying Lemma 8.17 one contradicts the strict convexity of f at l , as desired. \square

Proof. By Observation 4.14, $f^{h_\infty \Lambda}|_{[0, T_m]} \sim f^{h_m \Lambda}|_{[0, T_m]}$, where the implicit constants are independent of m, T . Hence, $|f_{H,i}^{h_m \Lambda}(t) - f_{H,i}^{h_\infty \Lambda}(t)| =$

$O(1)$ for every $0 \leq i \leq n$ and $t \in [0, T_m]$. By Remark 8.13, for every m for which T_m lies outside a $\Theta(C_1)$ -neighborhood of the vertices of f we have $L(\Gamma_\bullet^m) = L_f(T_m)$. Since f is C_2 -significant, if T_m is outside a (maybe larger) $O(C_1)$ -neighborhood of the vertices of f , $\partial^2 f(T_m) \in \{0\} \cup [\Theta(C_1), \infty)$. By Assertion (S.5) Eq. (8.5) and Assertion (S.6) Eq. (8.6) we can apply Corollary 8.24 to the filtration $\tilde{\Gamma}_\bullet^m$ of $\tilde{\Lambda}^m$ and deduce that $F_{\text{HN}}(\tilde{\Lambda}^m)$ contains $\tilde{\Gamma}_\bullet^m$ and that $F_{\text{HN}}(\tilde{\Lambda}^m)_{H,l} = f_{H,l}(T_m) + O(C_1)$ for every $0 \leq l \leq n$. Since $h^{m \rightarrow \infty} \tilde{\Lambda}^m = g_{T_m} h_\infty \Lambda$ and $d_H(h^{m \rightarrow \infty}, \text{Id}) = O(1)$ (see Subsection 8.3) it follows that for every m such that T_m is outside an $O(C_1)$ -neighborhood of the vertices of f we have $F_{\text{HN}}(g_{T_m} h_\infty \Lambda)_{H,l} = f_{H,l}(T_m) + O(C_1)$. Since for every $t \geq 0$ there is such an m with $t - T_m = O(C_1)$, we get that for every $t \geq 0$

$$(8.18) \quad F_{\text{HN}}(g_t h_\infty \Lambda)_{H,l} = f_{H,l}(t) + O(C_1).$$

This proves Condition 1 of Definition 2.23 of matching templates. As for condition 2, let $[a, b]$ be an interval such that $f_{E,l}(t) = E \in \mathcal{I}_l$ and $\partial^2 f_{H,l}(t) \geq C$ for every $t \in [a, b]$, for some $C > 0$ to be determined later. By Eq. (8.18), if $C = \Theta(C_1)$, then $F_{\text{HN}}(g_t h_\infty \Lambda)_{H,l} = \Theta(C)$ for every $t \in [a, b]$. It follows from Assertion (S.6) that $h_\infty^{-1} g_{-T_m} F_{\text{HN}}(g_{T_m} h_\infty \Lambda)_{\Gamma,l}$ is not a noise lattice at step m , and hence $\Gamma_l^m = h_\infty^{-1} g_{-T_m} F_{\text{HN}}(g_{T_m} h_\infty \Lambda)_{\Gamma,l}$. Assertion (S.5) and Eq. (8.4) now implies that $\text{span}(\tilde{\Gamma}_l^m) \in U_\varepsilon(\text{Gr}(l, n)_E^g)$. Using arguments similar to those used to prove Corollary 8.12, we conclude that for every $t \in [a + O(|\log \varepsilon|), b - O(|\log \varepsilon|)]$ we have $F_{\text{HN}}(g_{T_m} h_\infty \Lambda)_{\Gamma,l} \in U_\varepsilon(\text{Gr}(l, n)_E^g)$. This shows that $h_\infty \Lambda$ (ε, C) -matches f . \square

9. BOB'S STRATEGY

Bob's strategy is a function $\sigma_{\text{Bob}}^T : (T_0, h_0, A_1, h_1, \dots, A_{m+1}) \mapsto h_{m+1}$, where $h_{m+1} \in A_{m+1}$. As it turns out, the construction of Bob's strategy is simpler than the that of Alice's strategy. While Alice tries to create a good template, Bob's goal is to interfere with her attempts.

Let Λ be a lattice. At each step m , the set A_{m+1} satisfies

- $A_{m+1} \subseteq g_{-T_m} B_T g_{T_m} h_m$.
- $d_\varphi(h_{m+1}, h'_{m+1}) \geq \exp(-3^{\alpha_\varphi} T_{m+1})$ for every $h_{m+1}, h'_{m+1} \in A_{m+1}$.

Equivalently, denoting $\tilde{A}_{m+1} := g_{T_m} A_{m+1} h_m^{-1} g_{-T_m}$, we have

- $\tilde{A}_{m+1} \subseteq B_T$.
- $d_\varphi(h_{m+1}, h'_{m+1}) \geq \exp(-3^{\alpha_\varphi} T)$ for every $\tilde{h}_{m+1}, \tilde{h}'_{m+1} \in A_{m+1}$.

Bob needs to choose $h_{m+1} \in A_{m+1}$, or, alternatively, $\tilde{h}_{m+1} = g_{T_m} h_{m+1} h_m^{-1} g_{-T_m} \in \tilde{A}_{m+1}$.

In the end Alice and Bob generate $h_\infty := \lim_{m \rightarrow \infty} h_m \in H$. Bob's goal is to minimize $\Delta(f^{h_\infty \Lambda})$.

9.1. Point Sorting in a Height Sequences. To construct Bob's strategy, we need to introduce an order on the indices of a height sequence. Define $\rho(\zeta) = (\zeta l(n-l))_{l=0}^l$ for every $\zeta > 0$. For every height sequence a_\bullet , define the *vanishing number* $\zeta_l(a_\bullet)$ of any index $1 \leq l \leq n-1$ by

$$\zeta_l(a_\bullet) = \min\{\zeta > 0 : l \notin L(a_\bullet^{\rho(\zeta)})\}.$$

The number $\zeta_l(a_\bullet)$ behaves similarly to $\partial^2 a_l$: they vanish simultaneously in height sequences and one can show that $\zeta_l(a_\bullet) = \Theta(\partial^2 a_l)$.

9.2. The Strategy. For every m , consider the lattice $\tilde{\Lambda}^m := g_{T_m} h_m \Lambda$ and its Harder-Narasimhan filtration $F_{\text{HN}}(\tilde{\Lambda}^m)$. Let $\varepsilon > 0$ be sufficiently small to satisfy Theorem 2.7 and Lemma 6.3. Consider the subset of indices

$$L_0^m := \{l \in L(F_{\text{HN}}(\tilde{\Lambda}^m)) \setminus \{0, n\} : \exists E(m, l) \in \mathcal{I}_l, \text{span } \tilde{\Gamma}_l^m \in U_\varepsilon(\text{Gr}(l, n)_{E(m, l)}^g)\}.$$

By Lemma 6.3, the sequence $(E(m, l))_{l \in L_0^m}$ forms a direction filtration without the extremal multisets $\emptyset, E_{\text{all}}$. Sort the set L_0^m by the value of $\zeta_l(F_{\text{HN}}(\tilde{\Lambda}^m)_{H, \bullet})$ for $l \in L_0^m$, that is, $L_0^m = \{l_1^m, l_2^m, \dots, l_{k^m}^m\}$, where

$$\zeta_{l_1^m}(F_{\text{HN}}(\tilde{\Lambda}^m)_{H, \bullet}) \geq \zeta_{l_2^m}(F_{\text{HN}}(\tilde{\Lambda}^m)_{H, \bullet}) \geq \dots \geq \zeta_{l_{k^m}^m}(F_{\text{HN}}(\tilde{\Lambda}^m)_{H, \bullet}).$$

Let i^m denote the first index of l_\bullet^m with the following property: there is $\tilde{h}_{m+1} \in \tilde{A}_{m+1}$ such that for every $h' \in B_{d_\varphi}(h_{m+1}; 3^{\alpha_\varphi} \exp(-T))$ one has

$$\exists 1 \leq i \leq i^m, h' \Gamma_{i_k} \notin H_{\text{span } \tilde{\Gamma}_{l_i^m}^m \rightarrow E(m, l_i^m)}.$$

If such i^m exists, the strategy selects this \tilde{h}_{m+1} . In this case we call the step an **Interruption Step**, and say that Bob *interrupted* i^m indices. If no such i^m exists, let $\tilde{h}_{m+1} \in \tilde{A}_{m+1}$ be arbitrary, and call step m an **Idle Step**.

10. COMPUTATION OF THE DIMENSION — PROOF OF THEOREM 2.31

10.1. Computation of the Payoff. We fix a family \mathcal{F} of g -templates that is invariant to equivalence of g -templates. For every lattice Λ , we wish to prove that

$$\dim_{\mathbf{H}}(Y_{\Lambda, \mathcal{F}}; d_\varphi) = \sup_{f \in \mathcal{F}} \Delta_0(f).$$

We will use the Dimension Game introduced in Subsection 4.4.

To prove the lower bound, it is enough to fix $f_0 \in \mathcal{F}$ with $\Delta_0(f_0) \geq \sup_{f \in \mathcal{F}} \Delta_0(f) - \varepsilon$, and to prove that Alice has a strategy σ_{Alice}^T that guarantees that the payoff is at least D_{Alice}^T , for some D_{Alice}^T such that $\lim_{T \rightarrow \infty} D_{\text{Alice}}^T \geq \Delta_0(f)$. Any h_∞ attained with Alice's strategy will have $f^{h_\infty \Lambda} \sim f_0$ by 8.23.

To prove the upper bound, we have to prove that Bob has a strategy σ_{Bob}^T that guarantees that the payoff is at most D_{Bob}^T , for some D_{Bob}^T such that

$$\lim_{T \rightarrow \infty} D_{\text{Bob}}^T \leq \sup_{f \in \mathcal{F}} \Delta_0(f).$$

10.2. The Payoff Guaranteed by Alice's Strategy. Fix $\varepsilon > 0$ as in Section 8 and let $f \in \mathcal{F}$ with $\Delta_0(f) \geq \sup_{f \in \mathcal{F}} \Delta_0(f) - \varepsilon$. To apply Alice's strategy we have to choose $C_2 \ggg C_1 \ggg T$ and approximate f by a C_2 -separated and C_2 -significant g -template. By Lemma 5.9, there is an equivalent g -template $f' \sim f$ that is C_2 -separated and C_2 -significant. The construction of f' was carried out by constructing a proper shift sequence ρ and getting $f' = f^\rho$. The following claim will allow us to replace f by f' .

Claim 10.1. *For every g -template f and every shift sequence ρ we have $\Delta_0(f^\rho) \geq \Delta_0(f)$.*

Proof. To prove the result it is enough to show that $\delta(f_{E,\bullet}(t)) \leq \delta(f_{E,\bullet}^\rho(t))$ (see Definition 2.30 of shifting a height sequence). This inequality holds since δ is monotone nonincreasing with respect to adding multisets and the fact that each multiset in $f_{E,\bullet}^\rho(t)$ appears also in $f_{E,\bullet}(t)$. \square

We can thus assume w.l.o.g. that f is C_2 -significant and use Alice's strategy as defined in Section 8. Let $A_0, h_0, A_1, h_1, \dots$ be the choices made by Alice and Bob along the game. Note that Bob may play arbitrarily, not necessarily according to the strategy σ_{Bob}^T described in Section 9. By Lemma 8.23, the resulting matrix $h_\infty = \lim_{m \rightarrow \infty} h_m$ satisfies $f^{h_\infty \Lambda} \sim f$. It remains to compute the first option of Eq. (4.10), namely $\frac{1}{T} \liminf_{m \rightarrow \infty} \frac{1}{m} \sum_{k=1}^m \log \#A_k$.

The number of elements in A_{m+1} is larger than 1 only if m is a Standard Step. In every Standard Step we have $\#A_{m+1} = \Theta(\exp(T\delta(f_E^{\rho+\pi^m}(T_m))))$, and hence $\log \#A_{m+1} = T\delta(f_E^{\rho+\pi^m}(T_m)) + O(1)$. Thus,

$$\liminf_{m \rightarrow \infty} \frac{1}{Tm} \sum_{m'=0}^m \log \#A_{m'} = \liminf_{m \rightarrow \infty} \frac{1}{m} \sum_{\substack{m'=0 \\ m' \text{ is a Standard Step}}}^m \delta(f_E^{\rho+\pi^{m'}}(T_{m'})) + O(T^{-1}).$$

We will restrict the set of m' -s we sum over even further to obtain $f_E^{\rho+\pi^{m'}}(T_{m'}) = f_E(T_{m'})$.

For every $t > 0$ such that $\partial^2 f_{H,l}(t) \notin (0, 4C_1]$ for every $1 \leq l \leq n-1$ we have $f_E^{\rho+\pi^{m'}}(t) = f_E^{\rho+\pi^{m'}}(t)$. Since f is C_2 -significant, we have $\partial^2 f_{H,l}(t) \leq (0, 4C_1]$ only if t is in an $O(C_1)$ -neighborhood of a null vertex of f .

In addition, if T_m is not in an $O(C_1)$ -neighborhood of any vertex, m is a Standard Step.

Therefore,

$$\begin{aligned}
(10.1) \quad & \sum_{\substack{m' \\ m' \text{ is a Standard Step}}}^m \delta(f_E^{\rho+\pi^{m'}}(T_{m'})) \geq \frac{1}{T} \int_{J_m} \delta(f_E(t)) dt \\
& \geq \frac{1}{T} \int_0^{T_m} \delta(f_E(t)) dt - O(C_1 \cdot \#\{\text{vertices of } f \text{ until } T_m + O(C_1)\}) \\
& \geq \frac{1}{T} \int_0^{T_m} \delta(f_E(t)) dt - O(C_1 \cdot (2n^3 + O(T_m/C_2))) = O(C_1 \cdot (1 + T_m/C_2)),
\end{aligned}$$

where J_m is the set of points $0 \leq t \leq T_m$ such that t is not in an $O(C_1)$ -neighborhood of a vertex of J . Note that inequality (10.1) follows from Lemma 5.28. Consequently,

$$\liminf_{m \rightarrow \infty} \frac{1}{Tm} \sum_{m'=0}^m \log \#A_{m'} \geq \Delta_0(f) - O(TC_1/C_2).$$

If we choose a sequence of $C_2(T) \ggg C_1(T) \ggg T \rightarrow \infty$ such that $TC_1(T)/C_2(T) \xrightarrow{T \rightarrow \infty} 0$, we obtain the upper bound.

10.3. The Payoff Guaranteed by Bob's Strategy. In this subsection we prove the lower bound. To this end we fix $\varepsilon > 0$ and consider Bob's strategy σ_{Bob}^T , both described in Section 9. We then show that when Bob follows this strategy, no matter how Alice plays (not necessarily according to the strategy described in Section 8), the payoff is at most D_{Bob}^T , where $\lim_{T \rightarrow \infty} D_{\text{Bob}}^T = \sup_{f \in \mathcal{F}} \Delta_0(f)$.

As before, denote by $(T_0, h_0, A_1, h_1, \dots)$ the choices of Alice and Bob along the game. Denote $\bar{\Lambda}^m := g_{T_m} h_\infty \Lambda$, where $h_\infty := \lim_{m \rightarrow \infty} h_m$, and consider $f := f^{h_\infty \Lambda}$. We will bound $\Delta(f)$ using $\#A_{m+1}$. Let C be the constant given by Theorem 2.28 for ε . Assume that $h_\infty \Lambda$ (ε, C) -matches f . Fix $\zeta_1 > C$ and $\zeta_2 > 4C$ to be determined later. Let $C_1 > 0$, to be defined later as well.

Lemma 10.2. *Let $\zeta \gg T$ and let $m \geq 0$ be a step for which $f_{E,\bullet}^{\rho(\zeta)}(t) = f_{E,\bullet}^{\rho(\zeta+\zeta_2)}(t)$ is constant for every $t \in [T_m - C - C_1, T_{m+1} + C + C_1]$. Then $\#A_{m+1} = O(\exp(T\delta(f_{E,\bullet}^{\rho(\zeta)}(T_m))))$.*

Proof. Denote the distinct elements of $L(f_{E,\bullet}^{\rho(\zeta)}(T_m))$ by $\{0, l_1, \dots, l_r, n\}$ and fix $t \in [T_m - C - C_1, T_{m+1} + C + C_1]$. The elements $(\zeta_l(f_{H,\bullet}(t)))_{l=1}^r$ are the r largest elements in $\{\zeta_l(f_{H,\bullet}(t)) : l = 1, \dots, n-1\}$, and are larger than their successors by at least ζ_2 . Since $h\Lambda(\varepsilon, C)$ -matches f , it follows that $(\zeta_l(F_{\text{HN}}(g_{T_m} h\Lambda)_{H,\bullet}(t)))_{l=1}^r$ are the r largest elements in $\{\zeta_l F_{\text{HN}}(g_t h\Lambda)_{H,\bullet} : l = 1, \dots, n-1\}$ and are larger than their successors by at least $\zeta_2 - 4C$. Recall from Subsection 8.3 that $d_H(g_{T_m} h_\infty h_{m+1}^{-1} g_{-T_m}) \leq 1$. Hence, if $\zeta_2 - 4C$ is large enough, $(\zeta_l(F_{\text{HN}}(g_{T_m} h_m \Lambda)_{H,\bullet}))_{l=1}^r$ are the largest among all $\{\zeta_l(F_{\text{HN}}(g_{T_m} h_m \Lambda)) : 1 \leq l \leq n-1\}$.

Denote by $\Gamma_{l_i} \subseteq \Lambda$ the lattices that satisfy $g_t h \Gamma_{l_i} = F_{\text{HN}}(g_t h \Lambda)_{\Gamma, l_i}$ for $t = T_m, T_{m+1}$. Then $g_t h_{m+1} \Gamma_{l_i} = F_{\text{HN}}(g_t h_m \Lambda)_{\Gamma, l_i}$.

Fix $\varepsilon_1 > 0$, and choose C_1 such that if for some $1 \leq l \leq n-1$ and $V \in \text{Gr}(l, n)$ we have $g_t V \in U_\varepsilon(\text{Gr}(l, n)_{f_E}^g)$ for every $t \in [-C_1, C_1]$, then $g_t V \in U_{\varepsilon_1}(\text{Gr}(l, n)_{f_E}^g)$. Such a constant exists by Theorem 2.7. Since $h\Lambda(\varepsilon, C)$ -matches f , it follows that

$$\forall 1 \leq i \leq r, \text{span } g_t h \Gamma_{l_i} \in U_\varepsilon(\text{Gr}(l_i, n)_{f_{E, l_i}}^g) \text{ for } T_m - C_1 \leq t \leq T_{m+1} + C_1.$$

Consequently

$$\text{span } g_t h \Gamma_{l_i} \in U_{\varepsilon_1}(\text{Gr}(l_i, n)_{f_{E, l_i}}^g) \text{ for } t = T_m, T_{m+1}.$$

Let $M \in \text{SL}_n(\mathbb{R})$ be a matrix with $d_{\text{SL}_n(\mathbb{R})}(M, I) = O(\varepsilon_1)$ such that

$$\forall 1 \leq i \leq r, M \text{span } g_{T_{m+1}} h \Gamma_{l_i} \in \text{Gr}(l_i, n)_{f_{E, l_i}(T_{m+1})}^g.$$

Using Lemma 2.3 we get $h' \in B_{d_H}(\text{Id}; O(\varepsilon_1)) \subseteq B_{d_H}(\text{Id}; 1)$ such that

$$\forall 1 \leq i \leq r, h' \text{span } g_{T_{m+1}} h \Gamma_{l_i} \in \text{Gr}(l_i, n)_{\rightarrow f_{E, l_i}}^{O(\varepsilon_1).g}.$$

Then, $h'' := g_{-T} h' g_T \in B_{d_\varphi}(\text{Id}; \exp(-T))$ satisfies $h'' g_{T_m} h_\infty \Gamma \in \text{Gr}(l_i, n)_{\rightarrow f_{E, l_i}}^g$.

Since $d_\varphi(h_\infty, h_{m+1}) \leq \exp(-T_{m+1})$, we have $g_{T_m} h_\infty h_{m+1}^{-1} g_{-T_m} \in B_{d_\varphi}(\text{Id}; \exp(-T))$.

Observe that

$$d_\varphi(h'' \cdot g_{T_m} h h_{m+1}^{-1} g_{-T_m}, \text{Id}) \leq 2^{\alpha_\varphi}.$$

Since $\text{span } g_{T_m} h_\infty \Gamma_{l_i} \in U_{\varepsilon_1}(\text{Gr}(l_i, n)_{f_{E, l_i}}^g)$ we have

$$\text{span } g_{T_m} h_{m+1} \Gamma_{l_i} = (g_{T_m} h_\infty h_{m+1}^{-1} g_{-T_m})^{-1} g_{T_m} h \Gamma_{l_i} \in U_\varepsilon(\text{Gr}(l_i, n)_{f_{E, l_i}}^g),$$

provided that ε_1 is small enough as a function of ε , and T is large enough.

As a result, the m -th step is either an Idle Step or an Interruption Step, in which Bob interrupted more than r indices. Otherwise, Bob's choice for \tilde{h}_{m+1} contradicts the existence of h'' . By Lemma 7.13, we conclude that $\#A_{m+1} = O(\exp(T\delta(f_{E,\bullet}^{\rho(\zeta)}(t))))$. \square

Corollary 10.3. *There exists a $\zeta_1 \gg T$ such that the following is true. Let $\zeta_3 > 0$ and let $m \geq 0$ be a step for which $f^{\rho(\zeta_3)}$ has no non-null vertices $t \in [T_m - C - C_1, T_{m+1} + C + C_1]$. Then $\#A_{m+1} = O(\exp(T\delta(f_{E,\bullet}^{\rho(\zeta_1+\zeta_3)}(T_m))))$.*

Proof. For every $\zeta > 0$ consider the map

$$\psi_\zeta : [0, \zeta_2] \times [T_m - C - C_1, T_{m+1} + C + C_1] \rightarrow \{\text{height sequences}\},$$

which assigns

$$(\zeta_0, t) \mapsto f_{E,\bullet}^{\rho(\zeta+\zeta_0)}(t).$$

Note that if f has no non-null vertex in $[T_m - C - C_1, T_{m+1} + C + C_1]$, then the total length of $\{\zeta > 0 : \psi_\zeta \text{ is non-constant}\}$ is $O(T)$. We get that ψ_ζ is constant for some $\zeta_3 \leq \zeta \leq \zeta_1 + \zeta_3$ provided that ζ_3 is large enough. Hence, the conditions of Lemma 10.2 are met for $f^{\rho(\zeta_1)}$ and ζ . The result follows since $\delta(f_{E,\bullet}^{\rho(\zeta)}(T_m))$ is nondecreasing as we increase ζ to $\zeta_1 + \zeta_3$. \square

By Lemma 7.13, for the empty direction filtration $0 \subseteq E_{\text{all}}$ one has that $\#A_{m+1} = O(\exp(DT))$.

By Corollary 10.3, for every $\zeta_3 > 0$ we have

$$\begin{aligned} \sum_{m'=1}^m \#A_{m'} &\leq \int_0^{T_{m+1}} \log \delta(f_{E,\bullet}^{\rho(\zeta_1+\zeta_3)}(t)) dt + O(m) \\ &\quad + D(2C + T + 1) \#\{\text{non-null vertices of } f^{\rho(\zeta_3)} \text{ until } T_{m+1}\}. \end{aligned}$$

Since $t \mapsto \zeta_l(f_{H,\bullet}(t))$ is a Lipschitz function, $\max_{t \in J} \zeta_l(f_{H,\bullet}(t)) = O(\text{vol}(J))$ for every nontriviality interval J of f at l . Consequently, only connected components J of $U_l(f)$ with $\text{vol}(J) = \Omega(\zeta_3)$ are such that the function $\zeta_l(f_{H,\bullet}^{\rho(\zeta_3)}(t))|_J$ does not vanish. There are at most $O(Tm/\zeta_3 + 1)$ connected components of $U_l(f)$ that are larger than ζ_3 and intersect $[0, T_{m+1}]$. Each of them contains at most n^2 non-null vertices. It follows that the number of non-null vertices of $f^{\rho(\zeta_3)}$ until T_{m+1} is $O(Tm/\zeta_3 + 1)$. Consequently,

$$\sum_{m'=1}^m \#A_{m'} \leq \int_0^{T_{m+1}} \log \delta(f_{E,\bullet}^{\rho(\zeta_1+\zeta_3)}(t)) dt + O(m + T^2m/\zeta_3),$$

and hence for every $\zeta_3 > 0$ the payoff is

$$\begin{aligned} D_{\text{Bob}}^T &:= \liminf_{m \rightarrow \infty} \frac{1}{Tm} \sum_{m'=1}^m \#A_{m'} \\ &\leq \liminf_{m \rightarrow \infty} \int_0^{T_{m+1}} \log \delta(f_{E, \bullet}^{\rho(\zeta_1 + \zeta_3)}(t)) dt + O(T^{-1} + T/\zeta_3) \\ &= \Delta_0(f^{\rho(\zeta_1 + \zeta_3)}) + O(T^{-1} + T/\zeta_3). \end{aligned}$$

Therefore, $D_{\text{Bob}}^T \leq \Delta(f) + O(T^{-1})$, as desired.

11. THE DIMENSION OF THE SET OF DIVERGENT TRAJECTORIES

In this section we will prove Corollary 2.32. In view of Theorem 2.31 we will be interested in computing $\sup_{f \in \mathcal{F}} \Delta(f)$ for the set \mathcal{F} of g -templates that correspond to divergent trajectories. Recall that

$$D := \dim_{\mathbf{H}}(H; d_\varphi) = \delta(\emptyset \subseteq E_{\text{all}}) = \sum_{\eta, \eta' \in E_{\text{all}}} (\eta - \eta')^+.$$

Denote

$$\Xi = \sum_{\eta \in E_{\text{all}}} \eta^+.$$

The following two lemmas together with Theorem 2.31 imply Corollary 2.32. The first implies the upper bound and the second implies the lower bound. Let $\mathcal{F}_{\text{sing}} := \{g\text{-template } f : f_{H,1} \rightarrow \infty\}$. It follows that $\text{Sing}(H, \Lambda; g_t) = Y_{\Lambda, \mathcal{F}_{\text{sing}}}$.

Lemma 11.1. $\Delta_0(f) \leq D - \Xi$ for every $f \in \mathcal{F}_{\text{sing}}$.

Lemma 11.2. *There is a g -template $f \in \mathcal{F}_{\text{sing}}$ with $\Delta(f) = D - \Xi$, provided that $n \geq 3$.*

11.1. Single Dimension Analysis. We will study the behavior of the g -template f only on a single l . Fix $n_+, n_- \geq 1$ such that there are at most n_+ positive η_i -s, at most n_- negative η_i -s, and $n_+ + n_- = n$. (We use this ambiguous notation to preserve symmetry). Fix $1 \leq l \leq n-1$. Denote

$$\beta_l := \begin{cases} \frac{n_+}{l}, & \text{if } l \leq n_+, \\ \frac{n_-}{n-l}, & \text{if } l > n_+. \end{cases}$$

For $E \in \mathcal{I}_l$ denote $\delta(E) := \delta(\emptyset \subseteq E \subset E_{\text{all}})$.

Lemma 11.3. *For every $E \in \mathcal{I}_l$ we have*

$$\delta(E) - \beta_l \eta_E \leq D - \Xi$$

Lemma 11.3 is analogous to [7, Lemma 8.1] and contains the combinatorial essence of [12, Proposition 3.1].

Remark 11.4. The inequality is important because we need to integrate numbers of the form $\delta(f_{E,l}(t))$ in order to compute $\Delta_0(f)$ for a g -template f . Lemma 11.3 will enable us to transform the problem into bounding an integral of $\frac{d}{dt}f_{H,l}(t)$, which is more accessible.

Remark 11.5. In general equality occurs in Lemma 11.3 only if $l = 1, n - 1$. In the case $l = 1$, if $E_{\text{all}} \cap \{0\} = \emptyset$, then equality occurs at $E = \{\eta\}$, whenever η is the maximal negative or the minimal positive element in E_{all} . If $E_{\text{all}} \cap \{0\} \neq \emptyset$, then equality occurs for $E = \{0\}$, and perhaps at the maximal negative or the minimal positive elements in E_{all} , depending on the choice of n_+, n_- .

Proof. We reduce the proof to a polynomial inequality on integers, which will be proved in the appendix.

Note that

$$D - \delta(E) = \sum_{\substack{\eta \in E_{\text{all}} - E \\ \eta' \in E}} (\eta - \eta')^+.$$

Consequently, we need to show that

$$(11.1) \quad \sum_{\substack{\eta \in E_{\text{all}} - E \\ \eta' \in E}} (\eta - \eta')^+ + \beta_l \eta_E \geq \Xi.$$

Let $E' \subseteq \{1, \dots, n\}$ be a set for which $E = \{\eta_j : j \in E'\}$. Inequality (11.1) becomes

$$(11.2) \quad V(\eta_1, \dots, \eta_n) := \sum_{\substack{j \in \{1, \dots, n\} \setminus E' \\ j' \in E'}} (\eta_j - \eta_{j'})^+ + \beta_l \sum_{j \in E'} \eta_j - \sum_{j=n_-+1}^n \eta_j \geq 0.$$

In this form the inequality is symmetric with respect to the mapping

$$\eta_j \mapsto -\eta_{n-j+1}, \quad E \mapsto \{1 \leq j \leq n : n - j + 1 \notin E\}, \quad n_+ \leftrightarrow n_-,$$

and hence we may assume that $l \leq n_+$.

Inequality (11.2) is a linear homogeneous inequality in $(\eta_i)_{i=1}^n$, which satisfy the linear inequalities

$$(11.3) \quad \eta_1 \leq \dots \leq \eta_{n_-} \leq 0 \leq \eta_{n_-+1} \leq \dots \leq \eta_n,$$

$$(11.4) \quad \sum_{i=1}^n \eta_i = 0.$$

The set of η_i satisfying (11.3) and (11.4) is a cone defined by the rays

$$P_{a,b,c} := \left(\underbrace{-c, \dots, -c}_{a \text{ times}}, \underbrace{0, \dots, 0}_{b \text{ times}}, \underbrace{a, \dots, a}_{c \text{ times}} \right); \quad 1 \leq a \leq n_-, 1 \leq c \leq n_+, a+b+c = n.$$

It is left to prove that $V(P_{a,b,c}) \geq 0$ for every such a, b, c . Denote

$$a_0 = \#E' \cap \{1, \dots, a\}, b_0 = \#E' \cap \{a+1, \dots, a+b\}, c_0 = \#E' \cap \{a+b+1, \dots, n\}.$$

Since $\beta_l = \frac{n_+}{l} = \frac{n_+}{a_0+b_0+c_0}$, it remains to show that

$$(11.5) \quad \begin{aligned} V(P_{a,b,c}) &= (c - c_0)a_0(a + c) + (b - b_0)a_0c + (c - c_0)b_0a \\ &\quad + \frac{n_+}{a_0 + b_0 + c_0}(-a_0c + c_0a) - ac \geq 0. \end{aligned}$$

This inequality is proved in Appendix 12.1. \square

11.2. Upper Bound on Divergent Trajectories — Lemma 11.1.

Let f be a g -template on I . For every $t \in \bigcup_{l=1}^{n-1} U_l(f)$ denote by

$$\alpha_f(t) := \min_{l=1, \dots, n-1} \beta_l f_{H,l}(t).$$

This is a nonpositive continuous piecewise linear function, being the minimum of a family of such functions. Note that $\alpha_f(t) < 0$ for every $t \in \bigcup_{l=1}^{n-1} U_l(f)$. Since β_l is convex in l , the minimum must be attained at a point $l_f(t) \in L_f(t) \setminus \{0, n\}$.

Consequently, for every differentiable point $t \in \bigcup_{l=1}^{n-1} U_l(f)$ of α_f we have

$$\frac{d}{dt} \alpha_f(t) = \frac{d}{dt} \beta_{l_f(t)} f_{H,l_f(t)}(t) = \beta_{l_f(t)} \eta_{f_{E,l_f(t)}}(t).$$

The following lemma follows, and in turn implies Lemma 11.1.

Lemma 11.6. *For every interval $(t_0, t_1) \subseteq \bigcup_{l=1}^{n-1} U_l(f)$,*

$$\int_{t_0}^{t_1} \delta(f_{E,*}(t)) dt \leq (t_0 - t_1)(D - \Xi) + \alpha_f(t)|_{t=t_0}^{t_1}.$$

Proof. By Lemma 11.3,

$$(11.6) \quad \delta(f_{E,*}(t)) \leq \delta(f_{E,l_f(t)}(t)) \leq D - \Xi + \beta_{l_f(t)} \eta_{f_{E,l_f(t)}} = D - \Xi + \frac{d}{dt} \alpha_f(t).$$

Integrating Eq. (11.6) from t_0 to t_1 yields the result. \square

11.3. Construction of a Special g -Template — Proof of Lemma 11.2.

For the rest of the section assume $n \geq 3$.

Definition 11.7. We say that E_{all} is *standard* if its elements are proportional to $-(n-1), 1, \dots, 1$. This is the flow corresponding to the standard vector approximation. \heartsuit

The standard case was dealt with in [7, Section 9]. The arguments that we present below deals with the non-standard case, and can be adapted to handle the standard case as well. For the rest of the section assume E_{all} is not standard.

Lemma 11.8 (The connecting template). *Let η_- be the maximal negative weight, and η_+ be the minimal positive weight. There exist a g -template f^η and $C > 0$ such that*

- $f_{H,1}^\eta(t) = \begin{cases} \eta_+ t - 1, & \text{if } t \leq 0, \\ \eta_- t - 1, & \text{if } t \geq 0. \end{cases}$
- For every $t \in \mathbb{R} \setminus [-C, C]$,

$$f_{E,l}^\eta(t) = \begin{cases} \{\eta_+\}, & \text{if } l = 1, t < -C, \\ \{\eta_-\}, & \text{if } l = 1, t > C, \\ *, & \text{otherwise.} \end{cases}$$

Proof. Define

$$f_{E,1}^\eta(t) := \begin{cases} \{\eta_+\}, & \text{if } l = 1, t < -C, \\ \{\eta_-\}, & \text{if } l = 1, t > C, \\ *, & \text{otherwise.} \end{cases}$$

By definition, $t = 0$ is a null point of f^η at $l = 1$, hence must have $f_{E,l}^\eta(0) \neq *$ for some $1 < l < n$. We will use $l = 2$. Define $\tilde{f}_+^\eta(t) := (\eta_- + \eta_n)t - 2$ and let $t_+ := \frac{n}{(n-1)\eta_n + \eta_-} > 0$. Then t_+ is a solution of

$$\tilde{f}_{+,2}^\eta(t_+) = \frac{n-2}{n-1} f_{E,1}^\eta(t_+) = \frac{n-2}{n-1} (\eta_- t_+ - 1).$$

Since E_{all} is not standard, $(n-1)\eta_n + \eta_- > 0$, since instead of considering the sum of all η_i , we take $n-1$ times the largest weight together with another weight. Similarly, set $\tilde{f}_-^\eta(t) := (\eta_+ + \eta_1)t - 2$ and let $t_- := \frac{n}{(n-1)\eta_1 + \eta_+} < 0$. Then t_- is the solution of

$$\tilde{f}_{+,2}^\eta(t_-) = \frac{n-2}{n-1} f_{E,1}^\eta(t_-) = \frac{n-2}{n-1} (\eta_+ t_- - 1).$$

Define

$$f_{E,2}^\eta := \begin{cases} *, & \text{if } t < t_- \text{ or } t_+ < t, \\ \{\eta_+, \eta_1\}, & \text{if } t_- < t < 0, \\ \{\eta_-, \eta_m\}, & \text{if } 0 < t < t_+, \end{cases}$$

with the nontrivial connecting morphism $\{\eta_+, \eta_1\} \rightarrow \{\eta_-, \eta_m\}$ at 0.

Define

$$f_{H,2}^\eta(t) = \begin{cases} \tilde{f}_{-,2}^\eta, & \text{if } t_- \leq t \leq 0, \\ \tilde{f}_{+,2}^\eta, & \text{if } 0 \leq t \leq t_+, \\ \frac{n-2}{n-1} f_{H,1}^\eta(t), & \text{if } t \in (-\infty, t_-] \cup [t_+, \infty). \end{cases}$$

Extrapolate $(f_{H,l}^\eta)_{l=1,2}$ to $f_{H,\bullet}^\eta$ by convexity and the fact that no other $f_{E,l}^\eta$ is nontrivial. \square




FIGURE 8. The connecting g -template with $E_{\text{all}} = \{-1.2, 0.3, 0.9\}$.

We will use Lemma 11.8 to glue simple g -templates and thereby generate more complex ones.

Proof of Lemma 11.2. We will distinguish between two cases:

- (1) If $0 \notin E_{\text{all}}$.
- (2) If $0 \in E_{\text{all}}$.

Case (1): Let η_- be the largest negative weight and η_+ be the smallest positive weight. Let

$$E_{-,1} := \{\eta_-\}, \quad E_{+,1} := \{\eta_+\}, \quad E_{\pm,2} := \{\eta_+, \eta_-\}.$$

For every $t_0 < t_1$ denote by f^{t_0, t_1} the g -template on \mathbb{R} defined by:

$$f_{E, l}^{t_0, t_1}(t) := \begin{cases} E_{-, 1}, & \text{if } t_0 < t < t_{1/2}, l = 1, \\ E_{+, 1}, & \text{if } t_{1/2} < t < t_1, l = 1, \\ *, & \text{otherwise,} \end{cases}$$

with the non-null connecting morphism $E_{-, 1} \rightarrow E_{+, 1}$ and with $t_{1/2} = \frac{\eta_+ t_1 - \eta_- t_0}{\eta_+ - \eta_-}$ the only value for which the differential equation will define a $f_{H, 1}^{t_0, t_1}(t)$ that vanishes at $t = t_1$.

$$f_{H, 1}^{t_0, t_1}(t) := \begin{cases} \eta_-(t - t_0), & \text{if } t_0 \leq t \leq t_{1/2}, \\ \eta_+(t - t_1), & \text{if } t_{1/2} \leq t \leq t_1, \\ 0, & \text{otherwise,} \end{cases}$$

and extrapolate it by convexity to get $f_{H \bullet}^{t_0, t_1}(t)$ for every $t \in \mathbb{R}$.




FIGURE 9. The g -template f^{t_0, t_1} with $E_{\text{all}} = \{-1.2, 0.3, 0.9\}$.

The definition was selected because it satisfies the following equality:

$$\begin{aligned} \int_{t_0}^{t_1} \delta(f_{E, *}^{t_0, t_1}) dt &= (t_{1/2} - t_0)\delta(\{\eta_-\}) + (t_1 - t_{1/2})\delta(\{\eta_+\}) \\ &= (t_1 - t_0)D - (t_{1/2} - t_0) \sum_{\eta \neq \eta_-} (\eta - \eta_+)^+ - (t_1 - t_{1/2}) \sum_{\eta \neq \eta_+} (\eta - \eta_+)^+ \\ &= (t_1 - t_0)D - (t_{1/2} - t_0)(\Xi - n_+ \eta_-) - (t_1 - t_{1/2})(\Xi - n_+ \eta_+) \\ &= (t_1 - t_0)(D - \Xi). \end{aligned}$$

For every pair of intervals $t_0 < t_1, t'_0 < t'_1$ such that $t_{1/2} < t'_0 < t_1 < t'_{1/2}$, we would like to define the *merger* $f' = f^{t_0, t_1} \vee f^{t'_0, t'_1}$ of f^{t_0, t_1} and $f^{t'_0, t'_1}$. Unfortunately, the connecting template protrudes beyond $[t'_0, t_1]$, and hence we will need additional constraints on t_0, t_1, t'_0, t'_1 to define the merger.




FIGURE 10. The merger of f^{t_0, t_1} and $f^{t'_0, t'_1}$, where $t_0 = -5$, $t_1 = 0.5$, $t'_0 = -0.5$, $t'_1 = 12$, and $E_{\text{all}} = \{-1.2, 0.3, 0.9\}$.

Our goal is to ensure that the merger will satisfy $f'_{H,1}(t) = \max(f^{t_0, t_1}_{H,1}(t), f^{t'_0, t'_1}_{H,1}(t))$. Let $t''_0 := \frac{\eta_+ t_1 - \eta_- t'_0}{\eta_+ - \eta_-}$ denote the unique point $t''_0 \in [t_{1/2}, t'_{1/2}]$ for which

$$f^{t_0, t_1}_{H,1}(t''_0) = f^{t'_0, t'_1}_{H,1}(t''_0) = \frac{t_1 - t'_0}{\eta_+^{-1} - \eta_-^{-1}}.$$

Denote

$$\kappa := \frac{1}{\eta_+^{-1} - \eta_-^{-1}} \quad \text{and} \quad C' := C\kappa,$$

where C is the constant from Lemma 11.8. We define the merger $f^{t_0, t_1} \vee f^{t'_0, t'_1}$ only if the following inequality holds:

$$t''_{-1} := t'_0 - (t_1 - t'_0)C' > t_{1/2} \quad \text{and} \quad t''_1 := t_1 + (t_1 - t'_0)C < t'_{1/2}.$$

For $t \in \mathbb{R}$, define \tilde{f}^η by

$$\tilde{f}^\eta_{E,l}(t) := f^\eta_{E,l} \left(\frac{t - t''_0}{(t_1 - t'_0)\kappa} \right), \quad \tilde{f}^\eta_{H,l}(t) := (t_1 - t'_0)\kappa f^\eta_{H,l} \left(\frac{t - t''_0}{(t_1 - t'_0)\kappa} \right).$$

In other words, we translate f^η and apply a homothety so that the increasing and decreasing branches of $\tilde{f}_{H,1}^\eta$ will coincide with the corresponding branches of $f_{H,1}^{t_0,t_1}, f_{H,1}^{t'_0,t'_1}$ at t''_{-1}, t''_1 , respectively. We can then glue the three g -templates $f^{t_0,t_1}|_{(-\infty,t''_{-1}]}, \tilde{f}^\eta|_{[t''_{-1},t''_1]}, f^{t'_0,t'_1}|_{[t''_1,\infty)}$.

Using this method we can merge arbitrarily many g -templates f^{t_0,t_1} . Let $t_0^{(m)} < t_1^{(m)}, m \geq 0$, be an infinite sequence of pairs with $d^m := t_1^{(m)} - t_0^{(m+1)} > 0$ and

$$t_{1/2}^{(m)} \leq t_0^{(m+1)} - C'd_m < t_1^{(m)} + C'd_m < t_{1/2}^{(m+1)}.$$

We can merge the g -templates $(f^{t_0^{(m)},t_1^{(m)}})_{m \geq 0}$ into a single g -template $f = \bigvee_{m=0}^\infty f^{t_0^{(m)},t_1^{(m)}}$. Note that

$$f|_{[t_1^{(m)}+C'd_m, t_0^{(m+2)}-C'd_{m+1}]} = f^{t_0^{(m+1)}, t_1^{(m+1)}}.$$

Assume that

$$(11.7) \quad d_m \xrightarrow{m \rightarrow \infty} \infty,$$

$$(11.8) \quad \frac{d_m}{t_1^{(m)} - t_0^{(m)}} \xrightarrow{m \rightarrow \infty} 0,$$

$$(11.9) \quad \frac{t_1^{(m)} - t_0^{(m)}}{\sum_{m'=0}^m t_1^{(m')} - t_0^{(m')}} \xrightarrow{m \rightarrow \infty} 0.$$

For instance, the sequence $t_0^{(m)} = m^3 - m, t_1^{(m)} = (m+1)^3 + m$ works for m large enough as a function of C' . We now wish to compute

$$\Delta_0(f) = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \delta(f_{E,*}(t)) dt.$$

By Eq. (11.9), it is enough to compute the limit for $T = t_1^{(m)}$. By Eq. (11.8), the regions in which f is not one of the $(f^{t_0^{(m)}, t_1^{(m)}})_{m \geq 0}$ are

negligible. We have

$$\begin{aligned}
\Delta_0(f) &= \lim_{m \rightarrow \infty} \frac{1}{t_1^{(m)}} \int_0^{t_1^{(m)}} \delta(f_{E,*}(t)) dt \\
&= \lim_{m \rightarrow \infty} \frac{1}{t_1^{(m)}} \sum_{m'=0}^m \int_{t_0^{(m')}}^{t_1^{(m')}} \delta(f_{E,*}^{t_0^{(m')}, t_1^{(m')}}(t)) dt \\
&= \lim_{m \rightarrow \infty} \frac{1}{t_1^{(m)}} \sum_{m'=0}^m (t_1^{(m')} - t_0^{(m')})(D - \Xi) \\
&= \lim_{m \rightarrow \infty} \frac{1}{t_1^{(m)}} \sum_{m'=1}^m (t_1^{(m')} - t_1^{(m'-1)})(D - \Xi) \\
&= D - \Xi.
\end{aligned}$$

Similarly, the local minima of $f_{E,1}(t)$ are $(\kappa d_m)_{m \geq 0}$, which tend to infinity, and hence $f_{E,1}(t) \xrightarrow{t \rightarrow \infty} -\infty$, as desired.

Case (2): The construction is similar. We define $f^{t_0, t_1, \varepsilon}$ by

$$\begin{aligned}
f_{H,1}^{t_0, t_1, \varepsilon}(t) &:= \begin{cases} \max((t - t_0)\eta_-, t - t_1\eta_+, -\varepsilon(t_1 - t_0)), & \text{if } t \in [t_0, t_1], \\ 0, & \text{if } t \notin [t_0, t_1], \end{cases} \\
f_{E,l}^{t_0, t_1, \varepsilon}(t) &:= \begin{cases} \left\{ \frac{d}{dt} f_{H,1}^{t_0, t_1, \varepsilon}(t) \right\}, & \text{if } t \in (t_0, t_1) \text{ and } l = 1, \\ *, & \text{otherwise,} \end{cases}
\end{aligned}$$

and extrapolate $f_{H,l}^{t_0, t_1, \varepsilon}$ to $f_{H,\bullet}^{t_0, t_1, \varepsilon}$ by convexity. Let $t_{1/3} < t_{2/3}$ denote




FIGURE 11. The g -template $f_{H,l}^{t_0, t_1, \varepsilon}$ with $E_{\text{all}} = \{-1.2, 0, 0.3, 0.9\}$.

the two non-null vertices of $f^{t_0, t_1, \varepsilon}$ at 1. Then

$$\int_{t_{1/3}}^{t_{2/3}} \delta(f_{E,*}^{t_0, t_1, \varepsilon}(t)) dt = (t_{2/3} - t_{1/3})(D - \Xi).$$

Hence the merger of a sequence of such functions with $\varepsilon \rightarrow 0$ provides the desired function as in Case (1). \square

12. APPENDIX

12.1. Dimension Inequality. Here we will prove the following inequality that was used in the proof of Lemma 11.3.

Lemma 12.1. *Let $a \geq a_0 \geq 0, b \geq b_0 \geq 0, c \geq c_0 \geq 0$ be integers with $a_0 + b_0 + c_0 \geq 1$, and let $n_+ \in [c, c + b]$. Then*

$$\begin{aligned} V(a, b, c, a_0, b_0, c_0, n_+) &:= (c - c_0)a_0(a + c) + (b - b_0)a_0c + (c - c_0)b_0a \\ &\quad + \frac{n_+}{a_0 + b_0 + c_0}(-a_0c + c_0a) - ac \geq 0 \end{aligned}$$

provided that $a_0 + b_0 + c_0 \leq n_+$.

Proof. Since V is linear in n_+ we may assume n_+ is one of its boundary values:

- (1) $n_+ = c + b$,
- (2) $n_+ = c$, or
- (3) $n_+ = a_0 + b_0 + c_0$.

First we will show that we may decrease b until $b = b_0$ or $b > b_0$ and $b + c = n_+ = a_0 + b_0 + c_0$. We will show that

$$dV := V(a, b, c, a_0, b_0, c_0, n_+) - V(a, b - 1, c, a_0, b_0, c_0, n'_+) \geq 0,$$

where $n'_+ = n_+$ in cases 2, 3 or $n'_+ = n_+ - 1$ in case 1. In cases 2, 3 we see that $dV = a_0c \geq 0$. In case 1 we have $dV = \frac{-a_0c + c_0a}{a_0 + b_0 + c_0} + a_0c \geq 0$ as $a_0 + b_0 + c_0 \geq 1$.

The above three cases become four:

- (1) $n_+ = c + b, b = b_0$.
- (2) $n_+ = c, b = b_0$.
- (3) $n_+ = a_0 + b_0 + c_0, b = b_0$.
- (4) $n_+ = a_0 + b_0 + c_0 = c + b$.

Denote $a_1 = a - a_0, b_1 = b - b_0, c_1 = c - c_0$.

Case (1): Note that since $c + b = n_+ \geq a_0 + b_0 + c_0 = a_0 + b + c_0$ we get that $c_1 \geq a_0$. Since $a_0^2 - a_0, b_0^2 - b_0 \geq 0$. By simple algebraic

manipulations,

$$\begin{aligned}
& V(a, b, c, a_0, b_0, c_0, n_+) \cdot (a_0 + b_0 + c_0) \\
&= (a_0^2 - a_0)(a_0c_1 + a_1c_1 + 2c_0c_1 + a_1c_0 + c_0^2 + c_1^2 + a_0c_0 + b_0c_0 + 2b_0c_1) \\
&\quad + (c_1 - a_0)(a_0a_1c_0 + a_0c_0^2 + a_0c_0c_1 + a_0^2c_0 + a_0b_0c_0) \\
&\quad + (b_0^2 - b_0)a_1c_1 \\
&\quad + 2a_0a_1b_0c_1 + a_0b_0^2c_1 + a_1b_0c_0c_1 + a_0b_0c_1^2 + a_0b_0c_0c_1 \\
&\geq 0.
\end{aligned}$$

Case (2): Note that if $b_0 = 0$ then we were in the previous case, hence assume $b_0 \geq 1$. Since $c = n_+ \geq a_0 + b_0 + c_0$ we get that $c_1 \geq a_0 + b_0$. In particular, $c_1 \geq b_0 \geq 1$. By simple algebraic manipulations,

$$\begin{aligned}
& V(a, b, c, a_0, b_0, c_0, n_+) \cdot (a_0 + b_0 + c_0) \\
&= (a_0^2 - a_0)(a_0c_1 + b_0c_1 + c_1^2) \\
&\quad + (b_0 - 1)(a_0a_1c_1 + a_1b_0c_1 + 2a_0c_0c_1) \\
&\quad + (c_1 - 1)(a_0^2c_0 + a_0a_1c_0 + a_1b_0c_0 + a_0c_0^2 + a_0c_0c_1) \\
&\quad + (c_1 - b_0)a_0c_0 \\
&\quad + a_0^2a_1c_1 + a_0^2b_0c_1 + a_0a_1b_0c_1 + a_0b_0^2c_1 + a_0^2c_0c_1 + a_0b_0c_1^2 \\
&\geq 0.
\end{aligned}$$

Case (3): Note that $a_0 + b_0 + c_0 = n \in [c, c + b_0]$. If n was one of the endpoints, we would be in the previous cases. Hence, $c < a_0 + b_0 + c_0 < c + b_0$. Equivalently $a_0 < c_1 < a_0 + b_0$. In particular, $b_0 \geq 2, c_1 \geq 1$.

By simple algebraic manipulations,

$$\begin{aligned}
& V(a, b, c, a_0, b_0, c_0, n_+) \\
&= (b_0 - 2)(a_0c_1 + a_1c_1) \\
&\quad + (c_1 - 1)a_0c_0 \\
&\quad + a_1c_1 + a_0^2c_1 + a_0a_1c_1 + a_0c_1^2 \\
&\geq 0.
\end{aligned}$$

Case (4): The equation defining (4) implies $a_0 = c_1 + b_1$. If we had $b_1 = 0$ then we were in Case (1). Consequently, $a_0 \geq b_1 \geq 1$. By simple

algebraic manipulations,

$$\begin{aligned}
 &V(a, b, c, a_0, b_0, c_0, n_+) \\
 &= (b_1 - 1)(a_0c_0 + a_0c_1) \\
 &\quad + (a_0 - 1)(a_0c_1 + a_1c_1) \\
 &\quad + a_0b_0c_1 + a_1b_0c_1 + a_0c_0c_1 + a_0c_1^2 \\
 &\geq 0
 \end{aligned}$$

□

REFERENCES

- [1] J. An, L. Guan, A. Marnat, and R. Shi *Divergent trajectories on products of homogeneous spaces*, arXiv e-prints (2020).
- [2] A. Borel and J. P. Serre, *Corners and arithmetic groups*, Comment. Math. Helv. **48.1** (1973), 436–491.
- [3] J. W. S. Cassels, *An Introduction to the Geometry of Numbers*, Springer Science & Business Media (2012).
- [4] Y. Cheung, *Hausdorff dimension of the set of singular pairs*, Ann. of Math. (2011), 127–167.
- [5] Yitwah Cheung and Nicolas Chevallier, *Hausdorff dimension of singular vectors*, Duke Math. J. **165.12** (2016), 2273–2329.
- [6] S. G. Dani, *Divergent trajectories of flows on homogeneous spaces and Diophantine approximation*, J. Reine Angew. Math. **359** (1985), 55–89.
- [7] T. Das, L. Fishman, D. Simmons and M. Urbański, *A variational principle in the parametric geometry of numbers, with applications to metric Diophantine approximation*, C. R. Math. **355.8** (2017), 835–846.
- [8] M. Einsiedler and S. Kadyrov, *Entropy and escape of mass for $SL_3(\mathbb{R})/SL_3(\mathbb{Z})$* , Israel J. Math. **190** (2012), 253–288.
- [9] D. R. Grayson, *Reduction theory using semistability*, Comment. Math. Helv. **59** (1984), 600–634.
- [10] L. Guan and R. Shi. *Hausdorff dimension of divergent trajectories on homogeneous spaces*, Compos. Math. **156.2** (2020), 340–359.
- [11] G. Harder, M. S. Narasimhan, *On the cohomology groups of moduli spaces of vector bundles on curves*, Math. Ann. **212** (1974), 215–248.
- [12] S. Kadyrov, D. Kleinbock, E. Lindenstrauss, A. G. Margulis, *Singular systems of linear forms and non-escape of mass in the space of lattices*, J. Anal. Math. **133.1** (2017), 253–277.
- [13] D. Y. Kleinbock, *An extension of quantitative nondivergence and applications to Diophantine exponents*, Trans. Amer. Math. Soc. **360.12** (2008), 6497–6523.
- [14] D. Y. Kleinbock, *Badly approximable systems of affine forms*, J. Number Theory **79.1** (1999), 83–102.
- [15] D. Y. Kleinbock and G. A. Margulis. *Flows on homogeneous spaces and Diophantine approximation on manifolds*, Ann. of Math. (1998), 339–360.
- [16] P. Mattila, *Geometry of Sets and Measures in euclidean Spaces: Fractals and Rectifiability*, Cambridge Studies in Advanced Mathematics, No. 44. Cambridge University Press (1999).

- [17] L. Liao, R. Shi, O. N. Solan, and N. Tamam, *Hausdorff dimension of weighted singular vectors in \mathbb{R}^2* , J. Eur. Math. Soc. **22.3** (2019), 833–875.
- [18] D. Roy, *On Schmidt and Summerer parametric geometry of numbers*, Ann. of Math. (2015), 739–786.
- [19] W. M. Schmidt, *On badly approximable numbers and certain games*, Trans. Amer. Math. Soc. 123 (1966), 27–50.
- [20] W. M. Schmidt, *On parametric geometry of numbers*, Acta Arith. **195.4** (2020), 383–414.
- [21] W. M. Schmidt and L. Summerer, *Diophantine approximation and parametric geometry of numbers* Monatshefte für Mathematik **169.1** (2013), 51–104.
- [22] G. Tomanov and B. Weiss, *Closed orbits for actions of maximal tori on homogeneous spaces*, Duke Math. J. **199.2** (2003) 367–392.

INDEX

- $*$, 9
- $\#$, 7
- \trianglelefteq , 7
 - $E_{\bullet} \trianglelefteq E'_{\bullet}$, 57
- $\|\cdot\|$, 7
- \gg , 16
- a_{\bullet} , 6
- Aligned noise lattice, 73
- α_{φ} , 21
- A_m , 25
- \tilde{A}_{m+1} , 73
- Anchor, 74
- Basic intervals, 41
- $B_d(x; r)$, 16
- Blades, 7
- \mathcal{B}_l^n , 7
- $\text{Bl}\Gamma$, 7
- boundedly expanding action, 18
- B_T , 73
- Category flow, 10
- Complementary interval, 41
- Constructible graph, 33
- $\text{covol}\Gamma$, 5
- $\text{covol}(\Gamma_2/\Gamma_1/\Gamma_0)$, 66
- D , 15
- $\delta(E_{\bullet})$, 14
- $\Delta_0(f)$, 14
- $\Delta(f)$, 14
- $d_{\text{Gr}(l,n)}$, 6
- d_H , 5
- $\dim_{\mathbf{H}}$, 19
- Direction filtration, 6
- Doubling metric, 25
- d_{φ} , 21
- d_{SL_n} , 3
- d_{X_n} , 3
- $e_{a,b}$, 49
- E_{all} , 4
- E_{\bullet} , 6
- e_i , 49
- Elementary flip, 59
- Enticement, 32
 - compatible with a graph, 33
- finitary collection, 32
- satisfaction, 32
- solution, 32
- support, 32
- Equivalent trajectories, 4
- η_E , 8
- η_i , 3
- expansion semi-metric, 18
- f , 11
 - $f_{E,\bullet}$, 11
 - $f_{H,\bullet}$, 11
- \overline{F} , 21
- F_{HN} , 6
- Final multiset, 70
- Final noise lattice, 73
- Flag, 6
- f^{Λ} , 12
- \underline{F} , 21
- Game - (T, g) , 24
- Γ_{\bullet}^m , 72
- $\tilde{\Gamma}_{\bullet}^m$, 72
- γ_* , 72
- \mathcal{G}_f , 11
- $\text{Gr}(l, n)$, 6
 - $\text{Gr}(l, n)_E^g$, 6
 - $\text{Gr}(l, n)_{\rightarrow E}$, 51
 - $\text{Gr}(l, n)_{\rightarrow E}^{\varepsilon, g}$, 57
- g_t , 4
- g -template, 11
 - separated, 27
 - significant, 28
- H , 4
 - H_V , 48
 - $H_{V \rightarrow E}$, 51
- H^- , 4
- H^0 , 4
- H^{-0} , 4
- \check{h} , 55
- Harder-Narasimhan filtration, 5
- Hausdorff dimension, 19
 - with respect to a semi-metric, 20
- Height sequence, 6
- h_{∞} , 25
- h_m , 25

- $\tilde{h}^{m_0 \rightarrow m_1}$, 74
- \mathcal{I}_l^* , 9
- $\tilde{\mathcal{I}}_l$, 6
- Independent shift sequence, 30
- irregular point, 10
- \mathcal{I}_\bullet^* , 10
- $L(-)$, 6
- $\tilde{\Lambda}^m$, 72
- $L_f(t)$, 26
- Lower convex hull, 28
- $m_{H \rightarrow 0H}$, 4
- $\text{mult}(x; E)$, 59
- n , 3
- \mathbb{N}_0 , 60
- Noise lattice, 73
- Nontrivial place, 6
- Nontrivial point, 10
- Nontriviality interval, 11
- Null arrow, 9
- Null point, 10
- $O(-)$, 16
- O_g , 49
- $\Omega(-)$, 16
- p_Γ , 5
- $\partial^2 a_\bullet$, 6
- φ_t , 20
- $\tilde{\pi}^m$, 72
- $(-)^+$, 14
- P_m , 82
- \check{q} , 55
- Rectangle, 81
- Relative covolume, 66
- \mathbb{R}^\times , 7
- Scalar pair, 70
- Shift
 - of g -template f^ρ , 30
 - of height sequence a_\bullet^ρ , 29
 - sequence, 29
- $\text{SL}_n(\mathbb{R})$, 3
- $\text{SL}_n(\mathbb{Z})$, 3
- Standard weights, 98
- Steps of Alice
 - adding, 77
 - changing, 79
 - standard, 79
- Steps of Bob
 - idle, 90
 - interruption, 90
- Template, 11
- $\Theta(-)$, 16
- Transition arrow, 9
- U_ε , 7
- $U_i(f)$, 11
- $V:V_\eta$, 8
- Vanishing number, 90
- V_\bullet , 6
- Vertex, 26
 - impact, 28
 - non-null, 26
 - null, 26
- V^\rightarrow , 51
- $\bigwedge^l \mathbb{R}^n$, 7
- Ξ , 15
- X_n , 3
- $Y_{\Lambda, \mathcal{F}}$, 15
- $Y_{\Lambda, f}$, 16