
Learning Treatment Effects in Panels with General Intervention Patterns

Vivek F. Farias
Sloan School of Management
MIT
Cambridge, MA 02139
vivekf@mit.edu

Andrew A. Li
Tepper School of Business
Carnegie Mellon University
Pittsburgh, PA 15213
aali1@cmu.edu

Tianyi Peng
Department of Aeronautics and Astronautics
MIT
Cambridge, MA 02139
tianyi@mit.edu

Abstract

The problem of causal inference with panel data is a central econometric question. The following is a fundamental version of this problem: Let M^* be a low rank matrix and E be a zero-mean noise matrix. For a ‘treatment’ matrix Z with entries in $\{0, 1\}$ we observe the matrix O with entries $O_{ij} := M_{ij}^* + E_{ij} + \mathcal{T}_{ij}Z_{ij}$ where \mathcal{T}_{ij} are unknown, heterogenous treatment effects. The problem requires we estimate the average treatment effect $\tau^* := \sum_{ij} \mathcal{T}_{ij}Z_{ij} / \sum_{ij} Z_{ij}$. The synthetic control paradigm provides an approach to estimating τ^* when Z places support on a single row. This paper extends that framework to allow rate-optimal recovery of τ^* for general Z , thus broadly expanding its applicability. Our guarantees are the first of their type in this general setting. Computational experiments on synthetic and real-world data show a substantial advantage over competing estimators.

1 Introduction

Consider that we are given a sequence of T observations on each of n distinct ‘units’. In an econometric context, a unit might correspond to a geographic region with the associated sequence of observations corresponding to some economic time series of interest. In e-commerce, a unit may correspond to a customer, with the associated sequence of observations corresponding to site-visits for that customer over time. For each unit, some subset of its observations are potentially impacted by an intervention. This may correspond to a new economic policy in the econometric example, or the application of a new type of promotion in the e-commerce context. The question at hand is to estimate the ‘average treatment effect’ of this intervention. This is a core question in econometrics.

The problem above can be formalized as follows: Let $M^* \in \mathbb{R}^{n \times T}$ be a fixed, unknown matrix and E be a zero-mean random matrix; we refer to $M^* + E$ as the ‘counterfactual’ matrix with each row corresponding to a distinct ‘unit’. A known ‘treatment’ matrix Z with entries in $\{0, 1\}$ encodes observations impacted by an intervention. Specifically, we observe a matrix O with entries $O_{ij} = M_{ij}^* + E_{ij} + \mathcal{T}_{ij}Z_{ij}$ where the \mathcal{T}_{ij} are unknown, heterogenous treatment effects.¹ Our goal is to recover the average treatment effect $\tau^* = \sum_{ij} \mathcal{T}_{ij}Z_{ij} / \sum_{ij} Z_{ij}$.

¹Observable covariates on each unit may also be available; we suppress this aspect here for clarity

The ‘synthetic control’ paradigm [1, 2] has been a transformative approach to addressing this problem. That setup addresses the special case of this problem where a single unit (say, the first row) is treated; i.e. Z has support on a portion of the first row. The approach is conceptually simple: find a linear combination of the untreated units (i.e. all rows other than the first) that approximate the first row on its *untreated* entries. One then uses the same linear combination to impute counterfactual values i.e. the entries of M^* in the support of Z . Under the assumption that M^* is low rank (and other assumptions), one can show that this approach recovers τ^* .

Applications like the e-commerce setup alluded to earlier require we allow for more general intervention patterns, i.e. allow for *general* Z as opposed to the single row (or block) support required in synthetic control. A tantalizing possibility, first raised by [3], is treating entries of M^* in the support of Z as missing and applying matrix completion techniques to impute these counterfactual values. This viewpoint has the benefit of making no assumptions on the treatment effects and yields algorithms that do not require any special structure for Z . However, it is unclear that counterfactual recovery – which is equivalent to matrix recovery with general missingness patterns – is actually possible via such a method, so that recovery guarantees are typically unavailable.

1.1 This Paper

Succinctly, we develop an estimator that recovers the average treatment effect under provably minimal assumptions on Z . This is made possible by a new de-biasing identity and an extension of the entry-wise uncertainty quantification analysis of [4, 5] to general non-random missingness patterns that is of independent interest.

In providing context for our contributions, it is worth asking what one can hope for in this problem. In addition to requiring that M^* have low rank (say r), it is clear that we *cannot* in general expect to recover τ^* absent assumptions on the treatment matrix Z . For instance, we must rule out the existence of a rank r matrix M' , distinct from M^* for which $M' = M^* + \gamma Z$ for some $\gamma \neq 0$, or else identifying τ^* is impossible even if E is identically zero (we could rule this out if Z were not in the tangent space of M^*). Separately, unlike matrix completion, we actually *observe* $M_{ij}^* + \mathcal{T}_{ij} + E_{ij}$ on treated entries. If the heterogeneity in treatment effects is too large, however, it is unclear that these observations are of much value. Thus, in addition to benign assumptions on M^* and E , we make a set of assumptions on (A) the projection of Z onto the tangent space of M^* and (B) the heterogeneity in treatment effects. Against this backdrop, we make the following contributions:

Rate Optimal Estimator: We construct an estimator that achieves rate optimal guarantees for the recovery of the average treatment effect τ^* (Theorem 1) with general treatment patterns. We show under additional assumptions that our estimator is asymptotically normal (Theorem 2).

Minimality of Assumptions: Should the conditions we place on the projection of Z onto the tangent space of M^* be violated by an amount that grows small with problem size, we show *no algorithm* can recover τ^* even with homogeneous treatments (Proposition 2). Our assumptions on heterogeneity are also shown to be minimal, and satisfied by extant models in the synthetic control literature.

General Treatment Patterns Work: We show that our assumptions on Z are satisfied by general treatment patterns, e.g., requiring that at least a constant fraction of entries are *not* treated or matrices Z with rank polynomially larger than the rank of M^* .

Empirical Performance: We show both for synthetic and real data that our estimator provides a material improvement in empirical performance relative to available alternatives, including matrix completion based estimators and, where applicable, state-of-the-art synthetic control estimators.

1.2 Related Literature

The synthetic control literature pioneered by [1, 2] has grown to encompass sophisticated learning and inferential methods; see [6] for a review. [7, 8, 9] consider a variety of regularized regression techniques to learn the linear combination of untreated units that yields a synthetic control. [10, 11, 12] consider instead the use of principal component regression techniques. [13] propose alternative approaches to imputing counterfactuals by averaging across both untreated units (rows) and time (untreated columns). [14, 15] address inferential questions that arise in synthetic control with the latter providing a permutation test that is generally applicable.

Matrix completion methods present a means to allow for inference with *general* treatment patterns. [3] are among the first to study this, but provide no guarantees on recovering the average treatment

effect. Alternatively methods that do provide guarantees on the recovery of treatment effects via matrix completion tend to make strong assumptions: [16] effectively assume that Z has support on a block (so that traditional synthetic control techniques could also apply), [17] make stationarity assumptions on M^* and require it to be zero-mean and [18] assume that Z has i.i.d. entries (wherein a trivial estimator of the average treatment effect is also applicable).

A distinct (and common) setting for treatment effect estimation presents us with a single observation per unit and a rich set of observed covariates on these units. These observed covariates roughly ensure exchangeability of units across interventions within covariate strata. ML has been broadly used in this setting; eg. [19, 20, 21]. In contrast for our problem, one could view these covariates as latent.

Our estimation procedure begins with first computing a ‘rough’ estimate of the treatment effect via a natural convex estimator; [22, 23, 24] are empirical studies that have uses this estimator. Crucially, we provide a new de-biasing technique that allows for recovery guarantees and exhibits a significant performance improvement relative to this rough estimate. It is also worth noting that this convex estimator also finds application in the related problem of panel data regression; see [25, 26, 27]. State of the art methods there effectively require that Z is dense.

Whereas work on matrix completion with non-standard observation patterns [28, 29, 30, 31] exists this is by and large not obviously useful or applicable to our problem. Instead, we build on a recent program to bridge convex and non-convex formulations for matrix completion [4] and Robust-PCA [5]. That work has provided a pioneering analysis of entry-wise guarantees and uncertainty quantification for convex matrix completion estimators wherein entries remain missing at random. Our work here may be viewed as extending that program to a broad class of non-random missingness patterns, a contribution of important independent interest.

2 Model and Algorithm

We begin by formally defining our problem; we in fact present a generalization to the problem described in the previous section, allowing for *multiple* treatments. Let $M^* \in \mathbb{R}^{n \times n}$ be a fixed rank- r matrix with singular value decomposition (SVD) denoted by $M^* = U^* \Sigma^* V^{*\top}$, where $U^*, V^* \in \mathbb{R}^{n \times r}$ have orthonormal columns, and $\Sigma^* \in \mathbb{R}^{r \times r}$ is diagonal with diagonal entries $\sigma_1 \geq \dots \geq \sigma_r > 0$.² Let $\sigma_{\max} := \sigma_1$, $\sigma_{\min} := \sigma_r$ and $\kappa := \sigma_{\max}/\sigma_{\min}$ be the condition number of M^* . There are k treatments that can be applied to each entry, and for each treatment $m \in \{1, \dots, k\}$, we are given a *treatment matrix* $Z_m \in \{0, 1\}^{n \times n}$ which encodes the entries which have received the m^{th} treatment (0 meaning no treatment, and 1 meaning being treated).³ We then observe a single matrix of *outcomes*:

$$O := M^* + E + \sum_{m=1}^k \mathcal{T}_m \circ Z_m$$

(\circ is the Hadamard or ‘entrywise’ product), where each $\mathcal{T}_m \in \mathbb{R}^{n \times n}$ is an unknown matrix of *treatment effects*, and $E \in \mathbb{R}^{n \times n}$ is a (possibly heterogeneous) random noise matrix. Finally, let $\tau^* \in \mathbb{R}^k$ be the vector of *average treatment effects*, whose m^{th} value is defined as $\tau_m^* := \langle \mathcal{T}_m, Z_m \rangle / \|Z_m\|_1$ and let $\delta_m = \mathcal{T}_m \circ Z_m - \tau_m^* Z_m$ be associated ‘residual’ matrices. Our problem is to estimate τ^* after having observed O and Z_1, \dots, Z_k .

It is worth noting that the representation above is powerful: for instance, it subsumes the setting where the intervention on any entry is associated with a $\{0, 1\}^k$ -valued covariate vector, and the treatment effect observed on that entry is some linear function of this covariate vector plus idiosyncratic noise. Recovery of τ^* is then equivalent to recovering covariate dependent heterogeneous treatment effects.

Our problem also subsumes the synthetic control setting where $k = 1$ and Z_1 must place support on a single row; the focus of our later analysis will be the case where Z_1 is allowed to be general.

The assumptions that we will need to impose in order to state meaningful results can be divided into two groups. The first are assumptions on M^* and E that are, by this point, canonical in the matrix completion literature:

²Note that in contrast to the previous section, we are now assuming *square* matrices (i.e. $n = T$ in the notation of the previous section). This is purely to simplify the notation – for a rectangular n -by- T matrix, all of our theoretical guarantees hold if one swaps n with $\min\{n, T\}$.

³We allow for multiple treatments to be applied to an entry.

Assumption 1 (Random Noise). The entries of E are independent, mean-zero, sub-Gaussian random variables with sub-Gaussian norm bounded by σ : that is, $\|E_{ij}\|_{\psi_2} \leq \sigma$ for every $i, j \in [n]$.

Assumption 2 (Incoherence). M^* is μ -incoherent:

$$\|U^*\|_{2,\infty} \leq \sqrt{\mu r/n} \quad \text{and} \quad \|V^*\|_{2,\infty} \leq \sqrt{\mu r/n},$$

where $\|\cdot\|_{2,\infty}$ denotes the maximum ℓ_2 -norm of the rows of a matrix.

In addition to these standard conditions on M^* and E which we will assume throughout this paper, we will also need to impose conditions on the relationship between M^* and the Z_m 's. Loosely speaking, these conditions preclude treatment matrices Z_m that can be 'disguised' within M^* , in the sense that their projections onto the tangent spaces of M^* are large. Specifically, the formal statements relate to a particular decomposition of the linear space of $n \times n$ matrices, $\mathbb{R}^{n \times n} = T^* \oplus T^{*\perp}$, where T^* is the tangent space of M^* in the manifold consisting of matrices with rank no larger than $\text{rank}(M^*)$:

$$T^* = \{U^*A^\top + BV^{*\top} \mid A, B \in \mathbb{R}^{n \times \text{rank}(M^*)}\}.$$

Equivalently, the orthogonal space of T^* , denoted $T^{*\perp}$, is the subspace of $\mathbb{R}^{n \times n}$ whose columns and rows are orthogonal, respectively, to the spaces U^* and V^* . Let $P_{T^{*\perp}}(\cdot)$ denote the projection operator onto $T^{*\perp}$:

$$P_{T^{*\perp}}(A) = (I - U^*U^{*\top})A(I - V^*V^{*\top}).$$

We will defer the formal statements of the additional conditions to the next section, but suffice to say for now that they assume $P_{T^{*\perp}}(Z_m)$ to be sufficiently large (we will show that lower bounds on the size of $P_{T^{*\perp}}(Z_m)$, in the the precise form of our own conditions, are nearly necessary). In the remainder of this section, we will outline the core contribution of this paper, which is an estimator for τ^* with a provably rate-optimal guarantee.

2.1 A De-biased Convex Estimator

Our estimator is constructed in two steps, stated as Eqs. (1a) and (1b) below:

$$(\hat{M}, \hat{\tau}) \in \underset{M \in \mathbb{R}^{n \times n}, \tau \in \mathbb{R}^k}{\text{argmin}} \quad g(M, \tau) := \frac{1}{2} \left\| O - M - \sum_{m=1}^k \tau_m Z_m \right\|_{\text{F}}^2 + \lambda \|M\|_*, \quad (1a)$$

$$\tau^d := \hat{\tau} - D^{-1}\Delta. \quad (1b)$$

In Eq. (1b), define by $D \in \mathbb{R}^{k \times k}$ the Gram matrix with entries $D_{lm} = \langle P_{\hat{T}^\perp}(Z_l), P_{\hat{T}^\perp}(Z_m) \rangle$, and by $\Delta \in \mathbb{R}^k$ the 'error' vector with components $\Delta_l = \lambda \langle Z_l, \hat{U}\hat{V}^\top \rangle$, where we have let $\hat{M} = \hat{U}\hat{\Sigma}\hat{V}^\top$ be the SVD of \hat{M} , and let \hat{T} denote the tangent space of \hat{M} .⁴

The first step, Eq. (1a), is a natural convex optimization formulation that we use to compute a 'rough' estimate of the average treatment effects. The objective function's first term penalizes choices of M and τ which differ from the observed O , and the second term seeks to penalize the rank of M using the nuclear norm as a (convex) proxy. The tuning parameter $\lambda > 0$, which will be specified in our theoretical guarantees, encodes the relative weight of these two objectives.

After the first step, having $(\hat{M}, \hat{\tau})$ as a minimizer of Eq. (1a), we could simply use $\hat{\tau}$ as our estimator for τ^* . However, a brief analysis of the first-order optimality conditions for (1a) yields a simple, but powerful decomposition of $\hat{\tau} - \tau^*$ that suggests a first-order improvement to $\hat{\tau}$ via *de-biasing*:

Lemma 1. *Suppose $(\hat{M}, \hat{\tau})$ is a minimizer of (1a). Let $\hat{M} = \hat{U}\hat{\Sigma}\hat{V}^\top$ be the SVD of \hat{M} , and let \hat{T} denote the tangent space of \hat{M} . Denote $\hat{E} = E + \sum_m \delta_m \circ Z_m$. Then,*

$$D(\hat{\tau} - \tau^*) = \Delta^1 + \Delta^2 + \Delta^3, \quad (2)$$

where $\Delta^1, \Delta^2, \Delta^3 \in \mathbb{R}^k$ are vectors with components

$$\Delta_l^1 = \lambda \langle Z_l, \hat{U}\hat{V}^\top \rangle, \Delta_l^2 = \langle P_{\hat{T}^\perp}(Z_l), \hat{E} \rangle, \Delta_l^3 = \langle Z_l, P_{\hat{T}^\perp}(M^*) \rangle.$$

⁴Our definition implicitly assumes that D is invertible. We view this as a natural assumption on (the absence of) collinearity in treatments.

Consider this error decomposition, i.e. $\hat{\tau} - \tau^* = D^{-1}(\Delta^1 + \Delta^2 + \Delta^3)$ by Eq. (2), and note that $D^{-1}\Delta^1$ is entirely a function of observed quantities. Thus, it is known and *removable*. The second step of our algorithm, Eq. (1b), does exactly this. The resulting de-biased estimator, denoted τ^d , is the subject of this paper. As an aside, it is worth noting that while de-biased estimators for high-dimensional inference have received considerable attention recently, our de-biasing procedure is algorithmically distinct from existing notions of de-biasing, including those for problems closely related to our own (e.g. matrix completion [32, 18] and panel data regression [27]).

Our main results characterize the error $\tau^d - \tau^*$. The crux of this can be gleaned from the second and third terms of Eq. (2). If \hat{T} is sufficiently ‘close’ to T^* , then Δ^3 becomes negligible (because $P_{T^*\perp}(M^*) = 0$). Showing closeness of \hat{T} and T^* is the main technical challenge of this work. The remaining error, contributed by Δ^2 , can then be characterized as a particular ‘weighted average’ of the (independent) entries of E and the residual matrices δ_m which we show to be min-max optimal.

To conclude this section, we prove Lemma 1 for a single treatment ($k = 1$); the complete proof is a straightforward generalization, completed in Appendix A.

Proof of Lemma 1 for $k = 1$. Since $k = 1$, we suppress redundant subscripts. Consider the first-order optimality conditions of (1a):

$$\langle Z, O - \hat{M} - \hat{\tau}Z \rangle = 0 \quad (3a)$$

$$O - \hat{M} - \hat{\tau}Z = \lambda(\hat{U}\hat{V}^\top + W), \quad (3b)$$

$$\|W\| \leq 1 \quad (3c)$$

$$P_{\hat{T}\perp}(W) = W \quad (3d)$$

(W is called the ‘dual certificate’ in the matrix completion literature [33, 34, 35]). Combining Eq. (3b) and Eq. (3a), we have

$$\langle Z, O - \hat{M} - \hat{\tau}Z \rangle = 0 \implies \langle Z, \lambda(\hat{U}\hat{V}^\top + W) \rangle = 0 \implies \lambda \langle Z, \hat{U}\hat{V}^\top \rangle = -\langle Z, \lambda W \rangle. \quad (4)$$

Next, applying $P_{\hat{T}\perp}(\cdot)$ to both sides of Eq. (3b) and using Eq. (3d):

$$\begin{aligned} P_{\hat{T}\perp}(O - \hat{M} - \hat{\tau}Z) &= \lambda W \\ \implies P_{\hat{T}\perp}(M^*) + P_{\hat{T}\perp}(E + \delta \circ Z) - (\hat{\tau} - \tau^*)P_{\hat{T}\perp}(Z) &= \lambda W, \end{aligned} \quad (5)$$

where the implication is by definition: $O = M^* + E + \tau^*Z + \delta \circ Z$. Finally, substituting λW from Eq. (5) into Eq. (4), we obtain

$$\begin{aligned} \lambda \langle Z, \hat{U}\hat{V}^\top \rangle &= -\langle Z, P_{\hat{T}\perp}(M^*) + P_{\hat{T}\perp}(E + \delta \circ Z) - (\hat{\tau} - \tau^*)P_{\hat{T}\perp}(Z) \rangle \\ \implies (\hat{\tau} - \tau^*) \|P_{\hat{T}\perp}(Z)\|_{\text{F}}^2 &= \lambda \langle Z, \hat{U}\hat{V}^\top \rangle + \langle Z, P_{\hat{T}\perp}(E + \delta \circ Z) \rangle + \langle Z, P_{\hat{T}\perp}(M^*) \rangle \end{aligned}$$

This is equivalent to Eq. (2), completing the proof. \square

3 Theoretical Guarantees

Summarizing so far, our estimator is constructed in two steps: solve the convex program in Eq. (1a) to obtain an initial estimate $(\hat{M}, \hat{\tau})$, then de-bias according to Eq. (1b). While we have presented this estimator in a setting that allows for *multiple* treatments (i.e. $k \geq 1$), for the sake simplicity our results here (Theorems 1 and 2) are restricted to the single treatment (i.e. $k = 1$) setting. Recall that the work on synthetic control, which we seek to extend, is for a single treatment and a particular form of Z_1 (support on a single row); our results in this section are for a single treatment but *general* Z_1 . To ease notation, we will from here on suppress treatment-specific subscripts (Z_1, τ_1 , etc.)

As mentioned earlier, our results require a set of conditions that relate the treatment matrix Z to the tangent space T^* of M^* :

Assumption 3. There exist positive constants C_{r_1}, C_{r_2} such that

$$(a) \quad \|ZV^*\|_{\text{F}}^2 + \|Z^\top U^*\|_{\text{F}}^2 \leq (1 - C_{r_1}/\log(n)) \|Z\|_{\text{F}}^2.$$

$$(b) \left| \langle Z, U^* V^{*\top} \rangle \right| \|P_{T^* \perp}(Z)\| \leq (1 - C_{r_2}/\log(n)) \|P_{T^* \perp}(Z)\|_F^2.$$

Assumption 3 is nearly necessary for identifying τ^* (in a manner made formal by Proposition 2 below). These conditions are mild enough to allow for various treatment patterns that occur in practice, and broadly expand on the set of patterns possible in the synthetic control literature (as we discuss in Section 3.1).

Let $\delta = \mathcal{T} \circ Z - \tau^* Z$ be the matrix of treatment effect ‘residuals.’ Our first result establishes a bound on the error rate of τ^d . Note that δ is a zero-mean matrix and is zero outside the support of Z . Thus the requirement below that $\|\delta\| \leq C_\delta \sigma \sqrt{n}/r$ is mild. It is trivially met in synthetic control settings. It is also easily seen as met when δ has independent, sub-Gaussian entries. Finally as it turns out, the condition can also admit random sub-gaussian matrices with complex correlation patterns; see [36].

Theorem 1 (Optimal Error Rate). *Suppose $\frac{\sigma \sqrt{n}}{\sigma_{\min}} \leq C_1 \frac{1}{\kappa^2 r^2 \log^5(n)}$ and $\|\delta\| \leq C_\delta \sigma \sqrt{n}/r$. Then for any $C_2 > 0$, for sufficiently large n , with probability $1 - O(n^{-C_2})$, we have*

$$|\tau^d - \tau^*| \leq C_e \log(n) \max \left(\frac{\sigma}{\|Z\|_F} \frac{\kappa^3 r n \sigma \log^5(n)}{\sigma_{\min}}, \frac{|\langle P_{T^*}(\delta), P_{T^*}(Z) \rangle|}{\|Z\|_F^2} \right).$$

Here, C_e is a constant depending (polynomially) on $C_1, C_2, C_{r_1}, C_{r_2}, C_\delta$ (where C_{r_1} and C_{r_2} are the constants appearing in Assumption 3).

To begin parsing this result, consider a ‘typical’ scenario in which $\sigma, \kappa, r, \mu = O(1)$, and $\sigma_{\min} = \Omega(n)$. Then Theorem 1 implies that

$$|\tau^d - \tau^*| = \tilde{O} \left(\sigma / \|Z\|_F + |\langle P_{T^*}(\delta), P_{T^*}(Z) \rangle| / \|Z\|_F^2 \right). \quad (6)$$

This is minimax optimal (up to $\log n$ factors), as shown below:

Proposition 1 (Minimax Lower Bound). *For any estimator $\hat{\tau}$, there exists an instance with $\sigma, \kappa, r, \mu = \Theta(1)$, and $\sigma_{\min} = \Theta(n)$, on which, with probability at least $1/3$,*

$$|\hat{\tau} - \tau^*| \geq \max \left(\sigma / \|Z\|_F, |\langle P_{T^*}(\delta), P_{T^*}(Z) \rangle| / \|Z\|_F^2 \right).$$

Finally, it is worth considering some special cases under which Eq. (6) reduces further to

$$|\tau^d - \tau^*| = \tilde{O} \left(\sigma / \|Z\|_F \right), \quad (7)$$

which is the optimal rate (up to $\log n$) achievable even when M^* and δ are known. Any of the following are, alone, sufficient to imply Eq. (7):

Independent δ : Independent, sub-gaussian δ_{ij} with $O(1)$ sub-gaussian norm.

Synthetic control and block Z : $\|\delta\|_{\max} = O(1)$, Z consists of an $\ell \times c$ block that is sufficiently sparse: $\sqrt{\ell c}(\ell + c) = O(n)$. For comparison, state-of-the-art synthetic control results [13, 37] require the sparser condition $\ell c(\ell + c) = O(n)$ (though that condition enables asymptotic normality results).

Panel data regression: The conditions imposed in [27], the most notably that Z be sufficiently dense: $\|P_{T^* \perp}(Z)\|_F^2 = \Theta(n^2)$. This recovers their error guarantee (up to \log factors); see Appendix D.

Our second main result establishes asymptotic normality for our estimator. This naturally requires some additional control over the variability of δ . We consider the setting in which the $\delta_{ij} = \mathcal{T}_{ij} - \tau^*$ are independent variables.

Theorem 2 (Asymptotic Normality). *Suppose each δ_{ij} is a mean-zero, independent random variable with sub-Gaussian norm $\|\delta_{ij}\|_{\psi_2} = O(1)$. Assume $\kappa = r = \mu = \sigma = O(1)$ and $\sigma_{\min} = \Omega(n)$. Then with probability $1 - O(1/n^3)$,*

$$\tau^d - \tau^* = \frac{\langle E + (\delta \circ Z), P_{T^* \perp}(Z) \rangle}{\|P_{T^* \perp}(Z)\|_F^2} + O \left(\frac{\log^8(n)}{n} \right).$$

Consequently,

$$(\tau^d - \tau^*) / V_\tau^{1/2} \rightarrow \mathcal{N}(0, 1), \quad V_\tau = \sum_{ij} P_{T^* \perp}(Z)_{ij}^2 \text{Var}(E_{ij} + \delta_{ij} Z_{ij}) / \left(\sum_{ij} P_{T^* \perp}(Z)_{ij}^2 \right)^2,$$

provided that $V_\tau^{1/2} = \Omega(\log^9(n)/n)$.

Asymptotic normality is of econometric interest, as it enables *inference*. Specifically, inference can be performed using a ‘plug-in’ estimator \hat{V}_τ for V_τ , gotten by substituting \hat{T} for T^* and $O - M^d - \tau^d Z$ for $E + (\delta \circ Z)$, where M^d is a de-biased estimator for M^* (see Appendix L). This is a common procedure in the literature (e.g., see [18]), and it is straightforward to show that $\hat{V}_\tau \sim V_\tau$.

Proof Technique: The proofs of Theorems 1 and 2 are inspired by recent developments on bridging convex and non-convex formulations for matrix completion [4] and Robust-PCA [5]. Whereas that work assume a random, independent missingness pattern, our proof extends that program to deal with deterministic treatment patterns Z . As such, this analysis is likely of interest, in its own right, as a complement to the matrix completion literature [38, 39, 4, 5]. Broadly, we must address the issue that constructing a dual certificate to analyze the quality of our convex estimator directly is hard. Instead [4] show the existence of such a certificate non-constructively by studying a non-convex estimator and showing that a (fictitious) gradient descent algorithm applied to that estimator recovers a suitable dual certificate. We effectively extend that program to deterministic patterns Z , and provide entry-wise recovery guarantees on M^* in this setting that are of independent interest.

3.1 Applicability of Treatment Conditions

Having stated our results, we return to our assumptions on Z (Assumption 3) and discuss the extent to which they allow for treatment patterns that occur in practice. First however, we establish that Assumption 3 is necessary. Consider the proposition below, which establishes that should either of the condition of Assumption 3 be violated by an amount that grows negligible with n , then identification is rendered impossible so that no estimator can recover τ^* .

Proposition 2. *Let n be any positive even integer. There exists a matrix $Z \in \{0, 1\}^{n \times n}$ and a pair of rank-1 and $\mu = 2$ (incoherent) matrices $M_1, M_2 \in \mathbb{R}^{n \times n}$ with SVDs denoted by $M_i = U_i \Sigma_i V_i^\top$ and T_i being the tangent space of M_i , such that all three of the following statements hold:*

1. $\|ZV_i\|_F^2 + \|Z^\top U_i\|_F^2 = \|Z\|_F^2$, $i = 1, 2$
2. $|\langle Z, U_i V_i^\top \rangle| \|P_{T_i^* \perp}(Z)\| = \|P_{T_i^* \perp}(Z)\|_F^2$, $i = 1, 2$
3. $M_1 + Z = M_2$

We next discuss various treatment patterns that are admissible under Assumption 3:

1. *Rank grows faster than r :* Assumption 3 holds if $\sum_{i=1}^r \sigma_i(Z)^2 \leq (1 - C/\log n) \|Z\|_F^2 / (\sqrt{r} + 2)$. Loosely speaking, this requires that the rank of Z be strictly higher than r , and that less than $1/\sqrt{r}$ of its ‘mass’ lie in its first r components. Put another way, Z must be sufficiently different from any rank- r approximation. One common setting where this occurs is when there is sufficient randomness in generating Z , such as the case where the entries of Z are drawn independently, which is the canonical scenario in the matrix completion literature (e.g., [35, 38, 39]).

2. *Maximal number of ones in a row and column:* Let ℓ and c denote the maximum number of ones in a row and column, respectively, of Z . Assumption 3 holds if $\ell + c \leq (1 - C/\log n) n / (r^2 \mu)$. When $r, \mu = O(1)$, this allows $\ell, c = O(n)$ and generalizes the sparse block patterns studied in the literature (e.g. [40, 13]), where Z is a two-by-two block matrix with exactly one block equal to one.

3. *Single row or column (Synthetic Control):* Consider the case when Z is supported on a single row (or column, equivalently), as in synthetic control. Assumption 3 holds if $\|z^\top V^*\|_F^2 \leq (1 - C/\log n - \mu r/n) \|z\|^2$, where z^\top is the non-zero row of Z . This will easily hold, if allowing a negligible perturbation to either z_1 or the row space of M^* . It is also interesting to note that the identification assumption made in the canonical paper [2] (i.e., $T_0^{-1} \sigma_{\min}(\sum_{i=1}^{T_0} V_i^* V_i^{*\top}) > 0$ is bounded away from zero, where $T_0 = \max_{z_i=0} i$), together with $T_0 = \Omega(n/\log(n))$ and $\mu r = O(n/\log(n))$ (also implicitly assumed in [2] for an optimal guarantee), together imply Assumption 3.

We end this section by drawing a connection to the literature on panel data regression with interactive fixed effects [25, 36, 27]. That literature studies estimators similar in spirits to ours (e.g., [27] analyzed the performance of the convex estimator and a heuristic de-biasing approach). However, those approaches are only known to work if $\|P_{T^* \perp}(Z)\|_F = \Theta(n^2)$ ⁵. This is of course a substantially stronger assumption than Assumption 3 and rules out sparse treatment patterns (as in synthetic

⁵Paranthetically, the assumption $\|P_{T^* \perp}(Z)\|_F = \Theta(n^2)$ greatly simplifies analysis in [27] since a global bound on $\tau - \tau^*$ can be easily obtained. One of the main technical innovations in our paper, building on recent

control). In summary, our approach also has the potential to broaden the scope of problems addressed via panel data regression.

4 Experiments

We conducted a set of experiments on semi-synthetic datasets (the treatment is introduced artificially and thus ground-truth treatment-effect values are known) and real datasets (the treatment is real and ground-truth treatment-effect values are unknown). The results show that our estimator τ^d is more accurate than existing algorithms and its performance is robust to various treatment patterns.

The following four benchmarks were implemented: (i) Synthetic Difference-in-Difference (SDID) [13]; (ii) Matrix-Completion with Nuclear Norm Minimization (MC-NNM) [3] (iii) Robust Synthetic Control (RSC) [10] (iv) Ordinary Least Square (OLS): Selects $a, b \in \mathbb{R}^n, \tau \in \mathbb{R}$ to minimize $\|O - a1^T - 1b^T - \tau Z\|_F^2$, where $1 \in \mathbb{R}^n$ is the vector of ones. It is worth noting that SDID and RSC one apply to traditional synthetic control patterns (*block* and *stagger* below).

Semi-Synthetic Data (Tobacco). The first dataset consists of the annual tobacco consumption per capita for 38 states during 1970-2001, collected from the prominent synthetic control study [2] (the treated unit California is removed). Similar to [3], we view the collected data as M^* and introduce artificial treatments. We considered two families of patterns that are common in the economics literature: *block* and *stagger* [3]. Block patterns model simultaneous adoption of the treatment, while stagger patterns model adoption at different times. In both cases, treatment continues forever once adopted. Specifically, given the parameters (m_1, m_2) , a set of m_1 rows of Z are selected uniformly at random. On these rows, $Z_{ij} = 1$ if and only if $j \geq t_i$, where for block patterns, $t_i = m_2$, and for stagger patterns, t_i is selected uniformly from values greater than m_2 .

To model heterogenous treatment effects, let $\mathcal{T}_{ij} = \tau^* + \delta_i$ where δ_i is i.i.d and $\delta_i \sim \mathcal{N}(0, \sigma_\delta)$ characterizes the unit-specific effect. Then the observation is $O = M^* + \mathcal{T} \circ Z$. We fix $\tau^* = \sigma_\delta = \bar{M}^*/10$ through all experiments, where \bar{M}^* is the mean value of M^* ⁶. The hyperparameters for all algorithms were tuned using rank $r \sim 5$ (estimated via the spectrum of M^*).

Next, we compare the performances of the various algorithms on an ensemble of 1,000 instances with $m_1 \sim \text{Uni}[1, n_1], m_2 = \text{Uni}[1, n_2]$ for stagger patterns and $m_1 \sim \text{Uni}[1, 5], m_2 = 18$ for block patterns (matching the year 1988, where California passed its law for tobacco control). The results are reported in the first two rows of Table 1 in terms of the average normalized error $|\tau - \tau^*|/\tau^*$.

Note that the treatment patterns here are ‘home court’ for the SDID and RSC synthetic control methods but our approach nonetheless outperforms these benchmarks. One potential reason is that these methods do not leverage all of the available data for learning counterfactuals: MC-NNM and SDID ignore treated observations. RSC ignores even more: it in addition does not leverage some of the *untreated* observations in M^* on treated units (i.e. observations O_{ij} for $j < t_i$ on treated units).

Table 1: Comparison of our algorithm (De-biased Convex) to benchmarks on semi-synthetic datasets (Block and Stagger correspond to Tobacco dataset; Adaptive pattern corresponds to Sales dataset). Average normalized error $|\tau - \tau^*|/\tau^*$ is reported.

Pattern	De-biased Convex	SDID	MC-NNM	RSC	OLS
Block	0.31 (± 0.27)	0.46 (± 0.38)	0.54 (± 0.47)	0.60 (± 0.53)	0.75 (± 0.72)
Stagger	0.20 (± 0.20)	0.32 (± 0.37)	0.30 (± 0.32)	0.39 (± 0.54)	0.37 (± 0.38)
Adaptive	0.053 (± 0.04)	-	0.27 (± 0.19)	-	0.40 (± 0.37)

Semi-Synthetic Data (Sales). The second dataset consists of weekly sales of 167 products over 147 weeks, collected from a Kaggle competition [41]. In this application, treatment corresponds to various ‘promotions’ of a product (e.g. price reductions, advertisements, etc.). We introduced an artificial promotion Z , used the collected data as M^* ($\bar{M}^* \approx 12170$), and the goal was to estimate the average treatment effect given $O = M^* + \mathcal{T} \circ Z$ and Z (\mathcal{T} follows the same generation process as above with $\tau^* = \sigma = \bar{M}^*/10$).


advances in the matrix completion literature, is to conduct a refined ‘local’ analysis without the assumption on the density of Z .

⁶See Appendix E for estimating row-specific treatment effects

Now the challenge in these settings is that these promotions are often decided based on previous sales. Put another way, the treatment matrix Z is constructed *adaptively*. We considered a simple model for generating adaptive patterns for Z : Fix parameters (a, b) . If the sale of a product reaches its lowest point among the past a weeks, then we added promotions for the following b weeks (this models a common preference for promoting low-sale products). Across our instances, (a, b) was generated according to $a \in \text{Uni}[5, 25], b \in \text{Uni}[5, 25]$. This represents a treatment pattern where it is unclear how typical synthetic control approaches (SDID, RSC) might even be applied.

The rank of M^* is estimated via the spectrum with $r \sim 35$. See Table 1 for the results averaged over 1,000 instances. The average of $|\tau - \tau^*|/\tau^*$ is $\sim 5.3\%$ for our algorithm, versus 27.6% for MC-NNM. We conjecture that the reason for this is that highly structured missing-ness patterns are challenging for matrix-completion algorithms; we overcome this limitation by leveraging the treated data as well. Of course, there is a natural trade-off here: if the heterogeneity in δ were on the order of the variation in M^* (so that $\|\delta \circ Z\| \gg \sigma_r(M^*)$) then it is unclear that the treated data would help (and it might, in fact, hurt). But for most practical applications, the treatment effects we seek to estimate are typically small relative to the nominal observed values.

Real Data. This dataset consists of daily sales and promotion information of 571 drug stores over 942 days, collected from Rossmann Store Sales dataset [42]. The promotion dataset Z is binary (1 indicates a promo is running on that specific day and store). The *real* pattern is highly complex (see Fig. 1) and hence synthetic-control type methods (SDID, RSC) again do not apply. Our goal here is to estimate the average increase of sales τ^* brought by the promotion.



	τ	Test Error
De-biased Convex	118.2 (± 2.4)	0.04 (± 0.002)
MC-NNM	-49.4 (± 0.98)	0.07 (± 0.002)
OLS	-45.8 (± 1.24)	0.18 (± 0.003)


Figure 1: *Left:* The promotion pattern of the real data. *Right:* Estimation of τ and test errors.

The hyperparameters for all algorithms were tuned using rank $r \sim 70$ (estimated via cross validation). A test set Ω consisting of 20% of the treated entities is randomly sampled and hidden. The test error is then calculated by $\|P_{\Omega}(M + \tau Z - O)\|_{\mathbb{F}}^2 / \|P_{\Omega}(O - \bar{O})\|_{\mathbb{F}}^2$ where \bar{O} is the mean-value of O . Fig. 1 shows the results averaged over 100 instances. Our algorithm provides superior test error. This is potentially a conservative measure since it captures error in approximating both M^* and τ^* ; the variation contributed by M^* to observations is substantially larger than that contributed by τ^* . Now whereas the ground-truth for τ^* is not known here, the negative treatment effects estimated by MC-NNM and OLS seem less likely since store-wise promotions are typically associated with positive effects on sales.

Asymptotic Normality. The normality of our estimator is also verified, where the prediction from Theorem 2 is precise enough to enable inferential tasks such as constructing confidence intervals (CIs) for τ^* : our 95% CIs typically had “true” coverage rates in the range of 93-96% for a synthetic set of instances described in Appendix E. See Fig. 2.

References

- [1] Alberto Abadie and Javier Gardeazabal. The economic costs of conflict: A case study of the basque country. *American economic review*, 93(1):113–132, 2003.
- [2] Alberto Abadie, Alexis Diamond, and Jens Hainmueller. Synthetic control methods for comparative case studies: Estimating the effect of california’s tobacco control program. *Journal of the American statistical Association*, 105(490):493–505, 2010.
- [3] Susan Athey, Mohsen Bayati, Nikolay Doudchenko, Guido Imbens, and Khashayar Khosravi. Matrix completion methods for causal panel data models. *Journal of the American Statistical Association*, pages 1–41, 2021.



	$n_1 = 50$	100	150	200
$n_2/n_1 = 0.5$	0.916	0.957	0.956	0.942
1	0.953	0.946	0.954	0.939
2	0.946	0.947	0.945	0.957
4	0.94	0.934	0.949	0.944

Figure 2: Evaluation of our distributional characterization of τ^d on a synthetic ensemble where δ and E follow i.i.d Gaussian distribution. *Left:* Empirical Distribution of $(\tau^d - \tau^*)/V_\tau$ with $n = 100$, overlaid with the $\mathcal{N}(0, 1)$ density function as predicted by Theorem 2. *Right:* Coverage rates of 95% confidence intervals (the ‘correct’ coverage rate is 0.95) for different sizes (n_1, n_2) with $r = 10$. See Appendix E for the data generation processes in details.

- [4] Yuxin Chen, Yuejie Chi, Jianqing Fan, Cong Ma, and Yuling Yan. Noisy matrix completion: Understanding statistical guarantees for convex relaxation via nonconvex optimization. *arXiv preprint arXiv:1902.07698*, 2019.
- [5] Yuxin Chen, Jianqing Fan, Cong Ma, and Yuling Yan. Bridging convex and nonconvex optimization in robust pca: Noise, outliers, and missing data. *arXiv preprint arXiv:2001.05484*, 2020.
- [6] Alberto Abadie. Using synthetic controls: Feasibility, data requirements, and methodological aspects. *Journal of Economic Literature*, 2019.
- [7] Nikolay Doudchenko and Guido W Imbens. Balancing, regression, difference-in-differences and synthetic control methods: A synthesis. Technical report, National Bureau of Economic Research, 2016.
- [8] Kathleen T Li and David R Bell. Estimation of average treatment effects with panel data: Asymptotic theory and implementation. *Journal of Econometrics*, 197(1):65–75, 2017.
- [9] Eli Ben-Michael, Avi Feller, and Jesse Rothstein. The augmented synthetic control method. *Journal of the American Statistical Association*, (just-accepted):1–34, 2021.
- [10] Muhammad Amjad, Devavrat Shah, and Dennis Shen. Robust synthetic control. *The Journal of Machine Learning Research*, 19(1):802–852, 2018.
- [11] Muhammad Amjad, Vishal Misra, Devavrat Shah, and Dennis Shen. mrsc: Multi-dimensional robust synthetic control. *Proceedings of the ACM on Measurement and Analysis of Computing Systems*, 3(2):1–27, 2019.
- [12] Anish Agarwal, Devavrat Shah, Dennis Shen, and Dogyoon Song. On robustness of principal component regression. *Journal of the American Statistical Association*, (just-accepted):1–34, 2021.
- [13] Dmitry Arkhangelsky, Susan Athey, David A Hirshberg, Guido W Imbens, and Stefan Wager. Synthetic difference in differences, 2019.
- [14] Kathleen T Li. Statistical inference for average treatment effects estimated by synthetic control methods. *Journal of the American Statistical Association*, 115(532):2068–2083, 2020.
- [15] Victor Chernozhukov, Kaspar Wüthrich, and Yinchu Zhu. An exact and robust conformal inference method for counterfactual and synthetic controls. *Journal of the American Statistical Association*, (just-accepted):1–44, 2021.
- [16] Jushan Bai and Serena Ng. Matrix completion, counterfactuals, and factor analysis of missing data. *arXiv preprint arXiv:1910.06677*, 2019.
- [17] Ruoxuan Xiong and Markus Pelger. Large dimensional latent factor modeling with missing observations and applications to causal inference. *arXiv preprint arXiv:1910.08273*, 2019.
- [18] Yuxin Chen, Jianqing Fan, Cong Ma, and Yuling Yan. Inference and uncertainty quantification for noisy matrix completion. *arXiv preprint arXiv:1906.04159*, 2019.
- [19] Susan Athey and Guido Imbens. Recursive partitioning for heterogeneous causal effects. *Proceedings of the National Academy of Sciences*, 113(27):7353–7360, 2016.

- [20] Stefan Wager and Susan Athey. Estimation and inference of heterogeneous treatment effects using random forests. *Journal of the American Statistical Association*, 113(523):1228–1242, 2018.
- [21] Sören R Künzel, Jasjeet S Sekhon, Peter J Bickel, and Bin Yu. Metalearners for estimating heterogeneous treatment effects using machine learning. *Proceedings of the national academy of sciences*, 116(10):4156–4165, 2019.
- [22] Ruoxuan Xiong, Susan Athey, Mohsen Bayati, and Guido W Imbens. Optimal experimental design for staggered rollouts. *Available at SSRN*, 2019.
- [23] Laurent Gobillon and Thierry Magnac. Regional policy evaluation: Interactive fixed effects and synthetic controls. *Review of Economics and Statistics*, 98(3):535–551, 2016.
- [24] Laurent Gobillon and François-Charles Wolff. The local effects of an innovation: Evidence from the french fish market. *Ecological Economics*, 171:106594, 2020.
- [25] Jushan Bai. Panel data models with interactive fixed effects. *Econometrica*, 77(4):1229–1279, 2009.
- [26] Hyungsik Roger Moon and Martin Weidner. Dynamic linear panel regression models with interactive fixed effects. *Econometric Theory*, 33(1):158–195, 2017.
- [27] Hyungsik Roger Moon and Martin Weidner. Nuclear norm regularized estimation of panel regression models. *arXiv preprint arXiv:1810.10987*, 2018.
- [28] Sourav Chatterjee. A deterministic theory of low rank matrix completion. *IEEE Transactions on Information Theory*, 2020.
- [29] Simon Foucart, Deanna Needell, Reese Pathak, Yaniv Plan, and Mary Wootters. Weighted matrix completion from non-random, non-uniform sampling patterns. *arXiv preprint arXiv:1910.13986*, 2019.
- [30] Guangcan Liu, Qingshan Liu, and Xiaotong Yuan. A new theory for matrix completion. In *Advances in Neural Information Processing Systems*, pages 785–794, 2017.
- [31] Olga Klopp, Karim Lounici, and Alexandre B Tsybakov. Robust matrix completion. *Probability Theory and Related Fields*, 169(1-2):523–564, 2017.
- [32] Dong Xia. Confidence interval of singular vectors for high-dimensional and low-rank matrix regression. *arXiv preprint arXiv:1805.09871*, 2018.
- [33] John Wright, Arvind Ganesh, Shankar Rao, Yigang Peng, and Yi Ma. Robust principal component analysis: Exact recovery of corrupted low-rank matrices via convex optimization. In *Advances in neural information processing systems*, pages 2080–2088, 2009.
- [34] Benjamin Recht. A simpler approach to matrix completion. *Journal of Machine Learning Research*, 12(12), 2011.
- [35] Emmanuel J Candès and Benjamin Recht. Exact matrix completion via convex optimization. *Foundations of Computational mathematics*, 9(6):717, 2009.
- [36] Hyungsik Roger Moon and Martin Weidner. Linear regression for panel with unknown number of factors as interactive fixed effects. *Econometrica*, 83(4):1543–1579, 2015.
- [37] Anish Agarwal, Abdullah Alomar, Romain Cosson, Devavrat Shah, and Dennis Shen. Synthetic interventions. *arXiv preprint arXiv:2006.07691*, 2020.
- [38] Emmanuel Abbe, Jianqing Fan, Kaizheng Wang, and Yiqiao Zhong. Entrywise eigenvector analysis of random matrices with low expected rank. *arXiv preprint arXiv:1709.09565*, 2017.
- [39] Cong Ma, Kaizheng Wang, Yuejie Chi, and Yuxin Chen. Implicit regularization in nonconvex statistical estimation: Gradient descent converges linearly for phase retrieval, matrix completion, and blind deconvolution. *Foundations of Computational Mathematics*, pages 1–182, 2019.
- [40] Yiqing Xu. Generalized synthetic control method: Causal inference with interactive fixed effects models. *Political Analysis*, 25(1):57–76, 2017.
- [41] Predict future sales. <https://www.kaggle.com/c/competitive-data-science-predict-future-sales/overview>. Accessed: 2021-01-15.
- [42] Rossmann store sales. <https://www.kaggle.com/c/rossmann-store-sales>. Accessed: 2021-05-26.

- [43] Ji Chen, Dekai Liu, and Xiaodong Li. Nonconvex rectangular matrix completion via gradient descent without $l_{2,\infty}$ regularization. *IEEE Transactions on Information Theory*, 2020.
- [44] Rahul Mazumder, Trevor Hastie, and Robert Tibshirani. Spectral regularization algorithms for learning large incomplete matrices. *Journal of machine learning research*, 11(Aug):2287–2322, 2010.
- [45] Guido W Imbens and Jeffrey M Wooldridge. Recent developments in the econometrics of program evaluation. *Journal of economic literature*, 47(1):5–86, 2009.
- [46] Ali Taylan Cemgil. Bayesian inference for nonnegative matrix factorisation models. *Computational intelligence and neuroscience*, 2009, 2008.
- [47] Vivek F Farias, Andrew A Li, and Tianyi Peng. Near-optimal entrywise anomaly detection for low-rank matrices with sub-exponential noise. *arXiv preprint arXiv:2006.13126*, 2020.
- [48] Yi Yu, Tengyao Wang, and Richard J Samworth. A useful variant of the davis–kahan theorem for statisticians. *Biometrika*, 102(2):315–323, 2015.
- [49] Mark Rudelson, Roman Vershynin, et al. Hanson-wright inequality and sub-gaussian concentration. *Electronic Communications in Probability*, 18, 2013.

Appendix

A Proof of Lemma 1

We begin by stating the first order optimality conditions for Eq. (1a):

$$\left\langle Z_l, O - \hat{M} - \sum_{m=1}^k \hat{\tau}_m Z_m \right\rangle = 0 \quad \text{for } l = 1, 2, \dots, k \quad (8a)$$

$$\begin{aligned} O - \hat{M} - \sum_{m=1}^k \hat{\tau}_m Z_m &= \lambda(\hat{U}\hat{V}^\top + W), \\ \|W\| &\leq 1, \quad P_{\hat{T}^\perp}(W) = W, \end{aligned} \quad (8b)$$

Substituting Eq. (8b) in Eq. (8a) immediately yields

$$\langle Z_l, \hat{U}\hat{V}^\top \rangle = -\langle Z_l, W \rangle \quad (9)$$

Whereas projecting both sides of Eq. (8b) onto \hat{T}^\perp yields

$$P_{\hat{T}^\perp} \left(M^* + \hat{E} + \sum_{m=1}^k (\tau_m^* - \hat{\tau}_m) Z_m \right) = \lambda W$$

where we use the fact that $P_{\hat{T}^\perp}(\hat{M})$ and $P_{\hat{T}^\perp}(\hat{U}\hat{V}^\top)$ are identically zero, and $P_{\hat{T}^\perp}(W) = W$. Together with Eq. (9) this yields

$$\begin{aligned} -\lambda \langle Z_l, \hat{U}\hat{V}^\top \rangle &= \lambda \langle Z_l, W \rangle \\ &= \langle Z_l, P_{\hat{T}^\perp}(M^*) \rangle + \langle Z_l, P_{\hat{T}^\perp}(\hat{E}) \rangle + \sum_{m=1}^k (\tau_m^* - \hat{\tau}_m) \langle Z_l, P_{\hat{T}^\perp}(Z_m) \rangle \\ &= \langle Z_l, P_{\hat{T}^\perp}(M^*) \rangle + \langle P_{\hat{T}^\perp}(Z_l), \hat{E} \rangle + \sum_{m=1}^k (\tau_m^* - \hat{\tau}_m) \langle P_{\hat{T}^\perp}(Z_l), P_{\hat{T}^\perp}(Z_m) \rangle \end{aligned}$$

which is the result.

B Proof Outline of Theorem 1 and Theorem 2

We conclude by describing the main ideas and technical challenges in the proof of Theorem 1 and Theorem 2. The full details can be found in Appendix K and Appendix J. The first challenge is that a direct analysis of the minimizer of $g(M, \tau)$, defined in Eq. (1a), appears to be insufficient. So instead, we introduce the following non-convex *proxy* problem which plays a critical role in the analysis (note that our algorithm itself does *not* involve solving this proxy problem).

$$\underset{X \in \mathbb{R}^{n \times r}, Y \in \mathbb{R}^{n \times r}, \tau \in \mathbb{R}}{\text{minimize}} \quad f(X, Y; \tau) := \frac{1}{2} \|O - XY^\top - \tau Z\|_F^2 + \frac{\lambda}{2} \|X\|_F^2 + \frac{\lambda}{2} \|Y\|_F^2 \quad (10)$$

Eq. (10) can be viewed as minimizing the Euclidean error with a hard constraint $\text{rank}(XY^\top) = r$ and additional regularization terms that serve to maintain an important ‘balance’ between X and Y . The hope is that $X \approx X^*$, $Y \approx Y^*$ where $M^* = X^*Y^{*\top}$ with $X^* = U^*\Sigma^{*1/2}$, $Y^* = V^*\Sigma^{*1/2}$.

The proof of Theorems 1 and 2 involves two major parts: (i) characterizing the estimation error of the non-convex problem $f(X, Y; \tau)$ by an inductive argument, and (ii) showing that one of the critical points of f is extremely close to the optimal points of $g(M, \tau)$. Combining these two parts leads to the guarantees in Theorems 1 and 2. This larger framework is inspired by recent developments on bridging convex and non-convex problems [4, 5] for matrix completion and Robust-PCA. In contrast to those works, which assume a completely random pattern (i.e. independent across entries)

ALGORITHM 1: Gradient Descent of Non-convex Optimization

Input: the observation O and Z

1: **Initialization:** $X^0 = X^*, Y^0 = Y^*, \tau^0 = \frac{\langle Z, O - X^0 Y^{0\top} \rangle}{\|Z\|_{\mathbb{F}}^2}$ where

$$X^* = U^*(\Sigma^*)^{1/2}, Y^* = V^*(\Sigma^*)^{1/2}.$$

2: **Gradient updates:** for $t = 0, 1, \dots, t_* - 1$ **do**

$$\begin{aligned} X^{t+1} &= X^t - \eta[(X^t Y^{t\top} + \tau^t Z - O)Y^t + \lambda X^t]; \\ Y^{t+1} &= Y^t - \eta[(X^t Y^{t\top} + \tau^t Z - O)^\top X^t + \lambda Y^t]; \\ \tau^{t+1} &= \frac{\langle Z, O - X^{t+1} Y^{(t+1)\top} \rangle}{\|Z\|_{\mathbb{F}}^2} \end{aligned}$$

where η determines the learning rate.

of missing or corrupted entries, our deterministic treatment pattern Z necessitates a substantially more-careful analysis.

Below we present more concrete ideas for proving Theorem 2, where Theorem 1 can be viewed as a special case that follows the same proof framework.

Part 1: Properties of the Non-convex Optimizers: In the first part of the proof, we show that an alternating gradient descent algorithm (Algorithm 1) on f , initialized at (X^*, Y^*) , finds (X, Y, τ) with extremely small gradient (and hence is ‘approximately’ a critical point of f), and such that (X, Y) is ‘close’ to (X^*, Y^*) in a way that enables various guarantees on the closeness of T and T^* . Again, note that this gradient descent algorithm exists purely for the purpose of analyzing the properties of the convex optimizer of g , so this idealized choice of initialization is not illegitimate.

Our measure of the accuracy of (X^t, Y^t) should be rotation-invariant (since $X^t H^t H^{t\top} Y^{t\top} = X^t Y^{t\top}$ for any rotation/orthogonal matrix $H \in \mathcal{O}^{r \times r}$). Thus, the coming result is stated for the optimal rotation for aligning (X^t, Y^t) to (X^*, Y^*) :

$$H^t := \arg \min_{R \in \mathcal{O}^{r \times r}} \|X^t R - X^*\|_{\mathbb{F}}^2 + \|Y^t R - Y^*\|_{\mathbb{F}}^2.$$

Furthermore, defining the quantities $F^t, F^*, \nabla f(X, Y; \tau) \in \mathbb{R}^{2n \times r}$ greatly simplifies the presentation:

$$F^t := \begin{bmatrix} X^t \\ Y^t \end{bmatrix}, \quad F^* := \begin{bmatrix} X^* \\ Y^* \end{bmatrix}, \quad \text{and} \quad \nabla f(X, Y; \tau) := \begin{bmatrix} \nabla_X f(X, Y; \tau) \\ \nabla_Y f(X, Y; \tau) \end{bmatrix}.$$

Then, the following lemma provides guarantees for the iteration sequence of Algorithm 1. To simplify the analysis, we use the fact that δ can be absorbed as a part of noise E .

Lemma 2. *Assume Assumption 2 and Assumption 3(a) hold. Suppose $O = M^* + \tau^* Z + E$ where E_{ij} are independent sub-Gaussian random variables with $\|E_{ij}\|_{\psi_2} \leq \sigma$. Suppose*

$$\kappa^4 \mu^2 r^2 \log^2(n) \leq C_1 n \quad \text{and} \quad \frac{\sigma}{\sigma_{\min}} \sqrt{n} \leq C_2 \frac{1}{\kappa^3 \mu r^{4.5} \log^5(n)}.$$

Let $\lambda = C_\lambda \sigma \sqrt{n} \log^{1.5}(n)$, $\eta = \frac{C_\eta}{n^{20} \sigma_{\max}}$, $t_* = n^{C_t}$. For any $C_3 > 0$, for large enough n , with probability $1 - O(n^{-C_3})$, the iteration $(X^t, Y^t, \tau^t)_{0 \leq t \leq t_*}$ given in Algorithm 1 satisfies the following:

$$\|F^t H^t - F^*\|_{\text{F}} \leq C_{\text{F}} \left(\frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_{\text{F}} \quad (12a)$$

$$\|F^t H^t - F^*\|_{2,\infty} \leq C_{\infty} \left(\frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \right) \|F^*\|_{\text{F}} \quad (12b)$$

$$|\tau^t - \tau^*| \leq C_{\tau} \left(\frac{\sigma \mu r^{2.5} \kappa \log^{3.5} n}{\sqrt{n}} + \frac{\sigma \log^{1.5} n}{\|Z\|_{\text{F}}} \right) \quad (12c)$$

$$\min_{0 \leq t \leq t^*} \|\nabla f(X^t, Y^t, \tau^t)\|_{\text{F}} \leq \frac{\lambda \sqrt{\sigma_{\min}}}{n^{10}}. \quad (12d)$$

Furthermore, let T_t be the tangent space of $X^t Y^{t\top}$. With probability $1 - O(n^{-C_3})$, the following hold for all iterations $0 \leq t \leq t^*$ simultaneously:

$$\|P_{T_t^\perp}(M^*)\|_{\infty} \leq C_{T,1} \frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^7(n)}{\sigma_{\min}} \quad (13a)$$

$$\|P_{T_t^\perp}(E) - P_{T^{* \perp}}(E)\|_{\infty} \leq C_{T,2} \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^2 \log^4(n)}{\sigma_{\min}} \quad (13b)$$

$$\|P_{T_t^\perp}(Z) - P_{T^{* \perp}}(Z)\|_{\text{F}} \leq C_{T,3} \frac{r \kappa \sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \|Z\| \quad (13c)$$

$$\|P_{T_t^\perp}(Z) - P_{T^{* \perp}}(Z)\|_{\infty} \leq C_{T,4} \frac{r^{3.5} \kappa^{1.5} \sigma \mu^{1.5} \log^{3.5}(n)}{\sigma_{\min}} \left(\|Z\|_{2,\infty} + \|Z^\top\|_{2,\infty} \right). \quad (13d)$$

Here, $C_\lambda, C_\eta, C_t, C_{\text{F}}, C_\infty, C_\tau, C_{T,1}, C_{T,2}, C_{T,3}, C_{T,4}$ are constants depending (polynomially) on C_1, C_2, C_3, C_{r_1} (where C_{r_1} is the constant in Assumption 3(a)).

To provide some intuition, the goal of Lemma 2 is to establish that there exists (X, Y, τ) such that the gradient of the non-convex function f is extremely small (Eq. (17b)), while (X, Y, τ) is close to (X^*, Y^*, τ^*) (Eqs. (12b), (12c) and (17a)). In addition, there are various guarantees for T (Eq. (13)), which are desired given the results of Lemma 1.

Technical challenge. The proof of Lemma 2 is based on the framework of iteration analysis and leave-one-out techniques developed in [39, 4, 43] with significant improvement for this problem. The main difficulty here brought by the deterministic pattern of Z is the interdependence between $|\tau^t - \tau^*|$ and $\|F^t H^t - F^*\|_{\text{F}}, \|F^t H^t - F^*\|_{2,\infty}$. In particular, a direct application of [39] here will lead to either (i) strong assumptions on Z (e.g., entries in Z are i.i.d occurred) (ii) or divergence of the estimation errors (e.g., increasing $|\tau^t - \tau^*|$ leads to an increase of $\|F^{t+1} H^{t+1} - F^*\|_{\text{F}}$ and $\|F^{t+1} H^{t+1} - F^*\|_{2,\infty}$, and in turn $\|F^{t+1} H^{t+1} - F^*\|_{\text{F}}$ and $\|F^{t+1} H^{t+1} - F^*\|_{2,\infty}$ also increases $|\tau^{t+1} - \tau^*|$).

Decoupling idea. We tackle this issue by ‘decoupling’ this interdependence throughout a much more careful analysis on (i) mitigating the impact of $|\tau^t - \tau^*|$ on $\|F^t H^t - F^*\|_{\text{F}}$; (ii) establishing a new iterative analysis for $|\tau^t - \tau^*|$ without a strong requirement on $\|F^t H^t - F^*\|_{2,\infty}$. This minimizes the additional assumption for Z while maintaining a desired bound for the estimation errors. In addition to tackling the deterministic pattern of Z , we also put significant effort into obtaining various guarantees for the tangent space T^t from the guarantees on X^t, Y^t, τ^t . The full proof details of Lemma 2 can be found in Appendix I.

Part 2: Connecting the Convex and Non-convex Optimizers: Next, we show that the near-critical point of f provided by Lemma 2 is extremely close to the optimal points of the convex problem g . Recall that the first order conditions of the convex program for (M, τ) are: (i) $\tau \|Z\|_{\text{F}}^2 = \langle Z, O - M \rangle$ (Eq. (3a)); and (ii) there exists certificate W such that $O - M - \tau Z = \lambda(UV^\top + W)$ and $\|W\| \leq 1, P_{T^\perp}(W) = W$ (Eqs. (3b) to (3d)). To establish the connection between the two problems, consider

the exact conditions satisfied by the critical points of f :

$$\tau \|Z\|_{\mathbb{F}}^2 = \langle Z, O - XY^\top \rangle \quad (14a)$$

$$(O - XY^\top - \tau Z)Y = \lambda X \quad (14b)$$

$$(O - XY^\top - \tau Z)^\top X = \lambda Y. \quad (14c)$$

We establish a lemma showing that some approximately critical points of f satisfy the first-order conditions of g approximately. In particular, we have the following result. See the detailed proof in Appendix F.

Lemma 3. *Assume Assumption 3(a) and Assumption 3(b) hold, and $\frac{\sigma}{\sigma_{\min}}\sqrt{n} \leq C_1 \frac{1}{\kappa^2 r^2 \log^5(n)}$. Suppose $O = M^* + \tau^* Z + E$ for a deterministic E . Assume $\|E\| \leq C_2 \sigma \sqrt{n}$ with $|\langle E, P_{T^{\perp}}(Z) \rangle| \leq C_3 \sigma \sqrt{n/\log(n)} \|Z\|_{\mathbb{F}}$.*

Let $\lambda = C_\lambda \sigma \sqrt{n} \log^{1.5}(n)$, $X, Y \in \mathbb{R}^{n \times r}$, $\tau = \frac{\langle Z, O - XY^\top \rangle}{\|Z\|_{\mathbb{F}}^2}$, and T is the tangent space of XY^\top .

$$\|X - X^*\|_{\mathbb{F}} + \|Y - Y^*\|_{\mathbb{F}} \leq C_{\mathbb{F}} \left(\frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \right) \|X^*\|_{\mathbb{F}}, \quad (15a)$$

$$\|\nabla f(X, Y; \tau)\|_{\mathbb{F}} \leq \frac{\lambda \sqrt{\sigma_{\min}}}{\kappa n}. \quad (15b)$$

Let $R = \frac{1}{\lambda}(O - XY^\top - \tau Z - \lambda UV^\top)$ where $U\Sigma V^\top$ is the SVD of XY^\top . Then,

$$\|P_T(R)\|_{\mathbb{F}} \leq \frac{72\kappa}{\lambda \sqrt{\sigma_{\min}}} \|\nabla f(X, Y; \tau)\|_{\mathbb{F}} \quad \text{and} \quad \|P_{T^\perp}(R)\| \leq \left(1 - \frac{C}{\log n}\right). \quad (16)$$

Here, C is a constant depending (polynomially) on $C_1, C_2, C_3, C_\lambda, C_{\mathbb{F}}$.

We can view Eq. (16) as an approximated version of Eq. (3) since $\|P_T(R)\|_{\mathbb{F}} \approx 0$ implies $P_{T^\perp}(R) \approx R$ and then $\|P_{T^\perp}(R)\| \leq 1$ approximately implies $\|R\| \leq 1$. If (XY^\top, τ) satisfies the first-order conditions of g approximately, we can expect that (XY^\top, τ) is close to the optimizer of g . Based on the results of Lemma 3, we establish the following key property (see detailed proof in Appendix G).

Lemma 4. *Assume the same setup from Lemma 3. For any minimizer $(\hat{M}, \hat{\tau})$ of the convex program Eq. (1a), we have*

$$\begin{aligned} \|XY^\top - \hat{M}\|_{\mathbb{F}} &\leq C \frac{\kappa \log(n)}{\sqrt{\sigma_{\min}}} \|\nabla f(X, Y; \tau)\|_{\mathbb{F}} \\ |\tau - \hat{\tau}| &\leq C \frac{\kappa \log(n)}{\sqrt{\sigma_{\min}} \|Z\|_{\mathbb{F}}} \|\nabla f(X, Y; \tau)\|_{\mathbb{F}}. \end{aligned}$$

Furthermore, let \hat{T} be the tangent space of \hat{M} . For any matrix $A \in \mathbb{R}^{n \times n}$, we have

$$\begin{aligned} \|P_{\hat{T}^\perp}(A) - P_{T^\perp}(A)\|_{\mathbb{F}} &\leq C \|A\| \frac{\kappa \log(n)}{\sigma_{\min}^{1.5}} \|\nabla f(X, Y; \tau)\|_{\mathbb{F}} \\ \|P_{\hat{T}^\perp}(XY^\top)\|_{\mathbb{F}} &\leq C \frac{\kappa^3 \log^2(n)}{\sigma_{\min}^2} \|\nabla f\|_{\mathbb{F}}^2. \end{aligned}$$

Here, C is a constant depending (polynomially) on $C_1, C_2, C_3, C_\lambda, C_{\mathbb{F}}$.

The proofs of Lemma 3 and Lemma 4 are based on the framework in [4, 5], though we require a more-refined analysis to address the additional terms induced by τ and to limit the assumptions needed on Z . In particular, a direct application of [4] for Lemma 3 would require that $\|Z\| \ll n/\log(n)$ (noting that $1^\top Z 1$ is the number of entries in Z , this implies that Z would need to be sparse), which rules out many interesting patterns in causal inference. In contrast, Assumption 3(b) is notably weaker and almost necessary for the convex optimizers to be close to M^* . In addition, Lemma 4 requires utilizing the special form of τ , along with results for the soft-thresholding estimator [44] to show the claimed ‘closeness’ of \hat{T} to T .

Putting it Together: Given the results in Lemmas 1, 2 and 4, we are ready to finish the proof of Theorem 2. First, the bounds on $|\hat{\tau} - \tau^*|$ and $\|\hat{M} - M^*\|_\infty$. From Lemma 2, an approximate critical point (X, Y, τ) of f can be found with various guarantees, from which we can establish bounds on $|\tau - \tau^*|$ and $\|XY^T - M^*\|_\infty$ through Eqs. (12b) and (12c). Then by Lemma 4, we also obtain that $\tau \approx \hat{\tau}$ and $XY^T \approx \hat{M}$. The desired bounds on $|\hat{\tau} - \tau^*|$ and $\|\hat{M} - M^*\|_\infty$ follow directly.

Finally, in order to prove the guarantee on $\tau^d - \tau^*$ in Theorem 2, recall that by Lemma 1, we have

$$\begin{aligned} \tau^d - \tau^* &= \frac{\langle P_{\hat{T}^\perp}(Z), E \rangle}{\|P_{\hat{T}^\perp}(Z)\|_F^2} + \frac{\langle Z, P_{\hat{T}^\perp}(M^*) \rangle}{\|P_{\hat{T}^\perp}(Z)\|_F^2} \\ &\stackrel{(i)}{=} \frac{\langle P_{T^* \perp}(Z), E \rangle}{\|P_{\hat{T}^\perp}(Z)\|_F^2} + \frac{\langle Z, P_{\hat{T}^\perp}(E) - P_{T^* \perp}(E) \rangle + \langle Z, P_{\hat{T}^\perp}(M^*) \rangle}{\|P_{\hat{T}^\perp}(Z)\|_F^2} \end{aligned}$$

where in (i) we use the fact that for any $A, B \in \mathbb{R}^{n \times n}$, $\langle A, P_{T^* \perp}(B) \rangle = \langle P_{T^* \perp}(A), P_{T^* \perp}(B) \rangle = \langle P_{T^* \perp}(A), B \rangle$. The remainder boils down to controlling the error of $P_{\hat{T}^\perp}(M^*)$, $P_{\hat{T}^\perp}(E) - P_{T^* \perp}(E)$ and showing that $P_{\hat{T}^\perp}(Z) \approx P_{T^* \perp}(Z)$, which matches the bounds provided in Eq. (13) of Lemma 2. Combining the results in Eq. (13) and Lemma 4, together with Berry-Esseen inequalities, we obtain the desired result on $\tau^d - \tau^*$. See details in Appendix K.

Proof of Theorem 1: The proof framework of Theorem 1 is similar to the presented framework above, where the main difference is that we use the following Lemma 5 to replace Lemma 2.

Lemma 5. *Assume Assumption 3(a) holds and $\frac{\sigma}{\sigma_{\min}} \sqrt{n} \leq C_1 \frac{1}{\kappa^2 \tau^2 \log^5(n)}$. Suppose $O = M^* + \tau^* Z + E$ for a deterministic E . Assume $\|E\| \leq C_2 \sigma \sqrt{n}$ and $|\langle Z, E \rangle| \leq C_3 \sigma \sqrt{n} \|Z\|_F$.*

Let $\lambda = C_\lambda \sigma \sqrt{n} \log^{1.5}(n)$, $\eta = \frac{C_\eta}{\kappa^3 n^{20} \sigma_{\max}}$, $t^* = \kappa^5 n^{10000}$. Algorithm 1 satisfies the following:

$$\|F^t H^t - F^*\|_F \leq C_F \left(\frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_F \quad (17a)$$

$$\min_{0 \leq t \leq t^*} \|\nabla f(X^t, Y^t; \tau^t)\|_F \leq \frac{\lambda \sqrt{\sigma_{\min}}}{\kappa n^{10}}. \quad (17b)$$

Here, C_F is a constant depending (polynomially) on $C_1, C_2, C_3, C_\lambda, C_\eta, C_{r_1}$ (where C_{r_1} is the constant in Assumption 3(a)).

In contrast with Lemma 2, Lemma 5 does not assume E_{ij} is independent sub-Gaussian random variable. Although the guarantees in Lemma 5 are only for Frobenius norms, it is sufficient to establish the proof for Theorem 1 by combining Lemma 5 and Lemma 4. See details in Appendix J.

C Basis Transformation for Assumption 3

Here, we present an ‘‘intuitive’’ way to interpret Assumption 3(a) and Assumption 3(b) by a change of basis for Z .

Let $M^* = U^* \Sigma^* V^{*\top}$ be the SVD of M^* where $U^*, V^* \in \mathbb{R}^{n \times r}$ characterize the column and row space of M^* respectively. We consider the expansion of U^* and V^* . Note that $P_{T^* \perp}(Z) = (I - U^* U^{*\top}) Z (I - V^* V^{*\top})$ has the columns and rows orthogonal to the spaces of U^* and V^* . We take $P_{T^* \perp}(Z) = U^\perp \Sigma^\perp V^{\perp \top}$ be the SVD of $P_{T^* \perp}(Z)$. Then $\tilde{U} = [U^*, U^\perp]^\top$ constitutes the basis matrix for the space of \mathbb{R}^n (the same for $\tilde{V} = [V^*, V^\perp]$).

We then consider a basis transformation for Z (left and right basis respectively) based on \tilde{U} and \tilde{V} :

$$\tilde{U}^\top Z \tilde{V} = \begin{bmatrix} U^{*\top} Z V^* & U^{*\top} Z V^\perp \\ U^{\perp \top} Z V^* & U^{\perp \top} Z V^\perp \end{bmatrix} =: \begin{bmatrix} Z_A & Z_B \\ Z_C & Z_D \end{bmatrix}.$$

Here, the 2×2 block representation with $Z_A \in \mathbb{R}^{r \times r}$, $Z_B \in \mathbb{R}^{r \times (n-r)}$, $Z_C \in \mathbb{R}^{(n-r) \times r}$, $Z_D \in \mathbb{R}^{(n-r) \times (n-r)}$ is closely connected to $P_{T^* \perp}(Z)$ and $P_T(Z)$, which will provide nice interpretations

⁷Expand U^\perp to $n - r$ columns if it is not.

for our conditions (described momentarily). To begin, from the definition of $P_{T^*\perp}(Z)$, one can see that

$$\tilde{U}^\top P_{T^*\perp}(Z) \tilde{V} = \begin{bmatrix} U^{*\top} P_{T^*\perp}(Z) V^* & U^{*\top} P_{T^*\perp}(Z) V^\perp \\ U^{\perp\top} P_{T^*\perp}(Z) V^* & U^{\perp\top} P_{T^*\perp}(Z) V^\perp \end{bmatrix} =: \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & Z_D \end{bmatrix}.$$

Similarly, due to $P_{T^*}(Z) + P_{T^*\perp}(Z) = Z$, we have

$$\tilde{U}^\top P_{T^*}(Z) \tilde{V} =: \begin{bmatrix} Z_A & Z_B \\ Z_C & \mathbf{0} \end{bmatrix}.$$

In addition, this also provides a way to interpret $U^{*\top} Z$ and ZV^* by noting that

$$(U^{*\top} Z) \tilde{V} =: [Z_A \quad Z_B] \quad \tilde{U}^\top (ZV^*) =: \begin{bmatrix} Z_A \\ Z_C \end{bmatrix}$$

Then, we are ready to state the Assumption 3(a) in an equivalent way (See Lemma 9 for a generalized statement and a more algebraic proof).

Claim. *Assumption 3(a) is equivalent to the following condition: there exists a constant $C > 0$ such that*

$$\frac{C}{\log n} \|Z\|_F^2 \leq \|Z_D\|_F^2 - \|Z_A\|_F^2.$$

Proof. Note that $\|Z\|_F^2 = \|Z_A\|_F^2 + \|Z_B\|_F^2 + \|Z_C\|_F^2 + \|Z_D\|_F^2$, $\|ZV^*\|_F^2 = \|Z_A\|_F^2 + \|Z_C\|_F^2$, and $\|U^{*\top} Z\|_F^2 = \|Z_A\|_F^2 + \|Z_B\|_F^2$ due to the unitary invariance of the Frobenius norm. Then Condition 3(a) can be transformed into

$$\|Z\|_F^2 + \|Z_A\|_F^2 - \|Z_D\|_F^2 \leq \left(1 - \frac{C}{\log n}\right) \|Z\|_F^2$$

which completes the proof. \square

Since $\|Z_D\|_F^2 = \|P_{T^*\perp}(Z)\|_F^2$, the Assumption 3(a) is effectively a lower bound for requiring $\|P_{T^*\perp}(Z)\|_F^2$ being sufficiently large.

For Assumption 3(b), note that $\langle Z, U^* V^{*T} \rangle = \text{tr}(U^{*T} Z V^*) = \text{tr}(Z_A)$ and $\|P_{T^*\perp}(Z)\| = \|Z_D\|$ due to the unitary invariance of the operator norm. We have

Claim. *Assumption 3(b) is equivalent to the following condition: there exists a constant $C > 0$ such that*

$$|\text{tr}(Z_A)| \|Z_D\| \leq \left(1 - \frac{C}{\log n}\right) \|Z_D\|_F^2.$$

To see the looseness of these two conditions, note that Z_A is a $r \times r$ matrix and Z_D is a $(n-r) \times (n-r)$ matrix where $n \gg r$. Intuitively, some degree of non-overlapping between Z and M^* is enough to guarantee a small Z_A and large Z_D and sufficient for these two conditions. See Section 3.1 for a few examples to make this intuition precise.

D Discussions of Panel Data Regression

We provide a detailed discussion here for comparing assumptions and guarantees between [27] and our algorithm. To fit [27] into our model, we consider $k = 1$ and view Z as a regressor and δ as the idiosyncratic noise.

The key assumptions for Theorem 4 in [27] are (i) $\|\delta\| = O(\sqrt{n})$, (ii) $\langle P_{T^*\perp}(Z), \delta \rangle = O(n)$, (iii) $\|P_{T^*\perp}(Z)\|_F^2 = \Omega(n^2)$, (iv) $|\tau^0 - \tau^*| = o(1)$, and other low-rank and incoherence conditions. Then Theorem 4 in [27] provides an iterative estimator τ^t such that $|\tau^t - \tau^*| = O(1/n)$ for large enough t .

In contrast, our Eq. (6) provides $|\tau^d - \tau^*| = \tilde{O}(1/\|Z\|_F + |\langle P_{T^*}(\delta), P_{T^*}(Z) \rangle| / \|Z\|_F^2)$. Note that their assumption (ii) implies $\langle P_{T^*}(Z), P_{T^*}(\delta) \rangle = O(n)$ (since $\langle Z, \delta \rangle = 0$) and (iii) implies

$\|Z\|_F^2 \geq \|P_{T^*\perp}(Z)\|_F^2 = \Omega(n^2)$. Hence, our Eq. (6) can be simplified to $|\tau^d - \tau^*| = \tilde{O}(1/n)$, which recovers their results up to $\log(n)$ factors.

The assumption (iii) $\|P_{T^*\perp}(Z)\|_F^2 = \Omega(n^2)$ is probably the most restrictive one, which rules out many interesting scenarios (such as synthetic control). This assumption is also implicitly made in Theorem 1 and 2 in [27], where the bound $|\hat{\tau} - \tau^*| = O(1/\sqrt{n})$ is provided (recall $\hat{\tau}$ is the convex estimator). In particular, Theorem 1 [27] requires that $\text{rank}(Z) = O(1)$ and $\|P_{T^*\perp}(Z)\|_* - \|U^{*T}ZV^*\| = \Omega(n)$, hence implying $\sqrt{\text{rank}(Z)}\|P_{T^*\perp}(Z)\|_F \geq \|P_{T^*\perp}(Z)\|_* = \Omega(n)$, i.e., $\|P_{T^*\perp}(Z)\|_F = \Omega(n)$. Similarly, $\|P_{T^*\perp}(Z)\|_F = \Omega(n)$ is implied by their Assumption 1 and Theorem 2(iii) [27].

Technically, the assumption $\|P_{T^*\perp}(Z)\|_F = \Theta(n^2)$ greatly simplifies analysis in [27] since a global bound on $\hat{\tau} - \tau^*$ can be easily obtained. One of the main technical innovations in our paper, building on recent advances in the matrix completion literature, is to conduct a refined ‘local’ analysis without the assumption on the density of Z .

Other than technical challenges, another potential reason that panel data regression literature [25, 26, 27] did not loose the $\|P_{T^*\perp}(Z)\|_F = \Theta(n^2)$ assumption, is probably the regressor is usually viewed as a dense matrix (e.g., GDP, wages). In the contrast, the scenario we are considering is different, where Z characterizes the treatment pattern and $\|P_{T^*\perp}(Z)\|_F = \Theta(n^2)$ becomes a more problematic restriction.

Note that [27] also provides the results for k being a constant, where we think it is an interesting open question to study what are the minimal identification conditions for Z_m in that scenario.

E Additional Details of Experiments

In this section, we present more details for Section 4.

Computing Infrastructure. All experiments are done in a personal laptop equipped with 2.6 GHz 6-Core Intel Core i7 and 16 GB 2667 MHz DDR4. The operating system is macOS Catalina. For each instance, the running time for our algorithm is within seconds.

Algorithm Implementations. Recall that we implemented the following four benchmarks: (i) Matrix-Completion with Nuclear Norm Minimization (MC-NNM): [3] that applies matrix completion by viewing the treated entries as missing. (ii) Robust Synthetic Control (RSC): ‘Algorithm 1’ in [10] with linear regression used to recover the counterfactual results. This can be viewed as a robust variant of the well-known synthetic control method [1, 2]. (iii) Ordinary Least Square (OLS): Selects $a, b \in \mathbb{R}^n, \tau \in \mathbb{R}$ to minimize $\|O - a1^T - 1b^T - \tau Z\|_F^2$, where $1 \in \mathbb{R}^n$ is the vector of ones. This can be viewed as regression adapted to the difference-in-differences frameworks [45, 22]. (iv) Synthetic Difference-in-Difference (SDID): An algorithm proposed in [13] as a generalization of both synthetic control and difference-in-difference.

This section presents more details for algorithm implementations. All algorithms share the same input (O, Z) and r , where r is a pre-defined rank.

For implementing MC-NNM described in [3], let Ω be the set of observed control entries (Ω is the complement of Z). We optimize the following problem

$$\min_{M \in \mathbb{R}^{n \times n}, a \in \mathbb{R}^n, b \in \mathbb{R}^n} \|P_\Omega(O - a1^T - 1b^T - M)\|_F^2 + \lambda \|M\|_*$$

where a, b are for characterizing the fixed effects. Let $(\hat{M}, \hat{a}, \hat{b})$ be the optimizer for the above problem. The estimator of τ^* is then given by $\tau = \langle Z, O - \hat{M} - \hat{a}1^T - 1\hat{b}^T \rangle / \|Z\|_F^2$. To tune the hyper-parameter λ , we choose a large enough λ and then gradually decrease λ until the rank of \hat{M} achieves a pre-defined rank r .

For implementing RSC described in [10] for the block and stagger patterns, let S_1 be the set of treated units and S_2 be set of the control units. For each $i \in S_1$, we use the ‘Algorithm 1’ in [10] with the ‘linear regression ($\eta = 0$)’ to estimate the counterfactuals of the i -th row based on the control units S_2 . Let \hat{M} be their final estimate of the counterfactuals (combining all estimations across different rows). Let $\tau = \langle Z, O - \hat{M} \rangle / \|Z\|_F^2$ be the estimator of τ^* .

For OLS, we obtain the solution by a single linear regression. For SDID, we implement “Algorithm 1” in [13]. The SDID is designed for block patterns, to extend to stagger patterns, we adopt the suggestions in “Footnote 1” of [13].

For De-biased Convex (our algorithm), we implement an alternating optimization to solve the convex problem. Let $\tau = \tau^d$ be the estimator for τ^* . To tune the hyper-parameter λ , similar to the MC-NNM, we choose a large enough λ and then gradually decrease λ until the rank of \hat{M} achieves a pre-defined rank r .

The Choice of τ^* . For all algorithms we implemented, given (M^*, Z, E) , the error $\tau - \tau^*$ is in fact invariant from the change of τ^* . For MC-NNM, RSC, this is obvious since they do not use the information from treatment entries. In particular, one can check $\tau^* - \tau = \langle Z, M^* + E - \hat{M} \rangle / \|Z\|_F^2$ where \hat{M} is independent from the choice of τ^* in their algorithms.

For De-biased Convex, one can verify that if $(\hat{M}, \hat{\tau})$ satisfies the first-order conditions of the instance $O = M^* + Z * \tau^* + E$, then $(\hat{M}, \hat{\tau} + \Delta)$ satisfies the first order conditions of the instance $O = M^* + Z * (\tau^* + \Delta) + E$. This linear response together with the linear debias procedure implies that $\tau^* - \tau$ is independent from the choice of τ^* . Similarly, OLS and SDID also shares the same property. See Fig. 3 for the invariance of $\tau - \tau^*$ for a particular instance.




Figure 3: An example of showing that $|\tau - \tau^*|$ is invariant for different choice of τ^* for various algorithms.

Real Data. The data is collected from a Kaggle competition [42]. There are two types of promotion information: Promo Z_1 and Promo2 Z_2 , which are both binary matrices. Since Z_1 is store-independent (in a specific day, stores either all have promotions or all have no promotions), we discard Z_1 (viewing Z_1 being absorbed by latent features of days) and focus on estimating treatment effects of Z_2 . We also filter out stores that have no promotions for any days.

Asymptotic Normality. We consider a set of synthetic instances. For each instance, we used a typical procedure for generating low-rank non-negative matrices $M^* \in \mathbb{R}^{n_1 \times n_2}$ (e.g. [46, 47]). We select the rank r and mean-value \bar{M}^* . Given (r, \bar{M}^*) , we generate $U \in \mathbb{R}^{n_1 \times r}$ and $V \in \mathbb{R}^{n_2 \times r}$ with entries drawn i.i.d from Gamma(2, 1), and set $M^* = kUV^T$ with k chosen that the mean-value is indeed \bar{M}^* . For E and δ , its entries were drawn i.i.d from a Gaussian $\mathcal{N}(0, \sigma)$ and $\mathcal{N}(0, \sigma_\delta)$ respectively. The observation is given by $O = M^* + E + \tau^*Z + \delta \circ Z$.

Fig. 2 confirms the asymptotic normality of τ^d using an ensemble of 10,000 instances with $r = 10, \bar{M}^* = 10, \tau^* = \sigma = \sigma_\delta = 1$, and randomly generated stagger patterns Z where $m_1 \sim \text{Uni}[1, n_1], m_2 \sim \text{Uni}[n_2/2, n_2]$.⁸ For each instance, we computed $(\tau^d - \tau^*) / (\sigma / \|P_{\hat{T}^\perp}(Z)\|_F)$, which is predicted by Theorem 2 to be distributed according to $\mathcal{N}(0, 1)$. Fig. 2 (Left) shows the empirical distribution of δ_τ over instances with $n_1 = n_2 = 100$, overlaid with the true density function of $\mathcal{N}(0, 1)$. We also computed the *coverage rate*: the proportion of instances where τ^* is within the predicted 95% confidence interval of τ^d , calculated based on the estimated variance. Fig. 2 (Right) shows the desired coverage rates for different sizes of matrices. Both results suggest the effectiveness of our estimator for performing inference on τ^* .

⁸Experiments on other tested patterns (e.g., block patterns) also showed similar performances.

Row-specific Treatment Effects. We also test the performances of various algorithms for multiple treatment effects ($k > 1$). In particular, we reuse the setting for semi-synthetic (Tobacco) in Section 4, where (O, Z) is generated with the same distribution. Instead of estimating the overall average treatment effects, we now consider to estimate the unit-specific treatment effects $\vec{\tau}^*$, where $\vec{\tau}_i^* = \sum_j Z_{ij} \mathcal{T}_{ij} / \sum_j Z_{ij} = \tau^* + \delta_i$.

To estimate $\vec{\tau}^*$, we use De-biased Convex algorithm for multiple treatment effects specified in Eq. (1) (each Z_l corresponds to one row). We also extend RSC, MC-NNM, and OLS in this case, where the estimator $\vec{\tau}$ can be obtained directly based on their estimation of counterfactuals \hat{M} : $\vec{\tau}_i = \sum_j Z_{ij} (O_{ij} - \hat{M}_{ij}) / \sum_j Z_{ij}$.⁹ Table 2 reports the average $\|\vec{\tau} - \vec{\tau}^*\| / \|\vec{\tau}^*\|$ over 1000 instances. The results show the compelling performance of De-biased Convex algorithm compared to the state-of-the-arts.

Table 2: Comparison of our algorithm (De-biased Convex) to benchmarks on the semi-synthetic (Tobacco) dataset. Average normalized error $\|\vec{\tau} - \vec{\tau}^*\| / \|\vec{\tau}^*\|$ is reported.

Pattern	De-biased Convex	MC-NNM	RSC	OLS
Block	0.05 (± 0.06)	0.08 (± 0.13)	0.10 (± 0.12)	0.10 (± 0.16)
Stagger	0.03 (± 0.02)	0.05 (± 0.03)	0.06 (± 0.07)	0.07 (± 0.03)

F Proof of Lemma 3

Restating Lemma 3:

Lemma 3. Assume Assumption 3(a) and Assumption 3(b) hold, and $\frac{\sigma}{\sigma_{\min}} \sqrt{n} \leq C_1 \frac{1}{\kappa^2 r^2 \log^5(n)}$. Suppose $O = M^* + \tau^* Z + E$ for a deterministic E . Assume $\|E\| \leq C_2 \sigma \sqrt{n}$ with $|\langle E, P_{T^{\perp}}(Z) \rangle| \leq C_3 \sigma \sqrt{n / \log(n)} \|Z\|_F$.

Let $\lambda = C_\lambda \sigma \sqrt{n} \log^{1.5}(n)$, $X, Y \in \mathbb{R}^{n \times r}$, $\tau = \frac{\langle Z, O - XY^\top \rangle}{\|Z\|_F^2}$, and T is the tangent space of XY^\top .

$$\|X - X^*\|_F + \|Y - Y^*\|_F \leq C_F \left(\frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \right) \|X^*\|_F, \quad (15a)$$

$$\|\nabla f(X, Y; \tau)\|_F \leq \frac{\lambda \sqrt{\sigma_{\min}}}{\kappa n}. \quad (15b)$$

Let $R = \frac{1}{\lambda} (O - XY^\top - \tau Z - \lambda UV^\top)$ where $U \Sigma V^\top$ is the SVD of XY^\top . Then,

$$\|P_T(R)\|_F \leq \frac{72\kappa}{\lambda \sqrt{\sigma_{\min}}} \|\nabla f(X, Y; \tau)\|_F \quad \text{and} \quad \|P_{T^\perp}(R)\| \leq \left(1 - \frac{C}{\log n}\right). \quad (16)$$

Here, C is a constant depending (polynomially) on $C_1, C_2, C_3, C_\lambda, C_F$.

Let $XY^\top = U \Sigma V^\top$ be the singular value decomposition of XY^\top . Recall that T is the tangent space of XY^\top and $P_T(A)$ is the projection of A into the space T for any $A \in \mathbb{R}^{n \times n}$:

$$P_T(A) = A - (I_n - UU^\top)A(I_n - VV^\top).$$

By Lemma 10, for large enough n , the singular values of X, Y are in the intervals $[\sqrt{\frac{\sigma_{\min}}{2}}, \sqrt{2\sigma_{\max}}]$ and

$$\sigma_{\min}/2 \leq \sigma_{\min}(\Sigma) \leq \sigma_{\max}(\Sigma) \leq 2\sigma_{\max}.$$

We then consider $P_T(R)$ and $P_{T^\perp}(R)$ separately. The proof of $P_T(R)$ is similar to the proof of Claim 2 in [4], while the proof of $P_{T^\perp}(R)$ is based on a careful analysis of the property of τ .

⁹SDID is not easy to extend for multiple treatment effects here.

E1 Control of $P_T(R)$

By the definition of P_T , we have

$$\begin{aligned}
\|P_T(R)\|_F &= \|R - (I - UU^\top)R(I - VV^\top)\|_F \\
&= \|UU^\top R(I - VV^\top) + R(VV^\top)\|_F \\
&\stackrel{(i)}{\leq} \|U\| \|U^\top R\|_F \|I - VV^\top\| + \|RV\|_F \|V^\top\| \\
&\stackrel{(ii)}{\leq} \|U^\top R\|_F + \|RV\|_F
\end{aligned} \tag{18}$$

where (i) is due to $\|ABC\|_F \leq \|A\| \|B\|_F \|C\|$ and (ii) is due to $\|U\|, \|I - VV^\top\|, \|V\| \leq 1$.

Before showing bounds on $\|U^\top R\|_F$ and $\|RV\|_F$, consider an examination for the properties of X and Y . By Lemma 13, we can write $X = U\Sigma^{1/2}Q, Y = V\Sigma^{1/2}Q^{-\top}$ where $Q \in \mathbb{R}^{r \times r}$ is an invertible matrix and

$$\|\Sigma_Q - \Sigma_Q^{-1}\|_F \leq \frac{1}{\sigma_{\min}(\Sigma)} \|X^\top X - Y^\top Y\|_F, \tag{19}$$

where $U_Q \Sigma_Q V_Q^\top$ is the SVD of Q . In fact, we can show that $X^\top X \approx Y^\top Y$ due to that $\nabla f(X, Y; \tau) \approx 0$. To see this, set B_1, B_2 for the gradient $\nabla_X f(X, Y; \tau)$ and $\nabla_Y f(X, Y; \tau)$:

$$B_1 := (XY^\top + \tau Z - O)Y + \lambda X \tag{20}$$

$$B_2 := (XY^\top + \tau Z - O)^\top X + \lambda Y. \tag{21}$$

Left-multiplying X^\top on both sides of Eq. (20) and left-multiplying Y^\top on both sides of Eq. (21) and taking the transpose, we can obtain

$$\begin{aligned}
\lambda X^\top X &= X^\top B_1 - X^\top (\tau Z + XY^\top - O)Y \\
\lambda Y^\top Y &= B_2^\top Y - X^\top (\tau Z + XY^\top - O)Y.
\end{aligned}$$

This implies $\|X^\top X - Y^\top Y\|_F = \frac{1}{\lambda} \|X^\top B_1 - B_2^\top Y\|_F \leq \frac{1}{\lambda} \max(\|B_1\|_F, \|B_2\|_F)(\|X\| + \|Y\|)$. Combining this with Eq. (19), we obtain

$$\begin{aligned}
\|\Sigma_Q - \Sigma_Q^{-1}\|_F &\leq \frac{1}{\sigma_{\min}(\Sigma)} \|X^\top X - Y^\top Y\|_F \\
&\leq \frac{1}{\sigma_{\min}(\Sigma)\lambda} \max(\|B_1\|_F, \|B_2\|_F)(\|X\| + \|Y\|) \\
&\stackrel{(i)}{\leq} \frac{8\sqrt{\sigma_{\max}}}{\sigma_{\min}\lambda} \max(\|B_1\|_F, \|B_2\|_F) \\
&\leq \frac{8\sqrt{\sigma_{\max}}}{\sigma_{\min}\lambda} \|\nabla f(X, Y; \tau)\|_F \stackrel{(ii)}{\leq} 1
\end{aligned} \tag{22}$$

where (i) is due to $\sigma_{\min}(\Sigma) \geq \sigma_{\min}/2, \sigma_1(X) \leq 2\sigma_{\max}, \sigma_1(Y) \leq 2\sigma_{\max}$ and (ii) holds for large enough n . This also implies $\|\Sigma_Q\| \leq 2$. Intuitively speaking, $\nabla f(X, Y; \tau) \approx 0$ implies that $\Sigma_Q \approx I$ and hence Q is similar to an orthogonal matrix.

Next, we show how to control $\|RV\|_F$. Recall that

$$O - XY^\top - \tau Z = \lambda UV^\top + \lambda R. \tag{23}$$

Combining Eq. (20) and Eq. (23), we have

$$(-\lambda UV^\top - \lambda R)Y + \lambda X = B_1.$$

Therefore, $\lambda RY = \lambda X - B_1 - \lambda UV^\top Y$. Substituting X and Y with $X = U\Sigma^{1/2}Q, Y = V\Sigma^{1/2}Q^{-\top}$, we obtain

$$\lambda R(V\Sigma^{1/2}Q^{-\top}) = \lambda(U\Sigma^{1/2}Q) - B_1 - \lambda UV^\top(V\Sigma^{1/2}Q^{-\top}).$$

Right-multiplying $Q^\top \Sigma^{-1/2}$ on both sides and using $V^\top V = I$ leads to

$$\lambda R V = \lambda U \Sigma^{1/2} Q Q^\top \Sigma^{-1/2} - B_1 Q^\top \Sigma^{-1/2} - \lambda U.$$

Then this implies

$$\begin{aligned} \|R V\|_F &= \left\| U \Sigma^{1/2} (Q Q^\top - I) \Sigma^{-1/2} - \frac{1}{\lambda} B_1 Q^\top \Sigma^{-1/2} \right\|_F \\ &\leq \frac{1}{\lambda} \|B_1\|_F \|Q\| \|\Sigma^{-1/2}\| + \|U\| \|\Sigma^{1/2}\| \|\Sigma^{-1/2}\| \|I - Q Q^\top\|_F \\ &\stackrel{(i)}{\leq} \frac{1}{\lambda} \|B_1\|_F \|Q\| \frac{\sqrt{2}}{\sqrt{\sigma_{\min}}} + \sqrt{2\sigma_{\max}} \frac{\sqrt{2}}{\sqrt{\sigma_{\min}}} \|I - Q Q^\top\|_F \\ &\stackrel{(ii)}{\leq} \frac{2\sqrt{2}}{\lambda\sqrt{\sigma_{\min}}} \|B_1\|_F + 2\sqrt{\kappa} \|U_Q \Sigma_Q (\Sigma_Q^{-1} - \Sigma_Q) U_Q^\top\|_F \\ &\stackrel{(iii)}{\leq} \frac{2\sqrt{2}}{\lambda\sqrt{\sigma_{\min}}} \|\nabla f(X, Y; \tau)\|_F + 4\sqrt{\kappa} \|\Sigma_Q^{-1} - \Sigma_Q\|_F \\ &\stackrel{(iv)}{\leq} \left(\frac{2\sqrt{2}}{\lambda\sqrt{\sigma_{\min}}} + \frac{32\kappa}{\lambda\sqrt{\sigma_{\min}}} \right) \|\nabla f(X, Y; \tau)\|_F \\ &\leq \frac{36\kappa}{\lambda\sqrt{\sigma_{\min}}} \|\nabla f(X, Y; \tau)\|_F. \end{aligned}$$

where (i) is due to $\sigma_{\min}(\Sigma) \geq (1/2)\sigma_{\min}$, $\sigma_{\max}(\Sigma) \leq 2\sigma_{\max}$, (ii) is due to $\|Q\| = \|\Sigma_Q\| \leq 2$ and $Q = U_Q \Sigma_Q V_Q^\top$ is the SVD of Q , (iii) is due to $\|B_1\|_F \leq \|\nabla f(X, Y; \tau)\|_F$ and $\|U_Q\| = 1$, $\|\Sigma_Q\| \leq 2$, (iv) is due to the bound of $\|\Sigma_Q^{-1} - \Sigma_Q\|_F$ (Eq. (22)).

Similarly, one can obtain $\|U^\top R\|_F \leq \frac{36\kappa}{\lambda\sqrt{\sigma_{\min}}} \|\nabla f(X, Y; \tau)\|_F$. Combining these with Eq. (18) leads to the result:

$$\|P_T(R)\|_F \leq \frac{72\kappa}{\lambda\sqrt{\sigma_{\min}}} \|\nabla f(X, Y; \tau)\|_F. \quad (24)$$

This completes the proof for the bound of $P_T(R)$.

F.2 Control of $P_{T^\perp}(R)$

Next, we control $P_{T^\perp}(R)$. Using $O = M^* + E + \tau^* Z$ and applying P_{T^\perp} on both sides of $O - XY^\top - \tau Z = \lambda UV^\top + \lambda R$, we obtain

$$\begin{aligned} \lambda P_{T^\perp}(R) &\stackrel{(i)}{=} P_{T^\perp}(E) - P_{T^\perp}(XY^\top - M^*) - (\tau - \tau^*) P_{T^\perp}(Z) \\ &= P_{T^\perp}(E) - P_{T^\perp}((X - X^*)Y^\top + X(Y - Y^*)^\top) + P_{T^\perp}((X - X^*)(Y - Y^*)^\top) \\ &\quad - (\tau - \tau^*) P_{T^\perp}(Z) \\ &\stackrel{(ii)}{=} P_{T^\perp}(E) + P_{T^\perp}((X - X^*)(Y - Y^*)^\top) - (\tau - \tau^*) P_{T^\perp}(Z). \end{aligned} \quad (25)$$

where (i) is due to $P_{T^\perp}(UV^\top) = 0$, (ii) is due to $Y = V\Sigma^{1/2}Q$ and $X = U\Sigma^{1/2}Q^{-\top}$ by Lemma 13 and $P_{T^\perp}(UA^\top + BV^\top) = 0$ for any $A, B \in \mathbb{R}^{n \times r}$.

This implies that

$$\begin{aligned} \|P_{T^\perp}(R)\| &\leq \frac{1}{\lambda} \|P_{T^\perp}(E)\| + \frac{1}{\lambda} \|P_{T^\perp}((X - X^*)(Y - Y^*)^\top)\| + \frac{1}{\lambda} |\tau - \tau^*| \|P_{T^\perp}(Z)\| \\ &\stackrel{(i)}{\leq} \frac{1}{\lambda} \|E\| + \frac{1}{\lambda} \|X - X^*\| \|Y - Y^*\| + \frac{1}{\lambda} |\tau - \tau^*| \|P_{T^\perp}(Z)\| \end{aligned}$$

where (i) is due to $\|P_{T^\perp}(A)\| \leq \|I - UU^\top\| \|A\| \|I - VV^\top\| \leq \|A\|$ for any $A \in \mathbb{R}^{n \times n}$. Then it boils down to control $\frac{\|E\|}{\lambda}$, $\frac{\|X - X^*\| \|Y - Y^*\|}{\lambda}$, and $\frac{|\tau - \tau^*| \|P_{T^\perp}(Z)\|}{\lambda}$ separately.

Consider $\frac{\|E\|}{\lambda}$. Since we have the condition $\|E\| \lesssim \sigma\sqrt{n}$, then

$$\frac{\|E\|}{\lambda} \leq \frac{c\sigma\sqrt{n}}{C_\lambda\sigma\sqrt{n}\log^{1.5}(n)} \lesssim \frac{1}{\log^{1.5}(n)}. \quad (26)$$

Consider $\frac{\|X-X^*\| \|Y-Y^*\|}{\lambda}$. We have

$$\begin{aligned} \frac{\|X-X^*\| \|Y-Y^*\|}{\lambda} &\leq \frac{\|X-X^*\|_F \|Y-Y^*\|_F}{\lambda} \\ &\stackrel{(i)}{\leq} C_F^2 \frac{\sigma^2 n \log^5(n)}{\sigma_{\min}^2} \sigma_{\max}^r \frac{1}{C_\lambda \sigma \sqrt{n} \log^{1.5}(n)} \\ &\leq \frac{C_F^2 \sigma \sqrt{n} \log^{3.5}(n) \kappa r}{C_\lambda \sigma_{\min}} \\ &\stackrel{(ii)}{\lesssim} \frac{1}{\log^{1.5}(n)} \end{aligned} \quad (27)$$

where (i) is due to

$$\|X-X^*\|_F + \|Y-Y^*\|_F \leq C_F \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \sqrt{\sigma_{\max}^r}$$

with $\lambda = C_\lambda \sigma \sqrt{n} \log^{1.5}(n)$, and (ii) is due to $\frac{\sigma \sqrt{n}}{\sigma_{\min}} \lesssim \frac{1}{r^2 \kappa^2 \log^5(n)}$.

Consider $\frac{1}{\lambda} |\tau - \tau^*| \|P_{T^\perp}(Z)\|$. We have the following claim.

Claim 1. Suppose Assumption 3(a) and Assumption 3(b) hold. Suppose $\frac{\sigma \sqrt{n}}{\sigma_{\min}} \lesssim \frac{1}{r^2 \kappa^2 \log^5(n)}$. Then

$$\frac{1}{\lambda} |\tau - \tau^*| \|P_{T^\perp}(Z)\| \leq 1 - \frac{C}{\log(n)} \quad (28)$$

for some constant C .

Combining Eqs. (26) to (28), we complete the proof for controlling $P_{T^\perp}(R)$.

Proof of Claim 1. Similar to the proof of Lemma 1, we first consider the characterization for $\tau - \tau^*$. Since $\tau = \langle Z, O - XY^\top \rangle / \|Z\|_F^2$, we have $\langle O - XY^\top - \tau Z, Z \rangle = 0$. Combining this with $O - XY^\top - \tau Z = \lambda UV^\top + \lambda R$, we have

$$\begin{aligned} \langle Z, UV^\top + R \rangle &= 0 \\ \implies \langle Z, UV^\top \rangle &= -\langle Z, R \rangle \\ \implies \langle Z, UV^\top \rangle + \langle P_T(Z), P_T(R) \rangle &= -\langle P_{T^\perp}(Z), P_{T^\perp}(R) \rangle \end{aligned} \quad (29)$$

Substituting $P_{T^\perp}(R)$ in Eq. (29) by Eq. (25), we obtain

$$\begin{aligned} &\langle Z, UV^\top \rangle + \langle P_T(Z), P_T(R) \rangle \\ &= \frac{1}{\lambda} (\tau - \tau^*) \langle P_{T^\perp}(Z), P_{T^\perp}(Z) \rangle \\ &\quad - \frac{1}{\lambda} \langle P_{T^\perp}(Z), P_{T^\perp}(E) \rangle - \frac{1}{\lambda} \langle P_{T^\perp}(Z), P_{T^\perp}((X-X^*)(Y-Y^*)^\top) \rangle. \end{aligned}$$

Note that $\langle P_{T^\perp}(Z), P_{T^\perp}(Z) \rangle = \|P_{T^\perp}(Z)\|_F^2$. This implies

$$\begin{aligned} \frac{(\tau - \tau^*)}{\lambda} \|P_{T^\perp}(Z)\|_F^2 &= \langle Z, UV^\top \rangle + \langle P_T(Z), P_T(R) \rangle + \frac{1}{\lambda} \langle P_{T^\perp}(Z), P_{T^\perp}(E) \rangle \\ &\quad + \frac{1}{\lambda} \langle P_{T^\perp}(Z), P_{T^\perp}((X-X^*)(Y-Y^*)^\top) \rangle. \end{aligned}$$

This further implies

$$\begin{aligned}
& \frac{|\tau - \tau^*|}{\lambda} \|P_{T^\perp}(Z)\| \\
& \leq |\langle Z, UV^\top \rangle| \frac{\|P_{T^\perp}(Z)\|}{\|P_{T^\perp}(Z)\|_F^2} + |\langle P_T(Z), P_T(R) \rangle| \frac{\|P_{T^\perp}(Z)\|}{\|P_{T^\perp}(Z)\|_F^2} \\
& \quad + \frac{1}{\lambda} |\langle P_{T^\perp}(Z), P_{T^\perp}(E) \rangle| \frac{\|P_{T^\perp}(Z)\|}{\|P_{T^\perp}(Z)\|_F^2} \\
& \quad + \frac{1}{\lambda} |\langle P_{T^\perp}(Z), P_{T^\perp}((X - X^*)(Y - Y^*)^\top) \rangle| \frac{\|P_{T^\perp}(Z)\|}{\|P_{T^\perp}(Z)\|_F^2} \\
& \stackrel{(i)}{\leq} \underbrace{|\langle Z, UV^\top \rangle| \frac{\|P_{T^\perp}(Z)\|}{\|P_{T^\perp}(Z)\|_F^2}}_{A_0} + \underbrace{\frac{\|P_T(Z)\|_F}{\|P_{T^\perp}(Z)\|_F} \|P_T(R)\|_F}_{A_1} + \underbrace{\frac{|\langle P_{T^\perp}(Z), E \rangle|}{\lambda \|P_{T^\perp}(Z)\|_F}}_{A_2} \\
& \quad + \frac{1}{\lambda} \underbrace{\|P_{T^\perp}((X - X^*)(Y - Y^*)^\top)\|_F}_{A_3}
\end{aligned}$$

where (i) is due to $\|P_{T^\perp}(Z)\| \leq \|P_{T^\perp}(Z)\|_F$ and $|\langle A, B \rangle| \leq \|A\|_F \|B\|_F$ by Cauchy-Schwartz inequality. Then it boils down to control A_0, A_1, A_2, A_3 .

By Lemma 16, we have that

$$A_0 \leq 1 - \frac{C_{r_2}}{2 \log(n)}$$

$$\|P_{T^\perp}(Z)\|_F^2 \geq \frac{C_{r_1}}{2 \log(n)} \|Z\|_F^2 \quad (30)$$

$$\|P_{T^\perp}(Z) - P_{T^{*\perp}}(Z)\|_* \lesssim \frac{1}{r^{0.5} \log^{2.5}(n)} \|Z\|_F. \quad (31)$$

For A_1 , this implies

$$\begin{aligned}
A_1 &= \frac{\|P_T(Z)\|_F}{\|P_{T^\perp}(Z)\|_F} \|P_T(R)\|_F \\
&\leq \frac{\|P_T(Z)\|_F}{\|Z\|_F} \frac{2 \log(n)}{C_{r_1}} \|P_T(R)\|_F \\
&\leq \frac{2 \log(n)}{C_{r_1}} \|P_T(R)\|_F \\
&\stackrel{(i)}{\lesssim} \frac{1}{\log^2(n)}
\end{aligned}$$

where (i) is due to $\|P_T(R)\|_F \lesssim \frac{\kappa}{\lambda \sqrt{\sigma_{\min}}} \|\nabla f(X, Y; \tau)\|$.

For A_2 , with probability $1 - O(1/\text{poly}(n))$, we have

$$\begin{aligned}
|\langle P_{T^\perp}(Z), E \rangle| &\leq |\langle P_{T^\perp}(Z) - P_{T^{*\perp}}(Z), E \rangle| + |\langle P_{T^{*\perp}}(Z), E \rangle| \\
&\leq \|P_{T^\perp}(Z) - P_{T^{*\perp}}(Z)\|_* \|E\| + |\langle P_{T^{*\perp}}(Z), E \rangle| \\
&\stackrel{(i)}{\lesssim} \frac{\|Z\|_F}{r^{0.5} \log^{2.5}(n)} \sigma \sqrt{n} + |\langle P_{T^{*\perp}}(Z), E \rangle| \\
&\stackrel{(ii)}{\lesssim} \frac{\|Z\|_F}{r^{0.5} \log^{2.5}(n)} \sigma \sqrt{n} + \sigma \sqrt{n} \|Z\|_F.
\end{aligned}$$

Here, (i) is due to Eq. (31) and $\|E\| \lesssim \sigma\sqrt{n}$, and (ii) is due to $|\langle P_{T^{\perp}}(Z), E \rangle| \lesssim \sigma\sqrt{n/\log(n)} \|Z\|_{\text{F}}$. This implies

$$\begin{aligned} A_2 &= \frac{|\langle P_{T^{\perp}}(Z), E \rangle|}{\lambda \|P_{T^{\perp}}(Z)\|_{\text{F}}} \\ &\lesssim \frac{\sigma\sqrt{n} \|Z\|_{\text{F}}}{\sigma\sqrt{n} \log^2(n) \|P_{T^{\perp}}(Z)\|_{\text{F}}} \\ &\stackrel{(i)}{\lesssim} \frac{\|Z\|_{\text{F}}}{\log^{1.5}(n) \|Z\|_{\text{F}}} \\ &\lesssim \frac{1}{\log^{1.5}(n)}. \end{aligned}$$

Here, (i) is due to Eq. (30).

For A_3 , we have

$$\begin{aligned} A_3 &= \frac{1}{\lambda} \|P_{T^{\perp}}((X - X^*)(Y - Y^*)^{\top})\|_{\text{F}} \\ &\leq \frac{1}{\lambda} \|X - X^*\|_{\text{F}} \|Y - Y^*\|_{\text{F}} \stackrel{(i)}{\lesssim} \frac{1}{\log^{1.5}(n)}. \end{aligned}$$

Here, (i) is due to Eq. (27)

Combining all the results for A_0, A_1, A_2, A_3 , we have

$$\begin{aligned} \frac{|\tau - \tau^*|}{\lambda} \|P_{T^{\perp}}(Z)\| &\leq A_0 + A_1 + A_2 + A_3 \\ &\leq 1 - \frac{C_{r_2}}{2 \log(n)} + \frac{c_1}{\log^2(n)} + \frac{c_2}{\log^{1.5}(n)} + \frac{c_3}{\log^{1.5}(n)} \\ &\leq 1 - \frac{C}{\log(n)} \end{aligned}$$

for some constants c_1, c_2, c_3, C . This completes the proof of the claim. \square

G Proof of Lemma 4

Restating Lemma 4:

Lemma 4. *Assume the same setup from Lemma 3. For any minimizer $(\hat{M}, \hat{\tau})$ of the convex program Eq. (1a), we have*

$$\begin{aligned} \|XY^{\top} - \hat{M}\|_{\text{F}} &\leq C \frac{\kappa \log(n)}{\sqrt{\sigma_{\min}}} \|\nabla f(X, Y; \tau)\|_{\text{F}} \\ |\tau - \hat{\tau}| &\leq C \frac{\kappa \log(n)}{\sqrt{\sigma_{\min}} \|Z\|_{\text{F}}} \|\nabla f(X, Y; \tau)\|_{\text{F}}. \end{aligned}$$

Furthermore, let \hat{T} be the tangent space of \hat{M} . For any matrix $A \in \mathbb{R}^{n \times n}$, we have

$$\begin{aligned} \|P_{\hat{T}^{\perp}}(A) - P_{T^{\perp}}(A)\|_{\text{F}} &\leq C \|A\| \frac{\kappa \log(n)}{\sigma_{\min}^{1.5}} \|\nabla f(X, Y; \tau)\|_{\text{F}} \\ \|P_{\hat{T}^{\perp}}(XY^{\top})\|_{\text{F}} &\leq C \frac{\kappa^3 \log^2(n)}{\sigma_{\min}^2} \|\nabla f\|_{\text{F}}^2. \end{aligned}$$

Here, C is a constant depending (polynomially) on $C_1, C_2, C_3, C_{\lambda}, C_{\text{F}}$.

Proof of Lemma 4. Based on Lemma 3, recall that for $R = \frac{1}{\lambda}(O - XY^{\top} - \tau Z - \lambda UV^{\top})$, we have $\|P_T(R)\|_{\text{F}} \lesssim \frac{\kappa}{\lambda\sqrt{\sigma_{\min}}} \|\nabla f\|_{\text{F}}$ and $\|P_{T^{\perp}}(R)\| \leq 1 - C/\log(n)$.

Let $(M_{\text{cvx}}, \tau_{\text{cvx}})$ be one of the minimizers of g . Then we have

$$\frac{1}{2} \|O - M_{\text{cvx}} - \tau(M_{\text{cvx}})Z\|_{\text{F}}^2 + \lambda \|M_{\text{cvx}}\|_* \leq \frac{1}{2} \|O - XY^\top - \tau(XY^\top)Z\|_{\text{F}}^2 + \lambda \|XY^\top\|_* \quad (32)$$

where $\tau(M) := \frac{\langle Z, O-M \rangle}{\|Z\|_{\text{F}}^2}$.

Let $\Delta = M_{\text{cvx}} - XY^\top$. Let $\Delta' = \tau(M_{\text{cvx}})Z - \tau(XY^\top)Z$. Then Eq. (32) implies

$$\begin{aligned} & \frac{1}{2} \|O - XY^\top - \tau(XY^\top)Z - \Delta - \Delta'\|_{\text{F}}^2 + \lambda \|M_{\text{cvx}}\|_* \\ & \leq \frac{1}{2} \|O - XY^\top - \tau(XY^\top)Z\|_{\text{F}}^2 + \lambda \|XY^\top\|_* . \end{aligned}$$

Hence,

$$\frac{1}{2} \|\Delta + \Delta'\|_{\text{F}}^2 \leq \langle O - XY^\top - \tau(XY^\top)Z, \Delta + \Delta' \rangle - \lambda \|M_{\text{cvx}}\|_* + \lambda \|XY^\top\|_* .$$

Note that $\tau(XY^\top) = \langle Z, O - XY^\top \rangle / \langle Z, Z \rangle$, then $\langle O - XY^\top - \tau(XY^\top)Z, Z \rangle = 0$. Hence

$$\langle O - XY^\top - \tau(XY^\top)Z, \Delta' \rangle = 0 .$$

This implies that

$$0 \leq \frac{1}{2} \|\Delta + \Delta'\|_{\text{F}}^2 \leq \langle O - XY^\top - \tau(XY^\top)Z, \Delta \rangle - \lambda \|M_{\text{cvx}}\|_* + \lambda \|XY^\top\|_* .$$

Note that $\|\cdot\|_*$ is convex and $UV^\top + W$ is the subgradient of $\|XY^\top\|_*$ given that $W \in T^\perp$ and $\|W\| \leq 1$. This implies that $\|M_{\text{cvx}}\|_* \geq \|XY^\top\|_* + \langle UV^\top + W, \Delta \rangle$ due to the property of the convex function. Choose W such that $\langle W, \Delta \rangle = \|P_{T^\perp}(\Delta)\|_*$, we have

$$\begin{aligned} \frac{1}{2} \|\Delta + \Delta'\|_{\text{F}}^2 & \leq \langle O - XY^\top - \tau(XY^\top)Z, \Delta \rangle - \lambda \langle UV^\top, \Delta \rangle - \lambda \|P_{T^\perp}(\Delta)\|_* \\ & = \lambda \langle R, \Delta \rangle - \lambda \|P_{T^\perp}(\Delta)\|_* \\ & = \lambda \langle P_T(R), P_T(\Delta) \rangle + \lambda \langle P_{T^\perp}(R), P_{T^\perp}(\Delta) \rangle - \lambda \|P_{T^\perp}(\Delta)\|_* \\ & \leq \lambda \|P_T(R)\|_{\text{F}} \|P_T(\Delta)\|_{\text{F}} + \lambda \|P_{T^\perp}(R)\| \|P_{T^\perp}(\Delta)\|_* - \lambda \|P_{T^\perp}(\Delta)\|_* \\ & \stackrel{(i)}{\leq} \lambda \|P_T(R)\|_{\text{F}} \|P_T(\Delta)\|_{\text{F}} - \frac{C\lambda}{\log(n)} \|P_{T^\perp}(\Delta)\|_* . \end{aligned} \quad (33)$$

Here, (i) is due to $\|P_{T^\perp}(R)\| \leq 1 - C/\log(n)$. This implies that

$$\|P_{T^\perp}(\Delta)\|_* \leq \|P_T(R)\|_{\text{F}} \|P_T(\Delta)\|_{\text{F}} \frac{\log(n)}{C} \lesssim \frac{\kappa \log(n)}{\lambda \sqrt{\sigma_{\min}}} \|\nabla f\|_{\text{F}} \|P_T(\Delta)\|_{\text{F}} . \quad (34)$$

Note that $\Delta' = -\langle \Delta, Z \rangle / \langle Z, Z \rangle \cdot Z$. Investigate further on $\|\Delta + \Delta'\|_{\text{F}}^2$:

$$\begin{aligned} \|\Delta + \Delta'\|_{\text{F}}^2 & = \langle \Delta, \Delta \rangle - 2 \langle \Delta, Z \rangle^2 / \langle Z, Z \rangle + \langle \Delta, Z \rangle^2 / \langle Z, Z \rangle \\ & = \langle \Delta, \Delta \rangle - \langle \Delta, Z \rangle^2 / \langle Z, Z \rangle \\ & = \|\Delta\|_{\text{F}}^2 - (\langle P_T(\Delta), P_T(Z) \rangle + \langle P_{T^\perp}(\Delta), P_{T^\perp}(Z) \rangle)^2 / \|Z\|_{\text{F}}^2 \\ & \geq \|\Delta\|_{\text{F}}^2 - (\|P_T(\Delta)\|_{\text{F}} \|P_T(Z)\|_{\text{F}} + \|P_{T^\perp}(\Delta)\|_* \|P_{T^\perp}(Z)\|)^2 / \|Z\|_{\text{F}}^2 \\ & \stackrel{(i)}{\geq} \|\Delta\|_{\text{F}}^2 - \frac{\|P_T(\Delta)\|_{\text{F}}^2}{\|Z\|_{\text{F}}^2} \cdot \left(\|P_T(Z)\|_{\text{F}} + \frac{C' \kappa \log(n)}{\lambda \sqrt{\sigma_{\min}}} \|\nabla f\|_{\text{F}} \|P_{T^\perp}(Z)\| \right)^2 \\ & \stackrel{(ii)}{\geq} \|\Delta\|_{\text{F}}^2 \cdot \left(1 - \frac{1}{\|Z\|_{\text{F}}^2} \cdot \left(\|P_T(Z)\|_{\text{F}} + \frac{C' \kappa \log(n)}{\lambda \sqrt{\sigma_{\min}}} \|\nabla f\|_{\text{F}} \|P_{T^\perp}(Z)\|_{\text{F}} \right)^2 \right) \end{aligned}$$

for some constant C' . Here, (i) is due to Eq. (34) and (ii) is due to $\|P_T(\Delta)\|_{\text{F}} \leq \|\Delta\|_{\text{F}}$.

Note that $\|P_T(Z)\|_F^2 / \|Z\|_F^2 \leq 1 - \frac{C''}{\log(n)}$ for some constant C'' (by Lemma 16), and $\|\nabla f\|_F$ is small enough. This implies

$$\|\Delta + \Delta'\|_F^2 \gtrsim \|\Delta\|_F^2 \frac{1}{\log(n)}.$$

Combining this with Eq. (33), we have

$$\frac{1}{\log(n)} \|\Delta\|_F^2 \lesssim \lambda \|P_T(R)\|_F \|P_T(\Delta)\|_F \stackrel{(i)}{\lesssim} \frac{\kappa}{\sqrt{\sigma_{\min}}} \|\nabla f\|_F \|\Delta\|_F.$$

Here, (i) is due to the bound for $\|P_T(R)\|_F$ and $\|P_T(\Delta)\|_F \leq \|\Delta\|_F$. This implies

$$\|\Delta\|_F \lesssim \frac{\kappa \log(n)}{\sqrt{\sigma_{\min}}} \|\nabla f\|_F.$$

Furthermore,

$$\begin{aligned} |\tau(XY^\top) - \tau(M_{\text{cvx}})| &= |\langle Z, M_{\text{cvx}} - XY^\top \rangle| / \|Z\|_F^2 \\ &\leq \|Z\|_F \|\Delta\|_F / \|Z\|_F^2 \\ &\leq \|\Delta\|_F / \|Z\|_F \\ &\leq \frac{\kappa \log(n)}{\sqrt{\sigma_{\min}} \|Z\|_F} \|\nabla f\|_F. \end{aligned}$$

Let $M_{\text{cvx}} = U_{\text{cvx}} \Sigma_{\text{cvx}} V_{\text{cvx}}^\top$ be SVD of M_{cvx} . Let T_{cvx} be the tangent space of M_{cvx} . Next, we analyze the relation between T_{cvx} and T . We aim to show (i) $\text{rank}(M_{\text{cvx}}) = r$ based on the fact that M_{cvx} is obtained by soft-thresholding; (ii) establish the closeness between eigen-subspaces U_{cvx} (V_{cvx}) and U (V) by invoking the Davis-Kahan theorem; (iii) establish the closeness between T_{cvx} and T based on the closeness between eigen-subspaces.

We first aim to show that $\text{rank}(M_{\text{cvx}}) = r$. Note that

$$O - M_{\text{cvx}} - \tau_{\text{cvx}} Z = \lambda U_{\text{cvx}} V_{\text{cvx}}^\top + W_{\text{cvx}}$$

where $\|W_{\text{cvx}}\| \leq 1$, $P_{T_{\text{cvx}}}(W_{\text{cvx}}) = 0$. Let $\sigma_i(A)$ be the i -th largest singular values of A . Note that there is $O - \tau Z = U(\lambda I_r + \Sigma)V^\top + \lambda R$. By Weyl's inequality, for $1 \leq i \leq r$

$$\begin{aligned} \sigma_i(O - \tau Z) &\geq \sigma_i(U(\lambda I_r + \Sigma)V^\top) - \lambda \|R\| \\ &\geq \frac{\sigma_{\min}}{2} + \lambda - \lambda(\|P_{T^\perp}(R)\| + \|P_T(R)\|) \\ &\stackrel{(i)}{\geq} \frac{\sigma_{\min}}{2} \\ &\stackrel{(ii)}{>} 2\lambda. \end{aligned}$$

Here, (i) is due to the bound of $\|P_{T^\perp}(R)\|$ and $\|P_T(R)\| \leq \|P_T(R)\|_F$, (ii) is due to $\frac{\sigma_{\min}}{\sigma_{\min}} \lesssim \frac{1}{r^2 \kappa^2 \log^5(n)}$. Then, for $1 \leq i \leq r$,

$$\begin{aligned} \sigma_i(O - \tau_{\text{cvx}} Z) &\geq \sigma_i(O - \tau Z) - |\tau_{\text{cvx}} - \tau| \|Z\| \\ &\geq \sigma_i(O - \tau Z) - \frac{C\kappa \log(n)}{\sqrt{\sigma_{\min}} \|Z\|_F} \|\nabla f\|_F \|Z\| \\ &\geq \sigma_i(O - \tau Z) - \frac{C\kappa \log(n)}{\sqrt{\sigma_{\min}}} \|\nabla f\|_F \\ &\geq 2\lambda - \frac{\lambda \log(n)}{n^3} \\ &> \lambda. \end{aligned}$$

providing that $\|\nabla f\| \leq \frac{\lambda \sqrt{\sigma_{\min}}}{\kappa n}$ and $\log(n) \lesssim n$.

Meanwhile, for $r + 1 \leq i \leq n$, we have

$$\begin{aligned}\sigma_i(O - \tau Z) &\leq \sigma_i(U(\lambda I_r + \Sigma)V^\top) + \lambda \|R\| \\ &\leq \lambda \|R\| \\ &\leq (1 - C/\log(n))\lambda.\end{aligned}$$

Then for $r + 1 \leq i \leq n$, we have

$$\begin{aligned}\sigma_i(O - \tau_{\text{cvx}}Z) &\leq \sigma_i(O - \tau Z) + |\tau_{\text{cvx}} - \tau| \|Z\| \\ &\leq (1 - C/\log(n))\lambda + \frac{\lambda \kappa \log(n)}{n^2} \\ &< \lambda.\end{aligned}$$

Then, by soft-thresholding (M_{cvx} is obtained by truncating the singular values of $(O - \tau_{\text{cvx}}Z)$ by λ , see [44]), we have that $\text{rank}(M_{\text{cvx}}) = r$. Then, by Davis-Kahan's theorem [48], there exist rotation matrices $R_1, R_2 \in \mathbb{R}^{r \times r}$, where

$$\begin{aligned}\max(\|UR_1 - U_{\text{cvx}}\|_{\text{F}}, \|VR_2 - V_{\text{cvx}}\|_{\text{F}}) &\lesssim \frac{\|M_{\text{cvx}} - M\|_{\text{F}}}{\sigma_r(M)} \\ &\lesssim \frac{\kappa \log(n)}{\sigma_{\min}^{1.5}} \|\nabla f\|_{\text{F}}.\end{aligned}$$

This simply implies that $\|UU^\top - U_{\text{cvx}}U_{\text{cvx}}^\top\|_{\text{F}} \lesssim \|UR_1 - U_{\text{cvx}}\|_{\text{F}} \|U\| \lesssim \frac{\kappa \log(n)}{\sigma_{\min}^{1.5}} \|\nabla f\|_{\text{F}}$. Similarly, $\|VV^\top - V_{\text{cvx}}V_{\text{cvx}}^\top\|_{\text{F}} \lesssim \frac{\kappa \log(n)}{\sigma_{\min}^{1.5}} \|\nabla f\|_{\text{F}}$.

Then, for any matrix $A \in \mathbb{R}^{n \times n}$,

$$\begin{aligned}&\|P_{T^\perp}(A) - P_{T_{\text{cvx}}^\perp}(A)\|_{\text{F}} \\ &= \|(I - UU^\top)A(I - VV^\top) - (I - U_{\text{cvx}}U_{\text{cvx}}^\top)A(I - V_{\text{cvx}}V_{\text{cvx}}^\top)\|_{\text{F}} \\ &\leq \|(UU^\top - U_{\text{cvx}}U_{\text{cvx}}^\top)A(I - VV^\top)\|_{\text{F}} + \|(I - U_{\text{cvx}}U_{\text{cvx}}^\top)A(VV^\top - V_{\text{cvx}}V_{\text{cvx}}^\top)\|_{\text{F}} \\ &\lesssim \|A\| \max(\|UU^\top - U_{\text{cvx}}U_{\text{cvx}}^\top\|_{\text{F}}, \|VV^\top - V_{\text{cvx}}V_{\text{cvx}}^\top\|_{\text{F}}) \\ &\lesssim \|A\| \frac{\kappa \log(n)}{\sigma_{\min}^{1.5}} \|\nabla f\|_{\text{F}}.\end{aligned}$$

Furthermore, let $\Delta_U = U - U_{\text{cvx}}R_1^\top, \Delta_V = V - V_{\text{cvx}}R_2^\top$. Then $U\Sigma V^\top = U\Sigma R_2V_{\text{cvx}}^\top + U_{\text{cvx}}R_1^\top\Sigma\Delta_V^\top + \Delta_U\Sigma\Delta_V^\top$. We then have

$$\begin{aligned}&\|P_{T_{\text{cvx}}^\perp}(XY^\top)\|_{\text{F}} \\ &= \|(I - U_{\text{cvx}}U_{\text{cvx}}^\top)(U\Sigma V^\top)(I - V_{\text{cvx}}V_{\text{cvx}}^\top)\|_{\text{F}} \\ &= \|(I - U_{\text{cvx}}U_{\text{cvx}}^\top)(U\Sigma R_2V_{\text{cvx}}^\top + U_{\text{cvx}}R_1^\top\Sigma\Delta_V^\top + \Delta_U\Sigma\Delta_V^\top)(I - V_{\text{cvx}}V_{\text{cvx}}^\top)\|_{\text{F}} \\ &= \|(I - U_{\text{cvx}}U_{\text{cvx}}^\top)(\Delta_U\Sigma\Delta_V^\top)(I - V_{\text{cvx}}V_{\text{cvx}}^\top)\|_{\text{F}} \\ &\leq \|\Delta_U\Sigma\Delta_V^\top\|_{\text{F}} \\ &\leq \|\Delta_U\|_{\text{F}} \|\Delta_V\|_{\text{F}} \|\Sigma\| \\ &\lesssim \frac{\kappa^3 \log^2(n)}{\sigma_{\min}^2} \|\nabla f\|_{\text{F}}^2.\end{aligned}$$

This finishes the proof. □

H Analysis of Non-convex Estimators: Proof of Lemma 5

We introduce some notations to simplify the presentation. Let

$$F^t := \begin{bmatrix} X^t \\ Y^t \end{bmatrix} \in \mathbb{R}^{2n \times r} \quad \text{and} \quad F^* := \begin{bmatrix} X^* \\ Y^* \end{bmatrix} \in \mathbb{R}^{2n \times r}. \quad (35)$$

Note that the $X^t H^t H^{tT} Y^{tT} = X^t Y^{tT}$ for any rotation matrix $H \in \mathcal{O}^{r \times r}$. To avoid the ambiguity, we consider the optimal rotation for (X^t, Y^t) that is defined in the following way.

$$H^t := \arg \min_{R \in \mathcal{O}^{r \times r}} \|F^t R - F^*\|_F. \quad (36)$$

Note that we have the following facts for F^* .

$$\sigma_1(F^*) = \|F^*\| = \sqrt{2\sigma_{\max}}, \quad \sigma_r(F^*) = \sqrt{2\sigma_{\min}}. \quad (37a)$$

We use the mathematical induction to obtain the desired results. In particular, we aim to prove the following lemma.

Lemma 6. *Suppose Assumption 3(a) holds. Assume $O = M^* + \tau^* Z + E$ for a deterministic E . Suppose $\|E\| \lesssim \sigma\sqrt{n}$ and $|\langle Z, E \rangle| \lesssim \sigma\sqrt{n} \|Z\|_F$. Suppose $\frac{\sigma\sqrt{n}}{\sigma_{\min}} \lesssim \frac{1}{r^2 \kappa^2 \log^5(n)}$.*

Let $\lambda = C_\lambda \sigma\sqrt{n} \log^{1.5}(n)$, $\eta = \frac{C_\eta}{\kappa^3 n^{20} \sigma_{\max}}$. Suppose the following induction hypotheses Eq. (38) for $q = t$ hold with (pre-determined) constants C_F, C_B . Then we have the Eq. (38) for $q = t + 1$ holds.

$$\|F^q H^q - F^*\|_F \leq C_F \left(\frac{\sigma\sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_F \quad (38a)$$

$$\|X^{qT} X^q - Y^{qT} Y^q\|_F \leq C_B \frac{\sigma}{\kappa n^{15}}. \quad (38b)$$

Furthermore, if Eq. (38) holds for $q = t$, the following inequality holds.

$$f(X^{q+1}, Y^{q+1}; \tau^{q+1}) \leq f(X^q, Y^q; \tau^q) - \frac{\eta}{2} \|\nabla f(X^q, Y^q; \tau^q)\|_F^2. \quad (38c)$$

Based on Lemma 6, we can prove Lemma 5 which is restated below for convenience.

Lemma 5. *Assume Assumption 3(a) holds and $\frac{\sigma}{\sigma_{\min}} \sqrt{n} \leq C_1 \frac{1}{\kappa^2 r^2 \log^5(n)}$. Suppose $O = M^* + \tau^* Z + E$ for a deterministic E . Assume $\|E\| \leq C_2 \sigma\sqrt{n}$ and $|\langle Z, E \rangle| \leq C_3 \sigma\sqrt{n} \|Z\|_F$.*

Let $\lambda = C_\lambda \sigma\sqrt{n} \log^{1.5}(n)$, $\eta = \frac{C_\eta}{\kappa^3 n^{20} \sigma_{\max}}$, $t^ = \kappa^5 n^{10000}$. Algorithm 1 satisfies the following:*

$$\|F^t H^t - F^*\|_F \leq C_F \left(\frac{\sigma\sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_F \quad (17a)$$

$$\min_{0 \leq t \leq t^*} \|\nabla f(X^t, Y^t; \tau^t)\|_F \leq \frac{\lambda\sqrt{\sigma_{\min}}}{\kappa n^{10}}. \quad (17b)$$

Here, C_F is a constant depending (polynomially) on $C_1, C_2, C_3, C_\lambda, C_\eta, C_{r_1}$ (where C_{r_1} is the constant in Assumption 3(a)).

Proof of Lemma 5. Then we can prove the result based on above Lemma 6. Note that when $q = 0$, $X^0 = X^*$ and $Y^0 = Y^*$ easily satisfy Eq. (38).

Then by Lemma 6, we have Eq. (38) hold for $q = 0, 1, \dots, t^* - 1$. Telescoping the Eq. (38c) implies the following

$$f(X^{t^*}, Y^{t^*}; \tau^{t^*}) \leq f(X^0, Y^0; \tau^0) - \frac{\eta}{2} \sum_{t=0}^{t^*-1} \|\nabla f(X^t, Y^t; \tau^t)\|_F^2.$$

Hence,

$$\begin{aligned} \min_{0 \leq t < t^*} \|\nabla f(X^t, Y^t; \tau^t)\|_F^2 &\leq \frac{1}{t^*} \sum_{t=0}^{t^*-1} \|\nabla f(X^t, Y^t; \tau^t)\|_F^2 \\ &\leq \frac{2}{\eta t^*} (f(X^0, Y^0; \tau^0) - f(X^{t^*}, Y^{t^*}; \tau^{t^*})). \end{aligned}$$

Since $\tau^0 = \arg \min_{\tau} f(X^*, Y^*; \tau)$, hence $f(X^*, Y^*; \tau^0) \leq f(X^*, Y^*; \tau^*)$. Then

$$\begin{aligned}
& f(X^0, Y^0; \tau^0) - f(X^{t^*}, Y^{t^*}; \tau^{t^*}) \\
& \leq f(X^*, Y^*; \tau^*) - f(X^{t^*}, Y^{t^*}; \tau^{t^*}) \\
& = \frac{1}{2} \left(\|E\|_{\mathbb{F}}^2 + \lambda(\|X^*\|_{\mathbb{F}}^2 + \|Y^*\|_{\mathbb{F}}^2) \right) \\
& \quad - \frac{1}{2} \left(\|O - X^{t^*} Y^{t^* \top} - \tau^{t^*} Z\|_{\mathbb{F}}^2 + \lambda \|X^{t^*}\|_{\mathbb{F}}^2 + \lambda \|Y^{t^*}\|_{\mathbb{F}}^2 \right) \\
& \leq \|E\|_{\mathbb{F}}^2 + \lambda \left| \|X^*\|_{\mathbb{F}}^2 - \|X^{t^*}\|_{\mathbb{F}}^2 \right| + \lambda \left| \|Y^*\|_{\mathbb{F}}^2 - \|Y^{t^*}\|_{\mathbb{F}}^2 \right| \\
& = \|E\|_{\mathbb{F}}^2 + \lambda \left| \|X^*\|_{\mathbb{F}} - \|X^{t^*} H^{t^*}\|_{\mathbb{F}} \right| \left(\|X^*\|_{\mathbb{F}} + \|X^{t^*} H^{t^*}\|_{\mathbb{F}} \right) \\
& \quad + \lambda \left| \|Y^*\|_{\mathbb{F}} - \|Y^{t^*} H^{t^*}\|_{\mathbb{F}} \right| \left(\|Y^*\|_{\mathbb{F}} + \|Y^{t^*} H^{t^*}\|_{\mathbb{F}} \right) \\
& \lesssim n \|E\|^2 + \sigma \sqrt{n} \log^{1.5}(n) \|F^{t^*} H^{t^*} - F^*\|_{\mathbb{F}} \|F^*\|_{\mathbb{F}} \\
& \stackrel{(i)}{\lesssim} \sigma^2 n^2 + \sigma \sqrt{n} \log^{1.5}(n) \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \sigma_{\max} r \\
& \lesssim \sigma^2 n^2 + \sigma^2 n \log^4(n) r \kappa.
\end{aligned}$$

Here (i) is due to $\|E\| \lesssim \sigma \sqrt{n}$. This implies

$$\begin{aligned}
& \min_{0 \leq t < t^*} \|\nabla f(X^t, Y^t; \tau^t)\|_{\mathbb{F}} \\
& \leq \sqrt{\frac{2}{\eta_{t^*}}} (f(X^0, Y^0; \tau^0) - f(X^{t^*}, Y^{t^*}; \tau^{t^*})) \\
& \lesssim \sqrt{\frac{\sigma^2 n^2 + \sigma^2 n \log^4(n) r \kappa}{(\kappa^3 n^{20} \sigma_{\max})^{-1} t^*}} \\
& \stackrel{(i)}{\lesssim} \frac{\lambda \sqrt{\sigma_{\min}}}{\kappa n^{10}}.
\end{aligned}$$

Here (i) is obtained by taking $t^* = n^{10000} \kappa^5$. This completes the proof for Lemma 5. \square

The remainder is to finish the proof for Lemma 6. Under the hypothesis for $q = t$, we prove Eq. (38) and Eq. (38c) for $q = t + 1$ in the following subsections.

H.1 Direct implication of Eq. (38) for $q = t$

Under the hypothesis $q = t$, we first present a few direct implications that are helpful in the proof for $q = t + 1$. For large enough n , we have that the following results hold for F^t .

$$\begin{aligned}
\|F^t\|_{\mathbb{F}} & \leq 2 \|F^*\|_{\mathbb{F}} \lesssim \sqrt{\sigma_{\max} r} \\
\|F^t\| & \leq 2 \|F^*\| \lesssim \sqrt{\sigma_{\max}}.
\end{aligned}$$

In addition, we have

$$|\tau^t - \tau^*| \|Z\|_{\mathbb{F}} \lesssim \sigma \sqrt{n} \log^{2.5}(n) r^{0.5} \quad (39)$$

$$\|\nabla f(X^t, Y^t; \tau^t)\|_{\mathbb{F}} \lesssim \sigma r \kappa \log^{2.5}(n) \sqrt{n} \sqrt{\sigma_{\max}} \quad (40)$$

These bounds can be verified directly by the triangle inequality and Eq. (38). We may use them implicitly in the following subsections.

Consider the proof for the above bounds. It is easy to check that $\|F^*\|_{\mathbb{F}} \lesssim \sqrt{\sigma_{\max} r}$, $\|F^*\| \lesssim \sqrt{\sigma_{\max}}$. For $\|F^t\|_{\mathbb{F}}$, note that by Eq. (38a), we have

$$\begin{aligned}
\|F^t H^t - F^*\|_{\mathbb{F}} & \lesssim \left(\frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_{\mathbb{F}} \\
& \stackrel{(i)}{\leq} \frac{1}{2\sqrt{\kappa r}} \sqrt{\sigma_{\max} r} \\
& \leq \frac{\sqrt{\sigma_{\min}}}{2}.
\end{aligned}$$

where (i) is due to $\frac{\sigma\sqrt{n}}{\sigma_{\min}} \lesssim \frac{1}{r^2\kappa^2\log^5(n)}$ and holds for large enough n . Hence, by the triangle inequality,

$$\begin{aligned}\|F^t H^t\|_{\mathbb{F}} &\leq \|F^*\|_{\mathbb{F}} + \|F^t H^t - F^*\|_{\mathbb{F}} \\ &\leq \|F^*\|_{\mathbb{F}} + \frac{\sqrt{\sigma_{\min}}}{2} \\ &\stackrel{(i)}{\leq} 2\|F^*\|_{\mathbb{F}}.\end{aligned}$$

where (i) holds due to $\|F^*\|_{\mathbb{F}} \geq \sqrt{\sigma_{\min}r}$. Then $\|F^t\|_{\mathbb{F}} = \|F^t H^t H^{tT}\|_{\mathbb{F}} \leq \|F^t H^t\|_{\mathbb{F}} \|H^t\| \leq 2\|F^*\|_{\mathbb{F}}$.

Similarly, for $\|F^t H^t\|$, we have

$$\begin{aligned}\|F^t H^t\| &\leq \|F^*\| + \|F^t H^t - F^*\|_{\mathbb{F}} \\ &\leq \|F^*\| + \frac{\sqrt{\sigma_{\min}}}{2} \\ &\stackrel{(i)}{\leq} 2\|F^*\|_{\mathbb{F}}\end{aligned}$$

where (i) is due to $\|F^*\| \geq \sqrt{\sigma_{\min}}$. This implies that $\|F^t\| = \|F^t H^t H^{tT}\| \leq \|F^t H^t\| \|H^t\| \leq 2\|F^*\|$.

For $|\tau^t - \tau^*| \|Z\|_{\mathbb{F}}$, we have

$$\begin{aligned}|\tau^t - \tau^*| \|Z\|_{\mathbb{F}} &= \left| \frac{\langle Z, X^t Y^{tT} - X^* Y^{*T} \rangle}{\|Z\|_{\mathbb{F}}^2} - \frac{\langle Z, E \rangle}{\|Z\|_{\mathbb{F}}^2} \right| \|Z\|_{\mathbb{F}} \\ &\leq \|X^t Y^{tT} - X^* Y^{*T}\|_{\mathbb{F}} + |\langle Z, E \rangle| \frac{1}{\|Z\|_{\mathbb{F}}} \\ &\stackrel{(i)}{\lesssim} \|X^t H^t - X^*\|_{\mathbb{F}} \|Y\| + \|X^*\| \|Y^t H^t - Y^*\|_{\mathbb{F}} + \sigma\sqrt{n} \\ &\stackrel{(ii)}{\lesssim} \sigma\sqrt{n}\kappa r^{0.5} \log^{2.5}(n)\end{aligned}$$

where (i) is due to $|\langle Z, E \rangle| \lesssim \sigma\sqrt{n}$ and (ii) is due to $\|F^t H^t - F^*\|_{\mathbb{F}} \lesssim \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}\right) \sigma_{\max}^{0.5} r^{0.5}$ (Eq. (38a)).

For $\|\nabla f(X^t, Y^t; \tau^t)\|_{\mathbb{F}}$, note that

$$\begin{aligned}&\|\nabla_X f(X^t, Y^t; \tau^t)\|_{\mathbb{F}} \\ &= \|(X^t Y^{tT} - M^* - E + (\tau^t - \tau^* Z)) Y^t + \lambda X^t\|_{\mathbb{F}} \\ &\leq (\|X^t Y^{tT} - M^*\| + \|E\| + |\tau^t - \tau^*| \|Z\|) \|Y^t\|_{\mathbb{F}} + \lambda \|X^t\|_{\mathbb{F}} \\ &\leq (\|X^t H^t - X^*\| \|Y^t\| + \|X^*\| \|Y^t H^t - Y^*\| + \|E\| + |\tau^t - \tau^*| \|Z\|) \|Y^t\|_{\mathbb{F}} \\ &\quad + \lambda \|X^t\|_{\mathbb{F}} \\ &\stackrel{(i)}{\lesssim} \left(\left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \right) \sigma_{\max}^{0.5} r^{0.5} + \sigma\sqrt{n} + \sigma\sqrt{n}\kappa r^{0.5} \log^{2.5}(n) \right) \sqrt{\sigma_{\max}r} \\ &\quad + \sigma\sqrt{n} \log^{1.5}(n) \sqrt{\sigma_{\max}r}.\end{aligned}$$

In (i), we use that $\|F^t H^t - F^*\|_{\mathbb{F}} \lesssim \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}\right) \sigma_{\max}^{0.5} r^{0.5}$ (Eq. (38a)), $\|E\| \lesssim \sigma\sqrt{n}$ and $|\tau^t - \tau^*| \lesssim \sigma\sqrt{n}\kappa r^{0.5} \log^{2.5}(n)$ (Eq. (39)).

This implies

$$\|\nabla_X f(X^t, Y^t; \tau^t)\|_{\mathbb{F}} \lesssim \sigma\sqrt{n}r\kappa \log^{2.5}(n) \sqrt{\sigma_{\max}}. \quad (41)$$

Similarly, one can get $\|\nabla_Y f(X^t, Y^t; \tau^t)\|_{\mathbb{F}} \lesssim \sigma\sqrt{n}r\kappa \log^{2.5}(n) \sqrt{\sigma_{\max}}$. Hence

$$\begin{aligned}\|\nabla f(X^t, Y^t; \tau^t)\|_{\mathbb{F}} &\lesssim \|\nabla_X f(X^t, Y^t; \tau^t)\|_{\mathbb{F}} + \|\nabla_Y f(X^t, Y^t; \tau^t)\|_{\mathbb{F}} \\ &\lesssim \sigma\sqrt{n}r\kappa \log^{2.5}(n) \sqrt{\sigma_{\max}},\end{aligned}$$

which completes the proof.

H.2 Proof of Eq. (38a)

Recall that we want to show

$$\|F^{t+1}H^{t+1} - F^*\|_F \leq C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_F.$$

Write $\nabla f(X^t, Y^t; \tau^t)$ as $\nabla f(F^t; \tau^t)$ for simplification. Consider

$$\begin{aligned} \|F^{t+1}H^{t+1} - F^*\|_F^2 &\stackrel{(i)}{\leq} \|F^{t+1}H^t - F^*\|_F^2 \\ &\stackrel{(ii)}{=} \langle F^tH^t - F^* - \eta\nabla f(F^t; \tau^t)H^t, F^tH^t - F^* - \eta\nabla f(F^t; \tau^t)H^t \rangle. \end{aligned} \quad (42)$$

Here, (i) is due to that H^{t+1} is the optimal rotation to align F^{t+1} and F^* , and (ii) is due to $F^{t+1} = F^t - \eta\nabla f(F^t; \tau^t)$. Write $F^tH^t, X^tH^t, Y^tH^t, \tau^t$ as F, X, Y, τ if there is no ambiguity. Let $\Delta_X = X - X^*, \Delta_Y = Y - Y^*, \Delta_F = F - F^*$. Then we have

$$\begin{aligned} \|F^{t+1}H^t - F^*\|_F^2 &\stackrel{(i)}{=} \|\Delta_F\|_F^2 - 2\eta \langle \Delta_F, \nabla f(F; \tau)H^t \rangle + A_0 \\ &= \|\Delta_F\|_F^2 - 2\eta \langle \Delta_X, (XY^\top - M^* - E + (\tau - \tau^*)Z)Y + \lambda X \rangle \\ &\quad - 2\eta \langle \Delta_Y, (XY^\top - M^* - E + (\tau - \tau^*)Z)^\top X + \lambda Y \rangle + A_0 \end{aligned} \quad (43)$$

where in (i) A_0 is the term with η^2 coefficient

$$A_0 := \eta^2 \|\nabla f(F; \tau)\|_F^2.$$

Recall that $\|\nabla f(F; \tau)\|_F^2 \lesssim \sigma\sqrt{nr}\kappa \log^{2.5}(n)\sqrt{\sigma_{\max}}$ (Eq. (41)). Then we have

$$\begin{aligned} |A_0| &\lesssim \eta^2 (\sigma\sqrt{nr}\kappa \log^{2.5}(n)\sqrt{\sigma_{\max}})^2 \\ &\lesssim \eta \frac{1}{\kappa^3 n^{20} \sigma_{\max}} \sigma^2 r^2 \kappa^2 \log^5(n) n \sigma_{\max} \\ &\lesssim \eta \frac{\sigma^2}{n^{15}}. \end{aligned} \quad (44)$$

Continue the analysis on Eq. (43), by some algebra, we have

$$\begin{aligned} &\|F^{t+1}H^t - F^*\|_F^2 \\ &= \|\Delta_F\|_F^2 - 2\eta \langle \Delta_X, (XY^\top - M^* + (\tau - \tau^*)Z)Y \rangle \\ &\quad - 2\eta \langle \Delta_Y, (XY^\top - M^* + (\tau - \tau^*)Z)^\top X \rangle \\ &\quad + \underbrace{2\eta \langle \Delta_X, EY \rangle + 2\eta \langle \Delta_Y, E^\top X \rangle}_{A_1} + \underbrace{2\eta \langle \Delta_X, -\lambda X \rangle + 2\eta \langle \Delta_Y, -\lambda Y \rangle}_{A_2} + A_0 \\ &\leq \|\Delta_F\|_F^2 - 2\eta \langle XY^\top - M^* + (\tau - \tau^*)Z, \Delta_X Y^\top + X \Delta_Y^\top \rangle + A_0 + A_1 + A_2 \\ &\stackrel{(i)}{=} \|\Delta_F\|_F^2 - 2\eta \underbrace{\langle XY^\top - M^* + (\tau - \tau^*)Z, \Delta_X Y^{*\top} + X^* \Delta_Y^\top \rangle}_{A_3} + A_0 + A_1 + A_2 \\ &\quad - \underbrace{2\eta \langle XY^\top - M^* + (\tau - \tau^*)Z, 2\Delta_X \Delta_Y^\top \rangle}_{A_4}. \end{aligned} \quad (45)$$

where (i) is due to $Y = Y^* + \Delta_Y, X = X^* + \Delta_X$. For A_1, A_2 , and A_4 , we have the following claim.

Claim 2.

$$\begin{aligned} |A_1| &\lesssim \eta\sigma\sqrt{n} \|F^*\|_F \|\Delta_F\|_F \\ |A_2| &\lesssim \eta\sigma\sqrt{n} \log^{1.5}(n) \|F^*\|_F \|\Delta_F\|_F \\ |A_4| &\lesssim \eta\sigma\sqrt{n} \log(n) \|F^*\|_F \|\Delta_F\|_F \end{aligned}$$

For A_3 , note that $XY^\top - M^* = \Delta_X Y^{*\top} + X^* \Delta_Y^\top + \Delta_X \Delta_Y^\top$, $\tau - \tau^* = \frac{\langle Z, M^* - XY^\top + E \rangle}{\|Z\|_F^2}$, then we have

$$\begin{aligned}
A_3 &= \langle XY^\top - M^* + (\tau - \tau^*)Z, \Delta_X Y^{*\top} + X^* \Delta_Y^\top \rangle \\
&= \|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_F^2 + \underbrace{\langle \Delta_X \Delta_Y^\top, \Delta_X Y^{*\top} + X^* \Delta_Y^\top \rangle}_{B_0} \\
&\quad + \frac{\langle Z, M^* - XY^\top \rangle \langle Z, \Delta_X Y^{*\top} + X^* \Delta_Y^\top \rangle}{\|Z\|_F^2} + \underbrace{\frac{\langle Z, E \rangle \langle Z, \Delta_X Y^{*\top} + X^* \Delta_Y^\top \rangle}{\|Z\|_F^2}}_{B_1} \\
&\stackrel{(i)}{=} \|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_F^2 - \frac{\langle Z, \Delta_X Y^{*\top} + X^* \Delta_Y^\top \rangle \langle Z, \Delta_X Y^{*\top} + X^* \Delta_Y^\top \rangle}{\|Z\|_F^2} \\
&\quad - \underbrace{\frac{\langle Z, \Delta_X \Delta_Y^\top \rangle \langle Z, \Delta_X Y^{*\top} + X^* \Delta_Y^\top \rangle}{\|Z\|_F^2}}_{B_2} + B_0 + B_1
\end{aligned}$$

In (i), we use again $XY^\top - M^* = \Delta_X Y^{*\top} + X^* \Delta_Y^\top + \Delta_X \Delta_Y^\top$.

We then have the following claim to control B_0, B_1, B_2 .

Claim 3.

$$\begin{aligned}
|B_0| &\lesssim \sigma \sqrt{n} \|\Delta_F\| \|F^*\|_F \\
|B_1| &\lesssim \sigma \sqrt{n} \|\Delta_F\|_F \|F^*\|_F \\
|B_2| &\lesssim \sigma \sqrt{n} \|\Delta_F\| \|F^*\|_F
\end{aligned}$$

Let $T^* = \{BV^{*\top} + U^*A^\top \mid A, B \in \mathbb{R}^{n \times r}\}$ be the tangent space of M^* . Then $\Delta_X Y^{*\top} + X^{*\top} \Delta_Y^\top = \Delta_X \Sigma^{*1/2} V^{*\top} + U^* \Sigma^{*1/2} \Delta_Y^\top \in T^*$. Let P_{T^*} be the projection to T^* . Then we have

$$\begin{aligned}
|\langle Z, \Delta_X Y^{*\top} + X^{*\top} \Delta_Y^\top \rangle|^2 &= |\langle P_{T^*}(Z), \Delta_X Y^{*\top} + X^{*\top} \Delta_Y^\top \rangle|^2 \\
&\leq \|P_{T^*}(Z)\|_F^2 \|\Delta_X Y^{*\top} + X^{*\top} \Delta_Y^\top\|_F^2
\end{aligned}$$

This implies there exists a constant C , such that

$$\begin{aligned}
A_3 &\geq \|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_F^2 - \frac{\|P_{T^*}(Z)\|_F^2}{\|Z\|_F^2} \|\Delta_X Y^{*\top} + X^{*\top} \Delta_Y^\top\|_F^2 - B_2 + B_0 + B_1 \\
&\stackrel{(i)}{\geq} \frac{C}{\log(n)} \underbrace{\|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_F^2}_{B_3} - B_2 + B_0 + B_1. \tag{46}
\end{aligned}$$

where (i) is due to Assumption 3(a) and Lemma 9. We then have the following claim for connecting $\|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_F^2$ and $\|\Delta_F\|_F^2$.

Claim 4.

$$\|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_F^2 \geq \frac{\sigma_{\min}}{2} \|\Delta_F\|_F^2 - \frac{\sigma^2}{n^{13}}. \tag{47}$$

Combining Eq. (45), Eq. (46), and Eq. (47), we arrive at

$$\begin{aligned}
\|F^{t+1}H^t - F^*\|_F^2 &\stackrel{(i)}{=} \|\Delta_F\|_F^2 - 2\eta A_3 + A_0 + A_1 + A_2 + A_4 \\
&\stackrel{(ii)}{\leq} \|\Delta_F\|_F^2 - 2\eta \left(\frac{C}{\log(n)} \|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_F^2 - B_2 + B_0 + B_1 \right) \\
&\quad + A_0 + A_1 + A_2 + A_4 \\
&\stackrel{(iii)}{\leq} \|\Delta_F\|_F^2 - 2\eta \left(\frac{C}{\log(n)} \frac{\sigma_{\min}}{2} \|\Delta_F\|_F^2 - \underbrace{\frac{C\sigma^2}{\log(n)n^{13}}}_{B_3} - B_2 + B_0 + B_1 \right) \\
&\quad + A_0 + A_1 + A_2 + A_4
\end{aligned} \tag{48}$$

where (i) is due to Eq. (45), (ii) is due to Eq. (46), and (iii) is due to Eq. (47). Next, due to the bound on A_0 (Eq. (44)), the bound on A_1, A_2, A_4 (Claim 2), and the bound on B_0, B_1, B_2 (Claim 3), we have

$$\begin{aligned}
&|A_0| + |A_1| + |A_2| + |A_4| + \eta|B_0| + \eta|B_1| + \eta|B_2| + \eta|B_3| \\
&\lesssim \eta\sigma\sqrt{n}\log^{1.5}(n) \|F^*\|_F \|\Delta_F\|_F + \eta\frac{\sigma^2}{n^{13}}.
\end{aligned}$$

Continue the analysis for Eq. (48), we then have (for some constant C'),

$$\begin{aligned}
&\|F^{t+1}H^t - F^*\|_F^2 \\
&\leq \|\Delta_F\|_F^2 \left(1 - \eta \frac{C\sigma_{\min}}{\log n} \right) + |A_0| + |A_1| + |A_2| + |A_4| + 2\eta(|B_0| + |B_1| + |B_2| + |B_3|) \\
&\leq \|\Delta_F\|_F^2 \left(1 - \eta \frac{C\sigma_{\min}}{\log n} \right) + C'\eta\sigma\sqrt{n}\log^{1.5}(n) \|\Delta_F\|_F \|F^*\|_F + C'\eta\frac{\sigma^2}{n^{13}}.
\end{aligned} \tag{49}$$

Plug $\|\Delta_F\| \leq C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_F$ into above Eq. (49), we then get

$$\begin{aligned}
&\|F^{t+1}H^t - F^*\|_F^2 \\
&\leq \left(C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_F \right)^2 \left(1 - \eta \frac{C\sigma_{\min}}{\log n} \right) \\
&\quad + C'\eta\sigma\sqrt{n}\log^{1.5}(n) C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_F \|F^*\|_F + C'\eta\frac{\sigma^2}{n^{13}} \\
&= \left(C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_F \right)^2 - \underbrace{\eta \frac{C\sigma_{\min}}{\log n} \left(C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_F \right)^2}_{K_1} \\
&\quad + \underbrace{\eta C'\sigma\sqrt{n}\log^{1.5}(n) C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_F \|F^*\|_F}_{K_2} + \underbrace{\eta C'\frac{\sigma^2}{n^{13}}}_{K_3}.
\end{aligned}$$

Then, in order to show $\|F^{t+1}H^t - F^*\|_F^2 \leq \left(C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}\right) \|F^*\|_F\right)^2$, it is sufficient to show that $K_1 \geq K_2 + K_3$. Note that

$$\begin{aligned} K_1 - K_2 &= \frac{C\sigma_{\min}}{\log n} \left(C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}\right) \|F^*\|_F\right) \left(C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}\right) \|F^*\|_F\right) \\ &\quad - C'\sigma\sqrt{n}\log^{1.5}(n)C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}\right) \|F^*\|_F^2 \\ &= CC_F\sigma\sqrt{n}\log^{1.5}(n)C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}\right) \|F^*\|_F^2 \\ &\quad - C'\sigma\sqrt{n}\log^{1.5}(n)C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}\right) \|F^*\|_F^2 \\ &= (CC_F - C')\sigma\sqrt{n}\log^{1.5}(n)C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}\right) \|F^*\|_F^2. \end{aligned}$$

Then, take $C_F \geq \max(2C'/C, 1)$, we have

$$\begin{aligned} K_1 - K_2 &\geq C'\sigma\sqrt{n}\log^{1.5}(n) \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}\right) \sigma_{\max} \\ &= C'\sigma^2 n \log^4(n) \kappa \\ &\geq C' \frac{\sigma^2}{n^{13}} \\ &= K_3. \end{aligned}$$

This implies that $\|F^{t+1}H^t - F^*\|_F^2 \leq \left(C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}\right) \|F^*\|_F\right)^2$. Recall that (by Eq. (42))

$$\|F^{t+1}H^{t+1} - F^*\|_F \leq \|F^{t+1}H^t - F^*\|_F,$$

hence $\|F^{t+1}H^{t+1} - F^*\|_F \leq C_F \left(\frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}\right) \|F^*\|_F$, which completes the proof for Eq. (59c).

Proof of Claim 2. In this claim, we provide bounds for A_1, A_2, A_4 .

Recall that $A_1 = 2\eta \langle \Delta_X, EY \rangle + 2\eta \langle \Delta_Y, E^\top X \rangle$ and we want to show

$$|A_1| \lesssim \eta\sigma\sqrt{n} \|F^*\|_F \|\Delta_F\|_F.$$

This can be verified by the following.

$$\begin{aligned} |A_1| &\lesssim \eta \|\Delta_X\|_F \|EY\|_F + \eta \|\Delta_Y\|_F \|E^\top X\|_F \\ &\lesssim \eta \|\Delta_F\|_F \|E\| \|F\|_F \\ &\stackrel{(i)}{\lesssim} \eta\sigma\sqrt{n} \|\Delta_F\|_F \|F^*\|_F. \end{aligned}$$

where (i) is due to $\|E\| \lesssim \sigma\sqrt{n}$.

Recall that $A_2 = 2\eta \langle \Delta_X, -\lambda X \rangle + 2\eta \langle \Delta_Y, -\lambda Y \rangle$ and we want to show

$$|A_2| \lesssim \eta\sigma\sqrt{n}\log^{1.5}(n) \|F^*\|_F \|\Delta_F\|_F.$$

This can be verified by the following.

$$\begin{aligned} |A_2| &\lesssim \eta\lambda \|\Delta_X\|_F \|Y\|_F + \eta\lambda \|\Delta_Y\|_F \|X\|_F \\ &\lesssim \eta\lambda \|\Delta_F\|_F \|F\|_F \\ &\lesssim \eta\sigma\sqrt{n}\log^{1.5}(n) \|\Delta_F\|_F \|F^*\|_F. \end{aligned}$$

Recall that $A_4 = 2\eta \langle XY^\top - M^* + (\tau - \tau^*)Z, 2\Delta_X\Delta_Y^\top \rangle$ and we want to show

$$|A_4| \lesssim \eta\sigma\sqrt{n}\log(n) \|F^*\|_F \|\Delta_F\|_F.$$

This can be verified by

$$\begin{aligned}
|A_4| &\lesssim \eta \|\Delta_X \Delta_Y^\top\|_F (\|XY^\top - M^*\|_F + |\tau - \tau^*| \|Z\|_F) \\
&\leq \eta \|\Delta_X\|_F \|\Delta_Y\|_F (\|\Delta_X Y^\top\|_F + \|X^* \Delta_Y^\top\|_F + |\tau - \tau^*| \|Z\|_F) \\
&\stackrel{(i)}{\lesssim} \eta \|\Delta_F\|_F^2 (\|\Delta_F\|_F \|F^*\| + \sigma \sqrt{n} \kappa r^{0.5} \log^{2.5}(n)).
\end{aligned} \tag{50}$$

where (i) is due to $|\tau - \tau^*| \|Z\|_F \lesssim \sigma \sqrt{n} \kappa r^{0.5} \log^{2.5}(n)$ (by Eq. (39)).

Next, recall that $\|\Delta_F\|_F \lesssim \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \|F^*\|_F$. This implies that

$$\begin{aligned}
&\|\Delta_F\|_F \|F^*\| + \sigma \sqrt{n} \kappa r^{0.5} \log^{2.5}(n) \\
&\lesssim \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \|F^*\|_F \|F^*\| + \sigma \sqrt{n} \kappa r^{0.5} \log^{2.5}(n) \\
&\lesssim \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \sigma_{\max} \sqrt{r} + \sigma \sqrt{n} \kappa r^{0.5} \log^{2.5}(n) \\
&\lesssim \sigma \sqrt{n} \kappa r^{0.5} \log^{2.5}(n).
\end{aligned}$$

Combining this with Eq. (50), we obtain

$$\begin{aligned}
|A_4| &\lesssim \eta \|\Delta_F\|_F^2 \sigma \sqrt{n} \kappa r^{0.5} \log^{2.5}(n) \\
&\stackrel{(i)}{\lesssim} \eta \|\Delta_F\|_F \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \|F^*\|_F \sigma \sqrt{n} \kappa r^{0.5} \log^{2.5}(n) \\
&\stackrel{(ii)}{\lesssim} \eta \sigma \sqrt{n} \log(n) \|F^*\|_F \|\Delta_F\|_F
\end{aligned}$$

where in (i) we use $\|\Delta_F\|_F \lesssim \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \|F^*\|_F$ again, in (ii) we use $\frac{\sigma \sqrt{n}}{\sigma_{\min}} \lesssim \frac{1}{r^2 \kappa^2 \log^5(n)}$. This completes the proof for the bounds on A_1, A_2, A_4 . \square

Claim 3. In this claim, we aim to provide bounds on B_0, B_1, B_2 .

For B_0 , recall that $B_0 = \langle \Delta_X \Delta_Y^\top, \Delta_X Y^{*\top} + X^* \Delta_Y^\top \rangle$ and we want to show $|B_0| \lesssim \sigma \sqrt{n} \|\Delta_F\| \|F^*\|_F$. This can be verified by

$$\begin{aligned}
|B_0| &\lesssim \|\Delta_X\|_F \|\Delta_Y\|_F (\|\Delta_X\|_F \|Y^*\| + \|\Delta_Y\|_F \|X^*\|) \\
&\lesssim \|\Delta_F\|_F^3 \|F^*\| \\
&\lesssim \|\Delta_F\| \|F^*\|_F \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \sqrt{\sigma_{\max} r} \sqrt{\sigma_{\max}} \\
&\lesssim \|\Delta_F\| \|F^*\|_F \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \sigma \sqrt{n} \log^{2.5}(n) \kappa r^{0.5} \\
&\lesssim \sigma \sqrt{n} \|\Delta_F\| \|F^*\|_F
\end{aligned}$$

providing that $\frac{\sigma \sqrt{n}}{\sigma_{\min}} \lesssim \frac{1}{r^2 \kappa^2 \log^5(n)}$.

For B_1 , recall that $B_1 = \frac{\langle Z, E \rangle \langle Z, \Delta_X Y^{*\top} + X^* \Delta_Y^\top \rangle}{\|Z\|_F^2}$ and we want to show

$$|B_1| \lesssim \sigma \sqrt{n} \|\Delta_F\|_F \|F^*\|_F.$$

Note that $|\langle Z, E \rangle| / \|Z\|_F^2 \lesssim \sigma \sqrt{n} / \|Z\|_F$. Then

$$\begin{aligned}
|B_1| &\lesssim \frac{\sigma \sqrt{n}}{\|Z\|_F} \|Z\|_F \|\Delta_F\|_F \|F^*\| \\
&\lesssim \sigma \sqrt{n} \|\Delta_F\|_F \|F^*\|_F.
\end{aligned}$$

For B_2 , recall that $B_2 = \frac{\langle Z, \Delta_X \Delta_Y^\top \rangle \langle Z, \Delta_X Y^{*\top} + X^* \Delta_Y^\top \rangle}{\|Z\|_F^2}$ and we want to show $|B_2| \lesssim \sigma\sqrt{n} \|\Delta_F\| \|F^*\|_F$. Note that

$$\begin{aligned} |B_2| &\lesssim \frac{(\|Z\|_F \|\Delta_F\|_F \|\Delta_F\|_F)(\|Z\|_F \|\Delta_F\|_F \|F^*\|)}{\|Z\|_F^2} \\ &\lesssim \|\Delta_F\|_F^3 \|F^*\| \\ &\stackrel{(i)}{\lesssim} \sigma\sqrt{n} \|\Delta_F\| \|F^*\|_F \end{aligned}$$

where (i) is due to the similar analysis for B_0 . This completes the proof for the bounds on B_0, B_1, B_2 . \square

Proof of Claim 4. In this claim, we want to show $\|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_F^2 \geq \frac{\sigma_{\min}}{2} \|\Delta_F\|_F^2 - \frac{\sigma^2}{n^{13}}$.

Note that

$$\begin{aligned} \|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_F^2 &= \text{tr}((\Delta_X Y^{*\top} + X^* \Delta_Y^\top)(\Delta_X Y^{*\top} + X^* \Delta_Y^\top)^\top) \\ &= \|\Delta_X Y^{*\top}\|_F^2 + \|X^* \Delta_Y^\top\|_F^2 + \text{tr}(\Delta_X Y^{*\top} \Delta_Y X^{*\top} + X^* \Delta_Y^\top Y^* \Delta_X^\top) \\ &= \|\Delta_X Y^{*\top}\|_F^2 + \|X^* \Delta_Y^\top\|_F^2 + 2\text{tr}(X^* \Delta_Y^\top Y^* \Delta_X^\top). \end{aligned}$$

Consider

$$\|\Delta_X Y^{*\top}\|_F^2 = \sum_{i=1}^n \|(\Delta_X)_{i,\cdot} \cdot Y^{*\top}\|^2 \geq \sum_{i=1}^n \sigma_r^2(Y^*) \|(\Delta_X)_{i,\cdot}\|^2 = \sigma_{\min} \|\Delta_X\|_F^2.$$

Similarly, $\|X^* \Delta_Y^\top\|_F^2 \geq \sigma_{\min} \|\Delta_Y\|_F^2$. This implies that

$$\begin{aligned} \|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_F^2 &\geq \sigma_{\min} \|\Delta_X\|_F^2 + \sigma_{\min} \|\Delta_Y\|_F^2 + 2\text{tr}(X^* \Delta_Y^\top Y^* \Delta_X^\top) \\ &= \sigma_{\min} \|\Delta_F\|_F^2 + 2\text{tr}(\Delta_X^\top X^* \Delta_Y^\top Y^*). \end{aligned}$$

Furthermore, note that we can do the following derivation to reduce the problem to control $\|\Delta_X^\top X^* - Y^{*\top} \Delta_Y\|_F$.

$$\begin{aligned} &\|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_F^2 \\ &\geq \sigma_{\min} \|\Delta_F\|_F^2 + 2\text{tr}(\Delta_X^\top X^* \Delta_Y^\top Y^*) \\ &= \sigma_{\min} \|\Delta_F\|_F^2 + 2\text{tr}(Y^{*\top} \Delta_Y \Delta_Y^\top Y^*) + 2\text{tr}((\Delta_X^\top X^* - Y^{*\top} \Delta_Y) \Delta_Y^\top Y^*) \\ &= \sigma_{\min} \|\Delta_F\|_F^2 + 2\|Y^{*\top} \Delta_Y\|_F^2 + 2\text{tr}((\Delta_X^\top X^* - Y^{*\top} \Delta_Y) \Delta_Y^\top Y^*) \\ &\stackrel{(i)}{\geq} \sigma_{\min} \|\Delta_F\|_F^2 - 2\|\Delta_X^\top X^* - Y^{*\top} \Delta_Y\|_F \|Y^{*\top} \Delta_Y\|_F. \end{aligned} \tag{51}$$

Here, in (i) we use $\|Y^{*\top} \Delta_Y\|_F^2 \geq 0$ and $\text{tr}(AB^\top) \leq \|A\|_F \|B\|_F$ by Cauchy-Schwartz inequality.

Then it boils down to control $\|\Delta_X^\top X^* - Y^{*\top} \Delta_Y\|_F$. In order to show $\Delta_X^\top X^* \approx Y^{*\top} \Delta_Y$, we tend to invoke Lemma 11 and use the property that $X^\top X \approx Y^\top Y$. Note that by Lemma 11, we have $(F^t H^t)^\top F^*$ is symmetric since H^t is the optimal rotation to align F^t and F^* . Recall that $X = X^t H^t, Y = Y^t H^t$. This implies that $X^\top X^* + Y^\top Y^*$ is symmetric. Furthermore, since $X^{*\top} X^*, Y^{*\top} Y^*$ is symmetric, then

$$\Delta_X^\top X^* + \Delta_Y^\top Y^* = X^\top X^* + Y^\top Y^* - X^{*\top} X^* - Y^{*\top} Y^*$$

is also symmetric. Therefore $\Delta_X^\top X^* + \Delta_Y^\top Y^* = (\Delta_X^\top X^* + \Delta_Y^\top Y^*)^\top = X^{*\top} \Delta_X + Y^{*\top} \Delta_Y$, which further implies

$$\Delta_X^\top X^* - Y^{*\top} \Delta_Y = X^{*\top} \Delta_X - \Delta_Y^\top Y^*. \tag{52}$$

On the other hand, note that

$$\begin{aligned}
X^\top X - Y^\top Y &= (X^* + \Delta_X)^\top (X^* + \Delta_X) - (Y^* + \Delta_Y)^\top (Y^* + \Delta_Y) \\
&= X^{*\top} X - Y^{*\top} Y + \Delta_X^\top \Delta_X - \Delta_Y^\top \Delta_Y \\
&\quad + (\Delta_X^\top X^* - Y^{*\top} \Delta_Y) + (X^{*\top} \Delta_X - \Delta_Y^\top Y^*) \\
&\stackrel{(i)}{=} \Delta_X^\top \Delta_X - \Delta_Y^\top \Delta_Y + (\Delta_X^\top X^* - Y^{*\top} \Delta_Y) + (X^{*\top} \Delta_X - \Delta_Y^\top Y^*) \\
&\stackrel{(ii)}{=} \Delta_X^\top \Delta_X - \Delta_Y^\top \Delta_Y + 2(\Delta_X^\top X^* - Y^{*\top} \Delta_Y)
\end{aligned} \tag{53}$$

where (i) is due to $X^{*\top} X^* = Y^{*\top} Y^* = \Sigma^*$, and (ii) is due to Eq. (52). Then Eq. (53) implies that

$$\Delta_X^\top X^* - Y^{*\top} \Delta_Y = \frac{1}{2} (X^\top X - Y^\top Y - \Delta_X^\top \Delta_X + \Delta_Y^\top \Delta_Y).$$

Plug this equality into Eq. (51), we obtain

$$\begin{aligned}
&\|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_{\mathbb{F}}^2 \\
&\geq \sigma_{\min} \|\Delta_F\|_{\mathbb{F}}^2 - 2 \|\Delta_X^\top X^* - Y^{*\top} \Delta_Y\|_{\mathbb{F}} \|Y^{*\top} \Delta_Y\|_{\mathbb{F}} \\
&\geq \sigma_{\min} \|\Delta_F\|_{\mathbb{F}}^2 - \|X^\top X - Y^\top Y - \Delta_X^\top \Delta_X + \Delta_Y^\top \Delta_Y\|_{\mathbb{F}} \|Y^*\| \|\Delta_Y\|_{\mathbb{F}}.
\end{aligned}$$

Recall that $\|X^\top X - Y^\top Y\|_{\mathbb{F}} = \|H^{tT} (X^{tT} X^t - Y^{tT} Y^t) H^t\| \lesssim \frac{\sigma}{\kappa n^{15}}$ (by Eq. (38b)) and

$\|\Delta_F\|_{\mathbb{F}} \lesssim \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \sqrt{\sigma_{\max} r}$, then we have

$$\begin{aligned}
&\|X^\top X - Y^\top Y - \Delta_X^\top \Delta_X + \Delta_Y^\top \Delta_Y\|_{\mathbb{F}} \|Y^*\| \|\Delta_Y\|_{\mathbb{F}} \\
&\leq (\|X^\top X - Y^\top Y\|_{\mathbb{F}} + \|\Delta_X^\top \Delta_X\|_{\mathbb{F}} + \|\Delta_Y^\top \Delta_Y\|_{\mathbb{F}}) \sqrt{\sigma_{\max}} \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \sqrt{\sigma_{\max} r} \\
&\lesssim \left(\frac{\sigma}{\kappa n^{15}} + \|\Delta_F\|_{\mathbb{F}}^2 \right) \sigma \sqrt{n} \kappa r^{0.5} \log^{2.5}(n) \\
&\stackrel{(i)}{\lesssim} \|\Delta_F\|_{\mathbb{F}}^2 \frac{\sigma_{\min}}{\log^{0.5}(n)} + \frac{\sigma^2}{n^{13}}.
\end{aligned}$$

Here, (i) is due to $\frac{\sigma \sqrt{n}}{\sigma_{\min}} \lesssim \frac{1}{r^2 \kappa^2 \log^5(n)}$. Then this implies, for large enough n ,

$$\begin{aligned}
\|\Delta_X Y^{*\top} + X^* \Delta_Y^\top\|_{\mathbb{F}}^2 &\geq \sigma_{\min} \|\Delta_F\|_{\mathbb{F}}^2 - \|\Delta_F\|_{\mathbb{F}}^2 \frac{\sigma_{\min}}{2} - \frac{\sigma^2}{n^{13}} \\
&\geq \frac{\sigma_{\min}}{2} \|\Delta_F\|_{\mathbb{F}}^2 - \frac{\sigma^2}{n^{13}}
\end{aligned}$$

which completes the proof. \square

H.3 Proof of Eq. (38b)

Following the algebra of section D.8 in [4], one can verify the following equality

$$A^{t+1} = (1 - \lambda\eta)^2 A^t + \eta^2 (Y^{tT} D^{tT} D^t Y^t - X^{tT} D^t D^{tT} X^t)$$

where

$$\begin{aligned}
A^t &:= X^{tT} X^t - Y^{tT} Y^t \\
A^{t+1} &:= X^{(t+1)\top} X^{t+1} - Y^{(t+1)\top} Y^{t+1} \\
D^t &:= X^t Y^{tT} - M^* - E + (\tau^t - \tau^*) Z.
\end{aligned}$$

Similar to showing $\|\nabla f(X^t, Y^t; \tau^t)\| \lesssim \sigma \sqrt{n} r \kappa \log^{2.5}(n) \sqrt{\sigma_{\max}}$ in Eq. (41), one can verify that $\max(\|D^t Y^t\|_{\mathbb{F}}, \|D^{tT} X^t\|_{\mathbb{F}}) \lesssim \sigma \sqrt{n} r \kappa \log^{2.5}(n) \sqrt{\sigma_{\max}}$. Then there exists a constant C , where

$$\begin{aligned}
\|A^{t+1}\|_{\mathbb{F}} &\leq (1 - \lambda\eta)^2 \|A^t\|_{\mathbb{F}} + \eta^2 \left(\|D^t Y^t\|_{\mathbb{F}}^2 + \|D^{tT} X^t\|_{\mathbb{F}}^2 \right) \\
&\leq (1 - \lambda\eta) \|A^t\|_{\mathbb{F}} + C \eta^2 (\sigma \sqrt{n} r \kappa \log^{2.5}(n) \sqrt{\sigma_{\max}})^2 \\
&\leq C_B \frac{\sigma}{\kappa n^{15}} - C_B \lambda \eta \frac{\sigma}{\kappa n^{15}} + C \eta^2 (\sigma \sqrt{n} r \kappa \log^{2.5}(n) \sqrt{\sigma_{\max}})^2.
\end{aligned}$$

In order to show the desired bound $\|A^{t+1}\|_{\mathbb{F}} \leq C_B \frac{\sigma}{\kappa n^{15}}$, it is sufficient to show that $C_B \lambda \eta \frac{\sigma}{\kappa n^{15}} \geq C \eta^2 (\sigma \sqrt{nr} \kappa \log^{2.5}(n) \sqrt{\sigma_{\max}})^2$. This can be obtained by noting that

$$\begin{aligned} \frac{\kappa n^{15}}{\sigma \lambda \eta} \eta^2 (\sigma \sqrt{nr} \kappa \log^{2.5}(n) \sqrt{\sigma_{\max}})^2 &= \frac{\kappa n^{15}}{\sigma \cdot \sigma \sqrt{n} \log^{1.5}(n) \kappa^3 n^{20} \sigma_{\max}} \sigma^2 n r^2 \kappa^2 \log^5(n) \sigma_{\max} \\ &= \frac{\sqrt{nr}^2 \log^{3.5}(n)}{n^5} \\ &\ll 1. \end{aligned}$$

H.4 Proof of Eq. (38c)

Note that

$$f(F^{t+1}; \tau^t) = \frac{1}{2} \|X^{t+1} Y^{t+1T} - M^* - E + (\tau^t - \tau^*) Z\|_{\mathbb{F}}^2 + \frac{\lambda}{2} \|X^{t+1}\|_{\mathbb{F}}^2 + \frac{\lambda}{2} \|Y^{t+1}\|_{\mathbb{F}}^2$$

We write X^t, Y^t, τ^t as X, Y, τ if there is no ambiguity. Let $D := XY^\top - M^* - E + (\tau - \tau^*) Z$. Then

$$\begin{aligned} X^{t+1} &= X - \eta(DY + \lambda X) \\ Y^{t+1} &= Y - \eta(D^\top X + \lambda Y). \end{aligned}$$

Furthermore,

$$\begin{aligned} \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2 &= \|DY + \lambda X\|_{\mathbb{F}}^2 + \|D^\top X + \lambda Y\|_{\mathbb{F}}^2 \\ &= \text{tr}(Y^\top D^\top DY) + 2\lambda \text{tr}(X^\top DY) + \lambda^2 \text{tr}(X^\top X) \\ &\quad + \text{tr}(X^\top DD^\top X) + 2\lambda \text{tr}(Y^\top D^\top X) + \lambda^2 \text{tr}(Y^\top Y). \end{aligned}$$

Consider

$$\begin{aligned} &\|X^{t+1} Y^{t+1T} - M^* - E + (\tau^t - \tau^*) Z\|_{\mathbb{F}}^2 + \lambda \|X^{t+1}\|_{\mathbb{F}}^2 + \lambda \|Y^{t+1}\|_{\mathbb{F}}^2 \\ &= \|D - \eta((DY + \lambda X)Y^\top + X(X^\top D + \lambda Y^\top)) + \eta^2(DY + \lambda X)(D^\top X + \lambda Y)^\top\|_{\mathbb{F}}^2 \\ &\quad + \lambda \|X - \eta(DY + \lambda X)\|_{\mathbb{F}}^2 + \lambda \|Y - \eta(D^\top X + \lambda Y)\|_{\mathbb{F}}^2 \\ &= \|D\|_{\mathbb{F}}^2 + \lambda \|X\|_{\mathbb{F}}^2 + \lambda \|Y\|_{\mathbb{F}}^2 + \eta^2 A_0 \\ &\quad - 2\eta(\text{tr}(D^\top D Y Y^\top) + 2\lambda \text{tr}(D^\top X Y^\top) + \text{tr}(D^\top X X^\top D)) \\ &\quad - 2\eta(2\lambda \text{tr}(D Y X^\top) + \lambda^2(\text{tr}(X^\top X) + \text{tr}(Y^\top Y))) \\ &= 2f(F^t; \tau^t) - 2\eta \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2 + \eta^2 A_0 \end{aligned}$$

where A_0 is the term associated with the coefficient η^2 and one can verify that

$$\begin{aligned} \|A_0\|_{\mathbb{F}} &\leq 2\|D\|_{\mathbb{F}} \|DY + \lambda X\|_{\mathbb{F}} \|D^\top X + \lambda Y\|_{\mathbb{F}} + \lambda \|DY + \lambda X\|_{\mathbb{F}}^2 \\ &\quad + \lambda \|D^\top X + \lambda Y\|_{\mathbb{F}}^2 + (\|DY + \lambda X\|_{\mathbb{F}} \|Y^\top\|_{\mathbb{F}} + \|X\|_{\mathbb{F}} \|X^\top D + \lambda Y^\top\|_{\mathbb{F}})^2 \\ &\quad + 2\eta \|DY + \lambda X\|_{\mathbb{F}} \|Y^\top\|_{\mathbb{F}} (\|DY + \lambda X\|_{\mathbb{F}} \|D^\top X + \lambda Y\|_{\mathbb{F}}) \\ &\quad + 2\eta \|X\|_{\mathbb{F}} \|X^\top D + \lambda Y^\top\|_{\mathbb{F}} (\|DY + \lambda X\|_{\mathbb{F}} \|D^\top X + \lambda Y\|_{\mathbb{F}}) \\ &\quad + \eta^2 (\|DY + \lambda X\|_{\mathbb{F}} \|D^\top X + \lambda Y\|_{\mathbb{F}})^2. \end{aligned}$$

In order to show that $f(F^{t+1}; \tau^t) \leq f(F^t; \tau^t) - \frac{\eta}{2} \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2$, it is sufficient to show that $A_0 \eta \leq \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2$.

Note that $\max(\|DY + \lambda X\|_{\mathbb{F}}, \|D^\top X + \lambda Y\|_{\mathbb{F}}) \leq \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}$, recall that

$$\|\nabla f(F^t; \tau^t)\|_{\mathbb{F}} \lesssim \sigma \sqrt{nr} \kappa \log^{2.5}(n) \sqrt{\sigma_{\max}},$$

then we have

$$\begin{aligned}
& \eta \|A_0\|_{\mathbb{F}} \\
& \lesssim \eta \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2 (\|D\|_{\mathbb{F}} + \lambda + \|F\|_{\mathbb{F}}^2) + \eta^2 \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^3 \|F\|_{\mathbb{F}} + \eta^3 \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^4 \\
& \stackrel{(i)}{\lesssim} \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2 \eta (\sigma\sqrt{nr}\kappa \log^{2.5}(n) + \sigma_{\max} r) \\
& \quad + \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2 \eta^2 \sigma\sqrt{nr}\kappa \log^{2.5}(n) \sqrt{\sigma_{\max}} \sqrt{\sigma_{\max} r} \\
& \quad + \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2 \eta^3 (\sigma\sqrt{nr}\kappa \log^{2.5}(n) \sqrt{\sigma_{\max}})^2 \\
& \lesssim \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2 \frac{\sigma\sqrt{nr}\kappa \log^{2.5}(n) + \sigma_{\max} r}{\kappa^3 n^{20} \sigma_{\max}} \\
& \quad + \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2 \frac{\sigma\sqrt{nr}\kappa \log^{2.5}(n) \sigma_{\max}}{\kappa^6 n^{40} \sigma_{\max}^2} \\
& \quad + \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2 \frac{\sigma^2 n r^2 \kappa^2 \log^5(n) \sigma_{\max}}{\kappa^9 n^{60} \sigma_{\max}^3} \\
& \stackrel{(ii)}{\lesssim} \frac{\|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2}{n^{10}}.
\end{aligned}$$

In (i), we use that $\|D\|_{\mathbb{F}} \lesssim \sigma\sqrt{nr}^{0.5} \kappa \log^{2.5}(n)$ (shown in Eq. (41)). In (ii), we use that $\frac{\sigma\sqrt{n}}{\sigma_{\min}} \lesssim \frac{1}{r^2 \kappa^2 \log^5(n)}$. This implies $\eta A_0 \leq \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2$.

Therefore

$$f(F^{t+1}; \tau^t) \leq f(F^t; \tau^t) - \frac{\eta}{2} \|\nabla f(F^t; \tau^t)\|_{\mathbb{F}}^2.$$

By our choice of $\tau^{t+1} := \min_{\tau} f(F^{t+1}; \tau)$, we have $f(F^{t+1}; \tau^{t+1}) \leq f(F^{t+1}; \tau^t)$. This completes the proof.

I Analysis of Non-convex Estimators: Proof of Lemma 2

The proof of Lemma 2 is similar to Lemma 5, but with significant refined analyses on controlling $l_{2,\infty}$ -norm error.

Following the notations in the proof of Lemma 5, by the incoherence condition of M^* , we have the following facts for F^* .

$$\|F^*\|_{2,\infty} = \max\{\|X^*\|_{2,\infty}, \|Y^*\|_{2,\infty}\} \leq \sqrt{\mu r \sigma_{\max}/n}. \quad (54)$$

Before proceeding, inspired by the leave-one-out technique developed in [39, 38, 4], we introduce a set of auxiliary loss functions to facilitate the analysis of the gradient descent algorithm. In particular, for $1 \leq l \leq 2n$, let

$$f^{(l)}(X, Y; \tau) := \frac{1}{2} \|XY^{\top} + \tau Z + P_l(E) - O\|_{\mathbb{F}}^2 + \frac{\lambda}{2} \|X\|_{\mathbb{F}}^2 + \frac{\lambda}{2} \|Y\|_{\mathbb{F}}^2$$

where P_l is defined as the following

$$[P_l(B)]_{ij} = \begin{cases} B_{ij} & i = l \\ 0 & i \neq l \end{cases} \quad 1 \leq l \leq n \quad \text{and} \quad [P_l(B)]_{ij} = \begin{cases} B_{ij} & j = l - n \\ 0 & j \neq l - n \end{cases} \quad n + 1 \leq l \leq 2n.$$

Intuitively, $f^{(l)}$ removes the l -th row (or $(l-n)$ -th column for $l > n$) of the noise matrix E . We then consider the following gradient descent algorithm on the loss function $f^{(l)}$ for $1 \leq l \leq 2n$.

ALGORITHM 2: Leave-one-out Gradient Descent of non-convex optimization

Input: the observation O and Z

1: **Initialization:** $X^{0,(l)} = X^*, Y^{0,(l)} = Y^*, \tau^{0,(l)} = \frac{\langle Z, O - P_l(E) - X^* Y^{*\top} \rangle}{\|Z\|_F^2}$.

2: **Gradient updates:** for $t = 0, 1, \dots, t_* - 1$ do

$$X^{t+1,(l)} = X^{t,(l)} - \eta[(X^{t,(l)}(Y^{t,(l)})^\top + \tau^{t,(l)}Z + P_l(E) - O)Y^{t,(l)} + \lambda X^{t,(l)}]; \quad (55a)$$

$$Y^{t+1,(l)} = Y^{t,(l)} - \eta[(X^{t,(l)}(Y^{t,(l)})^\top + \tau^{t,(l)}Z + P_l(E) - O)^\top X^{t,(l)} + \lambda Y^{t,(l)}] \quad (55b)$$

$$\tau^{t+1,(l)} = \frac{\langle Z, O - P_l(E) - X^{t+1,(l)}Y^{t+1,(l)\top} \rangle}{\|Z\|_F^2}. \quad (55c)$$

where η determines the learning rate.

Let $F^{t,(l)} := \begin{bmatrix} X^{t,(l)} \\ Y^{t,(l)} \end{bmatrix} \in \mathbb{R}^{2n \times r}$. Similar to the definition of H^t in Eq. (36), we define the optimal rotation for aligning $F^{t,(l)}$ and F^* by the following

$$H^{t,(l)} := \arg \min_{A \in \mathcal{O}^{r \times r}} \|F^{t,(l)}A - F^*\|_F.$$

Furthermore, we are also interested in analyzing the relation between $F^{t,(l)}$ and F^t . Therefore, we introduce the following rotation to align $F^{t,(l)}$ and F^t .

$$R^{t,(l)} := \arg \min_{A \in \mathcal{O}^{r \times r}} \|F^{t,(l)}A - F^t H^t\|_F.$$

Next, consider $\tau^t = \frac{\langle Z, O - X^t Y^{tT} \rangle}{\|Z\|_F^2}$. Use $O = M^* + \tau^* Z + E$, we have

$$\tau^t = \frac{\langle Z, M^* - X^t Y^{tT} \rangle}{\|Z\|_F^2} + \frac{\langle Z, E \rangle}{\|Z\|_F^2} + \tau^*.$$

To analyze $\tau^t - \tau^*$, the main difficulty is to bound the error $\frac{\langle Z, M^* - X^t Y^{tT} \rangle}{\|Z\|_F^2}$ (note that $\langle Z, E \rangle / \|Z\|_F^2$ keeps the same for each iteration, hence easier to analyze) To establish the result, we introduce the ‘‘component separation’’ idea by analyzing each one of r components of the error separately. In particular, note that $M^* = X^* Y^{*\top}$. Let x_s^*, y_s^* be the s -column of X^* and Y^* respectively. Then

$$M^* = \sum_{s=1}^r x_s^* y_s^{*\top}.$$

In addition, note that $X^t Y^{tT} = (X^t H^t)(Y^t H^t)^\top$. Let $(X^t H^t)_s, (Y^t H^t)_s$ be the s -th column of $X^t H^t$ and $Y^t H^t$ respectively. Then

$$X^t Y^{tT} = \sum_{s=1}^r (X^t H^t)_s (Y^t H^t)_s^\top$$

We introduce the ‘‘component error’’ Δ_s^t defined as the following

$$\Delta_s^t := \frac{\langle Z, x_s^* y_s^{*\top} - (X^t H^t)_s (Y^t H^t)_s^\top \rangle}{\|Z\|_F^2}. \quad (56)$$

It is easy to see the connection between $\tau^t - \tau^*$ and Δ_s^t through the following.

$$\tau^t - \tau^* - \frac{\langle Z, E \rangle}{\|Z\|_F^2} = \frac{\langle Z, M^* - X^t Y^{tT} \rangle}{\|Z\|_F^2} = \sum_{s=1}^r \Delta_s^t. \quad (57)$$

In order to analyze the bounds for Δ_s^t , we introduce the ‘‘component coefficient’’ b_s . In particular, let u_s^*, v_s^* be the s -column of U^* and V^* respectively. We denote

$$b_s := \frac{\text{tr}(Z^\top Z v_s^* v_s^{*\top}) + \text{tr}(Z Z^\top u_s^* u_s^{*\top})}{\|Z\|_F^2} \quad (58)$$

One shall see the connection between b_s and Assumption 3(a). In fact, Assumption 3(a) implies that, for some constant C ,

$$\sum_{s=1}^r b_s \leq 1 - \frac{C}{\log(n)}$$

due to that $\sum_{s=1}^r v_s^* v_s^{*\top} = V^* V^{*\top}$, $\sum_{s=1}^r u_s^* u_s^{*\top} = U^* U^{*\top}$. The usefulness of b_s will be evident in bounding the component error Δ_s^t in Appendix I.2.

We use the mathematical induction to obtain the desired results. In particular, we aim to prove the following lemma.

Lemma 7. *Suppose Assumptions 1 and 2 and Assumption 3(a) hold. Suppose $O = M^* + \tau^* Z + E$. Suppose $\kappa^4 \mu^2 r^2 \log^2(n) \lesssim n$, $\frac{\sigma}{\sigma_{\min}} \sqrt{n} \lesssim \frac{1}{\kappa^3 r^{4.5} \log^5(n) \mu}$. Let $\lambda = C_\lambda \sigma \sqrt{n} \log^{1.5}(n)$, $\eta = \frac{C_\eta}{\kappa^3 n^{20} \sigma_{\max}}$. Suppose the following induction hypotheses Eq. (59) for $q = t$ hold with (pre-determined) constants $C_\tau, C_F, C_{l,1}, C_{l,2}, C_\infty$. Then, with probability at least $1 - O(n^{-10^6})$, we have the Eq. (59) for $q = t + 1$ holds.*

$$|\Delta_s^q| \leq C_\tau (C_0 + b_s r \log(n)) \left(\frac{\sigma \mu r^{1.5} \kappa \log^{2.5}(n)}{\sqrt{n}} + \frac{\sigma \log^{0.5}(n)}{r \|Z\|_F} \right) \quad (59a)$$

$$|\tau^q - \tau^*| \leq C_\tau \left(\frac{\sigma \mu r^{2.5} \kappa \log^{3.5} n}{\sqrt{n}} + \frac{\sigma \log^{1.5} n}{\|Z\|_F} \right) \quad (59b)$$

$$\|F^q H^q - F^*\|_F \leq C_F \left(\frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_F \quad (59c)$$

$$\max_{1 \leq l \leq 2n} \left\| F^q H^q - F^{q,(l)} R^{q,(l)} \right\|_F \leq C_{l,1} \frac{\sigma \log^{1.5} n}{\sigma_{\min}} \|F^*\|_F \quad (59d)$$

$$\max_{1 \leq l \leq 2n} \left\| (F^{q,(l)} H^{q,(l)} - F^*)_{l,\cdot} \right\|_2 \leq C_{l,2} \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F \quad (59e)$$

$$\|F^q H^q - F^*\|_{2,\infty} \leq C_\infty \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F \quad (59f)$$

$$\|X^{q\top} X^q - Y^{q\top} Y^q\|_F \leq C_B \frac{\sigma}{\kappa n^{15}}. \quad (59g)$$

Furthermore, let T_t be the tangent space of $X^t Y^{t\top}$. If Eq. (59) holds for $q = t$, the following inequality holds with probability $1 - O(n^{-10^6})$ for (pre-determined) constants $C_{T,1}, C_{T,2}, C_{T,3}$.

$$\left\| P_{T_t^\perp}(M^*) \right\|_\infty \leq C_{T,1} \frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^7(n)}{\sigma_{\min}} \quad (60a)$$

$$\left\| P_{T_t^\perp}(E) - P_{T^{*\perp}}(E) \right\|_\infty \leq C_{T,2} \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^{2.5} \log^4(n)}{\sigma_{\min}} \quad (60b)$$

$$\left\| P_{T_t^\perp}(Z) - P_{T^{*\perp}}(Z) \right\|_F \leq C_{T,3} \frac{r \kappa \sigma \sqrt{n} \log^2(n)}{\sigma_{\min}} \|Z\| \quad (60c)$$

$$f(X^{q+1}, Y^{q+1}; \tau^{q+1}) \leq f(X^q, Y^q; \tau^q) - \frac{\eta}{2} \|\nabla f(X^q, Y^q; \tau^q)\|_F^2. \quad (60d)$$

Note that based on Lemma 7, we can prove Lemma 2 in a straightforward way.

Proof of Lemma 2. Note that $\|E\| \leq C \sigma \sqrt{n}$ with probability $1 - O(n^{-10^6})$ for some constant C . Note that when $q = 0$, $X^0 = X^*$, $X^{0,(l)} = X^*$ and $Y^0 = Y^*$, $Y^{0,(l)} = Y^*$ easily satisfy Eqs. (59c)

to (59g) along with $\Delta_s^0 = 0$ satisfying Eq. (59a). For τ^0 , noting that $\tau^0 - \tau^* = \langle Z, E \rangle / \|Z\|_{\mathbb{F}}^2 \lesssim \sigma \sqrt{\log(n)} / \|Z\|_{\mathbb{F}}$ with probability $1 - O(n^{-10^6})$. This verifies the induction hypotheses for $q = 0$.

Take $t^* = \kappa^3 n^{10000}$, then by Lemma 7 and the union bound, with probability at least $1 - O(n^{-10^4})$, we have Eq. (59) and Eq. (60) hold for $q = 0, 1, \dots, t^* - 1$. By the same analysis in the proof of Lemma 5, we obtain the desired results of Lemma 2.¹⁰ \square

The remainder is to finish the proof for Lemma 7. Under the hypothesis for $q = t$, we prove Eq. (59) and Eq. (60) for $q = t + 1$ in the following subsections. Note that under the assumptions of Lemma 7, we can directly invoke Lemma 6, and hence Eqs. (59c), (59g) and (60d) directly hold.

I.1 Direct implication of Eq. (59) for $q = t$

Under the hypothesis $q = t$, we first present a few direct implications that are helpful in the proof for $q = t + 1$. For large enough n , we claim that the following results hold for F^t .

$$\begin{aligned} \|F^t\|_{\mathbb{F}} &\leq 2 \|F^*\|_{\mathbb{F}} \lesssim \sqrt{\sigma_{\max} r} \\ \|F^t\| &\leq 2 \|F^*\| \lesssim \sqrt{\sigma_{\max}} \\ \|F^t\|_{2,\infty} &\leq 2 \|F^*\|_{2,\infty} \lesssim \sqrt{\frac{\sigma_{\max} \mu r}{n}}. \end{aligned}$$

Similarly, for $l \in [2n]$, for large enough n , we also have

$$\begin{aligned} \|F^{t,(l)}\|_{\mathbb{F}} &\leq 2 \|F^*\|_{\mathbb{F}} \lesssim \sqrt{\sigma_{\max} r} \\ \|F^{t,(l)}\| &\leq 2 \|F^*\| \lesssim \sqrt{\sigma_{\max}} \\ \|F^{t,(l)}\|_{2,\infty} &\leq 2 \|F^*\|_{2,\infty} \lesssim \sqrt{\frac{\sigma_{\max} \mu r}{n}}. \end{aligned}$$

In addition, we have

$$\|\nabla f(X^t, Y^t; \tau^t)\|_{\mathbb{F}} \lesssim \sigma r \kappa \log^{2.5}(n) \sqrt{n} \sqrt{\sigma_{\max}} \quad (61)$$

The proof for $\|F^t\|_{\mathbb{F}}, \|F^t\|, \|\nabla f(X^t, Y^t; \tau^t)\|_{\mathbb{F}}$ have been shown in Appendix H.1. We show the bound for $\|F^t H^t\|_{2,\infty}$ below.

$$\begin{aligned} \|F^t H^t\|_{2,\infty} &\leq \|F^t H^t - F^*\|_{2,\infty} + \|F^*\|_{2,\infty} \\ &\stackrel{(i)}{\leq} C_{\infty} \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_{\mathbb{F}} + \|F^*\|_{2,\infty} \\ &\stackrel{(ii)}{\leq} 2C_{\infty} \frac{\sigma \sqrt{n} \mu r^2 \kappa \log^{3.5}(n)}{\sigma_{\min}} \cdot \sqrt{\frac{\sigma_{\max} r}{n}} + \|F^*\|_{2,\infty} \\ &\stackrel{(iii)}{\leq} \frac{\sqrt{\sigma_{\max}}}{2\sqrt{\kappa} \log(n)} \sqrt{\frac{r}{n}} + \|F^*\|_{2,\infty} \\ &\stackrel{(iv)}{\leq} 2 \|F^*\|_{2,\infty} \end{aligned}$$

Here, (i) is due to Eq. (59f), (ii) is due to $\|F^*\|_{\mathbb{F}} \leq \sqrt{2\sigma_{\max} r}$, (iii) is due to $\frac{\sigma}{\sigma_{\min}} \sqrt{n} \lesssim \frac{1}{\kappa^3 r^{4.5} \log^5(n) \mu}$ and holds with large enough n , and (iv) is due to $\|F^*\|_{2,\infty} \geq \sqrt{\frac{\sigma_{\min} r}{n}}$. This also implies $\|F^t\|_{2,\infty} = \|F^t H^t H^{tT}\|_{2,\infty} \leq \|F^t H^t\|_{2,\infty} \|H^t\| \leq 2 \|F^*\|_{2,\infty}$.

The similar bounds can be obtained for $F^{t,(l)}$ based on $\|F^t H^t - F^{t,(l)} R^{t,(l)}\|_{\mathbb{F}} \lesssim \frac{\sigma \log^{1.5} n}{\sigma_{\min}} \|F^*\|_{\mathbb{F}}$ (Eq. (59d)) and the triangle inequality. We omit the proof for brevity.

¹⁰Here, we consider all the bounds hold with probability $1 - O(n^{-10^4})$. The generalization to $1 - O(n^{-C})$, for some constant C , is straightforward.

I.2 Proof of Eqs.(59a) and (59b)

Recall that $\Delta_s^t = \frac{\langle Z, x_s^* y_s^{*\top} - (X^t H^t)_s (Y^t H^t)_s^\top \rangle}{\|Z\|_F^2}$. Then we have

$$\begin{aligned} \Delta_s^{t+1} &= \Delta_s^t + \frac{\langle Z, (X^t H^t)_s (Y^t H^t)_s^\top - (X^{t+1} H^{t+1})_s (Y^{t+1} H^{t+1})_s^\top \rangle}{\|Z\|_F^2} \\ &= \Delta_s^t + \frac{\langle Z, (X^t H^t)_s (Y^t H^t)_s^\top - (X^{t+1} H^t)_s (Y^{t+1} H^t)_s^\top \rangle}{\|Z\|_F^2} \\ &\quad + \underbrace{\frac{\langle Z, (X^{t+1} H^t)_s (Y^{t+1} H^t)_s^\top - (X^{t+1} H^{t+1})_s (Y^{t+1} H^{t+1})_s^\top \rangle}{\|Z\|_F^2}}_{A_0}. \end{aligned} \quad (62)$$

To control A_0 , we have the following claim to show that A_0 is negligible.

Claim 5.

$$\begin{aligned} \|H^t - H^{t+1}\|_F &\lesssim \eta \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} (\sigma \sqrt{n} \mu r^3 \kappa^2 \log^{3.5}(n)). \\ |A_0| &\lesssim \eta \sigma^2 \log^6(n) \kappa^3 \mu^2 r^4. \end{aligned}$$

Next, by the gradient updating rule,

$$X^{t+1} = X^t - \eta ((X^t Y^{tT} - M^* + (\tau^t - \tau^*)Z - E)Y^t + \lambda X^t).$$

Let A_s be the s -column of A for any matrix A . Suppose $x_s^t = (X^t H^t)_s, y_s^t = (Y^t H^t)_s$. Then, we have

$$\begin{aligned} (X^{t+1} H^t)_s &= (X^t H^t)_s - \eta ((X^t Y^{tT} - M^* + (\tau^t - \tau^*)Z - E)(Y^t H^t)_s + \lambda (X^t H^t)_s) \\ &= x_s^t - \eta (X^t Y^{tT} - M^* + (\tau^t - \tau^*)Z - E)y_s^t - \eta \lambda x_s^t. \end{aligned} \quad (63)$$

Similarly,

$$\begin{aligned} (Y^{t+1} H^t)_s &= (Y^t H^t)_s - \eta (X^t Y^{tT} - M^* + (\tau^t - \tau^*)Z - E)^\top (X^t H^t)_s + \lambda (Y^t H^t)_s \\ &= y_s^t - \eta (X^t Y^{tT} - M^* + (\tau^t - \tau^*)Z - E)^\top x_s^t - \eta \lambda y_s^t. \end{aligned} \quad (64)$$

To simplify the notation, we write $(XH)^t, (YH)^t, \tau^t, x_s^t, y_s^t$ by X, Y, τ, x_s, y_s if there is no ambiguity. Continuing the analysis for Δ_s^{t+1} in Eq. (62), we have

$$\begin{aligned} \Delta_s^{t+1} &= \Delta_s^t + A_0 + \frac{\langle Z, x_s y_s^\top - (X^{t+1} H^t)_s (Y^{t+1} H^t)_s^\top \rangle}{\|Z\|_F^2} \\ &= \Delta_s^t + A_0 + \frac{\langle Z, (x_s - (X^{t+1} H^t)_s) y_s^\top \rangle}{\|Z\|_F^2} + \frac{\langle Z, x_s (y_s - (Y^{t+1} H^t)_s)^\top \rangle}{\|Z\|_F^2} \\ &\quad - \frac{\langle Z, (x_s - (X^{t+1} H^t)_s) (y_s - (Y^{t+1} H^t)_s)^\top \rangle}{\|Z\|_F^2}. \end{aligned}$$

By expanding $x_s - (X^{t+1} H^t)_s$ and $y_s - (Y^{t+1} H^t)_s$ using Eq. (63) and Eq. (64), we then have

$$\begin{aligned} \Delta_s^{t+1} &= \Delta_s^t + A_0 + \frac{\eta}{\|Z\|_F^2} \underbrace{\langle Z, (XY^\top - M^* + (\tau - \tau^*)Z - E) y_s y_s^\top + \lambda x_s y_s^\top \rangle}_{A_1} \\ &\quad + \frac{\eta}{\|Z\|_F^2} \underbrace{\langle Z, x_s x_s^\top (XY^\top - M^* + (\tau - \tau^*)Z - E) + \lambda x_s y_s^\top \rangle}_{A_2} + A_3 \end{aligned} \quad (65)$$

where A_3 includes the term with coefficient η^2 :

$$A_3 := \frac{-\eta^2}{\|Z\|_F^2} \langle Z, D_1 D_2^\top \rangle$$

with

$$\begin{aligned} D_1 &:= (XY^\top - M^* + (\tau - \tau^*)Z - E)y_s + \lambda x_s \\ D_2 &:= (XY^\top - M^* + (\tau - \tau^*)Z - E)x_s + \lambda y_s. \end{aligned}$$

Note that η has been chosen small enough, one can verify that

$$\begin{aligned} |A_3| &\leq \frac{\eta^2}{\|Z\|_F^2} \|Z\|_F \|D_1\|_F \|D_2\|_F \\ &\leq \eta^2 \|D_1\|_F \|D_2\|_F \\ &\stackrel{(i)}{\lesssim} \eta \frac{1}{n^{20} \sigma_{\max}} (\sigma \sqrt{n} \mu r^3 \kappa \log^{3.5}(n) \sqrt{\sigma_{\max}})^2 \\ &= \eta \sigma^2 \frac{\mu^2 r^6 \kappa^2 \log^7(n)}{n^{19}} \\ &\stackrel{(ii)}{=} \eta \frac{\sigma^2}{n^{15}}. \end{aligned} \tag{66}$$

In (i), we use the bound $\|D_1\|_F + \|D_2\|_F \lesssim \sigma \sqrt{n} \mu r^3 \kappa \log^{3.5}(n) \sqrt{\sigma_{\max}}$ (the proof is the same as showing $\|\nabla f(X, Y; \tau)\|_F \lesssim \sigma \sqrt{n} \mu r^3 \kappa \log^{3.5}(n) \sqrt{\sigma_{\max}}$ in ?? and hence omitted here). In (ii), we use that $\kappa^4 \mu^2 r^2 \log^2(n) \lesssim n$.

Next, we analyze A_1 (A_2 is similar to A_1 by symmetry). For A_1 , to facilitate the analysis, let

$$\begin{aligned} B_0 &:= \langle Z, (XY^\top - M^* + (\tau - \tau^*)Z) (y_s y_s^\top - y_s^* y_s^{*\top}) \rangle \\ B_1 &:= \langle Z, \lambda x_s y_s^\top \rangle. \end{aligned}$$

Then, we have

$$\begin{aligned} A_1 &= \langle Z, (XY^\top - M^* + (\tau - \tau^*)Z) y_s^* y_s^{*\top} \rangle + B_0 + B_1 + \underbrace{\langle Z, -E y_s y_s^\top \rangle}_{B_2} \\ &= \langle Z, (XY^\top - M^*) y_s^* y_s^{*\top} \rangle + (\tau - \tau^*) \langle Z, Z y_s^* y_s^{*\top} \rangle + B_0 + B_1 + B_2. \end{aligned}$$

Recall that $y_s^* = \sqrt{\sigma_s^*} v_s^*$ where σ_s^* is the s -th largest singular value of M^* and v_s^* is the s -th corresponding right singular vector. Hence, for any $k \in [r]$, we have

$$y_k^{*\top} y_s^* = \sqrt{\sigma_s^*} \sqrt{\sigma_k^*} v_k^{*\top} v_s^* \stackrel{(i)}{=} \sigma_s^* \mathbb{1}\{k = s\}$$

where (i) is due to orthogonality between different singular vectors. This will help us simplify the analysis for A_1 . In particular,

$$\begin{aligned} A_1 &\stackrel{(i)}{=} \langle Z, (X - X^*) Y^{*\top} y_s^* y_s^{*\top} + X (Y - Y^*)^\top y_s^* y_s^{*\top} \rangle + (\tau - \tau^*) \sigma_s^* \text{tr}(Z^\top Z v_s^* v_s^{*\top}) \\ &\quad + B_2 + B_1 + B_0 \\ &\stackrel{(ii)}{=} \sigma_s^* \langle Z, (x_s - x_s^*) y_s^{*\top} \rangle + (\tau - \tau^*) \sigma_s^* \text{tr}(Z^\top Z v_s^* v_s^{*\top}) + \underbrace{\langle Z, X (Y - Y^*)^\top y_s^* y_s^{*\top} \rangle}_{B_3} \\ &\quad + B_2 + B_1 + B_0. \end{aligned}$$

where in (i) we use that $XY^\top - M^* = (X - X^*) Y^{*\top} + X (Y - Y^*)^\top$ and $y_s^* y_s^{*\top} = \sigma_s^* v_s^* v_s^{*\top}$, in (ii) we use the fact that y_s^* is orthogonal to y_k^* , $k \neq s$, which implies

$$(X - X^*) Y^{*\top} y_s^* = \sum_{k=1}^r (X - X^*) y_k^* y_k^{*\top} y_s^* = \sigma_s^* (x_s - x_s^*) y_s^{*\top}.$$

In order to bound B_0, B_1, B_2, B_3 , we have the following claim.

Claim 6. With probability $1 - O(n^{-10^7})$,

$$\begin{aligned}\frac{1}{\|Z\|_F^2}|B_0| &\lesssim \frac{\sigma\mu r^{1.5}\kappa \log^2(n)}{\sqrt{n}}\sigma_{\min} + \frac{\sigma \log^{0.5}(n)}{r\|Z\|_F}\sigma_{\min} \\ \frac{1}{\|Z\|_F^2}|B_1| &\lesssim \frac{\sigma\mu r^{1.5}\kappa \log^2(n)}{\sqrt{n}}\sigma_{\min} \\ \frac{1}{\|Z\|_F^2}|B_2| &\lesssim \frac{\sigma\mu r^{1.5}\kappa \log^2(n)}{\sqrt{n}}\sigma_{\min} \\ \frac{1}{\|Z\|_F^2}|B_3| &\lesssim \frac{\sigma\mu r^{1.5}\kappa \log^{2.5}(n)}{\sqrt{n}}\sigma_s^*.\end{aligned}$$

Let $\Delta = |B_0| + |B_1| + |B_2| + |B_3|$. Then we have

$$A_1 \leq \sigma_s^* \langle Z, (x_s - x_s^*)y_s^{*\top} \rangle + (\tau - \tau^*)\sigma_s^* \text{tr}(Z^\top Z v_s^* v_s^{*\top}) + \Delta.$$

Here,

$$\frac{1}{\|Z\|_F^2}\Delta = \frac{1}{\|Z\|_F^2}(|B_0| + |B_1| + |B_2| + |B_3|) \stackrel{(i)}{\lesssim} \frac{\sigma\mu r^{1.5}\kappa \log^{2.5}(n)}{\sqrt{n}}\sigma_s^* + \frac{\sigma \log^{0.5}(n)}{r\|Z\|_F}\sigma_{\min} \quad (67)$$

where (i) uses Claim 6 and $\sigma_{\min} = \sigma_r^* \leq \sigma_s^*$ for $s \in [r]$.

By symmetry, we can obtain the similar results for A_2 and in particular, we have

$$\begin{aligned}A_1 + A_2 &\leq \sigma_s^* \langle Z, (x_s - x_s^*)y_s^{*\top} \rangle + \sigma_s^* \langle Z, x_s^*(y_s - y_s^*)^\top \rangle \\ &\quad + (\tau - \tau^*)\sigma_s^* \text{tr}(Z^\top Z v_s^* v_s^{*\top}) + (\tau - \tau^*)\sigma_s^* \text{tr}(Z Z^\top u_s^* u_s^{*\top}) \\ &\quad + 2\Delta \\ &\stackrel{(i)}{\leq} \sigma_s^* \langle Z, x_s y_s^\top - x_s^* y_s^{*\top} \rangle - \sigma_s^* \langle Z, (x_s - x_s^*)(y_s - y_s^*)^\top \rangle + |\tau - \tau^*|\sigma_s^* b_s \|Z\|_F^2 + 2\Delta \\ &\stackrel{(ii)}{\leq} -\Delta_s^t \sigma_s^* \|Z\|_F^2 + \sigma_s^* \|Z\|_F^2 \|F - F^*\|_{2,\infty}^2 + |\tau - \tau^*|\sigma_s^* b_s \|Z\|_F^2 + 2\Delta\end{aligned}$$

where in (i) we use that $x_s y_s^\top - x_s^* y_s^{*\top} = (x_s - x_s^*)y_s^{*\top} + x_s^*(y_s - y_s^*)^\top + (x_s - x_s^*)(y_s - y_s^*)^\top$ and $b_s \|Z\|_F^2 = \text{tr}(Z^\top Z v_s^* v_s^{*\top}) + \text{tr}(Z Z^\top u_s^* u_s^{*\top})$, in (ii) we use that $\Delta_s^t \|Z\|_F^2 = \langle Z, x_s^* y_s^{*\top} - x_s y_s^\top \rangle$ and $|\langle Z, AB^\top \rangle| \leq \|Z\|_F^2 \|AB^\top\|_{\infty} \leq \|Z\|_F^2 \|A\|_{2,\infty} \|B\|_{2,\infty}$ for binary matrix Z .

Next we can bound Δ_s^{t+1} by using the bounds all above for A_0, A_1, A_2, A_3 in Eq. (65).

$$\begin{aligned}|\Delta_s^{t+1}| &\leq \left| \Delta_s^t + \eta \frac{1}{\|Z\|_F^2} (A_1 + A_2) + A_0 + A_3 \right| \\ &\leq |\Delta_s^t| (1 - \eta\sigma_s^*) + \eta \left(\sigma_s^* \|F - F^*\|_{2,\infty}^2 + |\tau - \tau^*|\sigma_s^* b_s + \frac{1}{\|Z\|_F^2} 2\Delta \right) + |A_0| + |A_3| \\ &\stackrel{(i)}{\leq} C_\tau (C_0 + b_s r \log(n)) \left(\frac{\sigma\mu r^{1.5}\kappa \log^{2.5}(n)}{\sqrt{n}} + \frac{\sigma \log^{0.5}(n)}{r\|Z\|_F} \right) (1 - \eta\sigma_s^*) \\ &\quad + \eta C_\tau \left(\frac{\sigma\mu r^{2.5}\kappa \log^{3.5} n}{\sqrt{n}} + \frac{\sigma \log^{1.5} n}{\|Z\|_F} \right) \sigma_s^* b_s \\ &\quad + \eta\sigma_s^* \|F - F^*\|_{2,\infty}^2 + \eta \frac{1}{\|Z\|_F^2} 2\Delta + |A_0| + |A_3| \quad (68)\end{aligned}$$

Here (i) is by the induction hypothesis Eq. (59a) and Eq. (59b). Let

$$D := \left(\frac{\sigma\mu r^{1.5}\kappa \log^{2.5}(n)}{\sqrt{n}} + \frac{\sigma \log^{0.5}(n)}{r\|Z\|_F} \right).$$

Then $D \log(n)r = \left(\frac{\sigma \mu r^{2.5} \kappa \log^{3.5} n}{\sqrt{n}} + \frac{\sigma \log^{1.5} n}{\|Z\|_F} \right)$. To simplify Eq. (68), by the direct algebra, we have

$$\begin{aligned} & C_\tau (C_0 + b_s r \log(n)) D(1 - \eta \sigma_s^*) + \eta C_\tau D \log(n) r \sigma_s^* b_s \\ &= C_\tau (C_0 + b_s r \log(n)) D - C_\tau (C_0 + b_s r \log(n)) D \eta \sigma_s^* + \eta C_\tau D \log(n) r \sigma_s^* b_s \\ &= C_\tau (C_0 + b_s r \log(n)) D - C_\tau C_0 D \eta \sigma_s^* - C_\tau b_s r \log(n) D \eta \sigma_s^* + \eta C_\tau D \log(n) r \sigma_s^* b_s \\ &= C_\tau (C_0 + b_s r \log(n)) D - C_\tau C_0 D \eta \sigma_s^*. \end{aligned}$$

To control $\eta \sigma_s^* \|F - F^*\|_{2,\infty}^2 + \eta \frac{1}{\|Z\|_F^2} 2\Delta + |A_0| + |A_3|$, recall that $\|F - F^*\|_{2,\infty} \lesssim \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n) \sqrt{\sigma_{\max} r}}{\sigma_{\min}}$ by Eq. (59f), $\frac{1}{\|Z\|_F^2} \Delta \lesssim \frac{\sigma \mu r^{1.5} \kappa \log^{2.5}(n)}{\sqrt{n}} \sigma_s^* + \frac{\sigma \log^{0.5}(n)}{r \|Z\|_F} \sigma_{\min}$ by Eq. (67), $|A_0| \lesssim \eta \sigma^2 \log^6(n) \kappa^3 \mu^2 r^4$ by Claim 5, and $|A_3| \lesssim \eta \frac{\sigma^2}{n^{15}}$. Then, we have, for some constant C ,

$$\begin{aligned} & \eta \sigma_s^* \|F - F^*\|_{2,\infty}^2 + \eta \frac{1}{\|Z\|_F^2} 2\Delta + |A_0| + |A_3| \\ & \lesssim \eta \left(\sigma_s^* \frac{\sigma^2 \mu^2 r^6 \log^7(n) \kappa^3}{\sigma_{\min}} + \frac{\sigma \mu r^{1.5} \kappa \log^{2.5}(n)}{\sqrt{n}} \sigma_s^* \right) \\ & \quad + \left(\frac{\sigma \log^{0.5}(n)}{r \|Z\|_F} \sigma_{\min} + \sigma^2 \log^6(n) \kappa^3 \mu^2 r^4 + \frac{\sigma^2}{n^{15}} \right) \\ & \stackrel{(i)}{\lesssim} \eta \left(\frac{\sigma \mu r^{1.5} \kappa \log^{2.5}(n)}{\sqrt{n}} \sigma_s^* + \frac{\sigma \log^{0.5}(n)}{r \|Z\|_F} \sigma_{\min} \right) \\ & \stackrel{(ii)}{\leq} C \eta \sigma_s^* \left(\frac{\sigma \mu r^{1.5} \kappa \log^{2.5}(n)}{\sqrt{n}} + \frac{\sigma \log^{0.5}(n)}{r \|Z\|_F} \right) \end{aligned}$$

where (i) is due to $\frac{\sigma}{\sigma_{\min}} \sqrt{n} \lesssim \frac{1}{\kappa^3 r^{4.5} \log^5(n) \mu}$ (hence some terms are negligible), and (ii) is due to $\sigma_{\min} \leq \sigma_s^*$.

This leads to the bound for $|\Delta_s^{t+1}|$.

$$\begin{aligned} |\Delta_s^{t+1}| & \leq C_\tau (C_0 + b_s r \log(n)) D - C_\tau C_0 D \eta \sigma_s^* \\ & \quad + C \eta \sigma_s^* \left(\frac{\sigma \mu r^{1.5} \kappa \log^{2.5}(n)}{\sqrt{n}} + \frac{\sigma \log^{0.5}(n)}{r \|Z\|_F} \right) \\ & \leq C_\tau (C_0 + b_s r \log(n)) D - C_\tau C_0 D \eta \sigma_s^* + C \eta \sigma_s^* D \\ & \stackrel{(i)}{\leq} C_\tau (C_0 + b_s r \log(n)) D. \end{aligned}$$

Here, (i) is obtained by choosing the constant C_τ large enough. This then completes the proof for the bound on Δ_s^{t+1} .

The bound on $|\tau^{t+1} - \tau^*|$ is a direct consequence of the bound on Δ_s^{t+1} and Assumption 3(a). Recall that $\tau^{t+1} - \tau^* - \frac{\langle Z, E \rangle}{\|Z\|_F^2} = \sum_{s=1}^r \Delta_s^{t+1}$ by Eq. (57). Then, we have

$$\begin{aligned} |\tau^{t+1} - \tau^*| & \leq \left| \sum_{s=1}^r \Delta_s^{t+1} \right| + \left| \frac{\langle Z, E \rangle}{\|Z\|_F^2} \right| \\ & \stackrel{(i)}{\leq} \left| \sum_{s=1}^r \Delta_s^{t+1} \right| + C \frac{\sigma \sqrt{\log(n)}}{\|Z\|_F} \\ & \leq \sum_{s=1}^r C_\tau (C_0 + b_s r \log(n)) \left(\frac{\sigma \mu r^{1.5} \kappa \log^{2.5}(n)}{\sqrt{n}} + \frac{\sigma \log^{0.5}(n)}{r \|Z\|_F} \right) + C \frac{\sigma \sqrt{\log(n)}}{\|Z\|_F} \\ & \stackrel{(ii)}{\leq} C_\tau \left(r C_0 + \left(1 - \frac{C_{r_1}}{\log(n)} \right) r \log(n) \right) \left(\frac{\sigma \mu r^{1.5} \kappa \log^{2.5}(n)}{\sqrt{n}} + \frac{\sigma \log^{0.5}(n)}{r \|Z\|_F} \right) \\ & \quad + C \frac{\sigma \sqrt{\log(n)}}{\|Z\|_F} \end{aligned}$$

where (i) is due to with probability $1 - O(n^{-10^7})$, $\left| \frac{\langle Z, E \rangle}{\|Z\|_F^2} \right| \leq C \frac{\sigma \sqrt{\log(n)}}{\|Z\|_F}$ and (ii) is due to $\sum_{s=1}^r b_s \leq 1 - C_{r_1} / \log(n)$ by Assumption 3(a). Note that $rC_0 + \left(1 - \frac{C_{r_1}}{\log(n)}\right) r \log(n) = rC_0 + r \log(n) - rC_{r_1}$ and choose $C_0 \leq C_{r_1}/2$, we have

$$\begin{aligned} |\tau^{t+1} - \tau^*| &\leq C_\tau (rC_0 + r \log(n) - rC_{r_1}) \left(\frac{\sigma \mu r^{1.5} \kappa \log^{2.5}(n)}{\sqrt{n}} + \frac{\sigma \log^{0.5}(n)}{r \|Z\|_F} \right) \\ &\quad + C \frac{\sigma \sqrt{\log(n)}}{\|Z\|_F} \\ &\stackrel{(i)}{\leq} C_\tau r \log(n) \left(\frac{\sigma \mu r^{1.5} \kappa \log^{2.5}(n)}{\sqrt{n}} + \frac{\sigma \log^{0.5}(n)}{r \|Z\|_F} \right) \\ &\leq C_\tau \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sqrt{n}} + C_\tau \frac{\sigma \log^{1.5} n}{\|Z\|_F} \end{aligned}$$

where (i) is obtained by choosing $C_0 \leq C_{r_1}/2$ and C_τ large enough. This completes the proof for Eq. (59b).

Proof of Claim 5. Recall that we want to show

$$\|H^t - H^{t+1}\|_F \lesssim \eta \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} (\sigma \sqrt{n} \mu r^3 \kappa^2 \log^{3.5}(n))$$

where $H^t = \arg \min_{A \in \mathcal{O}^{r \times r}} \|F^t A - F^*\|_F$, $H^{t+1} = \arg \min_{A \in \mathcal{O}^{r \times r}} \|F^{t+1} A - F^*\|_F$ with $F^{t+1} = [X^{t+1}; Y^{t+1}]$ and

$$\begin{aligned} X^{t+1} &= X^t - \eta((X^t Y^{tT} - M^* - E + (\tau^t - \tau^*)Z)Y^t + \lambda X^t) \\ Y^{t+1} &= Y^t - \eta((X^t Y^{tT} - M^* - E + (\tau^t - \tau^*)Z)^\top X^t + \lambda Y^t). \end{aligned}$$

The proof is similar to section D.4 in [4]. We intend to invoke Lemma 12 to control $\|H^t - H^{t+1}\|_F$. However, $\|F^t - F^{t+1}\|_F$ is too large to provide the desired bounds. We tackle this by constructing an auxiliary point \tilde{F}^{t+1} such that (i) H^t is also the optimal rotation to align \tilde{F}^{t+1} and F^* and (ii) $\|\tilde{F}^{t+1} - F^{t+1}\|_F \ll \|F^t - F^{t+1}\|_F$.

In particular, we construct the auxiliary point $\tilde{F}^{t+1} := \begin{bmatrix} \tilde{X}^{t+1} \\ \tilde{Y}^{t+1} \end{bmatrix}$ by substituting some of X^t, Y^t in F^{t+1} with $X^* H^{t\top}, Y^* H^{t\top}$ in the following

$$\begin{aligned} \tilde{X}^{t+1} &= X^t - \eta((X^t Y^{tT} - M^* - E + (\tau^t - \tau^*)Z)Y^* H^{t\top} + \lambda X^* H^{t\top}) \\ \tilde{Y}^{t+1} &= Y^t - \eta((X^t Y^{tT} - M^* - E + (\tau^t - \tau^*)Z)^\top X^* H^{t\top} + \lambda Y^* H^{t\top}). \end{aligned}$$

We then claim that H^t is also the optimal rotation to align \tilde{F}^{t+1} and F^* :

$$\left\| \tilde{F}^{t+1} H^t - F^* \right\|_F \leq \min_{A \in \mathcal{O}^{r \times r}} \left\| \tilde{F}^{t+1} A - F^* \right\|_F.$$

To verify this, by the property of Orthogonal Procrustes problem (Lemma 11), it is sufficient to show that $F^{*\top} \tilde{F}^{t+1} H^t$ is symmetric and positive semi-definite. The symmetry can be verified by direct algebra:

$$\begin{aligned} F^{*\top} \tilde{F}^{t+1} H^t &= X^{*\top} \tilde{X}^{t+1} H^t + Y^{*\top} \tilde{Y}^{t+1} H^t \\ &= X^{*\top} X^t H^t - \eta X^{*\top} (X^t Y^{tT} - M^* - E + (\tau^t - \tau^*)Z) Y^* - \eta \lambda X^{*\top} X^* \\ &\quad + Y^{*\top} Y^t H^t - \eta Y^{*\top} (X^t Y^{tT} - M^* - E + (\tau^t - \tau^*)Z)^\top X^* - \eta \lambda Y^{*\top} Y^* \\ &= F^{*\top} F^t H^t - \eta \lambda X^{*\top} X^* - \eta \lambda Y^{*\top} Y^* - \eta B_0 \end{aligned}$$

where

$$\begin{aligned} B_0 &:= (X^{*\top} (X^t Y^{tT} - M^* - E + (\tau^t - \tau^*)Z) Y^* \\ &\quad + Y^{*\top} (X^t Y^{tT} - M^* - E + (\tau^t - \tau^*)Z)^\top X^*). \end{aligned}$$

It is easy to see that $X^{*\top} X^*, Y^{*\top} Y^*$ are symmetric, together with $B_0 = B_0^\top$. Furthermore, note H^t is the optimal rotation for aligning F^t and F^* by definition. Then, by Lemma 11, $F^{*\top} F^t H^t$ is also symmetric. This leads to the result that $F^{*\top} \tilde{F}^{t+1} H^t$ is symmetric.

To verify the positive semi-definiteness, it is easy to check that

$$\begin{aligned} \left\| F^{*\top} \tilde{F}^{t+1} H^t - F^{*\top} F^* \right\| &\leq \|F^{*\top}\| \left\| \tilde{F}^{t+1} H^t - F^* \right\| \\ &\stackrel{(i)}{\leq} \sqrt{\sigma_{\max}} \|F^t H^t - F^* + \eta\delta\| \\ &\stackrel{(ii)}{\leq} \sigma_{\min}. \end{aligned}$$

Here, δ in (i) represents the term with coefficient η ; and (ii) is due to Eq. (59c) and that we choose η being sufficiently small.

Also note that $\lambda_{\min}(F^{*\top} F^*) = 2\sigma_{\min}$. By Weyl's inequality, this implies that the positive semi-definiteness of $F^{*\top} \tilde{F}^{t+1} H^t$.

$$\lambda_{\min}(F^{*\top} \tilde{F}^{t+1} H^t) \geq 2\sigma_{\min} - \left\| F^{*\top} \tilde{F}^{t+1} H^t - F^{*\top} F^* \right\| > 0.$$

Then, we can invoke Lemma 12 to bound $\|H^t - H^{t+1}\|$. Let $F_0 = F^*, F_1 = \tilde{F}^{t+1}, F_2 = F^{t+1}$. Recall that $\lambda \lesssim \sigma\sqrt{n} \log^{1.5}(n)$, $\|E\| \lesssim \sigma\sqrt{n}$, $|\tau^t - \tau^*| \|Z\| \lesssim \sigma\sqrt{n}\mu r^3 \kappa^2 \log^{3.5}(n)$ and

$$\|X^t Y^{tT} - M^*\| \lesssim \|F^* - F^t H^t\|_{\text{F}} \|F^*\| \lesssim \frac{\sigma\sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \sigma_{\max} \lesssim \sigma\sqrt{n} \log^{2.5}(n)\kappa.$$

Then

$$\begin{aligned} \|F_2 - F_1\|_{\text{F}} &\lesssim \eta \|F^* - F^t H^t\|_{\text{F}} (\lambda + \|X^t Y^{tT} - M^* - E + (\tau^t - \tau^*)Z\|) \\ &\lesssim \eta \|F^* - F^t H^t\|_{\text{F}} (\lambda + \|X^t Y^{tT} - M^*\| + \|E\| + |\tau^t - \tau^*| \|Z\|) \\ &\lesssim \eta \frac{\sigma\sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \|F^*\|_{\text{F}} (\sigma\sqrt{n}\mu r^{2.5} \kappa \log^{3.5}(n)). \end{aligned}$$

Then by Lemma 12, we have

$$\begin{aligned} \|H^t - H^{t+1}\|_{\text{F}} &\leq \frac{1}{\sigma_{\min}} \|F_2 - F_1\|_{\text{F}} \|F^*\| \\ &\lesssim \eta \frac{\sigma\sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} (\sigma\sqrt{n}\mu r^3 \kappa^2 \log^{3.5}(n)). \end{aligned}$$

This completes the control for $H^t - H^{t+1}$.

Next, to control A_0 , recall that $A_0 = \frac{\langle Z, (X^{t+1}H^t)_s(Y^{t+1}H^t)_s^\top - (X^{t+1}H^{t+1})_s(Y^{t+1}H^{t+1})_s^\top \rangle}{\|Z\|_F^2}$. Then, we have

$$\begin{aligned}
|A_0| &\leq \left| \frac{\langle Z, (X^{t+1}H^t)_s(Y^{t+1}H^t)_s^\top - (X^{t+1}H^{t+1})_s(Y^{t+1}H^{t+1})_s^\top \rangle}{\|Z\|_F^2} \right| \\
&\leq \frac{1}{\|Z\|_F^2} |\langle Z, (X^{t+1}H^t - X^{t+1}H^{t+1})_s(Y^{t+1}H^t)_s^\top \rangle| \\
&\quad + \frac{1}{\|Z\|_F^2} |\langle Z, (X^{t+1}H^{t+1})_s(Y^{t+1}H^t - Y^{t+1}H^{t+1})_s^\top \rangle| \\
&\stackrel{(i)}{\leq} \|(X^{t+1}H^t - X^{t+1}H^{t+1})_s\|_\infty \|(Y^{t+1}H^t)_s\|_\infty \\
&\quad + \|(X^{t+1}H^{t+1})_s\|_\infty \|(Y^{t+1}H^t - Y^{t+1}H^{t+1})_s\|_\infty \\
&\stackrel{(ii)}{\leq} \|X^{t+1}(H^t - H^{t+1})\|_{2,\infty} \|Y^{t+1}H^t\|_{2,\infty} \\
&\quad + \|X^{t+1}H^{t+1}\|_{2,\infty} \|Y^{t+1}(H^t - H^{t+1})\|_{2,\infty} \\
&\stackrel{(iii)}{\lesssim} \|X^{t+1}\|_{2,\infty} \|H^t - H^{t+1}\| \|Y^{t+1}\|_{2,\infty} \\
&\lesssim \eta \|F^*\|_{2,\infty}^2 \frac{\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} (\sigma\sqrt{n}\mu r^3 \kappa^2 \log^{3.5}(n)) \\
&\lesssim \eta \sigma^2 \log^6(n) \kappa^3 \mu^2 r^4.
\end{aligned}$$

Here, (i) is due to $\frac{1}{\|Z\|_F^2} |\langle Z, ab^\top \rangle| \leq \|ab^\top\|_\infty \leq \|a\|_\infty \|b\|_\infty$ for any $a, b \in \mathbb{R}^n$, (ii) is due to $\|(A)_s\|_\infty \leq \|A\|_{2,\infty}$ for any $A \in \mathbb{R}^{n \times r}$, and (iii) is due to $\|AH\|_{2,\infty} \leq \|A\|_{2,\infty} \|H\|$ for any matrices A and H . This completes the proof for A_0 . \square

Proof of Claim 6. Recall that we hope to show with probability $1 - O(n^{-10^7})$,

$$\begin{aligned}
\frac{1}{\|Z\|_F^2} |B_0| &\lesssim \frac{\sigma\mu r^{1.5}\kappa \log^2(n)}{\sqrt{n}} \sigma_{\min} + \frac{\sigma \log^{0.5}(n)}{r \|Z\|_F} \sigma_{\min} \\
\frac{1}{\|Z\|_F^2} |B_1| &\lesssim \frac{\sigma\mu r^{1.5}\kappa \log^2(n)}{\sqrt{n}} \sigma_{\min} \\
\frac{1}{\|Z\|_F^2} |B_2| &\lesssim \frac{\sigma\mu r^{1.5}\kappa \log^2(n)}{\sqrt{n}} \sigma_{\min} \\
\frac{1}{\|Z\|_F^2} |B_3| &\lesssim \frac{\sigma\mu r^{1.5}\kappa \log^{2.5}(n)}{\sqrt{n}} \sigma_s^*
\end{aligned}$$

where

$$\begin{aligned}
B_0 &= \langle Z, (XY^\top - M^* + (\tau - \tau^*)Z) (y_s y_s^\top - y_s^* y_s^{*\top}) \rangle \\
B_1 &= \langle Z, \lambda x_s y_s^\top \rangle \\
B_2 &= \langle Z, -E y_s y_s^\top \rangle \\
B_3 &= \langle Z, X(Y - Y^*)^\top y_s^* y_s^{*\top} \rangle.
\end{aligned}$$

To control B_0 , consider

$$\frac{1}{\|Z\|_F^2} |B_0| \leq \underbrace{\left| \frac{\langle Z, (XY^\top - M^*)(y_s y_s^\top - y_s^* y_s^{*\top}) \rangle}{\|Z\|_F^2} \right|}_{D_0} + \underbrace{\left| \frac{\langle Z, (\tau - \tau^*)Z(y_s y_s^\top - y_s^* y_s^{*\top}) \rangle}{\|Z\|_F^2} \right|}_{D_1}.$$

Note that $y_s y_s^\top - y_s^* y_s^{*\top} = y_s^* (y_s - y_s^*)^\top + (y_s - y_s^*) y_s^\top$. Then

$$\begin{aligned} \|y_s y_s^\top - y_s^* y_s^{*\top}\|_F &\leq (\|Y^*\| + \|Y\|) \|Y - Y^*\|_F \\ &\lesssim \sqrt{\sigma_{\max}} \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \|F^*\|_F \\ &\leq \sigma \sqrt{n} \log^{2.5}(n) \kappa r^{0.5}. \end{aligned}$$

This implies

$$\begin{aligned} D_1 &\leq |\tau - \tau^*| \frac{\|ZZ^\top\|_F}{\|Z\|_F^2} \|y_s y_s^\top - y_s^* y_s^{*\top}\|_F \\ &\stackrel{(i)}{\leq} |\tau - \tau^*| \sigma \sqrt{n} \log^{2.5}(n) \kappa r^{0.5} \end{aligned}$$

where (i) is due to $\frac{\|ZZ^\top\|_F}{\|Z\|_F^2} \leq \frac{\|Z\|_F \|Z^\top\|_F}{\|Z\|_F^2} \leq 1$.

Also, by $\frac{1}{\|Z\|_F^2} |\langle Z, A \rangle| \leq \|A\|_\infty$, one can verify that

$$\begin{aligned} D_0 &\leq \|(XY^\top - M^*)(y_s y_s^\top - y_s^* y_s^{*\top})\|_\infty \\ &\stackrel{(i)}{\leq} \|XY^\top - M^*\|_{2,\infty} \|y_s y_s^\top - y_s^* y_s^{*\top}\|_{2,\infty} \\ &\leq \|(X - X^*)Y^{*\top} + X(Y - Y^*)^\top\|_{2,\infty} \|y_s^* (y_s - y_s^*)^\top + (y_s - y_s^*) y_s^\top\|_{2,\infty} \\ &\stackrel{(ii)}{\leq} (\|X - X^*\|_{2,\infty} \|Y^*\| + \|X\|_{2,\infty} \|Y - Y^*\|) \\ &\quad \cdot (\|Y^*\|_{2,\infty} \|Y - Y^*\| + \|Y - Y^*\|_{2,\infty} \|Y\|). \end{aligned}$$

Here, (i) is by $\|AB^\top\|_\infty \leq \|A\|_{2,\infty} \|B\|_{2,\infty}$ for any matrices A, B , (ii) is by the triangle inequality and $\|AB\|_{2,\infty} \leq \|A\|_{2,\infty} \|B\|$. Recall that

$$\begin{aligned} \max(\|X - X^*\|_{2,\infty}, \|Y - Y^*\|_{2,\infty}) &\leq \|F - F^*\|_{2,\infty} \lesssim \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F \\ \max(\|X\|, \|Y\|, \|X^*\|, \|Y^*\|) &\lesssim \sqrt{\sigma_{\max}} \\ \max(\|X - X^*\|, \|Y - Y^*\|) &\leq \|F - F^*\|_F \lesssim \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \|F^*\|_F \\ \max(\|X\|_{2,\infty}, \|Y\|_{2,\infty}) &\lesssim \sqrt{\sigma_{\max}} \sqrt{\frac{\mu r}{n}}. \end{aligned}$$

We then have

$$\begin{aligned} D_0 &\lesssim \left(\frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F \sqrt{\sigma_{\max}} + \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \|F^*\|_F \sqrt{\sigma_{\max}} \sqrt{\frac{\mu r}{n}} \right)^2 \\ &\lesssim \left(\frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F \sqrt{\sigma_{\max}} \right)^2 \\ &\lesssim (\sigma \mu r^3 \kappa^2 \log^{3.5}(n))^2 \\ &\lesssim \sigma^2 \mu^2 r^6 \log^7(n) \kappa^4. \end{aligned}$$

This completes the bound for B_0 below.

$$\begin{aligned}
\frac{1}{\|Z\|_F^2} |B_0| &\leq |D_0| + |D_1| \\
&\lesssim \sigma^2 \mu^2 r^6 \log^7(n) \kappa^4 + |\tau - \tau^*| \sigma \sqrt{n} \log^{2.5}(n) \kappa r^{0.5} \\
&\lesssim \sigma^2 \mu^2 r^6 \log^7(n) \kappa^4 + \left(\frac{\sigma \mu r^{2.5} \kappa \log^{3.5} n}{\sqrt{n}} + \frac{\sigma \log^{1.5} n}{\|Z\|_F} \right) \sigma \sqrt{n} \log^{2.5}(n) \kappa r^{0.5} \\
&\stackrel{(i)}{\lesssim} \sigma \mu r^{1.5} \log^2(n) \kappa \cdot (\sigma \kappa^3 r^{4.5} \log^5(n) \mu) \\
&\quad + \left(\frac{\sigma \mu r^{2.5} \kappa \log^{3.5} n}{\sqrt{n}} + \frac{\sigma \log^{1.5} n}{\|Z\|_F} \right) \frac{\sigma_{\min}}{\kappa^2 r^4 \log^{2.5}(n)} \\
&\stackrel{(ii)}{\lesssim} \sigma \mu r^{1.5} \kappa \log^2(n) \frac{\sigma_{\min}}{\sqrt{n}} + \frac{\sigma \mu \log(n)}{\kappa r^{1.5}} \frac{\sigma_{\min}}{\sqrt{n}} + \frac{\sigma}{r^2 \kappa^4 \log(n) \|Z\|_F} \sigma_{\min} \\
&\lesssim \sigma \mu r^{1.5} \kappa \log^2(n) \frac{\sigma_{\min}}{\sqrt{n}} + \frac{\sigma \log^{0.5}(n)}{r \|Z\|_F} \sigma_{\min}
\end{aligned}$$

where (i) and (ii) are due to that $\frac{\sigma}{\sigma_{\min}} \sqrt{n} \lesssim \frac{1}{\kappa^3 r^{4.5} \log^5(n) \mu}$.

Next, consider B_1 . Recall that $B_1 = \langle Z, \lambda x_s y_s^\top \rangle$. We can directly obtain the desired bound $\frac{1}{\|Z\|_F^2} |B_1| \lesssim \frac{\sigma \mu r^{1.5} \kappa \log^2(n)}{\sqrt{n}} \sigma_{\min}$ by the following.

$$\begin{aligned}
\frac{1}{\|Z\|_F^2} |B_1| &\leq \|x_s\|_\infty \|y_s\|_\infty \lambda \\
&\lesssim \lambda \|F^*\|_{2,\infty}^2 \\
&\lesssim \sigma \sqrt{n} \log^{1.5}(n) \sigma_{\max} \frac{\mu r}{n} \\
&\lesssim \sigma \mu r \log^{1.5}(n) \frac{1}{\sqrt{n}} \sigma_{\max} \\
&\lesssim \frac{\sigma \mu r^{1.5} \kappa \log^2(n)}{\sqrt{n}} \sigma_{\min}.
\end{aligned}$$

Next, consider B_2 . Recall that $B_2 = \langle Z, -E y_s y_s^\top \rangle$ and we aim to show $\frac{1}{\|Z\|_F^2} |B_2| \lesssim \frac{\sigma \mu r^{1.5} \kappa \log^2(n)}{\sqrt{n}} \sigma_{\min}$. Note that

$$\frac{1}{\|Z\|_F^2} |B_2| \leq \|E y_s\|_\infty \|y_s\|_\infty.$$

To bound $\|E y_s\|_\infty$, let $y_s^{(l)}$ be the s -th column of $F^{t,(l)} R^{t,(l)}$ for $l \in [n]$. Let $E_{l,\cdot} \in \mathbb{R}^{1 \times n}$ be the l -th row of E . Then with probability $1 - O(n^{-10^7})$

$$\begin{aligned}
|(E y_s)_l| &= |E_{l,\cdot} y_s| \\
&\leq |E_{l,\cdot} y_s^{(l)}| + |E_{l,\cdot} (y_s^{(l)} - y_s)| \\
&\stackrel{(i)}{\lesssim} \sigma \|y_s^{(l)}\| \log^{0.5}(n) + \|E_{l,\cdot}\| \|y_s^{(l)} - y_s\| \\
&\lesssim \sigma \|F^{t,(l)} R^{t,(l)}\| \log^{0.5}(n) + \|E\| \|F^{t,(l)} R^{t,(l)} - F^t H^t\|_F
\end{aligned}$$

where (i) is due to independence between $E_{l,\cdot}$ and $y_s^{(l)}$ (Recall that the update of $F^{t,(l)}$ is independent from the l -th row of noise matrix E) and Hoeffding's inequality.

Recall that $\|F^{t,(l)}R^{t,(l)}\| \lesssim \|F^*\| = \sqrt{\sigma_{\max}}$, $\|E\| \lesssim \sigma\sqrt{n}$, and $\|F^{t,(l)}R^{t,(l)} - F^tH^t\|_{\mathbb{F}} \lesssim \frac{\sigma \log^{1.5} n}{\sigma_{\min}} \|F^*\|_{\mathbb{F}}$, we then have

$$\begin{aligned} |(Ey_s)_l| &\leq \sigma\sqrt{\sigma_{\max}} \log^{0.5}(n) + \sigma\sqrt{n} \frac{\sigma \log^{1.5} n}{\sigma_{\min}} \|F^*\|_{\mathbb{F}} \\ &\lesssim \sigma\sqrt{\sigma_{\max}} \log^{0.5}(n) + \sigma \frac{\sigma\sqrt{n} \log^{1.5}(n)\sqrt{r}}{\sigma_{\min}} \sqrt{\sigma_{\max}} \\ &\stackrel{(i)}{\lesssim} \sigma\sqrt{\sigma_{\max}} \log^{0.5}(n) + \sigma\sqrt{\sigma_{\max}} \\ &\lesssim \sigma\sqrt{\sigma_{\max}} \log^{0.5}(n). \end{aligned}$$

Here, (i) is due to $\frac{\sigma}{\sigma_{\min}}\sqrt{n} \lesssim \frac{1}{\kappa^3 r^{4.5} \log^5(n)\mu}$. Then this directly implies

$$\|Ey_s\|_{\infty} = \max_{l \in [n]} |(Ey_s)_l| \lesssim \sigma\sqrt{\sigma_{\max}} \log^{0.5}(n).$$

Therefore, we can obtain the desired bound for B_2 by the following.

$$\begin{aligned} \frac{1}{\|Z\|_{\mathbb{F}}^2} |B_2| &\leq \|Ey_s\|_{\infty} \|y_s\|_{\infty} \\ &\lesssim \sigma\sqrt{\sigma_{\max}} \log^{0.5}(n) \sqrt{\frac{\mu r \sigma_{\max}}{n}} \\ &\lesssim \frac{\sigma \mu r^{1.5} \kappa \log^2(n)}{\sqrt{n}} \sigma_{\min}. \end{aligned}$$

Finally, for B_3 , recall that $B_3 = \langle Z, X(Y - Y^*)^{\top} y_s^* y_s^{*\top} \rangle$. We can obtain the desired bound $\frac{1}{\|Z\|_{\mathbb{F}}^2} |B_3| \lesssim \frac{\sigma \mu r^{1.5} \kappa \log^{2.5}(n)}{\sqrt{n}} \sigma_s^*$ by the following.

$$\begin{aligned} \frac{1}{\|Z\|_{\mathbb{F}}^2} |B_3| &\stackrel{(i)}{\leq} \|X(Y - Y^*)^{\top}\|_{2,\infty} \|y_s^* y_s^{*\top}\|_{2,\infty} \\ &\stackrel{(ii)}{\lesssim} \|X\|_{2,\infty} \|Y - Y^*\| \|y_s^*\|_{2,\infty} \|y_s^*\| \\ &\lesssim \sqrt{\sigma_{\max}} \sqrt{\frac{\mu r}{n}} \frac{\sigma\sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \sqrt{\sigma_{\max} r} \sigma_s^* \sqrt{\frac{\mu r}{n}} \\ &\lesssim \sigma \kappa r^{1.5} \log^{2.5}(n) \mu \frac{\sigma_s^*}{\sqrt{n}}. \end{aligned}$$

Here, (i) is due to $\frac{1}{\|Z\|_{\mathbb{F}}^2} |\langle Z, AB^{\top} \rangle| \leq \|A\|_{2,\infty} \|B\|_{2,\infty}$ for any matrices A, B , and (ii) is due to $\|AB\|_{2,\infty} \leq \|A\|_{2,\infty} \|B\|$ for any matrices A, B . This completes the proof of the bounds for B_0, B_1, B_2, B_3 . \square

I.3 Proof of Eq. (59d)

Recall that we aim to obtain $\|F^{t+1}H^{t+1} - F^{t+1,(l)}R^{t+1,(l)}\|_{\mathbb{F}} \leq C_{l,1} \frac{\sigma \log^{1.5} n}{\sigma_{\min}} \|F^*\|_{\mathbb{F}}$. We will show the proof for $1 \leq l \leq n$ below. The scenario that $n < l \leq 2n$ is similar and omitted for brevity.

Note that $R^{t+1,(l)}$ is the optimal rotation to align $F^{t+1,(l)}$ and $F^{t+1}H^{t+1}$. In fact, we have

$$\left\| F^{t+1}H^{t+1} - F^{t+1,(l)}R^{t+1,(l)} \right\|_{\mathbb{F}} \leq \left\| F^{t+1}H^t - F^{t+1,(l)}R^{t,(l)} \right\|_{\mathbb{F}}.$$

Otherwise, by contradiction, suppose we have

$$\begin{aligned} \left\| F^{t+1}H^{t+1} - F^{t+1,(l)}R^{t+1,(l)} \right\|_{\mathbb{F}} &> \left\| F^{t+1}H^t - F^{t+1,(l)}R^{t,(l)} \right\|_{\mathbb{F}} \\ &\stackrel{(i)}{=} \left\| F^{t+1}H^t(H^{tT}H^{t+1}) - F^{t+1,(l)}R^{t,(l)}(H^{tT}H^{t+1}) \right\|_{\mathbb{F}} \\ &= \left\| F^{t+1}H^{t+1} - F^{t+1,(l)}R^{t,(l)}H^{tT}H^{t+1} \right\|_{\mathbb{F}}. \end{aligned}$$

Here, (i) is due to the unitary invariance of the $\|\cdot\|_{\mathbb{F}}$. This contradicts with that $R^{t+1,(l)} = \arg \min_{A \in \mathcal{O}^{r \times r}} \|F^{t+1}H^{t+1} - F^{t+1,(l)}A\|$. Hence we have

$$\left\| F^{t+1}H^{t+1} - F^{t+1,(l)}R^{t+1,(l)} \right\|_{\mathbb{F}} \leq \left\| F^{t+1}H^t - F^{t+1,(l)}R^{t,(l)} \right\|_{\mathbb{F}}.$$

Then it is sufficient to control $\|F^{t+1}H^t - F^{t+1,(l)}R^{t,(l)}\|_{\mathbb{F}}$.

First consider $\|X^{t+1}H^t - X^{t+1,(l)}R^{t,(l)}\|_{\mathbb{F}}$. Recall the updating rule for X^{t+1} and $X^{t+1,(l)}$ is

$$\begin{aligned} X^{t+1} &= X^t - \eta \left((X^t Y^{tT} - M^* - E + (\tau^t - \tau^*)Z) Y^t + \lambda X^t \right) \\ X^{t+1,(l)} &= X^{t,(l)} - \eta \left((X^{t,(l)} Y^{t,(l)T} - M^* - E + P_l(E) + (\tau^{t,(l)} - \tau^*)Z) Y^{t,(l)} + \lambda X^{t,(l)} \right) \end{aligned}$$

We then have

$$\begin{aligned} & X^{t+1,(l)}R^{t,(l)} - X^{t+1}H^t \\ &= \left(X^{t,(l)} - \eta \left((X^{t,(l)} Y^{t,(l)T} - M^* - E + P_l(E) + (\tau^{t,(l)} - \tau^*)Z) Y^{t,(l)} + \lambda X^{t,(l)} \right) \right) R^{t,(l)} \\ &\quad - \left(X^t - \eta \left((X^t Y^{tT} - M^* - E + (\tau^t - \tau^*)Z) Y^t + \lambda X^t \right) \right) H^t \\ &= X^{t,(l)}R^{t,(l)} - \eta \left(X^{t,(l)} Y^{t,(l)T} - M^* - E + P_l(E) + (\tau^{t,(l)} - \tau^*)Z \right) Y^{t,(l)} R^{t,(l)} \\ &\quad - \eta \lambda X^{t,(l)} R^{t,(l)} - \left(X^t H^t - \eta (X^t Y^{tT} - M^* - E + (\tau^t - \tau^*)Z) Y^t H^t - \eta \lambda X^t H^t \right) \\ &= (1 - \eta \lambda) (X^{t,(l)} R^{t,(l)} - X^t H^t) \\ &\quad + \underbrace{\eta \left(X^{t,(l)} Y^{t,(l)T} - M^* - E + P_l(E) + (\tau^{t,(l)} - \tau^*)Z \right) \left(Y^t H^t - Y^{t,(l)} R^{t,(l)} \right)}_{A_0} \\ &\quad - \eta \left(X^{t,(l)} Y^{t,(l)T} - X^t Y^{tT} + P_l(E) + (\tau^{t,(l)} - \tau^*)Z \right) Y^t H^t. \end{aligned}$$

To simplify the notation, we write $X^t H^t, Y^t H^t, F^t H^t, \tau^t$ as X, Y, F, τ if there is no ambiguity. Also, we write $X^{t,(l)} R^{t,(l)}, Y^{t,(l)} R^{t,(l)}, F^{t,(l)} R^{t,(l)}, \tau^{t,(l)}$ as $X^{(l)}, Y^{(l)}, F^{(l)}, \tau^{(l)}$. Furthermore, we write $\Delta_X = X^{(l)} - X, \Delta_Y = Y^{(l)} - Y, \Delta_F = F^{(l)} - F$. We have the following claim to control A_0 .

Claim 7. *With probability $1 - O(n^{-10^7})$,*

$$\|A_0\|_{\mathbb{F}} \lesssim \sigma \|F^*\|_{\mathbb{F}}.$$

Then

$$\begin{aligned} & \left\| X^{t+1,(l)}R^{t,(l)} - X^{t+1}H^t \right\|_{\mathbb{F}}^2 \\ &= \left\| (1 - \eta \lambda) \Delta_X - \eta \left(\left(X^{(l)} Y^{(l)T} - XY^T + P_l(E) + (\tau^{(l)} - \tau)Z \right) Y - A_0 \right) \right\|_{\mathbb{F}}^2 \\ &= (1 - \eta \lambda)^2 \|\Delta_X\|_{\mathbb{F}}^2 + A_1 \\ &\quad - 2(1 - \eta \lambda) \eta \underbrace{\left\langle \Delta_X, \left(\left(X^{(l)} Y^{(l)T} - XY^T + P_l(E) + (\tau^{(l)} - \tau)Z \right) Y - A_0 \right) \right\rangle}_{A_2} \end{aligned}$$

where A_1 includes the term with coefficient η^2 :

$$A_1 := \eta^2 \left\| \left(X^{(l)} Y^{(l)T} - XY^T + P_l(E) + (\tau^{(l)} - \tau)Z \right) Y - A_0 \right\|_{\mathbb{F}}^2.$$

Since we choose η sufficiently small, one can verify that $A_1 \lesssim \eta \frac{\sigma^2}{n^{15}}$ (similar as Eq. (66)).

Then, we proceed to analyze A_2 . Note that

$$\begin{aligned}
A_2 &= \left\langle \Delta_X, (X^{(l)}Y^{(l)\top} - XY^\top)Y \right\rangle + (\tau^{(l)} - \tau) \langle \Delta_X, ZY \rangle \\
&\quad + \underbrace{\langle \Delta_X, P_l(E)Y \rangle}_{B_0} - \langle \Delta_X, A_0 \rangle \\
&\stackrel{(i)}{=} \left\langle X^{(l)}Y^{(l)\top} - XY^\top, \Delta_X Y^\top \right\rangle + \frac{\langle Z, XY^\top - X^{(l)}Y^{(l)\top} \rangle \langle Z, \Delta_X Y^\top \rangle}{\|Z\|_F^2} \\
&\quad + \underbrace{\frac{\langle Z, -P_l(E) \rangle \langle Z, \Delta_X Y^\top \rangle}{\|Z\|_F^2}}_{B_1} + B_0 - \langle \Delta_X, A_0 \rangle \\
&\stackrel{(ii)}{=} \left\langle \Delta_X Y^\top + X \Delta_Y^\top, \Delta_X Y^\top \right\rangle + \underbrace{\langle \Delta_X \Delta_Y^\top, \Delta_X Y^\top \rangle}_{B_2} \\
&\quad - \frac{\langle Z, \Delta_X Y^\top + X \Delta_Y^\top \rangle \langle Z, \Delta_X Y^\top \rangle}{\|Z\|_F^2} \\
&\quad - \underbrace{\frac{\langle Z, \Delta_X \Delta_Y^\top \rangle \langle Z, \Delta_X Y^\top \rangle}{\|Z\|_F^2}}_{B_3} + B_1 + B_0 - \langle \Delta_X, A_0 \rangle
\end{aligned}$$

where (i) is due to

$$\begin{aligned}
\tau^{(l)} - \tau &= \frac{\langle Z, O - X^{(l)}Y^{(l)\top} - P_l(E) \rangle}{\|Z\|_F^2} - \frac{\langle Z, O - XY^\top \rangle}{\|Z\|_F^2} \\
&= \frac{\langle Z, XY^\top - X^{(l)}Y^{(l)\top} - P_l(E) \rangle}{\|Z\|_F^2},
\end{aligned}$$

and (ii) is due to $X^{(l)}Y^{(l)\top} - XY^\top = \Delta_X Y^\top + X \Delta_Y^\top + \Delta_X \Delta_Y^\top$.

We have the following claim to control B_0, B_1, B_2 and B_3 .

Claim 8. *With probability $1 - O(n^{-10^7})$,*

$$\begin{aligned}
|B_0| &\lesssim \sigma \log^{0.5}(n) \|\Delta_F\|_F \|F^*\|_F \\
|B_1| &\lesssim \sigma \log^{0.5}(n) \|\Delta_F\|_F \|F^*\|_F \\
|B_2| &\lesssim \sigma \|\Delta_F\|_F \|F^*\|_F \\
|B_3| &\lesssim \sigma \|\Delta_F\|_F \|F^*\|_F.
\end{aligned}$$

Then, note that η is choosing sufficiently small where $\eta\lambda \ll 1$, we have (for some constant C)

$$\begin{aligned}
&\left\| X^{t+1, (l)} R^{t, (l)} - X^{t+1} H^t \right\|_F^2 \\
&= (1 - \eta\lambda)^2 \|\Delta_X\|_F^2 + A_1 - 2(1 - \eta\lambda)\eta A_2 \\
&= (1 - \eta\lambda)^2 \|\Delta_X\|_F^2 + A_1 - 2(1 - \eta\lambda)\eta (B_0 + B_1 + B_2 - B_3 - \langle \Delta_X, A_0 \rangle) \\
&\quad - 2(1 - \eta\lambda)\eta \left(\langle \Delta_X Y^\top + X \Delta_Y^\top, \Delta_X Y^\top \rangle - \frac{\langle Z, \Delta_X Y^\top + X \Delta_Y^\top \rangle \langle Z, \Delta_X Y^\top \rangle}{\|Z\|_F^2} \right) \\
&\leq \|\Delta_X\|_F^2 + \eta C \|\Delta_F\|_F \sigma \log^{0.5}(n) \|F^*\|_F \\
&\quad - 2(1 - \eta\lambda)\eta \left(\langle \Delta_X Y^\top + X \Delta_Y^\top, \Delta_X Y^\top \rangle - \frac{\langle Z, \Delta_X Y^\top + X \Delta_Y^\top \rangle \langle Z, \Delta_X Y^\top \rangle}{\|Z\|_F^2} \right).
\end{aligned}$$

By the symmetry, we can similarly obtain the results for $Y^{t+1, (l)} R^{t, (l)} - Y^{t+1} H^t$

$$\begin{aligned} & \left\| Y^{t+1, (l)} R^{t, (l)} - Y^{t+1} H^t \right\|_{\mathbb{F}}^2 \\ & \leq \|\Delta_Y\|_{\mathbb{F}}^2 + \eta C \|\Delta_F\|_{\mathbb{F}} \sigma \log^{0.5}(n) \|F^*\|_{\mathbb{F}} \\ & \quad - 2(1 - \eta\lambda)\eta \left(\langle \Delta_X Y^\top + X \Delta_Y^\top, X \Delta_Y^\top \rangle - \frac{\langle Z, \Delta_X Y^\top + X \Delta_Y^\top \rangle \langle Z, X \Delta_Y^\top \rangle}{\|Z\|_{\mathbb{F}}^2} \right). \end{aligned}$$

Together implies that

$$\begin{aligned} & \left\| F^{t+1, (l)} R^{t, (l)} - F^{t+1} H^t \right\|_{\mathbb{F}}^2 \\ & \leq \|\Delta_F\|_{\mathbb{F}}^2 + 2\eta C \|\Delta_F\|_{\mathbb{F}} \sigma \log^{0.5}(n) \|F^*\|_{\mathbb{F}} \\ & \quad - 2(1 - \eta\lambda)\eta \left(\|\Delta_X Y^\top + X \Delta_Y^\top\|_{\mathbb{F}}^2 - \frac{(\langle Z, \Delta_X Y^\top + X \Delta_Y^\top \rangle)^2}{\|Z\|_{\mathbb{F}}^2} \right). \end{aligned}$$

Let T be the tangent space of XY^\top . We have (for some constant C')

$$\begin{aligned} & \|\Delta_X Y^\top + X \Delta_Y^\top\|_{\mathbb{F}}^2 - \frac{(\langle Z, \Delta_X Y^\top + X \Delta_Y^\top \rangle)^2}{\|Z\|_{\mathbb{F}}^2} \\ & = \|\Delta_X Y^\top + X \Delta_Y^\top\|_{\mathbb{F}}^2 - \frac{\langle P_T(Z), \Delta_X Y^\top + X \Delta_Y^\top \rangle^2}{\|Z\|_{\mathbb{F}}^2} \\ & \geq \|\Delta_X Y^\top + X \Delta_Y^\top\|_{\mathbb{F}}^2 \left(1 - \frac{\|P_T(Z)\|_{\mathbb{F}}^2}{\|Z\|_{\mathbb{F}}^2} \right) \\ & \stackrel{(i)}{\geq} \frac{C'}{\log(n)} \|\Delta_X Y^\top + X \Delta_Y^\top\|_{\mathbb{F}}^2. \end{aligned}$$

Here (i) is due to the implication of Assumption 3(a) for (X, Y) (by Lemma 16).

Then we have the following claim. The proof is similar to the analysis in Claim 4 by using the property that $\|X^{(l)\top} X - Y^{(l)\top} Y\|_{\mathbb{F}} \lesssim \frac{\sigma}{n^{15}}$ (similar to Eq. (59)). We omit the proof for the brevity.

Claim 9.

$$\|\Delta_X Y^\top + X \Delta_Y^\top\|_{\mathbb{F}}^2 \geq \frac{\sigma_{\min}}{4} \|\Delta_F\|_{\mathbb{F}}^2 - \frac{\sigma^2}{n^{13}}.$$

Then, we have

$$\begin{aligned} & \left\| F^{t+1, (l)} R^{t, (l)} - F^{t+1} H^t \right\|_{\mathbb{F}}^2 \\ & \leq \|\Delta_F\|_{\mathbb{F}}^2 \left(1 - \eta \frac{C' \sigma_{\min}}{4 \log(n)} \right) + 2\eta \frac{C' \sigma^2}{\log(n) n^{13}} + 2\eta C \|\Delta_F\|_{\mathbb{F}} \sigma \log^{0.5}(n) \|F^*\|_{\mathbb{F}} \\ & \leq \left(C_1 \frac{\sigma \log^{1.5} n}{\sigma_{\min}} \|F^*\|_{\mathbb{F}} \right)^2 - \underbrace{\eta \frac{C' \sigma_{\min}}{4 \log(n)} \left(C_1 \frac{\sigma \log^{1.5} n}{\sigma_{\min}} \|F^*\|_{\mathbb{F}} \right)^2}_{K_1} \\ & \quad + \underbrace{\eta 2 \frac{C' \sigma^2}{\log(n) n^{13}}}_{K_2} + \underbrace{\eta 2C \left(C_1 \frac{\sigma \log^{1.5} n}{\sigma_{\min}} \|F^*\|_{\mathbb{F}} \right) \sigma \log^{0.5}(n) \|F^*\|_{\mathbb{F}}}_{K_3}. \end{aligned}$$

In order to show $\|F^{t+1, (l)} R^{t, (l)} - F^{t+1} H^t\|_F^2 \leq \left(C_1 \frac{\sigma \log^{1.5} n}{\sigma_{\min}} \|F^*\|_F\right)^2$, it is sufficient to show that $K_1 \geq K_2 + K_3$. Take $C_1 \geq 4 + 8 \frac{C}{C'}$, one can verify that

$$\begin{aligned} K_1 - K_2 &= \left(\frac{C' C_1}{4} - 2C\right) \left(C_1 \frac{\sigma \log^{1.5} n}{\sigma_{\min}} \|F^*\|_F\right) \sigma \log^{0.5}(n) \|F^*\|_F \\ &\geq 2C' \frac{\sigma^2 \log^2(n)}{\sigma_{\min}} \|F^*\|_F^2 \\ &\geq 2C' \sigma^2 \log^2(n) \kappa \\ &\geq K_3. \end{aligned}$$

This completes the proof for Eq. (59d).

Proof of Claim 7. Recall that we want to show $|A_0| \lesssim \sigma \|F^*\|_F$ where $A_0 = (X^{t, (l)} Y^{t, (l)\top} - M^* - E + P_l(E) + (\tau^{t, (l)} - \tau^*) Z) (Y^t H^t - Y^{t, (l)} R^{t, (l)})$.

In fact, we have the following bounds for leave-one-out sequences similar to Eq. (59b) and Eq. (59c). The proof is exactly the same by viewing $E - P_l(E)$ as the noise. We omit it for brevity.

$$\begin{aligned} |\tau^{(l)} - \tau^*| &\lesssim \left(\frac{\sigma \mu r^{2.5} \kappa \log^{3.5} n}{\sqrt{n}} + \frac{\sigma \log^{1.5} n}{\|Z\|_F}\right) \\ \|F^{(l)} H^{(l)} - F^*\|_F &\lesssim C_F \left(\frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}}\right) \|F^*\|_F. \end{aligned}$$

Then, with probability $1 - O(n^{-10^7})$,

$$\begin{aligned} |A_0| &\leq \left(\|X^{(l)} Y^{(l)\top} - M^*\| + \|E\| + \|Z\| |\tau^{(l)} - \tau^*|\right) \|\Delta_F\|_F \\ &\lesssim \left(\|F^{(l)} H^{(l)} - F^*\|_F \|F^*\| + \sigma \sqrt{n} + \sigma \sqrt{n} \mu r^{2.5} \kappa \log^{3.5}(n)\right) \|\Delta_F\|_F \\ &\lesssim (\sigma \sqrt{n} \log^{2.5}(n) \kappa r + \sigma \sqrt{n} \mu r^{2.5} \kappa \log^{3.5}(n)) \|\Delta_F\|_F \\ &\lesssim \sigma \|F^*\|_F \end{aligned}$$

providing that $\frac{\sigma}{\sigma_{\min}} \sqrt{n} \lesssim \frac{1}{\kappa^3 r^{4.5} \log^5(n) \mu}$ and $\|\Delta_F\|_F \lesssim \frac{\sigma \log^{1.5}(n)}{\sigma_{\min}} \|F^*\|_F$. \square

Proof of Claim 8. For B_0 , recall that we want to show $|B_0| \lesssim \sigma \log^{0.5}(n) \|\Delta_F\|_F \|F^*\|_F$ where $B_0 = \langle \Delta_X, P_l(E) Y \rangle$. With probability $1 - O(n^{-10^7})$, we have

$$\begin{aligned} |B_0| &\leq \|\Delta_X\|_F \left\| P_l(E) Y^{(l)} \right\|_F + \|\Delta_X\|_F \|P_l(E)\| \left\| Y - Y^{(l)} \right\|_F \\ &\stackrel{(i)}{\lesssim} \|\Delta_F\|_F \sigma \log^{0.5}(n) \|F^*\|_F + \|\Delta_X\|_F \|P_l(E)\| \left\| Y - Y^{(l)} \right\|_F \\ &\lesssim \|\Delta_F\|_F \sigma \log^{0.5}(n) \|F^*\|_F + \|\Delta_F\|_F \sigma \sqrt{n} \frac{\sigma \log^{1.5}(n)}{\sigma_{\min}} \|F^*\|_F \\ &\lesssim \|\Delta_F\|_F \sigma \log^{0.5}(n) \|F^*\|_F \end{aligned}$$

where (i) is due to the independence between $P_l(E)$ and $Y^{(l)}$ and the Hanson-Wright inequality [49].

For B_1 , recall that we want to show $|B_1| \lesssim \sigma \log^{0.5}(n) \|\Delta_F\|_F \|F^*\|_F$ where $B_1 = \frac{\langle Z, -P_l(E) \rangle \langle Z, \Delta_X Y^\top \rangle}{\|Z\|_F^2}$. With probability $1 - O(n^{-10^7})$, we have

$$|\langle Z, P_l(E) \rangle| \lesssim \sigma \sqrt{\log(n)} \|P_l(Z)\|_F$$

by the Hoeffding's inequality, hence

$$\begin{aligned} |B_1| &\lesssim \sigma \sqrt{\log(n)} \|P_l(Z)\|_F \|Z\|_F \|\Delta_F\|_F \|F^*\|_F \frac{1}{\|Z\|_F^2} \\ &\lesssim \sigma \sqrt{\log(n)} \|\Delta_F\|_F \|F^*\|_F. \end{aligned}$$

For B_2 , recall that we want to show $|B_2| \lesssim \sigma \|\Delta_F\|_F \|F^*\|_F$ where $B_2 = \langle \Delta_X \Delta_Y^\top, \Delta_X Y^\top \rangle$. We have

$$\begin{aligned} |B_2| &\lesssim \|\Delta_F\|_F^3 \|F^*\|_F \\ &\lesssim \|\Delta_F\|_F \|F^*\|_F \left(\frac{\sigma \log^{1.5}(n)}{\sigma_{\min}} \|F^*\|_F \right)^2 \\ &\lesssim \|\Delta_F\|_F \|F^*\|_F \frac{\sigma^2 \log^3(n) r \kappa}{\sigma_{\min}} \\ &\lesssim \sigma \|\Delta_F\|_F \|F^*\|_F. \end{aligned}$$

For B_3 , recall that we want to show $|B_3| \lesssim \sigma \|\Delta_F\|_F \|F^*\|_F$ where $B_3 = \frac{\langle Z, \Delta_X \Delta_Y^\top \rangle \langle Z, \Delta_X Y^\top \rangle}{\|Z\|_F^2}$. Similar to B_2 , we have

$$\begin{aligned} |B_3| &\lesssim \|Z\|_F^2 \|\Delta_F\|_F^3 \|F^*\|_F \frac{1}{\|Z\|_F^2} \\ &\lesssim \sigma \|\Delta_F\|_F \|F^*\|_F. \end{aligned}$$

This completes the proof for B_0, B_1, B_2, B_3 . \square

I.4 Proof of Eqs. (59e) and (59f)

Recall that we want to show (Eq. (59e))

$$\max_{1 \leq l \leq 2n} \left\| (F^{t+1, (l)} H^{t+1, (l)} - F^*)_{l, \cdot} \right\|_2 \leq C_{l,2} \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F$$

and $\|F^{t+1} H^{t+1} - F^*\|_{2, \infty} \leq C_\infty \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F$ (Eq. (59f)).

For Eq. (59e), we consider $1 \leq l \leq n$, where the scenario $n+1 \leq l \leq 2n$ is similar. By triangle inequality,

$$\begin{aligned} \left\| (F^{t+1, (l)} H^{t+1, (l)} - F^*)_{l, \cdot} \right\|_2 &\leq \left\| (F^{t+1, (l)} H^{t, (l)} - F^*)_{l, \cdot} \right\|_2 \\ &\quad + \underbrace{\left\| (F^{t+1, (l)} (H^{t, (l)} - H^{t+1, (l)}))_{l, \cdot} \right\|_2}_{T_0}. \end{aligned}$$

We have the following claim.

Claim 10.

$$|T_0| \lesssim \eta \sigma \mu^{0.5} \|F^*\|_F.$$

For $(F^{t+1, (l)} H^{t, (l)} - F^*)_{l, \cdot}$, note that

$$\begin{aligned} &(F^{t+1, (l)} H^{t, (l)} - F^*)_{l, \cdot} \\ &= (X^{t+1, (l)} H^{t, (l)} - X^*)_{l, \cdot} \\ &= (X^{t, (l)} H^{t, (l)} - X^*)_{l, \cdot} \\ &\quad - \eta \left((X^{t, (l)} Y^{t, (l)\top} - M^* - E + P_l(E) + (\tau^{t, (l)} - \tau^*) Z) Y^{t, (l)} H^{t, (l)} + \lambda X^{t, (l)} H^{t, (l)} \right)_{l, \cdot}. \end{aligned}$$

For simplifying the notation, we write $X^{t, (l)} H^{t, (l)}$, $Y^{t, (l)} H^{t, (l)}$ as $X^{(l)}$, $Y^{(l)}$ and $F^{t, (l)} H^{t, (l)}$, $\tau^{t, (l)}$ as $F^{(l)}$, $\tau^{(l)}$. Furthermore, we write $\Delta_X = X^{(l)} - X^*$, $\Delta_Y = Y^{(l)} - Y^*$, $\Delta_F = F^{(l)} - F^*$. Then, we have

$$\begin{aligned} &\left\| (F^{t+1, (l)} H^{t, (l)} - F^*)_{l, \cdot} \right\|_2^2 \\ &= \left\| (\Delta_X)_{l, \cdot} \right\|_2^2 + A_0 \\ &\quad - 2\eta \underbrace{\left\langle (\Delta_X)_{l, \cdot}, (X^{(l)} Y^{(l)\top} - M^* - E + P_l(E) + (\tau^{(l)} - \tau^*) Z)_{l, \cdot} Y^{(l)} + \lambda X_{l, \cdot} \right\rangle}_{A_1}. \end{aligned}$$

where A_0 includes the term with η^2 :

$$A_0 := \eta^2 \left\| (X^{(l)} Y^{(l)\top} - M^* - E + P_l(E) + (\tau^{(l)} - \tau^*) Z)_{l,\cdot} Y^{(l)} + \lambda X_{l,\cdot} \right\|_2^2.$$

Since we choose η sufficiently small, one can verify that $|A_0| \lesssim \eta \frac{\sigma_{\min}^2}{n^{15}}$ (similar to Eq. (66)).

Next, consider A_1 . Note that $(P_l(E) - E)_{l,\cdot} = 0$, then

$$\begin{aligned} A_1 &= \left\langle (\Delta_X)_{l,\cdot}, (X^{(l)} Y^{(l)\top} - M^*)_{l,\cdot} Y^{(l)} \right\rangle \\ &\quad + \underbrace{\left\langle (\Delta_X)_{l,\cdot}, (\tau^{(l)} - \tau^*) Z_{l,\cdot} Y^{(l)} \right\rangle}_{B_0} + \underbrace{\left\langle (\Delta_X)_{l,\cdot}, \lambda X_{l,\cdot} \right\rangle}_{B_1} \\ &= \underbrace{\left\langle (\Delta_X)_{l,\cdot}, (\Delta_X Y^{(l)\top})_{l,\cdot} Y^{(l)} \right\rangle}_{B_2} + \underbrace{\left\langle (\Delta_X)_{l,\cdot}, (X^* \Delta_Y^\top)_{l,\cdot} Y^{(l)} \right\rangle}_{B_3} + B_0 + B_1. \end{aligned}$$

Note that

$$\begin{aligned} B_2 &= \text{tr}((\Delta_X)_{l,\cdot}^\top (\Delta_X)_{l,\cdot} Y^{(l)\top} Y^{(l)}) \\ &= \text{tr}((\Delta_X)_{l,\cdot}^\top (\Delta_X)_{l,\cdot} (Y^{(l)\top} Y^{(l)} - \sigma_r(Y^{(l)\top} Y^{(l)}) I_r)) + \sigma_r(Y^{(l)\top} Y^{(l)}) \text{tr}((\Delta_X)_{l,\cdot}^\top (\Delta_X)_{l,\cdot}) \\ &\stackrel{(i)}{\geq} \sigma_r(Y^{(l)\top} Y^{(l)}) \text{tr}((\Delta_X)_{l,\cdot}^\top (\Delta_X)_{l,\cdot}) \\ &\stackrel{(ii)}{\geq} \frac{\sigma_{\min}}{2} \|(\Delta_X)_{l,\cdot}\|_2^2. \end{aligned}$$

where (i) is due to $\text{tr}(AB) \geq 0$ if A, B are positive semi-definite matrices, (ii) is due to $\sigma_r(Y^{(l)}) \geq \sqrt{\frac{\sigma_{\min}}{2}}$. We also have the following claim for controlling B_0, B_1 and B_3 .

Claim 11.

$$\begin{aligned} |B_0| &\lesssim \sigma \mu r^{2.5} \log^{3.5}(n) \kappa \|\Delta_F\|_{2,\infty} \|F^*\|_F \\ |B_1| &\lesssim \sigma r^{0.5} \mu^{0.5} \kappa^{0.5} \log^{1.5}(n) \|\Delta_F\|_{2,\infty} \|F^*\|_F \\ |B_3| &\lesssim \sigma \kappa r^{1.5} \mu^{0.5} \log^{2.5}(n) \|\Delta_X\|_{2,\infty} \|F^*\|_F. \end{aligned}$$

Finally, take $C_2 > 4C$ large enough and note that $\|\Delta_F\|_{2,\infty} \lesssim C_2 \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F$, we arrive at

$$\begin{aligned} &\left\| (F^{t+1,(l)} H^{t,(l)} - F^*)_{l,\cdot} \right\|_2^2 \\ &= \|(\Delta_X)_{l,\cdot}\|_2^2 + A_0 - 2\eta A_1 \\ &= \|(\Delta_X)_{l,\cdot}\|_2^2 - 2\eta(B_2 + B_3 + B_0 + B_1) + A_0 \\ &\leq \|(\Delta_X)_{l,\cdot}\|_2^2 - 2\eta \frac{\sigma_{\min}}{2} \|(\Delta_X)_{l,\cdot}\|_2^2 + C\eta \sigma \mu r^{2.5} \log^{3.5}(n) \kappa \|\Delta_F\|_{2,\infty} \|F^*\|_F \\ &\leq \left(C_2 \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F \right)^2 (1 - \eta \sigma_{\min}) \\ &\quad + C\eta \left(C_2 \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F \right) \sigma \mu r^{2.5} \log^{3.5}(n) \|F^*\|_F \kappa \\ &\leq \left(C_2 \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F \right)^2 \left(1 - \frac{\sigma_{\min}}{4} \eta \right)^2. \end{aligned}$$

Furthermore

$$\begin{aligned}
& \left\| (F^{t+1,(l)} H^{t+1,(l)} - F^*)_{l,\cdot} \right\|_2 \\
& \leq \left\| (F^{t+1,(l)} H^{t,(l)} - F^*)_{l,\cdot} \right\|_2 + T_0 \\
& \leq \left(C_2 \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_{\mathbb{F}} \right) \left(1 - \frac{\sigma_{\min}}{4} \eta \right) + C \eta \sigma \mu^{0.5} \|F^*\|_{\mathbb{F}} \\
& \leq C_2 \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_{\mathbb{F}}
\end{aligned}$$

This completes the proof for Eq. (59e).

Then, for Eq. (59f), note that by triangle inequality, for each $l \in [n]$,

$$\begin{aligned}
\|F^{t+1} H^{t+1} - F^*\|_{2,\infty} & \leq \left\| F^{t+1,(l)} H^{t+1,(l)} - F^* \right\|_{2,\infty} + \left\| F^{t+1,(l)} H^{t+1,(l)} - F^{t+1} H^{t+1} \right\|_{\mathbb{F}} \\
& \leq C_2 \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_{\mathbb{F}} + \left\| F^{t+1,(l)} H^{t+1,(l)} - F^{t+1} H^{t+1} \right\|_{\mathbb{F}}.
\end{aligned}$$

Then it is sufficient to consider $\left\| F^{t+1,(l)} H^{t+1,(l)} - F^{t+1} H^{t+1} \right\|_{\mathbb{F}}$. Take $F_0 = F^*$, $F_1 = F^{t+1} H^{t+1}$, $F_2 = F^{t+1,(l)} R^{t+1,(l)}$. By Lemma 12 and the definition of H^{t+1} , $H^{t+1,(l)}$ and $R^{t+1,(l)}$, we then have

$$\begin{aligned}
\left\| F^{t+1,(l)} H^{t+1,(l)} - F^{t+1} H^{t+1} \right\|_{\mathbb{F}} & \leq \|F_1 - F_2\|_{\mathbb{F}} \\
& \lesssim \kappa \frac{\sigma \log^{1.5}(n)}{\sigma_{\min}} \|F^*\|_{\mathbb{F}}. \tag{69}
\end{aligned}$$

This implies, for large enough n ,

$$\begin{aligned}
\|F^{t+1} H^{t+1} - F^*\|_{2,\infty} & \leq C_2 \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_{\mathbb{F}} + C \kappa \frac{\sigma \log^{1.5}(n)}{\sigma_{\min}} \|F^*\|_{\mathbb{F}} \\
& \leq 2C_2 \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_{\mathbb{F}}
\end{aligned}$$

which completes the proof.

Proof of Claim 10. Recall that we want to show $|T_0| \lesssim \eta \sigma \mu^{0.5} \|F^*\|_{\mathbb{F}}$ where

$$T_0 = \left\| (F^{t+1,(l)} (H^{t,(l)} - H^{t+1,(l)}))_{l,\cdot} \right\|_2.$$

Note that

$$\begin{aligned}
|T_0| & \lesssim \left\| F^{t+1,(l)} \right\|_{2,\infty} \left\| H^{t,(l)} - H^{t+1,(l)} \right\| \\
& \lesssim \sqrt{\sigma_{\max} r} \sqrt{\frac{\mu r}{n}} \left\| H^{t,(l)} - H^{t+1,(l)} \right\| \\
& \stackrel{(i)}{\lesssim} \sqrt{\sigma_{\max} r} \sqrt{\frac{\mu r}{n}} \eta \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} (\sigma \sqrt{n} \mu r^3 \kappa^2 \log^{3.5}(n)) \\
& \lesssim \eta \sigma^2 \sqrt{n} \mu^{1.5} r^4 \kappa^{2.5} \log^5(n) \frac{1}{\sqrt{\sigma_{\min}}} \\
& \lesssim \eta \frac{\sigma^2 \sqrt{n} \mu^{1.5} r^{4.5} \kappa^{2.5} \log^5(n)}{\sigma_{\min}} \|F^*\|_{\mathbb{F}} \\
& \lesssim \eta \sigma \mu^{0.5} \|F^*\|_{\mathbb{F}}
\end{aligned}$$

where (i) is by the bound of

$$\left\| H^{t,(l)} - H^{t+1,(l)} \right\|_{\mathbb{F}} \lesssim \eta \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} (\sigma \sqrt{n} \mu r^3 \kappa^2 \log^{3.5}(n)).$$

This proof of the bound of $\left\| H^{t,(l)} - H^{t+1,(l)} \right\|_{\mathbb{F}}$ is exactly the same as the Claim 5 by viewing $E - P_l(E)$ as the noise matrix, where we omit for the brevity. \square

Proof of Claim 11. For B_0 , recall that we want to show

$$B_0 \lesssim \sigma \mu r^{2.5} \log^{3.5}(n) \kappa \|\Delta_F\|_{2,\infty} \|F^*\|_F$$

where $B_0 = \langle (\Delta_X)_{l,\cdot}, (\tau^{(l)} - \tau^*) Z_{l,\cdot} Y^{(l)} \rangle$. We have

$$\begin{aligned} |B_0| &\leq |\tau^{(l)} - \tau^*| \|\Delta_X\|_{2,\infty} \left\| Z Y^{(l)} \right\|_{2,\infty} \\ &\lesssim \left(\frac{\sigma \mu r^{2.5} \log^{3.5}(n) \kappa}{\sqrt{n}} + \frac{\sigma \log^{1.5}(n)}{\|Z\|_F} \right) \|\Delta_F\|_{2,\infty} \|Z\|_{2,\infty} \|Y^*\|_F \\ &\lesssim \left(\frac{\sigma \mu r^{2.5} \log^{3.5}(n) \kappa}{\sqrt{n}} + \frac{\sigma \log^{1.5}(n)}{\|Z\|_F} \right) \|\Delta_F\|_{2,\infty} \min(\|Z\|_F, \sqrt{n}) \|Y^*\|_F \\ &\lesssim \sigma \mu r^{2.5} \log^{3.5}(n) \kappa \|\Delta_F\|_{2,\infty} \|F^*\|_F. \end{aligned}$$

For B_1 , recall that we want to show $B_1 \lesssim \sigma r^{0.5} \mu^{0.5} \kappa^{0.5} \log^{1.5}(n) \|\Delta_F\|_{2,\infty} \|F^*\|_F$ where $B_1 = \langle (\Delta_X)_{l,\cdot}, \lambda X_{l,\cdot} \rangle$. We have

$$\begin{aligned} |B_1| &\leq \lambda \|\Delta_X\|_{2,\infty} \|X\|_{2,\infty} \\ &\lesssim \sigma \sqrt{n} \log^{1.5}(n) \|\Delta_X\|_{2,\infty} \sqrt{\sigma_{\max} r} \sqrt{\frac{\mu r}{n}} \\ &\lesssim \sigma r^{0.5} \mu^{0.5} \kappa^{0.5} \log^{1.5}(n) \|\Delta_X\|_{2,\infty} \sqrt{\sigma_{\min} r} \\ &\lesssim \sigma r^{0.5} \mu^{0.5} \kappa^{0.5} \log^{1.5}(n) \|\Delta_F\|_{2,\infty} \|F^*\|_F. \end{aligned}$$

For B_3 , recall that we want to show $B_3 \lesssim \sigma \kappa r^{1.5} \mu^{0.5} \log^{2.5}(n) \|\Delta_X\|_{2,\infty} \|F^*\|_F$ where $B_3 = \langle (\Delta_X)_{l,\cdot}, (X^* \Delta_Y^\top)_{l,\cdot} Y^{(l)} \rangle$. We have

$$\begin{aligned} |B_3| &\leq \|\Delta_X\|_{2,\infty} \left\| X^* \Delta_Y^\top Y^{(l)} \right\|_{2,\infty} \\ &\lesssim \|\Delta_X\|_{2,\infty} \|X^*\|_{2,\infty} \|\Delta_Y\|_F \left\| Y^{(l)} \right\| \\ &\lesssim \sqrt{\sigma_{\max} r} \sqrt{\frac{\mu r}{n}} \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \sqrt{\sigma_{\max} r} \|\Delta_X\|_{2,\infty} \|F^*\|_F \\ &\lesssim \sigma \kappa r^{1.5} \mu^{0.5} \log^{2.5}(n) \|\Delta_X\|_{2,\infty} \|F^*\|_F. \end{aligned}$$

This completes the proof for B_0, B_1, B_3 . \square

I.5 Proof of Eq. (60a)

Let T be the tangent space of $X^t Y^{tT}$. Recall we want to show $\|P_{T^\perp}(M^*)\|_\infty \leq C_{T,1} \frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^7(n)}{\sigma_{\min}}$.

Let $X = X^t H^t, Y = Y^t H^t, \Delta_X = X - X^*, \Delta_Y = Y - Y^*$ if there is no ambiguity. Then $T = \text{span}\{XA^\top + BY^\top \mid A, B \in \mathbb{R}^{n \times r}\}$.

Note that we have

$$\begin{aligned} P_{T^\perp}(M^*) &= P_{T^\perp}(X^* Y^{*\top}) \\ &= P_{T^\perp}((X - \Delta_X)(Y - \Delta_Y)^\top) \\ &\stackrel{(i)}{=} P_{T^\perp}(\Delta_X \Delta_Y^\top) \end{aligned}$$

where (i) is due to $P_{T^\perp}(XY^\top) = P_{T^\perp}(\Delta_X Y^\top) = P_{T^\perp}(X \Delta_Y^\top) = 0$.

Let $U \Sigma V^\top$ be the SVD of XY^\top . We also have that $U^* \Sigma^* V^{*\top}$ is the SVD of $X^* Y^{*\top}$. By Eq. (59c), Eq. (59f), and Eq. (59g), we can invoke Lemma 15 and obtain that there exists a rotation matrix

$R \in \mathcal{O}^{r \times r}$ such that

$$\max(\|UR - U^*\|_F, \|VR - V^*\|_F) \lesssim \frac{r\kappa\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \quad (70)$$

$$\max(\|UR - U^*\|_{2,\infty}, \|VR - V^*\|_{2,\infty}) \lesssim \frac{r^3\mu\kappa^{1.5}\sigma\log^{3.5}(n)}{\sigma_{\min}}. \quad (71)$$

Then consider

$$\begin{aligned} \|P_{T^\perp}(M^*)\|_\infty &= \|P_{T^\perp}(\Delta_X\Delta_Y^\top)\|_\infty \\ &= \|(I - UU^\top)\Delta_X\Delta_Y^\top(I - VV^\top)\|_\infty \\ &\leq \|(I - UU^\top)\Delta_X\|_{2,\infty} \|(I - VV^\top)\Delta_Y\|_{2,\infty}. \end{aligned}$$

From Eq. (71), we have $\|U\|_{2,\infty} \lesssim \|U^*\|_{2,\infty} + \frac{r^3\mu\kappa^{1.5}\sigma\log^{3.5}(n)}{\sigma_{\min}} \lesssim \sqrt{\frac{\mu r}{n}}$ due to the incoherence condition and $\frac{\sigma}{\sigma_{\min}}\sqrt{n} \lesssim \frac{1}{\kappa^3 r^{4.5} \log^5(n)\mu}$. Then

$$\begin{aligned} \|(I - UU^\top)\Delta_X\|_{2,\infty} &\leq \|\Delta_X\|_{2,\infty} + \|U\|_{2,\infty} \|U^\top\| \|\Delta_X\| \\ &\lesssim \frac{\sigma\mu r^3\kappa\log^{3.5}(n)\sqrt{\sigma_{\max}}}{\sigma_{\min}} + \sqrt{\frac{\mu r}{n}} \frac{\sigma\sqrt{n}\log^{2.5}(n)\sqrt{\sigma_{\max} r}}{\sigma_{\min}} \\ &\lesssim \frac{\sigma\mu r^3\kappa^{1.5}\log^{3.5}(n)}{\sqrt{\sigma_{\min}}}. \end{aligned}$$

Similarly, one can also obtain that $\|(I - VV^\top)\Delta_Y\|_{2,\infty} \lesssim \frac{\sigma\mu r^3\kappa^{1.5}\log^{3.5}(n)}{\sqrt{\sigma_{\min}}}$. This completes the proof for Eq. (60a):

$$\begin{aligned} \|P_{T^\perp}(M^*)\|_\infty &\lesssim \frac{\sigma\mu r^3\kappa^{1.5}\log^{3.5}(n)}{\sqrt{\sigma_{\min}}} \cdot \frac{\sigma\mu r^3\kappa^{1.5}\log^{3.5}(n)}{\sqrt{\sigma_{\min}}} \\ &\lesssim \frac{\sigma^2\mu^2 r^6\kappa^3\log^7(n)}{\sigma_{\min}}. \end{aligned}$$

I.6 Proof of Eq. (60b)

Recall that we want to show $\|P_{T_t^\perp}(E) - P_{T^{*\perp}}(E)\|_\infty \leq C_{T,2} \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^2 \log^4(n)}{\sigma_{\min}}$.

Let $U'\Sigma V'^\top$ be the SVD of $X^t Y^{tT}$, $U^*\Sigma^* V^{*\top}$ be the SVD of $X^* Y^{*\top}$. By Eq. (59c), Eq. (59f), and Eq. (59g), we can invoke Lemma 15 and obtain that there exists a rotation matrix $R \in \mathcal{O}^{r \times r}$ such that

$$\begin{aligned} \max(\|U'R - U^*\|_F, \|V'R - V^*\|_F) &\lesssim \frac{r\kappa\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \\ \max(\|U'R - U^*\|_{2,\infty}, \|V'R - V^*\|_{2,\infty}) &\lesssim \frac{r^3\mu\kappa^{1.5}\sigma\log^{3.5}(n)}{\sigma_{\min}}. \end{aligned}$$

One particular choice of R is $R = U_Q V_Q^\top$ where $U_Q \Sigma_Q V_Q^\top$ is the SVD of $\Sigma^{-1/2} U'^\top X^t H^t$.

Let $U = U'R, V = V'R$ for simplification. Note that

$$\begin{aligned} &\|P_{T_t^\perp}(E) - P_{T^{*\perp}}(E)\|_\infty \\ &\leq \|(I - UU^\top)E(I - VV^\top) - (I - U^*U^{*\top})E(I - V^*V^{*\top})\|_\infty \\ &\leq \|(U^*U^{*\top} - UU^\top)E(I - VV^\top)\|_\infty + \|(I - U^*U^{*\top})E(V^*V^{*\top} - VV^\top)\|_\infty. \end{aligned}$$

We have

$$\begin{aligned}
& \|(U^*U^{*\top} - UU^\top)E(I - VV^\top)\|_\infty \\
& \leq \|(U^*U^{*\top} - UU^\top)E\|_\infty + \|(U^*U^{*\top} - UU^\top)EVV^\top\|_\infty \\
& \leq \|(U^* - U)U^{*\top}E\|_\infty + \|U(U^* - U)^\top E\|_\infty \\
& \quad + \|(U^* - U)U^{*\top}EVV^\top\|_\infty + \|U(U^* - U)^\top EVV^\top\|_\infty \\
& \leq \underbrace{\|(U^* - U)\|_{2,\infty} \|E^\top U^*\|_{2,\infty}}_{A_0} + \underbrace{\|U\|_{2,\infty} \|E^\top (U^* - U)\|_{2,\infty}}_{A_1} \\
& \quad + \underbrace{\|(U^* - U)\|_{2,\infty} \|U^*\| \|E\| \|V\| \|V\|_{2,\infty} + \|U\|_{2,\infty} \|U - U^*\| \|E\| \|V\| \|V\|_{2,\infty}}_{A_2}.
\end{aligned}$$

For A_2 , given that $\|E\| \lesssim \sigma\sqrt{n}$, Eq. (70) and Eq. (71), it is easy to check that

$$A_2 \lesssim \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^{1.5} \log^{3.5}(n)}{\sigma_{\min}}.$$

For A_0 , one can verify that $\|E^\top U^*\|_{2,\infty} \lesssim \sigma\sqrt{r \log(n)}$ with probability $1 - O(n^{-10^6})$ since E and U^* are independent. This provides

$$A_0 \lesssim \frac{\sigma^2 r^{3.5} \mu \kappa^{1.5} \log^4(n)}{\sigma_{\min}}.$$

For A_1 , we intend to use the leave-one-out technique. Let $U^{(l)}\Sigma^{(l)}V^{(l)}$ be the SVD of $X^{(l),t}(Y^{(l),t})^\top$. By Lemma 15, we can obtain that there exists a rotation matrix $R^{(l)}$ such that

$$\|U^{(l)}R^{(l)} - U^*\|_{\text{F}} \lesssim \frac{r\kappa\sigma\sqrt{n} \log^{2.5}(n)}{\sigma_{\min}}.$$

In particular, $R^{(l)} = U_{Q^{(l)}}V_{Q^{(l)}}^\top$ where $U_{Q^{(l)}}\Sigma_{Q^{(l)}}V_{Q^{(l)}}^\top$ is the SVD of $\Sigma^{(l)-1/2}U^{(l)\top}X^{(l),t}H^{(l),t}$

Let $U^{(l)} = U^{(l)}R^{(l)}$. In fact, due to the construction of R and $R^{(l)}$, by Lemma 15, one can further verify that

$$\|U^{(l)}R^{(l)} - U^*R\|_{\text{F}} \lesssim \frac{\sqrt{\kappa}\sqrt{r}}{\sqrt{\sigma_{\min}}} \|F^t H^t - F^{t,(l)} H^{t,(l)}\|_{\text{F}} \stackrel{(i)}{\lesssim} \frac{\sigma r \kappa^2 \log^{1.5}(n)}{\sigma_{\min}}.$$

Here, (i) is due to Eq. (69). Then for $n < l \leq 2n$, we have with probability $1 - O(n^{10^{-6}})$,

$$\begin{aligned}
\|(E^\top(U^* - U))_{l-n,\cdot}\| & \leq \|(E^\top(U^{(l)} - U^*))_{l-n,\cdot}\| + \|E^\top(U^{(l)} - U)\|_{2,\infty} \\
& \stackrel{(i)}{\lesssim} \sigma \|U^{(l)} - U^*\|_{\text{F}} \sqrt{\log(n)} + \sigma\sqrt{n} \|U^{(l)} - U\|_{\text{F}} \\
& \lesssim \frac{\sigma^2 \sqrt{n} r \kappa^2 \log^3(n)}{\sigma_{\min}}
\end{aligned}$$

where (i) is due to the independence between E_{l-n}^\top and $U^{(l)}, U^*$. This implies that

$$\begin{aligned}
A_1 & \lesssim \sqrt{\frac{\mu r}{n}} \|E^\top(U^* - U)\|_{2,\infty} \\
& \lesssim \frac{\sigma^2 r^{1.5} \mu^{0.5} \kappa^2 \log^{3.5}(n)}{\sigma_{\min}}.
\end{aligned}$$

In conclusion,

$$\begin{aligned}
\|(U^*U^{*\top} - UU^\top)E(I - VV^\top)\|_\infty & \leq A_0 + A_1 + A_2 \\
& \lesssim \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^2 \log^4(n)}{\sigma_{\min}}.
\end{aligned}$$

This similar results hold for $\|(I - U^*U^{*\top})E(V^*V^{*\top} - VV^\top)\|_\infty$. Therefore,

$$\|P_{T_t^\perp}(E) - P_{T^{*\perp}}(E)\|_\infty \lesssim \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^2 \log^4(n)}{\sigma_{\min}},$$

which completes the proof.

I.7 Proof of Eq. (60c)

Follow the derivation for the proof of Eq. (60b), we have

$$\begin{aligned} & \|P_{T^\perp}(Z) - P_{T^{*\perp}}(Z)\|_F \\ & \leq \|(U^*U^{*\top} - UU^\top)Z(I - VV^\top)\|_F + \|(I - U^*U^{*\top})Z(V^*V^{*\top} - VV^\top)\|_F \\ & \leq \|U^*U^{*\top} - UU^\top\|_F \|Z\| + \|V^*V^{*\top} - VV^\top\|_F \|Z\| \\ & \lesssim (\|U - U^*\|_F + \|V - V^*\|_F) \|Z\| \\ & \lesssim \frac{r\kappa\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \|Z\|. \end{aligned}$$

For the entrywise norm bound, we also have

$$\begin{aligned} & \|P_{T^\perp}(Z) - P_{T^{*\perp}}(Z)\|_\infty \\ & \leq \|(U^*U^{*\top} - UU^\top)Z(I - VV^\top)\|_\infty + \|(I - U^*U^{*\top})Z(V^*V^{*\top} - VV^\top)\|_\infty. \end{aligned}$$

Similar to the bounds on Eq. (60b), note that

$$\begin{aligned} & \|(U^*U^{*\top} - UU^\top)Z(I - VV^\top)\|_\infty \\ & \leq \|(U^*U^{*\top} - UU^\top)Z\|_\infty + \|(U^*U^{*\top} - UU^\top)ZVV^\top\|_\infty \\ & \leq \|(U^* - U)U^{*\top}Z\|_\infty + \|U(U^* - U)^\top Z\|_\infty \\ & \quad + \|(U^* - U)U^{*\top}ZVV^\top\|_\infty + \|U(U^* - U)^\top ZVV^\top\|_\infty \\ & \leq \|U^* - U\|_{2,\infty} \|Z^\top\|_{2,\infty} \|U^*\| + \|U\|_{2,\infty} \|Z^\top\|_{2,\infty} \|U^* - U\| \\ & \quad + \|U^* - U\|_{2,\infty} \|U^*\| \|Z\| \|V\| \|V\|_{2,\infty} + \|U\|_{2,\infty} \|U - U^*\| \|Z\| \|V\| \|V\|_{2,\infty}. \end{aligned}$$

Based on

$$\begin{aligned} \max(\|U^* - U\|_{2,\infty}, \|V^* - V\|_{2,\infty}) & \lesssim \frac{r^3 \mu \kappa^{1.5} \sigma \log^{3.5}(n)}{\sigma_{\min}} \\ \max(\|U^* - U\|_F, \|V^* - V\|_F) & \lesssim \frac{r\kappa\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}}, \end{aligned}$$

we have

$$\begin{aligned} & \|(U^*U^{*\top} - UU^\top)Z(I - VV^\top)\|_\infty \\ & \lesssim \|Z\| \sqrt{\frac{\mu r}{n}} \frac{r^3 \mu \kappa^{1.5} \sigma \log^{3.5}(n)}{\sigma_{\min}} + \|Z^\top\|_{2,\infty} \frac{r^3 \mu \kappa^{1.5} \sigma \log^{3.5}(n)}{\sigma_{\min}} \\ & \lesssim \|Z^\top\|_{2,\infty} \frac{r^{3.5} \kappa^{1.5} \sigma \mu^{1.5} \log^{3.5}(n)}{\sigma_{\min}}. \end{aligned}$$

Similarly, one can obtain the bounds for $\|(I - U^*U^{*\top})Z(V^*V^{*\top} - VV^\top)\|_\infty$. This implies that

$$\|P_{T^\perp}(Z) - P_{T^{*\perp}}(Z)\|_\infty \lesssim \frac{r^{3.5} \kappa^{1.5} \sigma \mu^{1.5} \log^{3.5}(n)}{\sigma_{\min}} \left(\|Z\|_{2,\infty} + \|Z^\top\|_{2,\infty} \right).$$

J Proof of Theorem 1

To begin, let $E' = \delta + E$. We show a few properties about E' . Note that with probability $1 - O(n^{-C})$, we have $\|E\| \lesssim \sigma\sqrt{n}$. Hence

$$\|E'\| \leq \|\delta\| + \|E\| \lesssim \sigma\sqrt{n}.$$

Next, note that $\langle \delta, Z \rangle = 0$ by definition, then

$$|\langle Z, E' \rangle| \leq |\langle Z, \delta \rangle| + |\langle Z, E \rangle| \stackrel{(i)}{\lesssim} \sigma \sqrt{\log(n)} \|Z\|_{\mathbb{F}}$$

where (i) holds with probability $1 - O(n^{-C})$ by Chernoff bound (note that the sum of independent sub-Gaussian random variables is still sub-Gaussian).

Furthermore, consider

$$\begin{aligned} |\langle P_{T^*}(Z), E' \rangle| &\leq |\langle P_{T^*}(Z), E \rangle| + |\langle P_{T^*}(Z), \delta \rangle| \\ &\stackrel{(i)}{\lesssim} |\langle P_{T^*}(Z), E \rangle| + \sigma \log(n) \|Z\|_{\mathbb{F}} \\ &\stackrel{(ii)}{\lesssim} \sigma \sqrt{\log(n)} \|P_{T^*}(Z)\|_{\mathbb{F}} + \sigma \log(n) \|Z\|_{\mathbb{F}} \\ &\lesssim \sigma \log(n) \|Z\|_{\mathbb{F}} \end{aligned}$$

where (i) is due to the assumption of the theorem, (ii) holds with probability $1 - O(n^{-C})$ by Chernoff bound. This also implies

$$\begin{aligned} |\langle P_{T^*\perp}(Z), E' \rangle| &\leq |\langle P_{T^*}(Z), E' \rangle| + |\langle Z, E' \rangle| \\ &\lesssim \sigma \log(n) \|Z\|_{\mathbb{F}}. \end{aligned} \tag{72}$$

Then given Lemmas 1, 4 and 5, it is easy to finish the proof of Theorem 1. Based on Lemma 5, there exists a $0 \leq t < t^*$, such that

$$\begin{aligned} \|F^t H^t - F^*\|_{\mathbb{F}} &\leq C_{\mathbb{F}} \left(\frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_{\mathbb{F}} \\ \|\nabla f(X^t, Y^t; \tau^t)\|_{\mathbb{F}} &\leq \frac{\lambda \sqrt{\sigma_{\min}}}{\kappa n^{10}}. \end{aligned}$$

Let $X = X^t H^t, Y = Y^t H^t, \tau = \tau^t, T = T^t$. By Lemma 4, for any minimizer $(\hat{M}, \hat{\tau})$ of the convex program, we have $\|XY^{\top} - \hat{M}\|_{\mathbb{F}} \lesssim \frac{\lambda \log(n)}{n^{10}}$. This further implies

$$\begin{aligned} \|M^* - \hat{M}\|_{\mathbb{F}} &\lesssim \|M^* - XY^{\top}\|_{\mathbb{F}} + \|XY^{\top} - \hat{M}\|_{\mathbb{F}} \\ &\lesssim \frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \sqrt{\sigma_{\max} r} \sqrt{\sigma_{\max}} + \frac{\lambda \log(n)}{n^{10}} \\ &\lesssim \sigma \sqrt{n \kappa r^{0.5}} \log^{2.5}(n). \end{aligned}$$

Let $\hat{U} \hat{\Sigma} \hat{V}^{\top}$ be the SVD of \hat{M} . Then by Davis-Kahan theorem, we have

$$\begin{aligned} \max \left(\left\| \hat{U} \hat{U}^{\top} - U^* U^{*\top} \right\|_{\mathbb{F}}, \left\| U^* - \hat{U} \hat{U}^{\top} U^* \right\|_{\mathbb{F}} \right) &\lesssim \frac{\|M^* - \hat{M}\|_{\mathbb{F}}}{\sigma_r(\hat{M})} \\ &\lesssim \frac{\sigma \sqrt{n \kappa r^{0.5}} \log^{2.5}(n)}{\sigma_{\min}}. \end{aligned} \tag{73}$$

Similar bounds for V also hold by symmetry.

Recall by Lemma 1, we have

$$\begin{aligned} |\tau^d - \tau^*| &= \left| \frac{\langle P_{T^*\perp}(Z), E' \rangle}{\|P_{\hat{T}^*\perp}(Z)\|_{\mathbb{F}}^2} + \frac{\langle Z, P_{\hat{T}^*\perp}(E') - P_{T^*\perp}(E') \rangle + \langle Z, P_{\hat{T}^*\perp}(M^*) \rangle}{\|P_{\hat{T}^*\perp}(Z)\|_{\mathbb{F}}^2} \right| \\ &\leq \underbrace{\left| \frac{\langle P_{T^*\perp}(Z), E' \rangle}{\|P_{\hat{T}^*\perp}(Z)\|_{\mathbb{F}}^2} \right|}_{A_0} + \underbrace{\left| \frac{\langle Z, P_{\hat{T}^*\perp}(E') - P_{T^*\perp}(E') \rangle}{\|P_{\hat{T}^*\perp}(Z)\|_{\mathbb{F}}^2} \right|}_{A_1} + \underbrace{\left| \frac{\langle Z, P_{\hat{T}^*\perp}(M^*) \rangle}{\|P_{\hat{T}^*\perp}(Z)\|_{\mathbb{F}}^2} \right|}_{A_2}. \end{aligned}$$

For controlling A_0 , note that $|\langle P_{T^*\perp}(Z), E \rangle| \lesssim \sigma \log(n) \|Z\|_F$ by Eq. (72). We also have the facts that $\log(n) \|P_{\hat{T}^\perp}(Z)\|_F^2 \gtrsim \|Z\|_F^2$ by Lemma 16. This implies

$$\begin{aligned} A_0 &= \left| \frac{\langle P_{T^*\perp}(Z), E' \rangle}{\|P_{\hat{T}^\perp}(Z)\|_F^2} \right| \\ &\lesssim \frac{\sigma \log^2(n) \|Z\|_F}{\|Z\|_F^2} \\ &\lesssim \frac{\sigma \log^2(n)}{\|Z\|_F}. \end{aligned}$$

Next, we control A_1 . Note that

$$\begin{aligned} \|P_{\hat{T}^\perp}(E) - P_{T^*\perp}(E')\|_F &\leq \|(U^*U^{*\top} - UU^\top)E'(I - VV^\top)\|_F + \|(I - U^*U^{*\top})E'(V^*V^{*\top} - VV^\top)\|_F \\ &\leq \|U^*U^{*\top} - UU^\top\|_F \|E'\| + \|V^*V^{*\top} - VV^\top\|_F \|E'\| \\ &\stackrel{(i)}{\lesssim} \frac{\sigma\sqrt{n}\kappa r^{0.5} \log^{2.5}(n)}{\sigma_{\min}} \sigma\sqrt{n} \\ &\lesssim \frac{\sigma^2 n \kappa r^{0.5} \log^{2.5}(n)}{\sigma_{\min}} \end{aligned}$$

where (i) is due to Eq. (73) and $\|E'\| \lesssim \sigma\sqrt{n}$. Then we have

$$\begin{aligned} A_1 &= \frac{|\langle Z, P_{\hat{T}^\perp}(E') - P_{T^*\perp}(E') \rangle|}{\|P_{\hat{T}^\perp}(Z)\|_F^2} \\ &\leq \frac{\|P_{\hat{T}^\perp}(E') - P_{T^*\perp}(E')\|_F \|Z\|_F}{\|P_{\hat{T}^\perp}(Z)\|_F^2} \\ &\lesssim \frac{\sigma^2 n \kappa r^{0.5} \log^{2.5}(n) \|Z\|_F \log(n)}{\sigma_{\min} \|Z\|_F^2} \\ &\lesssim \frac{\sigma^2 n \kappa r^{0.5} \log^{3.5}(n)}{\sigma_{\min}} \frac{1}{\|Z\|_F}. \end{aligned}$$

Finally, we can control A_2 .

$$\begin{aligned} A_2 &= \frac{|\langle Z, P_{\hat{T}^\perp}(M^*) \rangle|}{\|P_{\hat{T}^\perp}(Z)\|_F^2} \\ &\leq \frac{\|P_{\hat{T}^\perp}(M^*)\|_F \|Z\|_F}{\|P_{\hat{T}^\perp}(Z)\|_F^2} \\ &= \left\| (U^* - \hat{U}\hat{U}^\top U^*) \Sigma^* (V^* - \hat{V}\hat{V}^\top V^*)^\top \right\| \frac{\|Z\|_F}{\|P_{\hat{T}^\perp}(Z)\|_F^2} \\ &\lesssim \|U^* - \hat{U}\hat{U}^\top U^*\|_F \|\Sigma^*\| \|V^* - \hat{V}\hat{V}^\top V^*\|_F \frac{\log(n)}{\|Z\|_F} \\ &\lesssim \frac{\sigma^2 n \kappa^3 r \log^6(n)}{\sigma_{\min}} \frac{1}{\|Z\|_F} \end{aligned}$$

Combining A_0, A_1, A_2 together, we have

$$|\tau^d - \tau^*| \lesssim \frac{\sigma}{\|Z\|_F} \left(\frac{\kappa^3 r n \sigma \log^6(n)}{\sigma_{\min}} + \log^2(n) \right)$$

which completes the proof.

K Proof of Theorem 2

In this section, we present the proof of an extended version of Theorem 2, which is stated below.

Theorem 3. *Assume Assumptions 1 to 3 hold. Suppose $O = M^* + \tau^* Z + E$. Suppose*

$$\kappa^4 \mu^2 r^2 \log^2(n) \leq C_1 n \quad \text{and} \quad \frac{\sigma}{\sigma_{\min}} \sqrt{n} \leq C_2 \frac{1}{\kappa^3 \mu r^{4.5} \log^5(n)}.$$

Let $(\hat{M}, \hat{\tau})$ be any minimizer of Eq. (1a) with $\lambda = C_\lambda \sigma \sqrt{n} \log^{1.5}(n)$, and let τ^d be defined as in Eq. (1b) based on $\hat{M}, \hat{\tau}$. Then for any $C_3 > 0$, for sufficiently large n , with probability $1 - O(n^{-C_3})$, we have

$$\tau^d - \tau^* = \frac{\langle P_{T^{*\perp}}(Z), E \rangle}{\|P_{T^{*\perp}}(Z)\|_F^2} + \delta, \quad (74)$$

where

$$|\delta| \leq C_\delta \left(\frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^8(n)}{\sigma_{\min}} + \frac{\sqrt{n}}{\|Z\|_F} \frac{r \kappa \sigma^2 \log^4(n)}{\sigma_{\min}} \right).$$

Here, $C_\lambda, C_\delta, C_\tau, C_\infty$ are constants depending (polynomially) on $C_1, C_2, C_3, C_{r_1}, C_{r_2}$ (where C_{r_1} and C_{r_2} are the constants in Assumptions 3(a) and 3(b)).

Given Lemmas 1 and 2 to 4, it is easy to finish the proof of Theorem 3. Based on Lemma 2, there exists a $0 \leq t < t^*$, such that

$$\begin{aligned} \|F^t H^t - F^*\|_F &\leq C_F \left(\frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \right) \|F^*\|_F \\ \|F^t H^t - F^*\|_{2,\infty} &\leq C_\infty \frac{\sigma \mu r^{2.5} \kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F \\ |\tau^t - \tau^*| &\leq C_\tau \left(\frac{\sigma \mu r^{2.5} \kappa \log^{3.5} n}{\sqrt{n}} + \frac{\sigma \log^{1.5} n}{\|Z\|_F} \right) \\ \|\nabla f(X^t, Y^t; \tau^t)\|_F &\leq \frac{\lambda \sqrt{\sigma_{\min}}}{n^{10}}. \end{aligned}$$

Furthermore,

$$\begin{aligned} \|P_{T_t^\perp}(M^*)\|_\infty &\leq C_{T,1} \frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^7(n)}{\sigma_{\min}} \\ \|P_{T_t^\perp}(E) - P_{T^{*\perp}}(E)\|_\infty &\leq C_{T,2} \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^2 \log^4(n)}{\sigma_{\min}} \\ \|P_{T_t^\perp}(Z) - P_{T^{*\perp}}(Z)\|_F &\leq C_{T,3} \frac{r \kappa \sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \|Z\| \\ \|P_{T_t^\perp}(Z) - P_{T^{*\perp}}(Z)\|_\infty &\leq C_{T,4} C_{T,4} \frac{r^{3.5} \kappa^{1.5} \sigma \mu^{1.5} \log^{3.5}(n)}{\sigma_{\min}} \left(\|Z\|_{2,\infty} + \|Z^\top\|_{2,\infty} \right). \end{aligned}$$

Let $X = X^t H^t, Y = Y^t H^t, \tau = \tau^t, T = T^t$. By Lemma 4, for any minimizer $(\hat{M}, \hat{\tau})$ of the convex program, we have

$$\begin{aligned} \|XY^\top - \hat{M}\|_F &\lesssim \frac{\lambda \kappa \log(n)}{n^{10}} \\ |\tau - \hat{\tau}| &\lesssim \frac{\lambda \kappa \log(n)}{n^{10} \|Z\|_F}. \end{aligned}$$

Furthermore, let \hat{T} be the tangent space of \hat{M} , for any $A \in \mathbb{R}^{n \times n}$, we have

$$\begin{aligned} \|P_{\hat{T}^\perp}(A) - P_{T^\perp}(A)\|_F &\lesssim \|A\| \frac{\lambda \kappa^2 \log(n)}{\sigma_{\min} n^{10}} \\ \|P_{\hat{T}^\perp}(XY^\top)\|_F &\lesssim \frac{\sigma^2 \kappa^5 \log^5(n)}{\sigma_{\min}} \frac{1}{n^{19}}. \end{aligned}$$

Given all above information, we can bound $\|\hat{M} - M^*\|_\infty$ by the triangle inequality.

$$\begin{aligned}
\|\hat{M} - M^*\|_\infty &\leq \|\hat{M} - XY^\top\|_F + \|XY^\top - M^*\|_\infty \\
&\lesssim \frac{\lambda\kappa \log(n)}{n^{10}} + \|X - X^*\|_{2,\infty} \|Y\|_{2,\infty} + \|X^*\|_{2,\infty} \|Y - Y^*\|_{2,\infty} \\
&\lesssim \frac{\lambda\kappa \log(n)}{n^{10}} + \frac{\sigma\mu r^{2.5}\kappa \log^{3.5}(n)}{\sigma_{\min}} \|F^*\|_F \|F^*\|_{2,\infty} \\
&\lesssim \frac{\lambda\kappa \log(n)}{n^{10}} + \frac{\sigma\mu r^{2.5}\kappa \log^{3.5}(n)}{\sigma_{\min}} \sqrt{\sigma_{\max} r} \sqrt{\frac{\mu r \sigma_{\max}}{n}} \\
&\lesssim \frac{\sigma\sqrt{n} \log^{2.5}(n)\kappa}{n^{10}} + \frac{\sigma\mu^{1.5} r^{3.5} \kappa^2 \log^{3.5}(n)}{\sqrt{n}} \\
&\lesssim \frac{\sigma\mu^{1.5} r^{3.5} \kappa^2 \log^{3.5}(n)}{\sqrt{n}}.
\end{aligned}$$

Similarly, by the triangle inequality, we can bound $|\hat{\tau} - \tau^*|$.

$$\begin{aligned}
|\hat{\tau} - \tau^*| &\leq |\hat{\tau} - \tau| + |\tau - \tau^*| \\
&\lesssim \frac{\lambda\kappa \log(n)}{n^{10} \|Z\|_F} + \left(\frac{\sigma\mu r^{2.5}\kappa \log^{3.5} n}{\sqrt{n}} + \frac{\sigma \log^{1.5} n}{\|Z\|_F} \right) \\
&\lesssim \frac{\sigma\mu r^{2.5}\kappa \log^{3.5} n}{\sqrt{n}} + \frac{\sigma \log^{1.5} n}{\|Z\|_F}.
\end{aligned}$$

Finally, we aim to provide a characterization for $\tau^d := \hat{\tau} - \frac{\langle Z, \hat{U}\hat{V}^\top \rangle}{\|P_{\hat{T}^\perp}(Z)\|_F^2}$ where $\hat{M} = \hat{U}\hat{\Sigma}\hat{V}^\top$ is the SVD of \hat{M} .

By the triangle inequality again, we can obtain the bounds for \hat{T} from T . We have

$$\begin{aligned}
&\|P_{\hat{T}^\perp}(M^*)\|_\infty \\
&\leq \|P_{\hat{T}^\perp}(M^*) - P_{T^\perp}(M^*)\|_\infty + \|P_{T^\perp}(M^*)\|_\infty \\
&\stackrel{(i)}{\leq} \|P_{\hat{T}^\perp}(M^* - XY^\top + XY^\top) - P_{T^\perp}(M^* - XY^\top)\|_\infty + \|P_{T^\perp}(M^*)\|_\infty \\
&\leq \|P_{\hat{T}^\perp}(M^* - XY^\top) - P_{T^\perp}(M^* - XY^\top)\|_F + \|P_{\hat{T}^\perp}(XY^\top)\|_F + \|P_{T^\perp}(M^*)\|_\infty \\
&\stackrel{(ii)}{\lesssim} \|M^* - XY^\top\| \frac{\lambda\kappa^2 \log(n)}{\sigma_{\min} n^{10}} + \frac{\sigma^2 \kappa^5 \log^5(n)}{\sigma_{\min}} \frac{1}{n^{19}} + \frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^7(n)}{\sigma_{\min}} \\
&\stackrel{(iii)}{\lesssim} \sigma\kappa r^{0.5} \sqrt{n} \log^{2.5}(n) \frac{\sigma\sqrt{n}\kappa^2 \log^{2.5}(n)}{\sigma_{\min} n^{10}} + \frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^7(n)}{\sigma_{\min}} \\
&\lesssim \frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^7(n)}{\sigma_{\min}}. \tag{75}
\end{aligned}$$

Here, (i) is due to $P_T^\perp(XY^\top) = 0$, (ii) is due to $\|P_{\hat{T}^\perp}(A) - P_{T^\perp}(A)\|_F \lesssim \|A\| \frac{\lambda\kappa^2 \log(n)}{\sigma_{\min} n^{10}}$ and $\|P_{\hat{T}^\perp}(XY^\top)\|_F \lesssim \frac{\sigma^2 \kappa^5 \log^5(n)}{\sigma_{\min}} \frac{1}{n^{19}}$, and (iii) is due to $\kappa^4 \mu^2 r^2 \log^2(n) \lesssim n$.

Similarly, one can obtain

$$\begin{aligned}
\|P_{\hat{T}^\perp}(E) - P_{T^{\ast\perp}}(E)\|_\infty &\leq \|P_{\hat{T}^\perp}(E) - P_{T^\perp}(E)\|_\infty + \|P_{T^\perp}(E) - P_{T^{\ast\perp}}(E)\|_\infty \\
&\lesssim \|P_{\hat{T}^\perp}(E) - P_{T^\perp}(E)\|_F + \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^{1.5} \log^4(n)}{\sigma_{\min}} \\
&\lesssim \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^{1.5} \log^4(n)}{\sigma_{\min}}. \tag{76}
\end{aligned}$$

Also,

$$\begin{aligned}
\|P_{\hat{T}^\perp}(Z) - P_{T^{*\perp}}(Z)\|_{\mathbb{F}} &\leq \|P_{\hat{T}^\perp}(Z) - P_{T^\perp}(Z)\|_{\mathbb{F}} + \|P_{T^\perp}(Z) - P_{T^{*\perp}}(Z)\|_{\mathbb{F}} \\
&\lesssim \|P_{\hat{T}^\perp}(Z) - P_{T^\perp}(Z)\|_{\mathbb{F}} + \frac{r\kappa\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \|Z\| \\
&\lesssim \frac{r\kappa\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \|Z\|_{\mathbb{F}}.
\end{aligned} \tag{77}$$

In fact, we can also obtain the entrywise norm bounds for $P_{\hat{T}^\perp}(Z) - P_{T^{*\perp}}(Z)$ here (will be used for characterizing the distribution of $M^d - M^*$).

$$\begin{aligned}
&\|P_{\hat{T}^\perp}(Z) - P_{T^{*\perp}}(Z)\|_{\infty} \\
&\leq \|P_{\hat{T}^\perp}(Z) - P_{T^\perp}(Z)\|_{\mathbb{F}} + \|P_{T^\perp}(Z) - P_{T^{*\perp}}(Z)\|_{\infty} \\
&\lesssim \|P_{\hat{T}^\perp}(Z) - P_{T^\perp}(Z)\|_{\mathbb{F}} + C_{T,4} \frac{r^{3.5}\kappa^{1.5}\sigma\mu^{1.5}\log^{3.5}(n)}{\sigma_{\min}} (\|Z\|_{2,\infty} + \|Z^\top\|_{2,\infty}) \\
&\lesssim \|Z\| \frac{\lambda\kappa^2\log(n)}{\sigma_{\min}n^{10}} + \frac{r^{3.5}\kappa^{1.5}\sigma\mu^{1.5}\log^{3.5}(n)}{\sigma_{\min}} (\|Z\|_{2,\infty} + \|Z^\top\|_{2,\infty}) \\
&\lesssim \frac{r^{3.5}\kappa^{1.5}\sigma\mu^{1.5}\log^{3.5}(n)}{\sigma_{\min}} (\|Z\|_{2,\infty} + \|Z^\top\|_{2,\infty}).
\end{aligned} \tag{78}$$

Recall by Lemma 1, we have

$$\begin{aligned}
\tau^d - \tau^* &= \frac{\langle P_{\hat{T}^\perp}(Z), E \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} + \frac{\langle Z, P_{\hat{T}^\perp}(M^*) \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} \\
&\stackrel{(i)}{=} \frac{\langle P_{T^{*\perp}}(Z), E \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} + \frac{\langle Z, P_{\hat{T}^\perp}(E) - P_{T^{*\perp}}(E) \rangle + \langle Z, P_{\hat{T}^\perp}(M^*) \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2}
\end{aligned}$$

where in (i) we use the fact that for any $A, B \in \mathbb{R}^{n \times n}$,

$$\langle A, P_{T^{*\perp}}(B) \rangle = \langle P_{T^{*\perp}}(A), P_{T^{*\perp}}(B) \rangle = \langle P_{T^{*\perp}}(A), B \rangle.$$

Let

$$\delta = \tau^d - \tau^* - \frac{\langle P_{T^{*\perp}}(Z), E \rangle}{\|P_{T^*}(Z)\|_{\mathbb{F}}^2}.$$

The following claim provides the bounds for δ , which completes the proof of Theorem 2.

Claim 12. *Providing Eqs. (75) to (77), we have*

$$|\delta| \lesssim \left(\frac{\sigma^2\mu^2r^6\kappa^3\log^8(n)}{\sigma_{\min}} + \frac{\sqrt{n}}{\|Z\|_{\mathbb{F}}} \frac{r\kappa\sigma^2\log^4(n)}{\sigma_{\min}} \right).$$

K.1 Proof of Claim 12

By the definition of δ , we have

$$\begin{aligned}
|\delta| &= \left| \frac{\langle P_{T^{*\perp}}(Z), E \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} - \frac{\langle P_{T^{*\perp}}(Z), E \rangle}{\|P_{T^*}(Z)\|_{\mathbb{F}}^2} + \frac{\langle Z, P_{\hat{T}^\perp}(E) - P_{T^{*\perp}}(E) \rangle + \langle Z, P_{\hat{T}^\perp}(M^*) \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} \right| \\
&\leq \underbrace{\left| \frac{\langle P_{T^{*\perp}}(Z), E \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} - \frac{\langle P_{T^{*\perp}}(Z), E \rangle}{\|P_{T^*}(Z)\|_{\mathbb{F}}^2} \right|}_{A_0} + \underbrace{\left| \frac{\langle Z, P_{\hat{T}^\perp}(E) - P_{T^{*\perp}}(E) \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} \right|}_{A_1} \\
&\quad + \underbrace{\left| \frac{\langle Z, P_{\hat{T}^\perp}(M^*) \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} \right|}_{A_2}.
\end{aligned}$$

For controlling A_0 , note that $|\langle P_{T^* \perp}(Z), E \rangle| \lesssim \sigma \|P_{T^* \perp}(Z)\|_{\mathbb{F}} \log^{0.5}(n)$ since the sum of independent sub-Gaussian random variables is still sub-Gaussian. We also have the facts that $\log(n) \|P_{T^* \perp}(Z)\|_{\mathbb{F}}^2 \gtrsim \|Z\|_{\mathbb{F}}^2$ by Lemma 16. This implies

$$\|P_{\hat{T} \perp}(Z) - P_{T^* \perp}(Z)\|_{\mathbb{F}} \lesssim \frac{r\kappa\sigma\sqrt{n}\log^{2.5}(n)}{\sigma_{\min}} \|Z\|_{\mathbb{F}} \lesssim \frac{r\kappa\sigma\sqrt{n}\log^3(n)}{\sigma_{\min}} \|P_{T^* \perp}(Z)\|_{\mathbb{F}}.$$

Then due to $\frac{\sigma}{\sigma_{\min}}\sqrt{n} \lesssim \frac{1}{\kappa^3 r^{4.5} \log^5(n)\mu}$, we have

$$\frac{1}{2} \|P_{T^* \perp}(Z)\|_{\mathbb{F}} \leq \|P_{\hat{T} \perp}(Z)\|_{\mathbb{F}} \leq 2 \|P_{T^* \perp}(Z)\|_{\mathbb{F}}.$$

Also, we have

$$\begin{aligned} \left| \frac{1}{\|P_{\hat{T} \perp}(Z)\|_{\mathbb{F}}^2} - \frac{1}{\|P_{T^* \perp}(Z)\|_{\mathbb{F}}^2} \right| &\leq \frac{|\|P_{T^* \perp}(Z)\|_{\mathbb{F}}^2 - \|P_{\hat{T} \perp}(Z)\|_{\mathbb{F}}^2|}{\|P_{T^* \perp}(Z)\|_{\mathbb{F}}^2 \|P_{\hat{T} \perp}(Z)\|_{\mathbb{F}}^2} \\ &\lesssim \frac{\|P_{\hat{T} \perp}(Z) - P_{T^* \perp}(Z)\|_{\mathbb{F}}}{\|P_{T^* \perp}(Z)\|_{\mathbb{F}}^3} \\ &\lesssim \frac{r\kappa\sigma\sqrt{n}\log^3(n)}{\sigma_{\min}} \frac{1}{\|P_{T^* \perp}(Z)\|_{\mathbb{F}}^2}. \end{aligned}$$

Then, we can control A_0 .

$$\begin{aligned} A_0 &\lesssim \sigma \|P_{T^* \perp}(Z)\|_{\mathbb{F}} \log^{0.5}(n) \cdot \frac{r\kappa\sigma\sqrt{n}\log^3(n)}{\sigma_{\min}} \frac{1}{\|P_{T^* \perp}(Z)\|_{\mathbb{F}}^2} \\ &\lesssim \frac{r\kappa\sigma^2\sqrt{n}\log^{3.5}(n)}{\sigma_{\min}} \frac{1}{\|P_{T^* \perp}(Z)\|_{\mathbb{F}}} \\ &\lesssim \frac{r\kappa\sigma^2\sqrt{n}\log^4(n)}{\sigma_{\min}} \frac{1}{\|Z\|_{\mathbb{F}}}. \end{aligned}$$

Next, we control A_1 . Note that $|\langle Z, A \rangle| = |\sum_{ij} A_{ij} Z_{ij}| \leq \|A\|_{\infty} \|Z\|_{\mathbb{F}}^2$ for binary matrix Z . We have

$$\begin{aligned} A_1 &= \frac{|\langle Z, P_{\hat{T} \perp}(E) - P_{T^* \perp}(E) \rangle|}{\|P_{\hat{T} \perp}(Z)\|_{\mathbb{F}}^2} \\ &\leq \frac{\|P_{\hat{T} \perp}(E) - P_{T^* \perp}(E)\|_{\infty} \|Z\|_{\mathbb{F}}^2}{\|P_{\hat{T} \perp}(Z)\|_{\mathbb{F}}^2} \\ &\lesssim \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^{1.5} \log^4(n)}{\sigma_{\min}} \frac{\|Z\|_{\mathbb{F}}^2}{\|P_{\hat{T} \perp}(Z)\|_{\mathbb{F}}^2} \\ &\lesssim \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^{1.5} \log^5(n)}{\sigma_{\min}}. \end{aligned}$$

Similarly, we can control A_2 .

$$\begin{aligned} A_2 &= \frac{|\langle Z, P_{\hat{T} \perp}(M - M^*) \rangle|}{\|P_{\hat{T} \perp}(Z)\|_{\mathbb{F}}^2} \\ &\leq \frac{\|P_{\hat{T} \perp}(M - M^*)\|_{\infty} \|Z\|_{\mathbb{F}}^2}{\|P_{\hat{T} \perp}(Z)\|_{\mathbb{F}}^2} \\ &\lesssim \frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^7(n)}{\sigma_{\min}} \frac{\|Z\|_{\mathbb{F}}^2}{\|P_{\hat{T} \perp}(Z)\|_{\mathbb{F}}^2} \\ &\lesssim \frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^8(n)}{\sigma_{\min}}. \end{aligned}$$

Combining A_0, A_1, A_2 together, we have

$$|\delta| \lesssim \frac{r\kappa\sigma^2\sqrt{n}\log^4(n)}{\sigma_{\min}} \frac{1}{\|Z\|_{\mathbb{F}}} + \frac{\sigma^2\mu^2r^6\kappa^3\log^8(n)}{\sigma_{\min}}$$

which completes the proof.

L Recovery of Counterfactuals

We also studied the following de-biased estimator M^d and derived the entry-wise characterization for $M^d - M^*$ (where the asymptotical normality follows directly). This particular way of de-biasing \hat{M} can be seen from the following lemma.

Lemma 8. *Suppose $(\hat{M}, \hat{\tau})$ is a minimizer of (1a). Let $\hat{M} = \hat{U}\hat{\Sigma}\hat{V}^\top$ be the SVD of \hat{M} , and let \hat{T} denote the tangent space of \hat{M} . Then,*

$$\begin{aligned} \hat{M} - M^* &= -\lambda\hat{U}\hat{V}^\top - \frac{\lambda\langle Z, \hat{U}\hat{V}^\top \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} P_{\hat{T}}(Z) \\ &\quad - \left(\frac{\langle P_{\hat{T}^\perp}(Z), E \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} + \frac{\langle Z, P_{\hat{T}^\perp}(M^*) \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} \right) P_{\hat{T}}(Z) \\ &\quad - P_{\hat{T}^\perp}(M^*) + P_{\hat{T}}(E). \end{aligned}$$

Proof Sketch of Lemma 8. From the first-order conditions of $(\hat{M}, \hat{\tau})$, we have $M^* - \hat{M} + (\tau^* - \hat{\tau})Z + E = \lambda\hat{U}\hat{V}^\top + \lambda W$. Substituting λW with $P_{\hat{T}^\perp}(M^*) + P_{\hat{T}^\perp}(E) - (\hat{\tau} - \tau^*)P_{\hat{T}^\perp}(Z) = \lambda W$ (Eq. (5)), we obtain

$$M^* - \hat{M} + (\tau^* - \hat{\tau})Z + E = \lambda\hat{U}\hat{V}^\top + P_{\hat{T}^\perp}(M^*) + P_{\hat{T}^\perp}(E) - (\hat{\tau} - \tau^*)P_{\hat{T}^\perp}(Z).$$

This implies

$$M^* + P_{\hat{T}}(E) + (\tau^* - \hat{\tau})P_{\hat{T}}(Z) = \hat{M} + \lambda\hat{U}\hat{V}^\top + P_{\hat{T}^\perp}(M^*)$$

Then substituting $\tau^* - \hat{\tau}$ with Lemma 1, we can obtain the desired decomposition. \square

Theorem 4. *Under the same setup in Theorem 2, let $M^d := \hat{M} + \lambda\hat{U}\hat{V}^\top + P_{\hat{T}}(Z) \cdot \lambda\langle Z, \hat{U}\hat{V}^\top \rangle / \|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2$. We have, for all $i, j \in [n]$,*

$$M_{ij}^d - M_{ij}^* = \left\langle P_{T^*}(e_i e_j^\top) - \frac{(P_{T^*}(Z))_{ij}}{\|P_{T^*}(Z)\|_{\mathbb{F}}^2} P_{T^*}(Z), E + \delta \circ Z \right\rangle + \tilde{O}\left(\frac{1}{n}\right)$$

where e_i (e_j) corresponds to the i -th (j -th) standard basis.

Proof. From Lemma 8, we have

$$\begin{aligned} M_{ij}^d - M_{ij}^* &= (P_{T^*}(E))_{ij} - \frac{\langle P_{T^*}(Z), E \rangle}{\|P_{T^*}(Z)\|_{\mathbb{F}}^2} (P_{T^*}(Z))_{ij} \\ &\quad - \underbrace{(P_{\hat{T}^\perp}(M^*))_{ij}}_{A_0} + \underbrace{(P_{\hat{T}}(E) - P_{T^*}(E))_{ij}}_{A_1} \\ &\quad - \underbrace{\left(\frac{\langle P_{\hat{T}^\perp}(Z), E \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} + \frac{\langle Z, P_{\hat{T}^\perp}(M^*) \rangle}{\|P_{\hat{T}^\perp}(Z)\|_{\mathbb{F}}^2} \right)}_{A_2} (P_{\hat{T}}(Z))_{ij} + \frac{\langle P_{T^*}(Z), E \rangle}{\|P_{T^*}(Z)\|_{\mathbb{F}}^2} (P_{T^*}(Z))_{ij}. \end{aligned}$$

Note that $(P_{T^*}(E))_{ij} = \langle P_{T^*}(E), e_i e_j^\top \rangle = \langle E, P_{T^*}(e_i e_j^\top) \rangle$. Hence, $\delta_{ij} = -A_0 + A_1 + A_2$. Then, it is sufficient to control A_0, A_1 and A_2 .

For A_0 , by Eq. (75), we have $|A_0| \leq \|P_{\hat{T}^\perp}(M^*)\|_\infty \lesssim \frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^7(n)}{\sigma_{\min}}$.

For A_1 , by Eq. (76), we have $|A_1| \leq \|P_{\hat{T}}(E) - P_{T^*}(E)\|_\infty = \|P_{\hat{T}^\perp}(E) - P_{T^*\perp}(E)\|_\infty \lesssim \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^{1.5} \log^4(n)}{\sigma_{\min}}$.

In order to control A_2 , we can further decompose A_2 :

$$A_2 = - \underbrace{\left(\frac{\langle P_{\hat{T}^\perp}(Z), E \rangle}{\|P_{\hat{T}^\perp}(Z)\|_F^2} + \frac{\langle Z, P_{\hat{T}^\perp}(M^*) \rangle}{\|P_{\hat{T}^\perp}(Z)\|_F^2} - \frac{\langle P_{T^*\perp}(Z), E \rangle}{\|P_{T^*\perp}(Z)\|_F^2} \right)}_{B_0} (P_{\hat{T}}(Z))_{ij} \\ + \underbrace{\frac{\langle P_{T^*\perp}(Z), E \rangle}{\|P_{T^*\perp}(Z)\|_F^2}}_{B_1} (P_{T^*}(Z) - P_{\hat{T}}(Z))_{ij}.$$

Then it boils down to control B_0 and B_1 . For B_1 , note that we have

$$\|P_{\hat{T}}(Z) - P_{T^*}(Z)\|_\infty = \|P_{\hat{T}^\perp}(Z) - P_{T^*\perp}(Z)\|_\infty \\ \lesssim \frac{r^{3.5} \kappa^{1.5} \sigma \mu^{1.5} \log^{3.5}(n)}{\sigma_{\min}} \left(\|Z\|_{2,\infty} + \|Z^\top\|_{2,\infty} \right)$$

by Eq. (78). We also have $\frac{|\langle P_{T^*\perp}(Z), E \rangle|}{\|P_{T^*\perp}(Z)\|_F^2} \lesssim \frac{\sigma \log(n)}{\|Z\|_F^2}$. Therefore,

$$|B_1| \lesssim \frac{r^{3.5} \kappa^{1.5} \sigma \mu^{1.5} \log^{3.5}(n)}{\sigma_{\min}} \left(\|Z\|_{2,\infty} + \|Z^\top\|_{2,\infty} \right) \cdot \frac{\sigma \log(n)}{\|Z\|_F} \\ \lesssim \frac{r^{3.5} \kappa^{1.5} \sigma^2 \mu^{1.5} \log^{4.5}(n)}{\sigma_{\min}}.$$

For B_0 , note that

$$\|P_{T^*}(Z)\|_\infty \leq \|U^* U^{*T} Z\|_\infty + \|Z V^* V^{*T}\|_\infty + \|U^* U^{*T} Z V^* V^{*T}\|_\infty \\ \leq \|U^*\|_{2,\infty} \|Z^T\|_{2,\infty} \|U^*\| + \|Z\|_{2,\infty} \|V^*\|_{2,\infty} \|V^*\| \\ + \|U^*\|_{2,\infty} \|V^*\|_{2,\infty} \|U^*\| \|V^*\| \|Z\| \\ \lesssim \sqrt{\frac{\mu r}{n}} \left(\|Z\|_{2,\infty} + \|Z^T\|_{2,\infty} \right) + \frac{\mu r}{n} \cdot \|Z\|. \quad (79)$$

By the triangle inequality, one can obtain the similar bound for $\|P_{\hat{T}}(Z)\|_\infty$. Furthermore, let

$$\delta' = \left(\frac{\langle P_{\hat{T}^\perp}(Z), E \rangle}{\|P_{\hat{T}^\perp}(Z)\|_F^2} + \frac{\langle Z, P_{\hat{T}^\perp}(M^*) \rangle}{\|P_{\hat{T}^\perp}(Z)\|_F^2} - \frac{\langle P_{T^*\perp}(Z), E \rangle}{\|P_{T^*\perp}(Z)\|_F^2} \right).$$

We can also obtain that $|\delta'| \lesssim \left(\frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^8(n)}{\sigma_{\min}} + \frac{\sqrt{n}}{\|Z\|_F} \frac{r \kappa \sigma^2 \log^4(n)}{\sigma_{\min}} \right)$ by Claim 12. This implies that

$$|B_0| \leq |\delta'| \|P_{\hat{T}}(Z)\|_\infty \\ \lesssim \left(\sqrt{\frac{\mu r}{n}} \left(\|Z\|_{2,\infty} + \|Z^T\|_{2,\infty} \right) + \frac{\mu r}{n} \cdot \|Z\| \right) \\ \cdot \left(\frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^8(n)}{\sigma_{\min}} + \frac{\sqrt{n}}{\|Z\|_F} \frac{r \kappa \sigma^2 \log^4(n)}{\sigma_{\min}} \right) \\ \lesssim \frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^8(n)}{\sigma_{\min}} \cdot \mu r + \frac{\mu r^2 \kappa \sigma^2 \log^4(n)}{\sigma_{\min}} \sqrt{\mu r} \\ \lesssim \frac{\sigma^2 \mu^3 r^7 \kappa^3 \log^8(n)}{\sigma_{\min}}.$$

Combining the bounds for A_0, A_1, B_0, B_1 together, we arrive at

$$\begin{aligned}
|\delta_{ij}| &\leq |A_0| + |A_1| + |B_0| + |B_1| \\
&\lesssim \frac{\sigma^2 \mu^2 r^6 \kappa^3 \log^7(n)}{\sigma_{\min}} + \frac{\sigma^2 r^{3.5} \mu^{1.5} \kappa^{1.5} \log^4(n)}{\sigma_{\min}} + \frac{r^{3.5} \kappa^{1.5} \sigma^2 \mu^{1.5} \log^{4.5}(n)}{\sigma_{\min}} \\
&\quad + \frac{\sigma^2 \mu^3 r^7 \kappa^3 \log^8(n)}{\sigma_{\min}} \\
&\lesssim \frac{\sigma^2 \mu^3 r^7 \kappa^3 \log^8(n)}{\sigma_{\min}}.
\end{aligned}$$

This completes the proof. \square

M Proof of Proposition 1 and Proposition 2

In this section, we present the proof of Proposition 1 and Proposition 2.

M.1 Proof of Proposition 1

To begin, let M_1, M_2 be the n -by- n matrices depicted below (for $n = 3k, k \in \mathbb{Z}$):

$$M_1 = \begin{bmatrix} -3t \cdot \mathbf{1}_{1 \times \frac{n}{3}} & \mathbf{0}_{1 \times \frac{2n}{3}} \\ \mathbf{1}_{(n-1) \times \frac{n}{3}} & \mathbf{0}_{(n-1) \times \frac{2n}{3}} \end{bmatrix} \quad M_2 = \begin{bmatrix} 3t \cdot \mathbf{1}_{1 \times \frac{n}{3}} & \mathbf{0}_{1 \times \frac{2n}{3}} \\ \mathbf{1}_{(n-1) \times \frac{n}{3}} & \mathbf{0}_{(n-1) \times \frac{2n}{3}} \end{bmatrix}, \quad (80)$$

where $\mathbf{1}_{a \times b}$ is a $a \times b$ block with all ones (similar for $\mathbf{0}$) and $t \in [0, 1/5]$ is a parameter. Furthermore, let Z be the matrix with all ones in the first row:

$$Z = \begin{bmatrix} \mathbf{1}_{1 \times n} \\ \mathbf{0}_{(n-1) \times n} \end{bmatrix}. \quad (81)$$

Then let δ_1, δ_2 be the n -by- n matrices below:

$$\delta_1 = \begin{bmatrix} 2t \cdot \mathbf{1}_{1 \times \frac{n}{3}} & -t \cdot \mathbf{1}_{1 \times \frac{2n}{3}} \\ \mathbf{0}_{(n-1) \times \frac{n}{3}} & \mathbf{0}_{(n-1) \times \frac{2n}{3}} \end{bmatrix} \quad \delta_2 = \begin{bmatrix} -2t \cdot \mathbf{1}_{1 \times \frac{n}{3}} & t \cdot \mathbf{1}_{1 \times \frac{2n}{3}} \\ \mathbf{0}_{(n-1) \times \frac{n}{3}} & \mathbf{0}_{(n-1) \times \frac{2n}{3}} \end{bmatrix} \quad (82)$$

Let $\tau_1^* = 0$ and $\tau_2^* = -2t$. Then one can verify that $\langle \delta_1, Z \rangle = 0, \langle \delta_2, Z \rangle = 0$ and

$$M_1 + \tau_1^* Z + \delta_1 \circ Z = M_2 + \tau_2^* Z + \delta_2 \circ Z,$$

i.e., $(M_1, \tau_1^*, \delta_1)$ and $(M_2, \tau_2^*, \delta_2)$ will be indistinguishable given the observation. Therefore, for any estimator $\hat{\tau}$, with probability at least $1/2$, either $|\hat{\tau} - \tau_1^*| > t$ holds or $|\hat{\tau} - \tau_2^*| > t$ holds.

Next, we examine the value of $\langle P_{T_1}(\delta_1), P_{T_1}(Z) \rangle$ where T_1 is the tangent space of M_1 . Easy to see that the SVD of M_1 is $M_1 = \sigma_1 u_1 v_1^\top$ where

$$v_1 := \sqrt{3} \left[\underbrace{\frac{1}{\sqrt{n}}, \dots, \frac{1}{\sqrt{n}}}_{\frac{n}{3}}, 0, \dots, 0 \right]^\top, \quad u_1 := \sqrt{\frac{n}{n-1+9t^2}} \left[\frac{-3t}{\sqrt{n}}, \frac{1}{\sqrt{n}}, \dots, \frac{1}{\sqrt{n}} \right]^\top$$

Then,

$$\begin{aligned}
P_{T_1}(Z) &= Z v_1 v_1^\top + u_1 u_1^\top Z (I - v_1 v_1^\top) \\
&= \begin{bmatrix} \mathbf{1}_{1 \times \frac{n}{3}} & \mathbf{0}_{1 \times \frac{2n}{3}} \\ \mathbf{0}_{(n-1) \times \frac{n}{3}} & \mathbf{0}_{(n-1) \times \frac{2n}{3}} \end{bmatrix} + u_1 u_1^\top Z (I - v_1 v_1^\top).
\end{aligned}$$

This implies

$$\begin{aligned}
\langle P_{T_1}(\delta_1), P_{T_1}(Z) \rangle &= \langle \delta_1, P_{T_1}(Z) \rangle \\
&= 2t \frac{n}{3} + \langle \delta_1, u_1 u_1^\top Z (I - v_1 v_1^\top) \rangle \\
&\leq \frac{2t}{3} n + \|u_1^\top \delta_1\| \|u_1^\top Z\| \|I - v_1 v_1^\top\| \\
&\stackrel{(i)}{\leq} tn.
\end{aligned}$$

where (i) is due to $t \leq 1/5$. Similarly, one can show that $\langle P_{T_2}(\delta_2), P_{T_2}(Z) \rangle \leq tn$. Note that $\|Z\|_{\mathbb{F}}^2 = n$. Then, with probability $1/2$, there exists $i \in \{1, 2\}$ such that

$$|\hat{\tau} - \tau_i^*| \geq \frac{\langle P_{T_i}(\delta_i), P_{T_i}(Z) \rangle}{\|Z\|_{\mathbb{F}}^2}$$

After introducing the noise E where $E_{ij} \sim \mathcal{N}(0, \sigma)$, it is also easy to see that $|\hat{\tau} - \tau^*| \geq C \frac{\sigma}{\|Z\|_{\mathbb{F}}^2}$ even when M^* and δ are known. In addition, it is easy to verify that $r = \kappa = 1, \mu = O(1), \sigma_{\min} = \Theta(n)$ for both M_1 and M_2 . Furthermore, $\|\delta_1\|, \|\delta_2\| = O(\sqrt{n})$ and Assumption 3 hold for both M_1, M_2 and Z . This then shows that Theorem 1 achieves the mini-max lower bound in this scenario.

M.2 Proof of Proposition 2

Consider the following simple example: fix any even-valued n , and let Z, M_1, M_2 be the n -by- n matrices depicted below:

$$Z = \begin{bmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{bmatrix} \quad M_1 = \begin{bmatrix} -\mathbf{1} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \quad M_2 = \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{bmatrix}, \quad (83)$$

where the boldfaced $\mathbf{1}$ represents the $n/2$ -by- $n/2$ matrix of all ones (and similarly for $\mathbf{0}$ and $-\mathbf{1}$).

Then, $Z + M_1 = M_2$ immediately follows. It is also easy to see that both M_1 and M_2 satisfy $r, \kappa = 1, \mu = O(1), \sigma_{\min} = \Theta(n)$.

Next, consider the SVD of M_1 : $M_1 = \sigma_1 u_1 v_1^\top$ where

$$v_1 := \sqrt{2} \left[\underbrace{\frac{1}{\sqrt{n}}, \dots, \frac{1}{\sqrt{n}}}_{\frac{n}{2}}, 0, \dots, 0 \right]^\top, \quad u_1 := \sqrt{2} \left[\underbrace{\frac{1}{\sqrt{n}}, \dots, \frac{1}{\sqrt{n}}}_{\frac{n}{2}}, 0, \dots, 0 \right]^\top$$

Then, one can verify that

$$\|Zv_1\|_{\mathbb{F}}^2 + \|Z^\top u_1\|_{\mathbb{F}}^2 = \frac{n}{4} + \frac{n}{4} = \|Z\|_{\mathbb{F}}^2.$$

Furthermore, let T_1 be the tangent space of M_1 . Note that $P_{T_1^\perp}(Z) = \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{bmatrix}$. Then,

$$|\langle Z, u_1 v_1^\top \rangle| \|P_{T_1^\perp}(Z)\| = \frac{n}{2} \cdot \frac{n}{2} = \frac{n^2}{4} = \|P_{T_1^\perp}(Z)\|_{\mathbb{F}}^2.$$

Similar equalities hold for M_2 by symmetry. This completes the proof.

N Technical Lemmas

Lemma 9 (Direct implication of Assumption 3(a)). *Given $Z \in \mathbb{R}^{n \times n}, U \in \mathbb{R}^{n \times r}, V \in \mathbb{R}^{n \times r}$ where $U^\top U = V^\top V = I_r$. Suppose*

$$\|ZV\|_{\mathbb{F}}^2 + \|Z^\top U\|_{\mathbb{F}}^2 \leq (1 - \eta) \|Z\|_{\mathbb{F}}^2 \quad (84)$$

for some $0 < \eta \leq 1$. Let $P_{T^\perp}(Z) = (I - UU^\top)Z(I - VV^\top)$. Then

$$\|U^\top ZV\|_{\mathbb{F}}^2 \leq \|P_{T^\perp}(Z)\|_{\mathbb{F}}^2 - \eta \|Z\|_{\mathbb{F}}^2.$$

This also implies $\|P_{T^\perp}(Z)\|_{\mathbb{F}}^2 \geq \eta \|Z\|_{\mathbb{F}}^2$ and $\|U^\top ZV\|_{\mathbb{F}} \leq (1 - \frac{\eta}{2}) \|P_{T^\perp}(Z)\|_{\mathbb{F}}$.

Proof. Note that $\|U^\top ZV\|_{\mathbb{F}}^2 = \text{tr}(UU^\top ZVV^\top Z^\top), \|P_{T^\perp}(Z)\|_{\mathbb{F}}^2 = \text{tr}((I - UU^\top)Z(I - VV^\top)Z^\top)$ since $\|A\|_{\mathbb{F}}^2 = \text{tr}(AA^\top)$ and $I - UU^\top, I - VV^\top$ are projection matrices. Then

$$\begin{aligned} \|P_{T^\perp}(Z)\|_{\mathbb{F}}^2 - \|U^\top ZV\|_{\mathbb{F}}^2 &= \text{tr}((I - UU^\top)Z(I - VV^\top)Z^\top) - \text{tr}(UU^\top ZVV^\top Z^\top) \\ &= \text{tr}(ZZ^\top) - \text{tr}(UU^\top ZZ^\top) - \text{tr}(ZVV^\top Z^\top) \\ &= \|Z\|_{\mathbb{F}}^2 - \|Z^\top U\|_{\mathbb{F}}^2 - \|ZV\|_{\mathbb{F}}^2 \\ &\geq \eta \|Z\|_{\mathbb{F}}^2 \end{aligned}$$

where the last inequality is by Eq. (84). To show $\|U^\top ZV\|_F \leq (1 - \frac{\eta}{2}) \|P_{T^\perp}(Z)\|_F$, note that

$$\begin{aligned} \left(1 - \frac{\eta}{2}\right)^2 \|P_{T^\perp}(Z)\|_F^2 &\geq (1 - \eta) \|P_{T^\perp}(Z)\|_F^2 \\ &\geq \|P_{T^\perp}(Z)\|_F^2 - \eta \|Z\|_F^2 \\ &\geq \|U^\top ZV\|_F^2. \end{aligned}$$

This finishes the proof. \square

Lemma 10. *Let $X^*, Y^* \in \mathbb{R}^{n \times r}$. Denote the i -th largest singular values of A by $\sigma_i(A)$. Suppose $\sigma_1(X^*) = \sigma_1(Y^*) = \sqrt{\sigma_{\max}}$, $\sigma_r(X^*) = \sigma_r(Y^*) = \sqrt{\sigma_{\min}}$ and $\kappa = \sigma_{\max}/\sigma_{\min}$. Consider $X, Y \in \mathbb{R}^{n \times r}$. Suppose there exists two constant C_F, C_2 such that*

$$\|X - X^*\|_F + \|Y - Y^*\|_F \leq C_F \left(\frac{\sigma \sqrt{n} \log^{2.5}(n)}{\sigma_{\min}} \right) \sqrt{\sigma_{\max} r}$$

and $\frac{\sigma \sqrt{n}}{\sigma_{\min}} \leq C_2 \frac{1}{\kappa^{0.5} r^{0.5} \log^3(n)}$. Then there exists N_0 , for $n \geq N_0$, one has for any $i \in [r]$,

$$\begin{aligned} \sigma_i(X) &\in \left[\sqrt{\frac{\sigma_{\min}}{2}}, \sqrt{2\sigma_{\max}} \right] \\ \sigma_i(Y) &\in \left[\sqrt{\frac{\sigma_{\min}}{2}}, \sqrt{2\sigma_{\max}} \right] \\ \sigma_i(XY^\top) &\in \left[\frac{\sigma_{\min}}{2}, 2\sigma_{\max} \right]. \end{aligned}$$

Proof. Note that $\|X^*\|_F \leq \|X^*\| \sqrt{r} = \sqrt{\sigma_{\max} r}$. Similarly, $\|Y^*\|_F \leq \sqrt{\sigma_{\max} r}$. Then one has

$$\begin{aligned} \|X - X^*\|_F + \|Y - Y^*\|_F &\leq C_F \frac{\sigma \sqrt{n}}{\sigma_{\min}} \log^{2.5}(n) \sqrt{\sigma_{\max} r} \\ &\leq C_F C_2 \frac{1}{\kappa^{0.5} r^{0.5} \log^3(n)} \log^{2.5}(n) \sqrt{\sigma_{\max} r} \\ &= C_F C_2 \frac{\sqrt{\sigma_{\min}}}{\log^{0.5}(n)}. \end{aligned}$$

When n goes large enough, $\|X - X^*\|_F \leq \sqrt{\frac{\sigma_{\min}}{8}}$, $\|Y - Y^*\|_F \leq \sqrt{\frac{\sigma_{\min}}{8}}$. Then, by Weyl's inequality,

$$\sigma_i(X) \leq \sigma_i(X^*) + \|X - X^*\| \leq \sigma_1(X^*) + \|X - X^*\|_F \leq \sqrt{2\sigma_{\max}}. \quad (85)$$

We also have $\sigma_i(X) \geq \sigma_i(X^*) - \|X - X^*\| \geq \sigma_r(X^*) - \|X - X^*\|_F \geq \sqrt{\frac{\sigma_{\min}}{2}}$. The similar results hold for $\sigma_i(Y)$: $\sigma_i(Y) \in \left[\sqrt{\frac{\sigma_{\min}}{2}}, \sqrt{2\sigma_{\max}} \right]$.

Then consider the bounds for $\sigma_i(XY^\top)$ where $i \in [r]$. Note that $\sigma_1(XY^\top) = \|XY^\top\| \leq \|X\| \|Y\| = \sigma_1(X) \sigma_1(Y) \leq 2\sigma_{\max}$. Furthermore, by the properties of SVD, for any $A \in \mathbb{R}^{n \times m}$ with rank r , we have

$$\sigma_r(A) = \min_{\|u\|=1, u \in \text{rowspan}(A)} \|Au\|$$

Here, $\text{rowspan}(A)$ is the row space of A . This implies

$$\begin{aligned} \sigma_r(XY^\top) &= \min_{\|u\|=1, u \in \text{rowspan}(XY^\top)} \|XY^\top u\| \\ &\stackrel{(i)}{\geq} \min_{\|u\|=1, u \in \text{rowspan}(XY^\top)} \|Y^\top u\| \\ &\stackrel{(ii)}{\geq} \sigma_r(X) \cdot \min_{\|u\|=1, u \in \text{rowspan}(Y^\top)} \|Y^\top u\| \\ &\geq \sigma_r(X) \sigma_r(Y) \\ &\geq \frac{\sigma_{\min}}{2} \end{aligned}$$

where (i) is due to $Y^\top u \in \text{rowspan}(X)$ since $X \in \mathbb{R}^{n \times r}$ has rank- r and (ii) is due to $\text{rowspan}(XY^\top) \subset \text{rowspan}(Y^\top)$. This completes the proof. \square

Lemma 11 (Orthogonal Procrustes Problem, Lemma 35 in [39]). *For $F_0, F_1 \in \mathbb{R}^{n \times m}$, H is the minimizer of the following optimization problem if and only if $F_0^\top F_1 H$ is symmetric and positive semidefinite:*

$$\underset{A \in \mathcal{O}^{r \times r}}{\text{minimize}} \|F_1 A - F_0\|_{\text{F}}.$$

Lemma 12. *Let $F_0, F_1, F_2 \in \mathbb{R}^{2n \times r}$ be three matrices. Suppose*

$$\|F_1 - F_0\| \|F_0\| \leq \sigma_r^2(F_0)/2 \quad \text{and} \quad \|F_1 - F_2\| \|F_0\| \leq \sigma_r^2(F_0)/4,$$

where $\sigma_i(F)$ is the i -th largest singular value of F . Let

$$R_1 := \arg \min_{R \in \mathcal{O}^{r \times r}} \|F_1 R - F_0\|_{\text{F}} \quad \text{and} \quad R_2 := \arg \min_{R \in \mathcal{O}^{r \times r}} \|F_2 R - F_0\|_{\text{F}}.$$

Then, the followings hold

$$\begin{aligned} \|R_1 - R_2\|_{\text{F}} &\leq \frac{2}{\sigma_r^2(F_0)} \|F_2 - F_1\|_{\text{F}} \|F_0\| & (86) \\ \|F_1 R_1 - F_2 R_2\| &\leq 5 \frac{\sigma_1^2(F_0)}{\sigma_r^2(F_0)} \|F_1 - F_2\| \\ \|F_1 R_1 - F_2 R_2\|_{\text{F}} &\leq 5 \frac{\sigma_1^2(F_0)}{\sigma_r^2(F_0)} \|F_1 - F_2\|_{\text{F}} \end{aligned}$$

Proof. The same as the proof of Lemma 37 in [39]. \square

Lemma 13 (Lemma 20 in [4]). *Let $U \Sigma V^\top$ be the SVD of a rank- r matrix XY^\top with $X, Y \in \mathbb{R}^{n \times r}$. Then there exists an invertible matrix $Q \in \mathbb{R}^{r \times r}$ such that $X = U \Sigma^{1/2} Q$ and $Y = V \Sigma^{1/2} Q^{-\top}$. In addition, one has*

$$\|\Sigma_Q - \Sigma_Q^{-1}\|_{\text{F}} \leq \frac{1}{\sigma_{\min}(\Sigma)} \|X^\top X - Y^\top Y\|_{\text{F}},$$

where $U_Q \Sigma_Q V_Q^\top$ is the SVD of Q . In particular, if $X^\top X - Y^\top Y = 0$, then Q must be a rotation matrix.

Lemma 14 (Lemma 13 in [18]). *Consider two symmetric matrices obeying $A_1 \succeq \mu_1 I$ and $A_2 \succeq \mu_2 I$ for some $\mu_1, \mu_2 > 0$. Let $R_1 \succeq 0$ ($R_2 \succeq 0$ respectively) be the (principal) matrix square root of A_1 (A_2 respectively). Then one has*

$$\|R_1 - R_2\| \leq \frac{1}{\sqrt{\mu_1} + \sqrt{\mu_2}} \|A_1 - A_2\|.$$

Lemma 15 (Perturbation on the row space and column space). *Suppose $X_1, X_2, Y_1, Y_2 \in \mathbb{R}^{n \times r}$ with singular values in the interval $[\sigma_a, \sigma_b]$ where $0 < \sigma_a \leq \sigma_b$. Suppose*

$$\max(\|X_1 - X_2\|_{\text{F}}, \|Y_1 - Y_2\|_{\text{F}}) \leq a \quad (87)$$

$$\max(\|X_1^\top X_1 - Y_1^\top Y_1\|_{\text{F}}, \|X_2^\top X_2 - Y_2^\top Y_2\|_{\text{F}}) \leq \epsilon. \quad (88)$$

Let $U_1 \Sigma_1 V_1^\top$ be the SVD of the matrix $X_1 Y_1^\top$, $U_2 \Sigma_2 V_2^\top$ be the SVD of the matrix $X_2 Y_2^\top$ where $U_1, U_2, V_1, V_2 \in \mathbb{R}^{n \times r}$, $\Sigma_1, \Sigma_2 \in \mathbb{R}^{r \times r}$. Then, there exists rotation matrices $R_1, R_2 \in \mathcal{O}^{r \times r}$ such that

$$\max(\|U_1 R_1 - U_2 R_2\|_{\text{F}}, \|V_1 R_1 - V_2 R_2\|_{\text{F}}) \leq 2\sqrt{r} \frac{\sigma_b}{\sigma_a} \left(a + \frac{2\sigma_b \epsilon}{\sigma_a^2} \right) \frac{1}{\sigma_a}. \quad (89)$$

Furthermore, if one has $\max(\|X_1 - X_2\|_{2, \infty}, \|Y_1 - Y_2\|_{2, \infty}) \leq b$, then there exists $R_1, R_2 \in \mathcal{O}^{r \times r}$ such that Eq. (89) and the following hold

$$\begin{aligned} &\max(\|U_1 R_1 - U_2 R_2\|_{2, \infty}, \|V_1 R_1 - V_2 R_2\|_{2, \infty}) \\ &\leq \left(b + \|U_2\|_{2, \infty} a \frac{\sigma_b}{\sigma_a} + \frac{2\sigma_b \epsilon}{\sigma_a^2} (1 + \|U_2\|_{2, \infty} \sigma_b / \sigma_a) \right) \frac{1}{\sigma_a}. \end{aligned} \quad (90)$$

One particular choice of R_1, R_2 satisfying above conditions is $R_1 = U_{Q_1} V_{Q_1}^\top, R_2 = U_{Q_2} V_{Q_2}^\top$ where $U_{Q_1} \Sigma_{Q_1} V_{Q_1}^\top$ is the SVD of the matrix $\Sigma_1^{-1/2} U_1^\top X_1$ and $U_{Q_2} \Sigma_{Q_2} V_{Q_2}^\top$ is the SVD of the matrix $\Sigma_2^{-1/2} U_2^\top X_2$.

Proof. Note that we have the following facts.

$$\sigma_{\max}(\Sigma_1) = \sigma_{\max}(X_1 Y_1^\top) \leq \sigma_{\max}(X_1) \sigma_{\max}(Y_1) \leq \sigma_b^2 \quad (91)$$

$$\sigma_{\min}(\Sigma_1) = \sigma_{\min}(X_1 Y_1^\top) \geq \sigma_{\min}(X_1) \sigma_{\min}(Y_1) \geq \sigma_a^2. \quad (92)$$

Similarly, $\sigma_{\max}(\Sigma_2) \leq \sigma_b^2, \sigma_{\min}(\Sigma_2) \geq \sigma_a^2$.

We first intend to invoke Lemma 13 to establish the connection between X_1, Y_1 (X_2, Y_2) and U_1, V_1 (U_2, V_2).

By Lemma 13, there exists invertible matrix Q_1 such that $X_1 = U_1 \Sigma_1^{1/2} Q_1, Y_1 = V_1 \Sigma_1^{1/2} Q_1^{-\top}$ and

$$\left\| \Sigma_{Q_1} - \Sigma_{Q_1}^{-1} \right\|_{\text{F}} \leq \frac{\|X_1^\top X - Y_1^\top Y\|_{\text{F}}}{\sigma_{\min}(\Sigma_1)} \stackrel{(i)}{\leq} \frac{\epsilon}{\sigma_a^2} \quad (93)$$

where $U_{Q_1} \Sigma_{Q_1} V_{Q_1}^\top$ is the SVD of Q_1 and (i) is due to Eq. (88) and $\sigma_{\min}(\Sigma_1) \geq \sigma_a^2$.

Similarly, we can also obtain that there exists $Q_2 \in \mathbb{R}^{r \times r}$ such that $X_2 = U_2 \Sigma_2^{1/2} Q_2, Y_2 = V_2 \Sigma_2^{1/2} Q_2^{-\top}$ and

$$\left\| \Sigma_{Q_2} - \Sigma_{Q_2}^{-1} \right\|_{\text{F}} \leq \frac{\epsilon}{\sigma_a^2}$$

where $U_{Q_2} \Sigma_{Q_2} V_{Q_2}^\top$ is the SVD of Q_2 .

We next intend to substitute Q_1, Q_2 by rotation matrices to simplify the analysis.

Let $R_1 = U_{Q_1} V_{Q_1}^\top, R_2 = U_{Q_2} V_{Q_2}^\top$. Note that $R_1, R_2 \in O^{r \times r}$ are rotation matrices. Let $X'_1 = X_1 Q_1^{-1} R_1 = U_1 \Sigma_1^{1/2} R_1$. In fact

$$\begin{aligned} X'_1 - X_1 &= X_1 (Q_1^{-1} R_1 - I_r) \\ &= X_1 (V_{Q_1} \Sigma_{Q_1}^{-1} U_{Q_1}^\top U_{Q_1} V_{Q_1}^\top - I_r) \\ &\stackrel{(i)}{=} X_1 (V_{Q_1} \Sigma_{Q_1}^{-1} V_{Q_1}^\top - I_r) \\ &\stackrel{(ii)}{=} X_1 (V_{Q_1} (\Sigma_{Q_1}^{-1} - I_r) V_{Q_1}^\top) \end{aligned}$$

where (i) is due to $U_{Q_1}^\top U_{Q_1} = I_r$ and (ii) is due to $V_{Q_1} V_{Q_1}^\top = I_r$. Then,

$$\begin{aligned} \|X'_1 - X_1\|_{\text{F}} &\stackrel{(i)}{\leq} \|X_1\| \|V_{Q_1}\| \left\| \Sigma_{Q_1}^{-1} - I_r \right\|_{\text{F}} \|V_{Q_1}^\top\| \\ &\stackrel{(ii)}{\leq} \|X_1\| \left\| \Sigma_{Q_1}^{-1} - I_r \right\|_{\text{F}} \\ &\stackrel{(iii)}{\leq} \sigma_b \frac{\epsilon}{\sigma_a^2} \end{aligned}$$

where (i) is due to $\|ABC\|_{\text{F}} \leq \|A\| \|B\|_{\text{F}} \|C\|$, (ii) is due to $\|V_{Q_1}\| = \|V_{Q_1}^\top\| = 1$, and (iii) is due to $\left\| \Sigma_{Q_1}^{-1} - I_r \right\|_{\text{F}} \leq \left\| \Sigma_{Q_1}^{-1} - \Sigma_{Q_1} \right\|_{\text{F}}$ and Eq. (93).

Let $Y'_1 = V_1 \Sigma_1^{1/2} R_1, X'_2 = U_2 \Sigma_2^{1/2} R_2, Y'_2 = V_2 \Sigma_2^{1/2} R_2$. Similarly, one can verify that

$$\max(\|Y'_1 - Y_1\|_{\text{F}}, \|X'_2 - X_2\|_{\text{F}}, \|Y'_2 - Y_2\|_{\text{F}}) \leq \sigma_b \frac{\epsilon}{\sigma_a^2}. \quad (94)$$

By triangle inequality, this guarantees that

$$\max(\|X'_1 - X'_2\|_{\text{F}}, \|Y'_1 - Y'_2\|_{\text{F}}) \leq a + \frac{2\sigma_b \epsilon}{\sigma_a^2}. \quad (95)$$

Note that we have the following decomposition for X'_1, X'_2 .

$$X'_1 = U_1 R_1 R_1^\top \Sigma_1^{1/2} R_1 \quad (96)$$

$$X'_2 = U_2 R_2 R_2^\top \Sigma_2^{1/2} R_2. \quad (97)$$

In order to establish a bound for $U_1 R_1 - U_2 R_2$ based on $X'_1 - X'_2$, we require an additional bound for $R_1^\top \Sigma_1^{1/2} R_1 - R_2^\top \Sigma_2^{1/2} R_2$. Notice that the following provides a bound for $R_1^\top \Sigma_1 R_1 - R_2^\top \Sigma_2 R_2$.

$$\begin{aligned} \|R_1^\top \Sigma_1 R_1 - R_2^\top \Sigma_2 R_2\|_F &\stackrel{(i)}{=} \|X_1'^\top X_1' - X_2'^\top X_2'\|_F \\ &\stackrel{(ii)}{\leq} \|(X_1'^\top - X_2'^\top)X_1'\|_F + \|X_2'^\top(X_1' - X_2')\|_F \\ &\stackrel{(iii)}{\leq} \|X_1'^\top - X_2'^\top\|_F \|X_1'\| + \|X_2'^\top\| \|X_1' - X_2'\|_F \\ &\stackrel{(iv)}{\leq} (a + \frac{2\sigma_b\epsilon}{\sigma_a^2}) \|X_1'\| + (a + \frac{2\sigma_b\epsilon}{\sigma_a^2}) \|X_2'\| \\ &\stackrel{(v)}{\leq} 2(a + \frac{2\sigma_b\epsilon}{\sigma_a^2})\sigma_b. \end{aligned} \quad (98)$$

where (i) is due to $X'_1 = U_1 \Sigma_1^{1/2} R_1, X'_2 = U_2 \Sigma_2^{1/2} R_2$, (ii) is due to the triangle inequality, (iii) is due to $\|ABC\|_F \leq \|A\| \|B\|_F \|C\|$, (iv) is due to Eq. (95), and (v) is due to $\|X'_1\| = \sigma_{\max}(\Sigma_1^{1/2}), \|X'_2\| = \sigma_{\max}(\Sigma_2^{1/2})$ and $\sigma_{\max}(\Sigma_1) \leq \sigma_b^2, \sigma_{\max}(\Sigma_2) \leq \sigma_b^2$ (see Eq. (91)).

By Lemma 14, this implies

$$\begin{aligned} \|R_1^\top \Sigma_1^{1/2} R_1 - R_2^\top \Sigma_2^{1/2} R_2\| &\leq \frac{1}{\sqrt{\sigma_{\min}(\Sigma_1)} + \sqrt{\sigma_{\min}(\Sigma_2)}} \|R_1^\top \Sigma_1 R_1 - R_2^\top \Sigma_2 R_2\| \\ &\stackrel{(i)}{\leq} \frac{1}{2\sigma_a} \|R_1^\top \Sigma_1 R_1 - R_2^\top \Sigma_2 R_2\|_F \\ &\stackrel{(ii)}{\leq} (a + \frac{2\sigma_b\epsilon}{\sigma_a^2}) \frac{\sigma_b}{\sigma_a} \end{aligned} \quad (99)$$

where (i) is due to $\sigma_{\min}(\Sigma_1) \geq \sigma_a^2, \sigma_{\min}(\Sigma_2) \geq \sigma_a^2$ and $\|A\| \leq \|A\|_F$ for any matrix A , and (ii) is due to Eq. (98).

Then we can establish the bound for $U_1 R_1 - U_2 R_2$. Note that by Eqs. (96) and (97), we have

$$X'_1 - X'_2 = (U_1 R_1 - U_2 R_2)(R_1^\top \Sigma_1^{1/2} R_1) + U_2 R_2 (R_1^\top \Sigma_1^{1/2} R_1 - R_2^\top \Sigma_2^{1/2} R_2)$$

which further implies

$$\begin{aligned} U_1 R_1 - U_2 R_2 &= (X'_1 - X'_2)(R_1^\top \Sigma_1^{-1/2} R_1) + U_2 R_2 (R_1^\top \Sigma_1^{1/2} R_1 - R_2^\top \Sigma_2^{1/2} R_2)(R_1^\top \Sigma_1^{-1/2} R_1). \end{aligned} \quad (100)$$

Therefore, we can provide a bound for $\|U_1 R_1 - U_2 R_2\|_F$ by the following.

$$\begin{aligned} \|U_1 R_1 - U_2 R_2\|_F &\stackrel{(i)}{\leq} \|X'_1 - X'_2\|_F \|R_1^\top \Sigma_1^{-1/2} R_1\| \\ &\quad + \|U_2 R_2\|_F \|R_1^\top \Sigma_1^{1/2} R_1 - R_2^\top \Sigma_2^{1/2} R_2\| \|R_1^\top \Sigma_1^{-1/2} R_1\| \\ &\stackrel{(ii)}{\leq} (a + \frac{2\sigma_b\epsilon}{\sigma_a^2}) \|R_1^\top \Sigma_1^{-1/2} R_1\| + \|U_2 R_2\|_F (a + \frac{2\sigma_b\epsilon}{\sigma_a^2}) \frac{\sigma_b}{\sigma_a} \|R_1^\top \Sigma_1^{-1/2} R_1\| \\ &\stackrel{(iii)}{\leq} (a + \frac{2\sigma_b\epsilon}{\sigma_a^2}) \sigma_{\max}(\Sigma_1^{-1/2}) + \sqrt{r} (a + \frac{2\sigma_b\epsilon}{\sigma_a^2}) \frac{\sigma_b}{\sigma_a} \sigma_{\max}(\Sigma_1^{-1/2}) \\ &\stackrel{(iv)}{\leq} 2\sqrt{r} \frac{\sigma_b}{\sigma_a} (a + \frac{2\sigma_b\epsilon}{\sigma_a^2}) \frac{1}{\sigma_a} \end{aligned}$$

where (i) is due to the triangle inequality, (ii) is due to Eq. (95) and Eq. (99), (iii) is due to $\left\|R_1^\top \Sigma_1^{-1/2} R_1\right\| = \sigma_{\max}(\Sigma_1^{-1/2})$ and $\|U_2 R_2\|_F = \|U_2\|_F = \sqrt{r}$, (iv) is due to $\sigma_{\max}(\Sigma_1^{-1/2}) = 1/\sqrt{\sigma_{\min}(\Sigma_1)} \leq 1/\sigma_a$. Similarly, one can obtain that $\|V_1 R_1 - V_2 R_2\|_F \leq 2\sqrt{r} \frac{\sigma_b}{\sigma_a} \left(a + \frac{2\sigma_b \epsilon}{\sigma_a^2}\right) \frac{1}{\sigma_a}$.

Note that $\|U_1 R_1 R_2^\top - U_2\|_F = \|U_1 R_1 - U_2 R_2\|_F$ and $\|V_1 R_1 R_2^\top - V_2\|_F = \|V_1 R_1 - V_2 R_2\|_F$, this completes the proof for Eq. (89).

In addition, if $\|X_1 - X_2\|_{2,\infty} \leq b$, we can obtain that

$$\begin{aligned} \|X'_1 - X'_2\|_{2,\infty} &\leq \|X_1 - X_2\|_{2,\infty} + \|X'_1 - X_1\|_F + \|X'_2 - X_2\|_F \\ &\leq b + \frac{2\sigma_b \epsilon}{\sigma_a^2}. \end{aligned}$$

Then based on Eq. (100), we have

$$\begin{aligned} \|U_1 R_1 - U_2 R_2\|_{2,\infty} &\stackrel{(i)}{\leq} \left\| (X'_1 - X'_2) R_1^\top \Sigma_1^{-1/2} R_1 \right\|_{2,\infty} \\ &\quad + \left\| U_2 R_2 (R_1^\top \Sigma_1^{1/2} R_1 - R_2^\top \Sigma_2^{1/2} R_2) R_1^\top \Sigma_1^{-1/2} R_1 \right\|_{2,\infty} \\ &\stackrel{(ii)}{\leq} \|X'_1 - X'_2\|_{2,\infty} \left\| R_1^\top \Sigma_1^{-1/2} R_1 \right\| \\ &\quad + \|U_2\|_{2,\infty} \|R_2\| \left\| R_1^\top \Sigma_1^{1/2} R_1 - R_2^\top \Sigma_2^{1/2} R_2 \right\| \left\| R_1^\top \Sigma_1^{-1/2} R_1 \right\| \\ &\stackrel{(iii)}{\leq} \left(b + \|U_2\|_{2,\infty} a \frac{\sigma_b}{\sigma_a} + \frac{2\sigma_b \epsilon}{\sigma_a^2} (1 + \|U_2\|_{2,\infty} \sigma_b / \sigma_a) \right) \frac{1}{\sigma_a} \end{aligned}$$

where (i) is due to triangle inequality, (ii) is due to $\|AB\|_{2,\infty} \leq \|A\|_{2,\infty} \|B\|$, (iii) follows the same derivation for $\|U_1 R_1 - U_2 R_2\|_F$. Notice that $\|U_1 R_1 R_2^\top - U_2\|_{2,\infty} = \|U_1 R_1 - U_2 R_2\|_{2,\infty}$. The similar results also can be obtained for $\|V_1 R_1 R_2^\top - V_2\|_{2,\infty}$. This completes the proof for Eq. (90). \square

The following lemma shows that Assumption 3(a) and Assumption 3(b) hold for a small ball around (X^*, Y^*) .

Lemma 16 (Implication of Assumption 3(a) and Assumption 3(b)). *Suppose $X^*, Y^*, X, Y \in \mathbb{R}^{n \times r}$. Suppose $X^* Y^{*\top} = U^* \Sigma^* V^{*\top}$, $XY^\top = U \Sigma V^\top$ is the SVD of $X^* Y^{*\top}$, XY^\top respectively. Suppose $X^* = U^* \Sigma^{*1/2}$, $Y^* = V^* \Sigma^{*1/2}$ and $\sigma_1(\Sigma^*) = \sigma_{\max}$, $\sigma_r(\Sigma^*) = \sigma_{\min}$, $\kappa = \sigma_{\max} / \sigma_{\min}$. Suppose $\frac{\sigma \sqrt{n}}{\sigma_{\min}} \leq C_2 \frac{1}{\kappa^2 r^2 \log^5(n)}$ and*

$$\|X - X^*\|_F + \|Y - Y^*\|_F \leq C_F \frac{\sigma \sqrt{n} \log^{2.5}(n) \sqrt{\sigma_{\max} r}}{\sigma_{\min}}.$$

Let T^* be the tangent space of $X^* Y^{*\top}$, T be the tangent space of XY^\top . If there exists a constant C_{r_1} such that $\|ZV^*\|_F^2 + \|Z^\top U^*\|_F^2 \leq \left(1 - \frac{C_{r_1}}{\log(n)}\right) \|Z\|_F^2$ (Assumption 3(a)), then for large enough n ,

$$\|ZV\|_F^2 + \|Z^\top U\|_F^2 \leq \left(1 - \frac{C_{r_1}}{2 \log(n)}\right) \|Z\|_F^2 \quad (101)$$

$$\|P_{T^\perp}(Z)\|_F^2 \geq \frac{C_{r_1}}{2 \log(n)} \|Z\|_F^2 \quad (102)$$

$$\|P_{T^\perp}(Z) - P_{T^{*\perp}}(Z)\|_* \lesssim \frac{1}{r^{0.5} \log^{2.5}(n)} \|Z\|_F. \quad (103)$$

In addition, if there exists a constant C_{r_2} such that $|\langle Z, U^* V^{*\top} \rangle| \|P_{T^{*\perp}}(Z)\| \leq \left(1 - \frac{C_{r_2}}{\log n}\right) \|P_{T^{*\perp}}(Z)\|_F^2$ (Assumption 3(b)), then for large enough n ,

$$|\langle Z, UV^\top \rangle| \|P_{T^\perp}(Z)\| \leq \left(1 - \frac{C_{r_2}}{2 \log n}\right) \|P_{T^\perp}(Z)\|_F^2.$$

Proof. From Lemma 10, we have $\sqrt{\frac{\sigma_{\min}}{2}} \leq \sigma_i(X) \leq \sqrt{2\sigma_{\max}}, \sqrt{\frac{\sigma_{\min}}{2}} \leq \sigma_i(Y) \leq \sqrt{2\sigma_{\max}}$ and $\frac{\sigma_{\min}}{2} \leq \sigma_i(\Sigma) \leq 2\sigma_{\max}$ for $i \in [r]$.

Also note that $X^{*\top}X^* = Y^{*\top}Y^* = \Sigma^*$, we have

$$\begin{aligned} & \|X^\top X - Y^\top Y\|_F \\ &= \|X^\top X - X^{*\top}X^* + Y^{*\top}Y^* - Y^\top Y\|_F \\ &\leq \|X^\top X - X^{*\top}X^*\|_F + \|Y^\top Y - Y^{*\top}Y^*\|_F \\ &\leq \|X^\top(X - X^*) + (X^\top - X^{*\top})X^*\|_F + \|Y^\top(Y - Y^*) + (Y^\top - Y^{*\top})Y^*\|_F \\ &\lesssim \|X - X^*\|_F \sqrt{\sigma_{\max}} + \|Y - Y^*\|_F \sqrt{\sigma_{\max}}. \end{aligned}$$

Then we can invoke Lemma 15 and obtain that, there exists rotation matrix $R \in \mathbb{R}^{r \times r}$ where

$$\|UR - U^*\|_F + \|VR - V^*\|_F \lesssim \frac{\sigma\sqrt{n} \log^{2.5}(n) r \kappa^2}{\sigma_{\min}} \stackrel{(i)}{\leq} \frac{C}{r \log^{2.5}(n)} \quad (104)$$

for some constant C . Here, (i) is due to $\frac{\sigma\sqrt{n}}{\sigma_{\min}} \leq C_2 \frac{1}{\kappa^2 r^2 \log^5(n)}$.

This implies

$$\begin{aligned} \|UU^\top - U^*U^{*\top}\|_F &= \|UR(UR)^\top - U^*U^{*\top}\|_F \\ &\leq \|UR - U^*\|_F \|UR\| + \|U^*\| \|UR - U^*\|_F \\ &\leq 2 \|UR - U^*\|_F \\ &\leq \frac{C}{r \log^{2.5}(n)}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \|VV^\top - V^*V^{*\top}\|_F &\leq \frac{C}{r \log^{2.5}(n)} \\ \|UV^\top - U^*V^{*\top}\|_F &\leq \frac{C}{r \log^{2.5}(n)}. \end{aligned} \quad (105)$$

Then, consider the proof of Eq. (101). We have

$$\begin{aligned} & \|ZV\|_F^2 + \|Z^\top U\|_F^2 \\ &= \text{tr}(Z^\top ZVV^\top) + \text{tr}(ZZ^\top UU^\top) \\ &= \text{tr}(Z^\top ZV^*V^{*\top}) + \text{tr}(Z^\top Z(VV^\top - V^*V^{*\top})) \\ &\quad + \text{tr}(ZZ^\top U^*U^{*\top}) - \text{tr}(ZZ^\top(UU^\top - U^*U^{*\top})) \\ &\stackrel{(i)}{\leq} \|ZV^*\|_F^2 + \|Z^\top U^*\|_F^2 + \|Z^\top Z\|_* \|VV^\top - V^*V^{*\top}\| + \|ZZ^\top\|_* \|UU^\top - U^*U^{*\top}\| \\ &\stackrel{(ii)}{\leq} \left(1 - \frac{C_{r_1}}{\log(n)}\right) \|Z\|_F^2 + \|Z\|_F^2 \frac{2C}{r \log^{2.5}(n)} \\ &\leq \left(1 - \frac{C_{r_1}}{2 \log(n)}\right) \|Z\|_F^2 \end{aligned}$$

where (i) is due to $\text{tr}(AB) \leq \|A\|_* \|B\|$, (ii) is due to $\|ZV^*\|_F^2 + \|Z^\top U^*\|_F^2 \leq \left(1 - \frac{C_{r_1}}{\log(n)}\right) \|Z\|_F^2$ and $\|Z^\top Z\|_* = \|ZZ^\top\|_* = \|Z\|_F^2$ together with

$$\max(\|UU^\top - U^*U^{*\top}\|_F, \|VV^\top - V^*V^{*\top}\|_F) \leq \frac{C}{r \log^{2.5}(n)}.$$

This completes the proof of Eq. (101).

Next, consider the proof of Eq. (102). This is in fact the simple implication of Eq. (101). Note that

$$\begin{aligned}
\|P_T(Z)\|_F^2 &= \|Z - (I - UU^\top)Z(I - VV^\top)\|_F^2 \\
&= \|UU^\top Z + (I - UU^\top)Z(VV^\top)\|_F^2 \\
&= \text{tr}(UU^\top ZZ^\top UU^\top) + \text{tr}((I - UU^\top)Z(VV^\top)(VV^\top)Z^\top(I - UU^\top)) \\
&= \text{tr}(UU^\top ZZ^\top) + \text{tr}(Z^\top ZVV^\top) - \text{tr}(UU^\top ZVV^\top Z^\top) \\
&\leq \|Z^\top U\|_F^2 + \|ZV\|_F^2 \leq \left(1 - \frac{C_{r_1}}{2\log n}\right) \|Z\|_F^2.
\end{aligned} \tag{106}$$

This implies that $\frac{C_{r_1}}{2\log n} \|Z\|_F^2 \leq \|Z\|_F^2 - \|P_T(Z)\|_F^2$. Note that $\|P_{T^\perp}(Z)\|_F^2 + \|P_T(Z)\|_F^2 = \|Z\|_F^2$, which completes the proof of Eq. (102).

Next, consider the proof of Eq. (103). Let $\Delta = P_{T^\perp}(Z) - P_{T^{*\perp}}(Z)$. We have

$$\begin{aligned}
\|\Delta\|_* &= \|(I - UU^\top)Z(I - VV^\top) - (I - U^*U^{*\top})Z(I - V^*V^{*\top})\|_* \\
&\leq \|(I - UU^\top)Z(I - VV^\top) - (I - U^*U^{*\top})Z(I - VV^\top)\|_* \\
&\quad + \|(I - U^*U^{*\top})Z(I - VV^\top) - (I - U^*U^{*\top})Z(I - V^*V^{*\top})\|_* \\
&\leq \|(U^*U^{*\top} - UU^\top)Z(I - VV^\top)\|_* + \|(I - U^*U^{*\top})Z(V^*V^{*\top} - VV^\top)\|_* \\
&\stackrel{(i)}{\leq} \|(U^*U^{*\top} - UU^\top)Z(I - VV^\top)\|_F \sqrt{\text{rank}(U^*U^{*\top} - UU^\top)} \\
&\quad + \|(I - U^*U^{*\top})Z(V^*V^{*\top} - VV^\top)\|_F \sqrt{\text{rank}(V^*V^{*\top} - VV^\top)} \\
&\leq \|Z\|_F (\|(U^*U^{*\top} - UU^\top)\| + \|(V^*V^{*\top} - VV^\top)\|) \sqrt{2r} \\
&\lesssim \frac{1}{r^{0.5} \log^{2.5}(n)} \|Z\|_F.
\end{aligned}$$

Here, (i) is due to $\|A\|_* \leq \|A\|_F \sqrt{\text{rank}(A)}$ and $\text{rank}(AB) \leq \text{rank}(A)$ for any matrices A and B . This completes the proof of Eq. (103).

Furthermore, we also have

$$\begin{aligned}
\|\Delta\|_F &\leq \|(I - UU^\top)Z(I - VV^\top) - (I - U^*U^{*\top})Z(I - V^*V^{*\top})\|_F \\
&\leq \|(U^*U^{*\top} - UU^\top)Z(I - VV^\top)\|_F + \|(I - U^*U^{*\top})Z(V^*V^{*\top} - VV^\top)\|_F \\
&\leq \|Z\|_F (\|(U^*U^{*\top} - UU^\top)\| + \|(V^*V^{*\top} - VV^\top)\|) \\
&\leq \frac{2C}{r \log^{2.5}(n)} \|Z\|_F.
\end{aligned} \tag{107}$$

Finally, consider the proof of Eq. (104). Recall the Assumption 3(b) provides the following.

$$\frac{C_{r_2}}{\log(n)} \|P_{T^{*\perp}}(Z)\|_F^2 \leq \|P_{T^{*\perp}}(Z)\|_F^2 - |\langle Z, U^*V^{*\top} \rangle| \|P_{T^{*\perp}}(Z)\|. \tag{108}$$

Consider substituting $P_{T^\perp}(Z)$ with $P_{T^\perp}(Z) = P_{T^{*\perp}}(Z) + \Delta$ in $\|P_{T^\perp}(Z)\|_F^2$, when $\frac{C_{r_2}}{2\log(n)} \leq 1$, we have

$$\begin{aligned}
&\left(1 - \frac{C_{r_2}}{2\log(n)}\right) \|P_{T^\perp}(Z)\|_F^2 \\
&= \left(1 - \frac{C_{r_2}}{2\log(n)}\right) \|P_{T^{*\perp}}(Z)\|_F^2 + \left(1 - \frac{C_{r_2}}{2\log(n)}\right) \|\Delta\|_F^2 + \left(1 - \frac{C_{r_2}}{2\log(n)}\right) 2\langle P_{T^{*\perp}}(Z), \Delta \rangle \\
&\stackrel{(i)}{\geq} \|P_{T^{*\perp}}(Z)\|_F^2 - \frac{C_{r_2}}{2\log(n)} \|P_{T^{*\perp}}(Z)\|_F^2 - 2\|P_{T^{*\perp}}(Z)\|_F \|\Delta\|_F \\
&\stackrel{(ii)}{\geq} \|P_{T^{*\perp}}(Z)\|_F^2 - \frac{C_{r_2}}{2\log(n)} \|P_{T^{*\perp}}(Z)\|_F^2 - \|Z\|_F^2 \frac{4C}{r \log^{2.5}(n)}
\end{aligned} \tag{109}$$

where (i) is due to $\|\Delta\|_{\mathbb{F}}^2 \geq 0$ and $|\langle A, B \rangle| \leq \|A\|_{\mathbb{F}} \|B\|_{\mathbb{F}}$, (ii) is due to Eq. (107).

Next, consider $|\langle Z, UV^{\top} \rangle| \|P_{T^{\perp}}(Z)\|$. Let $\Delta' = UV^{\top} - U^*V^{*\top}$. Then, we have

$$\begin{aligned}
& |\langle Z, UV^{\top} \rangle| \|P_{T^{\perp}}(Z)\| \\
&= |\langle Z, U^*V^{*\top} + \Delta' \rangle| \|P_{T^{*\perp}}(Z) + \Delta\| \\
&\leq |\langle Z, U^*V^{*\top} \rangle| \|P_{T^{*\perp}}(Z)\| + |\langle Z, \Delta' \rangle| \|P_{T^{*\perp}}(Z)\| + |\langle Z, U^*V^{*\top} \rangle| \|\Delta\| + |\langle Z, \Delta' \rangle| \|\Delta\| \\
&\leq |\langle Z, U^*V^{*\top} \rangle| \|P_{T^{*\perp}}(Z)\| + \|Z\|_{\mathbb{F}} \|\Delta'\|_{\mathbb{F}} \|P_{T^{*\perp}}(Z)\| \\
&\quad + \|Z\|_{\mathbb{F}} \|U^*V^{*\top}\|_{\mathbb{F}} \|\Delta\| + \|Z\|_{\mathbb{F}} \|\Delta'\|_{\mathbb{F}} \|\Delta\| \\
&\leq |\langle Z, U^*V^{*\top} \rangle| \|P_{T^{*\perp}}(Z)\| + \|Z\|_{\mathbb{F}}^2 \|\Delta'\|_{\mathbb{F}} + \|Z\|_{\mathbb{F}} \sqrt{r} \|\Delta\|_{\mathbb{F}} + \|Z\|_{\mathbb{F}} \|\Delta'\|_{\mathbb{F}} \|\Delta\| \\
&\stackrel{(i)}{\leq} |\langle Z, U^*V^{*\top} \rangle| \|P_{T^{*\perp}}(Z)\| + \|Z\|_{\mathbb{F}}^2 \left(\frac{C}{r \log^{2.5}(n)} + \frac{C}{r^{0.5} \log^{2.5}(n)} + \frac{C^2}{r^2 \log^3(n)} \right).
\end{aligned}$$

where (i) is due to Eq. (107) and Eq. (105). This implies

$$|\langle Z, UV^{\top} \rangle| \|P_{T^{\perp}}(Z)\| \leq |\langle Z, U^*V^{*\top} \rangle| \|P_{T^{*\perp}}(Z)\| + \|Z\|_{\mathbb{F}}^2 \frac{3C^2}{r^{0.5} \log^{2.5}(n)} \quad (110)$$

Combining Eq. (109) and Eq. (110), we have

$$\begin{aligned}
& \left(1 - \frac{C_{r_2}/2}{\log(n)}\right) \|P_{T^{\perp}}(Z)\|_{\mathbb{F}}^2 - |\langle Z, UV^{\top} \rangle| \|P_{T^{\perp}}(Z)\| \\
&\geq \|P_{T^{*\perp}}(Z)\|_{\mathbb{F}}^2 - \frac{C_{r_2}}{2 \log(n)} \|P_{T^{*\perp}}(Z)\|_{\mathbb{F}}^2 - \|Z\|_{\mathbb{F}}^2 \frac{4C}{r \log^{2.5}(n)} \\
&\quad - |\langle Z, U^*V^{*\top} \rangle| \|P_{T^{*\perp}}(Z)\| - \|Z\|_{\mathbb{F}}^2 \frac{3C^2}{r^{0.5} \log^{2.5}(n)} \\
&\stackrel{(i)}{\geq} \frac{C_{r_2}}{\log n} \|P_{T^{*\perp}}(Z)\|_{\mathbb{F}}^2 - \frac{C_{r_2}}{2 \log n} \|P_{T^{*\perp}}(Z)\|_{\mathbb{F}}^2 - \|Z\|_{\mathbb{F}}^2 \frac{4C}{r \log^{2.5}(n)} - \|Z\|_{\mathbb{F}}^2 \frac{3C^2}{r^{0.5} \log^{2.5}(n)} \\
&\stackrel{(ii)}{\geq} \frac{C_{r_2} C_{r_1}}{4 \log^2 n} \|Z\|_{\mathbb{F}}^2 - \frac{3C^2 + 4C}{\log^{2.5}(n)} \|Z\|_{\mathbb{F}}^2 \\
&\stackrel{(iii)}{\geq} 0
\end{aligned}$$

where (i) is by Eq. (108), (ii) is by $\|P_{T^{*\perp}}(Z)\|_{\mathbb{F}}^2 \geq \frac{C_{r_1}}{2 \log(n)} \|Z\|_{\mathbb{F}}^2$ due to the analysis in Eq. (106), and (iii) holds for large enough n . This implies $|\langle Z, UV^{\top} \rangle| \|P_{T^{\perp}}(Z)\| \leq \left(1 - \frac{C_{r_2}/2}{\log(n)}\right) \|P_{T^{\perp}}(Z)\|_{\mathbb{F}}^2$, which completes the proof. \square