

The basic distributional theory for the product of zero mean correlated normal random variables

Robert E. Gaunt*

Abstract

The product of two zero mean correlated normal random variables has received much attention in the statistics literature and appears in many application areas. However, many important distributional properties are yet to be tabulated. This paper fills this gap by providing the basic distributional theory for the product of two zero mean correlated normal random variables.

Keywords: Product of correlated normal random variables; distributional theory

AMS 2010 Subject Classification: Primary 60E05; 62E15

1 Introduction

Let (X, Y) be a bivariate normal random vector with zero mean vector, variances (σ_X^2, σ_Y^2) and correlation coefficient ρ . The distribution of the product $Z = XY$ arises in numerous applications, with recent examples including statistics of Lagrangian power in two-dimensional turbulence (Bandi and Connaughton [3]), product confidence limits for indirect effects (MacKinnon et al. [28]) and statistical mediation analysis (MacKinnon [27]). The mean $\bar{Z}_n = n^{-1}(Z_1 + Z_2 + \dots + Z_n)$, where Z_1, Z_2, \dots, Z_n are independent and identical copies of Z , has also found application in areas such as electrical engineering (Ware and Lad [42]), astrophysics (Mangilli, Plaszczyński and Tristram [29], Watts et al. [43]) and quantum cosmology (Grishchuk [18]).

The exact distribution of the product $Z = XY$ has been studied since 1936 (Craig [5]), with subsequent contributions coming from Aroian [1], Aroian, Taneja and Cornwell [2], Bandi and Connaughton [3], Haldane [20], Meeker et al. [31]; see Nadarajah and Pogány [32] for an overview of these and further contributions. Recently, Nadarajah and Pogány [32] used characteristic functions to find exact formulas for the probability density function (PDF) of the product Z and more generally the mean \bar{Z}_n ; these formulas are starting to become well-known and have found many recent applications. Independently, in the physics literature, Grishchuk [18] had obtained the formula for the PDF of Z and Watts et al. [43] then obtained the more general formula for the PDF of \bar{Z}_n . Much earlier, a formula for PDF of Z was given in the book of Springer [38, equation (4.8.22)], but there were an unfortunate typo.

*Department of Mathematics, The University of Manchester, Oxford Road, Manchester M13 9PL, UK, robert.gaunt@manchester.ac.uk

Despite the interest in the distributions of Z and \bar{Z}_n , many other important distributional properties beyond the PDF are not tabulated in the literature. A recent work of Gaunt [14] identified Z and \bar{Z}_n as variance-gamma (VG) random variables (leading to a simple independent derivation of their PDFs), which means that some of their distributional properties can be inferred from results for the VG distribution; see, for example, Chapter 4 of the book of Kotz, Kozubowski and Podgórski [25]. However, searching through the VG literature can be difficult; it is tedious to convert results for the VG distribution to results for the distributions of Z and \bar{Z} ; and results that exploit the special structure of the random variables Z and \bar{Z}_n are not available.

In this paper, we aim to fill a gap in the literature by providing the basic distributional theory for the product of two zero mean correlated normal random variables. We present results for the more general mean \bar{Z}_n , with results for the product Z following on setting $n = 1$. The end result is that many of the most important distributional properties of Z and \bar{Z}_n are now collected in a single handy reference. The properties covered include: formulas for the PDF (Section 2.1), list of related distributions (Section 2.2), the cumulative distribution function (Section 2.3), generating functions and infinite divisibility (Section 2.4), representations in terms of other random variables (Section 2.5), Stein characterisation (Section 2.6), moments and cumulants (Section 2.7), and mode and median (Section 2.8). Basic properties of the modified Bessel function of the second kind will be needed throughout the paper, and are collected in Appendix A.

2 Distributional theory for the product of zero mean correlated normal random variables

2.1 Probability density function

For independent derivations of the following formulas for the PDFs of Z and \bar{Z}_n we refer the reader to Watts et al. [43], Nadarajah and Pogány [32] and Gaunt [14, 15]. The formula for the PDF of Z was also earlier derived by Grishchuk [18] and Springer [38], with the latter reference having a typo in the formula. For $x \in \mathbb{R}$,

$$f_Z(x) = \frac{1}{\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \exp\left(\frac{\rho x}{\sigma_X\sigma_Y(1-\rho^2)}\right) K_0\left(\frac{|x|}{\sigma_X\sigma_Y(1-\rho^2)}\right), \quad (2.1)$$

and, for $n \geq 1$,

$$f_{\bar{Z}_n}(x) = \frac{n^{(n+1)/2} 2^{(1-n)/2} |x|^{(n-1)/2}}{(\sigma_X\sigma_Y)^{(n+1)/2} \sqrt{\pi(1-\rho^2)} \Gamma(n/2)} \exp\left(\frac{\rho n x}{\sigma_X\sigma_Y(1-\rho^2)}\right) K_{\frac{n-1}{2}}\left(\frac{n|x|}{\sigma_X\sigma_Y(1-\rho^2)}\right). \quad (2.2)$$

Here $K_\nu(x)$ is a modified Bessel function of the second kind; see Appendix A for a definition and some standard properties.

The modified Bessel function in the PDFs (2.1) and (2.2) make them difficult to parse on first inspection. We can gain some understanding from the following limiting forms.

Using the limiting form (A.30), we have that

$$f_{\bar{Z}_n}(x) \sim \begin{cases} -\frac{1}{\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \log|x|, & x \rightarrow 0, n = 1, \\ \frac{n(1-\rho^2)^{n/2-1} \Gamma((n-1)/2)}{2\sqrt{\pi}\sigma_X\sigma_Y \Gamma(n/2)}, & x \rightarrow 0, n \geq 2. \end{cases} \quad (2.3)$$

We see that the density has a singularity at the origin if $n = 1$. In fact, for all parameter values ($n \geq 1$, $-1 < \rho < 1$, $\sigma_X, \sigma_Y > 0$) the distribution of \bar{Z}_n is unimodal; see Section 2.8 for further details. For $n \geq 2$, the density is bounded. As observed by Watt et al. [43], the tail behaviour of the density is obtained by applying the limiting form (A.31):

$$f_{\bar{Z}_n}(x) \sim \frac{(n/2)^{n/2}}{(\sigma_X\sigma_Y)^{n/2}\Gamma(n/2)} x^{n/2-1} \exp\left(-\frac{nx}{\sigma_X\sigma_Y(1+\rho)}\right), \quad x \rightarrow \infty, \quad (2.4)$$

and

$$f_{\bar{Z}_n}(x) \sim \frac{(n/2)^{n/2}}{(\sigma_X\sigma_Y)^{n/2}\Gamma(n/2)} (-x)^{n/2-1} \exp\left(\frac{nx}{\sigma_X\sigma_Y(1-\rho)}\right), \quad x \rightarrow -\infty. \quad (2.5)$$

We observe that the distribution of \bar{Z}_n has semi-heavy tails.

In the case that n is even, we can take advantage of a standard simplification of the modified Bessel function of the second kind (see (A.28)) to obtain a representation of the PDF in terms of elementary functions:

$$\begin{aligned} f_{\bar{Z}_n}(x) &= \frac{(n/2)^{n/2}}{(\sigma_X\sigma_Y)^{n/2}\Gamma(n/2)} |x|^{n/2-1} \exp\left(\frac{n(\rho x - |x|)}{\sigma_X\sigma_Y(1-\rho^2)}\right) \times \\ &\times \sum_{j=0}^{n/2-1} \frac{(n/2-1+j)!}{(n/2-1-j)!j!} \left(\frac{\sigma_X\sigma_Y(1-\rho^2)}{2n|x|}\right)^j, \quad x \in \mathbb{R}, n \in 2\mathbb{Z}^+. \end{aligned} \quad (2.6)$$

2.2 Related distributions

1. Setting $n = 2$ in (2.6) yields

$$f_{\bar{Z}_2}(x) = \frac{1}{\sigma_X\sigma_Y} \exp\left(\frac{2(\rho x - |x|)}{\sigma_X\sigma_Y(1-\rho^2)}\right), \quad (2.7)$$

which is the PDF of the asymmetric Laplace distribution (see Chapter 3 of Kotz, Kozubowski and Podgórski [25] for a comprehensive account of its distributional theory).

2. As $\rho \uparrow 1$, \bar{Z}_n converges in distribution to a scaled chi-square random variable: $\bar{Z}_n \rightarrow_d (\sigma_X\sigma_Y/n)V$, where $V \sim \chi_{(n)}^2$, the chi-square distribution with n degrees of freedom. By symmetry, $\bar{Z}_n \rightarrow_d -(\sigma_X\sigma_Y/n)V$, as $\rho \downarrow -1$. This is hardly surprising given the definition of \bar{Z}_n , and can be confirmed rigorously by letting $\rho \rightarrow \pm 1$ in the formula (2.11) below for the characteristic function of \bar{Z}_n and comparing to the chi-square characteristic function using Lévy's continuity theorem.

3. By the central limit theorem, $\sqrt{n}(\bar{Z}_n - \rho\sigma_X\sigma_Y) \rightarrow_d N(0, \sigma_X^2\sigma_Y^2(1 + \rho^2))$, as $n \rightarrow \infty$. Here, $\mathbb{E}[Z] = \rho\sigma_X\sigma_Y$ and $\text{Var}(Z) = \sigma_X^2\sigma_Y^2(1 + \rho^2)$ (see (2.20) and (2.21)).

4. The variance-gamma (VG) distribution with parameters $r > 0$, $\theta \in \mathbb{R}$, $\sigma > 0$, $\mu \in \mathbb{R}$ has PDF

$$f_{\text{VG}}(x) = \frac{e^{\theta(x-\mu)/\sigma^2}}{\sigma\sqrt{\pi}\Gamma(r/2)} \left(\frac{|x-\mu|}{2\sqrt{\theta^2 + \sigma^2}} \right)^{\frac{r-1}{2}} K_{\frac{r-1}{2}} \left(\frac{\sqrt{\theta^2 + \sigma^2}}{\sigma^2} |x-\mu| \right), \quad x \in \mathbb{R}, \quad (2.8)$$

If a random variable V has PDF (2.8), we write $V \sim \text{VG}(r, \theta, \sigma, \mu)$. This parametrisation was given in Gaunt [11]. It is similar to the parametrisation given by Finlay and Seneta [9] and other parametrisations are given by Eberlein and Hammerstein [8] and Kotz, Kozubowski and Podgórski [25], with Chapter 4 of the latter reference giving the most comprehensive account of the distributional theory of the VG distribution in the literature. The VG distribution is also known as the Bessel function distribution or the McKay Type II distribution (McKay [30]), as well as the generalized Laplace distribution (Kotz, Kozubowski and Podgórski [25]).

It was noted in the thesis of Gaunt [10] that $Z \sim \text{VG}(1, \rho\sigma_X\sigma_Y, \sigma_X\sigma_Y\sqrt{1-\rho^2}, 0)$ and later by Gaunt [14] that also

$$\bar{Z}_n \sim \text{VG}(n, \rho\sigma_X\sigma_Y/n, \sigma_X\sigma_Y\sqrt{1-\rho^2}/n, 0). \quad (2.9)$$

5. An exact formula for the PDF of the product of correlated normal random variables with non-zero means was derived by Cui et al. [6]. The formula is more complicated than (2.1), involving a double sum of modified Bessel functions of the second kind. An exact formula for the PDF of $k \geq 1$ independent zero mean normal random variables was obtained by Springer and Thompson [39]. An exact formula for the PDF of the product of three or more correlated normal random variables is not available in the literature.

2.3 Cumulative distribution function

For general parameter values $n \geq 1$, $-1 < \rho < 1$ and $\sigma_X, \sigma_Y > 0$, a closed-form formula for the cumulative distribution function (CDF) of \bar{Z} is not available. We record some cases for which exact formulas are available. Let $F_{\bar{Z}_n}(x) = \mathbb{P}(\bar{Z}_n \leq x)$.

Suppose $\rho = 0$. Then, by the symmetry of the PDF of \bar{Z} , it follows that its CDF is given by $F_{\bar{Z}_n}(x) = 1/2 + \text{sgn}(x) \int_0^{|x|} f_{\bar{Z}_n}(t) dt$, where $\text{sgn}(x)$ denotes the sign of x . On calculating the integral using (A.33), we have that, for $x \in \mathbb{R}$,

$$F_{\bar{Z}_n}(x) = \frac{1}{2} + \frac{nx}{2\sigma_X\sigma_Y} \left[K_{\frac{n-1}{2}} \left(\frac{n|x|}{\sigma_X\sigma_Y} \right) \mathbf{L}_{\frac{n-3}{2}} \left(\frac{n|x|}{\sigma_X\sigma_Y} \right) + \mathbf{L}_{\frac{n-1}{2}} \left(\frac{n|x|}{\sigma_X\sigma_Y} \right) K_{\frac{n-3}{2}} \left(\frac{n|x|}{\sigma_X\sigma_Y} \right) \right],$$

where $\mathbf{L}_\nu(x)$ is a modified Struve function of the second kind. By using the fact that \bar{Z}_n follows the McKay Type II distribution (a VG distribution), other formulas for the CDF in the case $\rho = 0$ can be obtained from results of Jankov Maširević and Pogány [23] and Nadarajah, Srivastava and Gupta [33].

Now suppose that $n \in 2\mathbb{Z}^+$ and $-1 < \rho < 1$. Then, making use of the formula (2.6) for the PDF of \bar{Z}_n and calculating $F_{\bar{Z}_n}(x) = \int_{-\infty}^x f_{\bar{Z}_n}(t) dt$ for $x \leq 0$, and $F_{\bar{Z}_n}(x) = 1 - \int_x^{\infty} f_{\bar{Z}_n}(t) dt$ for $x > 0$, yields the following formulas. For $x \leq 0$,

$$F_{\bar{Z}_n}(x) = \frac{(1-\rho)^{n/2}}{2^{n/2}(n/2-1)!} \sum_{j=0}^{n/2-1} \frac{(n/2-1+j)!}{(n/2-1-j)!j!} \left(\frac{1+\rho}{2}\right)^j \Gamma\left(\frac{n}{2}-j, \frac{-nx}{\sigma_X\sigma_Y(1-\rho)}\right),$$

and, for $x > 0$,

$$F_{\bar{Z}_n}(x) = 1 - \frac{(1+\rho)^{n/2}}{2^{n/2}(n/2-1)!} \sum_{j=0}^{n/2-1} \frac{(n/2-1+j)!}{(n/2-1-j)!j!} \left(\frac{1-\rho}{2}\right)^j \Gamma\left(\frac{n}{2}-j, \frac{nx}{\sigma_X\sigma_Y(1+\rho)}\right),$$

where $\Gamma(a, x) = \int_x^{\infty} t^{a-1}e^{-t} dt$ is the upper incomplete gamma function. Nadarajah, Srivastava and Gupta [33] also gave an analogue of these formulas for the case $\rho = 0$ for the CDF of the McKay Type II distribution.

Suppose now that we are in the general setting $n \geq 1$, $-1 < \rho < 1$ and $\sigma_X, \sigma_Y > 0$. As a closed-form formula is not available for the CDF, the following asymptotic approximations for the tail probabilities are of interest. Let $\bar{F}_{\bar{Z}_n}(x) = 1 - F_{\bar{Z}_n}(x) = \mathbb{P}(\bar{Z}_n > x)$. Then, from (2.2) and the limiting form (A.35), we get that

$$\bar{F}_{\bar{Z}_n}(x) \sim \frac{n^{n/2-1}(1+\rho)}{2^{n/2}(\sigma_X\sigma_Y)^{n/2-1}\Gamma(n/2)} x^{n/2-1} \exp\left(-\frac{nx}{\sigma_X\sigma_Y(1+\rho)}\right), \quad x \rightarrow \infty, \quad (2.10)$$

and, by symmetry,

$$F_{\bar{Z}_n}(x) \sim \frac{n^{n/2-1}(1-\rho)}{2^{n/2}(\sigma_X\sigma_Y)^{n/2-1}\Gamma(n/2)} (-x)^{n/2-1} \exp\left(\frac{nx}{\sigma_X\sigma_Y(1-\rho)}\right), \quad x \rightarrow -\infty.$$

More generally, the tail behaviour of semi-heavy tailed distributions is described in Proposition 2 of Hammerstein [21]. Upper and lower bounds on $F_{\bar{Z}_n}(x)$ and $\bar{F}_{\bar{Z}_n}(x)$ can also be obtained by using bounds for the integral $\int_x^{\infty} e^{\beta t^\nu} K_\nu(t) dt$, $x > 0$, $-1 < \beta < 1$, $\nu > -1/2$, given in Gaunt [12, 13].

2.4 Generating functions and infinite divisibility

The moment generating function of Z was obtained by Craig [5] (without knowledge of the PDF (2.1)) and the moment generating (and characteristic) function of \bar{Z}_n can then be deduced from basic properties of moment generating (and characteristic) functions (see Nadarajah and Pogány [32]):

$$M_{\bar{Z}_n}(t) = \mathbb{E}[e^{t\bar{Z}_n}] = \left(1 - \frac{2\rho\sigma_X\sigma_Y t}{n} - \frac{\sigma_X^2\sigma_Y^2(1-\rho^2)t^2}{n^2}\right)^{-n/2},$$

which exists if $-n/(1-\rho) < \sigma_X\sigma_Y t < n/(1+\rho)$ (see (2.4) and (2.5)). The characteristic function is

$$\varphi_{\bar{Z}_n}(t) = \mathbb{E}[e^{it\bar{Z}_n}] = \left(1 - \frac{2\rho\sigma_X\sigma_Y it}{n} + \frac{\sigma_X^2\sigma_Y^2(1-\rho^2)t^2}{n^2}\right)^{-n/2}. \quad (2.11)$$

Alternatively, given the formula (2.2) for the PDF of \bar{Z}_n , the moment generating function is obtained from a simple integration of $M_{\bar{Z}_n}(t) = \int_{-\infty}^{\infty} e^{tx} f_{\bar{Z}_n}(x) dx$; see, for example, Eberlein and Hammerstein [8] for a calculation for the more general generalized hyperbolic distribution (which includes the VG distribution as a limiting case).

The cumulant generating function $K_{\bar{Z}_n}(t) = \log \mathbb{E}[e^{t\bar{Z}_n}]$, defined for $-n/(1-\rho) < \sigma_X \sigma_Y t < n/(1+\rho)$, is given by

$$\begin{aligned} K_{\bar{Z}_n}(t) &= -\frac{n}{2} \log \left(1 - \frac{2\rho\sigma_X\sigma_Y t}{n} - \frac{\sigma_X^2\sigma_Y^2(1-\rho^2)t^2}{n^2} \right) \\ &= -\frac{n}{2} \log \left(1 - \frac{\sigma_X\sigma_Y(\rho+1)t}{n} \right) - \frac{n}{2} \log \left(1 - \frac{\sigma_X\sigma_Y(\rho-1)t}{n} \right). \end{aligned} \quad (2.12)$$

For any $m \geq 1$, we can write $\bar{Z}_n \stackrel{d}{=} S_1 + S_2 + \dots + S_m$, where S_1, S_2, \dots, S_m are independent $\text{VG}(n/m, \rho\sigma_X\sigma_Y/n + \sigma_X\sigma_Y\sqrt{1-\rho^2}/n, 0)$ random variables. Thus, the distribution of \bar{Z}_n is infinitely divisible. This follows from the representation (2.9) of \bar{Z}_n as a VG distribution and the convolution property that if $V_1 \sim \text{VG}(r_1, \theta, \sigma, 0)$ and $V_2 \sim \text{VG}(r_2, \theta, \sigma, 0)$ are independent then $V_1 + V_2 \sim \text{VG}(r_1 + r_2, \theta, \sigma, 0)$ (Bibby and Sørensen [4]), which is easily verified by using the fact that $\varphi_V(t) = (1 - 2\theta t + \sigma^2 t^2)^{-r/2}$ is the characteristic function of $V \sim \text{VG}(r, \theta, \sigma, 0)$.

2.5 Representation in terms of other random variables

1. Let X_1, \dots, X_n and W_1, \dots, W_n be independent $N(0, 1)$ random variables. Then

$$\bar{Z}_n \stackrel{d}{=} \frac{\sigma_X\sigma_Y}{n} \sum_{j=1}^n (\sqrt{1-\rho^2}X_jW_j + \rho X_j^2).$$

It suffices to show that $Z \stackrel{d}{=} \sigma_X\sigma_Y(\sqrt{1-\rho^2}X_1W_1 + \rho X_1^2)$. For ease of notation, we suppose that $\sigma_X = \sigma_Y = 1$, with the general case following by rescaling. It is straightforward to verify that X and $W = (Y - \rho X)/\sqrt{1-\rho^2}$ are independent $N(0, 1)$ random variables. Thus, $Z = XY = X(\sqrt{1-\rho^2}W + \rho X) = \sqrt{1-\rho^2}XW + \rho X^2$, as required.

2. Suppose that $S \sim \chi_{(n)}^2$ and $T \sim N(0, 1)$ are independent. Then

$$\bar{Z}_n \stackrel{d}{=} \frac{\rho\sigma_X\sigma_Y}{n} S + \frac{\sigma_X\sigma_Y\sqrt{1-\rho^2}}{n} \sqrt{ST}. \quad (2.13)$$

The representation (2.13) follows from the representation (2.9) of \bar{Z}_n as a VG random variable and Proposition 4.1.2 of Kotz, Kozubowski and Podgórski [25], which states that if $S' \sim \Gamma(r/2, 1/2)$ and $T' \sim N(0, 1)$ are independent, then $\theta S' + \sigma\sqrt{S'T'} \sim \text{VG}(r, \theta, \sigma, 0)$.

3. Suppose that V and V' are independent $\chi_{(n)}^2$ random variables. Then

$$\bar{Z}_n \stackrel{d}{=} \frac{\sigma_X\sigma_Y}{2n} (1+\rho)V - \frac{\sigma_X\sigma_Y}{2n} (1-\rho)V'. \quad (2.14)$$

This representation of \overline{Z}_n is easily proved using a standard characteristic function argument with the formula (2.11) for the characteristic function of \overline{Z}_n and the standard formula for the characteristic function of the $\chi_{(n)}^2$ distribution, $\varphi_V(t) = (1 - 2it)^{-n/2}$.

From the representation (2.14) and the representation of the $\chi_{(n)}^2$ distribution as the sum of the squares of n independent $N(0, 1)$ random variables, it follows that

$$\overline{Z}_n =_d \rho\sigma_X\sigma_Y + \sum_{j=1}^{2n} \lambda_j(N_j^2 - 1),$$

where N_1, \dots, N_{2n} are independent $N(0, 1)$ random variables, and $\lambda_1 = \dots = \lambda_n = \sigma_X\sigma_Y(1 + \rho)/(2n)$ and $\lambda_{n+1} = \dots = \lambda_{2n} = \sigma_X\sigma_Y(\rho - 1)/(2n)$. Thus, \overline{Z}_n is a member of the second Wiener chaos (see Section 2.2 of Nourdin and Peccati [34]).

4. If $n \in 2\mathbb{Z}^+$, then the independent random variables $V \sim \chi_{(n)}^2$ and $V' \sim \chi_{(n)}^2$ can be expressed in terms of independent uniform $U(0, 1)$ random variables U_1, \dots, U_n as $V =_d \sum_{j=1}^{n/2} \log U_j$ and $V' =_d \sum_{j=n/2+1}^n \log U_j$. Therefore, from (2.14),

$$\overline{Z}_n =_d \frac{\sigma_X\sigma_Y}{2n}(1 + \rho) \sum_{j=1}^{n/2} \log U_j - \frac{\sigma_X\sigma_Y}{2n}(1 - \rho) \sum_{j=n/2+1}^n \log U_j. \quad (2.15)$$

The representation (2.15) is convenient for simulating the distribution of \overline{Z}_n when $n \in 2\mathbb{Z}^+$. If n is an odd integer, the distribution of \overline{Z}_n can be simulated using the representation (2.14) and simulating the chi-square distributions of V and V' using methods for simulating gamma distributions, as given in Chapter 9, Section 3 of Devroye [7].

2.6 Stein characterisation

Let W be a real-valued random variable. Then $W =_d \overline{Z}_n$ if and only if

$$\mathbb{E} \left[\frac{\sigma_X^2\sigma_Y^2(1 - \rho^2)}{n^2} W g''(W) + \frac{1}{n} (\sigma_X^2\sigma_Y^2(1 - \rho^2) + 2\rho\sigma_X\sigma_Y W) g'(W) + (\rho\sigma_X\sigma_Y - W) g(W) \right] = 0 \quad (2.16)$$

for all twice differentiable $g : \mathbb{R} \rightarrow \mathbb{R}$ such that the expectations $\mathbb{E}|g(\overline{Z})|$, $\mathbb{E}|\overline{Z}g(\overline{Z})|$, $\mathbb{E}|g'(\overline{Z})|$, $\mathbb{E}|\overline{Z}g'(\overline{Z})|$ and $\mathbb{E}|\overline{Z}g''(\overline{Z})|$ are all finite.

Necessity of the Stein characterisation (2.16) was established by Gaunt [15] (here we have corrected a typo in the characterising equation (2.16)). Let us give a simple proof of sufficiency. Suppose W is a real-valued random variable. To simplify notation, we set $\sigma_X = \sigma_Y = 1$, with the general case following by rescaling. Taking $g(x) = e^{itx}$ (which is twice differentiable and bounded) in (2.16) and setting $\varphi(t) = \mathbb{E}[e^{itW}]$ yields the differential equation

$$\left(\frac{(1 - \rho^2)}{n^2} t^2 - \frac{2\rho i}{n} t + 1 \right) \varphi'(t) + \left(\frac{(1 - \rho^2)}{n} t - \rho i \right) \varphi(t) = 0. \quad (2.17)$$

Note that $g(x) = e^{itx}$ is a complex-valued function; here we applied the characterising equation to the real and imaginary parts of g , which are real-valued functions. Solving (2.17) subject to the condition $\varphi(0) = 1$ gives that $\varphi(t) = (1 - 2\rho it/n + (1 - \rho^2)t^2/n^2)^{-n/2}$, which is the characteristic function (2.11) of \bar{Z}_n . Thus, $W =_d \bar{Z}_n$, which proves sufficiency.

Stein characterisations are most commonly used as part of Stein's method (Stein [40]) to prove quantitative limit theorems in probability theory; however, they can also be used to derive distributional properties. For example, setting $g_1(x) = x^k$ and $g_2(x) = (x - \mathbb{E}[\bar{Z}_n])^k = (x - \rho\sigma_X\sigma_Y)^k$ in (2.16) yields the following recursions for the k -th raw moment $\mu_k = \mathbb{E}[\bar{Z}_n^k]$ and the k -th central moment $\mu'_k = \mathbb{E}[(\bar{Z}_n - \mathbb{E}[\bar{Z}_n])^k]$:

$$\mu'_{k+1} = \frac{n + 2k}{n} \rho\sigma_X\sigma_Y\mu'_k + \frac{k(n + k - 1)}{n^2} \sigma_X^2 \sigma_Y^2 (1 - \rho^2) \mu'_{k-1}, \quad k \geq 1, \quad (2.18)$$

and

$$\begin{aligned} \mu_{k+1} &= \frac{2k}{n} \rho\sigma_X\sigma_Y\mu_k + \frac{k\sigma_X^2\sigma_Y^2}{n^2} (n + k - 1 + (n - k + 1)\rho^2) \mu_{k-1} \\ &\quad + \frac{k(k-1)}{n^2} \sigma_X^3\sigma_Y^3\rho(1 - \rho^2) \mu_{k-2}, \quad k \geq 2. \end{aligned} \quad (2.19)$$

These recurrences allow for the efficient computation of lower order raw (central) moments given just the first raw (first and second central) moments.

2.7 Moments and cumulants

The mean and variance of \bar{Z} can be easily calculated through several approaches; for example, using the standard method of extracting lower order moments via moment generating functions. We have

$$\mathbb{E}[\bar{Z}] = \rho\sigma_X\sigma_Y, \quad (2.20)$$

$$\text{Var}(\bar{Z}) = \frac{\sigma_X^2\sigma_Y^2}{n} (1 + \rho^2). \quad (2.21)$$

Lower order raw and central moments are then readily obtained from the recurrences (2.18) and (2.19). The first four raw moments are given by

$$\begin{aligned} \mu'_1 &= \rho\sigma_X\sigma_Y, \\ \mu'_2 &= \frac{\sigma_X^2\sigma_Y^2}{n} (1 + (n + 1)\rho^2), \\ \mu'_3 &= \frac{\rho\sigma_X^3\sigma_Y^3}{n^2} (3(n + 2) + (n + 1)(n + 2)\rho^2), \\ \mu'_4 &= \frac{\sigma_X^4\sigma_Y^4}{n^3} (3(n + 2) + 6(n + 2)(n + 3)\rho^2 + (n + 1)(n + 2)(n + 3)\rho^4), \end{aligned}$$

and the first four central moments are given by

$$\begin{aligned} \mu_1 &= 0, \\ \mu_2 &= \frac{\sigma_X^2\sigma_Y^2}{n} (1 + \rho^2), \end{aligned}$$

$$\begin{aligned}\mu_3 &= \frac{2\rho\sigma_X^3\sigma_Y^3}{n^2}(3 + \rho^2), \\ \mu_4 &= \frac{3\sigma_X^4\sigma_Y^4}{n^3}((n+2) + 2(n+6)\rho^2 + (n+2)\rho^4).\end{aligned}$$

We thus deduce that the skewness $\gamma_1 = \mu_3/\mu_2^{3/2}$ and kurtosis $\beta_2 = \mu_4/\mu_2^2$ are given by

$$\gamma_1 = \frac{2\rho(3 + \rho^2)}{\sqrt{n}(1 + \rho^2)^{3/2}}, \quad \beta_2 = \frac{3(n+2) + 6(n+6)\rho^2 + 3(n+2)\rho^4}{n(1 + \rho^2)^2},$$

whilst the excess kurtosis $\gamma_2 = \mu_4/\mu_2^2 - 3$ is

$$\gamma_2 = \frac{6 + 36\rho^2 + 6\rho^4}{n(1 + \rho^2)^2}.$$

A formula for the skewness of the product of two correlated normal variables with possibly non-zero means is given by Ware and Lad [42].

Higher order moments are easily calculated using the representation (2.14) of \bar{Z}_n and the moment formula $\mathbb{E}[V^k] = 2^k\Gamma(n/2 + k)/\Gamma(n/2) = n(n+2)(n+4)\cdots(n+2k-2)$, $k \geq 1$, for $V \sim \chi_{(n)}^2$. For $k \geq 1$,

$$\begin{aligned}\mu'_k &= \frac{\sigma_X^k\sigma_Y^k}{(2n)^k} \sum_{j=0}^k \binom{k}{j} (\rho+1)^j (\rho-1)^{k-j} \mathbb{E}[V^j] \mathbb{E}[V^{k-j}] \\ &= \frac{\sigma_X^k\sigma_Y^k}{n^k(\Gamma(n/2))^2} \sum_{j=0}^k \binom{k}{j} \Gamma\left(\frac{n}{2} + j\right) \Gamma\left(\frac{n}{2} + k - j\right) (\rho+1)^j (\rho-1)^{k-j}.\end{aligned}$$

In the case $n = 1$, the following formula is available (see Kan [24]):

$$\mu'_k = \begin{cases} \sigma_X^k\sigma_Y^k \frac{(k!)^2}{2^k} \sum_{j=0}^{k/2} \frac{(2\rho)^{2j}}{((k/2 - j)!)^2 (2j)!}, & \text{if } k \text{ is even,} \\ \sigma_X^k\sigma_Y^k \frac{(k!)^2}{2^k} \sum_{j=0}^{(k-1)/2} \frac{(2\rho)^{2j+1}}{((k/2 - 1/2 - j)!)^2 (2j+1)!}, & \text{if } k \text{ is odd.} \end{cases}$$

Returning to the general case $n \geq 1$, the raw moments of \bar{Z}_n can be expressed in terms of the hypergeometric function (see Chapter 15 of Olver et al. [35]). For $k \geq 1$,

$$\begin{aligned}\mu'_k &= \frac{\sigma_X^k\sigma_Y^k(n+k-1)!k!}{n^k(1-\rho^2)^{n/2}\Gamma(n/2+k+1)\Gamma(n/2)} \times \\ &\quad \times \left[(-1)^k(1-\rho)^{n+k} {}_2F_1\left(n+k, \frac{n}{2}; \frac{n}{2} + k + 1; -\frac{1-\rho}{1+\rho}\right) \right. \\ &\quad \left. + (1+\rho)^{n+k} {}_2F_1\left(n+k, \frac{n}{2}; \frac{n}{2} + k + 1; -\frac{1+\rho}{1-\rho}\right) \right].\end{aligned}$$

This formula is a re-parameterisation (using (2.9)) of the result of Theorem 2 of Holm and Alouini [22], which gives formulas for the raw moments of the McKay Type II distribution (a VG distribution). The proof of Holm and Alouini [22] involves writing $\mu'_k = \int_{-\infty}^{\infty} x^k f_{\bar{Z}_n}(x) dx = \int_{-\infty}^0 x^k f_{\bar{Z}_n}(x) dx + \int_0^{\infty} x^k f_{\bar{Z}_n}(x) dx = (-1)^k \int_0^{\infty} x^k f_{\bar{Z}_n}(-x) dx + \int_0^{\infty} x^k f_{\bar{Z}_n}(x) dx$, calculating the latter two integrals using (A.34) and then simplifying. The absolute moments can be calculated similarly by writing $\mathbb{E}|\bar{Z}_n|^k = \int_0^{\infty} x^k f_{\bar{Z}_n}(-x) dx + \int_0^{\infty} x^k f_{\bar{Z}_n}(x) dx$. This gives that, for $k \geq 1$,

$$\begin{aligned} \mathbb{E}|\bar{Z}_n|^k &= \frac{\sigma_X^k \sigma_Y^k (n+k-1)! k!}{n^k (1-\rho^2)^{n/2} \Gamma(n/2+k+1) \Gamma(n/2)} \times \\ &\times \left[(1-\rho)^{n+k} {}_2F_1\left(n+k, \frac{n}{2}; \frac{n}{2}+k+1; -\frac{1-\rho}{1+\rho}\right) \right. \\ &\quad \left. + (1+\rho)^{n+k} {}_2F_1\left(n+k, \frac{n}{2}; \frac{n}{2}+k+1; -\frac{1+\rho}{1-\rho}\right) \right]. \end{aligned}$$

Higher order central moments of \bar{Z}_n can be calculated using the representation (2.14) of \bar{Z}_n and the formula $\mathbb{E}[(V - \mathbb{E}[V])^k] = 2^k U(-k, 1-k-n/2, -n/2)$, where $U(a, b, x)$ is a confluent hypergeometric function of the second kind (see Weisstein [44]). We note that in the case $a = -m$, $m = 0, 1, 2, \dots$, the function $U(a, b, x)$ is a polynomial: $U(-m, b, x) = (-1)^m \sum_{j=0}^m \binom{m}{j} (b+j)_{m-j} (-x)^j$, and the Pochhammer symbol is $(a)_0 = 0$ and $(a)_j = a(a+1)(a+2) \cdots (a+j-1)$, $j \geq 1$ (see Section 13.2(i) of Olver et al. [35]). We have, for $k \geq 1$,

$$\begin{aligned} \mu_k &= \frac{\sigma_X^k \sigma_Y^k}{(2n)^k} \sum_{j=0}^k \binom{k}{j} (\rho+1)^j (\rho-1)^{k-j} \mathbb{E}[(V - \mathbb{E}[V])^j] \mathbb{E}[(V - \mathbb{E}[V])^{k-j}] \\ &= \frac{\sigma_X^k \sigma_Y^k}{n^k} \sum_{j=0}^k \binom{k}{j} U\left(-j, 1-j-\frac{n}{2}, -\frac{n}{2}\right) U\left(j-k, 1+j-k-\frac{n}{2}, -\frac{n}{2}\right) \times \\ &\quad \times (1+\rho)^j (\rho-1)^{k-j}. \end{aligned}$$

Suppose now that $\rho = 0$. Then, from (2.13), $\bar{Z}_n =_d (\sigma_X \sigma_Y / n) \sqrt{ST}$, where $S \sim \chi_{(n)}^2$ and $T \sim N(0, 1)$ are independent. If $k \geq 1$ is odd then $\mathbb{E}[T^k] = 0$, whilst for any $\alpha > 0$, $\mathbb{E}|T^\alpha| = \pi^{-1/2} 2^{\alpha/2} \Gamma((\alpha+1)/2)$ and $\mathbb{E}|S^\alpha| = \Gamma(n/2 + \alpha) / \Gamma(n/2)$. Therefore, for $k \geq 1$,

$$\mu'_k = \begin{cases} \frac{2^{k/2} \sigma_X^k \sigma_Y^k}{\sqrt{\pi} n^k} \frac{\Gamma((n+k)/2) \Gamma((k+1)/2)}{\Gamma(n/2)}, & \text{if } k \text{ is even,} \\ 0, & \text{if } k \text{ is odd.} \end{cases}$$

Suppose now that $-1 < \rho < 1$ and $n \geq 1$. The cumulants of \bar{Z}_n are readily calculated by either Taylor expanding the logarithms in (2.12), or by working with the representation (2.14) of \bar{Z}_n . In the latter approach, one uses the standard properties that for independent random variables S_1 and S_2 and constant c , the k -th cumulant satisfies $\kappa_k(cS_1) = c^k \kappa_k(S_1)$ and $\kappa_k(S_1 + S_2) = \kappa_k(S_1) + \kappa_k(S_2)$, and $\kappa_k(V) = 2^{k-1} (k-1)! n$ for $V \sim \chi_{(n)}^2$. For $k \geq 1$,

$$\kappa_k = \frac{\sigma_X^k \sigma_Y^k}{2n^{k-1}} (k-1)! [(1+\rho)^k + (\rho-1)^k].$$

In particular,

$$\begin{aligned}\kappa_1 &= \rho\sigma_X\sigma_Y, \\ \kappa_2 &= \frac{\sigma_X^2\sigma_Y^2}{n}(1 + \rho^2), \\ \kappa_3 &= \frac{2\rho\sigma_X^3\sigma_Y^3}{n^2}(3 + \rho^2), \\ \kappa_4 &= \frac{6\sigma_X^4\sigma_Y^4}{n^3}(1 + 6\rho^2 + \rho^4).\end{aligned}$$

2.8 Mode and median

A detailed study of the mode and median of the generalized hyperbolic and VG distributions was recently carried out by Gaunt and Merkle [16]; here we provide a synthesis of this work in the case of the product of correlated zero mean normal random variables.

The distribution of \bar{Z}_n is unimodal. This follows because \bar{Z}_n is self-decomposable (it is infinitely divisible) and self-decomposable distributions are unimodal (Yamazato [45]). Let M_n denote the mode of \bar{Z}_n . It is clear from (2.3) and (2.7) that

$$M_1 = M_2 = 0. \quad (2.22)$$

Suppose now that $n \geq 3$. Then, by applying the differentiation formula (A.32) to the PDF (2.2), we deduce that $M_n = \text{sgn}(\rho) \cdot x^*$, where x^* is the unique positive solution of the equation

$$K_{\frac{n-3}{2}}\left(\frac{nx}{\sigma_X\sigma_Y(1-\rho^2)}\right) = |\rho|K_{\frac{n-1}{2}}\left(\frac{nx}{\sigma_X\sigma_Y(1-\rho^2)}\right). \quad (2.23)$$

In the cases $n = 4$ and $n = 6$, we can apply the formulas in (A.29) to (2.23) to obtain simple algebraic equations for x^* which when solved yield the exact expressions

$$M_4 = \frac{\rho\sigma_X\sigma_Y}{4}(1 + |\rho|), \quad M_6 = \frac{\rho\sigma_X\sigma_Y}{12}(1 + |\rho|)\left(3 - \frac{1}{|\rho|} + \sqrt{\frac{1}{\rho^2} + \frac{6}{|\rho|} - 3}\right).$$

For $n = 8$ and $n = 10$, we can apply the formula (A.28) to (2.23) to obtain cubic and quartic equations for x^* , although the solutions to these equations are too complicated to be worth reporting. For other values of n , exact formulas for M_n are not available, except for the case $\rho = 0$ in which case the mode is equal to zero.

Whilst exact formulas are not available for M_n for general $n \geq 3$, simple and accurate upper and lower bounds can be derived. In what follows, we fix $\rho > 0$; bounds for the case $\rho < 0$ follow by symmetry. Applying the lower and upper bounds of (A.36) to (2.23) leads to simple algebraic equations for x^* which when solved lead to the upper and lower bounds of the following two-sided inequality, respectively:

$$\rho\sigma_X\sigma_Y(1 - 3/n) < M_n < \rho\sigma_X\sigma_Y(1 - 2/n), \quad n \geq 3. \quad (2.24)$$

There is equality in the upper bound when $n = 2$ (see (2.22)). Applying inequality (A.37) to (2.23) yields that, for $n \geq 4$,

$$M_n \geq \frac{\rho\sigma_X\sigma_Y}{2}\left[1 - \frac{2}{n} + \sqrt{\rho^2\left(1 - \frac{2}{n}\right)^2 + (1 - \rho^2)\left(1 - \frac{4}{n}\right)^2}\right], \quad (2.25)$$

with equality if and only if $n = 4$. Inequality (2.25) is more accurate than the lower bound in (2.24) for $n \geq 4$. The bounds in (2.24) and (2.25) were derived by Gaunt and Merkle [16] by similar considerations.

If $\rho = 0$, the median of \bar{Z}_n is equal to zero. The median of \bar{Z}_2 is given by

$$\text{Med}(\bar{Z}_2) = \begin{cases} \frac{\sigma_X \sigma_Y}{2} (1 + \rho) \log(1 + \rho), & \rho \geq 0, \\ -\frac{\sigma_X \sigma_Y}{2} (1 - \rho) \log(1 - \rho), & \rho < 0, \end{cases} \quad (2.26)$$

which follows from the well-known formula for the median of the asymmetric Laplace distribution (see Kozubowski and Podgórski [26]). Otherwise, an exact closed-form formula is not available for $\text{Med}(\bar{Z}_n)$, $n \geq 1$. Moreover, unlike for the mode, accurate upper and lower bounds for the median have yet to be worked out in the literature. Gaunt and Merkle [16] have, however, conjectured accurate bounds for the median of the VG distribution. As \bar{Z}_n follows the VG distribution (see (2.9)), we can present conjectured bounds for $\text{Med}(\bar{Z}_n)$. The numerical results of Table 1 support the conjectured bounds. The results were obtained with *Mathematica* via the same simple numerical procedure for numerically computing medians as described in pp. 16–17 of Gaunt and Merkle [16].

Conjecture 2.1. *Suppose $\rho > 0$. Then it is conjectured that*

$$\left(1 - \frac{1}{n}\right) \rho \sigma_X \sigma_Y < \text{Med}(\bar{Z}_n) < \rho \sigma_X \sigma_Y e^{-2/3n} < \left(1 - \frac{2}{3n} + \frac{2}{9n^2}\right) \rho \sigma_X \sigma_Y, \quad n \geq 1,$$

and

$$\text{Med}(\bar{Z}_n) \leq \left(1 - \frac{2(1 - \log 2)}{n}\right) \rho \sigma_X \sigma_Y, \quad n \geq 2. \quad (2.27)$$

Table 1: Median of \bar{Z}_n with $\sigma_X = \sigma_Y = 1$.

$n \backslash \rho$	0.1	0.3	0.5	0.7	0.9
1	0.0198	0.0813	0.164	0.265	0.386
3	0.0674	0.210	0.364	0.528	0.700
5	0.0802	0.245	0.416	0.594	0.777
7	0.0859	0.260	0.439	0.623	0.812
10	0.0901	0.272	0.457	0.646	0.838

We end by noting that because $\mathbb{E}[\bar{Z}_n] = \rho \sigma_X \sigma_Y$, it follows from (2.22) and (2.24) that, if $\rho > 0$, then $M_n < \mathbb{E}[\bar{Z}_n]$ for all $n \geq 1$. In fact, we have the two-sided inequality $2\sigma_X \sigma_Y \rho / n < \mathbb{E}[\bar{Z}_n] - M_n < 3\sigma_X \sigma_Y \rho / n$, $n \geq 3$, with equality in the lower bound if $n = 2$. If the conjectured median bounds hold, then it would follow that, for $\rho > 0$, $M_n < \text{Med}(\bar{Z}_n) < \mathbb{E}[\bar{Z}_n]$, meaning that \bar{Z}_n would satisfy the mean-median-mode inequality (Groeneveld and Meeden [19] and van Zwet [41]). We also note that as $n \rightarrow \infty$, M_n and $\text{Med}(\bar{Z}_n)$ converge to $\mathbb{E}[\bar{Z}_n] = \rho \sigma_X \sigma_Y$, which is to be expected given that \bar{Z}_n is approximately normally distributed (for which the mean, median and mode are equal) for large n .

A The modified Bessel function of the second kind

In this appendix, we collect some properties of the modified Bessel function of the second kind that are needed in this paper. Unless otherwise stated, these properties can be found in Olver et al. [35]. The *modified Bessel function of the second kind* $K_\nu(x)$ is defined, for $\nu \in \mathbb{R}$ and $x > 0$, by

$$K_\nu(x) = \int_0^\infty e^{-x \cosh(t)} \cosh(\nu t) dt.$$

For $x > 0$, the function $K_\nu(x)$ is positive for all $\nu \in \mathbb{R}$. For $\nu = m + 1/2$, $m = 0, 1, 2, \dots$, we have

$$K_{m+1/2}(x) = \sqrt{\frac{\pi}{2x}} \sum_{j=0}^m \frac{(m+j)!}{(m-j)!j!} (2x)^{-j} e^{-x}. \quad (\text{A.28})$$

In particular,

$$K_{1/2}(x) = \sqrt{\frac{\pi}{2x}} e^{-x}, \quad K_{3/2}(x) = \sqrt{\frac{\pi}{2x}} \left(1 + \frac{1}{x}\right) e^{-x}, \quad K_{5/2}(x) = \sqrt{\frac{\pi}{2x}} \left(1 + \frac{3}{x} + \frac{3}{x^2}\right) e^{-x}. \quad (\text{A.29})$$

The modified Bessel function of the second kind has the following asymptotic behaviour:

$$K_\nu(x) \sim \begin{cases} 2^{|\nu|-1} \Gamma(|\nu|) x^{-|\nu|}, & x \downarrow 0, \nu \neq 0, \\ -\log x, & x \downarrow 0, \nu = 0, \end{cases} \quad (\text{A.30})$$

$$K_\nu(x) \sim \sqrt{\frac{\pi}{2x}} e^{-x}, \quad x \rightarrow \infty, \nu \in \mathbb{R}. \quad (\text{A.31})$$

A differentiation formula is given by:

$$\frac{d}{dx} (x^\nu K_\nu(x)) = -x^\nu K_{\nu-1}(x). \quad (\text{A.32})$$

The following integral formulas can be found in Gradshteyn and Ryzhik [17]. For $a, \alpha > 0$ and $\nu > -1/2$,

$$\int_0^a t^\nu K_\nu(\alpha t) dt = \frac{\sqrt{\pi} 2^{\nu-1} \Gamma(\nu + 1/2)}{\alpha^\nu} a (K_\nu(\alpha a) \mathbf{L}_{\nu-1}(\alpha a) + K_{\nu-1}(\alpha a) \mathbf{L}_\nu(\alpha a)). \quad (\text{A.33})$$

For $\mu > |\nu|$ and $\beta < \alpha$,

$$\begin{aligned} \int_0^\infty t^{\mu-1} e^{\beta t} K_\nu(\alpha t) dt &= \frac{\sqrt{\pi} (2\alpha)^\nu}{(\alpha - \beta)^{\mu+\nu}} \frac{\Gamma(\mu + \nu) \Gamma(\mu - \nu)}{\Gamma(\mu + 1/2)} \times \\ &\times {}_2F_1\left(\mu + \nu, \nu + \frac{1}{2}; \mu + \frac{1}{2}; -\frac{\alpha + \beta}{\alpha - \beta}\right). \end{aligned} \quad (\text{A.34})$$

The following asymptotic approximation follows from a simple rescaling of limiting form (2.13) in Gaunt [13]. For $\beta < \alpha$ and $\nu > -1/2$,

$$\int_x^\infty e^{\beta t} t^\nu K_\nu(\alpha t) dt \sim \sqrt{\frac{\pi}{2\alpha}} \frac{1}{(\alpha - \beta)} x^{\nu-1/2} e^{-(\alpha-\beta)x}, \quad x \rightarrow \infty. \quad (\text{A.35})$$

The ratio $K_{\nu-1}(x)/K_\nu(x)$ satisfies the following inequalities. For $x > 0$,

$$\frac{x}{\nu - 1/2 + \sqrt{(\nu - 1/2)^2 + x^2}} < \frac{K_{\nu-1}(x)}{K_\nu(x)} < \frac{x}{\nu - 1 + \sqrt{(\nu - 1)^2 + x^2}}, \quad \nu > 1/2, \quad (\text{A.36})$$

and

$$\frac{K_{\nu-1}(x)}{K_\nu(x)} \leq \frac{x}{\nu - 1/2 + \sqrt{(\nu - 3/2)^2 + x^2}}, \quad \nu \geq 3/2, \quad (\text{A.37})$$

with equality if and only if $\nu = 3/2$. The lower and upper bounds in (A.36) were obtained by Segura [37] and Ruiz-Antolín and Segura [36], respectively, whilst inequality (A.37) can be found in Gaunt and Merkle [16].

Acknowledgements

The author is supported by a Dame Kathleen Ollerenshaw Research Fellowship.

References

- [1] Aroian, L. A. The probability function of the product of two normally distributed variables. *Ann. Math. Stat.* **18** (1947), pp. 265–271.
- [2] Aroian, L. A., Taneja, V. S. and Cornwell, L. W. Mathematical forms of the distribution of the product of two normal variables. *Commun. Stat. Theory* **7** (1978), pp. 165–172.
- [3] Bandi, M. M. and Connaughton, C. Craig’s XY distribution and the statistics of Lagrangian power in two-dimensional turbulence. *Phys. Rev. E* **77** (2008), 036318.
- [4] Bibby, B. M., and Sørensen, M. Hyperbolic Processes in Finance. In Rachev, S. (ed.), *Handbook of Heavy Tailed Distributions in Finance* (2003), pp. 211–248. Amsterdam: Elsevier Science.
- [5] Craig, C. C. On the Frequency Function of xy , *Ann. Math. Stat.* **7** (1936), pp. 1–15.
- [6] Cui, G., Yu, X. Iommelli, S. and Kong, L. Exact Distribution for the Product of Two Correlated Gaussian Random Variables. *IEEE Signal Process. Lett.* **23** (2016), pp. 1662–1666.
- [7] Devroye, L. *Non-Uniform Random Variate Generation*. New York: Springer-Verlag, 1986.
- [8] Eberlein, E. and Hammerstein E. Generalized Hyperbolic and Inverse Gaussian Distributions: Limiting Cases and Approximation of Processes. In: Dalang, R. C. Dozzi, M., Russo, F. (Eds.), Seminar on Stochastic Analysis, Random Fields and Applications IV, in: *Progress in Probability* **58** Birkhäuser Verlag, (2004), pp. 105–153.
- [9] Finlay, R. and Seneta, E. Option pricing with VG-like models. *Int. J. Theor. Appl. Finan.* **11** (2008), pp. 943–955.
- [10] Gaunt, R. E. *Rates of Convergence of Variance-Gamma Approximations via Stein’s Method*. DPhil thesis, University of Oxford, 2013.
- [11] Gaunt, R. E. Variance-Gamma approximation via Stein’s method. *Electron. J. Probab.* **19** no. 38 (2014), pp. 1–33.
- [12] Gaunt, R. E. Inequalities for modified Bessel functions and their integrals. *J. Math. Anal. Appl.* **420** (2014), pp. 373–386.
- [13] Gaunt, R. E. Inequalities for integrals of modified Bessel functions and expressions involving them. *J. Math. Anal. Appl.* **462** (2018), pp. 172–190.
- [14] Gaunt, R. E. A note on the distribution of the product of zero mean correlated normal random variables. *Stat. Neerl.* **73** (2019), pp. 176–179.

- [15] Gaunt, R. E. Stein's method and the distribution of the product of zero mean correlated normal random variables. *Commun. Stat. Theory* **50** (2021), pp. 280–285.
- [16] Gaunt, R. E. and Merkle, M. On bounds for the mode and median of the generalized hyperbolic and related distributions. *J. Math. Anal. Appl.* **493** (2021), Art. 124508, pp. 1–19.
- [17] Gradshteyn, I. S. and Ryzhik, I. M. *Table of Integrals, Series and Products*. seventh ed. Academic Press, 2007.
- [18] Grishchuk, L. P. Statistics of the Microwave Background Anisotropies Caused by the Squeezed Cosmological Perturbations. *Phys. Rev. D* **53** (1996), no. 6784.
- [19] Groeneveld, R. A. and Meeden, G. The Mode, Median, and Mean Inequality. *Amer. Stat.* **31** (1977), pp. 120–121.
- [20] Haldane, J. B. S. Moments of the distributions of powers and products of normal variates. *Biometrika* **32** (1942), pp. 226–242.
- [21] Hammerstein, E. A. V. Tail behaviour and tail dependence of generalized hyperbolic distributions. In: Kallsen, J., Papapantoleon, A. (Eds.) *Advanced Modelling in Mathematical Finance - A Festschrift in honour of Ernst Eberlein*, Springer (2016), pp. 3–40.
- [22] Holm, H. and Alouini, M.–S. Sum and Difference of two squared correlated Nakagami variates with the McKay distribution. *IEEE T. Commun.* **52** (2004), pp. 1367–1376.
- [23] Jankov Maširević, D. and Pogány, T. K. On new formulae for cumulative distribution function for McKay Bessel distribution. *Commun. Stat. Theory* **50** (2021), pp. 143–160.
- [24] Kan, R. From moments of sum to moments of product. *J. Multivariate Anal.* **99** (2008), pp. 542–554.
- [25] Kotz, S., Kozubowski, T. J. and Podgórski, K. *The Laplace Distribution and Generalizations: A Revisit with New Applications*. Springer, 2001.
- [26] Kozubowski, T. J. and Podgórski, K. Asymmetric Laplace Laws and Modeling Financial Data. *Math. Comput. Model.* **34** (2001), pp. 1003–1021.
- [27] MacKinnon, D. P. *Introduction to Statistical Mediation Analysis*. Routledge, New York, 2012.
- [28] MacKinnon, D. P., Fritz, M. S., Williams, J. and Lockwood, C. M. Distribution of the product confidence limits for the indirect effect: program PRODCLIN. *Behav. Res. Methods* **39** (2007), pp. 384–389.
- [29] Mangilli, A., Plaszczyński, S. and Tristram, M. Large-scale cosmic microwave background temperature and polarization cross-spectra likelihoods. *Mon. Not. R. Astron. Soc.* **453** (2015), pp. 3174–3189.
- [30] McKay, A. T. A Bessel function distribution. *Biometrika* **24** (1932), pp. 39–44.
- [31] Meeker, W. Q., Odeh, R. D., Cornwell, L. W. Aroian, L. A. and Kennedy, W. J. *Selected Tables in Mathematical Statistics: The Product of Two Normally Distributed Random Variables*. American Mathematical Society, Providence, RI, USA (1981).
- [32] Nadarajah, S. and Pogány, T. K. On the distribution of the product of correlated normal random variables. *C.R. Acad. Sci. Paris, Ser. I* **354** (2016), pp. 201–204.
- [33] Nadarajah, S., Srivastava, H. M. and Gupta, A. K. Skewed Bessel function distributions with application to rainfall data. *Statistics* **41** (2007), pp. 333–344.
- [34] Nourdin, I. and Peccati, G. Normal approximations with Malliavin calculus: from Stein's method to universality. Vol. 192. Cambridge University Press, 2012.
- [35] Olver, F. W. J., Lozier, D. W., Boisvert, R. F. and Clark, C. W. *NIST Handbook of Mathematical Functions*. Cambridge University Press, 2010.
- [36] Ruiz-Antolín, D. and Segura, J. A new type of sharp bounds for ratios of modified Bessel functions. *J. Math. Anal. Appl.* **443** (2016), pp. 1232–1246.
- [37] Segura, J. Bounds for ratios of modified Bessel functions and associated Turán-type inequalities. *J. Math. Anal. Appl.* **374** (2011), pp. 516–528.
- [38] Springer, M. D. *The Algebra of Random Variables*. John Wiley & Sons Inc, 1979.

- [39] Springer, M. D. and Thompson, W. E. The distribution of products of Beta, Gamma and Gaussian random variables. *SIAM J. Appl. Math.* **18** (1970), pp. 721–737.
- [40] Stein, C. A bound for the error in the normal approximation to the distribution of a sum of dependent random variables. In *Proc. Sixth Berkeley Symp. Math. Statis. Prob.* (1972), vol. 2, Univ. California Press, Berkeley, pp. 583–602.
- [41] van Zwet, W. R. Mean, median, mode II. *Stat. Neerl.* **33** (1979), pp. 1–5.
- [42] Ware, R. and Lad, F. Approximating the distribution for sums of products of normal variables. Working paper, Department of Mathematics and Statistics, University of Canterbury, New Zealand, 2013.
- [43] Watts, D. J., Addison, G. E., Bennett, C. L. and Weiland, J. L. Beyond Optical Depth: Future Determination of Ionization History from the Cosmic Microwave Background. *Astrophys. J.* **889**:130 (2020), pp. 1–9.
- [44] Weisstein, E. W. Chi-Squared Distribution. From MathWorld—A Wolfram Web Resource, <https://mathworld.wolfram.com/Chi-SquaredDistribution.html>. Last visited on 31/05/2021.
- [45] Yamazato, M. Unimodality of infinitely divisible distribution functions of class L. *Ann. Probab.* **6** (1978), pp. 523–531.