

# Identifiability of interaction kernels in mean-field equations of interacting particles

Quanjun Lang\*, Fei Lu\*

## Abstract

We completely characterize the identifiability of interaction kernels in mean-field equations for interacting particle systems. The key is to identify function spaces on which a probabilistic loss functional has a unique minimizer. We prove that identifiability holds on any subspace of two reproducing kernel Hilbert spaces (RKHS) that are intrinsic to the system and data-adaptive. Furthermore, the identifiability holds on  $L^2$  spaces if and only if the integral operators associated with the reproducing kernels are strictly positive. Thus, the inverse problem is ill-posed. Subsequently, we propose RKHS-based regularization and numerically demonstrate that it improves the accuracy of the estimator.

Keywords: mean-field, interacting particles, RKHS, regularization, inverse problem.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Main results</b>	<b>3</b>
2.1	Definition of identifiability . . . . .	3
2.2	The ambient function spaces of regression . . . . .	5
2.3	Main results . . . . .	6
<b>3</b>	<b>Radial interaction kernel</b>	<b>7</b>
3.1	The loss function . . . . .	7
3.2	The RKHS space and $L^2$ space . . . . .	9
3.3	The weighted $L^2$ space and RKHS . . . . .	11
<b>4</b>	<b>General interaction kernels</b>	<b>13</b>
4.1	The loss function . . . . .	13
4.2	The RKHS space and $L^2$ space . . . . .	14
4.3	The weighted $L^2$ space and RKHS . . . . .	15
4.4	Examples and general function spaces . . . . .	17
<b>5</b>	<b>Identifiability in computational practice</b>	<b>18</b>
5.1	Nonparametric regression in practice . . . . .	18
5.2	Identifiability and ill-conditioned normal matrix . . . . .	19
5.3	Two regularization norms . . . . .	20
5.4	Regularization and the RKHSs . . . . .	22
<b>A</b>	<b>Appendix</b>	<b>24</b>

---

\*Department of Mathematics, Johns Hopkins University. Email: qlang1@math.jhu.edu; feilu@math.jhu.edu

# 1 Introduction

We study the identifiability of the *interaction kernel*  $K_{true}$  from data consisting a solution of the mean-field equation of interacting particles

$$\begin{aligned} \partial_t u &= \nu \Delta u + \operatorname{div}[u(K_{true} * u)], \quad x \in D \subset \mathbb{R}^d \\ \partial u|_{\partial D} &= 0, \quad \nabla u|_{\partial D} = 0, \end{aligned} \quad (1.1)$$

where the domain  $D \subset \mathbb{R}^d$  is connected with smooth boundary and  $K_{true} = \nabla \Phi_{true} : \mathbb{R}^d \rightarrow \mathbb{R}^d$  with  $\Phi_{true}$  being the *interaction potential*. Here  $*$  denotes the convolution

$$(K_{true} * u)(x, t) = \int_{\mathbb{R}^d} K_{true}(y) u(x - y, t) dy.$$

In particular, if  $\Phi_{true}$  is radial, that is  $\Phi_{true}(x) = \Phi_{true}(|x|)$ , we have

$$K_{true}(x) = \nabla(\Phi_{true}(|x|)) = \phi_{true}(|x|) \frac{x}{|x|}, \quad \text{with } \phi_{true}(r) = \Phi'_{true}(r). \quad (1.2)$$

where  $\phi_{true}$  is also called *interaction kernel* for simplicity. In either case, estimating the interaction function  $K_{true} = \nabla \Phi_{true}$  is equivalent to estimating the potential  $\Phi_{true}$  up to a constant.

The mean-field equation (also called aggregation-diffusion equation [6]) describes the population density for large systems of interacting particles or agents:

$$\frac{d}{dt} \mathbf{X}_t^i = \frac{1}{N} \sum_{j=1}^N \phi_{true}(|\mathbf{X}_t^j - \mathbf{X}_t^i|) \frac{\mathbf{X}_t^j - \mathbf{X}_t^i}{|\mathbf{X}_t^j - \mathbf{X}_t^i|} + \sqrt{2\nu} d\mathbf{B}_t^i, \quad \text{for } i = 1, \dots, N, \quad (1.3)$$

where  $\mathbf{X}_t^i \in \mathbb{R}^d$  represents the position of agent  $i$  at time  $t$ . Denote by  $u^{(N)}(x, t)$  the probability density of particle  $\mathbf{X}^1$  at time  $t$ . Under suitable conditions on the potential  $\Phi_{true}$ , we have  $u^{(N)} \rightarrow u$  in relative entropy of the steady state as the number of particles  $N \rightarrow \infty$  (see e.g., [23, 22, 7, 12]).

Systems of interacting particles or agents are widely used in many areas of science and engineering (see [2, 31, 25, 1] and the references therein). Motivated by the applications, there has been increasing interests in inferring the interaction kernel (or the interaction potential), either parametrically [27] or in a nonparametric fashion for generality [5, 21, 19, 20, 16]. The pioneering work [5] minimizes a loss functional based on matching the true and predicted trajectories of particles (see also the recent development for mean-field games [4]). When the system has finitely many particles and when the data consist of multiple trajectories of all particles, the efforts [21, 19, 20, 18, 17] minimizes loss functionals based on likelihood, establishing computationally efficient algorithms that yield estimators achieving the minimax rate of convergence. When the number of particles is large (e.g.,  $10^6$  or  $10^{23}$ , the Avogadro number), it becomes impractical to collect trajectories of all particles, but one can often observe the population density, i.e., the solution of the mean-field equations. This leads to the inverse problem of inferring the interaction kernel of the mean-field equation from solutions of the PDE. In [16], based on a probabilistic loss functional, the authors introduced an efficient nonparametric regression algorithm: it totally avoids the costly numerical simulations of the mean-field equation or the particle system (which are necessary in a trajectory matching method), instead, it estimates the kernel simply by least squares regression. However, the fundamental issue of identifiability is left open.

In this study, we provide a complete characterization of the identifiability and discuss its implications to computational practice. Based on a probabilistic loss functional, we seek the function spaces on which it has a unique minimizer. We show that identifiability holds on any subspace of two reproducing kernel Hilbert spaces (RKHS), whose reproducing kernels are intrinsic to the system and

adaptive to data. Furthermore, identifiability holds on two ambient  $L^2$  spaces if and only if the integral operators associated with the reproducing kernels are strictly positive. Thus, the inverse problem is ill-posed in general because the compact integral operators have unbounded inverse.

In computational practice, this implies that the regression matrix becomes ill-conditioned as the dimension of the hypothesis space increases. Thus, regularization is necessary. The two ambient  $L^2$  spaces provides natural norms for the Tikhonov regularization. In the context of singular value decomposition (SVD) analysis, we demonstrate numerically that the weighted  $L^2$  norm is preferable over the the unweighted  $L^2$  space because it leads to larger eigenvalues.

Our results are applicable to both radial and general vector valued interaction kernels.

This study show that the identifiability is universal for mean-field equations and for systems with  $N$  interacting particles (which are SDEs) [18]. The key is the probabilistic loss functional from the likelihood. There are two technical differences: multiple-trajectory data are necessary for the SDEs (without using ergodicity) and a slightly more flexible lower bound with a factor  $\frac{-1}{N-2}$  (see [18, Proposition 2.1]). However, as both  $N$  and the number of trajectories go to infinity, the identifiability in [18] agrees with the identifiability for mean-field equations in this study.

A key feature in our identifiability theory is that the RKHSs and their reproducing kernels are intrinsic to the system and are data-adaptive. This is different from the widely-used kernel regression techniques [9, 26] where pre-selected parametric kernels are fitted to data. These intrinsic RKHSs invite further study to reveal the structure of the interaction kernels and applications for system classifications.

The exposition in our manuscript proceeds as follows. In Section 2, we provide a mathematical definition of identifiability and describe of the main results. Section 3 proves identifiability for radial interaction kernels and Section 4 extends the results to general non-radial cases. We discuss in Section 5 the implications of identifiability to computational practice. Appendix A provides a brief review of positive definite functions and reproducing kernel Hilbert spaces.

## 2 Main results

Throughout the paper, we assume that the data  $u$  is a regular solution:

**Assumption 2.1.** *Assume that  $u$  is  $C^2$  in space and  $C^1$  in time and bounded above by  $C_u$ .*

Such a solution exists when the interaction function is local Lipschitz with polynomial growth:

$$|K_{true}(x) - K_{true}(y)| \leq C(|x - y| \wedge 1)(1 + |x|^m + |y|^m), \forall x, y \in \mathbb{R}^d$$

for some constant  $C > 0$ . We refer to [30] for Lipschitz kernels, and [23, 22] for uniform convex kernels that lead to convergence to equilibrium, [7, 12, 13] and the references therein for general (including singular) kernels.

### 2.1 Definition of identifiability

Our goal is to study the identifiability of the interaction kernel  $K_{true}$  (or  $\phi_{true}$  in the radial case) from data consisting of a solution  $u$  on  $[0, T]$ . Since we know little about the interaction kernel in priori, it is natural to consider nonparametric approaches, in which one finds an estimator by minimizing a loss functional over a hypothesis space. Thus, we define the identifiability as the uniqueness of the minimizer of a loss functional over a proper function space.

**Definition 2.2** (Identifiability). *Given data consisting a solution  $(u(x, t), x \in \mathbb{R}^d, t \in [0, T])$  to the mean-field equation (1.1), we say that the interaction kernel  $K_{true}$  (or  $\phi_{true}$  when the potential is radial) is identifiable in a hypothesis space  $\mathcal{H}$  if there exists a data-based loss functional such that  $K_{true}$  is the unique minimizer of the loss functional on  $\mathcal{H}$ .*

The identifiability consists of three elements: a loss functional  $\mathcal{E}$ , a function space of learning (hypothesis space)  $\mathcal{H}$ , and uniqueness of the minimizer of the loss functional over the hypothesis space. In [16], the authors introduce a probabilistic loss functional based on the likelihood of the diffusion process of the mean-field equation. An oracle version using the true interaction kernel is as follows (see Theorem 3.2 and Theorem 4.1 for the practical version based solely on data).

**Definition 2.3** (Oracle loss function). *Let  $u$  be a solution of the mean-field equation (1.1) on  $[0, T]$ . For a candidate interaction potential  $\Psi : \mathbb{R}^d \rightarrow \mathbb{R}$ , we denote  $K_\psi(x) = \nabla \Psi(x)$ . If furthermore  $\Psi$  is radial, we denote  $\psi(r) = \Psi'(r)$  with an abuse of notation. In this case,  $K_\psi(x) = \psi(x) \frac{x}{|x|}$ . We consider the loss functional*

$$\begin{aligned} \text{Radial: } \quad \mathcal{E}(\psi) &= \langle\langle \psi, \psi \rangle\rangle - 2\langle\langle \psi, \phi_{true} \rangle\rangle, \\ \text{General: } \quad \mathcal{E}(K_\psi) &= \langle\langle K_\psi, K_\psi \rangle\rangle - 2\langle\langle K_\psi, K_{true} \rangle\rangle. \end{aligned} \tag{2.1}$$

where the bilinear form  $\langle\langle \cdot, \cdot \rangle\rangle$  is defined by

$$\begin{aligned} \text{Radial: } \quad \langle\langle \varphi, \psi \rangle\rangle &:= \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} u(K_\varphi * u) \cdot (K_\psi * u) \, dx \, dt \\ \text{General: } \quad \langle\langle K_\varphi, K_\psi \rangle\rangle &:= \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} u(K_\varphi * u) \cdot (K_\psi * u) \, dx \, dt \end{aligned} \tag{2.2}$$

assuming that the integrals are well-defined.

Note that in either case, the loss functional is quadratic. Hence its minimizer on a finite dimensional hypothesis space can be estimated by least squares.

**Remark 2.4** (Least squares estimator). *For any space  $\mathcal{H} = \text{span} \{\phi_1, \dots, \phi_n\}$  such that*

$$A_{ij} = \langle\langle \phi_i, \phi_j \rangle\rangle, \quad b_i = \langle\langle \phi_i, \phi_{true} \rangle\rangle \tag{2.3}$$

are well-defined, a minimizer of the loss functional  $\mathcal{E}$  on  $\mathcal{H}$  is given by least squares

$$\hat{\phi}_{\mathcal{H}} = \sum_{i=1}^n \hat{c}_i \phi_i, \quad \text{where } \hat{c} = \arg \min_{c \in \mathbb{R}^n} \mathcal{E}(c), \quad \mathcal{E}(c) = c^\top A c - 2c^\top b.$$

In particular, when  $A$  is invertible, we have  $\hat{c} = A^{-1}b$  and  $\hat{\phi}_{\mathcal{H}}$  is the unique minimizer of  $\mathcal{E}$  on  $\mathcal{H}$ .

By Remark 2.4, identifiability on a finite dimensional function space is equivalent to the non-degeneracy of the normal matrix  $A$ . In general, as the next lemma shows, identifiability is equivalent to the non-degeneracy of the bilinear form.

**Lemma 2.5.** *Identifiability holds on a linear space  $\mathcal{H}$  if the bilinear form  $\langle\langle \cdot, \cdot \rangle\rangle$  is well-defined and non-degenerate on  $\mathcal{H}$ . That is, the true kernel  $\phi_{true} \in \mathcal{H}$  (or  $K_{true} \in \mathcal{H}$ ) is the unique minimizer of the loss functional  $\mathcal{E}$  in (2.1) in  $\mathcal{H}$  iff  $\langle\langle h, h \rangle\rangle > 0$  for all  $h \in \mathcal{H}$  unless  $h = 0$ .*

*Proof.* Since the radial case and general case have the same bilinear form, we consider only the radial case. Note that

$$\mathcal{E}(\psi) = \langle\langle \psi - \phi_{true}, \psi - \phi_{true} \rangle\rangle - \langle\langle \phi_{true}, \phi_{true} \rangle\rangle. \tag{2.4}$$

Thus,  $\phi \in \mathcal{H}$  is the unique minimizer of  $\mathcal{E}$  iff  $\langle\langle h, h \rangle\rangle > 0$  for all  $h \in \mathcal{H}$  unless  $h = 0$ .  $\square$

The bilinear form is non-degenerate iff its Fréchet derivative, which is a linear operator on a suitable space, is invertible. Thus, our goal of studying identifiability is to find function spaces on which the bilinear form's derivative is invertible.

To find the function spaces, we first write the bilinear form  $\langle\langle \cdot, \cdot \rangle\rangle$  explicitly. In case of radial interaction, we have (see (3.7) for its derivation)

$$\langle\langle \varphi, \psi \rangle\rangle = \int_{\mathbb{R}^+} \int_{\mathbb{R}^+} \varphi(r) \psi(s) \overline{G}_T(r, s) dr ds, \quad (2.5)$$

with the integral kernel  $\overline{G}_T$  given by

$$\begin{aligned} \overline{G}_T(r, s) &= \frac{1}{T} \int_0^T G_t(r, s) dt, \quad \text{with} \\ G_t(r, s) &:= \int_{\mathbb{S}^d} \int_{\mathbb{S}^d} \int_{\mathbb{R}^d} \xi \cdot \eta (rs)^{d-1} u(x - r\xi, t) u(x - s\eta, t) u(x, t) dx d\xi d\eta, \end{aligned} \quad (2.6)$$

where  $\mathbb{S}^d$  denotes the unit sphere in  $\mathbb{R}^d$ . In the general case, we have

$$\langle\langle K_\varphi, K_\psi \rangle\rangle = \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} K_\varphi(y) \cdot K_\psi(z) \overline{F}_T(x, y) dy dz$$

with an integral kernel (see (4.2) for its derivation)

$$\overline{F}_T(y, z) = \frac{1}{T} \int_0^T F_t(y, z) dt, \quad \text{with } F_t(y, z) := \int_{\mathbb{R}^d} u(x - y, t) u(x - z, t) u(x, t) dx. \quad (2.7)$$

Hence the bilinear form's Fréchet derivative can be an integral operator either on  $L^2(\mathbb{R}^+)$  with integral kernel  $\overline{G}_T$  in the radial case or on  $L^2(\mathbb{R}^d)$  with integral kernel  $\overline{F}_T$  in the general case.

However, those spaces do not reflect information from data, not even the support of these kernels (note that either  $\overline{G}_T$  or  $\overline{F}_T$  is determined by the data  $(u(x, t), x \in \mathbb{R}^d, t \in [0, T])$ ). For example, when the integral kernel has a bounded support due to data, little can be said outside of the support in a nonparametric regression method using local basis. This promotes us to seek data-based function spaces of learning in the next section.

## 2.2 The ambient function spaces of regression

We introduce two ambient function spaces of learning (via a nonparametric regression with local basis): a weight  $L^2$  space with a data-based measure and an unweighted  $L^2$  space on the support of the measure.

The function space of learning should be data-based because we can only learn the function in the region where the data explore. Intuitively, the convolution  $K_\psi * u$  suggests that the independent variable of the interaction kernel  $\psi$  is  $|x - y|$  (or  $x - y$  for general kernel  $K$ ), with  $x$  and  $y$  from the data  $u(x, t)$  and  $u(y, t)$ . We show next that this intuition agrees with the probabilistic representation of the mean-field equation:  $\psi$  has independent variables  $|\overline{X}_t - \overline{Y}_t|$  (or  $\overline{X}_t - \overline{Y}_t$  for the general case), where  $\overline{X}_t$  and  $\overline{Y}_t$  are independent variables with probability density  $u(\cdot, t)$ .

Recall that Equation (1.1) is the Kolmogorov forward equation (also called the Fokker-Planck equation) of the following nonlinear stochastic differential equation

$$\begin{cases} d\overline{X}_t = - [K_{true} * u](\overline{X}_t, t) dt + \sqrt{2} dB_t, \\ \mathcal{L}(\overline{X}_t) = u(x, t) dx, \end{cases} \quad (2.8)$$

for all  $t \geq 0$ . Here  $\mathcal{L}(\overline{X}_t)$  denotes the law of  $\overline{X}_t$ , with probability density  $u(\cdot, t)$ .

Let  $(\overline{X}'_t, t \geq 0)$  be an independent copy of  $(\overline{X}_t, t \geq 0)$  and denote  $r_t = |\overline{X}_t - \overline{X}'_t|$ . We can write the convolution  $K_\psi * u$  as

$$[K_\psi * u](\overline{X}_t, t) = \mathbb{E}[K_\psi(\overline{X}_t - \overline{X}'_t) | \overline{X}_t] = \mathbb{E}[\psi(r_t) \frac{\overline{X}_t - \overline{X}'_t}{r_t} | \overline{X}_t]. \quad (2.9)$$

The above probabilistic representation of  $K_\psi * u$  indicates that the independent variable of  $\psi$  has the same distribution of  $\{|\bar{X}_t - \bar{X}'_t|, t \in [0, T]\}$  (or  $\{\bar{X}_t - \bar{X}'_t, t \in [0, T]\}$  for general kernel  $K$ ). Let  $\bar{\rho}_T$  denote the average of probability densities of  $\{|\bar{X}'_t - \bar{X}_t|, t \in [0, T]\}$  in the case of radial kernel, or  $\{\bar{X}'_t - \bar{X}_t, t \in [0, T]\}$  in the case of general kernel:

$$\begin{aligned} \text{Radial case: } \bar{\rho}_T(r) &= \frac{1}{T} \int_0^T \rho_t(r) dt = \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} \int_{\mathbb{S}^d} r^{d-1} u(y - r\xi, t) u(y, t) dy dt, \\ \text{General case: } \bar{\rho}_T(x) &= \frac{1}{T} \int_0^T \rho_t(x) dt = \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} u(y - x, t) u(y, t) dy dt, \end{aligned} \quad (2.10)$$

where  $\rho_t$  denotes the density of  $|\bar{X}'_t - \bar{X}_t|$  (or  $\bar{X}'_t - \bar{X}_t$  for the general case) for each  $t$ . Under Assumption 2.1, the probability density function  $\bar{\rho}_T$  is  $C^2$ . We denote the support of  $\bar{\rho}_T$  by  $\mathcal{X}$ :

$$\mathcal{X} := \text{supp}(\bar{\rho}_T).$$

Note that  $\mathcal{X}$  is bounded if  $u(x, t)$  has a bounded support uniformly for all  $t \in [0, T]$ .

Thus, the ambient function space of nonparametric regression is  $L^2(\mathcal{X}, \bar{\rho}_T(r) dr)$  for radial kernels and  $L^2(\mathcal{X}, \bar{\rho}_T(x) dx)$  for general kernels. For simplicity of notation, hereafter we denote both  $L^2(\mathcal{X}, \bar{\rho}_T(r) dr)$  and  $L^2(\mathcal{X}, \bar{\rho}_T(x) dx)$  by  $L^2(\bar{\rho}_T)$  as long as the context is clear.

Another natural function space of learning is  $L^2(\mathcal{X})$ , because the integral kernel  $\bar{G}_T$  in (2.6) (or  $\bar{F}_T$  in (2.7)) leads to an integral operator on  $L^2(\mathcal{X})$  which is the Fréchet derivative of the loss functional.

Note that  $L^2(\mathcal{X}) \subset L^2(\bar{\rho}_T)$ . We will show that both function spaces are reasonable for learning and they lead to different RKHS's on which the identifiability holds.

## 2.3 Main results

With the loss functional in Definition 2.3 and the ambient function spaces of learning in Section 2.2, we can now define the Fréchet derivative of the loss functional. The function spaces of identifiability are those on which the Fréchet derivative is an invertible operator.

We provide a complete characterization for the function spaces of identifiability in both  $L^2(\mathcal{X})$  and  $L^2(\bar{\rho}_T)$ . In the radial case, we have

- In  $L^2(\mathcal{X})$ , identifiability holds on the RKHS  $\mathcal{H}_G$  with reproducing kernel  $\bar{G}_T$  in (2.6). Furthermore, identifiability holds on  $L^2(\mathcal{X})$  iff  $\mathcal{H}_G$  is dense in it, equivalently, the integral operator  $\mathcal{L}_{\bar{G}_T}$  on  $L^2(\mathcal{X})$  with integral kernel  $\bar{G}_T$  is strictly positive. See Theorem 3.6– 3.7 in Section 3.2.
- In  $L^2(\bar{\rho}_T)$ , identifiability holds on the RKHS  $\mathcal{H}_R$  with reproducing kernel  $\bar{R}_T(r, s) = \frac{\bar{G}_T(r, s)}{\bar{\rho}_T(r)\bar{\rho}_T(s)}$ . Furthermore, identifiability holds on  $L^2(\bar{\rho}_T)$  iff  $\mathcal{H}_R$  is dense in it, equivalently, the integral operator  $\mathcal{L}_{\bar{R}_T}$  on  $L^2(\bar{\rho}_T)$  with integral kernel  $\bar{R}_T$  is strictly positive. See Theorem 3.9– 3.10 in Section 3.3.

Similar results holds in the general case (see Section 4).

We point out that identifiability is weaker than well-posedness. Identifiability only ensures that the loss functional has a unique minimizer on a hypothesis function space  $\mathcal{H}$ . It does not ensure well-posedness of the inverse problem unless the bilinear form satisfies a coercivity condition on  $\mathcal{H}$  (see Remark 3.11). When  $\mathcal{H}$  is finite dimensional, identifiability is equivalent to the invertibility of the normal matrix (see Remark 2.4), and identifiability implies well-posedness. However, when  $\mathcal{H}$  is infinite dimensional, the inverse problem is ill-posed because the inverse of the compact operator  $\mathcal{L}_{\bar{G}_T}$  or  $\mathcal{L}_{\bar{R}_T}$  is unbounded. Thus, identifiability is weaker than well-posedness.

We also discuss the implications of identifiability for the computational practice of estimating the interaction kernel from data. Identifiability theory implies that the regression matrix will become

Table 1: Notations.

	Radial case	General (non-radial) case
Interaction potential	$\Psi( x )$ and $\psi = \Psi'$ ;	$\Psi(x)$
Interaction kernel	$K_\psi(x) = \psi( x )\frac{x}{ x }$	$K_\psi(x) = \nabla\Psi(x)$
Loss functional	$\mathcal{E}(\psi)$	$\mathcal{E}(K)$
Density of regression measure	$\bar{\rho}_T, \mathcal{X} = \text{support}(\bar{\rho}_T)$ , in (2.10)	
Function space of learning	$L^2(\mathcal{X})$ and $L^2(\bar{\rho}_T)$	
Mercer kernel & RKHS in $L^2(\mathcal{X})$	$\bar{G}_T$ in (2.6) and $\mathcal{H}_G$	$\bar{F}_T$ in (2.7) and $\mathcal{H}_F$
Mercer kernel & RKHS in $L^2(\bar{\rho}_T)$	$\bar{R}_T$ in (3.11) and $\mathcal{H}_R$	$\bar{Q}_T$ in (4.6) and $\mathcal{H}_Q$

ill-conditioned as the dimension of the hypothesis space increases (see Theorem 5.1). Thus, regularization becomes necessary. We compare two regularization norms, the norms of  $L^2(\bar{\rho}_T)$  and  $L^2(\mathcal{X})$ , in the context of singular value decomposition (SVD) analysis. Numerical tests suggest that regularization based on  $L^2(\bar{\rho}_T)$  norm leads to slightly better conditioned inversion (see Figure 1). We also show that the regularization by truncated SVD is equivalent to learning on the low-frequency eigenspace of the RKHSs.

We shall use the notations in Table 1.

### 3 Radial interaction kernel

We start from the identifiability of radial interaction kernels. We will first derive the probabilistic loss functional in (3.4) (see Section 3.1). Then, in Section 3.2-3.3, we discuss the identifiability in  $L^2(\mathcal{X})$  and in  $L^2(\bar{\rho}_T)$ , in the ambient function spaces of learning defined in Section 2.2.

Through this section, we let  $u$  be a solution of the mean-field equation (1.1) on  $[0, T]$ . Let  $\psi \in L^2(\bar{\rho}_T)$  with  $\bar{\rho}_T$  in (2.10) and let  $\Psi : \mathbb{R}^+ \rightarrow \mathbb{R}$  be  $\Psi(r) = \int_0^r \psi(s) ds$ . Denote  $K_\psi(x) = \psi(|x|)\frac{x}{|x|}$ .

#### 3.1 The loss function

We show first that the radial loss functional in (2.1) is well-defined.

**Lemma 3.1.** *For any  $\varphi, \psi \in L^2(\bar{\rho}_T)$ , the bilinear form  $\langle\langle \varphi, \psi \rangle\rangle$  in (2.2) satisfies*

$$\langle\langle \varphi, \psi \rangle\rangle \leq \|\psi\|_{L^2(\rho_T)} \|\varphi\|_{L^2(\rho_T)}. \quad (3.1)$$

Also, for any  $\psi \in L^2(\bar{\rho}_T)$ , the radial loss functional in (2.1) is bounded above by

$$\mathcal{E}(\psi) \leq \|\psi - \phi_{true}\|_{L^2(\rho_T)}^2 - \langle\langle \phi_{true}, \phi_{true} \rangle\rangle. \quad (3.2)$$

*Proof.* Recall that  $u(\cdot, t)$  is the law of  $\bar{X}_t$  defined in (2.8). By definition in (2.2), we have

$$\begin{aligned} \langle\langle \varphi, \psi \rangle\rangle &= \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} (K_\varphi * u) \cdot (K_\psi * u) u(x, t) dx dt \\ &= \frac{1}{T} \int_0^T \mathbb{E}[K_\varphi * u(\bar{X}_t, t) \cdot K_\psi * u(\bar{X}_t, t)] dt, \end{aligned} \quad (3.3)$$

The integrand in time is controlled by

$$\begin{aligned}
& \mathbb{E}[K_\varphi * u(\bar{X}_t, t) \cdot K_\psi * u(\bar{X}_t, t)] \\
& \leq \mathbb{E}[|K_\varphi * u(\bar{X}_t, t)|^2]^{1/2} \mathbb{E}[|K_\psi * u(\bar{X}_t, t)|^2]^{1/2} \quad (\text{Cauchy-Schwartz}) \\
& \leq \mathbb{E}[\mathbb{E}[|K_\varphi(\bar{X}_t - \bar{X}'_t)|^2 | \bar{X}_t]]^{1/2} \mathbb{E}[\mathbb{E}[|K_\psi(\bar{X}_t - \bar{X}'_t)|^2 | \bar{X}_t]]^{1/2} \quad (\text{by (2.9)}) \\
& \leq \mathbb{E}[\mathbb{E}[|K_\varphi(\bar{X}_t - \bar{X}'_t)|^2 | \bar{X}_t]]^{1/2} \mathbb{E}[\mathbb{E}[|K_\psi(\bar{X}_t - \bar{X}'_t)|^2 | \bar{X}_t]]^{1/2} \quad (\text{by Jensen's inequality}) \\
& = \mathbb{E}[|K_\varphi(\bar{X}_t - \bar{X}'_t)|^2]^{1/2} \mathbb{E}[|K_\psi(\bar{X}_t - \bar{X}'_t)|^2]^{1/2} = \|\varphi\|_{L^2(\rho_t)} \|\psi\|_{L^2(\rho_t)}.
\end{aligned}$$

Then, we obtain (3.1).

The upper bound in (3.2) follows from (3.1) and (2.4).  $\square$

The next theorem shows that the radial loss functional is based on the maximal likelihood ratio the SDE (2.8) representation of the mean-field equation, and it provides a derivative-free loss functional in practice, which leads to an efficient and robust least square estimator (see [16]).

**Theorem 3.2** (Lost function). *The radial loss functional in (2.1) is the expectation of the average (in time) negative log-likelihood of the process  $(\bar{X}_t, t \in [0, T])$  in (2.8). If  $\Psi \in L^1(\bar{\rho}_T)$ , it equals to*

$$\mathcal{E}(\psi) = \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} [|K_\psi * u|^2 u + 2\partial_t u(\Psi * u) + 2\nu \nabla u \cdot (K_\psi * u)] dx dt. \quad (3.4)$$

Furthermore, if  $\nabla \cdot K_\psi \in L^1(\bar{\rho}_T)$ , we can replace the integrand  $\nabla u \cdot (K_\psi * u)$  by  $-u(\nabla \cdot K_\psi * u)$ .

*Proof.* We denote by  $\mathcal{P}_\phi$  the law of the process  $(\bar{X}_t)$  on the path space with initial condition  $\bar{X}_0 \sim u(\cdot, 0)$ , with the convention that  $\mathcal{P}_0$  denoting the Winner measure. Then, the average negative log-likelihood ratio of a trajectory  $\bar{X}_{[0, T]}$  from a law  $\mathcal{P}_\psi$  with respect to the Winner measure is (see e.g. [15, Section 1.1.4] and [14, Section 3.5])

$$\mathcal{E}_{\bar{X}_{[0, T]}}(\psi) = -\log \frac{d\mathcal{P}_\psi}{d\mathcal{P}_0}(\bar{X}_{[0, T]}) = \frac{1}{T} \int_0^T (|K_\psi * u(\bar{X}_t, t)|^2 dt - 2\langle K_\psi * u(\bar{X}_t, t), d\bar{X}_t \rangle), \quad (3.5)$$

where  $\frac{d\mathcal{P}_\psi}{d\mathcal{P}_0}$  is the Radon-Nikodym derivative.

Taking expectation and noting that  $d\bar{X}_t = K_{true} * u(\bar{X}_t, t) dt + \sqrt{2} dB_t$ , we obtain

$$\begin{aligned}
\mathbb{E} \mathcal{E}_{\bar{X}_{[0, T]}}(\psi) &= \frac{1}{T} \int_0^T \mathbb{E} [|K_\psi * u(\bar{X}_t, t)|^2 - 2K_\psi * u(\bar{X}_t, t) \cdot K_{true} * u(\bar{X}_t, t)] dt. \\
&= \langle \psi, \psi \rangle - 2\langle \phi_{true}, \psi \rangle = \mathcal{E}(\psi),
\end{aligned} \quad (3.6)$$

where the second equality follows from (3.3).

To replace the true interaction kernel  $K_{true}$  by data, we resort to the mean-field equation (1.1). Noticing that  $K_\psi * u = \nabla \Psi * u = \nabla(\Psi * u)$  for any  $\psi = \Psi'$ , we have

$$\begin{aligned}
& \int_0^T \int_{\mathbb{R}^d} u(K_{true} * u) \cdot (K_\psi * u) dx dt = \int_0^T \int_{\mathbb{R}^d} u(K_{true} * u) \cdot (\nabla \Psi * u) dx dt \\
&= - \int_0^T \int_{\mathbb{R}^d} (\nabla \cdot [u(K_{true} * u)]) \Psi * u dx dt \quad (\text{by integration by parts}) \\
&= - \int_0^T \int_{\mathbb{R}^d} (\partial_t u - \nu \Delta u) (\Psi * u) dx dt \quad (\text{by Equation (1.1)}) \\
&= - \int_0^T \int_{\mathbb{R}^d} \partial_t u (\Psi * u) dx dt - \nu \int_0^T \int_{\mathbb{R}^d} \nabla u \cdot (K_\psi * u) dx dt, \quad (\text{integration by parts}).
\end{aligned}$$

Combining it with (3.6) and (3.3), we obtain the radial loss functional in (3.4). At last, applying integration by parts again, we can replace the integrand  $\nabla u \cdot (K_\psi * u)$  by  $-u(\nabla \cdot K_\psi * u)$ .  $\square$



**Remark 3.3** (loss functional in terms of expectations). *Along with (2.9), we can write the loss functional in (3.4) in terms of expectations*

$$\mathcal{E}(\psi) = \frac{1}{T} \int_0^T (\mathbb{E}[|\mathbb{E}[K_\psi(\bar{X}_t - \bar{X}'_t)|\bar{X}_t]|^2] + \partial_t \mathbb{E}[\Psi(\bar{X}_t - \bar{X}'_t)] + 2\nu \mathbb{E}[\Delta \Psi(\bar{X}_t - \bar{X}'_t)]) dt,$$

where we used the fact that  $\mathbb{E}[\Psi(\bar{X}_t - \bar{X}'_t)] = \mathbb{E}[\mathbb{E}[\Psi(\bar{X}_t - \bar{X}'_t)|\bar{X}_t]]$  and

$$\int_{\mathbb{R}^d} \partial_t u(\Psi * u) dx = \frac{1}{2} \partial_t \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} u(x, t) \Psi(x - y) u(y, t) dy dx = \frac{1}{2} \partial_t \mathbb{E}[\mathbb{E}[\Psi(\bar{X}_t - \bar{X}'_t)|\bar{X}_t]].$$

Such a probabilistic loss functional can be approximated by Monte Carlo. This is particularly important in high-dimensional problems, when multidimensional integrals faces the well-known curse-of-dimensionality, and the mean-field solution (population density) can be efficiently approximated by the empirical distribution of random samples.

## 3.2 The RKHS space and $L^2$ space

Now we show that the interaction kernel is identifiable on the RKHS  $\mathcal{H}_G$  with reproducing kernel  $\bar{G}_T$  defined in (2.6), which is a subspace of  $L^2(\mathcal{X})$ . We show first that  $\bar{G}_T$  is a Mercer kernel. Then we connect its RKHS with  $L^2(\mathcal{X})$  through the integral operator of  $\bar{G}_T$ , which is the bilinear form's Fréchet derivative on  $L^2(\mathcal{X})$ .

Recall that the bilinear form leads to the function  $\bar{G}_T$  defined in (2.6):

$$\begin{aligned} \langle\langle \varphi, \psi \rangle\rangle &:= \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} (K_\varphi * u) \cdot (K_\psi * u) u(x, t) dx dt \\ &= \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} K_\varphi(y) \cdot K_\psi(z) \int_{\mathbb{R}^d} u(x - y, t) u(x - z, t) u(x, t) dx dy dz dt \\ &= \int_0^\infty \int_0^\infty \varphi(r) \psi(s) \underbrace{\frac{1}{T} \int_0^T \int_{\mathbb{S}^d} \int_{\mathbb{S}^d} \int_{\mathbb{R}^d} \xi \cdot \eta (rs)^{d-1} u(x - r\xi, t) u(x - s\eta, t) u(x, t) dx d\xi d\eta dr ds}_{\bar{G}_T(r, s)} \end{aligned} \quad (3.7)$$

where the third equality follows from a change of variable to polar coordinates, along with the notion that  $K_\varphi(y) = \varphi(|y|) \frac{y}{|y|} = \varphi(r) \xi$  with  $r = |y|$  and  $\xi = \frac{y}{|y|} \in \mathbb{S}^d$ .

**Lemma 3.4.** *The kernel  $\bar{G}_T$  is a Mercer kernel, i.e., it is symmetric, continuous and positive definite.*

*Proof.* The symmetry is clear from its definition and the continuity follows from the continuity of  $u$ . Following Definition A.1, for any  $\{r_1, \dots, r_k\} \subset \mathbb{R}^+$  and  $(c_1, \dots, c_k) \in \mathbb{R}^k$ , we have

$$\begin{aligned} \sum_{i, j} c_i c_j \bar{G}_T(r_i, r_j) &= \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} \sum_{i, j} c_i c_j (r_i r_j)^{d-1} \int_{\mathbb{S}^d} \int_{\mathbb{S}^d} \xi \cdot \eta u(x - r_i \xi, t) u(x - r_j \eta, t) d\xi d\eta u(x, t) dx dt \\ &= \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} \left| \sum_i c_i r_i^{d-1} \int_{\mathbb{S}^d} \xi u(x - r_i \xi, t) d\xi \right|^2 u(x, t) dx dt \geq 0. \end{aligned}$$

Thus, it is positive definite.  $\square$

Since  $\bar{G}_T$  is a Mercer kernel, it determines an RKHS  $\mathcal{H}_G$  with  $\bar{G}_T$  as reproducing kernel (see Appendix A). We show next that the identifiability holds on the RKHS  $\mathcal{H}_G$ , i.e.  $\langle\langle \cdot, \cdot \rangle\rangle$  is non-degenerate on  $\mathcal{H}_G$ , by studying the integral operator with kernel  $\bar{G}_T$ .

$$\mathcal{L}_{\bar{G}_T} f(r) = \int_{\mathbb{R}^+} \bar{G}_T(r, s) f(s) ds. \quad (3.8)$$

Note that by definition,

$$\langle\langle \varphi, \psi \rangle\rangle = \langle \varphi, \mathcal{L}_{\overline{G}_T} \psi \rangle_{L^2(\mathcal{X})}. \quad (3.9)$$

We start with a Lemma on the boundedness and integrability of  $\overline{G}_T$ .

**Lemma 3.5.** *The kernel function  $\overline{G}_T$  in (2.6) satisfies the following properties:*

- (a) *For all  $r, s \in \mathbb{R}^+$ ,  $\overline{G}_T(r, s) \lesssim \overline{\rho}_T(r)$ . Hence  $\overline{G}_T$  is supported in  $\mathcal{X} \times \mathcal{X}$ .*
- (b)  *$\overline{G}_T$  is in  $L^2(\mathcal{X} \times \mathcal{X})$ , and  $\overline{G}_T(r, \cdot) \in L^2(\mathcal{X})$  for every  $r \in \mathcal{X}$ .*

*Proof.* First we can write  $\overline{G}_T$  as

$$\overline{G}_T(r, s) = \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} \left( \int_{|y|=r} \frac{y}{|y|} u(x-y, t) dy \right) \cdot \left( \int_{|z|=s} \frac{z}{|z|} u(x-z, t) dz \right) u(x, t) dx dt.$$

Notice that  $\left( \int_{|z|=s} \frac{z}{|z|} u(x-z, t) dz \right)$  is uniformly bounded for all  $s \in \mathbb{R}^+$  because of the density property and the continuity of  $u$ . Also recall that  $\mathcal{X}$  is the support of  $\overline{\rho}_T$ , hence (a) is proved.

For (b), by Jensen's inequality and boundedness of  $u(\cdot, t)$  as a probability density,

$$\begin{aligned} & \int_{\mathbb{R}^+} \int_{\mathbb{R}^+} |\overline{G}_T(r, s)|^2 dr ds \\ & \leq \frac{1}{T^2} \int_{\mathbb{R}^+} \int_{\mathbb{R}^+} \int_0^T \int_{\mathbb{R}^d} \left( \int_{|y|=r} u(x-y, t)^2 dy \right) \cdot \left( \int_{|z|=s} u(x-z, t)^2 dz \right) u(x, t)^2 dx dt. \\ & \leq \frac{1}{T^2} \int_0^T \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |u(x-y, t) u(x-z, t) u(x, t)|^2 dx dy dz dt < \infty. \end{aligned}$$

The second property follows similarly.  $\square$

By Lemma 3.5 and Theorem A.3,  $\mathcal{L}_{\overline{G}_T}$  is a positive compact self-adjoint operator on  $L^2(\mathcal{X})$ , and it has countably many positive eigenvalues  $\{\lambda_i\}_{i=1}^\infty$  with orthonormal eigenfunctions  $\{\varphi_i\}_{i=1}^\infty$ . In particular,  $\{\sqrt{\lambda_i} \varphi_i\}_{i=1}^\infty$  is an orthonormal basis of  $\mathcal{H}_G$ . The following theorem follows directly.

**Theorem 3.6** (Identifiability on RKHS). *Identifiability holds on  $\mathcal{H}_G$ , the RKHS with reproducing kernel  $\overline{G}_T$  in (2.6).*

*Proof.* Since  $\{\sqrt{\lambda_i} \varphi_i\}$  is an orthonormal basis of  $\mathcal{H}_G$ , we can write each  $f \in \mathcal{H}_G$  as  $f = \sum_{i=1}^\infty c_i \sqrt{\lambda_i} \varphi_i$  and we have  $\langle f, f \rangle_{\mathcal{H}_G} = \sum_{i=1}^\infty c_i^2$ . Thus,  $\langle\langle f \rangle\rangle^2 = \sum_{i=1}^\infty c_i^2 \lambda_i^2 > 0$  for any  $f \neq 0$  in  $\mathcal{H}_G$ .  $\square$

The RKHS  $\mathcal{H}_G$  has the nice feature of being data-informed: its reproducing kernel  $\overline{G}_T$  depends solely on the data  $(u(x, t), x \in \mathbb{R}^d, t \in [0, T])$ . However, it is an implicit function space, so it is helpful to consider the explicit function space  $L^2(\mathcal{X})$ .

**Theorem 3.7** (Identifiability on  $L^2(\mathcal{X})$ ). *The following are equivalent.*

- (a) *Identifiability holds on  $L^2(\mathcal{X})$ .*
- (b)  *$\mathcal{L}_{\overline{G}_T}$  is strictly positive.*
- (c)  *$\mathcal{H}_G$  is dense in  $L^2(\mathcal{X})$ .*

Moreover, for any  $h = \sum_{i=1}^n c_i \varphi_i \in \mathcal{H} = \text{span}\{\varphi_1, \dots, \varphi_n\}$  with  $\{\varphi_i\}$  being orthonormal eigenfunctions of  $\mathcal{L}_{\overline{G}_T}$  corresponding to positive eigenvalues  $\{\lambda_i\}$ , we have

$$\langle\langle h \rangle\rangle^2 = \sum_{i=1}^n c_i^2 \lambda_i, \quad \|h\|_{\mathcal{H}_G}^2 = \sum_{i=1}^n c_i^2 \lambda_i^{-1}, \quad \|h\|_{L^2(\mathcal{X})}^2 = \sum_{i=1}^n c_i^2. \quad (3.10)$$

In particular, the bilinear form  $\langle\langle \cdot, \cdot \rangle\rangle$  satisfies the coercivity condition on  $\mathcal{H}$ :

$$\langle\langle h \rangle\rangle \geq \min\{\sqrt{\lambda_i}\}_{i=1}^n \|h\|_{L^2(\mathcal{X})}, \quad \forall h \in \mathcal{H}.$$

*Proof.* (a)  $\Rightarrow$  (b). Suppose  $\mathcal{L}_{\overline{G}_T}$  is not strictly positive, then there exists eigenfunction  $h \in L^2(\mathcal{X})$  corresponding to eigenvalue 0. But we would also have  $\langle\langle h \rangle\rangle = 0$ .

(b)  $\Rightarrow$  (a). If  $\overline{G}_T$  is strictly positive, then  $\{\varphi_i\}_{i=1}^\infty$  would be an orthonormal basis for  $L^2(\mathcal{X})$  and all eigenvalues of  $\mathcal{L}_{\overline{G}_T}$  are positive. Take  $h = \sum_{j=1}^\infty c_j \varphi_j$ . Then,  $\langle\langle h \rangle\rangle = 0$  implies

$$\left\langle \mathcal{L}_{\overline{G}_T} h, h \right\rangle_{L^2(\mathcal{X})} = \left\langle \sum_{j=1}^\infty c_j \lambda_j \varphi_j, \sum_{j=1}^\infty c_j \varphi_j \right\rangle_{L^2(\mathcal{X})} = \sum_{j=1}^\infty c_j^2 \lambda_j = 0.$$

Hence  $h = 0$  in  $L^2(\mathcal{X})$ .

(b)  $\Leftrightarrow$  (c). Note that  $\{\sqrt{\lambda_i} \varphi_i\}_{i=1}^\infty$  is a basis of  $\mathcal{H}_G$ . Thus,  $\mathcal{H}_G$  is dense in  $L^2(\mathcal{X})$  iff  $\{\varphi_i\}_{i=1}^\infty$  is a basis of  $L^2(\mathcal{X})$ , i.e.  $\mathcal{L}_{\overline{G}_T}$  is strictly positive.

At last, for any  $h = \sum_{j=1}^n c_j \varphi_j \in \mathcal{H} = \text{span} \{\varphi_1, \dots, \varphi_n\}$ , we have

$$\langle\langle h \rangle\rangle^2 = \left\langle \mathcal{L}_{\overline{G}_T} h, h \right\rangle_{L^2(\mathcal{X})} = \sum_{j=1}^n c_j^2 \lambda_j \geq \min\{\lambda_i\}_{i=1}^n \|h\|_{L^2(\mathcal{X})}^2.$$

Also, we have  $h = \sum_{j=1}^n c_j \lambda_j^{-1/2} \sqrt{\lambda_j} \varphi_j \in \mathcal{H}_G$  with  $\|h\|_{\mathcal{H}_G}^2 = \sum_{i=1}^n c_i^2 \lambda_i^{-1}$ ; □

### 3.3 The weighted $L^2$ space and RKHS

In this section, we study the identifiability on  $L^2(\overline{\rho}_T)$  through the RKHS whose reproducing kernel is a weighted integral kernel.

We define the following kernel  $\overline{R}_T$  on the set  $\mathcal{X} \times \mathcal{X}$ :

$$\overline{R}_T(x, y) = \frac{\overline{G}_T(r, s)}{\overline{\rho}_T(r) \overline{\rho}_T(s)}, \quad (r, s) \in \mathcal{X} \times \mathcal{X}. \quad (3.11)$$

Similar to Lemma 3.4, we can prove

**Lemma 3.8.** *The kernel  $\overline{R}_T$  defined in (3.11) is a Mercer kernel.*

With  $\overline{R}_T$  being a Mercer kernel, it leads to another RKHS  $\mathcal{H}_R$  (see Appendix A). To characterize the RKHS, we define the integral operator  $\mathcal{L}_{\overline{R}_T} : L^2(\overline{\rho}_T) \rightarrow L^2(\overline{\rho}_T)$ :

$$\mathcal{L}_{\overline{R}_T} \varphi(s) = \int_{\mathbb{R}^+} \overline{R}_T(r, s) \varphi(r) \overline{\rho}_T(r) dr. \quad (3.12)$$

Here we assume that  $\overline{R}_T \in L^2(\overline{\rho}_T \otimes \overline{\rho}_T)$ , so that  $\mathcal{L}_{\overline{R}_T}$  is a compact operator. This assumption holds true when  $\mathcal{X}$  is compact, which is typical in practice. Note that by definition,

$$\langle\langle \varphi, \psi \rangle\rangle = \langle \varphi, \mathcal{L}_{\overline{R}_T} \psi \rangle_{L^2(\overline{\rho}_T)}. \quad (3.13)$$

Under the above assumption,  $\mathcal{L}_{\overline{R}_T}$  is a positive and compact self-adjoint operator on  $L^2(\overline{\rho}_T)$ , and it has countably many positive eigenvalues  $\{\gamma_i\}_{i=1}^\infty$  with orthonormal eigenfunctions  $\{\psi_i\}_{i=1}^\infty$ . In particular,  $\{\sqrt{\gamma_i} \psi_i\}_{i=1}^\infty$  is an orthonormal basis of  $\mathcal{H}_R$ . Based on the eigenfunctions of  $\mathcal{L}_{\overline{R}_T}$  and by the same argument as in Theorem 3.6, we have

**Theorem 3.9** (Identifiability on RKHS). *Identifiability holds on  $\mathcal{H}_R$ , the RKHS with reproducing kernel  $\overline{R}_T$  in (3.11).*

Similar to Theorem 3.7, the identifiability holds on  $L^2(\overline{\rho}_T)$  if the RKHS  $\mathcal{H}_R$  is dense in it.

**Theorem 3.10** (Identifiability on  $L^2(\overline{\rho}_T)$ ). *The following are equivalent.*

- (a) Identifiability holds on  $L^2(\bar{\rho}_T)$ .
- (b)  $\mathcal{L}_{\bar{R}_T}$  in (3.12) is strictly positive.
- (c)  $\mathcal{H}_R$  is dense in  $L^2(\bar{\rho}_T)$ .

Moreover, for any  $h = \sum_{i=1}^n c_i \psi_i \in \mathcal{H} = \text{span}\{\psi_1, \dots, \psi_n\}$  with  $\{\psi_i\}$  being orthonormal eigenfunctions of  $\mathcal{L}_{\bar{R}_T}$  corresponding to eigenvalues  $\{\gamma_i > 0\}$ , we have

$$\langle\langle h \rangle\rangle^2 = \sum_{i=1}^n c_i^2 \gamma_i, \quad \|h\|_{\mathcal{H}_R}^2 = \sum_{i=1}^n c_i^2 \gamma_i^{-1}, \quad \|h\|_{L^2(\bar{\rho}_T)}^2 = \sum_{i=1}^n c_i^2. \quad (3.14)$$

In particular, the bilinear form  $\langle\langle \cdot, \cdot \rangle\rangle$  satisfies the coercivity condition on  $\mathcal{H}$ :

$$\langle\langle h \rangle\rangle \geq \min\{\sqrt{\gamma_i}\}_{i=1}^n \|h\|_{L^2(\bar{\rho}_T)}, \quad \forall h \in \mathcal{H}.$$

**Remark 3.11** (Relation to the coercivity condition of the bilinear form). *Recall that a bilinear form  $\langle\langle \cdot, \cdot \rangle\rangle$  is said to be coercive on a subspace  $\mathcal{H} \subset L^2(\bar{\rho}_T)$  if there exists a constant  $c_{\mathcal{H}} > 0$  such that  $\langle\langle \psi, \psi \rangle\rangle \geq c_{\mathcal{H}} \|\psi\|_{L^2(\bar{\rho}_T)}^2$  for all  $\psi \in \mathcal{H}$ . Such a coercivity condition has been introduced on subspaces of  $L^2(\bar{\rho}_T)$  in [5, 21, 19, 20, 18] for systems of finitely many particles or agents. Theorem 3.10 shows that, for any finite-dimensional hypothesis space  $\mathcal{H} = \text{span}\{\psi_1, \dots, \psi_n\}$ , the coercivity condition holds with  $c_{\mathcal{H}} = \min\{\gamma_i\}_{i=1}^n$ , but the coercivity constant vanishes as the dimension of  $\mathcal{H}$  increases to infinity.*

Now we have two RHKSs:  $\mathcal{H}_G$  and  $\mathcal{H}_R$ , which are function spaces of idnenifiably under different topologies. They are the images  $\mathcal{L}_{\bar{G}_T}^{1/2}(L^2(\mathcal{X}))$  and  $\mathcal{L}_{\bar{R}_T}^{1/2}(L^2(\bar{\rho}_T))$  (see Appendix A). The following remarks discuss the relation between them.

**Remark 3.12** (The two integral operators). *The integral operators  $\mathcal{L}_{\bar{G}_T}$  and  $\mathcal{L}_{\bar{R}_T}$  are derived from the same bilinear form: for any  $\varphi, \psi \in L^2(\mathcal{X}) \subset L^2(\bar{\rho}_T)$ , by (3.9) and (3.13), we have*

$$\langle\langle \varphi, \psi \rangle\rangle = \langle \mathcal{L}_{\bar{G}_T} \varphi, \psi \rangle_{L^2(\mathcal{X})} = \langle \mathcal{L}_{\bar{R}_T} \varphi, \psi \rangle_{L^2(\bar{\rho}_T)}.$$

Since  $L^2(\mathcal{X})$  is dense in  $L^2(\bar{\rho}_T)$ , the second equality implies that the null-space of and  $\mathcal{L}_{\bar{G}_T}$  is a subset of  $\mathcal{L}_{\bar{R}_T}$ . However, there is no correspondence between their eigenfunctions of nonzero eigenvalues. To see this, let  $\varphi_i \in L^2(\mathcal{X})$  be an eigenfunction of  $\mathcal{L}_{\bar{G}_T}$  with eigenvalue  $\lambda_i > 0$ . Then, it follows from the second equality that

$$\langle \lambda_i \varphi_i, \psi \rangle_{L^2(\mathcal{X})} = \langle \mathcal{L}_{\bar{G}_T} \varphi_i, \psi \rangle_{L^2(\mathcal{X})} = \langle \mathcal{L}_{\bar{R}_T} \varphi_i, \psi \rangle_{L^2(\bar{\rho}_T)} = \langle \bar{\rho}_T \mathcal{L}_{\bar{R}_T} \varphi_i, \psi \rangle_{L^2(\mathcal{X})}$$

for any  $\psi \in L^2(\mathcal{X})$ . Thus,  $\lambda_i \varphi_i = \bar{\rho}_T \mathcal{L}_{\bar{R}_T} \varphi_i$  in  $L^2(\mathcal{X})$ . Thus, neither  $\rho \varphi_i$  nor  $\frac{\varphi_i}{\rho}$  is an eigenfunction of  $\mathcal{L}_{\bar{R}_T}$ . In finite-dimensional approximation in Section 5, we will show that  $\mathcal{L}_{\bar{R}_T}$  leads to a generalized eigenvalue problem, while  $\mathcal{L}_{\bar{G}_T}$  leads to the normal matrix in regression.

**Remark 3.13** (Metrics on  $\mathcal{H}_G$  and  $\mathcal{H}_R$ ). *We have three metrics on  $\mathcal{H}_G \subset L^2(\mathcal{X})$ : the RKHS norm, the  $L^2(\mathcal{X})$  norm and the norm induced by the bilinear form. From (3.10), we have, for any  $h = \sum_{i=1}^{\infty} c_i \varphi_i \in \mathcal{H}_G$  with  $\{\varphi_i\}$  being the eigenfunctions  $\mathcal{L}_{\bar{G}_T}$ ,*

$$\langle\langle h \rangle\rangle^2 = \sum_{i=1}^{\infty} c_i^2 \lambda_i, \quad \|h\|_{\mathcal{H}_G}^2 = \sum_{i=1}^{\infty} c_i^2 \lambda_i^{-1}, \quad \|h\|_{L^2(\mathcal{X})}^2 = \sum_{i=1}^{\infty} c_i^2.$$

Thus, the three metrics satisfy  $\langle\langle h \rangle\rangle \leq \max_i \{\sqrt{\lambda_i}\} \|h\|_{L^2(\mathcal{X})} \leq \max_i \{\lambda_i\} \|h\|_{\mathcal{H}_G}$ . Similarly, there are three metrics on  $\mathcal{H}_R \subset L^2(\bar{\rho}_T)$  satisfying  $\langle\langle h \rangle\rangle \leq \max_i \{\sqrt{\gamma_i}\} \|h\|_{L^2(\bar{\rho}_T)} \leq \max_i \{\gamma_i\} \|h\|_{\mathcal{H}_R}$ .

## 4 General interaction kernels

We consider the case of general interaction kernel  $K_{true} : \mathbb{R}^d \rightarrow \mathbb{R}^d$ , which can be non-radial. In comparison to the radial case, the major difference is that  $K_{true} = \nabla\Phi$  is a vector-valued function on  $\mathbb{R}^d$ . The extension is straightforward and the elements are similar: the quadratic loss functional and the RKHS spaces of identifiability.

### 4.1 The loss function

Let  $K : \mathbb{R}^d \rightarrow \mathbb{R}^d$  be a general interaction kernel with potential  $\Psi$ , i.e.,  $K = \nabla\Psi$ . We will denote it by  $K_\psi$  to emphasize its relation with  $\Psi$ ; otherwise, we will use  $K$  for simplicity.

**Proposition 4.1** (Loss function). *Let  $u$  be a solution of the mean-field equation (1.1) on  $[0, T]$ . The general loss functional  $\mathcal{E}$  in (2.1) is the expectation of the average (in time) negative log-likelihood of the process  $(\bar{X}_t, t \in [0, T])$  in (2.8). If  $K_\psi = \nabla\Psi \in L^2(\bar{\rho}_T)$  with  $\bar{\rho}_T$  being the general density in (2.10), the loss functional can be written as*

$$\mathcal{E}(K_\psi) := \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} \left[ |K_\psi * u|^2 u + 2\partial_t u(\Psi * u) + 2\nu \nabla u \cdot (K_\psi * u) \right] dx dt. \quad (4.1)$$

Furthermore, if  $K_\psi \in L^1(\bar{\rho}_T)$ , we can replace the integrand  $\nabla u \cdot (K_\psi * u)$  by  $-u(\nabla \cdot K_\psi * u)$ .

*Proof.* The proof is the same as Theorem 3.2.  $\square$

The loss functional  $\mathcal{E}$  is quadratic. Thus, its Fréchet derivative (with respect to a suitable function space) is a linear operator and the uniqueness of its minimizer is equivalent to the non-degeneracy of its Fréchet derivative. Recall that for vector-valued functions  $K_1, K_2 : \mathbb{R}^d \rightarrow \mathbb{R}^d$ , we define the bilinear form in (2.2):

$$\begin{aligned} \langle\langle K_1, K_2 \rangle\rangle &= \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} (K_1 * u) \cdot (K_2 * u) u(x, t) dx dt \\ &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} K_1(y) \cdot K_2(z) \underbrace{\frac{1}{T} \int_0^T \int_{\mathbb{R}^d} u(x-y, t) u(x-z, t) u(x, t) dx dt}_{\bar{F}_T} dy dz \\ &= \sum_{i=1}^d \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} K_1^i(y) K_2^i(z) \bar{F}_T(y, z) dy dz =: \sum_{i=1}^d \langle\langle K_1^i, K_2^i \rangle\rangle \end{aligned} \quad (4.2)$$

where the scalar-valued kernel  $\bar{F}_T(y, z)$  is defined in (2.7), and in the last equality we abuse the notation of the bilinear form: we also denote it for the components of the vector-valued functions. We also denote  $\langle\langle K \rangle\rangle^2 := \langle\langle K, K \rangle\rangle$ .

The next lemma extends Lemma 2.5, highlighting the crucial role of the bilinear form.

**Lemma 4.2.** *Suppose  $\mathcal{H} = \prod_{i=1}^d \mathcal{H}_0$ . Then, identifiability holds in  $\mathcal{H}$  (or in  $\mathcal{H}_0$ ) if and only if  $\langle\langle \cdot \rangle\rangle^2$  is nondegenerate on  $\mathcal{H}_0$ , i.e.  $\langle\langle h \rangle\rangle^2 > 0$  for any  $h \neq 0$  in  $\mathcal{H}_0$ .*

*Proof.* By Definition 2.2, it suffices to show that the loss functional  $\mathcal{E}$  in (4.1) has a uniqueness minimizer on  $\mathcal{H}$ . With the bilinear functional (4.2), we can write the loss functional as  $\mathcal{E}(K) = \langle\langle K, K \rangle\rangle - 2\langle\langle K, K_{true} \rangle\rangle = \langle\langle K - K_{true} \rangle\rangle^2 - \langle\langle K_{true} \rangle\rangle^2$ . Let  $K - K_{true} = (k_1(x), \dots, k_d(x))$  with  $k_i \in \mathcal{H}_0$  for  $i = 1, \dots, d$ , then  $\langle\langle K - K_{true} \rangle\rangle^2 = \sum_{i=1}^d \langle\langle k_i \rangle\rangle^2$ . Thus,  $K_{true}$  is identifiable by  $\mathcal{E}$  in  $\mathcal{H}$  iff  $\langle\langle k \rangle\rangle^2 > 0$  for any  $k \neq 0$  in  $\mathcal{H}_0$ .  $\square$

**Remark 4.3** (Minimizer of the loss functional). Consider the hypothesis space  $\mathcal{H} = \text{span}\{K_1, \dots, K_n\}$  with  $K_i = \nabla \Phi_i \in L^2(\mathbb{R}^d, \mathbb{R}^d)$  such that  $A$  in (4.3) is invertible. Then, for any  $K = \sum_{i=1}^n c_i K_i \in \mathcal{H}$ , the loss functional  $\mathcal{E}$  becomes  $\mathcal{E}(K) = \mathcal{E}(c) = c^\top A c + b^\top c$  and it has a unique minimizer on  $\mathcal{H}$ :

$$\hat{K} = \sum_{i=1}^n \hat{c}_i K_i, \text{ with } \hat{c} = A^{-1}b, \quad (4.3)$$

where the normal matrix  $A$  and vector  $b$  are given by

$$A_{ij} = \langle\langle K_i, K_j \rangle\rangle = \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} (K_i * u) \cdot (K_j * u) u(x, t) dx dt,$$

$$b_i = \langle\langle K_i, K_{true} \rangle\rangle = -\frac{1}{T} \int_0^T \int_{\mathbb{R}^d} [\partial_t u (\Phi_i * u) + \nu \nabla u \cdot (K_i * u)] dx dt.$$

## 4.2 The RKHS space and $L^2$ space

We show next that the identifiability holds on the RKHS with reproducing kernel  $\bar{F}_T$  in (2.7), which is a subspace of  $L^2(\mathcal{X})$ . We show first that the  $\bar{F}_T$  is a Mercer kernel. Then, we connect its RKHS with  $L^2(\mathcal{X})$  through its integral operator on  $L^2(\mathcal{X})$ .

**Lemma 4.4.** *The function  $\bar{F}_T$  in (2.7) is a Mercer kernel.*

*Proof.* The symmetry of  $\bar{F}_T$  follows from its definition, and its continuity follows from the continuity of  $u$ . It is positive definite by its definition and Theorem A.2.  $\square$

Denote by  $\mathcal{H}_F$  the RKHS with reproducing kernel  $\bar{F}_T$  and denote its inner product by  $\langle \cdot, \cdot \rangle_{\mathcal{H}_F}$ . We show next that identifiability holds on the RKHS  $\mathcal{H}_F$ , i.e.,  $\langle\langle \cdot, \cdot \rangle\rangle$  is non-degenerate on  $\mathcal{H}_F$ , by studying the integral operator with kernel  $\bar{F}_T$ . We start with the next lemma on the boundedness and integrability of  $\bar{F}_T$ .

**Lemma 4.5.** *The symmetric function  $\bar{F}_T$  in (2.7) satisfies the following properties:*

- (a) For all  $x, y \in \mathbb{R}^d$ ,  $\bar{F}_T(x, y) \leq C_u \bar{\rho}_T(x)$ , where  $C_u = \sup_{x \in \mathbb{R}^d, t \in [0, T]} u(x, t)$ .
- (b)  $\bar{F}_T$  is in  $L^2(\mathbb{R}^d \times \mathbb{R}^d)$ , and  $\bar{F}_T(x, \cdot) \in L^2(\mathbb{R}^d)$  for every  $x \in \mathbb{R}^d$ .

*Proof.* The boundedness in (a) follows directly:

$$\bar{F}_T(x, y) \leq C_u \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} u(z - y, t) u(z, t) dz dt = C_u \bar{\rho}_T(y).$$

For (b), note that by symmetry, we have  $\bar{F}_T(x, y) \leq C_u \bar{\rho}_T(y)$  for any  $x, y \in \mathbb{R}^d$ . Then,

$$\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} (\bar{F}_T(x, y))^2 dx dy < C_u^2 \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \bar{\rho}_T(x) \bar{\rho}_T(y) dx dy = C_u^2,$$

and similarly,  $\int_{\mathbb{R}^d} (\bar{F}_T(x, y))^2 dy \leq C_u^2 \bar{\rho}_T(x)$  for every  $x$ .  $\square$

By Lemma 4.5, the integral operator  $\mathcal{L}_{\bar{F}_T} : L^2(\mathcal{X}) \rightarrow L^2(\mathcal{X})$ ,

$$\mathcal{L}_{\bar{F}_T} f(x) = \int_{\mathcal{X}} \bar{F}_T(y, x) f(y) dy. \quad (4.4)$$

is a bounded, positive and compact operator on  $L^2(\mathcal{X})$  (see e.g., [29, Proposition 1]). Hence  $\mathcal{L}_{\bar{F}_T}$  has at most countably many positive eigenvalues. We use the same notation  $\{\lambda_i\}_{i=1}^\infty$  and  $\{\varphi_i\}_{i=1}^\infty$  as  $\mathcal{L}_{\bar{G}_T}$

to denote the positive eigenvalues and the corresponding eigenfunctions of  $\mathcal{L}_{\bar{F}_T}$ , as long as there is no confusion in the context. Similarly,  $\{\varphi_i\}_{i=1}^\infty$  are orthonormal in  $L^2(\mathcal{X})$ . Moreover,  $\{\sqrt{\lambda_i}\varphi_i\}_{i=1}^\infty$  form an orthonormal basis of  $\mathcal{H}_F$  and  $\mathcal{H}_F = \mathcal{L}_{\bar{F}_T}^{1/2}(L^2(\mathcal{X}))$  (see e.g., [8, Section 4.4]). Note that

$$\langle\langle K_1, K_2 \rangle\rangle = \langle K_1, \mathcal{L}_{\bar{F}_T} K_2 \rangle_{L^2(\mathcal{X})}. \quad (4.5)$$

When  $\bar{\rho}_T$ 's support is bounded, then by Mercer's theorem,  $\bar{F}_T(x, y) = \sum_{i=1}^\infty \lambda_i \varphi_i(x) \varphi_i(y)$ , which converges uniformly on  $\mathcal{X} \times \mathcal{X}$ .

**Theorem 4.6** (Identifiability on RKHS). *Identifiability holds on  $\mathcal{H}_F$ .*

*Proof.* Since  $\{\sqrt{\lambda_i}\varphi_i\}$  is an orthonormal basis of  $\mathcal{H}_F$ , we can write each  $f \in \mathcal{H}_F$  as  $f = \sum_{i=1}^\infty c_i \sqrt{\lambda_i} \varphi_i$  and we have  $\langle f, f \rangle_{\mathcal{H}_F} = \sum_{i=1}^\infty c_i^2$ . Thus,  $\langle\langle f \rangle\rangle^2 = \sum_{i=1}^\infty c_i^2 \lambda_i^2 > 0$  for any  $f \neq 0$  in  $\mathcal{H}_F$ .  $\square$

The RKHS  $\mathcal{H}_F$  has the nice feature of being data-informed: its reproducing kernel  $\bar{F}_T$  depends on the data  $(u(x, t), x \in \mathbb{R}^d, t \in [0, T])$ . However, it is an implicit function space. Thus, it is helpful to consider the explicit function space  $L^2(\mathcal{X})$ .

**Theorem 4.7** (Identifiability on  $L^2(\mathcal{X})$ ). *The following are equivalent.*

- (a) *Identifiability holds on  $L^2(\mathcal{X})$ .*
- (b)  *$\mathcal{L}_{\bar{F}_T}$  in (4.4) is strictly positive.*
- (c)  *$\mathcal{H}_F$  is dense in  $L^2(\mathcal{X})$ .*

Moreover, for any finite dimensional space  $\mathcal{H} = \text{span}\{\varphi_1, \dots, \varphi_n\}$  with  $\{\varphi_i\}$  being eigenfunctions of  $\mathcal{L}_{\bar{F}_T}$  corresponding to eigenvalues  $\{\lambda_i > 0\}$ , we have  $\langle\langle k \rangle\rangle \geq \min\{\sqrt{\lambda_i}\}_{i=1}^n \|k\|_{L^2(\mathcal{X})}$ ,  $\forall k \in \mathcal{H}$ .

*Proof.* The proof is the same as Theorem 3.7.  $\square$

### 4.3 The weighted $L^2$ space and RKHS

In practice, the data  $(u(x, t), x \in \mathbb{R}^d, t \in [0, T])$  only explore a small region  $\mathcal{X}$  in  $\mathbb{R}^d$  with a weight  $\bar{\rho}_T$  defined in (2.10). Thus, it is important to consider the weighted space  $L^2(\bar{\rho}_T)$  (which denotes the space  $L^2(\bar{\rho}_T(x)dx)$  by abusing the notation) instead of  $L^2(\mathcal{X})$ . In this section, we study the identifiability on  $L^2(\bar{\rho}_T)$  through the RKHS whose reproducing kernel is a weighted kernel.

Consider the weighted integral kernel  $\bar{Q}_T$ :

$$\bar{Q}_T(x, y) = \frac{\bar{F}_T(x, y)}{\bar{\rho}_T(x)\bar{\rho}_T(y)}, \quad (x, y) \in \mathcal{X} \times \mathcal{X}. \quad (4.6)$$

With a proof similar to those for Lemma 4.4, we have

**Proposition 4.8.** *The kernel  $\bar{Q}_T$  defined in (4.6) is a Mercer kernel over  $(\mathcal{X}, \bar{\rho}_T)$ , which is symmetric, continuous and positive definite.*

With  $\bar{Q}_T$  being a Mercer kernel, we denote its RKHS space by  $\mathcal{H}_Q$  with inner product  $\langle \cdot, \cdot \rangle_{\mathcal{H}_Q}$ . Similar to Section 4.2, we connect the RKHS  $\mathcal{H}_Q$  with  $L^2(\bar{\rho}_T)$  through  $\bar{Q}_T$ 's integral operator. Here we assume that  $\bar{Q}_T \in L^2(\bar{\rho}_T \otimes \bar{\rho}_T)$ , so that its integral operator  $\mathcal{L}_{\bar{Q}_T} : L^2(\bar{\rho}_T) \rightarrow L^2(\bar{\rho}_T)$

$$\mathcal{L}_{\bar{Q}_T} f(x) = \int_{\mathbb{R}^d} \bar{Q}_T(y, x) f(y) \bar{\rho}_T(y) dy, \quad (4.7)$$

is a compact operator (actually, a Hilbert-Schmidt operator). This assumption holds true when  $\mathcal{X}$  is compact, which is the most typical case in practice. But it may not hold when  $\mathcal{X}$  is non-compact. The following two examples show that the assumption holds when the distribution of  $(\bar{X}_t, t \geq 0)$  is a Gaussian process, but it does not hold when the process is stationary with a Cauchy distribution.

**Example 4.9.** Suppose  $K(x) = x$  and  $d = 1$ . Then,  $U(x) = \frac{1}{\sqrt{2\pi}}e^{-\frac{x^2}{2\nu^2}}$  is a stationary solution to the mean-field equation (1.1) (equivalently,  $\mathcal{N}(0, \nu^2)$  is an invariant measure of the SDE (2.8)). Suppose that  $u(x, 0) = U(x)$ . Then, we have

$$\bar{F}_T(x, y) = \frac{1}{2\nu\sqrt{3\pi}}e^{-\frac{1}{3\nu^2}(x^2+y^2-xy)} \quad \text{and} \quad \bar{\rho}_T(x) = \frac{1}{2\nu\sqrt{\pi}}e^{-\frac{x^2}{4\nu^2}}.$$

The weighted integral kernel  $\bar{Q}_T$  is still square integrable due to the fast decay of  $\bar{F}_T$ :

$$t\|\bar{Q}_T\|_{L^2(\bar{\rho}_T \otimes \bar{\rho}_T)}^2 = \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{\bar{F}_T(x, y)^2}{\bar{\rho}_T(x)\bar{\rho}_T(y)} dx dy = \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{1}{3\pi} e^{-\frac{1}{12\nu^2}[4(x-y)^2+x^2+y^2]} dx dy < \infty.$$

**Example 4.10.** Suppose that the density of Cauchy distribution,  $U(x) = \frac{1}{\pi} \frac{1}{1+x^2}$ , is a steady solution to (1.1), and suppose that  $u(x, 0) = U(x)$ . Then we have

$$\bar{F}_T(x, y) = \frac{2}{\pi^2} \frac{(x^2 - xy + y^2 + 12)}{(x^2 + 4)(y^2 + 4)(x^2 - 2xy + y^2 + 4)} \quad \text{and} \quad \bar{\rho}_T(x) = \frac{2}{\pi} \frac{1}{(x^2 + 4)}$$

However,  $\bar{Q}_T$  is not square integrable:

$$\|\bar{Q}_T\|_{L^2(\bar{\rho}_T \otimes \bar{\rho}_T)}^2 = \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{\bar{F}_T(x, y)^2}{\bar{\rho}_T(x)\bar{\rho}_T(y)} dx dy = \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{(x^2 - xy + y^2 + 12)}{2(x^2 - 2xy + y^2 + 4)} dx dy = \infty.$$

Denote  $\{\gamma_i\}_{i=1}^{\infty}$  and  $\{\psi_i\}_{i=1}^{\infty}$  the positive eigenvalues and corresponding eigenfunctions of  $\mathcal{L}_{\bar{Q}_T}$ . Similarly,  $\{\psi_i\}_{i=1}^{\infty}$  are orthonormal in  $L^2(\bar{\rho}_T)$  and  $\{\sqrt{\gamma_i}\psi_i\}_{i=1}^{\infty}$  forms an orthonormal basis of  $\mathcal{H}_Q$ . Furthermore,  $\mathcal{H}_Q = \mathcal{L}_{\bar{Q}_T}^{1/2}(L^2(\bar{\rho}_T))$  and we can write the bilinear form (4.2) as

$$\langle\langle K_1, K_2 \rangle\rangle = \langle K_1, \mathcal{L}_{\bar{Q}_T} K_2 \rangle_{L^2(\bar{\rho}_T)}. \quad (4.8)$$

The following theorem on identifiability over the space  $\mathcal{H}_Q$  is a counterpart of Theorem 4.6.

**Theorem 4.11.** Let  $\mathcal{H} = \bigoplus_{i=1}^d \mathcal{H}_Q$ . Then identifiability holds on  $\mathcal{H}$ .

*Proof.* Let  $k \in \mathcal{H}_Q$  and write  $k = \sum_{i=1}^{\infty} c_i \sqrt{\gamma_i} \psi_i$ . Then  $\langle\langle k \rangle\rangle = 0$  implies

$$\begin{aligned} \langle\langle k \rangle\rangle &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} k(x)k(y)\bar{F}_T(x, y) dx dy = \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} k(x)k(y)\bar{Q}_T(x, y)\bar{\rho}_T(x)\bar{\rho}_T(y) dx dy \\ &= \left\langle \mathcal{L}_{\bar{Q}_T} k, k \right\rangle_{L^2(\bar{\rho}_T)} = \sum_{j=1}^{\infty} c_j^2 \gamma_j^2 = 0. \end{aligned} \quad (4.9)$$

Thus  $c_j = 0$  for all  $j$  and  $k(x) = 0$  a.e. in  $(\mathcal{X}, \bar{\rho}_T)$ . Suppose we have  $K \in \mathcal{H}$ , then we can apply the above discussion for the components of  $K$  and achieve the identifiability on  $\mathcal{H}$ .  $\square$

The following theorem is an  $L^2(\bar{\rho}_T)$  version of Theorem 4.7.

**Theorem 4.12** (Identifiability on  $L^2(\bar{\rho}_T)$ ). *The following are equivalent.*

- (a) Identifiability holds on  $\mathcal{H} = \bigoplus_{i=1}^d L^2(\bar{\rho}_T)$ .
- (b)  $\mathcal{L}_{\bar{Q}_T}$  in (4.4) is strictly positive.
- (c)  $\mathcal{H}_Q$  is dense in  $L^2(\bar{\rho}_T)$ .

Moreover, for any finite dimensional space  $\mathcal{H} = \text{span}\{\psi_1, \dots, \psi_n\}$  with  $\{\psi_i\}$  being eigenfunctions of  $\mathcal{L}_{\bar{Q}_T}$  corresponding to eigenvalues  $\{\gamma_i > 0\}$ , we have:  $\langle\langle K \rangle\rangle \geq \min\{\sqrt{\gamma_i}\}_{i=1}^n \|K\|_{L^2(\bar{\rho}_T)}$ ,  $\forall K \in \mathcal{H}$ .

*Proof.* The proof is identical to the proof in Theorem 4.7 and we omit it.  $\square$



## 4.4 Examples and general function spaces

We show a few examples that identifiability holds on  $L^2(\mathcal{X})$  or  $L^2(\bar{\rho}_T)$  when  $\mathcal{X} = \mathbb{R}^d$ , under mild conditions on the initial condition  $u_0$ . These examples are limited to the case when  $\mathcal{X} = \mathbb{R}^d$  because we prove their integral kernel to be strictly positive definite based on Fourier transformation.

Recall that  $F_t(x, y)$  in (2.7) is positive definite for all  $t \in [0, T]$ . The next lemma shows that we only need the integral kernel to be strictly positive definite at only a single time instant.

**Lemma 4.13.** *Assume that  $F_{t_0}(x, y)$  in (2.7) is strictly positive definite for some  $t_0 \in [0, T]$ , that is, for every function  $f \in L^2(\mathbb{R}^d)$ ,*

$$\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} f(x)f(y)F_{t_0}(x, y)dx dy = 0 \quad (4.10)$$

*implies  $f = 0$  a.e.. Then the identifiability holds on both  $L^2(\mathbb{R}^d)$  and  $L^2(\rho)$ .*

*Proof.* By the continuity of  $\bar{F}_T$ ,  $\langle\langle f \rangle\rangle = 0$  implies  $\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} f(x)f(y)F_{t_0}(x, y)dx dy = 0$ . Then  $f = 0$  a.e. by the assumption, also  $f = 0$  a.e. in  $\rho$  by the Radon-Nikodym theorem. Hence,  $L^2(\mathbb{R}^d)$  and  $L^2(\rho)$  have the identifiability.  $\square$

We show next that  $F_{t_0}(x, y)$  is strictly positive definite when  $u(x, t_0)$ 's Fourier transform satisfy suitable conditions. Without loss of generality, we assume  $t_0 = 0$  and consider the initial condition  $u_0$  in the following discussion.

**Theorem 4.14.** *Assume the initial value  $u_0$  is non-vanishing on  $\mathbb{R}^d$ .*

- (a) *Suppose there exists a Borel measure  $\mu$  on  $\mathbb{C}^d$  such that  $u_0(x) = \int_{\mathbb{R}^d} e^{ixy} d\mu(y)$ . If the carrier of  $\mu$  is the entire  $\mathbb{C}^d$ , then  $L^2(\mathbb{R}^d)$  and  $L^2(\bar{\rho}_T)$  have the identifiability.*
- (b) *Suppose there exists a Borel measure  $\mu$  on  $\mathbb{R}^d$  such that  $u_0(x) = \int_{\mathbb{R}^d} e^{ixy} d\mu(y)$ . If the carrier of  $\mu$  contains an open set, then the space of bounded and compactly supported functions  $M_{bc}(\mathbb{R}^d)$  has the identifiability.*
- (c) *Suppose  $u_0(x) = u_0(|x|)$ , i.e.  $u_0(x)$  is radial and there exists a Borel measure  $\nu$  on  $[0, \infty)$  such that  $u_0(x) = \int_0^\infty e^{-|x|^2 t} d\nu(t)$ . If  $\nu$  is not concentrated at zero, then the space of Borel probability measures  $\mathcal{P}(\mathbb{R}^d)$  has the identifiability.*

*Proof.* As for (a), by Lemma 4.2 and Lemma 4.13, we only need to show that for all  $f \in L^2(\mathbb{R}^d)$ , (4.10) implies  $f = 0$ . From the definition of  $F_t(x, y)$  in (2.7), we can write (4.10) as

$$\int_{\mathbb{R}^d} |u_0 * f(x)|^2 u_0(x) dx = 0,$$

which implies  $u_0 * f = 0$  in  $\mathbb{R}^d$  a.e. since  $u_0$  is supported on  $\mathbb{R}^d$ . Taking the Fourier transform on both sides, we have  $\mathfrak{F}\{f\}\mu = 0$ . Since  $\mu$  is supported on  $\mathbb{C}^d$ , we have  $\mathfrak{F}\{f\} = 0$  and then  $f = 0$  a.e..

The proof for (b) is due to [24]. By the previous steps, for  $f \in M_{bc}(\mathbb{R}^d)$  we have  $\mathfrak{F}\{f\}\mu = 0$ . Thus,  $\mathfrak{F}\{f\}$  vanishes on the open set supporting  $\mu$ . By Paley-Winner Theorem,  $\mathfrak{F}\{f\}$  is an entire function since  $f$  has compact support. Thus we have  $f = 0$  a.e..

The proof for (c) is due to [28] and [32]. In particular, Theorem 7.14 in [32] states that such  $u_0$  is strictly positive definite. Then, by Proposition 5 in [28],  $u_0$  must be characteristic in the sense that  $\mu \mapsto \int F_0(\cdot, y)\mu(dy)$  is injective on  $\mathcal{P}$ . Thus,  $\mathcal{P}$  has the identifiability. Note that in this case, we also cannot get the identifiability over  $L^2(\mathbb{R}^d)$  and  $L^2(\rho)$ .  $\square$

Note the condition for  $u_0$  is weakest in (b) and (a), (c) are stronger, hence the identifiable space in (a), (c) is larger than the one in (b). Also, in either case (b) or (c), we cannot get the identifiability over  $L^2(\mathbb{R}^d)$  and  $L^2(\rho)$ .

**Example 4.15.** Let  $u_0(x)$  be the density function of Gaussian distribution  $\mathcal{N}(\mu, \Sigma)$ . The Fourier transform, i.e., the characteristic function of  $u_0$  is  $e^{i\mu\xi - \frac{1}{2}\xi^\top \Sigma \xi}$ . Since it is nonvanishing on  $\mathbb{C}^d$ , it satisfies the condition in (a), and thus  $L^2(\mathbb{R}^d)$  and  $L^2(\rho)$  have the identifiability. By Theorem 4.7 and Theorem 4.12,  $\mathcal{H}_F$  is dense in  $L^2(\mathbb{R}^d)$  and  $\mathcal{H}_Q$  is dense in  $L^2(\rho)$ . Note that this density follows from the continuity of  $u$  and the initial distribution  $u_0$ .

**Example 4.16.** The condition in (c) for radial functions is equivalent to  $u_0$  being strictly positive [28]. Hence all the nonvanishing  $L^1(\mathbb{R}^d)$  integrable radial strictly positive functions will work, such as the radial Gaussian density of  $\mathcal{N}(0, I)$  and the Cauchy distribution  $u_0(x) \propto (c^2 + |x|^2)^{-\beta}$  with  $c > 0$ ,  $\beta > d/2$  (see [32, Theorem 6.13]). Actually, these density functions also satisfy the condition of (a). The Gaussian density is verified in the above example. Similarly, the characteristic function of the Cauchy distribution is  $e^{-|\beta|\xi}$ , which is nonzero for all  $\xi \in \mathbb{C}^d$ . Thus,  $L^2(\mathbb{R}^d)$  and  $L^2(\rho)$  have the identifiability,  $\mathcal{H}_F$  is dense in  $L^2(\mathbb{R}^d)$  and  $\mathcal{H}_Q$  is dense in  $L^2(\rho)$ .

We show next that identifiability hold on  $L^2(\mathbb{R}^d)$  and  $L^2(\rho)$  when  $u_0$  with support on  $\mathbb{R}^d$  is obtained from modifying a compactly supported probability density.

**Proposition 4.17.** Let  $v(x) \in C_c(\mathbb{R}^d)$  be a probability density, and  $\eta_\varepsilon$  be the density function of  $\mathcal{N}(0, \varepsilon I)$ . If the initial value  $u_0 = v * \eta_\varepsilon$ , then  $L^2(\mathbb{R}^d)$  and  $L^2(\rho)$  has the identifiability.

*Proof.* By Lemma 4.13, we show that  $f = 0$  a.e. if  $f \in L^2(\mathbb{R}^d)$  satisfy (4.10). Note that (4.10) implies that  $u_0 * f = 0$  a.e.. Taking Fourier transform, we have  $\mathfrak{F}\{v\}\mathfrak{F}\{\eta_\varepsilon\}\mathfrak{F}\{f\} = 0$ . Since  $v$  is a compactly supported probability density,  $\mathfrak{F}\{v\}$  is non-vanishing by Paley-Winner Theorem. Also,  $\mathfrak{F}\{\eta_\varepsilon\}$  is non-vanishing. Then, we must have  $\mathfrak{F}\{f\} = 0$ , and therefore  $f = 0$  a.e.  $\square$

## 5 Identifiability in computational practice

In this section, we discuss the implications of identifiability for the computational practice of estimating the interaction kernel from data. For simplicity, we consider only radial interaction kernels in the case that the spatial dimension is  $d = 1$ . We shall see that the regression matrix will become ill-conditioned as the dimension of the hypothesis space increases (see Theorem 5.1). Thus, regularization becomes necessary. We compare two regularization norms, the norms of  $L^2(\bar{\rho}_T)$  or  $L^2(\mathcal{X})$ , corresponding to the function spaces of learning, in the context of singular value decomposition (SVD) analysis. Numerical tests suggest that regularization based on  $L^2(\bar{\rho}_T)$  norm leads to slightly better conditioned inversion (see (5.8) and Figure 1). We also show that the regularization by truncated SVD is equivalent to learning on the low-frequency eigenspace of the RKHSs.

### 5.1 Nonparametric regression in practice

In computational practice, suppose we are given data  $u(x, t)$  on  $\mathbb{R} \times [0, T]$  on discrete space mesh grids, we find a minimizer of the loss functional by least squares as in Remark 2.4. We review only those fundamental elements and we refer to [16] for more details.

We first select a set of adaptive-to-data basis functions from the data-based measure  $\bar{\rho}_T$ , guided by the function space of learning (either  $L^2(\bar{\rho}_T)$  or  $L^2(\mathcal{X})$ ). The starting point is to find the measure  $\bar{\rho}_T$  using (2.10) on a fine mesh (the mesh that  $u$  is defined on) and obtain its support. Let  $\{r_i\}_{i=0}^n$  be a uniform partition of the support set  $\mathcal{X}$  and denote the width of each interval as  $\Delta r$ . They provide the knots for the B-spline basis functions of  $\mathcal{H}$ . Here we use piecewise constant basis functions to facilitate the rest discussions, that is,

$$\mathcal{H} = \text{span}\{\phi_1, \dots, \phi_n\}, \text{ with } \phi_i(r) = \mathbf{1}_{[r_{i-1}, r_i]}(r).$$

One may also use other partitions, for example, a partition with uniform probability for all intervals, as well as other basis functions, such as higher degree B-splines (often depending on the regularity of the true kernel) or weighted orthogonal polynomials (which are global basis functions).

With these basis functions, we let

$$A_{ij} = \langle\langle \phi_i, \phi_j \rangle\rangle = \frac{1}{T} \int_0^T \int_{\mathbb{R}^d} (K_{\phi_i} * u) \cdot (K_{\phi_j} * u) u(x, t) dx dt, \quad (5.1)$$

$$b_i = \langle\langle \phi_i, \phi_{true} \rangle\rangle = -\frac{1}{T} \int_0^T \int_{\mathbb{R}^d} [\partial_t u \Phi_i * u + \nu \nabla u \cdot (K_{\phi_i} * u)] dx dt. \quad (5.2)$$

where  $\Phi_i(r) = \int_0^r \phi_i(s) ds$  is an anti-derivative of  $\phi_i$ ,  $K_{\phi_i}(x) = \phi_i(|x|) \frac{x}{|x|}$  and the vector  $b \in \mathbb{R}^n$  comes from Theorem 3.2. The integrals are approximated from data by Riemann sum or trapezoid rule. We then find a minimizer of the loss functional over  $\mathcal{H}$  by least squares

$$\hat{\phi}_{\mathcal{H}} = \sum_{i=1}^n \hat{c}_i \phi_i, \quad \text{where } \hat{c} = \arg \min_{c \in \mathbb{R}^n} \mathcal{E}(c), \quad \mathcal{E}(c) = c^\top A c - 2c^\top b. \quad (5.3)$$

In practice, we compute minimizer by  $\hat{c} = A^{-1}b$  when the normal matrix  $A$  is well-conditioned. When  $A$  is ill-conditioned or singular, we use pseudo-inverse or regularization to obtain  $\hat{c} = (A + \lambda B)^{-1}b$  with  $\lambda B$  being a proper regularization. We select the optimal dimension of the hypothesis space that minimizes the loss functional so as to avoid undercutting or overfitting.

As we show in the next section, the normal matrix becomes ill-conditioned as the dimension of the hypothesis space increases. Thus, regularization becomes necessary. In Section 5.3, we will discuss regularizations by truncated SVD and the connection to the RKHSs.

## 5.2 Identifiability and ill-conditioned normal matrix

We show next that the normal matrix becomes ill-conditioned as the hypothesis space's dimension increases, because its smallest eigenvalue converges to zero.

**Theorem 5.1.** *Let  $\mathcal{H} = \text{span}\{\phi_i\}_{i=1}^n \subset L^2(\mathcal{X})$  and  $A$  be the matrix in (5.1). Then, restricted on  $\mathcal{H}$ ,*

- (a) *The operator  $\mathcal{L}_{\bar{G}_T} : L^2(\mathcal{X}) \rightarrow L^2(\mathcal{X})$  in (3.8) has the same eigenvalues as  $A$ , that is, if  $\mathcal{L}_{\bar{G}_T} \varphi = \lambda \varphi$  for some  $\varphi = \sum_{i=1}^n c_i \phi_i \in \mathcal{H}$ , then  $\lambda$  is an eigenvalue of  $A$ ;*
- (b) *The operator  $\mathcal{L}_{\bar{R}_T} : L^2(\bar{\rho}_T) \rightarrow L^2(\bar{\rho}_T)$  in (3.12) has eigenvalues solved by the generalized eigenvalue problem*

$$A c = \lambda P c, \quad \text{with } P = (\langle\langle \phi_i, \phi_j \rangle\rangle_{L^2(\bar{\rho}_T)}). \quad (5.4)$$

*In particular, the smallest (generalized) eigenvalue of  $A$  converges to zero as  $n \rightarrow \infty$ .*

*Proof.* Part (a) follows directly from (3.9). For Part (b), note that if  $\varphi = \sum_{i=1}^n c_i \phi_i$  is an eigenfunction of  $\mathcal{L}_{\bar{R}_T}|_{\mathcal{H}}$  with eigenvalue  $\lambda$ , then by definition of  $P$ , we have

$$\lambda (Pc)_k = \langle \lambda \varphi, \phi_k \rangle_{L^2(\bar{\rho}_T)} = \langle \mathcal{L}_{\bar{R}_T} \varphi, \phi_k \rangle_{L^2(\bar{\rho}_T)} = \sum_{i=1}^n c_i \langle \mathcal{L}_{\bar{R}_T} \phi_i, \phi_k \rangle_{L^2(\bar{\rho}_T)} = (Ac)_k,$$

where the last equality follows from (3.13).

The smallest (generalized) eigenvalue of  $A$  converges to zero since the integral operators are compact.  $\square$

**Remark 5.2** (Ill-conditioned normal matrix). *It follows from the above theorem that the normal matrix  $A$  becomes ill-conditioned as  $n$  increases. The rank of  $A$  is no larger than the dimension of either the RKHSs  $\mathcal{H}_G$  or  $\mathcal{H}_R$  because they are the images of the square root of the integral operators.*

Table 2: Notation of variables in the eigenvalue problems on  $\mathcal{H}$ .

	in $L^2(\mathcal{X})$	in $L^2(\bar{\rho}_T)$
integral kernel and operator	$\widehat{G}_T \approx \bar{G}_T, \mathcal{L}_{\widehat{G}_T} \approx \mathcal{L}_{\bar{G}_T}$	$\widehat{R}_T \approx \bar{G}_T, \mathcal{L}_{\widehat{R}_T} \approx \mathcal{L}_{\bar{R}_T}$
eigenfunction and eigenvalue	$\mathcal{L}_{\widehat{G}_T} \widehat{\varphi}_k = \widehat{\lambda}_k \widehat{\varphi}_k$	$\mathcal{L}_{\widehat{R}_T} \widehat{\psi}_k = \widehat{\gamma}_k \widehat{\psi}_k$
eigenvector and eigenvalue	$A \vec{\varphi}_k = \widehat{\lambda}_k \vec{\varphi}_k$	$A \vec{\psi}_k = \widehat{\gamma}_k P \vec{\psi}_k$

**Remark 5.3** (Well-posedness). *The inverse problem of identifying the interaction kernel is ill-posed in general: since the normal matrix becomes ill-conditioned as  $n$  increases, a small perturbation in the eigen-space of the smallest eigenvalues may lead to large errors in the estimator. More specifically, we are solving the inverse problem  $\hat{\phi} = \mathcal{L}_{\bar{R}_T}^{-1}(\mathcal{L}_{\bar{R}_T} \phi_{true})$  in  $L^2(\bar{\rho}_T)$ , where  $\mathcal{L}_{\bar{R}_T}^{-1}$  is an unbounded operator on  $\mathcal{H}_R$ . In practice, the normal matrix approximates the operator  $\mathcal{L}_{\widehat{R}_T}$ , and  $b$  approximates  $\mathcal{L}_{\bar{R}_T} \phi_{true}$ . On the other hand, if the error in  $b$  occurs only in the eigen-space of large eigenvalues, then the error will be controlled and the inverse problem is well-posed.*

We summarize the notations in these two eigenvalue problems in Table 2.

### 5.3 Two regularization norms

Theorem 5.1 suggests that the normal matrix  $A$  is, albeit invertible, often ill-conditioned, so  $A^{-1}$  will amplify the numerical error in  $b$ , leading to error-prone estimator  $\hat{c} = A^{-1}b$ . In this case, it is necessary to add regularization. The widely-used Tikhonov regularization [10] with  $H^1(\mathcal{X})$  norm and with the L-curve method [11] works well in [16]. There are two factors in the method, the regularization norm and the parameter. The L-curve method provides a way to select the parameter for a given regularization norm. However, the choice of regularization norm is problem dependent and is less explored.

In this section we compare two regularization norms: the  $L^2(\mathcal{X})$  and  $L^2(\bar{\rho}_T)$  norms. We show by numerical examples that the weighted norm leads to a better conditioned inverse problem.

**Tikhonov regularization** Instead of minimizing the loss functional  $\mathcal{E}(\psi)$ , Tikhonov regularization minimizes  $\mathcal{E}_\lambda(\psi) = \mathcal{E}(\psi) + \lambda \|\|\| \psi \|\|\|^2$ , where the regularization norm  $\|\|\| \cdot \|\|\|$  is to be specified later. Intuitively,  $\|\|\| \cdot \|\|\|$  gives an priori assumption on the regularity of the estimator. The minimizer of  $\mathcal{E}_\lambda$  over the hypothesis space  $\mathcal{H}$  is determined by

$$\hat{c}_\lambda = (A + \lambda B)^{-1}b \quad (5.5)$$

Here  $B$  is a  $n \times n$  matrix determined by the norm  $\|\|\| \cdot \|\|\|$ . The presence of  $B$  can reduce the condition number of the linear system (5.5), so that the error in  $b$  is not amplified by the inversion.

The major difficulty in Tikhonov regularization is to choose the regularization norm  $\|\|\| \cdot \|\|\|$  (hence the matrix  $B$ ) along with an optimal regularization parameter  $\lambda$ . Given a regularization norm  $\|\|\| \cdot \|\|\|$ , an optimal parameter aims to balance the decrement of the loss functional  $\mathcal{E}$  and the increment of the norm. Among various methods, the L-curve method [11] selects  $\lambda$  at where the largest curvature occurs on a parametric curve of  $(\log(\mathcal{E}(\hat{c}_\lambda)), \log(\|\|\| \hat{c}_\lambda \|\|\|))$ ; the truncated SVD methods [10] discard the smallest singular values of  $A$  and solve the modified least square problem.

We focus on comparing two regularization norms from the ambient function spaces of learning:

$$\begin{aligned} \text{Unweighted } L^2 \text{ norm: } \quad \|\|\| \cdot \|\|\| &= \|\cdot\|_{L^2(\mathcal{X})} & B_{ij} &= \langle \phi_i, \phi_j \rangle_{L^2(\mathcal{X})}, \\ \text{Weighted } L^2 \text{ norm: } \quad \|\|\| \cdot \|\|\| &= \|\cdot\|_{L^2(\bar{\rho}_T)} & B_{ij} &= \langle \phi_i, \phi_j \rangle_{L^2(\bar{\rho}_T)}. \end{aligned} \quad (5.6)$$

With the basis functions being piecewise constants on a uniform partition of  $\mathcal{X}$ , we have  $B = \Delta r I_n$  for the first case and  $B = P$  with  $P$  defined in (5.4). In particular, when we take  $\phi_i$  to be piecewise constants, the matrix  $P$  becomes a diagonal matrix

$$P = \text{diag}(\widehat{\rho}_T(r_1), \dots, \widehat{\rho}_T(r_n))\Delta r, \quad (5.7)$$

where  $\{\widehat{\rho}_T(r_n)\}$  is the average density on the interval  $[r_{i-1}, r_i]$ , i.e.  $\widehat{\rho}_T(r_i) = \frac{1}{\Delta r} \int_{r_{i-1}}^{r_i} \bar{\rho}_T(r) dr$ . Note that we can represent  $\bar{\rho}_T$  by  $\widehat{\rho}_T = \sum_{i=1}^n \widehat{\rho}_T(r_i) \phi_i$ ,

**Weighted and unweighted SVD** Singular value decomposition (SVD) analysis provides an effective tool for understanding the features of Tikhonov regularization [10, 11] so as to select an optimal parameter. The idea is to study the decay of the singular values (or eigenvalues for symmetric matrices) and compare it with the projection of  $b$  in the eigenspace. More precisely, let  $A = \sum_{i=1}^n \sigma_i \mathbf{u}_i \mathbf{u}_i^\top$ , where  $\sigma_i$  are the decreasingly-ordered eigenvalues of  $A$  and  $\mathbf{u}_i \in \mathbb{R}^n$  is the corresponding orthonormal eigenvector. The regularized regression solution to  $Ac = b$  is  $\widehat{c} = \sum_{i=1}^n f_i \frac{\mathbf{u}_i^\top b}{\sigma_i} \mathbf{u}_i$ , where the filter factors  $f_i$  from regularization aim to filter out the error-prone contributions from  $\frac{\mathbf{u}_i^\top b}{\sigma_i} \mathbf{u}_i$  when  $\sigma_i$  is small. This promotes the discrete Picard condition [10] which requires  $\mathbf{u}_i^\top b$  to decay faster than  $\sigma_i$ .




Figure 1: SVD analysis of the regression in three example potentials: cubic potential with  $\phi(r) = 3r^2$ , opinion dynamics with  $\phi$  being piecewise linear, and the attraction-repulsion potential with  $\phi(r) = r - r^{-1.5}$ . Here  $R$  represents the weighted SVD and  $G$  represents the unweighted SVD. In all three examples, the weighted SVD has larger eigenvalues than those of the unweighted SVD; and it has slightly smaller ratios  $\frac{\mathbf{u}_i^\top b}{\sigma_i}$ .

Here we demonstrate that the inverse problem is better conditioned on  $L^2(\bar{\rho}_T)$  than on  $L^2(\mathcal{X})$ , which corresponds to the two regularization norms in (5.6). These two function spaces lead to weighted and unweighted SVD, in which we compute the eigenvalues and eigenvectors as follows:

$$\begin{aligned} \text{Unweighted SVD: } & \Phi D_G \Phi^\top = A, \text{ with } \Phi^\top \Phi = I, \\ \text{Weighted SVD: } & \Psi D_R \Psi^\top = A, \text{ with } \Psi^\top P \Psi = I. \end{aligned} \quad (5.8)$$

Here  $D_G$  and  $D_R$  are diagonal matrices. The solution of the unweighted SVD,  $\Phi$  and  $D_G$ , provides orthonormal eigenvectors and eigenvalues of  $A = \mathcal{L}_{\bar{G}_T}|_{\mathcal{H}}$  on  $L^2(\mathcal{X})$ . In the weighted SVD, the generalized eigenvalue problem for  $(A, P)$  with solution  $\Psi$  and  $D_R$  are for  $A = \mathcal{L}_{\bar{R}_T}|_{\mathcal{H}}$  on  $L^2(\bar{\rho}_T)$ .

Figure 1 shows the decay of the singular values from the unweighted SVD (denoted by  $G$ ; and “G: b-projection” refers to  $|\mathbf{u}_i^\top b|$  with  $\mathbf{u}_i$  being the columns of  $\Phi$ ) and weighted SVD (denoted by  $R$ , and “R: b-projection” refers to  $|\mathbf{u}_i^\top b|$  with  $\mathbf{u}_i$  being the columns of  $\Psi$ ) for the three examples in [16]. These examples are cubic potential with  $\phi(r) = 3r^2$ , opinion dynamics with  $\phi$  being piecewise linear, and the attraction-repulsion potential with  $\phi(r) = r - r^{-1.5}$  (or equivalently  $\Phi(r) = \frac{r^2}{2} + 2r^{-0.5}$ ). In all these examples, the weighted SVD has larger eigenvalues than those of the unweighted SVD; and it has slightly smaller ratios of b-projection-to-eigenvalue. Thus, the weighted SVD leads to a better-posed inversion.

Figure 2 shows the regularized estimators via truncated SVD according to both the weighted and unweighted SVD. We pick a truncation level such that the sum of the top several singular values take 99% of the total summation of the singular values. The relative  $L^2(\rho)$  error is large, since we are using degree-0 spline basis, which is consistent to our theoretical analysis. One can obtain better results by using spline basis with higher order regularity [16]. Our goal is to illustrate that the regularization based on  $\mathcal{H}_R$  leads a better result. The corresponding basis are presented in Figure 3. When the kernel is smooth (i.e., the cubic potential and the attraction-repulsion potential), the weighted SVD leads to more accurate estimator than the unweighted SVD. When the kernel is discontinuous (i.e., the opinion dynamics), the two performed similarly.




Figure 2: Regularized estimators via unweighted ( $\mathcal{H}_G$ ) or weighted ( $\mathcal{H}_R$ ) SVD in three examples, superimposed with the exploration measure  $\rho$ . The un-regularized estimator (“all spline”) has a large error in each example. The weighted SVD has smaller errors than unweighted SVD, while both significantly improve the un-regularized estimate.

## 5.4 Regularization and the RKHSs

Here we connect the regularization with the RKHSs. In particular, we show that the regularization by truncated SVD is equivalent to selecting low-frequency orthogonal basis of the RKHSs.

**Unweighted SVD and the RKHS  $\mathcal{H}_G$**  First, we show that the integral kernel  $\overline{G}_T$  on the basis in  $\mathcal{H}$  is equivalent to  $A$  up to a multiplicative factor. We approximate  $\overline{G}_T$  by  $\widehat{\overline{G}}_T(r, s) = \sum_{i,j=1}^n \widehat{G}_T(r_i, r_j) \phi_i(r) \phi_j(s)$ , where  $\widehat{G}_T(r_i, r_j)$  is the average of  $\overline{G}_T$  over the region  $[r_{i-1}, r_i] \times [r_{j-1}, r_j]$ . Recall the definition of  $\overline{G}_T$  as in (2.6) and the bilinear form in (2.5), we have

$$\widehat{\overline{G}}_T(r_i, r_j) = \frac{1}{(\Delta r)^2} \int \int \phi_i(r) \phi_j(s) \overline{G}_T(r, s) dr ds = \frac{1}{(\Delta r)^2} \langle \phi_i, \phi_j \rangle = A_{i,j}.$$




Figure 3: Eigen-basis in the estimation via weighted and unweighted SVD in Figure 2. The weighted operator  $\mathcal{L}_R$  has smoother eigenfunctions than the unweighted operator  $\mathcal{L}_G$ .

Consider the eigenfunctions of the approximated operator  $\mathcal{L}_{\widehat{G}_T}$  given by  $\{\widehat{\varphi}_k\}$ , where  $\widehat{\varphi}_k(r) = \sum_{i=1}^n \widehat{\varphi}_k(r_i) \phi_i(r)$ . Then  $\mathcal{L}_{\widehat{G}_T} \widehat{\varphi}_k = \widehat{\lambda}_k \widehat{\varphi}_k$  implies

$$\begin{aligned}
\widehat{\lambda}_k \widehat{\varphi}_k(r) &= \int_{\mathbb{R}^d} \widehat{G}_T(r, s) \widehat{\varphi}_k(s) ds = \int_{\mathbb{R}^d} \left( \sum_{i,j=1}^n \widehat{G}_T(r_i, r_j) \phi_i(r) \phi_j(s) \right) \left( \sum_{i=1}^n \widehat{\varphi}_k(r_i) \phi_i(s) \right) ds \\
&= (\Delta r)^2 \sum_{j=1}^n \left( \sum_{i=1}^n \widehat{G}_T(r_i, r_j) \widehat{\varphi}_k(r_i) \right) \phi_j(r). \tag{5.9}
\end{aligned}$$

This is equivalent to say that,  $\vec{\varphi}_k = (\widehat{\varphi}_k(r_1), \dots, \widehat{\varphi}_k(r_n))$  for  $k = 1, \dots, n$  are eigenvectors of the matrix  $A$  corresponding to eigenvalues  $\{\widehat{\lambda}_k\}_{k=1}^n$  in the unweighted SVD in (5.8).

Thus, the discrete approximation of the RKHS  $\mathcal{H}_G$  is span  $\{\widehat{\varphi}_1, \dots, \widehat{\varphi}_m\}$ , the span of the eigenfunctions with positive eigenvalues. In particular, the low frequency eigenfunctions .

Next, we show that a truncated SVD regularization is a selection of low-frequency basis of the RKHS. Recall that in a truncated SVD, one truncates the eigenvalues of  $A$  to avoid amplifying the error in  $b$ . Keeping the first  $m$  eigenvalues above a threshold, we obtain the least squares solution

$$\widehat{c} = \sum_{k=1}^m \widehat{\lambda}_k^{-1} (\vec{\varphi}_k^\top b) \vec{\varphi}_k. \tag{5.10}$$

Equivalently, we can view the truncated SVD as finding a solution on  $\mathcal{H}_G^m = \text{span} \{\widehat{\varphi}_1, \dots, \widehat{\varphi}_m\}$ . Note that  $\mathcal{H}_G^m \subset \mathcal{H}_G$  is spanned by the first  $m$  eigenfunctions of the operator  $\mathcal{L}_{\widehat{G}_T}$ , which filters out the high-frequency functions. Suppose the estimator  $\widehat{\phi}$  in  $\mathcal{H}_G^m$  is given by  $\widehat{\phi}(r) = \sum_{k=1}^m \alpha_k \widehat{\varphi}_k(r) = \sum_{i=1}^n (\Phi_m \alpha)_i \phi_i(r)$ , where  $\alpha = (\alpha_1, \dots, \alpha_m)$  is a vector in  $\mathbb{R}^m$ ,  $\Phi_m$  is the matrix consists of the first  $m$  columns of  $\Phi$  and  $(\Phi_m \alpha)_i$  represents the  $i$ th component of the vector  $\Phi_m \alpha$ . Then the loss functional evaluated on  $\widehat{\phi}$  writes

$$\mathcal{E}(\widehat{\phi}) = \alpha^\top \Phi_m^\top A \Phi_m \alpha - 2b^\top \Phi_m \alpha.$$

The minimizer of  $\mathcal{E}(\hat{\phi})$  is determined by the  $\alpha$  such that  $\Phi_m^\top A \Phi_m \alpha = \Phi_m^\top b$ . Notice that  $\Phi_m^\top A \Phi_m = \text{diag}(\widehat{\lambda}_1, \dots, \widehat{\lambda}_m) = D_{G,m}$ , we have  $\alpha = D_{G,m}^{-1} \Phi_m^\top b$ . If we write  $\hat{\phi}(r) = \sum_{i=1}^n c_i \phi_i(r)$ , then

$$c = \Phi_m \alpha = \Phi_m D_{G,m}^{-1} \Phi_m^\top b$$

This is the same as the estimator in (5.10).

**Weighted SVD and the RKHS  $\mathcal{H}_R$**  We first show that solving eigenvalue of the operator  $\mathcal{L}_{\widehat{R}_T}$  over  $\mathcal{H}$  is equivalent to the generalized eigenvalue problem in (5.8). The approximation of  $\widehat{R}_T$  is given by  $\widehat{R}_T(r, s) = \sum_{i,j=1}^n \widehat{R}_T(r_i, r_j) \phi_i(r) \phi_j(s)$ , where  $\widehat{R}_T(r_i, r_j) = \frac{\widehat{G}_T(r_i, r_j)}{\widehat{\rho}_T(r_i) \widehat{\rho}_T(r_j)}$ . Then we can derive the discrete eigenfunctions of the operator  $\mathcal{L}_{\widehat{R}_T}$ . Suppose  $\widehat{\psi}_k(r) = \sum_{i=1}^n \psi_k(r_i) \phi_i(r)$  is an eigenfunction of  $\mathcal{L}_{\widehat{R}_T}$  with corresponding eigenvalue  $\widehat{\gamma}_k$ . Then we have, as in (5.9),

$$\widehat{\gamma}_k \left( \sum_{i=1}^n \widehat{\psi}_k(r_i) \phi_i(r) \right) = (\Delta r)^2 \sum_{j=1}^n \left( \sum_{i=1}^n \frac{\widehat{G}_T(r_i, r_j)}{\widehat{\rho}_T(r_i)} \widehat{\psi}_k(r_i) \right) \phi_j(r).$$

Denote  $\vec{\psi}_k = (\widehat{\psi}_k(r_1), \dots, \widehat{\psi}_k(r_n))$  for  $k = 1, \dots, n$ , with  $P$  in (5.7), we have

$$A \vec{\psi}_k = \widehat{\gamma}_k P \vec{\psi}_k.$$

Since both  $A$  and  $P$  are symmetric, it is well-defined that  $\{\vec{\psi}_k\}_{k=1}^n$  are the generalized eigenvectors of the pair  $(A, P)$  with eigenvalues  $\{\widehat{\gamma}_k\}_{k=1}^n$ . Let  $\{\vec{\psi}_k\}_{k=1}^n$  be the columns of  $\Psi$  and let  $D_R = \text{diag}(\widehat{\gamma}_1, \dots, \widehat{\gamma}_n)$ , we have the second equation in (5.8).

Next, consider the inference on the hypothesis space  $\mathcal{H}_R^m = \text{span}\{\widehat{\psi}_1, \dots, \widehat{\psi}_m\}$ . Suppose the estimator  $\hat{\phi}(r)$  in  $\mathcal{H}_R^m$  is given by  $\hat{\phi} = \sum_{k=1}^m \alpha_k \widehat{\psi}_k(r) = \sum_{i=1}^n (\Psi_m \alpha)_i \phi_i(r)$ , where  $\alpha = (\alpha_1, \dots, \alpha_m)$  is a vector in  $\mathbb{R}^m$ ,  $\Psi_m$  is the matrix consists of the first  $m$  columns of  $\Psi$  and  $(\Psi_m \alpha)_i$  represents the  $i$ th component of the vector  $\Psi_m \alpha$ . Then the loss functional evaluated on  $\hat{\phi}$  writes

$$\mathcal{E}(\hat{\phi}) = \alpha^\top \Psi_m^\top A \Psi_m \alpha - 2b^\top \Psi_m \alpha.$$

The minimizer of  $\mathcal{E}(\hat{\phi})$  is determined by the  $\alpha$  such that  $\Psi_m^\top A \Psi_m \alpha = \Psi_m^\top b$ . Notice that  $\Psi_m^\top A \Psi_m = \text{diag}(\widehat{\lambda}_1, \dots, \widehat{\lambda}_m) = D_{R,m}$ , we have  $\alpha = D_{R,m}^{-1} \Psi_m^\top b$ . If we write  $\hat{\phi}(r) = \sum_{i=1}^n c_i \phi_i(r)$ , then

$$\hat{c} = \Psi_m \alpha = \Psi_m D_{R,m}^{-1} \Psi_m^\top b.$$

This is equivalent to  $c = \sum_{k=1}^m \frac{1}{\widehat{\gamma}_k} \left( \vec{\psi}_k^\top b \right) \vec{\psi}_k$ , a truncated SVD with  $m$  eigenvectors.

## A Appendix

**Positive definite functions** We review the definitions and properties of positive definite kernels. The following is a real-variable version of the definition in [3, p.67].

**Definition A.1** (Positive definite function). *Let  $X$  be a nonempty set. A function  $G : X \times X \rightarrow \mathbb{R}$  is positive definite if and only if it is symmetric (i.e.  $G(x, y) = G(y, x)$ ) and  $\sum_{j,k=1}^n c_j c_k G(x_j, x_k) \geq 0$  for all  $n \in \mathbb{N}$ ,  $\{x_1, \dots, x_n\} \subset X$  and  $\mathbf{c} = (c_1, \dots, c_n) \in \mathbb{R}^n$ . The function  $\phi$  is strictly positive definite if the equality hold only when  $\mathbf{c} = \mathbf{0} \in \mathbb{R}^n$ .*



**Theorem A.2** (Properties of positive definite kernels). *Suppose that  $k, k_1, k_2 : X \times X \subset \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$  are positive definite kernels. Then*

- (a)  $k_1 k_2$  is positive definite. ([3, p.69])
- (b) Inner product  $\langle u, v \rangle = \sum_{j=1}^d u_j v_j$  is positive definite ([3, p.73])
- (c)  $f(u)f(v)$  is positive definite for any function  $f : X \rightarrow \mathbb{R}$  ([3, p.69]).

**RKHS and positive integral operators** We review the definitions and properties of Mercer kernel, RKHS, and related integral operator on compact domain (see e.g., [8]) and non-compact domain (see e.g., [29]).

Let  $(X, d)$  be a metric space and  $G : X \times X \rightarrow \mathbb{R}$  be continuous and symmetric. We say that  $G$  is a Mercer kernel if it is positive definite (as in Definition A.1). The reproducing kernel Hilbert space (RKHS)  $\mathcal{H}_G$  associated with  $G$  is defined to be closure of  $\text{span}\{G(x, \cdot) : x \in X\}$  with the inner product

$$\langle f, g \rangle_{\mathcal{H}_G} = \sum_{i=1, j=1}^{n, m} c_i d_j G(x_i, y_j)$$

for any  $f = \sum_{i=1}^n c_i G(x_i, \cdot)$  and  $g = \sum_{j=1}^m d_j G(x_j, \cdot)$ . It is the unique Hilbert space such that  $\text{span}\{G(\cdot, y), y \in X\}$  is dense in  $\mathcal{H}_G$  and having reproducing kernel property in the sense that for all  $f \in \mathcal{H}_G$  and  $x \in X$ ,  $f(x) = \langle G(x, \cdot), f \rangle_G$  (see [8, Theorem 2.9]).

By means of the Mercer Theorem, we can characterize the RKHS  $\mathcal{H}_G$  through the integral operator associated with the kernel. Let  $\mu$  be a nondegenerate Borel measure on  $(X, d)$  (that is,  $\mu(U) > 0$  for every open set  $U \subset X$ ). Define the integral operator  $L_G$  on  $L^2(X, \mu)$  by

$$L_G f(x) = \int_X G(x, y) f(y) d\mu(y).$$

The RKHS has the operator characterization (see e.g., [8, Section 4.4] and [29]):

**Theorem A.3.** *Assume that the  $G$  is a Mercer kernel and  $G \in L^2(X \times X, \mu \otimes \mu)$ . Then*

1.  $L_G$  is a compact positive self-adjoint operator. It has countably many positive eigenvalues  $\{\lambda_i\}_{i=1}^{\infty}$  and corresponding orthonormal eigenfunctions  $\{\phi_i\}_{i=1}^{\infty}$ .
2.  $\{\sqrt{\lambda_i} \phi_i\}_{i=1}^{\infty}$  is an orthonormal basis of the RKHS  $\mathcal{H}_G$ .
3. The RKHS is the image of the square root of the integral operator, i.e.,  $\mathcal{H}_G = L_G^{1/2} L^2(X, \mu)$ .

**Acknowledgements.** FL is grateful for supports from NSF-1913243, FA9550-20-1-0288 and DE-SC0021361. FL would like to thank Mauro Maggioni and P-E Jabin for helpful discussions.

## References

- [1] A. S. Baumgarten and K. Kamrin. A general constitutive model for dense, fine-particle suspensions validated in many geometries. *Proc Natl Acad Sci USA*, 116(42):20828–20836, 2019.
- [2] N. Bell, Y. Yu, and P. J. Mucha. Particle-based simulation of granular materials. In *Proceedings of the 2005 ACM SIGGRAPH/Eurographics Symposium on Computer Animation - SCA '05*, page 77, Los Angeles, California, 2005. ACM Press.
- [3] C. Berg, J. P. R. Christensen, and P. Ressel. *Harmonic analysis on semigroups: theory of positive definite and related functions*, volume 100. New York: Springer, 1984.
- [4] M. Bonafini, M. Fornasier, and B. Schmitzer. Data-driven entropic spatially inhomogeneous evolutionary games. *arXiv preprint arXiv:2103.05429*, 2021.

- [5] M. Bongini, M. Fornasier, M. Hansen, and M. Maggioni. Inferring interaction rules from observations of evolutive systems I: The variational approach. *Mathematical Models and Methods in Applied Sciences*, 27(05):909–951, 2017.
- [6] J. A. Carrillo, K. Craig, and Y. Yao. Aggregation-diffusion equations: dynamics, asymptotics, and singular limits. In *Active Particles, Volume 2*, pages 65–108. Springer, 2019.
- [7] J. A. Carrillo, M. DiFrancesco, A. Figalli, T. Laurent, and D. Slepčev. Global-in-time weak measure solutions and finite-time aggregation for nonlocal interaction equations. *Duke Math. J.*, 156(2):229–271, 2011.
- [8] F. Cucker and D. X. Zhou. *Learning theory: an approximation theory viewpoint*, volume 24. Cambridge University Press, 2007.
- [9] J. Fan and Q. Yao. *Nonlinear Time Series: Nonparametric and Parametric Methods*. Springer, New York, NY, 2003.
- [10] P. C. Hansen. REGULARIZATION TOOLS: A Matlab package for analysis and solution of discrete ill-posed problems. *Numer Algor*, 6(1):1–35, 1994.
- [11] P. C. Hansen. The L-curve and its use in the numerical treatment of inverse problems. pages 119–142, 2000.
- [12] P.-E. Jabin and Z. Wang. Mean Field Limit for Stochastic Particle Systems. In N. Bellomo, P. Degond, and E. Tadmor, editors, *Active Particles, Volume 1*, pages 379–402. Springer International Publishing, Cham, 2017.
- [13] P.-E. Jabin and Z. Wang. Quantitative estimates of propagation of chaos for stochastic systems with  $w^{-1,\infty}$  kernels. *Invent. math.*, 214(1):523–591, 2018.
- [14] I. Karatzas and S. E. Shreve. Brownian motion. In *Brownian Motion and Stochastic Calculus*. Springer, second edition, 1998.
- [15] Y. A. Kutoyants. *Statistical inference for ergodic diffusion processes*. Springer, 2004.
- [16] Q. Lang and F. Lu. Learning interaction kernels in mean-field equations of 1st-order systems of interacting particles. *to appear in SIAM J. Sci. Comput.*; *preprint arXiv:2010.15694*, 2020.
- [17] Z. Li and F. Lu. On the coercivity condition in the learning of interacting particle systems. *arXiv preprint arXiv:2011.10480*, 2020.
- [18] Z. Li, F. Lu, M. Maggioni, S. Tang, and C. Zhang. On the identifiability of interaction functions in systems of interacting particles. *Stochastic Processes and their Applications*, 132:135–163, 2021.
- [19] F. Lu, M. Maggioni, and S. Tang. Learning interaction kernels in heterogeneous systems of agents from multiple trajectories. *Journal of Machine Learning Research*, 22(32):1–67, 2021.
- [20] F. Lu, M. Maggioni, and S. Tang. Learning interaction kernels in stochastic systems of interacting particles from multiple trajectories. *to appear in Foundations of Computational Mathematics*, 2021.
- [21] F. Lu, M. Zhong, S. Tang, and M. Maggioni. Nonparametric inference of interaction laws in systems of agents from trajectory data. *Proceedings of the National Academy of Sciences of the United States of America*, 116(29):14424–14433, 2019.
- [22] F. Malrieu. Convergence to equilibrium for granular media equations and their Euler schemes. *Ann. Appl. Probab.*, 13(2):540–560, 2003.
- [23] S. Méléard. *Asymptotic Behaviour of Some Interacting Particle Systems; McKean-Vlasov and Boltzmann Models*, volume 1627, pages 42–95. Springer Berlin Heidelberg, Berlin, Heidelberg, 1996.

- [24] C. A. Micchelli, Y. Xu, and H. Zhang. Universal kernels. *J. Mach. Learn. Res.*, 7:2651–2667, Dec. 2006.
- [25] S. Mostch and E. Tadmor. Heterophilious Dynamics Enhances Consensus. *Siam Review*, 56(4):577 – 621, 2014.
- [26] C. E. Rasmussen. Gaussian processes in machine learning. In *Summer school on machine learning*, pages 63–71. Springer, 2003.
- [27] L. Sharrock, N. Kantas, P. Parpas, and G. A. Pavliotis. Parameter estimation for the mckean-vlasov stochastic differential equation. *arXiv preprint arXiv:2106.13751*, 2021.
- [28] B. K. Sriperumbudur, K. Fukumizu, and G. R. Lanckriet. Universality, characteristic kernels and rkhs embedding of measures. *Journal of Machine Learning Research*, 12(70):2389–2410, 2011.
- [29] H. Sun. Mercer theorem for rkhs on noncompact sets. *Journal of Complexity*, 21(3):337 – 349, 2005.
- [30] A.-S. Sznitman. *Topics in Propagation of Chaos*, volume 1464, pages 165–251. Springer Berlin Heidelberg, Berlin, Heidelberg, 1991.
- [31] T. Vicsek and A. Zafeiris. Collective motion. *Physics Reports*, 517:71 – 140, 2012.
- [32] H. Wendland. *Scattered Data Approximation*. Cambridge Monographs on Applied and Computational Mathematics. Cambridge University Press, 2004.