

# Thakurta metric does not describe a cosmological black hole

Tomohiro Harada<sup>a,\*</sup>, Hideki Maeda<sup>b,†</sup> and Takuma Sato<sup>a,‡</sup>

<sup>a</sup> *Department of Physics, Rikkyo University, Tokyo 171-8501, Japan and*

<sup>b</sup> *Department of Electronics and Information Engineering,  
Hokkai-Gakuen University, Sapporo 062-8605, Japan*

(Dated: January 8, 2022)

Recently, the Thakurta metric has been adopted as a model of primordial black holes by several authors. We show that the spacetime described by this metric has neither black-hole event horizon nor black-hole trapping horizon and involves the violation of the null energy condition as a solution of the Einstein equation. Therefore, this metric does not describe a cosmological black hole in the early universe.

PACS numbers: 04.70.Bw, 97.60.Lf, 04.20.Gz

Primordial black holes [1, 2] have attracted intense attention as it has been revealed that a considerable fraction of binary black holes discovered by gravitational wave observations by LIGO and Virgo have masses in the range of several tens of solar masses. In fact, it is proposed that such very massive black holes might be of cosmological origin [3–5]. Recently, from the data observed by LIGO and Virgo, Franciolini et al. [6] has concluded that the evidence for a primordial population is decisively favored compared to the null hypothesis. To discuss the physical properties of primordial black holes, Boehm et al. [7] and Picker [8] adopt the Thakurta metric [9, 10]. These works are criticized by Hütsi et al. [11] as this spacetime entails accretion history which is highly unlikely from a physical point of view. This controversy continues by a further comment [12] and a reply [13]. In this letter, we terminate the controversy on the Thakurta metric to turn the discussion to the right direction to seek for a mathematical model of primordial black holes. Note that the Thakurta metric is different from McVittie’s solutions [10, 14], whose causal structures are analyzed in detail in [15–18]. We use the units in which  $c = G = 1$  and follow the notation in Wald [19].

The Thakurta metric is given by

$$ds^2 = a^2(\eta) [-f(r)d\eta^2 + f(r)^{-1}dr^2 + r^2d\Omega^2], \quad (1)$$

$$f(r) := 1 - \frac{2M}{r}$$

where  $d\Omega^2$  is the metric on the unit 2-sphere and  $M$  is a constant [9, 10]. We assume  $M > 0$  throughout this letter. The spacetime approaches the flat Friedmann-Lemaître-Robertson-Walker (FLRW) solution with a conformal time  $\eta$  as  $r \rightarrow \infty$ . If the scale factor is given by  $a(\eta) = a_0\eta^\alpha$ , where  $a_0$  and  $\alpha$  are positive constants, the asymptotic metric is that of the flat FLRW solution with a perfect fluid that obeys the equation of state  $p = w\rho$  ( $w > -1/3$ ) through the relation  $\alpha = 2/(1+3w)$ , corresponding to a decelerated cosmological expansion in the domain  $\eta > 0$ .

The Kretschmann invariant  $K := R_{abcd}R^{abcd}$  of the

spacetime (1) is calculated to

$$K = \frac{4[3\alpha^2(1+\alpha^2)r^8 + 4M^2\eta^2\{3\eta^2(r-2M)^2 - \alpha^2r^4\}]}{a_0^4r^6(r-2M)^2\eta^{4(1+\alpha)}}, \quad (2)$$

which shows that there is a scalar polynomial curvature singularity at each of  $r = 0$  and  $\eta = 0$ .  $\eta = 0$  with  $r > 2M$  is a spacelike singularity. However, the property of  $r = 2M$  is subtle.

The line element (1) can be rewritten as

$$ds^2 = a^2(\eta) [f(r)(-d\eta^2 + dr^{*2}) + r^2d\Omega^2], \quad (3)$$

where  $r^* := r + 2M \ln[(r-2M)/2M]$ . Since  $r \rightarrow 2M$  corresponds to  $r^* \rightarrow -\infty$ , it is a null boundary in the Penrose diagram.

Now we show that future-directed ingoing radial null geodesics are incomplete at curvature singularity at  $(r, \eta) = (2M, \infty)$ . In the Thakurta spacetime (1), an ingoing radial null geodesic  $\gamma$  is given by  $u_+ = u_{+(0)} = \text{const}$ , where  $u_\pm := \eta \pm r^*$ . The Kretschmann invariant in the limit of  $r \rightarrow 2M$  along  $\gamma$  is then calculated to

$$\lim_{r \rightarrow 2M} K|_\gamma = \lim_{r^* \rightarrow -\infty} -\frac{\alpha^2 e^{-(r^*-2M)/M}}{M^4 a_0^4 (-r^*)^{2+4\alpha}} = -\infty. \quad (4)$$

Let  $k^\mu$  be a tangent vector of  $\gamma$ . Since the spacetime admits a conformal Killing vector  $\xi^a = (\partial/\partial\eta)^a$ , there exists a conserved quantity  $C = -\xi^a k_a = a^2(\eta)f(r)\dot{\eta}$  along  $\gamma$ , where the dot denotes the derivative with respect to the affine parameter  $\lambda$ . We can make  $C = 1$  by an affine transformation and then we find  $d\lambda = -a^2(\eta)dr$  through the relation  $k^a k_a = 0$ . The value of  $\lambda$  with which  $\gamma$  terminates at  $r = 2M$  is then given by

$$\lambda - \lambda_0 = -\int_{r_0^*}^{-\infty} a^2(u_{+(0)} - r^*)f(r)dr^*. \quad (5)$$

The interval of the integral on the right-hand side of Eq. (5) can be divided to  $(-\infty, -L]$  and  $[-L, r_0^*]$ . For sufficiently large but finite  $L$ , there exists  $\epsilon \in (0, 1/2)$  such that

$$(u_{+(0)} - r^*)^{2\alpha} f(r) < (2M)^{2\alpha} e^{(1-\epsilon)(r^*-2M)/(2M)} \quad (6)$$

for  $-r^* > L$ . Therefore, if  $a(\eta) = a_0\eta^\alpha$ , the integral over the first interval converges to a finite limit value, while that over the second is trivially finite. This means that the ingoing radial null geodesics are future incomplete at scalar polynomial curvature singularity at  $(r, \eta) = (2M, \infty)$ .




FIG. 1. Penrose diagrams of the Thakurta spacetime with  $a(\eta) \propto \eta^\alpha$  for  $0 < \alpha \leq 1$  (top) and  $\alpha > 1$  (bottom). Solid curves denote trapping horizons, which are past trapping horizons, and the shaded regions are past trapped. There is neither event horizon nor future outer trapping horizon in the spacetime.

As a result, the Penrose diagrams of the Thakurta spacetime (1) with  $a(\eta) \propto \eta^\alpha$  ( $\alpha > 0$ ) are given by Fig. 1. (See [20] for those of the corresponding FLRW spacetimes.) Clearly, there is no black-hole event horizon. Nevertheless, the spacetime might be interpreted as a black hole by an alternative definition in terms of a trapping horizon [21, 22] or an apparent horizon [19]. In fact, the Thakurta spacetime possesses a trapping horizon in the spherical foliation of the spacetime. However, this trapping horizon is not of black-hole type as we will see below.

To identify the location of a trapping horizon, it is useful to calculate the Misner-Sharp mass  $M_{\text{MS}}$  as [21–23]

$$M_{\text{MS}} := \frac{1}{2}S [1 - (\nabla_A S)(\nabla^A S)] = a \left( M + \frac{\mathcal{H}^2 r^3}{2f(r)} \right), \quad (7)$$

where  $S := ra$  is the areal radius and the suffix  $A$  takes  $A = 0, 1$ . We have put  $\mathcal{H} := a'/a$ , where the prime denotes the derivative with the argument  $\eta$ .

The location of a trapping horizon is determined by  $S = 2M_{\text{MS}}$ , or

$$f(r) = \mathcal{H}r, \quad (8)$$

where  $r > 2M$  is assumed. With  $\mathcal{H} = \alpha/\eta$ , Eq. (8) is solved for  $\eta$  to give

$$\eta = \frac{\alpha r^2}{r - 2M} =: \eta_{\text{TH}}(r). \quad (9)$$

The function  $\eta = \eta_{\text{TH}}(r)$  has a single local minimum  $\eta = 8\alpha M$  at  $r = 4M$ .

Among all the types of trapping horizons, it is a *future outer trapping horizon* that is associated with a black hole [21]. However, the Thakurta spacetime (1) with  $a(\eta) \propto \eta^\alpha$  ( $\alpha > 0$ ) admits only a *past* trapping horizon in the domain  $\eta > 0$ , which is associated with a white hole or a cosmological horizon. To see this, we introduce the outgoing and ingoing null expansions,  $\theta_\pm$ , defined by

$$\theta_\pm := \mathcal{A}^{-1} \partial_\pm \mathcal{A}, \quad (10)$$

where  $\partial_\pm := \partial/\partial u_\pm$  and  $\mathcal{A} := 4\pi S^2$ . Since Eq. (10) gives

$$\theta_\pm = \frac{1}{r} [\mathcal{H}r \pm f(r)], \quad (11)$$

$\theta_+ > 0$  and  $\theta_- = 0$  hold on the trapping horizon (8). For the further classification of a trapping horizon, we need to use

$$\partial_+ \theta_-|_{\theta_- = 0} = -\frac{f(r)[(1-\alpha)(r-2M) + 2\alpha M]}{2\alpha r^3}. \quad (12)$$

Depending on the signs of  $\theta_+|_{\theta_- = 0}$  and  $\partial_+ \theta_-|_{\theta_- = 0}$ , a trapping horizon is classified as in Table I.

TABLE I. Types of trapping horizons given by  $\theta_- = 0$ .

	+	-	0
$\theta_+ _{\theta_- = 0}$	Past	Future	Bifurcating
$\partial_+ \theta_- _{\theta_- = 0}$	Inner	Outer	Degenerate

Since  $\theta_+|_{\theta_- = 0} > 0$  holds, our trapping horizon (9) is a *past* trapping horizon. Then, by Eq. (12), it is an *outer* trapping horizon for  $0 < \alpha \leq 1$ . For  $\alpha > 1$ , it is an outer, degenerate, and inner trapping horizons in the regions of  $r - 2M < \beta_2$ ,  $r - 2M = \beta_2$ , and  $r - 2M > \beta_2$ , respectively, where

$$\beta_2 := \frac{2\alpha M}{\alpha - 1}. \quad (13)$$

In the region of  $0 < \eta < \eta_{\text{TH}}(r)$ , both  $\theta_+$  and  $\theta_-$  are positive, meaning that the region is *past* trapped. These results are summarized in Table II.

We also check the signature of the trapping horizon (9) as another important property. The line element on the hypersurface  $\eta = \eta_{\text{TH}}(r)$  is given by

TABLE II. Types of the trapping horizon in the Thakurta spacetime with  $a(\eta) \propto \eta^\alpha$  ( $\alpha > 0$ ) for the different ranges of  $r - 2M$ .

	Past outer	Past degenerate	Past inner
$0 < \alpha \leq 1$	$(0, \infty)$	$\emptyset$	$\emptyset$
$\alpha > 1$	$(0, \beta_2)$	$\beta_2$	$(\beta_2, \infty)$

$$ds_{\text{TH}}^2 = a^2(\eta_{\text{TH}}(r)) \left\{ \frac{[(1-\alpha)(r-2M) + 2\alpha M][(1+\alpha)(r-2M) - 2\alpha M]}{r^2 f^3(r)} dr^2 + r^2 d\Omega^2 \right\}. \quad (14)$$

Equation (14) clarifies the locations of timelike, null, and spacelike portions of the trapping horizon (9), which are summarized in Table III. We can see that the signature changes at  $r - 2M = \beta_1$  and  $\beta_2$ , where

$$\beta_1 := \frac{2\alpha M}{\alpha + 1}. \quad (15)$$

We notice that for  $\alpha = 1$ ,  $u_- \rightarrow -\infty$  and  $u_+ \rightarrow +\infty$  holds as  $r \rightarrow \infty$  along the trapping horizon  $\eta = \eta_{\text{TH}}(r)$ . This limit corresponds to spacelike infinity  $i^0$  in the Penrose diagram.

TABLE III. Signatures of the trapping horizon for the different ranges of  $r - 2M$ .

	Timelike	Null	Spacelike
$0 < \alpha \leq 1$	$(0, \beta_1)$	$\beta_1$	$(\beta_1, \infty)$
$1 < \alpha$	$(0, \beta_1), (\beta_2, \infty)$	$\beta_1, \beta_2$	$(\beta_1, \beta_2)$

Combining all the results obtained so far, we present  $r\eta$  planes for the Thakurta spacetime (1) in Fig. 2. The orbits of the trapping horizons are also presented in Fig. 1. We find that for all  $\alpha > 0$ , there exists a past outer trapping horizon located at  $0 < r - 2M < \beta_1$ , which is a timelike hypersurface. It is known that an outer (inner) trapping horizon is nontimelike (nonspacelike) under the null energy condition (NEC) in general relativity [21, 24]. Therefore, if we adopt the Thakurta spacetime (1) as a solution of the Einstein equation, the corresponding matter field violates the NEC on the trapping horizon located at  $0 < r - 2M < \beta_1$  for any  $\alpha > 0$ .

Here, we should emphasize that the Thakurta metric does not describe a cosmological black hole in spite of subtlety in defining cosmological black holes. Let us consider an observer O located outside a black-hole event horizon at finite distance and time, not at future null infinity nor timelike infinity. This is an observer usually preassumed in observational cosmology. By definition, such an observer never knows the existence of an event horizon (cf. [25]). Even though the causal nature of the  $r = 2M$  surface, such as whether it is finitely far or not and regular or not, crucially depends on the asymptotic




FIG. 2.  $r\eta$  planes for the Thakurta spacetime with  $a(\eta) \propto \eta^\alpha$  for (top)  $0 < \alpha \leq 1$  and (bottom)  $\alpha > 1$ . Several future light cones are put to clarify the signature of the trapping horizon  $\eta = \eta_{\text{TH}}(r)$ . The regions of  $0 < \eta < \eta_{\text{TH}}(r)$  with  $r > 2M$  are past trapped.

behavior of  $a(\eta)$  in the infinite future  $\eta \rightarrow \infty$ , it does not matter for O. What is even worse, if the universe is not asymptotically flat FLRW, it may not be possible to define future null infinity nor an event horizon. These facts suggest that, at least from an observational point of view, we should give up the definition of a cosmological black

hole in terms of an event horizon. Instead, for example, we may practically define a cosmological black hole by the existence of an “almost” future outer trapping horizon, a hypersurface foliated by “almost” marginal surfaces on which  $\theta_- < 0$ ,  $\theta_+$  is infinitesimally small positive, and  $\partial_- \theta_+ < 0$  are satisfied, that can be observed in principle by O. Even in such a rather broader definition, the Thakurta spacetime cannot be interpreted as a cosmological black hole because both  $\theta_+$  and  $\theta_-$  are always positive in the vicinity of  $r = 2M$  surface.

To conclude, the Thakurta metric (1) with  $a(\eta) \propto \eta^\alpha$  ( $\alpha > 0$ ) does not describe a cosmological black hole in the early universe, with a possible exceptional case, where the cosmological expansion is accelerated.

**Note added:** After uploading the present manuscript to the arXiv, the authors realized that the main content is included in [26], where the causal structures of the Thakurta metric are properly analysed in more general contexts. Nevertheless, they decided to keep this manuscript available on the arXiv because its spotlight on the case relevant to the discussion on the evolution of primordial black holes in the early universe may be useful in the community.

T. H. is grateful to N. Tanahashi for fruitful discussion. This work was partially supported by MEXT Grants-in-Aid for Scientific Research/JSPSKAKENHI Grants Nos. JP19K03876, JP19H01895, and 20H05853 (T.H.).

---

\* harada@rikkyo.ac.jp

† h-maeda@hgu.jp

‡ stakuma@rikkyo.ac.jp

- [1] B. J. Carr and S. W. Hawking, *Mon. Not. Roy. Astron. Soc.* **168** (1974), 399-415.
- [2] B. J. Carr, *Astrophys. J.* **201**, 1-19 (1975) doi:10.1086/153853
- [3] M. Sasaki, T. Suyama, T. Tanaka and S. Yokoyama, *Phys. Rev. Lett.* **117** (2016) no.6, 061101 [erratum: *Phys. Rev. Lett.* **121** (2018) no.5, 059901] doi:10.1103/PhysRevLett.117.061101 [arXiv:1603.08338 [astro-ph.CO]].
- [4] S. Clesse, and J. García-Bellido, *Phys. Dark Univ.* **15**, 142 (2017) doi:10.1016/j.dark.2016.10.002 [arXiv:1603.05234 [astro-ph.CO]].
- [5] S. Bird, I. Cholis, J. B. Muñoz, Y. Ali-Haïmoud, M. Kamionkowski, E. D. Kovetz, A. Raccanelli, and A. G. Riess, *Phys. Rev. Lett.* **116**, no. 20, 201301 (2016) doi:10.1103/PhysRevLett.116.201301 [arXiv:1603.00464 [astro-ph.CO]].
- [6] G. Franciolini, V. Baibhav, V. De Luca, K. K. Y. Ng, K. W. K. Wong, E. Berti, P. Pani, A. Riotto and S. Vitale, [arXiv:2105.03349 [gr-qc]].
- [7] C. Boehm, A. Kobakhidze, C. A. J. O’hare, Z. S. C. Picker and M. Sakellariadou, *JCAP* **03** (2021), 078 doi:10.1088/1475-7516/2021/03/078 [arXiv:2008.10743 [astro-ph.CO]].
- [8] Z. S. C. Picker, [arXiv:2103.02815 [astro-ph.CO]].
- [9] S. N. G. Thakurta, *Indian Journal of Physics* **55B**, 304 (1981).
- [10] V. Faraoni, *Universe* **4** (2018) no.10, 109 doi:10.3390/universe4100109 [arXiv:1810.04667 [gr-qc]].
- [11] G. Hütsi, T. Koivisto, M. Raidal, V. Vaskonen and H. Veermäe, [arXiv:2105.09328 [astro-ph.CO]].
- [12] C. Boehm, A. Kobakhidze, C. A. J. O’Hare, Z. S. C. Picker and M. Sakellariadou, [arXiv:2105.14908 [astro-ph.CO]].
- [13] G. Hütsi, T. Koivisto, M. Raidal, V. Vaskonen and H. Veermäe, [arXiv:2106.02007 [astro-ph.CO]].
- [14] G. C. McVittie, *Mon. Not. Roy. Astron. Soc.* **93** (1933), 325-339 doi:10.1093/mnras/93.5.325
- [15] B. C. Nolan, *Phys. Rev. D* **58** (1998), 064006 doi:10.1103/PhysRevD.58.064006 [arXiv:gr-qc/9805041 [gr-qc]].
- [16] B. C. Nolan, *Class. Quant. Grav.* **16** (1999), 1227-1254 doi:10.1088/0264-9381/16/4/012
- [17] B. C. Nolan, *Class. Quant. Grav.* **16** (1999), 3183-3191 doi:10.1088/0264-9381/16/10/310 [arXiv:gr-qc/9907018 [gr-qc]].
- [18] N. Kaloper, M. Kleban and D. Martin, *Phys. Rev. D* **81** (2010), 104044 doi:10.1103/PhysRevD.81.104044 [arXiv:1003.4777 [hep-th]].
- [19] R. M. Wald, “*General Relativity*” (Chicago: University of Chicago Press, 1983).
- [20] T. Harada, B. J. Carr and T. Igata, *Class. Quant. Grav.* **35** (2018) no.10, 105011 doi:10.1088/1361-6382/aab99f [arXiv:1801.01966 [gr-qc]].
- [21] S. A. Hayward, *Phys. Rev. D* **49** (1994), 6467-6474 doi:10.1103/PhysRevD.49.6467
- [22] S. A. Hayward, *Phys. Rev. D* **53** (1996) 1938 doi:10.1103/PhysRevD.53.1938 [gr-qc/9408002].
- [23] C. W. Misner and D. H. Sharp, *Phys. Rev.* **136** (1964), B571-B576 doi:10.1103/PhysRev.136.B571
- [24] M. Nozawa and H. Maeda, *Class. Quant. Grav.* **25** (2008), 055009 doi:10.1088/0264-9381/25/5/055009 [arXiv:0710.2709 [gr-qc]].
- [25] K. i. Nakao, C. M. Yoo and T. Harada, *Phys. Rev. D* **99** (2019) no.4, 044027 doi:10.1103/PhysRevD.99.044027 [arXiv:1809.00124 [gr-qc]].
- [26] M. M. C. Mello, A. Maciel and V. T. Zanchin, *Phys. Rev. D* **95** (2017) no.8, 084031 doi:10.1103/PhysRevD.95.084031 [arXiv:1611.05077 [gr-qc]].