

Gauge gravity vacuum in constraintless Clairaut-type formalism

Michael L. Walker

University of New South Wales, Kensington, NSW 2033, Australia
and

Steven Duplij

Center for Information Technology (WWU IT), University of Münster,
Röntgenstrasse 7-13, D-48149 Münster, Germany

Ver. 3: August 4, 2021; Ver. 2: July 25, 2021; Ver. 1: June 14, 2021

Abstract

A previously published covariant decomposition of the Levi-Civita tensor has demonstrated the strong mathematical parallel between gravity and $N = 2$ Yang-Mills theory. We use this to argue that the an R^2 type Lagrangian has a stable vacuum of lower energy than the perturbative vacuum. Using a Lagrangian in this form instead of the more conventional Hilbert-Einstein Lagrangian allows us to use the Clairaut-based formalism published by one of us previously to handle gravity's topological degrees of freedom. We study the effect on second quantisation and discuss the effects on, and of, the particle spectrum.

Contents

1	Introduction	3
2	A review of the covariant Abelian decomposition of gravity	4
2.1	Magnetic decomposition	4
2.2	Maximum Abelian decomposition	6
3	Application of Clairaut formalism to the Rotation-Boost decomposition of the gravitational connection	9
3.1	A review of the Hamiltonian-Clairaut formalism	9
3.2	The contribution of the Clairaut formalism	11
3.2.1	q^α curvature	11
3.2.2	Corrections to the equations of motion	12
3.2.3	Corrections to the commutation relations	13
4	Matter fields	16
5	Particle number and the Lorentz-monopole background	16
6	Discussion	18
A	The CDG decomposition in $SU(2)$ QCD	19
B	Some convenient identities	21
	References	21

1 Introduction

It is generally accepted that spacetime respects the Poincaré symmetry group, comprising both translation symmetry and Lorentz symmetry, so that attempts to construct a theory of gravity based on gauge symmetry are typically based on a localised Poincaré symmetry. While such a theory has been shown to be renormalisable [1] the approach is never-the-less problematic, for if the Lagrangian is taken to be quadratic in the gravitational curvature then third derivatives of the translation representation (vierbein) destabilise the theory with negative kinetic energy terms [2]. Alternately, a linear form may be taken such as the commonly used Hilbert-Einstein (H-E) Lagrangian.

A novel approach to stabilising the quadratic form was developed by Pak *et al.* [3, 4, 5], in a series of papers demonstrating that the energy could be bounded from below by quantum vacuum effects induced by a quantised contorsion. In this and similar models, typically based on Einstein-Cartan gravity, it is contorsion (or torsion) which is the quantised dynamic degree of freedom (DOF) while gravitational curvature manifests as a topological effect.

The topological contribution to Yang-Mills theories has also attracted considerable attention, especially in the context of confinement in QCD. Of particular importance is the notion of Abelian dominance, in which an Abelian subgroup of the gauge group is either seen or assumed to dominate at longer wavelengths. This reduction to a subgroup generates a chromo-monopole condensate leading to a dual Meissner effect which confines colour charge. Such magnetic confinement models date back to the work of Savvidy [6] and Nielsen and Olesen [7], but suffered the shortcoming of not being gauge covariant since specifying an Abelian, or any subgroup, breaks the gauge symmetry if care is not taken. In fact a gauge covariant way of identifying a subgroup of a gauge Yang-Mills theory, which automatically provides a potential for the corresponding chromo-monopole field, was developed by Cho [8] and also independently by Duan and Ge [9]. This Cho-Duan-Ge (CDG) decomposition, and another like it [10], was (re-)discovered [11] at about the turn of the century when several groups were readdressing the stability of the chromo-monopole condensate [12, 13, 14, 15, 16, 17, 18, 19]. The mathematical subtleties of the CDG decomposition were only understood after considerable effort by several independent groups, with some initial confusion regarding its DOFs and gauge-fixing. The interested reader is referred to [11, 12, 13, 14, 20] for further details (see, also [21, 22]).

The CDG decomposition has since been used to demonstrate other important results about QCD with three or more colours [18, 19, 23], the Wilson loop and confinement [24, 25, 26], and even the vacuum of supersymmetric theories [27]. More recently, Cho *et al.* adapted it to a Lorentz gauge group theory of gravity to show that its dynamics have a spin-1 component [28, 29] in the absence of matter fields. They also found the vacuum to have an $SU(2)$ monopole background of the Lorentz gauge field, although they did use the conventional H-E Lagrangian and did not discuss vacuum stability.

In this work we apply this decomposition to the vacuum of a Lorentz gauge theory. Throughout this paper we take "monopole" to be in reference to the Lorentz gauge field unless "chromo-" is specified to indicate QCD.

Previous analyses using the CDG decomposition in QCD [12] assumed the chromo-monopole condensate comprising the vacuum to provide a slow-moving vacuum background to the quantum DOFs, but in an earlier paper [30] the authors applied a new formalism to rigorously elucidate the dynamic DOFs from the topological. The Clairaut-type formulation, proposed by one of the authors (SD) [31, 32], is a constraintless generalization of the standard Hamiltonian formalism to include Hessians with zero determinant. It provides a rigorous treatment of the non-physical DOFs in the derivation of EOMs and the quantum commutation relations. Unfortunately this approach is not well-suited to the H-E Lagrangian, failing to find natural looking forms for the conjugate momenta. It has also been argued by Witten [33] that the H-E Lagrangian cannot represent a gauge theory in 3+1 dimensions. We therefore apply our Clairaut approach to the gauged Lorentz group [28, 29] theory with a Lagrangian quadratic in curvature. Rather than include contorsion, we observe the strong mathematical similarities with an $SU(2)$ Yang-Mills theory and use them to adapt results concerning the stability of that vacuum to this theory. To maintain this parallel, and to avoid third order derivatives of entering the EOMs, our theory does not include localised translation symmetry. We do not, therefore, consider it a full theory of gravity, but rather a toy model for providing insight.

A review of the covariant decomposition of the Lorentz group [28, 29] is given in Section 2. In Section 3 we give a brief overview of the Clairaut-Hamiltonian formalism and use it to study the quantisation of this theory, sorting dynamic from topological DOFs and identifying circumstances under which the particle interpretation is spoiled by background curvature. Matter fields are discussed in Section 4, while Section 5 gives an overview of the effective particle spectrum, including specific circumstances when the particle violating terms appear to vanish. We finish with a discussion in Section 6

2 A review of the covariant Abelian decomposition of gravity

2.1 Magnetic decomposition

Abelian dominance has proven to be vital in our understanding of the QCD vacuum, facilitating the demonstration of a chromo-monopole condensate in QCD. That a magnetic condensate suitable for colour confinement can have lower energy than the perturbative vacuum has been known since the 1970s [6, 7], but in early work the direction in internal space supporting the magnetic background could not be specified in a covariant manner and nor was there support for the magnetic condensate being due to chromo-monopoles.

These issues were rectified by the introduction of the CDG decomposition, whose application to QCD is described in technical detail in Appendix A. Cho *et al.* [28, 29] applied the CDG decomposition to a Lorentz gauge theory of gravity, where the connection forms the Lorentz gauge field. In fact he found two decompositions, one for each of the two isometries of Lorentz symmetry which he had characterised in an earlier paper [34], using its equivalence to a local gauge theory of Lorentz transformations. One, labelled *A2*, has a rotation-boost (non-lightlike) isometry and is suitable for our approach. The other, labelled *B2*, has a null (lightlike) isometry corresponding to light-front coordinates which we do not know how to treat in the Clairaut formalism. We shall therefore work exclusively with the *A2* formalism in this paper. Below we describe it just enough to establish our notation, and then apply the above approach from QCD to express the DOFs most succinctly.

The key physical insight is that the rotations are magnetically dual to the boosts along the corresponding axis in the same sense that $\epsilon_{\mu\nu\rho\lambda}F^{\rho\lambda}$ is dual to $F_{\mu\nu}$ [28, 29]. This duality has no direct equivalent in QCD. Equating the indices (1, 2, 3, 4, 5, 6) \equiv (23, 31, 12, 01, 02, 03), an adjoint representation of the Lorentz group is given by sextets of the form ¹

$$\mathbf{P} = \begin{pmatrix} \vec{m} \\ \vec{e} \end{pmatrix}, \quad (2.1)$$

where \vec{m}, \vec{e} are multiples of the rotation and boost generators \hat{e}, \hat{m} , respectively, where

$$\begin{aligned} \hat{m}_i^{ab} &= \epsilon_{0i}^{ab} \\ \hat{e}_i^{ab} &= \delta_0^a \delta_i^b - \delta_0^b \delta_i^a \end{aligned} \quad (2.2)$$

and the magnetic dual to \mathbf{P} is given by

$$\tilde{\mathbf{P}} = \begin{pmatrix} \vec{e} \\ -\vec{m} \end{pmatrix}. \quad (2.3)$$

where

$$\mathbf{P} \cdot \tilde{\mathbf{P}} = 2\vec{m} \cdot \vec{e}, \quad \mathbf{P} \times \tilde{\mathbf{P}} = 0. \quad (2.4)$$

\mathbf{P} 's components obey the relation

$$P^{ab} = -\frac{1}{2}P^{mn}(J_{mn})^{ab}. \quad (2.5)$$

The same indices define the invariant metric of the Lorentz group, given by

$$\begin{aligned} \delta_{AB} &= -f_{AC}^D f_{BD}^C \\ &= \text{diag}(+1, +1, +1, -1, -1, -1), \end{aligned} \quad (2.6)$$

¹We have replaced \mathbf{p} in Cho *et al.* with \mathbf{P} to avoid notation conflict later in the paper.

where f are the Lorentz group coefficients. The reader is referred to [28, 29] for complete details, many of which are not needed here as our toy model excludes local translation symmetry and does not treat the vierbein.

This notation allows $\mathbf{P}, \tilde{\mathbf{P}}$ to be naturally expressed with the basis

$$\mathbf{l}_i = \begin{pmatrix} \hat{e}_i \\ 0 \end{pmatrix}, \quad \mathbf{k}_i = \begin{pmatrix} 0 \\ \hat{e}_i \end{pmatrix} = -\tilde{\mathbf{l}}_i, \quad (2.7)$$

with the algebra

$$\mathbf{l}_i \cdot \mathbf{l}_j = \delta_{ij}, \quad \mathbf{l}_i \cdot \mathbf{k}_j = 0, \quad \mathbf{k}_i \cdot \mathbf{k}_j = -\delta_{ij}, \quad (2.8)$$

$$\mathbf{l}_i \times \mathbf{l}_j = \epsilon_{ijk} \mathbf{l}_k, \quad \mathbf{l}_i \times \mathbf{k}_j = \epsilon_{ijk} \mathbf{k}_k, \quad \mathbf{k}_i \times \mathbf{k}_j = -\epsilon_{ijk} \mathbf{l}_k. \quad (2.9)$$

2.2 Maximum Abelian decomposition

We use the CDG decomposition in which the embedding of a dominant direction $U(1)$ is denoted by \hat{n} which is defined by,

$$\hat{n}(x) \equiv \cos \theta(x) \sin \phi(x) \hat{e}_1 + \sin \theta(x) \sin \phi(x) \hat{e}_2 + \cos \phi(x) \hat{e}_3. \quad (2.10)$$

The following will prove useful:

$$\begin{aligned} \sin \phi(x) \hat{n}_\theta(x) &\equiv \int dy^4 \frac{d\hat{n}(x)}{d\theta(y)} = \sin \phi(x) (-\sin \theta(x) \hat{e}_1 + \cos \theta(x) \hat{e}_2), \\ \hat{n}_\phi(x) &\equiv \int dy^4 \frac{d\hat{n}(x)}{d\phi(y)} = \cos \theta(x) \cos \phi(x) \hat{e}_1 + \sin \theta(x) \cos \phi(x) \hat{e}_2 - \sin \phi(x) \hat{e}_3, \end{aligned} \quad (2.11)$$

for later convenience, we note that

$$\begin{aligned} \hat{n}_{\phi\phi} &= -\hat{n}, \quad \hat{n}_{\theta\theta} = -\sin \phi \hat{n} - \cos \phi \hat{n}_\phi, \\ \hat{n}_{\theta\phi} &= 0, \quad \hat{n}_{\phi\theta} = \cos \phi \hat{n}_\theta, \end{aligned} \quad (2.12)$$

and that the vectors $\hat{n}(x) = \hat{n}_\phi(x) \times \hat{n}_\theta(x)$ form an orthonormal basis of the internal space. Other useful identities are shown in Appendix B.

Using this decomposition, Cho *et al.* [28, 29] identify the covariant basis

$$\hat{\mathbf{l}} = \begin{pmatrix} \hat{n} \\ 0 \end{pmatrix}, \quad \tilde{\mathbf{l}} = \begin{pmatrix} 0 \\ -\hat{n} \end{pmatrix}. \quad (2.13)$$

We shall later have use for analogous definitions corresponding to $\hat{n}_\theta, \hat{n}_\phi, \hat{n}_{\phi\theta}, \hat{n}_{\phi\phi}, \hat{n}_{\theta\theta}$. As with $SU(2)$ Yang-Mills we have nominated an Abelian direction at each point in spacetime with the three-dimensional unit vector \hat{n} , finding it convenient to again use the derived unit vectors $\hat{n}_\theta, \hat{n}_\phi$ in subsequent calculations.

The most general connection which leaves the Abelian directions $\hat{\mathbf{I}}, \tilde{\mathbf{I}}$ covariant is given by

$$\begin{aligned}\hat{\Gamma}_\mu &\equiv c_\mu \hat{\mathbf{I}} - \tilde{c}_\mu \tilde{\mathbf{I}} + g^{-1} \partial_\mu \hat{\mathbf{I}} \times \hat{\mathbf{I}} \\ &= \mathbf{c}_\mu + \mathbf{C}_\mu,\end{aligned}\quad (2.14)$$

where

$$\mathbf{c}_\mu \equiv \begin{pmatrix} c_\mu \hat{n} \\ \tilde{c}_\mu \hat{n} \end{pmatrix}, \quad \mathbf{C}_\mu \equiv g^{-1} (\partial_\mu \hat{\mathbf{I}} \times \hat{\mathbf{I}}) = \begin{pmatrix} g^{-1} \partial_\mu \hat{n} \times \hat{n} \\ 0 \end{pmatrix}.$$
 (2.15)

We shall later have use for

$$\mathbf{H}_{\mu\nu} \equiv \partial_\mu \mathbf{C}_\nu - \partial_\nu \mathbf{C}_\mu + \mathbf{C}_\mu \times \mathbf{C}_\nu \equiv H_{\mu\nu} \hat{\mathbf{I}},$$
 (2.16)

and

$$\begin{aligned}\mathbf{F}_{\mu\nu} &\equiv (\partial_\mu c_\nu - \partial_\nu c_\mu) \mathbf{I} - (\partial_\mu \tilde{c}_\nu - \partial_\nu \tilde{c}_\mu) \tilde{\mathbf{I}} + \mathbf{H}_{\mu\nu} + g \mathbf{X}_\mu \times \mathbf{X}_\nu \\ &= (\hat{\mathbf{D}}_\mu \mathbf{c}_\nu - \hat{\mathbf{D}}_\nu \mathbf{c}_\mu) + \mathbf{H}_{\mu\nu} + g \mathbf{X}_\mu \times \mathbf{X}_\nu\end{aligned}\quad (2.17)$$

There are, similarly, two mutually dual valence components, $\vec{Z}_\mu, \tilde{Z}_\mu$, of the Lorentz connection which are always perpendicular to \hat{n} . The parallels with CDG Yang-Mills become even more striking if we absorb them into the above dualistic notation,

$$\mathbf{X}_\mu \equiv \begin{pmatrix} \vec{Z}_\mu \\ \tilde{Z}_\mu \end{pmatrix}.$$
 (2.18)

Substituting into the full connection coefficient we have

$$\Gamma_\mu \equiv \hat{\Gamma}_\mu + \mathbf{X}_\mu,$$
 (2.19)

which we use to define the covariant derivatives

$$\mathbf{D}_\mu \equiv \partial_\mu + g \Gamma_\mu \times,$$
 (2.20)

$$\hat{\mathbf{D}}_\mu \equiv \partial_\mu + g \hat{\Gamma}_\mu \times.$$
 (2.21)

The curvature is then given by

$$\mathbf{R}_{\mu\nu} = \frac{1}{2} \left(\hat{\mathbf{D}}_\mu \Gamma_\nu - \hat{\mathbf{D}}_\nu \Gamma_\mu + g \Gamma_\mu \times \Gamma_\nu \right).$$
 (2.22)

Hence we get that curvature has a topological non-dynamic part, and that this theory is formally very similar to two-colour Yang-Mills theory. Indeed, taking and substituting in the eqs (2.14-2.19) yields the Lagrangian

$$\mathcal{L} = -\frac{1}{4} \mathbf{R}_{\mu\nu} \mathbf{R}^{\mu\nu} = -\frac{1}{4} \mathbf{F}^{\mu\nu} \cdot \mathbf{F}_{\mu\nu} - \frac{1}{4} \left(\hat{\mathbf{D}}^\mu \mathbf{X}^\nu - \hat{\mathbf{D}}^\nu \mathbf{X}^\mu \right) \cdot \left(\hat{\mathbf{D}}_\mu \mathbf{X}_\nu - \hat{\mathbf{D}}_\nu \mathbf{X}_\mu \right),$$
 (2.23)

whose similarity to the decomposed $N = 2$ Yang-Mills Lagrangian [12, 13, 14, 15, 16, 20] is unmistakable.

The formal similarity to $SU(2)$ Yang-Mills theory extends to the gauge variation of the different components of the decomposition. We shall consider the rotation and boost transformations separately for the sake of clarity. Infinitesimal rotations are effected with a gauge parameter $\hat{\alpha} = \begin{pmatrix} \vec{\alpha} \\ 0 \end{pmatrix}$, where the $\hat{\ }$ notation is chosen in parallel with $\hat{\Gamma}$ and is not intended to imply a unit length in the usual sense. Under a gauged rotation transformation δ_r with $SU(2)$ parameter $\hat{\alpha}$

$$\begin{aligned}\delta_r \hat{\Gamma} &= \hat{\mathbf{D}}_\mu \hat{\alpha} \\ \delta_r \hat{n} &= \hat{n} \times \vec{\alpha}, \\ \delta_r \vec{Z}_\mu &= g \vec{Z}_\mu \times \vec{\alpha}, \\ \delta_r \tilde{Z}_\mu &= g \tilde{Z}_\mu \times \vec{\alpha}.\end{aligned}\tag{2.24}$$

These expressions describing the gauge transformation tell us two interesting things. The first is that the Abelian component \mathbf{c}_μ combined with the Cho connection \mathbf{C}_μ are enough to represent the full Lorentz symmetry even without the valence components \mathbf{X}_μ , Cho *et al.* [28, 29] described as the "restricted" theory. The second is that the valence components transform like a source transforms. There is a corresponding situation in $N = 2$ Yang-Mills where the valence gluons are interpreted as colour sources. The importance of this observation for this work is that we shall later discuss the possibility of mass generation for the valence gluons and this form for the gauge transformation leaves such mass terms covariant.

Infinitesimal boosts are effected by gauge parameters of the form $\tilde{\alpha} = \begin{pmatrix} 0 \\ \vec{\alpha} \end{pmatrix}$. Under a gauged boost transformation δ_b with $SU(2)$ parameter $\tilde{\alpha}$ the different components of the field transform according to

$$\begin{aligned}\delta_b \hat{\Gamma} &= \hat{\mathbf{D}}_\mu \tilde{\alpha} \\ \delta_b \hat{n} &= g \hat{n} \times \vec{\alpha}, \\ \delta_b \vec{Z}_\mu &= g \vec{Z}_\mu \times \vec{\alpha}, \\ \delta_b \tilde{Z}_\mu &= g \tilde{Z}_\mu \times \vec{\alpha}.\end{aligned}\tag{2.25}$$

The valence components $\vec{Z}_\mu, \tilde{Z}_\mu$, and by extension \mathbf{X}_μ , do not transform as gauge fields but as sources. Nonetheless their form must be constrained to ensure consistency with its being a massless gauge vector field [35], so we impose the gauge condition

$$\hat{\mathbf{D}}_\mu \mathbf{X}_\mu = 0,\tag{2.26}$$

whose similarity to a conventional gauge-fixing term is clear. Gauge-fixing the Abelian DOFs c_μ we begin with the condition

$$\begin{aligned}\hat{\mathbf{D}}_\mu \mathbf{c}_\mu &= 0 \\ \Rightarrow \hat{\mathbf{D}}_\mu(c_\mu \hat{\mathbf{1}}) &= 0, \quad \hat{\mathbf{D}}_\mu(\tilde{c}_\mu \tilde{\mathbf{1}}) = 0 \\ \Rightarrow \partial_\mu c_\mu &= 0, \quad \partial_\mu \tilde{c}_\mu = 0\end{aligned}\tag{2.27}$$

Hence the Abelian DOFs c_μ require only a straightforward Abelian gauge-fixing term and no ghosts.

The chromo-monopole background in $SU(2)$ QCD has long been argued to be energetically favourable to the perturbative vacuum [6, 7, 36] but demonstrating its stability took work by many authors over time [11, 12, 13, 14, 15, 16, 20]. The point of contention was the apparent existence of an imaginary part to the one-loop effective action, which was found to be an artifact of subtleties of regularisation [15, 16]. We observe that the form of the Lagrangian in eq. (2.23) is the same as that for $SU(2)$ QCD with the original CDG decomposition, as indicated by eqs. (1.9, 1.10) in Appendix A.

Working through the same calculations as in the analysis of $SU(2)$ Yang-Mills theory [15, 16], finds two copies of each of the contributing Feynman diagrams calculated by one of the authors (MLW) in conjunction with Cho [15] and also Pak [16], the other two being quadratically divergent and therefore not contributing after renormalisation. We find therefore that the analysis in these two papers holds in this effective theory yielding a stable vacuum condensate at one loop.

3 Application of Clairaut formalism to the Rotation-Boost decomposition of the gravitational connection

3.1 A review of the Hamiltonian-Clairaut formalism

Here we review the main ideas and formulae of the Clairaut-type formalism for singular theories [31, 37]. Let us consider a singular Lagrangian $L(q^A, v^A) = L^{\text{deg}}(q^A, v^A)$, $A = 1, \dots, n$, which is a function of $2n$ variables (n generalized coordinates q^A and n velocities $v^A = \dot{q}^A = dq^A/dt$) on the configuration space TM , where M is a smooth manifold, for which the Hessian's determinant is zero. Therefore, the rank of the Hessian matrix $W_{AB} = \frac{\partial^2 L(q^A, v^A)}{\partial v^B \partial v^C}$ is $r < n$, and we suppose that r is constant. We can rearrange the indices of W_{AB} in such a way that a nonsingular minor of rank r appears in the upper left corner. Then, we represent the index A as follows: if $A = 1, \dots, r$, we replace A with i (the ‘‘regular’’ index), and, if $A = r + 1, \dots, n$ we replace A with α (the ‘‘degenerate’’ index). Obviously, $\det W_{ij} \neq 0$, and $\text{rank } W_{ij} = r$. Thus any set of variables labelled by a single index splits as a disjoint union of two subsets. We call

those subsets regular (having Latin indices) and degenerate (having Greek indices). As was shown in [31, 37], the “physical” Hamiltonian can be presented in the form

$$H_{phys}(q^A, p_i) = \sum_{i=1}^r p_i V^i(q^A, p_i, v^\alpha) + \sum_{\alpha=r+1}^n B_\alpha(q^A, p_i) v^\alpha - L(q^A, V^i(q^A, p_i, v^\alpha), v^\alpha), \quad (3.1)$$

where the functions

$$B_\alpha(q^A, p_i) \stackrel{def}{=} \left. \frac{\partial L(q^A, v^A)}{\partial v^\alpha} \right|_{v^i=V^i(q^A, p_i, v^\alpha)} \quad (3.2)$$

are independent of the unresolved velocities v^α since $\text{rank } W_{AB} = r$. Also, the r.h.s. of (3.1) does not depend on the degenerate velocities v^α

$$\frac{\partial H_{phys}}{\partial v^\alpha} = 0, \quad (3.3)$$

which justifies the term “physical”. The Hamilton-Clairaut system which describes any singular Lagrangian classical system (satisfying the second order Lagrange equations) has the form

$$\frac{dq^i}{dt} = \{q^i, H_{phys}\}_{phys} - \sum_{\beta=r+1}^n \{q^i, B_\beta\}_{phys} \frac{dq^\beta}{dt}, \quad i = 1, \dots, r \quad (3.4)$$

$$\frac{dp_i}{dt} = \{p_i, H_{phys}\}_{phys} - \sum_{\beta=r+1}^n \{p_i, B_\beta\}_{phys} \frac{dq^\beta}{dt}, \quad i = 1, \dots, r \quad (3.5)$$

$$\begin{aligned} & \sum_{\beta=r+1}^n \left[\frac{\partial B_\beta}{\partial q^\alpha} - \frac{\partial B_\alpha}{\partial q^\beta} + \{B_\alpha, B_\beta\}_{phys} \right] \frac{dq^\beta}{dt} \\ &= \frac{\partial H_{phys}}{\partial q^\alpha} + \{B_\alpha, H_{phys}\}_{phys}, \quad \alpha = r+1, \dots, n \end{aligned} \quad (3.6)$$

where the “physical” Poisson bracket (in regular variables q^i, p_i) is

$$\{X, Y\}_{phys} = \sum_{i=1}^{n-r} \left(\frac{\partial X}{\partial q^i} \frac{\partial Y}{\partial p_i} - \frac{\partial Y}{\partial q^i} \frac{\partial X}{\partial p_i} \right). \quad (3.7)$$

Whether the variables $B_\alpha(q^A, p_i)$ have a nontrivial effect on the time evolution and commutation relations is equivalent to whether or not the so-called “ q^α -field strength”

$$\mathcal{F}_{\alpha\beta} = \frac{\partial B_\beta}{\partial q^\alpha} - \frac{\partial B_\alpha}{\partial q^\beta} + \{B_\alpha, B_\beta\}_{phys} \quad (3.8)$$

is non-zero. See [31, 32, 37] for more details.

3.2 The contribution of the Clairaut formalism

3.2.1 q^α curvature

Substituting in the notation of Section 2, the angles ϕ, θ are seen, in parallel with our previously published analysis [30], to have unresolved velocities. Their Clairaut-related quantities are

$$\begin{aligned}
B_\phi(x) &= \int dy^3 \frac{\delta \mathcal{L}}{\delta_x \partial_0 \phi(x)} \\
&= \int dy^3 \int dy^0 \delta(x^0 - y^0) \\
&\quad \times \left(\sin \phi(y) {}_y \partial^\mu \theta(y) \hat{\mathbf{l}}(y) + (\hat{\mathbf{l}}_\theta(y) \times \mathbf{X}^\mu)(y) \right) \cdot \mathbf{R}_{0\mu}(y) \delta^3(\vec{x} - \vec{y}) \\
&= \left(\sin \phi(x) \partial^\mu \theta(x) \hat{\mathbf{l}}(x) + (\hat{\mathbf{l}}_\theta(x) \times \mathbf{X}^\mu)(x) \right) \cdot \mathbf{R}_{0\mu}(x), \tag{3.9}
\end{aligned}$$

$$\begin{aligned}
B_\theta(x) &= \int dy^3 \frac{\delta \mathcal{L}}{\delta_x \partial_0 \theta(x)} \\
&= - \int dy^3 \int dy^0 \delta(x^0 - y^0) \sin \phi(y) \\
&\quad \times \left({}_y \partial^\mu \phi(y) \hat{\mathbf{l}}(y) + \sin \phi(y) (\hat{\mathbf{l}}_\phi(y) \times \mathbf{X}^\mu)(y) \right) \cdot \mathbf{R}_{0\mu}(y) \delta^3(\vec{x} - \vec{y}) \\
&= - \sin \phi(x) \left(\partial^\mu \phi(x) \hat{\mathbf{l}}(x) + (\hat{\mathbf{l}}_\phi(x) \times \mathbf{X}^\mu)(x) \right) \cdot \mathbf{R}_{0\mu}(x). \tag{3.10}
\end{aligned}$$

$$\frac{\delta B_\phi(x)}{\delta \theta(y)} = \left(\sin \phi(x) (\hat{\mathbf{l}}_{\theta\theta}(x) \times \mathbf{X}^\mu(x)) \cdot \mathbf{R}_{0\mu}(x) - T_\phi(x) \right) \delta^4(x - y), \tag{3.11}$$

$$\begin{aligned}
\frac{\delta B_\theta(x)}{\delta \phi(y)} &= - \left(\cos \phi(x) \left(\partial^\mu \phi(x) \hat{\mathbf{l}}(x) + (\hat{\mathbf{l}}_\phi(x) \times \mathbf{X}^\mu(x)) \right) \cdot \left(\mathbf{R}_{0\mu}(x) + \mathbf{H}_{0\mu}(x) \right) \right. \\
&\quad \left. + T_\theta(x) \right) \delta^4(x - y), \tag{3.12}
\end{aligned}$$

where now

$$\begin{aligned}
T_\phi(x) &= \partial^k \left[\sin \phi(x) \hat{\mathbf{l}}(x) \cdot \mathbf{R}_{0k}(x) \right. \\
&\quad \left. - \left(\sin \phi(x) \partial_k \theta(x) + (\hat{\mathbf{l}}_\theta(x) \times \mathbf{X}_k(x)) \cdot \mathbf{l}(x) \right) \partial_0 \phi(x) \right], \tag{3.13}
\end{aligned}$$

$$\begin{aligned}
T_\theta(x) &= -\partial^k \left[\sin \phi(x) \left(\hat{\mathbf{l}}(x) \cdot \mathbf{R}_{0k}(x) \right) \right. \\
&\quad \left. + \left(\partial_k \phi(x) + (\hat{\mathbf{l}}_\phi(x) \times \mathbf{X}_k(x)) \cdot \hat{\mathbf{l}}(x) \right) \partial_0 \theta(x) \right]. \tag{3.14}
\end{aligned}$$

This resulting q^α -curvature is given by equation (26) in [30], but simplifies further to

$$\begin{aligned}
\mathcal{F}_{\theta\phi}(x) &= \int dy^4 \left(\frac{\delta B_\theta(x)}{\delta\phi(y)} - \frac{\delta B_\phi(x)}{\delta\theta(y)} \right) \delta^4(x-y) + \{B_\phi(x), B_\theta(x)\}_{phys} \\
&= -\cos\phi(x) \left(\partial^\mu\phi(x) \hat{\mathbf{1}}(x) + \left(\hat{\mathbf{1}}_\phi(x) \times \mathbf{X}^\mu(x) \right) \right) \cdot \left(\mathbf{R}_{0\mu}(x) + \mathbf{H}_{0\mu}(x) \right) \\
&\quad - \sin\phi(x) \left(\hat{\mathbf{1}}_{\theta\theta}(x) \times \mathbf{X}^\mu(x) \right) \cdot \mathbf{R}_{\mu 0}(x) \\
&\quad + 2\partial^\mu \left(\sin\phi(x) \mathbf{R}_{0\mu}(x) \cdot \hat{\mathbf{1}}(x) + (\partial^\mu \mathbf{H}_{0\mu}(x) + g(\mathbf{C}_0(x) \times \mathbf{X}_\mu(x))) \cdot \mathbf{1}(x) \right).
\end{aligned} \tag{3.15}$$

3.2.2 Corrections to the equations of motion

Generalizing eqs. (7.1,7.3,7.5) in [31] (see also the discussion around eq. (3.1)),

$$\partial_0 \mathbf{q}(x) = \{ \mathbf{q}(x), H_{phys} \}_{new} = \frac{\delta H_{phys}}{\delta \mathbf{p}(x)} - \int dy^4 \sum_{\alpha=\phi,\theta} \frac{\delta B_\alpha(y)}{\delta \mathbf{p}(x)} \partial^0 \alpha(y), \tag{3.16}$$

the derivative of the Abelian component, complete with corrections from the Lorentz-monopole background is

$$\partial_0 \mathbf{c}_\sigma(x) = \frac{\delta H_{phys}}{\delta \mathbf{\Pi}^\sigma(x)} - \int dy^4 \sum_{\alpha=\phi,\theta} \frac{\delta B_\alpha(y)}{\delta \mathbf{\Pi}^\sigma(x)} \partial^0 \alpha(y), \tag{3.17}$$

where $\mathbf{\Pi}^\mu$ is the conjugate momentum of \mathbf{c}^μ . The effect of the second term is to remove the Lorentz-monopole contribution to $\frac{\delta H_{phys}}{\delta \mathbf{\Pi}^\sigma(x)}$. To see this, consider that, by construction, the Lorentz-monopole contribution to the Lagrangian and Hamiltonian is dependent on the time derivatives of θ, ϕ , so the monopole component of $\frac{\delta H_{phys}}{\delta \mathbf{\Pi}^\sigma(x)}$ is

$$\begin{aligned}
\frac{\delta}{\delta \mathbf{\Pi}^\sigma(x)} H_{phys}|_{\dot{\theta}\dot{\phi}} &= \frac{\delta}{\delta \mathbf{\Pi}^\sigma(x)} \left(\frac{\delta H_{phys}}{\delta \partial_0 \theta(x)} \partial_0 \theta(x) + \frac{\delta H_{phys}}{\delta \partial_0 \phi(x)} \partial_0 \phi(x) \right) \\
&= \frac{\delta}{\delta \mathbf{\Pi}^\sigma(x)} \left(\frac{\delta L_{phys}}{\delta \partial_0 \theta(x)} \partial_0 \theta(x) + \frac{\delta L_{phys}}{\delta \partial_0 \phi(x)} \partial_0 \phi(x) \right) \\
&= \frac{\delta}{\delta \mathbf{\Pi}^\sigma(x)} \left(B_\theta(x) \partial_0 \theta(x) + B_\phi(x) \partial_0 \phi(x) \right),
\end{aligned} \tag{3.18}$$

which is a consistency condition for eq. (3.17). This confirms the necessity of treating the monopole as a non-dynamic field.

We now observe that

$$\frac{\delta B_\theta(x)}{\delta \mathbf{c}^\sigma(y)} = \frac{\delta B_\phi(x)}{\delta \mathbf{c}^\sigma(y)} = 0, \tag{3.19}$$

from which it follows that the EOM of \mathbf{c}_σ receives no correction. However its $\{, \}_{phys}$ contribution, corresponding to the terms in the conventional EOM for the Abelian component, already contains a contribution from the monopole field strength.

Repeating the above steps for the valence components \mathbf{X}_μ , assuming $\sigma \neq 0$ and combining

$$\hat{D}_0 \tilde{\Pi}_\sigma(x) = \frac{\delta H}{\delta \mathbf{X}^\sigma(x)} - \int dy^4 \sum_{\alpha=\phi,\theta} \frac{\delta B_\alpha(y)}{\delta \mathbf{X}^\sigma(x)} \partial^0 \alpha(y). \quad (3.20)$$

with

$$\begin{aligned} \frac{\delta B_\phi(y)}{\delta \mathbf{X}^\sigma(x)} = & - \left(\left(\sin \phi(y) {}_y \partial_\sigma \theta(y) \hat{\mathbf{I}} + \hat{\mathbf{I}}_\theta(y) \times \mathbf{X}_\sigma(y) \right) \times \mathbf{X}_0(y) \right. \\ & \left. - \hat{\mathbf{I}}_\phi(y) \hat{\mathbf{I}}(y) \cdot \mathbf{R}_{0\sigma}(y) \right) \delta^4(x-y), \end{aligned} \quad (3.21)$$

$$\begin{aligned} \frac{\delta B_\theta(y)}{\delta \mathbf{X}^\sigma(x)} = & \left(\left({}_y \partial_\sigma \phi(y) \hat{\mathbf{I}}(y) + \sin \phi(y) \hat{\mathbf{I}}_\phi(y) \times \mathbf{X}_\sigma(y) \right) \times \mathbf{X}_0(y) \right. \\ & \left. - \sin \phi \hat{\mathbf{I}}_\theta(y) \hat{\mathbf{I}}(y) \cdot \mathbf{R}_{0\sigma}(y) \right) \delta^4(x-y), \end{aligned} \quad (3.22)$$

gives

$$\begin{aligned} \hat{D}_0 \tilde{\Pi}_\sigma(x) = & \frac{\delta H}{\delta \mathbf{X}^\sigma(x)} - \frac{1}{2} \left(\left(\sin \phi(x) (\partial_\sigma \phi(x) \partial_0 \theta(x) - \partial_\sigma \theta(x) \partial_0 \phi(x)) \hat{\mathbf{I}}(x) \right. \right. \\ & \left. \left. + \left(\sin \phi(x) \hat{\mathbf{I}}_\phi(x) \partial_0 \theta(x) - \hat{\mathbf{I}}_\theta(x) \partial_0 \phi(x) \right) \times \mathbf{X}_\sigma(x) \right) \times \mathbf{X}_0(x) \right) \\ = & \frac{\delta H}{\delta \mathbf{X}^\sigma(x)} - \frac{1}{2} g^2 \left(\mathbf{C}_\sigma(x) \times \mathbf{C}_0(x) + \mathbf{C}_0(x) \times \mathbf{X}_\sigma(x) \right) \times \mathbf{X}_0(x). \end{aligned} \quad (3.23)$$

This is the converse situation of the Abelian component, where their derivatives of \mathbf{X}_σ are uncorrected while their EOM receives a correction which cancels the monopole's electric contribution to $\{\hat{D}_0 \mathbf{X}_\sigma, H_{phys}\}_{phys}$. This is required by the conservation of topological current.

3.2.3 Corrections to the commutation relations

Corrections to the classical Poisson bracket correspond to corrections to the equal-time commutators in the quantum regime. Denoting conventional commutators as $[,]_{phys}$ and the corrected ones as $[,]_{new}$, for $\mu, \nu \neq 0$ we have

$$\begin{aligned} & [\mathbf{c}_\mu(x), \mathbf{c}_\nu(z)]_{new} \\ & = [\mathbf{c}_\mu(x), \mathbf{c}_\nu(z)]_{phys} \\ & \quad - \int dy^4 \left(\frac{\delta B_\theta(y)}{\delta \Pi^\mu(x)} \mathcal{F}_{\theta\phi}^{-1}(z) \frac{\delta B_\phi(y)}{\delta \Pi^\nu(z)} - \frac{\delta B_\phi(y)}{\delta \Pi^\mu(x)} \mathcal{F}_{\phi\theta}^{-1}(z) \frac{\delta B_\theta(y)}{\delta \Pi^\nu(z)} \right) \delta^4(x-z) \end{aligned} \quad (3.24)$$

From here the work is more transparent if we separate \mathbf{c}_μ into c_μ and \tilde{c}_μ , and $\mathbf{\Pi}_\mu$ into Π_μ and $\tilde{\Pi}_\mu$. We have

$$\begin{aligned}\frac{\delta B_\phi(y)}{\delta \mathbf{\Pi}_\nu(x)} &= \left(\sin \phi(y) {}_y\partial^\mu \theta(y) \hat{\mathbf{I}}(y) + (\hat{\mathbf{I}}_\theta(y) \times \mathbf{X}^\mu(y)) \right) \delta^4(x-y), \\ \frac{\delta B_\theta(y)}{\delta \mathbf{\Pi}_\nu(x)} &= -\sin \phi(y) \left({}_y\partial^\mu \phi(y) \hat{\mathbf{I}}(y) + (\hat{\mathbf{I}}_\phi(y) \times \mathbf{X}^\mu(y)) \right) \delta^4(x-y).\end{aligned}\quad (3.25)$$

Isolating the equations for Π_μ finds, $\tilde{\Pi}_\mu$ we find

$$\begin{aligned}\frac{\delta B_\phi(y)}{\delta \Pi_\nu(x)} &= \left(\sin \phi(y) {}_y\partial^\mu \theta(y) \hat{n}(y) + \hat{n}_\theta(y) \times \vec{Z}^\mu(y) \right) \delta^4(x-y), \\ \frac{\delta B_\theta(y)}{\delta \Pi_\nu(x)} &= -\sin \phi(y) \left({}_y\partial^\mu \phi(y) \hat{n}(y) + (\hat{n}_\phi(y) \times \vec{Z}^\mu(y)) \right) \delta^4(x-y),\end{aligned}\quad (3.26)$$

which upon substitution into eq. (3.24) gives us

$$\begin{aligned}& [c_\mu(x), c_\nu(z)]_{new} \\ &= [c_\mu(x), c_\nu(z)]_{phys} \\ &\quad - \sin \phi(x) \sin \phi(z) (\partial_\mu \phi(x) \partial_\nu \theta(z) - \partial_\nu \phi(z) \partial_\mu \theta(x)) \mathcal{F}_{\theta\phi}^{-1}(z) \delta^4(x-z).\end{aligned}\quad (3.27)$$

The second term on the final line, after integration over d^4z , clearly becomes

$$H_{\mu\nu}(x) \sin \phi(x) \mathcal{F}_{\theta\phi}^{-1}(x), \quad (3.28)$$

indicating the role of the monopole condensate in the correction. Repeating the calculation for \tilde{c}_μ

$$\begin{aligned}\frac{\delta B_\phi(x)}{\delta \tilde{\Pi}_\nu(y)} &= \hat{n}_\theta(x) \times \tilde{Z}^\mu(x) \delta^4(x-y), \\ \frac{\delta B_\theta(x)}{\delta \tilde{\Pi}_\nu(y)} &= -\sin \phi(y) \hat{n}_\phi(x) \times \tilde{Z}^\mu(x) \delta^4(x-y),\end{aligned}\quad (3.29)$$

we immediately see a significant difference, with the monopole condensate contributing no correctional terms to the Poisson bracket $[\tilde{c}_\mu(x), \tilde{c}_\nu(z)]_{phys}$. By contrast, the commutation relations

$$[c_\mu(x), \mathbf{\Pi}_\nu(z)]_{new} = [c_\mu(x), \mathbf{\Pi}_\nu(z)]_{phys}, \quad [\mathbf{\Pi}_\mu(x), \mathbf{\Pi}_\nu(z)]_{new} = [\mathbf{\Pi}_\mu(x), \mathbf{\Pi}_\nu(z)]_{phys}, \quad (3.30)$$

are unchanged. Nonetheless, the deviation from the canonical commutation shown in eq. (3.27) is inconsistent with the particle creation/annihilation operator formalism of conventional second quantization, so that, except under the very special circumstances

discussed in section 5, particle number is not well-defined for the c_μ field in general although it is for \tilde{c}_μ .

Repeating for the valence part,

$$\begin{aligned} & [\mathbf{\Pi}_\mu^a(x), \mathbf{\Pi}_\nu^b(z)]_{new} \\ &= [\mathbf{\Pi}_\mu^a(x), \mathbf{\Pi}_\nu^b(z)]_{phys} - \int dy^4 \left(\frac{\delta B_\theta(y)}{\delta \mathbf{X}_a^\mu(x)} \frac{\delta B_\phi(y)}{\delta \mathbf{X}_b^\nu(z)} - \frac{\delta B_\phi(y)}{\delta \mathbf{X}_a^\mu(x)} \frac{\delta B_\theta(y)}{\delta \mathbf{X}_b^\nu(z)} \right) \mathcal{F}_{\theta\phi}^{-1}(z) \end{aligned} \quad (3.31)$$

where now a, b are spatial indices and $\mathbf{\Pi}_\mu^a(x)$ is the a^{th} component of $\tilde{\mathbf{\Pi}}_\mu(x)$. We again find it illuminating to consider the rotation and boost components separately. For derivatives with respect to \vec{Z}_μ

$$\begin{aligned} \frac{\delta B_\phi(y)}{\delta \vec{Z}_a^\sigma(x)} &= - \left(\epsilon^{abc} \left(\sin \phi(y) {}_y\partial_\sigma \theta(y) \hat{n}_b(y) + \epsilon_{bde} \hat{n}_\theta^d(y) \times \vec{Z}_\sigma^e(y) \right) \times \vec{Z}_{c0}(y) \right. \\ &\quad \left. - \hat{n}_\phi^a \hat{\mathbf{I}}(y) \cdot \mathbf{R}_{0\sigma}(y) \right) \delta^4(x - y), \end{aligned} \quad (3.32)$$

$$\begin{aligned} \frac{\delta B_\theta(y)}{\delta \vec{Z}^\sigma(x)} &= \left(\epsilon^{abc} \left({}_y\partial_\sigma \phi(y) \hat{n}_b(y) + \epsilon_{bde} \sin \phi(y) \hat{n}_\phi^d(y) \times \vec{Z}_\sigma^e(y) \right) \times \vec{Z}_{c0}(y) \right. \\ &\quad \left. - \sin \phi \hat{n}_\theta^a(y) \hat{\mathbf{n}}(y) \cdot \mathbf{R}_{0\sigma}(y) \right) \delta^4(x - y), \end{aligned} \quad (3.33)$$

The second term on the final line integrates over d^4z to become

$$(n_\phi^a(x) n_\theta^b(x) - n_\theta^a(x) n_\phi^b(x)) \sin \phi(x) \hat{\mathbf{I}}(z) \cdot \mathbf{R}^{0\mu}(z) \hat{\mathbf{I}}(x) \cdot \mathbf{R}^{0\nu}(x) \mathcal{F}_{\theta\phi}^{-1}(x). \quad (3.34)$$

For the sake of completeness we expand the final term in these two equations, finding

$$\hat{\mathbf{I}}(y) \cdot \mathbf{R}_{0\nu}(y) = F_{0\sigma}(y) + \hat{n}(y) \cdot \left(\vec{Z}_\mu(y) \times \vec{Z}_0(y) - \vec{Z}_\mu(y) \times \vec{Z}_0(y) \right) \quad (3.35)$$

For derivatives with respect to \tilde{Z}_μ

$$\frac{\delta B_\phi(y)}{\delta \tilde{Z}_a^\sigma(x)} = -\delta^4(x - y) \epsilon^{abc} \left(\sin \phi(y) {}_y\partial_\sigma \theta(y) \hat{n}_b + \epsilon_{bde} \hat{n}_\theta^d(y) \times \tilde{Z}_\sigma^e(y) \right) \times \tilde{Z}_{c0}(y), \quad (3.36)$$

$$\frac{\delta B_\theta(y)}{\delta \tilde{Z}^\sigma(x)} = \delta^4(x - y) \epsilon^{abc} \left({}_y\partial_\sigma \phi(y) \hat{n}_b(y) + \sin \phi(y) \epsilon_{bde} \hat{l}_\phi^d(y) \times \tilde{Z}_\sigma^e(y) \right) \times \tilde{Z}_{c0}(y). \quad (3.37)$$

In parallel to the commutators of Abelian fields, we see correction to the commutator of \vec{Z}_μ but not of \tilde{Z}_μ .

Other relevant commutators are unchanged

$$[\mathbf{X}_\mu^a(x), \mathbf{\Pi}_\nu^b(z)]_{new} = [\mathbf{X}_\mu^a(x), \mathbf{\Pi}_\nu^b(z)]_{phys}, \quad [\mathbf{X}_\mu^a(x), \mathbf{X}_\nu^b(z)]_{new} = [\mathbf{X}_\mu^a(x), \mathbf{X}_\nu^b(z)]_{phys}. \quad (3.38)$$

4 Matter fields

There has been a strong parallel so far between the phenomenology of this theory and that of two-colour QCD. However their interactions with matter fields are different. Of particular note is that the Lorentz gauge field does not couple to scalars. We shall remedy this in future work when we incorporate the vierbein and translation symmetry, which are also required for gravitational coupling to fermions. The inclusion of fermions also requires a procedure for treating them within the Clairaut formalism, which we do not yet know how to do, so for now we can only consider vector fields.

To avoid the unhelpful complications of gauge fixing we consider a massive vector field V_μ whose kinetic terms are well-known to be

$$-\frac{1}{4}\vec{V}_{\mu\nu}\cdot\vec{V}^{\mu\nu}+\frac{1}{2}m^2V^\mu V_\nu=-\frac{1}{4}(\mathbf{D}_\mu V_\nu-\mathbf{D}_\nu V_\mu)\cdot(\mathbf{D}^\mu V^\nu-\mathbf{D}^\nu V^\mu)+\frac{1}{2}m^2V^\mu V_\nu, \quad (4.1)$$

where \mathbf{D}_μ is the covariant derivative and the conjugate momentum is

$$\mathbf{\Pi}_\nu=-\frac{1}{2}\mathbf{V}_{0\nu}. \quad (4.2)$$

The corresponding contributions to B_ϕ, B_θ are

$$\begin{aligned} B_\phi(x)|_{V_\nu}&=-\vec{V}^{0\nu}\cdot(\hat{\mathbf{l}}_\theta(x)V_\nu(x)), \\ B_\theta(x)|_{V_\nu}&=-\vec{V}^{0\nu}\cdot(\hat{\mathbf{l}}_\phi(x)V_\nu(x))\sin\phi(x), \end{aligned} \quad (4.3)$$

where we remind that $\hat{\mathbf{l}}, \hat{\mathbf{l}}_\theta, \hat{\mathbf{l}}_\phi$ are linear combinations of Lorentz matrices. This leads to a contribution of

$$\begin{aligned} \mathcal{F}_{\theta\phi}(x)|_{V_\nu}&=\mathbf{V}_{0\nu}\cdot\sin\phi(x)\hat{\mathbf{l}}_{\phi\theta}(x)V_\nu(x) \\ &=\mathbf{V}_{0\nu}\cdot\sin\phi(x)\cos\phi(x)\hat{\mathbf{l}}_\theta(x)V_\nu(x), \end{aligned} \quad (4.4)$$

where the last line follows from eq. (2.12) and because the different boosts are mutually orthogonal. The corrections to the commutation relations are proportional to the electric component of the vector field V_ν field strength.

5 Particle number and the Lorentz-monopole background

It is textbook knowledge that gravitational curvature spoils canonical quantisation, but our approach gives a detailed mechanism. It also provides some narrowly defined circumstances under which it may be salvaged. We saw in subsection 3.2.3 that among

the dynamic DOFs the rotation-related gauge fields c_μ, \vec{Z}_μ suffer corrections to their commutation relations inconsistent with a conventional particle interpretation while their boost-related counterparts \tilde{c}, \tilde{Z}_μ do not. Exactly how this affects the observed dynamics of the theory is not immediately obvious and beyond the scope of work done so far. A reasonable supposition is that the effect of linear momentum would dominate that of rotational momentum at large distances, as determined by some theory-relevant length scale.

We now consider the specific circumstances under which these particle-spoiling corrections might vanish. The form of eq. (3.27) indicates that they would arise for any polarisation of c_μ along either of the μ, ν directions. The only way to avoid this is if c_μ is polarised in the direction of the monopole field strength, requiring that the Abelian component of the connection propagate at a right-angle to the monopole field-strength. However, the form of the monopole field strength requires that a non-vanishing field must have a varying orientation in space, since it is proportional to the derivatives of the angles ϕ, θ . So even if the Abelian gauge component is propagating at a right-angle to the monopole field-strength with its polarisation in the direction of the field strength, in general this could not be assumed to continue as the orientation of the monopole field strength varied. However, if the variation were gradual over space in comparison to the wavelength of c_μ then it might continue to propagate while adjusting to the required orientations in a manner analogous to photon polarisation being rotated by successive, closely oriented, polarising filters. However, if the wavelength of c_μ is significant compared to the length scale of the field variation then such a mechanism could not act and the particle's energy would be either absorbed or deflected by the condensate, effectively suppressing the longer wavelengths and providing a measure of the background curvature. We remind the reader that c_μ represents the Abelian component of the Lorentz gauge field and has no ghost fields to consider due to its trivial gauge-fixing condition.

One important observation is that the background field is (Lorentz) magnetic, so that at any point in spacetime a reference frame exists where the monopole field and its associated potential lie entirely along the spatial directions.

The particle-inconsistent contribution from eq. (3.32) only occurs in the presence of a background electric component of the monopole field strength, vanishing when the polarisation of \vec{Z}_μ is orthogonal to the electric component of the background field. This restricts the polarisation for a transversally polarised field whose direction of propagation is not in the direction of this electric component, but not otherwise. Of course, the electric component of the background monopole field can always be removed by a suitable Lorentz transformation, but this still leaves the particle interpretation frame-dependent.

Some authors have argued that the valence gluons in two-colour QCD gain an effective mass term [13, 14] via their quartic interaction with the non-trivial monopole condensate. A similar mechanism could apply to the valence components of this theory.

Consider the following quartic term from the Lagrangian in eq. (2.23),

$$\begin{aligned} & \frac{g^2}{4} (\mathbf{C}_\mu \times \mathbf{X}_\nu) \cdot (\mathbf{C}^\mu \times \mathbf{X}^\nu) \\ &= \frac{g^2}{4} (\mathbf{C}_\mu \cdot \mathbf{C}^\mu \mathbf{X}_\nu \cdot \mathbf{X}^\nu - \mathbf{C}_\mu \cdot \mathbf{X}^\mu \mathbf{X}_\nu \cdot \mathbf{C}^\nu). \end{aligned} \quad (5.1)$$

Remembering that the Lorentz monopole field \mathbf{C}_μ has a non-zero condensate yields the term

$$\frac{g^2}{4} \langle \mathbf{C}_\mu \cdot \mathbf{C}^\mu \rangle \mathbf{X}_\nu \cdot \mathbf{X}^\nu, \quad (5.2)$$

so that the Lorentz-monopole condensate is seen to generate a mass term for the valence component. Such a mass term is covariant under the gauge transformation because, as shown in the discussion of eqs (2.24,2.25), the valence components transform as sources. In this case the valence components could also be longitudinally polarised. With longitudinal polarisation the only restriction is that the direction of propagation be orthogonal to the background electric component of the monopole field strength. The valence component might therefore enjoy a limited particle interpretation under a range of circumstances.

The particle spectrum of the Lorentz gauge field in the presence of the Lorentz monopole condensate would therefore contain the Abelian field \tilde{c}_μ and the valence field \tilde{Z}_μ corresponding to the boost symmetries. Corresponding particles for the rotation symmetries are not well-defined strictly speaking except for specific polarisations and directions of propagation as discussed above.

6 Discussion

We have considered the DOFs of gravitational curvature by applying the Clairaut formalism to Cho's adaptation of the CDG formalism for identifying the Abelian DOFs [29] in a Lorentz gauge theory quadratic in curvature \mathbf{R} . In this way we were able to rigorously separate physical from topological DOFs. Ours is not the only line of enquiry to consider gravitational phenomenology as topological rather than dynamic. Pak *et al.* [3, 4, 5] modelled gravitational curvature as a topological background with contorsion as the physical, confined, variable.

However our results indicate that gravity is not purely topological either but also contains dynamic DOFs, so that contorsion is not necessary for a stable background. As noted already, in the absence of matter fields the one-loop calculation of the monopole background in our theory is a doubling of the corresponding QCD calculation. This cannot be assumed to hold at higher order calculations when they become sensitive to the different consequences for rotation-related and boost-related components. Infra-red

analyses related to this issue were notoriously difficult in QCD and well beyond the scope of what we wish to address here.

It is well-known that the inclusion of light quark fields impacts the calculation of the β -function in QCD with consequences for asymptotic freedom. In particular, there is an upper limit on how many light quark flavours a QCD theory can have and remain asymptotically free. This number varies with the number of colours. If the stabilising background also holds here then the number of fermionic matter fields should also be limited. However the effect is offset by the bosonic fields, the lightest being photons and mesons, which would tend to strengthen the field. We simply note that the differences in their matter spectra suggest that this theory has significantly different infrared behaviour from that of $SU(2)$ QCD.

Our QCD analysis [30] demonstrated a breakdown of the second quantisation formalism. Similar terms also appeared in this analysis for the rotation-related gauge fields, but not for the boost-related ones. The physical implications of this observation must await later analysis but we speculate that effects coupling to linear momentum dominate those coupling to rotational momentum at low energy.

The particle-spoiling corrections were found to be neutralised by certain specific gauge choices and directions of propagation, and more robustly by the acquisition of mass. Hence massive particles were found to have a more robust particle number than massless ones. In the case of the valence components, parallels with two-colour QCD plausibly made this a consequence of the monopole field itself. For a massive vector field however, such convenient relationships among polarisation and background field strength did not occur. Future work will hopefully address these issues for fermionic matter fields.

We have left the inclusion of translation symmetry to subsequent work. We have side-stepped important technical issues by restricting our gauge theory to Lorentz symmetry and leaving out translational symmetry but a full gravitational theory must of course include the full Poincaré symmetry group. We suggest, however, that our Lorentz-only theory makes a sufficiently good approximation to indicate some relevant phenomenology.

A The CDG decomposition in $SU(2)$ QCD

Here we outline the application of the CDG decomposition in two-colour ($SU(2)$) QCD. The Lie group $SU(N)$ has $N^2 - 1$ generators $\lambda^{(a)}$ ($a = 1, \dots, N^2 - 1$), of which $N - 1$ are Abelian generators $\Lambda^{(i)}$ ($i = 1, \dots, N - 1$).

The gauge transformed Abelian directions (Cartan generators) are denoted as

$$\hat{n}_i(x) = U(x)^\dagger \Lambda^{(i)} U(x). \quad (1.1)$$

Gluon fluctuations in the $\hat{n}_i(x)$ directions are described by $c_\mu^{(i)}(x)$, where μ is the

Minkowski index. There is a covariant derivative which leaves the $\hat{n}_i(x)$ invariant,

$$\hat{D}_\mu \hat{n}_i(x) \equiv (\partial_\mu + g\vec{V}_\mu(x) \times) \hat{n}_i(x) = 0, \quad (1.2)$$

where $\vec{V}_\mu(x)$ is of the form

$$\vec{V}_\mu(x) = c_\mu^{(i)}(x) \hat{n}_i(x) + \vec{C}_\mu(x), \quad \vec{C}_\mu(x) = g^{-1} \partial_\mu \hat{n}_i(x) \times \hat{n}_i(x). \quad (1.3)$$

The vector notation refers to the internal space, and summation is implied over $i = 1, \dots, N - 1$. For later convenience we define

$$F_{\mu\nu}^{(i)}(x) = \partial_\mu c_\nu^{(i)}(x) - \partial_\nu c_\mu^{(i)}(x) \quad (1.4)$$

$$\vec{H}_{\mu\nu}(x) = \partial_\mu \vec{C}_\nu(x) - \partial_\nu \vec{C}_\mu(x) + g\vec{C}_\mu(x) \times \vec{C}_\nu(x) = H_{\mu\nu}^{(i)}(x) \hat{n}_i(x), \quad (1.5)$$

$$H_{\mu\nu}^{(i)}(x) = \vec{H}_{\mu\nu}(x) \cdot \hat{n}_i(x). \quad (1.6)$$

The vectors $\vec{X}_\mu(x)$ denote the dynamical components of the gluon field in the off-diagonal directions of the internal space, so if $\vec{A}_\mu(x)$ is the gluon field then

$$\vec{A}_\mu(x) = \vec{V}_\mu(x) + \vec{X}_\mu(x) = c_\mu^{(i)}(x) \hat{n}_i(x) + \vec{C}_\mu(x) + \vec{X}_\mu(x), \quad (1.7)$$

where

$$\vec{X}_\mu(x) \perp \hat{n}_i(x), \quad \forall 1 \leq i < N, \quad \vec{D}_\mu = \partial_\mu + g\vec{A}_\mu(x). \quad (1.8)$$

The Lagrangian density is still

$$\mathcal{L}_{gauge}(x) = -\frac{1}{4} \vec{F}_{\mu\nu}(x) \cdot \vec{F}^{\mu\nu}(x) \quad (1.9)$$

where the field strength tensor of QCD expressed in terms of the CDG decomposition is

$$\begin{aligned} \vec{F}_{\mu\nu}(x) &= (F_{\mu\nu}^{(i)}(x) + H_{\mu\nu}^{(i)}(x)) \hat{n}_i(x) \\ &+ (\hat{D}_\mu \vec{X}_\nu(x) - \hat{D}_\nu \vec{X}_\mu(x)) + g\vec{X}_\mu(x) \times \vec{X}_\nu(x). \end{aligned} \quad (1.10)$$

Henceforth we restrict ourselves to the $SU(2)$ theory, for which there is only one $\hat{n}(x)$ lying in a three dimensional internal space, and neglect the (i) indices. These results can be extended to larger $SU(N)$ gauge groups.

B Some convenient identities

The following identities concerning the unit vectors $\hat{n}, \hat{n}_\phi, \hat{n}_\theta$ and their derivatives may be of assistance in following the calculations:

$$\hat{n} = \hat{n}_\phi \times \hat{n}_\theta. \quad (2.1)$$

$$\begin{aligned} \partial_\mu \hat{n} \times \hat{n} &= \sin \phi \hat{n}_\phi \partial_\mu \theta - \hat{n}_\theta \partial_\mu \phi \\ \partial_\mu \hat{n} \times \hat{n}_\theta &= \hat{n} \partial_\mu \phi \\ \partial_\mu \hat{n} \times \hat{n}_\phi &= -\sin \phi \hat{n} \partial_\mu \theta - \hat{n}_\theta \partial_\mu \phi \end{aligned} \quad (2.2)$$

Corresponding identities hold for $\hat{\mathbf{I}}$ and its derivatives.

References

- [1] Kellogg S Stelle, "Renormalization of higher-derivative quantum gravity," *Physical Review D* **16**, 953 (1977).
- [2] Kenji Hayashi and Takeshi Shirafuji, "Gravity from Poincaré gauge theory of the fundamental particles. i: General formulation," *Progress of Theoretical Physics* **64**, 866 (1980).
- [3] S.-W. Kim and D. G. Pak, "Torsion as a dynamic degree of freedom of quantum gravity," *Classical and Quantum Gravity* **25**, 065011 (2008).
- [4] D.G. Pak, "Confinement, vacuum structure: from QCD to quantum gravity," *Nucl. Phys. A* **844**, 115c (2010), *Proceedings of the 4th International Symposium on Symmetries in Subatomic Physics*.
- [5] Y. M. Cho, D. G. Pak, and B. S. Park, "A minimal model of lorentz gauge gravity with dynamical torsion," *International Journal of Modern Physics A* **25**, 2867 (2010).
- [6] G. K. Savvidy, "Infrared instability of the vacuum state of gauge theories and asymptotic freedom," *Phys. Lett.* **B71**, 133 (1977).
- [7] N. K. Nielsen and P. Olesen, "An unstable Yang-Mills field mode," *Nucl. Phys.* **B144**, 376 (1978).
- [8] Y. M. Cho, "Colored monopoles," *Phys. Rev. Lett.* **44**, 1115 (1980).
- [9] Y. S. Duan and M. L. Ge, "SU(2) gauge theory and electrodynamics of N moving magnetic monopoles," *Sci. Sinica* **11**, 1072 (1979).
- [10] L. Faddeev and A. J. Niemi, "Partially dual variables in SU(2) Yang-Mills theory," *Phys. Rev. Lett.* **82**, 1624 (1999).

- [11] S. V. Shabanov, "An effective action for monopoles and knot solitons in Yang-Mills theory," *Phys. Lett.* **B458**, 322 (1999).
- [12] Y. M. Cho and D. G. Pak, "Monopole condensation in SU(2) QCD," *Phys. Rev.* **D65**, 074027 (2002).
- [13] K.-I. Kondo, T. Murakami, and T. Shinohara, "BRST symmetry of SU(2) Yang-Mills theory in Cho-Faddeev-Niemi decomposition," *Eur. Phys. J.* **C42**, 475 (2005).
- [14] K.-I. Kondo, "Gauge-invariant gluon mass, infrared Abelian dominance and stability of magnetic vacuum," *Phys. Rev.* **D74**, 125003 (2006).
- [15] Y. M. Cho and M. L. Walker, "Stability of monopole condensation in SU(2) QCD," *Mod. Phys. Lett.* **A19**, 2707 (2004).
- [16] Y. M. Cho, M. L. Walker, and D. G. Pak, "Monopole condensation and confinement of color in SU(2) QCD," *JHEP* **05**, 073 (2004).
- [17] D. Kay, A. Kumar, and R. Parthasarathy, "Savvidy vacuum in SU(2) Yang-Mills theory," *Mod. Phys. Lett.* **A20**, 1655 (2005).
- [18] M. L. Walker, "Stability of the magnetic monopole condensate in three- and four-colour QCD," *JHEP* **01**, 056 (2007).
- [19] M. L. Walker, "Higgs-free confinement hierarchy in five colour QCD," *Prog. Theor. Phys.* **119**, 139 (2007).
- [20] W. S. Bae, Y. M. Cho, and S. W. Kimm, "Qcd versus skyrme-faddeev theory," *Phys. Rev.* **D65**, 025005 (2002).
- [21] P. M. Lavrov and B. S. Merzlikin, "Legendre transformations and Clairaut-type equations," *Phys. Lett.* **B756**, 188 (2016).
- [22] J. Ren, H. Wang, Z. Wang, and F. Qu, "The Wu-Yang potential of magnetic skyrmion from $SU(2)$ flat connection," *Sci. China - Phys. Mech. Astron.* **62**, 950021 (2019).
- [23] M. L. Walker, "Extending SU(2) to SU(N) QCD," *Phys. Lett. B* **662**, 383 (2008).
- [24] K.-I. Kondo, "Wilson loop and magnetic monopole through a non-Abelian Stokes theorem," *Phys. Rev. D* **77**, 085029 (2008).
- [25] R. Matsudo and K.-I. Kondo, "Non-Abelian Stokes theorem for the Wilson loop operator in an arbitrary representation and its implication to quark confinement," *Phys. Rev. D* **92**, 125038 (2015).
- [26] K.-I. Kondo, S. Kato, A. Shibata, and T. Shinohara, "Quark confinement: dual superconductor picture based on a non-Abelian Stokes theorem and reformulations of Yang-Mills theory," *Phys. Repts.* **579**, 1 (2015).

- [27] M. L. Walker, "Applying the Cho-Duan-Ge decomposition to supersymmetric QCD," *J. Phys. G: Nuclear and Particle Physics* **40**, 095001 (2013).
- [28] Y. M. Cho, S. H. Oh, and Sang-Woo Kim, "Abelian dominance in Einstein's theory," *Classical and Quantum Gravity* **29**, 205007 (2012).
- [29] Y. M. Cho, S. H. Oh, and B. S. Park, "Abelian decomposition of Einstein's theory: Restricted gravity," *Grav. Cosm.* **21**, 257 (2015).
- [30] M. L. Walker and S. Duplij, "Cho-Duan-Ge decomposition of QCD in the constraintless Clairaut-type formalism," *Phys. Rev. D* **91**, 064022 (2015).
- [31] S. Duplij, "Generalized duality, Hamiltonian formalism and new brackets," *J. Math. Physics, Analysis, Geometry* **10**, 189 (2014).
- [32] S. Duplij, "Formulation of singular theories in a partial Hamiltonian formalism using a new bracket and multi-time dynamics," *Int. J. Geom. Methods in Modern Physics* **12**, 1550001 (2015).
- [33] Edward Witten, "2+ 1 dimensional gravity as an exactly soluble system," *Nuclear Physics B* **311**, 46 (1988).
- [34] Y. M. Cho, "Classification of gauge fields," *Journal of Mathematical Physics* **20**, 2605 (1979).
- [35] W. Bae, Y. M. Cho, and S. Kim, "QCD versus Skyrme-Faddeev theory," *Phys. Rev.* **D65**, 025005 (2001).
- [36] H. Flyvbjerg, "Improved QCD vacuum for gauge groups SU(3) and SU(4)," *Nucl. Phys.* **B176**, 379 (1980).
- [37] S. Duplij, "A new Hamiltonian formalism for singular Lagrangian theories," *J. Kharkov Univ., ser. Nuclei, Particles and Fields* **969**, 34 (2011).