

Reduced Phase Space Quantization of Black Hole, Path Integral, and Effective Dynamics

Cong Zhang^{*1}

¹*Faculty of Physics, University of Warsaw, Pasteura 5, 02-093 Warsaw, Poland*

We consider the loop quantum theory of the spherically symmetric model of gravity coupled to Gaussian dust fields, where the Gaussian dust fields provide a material reference frame of the space and time to deparameterize gravity. This theory, used to study the quantum features of the spherically symmetric BH, is constructed based on a 1-dimensional lattice $\gamma \subset \mathbb{R}$. Taking advantage of the path integral formulation, we investigate the quantum dynamics and obtain an effective action. With this action, we get an effective continuous description of this quantum lattice system which is not the same as the one described by the effective Hamiltonian used in [1], i.e. the classical Hamiltonian with the holonomy correction. It turns out that, the Hamiltonian derived in this paper can return to that used in [1] only for macro black holes, since the lattice γ should be chosen sufficiently fine. Moreover, the lattice is chosen sufficiently fine so that the continuous description can describe the underlying lattice theory well.

I. INTRODUCTION

Loop quantum gravity (LQG), as one of the most promising candidate for the theory of quantum gravity [2–5], has led to several significant achievements on the quantum features of spacetime [6–10]. However, there are still many unsettled open issues, particularly those on the dynamics of LQG. In spite of these efforts towards the full LQG dynamics [11–21], one applied the loop quantization procedures to symmetry-reduced models of gravity. The resulting models (see e.g. [22–28] for cosmological case, and see e.g. [1, 29–45] for black hole case), regarded as some symmetry-reduced sectors of LQG, yield valuable insight into the full LQG theory, regardless of the debates on embedding them into the full LQG [46–48].

There are two approaches to study the quantum nature of the spherically symmetric black hole (BH) with loop quantum method. An approach is to loop quantize the Schwarzschild interior. By virtue of homogeneity of the Schwarzschild interior, this approach leads to models of finite degrees of freedom [30–33, 36–44]. In this framework, the effective dynamics was detailed investigated based on various quantization strategies (see e.g. [32, 33] for μ_0 -scheme and $\bar{\mu}$ -scheme, and see [36, 38] for the modified scheme balancing the μ_0 -scheme and $\bar{\mu}$ -scheme). According to these works, the BH singularity is resolved. Particularly, the singularity is replaced by quantum bounce, which bridges the BH with a white hole in the framework of [32] for μ_0 -scheme and the framework of [38] for a modified scheme. However, for the $\bar{\mu}$ -scheme, according to the analysis in [32, 33], jumping over the quantum bounce, the classical BH gives birth to a baby BH which brings forth its own baby BH. This scenario continuous, giving the extended spacetime fractal structure, until eventually a Nariai type spacetime. Moreover, the quantum dynamics, in this framework, was also investigated by [42], which supports the existence of BH remnant by considering a general assumption on the quantization strategies. The other approach to investigate the quantum nature of BH is to consider the BH interior and exterior as a whole. This procedure was proposed and developed by [1, 34, 35, 45]. These works consider the spherical symmetry to reduce gravity, which leads to models of (1+1)-dimension field theory on \mathbb{R} . Moreover, they apply the $\bar{\mu}$ -scheme to do regularization, complemented by certain gauge fixing and choice of lapse function. An advantage of these models is to treat both the black hole interior and exterior in a unified manner.

The current work adapts the latter approach to the spherically symmetric model of gravity coupled to Gaussian dust fields as in [1]. In this model, gravity is deparameterized by the Gaussian dust fields which provide a material reference frame of the time and space. Thus its dynamics is governed by a physical Hamiltonian \mathbf{H} , which is identical to the Hamiltonian constraint of pure gravity with lapse function $N = 1$. Classically, this dynamics gives the Lemaître-Tolman-Bondi spacetimes. The quantum theory is obtained by applying the loop quantization procedure to this model. As the usual loop quantum theory, the states of this quantum model are constructed based on some graph γ . In this paper, γ is chosen to be a 1-dimensional lattice, i.e. $\gamma \subset \mathbb{R}$, whose vertices are denoted by $v \in V(\gamma)$. Each vertex carries some quantum numbers denoted by $\zeta(v)$ and $\mathbf{v}(v)$, where $\mathbf{v}(v)$ have the geometric interpretation of volume. To promote the physical Hamiltonian to an operator, some regularization procedure is necessary. Our work uses the $\bar{\mu}$ -scheme regularization strategy to get a regularized physical Hamiltonian \mathbf{H}_Δ . The Hamiltonian operator $\widehat{\mathbf{H}}_\Delta$ is thus derived by quantizing \mathbf{H}_Δ . To study the quantum dynamics, the path integral formulation is borrowed

* czhang(AT)fuw.edu.pl

to calculate the transition amplitude $A(\vec{\zeta}_i, \vec{\mathbf{v}}_i, \vec{\zeta}_f, \vec{\mathbf{v}}_f, T)$ from the initial state $|\vec{\zeta}_i, \vec{\mathbf{v}}_i\rangle$ at time $T_i = 0$ to the final state $|\vec{\zeta}_f, \vec{\mathbf{v}}_f\rangle$ at time $T_f = T$ [13, 49–51]. It turns out that this transition amplitude can be simplified in the standard path integral form, i.e.

$$A(\vec{\zeta}_i, \vec{\mathbf{v}}_i, \vec{\zeta}_f, \vec{\mathbf{v}}_f, T) = \int \prod_{v' \in V(\gamma)} \mathcal{D}[b(v')] \mathcal{D}[c(v')] \mathcal{D}[\ln(|\zeta(v')|)] \mathcal{D}[\mathbf{v}(v')] e^{i \frac{1}{\hbar} S(\vec{\zeta}, \vec{\mathbf{v}}, \vec{b}, \vec{c})} \quad (1.1)$$

where \vec{b} and \vec{c} are variables necessarily introduced during the derivation. With (1.1), the classical path is obtained by the method of stationary phase approximation, i.e. the equation $\delta S = 0$. $S(\vec{\zeta}, \vec{\mathbf{v}}, \vec{b}, \vec{c})$ is thus referred to as the effective action, from which the effective Hamiltonian $H(\vec{\zeta}, \vec{\mathbf{v}}, \vec{c}, \vec{b})$ is deduced.

We are now in a position to reconstruct an effective classical theory from the path integral formulation. This effective theory is a 1-dimensional lattice field theory whose dynamics is governed by $H(\vec{\zeta}, \vec{\mathbf{v}}, \vec{c}, \vec{b})$. An important result of our work is that $H(\vec{\zeta}, \vec{\mathbf{v}}, \vec{c}, \vec{b})$ is not the same \mathbf{H}_Δ , while the former can return to the latter if the volume of each vertex v , i.e. $\mathbf{v}(v)$, is much larger than the Planck volume. Here, it should be recalled that \mathbf{H}_Δ is the regularized Hamiltonian and its quantization gives the Hamiltonian operator $\widehat{\mathbf{H}}_\Delta$. In the usual work like [1], the continuous limit of \mathbf{H}_Δ , denoted by \tilde{H}_{eff} , is treated as the effective Hamiltonian which governs the improved dynamics of gravity with holonomy corrections. However, according to our analysis, this treatment is valid only if the following inequality holds,

$$\frac{2GM}{\ell_p} \gg \frac{4\pi \ell_p}{\alpha_0 \delta x} \quad (1.2)$$

where M is the ADM mass of Schwarzschild BH (the dynamics given by \tilde{H}_{eff} can recover the Schwarzschild solution in Lemaître coordinate according to [1]), δx , taking small values, is some specific coordinate length of edges in γ and α_0 is some constant. According to this inequality, since δx takes small values, this effective description by \tilde{H}_{eff} is valid for macro BH but not for small BH. In other words, the macro and small BHs behave differently once the loop quantum effects are considered.

Since the effective Hamiltonian coincides with that studied in [1] once (1.2) is satisfied, the effective dynamics for macro BH behaves the same as that in [1]. Particularly, the effective dynamics of the BH singularity is resolved by quantum bounce. In one regime far away from the bounce, the solution is semiclassical and reduces to the Schwarzschild spacetime in Lemaître coordinates. In the other asymptotic regime, the effective spacetime looks like the Nariai geometry which takes the metric of $\text{dS}_2 \times \mathbb{S}^2$. Moreover, according to the analysis in [1], both the area of \mathbb{S}^2 and the dS radius of the Nariai geometry obtained in this model are of quantum size. Thus, the current loop quantum model predicts a quantum final fate of Schwarzschild BH. However, it should be emphasized again that this is credible for the case where (1.2) holds.

It is worth noting that we are only concerned with the vacuum solution which can recover the Schwarzschild BH in this paper, even though Gaussian dusts are considered for deparameterization. For these solutions, the physical Hamiltonian, identical to the energy of the dusts field, vanishes. Thus they have no evolution with respect to the dust fields. From this point of view, a good analogy to these solutions in our situation is the ground-state solution of the quantum harmonic oscillator, regardless of the fact that “time” itself also disappears in our situation. Moreover, let us return to the original constraint theory where the deparameterization is not done. Then the solutions of the constraint equations, i.e., the physical states, are identical to the dynamical solutions derived from the deparameterized model. This identity tells us that, for the cases with non-vanishing dusts, one can pick the dust fields as time to give the physical states a relational evolution picture, while for the vanishing-dust case, one should choose other fields to do this. From these perspectives, it makes sense to consider the vacuum solution even in the deparameterization model.

This paper is organized as follows. In Sec. II, we review briefly the classical deparameterized model of spherically symmetric gravity coupled to Gaussian dust. Then the quantum kinematical theory of this model is introduced in Sec. III. Adapting the $\bar{\mu}$ -scheme, Sec. IV presents the physical Hamiltonian operator and its action. With these results, the quantum dynamics is studied by using the path integral formulation in Sec. V, where the transition amplitude is rewritten in the standard path integral form. Finally, the effective dynamics is studied in Sec. VI. Sec. VII summarizes the present work and gives some outlook on it.

II. CLASSICAL THEORY

A. Deparametrized gravity with Gaussian dust

In the reduced phase space formulation, gravity is coupled to some matter fields at classical level. The matter fields serve as the clock fields for deparametrizing the gravity dynamics. Here we focus on the scenario of gravity coupled to the Gaussian dust [52, 53]. The total action is given by

$$S = S_{\text{GR}} + S_{\text{GD}}, \quad (2.1)$$

where S_{GR} is the Host action of gravity [54]

$$S_{\text{GR}} [e_I^\mu, \Omega_{\mu\nu}^{IJ}] = \frac{1}{16\pi G} \int_M d^4y \sqrt{|\det(g)|} e_I^\mu e_J^\nu \left(\Omega_{\mu\nu}^{IJ} + \frac{1}{2\beta} \epsilon^{IJ}{}_{KL} \Omega_{\mu\nu}^{KL} \right) \quad (2.2)$$

where the tetrad e_I^μ determines the 4-metric by $g_{\mu\nu} = \eta_{IJ} e_I^\mu e_J^\nu$, and $\Omega_{\mu\nu}^{IJ}$ is the curvature of the $\mathfrak{so}(1,3)$ connection ω_μ^{IJ} . β is the Barbero-Immirzi parameter. S_{GD} is the action of the Gaussian dust:

$$S_{\text{GD}} [\rho, g_{\mu\nu}, T, S^a, W_b] = - \int_M d^4y \sqrt{|\det(g)|} \left[\frac{\rho}{2} (g^{\mu\nu} \partial_\mu T \partial_\nu T + 1) + g^{\mu\nu} \partial_\mu T (W_a \partial_\nu S^a) \right] \quad (2.3)$$

where $T, S^{a=1,2,3}$ are clock fields and defines time and space coordinates in the dust reference frame. ρ, W_a are Lagrange multipliers. The energy-momentum tensor of the Gaussian dust is

$$T_{\mu\nu} = \rho U_\mu U_\nu - U_{(\mu} W_{\nu)}, \quad U_\mu = -\partial_\mu T, \quad W_\nu = W_a \partial_\nu S^a. \quad (2.4)$$

which indicates that ρ is the energy density and W_μ relates to the heat-flow. See [52] for physical interpretation of the Gaussian dust theory.

We assume $M \simeq \mathbb{R} \times \Sigma$ and make Legendre transform of dust variables:

$$\begin{aligned} P &:= \frac{\delta S_{\text{GD}}}{\delta \dot{T}} = \sqrt{\det(q)} \{ \rho [\mathcal{L}_n T] + W_a [\mathcal{L}_n S^a] \} \\ P_a &:= \frac{\delta S_{\text{GD}}}{\delta \dot{S}^a} = \sqrt{\det(q)} W_a [\mathcal{L}_n T] \\ \pi &:= \frac{\delta S_{\text{GD}}}{\delta \dot{\rho}} = 0 \\ \pi^a &:= \frac{\delta S_{\text{GD}}}{\delta \dot{W}_a} = 0 \end{aligned} \quad (2.5)$$

where $q_{\mathbf{ab}}$ ($\mathbf{a}, \mathbf{b} = 1, 2, 3$) is the 3-metric and \mathcal{L}_n denotes the Lie derivative along the normal to the hypersurface Σ . The constraint analysis [52, 53] results in Hamiltonian and diffeomorphism constraints $\mathcal{C}^{\text{tot}}, \mathcal{C}_a^{\text{tot}}$ which are first-class constraints, and some second-class constraints whose solutions are

$$W_{\mathbf{a}} = \frac{P_{\mathbf{a}}}{\sqrt{\det(q)} (q^{\mathbf{ab}} T_{,\mathbf{a}} T_{,\mathbf{b}} + 1)^{1/2}} \quad (2.6)$$

$$\rho = \frac{P}{\sqrt{\det(q)} (q^{\mathbf{ab}} T_{,\mathbf{a}} T_{,\mathbf{b}} + 1)^{1/2}} - \frac{q^{\mathbf{ab}} T_{,\mathbf{a}} (P_c S_{,\mathbf{b}}^c)}{\sqrt{\det(q)} (q^{\mathbf{ab}} T_{,\mathbf{a}} T_{,\mathbf{b}} + 1)^{3/2}} \quad (2.7)$$

by a choice of sign in the ratio between W_a, P_a . The first class constraints $\mathcal{C}^{\text{tot}}, \mathcal{C}_a^{\text{tot}}$ reduce to

$$\mathcal{C}^{\text{tot}} = P + h, \quad h = \mathcal{C} \sqrt{1 + q^{\mathbf{ab}} T_{,\mathbf{a}} T_{,\mathbf{b}}} - q^{\mathbf{ab}} T_{,\mathbf{a}} \mathcal{C}_{\mathbf{b}}, \quad (2.8)$$

$$\mathcal{C}_a^{\text{tot}} = \mathcal{C}_a + P T_{,\mathbf{a}} + P_j S_{,\mathbf{a}}^j \quad (2.9)$$

where $\mathcal{C}, \mathcal{C}_a$ are pure gravity Hamiltonian and diffeomorphism constraints from S_{GR} .

S_{GR} leads to gravity canonical variables $A_{\mathbf{a}}^i(y), E_i^{\mathbf{a}}(y)$, where $A_{\mathbf{a}}^i(y)$ is the Ashtekar-Barbero connection and $E_i^{\mathbf{a}}(y) = \sqrt{\det q} e_i^{\mathbf{a}}(y)$ is the densitized triad. $i = 1, 2, 3$ is the Lie algebra index of $\mathfrak{su}(2)$. Dirac observables are constructed relationally by parametrizing (A, E) with values of dust fields $T(y) \equiv t, S^a(y) \equiv \sigma^a$, i.e.

$A_a^i(\sigma, t) = A_a^i(y)|_{T(y)\equiv t, S^a(y)\equiv\sigma^a}$ and $E_j^a(\sigma, t) = E_j^a(y)|_{T(y)\equiv t, S^a(y)\equiv\sigma^a}$, where σ, t are physical space and time coordinates in the dust reference frame. Here $a = 1, 2, 3$ is the dust coordinate index (e.g. $A_a(y) = A_a(y)\partial y^a/\partial\sigma^a$). $A_a^j(\sigma, t), E_j^a(\sigma, t)$ depending only on values of dust fields are independent of gauge choices of coordinates y . They are proven to be invariant (on the constraint surface) under gauge transformations generated by diffeomorphism and Hamiltonian constraints [55–57]. Moreover $A_a^j(\sigma, t), E_j^a(\sigma, t)$ satisfy the standard Poisson bracket in the dust frame:

$$\{E_i^a(\sigma, t), A_b^j(\sigma', t)\} = -\frac{1}{2}\kappa\beta \delta_i^j \delta_b^a \delta^3(\sigma, \sigma') \quad (2.10)$$

where β is the Barbero-Immirzi parameter and $\kappa = 16\pi G$.

The evolution in physical time t is generated by the physical Hamiltonian \mathbf{H}_0 given by integrating h on the constant $T(y) = t$ slice \mathcal{S} . The constant T slice \mathcal{S} is coordinated by the value of dust scalars $S^a = \sigma^a$ thus is referred to as the dust space [53, 57]. $T_{,a} = 0$ on \mathcal{S} leads to

$$\mathbf{H}_0 = \int_{\mathcal{S}} d^3\sigma \mathcal{C}(\sigma) \quad (2.11)$$

\mathbf{H}_0 formally coincides with smearing the gravity Hamiltonian \mathcal{C} with the unit lapse, while here $\mathcal{C}(\sigma)$ is in terms of Dirac observables $A_a^j(\sigma), E_j^a(\sigma)$. The classical dynamics is governed by the Hamilton's equation of \mathbf{H}_0 .

B. Spherical symmetry reduction

We assume the dust space $\mathcal{S} \simeq \mathbb{R} \times S^2$ and define the spherical coordinate $\sigma = (x, \theta, \phi)$. We further reduce the reduced phase space to the phase space Γ_{red} of spherical symmetric field configurations. The spherically symmetric conjugate variables (A_a^j, E_j^a) take the form [30, 34, 35]

$$\begin{aligned} A_a^j \tau_i d\sigma^a &= A_1(x) \tau_1 dx + \frac{1}{\sqrt{2}}(A_2(x) \tau_2 + A_3(x) \tau_3) d\theta + \frac{1}{\sqrt{2}}(A_2(x) \tau_3 - A_3(x) \tau_2) \sin(\theta) d\varphi + \cos(\theta) \tau_1 d\varphi \\ E_j^a \tau^j \frac{\partial}{\partial\sigma^a} &= E^1(x) \sin(\theta) \tau_1 \partial_x + \frac{1}{\sqrt{2}}(E^2(x) \tau_2 + E^3(x) \tau_3) \sin(\theta) \partial_\theta + \frac{1}{\sqrt{2}}(E^2 \tau_3 - E^3 \tau_2) \partial_\varphi \end{aligned} \quad (2.12)$$

where $\tau_j = -i/2(\text{Pauli matrix})_j$. Let Γ_{red} denote the reduced phase space of the spherically symmetric Ashtekar variables. The symplectic form Ω on Γ_{red} reads

$$\begin{aligned} \Omega(\delta_1, \delta_2) &= -\frac{1}{\kappa\beta} \int d^3\sigma \delta_1 A_a^j(\sigma) \wedge \delta_2 E_a^j(\sigma) \\ &= -\frac{1}{2G\beta} \int (\delta_1 A_1(x) \wedge \delta_2 E^1(x) + \delta_1 A_2(x) \wedge \delta_2 E^2(x) + \delta_1 A_3(x) \wedge \delta_2 E^3(x)) dx \end{aligned} \quad (2.13)$$

where δ_1 and δ_2 are two tangent vectors in the tangent space $T_{(A,E)}\Gamma_{\text{red}}$ and $\delta_1 F \wedge \delta_2 G \equiv \delta_1 F \delta_2 G - \delta_2 F \delta_1 G$. According to Ω , one has the Poisson bracket

$$\{A_I(x), E^J(y)\} = 2G\beta \delta(x, y) \delta_I^J, \quad I, J = 1, 2, 3. \quad (2.14)$$

The symmetry-reduced theory is an (1+1)-dimensional field theory.

The procedure of the deparametrization in Section II A does not solve the Gauss constraint. We still need to impose Gauss constraint to Γ_{red} . Substituting (2.12) into the general expression of the Gaussian constraint, one has the expression of $G[\lambda]$, associated to a $\mathfrak{su}(2)$ -valued smeared function $\lambda(x) = \lambda^i(x) \tau_i$ [29, 34],

$$G[\lambda] = 4\pi \int \lambda^1(x) [A_2(x) E^3(x) - A_3(x) E^2(x) + \partial_x E^1(x)] \quad (2.15)$$

Then the gauge transformation \mathcal{L}_G generated by the Gaussian constraint, i.e.,

$$\mathcal{L}_G(F) := e^{\{F, G[\lambda]\}} = 1 + \sum_{n=1}^{\infty} \frac{1}{n!} \underbrace{\{\dots \{F, G[\lambda]\}, G[\lambda]\} \dots}_{n \text{ folds}}, G[\lambda], \quad (2.16)$$

gives

$$\mathcal{L}_G(A_1(x)) = A_1(x) - \kappa\beta\partial_x\lambda^1(x) \quad (2.17)$$

$$\mathcal{L}_G(E^1(x)) = E^1(x) \quad (2.18)$$

$$\mathcal{L}_G(A_2(x) + iA_3(x)) = e^{i\kappa\beta\lambda^1(x)}(A_2 + iA_3) \quad (2.19)$$

$$\mathcal{L}_G(E^2(x) + iE^3(x)) = e^{i\kappa\beta\lambda^1(x)}(E^2 + iE^3). \quad (2.20)$$

That is, A_1 transforms as a U(1) gauge field and $A_2 + iA_3$ transforms as a U(1) Higgs field.

Given a state $(A_I, E^I) \in \Gamma_{\text{red}}$, one can always gauge transform it with some λ to vanish E^3 . Thus we introduce a gauge fixing condition

$$E^3(x) = 0. \quad (2.21)$$

The gauge fixed phase space then is coordinatized by $(A_I(x), E^I(x))$ with $I = 1, 2$ and, the Gaussian constraint (2.15) solves $A_3(x)$, leading to

$$A_3(x) = \frac{\partial_x E^1(x)}{E^2(x)}. \quad (2.22)$$

We introduce the following variables¹

$$\begin{aligned} K_x(x) &:= \frac{1}{2\beta}A_1(x), \quad E^x(x) = E^1(x), \\ K_\varphi(x) &:= \frac{1}{\sqrt{2}\beta}A_2(x), \quad E^\varphi(x) = \frac{1}{\sqrt{2}}E^2(x). \end{aligned} \quad (2.23)$$

The gauge-fixed reduced phase space $\mathcal{P} \subset \Gamma_{\text{red}}$ consist of canonical pairs $(K_x(x), E^x(x))$ and $(K_\varphi(x), E^\varphi(x))$. Restricting the symplectic form on \mathcal{P} , we get the non-vanishing Poisson brackets

$$\{K_I(y), E^I(z)\} = G\delta(y - z), \quad I = x, \varphi. \quad (2.24)$$

The classical physical Hamiltonian on \mathcal{P} is obtained from \mathbf{H}_0 in (2.11) by implementing the spherical symmetry, and adding a boundary term \mathbf{H}_{bdy}

$$\mathbf{H} = \frac{2\pi}{\kappa} \int dx \mathcal{C}(x) + \mathbf{H}_{\text{bdy}} \quad (2.25)$$

where $\mathcal{C}(x)$ is expressed as

$$\mathcal{C}(x) = \frac{\text{sgn}(E^\varphi)}{\sqrt{|E^x|}} \left(-\frac{E^{x'2}}{2E^\varphi} - 8E^x K_x K_\varphi - 2E^\varphi K_\varphi^2 - 2E^\varphi \right) + \left(\frac{2E^x E^{x'}}{\sqrt{|E^x||E^\varphi|}} \right)' \quad (2.26)$$

The equations of motion from \mathbf{H} is formally the same as the pure gravity dynamics with unit lapse, although the foliation is provided by the Gaussian dust.

The space of x is non-compact so boundary condition is needed. We follow [1] to discuss boundary conditions and corresponding boundary terms \mathbf{H}_{bdy} . Firstly, since we are going to discuss the quantum effective modification of the Schwarzschild black hole. It is useful to introduce the asymptotic boundary condition of the Schwarzschild at $x \rightarrow \infty$. A foliation of the Schwarzschild spacetime with unit lapse is the Lemaître coordinate. The asymptotic boundary condition in the Lemaître coordinate reads

$$E^x \sim \left(\frac{3}{2} \sqrt{R_s} x \right)^{4/3}, \quad K_x \sim \frac{R_s}{3 \times 12^{1/3} (\sqrt{R_s} x)^{4/3}}, \quad (2.27)$$

$$E^\varphi \sim \sqrt{R_2} \left(\frac{3}{2} \sqrt{R_s} x \right)^{1/3}, \quad K_\varphi \sim -\frac{(2/3)^{1/3} \sqrt{R_s}}{(\sqrt{R_s} x)^{1/3}}. \quad (2.28)$$

¹ $K_x(x), E^x(x), K_\varphi(x), E^\varphi(x)$ are the same as in e.g. [1, 35],

The following boundary Hamiltonian at $x \rightarrow \infty$ turns out to be suitable for this boundary condition for canceling boundary terms of $\delta\mathbf{H}$.

$$\mathbf{H}_{\text{bdy}} = -\frac{8\pi}{\kappa} \left(\frac{2E^x E^{x'}}{\sqrt{|E^x|}|E^\varphi|} - 2\sqrt{|E^x|} \right) \Big|_{\infty}. \quad (2.29)$$

Alternatively, instead of the above Schwarzschild asymptotic boundary condition, we may make an infrared cut-off at $x = L \gg 1$ and impose the Dirichlet boundary condition

$$\delta E^x|_{x=L} = 0 \quad (2.30)$$

In this case, \mathbf{H}_{bdy} is the same except for locating at the finite boundary $x = L$:

$$\mathbf{H}_{\text{bdy}} = -\frac{8\pi}{\kappa} \left(\frac{2E^x E^{x'}}{\sqrt{|E^x|}|E^\varphi|} - 2\sqrt{|E^x|} \right) \Big|_{x=L}. \quad (2.31)$$

In addition, on the other side $x \rightarrow -\infty$ or $x = -L$, we impose the Neumann boundary condition

$$E^{x'} = 0. \quad (2.32)$$

This boundary condition will be manifest in the effective dynamics of the black hole. We do not need any boundary Hamiltonian for the Neumann boundary condition. By using \mathbf{H}_{bdy} , the Hamiltonian \mathbf{H} can be simplified to the following form

$$\mathbf{H} = -\frac{4\pi}{\kappa} \int dx \mathfrak{h}(x) + \frac{1}{G} \sqrt{|E^x|} \Big|_{\text{bdy}} \quad (2.33)$$

with

$$\mathfrak{h}(x) = \frac{\text{sgn}(E^\varphi)}{\sqrt{|E^x|}} \left(\frac{E^{x'2}}{2E^\varphi} + 8E^x K_x K_\varphi + 2E^\varphi K_\varphi^2 + 2E^\varphi \right). \quad (2.34)$$

Let $\mathcal{V}(x)$ denote the diffeomorphism constraint of pure gravity. Then $\mathcal{V}(x)$ in the reduced model reads

$$\mathcal{V}(x) = K_x(x)E^{x'}(x) - E^\varphi(x)K'_\varphi(x). \quad (2.35)$$

It can be verified that $\{\mathbf{H}, \mathcal{V}(x)\} = 0$. Thus the time evolution by \mathbf{H} has infinitely many conserved charges $\mathcal{V}(x)$. It turns out that the Schwartzchild solution in Lemaître coordinates is obtained with setting the initial condition $\mathcal{V}(x) = 0$. Then $\{\mathbf{H}, \mathcal{C}(x)\} \propto \mathcal{V}(x)$ also vanishes, which gives another conserved charges $\mathcal{C}(x)$.

III. QUANTUM KINEMATICS

Discretize x -space to 1-dimensional equidistant lattice γ whose vertices are denoted by v . $e(v)$ denotes the edge starting at v and oriented toward the positive direction. The sets of edges and vertices in γ are denoted by $E(\gamma)$ and $V(\gamma)$ respectively. It is worth noting that, to make γ finite, we restrict ourselves in the interval $[-L, L]$ with $L \gg 1$ and consider the fields satisfying the boundary conditions aforementioned.

To do quantization, the fields K_I and E^I ($I = x, \varphi$) are first smeared by using γ as follows

$$\theta(v) = \int_{e(v)} dx K_x, \quad p(v) = E^x(\text{mid}_{e(v)}), \quad (3.1)$$

$$\Phi(v) = K_\varphi(\text{mid}_{e(v)}), \quad \Pi(v) = \int_{e(v)} dx E^\varphi, \quad (3.2)$$

where $\text{mid}_{e(v)}$ is the middle point within $e(v)$. Here, $K_x(x)$ is gauge fixed from the $U(1)$ gauge field and, thus, $e^{i\theta(v)}$ is its holonomy along $e(v)$. However, K_φ is gauge fixed from the $U(1)$ Higgs field, and, thus, $e^{i\Phi(v)}$ is its point holonomy at v .

With the variables $\theta(v)$ and $\Phi(v)$, the Hilbert space, constructed by the polymer quantization procedure, reads

$$\mathcal{H} = \overline{\left(L^2(\mathbb{R}_b, d\mu_h) \otimes L^2(\mathbb{R}_b, d\mu_h) \right)^{|V(\gamma)|}}. \quad (3.3)$$

where \mathbb{R}_b is the Bohr compactification of the real line \mathbb{R} and $d\mu_h$ is its Haar measure. To represent the elements in \mathcal{H} explicitly, let us define the space **Cyl** of almost periodic functions of $\vec{\theta} = \{\theta(v)\}_{v \in V(\gamma)}$ and $\vec{\Phi} = \{\Phi(v)\}_{v \in V(\gamma)}$. In other words, elements in **Cyl** are finite linear combination of the functions $T_{\vec{\mu}, \vec{\lambda}}$

$$T_{\vec{\mu}, \vec{\lambda}}(\vec{\theta}, \vec{\Phi}) = \prod_{v \in V(\gamma)} e^{i\lambda(v)\theta(v) + i\mu(v)\Phi(v)}. \quad (3.4)$$

Equipping **Cyl** with the inner product

$$\langle \Psi_1 | \Psi_2 \rangle = \left(\frac{1}{2T} \right)^{2|V(\gamma)|} \int \prod_{v \in V(\gamma)} d\theta(v) d\Phi(v) \overline{\Psi_1(\vec{\theta}, \vec{\Phi})} \Psi_2(\vec{\theta}, \vec{\Phi}). \quad (3.5)$$

we have

$$\langle T_{\{\vec{\lambda}, \vec{\mu}\}} | T_{\{\vec{\lambda}', \vec{\mu}'\}} \rangle = \delta_{\vec{\lambda}, \vec{\lambda}'} \delta_{\vec{\mu}, \vec{\mu}'}. \quad (3.6)$$

where the right hand side is the Kronecker delta. The Hilbert space \mathcal{H} is identical to the Cauchy completion of the space **Cyl** with respect this inner product. By using the Dirac bra-ket notions, we will denote

$$T_{\{\vec{\lambda}, \vec{\mu}\}} \equiv |\vec{\lambda}, \vec{\mu}\rangle. \quad (3.7)$$

(3.6) means that $|\vec{\lambda}, \vec{\mu}\rangle$ forms a family of orthonormal basis of \mathcal{H} .

In \mathcal{H} , there are the basic operators $e^{i\widehat{\lambda_o(v)\theta(v)}}$, $e^{i\widehat{\mu_o(v)\Phi(v)}}$, $\hat{p}(v)$ and $\hat{\Pi}(v)$ with constants $\lambda_o(v)$ and $\mu_o(v)$:

$$e^{i\widehat{\lambda_o(v)\theta(v)}} |\vec{\lambda}, \vec{\mu}\rangle = |\vec{\lambda} + \lambda_o(v)\vec{\delta}_v, \vec{\mu}\rangle \quad (3.8)$$

$$e^{i\widehat{\mu_o(v)\Phi(v)}} |\vec{\lambda}, \vec{\mu}\rangle = |\vec{\lambda}, \vec{\mu} + \mu_o(v)\vec{\delta}_v\rangle \quad (3.9)$$

$$\hat{p}(v) |\vec{\lambda}, \vec{\mu}\rangle = \ell_p^2 \lambda(v) |\vec{\lambda}, \vec{\mu}\rangle, \quad (3.10)$$

$$\hat{\Pi}(v) |\vec{\lambda}, \vec{\mu}\rangle = \ell_p^2 \mu(v) |\vec{\lambda}, \vec{\mu}\rangle. \quad (3.11)$$

where $\vec{\delta}_v$ is defined by

$$\vec{\delta}_v(v') = \begin{cases} 0, & v' \neq v, \\ 1, & v' = v. \end{cases} \quad (3.12)$$

It is remarkable that that operators $e^{i\widehat{\lambda_o(v)\theta(v)}}$ and $e^{i\widehat{\mu_o(v)\Phi(v)}}$ are not strongly continuous. Therefore, there is no operators $\hat{\theta}(v)$ and $\hat{\Phi}(v)$ such that $e^{i\widehat{\lambda_o(v)\theta(v)}} = \exp(i\lambda(v)\hat{\theta}(v))$ and $e^{i\widehat{\mu_o(v)\Phi(v)}} = \exp(i\mu(v)\hat{\Phi}(v))$ in this model.

Let V_v denoted the volume of the region $e(v) \times \mathbb{S}^2$. By definition, V_v is

$$V_v = \int_{e(v)} dx \int_0^\pi d\theta \int_0^{2\pi} d\varphi |E^\varphi| \sqrt{|E^x|} \sin(\theta) = 4\pi |\Pi(v)| \sqrt{|p(v)|}, \quad (3.13)$$

where we used that $\text{sgn}(E^\varphi)$ is a constant. Thus its corresponding operator $\hat{V}_v = 4\pi |\hat{\Pi}(v)| \sqrt{|\hat{p}(v)|}$ takes the basis $|\vec{\lambda}, \vec{\mu}\rangle$ as its eigenstate, i.e.

$$\hat{V}_v |\vec{\lambda}, \vec{\mu}\rangle = 4\pi \ell_p^3 |\mu(v)| \sqrt{|l(v)|} |\vec{\lambda}, \vec{\mu}\rangle. \quad (3.14)$$

For convenience \hat{V}_v as well as V_v will be referred to as the volume (operator) at v .

IV. THE PHYSICAL HAMILTONIAN OPERATOR

A. $\bar{\mu}$ -scheme regularization and quantization

Due to the absence of the operators $\hat{\theta}(v)$ and $\hat{\Phi}(v)$, the classical expression (2.34) of the physical Hamiltonian needs to be regularized in terms of $e^{i\lambda(v)\theta(v)}$ and $e^{i\mu(v)\theta(v)}$. One can refer to [1, 34] and references therein for the detailed derivations on the regularization. Generally speaking, the regularization procedure results in a family of regularized expressions $\mathbf{H}_{\bar{\lambda}, \bar{\mu}}$, depending on parameters $\bar{\lambda}$ and $\bar{\mu}$. Classically, it has

$$\lim_{\bar{\lambda}, \bar{\mu} \rightarrow 0} \mathbf{H}_{\bar{\lambda}, \bar{\mu}} = \mathbf{H}.$$

However, in the quantum theory the limit $\lim_{\bar{\lambda}, \bar{\mu} \rightarrow 0} \hat{\mathbf{H}}_{\bar{\lambda}, \bar{\mu}}$ fails to exist, where $\hat{\mathbf{H}}_{\bar{\lambda}, \bar{\mu}}$ denotes the quantization operator of $\mathbf{H}_{\bar{\lambda}, \bar{\mu}}$. To overcome the failure of the limit to exist, the idea proposed in loop quantum cosmology model [22, 24] is borrowed, which lead to the operator $\hat{\mathbf{H}}$, i.e. the quantization operator of \mathbf{H} , defined as

$$\hat{\mathbf{H}} = \lim_{\bar{\lambda} \rightarrow \bar{\lambda}, \bar{\mu} \rightarrow \bar{\mu}} \hat{\mathbf{H}}_{\bar{\lambda}, \bar{\mu}}$$

with some non-vanishing quantities $\bar{\lambda} = \{\lambda(v)\}$ and $\bar{\mu} = \{\mu(v)\}$. There are several strategies to choose $\bar{\lambda}$ and $\bar{\mu}$. The current work focus only on the $\bar{\mu}$ -scheme as in [1, 34] where $\bar{\lambda}$ and $\bar{\mu}$ are

$$\lambda(v) = \bar{\lambda}(v) := \frac{2\beta\sqrt{\Delta}\ell_p\sqrt{|p(v)|}}{\Pi(v)}, \quad \mu = \bar{\mu}(v) := \beta\frac{\sqrt{\Delta}\ell_p}{\text{sgn}(p(v))\sqrt{|p(v)|}}. \quad (4.1)$$

This strategy leads to the regularized physical Hamiltonian \mathbf{H}_Δ , referred as the improved Hamiltonian. Indeed, the improved Hamiltonian is equal to the effective Hamiltonian used in [1], but replacing the integral over x and, the variables K_I as well as E^I ($I = x, \varphi$), respectively, by discrete sum over $v \in V(\gamma)$ and, the corresponding smeared variables. Explicitly, \mathbf{H}_Δ reads

$$\mathbf{H}_\Delta = -\frac{1}{G} \sum_v \mathfrak{h}_{\Delta, v} + \frac{1}{G} \sqrt{|p(v_b)|} \quad (4.2)$$

with

$$\mathfrak{h}_{\Delta, v} = \frac{V_v \sin\left(\frac{\beta\sqrt{\Delta}\ell_p}{\text{sgn}(p(v))\sqrt{|p(v)|}}\Phi(v)\right) \sin\left(\frac{2\beta\sqrt{\Delta}\ell_p\sqrt{|p(v)|}}{\Pi(v)}\theta(v)\right)}{4\pi\beta^2\Delta\ell_p^2} \quad (4.3)$$

$$+ \frac{V_v \sin^2\left(\frac{\beta\sqrt{\Delta}\ell_p}{\text{sgn}(p(v))\sqrt{|p(v)|}}\Phi(v)\right)}{8\pi\beta^2\Delta\ell_p^2} + 2\pi\frac{\Pi(v)^2}{V_v} + \frac{\pi}{2}\frac{1}{V_v} [p(v+1) - p(v)]^2 \quad (4.4)$$

where v_b denotes the source of the edge e_b connecting to the boundary and $v+1$ denotes the target of $e(v)$. It is noted that $\sum_v \mathfrak{h}_{\Delta, v}$ reproduce $\int dx \mathfrak{h}(x)$ in the limits of low curvature and lattice refinement (e_v shrinks to zero size while vertices become dense).

The expression of $\mathfrak{h}_{\Delta, v}$ contains the inverse volume $1/V_v$ whose naive quantization $1/\hat{V}_v$ usually cause divergence at the zero eigenvalue of \hat{V}_v . Due to this difficulty, we need the trick

$$V(\mathcal{R})^{-1} = \frac{-4 \times 27}{6} \left(\frac{2}{\kappa\beta}\right)^3 \int_{\mathcal{R}} \left| \epsilon^{abc} \text{tr} \left(\{A_a(x), V(\mathcal{R})^{1/3}\} \{A_b(x), V(\mathcal{R})^{1/3}\} \{A_c(x), V(\mathcal{R})^{1/3}\} \right) \right|. \quad (4.5)$$

which can be verified by applying the Thiemann's trick [5]. Applying this formula to our current model, one finally get, by a straightforward calculation, that

$$\frac{1}{V_v} = \lim_{\lambda, \mu \rightarrow 0} \left(\frac{27}{8\pi^2 i G^3 \mu^2 \lambda} \right) \text{sgn}(p(v)) \mathcal{Q}_{\theta, \lambda}^{(1/3)}(v) \mathcal{Q}_{\Phi, \mu}^{(1/3)}(v) \mathcal{Q}_{\Phi, \mu}^{(1/3)}(v), \quad (4.6)$$

where

$$\mathcal{Q}_{\theta,\lambda}^{(r)}(v) = e^{i\lambda\theta(v)/2} \left\{ e^{-i\lambda\theta(v)/2}, V_v^r \right\} - e^{-i\lambda\theta(v)/2} \left\{ e^{i\lambda\theta(v)/2}, V_v^r \right\} \quad (4.7)$$

$$\mathcal{Q}_{\Phi,\mu}^{(r)}(v) = e^{i\mu\Phi(v)/2} \left\{ e^{-i\mu\Phi(v)/2}, V_v^r \right\} - e^{-i\mu\Phi(v)/2} \left\{ e^{i\mu\Phi(v)/2}, V_v^r \right\}. \quad (4.8)$$

Then the inverse volume operator can be quantized through replacing the Poisson brackets by the commutators. Because of the expression (4.3) of $\mathfrak{h}_{\Delta,v}$, we consider the new holonomy operators

$$\begin{aligned} \hat{h}_{\theta,\Delta}(v) &:= \exp \left[\widehat{i\bar{\lambda}(v)\theta(v)} \right] \\ \hat{h}_{\Phi,\Delta}(v) &:= \exp \left[\widehat{i\bar{\mu}(v)\Phi(v)} \right]. \end{aligned} \quad (4.9)$$

Because of the dependence of $\bar{\lambda}(v)$ and $\bar{\mu}(v)$ on $p(v)$ and $\Pi(v)$, to define these two operators, we use the idea proposed by [24] and set

$$\Phi(v) = i\ell_p^2 \frac{d}{d\Pi(v)}, \quad \theta(v) = i\ell_p^2 \frac{d}{dp(v)}. \quad (4.10)$$

Then the RHSs of (4.9) are written as

$$\begin{aligned} \hat{h}_{\theta,\Delta}(v)\psi(\vec{\lambda}, \vec{\mu}) &= \exp \left[-\frac{2\beta\sqrt{\Delta}\sqrt{|\lambda(v)|}}{\mu(v)} \frac{d}{d\lambda(v)} \right] \psi(\vec{\lambda}, \vec{\mu}) \\ \hat{h}_{\Phi,\Delta}(v)\psi(\vec{\lambda}, \vec{\mu}) &= \exp \left[-\frac{\beta\sqrt{\Delta}\ell_p}{\text{sgn}(p(v))\sqrt{|p(v)|}} \frac{d}{d\lambda(v)} \right] \psi(\vec{\lambda}, \vec{\mu}) \end{aligned} \quad (4.11)$$

Denoting

$$s(x) = \text{sgn}(x)\sqrt{|x|},$$

one can calculate the RHSs of (4.11) to get

$$\begin{aligned} \left(\hat{h}_{\theta,\Delta}(v)\psi \right) (\vec{\lambda}, \vec{\mu}) &= \psi(s^{-1}[s(\vec{\lambda}) - \beta\sqrt{\Delta}\mu(v)^{-1}\vec{\delta}_v], \vec{\mu}) \\ \left(\hat{h}_{\Phi,\Delta}(v)\psi \right) (\vec{\lambda}, \vec{\mu}) &= \psi(\vec{\lambda}, \vec{\mu} - \beta\sqrt{\Delta}s(\lambda(v))^{-1}\vec{\delta}_v) \end{aligned} \quad (4.12)$$

where $s(\vec{\lambda}) := \{s(\lambda(v))\}_v$ and s^{-1} denotes its inverse function. For convenience, we use $\vec{\zeta} = \frac{s(\vec{\lambda})}{\beta\sqrt{\Delta}}$ instead of $\vec{\lambda}$ to re-label the state $|\vec{\lambda}, \vec{\mu}\rangle$. Then we have

$$\text{sgn}(\hat{p}(v))\sqrt{|\hat{p}(v)|}|\vec{\zeta}, \vec{\mu}\rangle = \zeta(v)\beta\sqrt{\Delta}\ell_p|\vec{\zeta}, \vec{\mu}\rangle, \quad \hat{\Pi}(v)|\vec{\zeta}, \vec{\mu}\rangle = \ell_p^2\mu(v)|\vec{\zeta}, \vec{\mu}\rangle. \quad (4.13)$$

With the new conventions, (4.12) becomes

$$\begin{aligned} \hat{h}_{\theta,\Delta}(v)|\vec{\zeta}, \vec{\mu}\rangle &= |\vec{\zeta} + \mu(v)^{-1}\vec{\delta}_v, \vec{\mu}\rangle, \\ \hat{h}_{\Phi,\Delta}(v)|\vec{\zeta}, \vec{\mu}\rangle &= |\vec{\zeta}, \vec{\mu} + \zeta(v)^{-1}\vec{\delta}_v\rangle. \end{aligned} \quad (4.14)$$

Here, the derivation uses $(\hat{O}\psi)(\vec{\zeta}, \vec{\mu}) = \langle \vec{\zeta}, \vec{\mu} | \hat{O} | \psi \rangle$ for operator \hat{O} . Moreover, the action of the volume operator at v with the new convention is

$$\hat{V}_v|\vec{\zeta}, \vec{\mu}\rangle = 4\pi\ell_p^3\beta\sqrt{\Delta}|\mu(v)\zeta(v)||\vec{\zeta}, \vec{\mu}\rangle. \quad (4.15)$$

In the framework of the $\bar{\mu}$ -scheme, the inverse volume in (4.6) is quantized as

$$\begin{aligned} \widehat{1/V}_v &= \lim_{\substack{\lambda \rightarrow \bar{\lambda}(v) \\ \mu \rightarrow \bar{\mu}(v)}} (1/\widehat{V}_v)_{\vec{\lambda}, \vec{\mu}}, \\ (1/V_v)_{\vec{\lambda}, \vec{\mu}} &= \left(\frac{27}{8\pi^2 i G^3 \mu^2 \lambda} \right) \text{sgn}(p(v)) \mathcal{Q}_{\theta,\lambda}^{(1/3)}(v) \mathcal{Q}_{\Phi,\mu}^{(1/3)}(v) \mathcal{Q}_{\Phi,\mu}^{(1/3)}(v). \end{aligned} \quad (4.16)$$

This equation leads to the operators $\hat{\mathcal{Q}}_{\theta,\bar{\lambda}}^{(r)}$ and $\hat{\mathcal{Q}}_{\Phi,\bar{\mu}}^{(r)}$ defined as

$$\begin{aligned}\hat{\mathcal{Q}}_{\theta,\bar{\lambda}}^{(r)}(v) &:= \lim_{\lambda \rightarrow \bar{\lambda}} \hat{\mathcal{Q}}_{\theta,\lambda}^{(r)} = \frac{1}{i\hbar} e^{i\bar{\lambda}\theta(v)/2} \left[e^{-i\bar{\lambda}\theta(v)/2}, V_v^r \right] - \frac{1}{i\hbar} e^{-i\bar{\lambda}\theta(v)/2} \left[e^{i\bar{\lambda}\theta(v)/2}, V_v^r \right], \\ \hat{\mathcal{Q}}_{\Phi,\bar{\mu}}^{(r)}(v) &:= \lim_{\mu \rightarrow \bar{\mu}} \hat{\mathcal{Q}}_{\Phi,\mu}^{(r)} = \frac{1}{i\hbar} e^{i\bar{\mu}\Phi(v)/2} \left[e^{-i\bar{\mu}\Phi(v)/2}, V_v^r \right] - \frac{1}{i\hbar} e^{-i\bar{\mu}\Phi(v)/2} \left[e^{i\bar{\mu}\Phi(v)/2}, V_v^r \right]\end{aligned}$$

where the second equalities used $[\bar{\lambda}, V_v] = 0 = [\bar{\mu}, V_v]$. A straightforward calculation gives that

$$\hat{\mathcal{Q}}_{\theta,\bar{\lambda}}^{(r)}(v) = \hat{\mathcal{Q}}_{\Phi,\bar{\mu}}^{(r)}(v) \equiv \hat{\mathcal{Q}}^{(r)}(v) \quad (4.17)$$

with

$$\hat{\mathcal{Q}}^{(r)}(v) |\vec{\zeta}, \vec{\mu}\rangle = \frac{(4\pi\ell_p^3\beta\sqrt{\Delta})^r}{i\hbar} \mathcal{B}(\zeta(v), \mu(v), r) |\vec{\zeta}, \vec{\mu}\rangle$$

and

$$\mathcal{B}(\zeta(v), \mu(v), r) = \left| \mu(v)\zeta(v) + \frac{1}{2} \right|^r - \left| \mu(v)\zeta(v) - \frac{1}{2} \right|^r \quad (4.18)$$

Therefore, the inverse volume operator reads

$$\widehat{V}_v^{-1} |\zeta(v), \mu(v)\rangle = \left(\frac{27}{4\pi\ell_p^3\sqrt{\Delta}\beta} \right) B(\zeta(v)\mu(v)) |\zeta(v), \mu(v)\rangle \quad (4.19)$$

with

$$B(x) = |x| \left| |x + 1/2|^{1/3} - |x - 1/2|^{1/3} \right|^3. \quad (4.20)$$

Taking advantage of the blocks defined in the last section, we can now quantize the expression (4.3) to get the operator $\hat{\mathfrak{h}}_{\Delta,v}$

$$\begin{aligned}\hat{\mathfrak{h}}_{\Delta,v} &= \sqrt{\widehat{V}_v} \frac{\sin(\widehat{\bar{\mu}\Phi(v)}) \sin(\widehat{\bar{\lambda}\theta(v)})}{4\pi\beta^2\Delta\ell_p^2} \sqrt{\widehat{V}_v} \\ &+ \sqrt{\widehat{V}_v} \frac{\sin^2(\widehat{\bar{\mu}\Phi(v)})}{8\pi\beta^2\Delta\ell_p^2} \sqrt{\widehat{V}_v} + 2\pi\widehat{V}_v^{-1} \hat{\Pi}(v)^2 + \frac{\pi}{2} \widehat{V}_v^{-1} [\hat{p}(v+1) - \hat{p}(v)]^2.\end{aligned} \quad (4.21)$$

With this expression, the physical Hamiltonian operator $\widehat{\mathbf{H}}_{\Delta}$ reads

$$\widehat{\mathbf{H}}_{\Delta} = -\frac{1}{G} \sum_v \hat{\mathfrak{h}}_{\Delta,v}. \quad (4.22)$$

Since \widehat{V}_v annihilates $|\vec{\lambda}, \vec{\mu}\rangle$ when $\lambda(v) = 0$ or $\mu(v) = 0$, the operator orderings in the first and second terms ensure that $\hat{\mathfrak{h}}_{\Delta,v}$ is densely defined on the entire \mathcal{H} .

B. The action of the physical Hamiltonian operator

It is sufficient to present the action of $\hat{\mathfrak{h}}_{\Delta,v}$ instead of $\widehat{\mathbf{H}}_{\Delta}$ itself. To calculate the action, we convert the trigonometric functions in (4.21) to exponentials and define

$$\hat{\mathfrak{h}}_{\Delta,v} = \hat{\mathfrak{h}}_1(v) + \hat{\mathfrak{h}}_2(v) + \hat{\mathfrak{h}}_3(v) + \hat{\mathfrak{h}}_4(v), \quad (4.23)$$

with

$$\begin{aligned}
\hat{\mathfrak{h}}_1(v) &= -\sqrt{\hat{V}_v} \frac{(\hat{h}_{\Phi,\Delta}(v) - \hat{h}_{\Phi,\Delta}(v)^{-1}) (\hat{h}_{\theta,\Delta}(v) - \hat{h}_{\theta,\Delta}(v)^{-1})}{16\pi\beta^2\Delta\ell_p^2} \sqrt{\hat{V}_v} \\
\hat{\mathfrak{h}}_2(v) &= -\sqrt{\hat{V}_v} \frac{(\hat{h}_{\Phi,\Delta}(v) - \hat{h}_{\Phi,\Delta}(v)^{-1})^2}{32\pi\beta^2\Delta\ell_p^2} \sqrt{\hat{V}_v} \\
\hat{\mathfrak{h}}_3(v) &= 2\pi\widehat{V}_v^{-1}\hat{\Pi}(v)^2 \\
\hat{\mathfrak{h}}_4(v) &= \frac{\pi}{2}\widehat{V}_v^{-1} [\hat{p}(v+1) - \hat{p}(v)]^2.
\end{aligned} \tag{4.24}$$

Consider the action of operators $\hat{h}_{\Phi,\Delta}(v)^m \hat{h}_{\theta,\Delta}(v)^n$ and $\hat{h}_{\theta,\Delta}(v)^n \hat{h}_{\Phi,\Delta}(v)^m$ at first. We have

$$\begin{aligned}
\hat{h}_{\Phi,\Delta}(v)^m \hat{h}_{\theta,\Delta}(v)^n |\zeta(v), \mu(v)\rangle &= \left| \zeta(v) + \frac{n}{\mu(v)}, \frac{\mu(v)\zeta(v) + n + m}{\zeta(v) + n/\mu(v)} \right\rangle \\
\hat{h}_{\theta,\Delta}(v)^n \hat{h}_{\Phi,\Delta}(v)^m |\zeta(v), \mu(v)\rangle &= \left| \frac{\mu(v)\zeta(v) + m + n}{\mu(v) + m/\zeta(v)}, \mu(v) + \frac{m}{\zeta(v)} \right\rangle.
\end{aligned} \tag{4.25}$$

Because $\hat{h}_{\Phi,\Delta}(v)$ and $\hat{h}_{\theta,\Delta}(v)$ shift only the components of $\vec{\zeta}$ and $\vec{\mu}$ at v , i.e. $\zeta(v)$ and $\mu(v)$, $\hat{\mathfrak{h}}_1(v)$ and $\hat{\mathfrak{h}}_2(v)$ preserve the values of $\zeta(v')$ and $\mu(v')$ with $v' \neq v$. Thus we are only concerned on the actions of $\hat{\mathfrak{h}}_1(v)$ and $\hat{\mathfrak{h}}_2(v)$ on $|\zeta(v), \mu(v)\rangle$. A straightforward calculation gives that

$$\begin{aligned}
&\hat{\mathfrak{h}}_1(v)|\zeta(v), \mu(v)\rangle \\
&= -\frac{\ell_p}{4\beta\sqrt{\Delta}} \left(\sqrt{|\mu(v)\zeta(v)||\mu(v)\zeta(v) + 2|} \left| \zeta(v) + \frac{1}{\mu(v)}, \frac{\mu(v)\zeta(v) + 2}{\zeta(v) + 1/\mu(v)} \right\rangle \right. \\
&\quad - |\zeta(v)\mu(v)| \left| \zeta(v) - \frac{1}{\mu(v)}, \frac{\mu(v)\zeta(v)}{\zeta(v) - 1/\mu(v)} \right\rangle - |\zeta(v)\mu(v)| \left| \zeta(v) + \frac{1}{\mu(v)}, \frac{\mu(v)\zeta(v)}{\zeta(v) + 1/\mu(v)} \right\rangle \\
&\quad \left. + \sqrt{|\mu(v)\zeta(v)||\mu(v)\zeta(v) - 2|} \left| \zeta(v) - \frac{1}{\mu(v)}, \frac{\mu(v)\zeta(v) - 2}{\zeta(v) - 1/\mu(v)} \right\rangle \right).
\end{aligned} \tag{4.26}$$

Since $\hat{\mathfrak{h}}_1(v)$ is not symmetric, we also need to consider its adjoint $\hat{\mathfrak{h}}_1(v)^\dagger$, whose action reads

$$\begin{aligned}
&\hat{\mathfrak{h}}_1(v)^\dagger |\zeta(v), \mu(v)\rangle \\
&= -\frac{\ell_p}{4\beta\sqrt{\Delta}} \left(\sqrt{|\mu(v)\zeta(v)||\mu(v)\zeta(v) + 2|} \left| \frac{\mu(v)\zeta(v) + 2}{\mu(v) + 1/\zeta(v)}, \mu(v) + \frac{1}{\zeta(v)} \right\rangle \right. \\
&\quad - |\zeta(v)\mu(v)| \left| \frac{\mu(v)\zeta(v)}{\mu(v) + 1/\zeta(v)}, \mu(v) + \frac{1}{\zeta(v)} \right\rangle - |\zeta(v)\mu(v)| \left| \frac{\mu(v)\zeta(v)}{\mu(v) - 1/\zeta(v)}, \mu(v) - \frac{1}{\zeta(v)} \right\rangle \\
&\quad \left. + \sqrt{|\mu(v)\zeta(v)||\mu(v)\zeta(v) - 2|} \left| \frac{\mu(v)\zeta(v) - 2}{\mu(v) - 1/\zeta(v)}, \mu(v) - \frac{1}{\zeta(v)} \right\rangle \right).
\end{aligned} \tag{4.27}$$

For the operator $\mathfrak{h}_2(v)$, we have

$$\begin{aligned}
&\hat{\mathfrak{h}}_2(v)|\zeta(v), \mu(v)\rangle \\
&= -\frac{\ell_p}{8\beta\sqrt{\Delta}} \left(\sqrt{|\mu(v)\zeta(v)||\mu(v)\zeta(v) + 2|} \left| \zeta(v), \mu(v) + \frac{2}{\zeta(v)} \right\rangle - 2|\mu(v)\zeta(v)| \left| \zeta(v), \mu(v) \right\rangle + \right. \\
&\quad \left. \sqrt{|\mu(v)\zeta(v)||\mu(v)\zeta(v) - 2|} \left| \zeta(v), \mu(v) - \frac{2}{\zeta(v)} \right\rangle \right)
\end{aligned} \tag{4.28}$$

For the operators $\mathfrak{h}_3(v)$ and $\mathfrak{h}_4(v)$, the basis $|\vec{\mu}, \zeta\rangle$ are their eigenvectors. Their actions is

$$\hat{\mathfrak{h}}_3(v)|\zeta(v), \mu(v)\rangle = 27\ell_p \frac{\mu(v)^2}{2\beta\sqrt{\Delta}} B(\zeta(v)\mu(v)) |\zeta(v), \mu(v)\rangle, \tag{4.29}$$

and

$$\hat{\mathfrak{h}}_4(v)|\vec{\zeta}, \vec{\mu}\rangle = \frac{27\ell_p\beta^3\sqrt{\Delta}^3}{8}B(\zeta(v)\mu(v)) [\text{sgn}(\zeta((v+1))\zeta(v+1)^2 - \text{sgn}(\zeta(v))\zeta(v)^2)]^2 |\vec{\zeta}, \vec{\mu}\rangle. \quad (4.30)$$

Substituting the expression (4.26), (4.28), (4.29) and (4.30) into (4.25), one finally can get the action of $\hat{\mathfrak{h}}_{\Delta,v}$, which will not be written again.

It is worth noting that the quantity

$$\mathfrak{v}(v) := \zeta(v)\mu(v)$$

often appears in the RHSs of (4.26)–(4.30) as a whole. Then, these equations can be simplified in more compact forms. To do this, we will use $\vec{\mathfrak{v}} = \{\mathfrak{v}(v)\}_{v \in V(\gamma)}$ instead of $\vec{\mu}$ to re-label the basis $|\vec{\zeta}, \vec{\mu}\rangle$. However, this re-labeling can only be defined for those $|\vec{\zeta}, \vec{\mu}\rangle$ satisfying $\zeta(v) \neq 0$ for all v , because this condition guarantees the solvability of $\mu(v)$ through $\mathfrak{v}(v) = \zeta(v)\mu(v)$. Given a state $|\vec{\zeta}, \vec{\mathfrak{v}}\rangle$ with $\zeta(v) \neq 0, \forall v$, it has

$$\text{sgn}(\hat{p}(v))\sqrt{|\hat{p}(v)|}|\vec{\zeta}, \vec{\mathfrak{v}}\rangle = \beta\sqrt{\Delta}\ell_p\zeta(v)|\vec{\zeta}, \vec{\mathfrak{v}}\rangle, \quad \hat{V}_v|\vec{\zeta}, \vec{\mathfrak{v}}\rangle = 4\pi\beta\sqrt{\Delta}\ell_p^3|\mathfrak{v}(v)||\vec{\zeta}, \vec{\mathfrak{v}}\rangle \quad (4.31)$$

According to (4.14), one has

$$\begin{aligned} \hat{h}_{\theta,\Delta}(v)|\vec{\zeta}, \vec{\mathfrak{v}}\rangle &= |\vec{\zeta} + \zeta(v)\mathfrak{v}(v)^{-1}\vec{\delta}_v, \vec{\mathfrak{v}} + \vec{\delta}_v\rangle \\ \hat{h}_{\Phi,\Delta}(v)|\vec{\zeta}, \vec{\mathfrak{v}}\rangle &= |\vec{\zeta}, \vec{\mathfrak{v}} + \vec{\delta}_v\rangle. \end{aligned} \quad (4.32)$$

By this equation, the $\vec{\mu}$ -scheme holonomies are not densely defined on \mathcal{H} , since they cannot act on state $|\vec{\zeta}, \vec{\mathfrak{v}}\rangle$ when $\mathfrak{v}(v) = 0$. Moreover, they are not commutative:

$$\hat{h}_{\theta,\Delta}(v), \hat{h}_{\Phi,\Delta}(v)t|\vec{\zeta}, \vec{\mathfrak{v}}\rangle = |\vec{\zeta} + \zeta(v)(\mathfrak{v}(v) + 1)^{-1}\vec{\delta}_v, \vec{\mathfrak{v}} + 2\vec{\delta}_v\rangle - |\vec{\zeta} + \zeta(v)\mathfrak{v}(v)^{-1}\vec{\delta}_v, \vec{\mathfrak{v}} + 2\vec{\delta}_v\rangle. \quad (4.33)$$

Even though these holonomies $\hat{h}_{\theta,\Delta}(v)$ and $\hat{h}_{\Phi,\Delta}(v)$ are not well-defined for $|\vec{\zeta}, \vec{\mathfrak{v}}\rangle$ with $\mathfrak{v}(v) = 0$, the Hamiltonian operator $\hat{\mathfrak{h}}_{\Delta,v}$ does. For $\mathfrak{v}(v) = 0$, it has

$$\hat{\mathfrak{h}}_{\Delta,v}|\vec{\zeta}, \vec{\mathfrak{v}}\rangle = 0.$$

For $\mathbf{v}(v) \neq 0$, the formulas (4.26), (4.28), (4.29) and (4.30) can be rewritten as

$$\begin{aligned} \hat{\mathbf{h}}_1(v)|\zeta(v), \mathbf{v}(v)\rangle &= -\frac{\ell_p}{4\beta\sqrt{\Delta}} \left(\sqrt{|\mathbf{v}(v)||\mathbf{v}(v)+2|} \left| \zeta(v) \frac{\mathbf{v}(v)+1}{\mathbf{v}(v)}, \mathbf{v}(v)+2 \right\rangle \right. \\ &\quad - |\mathbf{v}(v)| \left| \zeta(v) \frac{\mathbf{v}(v)-1}{\mathbf{v}(v)}, \mathbf{v}(v) \right\rangle - |\mathbf{v}(v)| \left| \zeta(v) \frac{\mathbf{v}(v)+1}{\mathbf{v}(v)}, \mathbf{v}(v) \right\rangle \\ &\quad \left. + \sqrt{|\mathbf{v}(v)||\mathbf{v}(v)-2|} \left| \zeta(v) \frac{\mathbf{v}(v)-1}{\mathbf{v}(v)}, \mathbf{v}(v)-2 \right\rangle \right) \end{aligned} \quad (4.34)$$

$$\begin{aligned} \hat{\mathbf{h}}_1(v)^\dagger|\zeta(v), \mathbf{v}(v)\rangle &= -\frac{\ell_p}{4\beta\sqrt{\Delta}} \left(\sqrt{|\mathbf{v}(v)||\mathbf{v}(v)+2|} \left| \zeta(v) \frac{\mathbf{v}(v)+2}{\mathbf{v}(v)+1}, \mathbf{v}(v)+2 \right\rangle \right. \\ &\quad - |\mathbf{v}(v)| \left| \zeta(v) \frac{\mathbf{v}(v)}{\mathbf{v}(v)+1}, \mathbf{v}(v) \right\rangle - |\mathbf{v}(v)| \left| \zeta(v) \frac{\mathbf{v}(v)}{\mathbf{v}(v)-1}, \mathbf{v}(v) \right\rangle \\ &\quad \left. + \sqrt{|\mathbf{v}(v)||\mathbf{v}(v)-2|} \left| \zeta(v) \frac{\mathbf{v}(v)-2}{\mathbf{v}(v)-1}, \mathbf{v}(v)-2 \right\rangle \right) \end{aligned} \quad (4.35)$$

$$\begin{aligned} \hat{\mathbf{h}}_2(v)|\zeta(v), \mathbf{v}(v)\rangle &= -\frac{\ell_p}{8\beta\sqrt{\Delta}} \left(\sqrt{|\mathbf{v}(v)||\mathbf{v}(v)+2|} |\zeta(v), \mathbf{v}(v)+2\rangle - \right. \\ &\quad \left. 2|\mathbf{v}(v)| |\zeta(v), \mathbf{v}(v)\rangle + \sqrt{|\mathbf{v}(v)||\mathbf{v}(v)-2|} |\zeta(v), \mathbf{v}(v)-2\rangle \right) \end{aligned} \quad (4.36)$$

$$\hat{\mathbf{h}}_3(v)|\zeta(v), \mathbf{v}(v)\rangle = \frac{27\ell_p}{2\beta\sqrt{\Delta}} \frac{\mathbf{v}(v)^2}{\zeta(v)^2} B(\mathbf{v}(v)) |\zeta(v), \mathbf{v}(v)\rangle \quad (4.37)$$

$$\begin{aligned} \hat{\mathbf{h}}_4(v)|\vec{\zeta}, \vec{\mathbf{v}}\rangle &= \frac{27\ell_p\beta^3\sqrt{\Delta}^3}{8} B(\mathbf{v}(v)) \left[\text{sgn}(\zeta((v+1))\zeta(v+1))^2 - \right. \\ &\quad \left. \text{sgn}(\zeta(v))\zeta(v)^2 \right]^2 |\vec{\zeta}, \vec{\mathbf{v}}\rangle. \end{aligned} \quad (4.38)$$

V. PATH INTEGRAL FORMULATION

Let us define $\hat{h}_i = \sum_v \hat{h}_i(v)$. By (4.34)–(4.38), one has the matrix element of \hat{h}_i

$$\langle \vec{\zeta}_1, \vec{\mathbf{v}}_1 | \hat{h}_1 | \vec{\zeta}_2, \vec{\mathbf{v}}_2 \rangle = -\frac{\ell_p}{4\beta\sqrt{\Delta}} \sum_v \prod_{v' \neq v} \delta_{\zeta_1(v'), \zeta_2(v')} \delta_{\mathbf{v}_1(v'), \mathbf{v}_2(v')} \sqrt{|\mathbf{v}_1(v) \mathbf{v}_2(v)|} \times \quad (5.1)$$

$$\left(\delta_{\zeta_1(v)(\mathbf{v}_1(v)-2), \zeta_2(v)(\mathbf{v}_2(v)+1)} \delta_{\mathbf{v}_1(v), \mathbf{v}_2(v)+2} - \delta_{\zeta_1(v) \mathbf{v}_1(v), \zeta_2(v)(\mathbf{v}_2(v)-1)} \delta_{\mathbf{v}_1(v), \mathbf{v}_2(v)} - \delta_{\zeta_1(v) \mathbf{v}_1(v), \zeta_2(v)(\mathbf{v}_2(v)+1)} \delta_{\mathbf{v}_1(v), \mathbf{v}_2(v)} + \delta_{\zeta_1(v)(\mathbf{v}_1(v)+2), \zeta_2(v)(\mathbf{v}_2(v)-1)} \delta_{\mathbf{v}_1(v), \mathbf{v}_2(v)-2} \right)$$

$$\langle \vec{\zeta}_1, \vec{\mathbf{v}}_1 | \hat{h}_1^\dagger | \vec{\zeta}_2, \vec{\mathbf{v}}_2 \rangle = -\frac{\ell_p}{4\beta\sqrt{\Delta}} \sum_v \prod_{v' \neq v} \delta_{\zeta_1(v'), \zeta_2(v')} \delta_{\mathbf{v}_1(v'), \mathbf{v}_2(v')} \sqrt{|\mathbf{v}_2(v)| |\mathbf{v}_1(v)|} \times \quad (5.2)$$

$$\left(\delta_{\zeta_1(v)(\mathbf{v}_1(v)-1), \zeta_2(v)(\mathbf{v}_2(v)+2)} \delta_{\mathbf{v}_1(v), \mathbf{v}_2(v)+2} - \delta_{\zeta_1(v)(\mathbf{v}_1(v)+1), \zeta_2(v) \mathbf{v}_2(v)} \delta_{\mathbf{v}_1(v), \mathbf{v}_2(v)} - \delta_{\zeta_1(v)(\mathbf{v}_1(v)-1), \zeta_2(v) \mathbf{v}_2(v)} \delta_{\mathbf{v}_1(v), \mathbf{v}_2(v)} + \delta_{\zeta_1(v)(\mathbf{v}_1(v)+1), \zeta_2(v)(\mathbf{v}_2(v)-2)} \delta_{\mathbf{v}_1(v), \mathbf{v}_2(v)-2} \right)$$

$$\langle \vec{\zeta}_1, \vec{\mathbf{v}}_1 | \hat{h}_2 | \vec{\zeta}_2, \vec{\mathbf{v}}_2 \rangle = -\frac{\ell_p}{8\beta\sqrt{\Delta}} \sum_v \prod_{v' \neq v} \delta_{\zeta_1(v'), \zeta_2(v')} \delta_{\mathbf{v}_1(v'), \mathbf{v}_2(v')} \sqrt{|\mathbf{v}_1(v)| |\mathbf{v}_2(v)|} \times \quad (5.3)$$

$$\left(\delta_{\zeta_1(v), \zeta_2(v)} \delta_{\mathbf{v}_1(v), \mathbf{v}_2(v)+2} - 2\delta_{\zeta_1(v), \zeta_2(v)} \delta_{\mathbf{v}_1(v), \mathbf{v}_2(v)} + \delta_{\zeta_1(v), \zeta_2(v)} \delta_{\mathbf{v}_1(v), \mathbf{v}_2(v)-2} \right)$$

$$\langle \vec{\zeta}_1, \vec{\mathbf{v}}_1 | \hat{h}_3 | \vec{\zeta}_2, \vec{\mathbf{v}}_2 \rangle = \frac{27\ell_p}{2\beta\sqrt{\Delta}} \sum_v \frac{\mathbf{v}_2(v)^2}{\zeta_2(v)^2} B(\mathbf{v}_2(v)) \prod_{v'} \delta_{\zeta_1(v'), \zeta_2(v')} \delta_{\mathbf{v}_1(v'), \mathbf{v}_2(v')} \quad (5.4)$$

$$\langle \vec{\zeta}_1, \vec{\mathbf{v}}_1 | \hat{h}_4 | \vec{\zeta}_2, \vec{\mathbf{v}}_2 \rangle = \frac{27\ell_p\beta^3\sqrt{\Delta}^3}{8} \sum_v B(\mathbf{v}_2(v)) \left[\text{sgn}(\zeta_2((v+1))\zeta_2(v+1)^2 - \text{sgn}(\zeta_2(v))\zeta_2(v)^2) \right]^2 \quad (5.5)$$

$$\prod_{v'} \delta_{\zeta_1(v'), \zeta_2(v')} \delta_{\mathbf{v}_1(v'), \mathbf{v}_2(v')}.$$

In order to simplify the matrix elements of \hat{h}_i , consider the space of almost periodic functions and define a functional μ_h of these functions by

$$\mu_h(f) \equiv \int_{\mathbb{R}} d\mu_h(x) f(x) := \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T f(x) dx \quad (5.6)$$

Then, it can be verified that

$$\delta_{\lambda,0} = \int_{\mathbb{R}} d\mu_h(x) e^{i\lambda x}, \quad (5.7)$$

where $\delta_{\lambda,0}$ is the Kronecker delta. This formula provide an approach to rewrite the Kronecker-delta functions appearing in the expressions of the matrix elements of \hat{h}_i , so that we can simplify these expressions to get the final path integral formula. Before proceeding further, let us consider an issue on how to deal with the delta functions taking the form $\delta_{A(\mathbf{v}_1(v)) \zeta_1(v), B(\mathbf{v}_2(v)) \zeta_2(v)}$, which appears in the expressions (5.1)–(5.5). The deep idea is to re-express $\delta_{A(\mathbf{v}_1(v)) \zeta_1(v), B(\mathbf{v}_2(v)) \zeta_2(v)}$ such that it takes the form $\delta_{f(\zeta_1(v)) - f(\zeta_2(v)), g}$ with g independent of $\zeta_1(v)$ and $\zeta_2(v)$. Then, Lemma V.2 stated below can be used to rewrite the transition amplitude in the standard path integral form, i.e., the form in present of the Lebesgue measure. Thus, we need the identity

$$\delta_{A(\mathbf{v}_1(v)) \zeta_1(v), B(\mathbf{v}_2(v)) \zeta_2(v)} = \delta_{\text{sgn}(\zeta_1(v)), \text{sgn}(\zeta_2(v))} \delta_{\ln(|\zeta_2(v)|) - \ln(|\zeta_1(v)|), \ln(|B(\mathbf{v}_1(v))|/A(\mathbf{v}_2(v)))}. \quad (5.8)$$

Taking advantage of this identity and (5.7), we finally get

$$\langle \vec{\zeta}_1, \vec{\mathbf{v}}_1 | \sum_v \hat{h}_{\Delta, v} | \vec{\zeta}_2, \vec{\mathbf{v}}_2 \rangle = \left[\prod_v \delta_{\text{sgn}(\zeta_1(v)), \text{sgn}(\zeta_2(v))} \right] \int_{\mathbb{R}^{2|\mathcal{V}(\gamma)}} \prod_{v'} d\mu_h(b(v')) d\mu_h(c(v')) \times \quad (5.9)$$

$$\exp \left[\sum_{v'} i [\ln(\zeta_2(v')) - \ln(\zeta_1(v'))] c(v') + i(\mathbf{v}_2(v') - \mathbf{v}_1(v')) b(v') \right] H(\vec{\zeta}_1, \vec{\mathbf{v}}_1, \vec{\zeta}_2, \vec{\mathbf{v}}_2, \vec{b}, \vec{c})$$

where

$$\begin{aligned}
H(\vec{\zeta}_1, \vec{\mathbf{v}}_1, \vec{\zeta}_2, \vec{\mathbf{v}}_2, \vec{b}, \vec{c}) &= \sum_v \left\{ \left(-\frac{\ell_p \sqrt{|\mathbf{v}_1(v)\mathbf{v}_2(v)|}}{8\beta\sqrt{\Delta}} \right) \left(\exp \left[i \ln \left(\left| 1 + \frac{1}{\mathbf{v}_2(v)} \right| \right) c(v) + 2ib(v) \right] \right. \right. \\
&- \exp \left[i \ln \left(\left| 1 - \frac{1}{\mathbf{v}_2(v)} \right| \right) c(v) \right] - \exp \left[i \ln \left(\left| 1 + \frac{1}{\mathbf{v}_2(v)} \right| \right) c(v) \right] + \exp \left[i \ln \left(\left| 1 - \frac{1}{\mathbf{v}_2(v)} \right| \right) c(v) - 2ib(v) \right] \Big) + \\
&\left(-\frac{\ell_p \sqrt{|\mathbf{v}_1(v)\mathbf{v}_2(v)|}}{8\beta\sqrt{\Delta}} \right) \left(\exp \left[i \ln \left(\left| 1 + \frac{1}{\mathbf{v}_1(v)-1} \right| \right) c(v) + 2ib(v) \right] - \exp \left[i \ln \left(\left| 1 - \frac{1}{\mathbf{v}_1(v)+1} \right| \right) c(v) \right] \right. \\
&- \exp \left[i \ln \left(\left| 1 + \frac{1}{\mathbf{v}_1(v)-1} \right| \right) c(v) \right] + \exp \left[i \ln \left(\left| 1 - \frac{1}{\mathbf{v}_1(v)+1} \right| \right) c(v) - 2ib(v) \right] \Big) + \\
&\frac{\ell_p \sqrt{|\mathbf{v}_1(v)\mathbf{v}_2(v)|}}{2\beta\sqrt{\Delta}} \sin^2(b(v)) + \frac{27\ell_p \mathbf{v}_2(v)^2}{2\zeta_2(v)^2 \beta \sqrt{\Delta}} B(\mathbf{v}_2(v)) + \\
&\left. \frac{27\ell_p \beta^3 \sqrt{\Delta}^3}{8} B(\mathbf{v}_2(v)) \left[\text{sgn}(\zeta_2((v+1))\zeta_2(v+1)^2 - \text{sgn}(\zeta_2(v))\zeta_2(v)^2) \right]^2 \right\}. \tag{5.10}
\end{aligned}$$

Then the matrix element of $e^{-i\epsilon\hat{\mathbf{h}}}$, with $\hat{\mathbf{h}}_\Delta := \sum_v \hat{\mathbf{h}}_{\Delta,v}$, reads

$$\begin{aligned}
\langle \vec{\zeta}_1, \vec{\mathbf{v}}_1 | e^{-i\epsilon\hat{\mathbf{h}}_\Delta} | \vec{\zeta}_2, \vec{\mathbf{v}}_2 \rangle &= \langle \vec{\zeta}_1, \vec{\mathbf{v}}_1 | (1 - i\epsilon\hat{\mathbf{h}}) | \vec{\zeta}_2, \vec{\mathbf{v}}_2 \rangle \\
&= \left[\prod_v \delta_{\text{sgn}(\zeta_1(v)), \text{sgn}(\zeta_2(v))} \right] \int_{\mathbb{R}^{2|V(\gamma)}} \prod_{v'} d\mu_h(b(v')) d\mu_h(c(v')) \\
&\exp \left[\sum_{v'} i(\ln(|\zeta_2(v')|) - \ln(|\zeta_1(v')|))c(v') + i(\mathbf{v}_2(v') - \mathbf{v}_1(v'))b(v') \right] \times \\
&\left[1 - i\epsilon H(\vec{\zeta}_1, \vec{\mathbf{v}}_1, \vec{\zeta}_2, \vec{\mathbf{v}}_2, \vec{b}, \vec{c}) \right] + O(\epsilon^2). \tag{5.11}
\end{aligned}$$

To simplify the last equation, we claim the following theorem

Theorem V.1. *Given $F(\{\ln(|\zeta(v)|), \mathbf{v}(v)\}_{v \in V(\gamma)})$ a Schwartz function. Then*

$$\begin{aligned}
\tilde{F}(\{\ln(|\zeta(v)|), \mathbf{v}(v)\}_{v \in V(\gamma)}) &:= \sum_{\{\ln(|\zeta(v)|), \mathbf{v}(v)\}_{v \in V(\gamma)}} \langle \vec{\zeta}_1, \vec{\mathbf{v}}_1 | (1 - i\epsilon\hat{\mathbf{h}}) | \vec{\zeta}_2, \vec{\mathbf{v}}_2 \rangle F(\{\ln(|\zeta(v)|), \mathbf{v}(v)\}_{v \in V(\gamma)}) \\
&= \left[\prod_v \delta_{\text{sgn}(\zeta_1(v)), \text{sgn}(\zeta_2(v))} \right] \int_{\mathbb{R}^{4|V(\gamma)}} \prod_{v'} db(v') dc(v') \frac{d\zeta(v')}{\zeta(v')} d\mathbf{v}(v') \\
&\exp \left[\sum_{v'} i(\ln(|\zeta_2(v')|) - \ln(|\zeta_1(v')|))c(v') + i(\mathbf{v}_2(v') - \mathbf{v}_1(v'))b(v') \right] \times \\
&\left(1 - i\epsilon H(\vec{\zeta}_1, \vec{\mathbf{v}}_1, \vec{\zeta}_2, \vec{\mathbf{v}}_2, \vec{b}, \vec{c}) \right) F(\{\ln(|\zeta(v)|), \mathbf{v}(v)\}_{v \in V(\gamma)}) \tag{5.12}
\end{aligned}$$

with $db(v')$, $dc(v')$, $d\zeta(v')$, $d\mathbf{v}(v')$ being the Lebesgue measure.

This theorem can be deduced straightforwardly by the following lemmas V.2 and V.3.

Lemma V.2. *Given a function*

$$f(x, y) = \sum_{\vec{\lambda} \in \mathcal{L}} \mu(\vec{\lambda}, y) e^{ig(\vec{\lambda})x} \tag{5.13}$$

where $\mathcal{L} \subset \mathbb{R}^N$ is a finite lattice in \mathbb{R}^N , $g(\vec{\lambda})$ is a real function of $\vec{\lambda}$, and $\mu(\vec{\lambda}, y)$ satisfies that $\mu(\vec{\lambda}, \cdot)$ is a Schwartz function for each $\vec{\lambda}$. Then

$$\sum_y \int d\mu_h(x) e^{-i(y+\alpha)x} f(x, y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy e^{-i(y+\alpha)x} f(x, y). \tag{5.14}$$

with α being a constant.

Proof. By definition,

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy e^{-i(y+\alpha)x} f(x, y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \sum_{\vec{\lambda} \in \mathcal{L}} \mu(\vec{\lambda}, y) e^{i(g(\vec{\lambda})-y-\alpha)x} \quad (5.15)$$

Because \mathcal{L} is a finite lattice, one exchanges the order between the summation and the integrals to get

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy e^{-i(y+\alpha)x} f(x, y) = \frac{1}{2\pi} \sum_{\vec{\lambda} \in \mathcal{L}} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \mu(\vec{\lambda}, y) e^{i(g(\vec{\lambda})-y-\alpha)x}. \quad (5.16)$$

Since $\mu(\lambda, \cdot)$ is Schwartz function for each λ , by using the inverse Fourier inversion theorem, we have

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \mu(\vec{\lambda}, y) e^{i(g(\vec{\lambda})-y-\alpha)x} \quad (5.17)$$

$$= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dx e^{i(g(\vec{\lambda})-\alpha)x} \left(\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dy \mu(\vec{\lambda}, y) e^{-ixy} \right) \quad (5.18)$$

$$= \mu(\vec{\lambda}, g(\vec{\lambda}) - \alpha). \quad (5.19)$$

Thus

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy e^{-i(y+\alpha)x} f(x, y) = \sum_{\vec{\lambda} \in \mathcal{L}} \mu(\vec{\lambda}, g(\vec{\lambda}) - \alpha) \quad (5.20)$$

where the RHS is just $\sum_y \int d\mu_h(x) e^{-ixy} f(x, y)$ by definition of $\mu_h(x)$. \square

Lemma V.3. *Given a function $\mathcal{H}(x, y)$ taking the form*

$$\mathcal{H}(x, y_1, y_2) = \sum_{k=1}^N f_k(y_1, y_2) e^{ig_k x}, \quad (5.21)$$

where we assume that $f_k(y_1, y_2)$ are continuous and, there exist some α and β such that

$$\frac{f_k(y_1, y_2)}{(1 + (y_1)^2)^\alpha (1 + (y_2)^2)^\beta} < \infty \quad (5.22)$$

Let $F(y) \in \mathcal{S}(\mathbb{R})$, with $\mathcal{S}(\mathbb{R})$ being the space of Schwartz functions. Then

$$\begin{aligned} \tilde{F}(y_1) &= \sum_{y_2} \int d\mu_h(x) e^{-i(y_2-y_1)x} (1 + i\epsilon \mathcal{H}(x, y_1, y_2)) F(y_2) \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy_2 (1 + i\epsilon \mathcal{H}(x, y_1, y_2)) F(y_2) \end{aligned} \quad (5.23)$$

To prove this lemma, one only needs to verify $\mu(k, y_1, \cdot) \in \mathcal{S}(\mathbb{R})$ by (5.22). Then, Lemma V.2 can be applied to get (5.23).

According to Theorem V.1, $\langle \vec{\zeta}_1, \vec{\mathbf{v}}_1 | e^{i\epsilon \hat{h}} | \vec{\zeta}_2, \vec{\mathbf{v}}_2 \rangle$ can be written as

$$\begin{aligned} &\langle \vec{\zeta}_1, \vec{\mathbf{v}}_1 | e^{i\epsilon \hat{h}} | \vec{\zeta}_2, \vec{\mathbf{v}}_2 \rangle \\ &= \left[\prod_v \delta_{\text{sgn}(\zeta_1(v)), \text{sgn}(\zeta_2(v))} \right] \left(\frac{1}{2\pi} \right)^{|\mathcal{V}(\gamma)|} \int_{\mathbb{R}^{2|\mathcal{V}(\gamma)|}} \prod_{v'} db(v') dc(v') \\ &\exp \left[\sum_{v'} i(\ln(|\zeta_2(v')|) - \ln(|\zeta_1(v')|)) c(v') + i(\mathbf{v}_2(v') - \mathbf{v}_1(v')) b(v') \right] \times \\ &\left(1 + i\epsilon H(\vec{\zeta}_1, \vec{\mathbf{v}}_1, \vec{\zeta}_2, \vec{\mathbf{v}}_2, \vec{b}, \vec{c}) \right) + O(\epsilon^2). \end{aligned} \quad (5.24)$$

It is noted that the RHS is understood as a functional on the space of Schwartz functions of $\vec{\zeta}_1$ and \mathbf{v}_1 , as stated in Theorem V.1. With these formulas, let us consider the transition amplitude $A(\vec{\zeta}_i, \vec{\mathbf{v}}_i, \vec{\zeta}_f, \vec{\mathbf{v}}_f, T)$

$$\begin{aligned}
A(\vec{\zeta}_i, \vec{\mathbf{v}}_i, \vec{\zeta}_f, \vec{\mathbf{v}}_f, T) &= \langle \vec{\zeta}_f, \vec{\mathbf{v}}_f | e^{-i\frac{T}{G\hbar}\hat{b}} | \vec{\zeta}_i, \vec{\mathbf{v}}_i \rangle = \langle \vec{\zeta}_f, \vec{\mathbf{v}}_f | \left(e^{i\frac{T}{G\hbar N}\hat{b}} \right)^N | \vec{\zeta}_i, \vec{\mathbf{v}}_i \rangle \\
&= \lim_{N \rightarrow \infty} \sum_{\{\vec{\zeta}_n, \vec{\mathbf{v}}_n\}_{n=1}^N} \prod_{k=0}^N \langle \vec{\zeta}_{k+1}, \vec{\mathbf{v}}_{k+1} | e^{i\frac{T}{G\hbar N}\hat{b}} | \vec{\zeta}_k, \vec{\mathbf{v}}_k \rangle \\
&= \int \prod_{v'} \mathcal{D}[b(v')] \mathcal{D}[c(v')] \mathcal{D}[\ln(|\zeta(v')|)] \mathcal{D}[\mathbf{v}(v')] e^{i\frac{1}{\hbar}S(\vec{\zeta}, \vec{\mathbf{v}}, \vec{c}, \vec{b})}
\end{aligned} \tag{5.25}$$

where $\vec{\zeta}_0 = \vec{\zeta}_i$, $\vec{\zeta}_{N+1} = \vec{\zeta}_f$, $\vec{\mathbf{v}}_0 = \vec{\mathbf{v}}_i$ and $\vec{\mathbf{v}}_{N+1} = \vec{\mathbf{v}}_f$ and, with excluding the boundary term,

$$S(\vec{\zeta}, \vec{\mathbf{v}}, \vec{c}, \vec{b}) = \int_0^T dt \mathcal{L}(\vec{\zeta}, \vec{\mathbf{v}}, \vec{c}, \vec{b}) = \int_0^T dt \left(\sum_v -\frac{\hbar}{\zeta(v)} \frac{d\zeta(v)}{dt} c(v) - \hbar \frac{d\mathbf{v}(v)}{dt} b(v) + \frac{1}{G} H(\vec{\zeta}, \vec{\mathbf{v}}, \vec{c}, \vec{b}) \right) \tag{5.26}$$

$$\begin{aligned}
H(\vec{\zeta}, \vec{\mathbf{v}}, \vec{c}, \vec{b}) &= \sum_v \left\{ \left(-\frac{\ell_p |\mathbf{v}(v)|}{8\beta\sqrt{\Delta}} \right) \left(\exp \left[i \ln \left(\left| 1 + \frac{1}{\mathbf{v}(v)} \right| \right) c(v) + 2ib(v) \right] - \exp \left[i \ln \left(\left| 1 - \frac{1}{\mathbf{v}(v)} \right| \right) c(v) \right] \right. \right. \\
&\quad \left. \left. - \exp \left[i \ln \left(\left| 1 + \frac{1}{\mathbf{v}(v)} \right| \right) c(v) \right] + \exp \left[i \ln \left(\left| 1 - \frac{1}{\mathbf{v}(v)} \right| \right) c(v) - 2ib(v) \right] \right) + \right. \\
&\quad \left(-\frac{\ell_p |\mathbf{v}(v)|}{8\beta\sqrt{\Delta}} \right) \left(\exp \left[i \ln \left(\left| 1 + \frac{1}{\mathbf{v}(v) - 1} \right| \right) c(v) + 2ib(v) \right] - \exp \left[i \ln \left(\left| 1 - \frac{1}{\mathbf{v}(v) + 1} \right| \right) c(v) \right] \right. \\
&\quad \left. \left. - \exp \left[i \ln \left(\left| 1 + \frac{1}{\mathbf{v}(v) - 1} \right| \right) c(v) \right] + \exp \left[i \ln \left(\left| 1 - \frac{1}{\mathbf{v}(v) + 1} \right| \right) c(v) - 2ib(v) \right] \right) + \right. \\
&\quad \left. \left. \frac{|\mathbf{v}(v)|\ell_p}{2\beta\sqrt{\Delta}} \sin^2(b(v)) + \frac{27\ell_p \mathbf{v}(v)^2}{2\zeta(v)^2 \beta\sqrt{\Delta}} B(\mathbf{v}(v)) + \frac{27\ell_p \beta^3 \sqrt{\Delta}^3}{8} B(\mathbf{v}(v)) [\text{sgn}(\zeta((v+1))\zeta(v+1)^2 - \text{sgn}(\zeta(v))\zeta(v)^2]^2 \right\} \right. \\
&\quad \left. \right\} \tag{5.27}
\end{aligned}$$

VI. THE EFFECTIVE DYNAMICS

To consider the classical limit $\hbar \rightarrow 0$, we introduce the classical fields as

$$\begin{aligned}
\underline{\mathbf{v}}(v) &= 4\pi\beta\sqrt{\Delta}\ell_p^3 \mathbf{v}(v), \quad \underline{\zeta}(v) = \beta\sqrt{\Delta}\ell_p \zeta(v) \\
\underline{b}(v) &= \frac{1}{4\pi\beta\sqrt{\Delta}\ell_p} b(v), \quad \underline{c}(v) = \ell_p^2 c(v).
\end{aligned} \tag{6.1}$$

The non-vanishing Poisson brackets between these fields are

$$\{\underline{b}(v), \underline{\mathbf{v}}(v)\} = G, \quad \{\underline{c}(v), \underline{\zeta}(v)\} = G\underline{\zeta}(v). \tag{6.2}$$

In terms of the classical fields, H can be simplified as

$$\begin{aligned}
H(\vec{\zeta}, \vec{\mathbf{v}}, \vec{c}, \vec{b}) &= \sum_v \left\{ \left(-\frac{1}{16\pi\beta^2\Delta\ell_p^2} \right) |\mathbf{v}(v)| \times \right. \\
&\left(\cos \left[\ln \left(\left| 1 + \frac{4\pi\beta\sqrt{\Delta}\ell_p^3}{\mathbf{v}(v)} \right| \right) \frac{c(v)}{\ell_p^2} + 8\pi\beta\sqrt{\Delta}\ell_p b(v) \right] - \cos \left[\ln \left(\left| 1 - \frac{4\pi\beta\sqrt{\Delta}\ell_p^3}{\mathbf{v}(v)} \right| \right) \frac{c(v)}{\ell_p^2} \right] \right. \\
&\left. - \cos \left[\ln \left(\left| 1 + \frac{4\pi\beta\sqrt{\Delta}\ell_p^3}{\mathbf{v}(v)} \right| \right) \frac{c(v)}{\ell_p^2} \right] + \cos \left[\ln \left(\left| 1 - \frac{4\pi\beta\sqrt{\Delta}\ell_p^3}{\mathbf{v}(v)} \right| \right) \frac{c(v)}{\ell_p^2} - 8\pi\beta\sqrt{\Delta}\ell_p b(v) \right] \right) + \\
&\frac{|\mathbf{v}(v)|}{8\pi\beta^2\Delta\ell_p^2} \sin^2(4\pi\beta\sqrt{\Delta}\ell_p b(v)) + \frac{27\mathbf{v}(v)^2}{32\pi^2\beta\sqrt{\Delta}\zeta(v)^2\ell_p^3} B \left[\frac{\mathbf{v}(v)}{4\pi\beta\sqrt{\Delta}\ell_p^3} \right] + \\
&\left. \frac{27}{8\beta\sqrt{\Delta}\ell_p^3} B \left[\frac{\mathbf{v}(v)}{4\pi\beta\sqrt{\Delta}\ell_p^3} \right] \left[\text{sgn}(\zeta((v+1))\zeta(v+1)^2 - \text{sgn}(\zeta(v))\zeta(v)^2) \right]^2 \right\} \quad (6.3)
\end{aligned}$$

To investigate the effect of the holonomy correction but ignore the \hbar -order correction, we consider the limit $\ell_p \rightarrow 0$ but $\Delta\ell_p^2 = \text{constant}$. Then H can be simplified to the form

$$\begin{aligned}
H(\vec{\zeta}, \vec{\mathbf{v}}, \vec{c}, \vec{b}) &= \sum_v \left\{ \frac{1}{4\pi\beta^2\Delta\ell_p^2} |\mathbf{v}(v)| \sin \left[4\pi\beta\sqrt{\Delta}\ell_p b(v) \right] \sin \left[\frac{4\pi\beta\sqrt{\Delta}\ell_p}{\mathbf{v}(v)} c(v) + 4\pi\beta\sqrt{\Delta}\ell_p b(v) \right] + \right. \\
&\left. \frac{1}{8\pi\beta^2\Delta\ell_p^2} |\mathbf{v}(v)| \sin^2(4\pi\beta\sqrt{\Delta}\ell_p b(v)) + \frac{|\mathbf{v}(v)|}{8\pi\zeta(v)^2} + \frac{\pi}{2} \frac{1}{|\mathbf{v}(v)|} \left[\text{sgn}(\zeta(v+1))\zeta(v+1)^2 - \text{sgn}(\zeta(v))\zeta(v)^2 \right]^2 \right\} \quad (6.4)
\end{aligned}$$

where it is used that

$$\begin{aligned}
\frac{1}{\ell_p^2} \ln \left(\left| 1 \pm \frac{4\pi\beta\sqrt{\Delta}\ell_p^3}{\mathbf{v}(v)} \right| \right) &\cong \pm \frac{4\pi\beta\sqrt{\Delta}\ell_p}{\mathbf{v}(v)} \\
\frac{1}{\ell_p^3} B \left[\frac{\mathbf{v}(v)}{4\pi\beta\sqrt{\Delta}\ell_p^3} \right] &\cong \frac{4\pi\beta\sqrt{\Delta}}{27|\mathbf{v}(v)|}. \quad (6.5)
\end{aligned}$$

It is worth noting that the RHS of (6.4) returns to that of (4.3) with the assignment

$$\begin{aligned}
\theta(v) &= \frac{c(v) + b(v)\mathbf{v}(v)}{2\text{sgn}(\zeta(v))\zeta(v)^2}, & p(v) &= \text{sgn}(\zeta(v))\zeta(v)^2, \\
\Phi(v) &= 4\pi\zeta(v)b(v), & \Pi(v) &= \frac{\mathbf{v}(v)}{4\pi\zeta(v)}. \quad (6.6)
\end{aligned}$$

A. The equations of motion

According to the above discussions, we get an effective Hamiltonian

$$H_{\text{eff}}(\vec{\zeta}, \vec{\mathbf{v}}, \vec{c}, \vec{b}) = -\frac{1}{G} H(\vec{\zeta}, \vec{\mathbf{v}}, \vec{c}, \vec{b}). \quad (6.7)$$

Due to the Poisson brackets (6.2), the EOMs are

$$\begin{aligned}
\frac{d\underline{\mathbf{v}}(v)}{dt} &= \frac{\underline{\mathbf{v}}(v) (2 \sin [2\underline{\mathbf{b}}(v) + \underline{\mathbf{c}}(v)] + \sin [2\underline{\mathbf{b}}(v)])}{2\beta\sqrt{\Delta\ell_p}} \\
\frac{d\underline{\mathbf{b}}(v)}{dt} &= -\frac{\sin(\underline{\mathbf{b}}(v))}{\beta\sqrt{\Delta\ell_p}} \left[\frac{2 \sin [\underline{\mathbf{b}}(v) + \underline{\mathbf{c}}(v)] + \sin [\underline{\mathbf{b}}(v)]}{8\pi\beta\sqrt{\Delta\ell_p}} - \frac{\underline{\mathbf{c}}(v)}{\underline{\mathbf{v}}(v)} \cos [\underline{\mathbf{b}}(v) + \underline{\mathbf{c}}(v)] \right] \\
&\quad - \frac{1}{8\pi\underline{\zeta}(v)^2} + \frac{\pi}{2} \frac{1}{\underline{\mathbf{v}}(v)^2} (\underline{\zeta}(v)^2 - \underline{\zeta}(v+1)^2)^2 \\
\frac{d\underline{\zeta}(v)}{dt} &= \frac{\underline{\zeta}(v) \sin [\underline{\mathbf{b}}(v)] \cos [\underline{\mathbf{b}}(v) + \underline{\mathbf{c}}(v)]}{\beta\sqrt{\Delta\ell_p}} \\
\frac{d\underline{\mathbf{c}}(v)}{dt} &= \frac{\underline{\mathbf{v}}(v)}{4\pi\underline{\zeta}(v)^2} + 2\pi\underline{\zeta}(v)^2 \left(\frac{\underline{\zeta}(v+1)^2 - \underline{\zeta}(v)^2}{\underline{\mathbf{v}}(v)} - \frac{\underline{\zeta}(v)^2 - \underline{\zeta}(v-1)^2}{\underline{\mathbf{v}}(v-1)} \right)
\end{aligned} \tag{6.8}$$

where

$$\underline{\mathbf{b}}(v) = 4\pi\beta\sqrt{\Delta\ell_p}\underline{b}(v), \quad \underline{\mathbf{c}}(v) = 4\pi\beta\sqrt{\Delta\ell_p}\frac{\underline{\mathbf{c}}(v)}{\underline{\mathbf{v}}(v)}. \tag{6.9}$$

In (6.8), we have assumed that $\zeta(v) > 0$ without loss of generality, since, as seen below, $\text{sgn}(\zeta(v))$ is kept along the concerning dynamical trajectory.

In order to discuss the continuous limit of the EOMs (6.8), we introduce the continuous variables $\tilde{\mathbf{v}}(x)$, $\tilde{b}(x)$, $\tilde{\zeta}(x)$ and $\tilde{\mathbf{c}}(x)$, which are related to the classical variables $\underline{\mathbf{v}}(v)$, $\underline{b}(v)$, $\underline{\zeta}(v)$ and $\underline{\mathbf{c}}(v)$ via

$$\begin{aligned}
\underline{\mathbf{v}}(v) &= \int_{e(v)} dx \tilde{\mathbf{v}}(x), & \underline{b}(v) &= \tilde{b}(\text{mid}_{e(v)}) \\
\underline{\zeta}(v) &= \tilde{\zeta}(\text{mid}_{e(v)}), & \underline{\mathbf{c}}(v) &= \int_{e(v)} dx \tilde{\mathbf{c}}(x).
\end{aligned} \tag{6.10}$$

With the continuous variables, (6.6) is rewritten as

$$\begin{aligned}
K_x(x) &= \frac{\tilde{\mathbf{c}}(x) + \tilde{b}(x)\tilde{\mathbf{v}}(x)}{2\tilde{\zeta}(x)}, & E^x(x) &= \text{sgn}(\tilde{\zeta}(x))\tilde{\zeta}(x)^2, \\
K_\varphi(x) &= 4\pi\tilde{\zeta}(x)\tilde{b}(x), & E^\varphi(x) &= \frac{\tilde{\mathbf{v}}(x)}{4\pi\tilde{\zeta}(x)}.
\end{aligned} \tag{6.11}$$

Moreover, substituting the continuous variables into the EOMs (6.8) and considering the continuous limit of the lattice γ , we get

$$\begin{aligned}
\partial_t \tilde{\mathbf{v}} &= \frac{\tilde{\mathbf{v}} (2 \sin [2\tilde{\mathbf{b}} + \tilde{\mathbf{c}}] + \sin [2\tilde{\mathbf{b}}])}{2\beta\sqrt{\Delta\ell_p}} \\
\partial_t \tilde{b} &= -\frac{\sin(\tilde{\mathbf{b}})}{\beta\sqrt{\Delta\ell_p}} \left(\frac{2 \sin [\tilde{\mathbf{b}} + \tilde{\mathbf{c}}] + \sin [\tilde{\mathbf{b}}]}{8\pi\beta\sqrt{\Delta\ell_p}} - \frac{\tilde{\mathbf{c}}}{\tilde{\mathbf{v}}} \cos [\tilde{\mathbf{b}} + \tilde{\mathbf{c}}] \right) - \frac{1}{8\pi\tilde{\zeta}^2} + \frac{\pi}{2} \frac{(\partial_x(\tilde{\zeta}^2))^2}{\tilde{\mathbf{v}}^2} \\
\partial_t \tilde{\zeta} &= \frac{\tilde{\zeta} \sin [\tilde{\mathbf{b}}] \cos [\tilde{\mathbf{b}} + \tilde{\mathbf{c}}]}{\beta\sqrt{\Delta\ell_p}} \\
\partial_t \tilde{\mathbf{c}} &= \frac{\tilde{\mathbf{v}}}{4\pi\tilde{\zeta}^2} + 2\pi\tilde{\zeta}^2 \partial_x \left(\frac{1}{\tilde{\mathbf{v}}} \partial_x(\tilde{\zeta}^2) \right)
\end{aligned} \tag{6.12}$$

where

$$\tilde{\mathbf{b}}(x) = 4\pi\beta\sqrt{\Delta\ell_p}\tilde{b}(x), \quad \tilde{\mathbf{c}}(x) = 4\pi\beta\sqrt{\Delta\ell_p}\frac{\tilde{\mathbf{c}}(x)}{\tilde{\mathbf{v}}(x)}. \tag{6.13}$$

The same EOMs as (6.12) can be obtained if one consider a system of the phase space of $(\tilde{\mathbf{u}}(x), \tilde{\mathbf{b}}(x), \tilde{\zeta}(x), \tilde{c}(x))$, with the non-vanishing Poisson brackets

$$\{\tilde{\mathbf{b}}(x), \tilde{\mathbf{u}}(y)\} = G\delta(x, y), \quad \{\tilde{c}(x), \tilde{\zeta}(y)\} = G\tilde{\zeta}(x)\delta(x, y) \quad (6.14)$$

and the Hamiltonian

$$\begin{aligned} \tilde{H}_{\text{eff}} = & -\frac{1}{G} \int_{\mathbb{R}} \left\{ \frac{1}{4\pi\beta^2\Delta\ell_p^2} |\tilde{\mathbf{u}}(x)| \sin \left[4\pi\beta\sqrt{\Delta}\ell_p \tilde{\mathbf{b}}(x) \right] \sin \left[\frac{4\pi\beta\sqrt{\Delta}\ell_p}{\tilde{\mathbf{u}}(x)} \tilde{c}(x) + 4\pi\beta\sqrt{\Delta}\ell_p \tilde{\mathbf{b}}(x) \right] + \right. \\ & \left. \frac{1}{8\pi\beta^2\Delta\ell_p^2} |\tilde{\mathbf{u}}(x)| \sin^2(4\pi\beta\sqrt{\Delta}\ell_p \tilde{\mathbf{b}}(x)) + \frac{|\tilde{\mathbf{u}}(x)|}{8\pi\tilde{\zeta}(x)^2} + \frac{\pi}{2} \frac{1}{|\tilde{\mathbf{u}}(x)|} \left(\partial_x \left(\text{sgn}(\tilde{\zeta}(x)) \tilde{\zeta}(x)^2 \right) \right)^2 \right\}. \end{aligned} \quad (6.15)$$

In this sense, this system described by (6.14) and (6.15) are the continuous limit of effective dynamics based on the lattice γ and encoded in (6.2) and (6.7). Since (6.15) is the same as the effective Hamiltonian used in [1] if (6.11) is substituted, (6.12) is the same as the EOMs used in [1].

B. A solution to the EOMs

The continuous effective Hamiltonian (6.15) returns to the classical Hamiltonian (2.33) in the low curvature region where $\tilde{\mathbf{b}}(x) \ll 1$ and $\tilde{c}(c)/\tilde{\mathbf{u}}(x) \ll 1$. As is known in [1], solving the EOMs generated by the classical Hamiltonian (2.33) gives the Schwarzschild metric in Lemâitre coordinates. For this solution, all of the dynamical variables depend only on $x - t$, which represents homogeneity of the interior and the static feature of the exterior of the Schwarzschild solution. In this section, we are also concerned about the solutions depending only one $x - t$, that is, solutions taking the form

$$\tilde{\mathbf{u}}(x, t) = \tilde{\mathbf{u}}(x - t), \quad \tilde{\mathbf{b}}(x, t) = \tilde{\mathbf{b}}(x - t), \quad \tilde{\zeta}(x, t) = \tilde{\zeta}(x - t), \quad \tilde{c}(x, t) = \tilde{c}(x - t). \quad (6.16)$$

Before proceeding further, it is helpful to investigate constants of motion in the effective dynamics. As mentioned in (2.35), $\mathcal{V}(x)$ is a conserved charge in the classical dynamics. Fortunately, this feature is kept in the effective dynamics, because of $0 = \{\mathcal{V}(x), \tilde{H}_{\text{eff}}\}$. In terms of the continuous variables, $\mathcal{V}(x)$ is

$$\mathcal{V}(x) = \frac{\tilde{c}(x)\partial_x\tilde{\zeta}(x)}{\tilde{\zeta}(x)} - \tilde{\mathbf{u}}(x)\partial_x\tilde{\mathbf{b}}(x) \quad (6.17)$$

Since we are concern on the effective solutions which can recover the Schwarzschild metric in the low curvature region, it has

$$\mathcal{V}(x) = \frac{\tilde{c}(x)\partial_x\tilde{\zeta}(x)}{\tilde{\zeta}(x)} - \tilde{\mathbf{u}}(x)\partial_x\tilde{\mathbf{b}}(x) = 0 \quad (6.18)$$

with recalling the statements below (2.35).

Substitute the ansatz (6.16) in to the EOMs (6.12) and (6.18). With denoting $y = x - t$, we have

$$-\frac{d\tilde{\mathbf{u}}(y)}{dy} = \frac{\tilde{\mathbf{u}}(y) \left(2 \sin \left[2\tilde{\mathbf{b}}(y) + \tilde{c}(y) \right] + \sin \left[2\tilde{\mathbf{b}}(y) \right] \right)}{4\beta\sqrt{\Delta}\ell_p} \quad (6.19)$$

$$\begin{aligned} -\frac{d\tilde{\mathbf{b}}(y)}{dy} = & -\frac{\sin(\tilde{\mathbf{b}}(y))}{\beta\sqrt{\Delta}\ell_p} \left(\frac{2 \sin \left[\tilde{\mathbf{b}}(y) + \tilde{c}(y) \right] + \sin \left[\tilde{\mathbf{b}}(y) \right]}{8\pi\beta\sqrt{\Delta}\ell_p} - \frac{\tilde{c}(y)}{\tilde{\mathbf{u}}(y)} \cos \left[\tilde{\mathbf{b}}(y) + \tilde{c}(y) \right] \right) \\ & - \frac{1}{8\pi\tilde{\zeta}(y)^2} + \frac{\pi}{2} \frac{1}{\tilde{\mathbf{u}}(y)^2} \left(\frac{d\tilde{\zeta}(y)^2}{dy} \right)^2 \end{aligned} \quad (6.20)$$

$$-\frac{d\tilde{\zeta}(y)}{dy} = \frac{\tilde{\zeta}(y) \sin \left[\tilde{\mathbf{b}}(y) \right] \cos \left[\tilde{\mathbf{b}}(y) + \tilde{c}(y) \right]}{\beta\sqrt{\Delta}\ell_p} \quad (6.21)$$

$$-\frac{d\tilde{c}(y)}{dy} = \frac{\tilde{\mathbf{u}}(y)}{4\pi\tilde{\zeta}(y)^2} + 2\pi\tilde{\zeta}(y)^2 \frac{d}{dy} \left(\frac{1}{\tilde{\mathbf{u}}(y)} \frac{d\tilde{\zeta}(y)^2}{dy} \right) \quad (6.22)$$




FIG. 1: Solutions to (6.19), (6.21), (6.22) and (6.23). The initial data are chosen such that $\tilde{\mathbf{u}}(y_0) = 6\pi F_0 y_0$, $\tilde{\mathbf{b}}(y_0) = -\frac{1}{6\pi y_0}$, $\tilde{\zeta}(y_0) = (\frac{3}{2}\sqrt{F_0 y_0})^{2/3}$, $\tilde{\mathbf{c}}(y_0) = \frac{3F_0}{2}$ and $\tilde{\zeta}'(y_0) = \frac{2}{3}(\frac{3}{2}\sqrt{F_0 y_0})^{-1/3}$ with $y_0 = 10^5 \ell_p$ and $F_0 = 10^4 \ell_p$. The initial data are chosen by considering the Schwarzschild solution with $2GM = F_0$ in Lemâitre coordinate at $x - t = y_0$. The parameters are set to be $\Delta = 0.1$ and $\beta = 0.2375$.

and

$$\frac{\tilde{\mathbf{c}}(y)}{\tilde{\zeta}(y)} \frac{d\tilde{\zeta}(y)}{dy} - \tilde{\mathbf{u}}(y) \frac{d\tilde{\mathbf{b}}(y)}{dy} = 0 \quad (6.23)$$

To get the solutions, the equations (6.19), (6.21), (6.22) and (6.23) are chosen. A set of numerical results is shown in Fig. 1. Since (6.20) is used to get the solution, we can use it to check the accuracy of our solutions. Substituting the numerical solutions achieved from (6.19), (6.21), (6.22) and (6.23) into (6.20), we have the residuals plotted in Fig. 2. According to the numerical results, for $y = x - t \rightarrow -\infty$, the variables behaves as

$$\tilde{\mathbf{u}}(x - t) = e^{\mathbf{a} + \mathbf{b}|x-t|\ell_p^{-1}}, \quad \tilde{\mathbf{b}}(x - t) = \mathbf{c}, \quad \tilde{\zeta}(x - t) = r_0, \quad \tilde{\mathbf{c}}(x - t) = \mathbf{d} \quad (6.24)$$

where \mathbf{a} , \mathbf{b} , \mathbf{c} , r_0 and \mathbf{d} are constants. Thus, at $y \rightarrow -\infty$ is the spacetime is diffeomorphism to Nariai geometry taking $dS_2 \times S^2$ metric

$$ds^2 = -dt^2 + \left(\frac{\tilde{\mathbf{u}}(x, t)}{\tilde{\zeta}^2} \right)^2 dx^2 + \tilde{\zeta}^2 d\Omega^2, \quad (6.25)$$

where the metric in terms of the variables $\tilde{\mathbf{u}}$ and $\tilde{\zeta}$ is given in [1, 34]. This result is the same as the results given in [1], which is not surprising because the equations (6.19), (6.20), (6.21) and (6.22) are the same as that used in [1]. Thus one can refer to [1] for more details on this solution. According to the analysis there, both the area of S^2 and the dS radius of the Nariai geometry obtained in this model are of quantum size. In other words, the current model predicts a quantum final fate of Schwarzschild BH. However, it should be emphasized that this is not credible always, which will be shown below by discussing the scope of the continuous effective descriptions.

Let us discuss the scope of continuous effective differential equations (6.19)–(6.22). At first, the derivation from (6.3) to (6.4) requires

$$\mathbf{u}(v) \gg 4\pi\beta\sqrt{\Delta}\ell_p^3, \quad \forall v \in V(\gamma). \quad (6.26)$$

According to (6.10), this means

$$\int_{\varepsilon(v)} dx \tilde{\mathbf{u}}(x) \gg 4\pi\beta\sqrt{\Delta}\ell_p^3, \quad \forall v \in V(\gamma). \quad (6.27)$$




FIG. 2: The residuals of (6.20) by substituting the numerical solutions shown in Fig. 1

Let $\tilde{\mathbf{v}}_{\min}$ be the smallest value of $\tilde{\mathbf{v}}(x, t)$. Since $\int_{e(v)} dx \tilde{\mathbf{v}}(x) > \delta x \tilde{\mathbf{v}}_{\min}$ where δx denotes the coordinate length of $e(v)$, it is concluded that (6.27) can be satisfied provided that

$$\delta x \tilde{\mathbf{v}}_{\min} \gg 4\pi\beta\sqrt{\Delta} \ell_p^3. \quad (6.28)$$




FIG. 3: Dependence of $\tilde{\mathbf{v}}_{\min}$ on M for various values of Δ . As shown in the figure, for $M \gg 1$, $\tilde{\mathbf{v}}_{\min}$ can be well approximated by (6.29). In this plot, the values of α_i , corresponding to Δ_i , with $i = 1, 2$ are computed by (6.30). The parameters are chosen as $\beta = 0.2375$.

The solutions to (6.19)–(6.22) is determined by the initial data. Meanwhile, the initial data are chosen such that in the low curvature region the effective dynamics returns to the Schwarzschild metric which depends only on its mass M . As a consequence, the value $\tilde{\mathbf{v}}_{\min}$, as the minimal value of the solution, is related to the mass of the Schwarzschild spacetime to set the initial data. According to our numerical computation shown in Fig. 3, the correlation between $\tilde{\mathbf{v}}_{\min}$ and M , for $M \gg 1$, can be well approximated by a formula of the form

$$\tilde{\mathbf{v}}_{\min} \cong \alpha(2GM)\ell_p. \quad (6.29)$$

where α is some coefficient depends only on $\beta\sqrt{\Delta}\ell_p$, i.e. $\alpha = \alpha(\beta\sqrt{\Delta}\ell_p)$, because the coefficients in (6.19)–(6.22) depends on $\beta\sqrt{\Delta}\ell_p$ merely. The dependence of α on $\beta\sqrt{\Delta}\ell_p$ can be investigated numerically, which, as shown in Fig. 4, is

$$\alpha(\beta\sqrt{\Delta}\ell_p) \cong \alpha_0\beta\sqrt{\Delta} \quad (6.30)$$




FIG. 4: Plot of the values α , introduced in (6.31), depending on $\beta\sqrt{\Delta}\ell_p$. As shown in the figure, α depends on $\beta\sqrt{\Delta}$ linearly. More explicitly, $\alpha = \alpha_0\beta\sqrt{\Delta}$ with $\alpha_0 = 3.0000$ according to the numerical results shown here.

with $\alpha_0 \cong 3.0000$. Therefore, for $M \gg 1$, $\tilde{\mathbf{v}}_{\min}$ is approximated by

$$\tilde{\mathbf{v}}_{\min} \cong \alpha_0\beta\sqrt{\Delta}(2GM)\ell_p. \quad (6.31)$$

Substituting (6.31) into (6.28), we finally obtain that

$$\frac{2GM}{\ell_p} \gg \frac{4\pi}{\alpha_0} \frac{\ell_p}{\delta x} \quad (6.32)$$

which presents the scope of the continuous effective EOMs. It should be emphasized that δx is the coordinate length of each edge in a specific coordinate such that the effective metric depends only on $x - t$ and, takes the limit of Schwarzschild metric in Lemâitre coordinate in the low curvature region. Moreover, the above discussion assumed that the lattice γ is equidistant in this specific coordinate.

According to (6.32), δx should take large values to enlarge the scope of the continuous effective EOMs. However, since we used differentials to approximate the differences in (6.8), a tension arises that δx cannot be too large. To see more explicitly how this tension limits δx , one notices that the approximations of differences by differentials in (6.8) are

$$\begin{aligned} \frac{1}{\underline{\mathbf{v}}(v)} (\underline{\zeta}(v+1)^2 - \zeta(v)^2) &\cong \frac{1}{\tilde{\mathbf{v}}(x)} \partial_x (\tilde{\zeta}(x)^2) \Big|_{x=\text{mid}_{e(v)}} \\ \left(\frac{\underline{\zeta}(v+1)^2 - \underline{\zeta}(v)^2}{\underline{\mathbf{v}}(v)} - \frac{\underline{\zeta}(v)^2 - \underline{\zeta}(v-1)^2}{\underline{\mathbf{v}}(v-1)} \right) &\cong \partial_x \left(\frac{1}{\tilde{\mathbf{v}}(x)} \partial_x (\tilde{\zeta}(x)^2) \right) \Big|_{x=\text{mid}_{e(v)}}. \end{aligned} \quad (6.33)$$

The last equations omit the dependence of all variables on t for simplicity. That is, we are discussing the validity these two equations for a fixed t . Consider the first equation as an example. According to (6.10), one has

$$\frac{1}{\underline{\mathbf{v}}(v)} (\underline{\zeta}(v+1)^2 - \zeta(v)^2) = \left(\frac{\partial_x (\tilde{\zeta}^2)(\text{mid}_{e(v)})}{\tilde{\mathbf{v}}(\text{mid}_{e(v)})} + \frac{1}{2} \frac{\partial_x^2 (\tilde{\zeta}^2)(c') \delta x}{\tilde{\mathbf{v}}(\text{mid}_{e(v)})} \right) \frac{1}{1 + \frac{\partial_x \tilde{\mathbf{v}}(c) \delta x}{2\tilde{\mathbf{v}}(\text{mid}_{e(v)})}} \quad (6.34)$$

where c and c' are some points in $e(v)$ and the mean-value form of the remainder of the Taylor series is applied. As mentioned above, in the region $x - t \gg 1$, the solutions to the effective continuous EOMs converge to the Schwarzschild metric in Lemâitre coordinate, i.e.

$$\begin{aligned} \tilde{\zeta}(x, t) &= \left(\frac{3}{2} \sqrt{F}(x - t) \right)^{2/3} \\ \tilde{\mathbf{v}}(x, t) &= 6\pi F(x - t) \end{aligned} \quad (6.35)$$

with some constant F . Substituting these expressions into (6.34), we have

$$\frac{1}{\underline{\mathbf{v}}(v)} (\underline{\zeta}(v+1)^2 - \underline{\zeta}(v)^2) = \frac{\partial_x(\tilde{\zeta}^2)(\text{mid}_{e(v)})}{\underline{\mathbf{v}}(\text{mid}_{e(v)})} + o(\delta x/(x-t)). \quad (6.36)$$

Thus in the region $x-t \gg 1$, the differentials can well approximate the differences if $\delta x \sim 1$. Moreover, because $\tilde{\zeta}(x,t)$ keeps constant in the region $x-t \ll -1$, the differentials can well approximate the differences in this region for any value of δx . As a summary, the approximation of the differences by differentials is well behaved in the region $|x-t| \gg 1$. Thus, we only need to check the validity of this approximation in the region of $|x-t| \sim 1$. To do this, we substitute (6.10) into (6.8) and numerically compute the following dimensionless variables

$$\begin{aligned} \text{err}_{\underline{\mathbf{v}}}(v,t) &:= -\frac{\ell_p}{\underline{\mathbf{v}}(v,t)} \frac{d\underline{\mathbf{v}}(v,t)}{dt} + \frac{(2 \sin [2\mathbf{b}(v,t) + \mathbf{c}(v,t)] + \sin [2\mathbf{b}(v,t)])}{2\beta\sqrt{\Delta}}, \\ \text{err}_{\underline{\mathbf{b}}}(v,t) &:= -\ell_p^2 \frac{d\underline{\mathbf{b}}(v,t)}{dt} - \frac{\sin(\mathbf{b}(v,t))}{\beta\sqrt{\Delta}} \left[\frac{2 \sin [\mathbf{b}(v,t) + \mathbf{c}(v,t)] + \sin [\mathbf{b}(v,t)]}{8\pi\beta\sqrt{\Delta}} - \ell_p \frac{\underline{\mathbf{c}}(v,t)}{\underline{\mathbf{v}}(v,t)} \cos [\mathbf{b}(v,t) + \mathbf{c}(v,t)] \right] \\ &\quad - \frac{\ell_p^2}{8\pi\underline{\zeta}(v,t)^2} + \frac{\pi}{2} \frac{\ell_p^2}{\underline{\mathbf{v}}(v,t)^2} (\underline{\zeta}(v,t)^2 - \underline{\zeta}(v+1,t)^2)^2, \\ \text{err}_{\underline{\zeta}}(v,t) &:= -\frac{\ell_p}{\underline{\zeta}(v,t)} \frac{d\underline{\zeta}(v,t)}{dt} + \frac{\sin [\mathbf{b}(v,t)] \cos [\mathbf{b}(v,t) + \mathbf{c}(v,t)]}{\beta\sqrt{\Delta}}, \\ \text{err}_{\underline{\mathbf{c}}}(v,t) &:= -\frac{\ell_p^2}{\underline{\mathbf{v}}(v,t)} \frac{d\underline{\mathbf{c}}(v,t)}{dt} + \frac{\ell_p^2}{4\pi\underline{\zeta}(v,t)^2} + 2\pi\ell_p^2 \frac{\underline{\zeta}(v,t)^2}{\underline{\mathbf{v}}(v,t)} \left(\frac{\underline{\zeta}(v+1,t)^2 - \underline{\zeta}(v,t)^2}{\underline{\mathbf{v}}(v,t)} - \frac{\underline{\zeta}(v,t)^2 - \underline{\zeta}(v-1,t)^2}{\underline{\mathbf{v}}(v-1,t)} \right). \end{aligned} \quad (6.37)$$

Fig 5 plots the values of $\max |\text{err}_{\underline{\mathbf{v}}}|$, $\max |\text{err}_{\underline{\mathbf{b}}}|$, $\max |\text{err}_{\underline{\zeta}}|$ and $\max |\text{err}_{\underline{\mathbf{c}}}|$ which are defined by

$$\begin{aligned} \max |\text{err}_{\underline{\mathbf{v}}}| &:= \max_{t \in \mathbb{R}, v \in V(\gamma)} |\text{err}_{\underline{\mathbf{v}}}(v,t)|, \\ \max |\text{err}_{\underline{\mathbf{b}}}| &:= \max_{t \in \mathbb{R}, v \in V(\gamma)} |\text{err}_{\underline{\mathbf{b}}}(v,t)|, \\ \max |\text{err}_{\underline{\zeta}}| &:= \max_{t \in \mathbb{R}, v \in V(\gamma)} |\text{err}_{\underline{\zeta}}(v,t)|, \\ \max |\text{err}_{\underline{\mathbf{c}}}| &:= \max_{t \in \mathbb{R}, v \in V(\gamma)} |\text{err}_{\underline{\mathbf{c}}}(v,t)|. \end{aligned} \quad (6.38)$$

According to (6.32), the minimal values of $2GM\ell_p^{-1}$ for a fixed δx is $(2GM\ell_p^{-1})_{\min} \cong 4.19/(\delta x\ell_p^{-1})$, which is 4.19×10^{-4} for the case plotted in Fig. 5. Thus, the inequality of (6.32) is satisfied for $2GM \geq 10^6\ell_p$. As shown in this figure, the values of $\max |\text{err}_{\underline{\mathbf{v}}}|$, $\max |\text{err}_{\underline{\mathbf{b}}}|$, $\max |\text{err}_{\underline{\zeta}}|$ and $\max |\text{err}_{\underline{\mathbf{c}}}|$ are tiny. That is, the numerical results support that the solutions to the differential equations (6.12) approximate well the solutions to the difference EOMs (6.8) for large $2GM$, with noting that for large $2GM$, δx can be chosen small. However, for small $2GM$, this conclusion is no longer valid due to the large value of the allowed δx in this case. This can be seen intuitively from the numerical results. Fig. 6 plots the values of $\max |\text{err}_{\underline{\mathbf{v}}}|$, $\max |\text{err}_{\underline{\mathbf{b}}}|$, $\max |\text{err}_{\underline{\zeta}}|$ and $\max |\text{err}_{\underline{\mathbf{c}}}|$ depending on δx for $2GM = 100\ell_p$. For this case, the allowed values of δx by (6.32) are those satisfying $\delta x\ell_p^{-1} \gg 4.19 \times 10^{-2}$ which are indicated by the red lines in Fig. 6. By the numerical results, the values of $\max |\text{err}_{\underline{\mathbf{v}}}|$, $\max |\text{err}_{\underline{\mathbf{b}}}|$, $\max |\text{err}_{\underline{\zeta}}|$ and $\max |\text{err}_{\underline{\mathbf{c}}}|$ are no longer small for the allowed δx . That is, the solutions to the differential equations (6.12) no longer well approximate the solutions to the difference EOMs (6.8).

By this analysis, the discrete EOMs (6.8) have correct classical limit only if the inequality (6.32) is satisfied and simultaneously δx is small enough, so that solutions to (6.8) are compatible with that of the continuous effective dynamics, and, thus, the classical Schwarzschild solutions are recovered asymptotically. However, these two conditions lead to opposite tendencies for the values of δx . According to (6.32), δx should be large so that (6.8) can be valid for a large scope, which is contradictory to the second requirement that δx should be small such that the continuous EOMs can well approximate the discrete ones. This tension, on one hand, prevent the value of δx to vanishing, because for vanishing δx , the effective EOMs are no longer valid for any $2GM$. On the other hand, this tension requires δx to be small enough, because, otherwise the continuous EOMs cannot give well-approximated solutions to the discrete EOMs. In other words, δx should be some finitely small value. Since δx is interpreted as the length of the edges in γ , we conclude that the current model does not allow γ to shrink to a continuous line, even though the classical theory is presented based on a continuous line. This implies that the lattice structure underlying the current model is fundamental. However, this conclusion does not mean that the current lattice model never owns continuous limit.




FIG. 5: Plots of the values of $\max|\text{err}_b|$, $\max|\text{err}_b|$, $\max|\text{err}_\zeta|$ and $\max|\text{err}_\zeta|$ depending on $2GM$ for $\delta x = 10^{-4}\ell_p$. By (6.32), the minimal values of $(2GM)\ell_p^{-1}$ in this case is $4\pi/(\alpha_0\delta x\ell_p^{-1}) \approx 4.19 \times 10^4$.




FIG. 6: The values of $\max|\text{err}_b|$, $\max|\text{err}_b|$, $\max|\text{err}_\zeta|$ and $\max|\text{err}_\zeta|$ depending on δx for $2GM = 100\ell_p$ (black dots). $\delta x \ell_p^{-1} = 4\pi/(\alpha_0(2GM)\ell_p^{-1})$ is plotted by the red lines which indicate the allowed values of $\delta x \ell_p^{-1}$ by (6.32).

Since δx is small, the discrete EOMs can be well approximated by the differential EOMs (6.12), which can be derived from a continuous Hamiltonian (6.15) together with the Poisson brackets (6.14). Thus, the continuous description of the lattice model is the model whose phase space, consisting of fields $(\tilde{\mathbf{u}}(x), \tilde{\mathbf{b}}(x), \tilde{\zeta}(x), \tilde{\mathcal{Z}}(x))$ on \mathbb{R} , is endowed with the Poisson brackets (6.14) and the Hamiltonian (6.15). This continuous description is usually referred to as the effective dynamics and has been studied in [1]. However, according to our analysis, this continuous description is only valid for black holes with macro masses satisfying (6.32) but not for small BHs. In other words, macro BHs and small BHs behave differently if the LQG effects are considered.

VII. CONCLUSION AND OUTLOOK

This paper considers the loop quantum theory of spherically symmetric gravity coupled to Gaussian dusts where the Gaussian dusts provide a material reference frame of the space and time to deparameterize gravity. Classically, the dynamics of this model can give the Schwarzschild solution in Lemaître coordinate. Thus this model provides an approach to study the spherically symmetric BH. As in the usual loop quantum theory, the present model is constructed based on some graph γ which is a 1-dimensional lattice here. By using the $\bar{\mu}$ -scheme to regularize the physical Hamiltonian, we obtain a Hamiltonian operator which governs the quantum dynamics. Then the quantum dynamics is studied by the path integral formulation and an effective action is obtained. With this effective action, an effective continuous description of the quantum lattice model is derived which is encoded in (6.15). Note that (6.15) is indeed the classical Hamiltonian with the holonomy correction which is usually used in literature like [1]. However, according to our analysis, this effective continuous description is valid only if (6.32) holds. In other works, this effective description, roughly speaking, is valid for macro BHs but not for small BHs. Therefore, it is natural to ask questions on the dynamics of small BHs. Moreover, it is also interesting to introduce BH evaporation to the current model and consider the issue on the final fate of the evaporation. These topics will be left for our further studying.

Acknowledgements

This work is benefited greatly from the numerous discussions with Muxin Han and Honguang Liu, and supported by the Polish Narodowe Centrum Nauki, Grant No. 2018/30/Q/ST2/00811.

-
- [1] M. Han and H. Liu, *Improved effective dynamics of loop-quantum-gravity black hole and nariai limit*, *arXiv preprint arXiv:2012.05729* (2020).
 - [2] A. Ashtekar and J. Lewandowski, *Background independent quantum gravity: A Status report*, *Class.Quant.Grav.* **21** (2004) R53, [[gr-qc/0404018](#)].
 - [3] C. Rovelli, *Quantum Gravity*. Cambridge University Press, 2004.
 - [4] M. Han, W. Huang, and Y. Ma, *Fundamental structure of loop quantum gravity*, *Int.J.Mod.Phys.* **D16** (2007) 1397–1474, [[gr-qc/0509064](#)].
 - [5] T. Thiemann, *Modern Canonical Quantum General Relativity*. Cambridge University Press, 2007.
 - [6] C. Rovelli and L. Smolin, *The Physical Hamiltonian in nonperturbative quantum gravity*, *Phys. Rev. Lett.* **72** (1994) 446–449, [[gr-qc/9308002](#)].
 - [7] C. Rovelli and L. Smolin, *Knot Theory and Quantum Gravity*, *Physical Review Letters* **61** (Sept., 1988) 1155–1158.
 - [8] C. Rovelli and L. Smolin, *Discreteness of area and volume in quantum gravity*, *Nuclear Physics B* **442** (May, 1995) 593–619.
 - [9] A. Ashtekar and J. Lewandowski, *Quantum theory of geometry. 1: Area operators*, *Class.Quant.Grav.* **14** (1997) A55–A82, [[gr-qc/9602046](#)].
 - [10] A. Ashtekar and J. Lewandowski, *Quantum theory of geometry. 2. Volume operators*, *Adv.Theor.Math.Phys.* **1** (1998) 388–429, [[gr-qc/9711031](#)].
 - [11] T. Thiemann, *The Phoenix Project: master constraint programme for loop quantum gravity*, *Classical and Quantum Gravity* **23** (Apr., 2006) 2211–2247.
 - [12] T. Thiemann, *Quantum spin dynamics: VIII. The master constraint*, *Classical and Quantum Gravity* **23** (Apr., 2006) 2249–2265.
 - [13] J. Engle, E. Livine, R. Pereira, and C. Rovelli, *LQG vertex with finite Immirzi parameter*, *Nucl.Phys.* **B799** (2008) 136–149, [[arXiv:0711.0146](#)].
 - [14] M. Domagala, K. Giesel, W. Kaminski, and J. Lewandowski, *Gravity quantized: Loop Quantum Gravity with a Scalar Field*, *Phys. Rev.* **D82** (2010) 104038, [[arXiv:1009.2445](#)].

- [15] V. Husain and T. Pawłowski, *Time and a physical Hamiltonian for quantum gravity*, *Phys. Rev. Lett.* **108** (2012) 141301, [arXiv:1108.1145].
- [16] J. Yang and Y. Ma, *New Hamiltonian constraint operator for loop quantum gravity*, *Phys. Lett.* **B751** (2015) 343–347, [arXiv:1507.00986].
- [17] E. Alesci, M. Assanioussi, J. Lewandowski, and I. Mäkinen, *Hamiltonian operator for loop quantum gravity coupled to a scalar field*, *Phys. Rev.* **D91** (2015), no. 12 124067, [arXiv:1504.02068].
- [18] A. Dapor and K. Liegener, *Cosmological Effective Hamiltonian from full Loop Quantum Gravity Dynamics*, *Phys. Lett.* **B785** (2018) 506–510, [arXiv:1706.09833].
- [19] T. Lang, K. Liegener, and T. Thiemann, *Hamiltonian renormalisation I: derivation from Osterwalder-Schrader reconstruction*, *Class. Quant. Grav.* **35** (2018), no. 24 245011, [arXiv:1711.05685].
- [20] C. Zhang, J. Lewandowski, and Y. Ma, *Towards the self-adjointness of a Hamiltonian operator in loop quantum gravity*, *Phys. Rev.* **D98** (2018), no. 8 086014, [arXiv:1805.08644].
- [21] C. Zhang, J. Lewandowski, H. Li, and Y. Ma, *Bouncing evolution in a model of loop quantum gravity*, *Phys. Rev.* **D99** (2019), no. 12 124012, [arXiv:1904.07046].
- [22] A. Ashtekar, M. Bojowald, J. Lewandowski, et al., *Mathematical structure of loop quantum cosmology*, *Advances in Theoretical and Mathematical Physics* **7** (2003), no. 2 233–268.
- [23] M. Bojowald, *Loop quantum cosmology*, *Living Rev. Rel.* **8** (2005) 11, [gr-qc/0601085].
- [24] A. Ashtekar, T. Pawłowski, and P. Singh, *Quantum Nature of the Big Bang: Improved dynamics*, *Phys. Rev.* **D74** (2006) 084003, [gr-qc/0607039].
- [25] A. Ashtekar, *Loop Quantum Cosmology: An Overview*, *Gen. Rel. Grav.* **41** (2009) 707–741, [arXiv:0812.0177].
- [26] J. Yang, Y. Ding, and Y. Ma, *Alternative quantization of the Hamiltonian in loop quantum cosmology II: Including the Lorentz term*, *Phys. Lett.* **B682** (2009) 1–7, [arXiv:0904.4379].
- [27] A. Ashtekar and E. Wilson-Ewing, *Loop quantum cosmology of Bianchi I models*, *Phys. Rev.* **D79** (2009) 083535, [arXiv:0903.3397].
- [28] I. Agullo and P. Singh, *Loop Quantum Cosmology*, in *Loop Quantum Gravity: The First 30 Years* (A. Ashtekar and J. Pullin, eds.), pp. 183–240. WSP, 2017. arXiv:1612.01236.
- [29] M. Bojowald, *Spherically symmetric quantum geometry: states and basic operators*, *Classical and Quantum Gravity* **21** (2004), no. 15 3733.
- [30] A. Ashtekar and M. Bojowald, *Quantum geometry and the Schwarzschild singularity*, *Class. Quant. Grav.* **23** (2006) 391–411, [gr-qc/0509075].
- [31] L. Modesto, *Loop quantum black hole*, *Class. Quant. Grav.* **23** (2006) 5587–5602, [gr-qc/0509078].
- [32] C. G. Boehmer and K. Vandersloot, *Loop Quantum Dynamics of the Schwarzschild Interior*, *Phys. Rev. D* **76** (2007) 104030, [arXiv:0709.2129].
- [33] D.-W. Chiou, *Phenomenological loop quantum geometry of the schwarzschild black hole*, *Physical Review D* **78** (2008), no. 6 064040.
- [34] D.-W. Chiou, W.-T. Ni, and A. Tang, *Loop quantization of spherically symmetric midisuperspaces and loop quantum geometry of the maximally extended Schwarzschild spacetime*, arXiv:1212.1265.
- [35] R. Gambini, J. Olmedo, and J. Pullin, *Quantum black holes in Loop Quantum Gravity*, *Class. Quant. Grav.* **31** (2014) 095009, [arXiv:1310.5996].
- [36] A. Corichi and P. Singh, *Loop quantization of the schwarzschild interior revisited*, *Classical and Quantum Gravity* **33** (2016), no. 5 055006.
- [37] J. Olmedo, S. Saini, and P. Singh, *From black holes to white holes: a quantum gravitational, symmetric bounce*, *Class. Quant. Grav.* **34** (2017), no. 22 225011, [arXiv:1707.07333].
- [38] A. Ashtekar, J. Olmedo, and P. Singh, *Quantum extension of the Kruskal spacetime*, *Phys. Rev.* **D98** (2018), no. 12 126003, [arXiv:1806.02406].
- [39] A. Ashtekar, J. Olmedo, and P. Singh, *Quantum Transfiguration of Kruskal Black Holes*, *Phys. Rev. Lett.* **121** (2018), no. 24 241301, [arXiv:1806.00648].
- [40] J. Ben Achour, F. Lamy, H. Liu, and K. Noui, *Polymer Schwarzschild black hole: An effective metric*, *EPL* **123** (2018), no. 2 20006, [arXiv:1803.01152].
- [41] N. Bodendorfer, F. M. Mele, and J. Münch, *Effective Quantum Extended Spacetime of Polymer Schwarzschild Black Hole*, *Class. Quant. Grav.* **36** (2019), no. 19 195015, [arXiv:1902.04542].
- [42] C. Zhang, Y. Ma, S. Song, and X. Zhang, *Loop quantum schwarzschild interior and black hole remnant*, *Physical Review D* **102** (2020), no. 4 041502(R).
- [43] W.-C. Gan, N. O. Santos, F.-W. Shu, and A. Wang, *Properties of the spherically symmetric polymer black holes*, *Physical Review D* **102** (2020), no. 12 124030.
- [44] S. Rastgoo, P. Bosso, O. J. Obregon, and W. Yupanqui, *Deformed algebra and the effective dynamics of the interior of black holes*, *Classical and Quantum Gravity* (2021).
- [45] R. Gambini, J. Olmedo, and J. Pullin, *Spherically symmetric loop quantum gravity: analysis of improved dynamics*, arXiv:2006.01513.
- [46] J. Engle, *Relating loop quantum cosmology to loop quantum gravity: Symmetric sectors and embeddings*, *Class. Quant. Grav.* **24** (2007) 5777–5802, [gr-qc/0701132].
- [47] N. Bodendorfer, *An embedding of loop quantum cosmology in (b, v) variables into a full theory context*, *Class. Quant. Grav.* **33** (2016), no. 12 125014, [arXiv:1512.00713].
- [48] N. Bodendorfer, *State refinements and coarse graining in a full theory embedding of loop quantum cosmology*,

arXiv:1607.06227.

- [49] A. Ashtekar, M. Campiglia, and A. Henderson, *Loop Quantum Cosmology and Spin Foams*, *Phys. Lett.* **B681** (2009) 347–352, [arXiv:0909.4221].
- [50] A. Ashtekar, M. Campiglia, and A. Henderson, *Casting Loop Quantum Cosmology in the Spin Foam Paradigm*, *Class. Quant. Grav.* **27** (2010) 135020, [arXiv:1001.5147].
- [51] M. Han and H. Liu, *Effective Dynamics from Coherent State Path Integral of Full Loop Quantum Gravity*, *Phys. Rev.* **D101** (2020), no. 4 046003, [arXiv:1910.03763].
- [52] K. V. Kuchar and C. G. Torre, *Gaussian reference fluid and interpretation of quantum geometrodynamics*, *Phys. Rev.* **D43** (1991) 419–441.
- [53] K. Giesel and T. Thiemann, *Scalar Material Reference Systems and Loop Quantum Gravity*, *Class. Quant. Grav.* **32** (2015) 135015, [arXiv:1206.3807].
- [54] S. Holst, *Barbero’s Hamiltonian derived from a generalized Hilbert-Palatini action*, *Phys.Rev.* **D53** (1996) 5966–5969, [gr-qc/9511026].
- [55] B. Dittrich, *Partial and complete observables for Hamiltonian constrained systems*, *Gen. Rel. Grav.* **39** (2007) 1891–1927, [gr-qc/0411013].
- [56] T. Thiemann, *Reduced phase space quantization and Dirac observables*, *Class. Quant. Grav.* **23** (2006) 1163–1180, [gr-qc/0411031].
- [57] K. Giesel and T. Thiemann, *Algebraic quantum gravity (AQG). IV. Reduced phase space quantisation of loop quantum gravity*, *Class. Quant. Grav.* **27** (2010) 175009, [arXiv:0711.0119].