

# An odd feature of the ‘most classical’ states of $SU(2)$ invariant quantum mechanical systems

László B. Szabados  
Wigner Research Centre for Physics,  
H-1525 Budapest 114, P. O. Box 49, EU

June 17, 2021

## Abstract

Complex and spinorial techniques of general relativity are used to determine all the states of the  $SU(2)$  invariant quantum mechanical systems in which the equality holds in the uncertainty relations for the components of the angular momentum vector operator in two given directions. The expectation values depend on a discrete ‘quantum number’ and two parameters, one of them is the angle between the two angular momentum components and the other is the quotient of the two standard deviations. It is shown that although the standard deviations change continuously, one of the expectation values changes *discontinuously* on this parameter space. Since physically neither of the angular momentum components is distinguished over the other, this discontinuity suggests that the genuine parameter space must be a *Riemann surface* known in connection with the complex function  $\sqrt{z}$ . Moreover, the angle between the angular momentum components plays the role of the parameter of an interpolation between the continuous range of the expectation values found in the special case of the orthogonal angular momentum components by Aragone *et al* (J. Phys. A. **7** L149 (1974)) and the discrete point spectrum of one angular momentum component. The consequences in the *simultaneous* measurements of these angular momentum components are also discussed briefly.

## 1 Introduction

If the algebra of observables of the quantum system is the Heisenberg algebra, then the so-called canonical coherent states, defined to be the eigenstates of the annihilation operator, can be characterized in various equivalent ways (see e.g. [1, 2]). E.g. these are the smallest uncertainty states for the basic canonically conjugate observables, and, with the Hamiltonian of the harmonic oscillator, they do not spread out in space during their time evolution. Thus, these states are usually interpreted as the ‘most classical’ states of the quantum system. These states have particular significance e.g. in the characterization of the resulting state after the most accurate *simultaneous* measurement of *two non-commuting* observables [3, 4]; or in the characterization of the system in the classical limit.

The notion of coherent states has already been introduced for systems whose algebra of observables is more complicated than the Heisenberg algebra, e.g. when it is  $su(2)$  (see

e.g. [5]-[8]). These states could have particular significance e.g. in the characterization of the classical limit of the spin systems [6], or, in particular, in the determination of the large spin limit of the spin network model of the quantum spacetime suggested by Penrose in [9]. (A further potential application of these ideas in the somewhat related investigations will be given elsewhere [10].)

Our aim in the present paper is to determine *all* the ‘most classical’ states of the general  $SU(2)$ -invariant quantum mechanical systems *even with given intrinsic spin* in the sense that, in these states, *the equality holds in the uncertainty relation for the components of the angular momentum vector operator in two given directions*. To avoid confusion it should be stressed that these states refer to *pairs of observables*, actually to pairs of components of the angular momentum vector operator, rather than to the whole algebra of observables. The analogous problem was solved by Aragone *et al* in [7] in the framework of the spin coherent states of Radcliffe [5] when the two directions were orthogonal to one another.

We show that, allowing the angle between the two angular momentum components to be arbitrary, the expectation values depend on one discrete ‘quantum number’ and two continuous parameters; and since the two expectation values are the real and imaginary parts of the square root of a complex function on this two-dimensional parameter space, one of the expectation values change continuously but the other *discontinuously*. A possible resolution of this asymmetry between the angular momentum components is that *the genuine parameter space is homeomorphic to the Riemann surface appearing in connection with the function  $\sqrt{z}$  in complex analysis*. As far as we know, this is a new and typical quantum mechanical phenomenon. A potential experimental verification of this phenomenon, based on the use of the entangled coherent states [8] is raised.

Also, the angle between the two directions plays the role of the parameter of an *interpolation* between the continuous range of the expectation values found in [7] and the discrete point spectrum of one angular momentum component. We also determine the eigenfunctions both as series and in closed form. The analogous investigations for the  $E(3)$  (i.e. Euclidean) invariant elementary quantum mechanical systems will be presented in [11].

In deriving the above results, complex and spinorial techniques of general relativity are used. These techniques make it possible to determine the spectrum and the eigenfunctions of the relevant non-self-adjoint operator in explicit, *closed form* in a relatively straightforward (but technically a bit involved) way. It is this technique that can be (and, in fact, have been) used to carry out the analogous investigations in considerably more complicated systems [11]. To make the present paper self-contained, these ideas and techniques are summarized in Appendix 4.1. For a more detailed discussion of them, see e.g. [12, 13, 14, 15, 16, 17].

In section 2, we determine the ‘most classical’ states of the  $SU(2)$ -invariant systems and discuss its properties, namely we calculate the expectation values, the standard deviations and find the wave functions. There we also raise the possibility that the proper parameter space for the expectation value is the Riemann surface above. Section 3 is devoted to a brief discussion of how could it be possible to decide experimentally, at least in principle, that the genuine parameter space is this Riemann surface. Several technical details of the analyses in section 2 are presented in Appendix 4.2.

Our conventions are mostly those of [12, 13], except that the signature of metric of the Euclidean 3-space is positive definite. Here we do *not* use abstract indices: every index is a concrete (name, or component) index referring to some basis. Round brackets

around indices denote symmetrization.

## 2 The most classical states

As is well known (see e.g. [18]), using the Cauchy–Schwarz inequality, for the standard deviation (or uncertainty)  $\Delta_\phi \mathbf{A}$  and  $\Delta_\phi \mathbf{B}$  of any two observables  $\mathbf{A}$  and  $\mathbf{B}$  in any normalized state  $\phi$ , respectively, Schrödinger derived the inequality

$$(\Delta_\phi \mathbf{A})^2 (\Delta_\phi \mathbf{B})^2 \geq \frac{1}{4} | \langle [\mathbf{A}, \mathbf{B}] \rangle_\phi |^2 + \left( \langle \frac{1}{2} (\mathbf{A}\mathbf{B} + \mathbf{B}\mathbf{A}) \rangle_\phi - \langle \mathbf{A} \rangle_\phi \langle \mathbf{B} \rangle_\phi \right)^2. \quad (2.1)$$

Here, e.g.  $\langle \mathbf{A} \rangle_\phi$  is the expectation value of  $\mathbf{A}$ , and the standard deviation in the state  $\phi$  is given by  $\Delta_\phi \mathbf{A} = \sqrt{\langle \mathbf{A}^2 \rangle_\phi - (\langle \mathbf{A} \rangle_\phi)^2}$ . The necessary and sufficient condition of the equality in (2.1) is well known to be the vanishing of the second term on the right, which is the first order correlation between the two expectation values, and the equality in the Cauchy–Schwarz inequality. These two conditions together yield that the state  $\phi$  must be a solution of the eigenvalue equation

$$(\mathbf{A} - i\lambda \mathbf{B})\phi = (\langle \mathbf{A} \rangle_\phi - i\lambda \langle \mathbf{B} \rangle_\phi)\phi, \quad (2.2)$$

where, for *non-commuting* observables,  $\lambda$  is a nonzero real constant. Then, as (2.2) immediately implies,  $|\lambda| = \Delta_\phi \mathbf{A} / \Delta_\phi \mathbf{B}$  holds; and, for the sake of simplicity,  $\lambda$  will be chosen to be *positive*. These states are all among the smallest uncertainty states (i.e. for which  $F(\phi) := (\Delta_\phi \mathbf{A})(\Delta_\phi \mathbf{B})$ , as a real function on the unit sphere in the Hilbert space of the states, has a minimum at  $\phi$ ), but it is known [7] that the smallest uncertainty states do *not* necessarily saturate the inequality (2.1). The ‘most classical’ states are, in general, more special than the smallest uncertainty states. (Equation (2.2) shows also why it is almost impossible to find a non-trivial state in which the equality in (2.1) would hold for any two of *three* non-commuting observables  $\mathbf{A}$ ,  $\mathbf{B}$  and  $\mathbf{C}$ : Such a  $\phi$  would have to be a simultaneous eigenvector of the *three* non-commuting operators  $\mathbf{A} - i\lambda_1 \mathbf{B}$ ,  $\mathbf{B} - i\lambda_2 \mathbf{C}$  and  $\mathbf{C} - i\lambda_3 \mathbf{A}$  with the corresponding three  $\lambda$ ’s.)

In the present paper, we use these general ideas to find the ‘most classical states’ with a given intrinsic spin when the observables form the  $su(2)$  Lie algebra. In our next paper [11], we consider systems for which this is the Lie algebra  $e(n)$  of the Euclidean groups  $E(n)$  for  $n = 2, 3$ .

### 2.1 $SU(2)$ invariant elementary systems

The generators of the Lie algebra  $su(2)$  are denoted by  $\mathbf{J}_i$ ,  $i = 1, 2, 3$ , satisfying  $[\mathbf{J}_i, \mathbf{J}_j] = i\hbar \varepsilon_{ijk} \mathbf{J}^k$ , where the lowering and raising of the small Latin indices are defined by  $\delta_{ij}$  and its inverse, and  $\varepsilon_{ijk}$  is the alternating Levi-Civita symbol. Its only Casimir operator is  $\mathbf{J}_i \mathbf{J}^i$ .

We search for the general ‘most classical states’ in the form of superpositions of such states belonging to the *irreducible* representations of  $SU(2)$  with given, but arbitrary intrinsic spin. The finite dimensional irreducible representations of  $SU(2)$  and  $su(2)$  in their traditional, purely algebraic bra-ket formalism by unitary and self-adjoint operators, respectively, are well known. Nevertheless, the representations that we use here are less well known outside the general relativity community, and they are based on the complex line bundles  $\mathcal{O}(-2s)$  over the unit 2-sphere  $\mathcal{S} \approx S^2$ , where  $2s \in \mathbb{Z}$  is fixed. The integral of

the first Chern class of  $\mathcal{O}(-2s)$  is  $2s$ . This is a topological invariant and characterizing the global non-triviality (or ‘twist’) of the bundle, and it represents the *intrinsic spin* of the quantum mechanical system. The carrier space of such a representation is an appropriate *finite* dimensional subspace of the Hilbert space  $\mathcal{H} = L_2(\mathcal{S}, d\mathcal{S})$  of the square-integrable cross sections  $\phi$  of the line bundle, where the measure  $d\mathcal{S}$  is the natural metric area element on  $\mathcal{S}$ . The reason why we use this less well known form of the unitary, irreducible representations of  $su(2)$  is twofold: First, it is this form that fits smoothly to the most convenient representation of the  $E(3)$ -invariant elementary quantum mechanical systems that we will consider in our next paper [11], where the results of the present paper will also be used; and, second, the *intrinsic spin* of the system appears naturally in this formalism (see Appendix 4.1.4), though it is not linked to any Casimir operator.

The form of the angular momentum operators that we use are (densely defined) self-adjoint differential operators acting on the smooth cross sections of  $\mathcal{O}(-2s)$ . Explicitly, they are

$$\mathbf{J}^i \phi = s\hbar n^i \phi + \hbar(m^i \delta' \phi - \bar{m}^i \delta \phi), \quad (2.3)$$

where  $n^i$  are the Cartesian components of the unit normal of  $\mathcal{S}$  in  $\mathbb{R}^3$ ,  $m^i$  and  $\bar{m}^i$  are those of the complex null tangents of  $\mathcal{S}$ , and  $\delta$  and  $\delta'$ , the so-called edth operators, are the covariant directional derivative operators in the corresponding complex null directions. The spin weight  $s$  of  $\phi$ , i.e. the intrinsic spin of the system, will be fixed in what follows, and may take any integer or half-odd-integer value.

In this representation, the  $su(2)$  Casimir operator is  $\mathbf{J}_i \mathbf{J}^i \phi = s^2 \hbar^2 \phi - \hbar^2 (\delta \delta' + \delta' \delta) \phi$ , where  $(\delta \delta' + \delta' \delta)$  is just the familiar Laplacian on the unit 2-sphere. In Appendix 4.1, we summarize the construction and the key properties of the null tangents, the line bundles  $\mathcal{O}(-2s)$ , the operators  $\delta$  and  $\delta'$  and the related concepts, as well as the derivation of (2.3) and the form of the Casimir operator (see, in particular, equations (4.14)-(4.15)). A more detailed discussion of the line bundles  $\mathcal{O}(-2s)$  and the related concepts are given e.g. in [14] (see also [12, 13]).

## 2.2 The spectrum

For any  $\alpha^i, \beta^i \in \mathbb{R}^3$  satisfying  $\alpha^i \alpha^j \delta_{ij} = \beta^i \beta^j \delta_{ij} = 1$ , we form the operators  $\mathbf{J}(\alpha) := \alpha^i \mathbf{J}_i$  and  $\mathbf{J}(\beta) := \beta^i \mathbf{J}_i$ , i.e. the components of the angular momentum vector operator determined by the directions  $\alpha^i$  and  $\beta^i$ . However, without loss of generality, we may assume that e.g.  $\beta^1 = \beta^2 = 0$  and  $\beta^3 = 1$ , because by an appropriate rotation of the Cartesian coordinate system this can always be achieved. Moreover, we assume that  $\alpha_3 \neq \pm 1$ , because otherwise  $\alpha^i = \pm \beta^i$  would be allowed, and for these  $\mathbf{J}(\alpha) = \pm \mathbf{J}(\beta)$  would hold. Then by (2.1) and the commutation relations we have  $\Delta_\phi \mathbf{J}(\alpha) \Delta_\phi \mathbf{J}(\beta) \geq \frac{1}{2} \hbar |\alpha^i \beta^j \varepsilon_{ijk} \langle \mathbf{J}^k \rangle_\phi|$ , in which, by (2.2), the equality holds precisely when  $\phi$  solves the eigenvalue equation

$$(\mathbf{J}(\alpha) - i\lambda \mathbf{J}(\beta))\phi = \left( \langle \mathbf{J}(\alpha) \rangle_\phi - i\lambda \langle \mathbf{J}(\beta) \rangle_\phi \right) \phi =: \hbar C \phi \quad (2.4)$$

for some  $\lambda > 0$ . By (2.1) and (2.3) this eigenvalue equation is

$$(\alpha^i - i\lambda \beta^i) \left( m_i \delta' \phi - \bar{m}_i \delta \phi + s n_i \phi \right) = C \phi. \quad (2.5)$$

We solve this eigenvalue problem in two steps: First, using spinorial techniques of general relativity, we determine the spectrum of the operator  $\mathbf{J}(\alpha) - i\lambda \mathbf{J}(\beta)$ , and then, in the next subsection, we determine the eigenfunctions both as series and *in a closed form*, too.

Since  $\mathbf{J}_i \mathbf{J}^i$  and  $\mathbf{J}(\alpha) - i\lambda \mathbf{J}(\beta)$  are commuting, they have a system of common eigenfunctions. Hence, it might appear to be natural to search for the eigenfunctions of the latter in the form of a linear combination of the spin weighted spherical harmonics  ${}_s Y_{j,m}$ , where  $j = |s|, |s| + 1, |s| + 2, \dots$  and  $m = -j, -j + 1, \dots, j$  (see Appendix 4.1.1-4.1.3). However, the direct use of these harmonics in (2.5) would yield a coupled system of algebraic equations even for given  $j$ , and it would be difficult to get its solutions in a simple, closed form. Our strategy is based on the trick that,<sup>1</sup> instead of the Cartesian spinor basis (see Appendix 4.1.1), we use the *principal spinors* of the (unitary spinor form  $\gamma_{\underline{A}\underline{B}}$  of the) purely spatial complex vector  $\alpha^i - i\lambda\beta^i$ . This makes the calculation of the spectrum of  $\mathbf{J}(\alpha) - i\lambda\mathbf{J}(\beta)$  easier. Nevertheless, the basis of the spin space that these principal spinors define is *boosted* with respect to the Cartesian spin frame.

Thus, we form  $\gamma_{\underline{A}\underline{A}'} := (\alpha_i - i\lambda\beta_i)\sigma_{\underline{A}\underline{A}'}^i$ , and, since  $\alpha_i - i\lambda\beta_i$  is a spatial vector, its unitary spinor form,  $\gamma_{\underline{A}\underline{B}} := \gamma_{\underline{A}\underline{A}'}\sqrt{2}\sigma_{\underline{A}'\underline{B}}^0$ , is symmetric. (Here,  $\sigma_{\underline{A}\underline{A}'}^a = (\sigma_{\underline{A}\underline{A}'}^0, \sigma_{\underline{A}\underline{A}'}^i)$  are the standard  $SL(2, \mathbb{C})$  Pauli matrices. For our conventions, see Appendix 4.1.1.) Then (2.5) takes the form

$$\begin{aligned} C\phi &= \gamma_{\underline{A}\underline{A}'} o^{\underline{A}} \bar{t}^{\underline{A}'} \delta' \phi - \gamma_{\underline{A}\underline{A}'} t^{\underline{A}} \bar{o}^{\underline{A}'} \delta \phi + \sqrt{2}s \gamma_{\underline{A}\underline{A}'} o^{\underline{A}} \bar{o}^{\underline{A}'} \phi = \\ &= -\gamma_{\underline{A}\underline{B}} o^{\underline{A}} o^{\underline{B}} \delta' \phi - \gamma_{\underline{A}\underline{B}} t^{\underline{A}} t^{\underline{B}} \delta \phi + \sqrt{2}s \gamma_{\underline{A}\underline{B}} o^{\underline{A}} t^{\underline{B}} \phi. \end{aligned} \quad (2.6)$$

Next, for any given  $j$ , let us consider the function

$$\phi = \phi_{\underline{A}_1 \dots \underline{A}_{2j}} o^{\underline{A}_1} \dots o^{\underline{A}_{j+s}} t^{\underline{A}_{j+s+1}} \dots t^{\underline{A}_{2j}},$$

where the coefficients  $\phi_{\underline{A}_1 \dots \underline{A}_{2j}}$  are constant and completely symmetric in its indices. Recalling how the operators  $\delta$  and  $\delta'$  act on the spinors  $o^{\underline{A}}$  and  $t^{\underline{A}}$  (see Appendix 4.1.3) and how the operator  $\mathbf{J}_i \mathbf{J}^i$  is built from  $\delta$  and  $\delta'$  (see Appendix 4.1.4), it is easy to check that these functions are, in fact, eigenfunctions of  $\mathbf{J}_i \mathbf{J}^i$  with eigenvalue  $j(j+1)\hbar^2$ . Substituting this  $\phi$  into (2.6) and using that  $\varepsilon^{\underline{A}\underline{B}} = o^{\underline{A}} t^{\underline{B}} - t^{\underline{A}} o^{\underline{B}}$ , after some algebra we obtain

$$\begin{aligned} C\phi_{\underline{A}_1 \dots \underline{A}_{2j}} o^{\underline{A}_1} \dots o^{\underline{A}_{j+s}} t^{\underline{A}_{j+s+1}} \dots t^{\underline{A}_{2j}} &= \\ &= -\frac{1}{\sqrt{2}} \left( (j+s)\gamma_{\underline{B}\underline{A}_1} \phi_{\underline{A}_2 \dots \underline{A}_{2j}}^{\underline{B}} + (j-s)\gamma_{\underline{B}\underline{A}_{2j}} \phi_{\underline{A}_1 \dots \underline{A}_{2j-1}}^{\underline{B}} \right) o^{\underline{A}_1} \dots o^{\underline{A}_{j+s}} t^{\underline{A}_{j+s+1}} \dots t^{\underline{A}_{2j}}. \end{aligned} \quad (2.7)$$

Since  $\gamma_{\underline{A}\underline{B}}$  is symmetric, there are spinors  $\mu_{\underline{A}}$  and  $\nu_{\underline{A}}$ , the so-called principal spinors of  $\gamma_{\underline{A}\underline{B}}$ , such that  $\gamma_{\underline{A}\underline{B}} = \sqrt{2}\mu_{(\underline{A}}\nu_{\underline{B})}$ . Substituting this into (2.7) we find

$$C\phi_{\underline{A}_1 \dots \underline{A}_{2j}} = -j \left( \mu_{\underline{A}} \phi_{(\underline{A}_1 \dots \underline{A}_{2j-1}}^{\underline{A}} \nu_{\underline{A}_{2j})} + \nu_{\underline{A}} \phi_{(\underline{A}_1 \dots \underline{A}_{2j-1}}^{\underline{A}} \mu_{\underline{A}_{2j})} \right). \quad (2.8)$$

We solve this equation when  $\gamma_{\underline{A}\underline{B}}$  is not null ('generic case'), i.e. if  $\mu_{\underline{A}}$  and  $\nu_{\underline{A}}$  are not proportional to each other, and when  $\gamma_{\underline{A}\underline{B}}$  is null ('exceptional case') separately.

If  $\gamma_{\underline{A}\underline{B}}$  is not null, then  $0 \neq (\nu_{\underline{A}} \mu^{\underline{A}})^2 = -\gamma_{\underline{A}\underline{B}} \gamma^{\underline{A}\underline{B}} = -\gamma_{\underline{A}\underline{A}'} \gamma^{\underline{A}\underline{A}'} = (\alpha_i - i\lambda\beta_i)(\alpha^i - i\lambda\beta^i) = 1 - \lambda^2 - 2i\lambda\alpha_3$ , and  $\mu_{\underline{A}}$  and  $\nu_{\underline{A}}$  span the space of spinors. Hence the spinors of the form

$$\phi_{\underline{A}_1 \dots \underline{A}_{2j}} = \mu_{(\underline{A}_1} \dots \mu_{\underline{A}_{j-m}} \nu_{\underline{A}_{j-m+1}} \dots \nu_{\underline{A}_{2j})}, \quad (2.9)$$

---

<sup>1</sup>The idea of using the principal spinors of  $\gamma_{\underline{A}\underline{B}}$  instead of the Cartesian spinors is due to Paul Tod [19], and the next two paragraphs are based on his ideas and calculations.

$m = -j, -j + 1, \dots, j$ , form a basis in the space of the totally symmetric spinors of rank  $2j$  (see also Appendix 4.1.1). Substituting this into (2.8), we obtain

$$C\mu_{(\underline{A}_1 \cdots \underline{A}_{j-m} \nu_{\underline{A}_{j-m+1}} \cdots \nu_{\underline{A}_{2j}})} = m(\nu_{\underline{A}} \mu^{\underline{A}}) \mu_{(\underline{A}_1 \cdots \underline{A}_{j-m} \nu_{\underline{A}_{j-m+1}} \cdots \nu_{\underline{A}_{2j}})};$$

i.e. the eigenvalues are

$$C = m(\nu_{\underline{A}} \mu^{\underline{A}}) = \pm m \sqrt{1 - \lambda^2 - 2i\lambda\alpha_3}, \quad (2.10)$$

and the corresponding (not normalized) eigenfunctions are

$$\phi_{s,j,m} := \mu_{(\underline{A}_1 \cdots \underline{A}_{j-m} \nu_{\underline{A}_{j-m+1}} \cdots \nu_{\underline{A}_{2j}})} o^{\underline{A}_1} \dots o^{\underline{A}_{j+s}} \iota^{\underline{A}_{j+s+1}} \dots \iota^{\underline{A}_{2j}}. \quad (2.11)$$

For the sake of concreteness, in the rest of the paper, we will choose the *upper sign* in (2.10).

The principal spinors of  $\gamma_{\underline{A}\underline{B}}$  are determined only up to the scale ambiguity,  $(\mu_{\underline{A}}, \nu_{\underline{A}}) \mapsto (\chi\mu_{\underline{A}}, \chi^{-1}\nu_{\underline{A}})$ , where  $\chi$  is any non-zero complex constant; and up to their order. However,  $\nu_{\underline{A}}\mu^{\underline{A}}$  is invariant under the rescalings by  $\chi$ , while under the interchange of the principal spinors both  $\nu_{\underline{A}}\mu^{\underline{A}}$  and  $m$  change sign (see (2.9)). Therefore, the eigenvalue  $C$  is independent of these ambiguities (as it should be). Under such transformations the eigenfunctions change according to  $\phi_{s,j,m} \mapsto \chi^{-2m}\phi_{s,j,m}$  and  $\phi_{s,j,m} \mapsto \phi_{s,j,-m}$ , respectively. The first of these ambiguities is reduced to a phase ambiguity by the normalization of the eigenfunctions.

If  $\gamma_{\underline{A}\underline{B}} = \sqrt{2}\mu_{\underline{A}}\mu_{\underline{B}}$  (i.e. when  $\gamma_{\underline{A}\underline{B}}$  is null), then let us consider the spinor of the form

$$\phi_{\underline{A}_1 \dots \underline{A}_{2j}} = \mu_{(\underline{A}_1 \cdots \underline{A}_n \chi_{\underline{A}_{n+1} \dots \underline{A}_{2j}})},$$

where  $\chi_{\underline{A}_1 \dots \underline{A}_{2j-n}}$  is some totally symmetric spinor for which  $\mu^{\underline{A}_1} \chi_{\underline{A}_1 \dots \underline{A}_{2j-n}} \neq 0$ ,  $n = 0, 1, \dots, 2j$ . Substituting this into (2.8) we obtain

$$\begin{aligned} C\mu_{(\underline{A}_1 \cdots \underline{A}_n \chi_{\underline{A}_{n+1} \dots \underline{A}_{2j}})} &= (2j+1)\mu_{(\underline{A}_1 \cdots \underline{A}_{n+1} \chi_{\underline{A}_{n+2} \dots \underline{A}_{2j}\underline{A}})} \mu^{\underline{A}} = \\ &= (2j-n)\mu_{(\underline{A}_1 \cdots \underline{A}_{n+1} \chi_{\underline{A}_{n+2} \dots \underline{A}_{2j}\underline{A}})} \mu^{\underline{A}}. \end{aligned}$$

Since, however, the number of the spinors  $\mu_{\underline{A}}$  with free index on the left hand side is  $n$  while on the right it is  $n+1$ , this equation can hold true precisely when  $n = 2j$ ; i.e. the eigenvalue is zero,  $C = 0$ , the constant spinor  $\phi_{\underline{A}_1 \dots \underline{A}_{2j}}$  is null with  $\mu_{\underline{A}}$  as its  $2j$ -fold principal spinor, and the corresponding (not normalized) eigenfunction is

$$\phi_{s,j} := \mu_{\underline{A}_1} \cdots \mu_{\underline{A}_{2j}} o^{\underline{A}_1} \dots o^{\underline{A}_{j+s}} \iota^{\underline{A}_{j+s+1}} \dots \iota^{\underline{A}_{2j}}. \quad (2.12)$$

Thus, the solution of (2.8) in the exceptional case is *not* the special case of that in the generic case: while the eigenvalue is the  $m = 0$  special case of (2.10), the eigenfunction is the  $m = -j$  special case of (2.11).

Remarkably enough, the eigenvalue (2.10) does *not* depend directly on  $j$  or on  $s$ ; it depends on them only indirectly through  $m$  since  $m = -j, -j + 1, \dots, j$  and  $j = |s|, |s| + 1, \dots$ . Moreover, we got *no* restriction on the parameters  $\alpha_3$  and  $\lambda$ . Therefore, there are infinitely many eigenfunctions, labelled by  $j$ , with the given eigenvalue (2.10). Let  $\phi$  be *any* normalized solution of the eigenvalue equation (2.4) with the given eigenvalue. Then by (2.10)  $\langle \mathbf{J}(\alpha) \rangle_{\phi} - i\lambda \langle \mathbf{J}(\beta) \rangle_{\phi} = m\hbar\sqrt{1 - \lambda^2 - 2i\lambda\alpha_3}$  holds. Since both  $\mathbf{J}(\alpha)$  and  $\mathbf{J}(\beta)$

are self-adjoint, their expectation value must be real, and hence an elementary algebraic calculation yields that

$$\langle \mathbf{J}(\alpha) \rangle_\phi = m \frac{\hbar}{\sqrt{2}} \sqrt{1 - \lambda^2 + \sqrt{(1 - \lambda^2)^2 + 4\lambda^2 \alpha_3^2}}, \quad (2.13)$$

$$\lambda \langle \mathbf{J}(\beta) \rangle_\phi = m \operatorname{sign}(\alpha_3) \frac{\hbar}{\sqrt{2}} \sqrt{\lambda^2 - 1 + \sqrt{(1 - \lambda^2)^2 + 4\lambda^2 \alpha_3^2}} \quad (2.14)$$

if  $\alpha_3 \neq 0$ ; these are  $m\hbar\sqrt{1 - \lambda^2}$  and 0, respectively, if  $\alpha_3 = 0$  and  $\lambda < 1$ ; while these are 0 and  $m\hbar\sqrt{\lambda^2 - 1}$ , respectively, if  $\alpha_3 = 0$  and  $\lambda > 1$ . (Here, the overall  $\pm$  sign ambiguity coming from taking the square root of  $1 - \lambda^2 - 2i\lambda\alpha_3$  has been included in the ‘quantum number’  $m$ .) In the exceptional case both these expectation values are vanishing.

Therefore, apart from the discrete ‘quantum number’  $m$ , the expectation values are functions of two parameters:  $\alpha_3 \in (-1, 1)$ , as we expected, and  $\lambda \in (0, \infty)$ . The appearance of  $\lambda$  in the *expectation values* is a bit surprising, because, by its very meaning, it characterizes *the ratio of the two standard deviations*. Now we discuss the behaviour of the expectation values as functions of these two parameters.

First, for fixed non-zero  $m$ ,  $\langle \mathbf{J}(\alpha) \rangle_\phi$  and  $\langle \mathbf{J}(\beta) \rangle_\phi$ , as functions of  $\lambda$ , behave in a symmetric, but complementary way:  $|\langle \mathbf{J}(\alpha) \rangle_\phi| = |\langle \mathbf{J}(\beta) \rangle_\phi|$  precisely when  $\lambda = 1$ ; and in the  $\lambda \rightarrow 0$  limit  $\langle \mathbf{J}(\alpha) \rangle_\phi \rightarrow m\hbar$  and  $\langle \mathbf{J}(\beta) \rangle_\phi \rightarrow m\hbar\alpha_3$ ; while in the  $\lambda \rightarrow \infty$  limit  $\langle \mathbf{J}(\alpha) \rangle_\phi \rightarrow m\hbar|\alpha_3|$  and  $\langle \mathbf{J}(\beta) \rangle_\phi \rightarrow \operatorname{sign}(\alpha_3)m\hbar$ . The expectation values can be zero only if  $\alpha_3 = 0$  (see above); and, for  $\alpha_3 \neq 0$  and  $m \neq 0$ ,  $\langle \mathbf{J}(\alpha) \rangle_\phi/m$ , as a function of  $\lambda$ , is strictly monotonically *decreasing* from  $\hbar$  to  $\hbar\sqrt{|\alpha_3|}$ , while  $\operatorname{sign}(\alpha_3)\langle \mathbf{J}(\beta) \rangle_\phi/m$  is strictly monotonically *increasing* from  $\hbar\sqrt{|\alpha_3|}$  to  $\hbar$ . Both expectation values are smooth functions of  $\lambda$ .

On the other hand, also for fixed non-zero  $m$ , although  $\langle \mathbf{J}(\alpha) \rangle_\phi$  is a continuous function of  $\alpha_3$ , but it is *not differentiable* on the  $\lambda > 1$  portion of the  $\alpha_3 = 0$  line (on which it is vanishing).  $\langle \mathbf{J}(\beta) \rangle_\phi$  behaves just in the opposite way: it is *not continuous* at  $\alpha_3 = 0$  for  $\lambda > 1$ , its left/right limit is  $\mp m\hbar\sqrt{\lambda^2 - 1}/\lambda$ ; but its derivative with respect to  $\alpha_3$  is zero there both from the  $\alpha_3 > 0$  and the  $\alpha_3 < 0$  directions. Thus,  $\langle \mathbf{J}(\beta) \rangle_\phi$  *has a jump*, while its square,  $(\langle \mathbf{J}(\beta) \rangle_\phi)^2$ , is already differentiable.

It follows from these properties that, for given  $j$ , the range of the expectation values  $\langle \mathbf{J}(\alpha) \rangle_\phi$  and  $\langle \mathbf{J}(\beta) \rangle_\phi$  is a connected interval of  $\mathbb{R}$ , parameterized by  $\lambda$ , precisely when the directions  $\alpha^i$  and  $\beta^i$  are orthogonal to each other (see above). This case was considered by Aragone *et al* in [7]. If  $j$  may take any allowed value, then the range of the expectation values is the whole  $\mathbb{R}$ . If the two directions are not orthogonal to each other, then the range of them is a *union of intervals* of length  $|m|\hbar(1 - \sqrt{|\alpha_3|})$ ,  $m = -j, -j + 1, \dots, j$ . In the limit  $|\alpha_3| \rightarrow 1$  (i.e. when  $\mathbf{J}(\alpha) \rightarrow \pm\mathbf{J}(\beta)$ ), these intervals are getting to be disjoint and, ultimately, shrink to points. Thus, in this limit, the range of the expectation values reduces to the discrete (point) spectrum of the operator  $\mathbf{J}(\beta)$ . Therefore,  $\alpha_3$  provides the parameter in an *interpolation* between the purely continuous, connected range of the expectation values and the completely discrete spectrum of  $\mathbf{J}(\beta)$ .

We return to the discussion of the expectation values and the structure of the parameter space in subsection 2.5 and section 3.

## 2.3 The eigenfunctions

The eigenfunctions  $\phi_{s,j,m}$  and  $\phi_{s,j}$  given by (2.11) and (2.12) in the generic and exceptional cases, respectively, are not normalized. The corresponding normalized eigenfunctions

will be denoted by  $W_{s,j,m} := N_{s,j,m} \phi_{s,j,m}$  and  $W_{s,j} := N_{s,j} \phi_{s,j}$ , respectively, where the factors of normalization,  $N_{s,j,m}$  and  $N_{s,j}$ , (as well as the explicit coordinate form of the eigenfunctions) will be determined in Appendix 4.2.1 and 4.2.2. However, as we will see there, although the functions  $W_{s,j,m}$  and  $W_{s',j',m'}$  are  $L_2$ -orthogonal to each other if  $s$  and  $s'$  or  $j$  and  $j'$  are different, but  $W_{s,j,m}$  and  $W_{s,j,m'}$  are, in general, not. In fact, since (in contrast to  $\mathbf{J}_3$  or  $\mathbf{J}_i \mathbf{J}^i$ ) the operator  $\mathbf{J}(\alpha) - i\lambda \mathbf{J}(\beta)$  is *not* self-adjoint (and, in fact, not even normal), the orthogonality of its eigenfunctions with different eigenvalues does not follow.

Since the eigenfunctions  $W_{s,j,m}$  and  $W_{s,j}$  belong to the eigenvalue labelled by  $m$  and zero, respectively, the general solution  $\phi$  of the eigenvalue problem (2.5) with given  $s$  can be written as a combination of these functions:

$$\phi = \sum_{j=\max\{|s|,|m|\}}^{\infty} c^j W_{s,j,m}, \quad \text{or} \quad \sum_{j=|s|}^{\infty} c^j W_{s,j}, \quad (2.15)$$

respectively, where  $\sum_j |c^j|^2 = 1$ . Next we determine the eigenfunctions of the eigenvalue equation (2.5) *in closed form*.

Since, as we will see, all the functions in (2.5) are polynomial in the complex stereographic coordinates, and polynomial equations are easier to solve than trigonometrical ones, we search for its solutions in these coordinates. Also, since  $\alpha_i \alpha^i = 1$ , we can (and will) write  $(\alpha_1, \alpha_2) = \sqrt{1 - \alpha_3^2} (\cos \alpha, \sin \alpha)$ , where  $\alpha \in [0, 2\pi)$ .

Using  $\beta^i = (0, 0, 1)$  and the explicit form of  $m^i$  and the edth operators (see Appendix 4.1.2 and 4.1.3), equation (2.5) in the coordinate system  $(\zeta, \bar{\zeta})$  takes the form

$$\begin{aligned} & \left( \frac{1}{2} \sqrt{1 - \alpha_3^2} (1 - \exp[-2i\alpha] \zeta^2) + (\alpha_3 - i\lambda) \exp[-i\alpha] \zeta \right) \exp[i\alpha] \frac{\partial \ln \phi}{\partial \zeta} - \\ & - \left( \frac{1}{2} \sqrt{1 - \alpha_3^2} (1 - \exp[2i\alpha] \bar{\zeta}^2) + (\alpha_3 - i\lambda) \exp[i\alpha] \bar{\zeta} \right) \exp[-i\alpha] \frac{\partial \ln \phi}{\partial \bar{\zeta}} + \\ & + \frac{1}{2} s \left( \sqrt{1 - \alpha_3^2} (\exp[i\alpha] \bar{\zeta} + \exp[-i\alpha] \zeta) - 2(\alpha_3 - i\lambda) \right) = C. \end{aligned}$$

Thus, it seems useful to rotate the complex coordinates according to  $\zeta \mapsto \xi := \exp[-i\alpha] \zeta$ , and in the new coordinates  $(\xi, \bar{\xi})$  this equation simplifies to

$$\begin{aligned} & \left( \frac{1}{2} \sqrt{1 - \alpha_3^2} (1 - \xi^2) + (\alpha_3 - i\lambda) \xi \right) \frac{\partial \ln \phi}{\partial \xi} - \\ & - \left( \frac{1}{2} \sqrt{1 - \alpha_3^2} (1 - \bar{\xi}^2) + (\alpha_3 - i\lambda) \bar{\xi} \right) \frac{\partial \ln \phi}{\partial \bar{\xi}} = C + s(\alpha_3 - i\lambda) - \frac{1}{2} s \sqrt{1 - \alpha_3^2} (\xi + \bar{\xi}). \end{aligned}$$

Denoting the *complex* vector field on the left of this equation by  $X$ , this equation can be written in the compact form<sup>2</sup>

$$X(\ln \phi) = C - \frac{1}{2} s \left( \sqrt{1 - \alpha_3^2} \xi - \alpha_3 + i\lambda \right) - \frac{1}{2} s \left( \sqrt{1 - \alpha_3^2} \bar{\xi} - \alpha_3 + i\lambda \right), \quad (2.16)$$

i.e. the directional derivative of  $\ln \phi$  in the direction  $X$  is a given function on  $\mathcal{S}$ .

<sup>2</sup>Rewriting  $X$  in the polar coordinate system, it becomes clear that its real part is  $\lambda$ -times the rotation generator about  $\beta^i$ , while its imaginary part is the rotation generator about  $\alpha^i$ . Since  $\alpha^i \neq \pm \beta^i$ ,  $X$  does not have any zero on  $\mathcal{S}$ .



To find its solution, let us observe that

$$\begin{aligned} -(\sqrt{1 - \alpha_3^2} \xi - \alpha_3 + i\lambda) &= \frac{\partial}{\partial \xi} \left( \frac{1}{2} \sqrt{1 - \alpha_3^2} (1 - \xi^2) + (\alpha_3 - i\lambda) \xi \right) = \\ &= X \left( \ln \left( \frac{1}{2} \sqrt{1 - \alpha_3^2} (1 - \xi^2) + (\alpha_3 - i\lambda) \xi \right) \right), \end{aligned}$$

and hence the last two terms on the right of (2.16) can be written in the form

$$X \left( \ln \left( \frac{\sqrt{1 - \alpha_3^2} (1 - \xi^2) + 2(\alpha_3 - i\lambda) \xi}{\sqrt{1 - \alpha_3^2} (1 - \bar{\xi}^2) + 2(\alpha_3 - i\lambda) \bar{\xi}} \right)^{s/2} \right).$$

Introducing the notation

$$\xi_{\pm} := \frac{\alpha_3 - i\lambda \pm \sqrt{1 - \lambda^2 - 2i\lambda\alpha_3}}{\sqrt{1 - \alpha_3^2}}, \quad (2.17)$$

this can be rewritten into the remarkably simple form

$$X \left( \ln \left( \frac{\sqrt{1 - \alpha_3^2} (1 - \xi^2) + 2(\alpha_3 - i\lambda) \xi}{\sqrt{1 - \alpha_3^2} (1 - \bar{\xi}^2) + 2(\alpha_3 - i\lambda) \bar{\xi}} \right)^{s/2} \right) = X \left( \ln \left( \frac{(\xi - \xi_+)(\xi - \xi_-)}{(\bar{\xi} - \xi_+)(\bar{\xi} - \xi_-)} \right)^{s/2} \right); \quad (2.18)$$

while the vector field  $X$  is

$$X = -\frac{1}{2} \sqrt{1 - \alpha_3^2} \left( (\xi - \xi_+)(\xi - \xi_-) \frac{\partial}{\partial \xi} - (\bar{\xi} - \xi_+)(\bar{\xi} - \xi_-) \frac{\partial}{\partial \bar{\xi}} \right). \quad (2.19)$$

It might be worth noting that  $\xi_{\pm}$  is in a one-to-one correspondence with the expectation value given by (2.14):

$$|\xi_{\pm}|^2 = \frac{m\hbar \pm \langle \mathbf{J}_3 \rangle_{\phi}}{m\hbar \mp \langle \mathbf{J}_3 \rangle_{\phi}}, \quad (2.20)$$

which can be inverted to express  $\langle \mathbf{J}_3 \rangle_{\phi}$  by  $|\xi_{\pm}|^2$ .

If  $u$  were a function on  $\mathcal{S}$  for which  $X(u) = 1$  held, then the first term on the right of (2.16) would have the form  $X(Cu)$ , and hence we already would have found a particular solution of the *inhomogeneous equation* (2.16). We also need the general solution  $\phi_0$  of the corresponding *homogeneous equation*  $X(\ln \phi_0) = 0$ , i.e. of  $X(\phi_0) = 0$ . To find this  $u$ , let us observe that if the functions  $u_1 = u_1(\xi)$  and  $u_2 = u_2(\bar{\xi})$  solve

$$\frac{du_1}{d\xi} = -\frac{1}{\sqrt{1 - \alpha_3^2}} \frac{1}{(\xi - \xi_+)(\xi - \xi_-)}, \quad \frac{du_2}{d\bar{\xi}} = -\frac{1}{\sqrt{1 - \alpha_3^2}} \frac{1}{(\bar{\xi} - \xi_+)(\bar{\xi} - \xi_-)}, \quad (2.21)$$

respectively, then  $X(u_1 - u_2) = 1$  (and  $X(u_1 + u_2) = 0$ , too). However, since

$$\frac{d}{d\xi} \ln \left( \frac{\xi - \xi_+}{\xi - \xi_-} \right) = \frac{\xi_+ - \xi_-}{(\xi - \xi_+)(\xi - \xi_-)},$$

we find that

$$u_1 = -\frac{1}{2\sqrt{1 - \lambda^2 - 2i\lambda\alpha_3}} \ln \frac{\xi - \xi_+}{\xi - \xi_-}, \quad u_2 = -\frac{1}{2\sqrt{1 - \lambda^2 - 2i\lambda\alpha_3}} \ln \frac{\bar{\xi} - \xi_+}{\bar{\xi} - \xi_-}, \quad (2.22)$$

provided that  $1 - \lambda^2 - 2i\lambda\alpha_3 \neq 0$ , i.e. when  $(1 - \lambda)^2 + \alpha_3^2 > 0$  ('generic case'). Therefore, in the generic case,  $u = u_1 - u_2$  and the general *local* solution of (2.16) is

$$\phi = \phi_0 \left( \frac{(\xi - \xi_-)(\bar{\xi} - \xi_+)}{(\xi - \xi_+)(\bar{\xi} - \xi_-)} \right)^{m/2} \left( \frac{(\xi - \xi_+)(\xi - \xi_-)}{(\bar{\xi} - \xi_+)(\bar{\xi} - \xi_-)} \right)^{s/2}, \quad (2.23)$$

where we used the expression (2.10) (with the upper sign) for the eigenvalue  $C$ , and  $\phi_0$  is the general solution of the homogeneous equation. Clearly, an arbitrary complex function  $\phi_0$  of  $v := u_1 + u_2$  solves  $X(\phi_0) = 0$ ; and it is easy to see that this is, in fact, the *general* solution of  $X(\phi_0) = 0$ . Indeed, let us define  $\psi_0(u_1, u_2) := \phi_0(\xi, \bar{\xi})$  and observe that by (2.19)  $X(\phi_0) = 0$  is equivalent to

$$0 = (\xi - \xi_+)(\xi - \xi_-) \frac{\partial \phi_0}{\partial \xi} - (\bar{\xi} - \xi_+)(\bar{\xi} - \xi_-) \frac{\partial \phi_0}{\partial \bar{\xi}} = -\frac{\xi_+ - \xi_-}{2\sqrt{1 - \lambda^2 - 2i\lambda\alpha_3}} \left( \frac{\partial \psi_0}{\partial u_1} - \frac{\partial \psi_0}{\partial u_2} \right).$$

However, its general solution is an arbitrary complex valued smooth function of  $v := u_1 + u_2$ , i.e.  $\phi_0$  is an arbitrary complex valued smooth function of

$$w := \frac{(\xi - \xi_+)(\bar{\xi} - \xi_+)}{(\xi - \xi_-)(\bar{\xi} - \xi_-)}.$$

In particular,

$$\omega := \frac{(\xi - \xi_+)^a (\bar{\xi} - \xi_+)^a (\xi - \xi_-)^b (\bar{\xi} - \xi_-)^b}{(1 + \xi\bar{\xi})^{a+b}} = w^a \left( \frac{\xi_+ - \xi_-}{\xi_+ - \xi_- w} \right)^{a+b}$$

is a solution of  $X(\phi_0) = 0$  even with *arbitrary* real  $a$  and  $b$ . But with such a general  $\phi_0$  (2.23) is only a *local* solution of (2.5): We still have to ensure that  $\phi$  be well defined even on small circles surrounding the poles, and be square integrable, too. These requirements restrict the structure of  $\phi_0$ , and, in particular,  $a$  and  $b$  in  $\omega$  are restricted to be only integer or half-odd-integer. In particular, by (4.18) and (4.22)-(4.23) the eigenfunctions  $W_{s,j,m}$  are, in fact, combinations of functions of the form (2.23) with  $\phi_0 = \omega$  in which  $a$  and  $b$  are integer or half-odd-integer such that  $a + b = j$ .

If  $\lambda = 1$  and  $\alpha_3 = 0$  ('exceptional case'), then  $\xi_{\pm} = -i$  and the solution of equation (2.21) is given by

$$u_1 = \frac{1}{i + \xi}, \quad u_2 = \frac{1}{i + \bar{\xi}}. \quad (2.24)$$

Thus  $u = u_1 - u_2$ , and the *general* solution of  $X(\phi_0) = 0$  is an arbitrary complex function of  $v := u_1 + u_2$ . Hence, in the exceptional case, the *local* solution of (2.16) is

$$\phi = \phi_0 \left( \frac{i + \xi}{i + \bar{\xi}} \right)^s, \quad (2.25)$$

where we have used that in the exceptional case the eigenvalue  $C$  in (2.10) is zero. In particular,

$$\omega := \left( \frac{(\xi + i)(\bar{\xi} + i)}{(1 + \xi\bar{\xi})} \right)^a = \frac{1}{(1 - iv)^a}$$

is a solution of  $X(\phi_0) = 0$  even with *arbitrary* real  $a$ . This  $a$  is restricted to be integer or half-odd-integer by the requirement that the corresponding  $\phi$  be well defined and square integrable. In particular, by (4.19) and (4.24) the eigenfunctions  $W_{s,j}$  are combinations of functions of the form (2.25) with  $\phi_0 = \omega$  in which  $a = j$ .

Therefore, the eigenfunctions even with fixed eigenvalue depend on one (almost completely) free function; or, in their series expansion, on infinitely many freely specifiable complex constants  $c^j$  (see (2.15)).

## 2.4 The standard deviations

Taking the  $L_2$ -scalar product of the eigenvalue equation  $\mathbf{J}(\alpha)\phi = \hbar C\phi + i\lambda\mathbf{J}(\beta)\phi$  with itself and using the consequence  $\langle\phi, \mathbf{J}(\alpha)\phi\rangle - i\lambda\langle\phi, \mathbf{J}(\beta)\phi\rangle = \hbar C$  of the eigenvalue equation, we obtain

$$\langle\phi, (\mathbf{J}(\alpha))^2\phi\rangle = \langle\mathbf{J}(\alpha)\phi, \mathbf{J}(\alpha)\phi\rangle = \langle\phi, \mathbf{J}(\alpha)\phi\rangle^2 - \lambda^2\langle\phi, \mathbf{J}(\beta)\phi\rangle^2 + \lambda^2\langle\phi, (\mathbf{J}(\beta))^2\phi\rangle.$$

This does, in fact, imply that  $\Delta_\phi\mathbf{J}(\alpha) = \lambda\Delta_\phi\mathbf{J}(\beta)$ , yielding  $\Delta_\phi\mathbf{J}(\alpha)\Delta_\phi\mathbf{J}(\beta) = \lambda(\Delta_\phi\mathbf{J}(\beta))^2$ , too. Hence, we need to compute only  $\langle(\mathbf{J}(\beta))^2\rangle_\phi$ . Thus, let  $\phi$  be given by (2.15). Then

$$\langle(\mathbf{J}(\beta))^2\rangle_\phi = \langle\mathbf{J}(\beta)\phi, \mathbf{J}(\beta)\phi\rangle = \sum_{j,j'=\max\{|s|,|m|\}}^{\infty} \bar{c}^{j'} c^j N_{s,j',m} N_{s,j,m} \langle\mathbf{J}_3\phi_{s,j',m}, \mathbf{J}_3\phi_{s,j,m}\rangle.$$

Therefore, we should calculate  $\langle\mathbf{J}_3\phi_{s,j',m}, \mathbf{J}_3\phi_{s,j,m}\rangle$ . This, with the formal substitution  $m = -j$ , will give the general form of  $\langle\mathbf{J}_3\phi_{s,j'}, \mathbf{J}_3\phi_{s,j}\rangle$  in the exceptional case, too.

To calculate  $\langle\mathbf{J}_3\phi_{s,j',m}, \mathbf{J}_3\phi_{s,j,m}\rangle$ , we need to know the action of  $\mathbf{J}_3$  on the eigenfunctions  $\phi_{s,j,m}$ . Repeating the analysis behind the equations (2.7)-(2.8), we obtain that

$$\begin{aligned} \mathbf{J}_i\phi_{s,j,m} = \frac{\hbar}{\sqrt{2}} \left\{ (j+s)\sigma_i^B \underline{A} \phi_{B\underline{A}_2\dots\underline{A}_{2j}} o^A o^{A_2} \dots o^{A_{j+s}} t^{A_{j+s+1}} \dots t^{A_{2j}} + \right. \\ \left. + (j-s)\sigma_i^B \underline{A} \phi_{B\underline{A}_2\dots\underline{A}_{2j}} t^A o^{A_2} \dots o^{A_{j+s+1}} t^{A_{j+s+2}} \dots t^{A_{2j}} \right\}. \end{aligned} \quad (2.26)$$

As a by-product, this formula makes it possible to calculate all the expectation values  $\langle\mathbf{J}_i\rangle_\phi$ , too. In fact, using the techniques of Appendix 4.2.2, a straightforward calculation yields

$$\langle\phi_{s,j',m}, \mathbf{J}_i\phi_{s,j,m}\rangle = -(-)^{2j} \delta_{j'j} \sqrt{2} \hbar j 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} \phi_{B\underline{A}_2\dots\underline{A}_{2j}}^\dagger \sigma_i^B \underline{C} \phi_{C\underline{A}_2\dots\underline{A}_{2j}}, \quad (2.27)$$

where the adjoint  $\phi_{\underline{A}_1\dots\underline{A}_{2j}}^\dagger$  of the spinor  $\phi_{\underline{A}_1\dots\underline{A}_{2j}}$  is defined according to the convention  $\phi_{\underline{A}}^\dagger := \bar{\phi}_{\underline{A}}' \sqrt{2} \sigma_{0\underline{A}}^A$ . Since this is non-zero only if  $j' = j$ , the expectation value of  $\mathbf{J}_i$  in the state  $\phi$  given by (2.15) is simply  $\langle\mathbf{J}_i\rangle_\phi = \sum_{j=\max\{|s|,|m|\}}^{\infty} |c^j|^2 N_{s,j,m}^2 \langle\phi_{s,j,m}, \mathbf{J}_i\phi_{s,j,m}\rangle$ . Using (2.26) and the techniques of Appendix 4.2.2, the expectation values  $\langle\phi_{s,j,m}, \mathbf{J}_i\phi_{s,j,m}\rangle$  can be expressed as

$$\begin{aligned} \langle\phi_{s,j,m}, (\mathbf{J}_1 + i\mathbf{J}_2)\phi_{s,j,m}\rangle &= \hbar 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} \sum_{k=0}^{2j} (2j-k) \binom{2j}{k} \Phi_{k,j,m} \bar{\Phi}_{k+1,j,m}, \\ \langle\phi_{s,j,m}, \mathbf{J}_3\phi_{s,j,m}\rangle &= -\hbar 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} \sum_{k=0}^{2j} (j-k) \binom{2j}{k} |\Phi_{k,j,m}|^2, \end{aligned} \quad (2.28)$$

where, analogously to (2.11) (see also Appendix 4.1.1), we introduced the notation

$$\Phi_{k,j,m} := \phi_{\underline{A}_1\dots\underline{A}_{2j}} O^{\underline{A}_1} \dots O^{\underline{A}_k} I^{\underline{A}_{k+1}} \dots I^{\underline{A}_{2j}}. \quad (2.29)$$

These, together with the expression (4.28) for the factor of normalization and the explicit form (4.29) for  $\Phi_{k,j,m}$  given in Appendix 4.2.2, yield the general form for the expectation value of the operators  $\mathbf{J}_i$  in the state  $W_{s,j,m}$ . In particular, the latter for  $\mathbf{J}_3$  yields (2.14)

in that particular state. We will use this form of the expectation value of  $\mathbf{J}_3$  in Appendix 4.2.4.

Using (2.26), a bit longer but straightforward spinorial calculation, similar to that behind (2.27), yields that  $\langle \mathbf{J}_3 \phi_{s,j',m}, \mathbf{J}_3 \phi_{s,j,m} \rangle$  is zero unless  $j' = j$ , and that

$$\begin{aligned} \langle \mathbf{J}_3 \phi_{s,j,m}, \mathbf{J}_3 \phi_{s,j,m} \rangle &= (-)^{2j} j \hbar^2 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} \left( \beta_{\underline{C}}^B \beta_{\underline{D}}^C \phi_{\underline{B}A_2 \dots A_{2j}}^\dagger \phi^{\underline{D}A_2 \dots A_{2j}} + \right. \\ &\quad \left. + (2j-1) \beta_{\underline{D}}^B \phi_{\underline{E}CA_3 \dots A_{2j}}^\dagger \beta_{\underline{E}}^C \phi^{\underline{D}EA_3 \dots A_{2j}} \right) = \\ &= (-)^{2j} j \hbar^2 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} \left( j \phi_{\underline{A}_1 \dots A_{2j}}^\dagger \phi^{\underline{A}_1 \dots A_{2j}} + \right. \\ &\quad \left. + 2(2j-1) O^{\underline{A}} I_{\underline{A}B}^B \phi_{\underline{A}BA_3 \dots A_{2j}}^\dagger O_{\underline{C}} I_{\underline{D}} \phi^{\underline{C}DA_3 \dots A_{2j}} \right). \end{aligned}$$

Here, in the second step, we used  $I^{\underline{A}} O_{\underline{B}} - O^{\underline{A}} I_{\underline{B}} = \delta_{\underline{B}}^{\underline{A}}$  and  $2\beta_{\underline{A}C}^C \beta_{\underline{C}B} = -\delta_{\underline{B}}^{\underline{A}}$ , where  $\beta_{\underline{A}B}$  is the unitary spinor form of  $\beta_i$ . Then, using the definition of the adjoint  $\phi_{\underline{A}_1 \dots A_{2j}}^\dagger$ , we obtain

$$\begin{aligned} \langle \mathbf{J}_3 \phi_{s,j,m}, \mathbf{J}_3 \phi_{s,j,m} \rangle &= \\ &= \hbar^2 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} j^2 \sum_{k=0}^{2j} \binom{2j}{k} |\phi_{\underline{A}_1 \dots A_{2j}} O^{\underline{A}_1} \dots O^{\underline{A}_k} I^{\underline{A}_{k+1}} \dots I^{\underline{A}_{2j}}|^2 - \\ &\quad - \hbar^2 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} 2j(2j-1) \sum_{k=0}^{2j-2} \binom{2j-2}{k} \\ &\quad \times |O^{\underline{A}} I_{\underline{A}B}^B \phi_{\underline{A}BA_1 \dots A_{2j-2}} O^{\underline{A}_1} \dots O^{\underline{A}_k} I^{\underline{A}_{k+1}} \dots I^{\underline{A}_{2j-2}}|^2 = \\ &= \hbar^2 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} \sum_{k=0}^{2j} (j-k)^2 \binom{2j}{k} |\Phi_{k,j,m}|^2. \end{aligned} \quad (2.30)$$

Since for any normalized state  $\Delta_\phi \mathbf{J}(\alpha) \Delta_\phi \mathbf{J}(\beta) = \lambda (\Delta_\phi \mathbf{J}(\beta))^2$  holds, the normalization condition  $\sum_{j=\max\{|s|, |m|\}}^\infty |c^j|^2 = 1$  and equation (2.14) give that the ‘product uncertainty’ in the state  $\phi$  is

$$\Delta_\phi \mathbf{J}(\alpha) \Delta_\phi \mathbf{J}(\beta) = \sum_{j=\max\{|s|, |m|\}}^\infty |c^j|^2 \lambda \left( \langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle - \langle W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle^2 \right). \quad (2.31)$$

Hence, it is enough to evaluate the product uncertainty only in the normalized eigenstates  $W_{s,j,m}$ , i.e., by (2.30), to clarify the properties of  $\Phi_{k,j,m}$  only. This will be done in Appendix 4.2.2-4.2.3. In particular, in Appendix 4.2.3 we show that  $\langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$  does not depend on the sign of  $m$ , while  $\langle W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$  changes sign if  $m$  is replaced by  $-m$ . Moreover,  $\langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$  is continuous even at  $\alpha_3 = 0$  for  $\lambda > 1$ , while  $\langle W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$  has a jump there. Also, in Appendix 4.2.4, we determine the asymptotic behaviour of the standard deviations in the  $\lambda \rightarrow 0$  and  $\lambda \rightarrow \infty$  limits. We will see that the standard deviation for  $\mathbf{J}(\beta)$  is finite and that for  $\mathbf{J}(\alpha)$  tends to zero as  $\lambda$  if  $\lambda \rightarrow 0$ ; while the standard deviation for  $\mathbf{J}(\beta)$  tends to zero as  $1/\lambda$  and that for  $\mathbf{J}(\alpha)$  is finite if  $\lambda \rightarrow \infty$ . Hence, the product uncertainty tends to zero in both limits. Therefore, the uncertainties do *not* indicate any asymmetry between the  $\mathbf{J}(\alpha)$  and  $\mathbf{J}(\beta)$  angular momentum components, just as we expect it on physical grounds.

Finally, in the exceptional case, (2.30) together with the expression of  $\mu_A O^A$  and  $\mu_A I^A$  given in Appendix 4.2.2 yield  $\langle \mathbf{J}_3 W_{s,j}, \mathbf{J}_3 W_{s,j} \rangle = \frac{1}{2} j \hbar^2$ . Since  $\langle W_{s,j}, \mathbf{J}_3 W_{s,j} \rangle = 0$ , we find that the product uncertainty in the state  $W_{s,j}$  is  $\frac{1}{2} j \hbar^2$ . Hence, in the general eigenstate  $\phi = \sum_{j=|s|}^{\infty} c^j W_{s,j}$ , the product uncertainty is  $\frac{1}{2} \hbar^2 \sum_{j=|s|}^{\infty} j |c^j|^2$ , which can be zero only for  $j = |s| = 0$ .

## 2.5 The extension of the parameter space

Since physically  $\mathbf{J}(\alpha)$  and  $\mathbf{J}(\beta)$  are on equal footing, neither of them may be distinguished over the other. Indeed, the behaviour of the standard deviations do not break this symmetry between the two observables. However, the behaviour of the expectation values  $\langle \mathbf{J}(\alpha) \rangle_{\phi}$  and  $\langle \mathbf{J}(\beta) \rangle_{\phi}$  as functions on the *classical parameter space*  $\mathcal{P} := \{(\lambda, \alpha_3) | \lambda \in (0, \infty), \alpha_3 \in (-1, 1)\}$  apparently *does* break this symmetry. Hence, this odd behaviour indicates that either the expectation values should be extended to be *double valued* on  $\mathcal{P}$ , or they should in fact be functions on a *non-trivial Riemann surface*  $\mathcal{R}$  larger than  $\mathcal{P}$ , which must be homeomorphic to the one known in elementary complex analysis in connection with the function  $\sqrt{z}$ .

In fact, this second possibility seems more natural, because, for fixed  $m$  and apart from the factor  $\hbar$ , the complex eigenvalue of  $\mathbf{J}(\alpha) - i\lambda \mathbf{J}(\beta)$  is the *square root* of the complex function  $1 - \lambda^2 - 2i\lambda\alpha_3$  on  $\mathcal{P}$  (see equation (2.10)). The corresponding Riemann surface  $\mathcal{R}$  is obtained from  $\mathcal{P}$  by cutting it along the  $\lambda \geq 1$  part of the  $\alpha_3 = 0$  axis, and identifying the resulting edges for  $\lambda > 1$  with the opposite edges of another copy of  $\mathcal{P}$  that has been cut in the similar way. The branch point of the resulting Riemann surface is at  $\lambda = 1, \alpha_3 = 0$ , which corresponds just to the exceptional case of subsections 2.2 and 2.3. Then the two expectation values are extended from the first to the second copy of  $\mathcal{P}$  in  $\mathcal{R}$  just to be  $-1$  times the ones on the original ‘classical’ copy  $\mathcal{P}$ . With this extension the resulting expectation values behave in the same way, and, in particular, will be differentiable on the whole of  $\mathcal{R}$ .

## 3 On the simultaneous measurement of $\mathbf{J}(\alpha)$ and $\mathbf{J}(\beta)$

In the pioneering work [3], Arthurs and Kelly raised the possibility of the *simultaneous* measurement of the conjugate, and hence *not commuting*, observables in a Heisenberg system. Soon later, it was argued [4] that *every* measurement of the momentum or of the position is, in fact, such a simultaneous measurement, though while we measure one quantity precisely, we measure the conjugate one rather imprecisely. E.g. when we do a precise measurement of the momentum of a particle, then the particle must be in the measuring apparatus (or at least in the laboratory), so we do a rude position measurement as well; and when we measure the position of the particle accurately and the measuring apparatus is not destroyed in the measuring procedure, then we can be sure that the particle’s momentum could not have arbitrarily large momentum, so we did a rude momentum measurement, too. This idea of simultaneous measurement of the conjugate observables was discussed further e.g. in [4, 20, 21], clarifying, in particular, the precise relationship between the errors of the measurements and the standard deviations, deriving inequalities for the former, etc.

The basis of the possibility of such simultaneous measurements is that the non-commuting observables are *different* kinds of quantities: to measure them we need different devices, and during the simultaneous running of them one does not disturb the other.

If, however, the two non-commuting observables are of the *same* kind, e.g. the  $\mathbf{J}(\alpha)$  and  $\mathbf{J}(\beta)$  components of angular momentum, then it does not seem to be possible to carry out such a measurement *in a direct way*. For example, in the Stern–Gerlach apparatus we cannot have two different magnetic fields at the same time to measure the spin of the particle in the corresponding two different directions.

Nevertheless, the phenomenon of the quantum entanglement makes it possible, at least in principle, to measure  $\mathbf{J}(\alpha)$  and  $\mathbf{J}(\beta)$  simultaneously *in an indirect way*. In fact, it is known that even the coherent spin states can be entangled [8], and one particle in its most classical state can be maximally entangled with a second one. Thus, preparing two such particles to be maximally entangled and then spatially separating them, we can measure  $\mathbf{J}(\alpha)$  for one, and  $\mathbf{J}(\beta)$  for the second in two independent Stern–Gerlach apparatus.

In this way, in principle, it can be experimentally verify whether or not the proper parameter space preferred by the Nature is the classical  $\mathcal{P}$  or the Riemann surface  $\mathcal{R}$ . (The parameter  $\alpha_3$  has the obvious meaning in the experimental apparatus, and the  $\lambda$  can also be controlled in an indirect way since it is the quotient of the two standard deviations.) Indeed, let us consider the closed path in the classical parameter space  $\mathcal{P}$  consisting of the following four straight line segments:

1. the initial point is given by  $\lambda = 2$  and  $\alpha_3 = 1/\sqrt{2}$  (i.e.  $\Delta_\phi \mathbf{J}(\alpha)$  is just twice of  $\Delta_\phi \mathbf{J}(\beta)$  and the angle between the directions  $\alpha^i$  and  $\beta^i$  is  $\pi/4$ ), and the end point is defined by  $\lambda = 1/2$  and  $\alpha_3 = 1/\sqrt{2}$ ;
2. the end point of the second segment is at  $\lambda = 1/2$ ,  $\alpha_3 = -1/\sqrt{2}$  (i.e. we enlarge the angle between  $\alpha^i$  and  $\beta^i$  from  $\pi/4$  to  $3\pi/4$  while keeping  $\lambda$  to be  $1/2$ );
3. the end point of the third segment is at  $\lambda = 2$ ,  $\alpha_3 = -1/\sqrt{2}$ ;
4. the fourth segment closes the path, returning to the initial point of the first segment.

Next consider a 1-parameter family of the most classical states with fixed  $m$  and parameterized by the parameter of this closed path, and measure the two expectation values. If we find that *both* expectation values change continuously along the closed part from their initial value at the initial point of the path to their own negative at the end point of the path, then the genuine parameter space is the Riemann surface  $\mathcal{R}$ ; but if *only one* changes continuously along the path but the other had a discontinuity (or, equivalently, one does *not* change sign along the closed path but the other does), then the Nature’s parameter space is the classical  $\mathcal{P}$ .

## 4 Appendix

### 4.1 The irreducible representations of $su(2)$ by spin weighted functions

#### 4.1.1 An algebraic introduction of the spin weighted spherical harmonics

As is well known, any finite dimensional irreducible (and hence unitary) representation of  $SU(2)$  can be labelled by a non-negative integer or half-odd-integer  $j$ , i.e. for which  $2j = 0, 1, 2, \dots$ , and the carrier space  $\mathcal{H}_j$  of such a representation is a  $2j + 1$  dimensional

Hilbert space, in which the vectors of an orthonormal basis is usually denoted by  $|j, m\rangle$ , where  $m = -j, -j + 1, \dots, j$ .

One concrete realization of this abstract Hilbert space is the space  $\mathbb{S}_{\underline{A}_1 \dots \underline{A}_{2j}}$  of the completely symmetric spinors  $\phi_{\underline{A}_1 \dots \underline{A}_{2j}}$  of rank  $2j$ , i.e. for which  $\phi_{\underline{A}_1 \dots \underline{A}_{2j}} = \phi_{(\underline{A}_1 \dots \underline{A}_{2j})}$ . If  $\{O_{\underline{A}}, I_{\underline{A}}\}$  is any basis in  $\mathbb{S}_{\underline{A}}$ , then the spinors of the form  $Z(j, m)_{\underline{A}_1 \dots \underline{A}_{2j}} := O_{(\underline{A}_1 \dots \underline{A}_{j-m}} I_{\underline{A}_{j-m+1}} \dots I_{\underline{A}_{2j})}$  form a basis in  $\mathbb{S}_{\underline{A}_1 \dots \underline{A}_{2j}}$ . If the basis  $\{O_{\underline{A}}, I_{\underline{A}}\}$  is chosen to be normalized with respect to the (anti-symmetric) symplectic metric  $\varepsilon^{\underline{A}\underline{B}}$ , i.e. for which  $O_{\underline{A}} I_{\underline{A}} = \varepsilon^{\underline{A}\underline{B}} O_{\underline{A}} I_{\underline{B}} = 1$ , then with the definition  $\sqrt{2}T_{\underline{A}\underline{A}'} := O_{\underline{A}} \bar{O}_{\underline{A}'} + I_{\underline{A}} \bar{I}_{\underline{A}'}$  one has  $T_{\underline{A}\underline{A}'} \bar{T}^{\underline{A}\underline{A}'} = 1$ ,  $\sqrt{2}T^{\underline{A}\underline{A}'}$  plays the role of a positive definite Hermitean metric on  $\mathbb{S}_{\underline{A}}$ , and  $T_{\underline{A}\underline{A}'} \bar{O}^{\underline{A}'} = -\frac{1}{\sqrt{2}} I_{\underline{A}}$  and  $T_{\underline{A}\underline{A}'} \bar{I}^{\underline{A}'} = \frac{1}{\sqrt{2}} O_{\underline{A}}$  hold. With the choice  $O^{\underline{A}} = \delta_0^{\underline{A}}$ ,  $I^{\underline{A}} = \delta_1^{\underline{A}}$  this Hermitean metric is just the unit matrix  $\sqrt{2}\sigma_{\underline{A}\underline{A}'}^0$ , where  $\sigma_{\underline{A}\underline{A}'}^0$  denotes the zeroth of the four  $SL(2, \mathbb{C})$  Pauli matrices  $\sigma_{\underline{A}\underline{A}'}^a$  (including the factor  $1/\sqrt{2}$ ), according to the conventions of [12]. Hence, with respect to this scalar product, the basis  $\{O_{\underline{A}}, I_{\underline{A}}\}$  is orthonormal. We call such a basis a Cartesian spin frame. This scalar product makes it possible to normalize the basis vectors  $Z(j, m)_{\underline{A}_1 \dots \underline{A}_{2j}}$ . In fact, the vectors

$$\sqrt{\frac{(2j)!}{(j+m)!(j-m)!}} Z(j, m)_{\underline{A}_1 \dots \underline{A}_{2j}}$$

have not only unit norm with respect to  $\sqrt{2}T^{\underline{A}\underline{A}'}$ , but they are orthogonal to each other as well. Thus they can be identified with the abstract basis vectors  $|j, m\rangle$  in  $\mathcal{H}_j$  above.

Another basis in the space  $\mathbb{S}_{\underline{A}_1 \dots \underline{A}_{2j}}$  is provided by the so-called Newman–Penrose spin frame  $\{o_{\underline{A}}, \iota_{\underline{A}}\}$ . Such a frame can be chosen to be

$$o_{\underline{A}} = \frac{-i}{\sqrt{1 + \zeta\bar{\zeta}}} (\zeta O_{\underline{A}} + I_{\underline{A}}), \quad \iota_{\underline{A}} = \frac{-i}{\sqrt{1 + \zeta\bar{\zeta}}} (O_{\underline{A}} - \bar{\zeta} I_{\underline{A}}), \quad (4.1)$$

where  $\zeta \in \mathbb{C}$  (see e.g. [12]). The matrix of the basis transformation  $\{O_{\underline{A}}, I_{\underline{A}}\} \mapsto \{o_{\underline{A}}, \iota_{\underline{A}}\}$  is an  $SU(2)$  matrix. Hence, in particular,  $\sqrt{2}t_{\underline{A}\underline{A}'} := o_{\underline{A}} \bar{o}_{\underline{A}'} + \iota_{\underline{A}} \bar{\iota}_{\underline{A}'} = O_{\underline{A}} \bar{O}_{\underline{A}'} + I_{\underline{A}} \bar{I}_{\underline{A}'} = \sqrt{2}T_{\underline{A}\underline{A}'}$ . The frame  $\{o_{\underline{A}}, \iota_{\underline{A}}\}$  is a two-real-parameter family of bases, which can be considered to be defined on the (e.g. unit) 2-sphere  $\mathcal{S}$ , where  $\zeta$  is the complex stereographic coordinate on the 2-sphere (see the next subsection). Hence, the spinors of the form  $z(j, s)_{\underline{A}_1 \dots \underline{A}_{2j}} := o_{(\underline{A}_1 \dots \underline{A}_{j+s}} \iota_{\underline{A}_{j+s+1}} \dots \iota_{\underline{A}_{2j})}$ ,  $s = -j, -j + 1, \dots, j$ , form another basis in  $\mathbb{S}_{\underline{A}_1 \dots \underline{A}_{2j}}$ . These basis vectors can also be normalized, and

$$\sqrt{\frac{(2j)!}{(j-s)!(j+s)!}} z(j, s)_{\underline{A}_1 \dots \underline{A}_{2j}}$$

form another orthonormal basis with respect to  $\sqrt{2}T^{\underline{A}\underline{A}'}$ . Hence, the contractions

$$U(j)_{m,s} := \frac{(2j)!}{\sqrt{(j+m)!(j-m)!(j-s)!(j+s)!}} Z(j, m)_{\underline{A}_1 \dots \underline{A}_{2j}} z(j, s)^{\underline{A}_1 \dots \underline{A}_{2j}}$$

form a  $(2j + 1) \times (2j + 1)$  unitary matrix, taking one orthonormal basis of  $\mathbb{S}_{\underline{A}_1 \dots \underline{A}_{2j}}$  to another one. Apart from a numerical coefficient, the familiar spin weighted spherical harmonics are just the components of this matrix:

$${}_s Y_{j,m} := (-)^{j+m} \sqrt{\frac{2j+1}{4\pi}} U(j)_{m,s}, \quad (4.2)$$

where the numerical coefficient makes these to be normalized with respect to the  $L_2$ -scalar product on the unit sphere. For further properties of these harmonics, see [15, 16, 12, 13].

### 4.1.2 Spinorial coordinates on $\mathcal{S}$

Let  $p^i$ ,  $i = 1, 2, 3$ , denote Cartesian coordinates in  $\mathbb{R}^3$ , in which the components of the independent rotation Killing vector fields are  $k_{mn}^i = p^j(\delta_{jn}\delta_m^i - \delta_{jm}\delta_n^i)$ ,  $m, n = 1, 2, 3$ . Thus here  $i$  is the *vector* index, while  $m$  and  $n$  are the *name* indices of the Killing fields. These are tangent to the 2-spheres of radius  $P$ ,  $\mathcal{S} := \{p^i \in \mathbb{R}^3 | P^2 := \delta_{ij}p^i p^j = \text{const}\}$ , and vanish at the point  $p^i = 0$ . These Killing fields with the Lie bracket generate the Lie algebra  $su(2)$ .

The complex stereographic coordinates, projected from the *north* pole, are defined on  $U_n := \mathcal{S} - \{(0, 0, P)\}$ , the sphere minus its *north* pole, by  $\zeta := \exp(i\varphi) \cot(\theta/2)$ , where  $(\theta, \varphi)$  are the standard spherical polar coordinates. In terms of  $(\zeta, \bar{\zeta})$ , the Cartesian coordinates of the point  $p^i \in U_n$  are

$$p^i = P \left( \frac{\bar{\zeta} + \zeta}{1 + \zeta\bar{\zeta}}, i \frac{\bar{\zeta} - \zeta}{1 + \zeta\bar{\zeta}}, \frac{\zeta\bar{\zeta} - 1}{1 + \zeta\bar{\zeta}} \right). \quad (4.3)$$

The outward pointing unit normal to  $\mathcal{S}$  at the point  $p^i$  is  $n^i := p^i/P$ . This normal is completed to be a basis by the complex vector field

$$m^i := \frac{1}{\sqrt{2}} \left( \frac{1 - \zeta^2}{1 + \zeta\bar{\zeta}}, i \frac{1 + \zeta^2}{1 + \zeta\bar{\zeta}}, \frac{2\zeta}{1 + \zeta\bar{\zeta}} \right) \quad (4.4)$$

and its complex conjugate  $\bar{m}^i$ . These are orthogonal to  $n^i$ , null (i.e.  $m^i m_i = 0$ ), normalized with respect to each other (i.e.  $m^i \bar{m}_i = 1$ ), and  $p^i m^j \bar{m}^k \varepsilon_{ijk} = iP$  holds. (Recall that, in the present paper, the metric on  $\mathbb{R}^3$  is chosen to be the *positive definite*  $\delta_{ij}$ , rather than the negative definite spatial part of the Minkowski metric  $\eta_{\underline{a}\underline{b}} := \text{diag}(1, -1, -1, -1)$ , where  $\underline{a}, \underline{b} = 0, i$ .) In fact, apart from a phase factor, the complex null vectors  $m^i$  and  $\bar{m}^i$  are uniquely determined by the intrinsic complex structure of  $\mathcal{S}$  (see e.g. [13]). The vector field  $m^i$ , as a differential operator, is given by

$$m^i \left( \frac{\partial}{\partial p^i} \right) = \frac{1}{\sqrt{2}P} (1 + \zeta\bar{\zeta}) \left( \frac{\partial}{\partial \zeta} \right). \quad (4.5)$$

Hence,  $\zeta$  is a local *anti-holomorphic* coordinate on  $U_n$ . Also in these coordinates, the line element of the metric and the corresponding area element on  $\mathcal{S}$  of radius  $P$ , respectively, are

$$dh^2 = \frac{4P^2}{(1 + \zeta\bar{\zeta})^2} d\zeta d\bar{\zeta}, \quad d\mathcal{S} = \frac{-2iP^2}{(1 + \zeta\bar{\zeta})^2} d\zeta \wedge d\bar{\zeta}. \quad (4.6)$$

These are just the metric and area element inherited from the metric and volume element of  $\mathbb{R}^3$ , respectively. There are analogous constructions on  $U_s := \mathcal{S} - \{(0, 0, -P)\}$ , on the 2-sphere minus the *south* pole, too; and the structures defined on  $U_n$  are related to those introduced on  $U_s$  smoothly on the overlap  $U_n \cap U_s$ .

Considering  $\mathbb{R}^3$  to be the  $p^0 = P$  hyperplane of the Minkowski space  $\mathbb{R}^{1,3}$  with the Cartesian coordinates  $p^{\underline{a}} = (p^0, p^i)$  and the flat metric  $\eta_{\underline{a}\underline{b}}$ , the 2-sphere  $\mathcal{S}$  is just the intersection of the  $p^0 = P$  hyperplane with the null cone of the origin  $p^{\underline{a}} = 0$  in  $\mathbb{R}^{3,1}$ . Hence, for any  $p^i \in \mathcal{S}$ , there is a spinor  $\pi^{\underline{A}}$ , the ‘spinor constituent’ of  $p^i$ , such that  $p^i = \sigma_{\underline{A}\underline{A}}^i \pi^{\underline{A}} \bar{\pi}^{\underline{A}'}$ , and  $\pi^{\underline{A}}$  is unique only up to the phase ambiguity  $\pi^{\underline{A}} \mapsto \exp(i\gamma) \pi^{\underline{A}}$ ,  $\gamma \in [0, 2\pi)$ . (Note that we lower and raise the small Latin indices by the *positive definite*  $\delta_{ij}$  and its inverse.) Since  $P^2 = \delta_{ij} p^i p^j = -\eta_{ij} \sigma_{\underline{A}\underline{A}}^i \sigma_{\underline{B}\underline{B}}^j \pi^{\underline{A}} \bar{\pi}^{\underline{A}'} \pi^{\underline{B}} \bar{\pi}^{\underline{B}'} = -(\eta_{\underline{a}\underline{b}} \sigma_{\underline{A}\underline{A}}^{\underline{a}} \sigma_{\underline{B}\underline{B}}^{\underline{b}} - \sigma_{\underline{A}\underline{A}}^0 \sigma_{\underline{B}\underline{B}}^0) \pi^{\underline{A}} \bar{\pi}^{\underline{A}'} \pi^{\underline{B}} \bar{\pi}^{\underline{B}'} = (\sigma_{\underline{A}\underline{A}}^0 \pi^{\underline{A}} \bar{\pi}^{\underline{A}'})^2$ , the norm of  $\pi^{\underline{A}}$  with respect to  $\sigma_{\underline{A}\underline{A}}^0$ , i.e. to



$T_{\underline{A}\underline{A}'}$  of the previous subsection, is  $\sqrt{P}$ . However,  $\pi^{\underline{A}}$  as a spinor *field* is well defined only on  $\mathcal{S}$  *minus one point* (see e.g. [13, 14]). In particular, on  $U_n$ , this spinor field, up to a phase, is  $\pi^{\underline{A}} = (\sqrt{2}P)^{1/2}o^{\underline{A}}$ , where  $o^{\underline{A}}$  is given by (4.1). Thus  $\pi^{\underline{A}}$  on  $U_n$  and the analogous one on  $U_s$  are only *locally defined* ‘spinorial coordinates’ on  $\mathcal{S}$ .

Using the Newman–Penrose spinor basis  $\{o^{\underline{A}}, \iota^{\underline{A}}\}$ , given explicitly on  $U_n$  by (4.1) with the choice  $O^{\underline{A}} = \delta_0^{\underline{A}}$  and  $I^{\underline{A}} = \delta_1^{\underline{A}}$ , it is easy to verify that  $m^i \sigma_i^{\underline{A}\underline{A}'} = -o^{\underline{A}}\bar{\iota}^{\underline{A}'}$  and  $p^i \sigma_i^{\underline{A}\underline{A}'} = P(\iota^{\underline{A}}\bar{\iota}^{\underline{A}'} - o^{\underline{A}}\bar{o}^{\underline{A}'})/\sqrt{2}$  also hold.

### 4.1.3 The line bundles $\mathcal{O}(-2s)$ over $\mathcal{S}$

One way of defining the complex line bundles  $\mathcal{O}(-2s)$  over  $\mathcal{S}$  can be based on the concept of the bundle of totally symmetric N-type spinors of rank  $2|s|$  on  $\mathcal{S}$ : if  $s = -|s| \leq 0$  then these spinors are *unprimed* and their principal spinor at the point  $p^i = \sigma_{\underline{A}\underline{A}'}^i \pi^{\underline{A}} \bar{\pi}^{\underline{A}'}$  is  $\pi^{\underline{A}}$ ; and if  $s = |s| > 0$  then the spinors are *primed* and their principal spinor is  $\bar{\pi}^{\underline{A}'}$ . (Recall that e.g.  $\lambda^{\underline{A}}$  is a  $2|s|$ -fold principal spinor of the totally symmetric spinor  $\phi^{\underline{A}_1 \dots \underline{A}_{2|s|}}$  if  $\phi^{\underline{A}_1 \dots \underline{A}_{2|s|}} \lambda_{\underline{A}_1} = 0$  holds, in which case  $\phi^{\underline{A}_1 \dots \underline{A}_{2|s|}}$  necessarily has the form  $\phi \lambda^{\underline{A}_1} \dots \lambda^{\underline{A}_{2|s|}}$  for some  $\phi$ ; and the Penrose–Petrov type of the spinor is called null or of type N, see e.g. [12, 13].) Hence, e.g. on the domain  $U_n$ , these spinors have the form  $\phi^{\underline{A}_1 \dots \underline{A}_{2|s|}} = \phi \pi^{\underline{A}_1} \dots \pi^{\underline{A}_{2|s|}}$  and  $\chi^{\underline{A}'_1 \dots \underline{A}'_{2|s|}} = \chi \bar{\pi}^{\underline{A}'_1} \dots \bar{\pi}^{\underline{A}'_{2|s|}}$ , where  $\phi$  and  $\chi$  are complex functions on  $U_n$ . Thus, the fibers of these bundles are one complex dimensional, and the line bundle  $\mathcal{O}(-2s)$  is just the abstract bundle of these fibers over  $\mathcal{S}$ .  $U_n$  and  $U_s$  are local trivialization domains of  $\mathcal{O}(-2s)$ , and the functions  $\phi$  for  $s = -|s|$  (and  $\chi$  for  $s = |s|$ ) are *local* cross sections of  $\mathcal{O}(-2s)$  on  $U_n$ .  $\mathcal{O}(-2s)$  is globally trivializable precisely when  $s = 0$ .

The phase ambiguity  $\pi^{\underline{A}} \mapsto \exp(i\gamma)\pi^{\underline{A}}$  in the principal spinor yields the ambiguity  $\phi \mapsto \exp(-2i|s|\gamma)\phi$ , where  $\gamma$  is an arbitrary  $[0, 2\pi)$ -valued locally defined function. The analogous ambiguity in the function  $\chi$  is  $\chi \mapsto \exp(2is\gamma)\chi$ . Therefore, in spite of this ambiguity, the Hermitian scalar product of any two global cross sections, representing e.g.  $\phi^{\underline{A}_1 \dots \underline{A}_{2|s|}}$  and  $\psi^{\underline{A}_1 \dots \underline{A}_{2|s|}}$  and given by

$$\langle \phi^{\underline{A}_1 \dots \underline{A}_{2|s|}}, \psi^{\underline{A}_1 \dots \underline{A}_{2|s|}} \rangle_s := \int_{\mathcal{S}} \bar{\phi} \psi d\mathcal{S}, \quad (4.7)$$

is well defined. The space of the square-integrable cross sections of  $\mathcal{O}(-2s)$  is a Hilbert space, and is also denoted by  $\mathcal{H}$ . An alternative, and perhaps more familiar form of this scalar product can be given in terms of the spinor fields themselves. To rewrite (4.7) in this form, let us recall from subsection 4.1.2 that  $\sigma_{\underline{A}\underline{A}'}^0 = (O_{\underline{A}} \bar{O}_{\underline{A}'} + I_{\underline{A}} \bar{I}_{\underline{A}'})/\sqrt{2} = (o_{\underline{A}} \bar{o}_{\underline{A}'} + \iota_{\underline{A}} \bar{\iota}_{\underline{A}'})/\sqrt{2}$ . Using this, the integrand of (4.7) can be rewritten as

$$\begin{aligned} \bar{\phi} \psi &= (\sqrt{2}P)^{-2|s|} \bar{\iota}_{\underline{A}'_1} \dots \bar{\iota}_{\underline{A}'_{2|s|}} \bar{\phi}^{\underline{A}'_1 \dots \underline{A}'_{2|s|}} \iota_{\underline{A}_1} \dots \iota_{\underline{A}_{2|s|}} \psi^{\underline{A}_1 \dots \underline{A}_{2|s|}} = \\ &= P^{-2|s|} \sigma_{\underline{A}_1 \underline{A}'_1}^0 \dots \sigma_{\underline{A}_{2|s|} \underline{A}'_{2|s|}}^0 \bar{\phi}^{\underline{A}'_1 \dots \underline{A}'_{2|s|}} \psi^{\underline{A}_1 \dots \underline{A}_{2|s|}}. \end{aligned} \quad (4.8)$$

If we think of  $\sigma_{\underline{A}\underline{A}'}^0$  as the components of the spinor form of the timelike vector of the orthonormal vector basis in a Lorentzian vector space whose spatial vectors span  $\mathbb{R}^3$ , then by (4.8) the scalar product (4.7) is just  $(\sqrt{2}P)^{-2|s|}$  times the familiar, standard  $L_2$ -scalar product of the two spinor fields (see [22]).

Recalling that a scalar  $\phi$  is said to have the spin weight  $\frac{1}{2}(p-q)$  if under the rescaling  $\{o^{\underline{A}}, \iota^{\underline{A}}\} \mapsto \{\lambda o^{\underline{A}}, \lambda^{-1} \iota^{\underline{A}}\}$ , where  $\lambda$  is any nowhere vanishing complex function on the

domain of the spin frame, the scalar  $\phi$  transforms as  $\phi \mapsto \lambda^p \bar{\lambda}^q \phi$  (see e.g. [12, 13]), we can see that  $\mathcal{O}(-2s)$  is just the bundle of spin weighted scalars of weight  $s$  on  $\mathcal{S}$ . In particular, the components of the vectors  $m^i$  and  $\bar{m}^i$  are of type  $(1, -1)$  and  $(-1, 1)$ , respectively, while  $p^i$  is a sum of a  $(1, 1)$  and a  $(-1, -1)$  type scalar. Thus, the spin weight of  $m^i$ ,  $\bar{m}^i$  and  $p^i$  is 1,  $-1$  and 0, respectively; and the spin weight of the spherical harmonics  ${}_s Y_{j,m}$ , defined by (4.2), is  $s$ .

If  $\delta_i$  denotes the (Cartesian components of the) covariant derivative operator of the induced Levi-Civita connection acting on the spinor fields on  $\mathcal{S}$ , then e.g. for  $s = -|s|$  the  $\iota^{\underline{A}}$ -spinor components of the covariant directional derivatives of the spinor fields,  $\delta_s \phi := m^i \delta_i ((\sqrt{2}P)^{-|s|} \phi_{\underline{A}_1 \dots \underline{A}_{2|s|}}) \iota^{\underline{A}_1} \dots \iota^{\underline{A}_{2|s|}}$  and  $\delta'_s \phi := \bar{m}^i \delta_i ((\sqrt{2}P)^{-|s|} \phi_{\underline{A}_1 \dots \underline{A}_{2|s|}}) \iota^{\underline{A}_1} \dots \iota^{\underline{A}_{2|s|}}$ , give just the edth and edth-prime operators of Newman and Penrose [15] acting on the appropriate line bundles (see also [12, 13, 14, 16]).  $\delta_s$  and  $\delta'_s$  acting on cross sections of  $\mathcal{O}(-2s)$  for  $s = |s|$  are defined analogously.  $\delta_s$  increases, while  $\delta'_s$  decreases the spin weight by one. In the complex stereographic coordinates on  $U_n$ , the explicit form of these operators, acting on a function  $\phi$  of spin weight  $s$ , is

$$\delta_s \phi = \frac{1}{\sqrt{2}P} \left( (1 + \zeta \bar{\zeta}) \frac{\partial \phi}{\partial \zeta} + s \zeta \phi \right), \quad \delta'_s \phi = \frac{1}{\sqrt{2}P} \left( (1 + \zeta \bar{\zeta}) \frac{\partial \phi}{\partial \zeta} - s \bar{\zeta} \phi \right). \quad (4.9)$$

If no confusion arises, simply we write  $\delta$  and  $\delta'$  instead of  $\delta_s$  and  $\delta'_s$ . These operators link the spinors  $o^{\underline{A}}$  and  $\iota^{\underline{A}}$ :  $\delta o^{\underline{A}} = 0$ ,  $\delta' o^{\underline{A}} = \iota^{\underline{A}} / (\sqrt{2}P)$ ,  $\delta \iota^{\underline{A}} = -o^{\underline{A}} / (\sqrt{2}P)$  and  $\delta' \iota^{\underline{A}} = 0$ ; which imply  $\delta p^i = m^i$ ,  $\delta m^i = 0$  and  $\delta \bar{m}^i = -p^i / P^2$ .

The spin weighted spherical harmonics  ${}_s Y_{j,m}$  can also be defined (up to phase and normalization) by the pair of equations  $\delta {}_s Y_{j,m} = -\frac{1}{\sqrt{2}P} \sqrt{(j+s+1)(j-s)} {}_{s+1} Y_{j,m}$  and  $\delta' {}_s Y_{j,m} = \frac{1}{\sqrt{2}P} \sqrt{(j-s+1)(j+s)} {}_{s-1} Y_{j,m}$ . These imply that the harmonics are eigenfunctions of  $\delta \delta'$  and  $\delta' \delta$ , and also of the Laplacian:  $(\delta \delta' + \delta' \delta) {}_s Y_{j,m} = -P^{-2} (j(j+1) - s^2) {}_s Y_{j,m}$ . (For more details, see [15, 16, 12, 13].)

For the general, abstract definition of the bundles  $\mathcal{O}(-2s)$  over complex projective spaces, see e.g. [12, 13, 14]. For their introduction and a discussion of some of the global topological properties of the operators  $\delta$  and  $\delta'$  on closed metric 2-surfaces even with any genus, see [17].

#### 4.1.4 The angular momentum operators on $\mathcal{O}(-2s)$

In this subsection, we determine the specific form of the angular momentum operators acting on spin weighted functions. This geometrical form is more natural than the usual one when the algebra of quantum observables  $su(2)$  is considered to be a subalgebra of the Euclidean algebra  $e(3)$ , which will be considered in [11]. The form of this representation is similar to that of the generators of the Poincaré algebra of relativistic quantum systems [22], too.

The action of the group  $SU(2)$  on  $\mathbb{R}^3$  is defined by  $p^i \rightarrow \Lambda^i_j(A) p^j$ , where  $\Lambda^i_j(A) := -\sigma^i_{\underline{A}\underline{A}'} A^{\underline{A}}_{\underline{B}} \bar{A}^{\underline{A}'}_{\underline{B}'} \sigma_j^{\underline{B}\underline{B}'}$ , in which  $A^{\underline{A}}_{\underline{B}} \in SU(2)$  and over-bar denotes complex conjugation. (The  $(-)$  sign in the expression of  $\Lambda^i_j(A)$  is due to our convention that we lower and raise the small Latin indices by the *positive definite*  $\delta_{ij}$  and its inverse.) Thus  $SU(2)$  is the (universal covering group of the) group of those *isometries* of the *flat Riemannian 3-manifold*  $(\mathbb{R}^3, \delta_{ij})$  that leave the origin  $p^i = 0$  fixed. The surfaces of transitivity of  $SU(2)$  are the 2-spheres  $\mathcal{S}$  with radius  $P > 0$  and the origin  $p^i = 0$ . The latter case is uninteresting for us in the present paper, because that yields the trivial representation.

One way of determining the irreducible representations of  $SU(2)$  is by means of the method of induced representations. In this way, first we should find the representations of the stabilizer subgroup for a point  $p^i \in \mathcal{S}$  in  $SU(2)$ . This is  $U(1) \subset SU(2)$ , and, by Schur's lemma, all of its irreducible representations are one-dimensional, and these are labelled by  $s = 0, \pm\frac{1}{2}, \pm 1, \dots$ . If  $\overset{\circ}{\pi}^{\underline{A}}$  is the spinor constituent of  $p^i$  (see Appendix 4.1.2), then this one-dimensional representation space is chosen to be spanned by the spinor of the form  $\overset{\circ}{\pi}^{\underline{A}_1} \dots \overset{\circ}{\pi}^{\underline{A}_{2|s|}}$  if  $s = -|s| \leq 0$ , and  $\overset{\circ}{\pi}^{\underline{A}'_1} \dots \overset{\circ}{\pi}^{\underline{A}'_{2s}}$  if  $s = |s| > 0$ . The next step is the generation of the representation space for the whole group  $SU(2)$  from this one dimensional space by the elements of  $SU(2)$  that do not leave  $p^i$  fixed.

Geometrically, the above method (by using the group action from  $p^i$ ) is the construction of the bundle of totally symmetric unprimed  $N$ -type spinors  $\phi^{\underline{A}_1 \dots \underline{A}_{2|s|}}$  on  $\mathcal{S}$  if  $s = -|s|$ , and of the totally symmetric primed  $N$ -type spinors  $\chi^{\underline{A}'_1 \dots \underline{A}'_{2s}}$  on  $\mathcal{S}$  if  $s = |s|$ . (Clearly, these are equivalent to the line bundles  $\mathcal{O}(-2s)$  with the corresponding  $s$ .) The  $2|s|$ -fold principal spinor of them at the point  $p^i = \sigma^i_{\underline{A}\underline{A}'} \pi^{\underline{A}} \bar{\pi}^{\underline{A}'}$  is  $\pi^{\underline{A}}$  and  $\bar{\pi}^{\underline{A}'}$ , respectively.

To determine the explicit form of the representation of  $SU(2)$  by operators  $\mathbf{U}(A)$  acting on the spinor fields, let us recall that the rotation Killing vectors  $k_{mn}^i$  are tangent to  $\mathcal{S}$  and generate its isometries, too. Then the action of  $SU(2)$  e.g. on any  $\phi^{\underline{A}_1 \dots \underline{A}_{2|s|}}$  is defined by  $(\mathbf{U}(A)\phi)^{\underline{A}_1 \dots \underline{A}_{2|s|}}(p^i) := A^{\underline{A}_1}_{\underline{B}_1} \dots A^{\underline{A}_{2|s|}}_{\underline{B}_{2|s|}} \phi^{\underline{B}_1 \dots \underline{B}_{2|s|}}(\Lambda(A^{-1})^i_j p^j)$ . In particular,  $(\mathbf{U}(A)\pi)^{\underline{A}} = \pi^{\underline{A}}$ , i.e. the spinor constituent of the position vector field is  $SU(2)$ -invariant. Hence, any  $2|s|$ -rank spinor field that belongs to the carrier space of an irreducible representation of  $SU(2)$  is necessarily  $N$ -type with  $\pi^{\underline{A}}$  and  $\bar{\pi}^{\underline{A}'}$  as its  $2|s|$ -fold principal spinor for  $s = -|s|$  and  $s = |s|$ , respectively. Note that the function  $\phi$  appearing in  $\phi^{\underline{A}_1 \dots \underline{A}_{2|s|}} = \phi \pi^{\underline{A}_1} \dots \pi^{\underline{A}_{2|s|}}$  has spin weight  $s$ . Using (4.8), it is straightforward to check that the operator  $\mathbf{U}(A)$  is unitary with respect to the scalar product (4.7). As we will see in [11], the  $L_2$  space  $\mathcal{H}$  of the  $N$ -type spinor fields with given  $s$  provides the carrier space of an *irreducible* representation for the  $E(3)$  group, but it is *not* irreducible for the  $SU(2)$ .

In this representation, still for  $s = -|s| \leq 0$ , the operators  $\mathbf{J}_{ij} := \varepsilon_{ijk} \mathbf{J}^k$  are defined to be the densely defined self-adjoint generators of these transformations: Let  $A^{\underline{A}}_{\underline{B}}(u)$  be a 1-parameter subgroup in  $SU(2)$  generated by  $\lambda^{\underline{A}}_{\underline{B}} \in su(2)$ . Then its trajectories on  $\mathcal{S}$  are necessarily integral curves of some rotation Killing vector field, say of  $M^{mn} k_{mn}^i$  for some constants  $M^{mn} = -M^{nm}$ . Then  $\mathbf{J}_{ij}$  is defined by  $(i/\hbar) M^{ij} \mathbf{J}_{ij} \phi^{\underline{A}_1 \dots} := \frac{d}{du} ((\mathbf{U}(A(u))\phi)^{\underline{A}_1 \dots})|_{u=0}$ . This is just minus the Lie derivative of the spinor field along the Killing vector  $M^{mn} k_{mn}^i$  (see e.g. [12, 13]). Here the limit in the definition of the derivative is meant in the strong topology of  $\mathcal{H}$ . To evaluate this, let us calculate the tangent of the trajectories  $\Lambda^i_j(A(u)) p^j$  at  $p^i$ . This is  $k^i = -\sigma^i_{\underline{A}\underline{A}'} (\lambda^{\underline{A}}_{\underline{B}} \delta^{\underline{A}'}_{\underline{B}'} + \delta^{\underline{A}}_{\underline{B}} \bar{\lambda}^{\underline{A}'}_{\underline{B}'}) \sigma_j^{\underline{B}\underline{B}'} p^j$ , which must coincide with  $M^{mn} k_{mn}^i = 2M^i_j p^j$ . Since  $\lambda^{\underline{A}}_{\underline{B}} \in su(2)$ , its complex conjugate is not independent of  $\lambda^{\underline{A}}_{\underline{B}} \in su(2)$ , because  $\lambda_{\underline{A}\underline{B}} = 2\bar{\lambda}^{\underline{A}'}_{\underline{B}'} \sigma^0_{\underline{A}'\underline{A}} \sigma^0_{\underline{B}'\underline{B}}$  holds. Thus, introducing the standard  $SU(2)$  Pauli matrices  $\sigma^{\underline{A}}_{\underline{B}}$  (including the factor  $1/\sqrt{2}$ ) as the unitary spinor form of the three non-trivial  $SL(2, \mathbb{C})$  Pauli matrices (i.e. which are given explicitly by  $\sigma^{\underline{A}}_{\underline{B}} = \delta_{ij} \varepsilon^{\underline{A}\underline{C}} \sigma^j_{\underline{C}\underline{B}'} \sqrt{2} \sigma^0_{\underline{B}'\underline{B}}$ ), we can express  $\lambda^{\underline{A}}_{\underline{B}}$  in terms of  $M^{ij}$ :

$$\lambda^{\underline{A}}_{\underline{B}} = -\frac{i}{\sqrt{2}} M^{ij} \varepsilon_{ij}{}^k \sigma_k^{\underline{A}}_{\underline{B}}.$$

In deriving this formula we also used the identity

$$\sigma^{\underline{A}}_{\underline{B}} \sigma^{\underline{B}}_{\underline{C}} = -\frac{i}{\sqrt{2}} \varepsilon_{ij}{}^k \sigma_k^{\underline{A}}_{\underline{C}} + \frac{1}{2} \delta_{ij} \delta_{\underline{C}}^{\underline{A}}.$$

Now we are able to evaluate the equation defining the operator  $\mathbf{J}_{ij}$ . We find that, still for  $s = -|s|$ ,

$$\mathbf{J}_{ij}\phi^{A_1\dots} = i\hbar\left(p_j\frac{\partial}{\partial p^i} - p_i\frac{\partial}{\partial p^j}\right)\phi^{A_1\dots} + \sqrt{2}\hbar s \varepsilon_{ij}{}^k \sigma_k^{(A_1}{}_{(B_1} \delta_{B_2}^{A_2} \dots \delta_{B_{2|s|}}^{A_{2|s|})} \phi^{B_1\dots}. \quad (4.10)$$

Thus,  $\mathbf{J}_{ij}$  is  $i\hbar$ -times the Lie derivative operator along the Killing vector  $k_{ij}^k$ , and hence is well defined on the *dense subspace* of the smooth spinor fields in  $\mathcal{H}$ . Then it is a straightforward calculation to check that these operators do, indeed, satisfy the defining commutation relations of  $su(2)$  on the appropriate dense subspaces, i.e. provide a representation of the Lie algebra of the rotation Killing fields. Repeating the analogous analysis for  $s = |s|$ , we obtain the same expression (4.10) for  $\mathbf{J}_{ij}$ .

Next, using the original form of the angular momentum operator,  $\mathbf{J}_i = \frac{1}{2}\varepsilon_i{}^{jk}\mathbf{J}_{jk}$ , we determine its contraction with the basis vectors  $p^i$ ,  $m^i$  and  $\bar{m}^i$ . First, recalling from Appendix 4.1.2 that  $p^i\sigma_i^{AA'} = P(\iota^A\bar{\iota}^{A'} - o^A\bar{o}^{A'})/\sqrt{2}$ , for its unitary spinor form we obtain

$$p^i\sigma_i^A{}_{\underline{B}} := p^i\sigma_i^A{}_{\underline{A}'}\sqrt{2}\sigma_0^A{}_{\underline{B}} = -\frac{P}{\sqrt{2}}(\iota^A\bar{\iota}^{A'} - o^A\bar{o}^{A'}) (\bar{o}_{A'}o_{\underline{A}} + \bar{\iota}_{A'}\iota_{\underline{A}}) = -\frac{P}{\sqrt{2}}(\iota^A o_{\underline{B}} + o^A \iota_{\underline{B}}).$$

Since on the domain  $U_n$  the spinor field has the form  $\phi^{A_1\dots A_{2|s|}} = \phi\pi^{A_1}\dots\pi^{A_{2|s|}}$ , we find that

$$p^i\mathbf{J}_i\phi^{A_1\dots A_{2|s|}} = sP\hbar\phi^{A_1\dots A_{2|s|}}. \quad (4.11)$$

It might be worth noting that in the *unitary, irreducible* representation of the Euclidean group  $E(3)$  exactly the same expression emerges as the analog of the Pauli–Lubanski spin operator [11], which is one of the two Casimir operators of  $e(3)$ .

Similarly, one can show that  $m^i\sigma_i^A{}_{\underline{B}} = o^A o_{\underline{B}} = \frac{1}{\sqrt{2}P}\pi^A\pi_{\underline{B}}$  and  $\bar{m}^i\sigma_i^A{}_{\underline{B}} = -\iota^A\iota_{\underline{B}}$ . Then, using  $m^i p^j \varepsilon_{ijk} = -p^i m^j \bar{m}^l \varepsilon_{ijl} m_k = -iPm_k$  and  $\bar{m}^i p^j \varepsilon_{ijk} = iP\bar{m}_k$  (see Appendix 4.1.2), as well as the definition of the  $\delta$  and  $\delta'$  operators (Appendix 4.1.3), we find

$$m^i\mathbf{J}_i\phi^{A_1\dots A_{2|s|}} = -P\hbar(\delta\phi)\pi^{A_1}\dots\pi^{A_{2|s|}}. \quad (4.12)$$

$$\bar{m}^i\mathbf{J}_i\phi^{A_1\dots A_{2|s|}} = P\hbar(\delta'\phi)\pi^{A_1}\dots\pi^{A_{2|s|}}. \quad (4.13)$$

Hence, by (4.11)-(4.13),

$$\begin{aligned} \mathbf{J}_i\phi^{A_1\dots A_{2|s|}} &= (m_i\bar{m}^j + \bar{m}_i m^j + \frac{1}{P^2}p_i p^j)\mathbf{J}_j\phi^{A_1\dots A_{2|s|}} = \\ &= P\hbar\left(m_i\delta'\phi - \bar{m}_i\delta\phi + s\frac{p_i}{P^2}\phi\right)\pi^{A_1}\dots\pi^{A_{2|s|}}. \end{aligned}$$

Therefore, defining the action of  $\mathbf{J}_i$  on the spin weighted function  $\phi$  with spin weight  $s$  simply by  $\mathbf{J}_i\phi := (\sqrt{2}P)^{-|s|}(\mathbf{J}_i\phi_{\underline{A}_1\dots\underline{A}_{2|s|}})\iota^{A_1}\dots\iota^{A_{2|s|}}$ , we obtain

$$\mathbf{J}_i\phi = P\hbar\left(m_i\delta'\phi - \bar{m}_i\delta\phi + s\frac{n_i}{P}\phi\right). \quad (4.14)$$

In  $E(3)$  invariant elementary quantum mechanical systems the same expression is interpreted as the decomposition of the total angular momentum into its orbital and spin parts, where  $P$  and  $s$  are fixed by the two Casimir operators of  $e(3)$  [11].

Then by (4.14) it is easy to compute the only Casimir operator  $\mathbf{J}_i\mathbf{J}^i$  of the  $su(2)$  algebra. It is

$$\mathbf{J}_i\mathbf{J}^i\phi = \hbar^2\left(-P^2(\delta\delta' + \delta'\delta)\phi + s^2\phi\right); \quad (4.15)$$

where, as we noted in Appendix 4.1.3,  $\delta\delta' + \delta'\delta$  is just the Laplace operator on  $\mathcal{S}$ . Hence the spectrum of  $\mathbf{J}_i\mathbf{J}^i$  is  $j(j+1)\hbar^2$ ,  $j = |s|, |s|+1, \dots$ , as we expected, and, for fixed  $s$  and  $j$ , the  $SU(2)$ -irreducible subspace in  $\mathcal{H}$  is spanned by the spin weighted spherical harmonics  ${}_sY_{j,m}$ ,  $m = -j, -j+1, \dots, j$ .

## 4.2 The explicit form of the eigenfunctions $W_{s,j,m}$ and the asymptotics of the standard deviations

### 4.2.1 The explicit form of the eigenfunctions $\phi_{s,j,m}$

The strategy of the determination of the explicit coordinate form of  $W_{s,j,m}$  and  $W_{s,j}$  (and in particular the factors of normalization,  $N_{s,j,m}$  and  $N_{s,j}$ ) is just that for the spin weighted spherical harmonics  ${}_sY_{j,m}$  given in [12]: The eigenfunction

$$\phi_{s,j,m} = \mu_{\underline{A}_1} \cdots \mu_{\underline{A}_{j-m}} \nu_{\underline{A}_{j-m+1}} \cdots \nu_{\underline{A}_{2j}} o^{\underline{A}_1} \cdots o^{\underline{A}_{j+s}} t^{\underline{A}_{j+s+1}} \cdots t^{\underline{A}_{2j}} \quad (4.16)$$

is the combination of terms of the form

$$\frac{1}{(2j)!} (\mu_{\underline{A}} o^{\underline{A}})^r (\mu_{\underline{B}} t^{\underline{B}})^k (\nu_{\underline{C}} o^{\underline{C}})^l (\nu_{\underline{D}} t^{\underline{D}})^{2j-r-k-l}$$

with non-negative integer coefficients. But in such a term the number of the  $o^{\underline{A}}$  spinors must be  $j+s$ , and the number of the  $t^{\underline{A}}$  spinors must be  $j-s$ , and hence  $l = j+s-r$ . In a similar way, the number of the  $\mu_{\underline{A}}$  spinors is  $j-m$  and that of  $\nu_{\underline{A}}$  is  $j+m$ , and hence  $k = j-m-r$ . Therefore, for given  $r$ , the structure of the terms in (4.16) is

$$\frac{1}{(2j)!} (\mu_{\underline{A}} o^{\underline{A}})^r (\mu_{\underline{B}} t^{\underline{B}})^{j-m-r} (\nu_{\underline{C}} o^{\underline{C}})^{j+s-r} (\nu_{\underline{D}} t^{\underline{D}})^{r+m-s}. \quad (4.17)$$

$r$  takes its minimum value when all the  $\nu_{\underline{A}}$  spinors are contracted with  $o^{\underline{A}}$  (but no  $t^{\underline{A}}$ ) spinors; and in this case, if  $s-m \geq 0$ , the number of the  $o^{\underline{A}}$  spinors that remain to contract with the  $\mu_{\underline{A}}$  spinors is  $s-m$ . Hence,  $r \geq \max\{0, s-m\}$ . The maximum value of  $r$  is the maximal number of contractions  $\mu_{\underline{A}} o^{\underline{A}}$ , which is  $j+s$  if  $j-m \geq j+s$ , and it is  $j-m$  if  $j+s \geq j-m$ . Therefore,  $r \leq \min\{j-m, j+s\}$ . Finally, to determine the number of the terms of the form (4.17) in (4.16), let us rewrite (4.17) in the form

$$\begin{aligned} & \frac{1}{(2j)!} \left( \mu_{\underline{A}_1} \cdots \mu_{\underline{A}_r} \nu_{\underline{A}_{r+1}} \cdots \nu_{\underline{A}_{j+s}} \right) o^{\underline{A}_1} \cdots o^{\underline{A}_{j+s}} \\ & \times \left( \mu_{\underline{B}_1} \cdots \mu_{\underline{B}_{j-m-r}} \nu_{\underline{B}_{j-m-r+1}} \cdots \nu_{\underline{B}_{j-s}} \right) t^{\underline{B}_1} \cdots t^{\underline{B}_{j-s}}. \end{aligned}$$

The number of ways in which the  $\mu_{\underline{A}}$  and the  $\nu_{\underline{A}}$  spinors can be chosen in this manner is  $\binom{j-m}{r}$  and  $\binom{j+m}{r+m-s}$ , respectively; and each of these choices can be contracted with the  $o^{\underline{A}}$  spinors in  $(j+s)!$  ways and with the  $t^{\underline{B}}$  spinors in  $(j-s)!$  ways. Thus, their total number is  $\binom{j-m}{r} \binom{j+m}{r+m-s} (j+s)!(j-s)!$ . Hence

$$\begin{aligned} \phi_{s,j,m} &= \frac{(j+m)!(j-m)!(j+s)!(j-s)!}{(2j)!} \\ & \times \sum_r \frac{(\mu_{\underline{A}} o^{\underline{A}})^r (\mu_{\underline{B}} t^{\underline{B}})^{j-m-r} (\nu_{\underline{C}} o^{\underline{C}})^{j+s-r} (\nu_{\underline{D}} t^{\underline{D}})^{r+m-s}}{r!(j-m-r)!(j+s-r)!(r+m-s)!}, \end{aligned} \quad (4.18)$$

where  $\max\{0, s - m\} \leq r \leq \min\{j - m, j + s\}$ . In the exceptional case  $r = j + s$ , and hence (4.18) reduces to

$$\phi_{s,j} = (\mu_{\underline{A}} o^{\underline{A}})^{j+s} (\mu_{\underline{B}} \iota^{\underline{B}})^{j-s}; \quad (4.19)$$

which can be derived directly even from (4.16), too.

Next, we determine the explicit form of the contractions  $\mu_{\underline{A}} o^{\underline{A}}$ ,  $\nu_{\underline{A}} o^{\underline{A}}$ ,  $\mu_{\underline{A}} \iota^{\underline{A}}$  and  $\nu_{\underline{A}} \iota^{\underline{A}}$ . (These are not only the factors in (4.18), but also these are the eigenfunctions  $\phi_{s,\frac{1}{2},m}$  with  $(s, m) = (\frac{1}{2}, -\frac{1}{2}), (\frac{1}{2}, \frac{1}{2}), (-\frac{1}{2}, -\frac{1}{2})$  and  $(-\frac{1}{2}, \frac{1}{2})$ , respectively.) First, suppose that  $\gamma_{\underline{A}\underline{B}}$  is not null. Then

$$\gamma_{\underline{A}\underline{B}} = \frac{1}{\sqrt{2}} \begin{pmatrix} -\alpha_1 + i\alpha_2 & \alpha_3 - i\lambda \\ \alpha_3 - i\lambda & \alpha_1 + i\alpha_2 \end{pmatrix},$$

and hence the solution of  $\sqrt{2}\gamma_{\underline{A}\underline{B}} = \mu_{\underline{A}}\nu_{\underline{B}} + \mu_{\underline{B}}\nu_{\underline{A}}$  is given by

$$\mu_{\underline{A}} = \mu_1 O_{\underline{A}} - \mu_0 I_{\underline{A}} = -(\xi_{\mp} \exp[i\alpha] O_{\underline{A}} + I_{\underline{A}}) \mu_0, \quad (4.20)$$

$$\nu_{\underline{A}} = \nu_1 O_{\underline{A}} - \nu_0 I_{\underline{A}} = \frac{1}{2} \sqrt{1 - \alpha_3^2} (\xi_{\pm} O_{\underline{A}} + \exp[-i\alpha] I_{\underline{A}}) \mu_0^{-1}; \quad (4.21)$$

where  $\mu_0 := \mu_{\underline{A}} O^{\underline{A}}$  is an arbitrary nonzero constant and  $\xi_{\pm}$  has been defined by (2.17). However, this solution gives  $\nu_{\underline{A}} \mu^{\underline{A}} = \mp \sqrt{1 - \lambda^2 - 2i\lambda\alpha_3}$ , and hence, to be compatible with our sign convention in (2.10), we must choose  $\xi_+$  in  $\mu_1$  (and hence  $\xi_-$  in  $\nu_1$ ). Therefore,

$$\mu_{\underline{A}} o^{\underline{A}} = -i \frac{\xi_- - \xi_+}{\sqrt{1 + \xi\xi}} \mu_0 \exp[i\alpha], \quad \nu_{\underline{A}} o^{\underline{A}} = \frac{i}{2} \sqrt{1 - \alpha_3^2} \frac{\xi_- - \xi_+}{\sqrt{1 + \xi\xi}} \mu_0^{-1}, \quad (4.22)$$

$$\mu_{\underline{A}} \iota^{\underline{A}} = -i \xi_+ \frac{\bar{\xi} - \xi_-}{\sqrt{1 + \xi\xi}} \mu_0, \quad \nu_{\underline{A}} \iota^{\underline{A}} = \frac{i}{2} \xi_- \sqrt{1 - \alpha_3^2} \frac{\bar{\xi} - \xi_+}{\sqrt{1 + \xi\xi}} (\mu_0 \exp[i\alpha])^{-1} \quad (4.23)$$

In the exceptional case, i.e. when  $\lambda = 1$  and  $\alpha_3 = 0$ , the components of the principal spinor  $\mu_{\underline{A}}$  are  $\mu_0 := \mu_{\underline{A}} O^{\underline{A}} = \pm \frac{i}{\sqrt{2}} \exp[-\frac{i}{2}\alpha]$  and  $\mu_1 := \mu_{\underline{A}} I^{\underline{A}} = \mp \frac{1}{\sqrt{2}} \exp[\frac{i}{2}\alpha]$ . Hence, the corresponding solutions  $\phi_{\frac{1}{2},\frac{1}{2}}$  and  $\phi_{-\frac{1}{2},\frac{1}{2}}$ , respectively, are

$$\mu_{\underline{A}} o^{\underline{A}} = \pm \frac{1}{\sqrt{2}} \frac{\xi + i}{\sqrt{1 + \xi\xi}} \exp[\frac{i}{2}\alpha], \quad \mu_{\underline{A}} \iota^{\underline{A}} = \mp \frac{i}{\sqrt{2}} \frac{\bar{\xi} + i}{\sqrt{1 + \xi\xi}} \exp[-\frac{i}{2}\alpha]. \quad (4.24)$$

Therefore, the eigenfunctions  $\phi_{s,j,m}$  and  $\phi_{s,j}$  are polynomials in  $\xi$  and  $\bar{\xi}$  (with the overall order  $2j$ ) and divided by  $(1 + \xi\bar{\xi})^j$ . Thus, in particular, all these are bounded on  $\mathcal{S}$  and smooth on  $\mathcal{S}$  minus the ‘north pole’ of the coordinate system  $(\xi, \bar{\xi})$ .

#### 4.2.2 Orthogonality properties and the factors of normalization

To find the factor of normalization, and also to clarify whether or not the functions  $\phi_{s,j,m}$  and  $\phi_{s,j}$  form orthogonal systems, let us recall from [12] (Lemma 4.15.86) that

$$\int o_{\underline{A}_1} \cdots o_{\underline{A}_k} \iota^{\underline{B}_1} \cdots \iota^{\underline{B}_k} d\mathcal{S} = \frac{4\pi}{k+1} \delta_{\underline{A}_1}^{(\underline{B}_1} \cdots \delta_{\underline{A}_k}^{\underline{B}_k)}; \quad (4.25)$$

but the integral is zero if the number of the  $o_{\underline{A}}$  and of the  $\iota^{\underline{B}}$  spinors under the integral is different. Here, and in the rest of this Appendix,  $\mathcal{S}$  is assumed to be a *unit* sphere.

Let  $\phi_{s,j,m}$  and  $\phi_{s',j',m'}$  be two eigenfunctions, where the latter is built from  $\phi'_{\underline{A}_1 \dots \underline{A}_{2j'}}$  given by (2.9) but in which  $m$  is replaced by  $m'$ . Recalling that  $2\sigma_0^{\underline{A}'\underline{B}'}\sigma_{\underline{B}'\underline{B}'}^0 = \delta_{\underline{B}'}^{\underline{A}'}$  and that the Hermitean adjoint e.g. of a spinor  $\phi_{\underline{A}}$  is defined by  $\phi_{\underline{A}}^\dagger := \bar{\phi}_{\underline{A}'}\sqrt{2}\sigma_0^{\underline{A}'\underline{A}}$ , we have that

$$\begin{aligned} & \langle \phi_{s',j',m'}, \phi_{s,j,m} \rangle = \\ & = \int \bar{\phi}'_{\underline{D}'_1 \dots \underline{D}'_{2j'}} 2\sigma_0^{\underline{D}'_1 \underline{D}_1} \sigma_{\underline{D}_1 \underline{A}'_1}^0 \dots 2\sigma_0^{\underline{D}'_{2j'} \underline{D}_{2j'}} \sigma_{\underline{D}_{2j'} \underline{A}'_{2j'}}^0 \bar{o}^{\underline{A}'_1} \dots \bar{o}^{\underline{A}'_{j'+s'}} \bar{t}^{\underline{A}'_{j'+s'+1}} \dots \bar{t}^{\underline{A}'_{2j'}} \\ & \quad \times \phi_{\underline{A}_1 \dots \underline{A}_{2j}} o^{\underline{A}_1} \dots o^{\underline{A}_{j+s}} t^{\underline{A}_{j+s+1}} \dots t^{\underline{A}_{2j}} d\mathcal{S} = \\ & = (-)^{j+s} \phi_{\underline{D}_1 \dots \underline{D}_{j'+s'}}^{\dagger} \bar{\phi}'_{\underline{D}'_{j'+s'+1} \dots \underline{D}'_{2j'}} o^{\underline{A}_1 \dots \underline{A}_{j+s}} \phi_{\underline{A}_{j+s+1} \dots \underline{A}_{2j}} \\ & \quad \times \int o_{\underline{A}_1} \dots o_{\underline{A}_{j+s}} o_{\underline{D}'_{j'+s'+1}} \dots o_{\underline{D}'_{2j'}} t^{\underline{D}_1} \dots t^{\underline{D}_{j'+s'}} t^{\underline{A}_{j+s+1}} \dots t^{\underline{A}_{2j}} d\mathcal{S}. \end{aligned}$$

However, as we noted, this integral can be non-zero only if the number of the  $o_{\underline{A}}$  and of the  $t^{\underline{D}}$  spinors is the same, i.e. if  $s' = s$ . Hence, by (4.25), this integral is

$$(-)^{j+s} \delta_{ss'} \phi_{\underline{D}_1 \dots \underline{D}_{j'+s}}^{\dagger} o^{\underline{A}_{j+s+1} \dots \underline{A}_{j+j'}} \phi_{\underline{A}_1 \dots \underline{A}_{j+s}} \bar{\phi}'_{\underline{D}'_{j'+s'+1} \dots \underline{D}'_{j+j'}} \frac{4\pi}{j+j'+1} \delta_{\underline{A}_1}^{(\underline{D}_1 \dots \underline{D}_{j+j'})}.$$

But since both  $\phi_{\underline{A}_1 \dots \underline{A}_{2j}}$  and  $\phi'_{\underline{A}_1 \dots \underline{A}_{2j'}}$  are totally symmetric, a term in this expression is non-zero precisely when every index of one spinor is contracted with some index of the other spinor. Hence,  $j = j'$ . Therefore,

$$\langle \phi_{s',j',m'}, \phi_{s,j,m} \rangle = (-)^{2j} \delta_{ss'} \delta_{jj'} 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} \phi_{\underline{A}_1 \dots \underline{A}_{2j}}^{\dagger} \phi_{\underline{A}_1 \dots \underline{A}_{2j}}. \quad (4.26)$$

Finally, rewriting  $\phi_{\underline{A}_1 \dots \underline{A}_{2j}}^{\dagger}$  by the complex conjugate spinor and recalling that  $\sqrt{2}\sigma_0^{\underline{A}\underline{A}'} = O^{\underline{A}}\bar{O}^{\underline{A}'} + I^{\underline{A}}\bar{I}^{\underline{A}'}$ , we obtain

$$\begin{aligned} \langle \phi_{s',j',m'}, \phi_{s,j,m} \rangle & = \delta_{ss'} \delta_{jj'} 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} \\ & \quad \times \bar{\phi}'_{\underline{A}'_1 \dots \underline{A}'_{2j}} (\bar{O}^{\underline{A}'_1} O^{\underline{A}_1} + \bar{I}^{\underline{A}'_1} I^{\underline{A}_1}) \dots (\bar{O}^{\underline{A}'_{2j}} O^{\underline{A}_{2j}} + \bar{I}^{\underline{A}'_{2j}} I^{\underline{A}_{2j}}) \phi_{\underline{A}_1 \dots \underline{A}_{2j}} = \\ & = \delta_{ss'} \delta_{jj'} 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} \sum_{k=0}^{2j} \binom{2j}{k} \Phi_{k,j,m} \bar{\Phi}_{k,j,m'}, \end{aligned} \quad (4.27)$$

where  $\Phi_{k,j,m}$  was defined by (2.29). If in  $\phi_{\underline{A}_1 \dots \underline{A}_{2j}}$  the spinors  $\mu_{\underline{A}}$  and  $\nu_{\underline{A}}$  were  $O_{\underline{A}}$  and  $I_{\underline{A}}$ , respectively, then  $\langle \phi_{s',j',m'}, \phi_{s,j,m} \rangle$  would be proportional to  $\delta_{mm'}$ , too, and the whole expression would reduce to the expression of the norm of the not normalized spin weighted spherical harmonics  ${}_s Z(j, m)$  in [12]. However, with our  $\mu_{\underline{A}}$  and  $\nu_{\underline{A}}$ , the eigenfunctions  $\phi_{s,j,m}$  and  $\phi_{s,j,m'}$ ,  $m \neq m'$ , are *not* orthogonal to each other. Nevertheless, (4.27) provides the factor of normalization  $N_{s,j,m}$ :

$$N_{s,j,m}^{-2} := \langle \phi_{s,j,m}, \phi_{s,j,m} \rangle = 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} \sum_{k=0}^{2j} \binom{2j}{k} |\Phi_{k,j,m}|^2. \quad (4.28)$$

To see its dependence on the parameters  $\lambda$  and  $\alpha_3$ , we should find the explicit form of  $\Phi_{k,j,m}$ . We determine this in the next subsection.

The previous analysis is valid even in the exceptional case. The only difference between the generic and exceptional cases is that now  $m = m' = -j$ , and hence, according to (4.27), the eigenfunctions  $\phi_{s,j}$  form an orthogonal system. Since, in the exceptional case, the modulus of the spinor components,  $\mu_0 := \mu_{\underline{A}} O^{\underline{A}}$  and  $\mu_1 := \mu_{\underline{A}} I^{\underline{A}}$ , is  $1/\sqrt{2}$ , the factor of normalization  $N_{s,j} = N_{s,j,-j}$ , given by (4.28), can be written as

$$N_{s,j}^{-2} = 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!} \sum_{k=0}^{2j} \binom{2j}{k} |\mu_0|^{2k} |\mu_1|^{2(2j-k)} = 4\pi \frac{(j-s)!(j+s)!}{(2j+1)!}, \quad (4.29)$$

which is a pure number, independently of the (only) free parameter  $\alpha$ .

### 4.2.3 The continuity of the standard deviations

(4.18) gives the structure of  $\Phi_{k,j,m}$ , too, if  $\sigma^{\underline{A}}$  is replaced by  $O^{\underline{A}}$ ,  $\iota^{\underline{A}}$  by  $I^{\underline{A}}$ , and  $s$  by  $k-j$ . Thus, using (4.20)-(4.21) and taking into account  $\xi_+\xi_- = -1$  (see (2.17)), we find that

$$\begin{aligned} \Phi_{k,j,m} &= (-)^k \mu_0^{-2m} \left(\frac{1}{2}\sqrt{1-\alpha_3^2}\right)^{j+m} (\exp[i\alpha])^{j-m-k} (\xi_-)^{2m-k} \\ &\quad \times \frac{k!(2j-k)!}{(2j)!} \sum_r (-)^r \binom{j-m}{r} \binom{j+m}{k-r} (\xi_-)^{2r}, \end{aligned} \quad (4.30)$$

where  $\max\{0, k-j-m\} \leq r \leq \min\{j-m, k\}$ . However, by equations (4.28), (2.28) and (2.30), in the expression of  $\langle \mathbf{J}(\beta) \rangle_\phi$  and  $\langle (\mathbf{J}(\beta))^2 \rangle_\phi$  it is only the absolute value of

$$S_{k,j,m} := (\xi_-)^{-k} \sum_r (-)^r \binom{j-m}{r} \binom{j+m}{k-r} (\xi_-)^{2r} \quad (4.31)$$

that appears. In fact, in terms of this,  $\langle W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$  and  $\langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$  take the form

$$\langle W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle = -\hbar \frac{\sum_{k=0}^{2j} (j-k)k!(2j-k)! |S_{k,j,m}|^2}{\sum_{k=0}^{2j} k!(2j-k)! |S_{k,j,m}|^2}, \quad (4.32)$$

$$\langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle = \hbar^2 \frac{\sum_{k=0}^{2j} (j-k)^2 k!(2j-k)! |S_{k,j,m}|^2}{\sum_{k=0}^{2j} k!(2j-k)! |S_{k,j,m}|^2}. \quad (4.33)$$

These depend on  $\lambda$  and  $\alpha_3$  only through  $\xi_-$  according to (2.17) via  $S_{k,j,m}$ , but they do not depend on  $\alpha$  or  $\mu_0$  (as they should not).

To see the structure of  $S_{k,j,m}$  and its dependence on  $\lambda$  and  $\alpha_3$ , let us recall that the range of the summation in (4.31) is  $\max\{0, k-j-m\} \leq r \leq \min\{j-m, k\}$ . Hence, we should split the range of  $k$  in (4.32)-(4.33) into three disjoint domains: i. when  $0 \leq k < j - |m|$ , in which case  $r = 0, \dots, k$ ; ii.a. when  $j - |m| \leq k \leq j + |m|$  and  $m \geq 0$ , then  $r = 0, \dots, j - m$ , while ii.b. if  $j - |m| \leq k \leq j + |m|$  and  $m < 0$ , then  $r = k - j - m, \dots, k$ ; iii. when  $j + |m| < k \leq 2j$ , in which case  $r = k - j - m, \dots, j - m$ .



Hence, redefining  $r$  and its range if needed, in the respective cases

$$\text{i. } S_{k,j,m} = (\xi_-)^{-k} \sum_{r=0}^k (-)^r \binom{j-m}{r} \binom{j+m}{k-r} (\xi_-)^{2r}, \quad (4.34)$$

$$\text{ii.a. } S_{k,j,m} = (\xi_-)^{-k} \sum_{r=0}^{j-m} (-)^r \binom{j-m}{r} \binom{j+m}{k-r} (\xi_-)^{2r}, \quad (4.35)$$

$$\text{ii.b. } S_{k,j,m} = (-)^{k-j-m} (\xi_-)^{k-2(j+m)} \sum_{r=0}^{j+m} (-)^r \binom{j-m}{2j-k-r} \binom{j+m}{r} (\xi_-)^{2r}, \quad (4.36)$$

$$\text{iii. } S_{k,j,m} = (-)^{k-j-m} (\xi_-)^{k-2(j+m)} \sum_{r=0}^{2j-k} (-)^r \binom{j-m}{2j-k-r} \binom{j+m}{r} (\xi_-)^{2r}. \quad (4.37)$$

Next we show that these admit a rather non-trivial, hidden symmetry: For given  $k, j, m$  let us define  $\tilde{k} := 2j - k$ , and clearly  $k = 0, \dots, j - |m| - 1$  precisely when  $\tilde{k} = j + |m| + 1, \dots, 2j$ ; and  $k = j - |m|, \dots, j + |m|$  precisely when  $\tilde{k} = j - |m|, \dots, j + |m|$ . Then e.g. for  $k = 0, \dots, j - |m| - 1$  and  $m \geq 0$  by (4.34) and (4.37) with the notation  $\bar{r} := r - k + j - m$  we have that

$$\begin{aligned} S_{k,j,-m} &= (\xi_-)^{-k} \sum_{r=0}^k (-)^r \binom{j+m}{r} \binom{j-m}{k-r} (\xi_-)^{2r} \\ &= (\xi_-)^{\tilde{k}-2j} \sum_{\bar{r}=\tilde{k}-j-m}^{j-m} (-)^{\bar{r}+j+m-\tilde{k}} \binom{j+m}{\bar{r}-\tilde{k}+j+m} \binom{j-m}{j-m-\bar{r}} (\xi_-)^{2(\bar{r}-\tilde{k}+j+m)} \\ &= (-)^{j+m-\tilde{k}} (\xi_-)^{2m-\tilde{k}} \sum_{\bar{r}=\tilde{k}-j-m}^{j-m} (-)^{\bar{r}} \binom{j+m}{\tilde{k}-\bar{r}} \binom{j-m}{\bar{r}} (\xi_-)^{2\bar{r}} \\ &= (-)^{j+m-\tilde{k}} (\xi_-)^{2m} S_{\tilde{k},j,m}. \end{aligned} \quad (4.38)$$

Analogous calculations in the other cases show that  $S_{k,j,-m} = (-)^{j+m-\tilde{k}} (\xi_-)^{2m} S_{\tilde{k},j,m}$  holds in general.

This symmetry of  $S_{k,j,m}$  yields that the common denominator in (4.32) and (4.33) is

$$\sum_{k=0}^{2j} k!(2j-k)! |S_{k,j,-m}|^2 = |\xi_-|^{4m} \left( \sum_{k=0}^{2j} k!(2j-k)! |S_{k,j,m}|^2 \right)^2,$$

while their numerators, respectively, are

$$\begin{aligned} \sum_{k=0}^{2j} (j-k)k!(2j-k)! |S_{k,j,-m}|^2 &= -|\xi_-|^{4m} \left( \sum_{k=0}^{2j} (j-k)k!(2j-k)! |S_{k,j,m}|^2 \right)^2, \\ \sum_{k=0}^{2j} (j-k)^2 k!(2j-k)! |S_{k,j,-m}|^2 &= |\xi_-|^{4m} \left( \sum_{k=0}^{2j} (j-k)^2 k!(2j-k)! |S_{k,j,m}|^2 \right)^2. \end{aligned}$$

These immediately imply that  $\langle W_{s,j,-m}, \mathbf{J}_3 W_{s,j,-m} \rangle = -\langle W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$ , as we expected, and  $\langle \mathbf{J}_3 W_{s,j,-m}, \mathbf{J}_3 W_{s,j,-m} \rangle = \langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$ .

Next we show that  $\langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$  is a continuous function of  $\alpha_3$ , although  $\langle W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$  has a jump at  $\alpha_3 = 0$  for  $\lambda > 1$ .

$\xi_-$ , as a function of  $\alpha_3$  (see (2.17)), is *not* continuous at  $\alpha_3 = 0$  for  $\lambda > 1$ : Its  $\alpha_3 \rightarrow 0$  limit from the left is  $-i(\lambda + \sqrt{\lambda^2 - 1})$ , while from the right it is  $-i(\lambda - \sqrt{\lambda^2 - 1})$ . Hence,  $S_{k,j,m}$  is not continuous there. Now we calculate the left limit of  $S_{k,j,m}$ , denoted by  $S_{k,j,m}^-$ , and relate it to the right limit  $S_{k,j,m}^+$  of  $S_{k,j,m}$ . The key observation is that  $\lambda + \sqrt{\lambda^2 - 1} = (\lambda - \sqrt{\lambda^2 - 1})^{-1}$ , but the strategy of the calculation is the same that we followed in the derivation of (4.38). In particular, for  $k = 0, \dots, j - m - 1$ ,  $m \geq 0$ , with the notation  $\bar{r} := j - m - r$  the main steps are

$$\begin{aligned}
S_{k,j,m}^- &= (-i)^{-k} (\lambda + \sqrt{\lambda^2 - 1})^{-k} \sum_{r=0}^k (-)^r \binom{j-m}{r} \binom{j+m}{k-r} (-i(\lambda + \sqrt{\lambda^2 - 1}))^{2r} \\
&= (-i)^{\bar{k}-2m} (\lambda - \sqrt{\lambda^2 - 1})^{-\bar{k}+2m} \sum_{\bar{r}=j-m}^{\bar{k}-j-m} (-)^{\bar{r}} \binom{j-m}{\bar{r}} \binom{j+m}{\bar{k}-\bar{r}} (-i(\lambda - \sqrt{\lambda^2 - 1}))^{2\bar{r}} \\
&= (-)^{j-k} (\lambda - \sqrt{\lambda^2 - 1})^{2m} S_{k,j,m}^+. \tag{4.39}
\end{aligned}$$

One can check that  $S_{k,j,m}^- = (-)^{j-k} (\lambda - \sqrt{\lambda^2 - 1})^{2m} S_{k,j,m}^+$  holds in the other cases, too. Hence the  $\alpha_3 \rightarrow 0$  limit from the left of both the denominator and the numerator in (4.33) is  $(\lambda - \sqrt{\lambda^2 - 1})^{4m}$  times their  $\alpha_3 \rightarrow 0$  limit from the right, and hence the left and right limits of  $\langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$  coincide. Similar argumentation confirms that  $\langle W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$  changes sign at  $\alpha_3 = 0$  for  $\lambda > 1$ , i.e. it jumps there.

Since  $\langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$  and  $(\langle W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle)^2$  are continuous and  $\Delta_\phi \mathbf{J}(\alpha) = \lambda \Delta_\phi \mathbf{J}(\beta)$  holds, (2.31) implies that the standard deviations, both for  $\mathbf{J}(\alpha)$  and  $\mathbf{J}(\beta)$ , are continuous functions of the parameters  $\lambda$  and  $\alpha_3$ .

#### 4.2.4 The $\lambda \rightarrow 0$ and $\lambda \rightarrow \infty$ limits of the standard deviations

First consider the  $\lambda \rightarrow 0$  limit. By (2.17), in this limit,

$$\xi_- = -\sqrt{\frac{1 - \alpha_3}{1 + \alpha_3}} + O(\lambda^2) + iO(\lambda).$$

Hence, by (4.31), all the  $S_{k,j,m}$  are bounded in this limit, implying that  $\langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle = \langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle|_{\lambda=0} + O(\lambda)$ . Since, as we saw in subsection 2.2, the expectation value is also finite in this limit, the standard deviation for  $\mathbf{J}(\beta)$  is finite. But then  $\Delta_\phi \mathbf{J}(\alpha) = \lambda \Delta_\phi \mathbf{J}(\beta)$  implies that both the standard deviation for  $\mathbf{J}(\alpha)$  and the product uncertainty,  $\Delta_\phi \mathbf{J}(\alpha) \Delta_\phi \mathbf{J}(\beta)$ , tend to zero as  $\lambda$  in the  $\lambda \rightarrow 0$  limit.

In the  $\lambda \rightarrow \infty$  limit, (2.17) gives that

$$\xi_- = -\frac{1}{2} \sqrt{1 - \alpha_3^2} (i\lambda^{-1} + \alpha_3 \lambda^{-2} + O(\lambda^{-3})), \quad \text{if } \alpha_3 > 0, \tag{4.40}$$

$$\xi_- = -\frac{2}{\sqrt{1 - \alpha_3^2}} (i\lambda + |\alpha_3| + O(\lambda^{-1})). \quad \text{if } \alpha_3 \leq 0. \tag{4.41}$$

To calculate the standard deviations in this limit, let us recall from the previous subsection that  $\langle W_{s,j,-m}, \mathbf{J}_3 W_{s,j,-m} \rangle = -\langle W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$  and  $\langle \mathbf{J}_3 W_{s,j,-m}, \mathbf{J}_3 W_{s,j,-m} \rangle = \langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle$ , and hence the standard deviation of  $\mathbf{J}_3$  in the states  $W_{s,j,m}$  and  $W_{s,j,-m}$  is the same. Therefore, it is enough to consider the  $m = 0$  and  $m > 0$  disjoint cases.

First suppose that  $m = 0$ . Then, with the notations of the previous subsection, the hidden symmetry (4.38) yields that  $S_{k,j,0} = (-)^{k-j} S_{k,j,0}$ . Hence the denominator and the numerator, respectively, in (4.33) are

$$(j!)^2 |S_{j,j,0}|^2 + \sum_{k=0}^{j-1} k!(2j-k)! |S_{k,j,0}|^2, \quad 2 \sum_{k=0}^{j-1} (j-k)^2 k!(2j-k)! |S_{k,j,0}|^2;$$

while the numerator in (4.32) is vanishing. Therefore, the square of the standard deviation of  $\mathbf{J}_3$  in the states  $W_{s,j,0}$  is simply the quotient of these two. Thus the only thing to do is to determine the leading order terms of these two expressions in the cases (4.40) and (4.41). However, in both cases

$$S_{k,j,0} = (\xi_-)^{-k} \sum_{r=0}^k (-)^r \binom{j}{r} \binom{j}{k-r} (\xi_-)^{2r} = \left( \frac{2i}{\sqrt{1-\alpha_3^2}} \right)^k \binom{j}{k} \lambda^k + O(\lambda^{k-1}),$$

yielding that for the leading order term in the numerator is at  $k = j - 1$ , and in the denominator it is at  $k = j$ . Therefore,

$$\langle \mathbf{J}_3 W_{s,j,0}, \mathbf{J}_3 W_{s,j,0} \rangle = \frac{1}{2} j(j+1)(1-\alpha_3^2) \lambda^{-2} + O(\lambda^{-3});$$

and hence, by  $m = 0$ , the standard deviation for  $\mathbf{J}(\beta)$  tends to zero as  $1/\lambda$ , and that for  $\mathbf{J}(\alpha)$  remains finite. Thus, by (2.31) in the states  $\phi$  with eigenvalue  $m = 0$ , the product uncertainty  $\Delta_\phi \mathbf{J}(\alpha) \Delta_\phi \mathbf{J}(\beta) = \lambda (\Delta_\phi \mathbf{J}(\beta))^2$  tends to zero as  $1/\lambda$  in the  $\lambda \rightarrow \infty$  limit.

Next suppose that  $m > 0$ . If  $j$  is integer, then the common denominator in (4.32) and (4.33) is

$$\begin{aligned} D := & \sum_{k=0}^{j-m-1} k!(2j-k)! |S_{k,j,m}|^2 + \sum_{k=j-m}^{j-1} k!(2j-k)! |S_{k,j,m}|^2 \\ & + (j!)^2 |S_{j,j,m}|^2 + \sum_{k=j+1}^{j+m} k!(2j-k)! |S_{k,j,m}|^2 + \sum_{k=j+m+1}^{2j} k!(2j-k)! |S_{k,j,m}|^2. \end{aligned}$$

Then, using (4.34), (4.35), (4.37) and (4.40), it is a tedious but routine calculation to determine the leading order terms in  $D$  when  $\alpha_3 > 0$ . It is

$$D = (j-m)!(j+m)! \left( \frac{2}{1-\alpha_3^2} \right)^{j+m} \lambda^{2(j+m)} + O(\lambda^{2(j+m-1)}).$$

We obtain the same expression when  $j$  is half-odd-integer. Similar calculations show that the numerators  $N_1$  and  $N_2$  in (4.32) and (4.33), respectively, are

$$\begin{aligned} N_1 &= -m(j+m)!(j-m)! \left( \frac{2}{1-\alpha_3^2} \right)^{j+m} \lambda^{2(j+m)} + O(\lambda^{2(j+m-1)}), \\ N_2 &= m^2(j+m)!(j-m)! \left( \frac{2}{1-\alpha_3^2} \right)^{j+m} \lambda^{2(j+m)} + O(\lambda^{2(j+m-1)}). \end{aligned}$$

If  $\alpha_3 \leq 0$ , then, using (4.40), analogous calculations yield the same expressions for  $D$ ,  $N_1$  and  $N_2$  except that  $m$  should be replaced by  $-m$ . Hence,

$$\langle W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle = \text{sign}(\alpha_3) m \hbar + O(\lambda^{-2}), \quad \langle \mathbf{J}_3 W_{s,j,m}, \mathbf{J}_3 W_{s,j,m} \rangle = m^2 \hbar^2 + O(\lambda^{-2});$$

which, by (2.31), imply that, in the states  $\phi$  with eigenvalue  $m > 0$ , the standard deviation for  $\mathbf{J}(\beta)$  tends to zero as  $1/\lambda$ , and that for  $\mathbf{J}(\alpha)$  remains finite. These imply that the product uncertainty  $\Delta_\phi \mathbf{J}(\alpha) \Delta_\phi \mathbf{J}(\beta) = \lambda(\Delta_\phi \mathbf{J}(\beta))^2$  tends to zero as  $1/\lambda$  in the  $\lambda \rightarrow \infty$  limit.

## Acknowledgment

Thanks are due to Paul Tod for the idea how the spectrum of the operator  $\mathbf{J}(\alpha) - i\lambda\mathbf{J}(\beta)$  of subsection 2.2 can be calculated in the most economical way, as well as for his helpful remarks and comments; and also to Péter Vecsernyés for his remarks on the classical limit of non-relativistic quantum systems.

## References

- [1] J. R. Klauder, B.-S. Kagerstam, Coherent states, Applications in physics and mathematical physics, in *Coherent States*, World Scientific, Singapore 1985
- [2] W.-M. Zhang, D. H. Feng, R. Gilmore, Coherent states: Theory and some applications, *Rev. Mod. Phys.* **62** 867-927 (1990)
- [3] E. Arthurs, J. L. Kelly, On the simultaneous measurements of a pair of conjugate observables, *Bell System Technical J.* **44** 725-729 (1965)
- [4] C. Y. She, H. Heffner, Simultaneous measurement of noncommuting observables, *Phys. Rev.* **152** 1103-1110 (1966)
- [5] J. M. Radcliffe, Some properties of coherent spin states, *J. Phys. A: Gen. Phys.* **4** 313-323 (1971)
- [6] E. H. Lieb, The classical limit of quantum spin systems, *Commun. Math. Phys.* **31** 327-340 (1973)
- [7] C. Aragone, G. Guerri, S. Salamo, J. L. Tani, Intelligent spin states, *J. Phys. A: Math., Nucl. Gen.*, **7** L149-L151 (1974)
- [8] X. Wang, B. C. Sanders, S. Pan, Entangled coherent states for systems with  $SU(2)$  and  $SU(1, 1)$  symmetries, *J. Phys. A: Math. Gen.* **33** 7451-7467 (2000)
- [9] R. Penrose, Combinatorial quantum theory and quantized directions, in *Advances in Twistor Theory*, Eds. L. P. Houghston, R. S. Ward, Pitman Publ. Ltd, London 1979  
R. Penrose, Angular momentum: An approach to combinatorial spacetime, in *Quantum Theory and Beyond*, Ed. T. Bastin, Cambridge Univ. Press, Cambridge 1971  
R. Penrose, On the nature of quantum geometry, in *Magic without Magic*, Ed. J. Klauder, Freeman, San Francisco 1972
- [10] L. B. Szabados, Three-space from quantum mechanics, (in preparation)
- [11] L. B. Szabados, The ‘most classical states’ of Euclidean invariant elementary quantum mechanical systems, (in preparation)

- [12] R. Penrose, W. Rindler, *Spinors and Spacetime*, vol 1, Cambridge University Press, Cambridge 1984
- [13] S. A. Hugget, K. P. Tod, *An Introduction to Twistor Theory*, London Mathematical Society Student Texts 4, 2nd edition, Cambridge University Press, Cambridge 1994
- [14] M. Eastwood, P. Tod, Edth – a differential operator on the sphere, *Math. Proc. Camb. Phil. Soc.* **92** 317-330 (1982)
- [15] E. T. Newman, R. Penrose, Note on the Bondi–Metzner–Sachs group, *J. Math. Phys.* **7** 863-879 (1966)
- [16] J. N. Goldberg, A. J. Macfarlane, E. T. Newman, F. Rochlich, E. C. G. Sudarshan, Spin-s spherical harmonics and  $\delta$ , *J. Math. Phys.* **8** 2155-2161 (1967)
- [17] J. Frauendiener, L. B. Szabados, The kernel of the edth operators on higher-genus spacelike 2-surfaces, *Class. Quantum Grav.* **18** 1003-1014 (2001), gr-qc/0010089
- [18] N. Wheeler, Simultaneous measurement of noncommuting quantum observables, (manuscript) 2012, URL: <https://www.reed.edu/physics/faculty/wheeler/documents/index.html>
- [19] K. P. Tod, (private communication, 2021 January)
- [20] S. Stenholm, Simultaneous measurement of conjugate variables, *Ann. Phys. (N.Y.)* **218** 233-254 (1992)
- [21] M. G. Raymer, Uncertainty principle for joint measurement of noncommuting variables, *Am. J. Phys.* **62** 986-993 (1994)
- [22] R. F. Streater, A. S. Wightman, *PCT, Spin and Statistics, and All That*, W. A. Benjamin, INC, New York 1964