

Search for gravitational waves from the coalescence of sub-solar mass binaries in the first half of Advanced LIGO and Virgo’s third observing run

Alexander H. Nitz* and Yi-Fan Wang

Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), D-30167 Hannover, Germany and Leibniz Universität Hannover, D-30167 Hannover, Germany

(Dated: June 21, 2021)

We present a search for gravitational waves from the coalescence of sub-solar mass black hole binaries using data from the first half of Advanced LIGO and Virgo’s third observing run. The observation of a sub-solar mass black hole merger is a clear indication of primordial origin; primordial black holes may contribute to the dark matter distribution. We search for black hole mergers where the primary mass is $0.1 - 7M_{\odot}$ and the secondary mass is $0.1 - 1M_{\odot}$. A variety of models predict the production and coalescence of binaries containing primordial black holes; some involve dynamical assembly which may allow for residual eccentricity to be observed. For component masses $> 0.5M_{\odot}$, we also search for sources in eccentric orbits, measured at a reference gravitational-wave frequency of 10 Hz, up to $e_{10} \sim 0.3$. We find no convincing candidates and place new upper limits on the rate of primordial black hole mergers. The merger rate of 0.5-0.5 (1.0-1.0) M_{\odot} sources is < 7100 (1200) $\text{Gpc}^{-3}\text{yr}^{-1}$. Our limits are $\sim 3 - 4\times$ more constraining than prior analyses. Finally, we demonstrate how our limits can be used to constrain arbitrary models of the primordial black hole mass distribution and merger rate.

I. INTRODUCTION

Gravitational-wave astronomy has entered an era of routine observations. The Advanced LIGO [1] and Virgo [2] observatories have now completed three observing runs (O1, O2, and O3); each was accompanied by significant increases in sensitivity [3]. To date, over 50 binary black hole mergers have been reported [4–6]. These observations have had significant impact on the study of the merger rate and population of compact objects [7]; notable events confirm the likely existence of black holes with component masses in the pair-instability gap ($> 50 M_{\odot}$) [4, 8] or in the region 3-5 M_{\odot} [9]. The possibility that these extremal parts of the distribution [10–13] or a fraction of the bulk of observed mergers [14–22] may be due to the coalescence of primordial black holes (PBHs) is under active investigation.

Currently, there is no clear observational evidence for the existence of PBHs. However, in addition to providing an explanation for some of the observed LIGO and Virgo mergers [11], primordial black holes may be the origin of some observed microlensing incidents [23], excess cross-correlation between cosmic x-ray and cosmic microwave background [24], the current excess in gravitational-wave background observed by NANOGrav [25], and the seeds for galaxy and supermassive black hole formation [26–28]. Many of these observations are also consistent with more mundane explanations and standard stellar-formation scenarios [7]. In contrast, there are no known mechanisms through standard stellar evolution to produce sub-solar mass black holes; the observation of a single sub-solar mass black hole would be decisive for the existence of primordial black holes.

Several searches for gravitational waves from the coalescence of sub-solar mass mergers have already been conducted using data from LIGO’s first two observing runs (O1, O2); these include searches for comparable mass binary black holes [29–31], eccentric mergers [32], and high-mass-ratio sources [33]. No likely candidates have been found. In this paper we report a search for gravitational waves from the coalescence of black holes with primary mass $0.1 - 7M_{\odot}$ and secondary mass $0.1 - 1M_{\odot}$ using the open data from the first half of the third observing (O3a) run of Advanced LIGO and Virgo. The parameter space for our search region is shown in Fig. 1. The most significant candidate in our search has a false alarm rate of 1 per O(month). Given the time observed, we consider our results consistent with a null observation and place new limits on the rate of sub-solar mass mergers which are $3 - 4\times$ more stringent than prior analyses.

Our limits on the rate of sub-solar mass mergers can be related to constraints on the fraction of dark matter composed of PBHs; this requires a model of the binary’s formation to predict the PBH abundance from the observed merger rate. Current astrophysical models have large uncertainties in their predictions of both the black hole mass function and binary formation rate [18–20, 34–37]. Primordial black holes may form binaries in the early universe if they can decouple from the cosmic expansion. However, it is under investigation what fraction of binaries would be disrupted in the following evolution. Ref. [38] shows with N-body simulations that a significant fraction would be disrupted if the fraction that primordial black hole contribute to the dark matter is $f_{\text{PBH}} = 100\%$. Primordial black holes can also form binaries in the late universe by dynamical capture due to gravitational-wave dissipation [18, 39, 40]; this scenario may also lead to residual eccentricity by the time it is observable by gravitational-wave detectors [41]. However, the event rate of binaries formed in the early universe

* alex.nitz@aei.mpg.de




FIG. 1. The regions searched by recent gravitational-wave analyses of the LIGO and Virgo data as a function of detector-frame primary and secondary mass. The region we search for non-eccentric sources (green) and the region sensitive to sources with eccentricity up to $e_{10} \sim 0.3$ (red) are shown. For comparison, we include the search region of the most recent sub-solar mass search by the LVC [30] (dashed), the search for high-mass-ratio mergers [33], and searches for standard stellar-mass sources [4, 5]. The axes are truncated at $200 M_{\odot}$.

is expected to be dominant compared with the late universe channel, depending on the intensity of binary disruption [38].

Due to the wide variety of models, in this paper, we consider a fiducial model which assumes a monochromatic distribution of primordial black hole mass. The same model was used in past sub-solar mass searches [29–33], and we include for comparison purposes. We also consider the uncertainty on the rate estimates arising from the fraction of binaries that are disrupted after formation [38]. Constraints for specific models with broad mass distributions can be derived from our observational constraints.

II. SEARCH

We conduct our analysis in a similar manner to our previously presented search of the first two observing runs [32]; however, for the first time we include data from the Virgo observatory. We use the open-source PyCBC toolkit [42, 43] to conduct a matched-filtering based search [44]. Matched filtering allows us to extract a potential signal using the predicted gravitational waveform as a template. Potential candidates are assessed for consistency between the operating observatories [45] and against the expected morphology of the gravitational waveform [46, 47]. Each candidate is assigned a ranking statistic value that takes into account these factors in addition to the measured noise variance [48, 49].

The statistical significance of each candidate is as-

essed by comparing to an empirically measured distribution of false alarms [42]. The distribution is measured by conducting numerous fictitious analyses whereby we offset detectors’ data in time. This procedure purposefully violate the time-of-flight constraints between the detectors to remove coincident astrophysical sources and create analyses containing only false alarms [50–52].

As matched filtering requires accurate models of the expected gravitational-wave signal, we use a combination of the TaylorF2 post-Newtonian approximant accurate to 3.5PN [53–56] and the TaylorF2e model [57–59]. TaylorF2e is an extension of TaylorF2 which includes corrections for moderate eccentricity. Both TaylorF2 and TaylorF2e model only the inspiral portion of a gravitational-wave signal and do not account for the phase where the binary finally merges. The merger can be safely neglected as we search for sources only up to a total mass of $8 M_{\odot}$. For these sources, the merger occurs at a frequency above the most sensitive band of the instruments.

To search for a broad region, we create a discrete bank of templates designed to ensure that we recover $> 95\%$ of a signal’s signal-to-noise ratio if it has parameters within the boundaries of our search. Our bank is designed to recover binaries in quasi-circular orbits where the primary mass is $0.1 - 7 M_{\odot}$ and the secondary mass is $0.1 - 1 M_{\odot}$. In addition, for sources with component masses $> 0.5 M_{\odot}$, the bank is designed to recover sources with eccentric orbits up to $e_{10} \sim 0.3$, where e_{10} is the eccentricity at the fiducial dominant-mode gravitational-wave frequency of 10 Hz. We assume that PBHs will have negligible spin; this is consistent with the predictions of PBH spin distributions [60–64]. To save on computational cost, we limit the starting frequency of each template so that its duration is $< 512s$; otherwise a cutoff at a gravitational-wave frequency of 20 Hz is used.

III. OBSERVATIONAL RESULTS

We search for gravitational waves from the coalescence of sub-solar mass compact binaries using the public LIGO and Virgo data from the first half of the third observing run (O3a) [65, 66]; data from the second half of the observing run is not yet available. We analyze the nearly 150 days of data where at least two observatories were operating; the twin LIGO observatories were operating for ~ 100 days of this period. In comparison to the previous observing run, the LIGO instruments had $\sim 30 - 50\%$ greater range [3].

The most significant candidates from our search are identified at a false alarm rate of one per O(month); this is consistent with a null observation and our expectation that the noise candidates follow a Poisson distribution. Under the assumption of a null detection, we place limits on the rate of binary mergers at 90% confidence using the loudest event method of Ref. [67]. The limit on the




FIG. 2. Upper limit on the rate of mergers at 90% confidence (R_{90}) for our search (purple). For comparison, we show the previous limits from our search of the O1 and O2 data (blue dashed) [32] in addition to the most recent results from sub-solar masses searches conducted by the LVC of the O2 data (black dotted) [30]. These all assume that sources are quasi-circular; constraints assuming a uniform distribution of $e_{10} \in [0, 0.3]$ are also shown (green). As expected, the limit for eccentric sources closely matches that for quasi-circular binaries where the component masses are $> 0.5M_{\odot}$. Shaded regions show the one sigma uncertainty on the rate due to the monte-carlo estimation of the search’s surveyed volume-time.

merger rate R_{90} is given as

$$R_{90} = \frac{2.3}{\sqrt{V}T} \quad (1)$$

where V is the estimated sensitive volume of the analysis assessed at the false alarm rate of the most significant observed candidate and T is the duration of the observation period. We estimate the surveyed volume-time of our analysis by measuring our analysis’ response to a simulated population of $O(10^5)$ sources.

The resulting upper limit on the merger rate, combined with the limit from the prior analysis of O1 and O2 [32], is shown as a function of chirp mass in Fig. 2, where chirp mass is defined as $\mathcal{M} = (m_1 m_2)^{3/5} / (m_1 + m_2)^{1/5}$ and $m_{1,2}$ are the masses of the two components of a binary. For sources within our target parameter space, this limit also holds for sources with varied mass ratios, but the same chirp mass; a similar conclusion was noted in Ref. [30]. Our results limit the merger rate for 0.1-0.1, 0.5-0.5, and 1.0-1.0 M_{\odot} sources to < 670000 , 7100, and 1200 $\text{Gpc}^{-3}\text{yr}^{-1}$, respectively. This is an improvement between $3 - 4\times$ over previous analyses which only used data from the first two observing runs.

IV. IMPLICATIONS FOR PRIMORDIAL BLACK HOLE ABUNDANCE

We use the observational upper limits on the sub-solar mass compact binary merger rate to constrain models of primordial black hole binary formation. Existing models have significant uncertainties on the primordial black hole mass distribution and the binary formation rate [10, 18–20, 34–37]. Constraints on models can be derived from the observational event rate upper limits given a specific primordial black hole mass distribution. We consider a fiducial mass distribution, a delta distribution following Refs. [29, 30] and our previous work [32, 33] to allow for a consistent comparison.

For the binary formation rate, we consider the mechanism initially proposed by [34] and developed by [20, 22, 36–38, 68] where primordial black holes form bounded binaries in the early universe and merge recently. The binaries form in the late universe is subdominant [41], therefore we use the search results from circular binaries to constrain the early universe formation model. The binary merger rate is given by Refs. [36, 37] for a general mass distribution $P(m)$

$$\begin{aligned} R(\tilde{f}_{\text{PBH}}, m_1, m_2) &= 3 \times 10^6 \tilde{f}_{\text{PBH}}^2 (0.7 \tilde{f}_{\text{PBH}}^2 + \sigma_{\text{eq}}^2)^{-\frac{21}{4}} \\ &\times (m_1 m_2)^{\frac{3}{37}} (m_1 + m_2)^{\frac{36}{37}} \min\left(\frac{P(m_1)}{m_1}, \frac{P(m_2)}{m_2}\right) \\ &\times \left(\frac{P(m_1)}{m_1} + \frac{P(m_2)}{m_2}\right) \text{Gpc}^{-3}\text{yr}^{-1}, \quad (2) \end{aligned}$$

where $R \cdot dm_1 dm_2$ is the event rate at the binary component masses $m_{1/2} M_{\odot}$. The normalization for mass distribution is $\int P(m) dm = 1$. The parameter $\sigma_{\text{eq}} = 0.005$ is the variance of dark matter density perturbation at the matter radiation equality epoch [37], and we keep this factor separate from other binary disruption effects to perform consistent comparison with previous work. Ref. [38] has used an N-body simulation to show most binaries after formation would be disrupted by their environment if $f_{\text{PBH}} = 100\%$, and thus introduced a suppression factor S with value < 1 which accounts for the disruption as a function of primordial black hole mass and fraction. For a relatively narrow mass distribution, Ref. [22] shows S (referred to as S_2 in Ref. [22]) is estimated to be 1% for $f_{\text{PBH}} = 100\%$ and ~ 1 for $f_{\text{PBH}} < 1\%$ and the suppression is only a function of f_{PBH} . However, underestimation by one order of magnitude may exist for $f_{\text{PBH}} = 100\%$ if the merger of binaries perturbed by the environment after formation is included [22, 31, 68]. To account for this uncertainty, we define an effective fraction which relates to the true fraction by $\tilde{f}_{\text{PBH}}^{53/37} = S f_{\text{PBH}}^{53/37}$. Constraints in our past work [32, 33] on f_{PBH} should also be understood as constraints on \tilde{f}_{PBH} ; they implicitly assume the suppression factor is unity. The true fraction f_{PBH} can be recovered with a well-understood estimation of S .

For a delta distribution of mass, Eq. 3 is reduced to

$$R(\tilde{f}_{\text{PBH}}, m) = 3 \cdot 10^6 \tilde{f}_{\text{PBH}}^2 (0.7 \tilde{f}_{\text{PBH}}^2 + \sigma_{\text{eq}}^2)^{-\frac{21}{74}} m^{-\frac{32}{37}} \quad (3)$$

Using the observational rate limit given in Fig. 2, the upper limit on the effective fraction of primordial black hole with a delta distribution of mass is shown in Fig. 3. A comparison with previous search results for sub-solar mass compact binaries is also plotted. The results should be interpreted as constraints on the combined effect on the binary merger rate from the abundance of primordial black holes and the merger rate suppression due to environmental interaction. Future improvements on theoretical modeling and observational results can resolve the entanglement. Overall, results in Fig. 3 show that our constraints are $\sim 2 - 3$ times tighter than [32] using O1 and O2 data.

Constraints can also be derived on primordial black hole scenarios which predict broad mass distributions. By extending Eq. 1, the corresponding 90% upper limits can be obtained by requiring

$$\int R(\vec{\theta}, m_1, m_2) VT(m_1, m_2) dm_1 dm_2 = 2.3, \quad (4)$$

where R is the model-predicted merger rate density as a function of the component masses and may include additional model parameters $\vec{\theta}$. VT is the surveyed volume-time of our analysis as a function of the component masses; this is available as part of our data release.

To illustrate, we consider the scenario that primordial black holes are produced in the early Universe QCD phase transition era [10]. This model is shown to have a peak at $\sim 1M_{\odot}$ and to be able to account for the origin of the extremal parts of the already observed binary black hole merger distribution. We choose the mass distribution $P(m)$ by following Ref. [10] from the early universe scalar perturbation spectral index 0.96 and use the binary formation rate of Eq. 2. We assume a mass-independent suppression factor may be applicable here given the narrow peak of $P(m)$. The observational upper limits on event rate from our search require $\tilde{f}_{\text{PBH}} \leq 1\%$ in this scenario. For more general mass distributions, Eq. 4 is capable of constraining the PBH abundance or other model parameters given a specific primordial black hole mass distribution and prescription which can predict the resulting merger rate density $R(\vec{\theta}, m_1, m_2)$.

V. CONCLUSIONS

We conduct a search for gravitational-waves from the coalescence of sub-solar mass black holes using data from the first half of the third observing run of Advanced LIGO and Virgo. We find no clear detections and so place new limits on the rate of mergers. The increased sensitivity of the O3a data allows us to improve upon the state-of-the-art limits by $3 - 4\times$. The second half of




FIG. 3. The upper limits on the effective fraction of the primordial black hole contribution to dark matter, \tilde{f}_{PBH} , from the search for sub-solar mass compact binaries in LVC O1-O3a data. As a comparison, we also plot the constraints from O1-O2 data [32] for the monochromatic mass distribution and for the two-point mass distribution of high-mass-ratio binaries [33] where the primary mass is fixed to the average of the 2-OGC events $\simeq 37M_{\odot}$. Note that this latter limit requires the additional assumption that the primary mass abundance be consistent with accounting for the majority of LIGO-observed BBH mergers.

the O3 data (O3b) would be expected to improve these limits by another factor of $1.5 - 2\times$.

We apply our rate limits to a fiducial monochromatic mass distribution and compare our results to the limits from prior analyses. Overall the constraints on the effective fraction of primordial black holes in dark matter are 2-3 tighter than the previous results. Our observational results can also help resolve the event rate modeling uncertainties by constraining model parameters. If we assume that dark matter consisted entirely of black holes, we constrain the suppression factor to $S \leq 0.1\%$ for $0.5-0.5 M_{\odot}$ mergers.

Lastly, we demonstrate how to apply our limits to models that can predict the rate of PBH mergers. To aid in comparing our rate limits to the various formation scenarios leading to sub-solar mass mergers, we make the detailed constraints available at <https://github.com/gwastro/subsolar-o3a-search>. In addition, we make available the configuration files and template bank necessary to reproduce the analysis.

ACKNOWLEDGMENTS

We acknowledge the Max Planck Gesellschaft. We thank the computing team from AEI Hannover for their significant technical support. This research has made use of data from the Gravitational Wave Open Science Center (<https://www.gw-openscience.org>), a service of LIGO

Laboratory, the LIGO Scientific Collaboration and the Virgo Collaboration. LIGO is funded by the U.S. National Science Foundation. Virgo is funded by the French Centre National de Recherche Scientifique (CNRS), the Italian Istituto Nazionale della Fisica Nucleare (INFN) and the Dutch Nikhef, with contributions by Polish and Hungarian institutes.

-
- [1] J. Aasi *et al.* (LIGO Scientific Collaboration), “Advanced LIGO,” *Class. Quantum Grav.* **32**, 074001 (2015), [arXiv:1411.4547 \[gr-qc\]](https://arxiv.org/abs/1411.4547).
- [2] F. Acernese *et al.* (VIRGO), “Advanced Virgo: a second-generation interferometric gravitational wave detector,” *Class. Quantum Grav.* **32**, 024001 (2015), [arXiv:1408.3978 \[gr-qc\]](https://arxiv.org/abs/1408.3978).
- [3] Aaron Buikema *et al.* (aLIGO), “Sensitivity and performance of the Advanced LIGO detectors in the third observing run,” *Phys. Rev. D* **102**, 062003 (2020), [arXiv:2008.01301 \[astro-ph.IM\]](https://arxiv.org/abs/2008.01301).
- [4] R. Abbott *et al.* (LIGO Scientific, Virgo), “GWTC-2: Compact Binary Coalescences Observed by LIGO and Virgo During the First Half of the Third Observing Run,” *Phys. Rev. X* **11**, 021053 (2021), [arXiv:2010.14527 \[gr-qc\]](https://arxiv.org/abs/2010.14527).
- [5] Alexander H. Nitz, Collin D. Capano, Sumit Kumar, Yi-Fan Wang, Shilpa Kastha, Marlin Schäfer, Rahul Dhurkunde, and Miriam Cabero, “3-OGC: Catalog of gravitational waves from compact-binary mergers,” (2021), [arXiv:2105.09151 \[astro-ph.HE\]](https://arxiv.org/abs/2105.09151).
- [6] Tejaswi Venumadhav, Barak Zackay, Javier Roulet, Liang Dai, and Matias Zaldarriaga, “New Binary Black Hole Mergers in the Second Observing Run of Advanced LIGO and Advanced Virgo,” (2019), [arXiv:1904.07214 \[astro-ph.HE\]](https://arxiv.org/abs/1904.07214).
- [7] R. Abbott *et al.* (LIGO Scientific, Virgo), “Population Properties of Compact Objects from the Second LIGO-Virgo Gravitational-Wave Transient Catalog,” *Astrophys. J. Lett.* **913**, L7 (2021), [arXiv:2010.14533 \[astro-ph.HE\]](https://arxiv.org/abs/2010.14533).
- [8] R. Abbott *et al.* (LIGO Scientific, Virgo), “GW190521: A Binary Black Hole Merger with a Total Mass of $150 M_{\odot}$,” *Phys. Rev. Lett.* **125**, 101102 (2020), [arXiv:2009.01075 \[gr-qc\]](https://arxiv.org/abs/2009.01075).
- [9] R. Abbott *et al.* (LIGO Scientific, Virgo), “GW190814: Gravitational Waves from the Coalescence of a 23 Solar Mass Black Hole with a 2.6 Solar Mass Compact Object,” *Astrophys. J. Lett.* **896**, L44 (2020), [arXiv:2006.12611 \[astro-ph.HE\]](https://arxiv.org/abs/2006.12611).
- [10] Bernard Carr, Sebastien Clesse, Juan García-Bellido, and Florian Kühnel, “Cosmic conundra explained by thermal history and primordial black holes,” *Phys. Dark Univ.* **31**, 100755 (2021), [arXiv:1906.08217 \[astro-ph.CO\]](https://arxiv.org/abs/1906.08217).
- [11] Sebastien Clesse and Juan Garcia-Bellido, “GW190425, GW190521 and GW190814: Three candidate mergers of primordial black holes from the QCD epoch,” (2020), [arXiv:2007.06481 \[astro-ph.CO\]](https://arxiv.org/abs/2007.06481).
- [12] Kyriakos Vattis, Isabelle S. Goldstein, and Savvas M. Koushiappas, “Could the 2.6 M_{\odot} object in GW190814 be a primordial black hole?” *Phys. Rev. D* **102**, 061301 (2020), [arXiv:2006.15675 \[astro-ph.HE\]](https://arxiv.org/abs/2006.15675).
- [13] V. De Luca, V. Desjacques, G. Franciolini, P. Pani, and A. Riotto, “GW190521 Mass Gap Event and the Primordial Black Hole Scenario,” *Phys. Rev. Lett.* **126**, 051101 (2021), [arXiv:2009.01728 \[astro-ph.CO\]](https://arxiv.org/abs/2009.01728).
- [14] Gabriele Franciolini, Vishal Baibhav, Valerio De Luca, Ken K. Y. Ng, Kaze W. K. Wong, Emanuele Berti, Paolo Pani, Antonio Riotto, and Salvatore Vitale, “Evidence for primordial black holes in LIGO/Virgo gravitational-wave data,” (2021), [arXiv:2105.03349 \[gr-qc\]](https://arxiv.org/abs/2105.03349).
- [15] V. De Luca, G. Franciolini, P. Pani, and A. Riotto, “Bayesian Evidence for Both Astrophysical and Primordial Black Holes: Mapping the GWTC-2 Catalog to Third-Generation Detectors,” *JCAP* **05**, 003 (2021), [arXiv:2102.03809 \[astro-ph.CO\]](https://arxiv.org/abs/2102.03809).
- [16] Karsten Jedamzik, “Consistency of Primordial Black Hole Dark Matter with LIGO/Virgo Merger Rates,” *Phys. Rev. Lett.* **126**, 051302 (2021), [arXiv:2007.03565 \[astro-ph.CO\]](https://arxiv.org/abs/2007.03565).
- [17] Juan García-Bellido, José Francisco Nuño Siles, and Ester Ruiz Morales, “Bayesian analysis of the spin distribution of LIGO/Virgo black holes,” *Phys. Dark Univ.* **31**, 100791 (2021), [arXiv:2010.13811 \[astro-ph.CO\]](https://arxiv.org/abs/2010.13811).
- [18] Simeon Bird, Ilias Cholis, Julian B. Muñoz, Yacine Ali-Haïmoud, Marc Kamionkowski, Ely D. Kovetz, Alvise Raccanelli, and Adam G. Riess, “Did LIGO Detect Dark Matter?” *Phys. Rev. Lett.* **116**, 201301 (2016), [arXiv:1603.00464 \[astro-ph.CO\]](https://arxiv.org/abs/1603.00464).
- [19] Sebastien Clesse and Juan García-Bellido, “The clustering of massive Primordial Black Holes as Dark Matter: measuring their mass distribution with Advanced LIGO,” *Phys. Dark Univ.* **15**, 142–147 (2017), [arXiv:1603.05234 \[astro-ph.CO\]](https://arxiv.org/abs/1603.05234).
- [20] Misao Sasaki, Teruaki Suyama, Takahiro Tanaka, and Shuichiro Yokoyama, “Primordial Black Hole Scenario for the Gravitational-Wave Event GW150914,” *Phys. Rev. Lett.* **117**, 061101 (2016), [arXiv:1603.08338 \[astro-ph.CO\]](https://arxiv.org/abs/1603.08338).
- [21] V. De Luca, G. Franciolini, P. Pani, and A. Riotto, “Primordial Black Holes Confront LIGO/Virgo data: Current situation,” *JCAP* **06**, 044 (2020), [arXiv:2005.05641 \[astro-ph.CO\]](https://arxiv.org/abs/2005.05641).
- [22] Gert Hütsi, Martti Raidal, Ville Vaskonen, and Hardi Veermäe, “Two populations of LIGO-Virgo black holes,” *JCAP* **03**, 068 (2021), [arXiv:2012.02786 \[astro-ph.CO\]](https://arxiv.org/abs/2012.02786).
- [23] Hiroko Niikura, Masahiro Takada, Shuichiro Yokoyama, Takahiro Sumi, and Shogo Masaki, “Constraints on Earth-mass primordial black holes from OGLE 5-year microlensing events,” *Phys. Rev. D* **99**, 083503 (2019), [arXiv:1901.07120 \[astro-ph.CO\]](https://arxiv.org/abs/1901.07120).
- [24] G. Hasinger, “Illuminating the dark ages: Cosmic backgrounds from accretion onto primordial black hole dark

- matter,” *JCAP* **07**, 022 (2020), [arXiv:2003.05150 \[astro-ph.CO\]](#).
- [25] V. De Luca, G. Franciolini, and A. Riotto, “Nanograv data hints at primordial black holes as dark matter,” *Phys. Rev. Lett.* **126**, 041303 (2021).
- [26] Sergei G. Rubin, Alexander S. Sakharov, and Maxim Yu. Khlopov, “The Formation of primary galactic nuclei during phase transitions in the early universe,” *J. Exp. Theor. Phys.* **91**, 921–929 (2001), [arXiv:hep-ph/0106187](#).
- [27] M. Yu. Khlopov, S. G. Rubin, and A. S. Sakharov, “Strong primordial inhomogeneities and galaxy formation,” (2002), [arXiv:astro-ph/0202505](#).
- [28] Maxim Yu. Khlopov, Sergei G. Rubin, and Alexander S. Sakharov, “Primordial structure of massive black hole clusters,” *Astropart. Phys.* **23**, 265 (2005), [arXiv:astro-ph/0401532](#).
- [29] B.P. Abbott *et al.* (LIGO Scientific, Virgo), “Search for Subsolar-Mass Ultracompact Binaries in Advanced LIGO’s First Observing Run,” *Phys. Rev. Lett.* **121**, 231103 (2018), [arXiv:1808.04771 \[astro-ph.CO\]](#).
- [30] B.P. Abbott *et al.* (LIGO Scientific, Virgo), “Search for Subsolar Mass Ultracompact Binaries in Advanced LIGO’s Second Observing Run,” *Phys. Rev. Lett.* **123**, 161102 (2019), [arXiv:1904.08976 \[astro-ph.CO\]](#).
- [31] Khun Sang Phukon, Gregory Baltus, Sarah Caudill, Sebastien Clesse, Antoine Depasse, Maxime Fays, Heather Fong, Shasvath J. Kapadia, Ryan Magee, and Andres Jorge Tanasijczuk, “The hunt for sub-solar primordial black holes in low mass ratio binaries is open,” (2021), [arXiv:2105.11449 \[astro-ph.CO\]](#).
- [32] Alexander H. Nitz and Yi-Fan Wang, “Search for gravitational waves from the coalescence of sub-solar mass and eccentric compact binaries,” (2021), [arXiv:2102.00868 \[astro-ph.HE\]](#).
- [33] Alexander Harvey Nitz and Yi-Fan Wang, “Search for gravitational waves from high-mass-ratio compact-binary mergers of stellar mass and sub-solar mass black holes,” *Phys. Rev. Lett.* **126**, 021103 (2021), [arXiv:2007.03583 \[astro-ph.HE\]](#).
- [34] Takashi Nakamura, Misao Sasaki, Takahiro Tanaka, and Kip S. Thorne, “Gravitational waves from coalescing black hole macho binaries,” *Astrophys. J.* **487**, L139 (1997).
- [35] Hiroya Nishikawa, Ely D. Kovetz, Marc Kamionkowski, and Joseph Silk, “Primordial-black-hole mergers in dark-matter spikes,” *Phys. Rev. D* **99**, 043533 (2019).
- [36] Zu-Cheng Chen and Qing-Guo Huang, “Merger Rate Distribution of Primordial-Black-Hole Binaries,” *Astrophys. J.* **864**, 61 (2018), [arXiv:1801.10327 \[astro-ph.CO\]](#).
- [37] Yacine Ali-Haïmoud, Ely D. Kovetz, and Marc Kamionkowski, “Merger rate of primordial black-hole binaries,” *Phys. Rev. D* **96**, 123523 (2017), [arXiv:1709.06576 \[astro-ph.CO\]](#).
- [38] Martti Raidal, Christian Spethmann, Ville Vaskonen, and Hardi Veermäe, “Formation and Evolution of Primordial Black Hole Binaries in the Early Universe,” *JCAP* **02**, 018 (2019), [arXiv:1812.01930 \[astro-ph.CO\]](#).
- [39] Saeed Fakhry, Javad T. Firouzjaee, and Mehrdad Farhodi, “Primordial Black Hole Merger Rate in Ellipsoidal-Collapse Dark Matter Halo Models,” *Phys. Rev. D* **103**, 123014 (2021), [arXiv:2012.03211 \[astro-ph.CO\]](#).
- [40] Saeed Fakhry, Mahdi Naseri, Javad T. Firouzjaee, and Mehrdad Farhodi, “Primordial Black Hole Merger Rate in Self-Interacting Dark Matter Halo Models,” (2021), [arXiv:2106.06265 \[astro-ph.CO\]](#).
- [41] Yi-Fan Wang and Alexander H. Nitz, “Prospects for detecting gravitational waves from eccentric sub-solar mass compact binaries,” (2021), [arXiv:2101.12269 \[astro-ph.HE\]](#).
- [42] Samantha A. Usman *et al.*, “The PyCBC search for gravitational waves from compact binary coalescence,” *Class. Quant. Grav.* **33**, 215004 (2016), [arXiv:1508.02357 \[gr-qc\]](#).
- [43] Alexander H. Nitz, Ian W. Harry, Joshua L. Willis, Christopher M. Biwer, Duncan A. Brown, Larne P. Pekowsky, T. Dal Canton, Andrew R. Williamson, Thomas Dent, Collin D. Capano, Thomas J. Massinger, Amber K. Lenon, Alex B. Nielsen, and Miriam Cabero, “PyCBC Software,” <https://github.com/gwastro/pycbc> (2018).
- [44] Bruce Allen, Warren G. Anderson, Patrick R. Brady, Duncan A. Brown, and Jolien D. E. Creighton, “FIND-CHIRP: An Algorithm for detection of gravitational waves from inspiraling compact binaries,” *Phys. Rev. D* **85**, 122006 (2012), [arXiv:0509116 \[gr-qc\]](#).
- [45] Alexander H. Nitz, Thomas Dent, Tito Dal Canton, Stephen Fairhurst, and Duncan A. Brown, “Detecting binary compact-object mergers with gravitational waves: Understanding and Improving the sensitivity of the PyCBC search,” *Astrophys. J.* **849**, 118 (2017), [arXiv:1705.01513 \[gr-qc\]](#).
- [46] Alexander Harvey Nitz, “Distinguishing short duration noise transients in LIGO data to improve the PyCBC search for gravitational waves from high mass binary black hole mergers,” *Class. Quant. Grav.* **35**, 035016 (2018), [arXiv:1709.08974 \[gr-qc\]](#).
- [47] Bruce Allen, “A χ^2 time-frequency discriminator for gravitational wave detection,” *Phys. Rev. D* **71**, 062001 (2005), [arXiv:gr-qc/0405045 \[gr-qc\]](#).
- [48] Gareth S. Davies, Thomas Dent, Márton Tápai, Ian Harry, Connor Mclsaac, and Alexander H. Nitz, “Extending the pycbc search for gravitational waves from compact binary mergers to a global network,” *Phys. Rev. D* **102**, 022004 (2020), [arXiv:2002.08291 \[astro-ph.HE\]](#).
- [49] S. Mozzon, L. K. Nuttall, A. Lundgren, T. Dent, S. Kumar, and A. H. Nitz, “Dynamic Normalization for Compact Binary Coalescence Searches in Non-Stationary Noise,” *Class. Quant. Grav.* **37**, 215014 (2020), [arXiv:2002.09407 \[astro-ph.IM\]](#).
- [50] Michal Was, Marie-Anne Bizouard, Violette Brisson, Fabien Cavalier, Michel Davier, *et al.*, “On the background estimation by time slides in a network of gravitational wave detectors,” *Class. Quantum Grav.* **27**, 015005 (2010), [arXiv:0906.2120 \[gr-qc\]](#).
- [51] C. Capano, T. Dent, C. Hanna, M. Hendry, C. Messenger, Y.-M. Hu, and J. Veitch, “Systematic errors in estimation of gravitational-wave candidate significance,” *Phys. Rev. D* **96**, 082002 (2017), [arXiv:1708.06710 \[astro-ph.IM\]](#).
- [52] B. P. Abbott *et al.* (Virgo, LIGO Scientific), “Characterization of transient noise in Advanced LIGO relevant to gravitational wave signal GW150914,” *Class. Quant. Grav.* **33**, 134001 (2016), [arXiv:1602.03844 \[gr-qc\]](#).
- [53] B. S. Sathyaprakash and S. V. Dhurandhar, “Choice of filters for the detection of gravitational waves from coalescing binaries,” *Phys. Rev. D* **44**, 3819–3834 (1991).

- [54] Serge Droz, Daniel J. Knapp, Eric Poisson, and Benjamin J. Owen, “Gravitational waves from inspiraling compact binaries: Validity of the stationary-phase approximation to the fourier transform,” *Phys. Rev. D* **59**, 124016 (1999).
- [55] Luc Blanchet, “Gravitational radiation from post-Newtonian sources and inspiralling compact binaries,” *Living Rev. Rel.* **5**, 3 (2002), [arXiv:gr-qc/0202016](#).
- [56] Guillaume Faye, Sylvain Marsat, Luc Blanchet, and Bala R. Iyer, “The third and a half post-Newtonian gravitational wave quadrupole mode for quasi-circular inspiralling compact binaries,” *Class. Quant. Grav.* **29**, 175004 (2012), [arXiv:1204.1043 \[gr-qc\]](#).
- [57] Blake Moore and Nicolás Yunes, “A 3PN Fourier domain waveform for non-spinning binaries with moderate eccentricity,” *Classical and Quantum Gravity* **36**, 185003 (2019), [arXiv:1903.05203 \[gr-qc\]](#).
- [58] Blake Moore and Nicolás Yunes, “Data Analysis Implications of Moderately Eccentric Gravitational Waves,” *arXiv e-prints*, [arXiv:1910.01680](#) (2019), [arXiv:1910.01680 \[gr-qc\]](#).
- [59] Blake Moore, Travis Robson, Nicholas Loutrel, and Nicolás Yunes, “Towards a Fourier domain waveform for non-spinning binaries with arbitrary eccentricity,” *Classical and Quantum Gravity* **35**, 235006 (2018), [arXiv:1807.07163 \[gr-qc\]](#).
- [60] Takeshi Chiba and Shuichiro Yokoyama, “Spin Distribution of Primordial Black Holes,” *PTEP* **2017**, 083E01 (2017), [arXiv:1704.06573 \[gr-qc\]](#).
- [61] V. De Luca, V. Desjacques, G. Franciolini, A. Malhotra, and A. Riotto, “The initial spin probability distribution of primordial black holes,” *JCAP* **05**, 018 (2019), [arXiv:1903.01179 \[astro-ph.CO\]](#).
- [62] V. De Luca, G. Franciolini, P. Pani, and A. Riotto, “The Evolution of Primordial Black Holes and their Final Observable Spins,” *JCAP* **04**, 052 (2020), [arXiv:2003.02778 \[astro-ph.CO\]](#).
- [63] Mehrdad Mirbabayi, Andrei Gruzinov, and Jorge Noreña, “Spin of Primordial Black Holes,” *JCAP* **03**, 017 (2020), [arXiv:1901.05963 \[astro-ph.CO\]](#).
- [64] K. Postnov, A. Kuranov, and N. Mitichkin, “Spins of black holes in coalescing compact binaries,” *Phys. Usp.* **62**, 1153–1161 (2019), [arXiv:1907.04218 \[astro-ph.HE\]](#).
- [65] Michele Vallisneri, Jonah Kanner, Roy Williams, Alan Weinstein, and Branson Stephens, “The LIGO Open Science Center,” *Proceedings, 10th International LISA Symposium: Gainesville, Florida, USA, May 18-23, 2014*, *J. Phys. Conf. Ser.* **610**, 012021 (2015), [arXiv:1410.4839 \[gr-qc\]](#).
- [66] Rich Abbott *et al.* (LIGO Scientific, Virgo), “Open data from the first and second observing runs of Advanced LIGO and Advanced Virgo,” *SoftwareX* **13**, 100658 (2021), [arXiv:1912.11716 \[gr-qc\]](#).
- [67] Rahul Biswas, Patrick R. Brady, Jolien D. E. Creighton, and Stephen Fairhurst, “The Loudest Event Statistic: General Formulation, Properties and Applications,” *Class. Quantum Grav.* **26**, 175009 (2009), [arXiv:arXiv:0710.0465 \[gr-qc\]](#).
- [68] Ville Vaskonen and Hardi Veermäe, “Lower bound on the primordial black hole merger rate,” *Phys. Rev. D* **101**, 043015 (2020), [arXiv:1908.09752 \[astro-ph.CO\]](#).