

YET ANOTHER WAY OF CALCULATING MOMENTS OF THE KESTEN'S DISTRIBUTION AND ITS CONSEQUENCES

PAWEŁ J. SZABŁOWSKI

ABSTRACT. We calculate moments of the so-called Kesten distribution by means of the expansion of the nominator of the density of this distribution and then integrating all summands with respect to the semicircle distribution. By comparing so obtained expression with the formulae for the moments of Kesten's distribution obtained by other means, we obtain identities involving polynomials whose power coefficients are closely related to Catalan numbers, Catalan triangles binomial coefficients.

1. INTRODUCTION

The purpose of this note is to calculate a sequence of moments of the Kesten's distribution and thus by comparison with the existing formulae to obtain some polynomial type identities involving Catalan and related to them some sequences of numbers (see Proposition 1). In 2015 in [5] and in 2020 in [6] Szabłowski calculated in two different ways the moments of Kesten distribution. Later T. Hasegawa and S. Saito in [1] calculated these moments in some other ways and equating the results obtained interesting identities involving Catalan and related numbers. So in this note, we will calculate these moments in yet another way and obtain some other identities.

2. BASIC INGREDIENTS

We start with the modified semicircle distribution i.e. distribution with the density

$$f_S(x) = \begin{cases} \frac{1}{2\pi}\sqrt{4-x^2} & \text{if } |x| \leq 2 \\ 0 & \text{if } |x| > 2 \end{cases}.$$

It is well known that the moment sequence of f_S are the famous Catalan numbers. More precisely, we have:

$$\int_{-2}^2 x^{2n} f_S(x) dx = C_n = \frac{1}{n+1} \binom{2n}{n}.$$

We know also that the moment generating function see .e.g. [2] (4.8 p. 107) of this distribution is after inserting proper values of parameters equal to

$$g_S(z) = \frac{2}{\sqrt{1-4z^2+1}},$$

Date: June, 2021.

2020 Mathematics Subject Classification. Primary 05A19, Secondary 60E10, 62E10.

Key words and phrases. sequence of moments, Kesten distribution, Catalan numbers.

for $|z| \leq 1/2$. One can easily notice also that

$$\sum_{k \geq 0} t^k C_k = \frac{2}{\sqrt{1-4t}+1},$$

for $|t| \leq 1/4$.

The other ingredient is the definition of the Kesten distribution. It was considered in many papers including [2], [5], [6], and recently in [1] with different parametrization. The following distribution chooses one of these parametrizations. Let us consider distribution with the following density

$$(2.1) \quad f_K(x|p, r) = \begin{cases} 0 & \text{if } |x| > 2/\sqrt{r} \\ \frac{p}{2\pi} \frac{\sqrt{4r-x^2}}{p^2-(p-r)x^2} & \text{if } |x| \leq 2/\sqrt{r} \end{cases},$$

for $0 \leq p \leq 2r$.

Now notice that, if $p = r$, then

$$f_K(x|r, r) = \frac{1}{\sqrt{r}} f_S(x/\sqrt{r}).$$

Thus, we have

$$\int_{-2}^2 x^{2n} f_K(x|r, r) dx = r^n C_n = \frac{r^n}{n+1} \binom{2n}{n}.$$

Now notice that for

$$(2.2) \quad \left| \frac{p-r}{p^2} \right| x^2 \leq \left| \frac{p-r}{p^2} \right| 4r < 1,$$

we have the following expansion:

$$f_K(x|p, r) = \frac{r}{p} \frac{\sqrt{4r-x^2}}{2\pi r} \sum_{k \geq 0} \left(\frac{p-r}{p^2} \right)^k x^{2k}.$$

Notice, that for $|x| \leq 2\sqrt{r}$ and parameters satisfying 2.2 the convergence of the series above is uniform and one can integrate part by part. Then, we will get :

$$(2.3) \quad M_{2m}(p, r) = \int_{-2/\sqrt{r}}^{2/\sqrt{r}} x^{2m} f_K(x|p, r) dx = \frac{r}{p} \sum_{k \geq 0} \left(\frac{p-r}{p^2} \right)^k r^{k+m} C_{k+m}.$$

Let us introduce a new auxiliary variable

$$(2.4) \quad t = \frac{r}{p}.$$

General conditions on p and r require that $t \geq 1/2$. Recall that we must also have $|t(1-t)| < 1/4$, which leads to the following condition

$$(2.5) \quad 1/2 \leq t < (1 + \sqrt{2})/2$$

Summarizing, we get the following result.

Theorem 1. *Let Kesten distribution be defined by the density f_K given by (2.1). Let parameters p and r satisfy the following relationship*

$$2r \geq p \geq 2(\sqrt{2}-1)r,$$

then we have:

$$(2.6) \quad M_{2m}(p, r) = p^m \frac{t}{(1-t)^m} \left(\frac{1}{t} - \sum_{k=0}^{m-1} t^k (1-t)^k C_k \right),$$

with z given by (2.4), with an obvious condition $|z| < 1/4$ and $z \neq 0$. For $z = 0$ we obviously have $M_{2m}(r, r) = r^m C_m$.

Proof. First let us analyze when the series (2.3) is convergent. Namely it is absolutely convergent if

$$\left| \frac{r}{p} \left(1 - \frac{r}{p} \right) \right| < \frac{1}{4},$$

that is when

$$-\frac{1}{4} < \frac{r}{p} - \left(\frac{r}{p} \right)^2 < \frac{1}{4}.$$

The inequality $\frac{r}{p} - \left(\frac{r}{p} \right)^2 < \frac{1}{4}$ is equivalent to the following $\left(\frac{1}{2} - \frac{r}{p} \right)^2 > 0$ which is true for all $2r > p$. The second one leads to inequality

$$\left(\frac{r}{p} \right)^2 - \frac{r}{p} - \frac{1}{4} < 0,$$

which leads to $t < \frac{1+\sqrt{2}}{2}$ or equivalently $p > 2(\sqrt{2} - 1)r$ as shown. Since now on let us work with the new variable t introduced by (2.4)

Now, we have

$$\sum_{n \geq 0} t^n (1-t)^n C_n = \frac{1}{t},$$

Analyzing (2.3) we get, for $|z| < 1/4$ and $z \neq 0$.

$$\begin{aligned} M_{2m}(p, r) &= \frac{r}{p} \frac{p^{2m}}{(p-r)^m} \sum_{k \geq 0} t^{k+m} (1-t)^{k+m} C_{k+m} \\ &= \frac{r}{p} \frac{p^m}{\left(1 - \frac{r}{p}\right)^m} \left(\sum_{k \geq 0} t^k (1-t)^k C_k - \sum_{k \neq 0}^{m-1} t^k (1-t)^k C_k \right) \\ &= p^m \frac{t}{(1-t)^m} \left(\frac{1}{t} - \sum_{k \neq 0}^{m-1} t^k (1-t)^k C_k \right). \end{aligned}$$

□

Let us now compare this result with known formulae for the moments of Kesten distribution.

In [5] the following result has been obtained, after necessary change of parameters, $\rho^2 = 1 - p/r$, $y = 0$, $q = 0$ and x replaced by x/\sqrt{r} :

$$M_{2m}(p, r) = r^m \sum_{k=0}^m \left(1 - \frac{p}{r} \right)^{m-k} S_{m,k},$$

where $S_{m,k} = \binom{2m}{k} - \binom{2m}{k-1}$, with understanding that $\binom{2m}{-1} = 0$. Hence in terms of our parameter t we have:

$$\begin{aligned} M_{2m}(p, r) &= p^m \sum_{k=0}^m \left(-\frac{r}{p} + 1 \right)^{m-k} \left(\frac{r}{p} \right)^k S_{m,k} \\ (2.7) \quad &= p^m \sum_{k=0}^m t^k (1-t)^{m-k} S_{m,k}. \end{aligned}$$

In [1] two other expressions for the moments of Kesten distribution can be found. Namely the following formulae:

$$(2.8) \quad M_{2m}(p, r) = p \sum_{j=0}^{m-1} p^{k-1-j} r^j T_{k-1,j},$$

$$(2.9) \quad M_{2m}(p, r) = p \sum_{j=0}^{m-1} (p-r)^j r^{k-1-j} B_{k,j+1},$$

where numbers $T_{k,j}$ and $B_{k,j}$ are called, depending on the author, Catalan triangles. The the first one is given by

$$T_{k,j} = \frac{k-j+1}{k+1} \binom{k+i}{k}$$

and in Sloane's Encyclopedia [3] is denoted as <https://oeis.org/A009766> and the second was introduced by L.W. Shapiro in [4] and which can be found in Sloane's Encyclopedia as <https://oeis.org/A039598> and is given by:

$$B_{k,j} = \frac{j}{k} \binom{2k}{k-j},$$

$k \geq j \geq 1$. Hence, we have the following result:

Proposition 1. *For all $m \geq 1$ and $t \in \mathbb{C}$ we have:*

$$(2.10) \quad (1-t)^m \sum_{k=0}^m t^k (1-t)^{m-k} S_{m,k} = 1 - \sum_{k=0}^{m-1} t^{k+1} (1-t)^k C_k =$$

$$(2.11) \quad (1-t)^m \sum_{j=0}^{m-1} t^j T_{k-1,j} = (1-t)^m \sum_{j=0}^{m-1} (1-t)^j t^{m-1-j} B_{k,j+1}.$$

Proof. Firstly, for t satisfying (2.5) we equate (2.6) with (2.7) and divide both sides by p^m and multiply also both sides by $(1-t)^m$, we obtain (2.10). Secondly, we multiply and divide both (2.8) and (2.9) by p^{m-1} enabling introduction of parameter t . Then we divide obtained expressions by p^m equate obtained expressions with (2.6) and then multiply both sides by $(1-t)^m$. This way we obtain 3 equations involving polynomials. But when dealing with polynomials we can extend the domain from any segment, to all complex numbers. \square

Now we can use the above-mentioned identities in different ways for different purposes concerning relationships between sets of Catalan numbers and Catalan triangles and binomial coefficients.

In doing so the following new parameterization might help. Namely, if we set $z = t(1-t)$, then $t = (1 + \sqrt{1-4z})/2$ and $1-t = (1 - \sqrt{1-4z})/2$. Thus for example if $z = 1$ then $t = (1 + i\sqrt{3})/2$ and $1-t = (1 - i\sqrt{3})/2$, or $z = -1$ then $t = (1 + \sqrt{5})/2$ and $1-t = (1 - \sqrt{5})/2$. Finally if $z = -2$, then $t = 2$ and $1-t = -1$. Thus we can express sums or alternating sums of Catalan numbers with the help of binomial coefficient or numbers forming Catalan triangles.

REFERENCES

- [1] Hasegawa, T and Saito, S. A note on the moments of the Kesten distribution, *Discrete Math.* (in print)

- [2] Kesten, Harry. Symmetric random walks on groups. *Trans. Amer. Math. Soc.* **92** (1959), 336–354. MR0109367
- [3] Sloane, N., J., A. The on-line Encyclopedia of Integer Sequences, <https://oeis.org>.
- [4] Shapiro, L. W. A Catalan triangle. *Discrete Math.* **14** (1976), no. 1, 83–90. MR0387069
- [5] Szablowski, Paweł, J., Moments of q -Normal and conditional q -Normal distributions, *Stoch. and Probab. Letters*, **106** (2015), 65-72
- [6] Szablowski, Paweł J. On the generalized Kesten–McKay distributions. *ESAIM Probab. Stat.* **24** (2020), 56–68. MR4069296, ArXiv:<http://arxiv.org/abs/1507.03191>

EMERITUS IN DEPARTMENT OF MATHEMATICS AND INFORMATION SCIENCES,, WARSAW UNIVERSITY OF TECHNOLOGY, UL KOSZYKOWA 75, 00-662 WARSAW, POLAND