

THE SPECTRUM OF THE EXPONENTS OF REPETITION

DEOKWON SIM

ABSTRACT. For an infinite word \mathbf{x} , Bugeaud and Kim introduced a new complexity function $\text{rep}(\mathbf{x})$ which is called the exponent of repetition of \mathbf{x} . They showed $1 \leq \text{rep}(\mathbf{x}) \leq \sqrt{10} - \frac{3}{2}$ for any Sturmian word \mathbf{x} . Ohnaka and Watanabe found a gap in the set of the exponents of repetition of Sturmian words. For an irrational number $\theta \in (0, 1)$, let

$$\mathcal{L}(\theta) := \{\text{rep}(\mathbf{x}) : \mathbf{x} \text{ is an Sturmian word of slope } \theta\}.$$

In this article, we look into $\mathcal{L}(\theta)$. The minimum of $\mathcal{L}(\theta)$ is determined where θ has bounded partial quotients in its continued fraction expression. In particular, we find out the maximum and the minimum of $\mathcal{L}(\varphi)$ where $\varphi := \frac{\sqrt{5}-1}{2}$ is the fraction part of the golden ratio. Furthermore, we show that the three largest values are isolated points in $\mathcal{L}(\varphi)$ and the fourth largest point is a limit point of $\mathcal{L}(\varphi)$.

1. INTRODUCTION

A word \mathbf{x} over \mathcal{A} is a finite or infinite sequence of elements of a finite set \mathcal{A} . An element of \mathcal{A} is called a *letter*. For each integer $n \geq 1$, $p(n, \mathbf{x})$ is defined by the number of distinct subwords of length n appearing in the word \mathbf{x} and is called by the *subword complexity* of \mathbf{x} . Morse and Hedlund showed that an infinite word is eventually periodic if and only if its subword complexity is bounded [16]. Thus, a non-eventually periodic word \mathbf{x} with the smallest subword complexity satisfies $p(n, \mathbf{x}) = n + 1$ for all $n \geq 1$.

Definition 1.1. A *Sturmian word* is an infinite word \mathbf{x} over $\mathcal{A} = \{0, 1\}$ satisfying $p(n, \mathbf{x}) = n + 1$ for all $n \geq 1$.

Let us recall a classical result on Sturmian words [15, Chapter 2]. For a Sturmian word $\mathbf{x} = x_1x_2\dots$, there exist an irrational number $\theta \in (0, 1)$ and a real number ρ such that either

$$x_n = \lfloor (n+1)\theta + \rho \rfloor - \lfloor n\theta + \rho \rfloor \text{ for all } n \geq 1$$

or

$$x_n = \lceil (n+1)\theta + \rho \rceil - \lceil n\theta + \rho \rceil \text{ for all } n \geq 1.$$

Conversely, for an irrational number $\theta \in (0, 1)$ and a real number ρ , the infinite words defined by

$$\mathbf{s}_{\theta, \rho} := s_1s_2\dots, \mathbf{s}'_{\theta, \rho} := s'_1s'_2\dots$$

are Sturmian where $s_n = \lfloor (n+1)\theta + \rho \rfloor - \lfloor n\theta + \rho \rfloor$ and $s'_n = \lceil (n+1)\theta + \rho \rceil - \lceil n\theta + \rho \rceil$ for $n \geq 1$. They are called the Sturmian words *of slope θ and intercept ρ* .

Sturmian words have been studied in many different areas [4, 9, 14, 15]. Various complexities have been looked into characterize Sturmian words such as Cassaigne's recurrence function, rectangle complexity [5, 11]. Bugeaud and Kim recently suggested a new complexity function $r(n, \mathbf{x})$ and characterized Sturmian words and eventually periodic words in terms of $r(n, \mathbf{x})$ [8].

Definition 1.2. Given an infinite word $\mathbf{x} = x_1x_2\dots$, let $r(n, \mathbf{x})$ denote the length of the smallest prefix in which some subword of length n occurs twice. More precisely,

$$r(n, \mathbf{x}) := \min\{m : x_jx_{j+1}\dots x_{j+n-1} = x_{m-n+1}x_{m-n+2}\dots x_m \text{ for some } 1 \leq j \leq m - n\}.$$

The *exponent of repetition* of \mathbf{x} is defined by

$$\text{rep}(\mathbf{x}) := \liminf_{n \rightarrow \infty} \frac{r(n, \mathbf{x})}{n}.$$

By definition, $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for $n \geq 1$ and $\text{rep}(c\mathbf{x}) = \text{rep}(\mathbf{x})$ for any finite word c . Theorem 2.3 and 2.4 in [8] says that for an infinite word \mathbf{x} , the following statements hold:

- (i) \mathbf{x} is eventually periodic if and only if $r(n, \mathbf{x}) \leq 2n$ for all sufficiently large integers n .
- (ii) \mathbf{x} is a Sturmian word if and only if $r(n, \mathbf{x}) \leq 2n + 1$ for all $n \geq 1$ and equality holds for infinitely many n .

A number of spectrums such as Lagrange spectrum and Markoff spectrum have been studied for a long time. Spectrums connected with Diophantine approximation are one of main topics in Number theory, Combinatorics, Dynamics [3, 12, 13]. In particular, the set of the irrationality exponents appears in many references [2, 7]. Theorem 4.5 in [8] says that for a Sturmian word $\mathbf{x} = x_1x_2\dots$, an integer $b \geq 2$, and a Sturmian number $r_{\mathbf{x}} := \sum_{k \geq 1} \frac{x_k}{b^k}$, the exponent of repetition $\text{rep}(\mathbf{x})$ and the irrationality exponent of $r_{\mathbf{x}}$, denoted by $\sup\{\mu : |r_{\mathbf{x}} - \frac{p}{q}| < \frac{1}{q^\mu}$ for infinitely many $p, q\}$, have a special relation: $\mu(r_{\mathbf{x}}) = \frac{\text{rep}(\mathbf{x})}{\text{rep}(\mathbf{x})-1}$ where $\mu(r_{\mathbf{x}})$ is the irrationality exponent of $r_{\mathbf{x}}$. Thus, we look into the spectrum of the exponents of repetition of Sturmian words. From now on, we call $r_{\mathbf{x}}$ a *Sturmian number of slope θ* if \mathbf{x} is a Sturmian word of slope θ .

Let us set up some notations on continued fraction expansion. Any irrational number θ has the unique continued fraction expansion of the form

$$[a_0; a_1, a_2, \dots] := a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots}}$$

where $a_0 \in \mathbb{Z}$ and $a_n \in \mathbb{Z}_{>0}$ for $n \geq 1$. Each a_n is called a *partial quotient* of θ . We say that θ has *bounded partial quotients* if all of partial quotients of θ are bounded. Let $k \geq 0$. We call $[a_0; a_1, a_2, \dots, a_k]$ the *kth principal convergent* of θ . Denote $\frac{p_k}{q_k} = [a_0; a_1, a_2, \dots, a_k]$. Let $p_{-1} = 1, q_{-1} = 0$. It is known that $p_{k+1} = a_{k+1}p_k + p_{k-1}$ and $q_{k+1} = a_{k+1}q_k + q_{k-1}$ for all $k \geq 0$.

Let us denote by \mathbf{St} the set of Sturmian words over $\mathcal{A} = \{0, 1\}$ and by $\text{rep}(\mathbf{St})$ the set of $\text{rep}(\mathbf{x})$ for all $\mathbf{x} \in \mathbf{St}$. In [8], Bugeaud and Kim found the maximum of $\text{rep}(\mathbf{St})$ and a necessary and sufficient condition for the maximum of $\text{rep}(\mathbf{St})$. They also gave an alternative proof of [1, Proposition 11.1], which is a sufficient condition for the minimum of $\text{rep}(\mathbf{St})$.

Theorem 1.1. ([8], Theorem 3.3 and 3.4) For any Sturmian word \mathbf{x} ,

$$1 \leq \text{rep}(\mathbf{x}) \leq \sqrt{10} - \frac{3}{2}$$

Moreover, a Sturmian word \mathbf{x} satisfies $\text{rep}(\mathbf{x}) = r_{\max} := \sqrt{10} - \frac{3}{2}$ if and only if its slope is of the form $[0; a_1, a_2, \dots, a_K, \overline{2, 1, 1}]$ for some K . A Sturmian word \mathbf{x} satisfies $\text{rep}(\mathbf{x}) = 1$ if its slope has unbounded partial quotients.

On the other hand, Ohnaka and Watanabe recently discovered the second largest value r_1 of $\text{rep}(\mathbf{St})$ and proved that r_1 is a limit point of $\text{rep}(\mathbf{St})$.

Theorem 1.2. ([17], Theorem 0.4) For $r_1 = \frac{48+\sqrt{10}}{31} = 1.650\dots$,

$$(r_1, r_{\max}) \cap \text{rep}(\mathbf{St}) = \emptyset$$

Moreover, r_1 is a limit point of $\text{rep}(\mathbf{St})$.

For any irrational number $\theta \in (0, 1)$, let $\mathcal{L}(\theta)$ be the set of the exponents of repetition of Sturmian words of slope θ , i.e.

$$\mathcal{L}(\theta) = \{\text{rep}(\mathbf{x}) : \mathbf{x} \text{ is a Sturmian word of slope } \theta\}.$$

In this article, we mainly study $\mathcal{L}(\theta)$. Theorem 3.3 in [8] gives $\text{rep}(\mathbf{x}) = 1$ where the slope of \mathbf{x} has unbounded partial quotients. We find the minimum of $\mathcal{L}(\theta)$ where θ has bounded partial quotients.

Theorem 1.3. Let $\theta = [0; a_1, a_2, \dots]$ have bounded partial quotients. We have

$$\min \mathcal{L}(\theta) = \liminf_{k \rightarrow \infty} [1; 1 + a_k, a_{k-1}, a_{k-2}, \dots, a_1].$$

In particular, for $\varphi := [0; \overline{1}]$, we find the minimum μ_{\min} and the maximum μ_{\max} of $\mathcal{L}(\varphi)$. In addition, we determine Sturmian words \mathbf{x} satisfying $\text{rep}(\mathbf{x}) = \mu_{\min}$ or μ_{\max} in terms of the locating chain of \mathbf{x} , to be defined in Section 2 and 4.

Theorem 1.4. Let \mathbf{x} be a Sturmian word of slope φ . Then, $\mu_{\min} \leq \text{rep}(\mathbf{x}) \leq \mu_{\max}$. Moreover, the locating chain of \mathbf{x} is $u\bar{a}$ or $v\bar{b}$ for some finite words u, v if and only if $\text{rep}(\mathbf{x}) = \mu_{\max}$. We have $\mathbf{x} \in S_d$ for any $d \geq 1$ if and only if $\text{rep}(\mathbf{x}) = \mu_{\min}$.

We call an interval (c, d) a *maximal gap* in $\mathcal{L}(\varphi)$ if $(c, d) \cap \mathcal{L}(\varphi) = \emptyset$ and there exist Sturmian words \mathbf{x}, \mathbf{y} such that $\text{rep}(\mathbf{x}) = c$, $\text{rep}(\mathbf{y}) = d$. We show that the three largest points μ_{\max}, μ_2, μ_3 are isolated points in $\mathcal{L}(\varphi)$ and the fourth largest point μ_4 is an accumulation point of $\mathcal{L}(\varphi)$. Moreover, we determine Sturmian words \mathbf{x} satisfying $\text{rep}(\mathbf{x}) = \mu_2, \mu_3$, or μ_4 in terms of the locating chain of \mathbf{x}

Theorem 1.5. The intervals $(\mu_2, \mu_{\max}), (\mu_3, \mu_2)$ are maximal gaps in $\mathcal{L}(\varphi)$. Moreover, the locating chain of \mathbf{x} is $u\bar{a}\bar{b}$ for some finite word u if and only if $\text{rep}(\mathbf{x}) = \mu_2$. The locating chain of \mathbf{x} is $v\bar{b}^2\bar{a}^2$ for some finite word v if and only if $\text{rep}(\mathbf{x}) = \mu_3$.

Theorem 1.6. The interval (μ_4, μ_3) is a maximal gap in $\mathcal{L}(\varphi)$. Moreover, $\text{rep}(\mathbf{x}) = \mu_4$ if and only if $\mathbf{x} \in S_2 \cap S_3^c$ satisfies the following two conditions:

- 1) The locating chain of \mathbf{x} is $u(b^2a^2)^{e_1}ba(b^2a^2)^{e_2}ba\dots$ for some finite word u and integers $e_i \geq 1$.
- 2) $\limsup_{i \geq 1} \{e_i\} = \infty$.

Furthermore, μ_4 is a limit point of $\mathcal{L}(\varphi)$.

Indeed,

$$\mu_{max} = 1 + \varphi = 1.618\dots, \quad \mu_2 = 1 + 2\varphi^3 = 1.472\dots, \quad \mu_3 = 1 + \frac{\varphi^2(\varphi^4 + \varphi^2 + 1)}{\varphi^5 + \varphi^3 + 1} = 1.440\dots,$$

$$\mu_4 = 1 + \frac{1 - \varphi^6}{1 + 2\varphi - 2\varphi^7 + \varphi^{11}} = 1.435\dots, \quad \mu_{min} = 1 + \varphi^2 = 1.382\dots$$



FIGURE 1. $\mu_{max}, \mu_2, \mu_3, \mu_4, \mu_{min}$ in $\mathcal{L}(\varphi)$

For $\alpha \in \{\mu_{max}, \mu_2, \mu_3, \mu_4, \mu_{min}\}$, we obtain the cardinality of the set of Sturmian words \mathbf{x} satisfying $\text{rep}(\mathbf{x}) = \alpha$.

Proposition 1.6.1. For $\alpha \in \{\mu_{max}, \mu_2, \mu_3\}$, there are only countably many Sturmian words \mathbf{x} of slope φ satisfying $\text{rep}(\mathbf{x}) = \alpha$. For $\beta \in \{\mu_4, \mu_{min}\}$, there are uncountably many Sturmian words \mathbf{x} of slope φ satisfying $\text{rep}(\mathbf{x}) = \beta$.

The article is organized as follows. In Section 2, we define basic notations in Sturmian words and the locating chain of a Sturmian word. In Section 3, we investigate the set $\{n : r(n, \mathbf{x}) = 2n + 1\}$ for a Sturmian word \mathbf{x} . We prove Theorem 1.3 in Theorem 3.3. In Section 4, we study $\text{rep}(\mathbf{x})$ for a Sturmian word \mathbf{x} of slope φ . We establish Theorem 1.4, 1.5, and 1.6 in Theorem 4.4, 4.5, and 4.6, respectively. We prove Proposition 1.6.1 in Proposition 4.6.1.

At the final stage of writing the paper, the author found an independent result by Bugeaud and Laurent [10]: Theorem 1.3 is equivalent to Theorem 2.6 in [10]. For a given irrational number $\theta \in (0, 1)$, Bugeaud and Laurent found the maximum of the irrationality exponents of Sturmian numbers of slope θ . Using Theorem 4.5 in [8] and Section 4 in [6], Theorem 1.3 is induced.

2. BASIC DEFINITIONS AND EXAMPLES

Let us begin by some notations in Sturmian words of slope θ where $\theta \in (0, 1)$ is an irrational number. Let $\theta := [0; a_1, a_2, \dots]$. We define a sequence $\{M_k\}_{k \geq 0}$ in the following way: Let us define $M_0 := 0$, $M_1 := 0^{a_1-1}1$, and $M_{k+1} := M_k^{a_{k+1}}M_{k-1}$ for $k \geq 1$. Let us denote the numerator and denominator of the k th principal convergent of θ by p_k, q_k , respectively. We denote the length of M_k by $|M_k|$. Since $|M_1| = a_1 = q_1$, $|M_0| = 1 = q_0$, and $|M_{k+1}| = a_{k+1}|M_k| + |M_{k-1}|$

for $k \geq 1$, it is obvious that $|M_k| = q_k$. The *characteristic Sturmian word* of slope θ is obtained by $\mathbf{c}_\theta := \lim_{k \rightarrow \infty} M_k$.

For a non-empty finite word V , let us denote by V^- the word V with the last letter removed. Let $k \geq 1$. Note that $M_k M_{k-1}$ and $M_{k-1} M_k$ are identical, except for the last two letters [15, Proposition 2.2.2]. Let $\widetilde{M}_k := M_k M_{k-1}^- = M_{k-1} M_k^-$. The last two letters of $M_k M_{k-1}$ is 01 (resp., 10) if and only if the last two letters of $M_{k-1} M_k$ is 10 (resp., 01). We denote by D_k, D'_k the last two letters such that $M_k M_{k-1} = \widetilde{M}_k D_k, M_{k-1} M_k = \widetilde{M}_k D'_k$, respectively.

In this section, let $\mathbf{x} = x_1 x_2 \dots$ be a Sturmian word of slope θ . By Lemma 7.2 in [8], for any $k \geq 1$, there exists a unique word W_k satisfying one of the following cases

- (i) $\mathbf{x} = W_k M_k \widetilde{M}_k \dots$, where W_k is a non-empty suffix of M_k ,
- (ii) $\mathbf{x} = W_k M_{k-1} M_k \widetilde{M}_k \dots$, where W_k is a non-empty suffix of M_k ,
- (iii) $\mathbf{x} = W_k M_k \widetilde{M}_k \dots$, where W_k is a non-empty suffix of M_{k-1} .

For case (i) and case (ii), there exist q_k non-empty suffices of M_k . For case (iii), there exist q_{k-1} non-empty suffices of M_{k-1} . Lemma 7.2 in [8] also gives that all the $(2q_k + q_{k-1})$ cases are mutually exclusive. For each $k \geq 1$, we say that \mathbf{x} *belongs to case (i), (ii), (iii) at level k* if W_k satisfies case (i), (ii), (iii), respectively. We denote by $\mathcal{C}_k^{(i)}, \mathcal{C}_k^{(ii)}, \mathcal{C}_k^{(iii)}$ the set of Sturmian words which belong to case (i), (ii), (iii) at level k , respectively. For each \mathbf{x} , we have an infinite sequence of (i), (ii) and (iii) for which case \mathbf{x} belongs to at level $1, 2, \dots$, called the *locating chain* of \mathbf{x} . In the locating chain of \mathbf{x} , let $u^d := \underbrace{uu \dots u}_d$ where u is a finite word of (i),(ii),(iii).

Example 2.1. Let $\mathbf{x} = \mathbf{c}_\varphi$, i.e. the characteristic Sturmian word of slope $\varphi = [0; \overline{1}]$. Since \mathbf{x} starts with $M_{k+1} M_k M_{k+1} = M_k M_{k-1} M_k \widetilde{M}_k D_k$ for any $k \geq 1$, $\mathbf{x} \in \mathcal{C}_k^{(ii)}$ and $W_k = M_k$ for all $k \geq 1$. Hence, the locating chain of \mathbf{x} is $\overline{(ii)}$.

Example 2.2. Let $\mathbf{x} = 1\mathbf{c}_\varphi$. Since \mathbf{x} starts with $1M_{k+2} = 1M_k M_{k-1} M_k = 1M_k \widetilde{M}_k D'_k$ for any $k \geq 1$, $W_k = 1$ for $k \geq 1$. Moreover, $\mathbf{x} \in \mathcal{C}_k^{(i)}$ if k is odd, and $\mathbf{x} \in \mathcal{C}_k^{(iii)}$ if k is even. Hence, the locating chain of \mathbf{x} is $\overline{(i)(iii)}$.

Example 2.3. Let $\mathbf{x} = 10101M_4 M_5 \dots$. Since \mathbf{x} starts with $W_1 M_0 M_1 = 101$, $\mathbf{x} \in \mathcal{C}_1^{(ii)}$ and $W_2 = W_1 M_0 = 10$. Since \mathbf{x} starts with $W_2 M_2 \widetilde{M}_2 = 10101$, $\mathbf{x} \in \mathcal{C}_2^{(i)}$ and $W_3 = W_2 = 10$. Since \mathbf{x} starts with $W_3 M_3 M_4 = W_3 M_3 \widetilde{M}_3 D_3$, $\mathbf{x} \in \mathcal{C}_3^{(iii)}$ and $W_4 = W_3 = 10$. Since \mathbf{x} starts with $W_4 M_4 M_5 \widetilde{M}_5$, $\mathbf{x} \in \mathcal{C}_4^{(ii)}$ and $W_5 = W_4 M_3 = 10101$. Moreover, for $k \geq 5$, \mathbf{x} starts with $W_k M_{k-1} M_k \widetilde{M}_k$ and $W_{k+1} = W_k M_{k-1}$ where W_k is a non-empty suffix of M_k . Hence, $\mathbf{x} \in \mathcal{C}_k^{(ii)}$ for $k \geq 5$. Therefore, the locating chain of $\mathbf{x} = \overline{(ii)(i)(iii)(ii)}$.

Theorem 2.4 in [8] says that $r(n, \mathbf{x}) \leq 2n + 1$ for all $n \geq 1$ and equality holds for infinitely many n . Let

$$\Lambda(\mathbf{x}) := \{n \in \mathbb{N} : r(n, \mathbf{x}) = 2n + 1\}.$$

We have $\Lambda(\mathbf{x}) = \{n_1, n_2, \dots\}$ for an increasing sequence $\{n_i\}_{i \geq 1}$. From Lemma 5.3 in [8], $r(n, \mathbf{x}) \leq 2n$ implies $r(n, \mathbf{x}) \leq r(n-1, \mathbf{x}) + 1$. Thus,

$$r(n, \mathbf{x}) = r(n-1, \mathbf{x}) + 1 \quad \text{if } n \notin \Lambda(\mathbf{x}).$$

Hence, the sequence $\left\{ \frac{r(n, \mathbf{x})}{n} \right\}_{n \geq 1}$ is decreasing on each interval $[n_i, n_{i+1} - 1]$. It gives

$$\text{rep}(\mathbf{x}) = \liminf_{i \rightarrow \infty} \left(1 + \frac{n_i}{n_{i+1}} \right).$$

3. THE EXPONENTS OF REPETITION OF STURMIAN WORDS

Let \mathbf{x} be a Sturmian word of slope $\theta := [0; a_1, a_2, \dots]$. For $k \geq 1$, there is a relation between cases which \mathbf{x} belongs to at level k and $k + 1$.

Lemma 3.1. Let $k \geq 1$. The following statements are satisfied.

- (1) If $\mathbf{x} \in \mathcal{C}_k^{(i)}$ and $\mathbf{x} \in \mathcal{C}_{k+1}^{(i)} \cup \mathcal{C}_{k+1}^{(ii)}$, then $W_{k+1} = W_k M_k^t M_{k-1}$ for some $1 \leq t \leq a_{k+1} - 1$.
- (2) If $\mathbf{x} \in \mathcal{C}_k^{(i)}$ and $\mathbf{x} \in \mathcal{C}_{k+1}^{(iii)}$, then $W_{k+1} = W_k$.
- (3) If $\mathbf{x} \in \mathcal{C}_k^{(ii)}$, then $\mathbf{x} \in \mathcal{C}_{k+1}^{(i)} \cup \mathcal{C}_{k+1}^{(ii)}$ and $W_{k+1} = W_k M_{k-1}$.
- (4) If $\mathbf{x} \in \mathcal{C}_k^{(iii)}$, then $\mathbf{x} \in \mathcal{C}_{k+1}^{(i)} \cup \mathcal{C}_{k+1}^{(ii)}$ and $W_{k+1} = W_k$.

Proof. In this proof, for all $k \geq 1$, let W_k be the unique prefix of \mathbf{x} defined in which case \mathbf{x} belongs to at level k . Note that \widetilde{M}_k is a prefix of M_{k+1}^- by definition.

(1) Let $\mathbf{x} \in \mathcal{C}_{k+1}^{(i)} \cup \mathcal{C}_{k+1}^{(ii)}$. Note that \mathbf{x} starts with $W_{k+1} M_k \widetilde{M}_k$ for the suffix W_{k+1} of M_{k+1} . If W_{k+1} is a non-empty suffix of M_{k-1} , then $\mathbf{x} \in \mathcal{C}_k^{(iii)}$. It is a contradiction. If $W_{k+1} = W'_k M_{k-1}$ for some non-empty suffix W'_k of M_k , then \mathbf{x} starts with $W'_k M_{k-1} M_k \widetilde{M}_k$. Thus, $\mathbf{x} \in \mathcal{C}_k^{(ii)}$. It is a contradiction. Hence, $a_{k+1} > 1$ and $W_{k+1} = W''_k M_k^t M_{k-1}$ for some $1 \leq t \leq a_{k+1} - 1$ and some non-empty suffix W''_k of M_k . Consequently, \mathbf{x} starts with $W''_k M_k \widetilde{M}_k$. By the uniqueness of W_k , $W_k = W''_k$. It gives $W_{k+1} = W_k M_k^t M_{k-1}$.

(2) Let $\mathbf{x} \in \mathcal{C}_{k+1}^{(iii)}$. We have $\mathbf{x} = W_{k+1} M_{k+1} \widetilde{M}_{k+1}$ for a non-empty suffix W_{k+1} of M_k . Since $M_{k+1} \widetilde{M}_{k+1} = M_k M_{k+1}^- D_{k+1} M_{k+1}^-$, \mathbf{x} starts with $W_{k+1} M_k \widetilde{M}_k$. By the uniqueness of W_k , $W_{k+1} = W_k$.

(3) Let $\mathbf{x} \in \mathcal{C}_k^{(ii)}$. Note that \mathbf{x} starts with $W_k M_{k-1} M_k \widetilde{M}_k$ for a non-empty suffix W_k of M_k . Assume that $\mathbf{x} \in \mathcal{C}_{k+1}^{(iii)}$. Since \mathbf{x} starts with $W_{k+1} M_{k+1} \widetilde{M}_{k+1}$ for the suffix W_{k+1} of M_k , \mathbf{x} starts with $W_{k+1} M_k \widetilde{M}_k$. Hence, $\mathbf{x} \in \mathcal{C}_k^{(i)}$. It is a contradiction. Hence, $\mathbf{x} \in \mathcal{C}_{k+1}^{(i)} \cup \mathcal{C}_{k+1}^{(ii)}$. Thus, \mathbf{x} starts with $W_{k+1} M_k \widetilde{M}_k$ for the suffix W_{k+1} of M_{k+1} . If W_{k+1} is a non-empty suffix of M_{k-1} , then $\mathbf{x} \in \mathcal{C}_k^{(iii)}$. It is a contradiction. If $W_{k+1} = W'_k M_k^t M_{k-1}$ for some $1 \leq t \leq a_{k+1} - 1$ and some non-empty suffix W'_k of M_k , then \mathbf{x} starts with $W'_k M_k \widetilde{M}_k$. Thus, $\mathbf{x} \in \mathcal{C}_k^{(i)}$. It is a contradiction. Hence, $W_{k+1} = W'_k M_{k-1}$ for some non-empty suffix W'_k of M_k . By the uniqueness of W_k , $W_{k+1} = W_k M_{k-1}$.

(4) Let $\mathbf{x} \in \mathcal{C}_{k+1}^{(iii)}$. Note that \mathbf{x} starts with $W_k M_k \widetilde{M}_k$ for the suffix W_k of M_{k-1} . Assume that $\mathbf{x} \in \mathcal{C}_{k+1}^{(iii)}$. Since \mathbf{x} starts with $W_{k+1} M_{k+1} \widetilde{M}_{k+1}$ for the suffix W_{k+1} of M_k , \mathbf{x} starts with $W_{k+1} M_k \widetilde{M}_k$. Hence, $\mathbf{x} \in \mathcal{C}_k^{(i)}$. It is a contradiction. Hence, $\mathbf{x} \in \mathcal{C}_{k+1}^{(i)} \cup \mathcal{C}_{k+1}^{(ii)}$. Thus, \mathbf{x} starts with $W_{k+1} M_k \widetilde{M}_k$ for the suffix W_{k+1} of M_{k+1} . If $W_{k+1} = W'_k M_k^t M_{k-1}$ for $0 \leq t \leq a_{k+1} - 1$ and some non-empty suffix W'_k of M_k , then \mathbf{x} starts with $W'_k M_{k-1} M_k \widetilde{M}_k$ or $W'_k M_k \widetilde{M}_k$. Thus, $\mathbf{x} \in \mathcal{C}_k^{(i)} \cup \mathcal{C}_k^{(ii)}$. It is a contradiction. Hence, $W_{k+1} = W'_{k-1}$ for some non-empty suffix W'_{k-1} of M_{k-1} . By the uniqueness of W_k , $W_{k+1} = W_k$. \square

k	$k + 1$	The relation between W_{k+1} and W_k
case (i)	case (i) case (ii)	$W_{k+1} = W_k M_k^t M_{k-1}$ ($1 \leq t \leq a_{k+1} - 1$)
	case (iii)	$W_{k+1} = W_k$
case (ii)	case (i) case (ii)	$W_{k+1} = W_k M_{k-1}$
case (iii)	case (i) case (ii)	$W_{k+1} = W_k$

TABLE 1. The relation between W_{k+1} and W_k

Let

$$\begin{aligned} u_{t,k} &= tq_k + q_{k-1} - 1, & v_{t,k} &= |W_k| + tq_k + q_{k-1} - 1, \\ u'_k &= q_{k+1} - 1, & v'_k &= |W_k| + q_{k+1} - 1. \end{aligned}$$

The following lemma shows that all of elements in $\Lambda(\mathbf{x}) \cap [u_{1,1}, \infty)$ are expressed in terms of q_k 's and $|W_k|$'s.

Lemma 3.2. Let $k \geq 1$.

(1) If $\mathbf{x} \in \mathcal{C}_k^{(i)} \cap \mathcal{C}_{k+1}^{(i)}$, then

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+1} - 1] = \begin{cases} \{v_{t,k}, u'_k\} & \text{for } t = a_{k+1} - 1, \\ \{v_{t,k}, u_{t+1,k}, v_{t+1,k}, u'_k\} & \text{for } t < a_{k+1} - 1 \end{cases}$$

where t satisfies $W_{k+1} = W_k M_k^t M_{k-1}$.

(2) If $\mathbf{x} \in \mathcal{C}_k^{(i)} \cap \mathcal{C}_{k+1}^{(ii)}$, then

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+1} - 1] = \{v_{t,k}, u_{t+1,k}, v_{t+1,k}\}$$

where t satisfies $W_{k+1} = W_k M_k^t M_{k-1}$.

(3) If $\mathbf{x} \in \mathcal{C}_k^{(i)} \cap \mathcal{C}_{k+1}^{(iii)} \cap \mathcal{C}_{k+2}^{(i)}$, then

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+2} - 1] = \{v'_k, u'_{k+1}\}.$$

(4) If $\mathbf{x} \in \mathcal{C}_k^{(i)} \cap \mathcal{C}_{k+1}^{(iii)} \cap \mathcal{C}_{k+2}^{(ii)}$, then

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+2} - 1] = \{v'_k\}.$$

(5) If $\mathbf{x} \in \mathcal{C}_k^{(ii)} \cap \mathcal{C}_{k+1}^{(i)}$, then

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+1} - 1] = \begin{cases} \{u_{1,k}\} & \text{for } a_{k+1} = 1, \\ \{u_{1,k}, v_{1,k}, u'_k\} & \text{for } a_{k+1} > 1. \end{cases}$$

(6) If $\mathbf{x} \in \mathcal{C}_k^{(ii)} \cap \mathcal{C}_{k+1}^{(ii)}$, then

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+1} - 1] = \{u_{1,k}, v_{1,k}\}.$$

Proof. For $\mathbf{x} = x_1x_2\dots$, let $x_i^j := x_ix_{i+1}\dots x_j$.

(1) Suppose that $\mathbf{x} \in \mathcal{C}_k^{(i)} \cap \mathcal{C}_{k+1}^{(i)}$. Since M_kM_{k-1} is primitive, Lemma 7.1 in [8] implies that for

$$\mathbf{x} = W_kM_k\widetilde{M}_k\cdots = W_kM_kM_{k-1}M_k^{--} = W_kM_{k-1}M_k^{--}D_kM_{k-1}^{--}\cdots,$$

the first q_k subwords of length $u_{1,k}$ are mutually distinct. From $x_1^{u_{1,k}} = x_{q_k+1}^{u_{2,k}}$, $r(u_{1,k}, \mathbf{x}) = u_{2,k}$. Note that

$$\begin{aligned} \mathbf{x} &= W_{k+1}M_{k+1}\widetilde{M}_{k+1}\cdots = W_kM_k^tM_{k-1}M_k^{a_{k+1}}M_{k-1}\widetilde{M}_{k+1}\cdots \\ &= W_kM_k^{t+1}M_{k-1}^{--}D'_kM_k^{a_{k+1}-1}M_{k-1}\widetilde{M}_{k+1}\cdots \end{aligned}$$

Since $x_1^{v_{t,k}-1} = x_{q_k+1}^{v_{t+1,k}-1}$, $r(v_{t,k} - 1, \mathbf{x}) \leq v_{t,k} - 1$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + q_k$$

for $u_{1,k} \leq n \leq v_{t,k} - 1$. Moreover, we have $r(v_{t,k}, \mathbf{x}) \geq r(v_{t,k} - 1, \mathbf{x}) + 2$. Hence, $r(v_{t,k}, \mathbf{x}) = 2v_{t,k} + 1$ by Theorem 2.4 and Lemma 5.3 in [8]. Since $x_1^{u_{t+1,k}-1} = x_{v_{t,k}+2}^{v_{2t+1,k}+q_{k-1}-1}$, we have $r(u_{t+1,k} - 1, \mathbf{x}) \leq v_{2t+1,k} + q_{k-1} - 1$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + v_{t,k} + 1$$

for $v_{t,k} \leq n \leq u_{t+1,k} - 1$. Note that

$$\begin{aligned} \mathbf{x} &= W_{k+1}M_{k+1}\widetilde{M}_{k+1}\cdots = W_kM_k^tM_{k-1}M_k^{a_{k+1}}M_{k-1}\widetilde{M}_{k+1}\cdots \\ &= W_kM_k^{t+1}M_{k-1}^{--}D'_kM_k^{a_{k+1}-1}M_{k-1}\widetilde{M}_{k+1}\cdots = W_kM_k^tM_{k-1}M_k^{t+1}M_{k-1}\cdots \end{aligned}$$

It gives $r(u_{t+1,k}, \mathbf{x}) \geq r(u_{t+1,k} - 1, \mathbf{x}) + 2$. Thus, we have $r(u_{t+1,k}, \mathbf{x}) = 2u_{t+1,k} + 1$ by Theorem 2.4 and Lemma 5.3 in [8]. If $t = a_{k+1} - 1$, then $(t+1)q_k + q_{k-1} = q_{k+1}$. By the argument used at level k , $r(u_{1,k+1} - 1, \mathbf{x}) = u_{2,k+1} - 1$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + u'_k + 1$$

for $u'_k \leq n \leq u_{1,k+1} - 1$. It follows that

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+1} - 1] = \{v_{t,k}, u'_k\}.$$

Now, let $t < a_{k+1} - 1$. Note that

$$\begin{aligned} \mathbf{x} &= W_{k+1}M_{k+1}\widetilde{M}_{k+1}\cdots = W_kM_k^tM_{k-1}M_k^{a_{k+1}}M_{k-1}M_kM_{k+1}^{--}\cdots \\ &= W_kM_k^tM_{k-1}M_kM_k^{a_{k+1}-2}M_kM_{k-1}M_kM_{k+1}^{--}\cdots \\ &= W_kM_k^{t+1}M_{k-1}^{--}D'_kM_k^{a_{k+1}-1}M_kM_{k-1}^{--}D'_kM_{k+1}^{--}\cdots \end{aligned}$$

Since $x_1^{v_{t+1,k}-1} = x_{u_{t+1,k}+2}^{|W_k|+tq_k+q_{k-1}+((t+2)q_k+q_{k-1}-2)}$, we have $r(v_{t+1,k} - 1, \mathbf{x}) \leq v_{2t+2,k} + q_{k-1} - 1$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + u_{t+1,k} + 1$$

for $u_{t+1,k} \leq n \leq v_{t+1,k} - 1$. Note that

$$\begin{aligned} \mathbf{x} &= W_{k+1}M_{k+1}\widetilde{M}_{k+1}\cdots = W_kM_k^tM_{k-1}M_k^{a_{k+1}}M_{k-1}M_kM_{k+1}^{--}\cdots \\ &= W_kM_k^{t+1}M_{k-1}^{--}D'_kM_k^{a_{k+1}-1}M_kM_{k-1}^{--}D'_kM_{k+1}^{--}\cdots \\ &= W_kM_k^tM_{k-1}M_k^{t+2}M_{k-1}\cdots \end{aligned}$$

It gives $r(v_{t+1,k}, \mathbf{x}) \geq r(v_{t+1,k} - 1, \mathbf{x}) + 2$. Hence, we have $r(v_{t+1,k}, \mathbf{x}) = 2v_{t+1,k} + 1$ from Theorem 2.4 and Lemma 5.3 in [8]. On the other hand, from

$$\begin{aligned} \mathbf{x} &= W_{k+1}M_{k+1}\widetilde{M}_{k+1}\cdots = W_kM_k^tM_{k-1}M_kM_k^{a_{k+1}-1}M_{k-1}M_kM_{k+1}^{--}\cdots \\ &= W_kM_k^tM_{k-1}M_kM_k^{a_{k+1}}M_{k-1}^{--}D'_kM_{k+1}^{--}\cdots, \end{aligned}$$

we have $x_{|W_k|+tq_k+q_{k-1}+(a_{k+1}q_k+q_{k-1}-2)}^{|W_k|+tq_k+q_{k-1}+1} = x_{|W_k|+(t+1)q_k+q_{k-1}+(a_{k+1}q_k+q_{k-1}-2)}^{|W_k|+(t+1)q_k+q_{k-1}+1}$. It gives $r(u'_k - 1, \mathbf{x}) \leq (v_{t+1,k} + 1) + (u'_k - 1)$. The fact that $r(n + 1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + v_{t+1,k} + 1$$

for $v_{t+1,k} \leq n \leq u'_k - 1$. Moreover, $r(u'_k, \mathbf{x}) \geq r(u'_k - 1, \mathbf{x}) + 2$. From Theorem 2.4 and Lemma 5.3 in [8], $r(u'_k, \mathbf{x}) = 2u'_k + 1$. Note that $\mathbf{x} \in \mathcal{C}_{k+1}^{(i)}$. By the argument used at level k , $r(u_{1,k+1} - 1, \mathbf{x}) = u_{2,k+1} - 1$. The fact that $r(n + 1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + u'_k + 1$$

for $u'_k \leq n \leq u_{1,k+1} - 1$. It follows that

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+1} - 1] = \{v_{t,k}, u_{t+1,k}, v_{t+1,k}, u'_k\}.$$

(2) Suppose that $\mathbf{x} \in \mathcal{C}_k^{(i)} \cap \mathcal{C}_{k+1}^{(ii)}$. Note that

$$\mathbf{x} = W_{k+1}M_k\cdots = W_kM_k^tM_{k-1}M_k\cdots = W_kM_k^{t+1}M_{k-1}^{--}D'_k\cdots$$

Use the argument used at level k in (1). Since $x_1^{v_{t,k}-1} = x_{q_k+1}^{v_{t+1,k}-1}$,

$$r(n, \mathbf{x}) = n + q_k$$

for $u_{1,k} \leq n \leq v_{t,k} - 1$. Note that

$$\begin{aligned} \mathbf{x} &= W_{k+1}M_kM_{k+1}\widetilde{M}_{k+1}\cdots = W_kM_k^tM_{k-1}M_kM_k^{a_{k+1}}M_{k-1}\widetilde{M}_{k+1}\cdots \\ &= W_kM_k^{t+1}M_{k-1}^{--}D'_kM_k^{a_{k+1}}M_{k-1}\widetilde{M}_{k+1}\cdots = W_kM_k^tM_{k-1}M_k^{t+1}M_{k-1}\cdots \end{aligned}$$

Since $x_{|W_k|+u_{t+1,k}-1}^{|W_k|+u_{t+1,k}-1} = x_{v_{t,k}+1}^{v_{t,k}+1+u_{t+1,k}-1}$, $r(u_{t+1,k} - 1, \mathbf{x}) \leq v_{2t+1,k} + q_{k-1} - 1$. Moreover, $r(v_{t,k}, \mathbf{x}) \geq r(v_{t,k} - 1, \mathbf{x}) + 2$. From Theorem 2.4 and Lemma 5.3 in [8], $r(v_{t,k}, \mathbf{x}) = 2v_{t,k} + 1$. The fact that $r(n + 1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + v_{t,k} + 1$$

for $v_{t,k} \leq n \leq u_{t+1,k} - 1$. Moreover, $r(u_{t+1,k}, \mathbf{x}) \geq r(u_{t+1,k} - 1, \mathbf{x}) + 2$. From Theorem 2.4 and Lemma 5.3 in [8], $r(u_{t+1,k}, \mathbf{x}) = 2u_{t+1,k} + 1$. On the other hand, note that

$$\mathbf{x} = W_{k+1}M_kM_{k+1}\widetilde{M}_{k+1}\cdots = W_kM_k^tM_{k-1}M_kM_k^{a_{k+1}}M_{k-1}M_k\cdots$$

$$= W_k M_k^t M_{k-1} M_k M_k^{a_{k+1}+1} M_{k-1}^{--} \dots = W_k M_k^{t+1} M_{k-1}^{--} D'_k M_k^{a_{k+1}+1} M_{k-1}^{--} \dots$$

Since $x_1^{v_{t+1,k}-1} = x_{u_{t+1,k}+2}^{u_{t+1,k}+1+v_{t+1,k}-1}$, $r(v_{t+1,k}-1, \mathbf{x}) \leq v_{2t+2,k} + q_{k-1} - 1$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + u_{t+1,k} + 1$$

for $u_{t+1,k} \leq n \leq v_{t+1,k} - 1$. Moreover, $r(v_{t+1,k}, \mathbf{x}) \geq r(v_{t+1,k} - 1, \mathbf{x}) + 2$. From Theorem 2.4 and Lemma 5.3 in [8], $r(v_{t+1,k}, \mathbf{x}) = 2v_{t+1,k} + 1$. On the other hand, note that

$$\begin{aligned} \mathbf{x} &= W_{k+1} M_k M_{k+1} \widetilde{M}_{k+1} \dots = W_k M_k^t M_{k-1} M_k M_k^{a_{k+1}} M_{k-1} M_k \dots \\ &= W_k M_k^t M_{k-1} M_k M_k^{a_{k+1}+1} M_{k-1}^{--} D'_k \dots \end{aligned}$$

Since $x_{v_{t,k}+2}^{v_{t,k}+1+u_{a_{k+1}+1,k}-1} = x_{v_{t+1,k}+2}^{v_{t+1,k}+1+u_{a_{k+1}+1,k}-1}$, $r(u_{1,k+1} - 1, \mathbf{x}) \leq u_{1,k+1} - 1 + v_{t+1,k} + 1$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + v_{t+1,k} + 1$$

for $v_{t+1,k} \leq n \leq u_{1,k+1} - 1$. Hence,

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+1} - 1] = \{v_{t,k}, u_{t+1,k}, v_{t+1,k}\}.$$

(3) Suppose that $\mathbf{x} \in \mathcal{C}_k^{(i)} \cap \mathcal{C}_{k+1}^{(iii)} \cap \mathcal{C}_{k+2}^{(i)}$. Since $\mathbf{x} \in \mathcal{C}_k^{(i)}$, the argument used at level k in (1) yields $r(u_{1,k}, \mathbf{x}) = u_{2,k}$. Note that

$$\mathbf{x} = W_{k+1} M_{k+1} \widetilde{M}_{k+1} \dots = W_k M_k^{a_{k+1}} M_{k-1} M_k M_{k+1}^{--} \dots = W_k M_k^{a_{k+1}} \widetilde{M}_k D'_k M_{k+1}^{--} \dots$$

Since $x_1^{v_{a_{k+1},k}-1} = x_{q_k+1}^{v_{a_{k+1},k}+q_k-1}$, $r(v'_k - 1, \mathbf{x}) \leq v'_k + q_k - 1$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + q_k$$

for $u_{1,k} \leq n \leq v'_k - 1$. Moreover, $r(v'_k, \mathbf{x}) \geq r(v'_k - 1, \mathbf{x}) + 2$. From Theorem 2.4 and Lemma 5.3 in [8], $r(v'_k, \mathbf{x}) = 2v'_k + 1$. Note that

$$\begin{aligned} \mathbf{x} &= W_{k+2} M_{k+2} \widetilde{M}_{k+2} \dots = W_k M_{k+1} M_{k+1}^{a_{k+2}-1} M_k \widetilde{M}_{k+2} \dots \\ &= W_k M_{k+1} M_{k+1}^{a_{k+2}} M_k^{--} D'_{k+1} M_{k+2}^{--} \dots \end{aligned}$$

Since $x_{|W_k|+1}^{|W_k|+u_{a_{k+2},k+1}-1} = x_{|W_k|+q_{k+1}+1}^{|W_k|+u_{a_{k+2}+1,k+1}-1}$, $r(u'_{k+1} - 1, \mathbf{x}) \leq |W_k| + q_{k+1} + u'_{k+1} - 1$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + v'_k + 1$$

for $v'_k \leq n \leq u'_{k+1} - 1$. Moreover, $r(u'_{k+1}, \mathbf{x}) \geq r(u'_{k+1} - 1, \mathbf{x}) + 2$. From Theorem 2.4 and Lemma 5.3 in [8], $r(u'_{k+1}, \mathbf{x}) = 2u'_{k+1} + 1$. Since $\mathbf{x} \in \mathcal{C}_{k+2}^{(i)}$, the argument used at level k in (1) implies $r(u_{1,k+2}, \mathbf{x}) = u_{2,k+2}$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + u'_{k+1} + 1$$

for $u'_{k+1} \leq n \leq u_{1,k+2}$. Hence,

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+2} - 1] = \{v'_k, u'_{k+1}\}.$$

(4) Suppose that $\mathbf{x} \in \mathcal{C}_k^{(i)} \cap \mathcal{C}_{k+1}^{(iii)} \cap \mathcal{C}_{k+2}^{(ii)}$. Use the same argument with (3). Since $\mathbf{x} \in \mathcal{C}_k^{(i)} \cap \mathcal{C}_{k+1}^{(iii)}$, we have

$$r(n, \mathbf{x}) = n + q_k$$

for $u_{1,k} \leq n \leq v'_k - 1$ and $r(v'_k, \mathbf{x}) = 2v'_k + 1$. On the other hand, note that

$$\begin{aligned} \mathbf{x} &= W_{k+2}M_{k+1}M_{k+2}\widetilde{M}_{k+2}\cdots = W_kM_{k+1}M_{k+1}^{a_{k+2}}M_k\widetilde{M}_{k+2}\cdots \\ &= W_kM_{k+1}M_{k+1}^{a_{k+2}+1}M_k^{--}D'_{k+1}M_{k+2}^{--}\cdots \end{aligned}$$

Since $x_{|W_k|+u_{a_{k+2}+1,k+1}-1}^{|W_k|+u_{a_{k+2}+1,k+1}-1} = x_{|W_k|+q_{k+1}+u_{a_{k+2}+1,k+1}-1}^{|W_k|+q_{k+1}+u_{a_{k+2}+1,k+1}-1}$, $r(u_{1,k+2} - 1, \mathbf{x}) \leq |W_k| + q_{k+1} + u_{1,k+2} - 1$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + v'_k + 1$$

for $v'_k \leq n \leq u_{1,k+2} - 1$. Hence,

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+2} - 1] = \{v'_k\}.$$

(5) Suppose that $\mathbf{x} \in \mathcal{C}_k^{(ii)} \cap \mathcal{C}_{k+1}^{(i)}$. Note that

$$\mathbf{x} = W_kM_{k-1}M_k\widetilde{M}_k\cdots = W_kM_{k-1}M_kM_{k-1}M_k^{--}\cdots$$

Since M_kM_{k-1} is primitive, Lemma 7.1 in [8] implies that the first $(u_{1,k} + 1)$ subwords of length $u_{1,k}$ are mutually distinct. Thus, $r(u_{1,k}, \mathbf{x}) = 2u_{1,k} + 1$. Since $x_1^{v_{1,k}-1} = x_{q_k+q_{k-1}+v_{1,k}-1}^{q_k+q_{k-1}+v_{1,k}-1}$, $r(v_{1,k} - 1, \mathbf{x}) \leq q_k + q_{k-1} + v_{1,k} - 1$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + u_{1,k} + 1$$

for $u_{1,k} \leq n \leq v_{1,k} - 1$. Note that

$$\mathbf{x} = W_{k+1}M_{k+1}\widetilde{M}_{k+1}\cdots = W_kM_{k-1}M_k^{a_{k+1}}M_{k-1}M_kM_{k+1}^{--}\cdots$$

If $a_{k+1} = 1$, then $x_1^{v_{2,k}+q_{k-1}-1} = x_{q_k+q_{k-1}+v_{2,k}+q_{k-1}-1}^{q_k+q_{k-1}+v_{2,k}+q_{k-1}-1}$. It implies $r(v_{2,k} + q_{k-1} - 1, \mathbf{x}) \leq v_{3,k} + 2q_{k-1} - 1$. Since $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$,

$$r(n, \mathbf{x}) = n + u_{1,k} + 1$$

for $u_{1,k} \leq n \leq v'_k + q_{k+1} - 1$. Hence,

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+1} - 1] = \{u_{1,k}\}.$$

Now, let $a_{k+1} > 1$. Since $\mathbf{x} = W_kM_{k-1}M_kM_kM_{k-1}\cdots$, $r(v_{1,k}, \mathbf{x}) \geq r(v_{1,k} - 1, \mathbf{x}) + 2$. From Theorem 2.4 and Lemma 5.3 in [8], $r(v_{1,k}, \mathbf{x}) = 2v_{1,k} + 1$. On the other hand, note that

$$\begin{aligned} \mathbf{x} &= W_{k+1}M_{k+1}\widetilde{M}_{k+1}\cdots = W_kM_{k-1}M_k^{a_{k+1}}M_{k-1}\widetilde{M}_{k+1}^{--}\cdots \\ &= W_kM_{k-1}M_kM_k^{a_{k+1}-1}M_{k-1}\widetilde{M}_{k+1}^{--}\cdots = W_kM_{k-1}M_kM_k^{a_{k+1}}M_{k-1}^{--}D'_kM_{k+1}^{--}\cdots \end{aligned}$$

Since $x_{|W_k|+q_{k-1}+u_{a_{k+1},k}-1}^{|W_k|+q_{k-1}+u_{a_{k+1},k}-1} = x_{|W_k|+q_k+q_{k-1}+u_{a_{k+1},k}-1}^{|W_k|+q_k+q_{k-1}+u_{a_{k+1},k}-1}$, $r(u'_k - 1, \mathbf{x}) \leq |W_k| + q_k + q_{k-1} + u'_k - 1$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + v_{1,k} + 1$$

for $v_{1,k} \leq n \leq u'_k - 1$. Moreover, $r(u'_k, \mathbf{x}) \geq r(u'_k - 1, \mathbf{x}) + 2$. From Theorem 2.4 and Lemma 5.3 in [8], $r(u'_k, \mathbf{x}) = 2u'_k + 1$. Since $\mathbf{x} \in \mathcal{C}_{k+1}^{(i)}$, the argument used at level k in (1) implies $r(u_{1,k+1}, \mathbf{x}) = u_{2,k+1}$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + u'_k + 1$$

for $u'_k \leq n \leq u_{1,k+1}$. Hence,

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+1} - 1] = \{u_{1,k}, v_{1,k}, u'_k\}.$$

(6) Suppose that $\mathbf{x} \in \mathcal{C}_k^{(ii)} \cap \mathcal{C}_{k+1}^{(ii)}$. Use the same argument with (5). Since $\mathbf{x} \in \mathcal{C}_k^{(ii)}$, we have

$$r(n, \mathbf{x}) = n + u_{1,k} + 1$$

for $u_{1,k} \leq n \leq v_{1,k} - 1$. Note that

$$\begin{aligned} \mathbf{x} &= W_{k+1} M_k M_{k+1} \widetilde{M}_{k+1} \cdots = W_k M_{k-1} M_k M_k^{a_{k+1}} M_{k-1} M_k M_{k+1} \overline{\overline{\cdots}} \\ &= W_k M_{k-1} M_k M_k^{a_{k+1}+1} M_{k-1} \overline{\overline{D'_k M_{k+1} \overline{\overline{\cdots}}}} \end{aligned}$$

Since $\mathbf{x} = W_k M_{k-1} M_k M_k M_{k-1} \cdots$, $r(v_{1,k}, \mathbf{x}) \geq r(v_{1,k} - 1, \mathbf{x}) + 2$. From Theorem 2.4 and Lemma 5.3 in [8], $r(v_{1,k}, \mathbf{x}) = 2v_{1,k} + 1$. Moreover, since $x_{|W_k|+q_{k-1}+u_{a_{k+1}+1, k-1}} = x_{v_{1,k}+1+u_{a_{k+1}+1, k-1}} = x_{v_{1,k}+2}$, $r(u_{1,k+1} - 1, \mathbf{x}) \leq |W_k| + q_k + q_{k-1} + u_{1,k+1} - 1$. The fact that $r(n+1, \mathbf{x}) \geq r(n, \mathbf{x}) + 1$ for any $n \geq 1$ gives

$$r(n, \mathbf{x}) = n + v_{1,k} + 1$$

for $v_{1,k} \leq n \leq u_{1,k+1} - 1$. Hence,

$$\Lambda(\mathbf{x}) \cap [u_{1,k}, u_{1,k+1} - 1] = \{u_{1,k}, v_{1,k}\}.$$

□

Remark. When $\mathbf{x} \in \mathcal{C}_1^{(iii)}$, Lemma 3.2 does not determine the elements in $\Lambda(\mathbf{x}) \cap [u_{1,1}, u_{1,2} - 1]$. Thus, we should check how the elements of $\Lambda(\mathbf{x}) \cap [u_{1,1}, u_{1,2} - 1]$ are expressed in terms of q_k 's and $|W_k|$'s. If $\mathbf{x} \in \mathcal{C}_1^{(iii)}$, then \mathbf{x} starts with $0^{a_1} 10^{a_1-1}$. Thus, $r(n, \mathbf{x}) = n + 1$ for $1 \leq n \leq v'_0 - 1$. Moreover, since $\mathbf{x} \in \mathcal{C}_2^{(i)} \cup \mathcal{C}_2^{(ii)}$, \mathbf{x} starts with $0^{a_1} 10^{a_1}$ or $0^{a_1} 10^{a_1-1} 1$. Thus, $r(v'_0, \mathbf{x}) = 2v'_0 + 1$. Hence, we can follow the proof of Lemma 3.2 (3) and (4). We have $\Lambda(\mathbf{x}) \cap [u_{1,1}, u_{1,2} - 1] = \{v'_0, u'_1\}$ for $\mathbf{x} \in \mathcal{C}_2^{(i)}$ and $\Lambda(\mathbf{x}) \cap [u_{1,1}, u_{1,2} - 1] = \{v'_0\}$ for $\mathbf{x} \in \mathcal{C}_2^{(ii)}$.

For $l = 1, 2, \dots, 6$, define

$$\Lambda_l(\mathbf{x}) := \{n \in \Lambda(\mathbf{x}) : n \text{ appears in } (l) \text{ of Lemma 3.2}\}.$$

It is obvious that $\Lambda_l(\mathbf{x})$'s are mutually distinct and $\cup_{l=1}^6 \Lambda_l(\mathbf{x}) = \Lambda(\mathbf{x}) \cap [u_{1,1}, \infty)$. Now, we find the minimum of $\mathcal{L}(\theta)$ where θ has bounded partial quotients.

Theorem 3.3. Let $\theta = [0; a_1, a_2, \dots]$ have bounded partial quotients. We have

$$\min \mathcal{L}(\theta) = \liminf_{k \rightarrow \infty} [1; 1 + a_k, a_{k-1}, a_{k-2}, \dots, a_1].$$

Proof. Let \mathbf{x} be a Sturmian word of slope θ . For $k \geq 1$, set $\eta_k := \frac{q_{k-1}}{q_k}$, $t_k := \frac{|W_k|}{q_k}$, $\epsilon_k := \frac{1}{q_k}$. Note that $t_k \leq 1$, $\lim_{k \rightarrow \infty} \epsilon_k = 0$, and $\eta_k \geq \epsilon_k$ for $k \geq 1$. Set $m_\theta := \underline{\lim}_{i \rightarrow \infty} a_i$, $M_\theta := \overline{\lim}_{i \rightarrow \infty} a_i$. Let $\liminf_{k \rightarrow \infty} \eta_k = [0; b_1, b_2, \dots]$.

First, assume that $a_i > 1$ for infinitely many i . Since $\eta_k = [0; a_k, a_{k-1}, \dots, a_1]$ and $M_\theta \geq 2$, we have $b_1 = M_\theta$, $\liminf_{k \rightarrow \infty} \eta_k < \frac{1}{2}$. Using Lemma 3.2, let us prove that $\liminf_{n_i \in \Lambda(\mathbf{x})} \frac{n_i}{n_{i+1}} \geq$

$\liminf_{k \rightarrow \infty} [0; 1 + a_{k+1}, a_k, \dots]$ through the 7 cases below. Note that $\liminf_{k \rightarrow \infty} \eta_k > \frac{\liminf_{k \rightarrow \infty} \eta_k}{1 + \liminf_{k \rightarrow \infty} \eta_k} =$

$\liminf_{k \rightarrow \infty} \frac{\eta_k}{1 + \eta_k} = \liminf_{k \rightarrow \infty} [0; 1 + a_k, a_{k-1}, \dots]$.

(1) For any $n_i \in \Lambda_1(\mathbf{x})$ with $n_{i+1} \in \Lambda_1(\mathbf{x}) \cup \Lambda_2(\mathbf{x})$, $\frac{n_i}{n_{i+1}}$ is

$$\frac{t_k + t + \eta_k - \epsilon_k}{t + 1 + \eta_k - \epsilon_k}, \frac{t + 1 + \eta_k - \epsilon_k}{t + 1 + t_k + \eta_k - \epsilon_k}, \frac{\eta_{k+1}(t + 1 + t_k + \eta_k - \epsilon_k)}{1 - \epsilon_{k+1}}, \text{ or } \frac{1 - \epsilon_{k+1}}{t' + t_{k+1} + \eta_{k+1} - \epsilon_{k+1}}$$

for some $k = k(i)$ and t, t' satisfying $W_{k+1} = W_k M_k^t M_{k-1}$, $W_{k+2} = W_{k+1} M_{k+1}^{t'} M_k$. We have

$$\liminf_{k \rightarrow \infty} \frac{t_k + t + \eta_k - \epsilon_k}{t + 1 + \eta_k - \epsilon_k}, \liminf_{k \rightarrow \infty} \frac{t + 1 + \eta_k - \epsilon_k}{t + 1 + t_k + \eta_k - \epsilon_k} \geq \frac{t}{t + 1} \geq \frac{1}{2} > \liminf_{k \rightarrow \infty} \eta_k,$$

$$\liminf_{k \rightarrow \infty} \frac{\eta_{k+1}}{1 - \epsilon_{k+1}} (t + 1 + t_k + \eta_k - \epsilon_k) \geq \liminf_{k \rightarrow \infty} 2\eta_{k+1} \geq \liminf_{k \rightarrow \infty} \eta_k,$$

and

$$\liminf_{k \rightarrow \infty} \frac{1 - \epsilon_{k+1}}{t_{k+1} + t' + \eta_{k+1} - \epsilon_{k+1}} = \liminf_{k \rightarrow \infty} \frac{1}{t_{k+1} + t' + \eta_{k+1}} \geq \liminf_{k \rightarrow \infty} \frac{1}{a_{k+2} + \eta_{k+1}} = \liminf_{k \rightarrow \infty} \eta_{k+2}.$$

(2) For any $n_i \in \Lambda_2(\mathbf{x})$, $\frac{n_i}{n_{i+1}}$ is

$$\frac{t_k + t + \eta_k - \epsilon_k}{t + 1 + \eta_k - \epsilon_k}, \frac{t + 1 + \eta_k - \epsilon_k}{t + 1 + t_k + \eta_k - \epsilon_k}, \text{ or } \frac{\eta_{k+1}(t + 1 + t_k + \eta_k - \epsilon_k)}{1 + \eta_{k+1} - \epsilon_{k+1}}$$

for some $k = k(i)$ and t satisfying $W_{k+1} = W_k M_k^t M_{k-1}$. From the previous case,

$$\liminf_{k \rightarrow \infty} \frac{t_k + t + \eta_k - \epsilon_k}{t + 1 + \eta_k - \epsilon_k}, \liminf_{k \rightarrow \infty} \frac{t + 1 + \eta_k - \epsilon_k}{t + 1 + t_k + \eta_k - \epsilon_k} \geq \frac{1}{2} > \liminf_{k \rightarrow \infty} \eta_k.$$

We also have

$$\liminf_{k \rightarrow \infty} \frac{\eta_{k+1}}{1 + \eta_{k+1} - \epsilon_{k+1}} (t + 1 + t_k + \eta_k - \epsilon_k) \geq \liminf_{k \rightarrow \infty} \frac{2\eta_{k+1}}{1 + \eta_{k+1}} \geq \liminf_{k \rightarrow \infty} \eta_k.$$

(3) For any $n_i \in \Lambda_5(\mathbf{x})$ with $n_{i+1} \in \Lambda_1(\mathbf{x}) \cup \Lambda_2(\mathbf{x}) \cup \Lambda_5(\mathbf{x})$, $\frac{n_i}{n_{i+1}}$ is

$$\frac{1 + \eta_k - \epsilon_k}{1 + t_k + \eta_k - \epsilon_k}, \frac{\eta_{k+1}}{1 - \epsilon_{k+1}} (1 + t_k + \eta_k - \epsilon_k), \text{ or } \frac{1 - \epsilon_{k+1}}{t + t_{k+1} + \eta_{k+1} - \epsilon_{k+1}}$$

for some $k = k(i)$ and t satisfying $W_{k+2} = W_{k+1} M_{k+1}^t M_k$. We have

$$\liminf_{k \rightarrow \infty} \frac{1 + \eta_k - \epsilon_k}{1 + t_k + \eta_k - \epsilon_k} \geq \liminf_{k \rightarrow \infty} \frac{1}{1 + t_k} \geq \frac{1}{2} > \liminf_{k \rightarrow \infty} \eta_k,$$

$$\liminf_{k \rightarrow \infty} \frac{\eta_{k+1}}{1 - \epsilon_{k+1}} (1 + t_k + \eta_k - \epsilon_k) \geq \liminf_{k \rightarrow \infty} \eta_{k+1},$$

and

$$\liminf_{k \rightarrow \infty} \frac{1 - \epsilon_{k+1}}{t_{k+1} + t + \eta_{k+1} - \epsilon_{k+1}} = \liminf_{k \rightarrow \infty} \frac{1}{t_{k+1} + t + \eta_{k+1}} \geq \liminf_{k \rightarrow \infty} \frac{1}{a_{k+2} + \eta_{k+1}} = \liminf_{k \rightarrow \infty} \eta_{k+2}.$$

(4) For any $n_i \in \Lambda_6(\mathbf{x})$, $\frac{n_i}{n_{i+1}}$ is

$$\frac{1 + \eta_k - \epsilon_k}{1 + t_k + \eta_k - \epsilon_k} \text{ or } \frac{\eta_{k+1}}{1 + \eta_{k+1} - \epsilon_{k+1}} (1 + t_k + \eta_k - \epsilon_k)$$

for some $k = k(i)$. From the previous case,

$$\liminf_{k \rightarrow \infty} \frac{1 + \eta_k - \epsilon_k}{1 + t_k + \eta_k - \epsilon_k} > \liminf_{k \rightarrow \infty} \eta_k.$$

We also have

$$\begin{aligned} & \liminf_{k \rightarrow \infty} \frac{\eta_{k+1}}{1 + \eta_{k+1} - \epsilon_{k+1}} (1 + t_k + \eta_k - \epsilon_k) \geq \liminf_{k \rightarrow \infty} \frac{\eta_{k+1}}{1 + \eta_{k+1}} \liminf_{k \rightarrow \infty} (1 + t_k + \eta_k) \\ & \geq \frac{\liminf_{k \rightarrow \infty} \eta_{k+1}}{1 + \liminf_{k \rightarrow \infty} \eta_{k+1}} (1 + \liminf_{k \rightarrow \infty} \eta_k) = \liminf_{k \rightarrow \infty} \eta_{k+1}. \end{aligned}$$

(5) For any $n_i \in \Lambda_3(\mathbf{x})$ with $n_{i+1} \in \Lambda_1(\mathbf{x}) \cup \Lambda_2(\mathbf{x}) \cup \Lambda_3(\mathbf{x})$, $\frac{n_i}{n_{i+1}}$ is

$$(1 + t_k \eta_{k+1} - \epsilon_{k+1}) \frac{\eta_{k+2}}{1 - \epsilon_{k+2}} \text{ or } \frac{1 - \epsilon_{k+2}}{t + t_{k+2} + \eta_{k+2} - \epsilon_{k+2}}$$

for some $k = k(i)$ and t satisfying $W_{k+3} = W_{k+2} M_{k+2}^t M_{k+1}$. We have

$$\liminf_{k \rightarrow \infty} \frac{(1 + t_k \eta_{k+1} - \epsilon_{k+1}) \eta_{k+2}}{1 - \epsilon_{k+2}} \geq \liminf_{k \rightarrow \infty} \frac{(1 - \epsilon_{k+1}) \eta_{k+2}}{1 - \epsilon_{k+2}} = \liminf_{k \rightarrow \infty} \eta_{k+2}$$

and

$$\liminf_{k \rightarrow \infty} \frac{1 - \epsilon_{k+2}}{t_{k+2} + t + \eta_{k+2} - \epsilon_{k+2}} = \liminf_{k \rightarrow \infty} \frac{1}{t_{k+2} + t + \eta_{k+2}} \geq \liminf_{k \rightarrow \infty} \frac{1}{a_{k+3} + \eta_{k+2}} = \liminf_{k \rightarrow \infty} \eta_{k+3}.$$

(6) For any $n_i \in \Lambda_1(\mathbf{x}) \cup \Lambda_3(\mathbf{x}) \cup \Lambda_5(\mathbf{x})$ with $n_{i+1} \in \Lambda_3(\mathbf{x}) \cup \Lambda_4(\mathbf{x})$, $\frac{n_i}{n_{i+1}}$ is

$$\frac{\eta_{k+1} - \epsilon_{k+1}}{1 + t_k \eta_{k+1} - \epsilon_{k+1}}$$

for some $k = k(i)$. We have

$$\liminf_{k \rightarrow \infty} \frac{\eta_{k+1} - \epsilon_{k+1}}{1 + t_k \eta_{k+1} - \epsilon_{k+1}} \geq \liminf_{k \rightarrow \infty} \frac{\eta_{k+1}}{1 + \eta_{k+1}} = \liminf_{k \rightarrow \infty} [0; 1 + a_{k+1}, a_k, a_{k-1}, \dots].$$

(7) For any $n_i \in \Lambda_4(\mathbf{x})$, $\frac{n_i}{n_{i+1}}$ is

$$\frac{\eta_{k+2}}{1 + \eta_{k+2} - \epsilon_{k+2}} (1 + t_k \eta_{k+1} - \epsilon_{k+1})$$

for some $k = k(i)$. We have

$$\liminf_{k \rightarrow \infty} \frac{\eta_{k+2} (1 + t_k \eta_{k+1} - \epsilon_{k+1})}{1 + \eta_{k+2} - \epsilon_{k+2}} \geq \liminf_{k \rightarrow \infty} \frac{\eta_{k+2}}{1 + \eta_{k+2}} = \liminf_{k \rightarrow \infty} [0; 1 + a_{k+2}, a_{k+1}, a_k, \dots].$$

Hence, from (1)-(7),

$$\text{rep}(\mathbf{x}) \geq \liminf_{k \rightarrow \infty} [1; 1 + a_k, a_{k-1}, \dots]$$

where $a_i > 1$ for infinitely many i .

Now, assume that there exists an integer $I > 0$ such that $a_i = 1$ for $i \geq I$. The assumptions of Lemma 3.1 (1) and (2) are not satisfied for any level $k \geq I$. In other words, $\Lambda_1(\mathbf{x}) \cup \Lambda_2(\mathbf{x})$ is finite. Thus, it is sufficient to check (3)-(7). Note that $\liminf_{k \rightarrow \infty} \eta_k = [0; \bar{1}] = \varphi$. Using $1 \geq t_k$,

$$\lim_{k \rightarrow \infty} \epsilon_k = 0,$$

$$\liminf_{k \rightarrow \infty} \frac{1 + \eta_k - \epsilon_k}{1 + t_k + \eta_k - \epsilon_k} \geq \liminf_{k \rightarrow \infty} \frac{1 + \eta_k}{2 + \eta_k} = \frac{1 + \liminf_{k \rightarrow \infty} \eta_k}{2 + \liminf_{k \rightarrow \infty} \eta_k} = \varphi = \liminf_{k \rightarrow \infty} \eta_k,$$

(3)-(7) are similarly proved. Hence, $\text{rep}(\mathbf{x}) \geq [1; 2, \bar{1}]$ for a Sturmian word \mathbf{x} of slope φ . Therefore,

$$\text{rep}(\mathbf{x}) \geq \liminf_{k \rightarrow \infty} [1; 1 + a_k, a_{k-1}, \dots]$$

for a Sturmian word \mathbf{x} of slope θ .

The equality holds in the following setting. Choose the sequence $\{k_j\}$ such that $\eta_{k_j} \rightarrow \liminf_{k \rightarrow \infty} \eta_k$, $\lim_{j \rightarrow \infty} (k_{j+1} - k_j) = \infty$, and $k_{j+1} - k_j$ is odd for all j . Let $\mathbf{x} \in \mathcal{C}_{k_j}^{(ii)}$ for all k_j and $\mathbf{x} \in \mathcal{C}_{k_j+2l-1}^{(i)} \cap \mathcal{C}_{k_j+2l}^{(iii)}$ for all $0 < l \leq \frac{k_{j+1}-k_j-1}{2}$. Thus, $W_{k_j+1} = W_{k_j}M_{k_j-1}$ for all k_j and $W_{k+1} = W_k$ for all $k \neq k_j$. In the proof of (7), we have $\lim_{j \rightarrow \infty} t_{k_j} = 0$ and $\text{rep}(\mathbf{x}) = \underline{\lim}_{k \rightarrow \infty} [1; 1 + a_k, a_{k-1}, a_{k-2}, \dots]$. In conclusion, $\min \mathcal{L}(\theta) = \underline{\lim}_{k \rightarrow \infty} [1; 1 + a_k, a_{k-1}, a_{k-2}, \dots]$. \square

4. THE SPECTRUM OF THE EXPONENTS OF REPETITION OF FIBONACCI WORDS

In this section, we investigate $\mathcal{L}(\varphi)$ where $\varphi := \frac{\sqrt{5}-1}{2} = [0; \bar{1}]$. In what follows, assume that \mathbf{x} is a Sturmian word of slope φ , called a *Fibonacci word*. Note that $M_{k+1} = M_k^{a_{k+1}}M_{k-1} = M_kM_{k-1}$ for all $k \geq 1$. The following lemma is a special case of Lemma 3.1.

Lemma 4.1. Let $k \geq 1$. The following statements hold.

- (1) If $\mathbf{x} \in \mathcal{C}_k^{(i)}$, then $\mathbf{x} \in \mathcal{C}_{k+1}^{(iii)}$ and $W_{k+1} = W_k$.
- (2) If $\mathbf{x} \in \mathcal{C}_k^{(ii)}$, then $\mathbf{x} \in \mathcal{C}_{k+1}^{(i)} \cup \mathcal{C}_{k+1}^{(ii)}$ and $W_{k+1} = W_kM_{k-1}$.
- (3) If $\mathbf{x} \in \mathcal{C}_k^{(iii)}$, then $\mathbf{x} \in \mathcal{C}_{k+1}^{(i)} \cup \mathcal{C}_{k+1}^{(ii)}$ and $W_{k+1} = W_k$.

Proof. For $k \geq 1$, let W_k be the unique non-empty prefix of \mathbf{x} defined in which case \mathbf{x} belongs to at level k .

(1) Since $a_k = 1$ for all $k \geq 1$, the assumption of (1) in Lemma 3.1 cannot be satisfied. Hence, $\mathbf{x} \in \mathcal{C}_{k+1}^{(iii)}$. Since \mathbf{x} starts with $W_{k+1}M_{k+1}\widetilde{M}_{k+1} = W_{k+1}M_k\widetilde{M}_kD'_kM_{k+1}^-$ for the suffix W_{k+1} of M_k , $W_{k+1} = W_k$ by the uniqueness of W_k .

(2) and (3) are equivalent to (3) and (4) in Lemma 3.1 respectively. \square

By Lemma 4.1, only (iii) should follow (i) in the locating chain of \mathbf{x} : $\mathbf{x} \in \mathcal{C}_k^{(i)}$ implies $\mathbf{x} \in \mathcal{C}_{k+1}^{(iii)}$. Hence, if $\mathbf{x} \notin \mathcal{C}_1^{(iii)}$, then the locating chain of \mathbf{x} can be expressed as an infinite sequence of (i)(iii) and (ii). If $\mathbf{x} \in \mathcal{C}_1^{(iii)}$, then the locating chain of \mathbf{x} is an infinite sequence of (i)(iii) and (ii), except for the first letter (iii). Let us denote (i)(iii) and (ii) by a and b , respectively.

Since only (iii) should follow (i) in the locating chain of \mathbf{x} , the assumptions of Lemma 3.2 (1) and (2) cannot be satisfied. We have the following lemma corresponding to Lemma 3.2. Using

k	$k + 1$	The relation between W_{k+1} and W_k
case (i)	case (iii)	$W_{k+1} = W_k$
case (ii)	case (i) case (ii)	$W_{k+1} = W_k M_{k-1}$
case (iii)	case (i) case (ii)	$W_{k+1} = W_k$

TABLE 2. The relation between W_{k+1} and W_k

$q_{k+1} = q_k + q_{k-1}$ for all $k \geq 1$, Lemma 3.2 (3)-(6) are equivalent to (1)-(4) of the following lemma, respectively.

Lemma 4.2. Let $k \geq 1$.

- (1) If $\mathbf{x} \in \mathcal{C}_k^{(i)} \cap \mathcal{C}_{k+2}^{(i)}$, then $\Lambda(\mathbf{x}) \cap [q_{k+1} - 1, q_{k+3} - 2] = \{q_{k+1} + |W_k| - 1, q_{k+2} - 1\}$.
- (2) If $\mathbf{x} \in \mathcal{C}_k^{(i)} \cap \mathcal{C}_{k+2}^{(ii)}$, then $\Lambda(\mathbf{x}) \cap [q_{k+1} - 1, q_{k+3} - 2] = \{q_{k+1} + |W_k| - 1\}$.
- (3) If $\mathbf{x} \in \mathcal{C}_k^{(ii)} \cap \mathcal{C}_{k+1}^{(i)}$, then $\Lambda(\mathbf{x}) \cap [q_{k+1} - 1, q_{k+2} - 2] = \{q_{k+1} - 1\}$.
- (4) If $\mathbf{x} \in \mathcal{C}_k^{(ii)} \cap \mathcal{C}_{k+1}^{(ii)}$, then $\Lambda(\mathbf{x}) \cap [q_{k+1} - 1, q_{k+2} - 2] = \{q_{k+1} - 1, q_{k+1} + |W_k| - 1\}$.

For $l = 1, 2, 3, 4$, define

$$\Lambda'_l(\mathbf{x}) := \{n \in \Lambda(\mathbf{x}) : n \text{ appears in } (l) \text{ of Lemma 4.2}\}.$$

It is obvious that $\Lambda'_l(\mathbf{x})$'s are mutually distinct and $\cup_{l=1}^4 \Lambda'_l(\mathbf{x}) = \Lambda(\mathbf{x}) \cap [q_2 - 1, \infty) = \Lambda(\mathbf{x})$. From the definition of $\Lambda_l(\mathbf{x})$, $\Lambda'_l(\mathbf{x}) = \Lambda_{l+2}(\mathbf{x})$ for $l = 1, 2, 3, 4$ where the slope of \mathbf{x} is φ .

Note that $\text{rep}(\mathbf{x})$ is the limit infimum of $(1 + \frac{n_i}{n_{i+1}})$'s for $n_i \in \cup_{l=1}^4 \Lambda'_l(\mathbf{x})$. The following lemma says that it is enough to consider the elements of $\Lambda'_2(\mathbf{x})$ and $\Lambda'_3(\mathbf{x})$ to obtain $\text{rep}(\mathbf{x})$.

Lemma 4.3. Suppose that both a and b appear infinitely many in the locating chain of \mathbf{x} . Then,

$$\text{rep}(\mathbf{x}) = \liminf_{n_i \in \Lambda'_2(\mathbf{x}) \cup \Lambda'_3(\mathbf{x})} \left(1 + \frac{n_i}{n_{i+1}} \right).$$

Proof. First, for each k satisfying $\mathbf{x} \in \mathcal{C}_k^{(ii)} \cap \mathcal{C}_{k+1}^{(i)}$, there exists $d(k) > 0$ such that $\mathbf{x} \in \mathcal{C}_k^{(ii)} \cap \mathcal{C}_{k+2d(k)+1}^{(ii)}$ and $\mathbf{x} \in \mathcal{C}_{k+2d-1}^{(i)} \cap \mathcal{C}_{k+2d}^{(iii)}$ for $1 \leq d \leq d(k)$. By Lemma 4.2, $\mathcal{J}_k := \Lambda(\mathbf{x}) \cap [q_{k+1} - 1, q_{k+2d(k)+2} - 2] \subset \Lambda'_1(\mathbf{x}) \cup \Lambda'_2(\mathbf{x}) \cup \Lambda'_3(\mathbf{x})$. Note that $W_{k+1} = \dots = W_{k+2d(k)+1}$ and

$$\frac{q_{j+1} + |W_j| - 1}{q_{j+2} - 1} \geq \frac{q_{j+1} - 1}{q_{j+2} + |W_{j+1}| - 1}$$

for $j = k + 1, k + 3, \dots, k + 2d(k) - 1$. Since

$$\frac{q_{j+1} - 1}{q_{j+2} + |W_{j+1}| - 1}$$

is increasing for $j = k, k + 1, \dots, k + 2d(k) - 2$,

$$\min_{n_i \in \mathcal{J}_k} \frac{n_i}{n_{i+1}} = \min \left\{ \frac{q_{k+1} - 1}{q_{k+2} + |W_{k+1}| - 1}, \frac{q_{k+2d(k)} + |W_{k+2d(k)-1}| - 1}{q_{k+2d(k)+2} - 1} \right\} = \min_{n_i \in (\Lambda'_2(\mathbf{x}) \cup \Lambda'_3(\mathbf{x})) \cap \mathcal{J}_k} \frac{n_i}{n_{i+1}}.$$

Second, for each l satisfying $\mathbf{x} \in \mathcal{C}_l^{(\text{iii})} \cap \mathcal{C}_{l+1}^{(\text{ii})}$, there exists $d'(l) > 0$ such that $\mathbf{x} \in \mathcal{C}_l^{(\text{iii})} \cap \mathcal{C}_{l+d'(l)+1}^{(\text{i})}$ and $\mathbf{x} \in \mathcal{C}_j^{(\text{ii})}$ for $l+1 \leq j \leq l+d'(l)$. By Lemma 4.2, $\mathcal{J}_l' := \Lambda(\mathbf{x}) \cap [ql-1, ql+d'(l)+2-2] \subset \Lambda_2'(\mathbf{x}) \cup \Lambda_3'(\mathbf{x}) \cup \Lambda_4'(\mathbf{x})$. Note that

$$\frac{ql + |W_{l-1}| - 1}{ql+2-1} \leq \frac{ql+1-1}{ql+2-1} \text{ and } \frac{q_{j+1}-1}{q_{j+2}-1} \leq \frac{q_{j+1}-1}{q_{j+1} + |W_j| - 1}, \quad \frac{q_{j+1} + |W_j| - 1}{q_{j+2}-1}$$

for $j = l+1, \dots, l+d'(l)-1$. Since

$$\frac{q_j-1}{q_{j+1}-1}$$

is increasing for $j = l+1, \dots, l+d'(l)$,

$$\min_{n_i \in \mathcal{J}_l'} \frac{n_i}{n_{i+1}} = \min \left\{ \frac{ql + |W_{l-1}| - 1}{ql+2-1}, \frac{ql+d'(l)+1-1}{ql+d'(l)+2 + |W_{l+d'(l)+1}| - 1} \right\} = \min_{n_i \in (\Lambda_2'(\mathbf{x}) \cup \Lambda_3'(\mathbf{x})) \cap \mathcal{J}_l'} \frac{n_i}{n_{i+1}}.$$

Since $\Lambda(\mathbf{x})$ is the union of \mathcal{J}_k 's and \mathcal{J}_l' 's, $\text{rep}(\mathbf{x}) = \liminf_{n_i \in \Lambda_2'(\mathbf{x}) \cup \Lambda_3'(\mathbf{x})} \left(1 + \frac{n_i}{n_{i+1}}\right)$. \square

Let d be a positive integer. We define *a-chain* to be a subword $aa \cdots a$ in the locating chain of \mathbf{x} which b appears before and after. For example, if the locating chain of $\mathbf{x} = abbaaabaab \dots$, then *a-chains* are a, aaa, aa, \dots . Similarly, *b-chain* is defined by a subword $bb \cdots b$ in the locating chain of \mathbf{x} which a appears before and after. We say that an *a-chain* or a *b-chain* is a *chain*. From the definition of a chain, *a-chains* and *b-chains* alternatively appear in the locating chain of \mathbf{x} . We can choose two sequences $\{m_i(\mathbf{x})\}_{i \geq 1}$ and $\{l_j(\mathbf{x})\}_{j \geq 1}$ defined as follows: Let $m_i(\mathbf{x})$ (resp., $l_j(\mathbf{x})$) be the length of the i th *a-chain* (resp., the j th *b-chain*) in the locating chain of \mathbf{x} . Let $b^{l_1(\mathbf{x})}$ follow $a^{m_1(\mathbf{x})}$. In other words, the locating chain of \mathbf{x} is $c(\mathbf{x})a^{m_1(\mathbf{x})}b^{l_1(\mathbf{x})}a^{m_2(\mathbf{x})}b^{l_2(\mathbf{x})} \dots$ for the unique finite word $c(\mathbf{x})$. For example, if the locating chain of \mathbf{x} is (iii)bbabbabaabba \dots , then $c(\mathbf{x}) = (\text{iii})bb$, $m_1(\mathbf{x}) = 1$, $m_2(\mathbf{x}) = 1$, $m_3(\mathbf{x}) = 2$, $l_1(\mathbf{x}) = 2$, $l_2(\mathbf{x}) = 1$, $l_3(\mathbf{x}) = 3$. Let

$$S_d := \{\mathbf{x} : m_i(\mathbf{x}) \geq d \text{ for infinitely many } i \text{ or } l_j(\mathbf{x}) \geq d \text{ for infinitely many } j\}.$$

By definition, $S_{d+1} \subset S_d$. In what follows, we will write $m_i(\mathbf{x})$ and $l_j(\mathbf{x})$ simply m_i and l_j , when no confusion can arise. Now, let us prove Theorem 1.4, 1.5, 1.6 and Proposition 1.6.1.

Theorem 4.4. Let \mathbf{x} be a Sturmian word of slope φ . Then, $\mu_{\min} \leq \text{rep}(\mathbf{x}) \leq \mu_{\max}$. Moreover, the locating chain of \mathbf{x} is $u\bar{a}$ or $v\bar{b}$ for some finite words u, v if and only if $\text{rep}(\mathbf{x}) = \mu_{\max}$. We have $\mathbf{x} \in S_d$ for any $d \geq 1$ if and only if $\text{rep}(\mathbf{x}) = \mu_{\min}$.

Proof. First, assume that there exists a constant K such that $\mathbf{x} \in \mathcal{C}_k^{(\text{ii})}$ for all $k \geq K$. By Lemma 4.2, $\Lambda(\mathbf{x}) \cap [q_{K+1}-1, \infty) = \{q_k-1, q_k + |W_{k-1}| - 1 : k \geq K+1\}$. Since $W_{k+1} = W_k M_{k-1}$ for any $k \geq K$,

$$\begin{aligned} \liminf_{n_i \in \Lambda(\mathbf{x})} \frac{n_i}{n_{i+1}} &= \liminf_{k \geq K+1} \left\{ \frac{q_k-1}{q_k + |W_{k-1}| - 1}, \frac{q_k + |W_{k-1}| - 1}{q_{k+1}-1} \right\} \\ &= \min \left\{ \liminf_{k \geq K+1} \left(\frac{q_k-1}{q_k + |W_{k-1}| - 1} \right), \liminf_{k \geq K+1} \left(\frac{q_k + |W_{k-1}| - 1}{q_{k+1}-1} \right) \right\} = \min\{\varphi, 1\} = \varphi. \end{aligned}$$

Hence, $\text{rep}(\mathbf{x}) = \liminf_{n_i \in \Lambda(\mathbf{x})} \left(1 + \frac{n_i}{n_{i+1}}\right) = \mu_{\max}$.

Second, assume that there exists a constant K such that $\mathbf{x} \in \mathcal{C}_{K+2l}^{(i)} \cap \mathcal{C}_{K+2l+1}^{(iii)}$ for all $l \geq 0$. By Lemma 4.2, $\Lambda(\mathbf{x}) \cap [q_{K+1} - 1, \infty) = \{q_{K+2l+1} + |W_{K+2l}| - 1, q_{K+2l+2} - 1 : l \geq 0\}$. Since $|W_k|$ is constant for $k \geq K$,

$$\begin{aligned} \liminf_{n_i \in \Lambda(\mathbf{x})} \frac{n_i}{n_{i+1}} &= \liminf_{l \geq 0} \left\{ \frac{q_{K+2l+1} + |W_{K+2l}| - 1}{q_{K+2l+2} - 1}, \frac{q_{K+2l+2} - 1}{q_{K+2l+3} + |W_{K+2l+2}| - 1} \right\} \\ &= \min \left\{ \liminf_{l \geq 0} \left(\frac{q_{K+2l+1} + |W_{K+2l}| - 1}{q_{K+2l+2} - 1} \right), \liminf_{l \geq 1} \left(\frac{q_{K+2l} - 1}{q_{K+2l+1} + |W_{K+2l}| - 1} \right) \right\} = \varphi. \end{aligned}$$

Hence, $\text{rep}(\mathbf{x}) = \liminf_{n_i \in \Lambda(\mathbf{x})} \left(1 + \frac{n_i}{n_{i+1}} \right) = \mu_{max}$.

Now, let both a and b occur infinitely many in the locating chain of \mathbf{x} . Since ba appears infinitely many in the locating chain of \mathbf{x} , we can choose an infinite sequence $\{n_{i(j)}\}_{j \geq 1} \subset \Lambda'_3(\mathbf{x})$. For each $j \geq 1$, Lemma 4.2 gives $n_{i(j)} = q_{k+1} - 1, n_{i(j)+1} = q_{k+2} + |W_{k+1}| - 1$ for some $k = k(j)$. Note that $W_{k(j)+1} = W_{k(j)}M_{k(j)-1}$. By definition,

$$\begin{aligned} \text{rep}(\mathbf{x}) &\leq \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + |W_{k(j)+1}| - 1} \right) \\ &\leq \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + q_{k(j)-1} - 1} \right) = 1 + \frac{\varphi}{1 + \varphi^3} < \mu_{max}. \end{aligned}$$

Hence, $\text{rep}(\mathbf{x}) < \mu_{max}$. In other words, $\text{rep}(\mathbf{x}) = \mu_{max}$ implies that the locating chain of \mathbf{x} is $u\bar{a}$ or $v\bar{b}$ for some finite words u, v .

Finally, let us use Lemma 4.3 to find out the minimum of $\mathcal{L}(\varphi)$. For $n_i \in \Lambda'_2(\mathbf{x})$, let $n_i = q_k + |W_{k-1}| - 1, n_{i+1} = q_{k+2} - 1$ for some $k = k(i)$. For $n_i \in \Lambda'_3(\mathbf{x})$, let $n_i = q_{k'+1} - 1, n_{i+1} = q_{k'+2} + |W_{k'+1}| - 1$ for some $k' = k'(i)$. From Lemma 4.3,

$$\begin{aligned} \text{rep}(\mathbf{x}) &= \min \left\{ \liminf_{n_i \in \Lambda'_2(\mathbf{x})} \left(1 + \frac{n_i}{n_{i+1}} \right), \liminf_{n_i \in \Lambda'_3(\mathbf{x})} \left(1 + \frac{n_i}{n_{i+1}} \right) \right\} \\ &= \min \left\{ \liminf_{i \rightarrow \infty} \left(1 + \frac{q_{k(i)} + |W_{k(i)-1}| - 1}{q_{k(i)+2} - 1} \right), \liminf_{i \rightarrow \infty} \left(1 + \frac{q_{k'(i)+1} - 1}{q_{k'(i)+2} + |W_{k'(i)+1}| - 1} \right) \right\} \\ &\geq \min \left\{ \liminf_{i \rightarrow \infty} \left(1 + \frac{q_{k(i)} - 1}{q_{k(i)+2} - 1} \right), \liminf_{i \rightarrow \infty} \left(1 + \frac{q_{k'(i)+1} - 1}{q_{k'(i)+2} + q_{k'(i)+1} - 1} \right) \right\} \\ &= \min \left\{ 1 + \varphi^2, 1 + \frac{\varphi}{\varphi + 1} \right\} = \mu_{min}. \end{aligned}$$

Hence, $\text{rep}(\mathbf{x}) \geq \mu_{min}$. Moreover, if $\text{rep}(\mathbf{x}) = \mu_{min}$, then

$$\liminf_{i \rightarrow \infty} \frac{|W_{k(i)-1}|}{q_{k(i)}} = 0 \text{ or } \limsup_{i \rightarrow \infty} \frac{|W_{k'(i)+1}|}{q_{k'(i)+1}} = 1.$$

Thus, arbitrarily long sequence $aa \dots a$ or $bb \dots b$ should occur in the locating chain of \mathbf{x} , i.e. $\mathbf{x} \in S_d$ for any $d \geq 1$. \square

Theorem 4.5. The intervals $(\mu_2, \mu_{max}), (\mu_3, \mu_2)$ are maximal gaps in $\mathcal{L}(\varphi)$. Moreover, the locating chain of \mathbf{x} is $u\bar{a}b$ for some finite word u if and only if $\text{rep}(\mathbf{x}) = \mu_2$. The locating chain of \mathbf{x} is $v\bar{b}^2a^2$ for some finite word v if and only if $\text{rep}(\mathbf{x}) = \mu_3$.

Proof. If $\text{rep}(\mathbf{x}) < \mu_{max}$, then $\mathbf{x} \in S_1$ by Theorem 1.4.

First, let $\mathbf{x} \in S_1 \cap S_2^c$. Since any chains of length greater than 1 occur at most finitely many in the locating chain of \mathbf{x} , there exists an integer $I > 0$ satisfying $m_j = l_j = 1$ for $j \geq I$. Thus, the locating chain of \mathbf{x} is $u\bar{a}\bar{b}$ for some finite word u . Using Lemma 4.2 and 4.3, we obtain $\text{rep}(\mathbf{x}) = \mu_2$.

Second, let $\mathbf{x} \in S_2$. Using Lemma 4.2 and 4.3, $\text{rep}(\mathbf{x}) = \mu_3$ where the locating chain of \mathbf{x} is $\overline{va^2b^2}$ for some finite word v . If $m_j \geq 3$ for infinitely many j , then there exists an infinite sequence $\{k(j)\}$ satisfying $\mathbf{x} \in \mathcal{C}_{k(j)-6}^{(i)} \cap \mathcal{C}_{k(j)-4}^{(i)} \cap \mathcal{C}_{k(j)-2}^{(i)} \cap \mathcal{C}_{k(j)}^{(ii)}$ for all j . By Lemma 4.3,

$$\text{rep}(\mathbf{x}) \leq \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)-1} + |W_{k(j)-6}| - 1}{q_{k(j)+1} - 1} \right) \leq 1 + \varphi^2 + \varphi^7$$

where $W_{k(j)} = W_{k(j)-6}$ for all j . If $l_j \geq 3$ for infinitely many j , then there exists an infinite sequence $\{k(j)\}$ satisfying $\mathbf{x} \in \mathcal{C}_{k(j)-2}^{(ii)} \cap \mathcal{C}_{k(j)-1}^{(ii)} \cap \mathcal{C}_{k(j)}^{(ii)} \cap \mathcal{C}_{k(j)+1}^{(i)}$ for all j . By Lemma 4.3,

$$\begin{aligned} \text{rep}(\mathbf{x}) &\leq \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + |W_{k(j)+1}| - 1} \right) \\ &= \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + q_{k(j)-1} + q_{k(j)-2} + q_{k(j)-3} + |W_{k(j)-2}| - 1} \right) \\ &\leq \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + 2q_{k(j)-1} - 1} \right) = 1 + \frac{\varphi}{1 + 2\varphi^3}. \end{aligned}$$

Since $1 + \varphi^2 + \varphi^7, 1 + \frac{\varphi}{1 + 2\varphi^3} < \mu_3$, $\text{rep}(\mathbf{x}) < \mu_3$ for $\mathbf{x} \in S_3$. Now, let $\mathbf{x} \in S_2 \cap S_3^c$. By definition, there exists an integer $I > 0$ such that $m_j, l_j \leq 2$ for $j \geq I$. If $l_j = 1, m_{j+1} = 2$ for infinitely many j , then there exists an infinite sequence $\{k(j)\}$ satisfying $\mathbf{x} \in \mathcal{C}_{k(j)-7}^{(i)} \cap \mathcal{C}_{k(j)-5}^{(ii)} \cap \mathcal{C}_{k(j)-4}^{(i)} \cap \mathcal{C}_{k(j)-2}^{(i)} \cap \mathcal{C}_{k(j)}^{(ii)}$ for all j . By Lemma 4.3,

$$\text{rep}(\mathbf{x}) \leq \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)-1} + q_{k(j)-6} + |W_{k(j)-7}| - 1}{q_{k(j)+1} - 1} \right) \leq 1 + \varphi^2 + \varphi^7 + \varphi^8$$

where $W_{k(j)} = W_{k(j)-4}, W_{k(j)-5} = W_{k(j)-7}$, and $W_{k(j)-4} = W_{k(j)-5}M_{k(j)-6}$ for all j . If $m_j = 1, l_j = 2$ for infinitely many j , then there exists an infinite sequence $\{k(j)\}$ satisfying $\mathbf{x} \in \mathcal{C}_{k(j)-4}^{(ii)} \cap \mathcal{C}_{k(j)-3}^{(i)} \cap \mathcal{C}_{k(j)-1}^{(ii)} \cap \mathcal{C}_{k(j)}^{(ii)} \cap \mathcal{C}_{k(j)+1}^{(i)}$ for all j . By Lemma 4.3,

$$\begin{aligned} \text{rep}(\mathbf{x}) &\leq \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + |W_{k(j)+1}| - 1} \right) \\ &= \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + q_{k(j)-1} + q_{k(j)-2} + |W_{k(j)-3}| - 1} \right) \\ &\leq \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + q_{k(j)} + q_{k(j)-5} - 1} \right) = 1 + \frac{\varphi}{\varphi^7 + \varphi^2 + 1} \end{aligned}$$

where $W_{k(j)-1} = W_{k(j)-3}$ and $W_{k(j)-3} = W_{k(j)-4}M_{k(j)-5}$ for all j . Since $1 + \varphi^2 + \varphi^7 + \varphi^8, 1 + \frac{\varphi}{\varphi^7 + \varphi^2 + 1} < \mu_3$, $\text{rep}(\mathbf{x}) < \mu_3$ where the locating chain of \mathbf{x} is not $\overline{va^2b^2}$ for some finite word v . Hence, $\max\{\text{rep}(\mathbf{x}) : \mathbf{x} \in S_2\} = \mu_3$. Moreover, $\text{rep}(\mathbf{x}) = \mu_3$ if and only if the locating chain

of \mathbf{x} is $\overline{va^2b^2}$ for some finite word v . Therefore, two intervals (μ_2, μ_{max}) , (μ_3, μ_2) are maximal gaps in $\mathcal{L}(\varphi)$. On the other hand, by Theorem 1.4, $\text{rep}(\mathbf{x}) = \mu_{max}$ if and only if $\mathbf{x} \in S_1^c$. In the proof above, $\text{rep}(\mathbf{x}) = \mu_2$ for $\mathbf{x} \in S_1 \cap S_2^c$, and $\text{rep}(\mathbf{x}) \leq \mu_3$ for $\mathbf{x} \in S_2$. Hence, $\text{rep}(\mathbf{x}) = \mu_2$ if and only if $\mathbf{x} \in S_1 \cap S_2^c$. \square

Theorem 4.6. The interval (μ_4, μ_3) is a maximal gap in $\mathcal{L}(\varphi)$. Moreover, $\text{rep}(\mathbf{x}) = \mu_4$ if and only if $\mathbf{x} \in S_2 \cap S_3^c$ satisfies the following two conditions:

- 1) The locating chain of \mathbf{x} is $u(b^2a^2)^{e_1}ba(b^2a^2)^{e_2}ba\dots$ for some finite word u and integers $e_i \geq 1$.
- 2) $\limsup_{i \geq 1} \{e_i\} = \infty$.

Furthermore, μ_4 is a limit point of $\mathcal{L}(\varphi)$.

Proof. From Lemma 4.2 and 4.3, $\text{rep}(\mathbf{x}) = \mu_4$ if $x \in S_2 \cap S_3^c$ satisfies the above two conditions 1) and 2). Assume that a Sturmian word \mathbf{x} satisfies $\text{rep}(\mathbf{x}) \in (\mu_4, \mu_3)$. In the proof of Theorem 1.5, $\text{rep}(\mathbf{x}) \geq \mu_2$ for $\mathbf{x} \in S_2^c$, and $\text{rep}(\mathbf{x}) \leq \min\{1 + \varphi^2 + \varphi^7, 1 + \frac{\varphi}{2\varphi^3 + 1}\} < \mu_4$ for $\mathbf{x} \in S_3$. Thus, $\mathbf{x} \in S_2 \cap S_3^c$. By definition, there exists an integer $I > 0$ such that $m_j, l_j \leq 2$ for all $j \geq I$. Moreover, $\text{rep}(\mathbf{x}) < \mu_3$ implies that $\{j : m_j = 1\} \cup \{j : l_j = 1\}$ is infinite. Hence, $\{j : m_j = 1, l_j = 2\} \cup \{j : l_j = 1, m_{j+1} = 2\}$ is infinite.

First, if $\{j : l_j = 1, m_{j+1} = 2\}$ is infinite, then $\text{rep}(\mathbf{x}) < 1.432 < \mu_4$. Thus, $\{j : l_j = 1, m_{j+1} = 2\}$ is finite. In other words, there exists an integer $I' > 0$ such that $m_j = 2$ implies $l_{j-1} = 2$ for all $j > I'$. Since $\{j : m_j = 1, l_j = 2\} \cup \{j : l_j = 1, m_{j+1} = 2\}$ is infinite, $\{j : m_j = 1, l_j = 2\}$ is infinite. Now, let us show that both $\{j : m_j = 1, l_{j-1} = l_j = 2\}$ and $\{j : m_{j-1} = m_j = 1, l_{j-1} = 1, l_j = 2\}$ are finite. If $\{j : m_j = 1, l_{j-1} = l_j = 2\}$ is infinite, then there exists an infinite sequence $\{k(j)\}$ such that $\mathbf{x} \in \mathcal{C}_{k(j)-5}^{(ii)} \cap \mathcal{C}_{k(j)-4}^{(ii)} \cap \mathcal{C}_{k(j)-3}^{(i)} \cap \mathcal{C}_{k(j)-1}^{(ii)} \cap \mathcal{C}_{k(j)}^{(ii)} \cap \mathcal{C}_{k(j)+1}^{(i)}$ for all j . Hence,

$$\begin{aligned} \text{rep}(\mathbf{x}) &\leq \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + |W_{k(j)+1}| - 1} \right) \\ &= \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + q_{k(j)-1} + q_{k(j)-2} + q_{k(j)-5} + q_{k(j)-6} + |W_{k(j)-5}| - 1} \right) \\ &\leq \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + q_{k(j)-1} + q_{k(j)-2} + q_{k(j)-5} + q_{k(j)-6} - 1} \right) \\ &= 1 + \frac{\varphi}{1 + \varphi^3 + \varphi^4 + \varphi^7 + \varphi^8} < \mu_4. \end{aligned}$$

It follows that $\{j : m_j = 1, l_{j-1} = l_j = 2\}$ is finite.

If $\{j : m_{j-1} = m_j = 1, l_{j-1} = 1, l_j = 2\}$ is infinite, then there exists an infinite sequence $\{k(j)\}$ such that $\mathbf{x} \in \mathcal{C}_{k(j)-7}^{(ii)} \cap \mathcal{C}_{k(j)-6}^{(i)} \cap \mathcal{C}_{k(j)-4}^{(ii)} \cap \mathcal{C}_{k(j)-3}^{(i)} \cap \mathcal{C}_{k(j)-1}^{(ii)} \cap \mathcal{C}_{k(j)}^{(ii)} \cap \mathcal{C}_{k(j)+1}^{(i)}$ for all j . Hence,

$$\begin{aligned} \text{rep}(\mathbf{x}) &\leq \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + |W_{k(j)+1}| - 1} \right) \\ &= \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + q_{k(j)-1} + q_{k(j)-2} + q_{k(j)-5} + q_{k(j)-8} + |W_{k(j)-7}| - 1} \right) \end{aligned}$$

$$\begin{aligned} &\leq \liminf_{j \rightarrow \infty} \left(1 + \frac{q_{k(j)+1} - 1}{q_{k(j)+2} + q_{k(j)-1} + q_{k(j)-2} + q_{k(j)-5} + q_{k(j)-8} - 1} \right) \\ &= 1 + \frac{\varphi}{1 + \varphi^3 + \varphi^4 + \varphi^7 + \varphi^{10}} < \mu_4. \end{aligned}$$

It follows that $\{j : m_{j-1} = m_j = 1, l_{j-1} = 1, l_j = 2\}$ is finite. Therefore, both $\{j : m_j = 1, l_{j-1} = l_j = 2\}$ and $\{j : m_{j-1} = m_j = 1, l_{j-1} = 1, l_j = 2\}$ are finite.

Next, let us prove that $\{j : m_j = m_{j+1} = 1\}$ is finite. Suppose that $\{j : m_j = m_{j+1} = 1\}$ is infinite. Then, $\{j : m_j = m_{j+1} = 1, l_j = 1\} \cup \{j : m_j = m_{j+1} = 1, l_j = 2\}$ is infinite. Note that $\{j : l_j = 1, m_{j+1} = 2\}$ is finite. In other words, there exists an integer $I'' > 0$ such that $l_j = 1$ implies $m_{j+1} = 1$ for all $j > I''$. Thus, if $\{j : m_j = m_{j+1} = 1, l_j = 1\}$ is infinite, then $\{j : m_j = m_{j+1} = 1, l_j = 1, l_{j+1} = 2\} \cup \{j : m_j = m_{j+1} = m_{j+2} = 1, l_j = l_{j+1} = 1\}$ is infinite. Since $\{j : m_j = m_{j+1} = 1, l_j = 1, l_{j+1} = 2\}$ is finite, $\{j : m_j = m_{j+1} = m_{j+2} = 1, l_j = l_{j+1} = 1\}$ is infinite. Thus, $\{j : m_j = m_{j+1} = m_{j+2} = 1, l_j = l_{j+1} = 1, l_{j+2} = 2\} \cup \{j : m_j = m_{j+1} = m_{j+2} = m_{j+3} = 1, l_j = l_{j+1} = l_{j+2} = 1\}$ is infinite. Since $\{j : m_j = m_{j+1} = 1, l_j = 1, l_{j+1} = 2\}$ is finite, $\{j : m_j = m_{j+1} = m_{j+2} = m_{j+3} = 1, l_j = l_{j+1} = l_{j+2} = 1\}$ is infinite. By the same argument, it follows that the locating chain of \mathbf{x} is $u\bar{a}b$ for some finite word u , which leads a contradiction with $x \in S_2$. Hence, $\{j : m_j = m_{j+1} = 1, l_j = 1\}$ is finite. Similarly, we use the same argument to induce that $\{j : m_j = m_{j+1} = 1, l_j = 2\}$ is finite. Therefore, $\{j : m_j = m_{j+1} = 1\}$ is finite.

From the above arguments, we have the locating chain of \mathbf{x} is $u(b^2a^2)^{e_1}b^{f_1}a(b^2a^2)^{e_2}b^{f_2}a\dots$ for integers $e_i \geq 1$, $f_j = 1$ or 2 , and some finite word u . Since $\{j : m_j = 1, l_{j-1} = l_j = 2\}$ is finite, we can assume that $f_j = 1$ for all j . The locating chain of \mathbf{x} is $u(b^2a^2)^{e_1}ba(b^2a^2)^{e_2}ba\dots$. Moreover, if $d = \limsup_{i \geq 1} \{e_i\} < \infty$, then

$$\text{rep}(\mathbf{x}) \leq 1 + \left(\frac{1}{\varphi} + \varphi + \frac{\varphi^6}{1 - \varphi^{6d+3}} + \frac{\varphi^{10}}{1 - \varphi^6} \frac{1 - \varphi^{6d}}{1 - \varphi^{6d+3}} \right)^{-1} < \mu_4.$$

It follows $\limsup_{i \geq 1} \{e_i\} = \infty$. However, for a Sturmian word \mathbf{x} such that the locating chain of \mathbf{x} is $u(b^2a^2)^{e_1}ba(b^2a^2)^{e_2}ba\dots$ and $\limsup_{i \geq 1} \{e_i\} = \infty$, $\text{rep}(\mathbf{x}) = \mu_4$. It implies that there does not exist a Sturmian word \mathbf{x} satisfying $\text{rep}(\mathbf{x}) \in (\mu_4, \mu_3)$. Hence, (μ_4, μ_3) is a maximal gap in $\mathcal{L}(\varphi)$. Furthermore,

$$\begin{aligned} \text{rep}(\dots \overline{(b^2a^2)^d ba}) &= 1 + \left(\frac{1}{\varphi} + \varphi + \frac{\varphi^6}{1 - \varphi^{6d+3}} + \frac{\varphi^{10}}{1 - \varphi^6} \frac{1 - \varphi^{6d}}{1 - \varphi^{6d+3}} \right)^{-1} \\ &\rightarrow \mu_4 \text{ as } d \rightarrow \infty. \end{aligned}$$

Hence, μ_4 is a limit point of $\mathcal{L}(\theta)$. □

Proposition 4.6.1. For $\alpha \in \{\mu_{max}, \mu_2, \mu_3\}$, there are only countably many Sturmian words \mathbf{x} of slope φ satisfying $\text{rep}(\mathbf{x}) = \alpha$. For $\beta \in \{\mu_4, \mu_{min}\}$, there are uncountably many Sturmian words \mathbf{x} of slope φ satisfying $\text{rep}(\mathbf{x}) = \beta$.

Proof. From Theorem 1.4, $\text{rep}(\mathbf{x}) = \mu_{max}$ if and only if the locating chain of \mathbf{x} is $u\bar{a}$ or $v\bar{b}$ for some finite words u, v . Thus, \mathbf{x} satisfying $\text{rep}(\mathbf{x}) = \mu_{max}$ is completely determined by the choice of u or v . Hence, there exist only countably many Sturmian words \mathbf{x} of slope φ with $\text{rep}(\mathbf{x}) = \mu_{max}$. Theorem 1.4 also implies that $\text{rep}(\mathbf{x}) = \mu_{min}$ if and only if $\mathbf{x} \in S_d$ for any $d \geq 1$. Hence, it is possible to choose 2^{\aleph_0} sequences $\{m_i\} \cup \{l_j\}$. Namely, there exist uncountably many Sturmian words \mathbf{x} of slope φ with $\text{rep}(\mathbf{x}) = \mu_{min}$.

On the other hand, Theorem 1.5 implies that $\text{rep}(\mathbf{x}) = \mu_2$ if and only if the locating chain of \mathbf{x} is $u\bar{b}a$ for some finite word u . Thus, \mathbf{x} satisfying $\text{rep}(\mathbf{x}) = \mu_2$ is completely determined by the choice of u . Hence, there exist only countably many Sturmian words \mathbf{x} of slope φ with $\text{rep}(\mathbf{x}) = \mu_2$. Theorem 1.5 also implies that $\text{rep}(\mathbf{x}) = \mu_3$ if and only if the locating chain of \mathbf{x} is $u\overline{b^2a^2}$ for some finite word u . Thus, \mathbf{x} satisfying $\text{rep}(\mathbf{x}) = \mu_3$ is completely determined by the choice of u . Hence, there exist only countably many Sturmian words \mathbf{x} of slope φ with $\text{rep}(\mathbf{x}) = \mu_3$.

Finally, Theorem 1.6 implies that $\text{rep}(\mathbf{x}) = \mu_4$ if and only if the locating chain of \mathbf{x} is $u(b^2a^2)^{e_1}ba(b^2a^2)^{e_2}ba\dots$ for some finite word u and integers $e_i \geq 1$ satisfying $\limsup_{i \geq 1} \{e_i\} = \infty$. Hence, it is possible to choose 2^{\aleph_0} sequences $\{e_i\}$. Namely, there exist uncountably many Sturmian words \mathbf{x} of slope φ with $\text{rep}(\mathbf{x}) = \mu_4$. \square

ACKNOWLEDGEMENT

The author wishes to thank Dong Han Kim and Seonhee Lim for introducing me to the subject and very helpful comments.

REFERENCES

- [1] B. Adamczewski and Y. Bugeaud, *Nombres réels de complexité sous-linéaire: mesures d'irrationalité et de transcendance*, Journal für die reine und angewandte Mathematik **658** (2011), 65–98.
- [2] B. Adamczewski and T. Rivoal, *Irrationality measures for some automatic real numbers*, Mathematical Proceedings of The Cambridge Philosophical Society **147** (2009), no. 3, 659–678.
- [3] S. Akiyama and H. Kaneko, *Multiplicative analogue of Markoff-Lagrange spectrum and Pisot numbers*, Advances in Mathematics **380** (2021), no. 107547.
- [4] P. Alessandri and V. Berthé, *Three distance theorems and combinatorics on words*, Enseignement Mathématique **44** (1998), 103–132.
- [5] V. Berthé and L. Vuillon, *A two-dimensional generalization of Sturmian sequences: tilings and rotations* **223** (2000), no. 1-3, 27–53.
- [6] P. E. Böhmer, *Über die Transzendenz gewisser dyadischer Brüche*, Math. Ann. **96** (1927), 367–377.
- [7] Y. Bugeaud, *Distribution modulo one and Diophantine approximation*, Cambridge Tracts in Mathematics, vol. 193, Cambridge University Press, 2012.
- [8] Y. Bugeaud and D. H. Kim, *A new complexity function, repetitions in Sturmian words, and irrationality exponents of Sturmian numbers*, Transactions of the American Mathematical Society **371** (2019), no. 5, 3281–3308.
- [9] Y. Bugeaud, D. H. Kim, M. Laurent, and A. Nogueira, *On the Diophantine nature of the elements of Cantor sets arising in the dynamics of contracted rotations*, Annali della Scuola Normale Superiore di Pisa, Classe di Scienze. available at arxiv.org/abs/2001.00380.

- [10] Y. Bugeaud and M. Laurent, *Combinatorial structure of Sturmian words and continued fraction expansions of Sturmian numbers*. available at arxiv.org/abs/2104.09239.
- [11] J. Cassaigne, *Sequences with grouped factors*, *Developments in language theory* **3** (1997), 211–222.
- [12] T. W. Cusick and M. E. Flahive, *The Markoff and Lagrange spectra*, *Mathematical Surveys and Monographs*, vol. 30, American Mathematical Society, Providence, RI, 1989.
- [13] S. Ferenczi, *Dynamical generalizations of the Lagrange spectrum*, *Journal d'Analyse Mathématique* **118** (2012), no. 1, 19–53.
- [14] R. Kenyon, *The construction of self-similar tilings*, *Geometric & Functional Analysis* **6** (1996), no. 3, 471–488.
- [15] M. Lothaire, *Algebraic combinatorics on words*, Vol. 90, Cambridge university press, 2002.
- [16] M. Morse and G. A. Hedlund, *Symbolic dynamics II. Sturmian trajectories*, *American Journal of Mathematics* **62** (1940), no. 1, 1–42.
- [17] S. Ohnaka and T. Watanabe, *A gap of the exponents of repetitions of Sturmian words*. available at arxiv.org/abs/2003.10721.