


## A VECTOR IDENTITY FOR QUADRILATERALS

CHRISTIAN AEBI AND GRANT CAIRNS

Consider a quadrilateral  $ABCD$  in the plane. We let  $K_{ABCD}$  denote its area and we use obvious similar notation for the area of various triangles. Consider the position vectors  $\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}$  of the vertices. Then one has the following result, which one might call the *Jacobi identity for quadrilaterals*.

**Theorem.**  $K_{BCD} \mathbf{A} - K_{ACD} \mathbf{B} + K_{ABD} \mathbf{C} - K_{ABC} \mathbf{D} = 0$ .



*Proof.* First notice that the identity is unchanged by translation. Indeed, for a translation by a vector  $\mathbf{X}$ , the areas are unchanged and

$$\begin{aligned} K_{BCD}(\mathbf{A} + \mathbf{X}) - K_{ACD}(\mathbf{B} + \mathbf{X}) + K_{ABD}(\mathbf{C} + \mathbf{X}) - K_{ABC}(\mathbf{D} + \mathbf{X}) \\ = K_{BCD} \mathbf{A} - K_{ACD} \mathbf{B} + K_{ABD} \mathbf{C} - K_{ABC} \mathbf{D}, \end{aligned}$$

since  $K_{BCD} + K_{ABD} = K_{ABCD} = K_{ACD} + K_{ABC}$ . So we may take  $D$  to be the origin,  $O$ . Then our desired identity is

$$K_{BCO} \mathbf{A} - K_{ACO} \mathbf{B} + K_{ABO} \mathbf{C} = 0.$$

Observe that the vector cross product  $\mathbf{B} \times \mathbf{C}$  has magnitude  $2K_{BCO}$  and direction  $\mathbf{k}$  perpendicular to the plane. Further,  $\mathbf{k} \times \mathbf{A}$  is the vector obtained from  $\mathbf{A}$  by rotating it through angle  $\frac{\pi}{2}$  in the positive direction; let us denote this vector  $\mathbf{A}'$ . Similarly, we define  $\mathbf{B}'$  and  $\mathbf{C}'$ . Now, rotating the vector  $K_{BCO} \mathbf{A} - K_{ACO} \mathbf{B} + K_{ABO} \mathbf{C}$  through  $\frac{\pi}{2}$ , and multiplying by 2, we obtain

$$\begin{aligned} 2(K_{BCO} \mathbf{A}' - K_{ACO} \mathbf{B}' + K_{ABO} \mathbf{C}') \\ = (\mathbf{B} \times \mathbf{C}) \times \mathbf{A} - (\mathbf{A} \times \mathbf{C}) \times \mathbf{B} + (\mathbf{A} \times \mathbf{B}) \times \mathbf{C} = 0, \end{aligned}$$

by the Jacobi vector triple product identity. Hence  $K_{BCO} \mathbf{A} - K_{ACO} \mathbf{B} + K_{ABO} \mathbf{C} = 0$ , as desired.  $\square$

It remains for us to clarify what we mean by “quadrilateral”. In fact, the word does not have an entirely universal meaning; see [2]. However, the astute reader

will have noticed that the above proof doesn't require any property of  $ABCD$ ; it is actually true for every choice of 4 points  $A, B, C, D$  in the plane, provided one interprets the numbers  $K_{ABC}$  etc as *signed areas*; that is,  $K_{ABC}$  is positive if  $A, B, C$  are positively oriented, and negative otherwise.

For more on the wonderful world of quadrilaterals, see the delightful book by Alsina and Nelsen [1].

#### REFERENCES

1. Claudi Alsina and Roger B. Nelsen, *A Cornucopia of quadrilaterals*, Mathematical Association of America, Dolciani Mathematical Expositions, 2020.
2. Zalman Usiskin and Jennifer Griffin, *The classification of quadrilaterals : a study of definition*, Research in Mathematics Education, IAP, Charlotte, 2008.

COLLÈGE CALVIN, GENEVA, SWITZERLAND 1211  
*Email address:* christian.aebi@edu.ge.ch

DEPARTMENT OF MATHEMATICS, LA TROBE UNIVERSITY, MELBOURNE, AUSTRALIA 3086  
*Email address:* G.Cairns@latrobe.edu.au