

## Graph space: using both geometric and probabilistic structure to evaluate statistical graph models

LOUIS DUVIVIER

*Univ Lyon, INSA Lyon, CNRS, LIRIS UMR5205, F-69621 France.*  
louis.duvivier@insa-lyon.fr

AND

RÉMY CAZABET

*Univ Lyon, Université Lyon 1, CNRS, LIRIS UMR5205, F-69622 France.*  
remy.cazabet@univ-lyon1.fr

AND

CÉLINE ROBARDET

*Univ Lyon, INSA Lyon, CNRS, LIRIS UMR5205, F-69621 France.*  
celine.robardet@insa-lyon.fr

[Received on 14 September 2021]

Statistical graph models aim at modeling graphs as random realization among a set of possible graphs. One issue is to evaluate whether or not a graph is likely to have been generated by one particular model. In this paper we introduce the edit distance expected value (EDEV) and compare it with other methods such as entropy and distance to the barycenter. We show that contrary to them, EDEV is able to distinguish between graphs that have a typical structure with respect to a model, and those that do not. Finally we introduce a statistical hypothesis testing methodology based on this distance to evaluate the relevance of a candidate model with respect to an observed graph.

*Keywords:* statistical graph model, graph ensemble, entropy, statistical hypothesis testing, edit distance

### 1. Introduction

The study of large and complex systems in domains as diverse as physics, biology, computer science or social science brings forward interaction networks which are neither regular, nor random. The interaction structure follows no clear pattern, as in a grid for example, but it often presents remarkable properties such as a heterogeneous degree distribution (scale-free) [1], small average distances (the small-world phenomenon) [16], a high level of transitivity and community structure [11], which are not present in random graphs. This suggests that interactions are the result of a random process following some constraints which shape the resulting network.

Various statistical models have been proposed to describe such constrained random graphs: the configuration model, which preserves degree distribution; the stochastic blockmodel, which preserves local densities; spatial models, which focus on distance between nodes [15]. A review of statistical graph models can be found in [6] and an introduction to their formalism in [4].

This diversity of parametric models raises the issue of model selection: considering one specific graph, which model is the most relevant? And with which set of parameters? Considering models as

ensembles makes it possible to leverage results from statistical physics and information theory to perform model selection [5], in particular using the minimum description length [7]. In this framework, a model is considered to be good for a given graph if it generates it with high probability, *i.e.* if the associated ensemble has a small entropy. Therefore, many papers have focused on computing the entropy of various graph ensemble [3], [13], [18]. A whole methodology for community detection based on those principles has been developed in the case of stochastic blockmodels [14]. Yet, by considering models as unstructured sets, this approach neglects graphs topology.

Indeed, graphs are also geometrical objects, in the sense that one can define distances between them. Such a distance induces a structure on a model's ensemble. Much work has been devoted to quantifying how similar two graphs are [17], especially from a topological point of view [10], [9]. These distances between two graphs can be generalized to evaluate the quality of a model by computing a distance between an observed graph and the graph ensemble associated to a model. For example, the widely used measure for community detection known as modularity [12] evaluate the quality of a partition by comparing the edge weight in the observed graph with the expected edge weight of the graph in the configuration model, as developed in section 2.2. In this case, the problem is to evaluate the statistical significance of the results, in order not to mistake noise for structure [8].

In this paper, after reviewing in section 2.1 and 2.2 existing techniques to measure the relevance of a model with respect to a graph, we introduce in section 3 the edit distance expected value, a measure which takes into account both the geometric and the probabilistic structure of the graph ensemble. Finally, we show how this measure can be used to evaluate a model relevance with respect to a given graph in section 4.

## 2. Microcanonical ensemble

The main point of statistical modeling is to describe the structure of a graph  $G$  based on a set of global properties of that graph  $P_M$  [4]. Some examples of common models and their associated properties are given in table 1. Apart from those properties, the graph is considered to be random. In practice, this means that  $G$  is considered to have been chosen from the set:

$$\Omega_M = \{H \mid P_M(H) = P_M(G)\}$$

This set is called the microcanonical ensemble, in reference to statistical physics in which it was first introduced to represent all the possible states of a system corresponding to a global property. It can be defined with directed or undirected graphs, weighted or not, with or without self-loops. In the rest of the article, we will consider labelled directed multigraphs with self-loops. Although they are not the most widely used in practice, it makes probability derivations easier, especially for the configuration model.

Table 1. Common statistical graph models and their associated properties

Model $M$	Properties $P_M$
Erdős-Renyi	number of nodes and number of edges
Configuration model	degree distribution
Stochastic block model	block to block density
Gravity model	node position, strength, and deterrence function
Radiation model	node position and strength

## 2.1 Entropy

Information theory shows that, in order to ensure that no additional information bias the results, the probability distribution  $\mathbb{P}$  on  $\Omega_M$  has to be the one that maximises Shannon's entropy  $S = -\sum_{H \in \Omega_M} \mathbb{P}(H) \log(\mathbb{P}(H))$ . With no additional constraint, this optimal distribution is simply the uniform one:

$$\forall H \in \Omega_M, \mathbb{P}(H) = \frac{1}{|\Omega_M|}$$

whose entropy is  $\log(|\Omega_M|)$ . Thus, computing the probability to choose  $G$  among all possible graphs in  $\Omega_M$  boils down to counting the number of graphs it contains. This has been done for different models in [13].

The microcanonical ensemble and its entropy provide a common formulation for various statistical models. This is useful to perform model selection. Indeed, entropy is directly related to the likelihood of a given model. If we observe a graph  $G$  and consider a set of models  $\mathcal{M} = \{M_1, \dots, M_p\}$ , we can find which model  $G$  has most likely been sampled from by maximising its likelihood

$$M^* = \operatorname{argmax}_{M_i \in \mathcal{M}} \mathbb{P}[M_i|G]$$

This maximisation can be done using Bayes theorem

$$\mathbb{P}[M_i|G] = \frac{\mathbb{P}[G|M_i] \times \mathbb{P}[M_i]}{\mathbb{P}[G]}$$

If we assume a non-informative uniform prior distribution on all models  $\mathbb{P}[M_i] = \frac{1}{p}$ , as  $\mathbb{P}[G]$  is a constant, maximising the likelihood is equivalent to maximising  $\mathbb{P}[G|M_i] = \frac{1}{|\Omega_{M_i}|}$  or minimising the entropy  $\log(|\Omega_{M_i}|)$ . This idea is developed and applied to the case of stochastic blockmodels in [14].

Considering the common situation in which one is interested in modeling the global topology of  $G$  rather than its precise edge list, likelihood maximisation appears insufficient as a model selection criterion. For example, if we consider the three graphs on figure 1, with  $G_1$  as a reference, both  $G_2$  and  $G_3$  are different from  $G_1$ , but the topology of  $G_1$  and  $G_2$  is almost the same. Therefore, a model which produces mostly  $G_3$ -like graphs cannot be considered as good a model for  $G_1$  as one which produces  $G_2$ -like ones.

## 2.2 Distance to the barycenter

On the other hand, one may measure how typical  $G$  is with respect to  $\Omega_M$  by comparing it with an appropriate representative of this ensemble, for example its barycenter:

$$G_M = \sum_{H \in \Omega_M} \mathbb{P}(H)H$$

If we denote  $W_G$  the weight matrix of graph  $G$ , it can easily be derived that

$$\forall (i, j) \in V^2, W_{G_M}(i, j) = \mathbb{E}[W_H(i, j)] \quad (2.1)$$

REMARK 2.1  $G_M$  does not necessarily belong to  $\Omega_M$ . In particular, even if all graphs in  $\Omega_M$  have whole weight, it does not imply that  $G_M$ 's edge weights are integers. Examples of barycenter weights for common models are given in table 2




FIG. 1. **Three graphs with 8 nodes and 10 edges.**  $G_1$  and  $G_2$  (leftmost) share the same topology even if they are not strictly equal, which is not the case of  $G_3$  (rightmost).

Table 2. Statistical graph models' barycenter weight

Model $M$	Parameters	$G_M(i, j)$	$m_{G_M}$
Erdős-Renyi	$n, m$	$\frac{m}{n^2}$	$m$
Configuration model	$(k_i^{in}, k_i^{out})_{i \in V}$	$\frac{k_i^{out} k_j^{in}}{m}$	$m$
Stochastic block model	$(m_{r,s})_{r,s}$	$\frac{m_{c_i, c_j}}{ c_i   c_j }$	$m$
Gravity model	$(k_i)_{i \in V}, f$	$k_i k_j f(d(i, j))$	$m$
Radiation model	$(k_i^{in}, k_i^{out})_{i \in V}$	$\frac{k_i^{out} k_i^{in} k_j^{in}}{(k_i^{in} + s_{ij})(k_j^{in} + s_{ij})}$	$m$

The famous modularity function to evaluate the quality of a node partition  $B = (b_1, \dots, b_p)$  on a graph  $G = (V, E)$  with weight matrix  $W_G$  is defined as the difference between the number of edges inside each cluster and the expected number for a random graph with the same degree distribution (*i.e.* following the configuration model). It can be understood as a comparison with the barycenter  $G_M$  of the corresponding configuration model.

$$\begin{aligned}
 Q(G, B) &= \frac{1}{2m} \sum_{i=1}^p \sum_{u, v \in b_i} \left( W_G(u, v) - \frac{k_u^{out} k_v^{in}}{m} \right) \\
 &= \frac{1}{2m} \sum_{i=1}^p \sum_{u, v \in b_i} (W_G(u, v) - W_{G_M}(i, j)) \\
 &= d(G, G_M)
 \end{aligned}$$

A problem is that  $G$  is compared with a single graph  $G_M$  which is supposed to account for the whole graph ensemble  $\Omega_M$ . In particular, all information about the dispersion around the barycenter is lost, which undermines any attempt to interpret statistically the results.

### 3. Graph space and the edit distance expected value

As we have seen, existing techniques to compare a graph  $G$  and a model  $M$  exploit in different ways the ensemble  $\Omega_M$ . Entropy based techniques described in section 2.1 focus on its cardinal, but they neglect the topological similarities of graphs inside the ensemble. On the other hand, as described in section 2.2, an objective function such as the modularity accounts for these similarities, but it does so with a single graph which is supposed to represent the whole set. Reality is more complex:  $\Omega_M$  is a set of graphs with a probability distribution, and it can be further structured with a metric, making it a graph space. Both aspects, probabilistic and geometric, should be taken into account in order to understand the structure of  $\Omega_M$ , and the plausibility that a graph  $G$  was generated by the associated model  $M$ .

Many different measures exist to compute a similarity score between two graphs  $G$  and  $H$  [17]. One of the simplest is the edit distance. For two graphs on the same vertex sets  $G_1 = (V, E_1)$  and  $G_2 = (V, E_2)$ , it counts the number of differences between their respective sets of edges.

$$\text{ed}(G_1, G_2) = \sum_{(i,j) \in V^2} |W_{G_1}(i, j) - W_{G_2}(i, j)|$$

As expected from its name, edit distance is a distance between graphs. Indeed, if we consider the weight matrix  $W_G$  of a graph  $G$  as a point in  $\mathbb{R}^{n^2}$ , the edit distance corresponds to the  $\mathcal{L}_1$  distance and for any model  $M$ ,  $\Omega_M$  is a subset of  $\mathbb{R}^{n^2}$ . The dimension prevents any direct drawing of it for graphs with more than 2 nodes, but it is possible to obtain some intuition about its shape.

In the following, we will use a normalized version of the edit distance which can be interpreted as the fraction of different edges between  $G_1$  and  $G_2$ .

$$\text{ned}(G_1, G_2) = \frac{1}{2m} \sum_{(i,j) \in V^2} |W_{G_1}(i, j) - W_{G_2}(i, j)|$$

This normalized edit distance is no longer a distance on  $\mathbb{R}^{n^2}$ . Yet, for all models  $M$  considered here, the number of edges  $m$  is constant over the set  $\Omega_M$ . Thus, the normalized edit distance is equivalent to edit distance inside  $\Omega_M$  and it allows to compare more easily results between various models, because whatever the model  $M$ , the distance between any two graphs  $G_1$  and  $G_2$  in  $\Omega_M$  is at most 1.

#### 3.1 Edit distance to the barycenter

$\Omega_M$  barycenter has already been introduced in section 2.2, where it was used as a proxy for the whole space. Using normalized edit distance, it is possible to check how much graphs in  $\Omega_M$  are similar to the barycenter  $G_M$ . We consider six different models:

1. EM: Erdős-Renyi with 50 nodes and 1000 edges
2. CFM cst: configuration model with 50 nodes and a constant degree distribution ( $k_i^{\text{in}} = k_i^{\text{out}} = 20$ )
3. CFM arith: configuration model with 50 nodes and an arithmetic degree distribution ( $k_i^{\text{in}} = k_i^{\text{out}} = i + 1$ )
4. SBM hom: stochastic block model with 50 nodes and 5 communities, each having internal density 1.2, and external density 0.2.
5. SBM het: stochastic block model with 50 nodes and 5 communities, with internal density 0.4, 0.8, 1.2, 1.6, 2, and external density 0.2.

For each model  $M$ , we pick a random sample  $\mathcal{S}_M$  of 100 graphs in  $\Omega_M$  and for all  $G \in \mathcal{S}_M$  we compute the normalized edit distance to the barycenter  $\text{ned}(G, G_M)$ . Results are shown in figure 2.




FIG. 2. **Edit distance to the barycenter for 6 different models.** For each model  $M$  in ordinate, we draw 100 graphs  $G$  at random from  $\Omega_M$  and compute for each of them  $\text{ned}(G, G_M)$ . The distribution of results is then plotted as a boxplot.

The first thing to underline is that whatever the model,  $\text{ned}(G, G_M)$  is greater than 0.5, which means that most graphs in  $\Omega_M$  have at most half of their edges in common with  $G_M$ . This observation shows that for those models, the graph space is not concentrated around its barycenter. On the contrary, most graphs in  $\Omega_M$  seem to be at a specific distance from its barycenter, as would happen for a sphere with a radius depending on the model: 0.67 for ER and CFM cst, 0.55 for CFM arith, 0.69 for SBM hom and 0.71 for SBM het.

All models were chosen to have similar entropy, as shown in table 3, yet their characteristic distance to the barycenter vary greatly. Furthermore, we observe that these quantities are not positively correlated: CFM arith, which is the model with the larger entropy is also the one which is the most concentrated around its barycenter. This means that even if this model can generate a higher number of different graphs, the graphs it produces tend to be more similar one to the other than for other models. This is logical as this model preserves a degree distribution, which enforces more constraints on edges' distribution than an Erdős-Renyi or a stochastic block model.

Model	Characteristic distance	Entropy
ER	0.67	2050
CFM cst	0.67	2500
CFM arith	0.55	3300
SBM hom	0.69	1840
SBM het	0.71	1840

Table 3. **Edit distance to the barycenter and entropy.**

This concentration of graphs at a specific distance from the barycenter is a consequence of the dimensionality of the vector space. Let's denote  $\mathcal{B}(G, r)$  the ball of center  $G$  and radius  $r$  in  $(\mathbb{R}^{n^2}, \text{ed})$ .

We consider the set

$$\begin{aligned}\Omega_M(r) &= \{G \in \Omega_M \mid \text{ned}(G, G_M) \leq r\} \\ &= \{G \in \Omega_M \mid \text{ed}(G, G_M) \leq 2mr\} \\ &= \Omega_M \cap \mathcal{B}(G_M, 2mr)\end{aligned}$$

The volume  $V_n(r)$  of  $\mathcal{B}(G_M, 2mr)$  is proportional to  $r^{n^2}$ , therefore

$$\forall r < 1, \frac{V_n(r)}{V_n(1)} \xrightarrow{n \rightarrow \infty} 0 \quad (3.1)$$

The volume of the ball concentrates quickly at its periphery as the dimension increases, and so does the volume of  $\Omega_M$ . The additional constraints on  $\Omega_M$  modify its shape in such a way that graphs too far away from the barycenter are rare, which explains why the concentration does not happen at distance 1 from the barycenter. Still, this phenomenon is strong enough to imply that even graphs generated according to a model  $M$  will share only a relatively small fraction of their edges with the barycenter of the model.

### 3.2 Edit distance expected value

The previous observations on the structure of graph spaces show that in order to compare a graph  $G$  with a model  $M$ , one should consider more than the mere cardinal of  $\Omega_M$ . One possibility to evaluate how similar to  $G$  are the graphs in  $\Omega_M$  is to compute the expected value of the normalized edit distance:

$$\text{EDEV}(G, M) = \mathbb{E}_{H \in \Omega_M} \left[ \frac{1}{2m} \sum_{(i,j) \in V^2} |W_G(i, j) - W_H(i, j)| \right]$$

To illustrate how EDEV provides further information on the place of  $G$  within the graph space, we compare it with entropy for different synthetic graphs. A low value indicates that  $G$  is close to other graphs in  $\Omega_M$ , and thus that it is typical of the model, while a high value shows that it is an outlier. As a case study, we consider the Erdős-Renyi model. Let's recall that we consider multigraphs, which implies that we allow for densities rising above 1. The extension of Erdős-Renyi model to multigraphs is straightforward,  $\Omega_{ER(n,m)}$  contains all multigraphs with  $n$  nodes and  $m$  edges and each multigraph is generated with the same probability  $\frac{1}{|\Omega_{ER(n,m)}|}$ . In practice, we consider models with  $n = 100$  nodes and a number of edges  $m$  ranging from 100 to 500000. For each, we consider three graphs:

- $G_1(m)$ , picked uniformly at random inside  $\Omega_{ER(n,m)}$
- $G_2(m)$ , a graph made of two equal communities, each with  $\frac{n}{2}$  nodes and  $\frac{m}{2}$  edges, perfectly separated.
- $G_3(m)$ , the graph where all edges are between nodes 0 and 1.

Results are shown on figure 3.

For each value of  $m$ , all three graphs belong to the graph ensemble  $\Omega_{ER(n,m)}$ . We observe that as density increases  $|\Omega_{ER(n,m)}|$  grows exponentially, which implies that the probability to pick at random




FIG. 3. **Entropy and edit distance expected value against density.** For each density, three graphs are generated.  $G_1(m)$  is random,  $G_2(m)$  is made of two random communities and  $G_3(m)$  has its edges concentrated on a single pair of nodes of weight  $m$ . On the top plot, the entropy  $\log(|\Omega_{ER(n,m)}|)$  is plotted against the density  $\frac{m}{n^2}$ . As all graphs belong to the same graph ensemble  $\Omega_{ER(n,m)}$ , the three curves are the same. On the bottom plot the edit distance expected value  $EDEV(G(m), \Omega_{ER(n,m)})$  is plotted against density.

$G_1(m)$ ,  $G_2(m)$  or  $G_3(m)$  becomes even less probable. Yet, in the case of the random graph  $G_1(m)$  this is counter-intuitive: as density grows and becomes higher than 1, most graph in  $\Omega_{ER(n,m)}$  become complete graphs with each edge having weight about  $\frac{m}{n^2}$ . This is the case of  $G_1$  too with a high probability, so ER( $n, m$ ) is very likely to produce graphs similar to  $G_1(m)$ , even if it is very unlikely to produce  $G_1(m)$  itself.

On the other hand, edit distance expected value is able to capture this phenomenon. While it is close to 1 for all three types of graphs when density is low because in this situation a random model can hardly predict correctly which edge is present in any graph, it decreases quickly towards 0 when density rises above 0.1 for  $G_1(m)$ . For  $G_2(m)$  we have an intermediate situation: edit distance decreases too, but it reaches its minimum around 0.5, indicating that even when it is densely populated, the model is only able to reproduce correctly half of its edges. This is normal as  $G_2(m)$  concentrates them inside the communities, which means on half of all possible node pairs. These observations are actually a particular case of a more general result, which can be stated as:

**THEOREM 3.1** Let  $B_1$  and  $B_2$  be two partition on  $\llbracket 1, n \rrbracket$ , with  $p_1$  and  $p_2$  blocks respectively. Let

$M_1 \in \mathcal{M}_{p_1}(\mathbb{N})$  and  $M_2 \in \mathcal{M}_{p_2}(\mathbb{N})$  be two block adjacency matrices such that

$$\sum_{i,j \in [1,p_1]^2} M_1[i,j] = \sum_{k,l \in [1,p_2]^2} M_2[k,l] = m$$

Let's consider two series of stochastic blockmodels defined as  $S_1(k) = (B_1, k \cdot M_1)$  and  $S_2(k) = (B_2, k \cdot M_2)$ , whose barycenters are denoted  $G_1(k)$  and  $G_2(k)$ . We have that

1. There exists  $d \in \mathbb{R}, \forall k \in \mathbb{N}, \text{ed}(G_1(k), G_2(k)) = d$
2. Let  $(G_k)_{k \in \mathbb{N}}$  be a series of random graphs, each drawn following model  $S_1(k)$ .

$$\text{EDEV}(G_k, S_2(k)) \xrightarrow[k \rightarrow \infty]{\mathbb{P}} d \quad (3.2)$$

(proof in Appendix)

REMARK 3.1 In particular, if  $M_1 = M_2$ , this theorem means that the normalized edit distance expected value between a graph picked at random and the barycenter of the stochastic blockmodel converges toward 0:  $\Omega_{S(k)}$  shrinks around  $G_{S(k)}$ . This is what we observe with  $G_1(m)$ . Yet, we also observe on figure 3 that the normalized edit distance converges toward 0 only as density rises above 1. Thus, in practice, the vast majority of graphs are too sparse for this assumption to hold true and most graphs in  $\Omega_{S(k)}$  are far from  $G_{S(k)}$ , as developed in section 3.1

#### 4. Model likelihood

As the distance to the barycenter, the expected value of the normalized edit distance is characteristic of a model. For a model  $M$ , the values of  $\text{EDEV}(H, M)$  for graphs  $H$  in  $\Omega_M$  are concentrated around a specific value  $d_M$ . We can use this fact to rule out models which fit badly on an observed graph  $G$ .

For example, let's consider the configuration model  $CFMD(n, k_i^{out}, k_i^{in})$

$$n = 50 \\ \forall i \in \llbracket 0, n-1 \rrbracket, k_i^{out} = k_i^{in} = i$$

We use this model to generate a graph  $G_i$ .  $CFMD(n, k_i^{out}, k_i^{in})$  will thus be called the *generative* model, and  $G_i$  the *observed* graph. On this observed graph, we test the stochastic blockmodel  $SBM_i$  obtained by partitioning its nodes in two blocks:  $B_0$  contains even nodes and  $B_1$  odd nodes (this way we avoid to put all high-degree nodes in the same block) and learning the block adjacency matrix on  $G_i$ . We call  $SBM_i$  the *candidate* model. We generate a sample  $\mathcal{S}_i$  of 100 test graphs with the candidate model  $SBM_i$  and compare the normalized edit distance  $\text{EDEV}(G_i, \Omega_{SBM_i})$  of the observed graph to the candidate model with  $\text{EDEV}(H, \Omega_{SBM_i})$  for all test graphs  $H \in \mathcal{S}_i$ . This experiment is performed 5 times, and results are shown on figure 4.

We observe that for all five experiments, the normalized edit distance expected value to the candidate model  $SBM_i$  for the observed graph  $G_i$  is around 0.74, while for the test graphs generated by  $SBM_i$  it is between 0.53 and 0.56. This shows that the normalized edit distance expected value to  $SBM_i$  is significantly different for the observed graphs, which are generated by  $CFMD(n, k_i^{out}, k_i^{in})$ , and for the




FIG. 4. 5 graphs  $G_i$  were generated with a generative model  $CFMD(n, k_i^{out}, k_i^{in})$ . Their normalized edit distance expected value with respect to a candidate model  $SBM_i$  is plotted as diamond. As a comparison point, the distribution of the normalized edit distance expected value for 100 test graphs generated with  $SBM_i$  is plotted as dots and boxplot.




FIG. 5. 5 graphs  $G_i$  were generated with a generative model  $SBM(n, B, M)$ . Their normalized edit distance expected value with respect to a test model  $CFMD_i$  is plotted as diamond. As a comparison point, the distribution of the normalized edit distance expected value for 100 test graphs generated with  $CFMD_i$  is plotted as dots and boxplot.

test graphs generated by  $SBM_i$ . It is thus very unlikely that the observed graph  $G_i$  was generated by the candidate model  $SBM_i$ .

We then perform the same experiment the other way round, by considering as generative model a stochastic blockmodel  $SBM_0(n, B, M)$  defined by

$$\begin{aligned} n &= 50 \\ B &= [[0, 24], [25, 49]] \\ M &= \begin{bmatrix} 500 & 0 \\ 0 & 500 \end{bmatrix} \end{aligned}$$

5 graphs  $G_i$  are generated using this stochastic blockmodel. As candidate model, we consider a configuration model  $CFMD_i$  obtained by learning the degree sequence of  $G_i$ . A sample  $\mathcal{S}'$  of 100 test graphs is randomly picked in  $\Omega_{CFMD_i}$  and we compare  $EDEV(G_i, \Omega_{CFMD_i})$  with  $EDEV(H, \Omega_{CFMD_i})$  for all  $H \in \mathcal{S}'$ . Results are shown on figure 5. Once again, we observe that the normalized edit distance expected value to the candidate model  $CFMD_i$  is significantly different for the observed graphs, which were generated by  $SBM(n, B, M)$ , and for the test graphs generated by  $CFMD_i$ . This allows us to reject the hypothesis that the observed graph  $G_i$  was generated by the candidate model  $CFMD_i$ .

#### 4.1 Statistical hypothesis testing

This methodology can be formalized using statistical hypothesis testing. Let's say we have a graph  $G$  and a model  $M$  (possibly obtained by fitting some parameters on  $G$ ). We want to test the hypothesis  $\mathcal{H}$ : "the observed graph  $G$  has been generated by the candidate model  $M$ ". We do so in the following way:

1. Choose a confidence level  $\delta$ , for example 0.01.
2. Generate a random sample from the candidate model  $\mathcal{S} \subset \Omega_M$ , of  $n_b$  test graphs.
3. Infer the probability distribution  $\mathbb{P}_{ed}$  of  $EDEV(H, \Omega_M)$  for  $H \in \Omega_M$  from the sample  $\mathcal{S}$ .
4. Compute the probability

$$p = \mathbb{P}_{ed}[|EDEV(H, \Omega_M) - d_M| \geq |EDEV(G, \Omega_M) - d_M|]$$

5. If  $p < \delta$ , it means that the probability that  $G$  was generated by  $M$  is inferior to  $\delta$  and the hypothesis  $\mathcal{H}$  can be rejected.

In practice, to infer the probability distribution  $\mathbb{P}_{ed}$ , we assume that this distribution is normal, so we only need to infer the mean  $d_M$  and the standard deviation  $\sigma_M$  of the distribution. This assumption is verified on the models described in the previous subsection, by comparing the cumulative distribution function of the sample with the one of the corresponding normal distribution. An example is shown in figure 6. The shapiro-wilk test returned a p-value below 0.05 (around 0.01) for 2 of the 10 models. This means that the normal distribution can only be considered a rough estimate in general, but we found it good enough as a first approximation.




FIG. 6. Comparison between the empirical distribution of the edit distance expected value for graphs generated with a configuration model ( $CFMD_0$ ) and the normal distribution with the same mean and standard deviation.




FIG. 7. Spatial graph generated using the waxman random geometric model. 100 nodes are randomly distributed in a  $[0, 1] \times [0, 1]$  square. They are then connected with a probability depending on the distance  $d$  between nodes:  $p(d) = 10 \exp\left(\frac{d}{0.05L}\right)$ , with  $L$  the maximum distance between two nodes. Communities are computed using graph tools<sup>2</sup>.

<sup>2</sup><https://graph-tool.skewed.de/>

Let's consider a situation in which one wishes to evaluate the relevance of the block structure computed on a graph  $G$ , with a weight matrix  $W_G$ . Whatever the graph, and the partition  $B = (b_1, \dots, b_p)$  of its  $n$  nodes, it is always possible to define  $M \in \mathcal{M}_p(\mathbb{N})$  as

$$\forall i, j \in \llbracket 1, p \rrbracket, M[i, j] = \sum_{u \in b_i, v \in b_j} W_G[u, v]$$

such that  $G \in \Omega_{SBM(n, B, M)}$ . The objective is to evaluate whether this stochastic blockmodel is a relevant model of  $G$ . An even trickier question is to evaluate whether any stochastic blockmodel can be a relevant model. In particular, spatial models can generate graphs with groups of nodes densely connected due to their position rather than to block membership. An example of such a graph is shown in figure 7. It may then be hard to tell whether the blocks found are indeed a legitimate model of the observed graph or should be considered as artefacts, consequences of the underlying spatial structure.

To illustrate how statistical hypothesis testing allows to address this issue, we consider eight models: four stochastic blockmodels and four waxman model for random geometric graphs<sup>3</sup> with different sets of parameters. The waxman model for spatial graphs allows to easily control the strength of the spatial structure, by tuning the speed at which edge probability decays as the distance between nodes rises. The number of nodes is fixed to  $n = 100$  and the parameters are fixed such as to ensure a density  $d$  around 0.036. All stochastic blockmodels use a node partition in four blocks of 25 nodes, with a block adjacency matrix of the form:

$$\begin{bmatrix} m_{int} & m_{ext} & m_{ext} & m_{ext} \\ m_{ext} & m_{int} & m_{ext} & m_{ext} \\ m_{ext} & m_{ext} & m_{int} & m_{ext} \\ m_{ext} & m_{ext} & m_{ext} & m_{int} \end{bmatrix}$$

The four stochastic blockmodels are then defined by:

1.  $M_0$ :  $m_{int} = 90, m_{ext} = 0$
2.  $M_1$ :  $m_{int} = 75, m_{ext} = 5$
3.  $M_2$ :  $m_{int} = 60, m_{ext} = 10$
4.  $M_3$ :  $m_{int} = 45, m_{ext} = 15$

This way, the graphs generated using  $SBM_0$  are made of perfectly separated blocks of nodes, while those generated by  $SBM_3$  have blocks with as many internal and external edges.

For the waxman models, we also consider four parameter sets:

1.  $M_4$ :  $\alpha = 0.1, \beta = 1$
2.  $M_5$ :  $\alpha = 0.08, \beta = 1.6$
3.  $M_6$ :  $\alpha = 0.06, \beta = 2.7$
4.  $M_7$ :  $\alpha = 0.04, \beta = 8.5$

<sup>3</sup>[https://networkx.org/documentation/stable/reference/generated/networkx.generators.geometric.waxman\\_graph.html](https://networkx.org/documentation/stable/reference/generated/networkx.generators.geometric.waxman_graph.html)




FIG. 8. For each model  $M_i$  in ordinate, and each graph number  $i$  in absciss, we compute the probability  $p_{i,j}$  that a graph generated using  $SBM_{i,j}$  has a normalized edit distance expected value to  $SBM_{i,j}$  further away from the mean value  $d_{i,j}$  than  $G_{i,j}$ . The probability is plotted in heatmap.

The lower the value of  $\alpha$ , the stronger the spatial structure.

With each model  $M_i$ , we generate 8 graphs  $(G_{i,j})_{j \in \llbracket 0,7 \rrbracket}$ . On each of those observed graph, we find the minimum entropy node partition  $B_{i,j}$  using graph tools, and fit a candidate stochastic blockmodel  $SBM_{i,j}$  on  $G_{i,j}$  based on this node partition. We then evaluate the relevance of this stochastic blockmodel using the previously described methodology. We use a confidence level  $\delta$  of 0.01 and a sample size  $n_b = 200$ . The probabilities  $p_{i,j}$  obtained are plotted in figure 8.

We observe that for all graphs generated by stochastic block models but one,  $p_{i,j} > 10^{-2} = \delta$ . On the other hand, for all spatial graphs  $p_{i,j} < \delta$ . This means that the hypothesis "  $G_{i,j}$  has been generated by the candidate stochastic block model  $SBM_{i,j}$  " is rejected for all 32 spatial graphs, and for the 8<sup>th</sup> graph generated by  $SBM_2$ . Let's stress again that a probability  $p_{i,j}$  superior to  $\delta$  does not mean that  $SBM_{i,j}$  is the right model for  $G_{i,j}$ . It only means that there is not enough statistical evidence to reject it.

These results show that, on 32 spatial graphs generated with various sets of parameters, the statistical hypothesis testing methodology is able to correctly identify that the block structure found is not a relevant model, despite being the solution maximizing the likelihood. This result is not trivial, as illustrated by the graph depicted on figure 7. What is more, this methodology manages to reject the block structure for all spatial graphs while spuriously rejecting it for only one out of 32 graphs originally generated with a stochastic blockmodel. In other words, there is no false negative, and only one false positive.

Strictly speaking, these results only allow to rule out one node partition. As the block structure tested were fitted on the observed graphs using minimum entropy, one could argue that ruling out this partition implies that no other node partition can lead to a relevant model. However, for most real graphs, there is

more than one plausible node partition and minimizing the entropy of a partition is a stochastic process. Therefore, the experiment should be performed more than once to conclude that the observed graph has no block structure.

## 5. Conclusion

As a conclusion, we have seen that widely used quality measures for graph models rely either on the number of different graphs they can produce, which neglects the geometric structure of the graph space, or on a direct comparison with the barycenter of those graphs, which discards information about the distribution around this barycenter. Because of these restrictions, they are unable to distinguish between graphs which have a typical structure of a model and graphs which may be generated by this model but as outliers.

In this paper, we show how edit distance can provide additional information on the structure of graphs generated by a model, which is captured neither by the entropy nor the barycenter. By computing the expected value of the normalized edit distance for a given graph, we obtain a criterion which can be used to evaluate the model quality with respect to this graph. Finally, we incorporate this criterion to a statistical hypothesis testing methodology to perform model selection.

Graph space is a theoretical framework which can be used for any statistical model, and particularly spatial models. It allows to compare them with SBM or configuration model, and perform model selection between models of different nature. What is more, statistical hypothesis testing provides a statistically rigorous methodology to evaluate the relevance of a candidate model to an observed graph.

Any graph property could be used for such a test: average path length, clustering coefficient, etc. The main requirement for such a property to be used in statistical testing is that the distribution of its possible values for a given model is concentrated around its mean. If it is not, no value measured on the observed graph will allow to reject the candidate model.

Apart from its simplicity to compute and interpret, one important advantage of the normalized edit distance expected value is that for most models its values concentrate quickly around the mean. It should be underlined that this fundamental property is a consequence of a geometrical result (the volume of a ball in  $n$  dimensions), which highlights the benefits of considering the geometric structure of graph ensembles. However, the edit distance is not the only distance that can be used. Considering other metrics which are more sensitive to the global topology of the network, like the perturbation-resistance distance or spectral distances, could provide additional insight on the structure of the graph space.

## 6. Appendix

### 6.1 Barycenter graph weight of various statistical models

We have defined the barycenter of a graph model as

$$G_M = \sum_{H \in \Omega_M} \mathbb{P}(H) \cdot H$$

Which means that

$$\begin{aligned} \forall (i, j) \in V^2, W_{G_M}(i, j) &= \sum_{H \in \Omega_M} \mathbb{P}(H) \times W_H(i, j) \\ &= \mathbb{E}[W_H(i, j)] \end{aligned}$$

Let's illustrate how this can be computed for some classical models.

**ERDŐS-RÉNYI MODEL** The simplest graph model is the Erdős-Rényi model for random graphs. It's associated microcanonical ensemble can be defined as:

$$\Omega_{ER(n,m)} = \{H = (V, E) \mid |V| = n \wedge |E| = m\}$$

Let's recall that for the sake of simplicity, we chose to consider multigraphs with self loops. Thus, the computation of  $\mathbb{E}[W_H(i, j)]$  is particularly simple. Indeed, if for each pair of node  $(i, j) \in V^2$  and each  $k \in [1, m]$  we define the random variable  $X_{i,j,k}$  which is equal to 1 if the  $k^{\text{th}}$  edge is  $i \rightarrow j$  and to 0 else, then we have that  $W_H(i, j) = \sum_{k=1}^m X_{i,j,k}$ . It is a sum of independent Bernouillis' random variable so it follows a binomial law of parameters  $m$  and  $\frac{1}{n^2}$ , and thus

$$W_{G_{ER(n,m)}}(i, j) = \mathbb{E}_{\Omega_{ER}} [W_H(i, j)] = \frac{m}{n^2} \quad (6.1)$$

**CONFIGURATION MODEL** For the configuration model, all graphs in the microcanonical ensemble must have the same degree distribution. Let's consider the directed version.

$$\Omega_{CFMD} = \{G \mid \forall i \in V, \deg_G^{\text{out}}(i) = k_i^{\text{out}} \wedge \deg_G^{\text{in}}(i) = k_i^{\text{in}}\}$$

To compute the weight of the barycenter graph's edges, we consider that each node  $i$  has  $k_i^{\text{out}}$  outgoing stubs and  $k_i^{\text{in}}$  ingoing stubs. Any graph in  $\Omega_{CFMD}$  is characterized by a configuration of connections of outgoing stubs with ingoing stubs. For every pair of nodes  $i, j \in V^2$  and any pair of stub  $k \in [1, k_i^{\text{out}}]$ ,  $l \in [1, k_j^{\text{in}}]$ , we define the random variable  $X_{i,j,k,l}$  which is equal to 1 if the  $k^{\text{th}}$  outgoing stub of  $i$  is connected to the  $l^{\text{th}}$  ingoing stub of  $j$ , and to 0 otherwise. Then

$$W_H(i, j) = \sum_{k=1}^{k_i^{\text{out}}} \sum_{l=1}^{k_j^{\text{in}}} X_{i,j,k,l}$$

As each outgoing stub of  $i$  has the same probability to be connected to any of the  $m$  ingoing stubs

$$\forall i, j, k, l, \mathbb{P}[X_{i,j,k,l} = 1] = \frac{1}{m}$$

Thus,  $W_H(i, j)$  follows a binomial law of parameters  $\frac{1}{m}$  and  $k_i^{\text{out}} \times k_j^{\text{in}}$ . Finally

$$W_{G_{CFMD}}(i, j) = \mathbb{E}_{\Omega_{CFMD}} [W_H(i, j)] = \frac{k_i^{\text{out}} \times k_j^{\text{in}}}{m} \quad (6.2)$$

**STOCHASTIC BLOCKMODEL** The case of the stochastic blockmodel can be treated in the same way as erdős-rényi. It is defined, considering a partition of the nodes  $B = (b_1, \dots, b_q)$  and a block adjacency matrix  $M \in \mathcal{M}_q(\mathbb{N})$  by

$$\Omega_{SBM} = \left\{ H \mid \forall b_k, b_l, \sum_{i \in b_k} \sum_{j \in b_l} W_H(i, j) = M(k, l) \right\}$$

So, for any pair of nodes  $i \in b_k, j \in b_l$ ,  $W_H(i, j)$  follows a binomial law of parameters  $(M(k, l), |b_k||b_l|)$ . Thus

$$W_{G_{SBM}}(i, j) = \mathbb{E}_{\Omega_{SBM}} [W_H(i, j)] = \frac{M(k, l)}{|b_k||b_l|} \quad (6.3)$$

**SPATIAL MODELS** References for the gravitational model and the radiation model can be found in [2] and [15]. In both cases, they are constructed in such a way that edges weight have a given expected value. In the case of the gravitational model, it is

$$W_{G_{\text{grav}}}(i, j) = f(d(i, j)) \times k_i^{\text{out}} \times k_j^{\text{in}} \quad (6.4)$$

where  $d(i, j)$  is the distance from node  $i$  to node  $j$ , and  $f$  is a deterence function.

Finally, in the case of the radiation model, it is

$$W_{G_{\text{rad}}}(i, j) = \frac{k_i^{\text{out}} \times k_i^{\text{in}} \times k_j^{\text{in}}}{(k_i^{\text{in}} + s_{ij}) \times (k_i^{\text{in}} + k_j^{\text{in}} + s_{ij})} \quad (6.5)$$

with  $s_{ij} = \sum_{u \in \mathcal{C}(i, j)} k_u^{\text{in}}$  and  $\mathcal{C}(i, j) = \{u \in V \mid 0 < d(i, u) < d(i, j)\}$ .

## 6.2 Convergence proof

First of all, let's prove the following lemma

LEMMA 6.1 Let  $B$  be a partition of  $\llbracket 1, n \rrbracket$  with  $p$  blocks. Let  $M \in \mathcal{M}_p(\mathbb{N})$  be a block adjacency matrix. For all  $k \in \mathbb{N}$ , we define the stochastic blockmodel  $S(k) = (B, k \cdot M)$ , and its barycenter  $G_{S(k)}$ . We consider a sequence of random graphs  $(G_k)_{k \in \mathbb{N}}$ , each drawn from  $S(k)$ . We have that

$$\text{ed}(G_k, G_{S(k)}) \xrightarrow[k \rightarrow \infty]{\mathbb{P}} 0$$

Given the notation above, we want to prove that:

$$\forall \alpha > 0, \mathbb{P}[\text{ed}(G_k, G_{S(k)}) > \alpha] \xrightarrow[k \rightarrow \infty]{} 0$$

Let  $\alpha > 0$ . Let's denote  $m = \sum_{i,j} M_{i,j}$  the number of edges of graphs in  $\Omega_{S(1)}$ . By definition,

$$\text{ed}(G_k, G_{S(k)}) = \frac{1}{2km} \sum_{u,v} |W_{G_k}(u,v) - W_{G_{S(k)}}(u,v)|$$

Thus,

$$\text{ed}(G_k, G_{S(k)}) > \alpha \Rightarrow \exists (u,v), \left| \frac{W_{G_k}(u,v) - W_{G_{S(k)}}(u,v)}{2km} \right| > \frac{\alpha}{n^2}$$

and

$$\mathbb{P}[\text{ed}(G_k, G_{S(k)}) > \alpha] \leq \sum_{u,v} \mathbb{P} \left[ \left| \frac{W_{G_k}(u,v) - W_{G_{S(k)}}(u,v)}{2km} \right| > \frac{\alpha}{n^2} \right]$$

Let's consider two blocks  $b_i$  and  $b_j$  in  $B$ . We know that  $\forall u \in b_i, v \in b_j, W_{G_{S(k)}}(u,v) = k \cdot \frac{M_{i,j}}{|b_i||b_j|}$  and  $W_{G_k}(u,v) \sim \mathcal{B}(k \cdot M_{i,j}, p_{i,j})$  with  $p_{i,j} = \frac{1}{|b_i||b_j|}$ . Therefore, according to the Bienaymé-Tchebychev inequality:

$$\begin{aligned} \mathbb{P} \left[ \left| \frac{W_{G_k}(u,v) - W_{G_{S(k)}}}{2km} \right| > \frac{\alpha}{n^2} \right] &\leq \frac{k \times M_{i,j} \times p_{i,j} \times (1 - p_{i,j}) \times n^2}{4 \times k^2 \times m^2 \times \alpha} \\ &\leq \frac{M_{i,j} \times p_{i,j} \times (1 - p_{i,j}) \times n^2}{4 \times k \times m^2 \times \alpha} \\ &\xrightarrow[k \rightarrow \infty]{} 0 \end{aligned}$$

Thus,

$$\mathbb{P}[\text{ed}(G_k, G_{S(k)}) > \alpha] \xrightarrow[k \rightarrow \infty]{} 0 \tag{6.6}$$

Which proves the lemma.

We can now prove the theorem

**THEOREM 6.1** Let  $B_1$  and  $B_2$  be two partition on  $\llbracket 1, n \rrbracket$ , with  $p_1$  and  $p_2$  blocks respectively. Let  $M_1 \in \mathcal{M}_{p_1}(\mathbb{N})$  and  $M_2 \in \mathcal{M}_{p_2}(\mathbb{N})$  be two block adjacency matrices such that

$$\sum_{i,j \in [1, p_1]^2} M_1[i, j] = \sum_{k, l \in [1, p_2]^2} M_2[k, l] = m$$

Let's consider two series of stochastic blockmodels defined as  $S_1(k) = (B_1, k \cdot M_1)$  and  $S_2(k) = (B_2, k \cdot M_2)$ , whose barycenters are denoted  $G_1(k)$  and  $G_2(k)$ . We have that

1. There exists  $d \in \mathbb{R}, \forall k \in \mathbb{N}, \text{ed}(G_1(k), G_2(k)) = d$
2. Let  $(G_k)_{k \in \mathbb{N}}$  be a serie of random graph each drawn following model  $S_1(k)$ .

$$\text{EDEV}(G_k, S_2(k)) \xrightarrow[k \rightarrow \infty]{\mathbb{P}} d$$

For any pair of nodes  $i, j$ , belonging to blocks  $b(i)$  and  $b(j)$  in  $B_1$  (resp.  $B_2$ ), the weight of the edge  $i \rightarrow j$  in  $G_1(k)$  (resp.  $G_2(k)$ ) is given by:

$$W_{G_1(k)}[i, j] = k \cdot \frac{M[b(i), b(j)]}{|b(i)||b(j)|}$$

Therefore, the edit distance between  $G_1(k)$  and  $G_2(k)$  is

$$\begin{aligned} \text{ed}(G_1(k), G_2(k)) &= \frac{1}{2km} \sum_{i,j \in [1, n]^2} |W_{G_1(k)}[i, j] - W_{G_2(k)}[i, j]| \\ &= \frac{1}{2km} \sum_{i,j \in [1, n]^2} \left| k \cdot \frac{M_1[b_1(i), b_1(j)]}{|b_1(i)||b_1(j)|} - k \cdot \frac{M_2[b_2(i), b_2(j)]}{|b_2(i)||b_2(j)|} \right| \\ &= \frac{1}{2m} \sum_{i,j \in [1, n]^2} \left| \frac{M_1[b_1(i), b_1(j)]}{|b_1(i)||b_1(j)|} - \frac{M_2[b_2(i), b_2(j)]}{|b_2(i)||b_2(j)|} \right| \end{aligned}$$

which is constant with respect to  $k$ . In the following we will denote this distance  $d$  for the sake of conciseness. We want to show that

$$\text{EDEV}(G_k, S_2(k)) \xrightarrow[k \rightarrow \infty]{\mathbb{P}} d$$

We start by noticing that

$$\begin{aligned} \text{EDEV}(G_k, S_2(k)) - d &= \mathbb{E}_{H \in S_2(k)} [\text{ed}(G_k, H)] - d \\ &\leq \mathbb{E}_{H \in S_2(k)} [\text{ed}(G_k, G_1(k)) + \text{ed}(G_1(k), G_2(k)) + \text{ed}(G_2(k), H)] - d \\ &\leq \text{ed}(G_k, G_1(k)) + \mathbb{E}_{H \in S_2(k)} [\text{ed}(G_2(k), H)] \end{aligned}$$

On the other hand,

$$\begin{aligned}
 d - \text{EDEV}(G_k, S_2(k)) &= \mathbb{E}_{H \in S_2(k)} [\text{ed}(G_1(k), G_2(k)) - \text{ed}(G_k, H)] \\
 &\leq \mathbb{E}_{H \in S_2(k)} [\text{ed}(G_1(k), G_k) + \text{ed}(G_k, H) + \text{ed}(H, G_2(k)) - \text{ed}(G_k, H)] \\
 &\leq \text{ed}(G_k, G_1(k)) + \mathbb{E}_{H \in S_2(k)} [\text{ed}(G_2(k), H)]
 \end{aligned}$$

Thus

$$|\text{EDEV}(G_k, S_2(k)) - d| \leq \text{ed}(G_k, G_1(k)) + \mathbb{E}_{H \in \Omega_{S_2(k)}} [\text{ed}(G_2(k), H)] \quad (6.7)$$

Because  $G_k$  is generated following  $S_1(k)$ , a direct application of lemma 6.1 is that

$$\text{ed}(G_k, G_1(k)) \xrightarrow[k \rightarrow \infty]{\mathbb{P}} 0$$

What is more, if  $H$  is generated following  $S_2(k)$ , we also have that

$$\text{ed}(H, G_2(k)) \xrightarrow[k \rightarrow \infty]{\mathbb{P}} 0$$

which implies that  $\text{ed}(H, G_2(k)) \xrightarrow[k \rightarrow \infty]{\mathcal{L}} 0$  and in particular

$$\mathbb{E}_{H \in \Omega_{S_2(k)}} [H, G_2(k)] \xrightarrow[k \rightarrow \infty]{} 0$$

Finally, we obtain that

$$\text{ed}(G_k, G_1(k)) + \mathbb{E}_{H \in \Omega_{S_2(k)}} [H, G_2(k)] \xrightarrow[k \rightarrow \infty]{\mathbb{P}} 0$$

And thanks to equation 6.7:

$$\text{EDEV}(G_k, S_2(k)) \xrightarrow[k \rightarrow \infty]{\mathbb{P}} d \quad (6.8)$$

#### REFERENCES

1. Barabási, A.-L. & Albert, R. (1999) Emergence of scaling in random networks. *science*, **286**(5439), 509–512.
2. Barthélemy, M. (2011) Spatial networks. *Physics Reports*, **499**(1-3), 1–101.
3. Bianconi, G. (2009) Entropy of network ensembles. *Physical Review E*, **79**(3), 036114.
4. Cimini, G., Squartini, T., Saracco, F., Garlaschelli, D., Gabrielli, A. & Caldarelli, G. (2018) The Statistical Physics of Real-World Networks. *arXiv:1810.05095 [cond-mat, physics:physics]*. arXiv: 1810.05095.
5. Ding, J., Tarokh, V. & Yang, Y. (2018) Model Selection Techniques – An Overview. *arXiv:1810.09583 [physics, stat]*. arXiv: 1810.09583.
6. Goldenberg, A., Zheng, A. X., Fienberg, S. E., Airolidi, E. M. et al. (2010) A survey of statistical network models. *Foundations and Trends® in Machine Learning*, **2**(2), 129–233.
7. Grünwald, P. (2005) A tutorial introduction to the minimum description length principle. *Advances in minimum description length: Theory and applications*, pages 3–81.

8. Guimera, R., Sales-Pardo, M. & Amaral, L. A. N. (2004) Modularity from fluctuations in random graphs and complex networks. *Physical Review E*, **70**(2), 025101.
9. Koutra, D., Vogelstein, J. T. & Faloutsos, C. (2013) Deltacon: A principled massive-graph similarity function. *Proceedings of the 2013 SIAM International Conference on Data Mining*, pages 162–170.
10. Monnig, N. D. & Meyer, F. G. (2018) The resistance perturbation distance: A metric for the analysis of dynamic networks. *Discrete Applied Mathematics*, **236**, 347–386.
11. Newman, M. E. (2003) The structure and function of complex networks. *SIAM review*, **45**(2), 167–256.
12. Newman, M. E. (2006) Modularity and community structure in networks. *Proceedings of the national academy of sciences*, **103**(23), 8577–8582.
13. Peixoto, T. P. (2012) Entropy of stochastic blockmodel ensembles. *Physical Review E*, **85**(5), 056122.
14. Peixoto, T. P. (2017) Bayesian stochastic blockmodeling. *arXiv preprint arXiv:1705.10225*.
15. Simini, F., González, M. C., Maritan, A. & Barabási, A.-L. (2012) A universal model for mobility and migration patterns. *Nature*, **484**(7392), 96.
16. Watts, D. J. & Strogatz, S. H. (1998) Collective dynamics of ‘small-world’ networks. *nature*, **393**(6684), 440.
17. Wills, P. & Meyer, F. G. (2019) Metrics for Graph Comparison: A Practitioner’s Guide. *arXiv:1904.07414 [physics, q-bio, stat]*. arXiv: 1904.07414.
18. Zingg, C., Casiraghi, G., Vaccario, G. & Schweitzer, F. (2019) What is the Entropy of a Social Organization?. *arXiv:1905.09772 [physics]*. arXiv: 1905.09772.

### Funding

Work supported by the ACADEMICS grant of the IDEXLYON, project of the Université de Lyon, PIA operated by ANR-16-IDEX-0005