

WEIGHTED UNARY-BINARY TREES, HEX-TREES, AND HORTON-STRAHLER NUMBERS REVISITED

HELMUT PRODINGER

ABSTRACT. Hex-trees are identified as a particular instance of weighted unary-binary trees. The Horton-Strahler numbers of these objects are revisited, and, thanks to a substitution that is not very intuitive, explicit results are possible. They are augmented by asymptotic evaluations as well.

1. BINARY TREES AND HORTON-STRAHLER NUMBERS

Binary trees may be expressed by the following symbolic equation, which says that they include the empty tree and trees recursively built from a root followed by two subtrees, which are binary trees:


$$\mathcal{B} = \square + \begin{array}{c} \circ \\ \swarrow \quad \searrow \\ \mathcal{B} \quad \mathcal{B} \end{array}$$

Binary trees are counted by Catalan numbers and there is an important parameter reg , which in Computer Science is called the register function. It associates to each binary tree (which is used to code an arithmetic expression, with data in the leaves and operators in the internal nodes) the minimal number of extra registers that is needed to evaluate the tree. The optimal strategy is to evaluate difficult subtrees first, and use one register to keep its value, which does not hurt, if the other subtree requires less registers. If both subtrees are equally difficult, then one more register is used, compared to the requirements of the subtrees. This natural parameter is among Combinatorialists known as the Horton-Strahler numbers, and we will adopt this name throughout this paper.

There is a recursive description of this function: $\text{reg}(\square) = 0$, and if tree t has subtrees t_1 and t_2 , then

$$\text{reg}(t) = \begin{cases} \max\{\text{reg}(t_1), \text{reg}(t_2)\} & \text{if } \text{reg}(t_1) \neq \text{reg}(t_2), \\ 1 + \text{reg}(t_1) & \text{otherwise.} \end{cases}$$

The recursive description attaches numbers to the nodes, starting with 0's at the leaves and then going up; the number appearing at the root is the Horton-Strahler number of the tree.



Let \mathcal{R}_p denote the family of trees with Horton-Strahler number = p , then one gets immediately from the recursive definition:

$$\mathcal{R}_p = \begin{array}{c} \circ \\ \swarrow \quad \searrow \\ \mathcal{R}_{p-1} \quad \mathcal{R}_{p-1} \end{array} + \begin{array}{c} \circ \\ \swarrow \quad \searrow \\ \mathcal{R}_p \quad \sum_{j < p} \mathcal{R}_j \end{array} + \begin{array}{c} \circ \\ \swarrow \quad \searrow \\ \sum_{j < p} \mathcal{R}_j \quad \mathcal{R}_p \end{array}$$

In terms of generating functions, these equations read as

$$R_p(z) = zR_{p-1}^2(z) + 2zR_p(z) \sum_{j < p} R_j(z);$$

the variable z is used to mark the size (i. e., the number of internal nodes) of the binary tree.

A historic account of these concepts, from the angle of Philippe Flajolet, who was one of the pioneers is [8], compare also [7].

Amazingly, the recursion for the generating functions $R_p(z)$ can be solved explicitly! The substitution

$$z = \frac{u}{(1+u)^2}$$

that de Bruijn, Knuth, and Rice [1] also used, produces the nice expression

$$R_p(z) = \frac{1-u^2}{u} \frac{u^{2^p}}{1-u^{2^{p+1}}}.$$

Of course, once this is *known*, it can be proved by induction, using the recursive formula. For the readers benefit, this will be sketched now.

We start with the auxiliary formula

$$\sum_{0 \leq j < p} \frac{u^{2^j}}{1-u^{2^{j+1}}} = \frac{u}{1-u} - \frac{u^{2^p}}{1-u^{2^p}},$$

which is easy to prove by induction: For $p = 0$, the formula $0 = \frac{u}{1-u} - \frac{u}{1-u}$ is correct, and then

$$\begin{aligned} \sum_{0 \leq j < p+1} \frac{u^{2^j}}{1-u^{2^{j+1}}} &= \frac{u}{1-u} - \frac{u^{2^p}}{1-u^{2^p}} + \frac{u^{2^p}}{1-u^{2^{p+1}}} \\ &= \frac{u}{1-u} - \frac{u^{2^p}(1+u^{2^p})}{1-u^{2^{p+1}}} + \frac{u^{2^p}}{1-u^{2^{p+1}}} = \frac{u}{1-u} - \frac{u^{2^{p+1}}}{1-u^{2^{p+1}}}. \end{aligned}$$

Now the formula for $R_p(z)$ can also be proved by induction. First, $R_0(z) = \frac{1-u^2}{u} \frac{u}{1-u^2} = 1$, as it should, and

$$zR_{p-1}^2(z) + 2zR_p(z) \sum_{j < p} R_j(z)$$

$$\begin{aligned}
 &= \frac{u}{(1+u)^2} \frac{(1-u^2)^2}{u^2} \frac{u^{2p}}{(1-u^{2p})^2} + \frac{2u}{(1+u)^2} R_p(z) \sum_{j < p} \frac{1-u^2}{u} \frac{u^{2j}}{1-u^{2^{j+1}}} \\
 &= \frac{u^{2p-1}(1-u)^2}{(1-u^{2p})^2} + \frac{2(1-u)}{(1+u)} R_p(z) \sum_{j < p} \frac{u^{2j}}{1-u^{2^{j+1}}}.
 \end{aligned}$$

Solving

$$R_p(z) = \frac{u^{2p-1}(1-u)^2}{(1-u^{2p})^2} + \frac{2(1-u)}{(1+u)} R_p(z) \left[\frac{u}{1-u} - \frac{u^{2p}}{1-u^{2p}} \right]$$

leads to

$$R_p(z) \frac{1-u}{1+u} \left[1 + 2 \frac{u^{2p}}{1-u^{2p}} \right] = \frac{u^{2p-1}(1-u)^2}{(1-u^{2p})^2},$$

or, simplified

$$R_p(z) = \frac{u^{2p-1}(1-u^2)}{(1-u^{2p})(1+u^{2p})} = \frac{1-u^2}{u} \frac{u^{2p}}{(1-u^{2^{p+1}})},$$

which is the formula that we needed to prove. □

2. UNARY-BINARY TREES AND HEX-TREES

The family of unary-binary trees \mathcal{M} might be defined by the symbolic equation

$$\mathcal{M} = \square + \begin{array}{c} \circ \\ | \\ \mathcal{M} \setminus \{\square\} \end{array} + \begin{array}{c} \circ \\ / \quad \backslash \\ \mathcal{M} \quad \mathcal{M} \end{array}$$

The equation for the generating function is

$$M = 1 + z(M-1) + zM^2$$

with the solution

$$M = M(z) = \frac{1-z-\sqrt{1-6z+5z^2}}{2z} = 1+z+3z^2+10z^3+36z^4+\dots;$$

the coefficients form sequence A002212 in [9] and enumerate Schröder paths, among many other things. We will come to equivalent structures a bit later.

In the instance of unary-binary trees, we can also work with a substitution: Set $z = \frac{u}{1+3u+u^2}$, then $M(z) = 1+u$. Unary-binary trees and the register function were investigated in [3], but the present favourable substitution was not used. Therefore, in this previous paper, asymptotic results were available but no explicit formulae.

This works also with a weighted version, where we allow unary edges with a different colours. Then

$$\mathcal{N} = \square + a \cdot \begin{array}{c} \circ \\ | \\ \mathcal{N} \setminus \{\square\} \end{array} + \begin{array}{c} \circ \\ / \quad \backslash \\ \mathcal{N} \quad \mathcal{N} \end{array}$$

and with the substitution $z = \frac{u}{1+(a+2)u+u^2}$, the generating function is beautifully expressed as $N(z) = 1+u$. For $a = 0$, this covers also binary trees.

We will consider the Horton-Strahler numbers of unary-binary trees in the sequel. The definition is naturally extended by

$$\text{reg}\left(\begin{array}{c} \circ \\ | \\ t \end{array}\right) = \text{reg}(t).$$

Now we can move again to $R_p(z)$, the generating function of (generalized) unary-binary trees with Horton-Strahler number $= p$. The recursion (for $p \geq 1$) is

$$\mathcal{R}_p = \begin{array}{c} \circ \\ / \quad \backslash \\ \mathcal{R}_{p-1} \quad \mathcal{R}_{p-1} \end{array} + \begin{array}{c} \circ \\ / \quad \backslash \\ \mathcal{R}_p \quad \sum_{j < p} \mathcal{R}_j \end{array} + \begin{array}{c} \circ \\ / \quad \backslash \\ \sum_{j < p} \mathcal{R}_j \quad \mathcal{R}_p \end{array} + a \cdot \begin{array}{c} \circ \\ | \\ \mathcal{R}_p \end{array}$$

In terms of generating functions, these equations read as

$$R_p(z) = zR_{p-1}^2(z) + 2zR_p(z) \sum_{j < p} R_j(z) + azR_p(z), \quad p \geq 1; \quad R_0(z) = 1.$$

Amazingly, with the substitution $z = \frac{u}{1+(a+2)u+u^2}$, formally we get the *same* solution as in the binary case:

$$R_p(z) = \frac{1-u^2}{u} \frac{u^{2p}}{1-u^{2p+1}}.$$

The proof by induction is as before. One sees another advantage of the substitution: On a formal level, many manipulations do not need to be repeated. Only when one switches back to the z -world, things become different.

Now we move to Hex-trees.

$$\mathcal{H} = \square + \circ + \begin{array}{c} \circ \\ / \quad \backslash \\ \mathcal{H} \setminus \{\square\} \quad \mathcal{H} \setminus \{\square\} \end{array} + \begin{array}{c} \circ \\ / \quad \backslash \\ \mathcal{H} \setminus \{\square\} \quad \square \end{array} + \begin{array}{c} \circ \\ | \\ \mathcal{H} \setminus \{\square\} \end{array} + \begin{array}{c} \circ \\ / \quad \backslash \\ \square \quad \mathcal{H} \setminus \{\square\} \end{array}$$

Hex trees either have two non-empty successors, or one of 3 types of unary successors (called left, middle, right). The author has seen this family first in [6], but one can find older literature following the references and the usual search engines.

The generating function satisfies

$$H(z) = 1 + z(H(z) - 1)^2 + z + 3z(H(z) - 1) = \frac{1 - z - \sqrt{(1-z)(1-5z)}}{2z}.$$

$$1 + z + 3z^2 + 10z^3 + 36z^4 + 137z^5 + 543z^6 + 2219z^7 + 9285z^8 + 39587z^9 + 171369z^{10} + \dots$$

The same generating function also appears in [5], and it is again sequence A002212 in [9]. One can rewrite the symbolic equation as

$$\mathcal{H} = \square + \begin{array}{c} \circ \\ / \quad \backslash \\ \mathcal{H} \quad \mathcal{H} \end{array} + \begin{array}{c} \circ \\ | \\ \mathcal{H} \setminus \{\square\} \end{array}$$

and sees in this way that the Hex trees are unary-binary trees (with parameter $a = 1$).

3. CONTINUING WITH ENUMERATIONS

First, we will enumerate the number of (generalized) unary-binary trees with n (internal) nodes. For that we need the notion of generalized trinomial coefficients, viz.

$$\binom{n; 1, a, 1}{k} := [z^k](1 + az + z^2)^n.$$

Of course, for $a = 2$, this simplifies to a binomial coefficient $\binom{2n}{k}$. We will use contour integration to pull out coefficients, and the contour of integer, in whatever variable, is a small circle (or equivalent) around the origin. The desired number is

$$\begin{aligned}
 [z^n](1+u) &= \frac{1}{2\pi i} \oint \frac{dz}{z^{n+1}}(1+u) \\
 &= \frac{1}{2\pi i} \oint \frac{du(1-u^2)(1+(a+2)u+u^2)^{n+1}}{(1+(a+2)u+u^2)^2 u^{n+1}}(1+u) \\
 &= [u^{n+1}](1-u)(1+u)^2(1+(a+2)u+u^2)^{n-1} \\
 &= \binom{n-1; 1, a+2, 1}{n+1} + \binom{n-1; 1, a+2, 1}{n} \\
 &\quad - \binom{n-1; 1, a+2, 1}{n-1} - \binom{n-1; 1, a+2, 1}{n-2}.
 \end{aligned}$$

Then we introduce $S_p(z) = R_p(z) + R_{p+1}(z) + R_{p+2}(z) + \dots$, the generating function of trees with Horton-Strahler number $\geq p$. Using the summation formula proved earlier, we get

$$S_p(z) = \frac{1-u^2}{u} \frac{u^{2p}}{1-u^{2p}} = \frac{1-u^2}{u} \sum_{k \geq 1} u^{k2^p}.$$

Further,

$$[z^n]S_p(z) = \sum_{k \geq 1} \frac{1}{2\pi i} \oint \frac{dz}{z^{n+1}} \frac{1-u^2}{u} u^{k2^p}.$$

4. ASYMPTOTICS

We start by deriving asymptotics for the number of (generalized) unary-binary trees with n (internal) nodes. This is a standard application of singularity analysis of generating, as described in [2] and [4].

We start from the generating function

$$N(z) = \frac{1 - az - \sqrt{1 - 2(a+2)z + a(a+4)z^2}}{2z}$$

and determine the singularity closest to the origin, which is the value making the square root disappear: $z = \frac{1}{a+4}$. After that, the local expansion of $N(z)$ around this singularity is determined:

$$N(z) \sim 2 - \sqrt{a+4} \sqrt{1 - (a+4)z}.$$

The translation lemmas given in [2] and [4] provide the asymptotics:

$$\begin{aligned}
 [z^n]N(z) &\sim [z^n] \left(2 - \sqrt{a+4} \sqrt{1 - (a+4)z} \right) \\
 &= -\sqrt{a+4} (a+4)^n \frac{n^{-3/2}}{\Gamma(-\frac{1}{2})} = (a+4)^{n+1/2} \frac{1}{2\sqrt{\pi} n^{3/2}}.
 \end{aligned}$$

Just note that $a = 0$ is the well-known formula for binary trees with n nodes.

Now we move to the generating function for the average number of registers. Apart from normalization it is

$$\begin{aligned} \sum_{p \geq 1} pR_p(z) &= \sum_{p \geq 1} S_p(z) = \frac{1-u^2}{u} \sum_{p \geq 1} \sum_{k \geq 1} u^{k2^p} \\ &= \frac{1-u^2}{u} \sum_{n \geq 1} v_2(n)u^n, \end{aligned}$$

where $v_2(n)$ is the highest exponent k such 2^k divides n .

This has to be studied around $u = 1$, which, upon setting $u = e^{-t}$, means around $t = 0$. Eventually, and that is the only thing that is different here, is to retranslate that into a singular expansion of z around its singularity, which depends on the parameter a .

For the reader's convenience, we also repeat the steps that were known before. The first factor is elementary:

$$\frac{1-u^2}{u} \sim 2t + \frac{1}{3}t^3 + \dots$$

For

$$\sum_{p \geq 1} \sum_{k \geq 1} e^{-k2^p t},$$

one applied the Mellin transform, with the result

$$\frac{\Gamma(s)\zeta(s)}{2^s - 1}.$$

Applying the inversion formula, one finds

$$\sum_{p \geq 1} \sum_{k \geq 1} e^{-k2^p t} = \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} t^{-s} \frac{\Gamma(s)\zeta(s)}{2^s - 1} ds.$$

Shifting the line of integration to the left, the residues at the poles $s = 1$, $s = 0$, $s = \chi_k = \frac{2k\pi i}{\log 2}$, $k \neq 0$ provide enough terms for our asymptotic expansion.

$$\frac{1}{t} + \frac{\gamma}{2 \log 2} - \frac{1}{4} - \frac{\log \pi}{2 \log 2} + \frac{\log t}{2 \log 2} + \frac{1}{\log 2} \sum_{k \neq 0} \Gamma(\chi_k) \zeta(\chi_k) t^{-\chi_k}.$$

Combined with the elementary factor, this leads to

$$2 + \left(\frac{\gamma}{\log 2} - \frac{1}{2} - \frac{\log \pi}{\log 2} + \frac{\log t}{\log 2} \right) t + \frac{2t}{\log 2} \sum_{k \neq 0} \Gamma(\chi_k) \zeta(\chi_k) t^{-\chi_k} + O(t^2 \log t).$$

Now we want to translate into the original z -world. Since $z = \frac{u}{1+(a+2)u+u^2}$, $u = 1$ translates into the singularity $z = \frac{1}{4+a}$. Further,

$$t \sim \sqrt{4+a} \cdot \sqrt{1-z(4+a)},$$

let us abbreviate $A = 4+a$, then for singularity analysis we must consider

$$\begin{aligned} &\frac{\sqrt{A} \cdot \sqrt{1-zA} \log(1-zA)}{2 \log 2} \\ &+ \left(\frac{\gamma}{\log 2} - \frac{1}{2} - \frac{\log \pi}{\log 2} + \frac{\log A}{2 \log 2} \right) \sqrt{A} \cdot \sqrt{1-zA} \end{aligned}$$

$$+ \frac{2}{\log 2} \sum_{k \neq 0} \Gamma(\chi_k) \zeta(\chi_k) A^{\frac{1-\chi_k}{2}} (1-zA)^{\frac{1-\chi_k}{2}}.$$

The formula that is perhaps less known and needed here is

$$[z^n] \log(1-z) \sqrt{1-z} \sim \frac{n^{-3/2} \log n}{2\sqrt{\pi}} + \frac{n^{-3/2}}{2\sqrt{\pi}} (-2 + \gamma + 2 \log 2);$$

furthermore we need

$$[z^n] (1-z)^\alpha \sim \frac{n^{-\alpha-1}}{\Gamma(-\alpha)}.$$

We start with the most complicated term:

$$\begin{aligned} \frac{[z^n] \frac{\sqrt{A} \sqrt{1-zA} \log(1-zA)}{2 \log 2}}{[z^n] N(z)} &\sim \frac{\sqrt{A} A^n \left(\frac{n^{-3/2} \log n}{2\sqrt{\pi}} + \frac{n^{-3/2}}{2\sqrt{\pi}} (-2 + \gamma + 2 \log 2) \right)}{2 \log 2 \frac{A^{n+1/2}}{2\sqrt{\pi n^{3/2}}}} \\ &= \log_4 n + 1 + \frac{\gamma}{2 \log 2} - \frac{1}{\log 2}. \end{aligned}$$

The next term we consider is

$$\begin{aligned} \left(\frac{\gamma}{\log 2} - \frac{1}{2} - \frac{\log \pi}{\log 2} + \frac{\log A}{2 \log 2} \right) \sqrt{A} \frac{[z^n] \sqrt{1-zA}}{[z^n] N(z)} \\ \sim \left(\frac{\gamma}{\log 2} - \frac{1}{2} - \frac{\log \pi}{\log 2} + \frac{\log A}{2 \log 2} \right) \sqrt{A} \frac{[z^n] \sqrt{1-zA}}{-\sqrt{A} [z^n] \sqrt{1-zA}} \\ = -\frac{\gamma}{\log 2} + \frac{1}{2} + \frac{\log \pi}{\log 2} - \frac{\log A}{2 \log 2}. \end{aligned}$$

The last term we consider is

$$\begin{aligned} \frac{2}{\log 2} \Gamma(\chi_k) \zeta(\chi_k) A^{\frac{1-\chi_k}{2}} \frac{[z^n] (1-zA)^{\frac{1-\chi_k}{2}}}{-\sqrt{A} [z^n] \sqrt{1-zA}} \\ \sim -\frac{4\sqrt{\pi}}{\log 2} \frac{\Gamma(\chi_k) \zeta(\chi_k)}{\Gamma(\frac{\chi_k-1}{2})} A^{\frac{1-\chi_k}{2}} n^{\chi_k/2}. \end{aligned}$$

Eventually we have evaluated the average value of the Horton-Strahler numbers:

Theorem 1.

$$\begin{aligned} \log_4 n - \frac{\gamma}{2 \log 2} - \frac{1}{\log 2} + \frac{3}{2} + \frac{\log \pi}{\log 2} - \frac{\log A}{2 \log 2} - \frac{4\sqrt{\pi A}}{\log 2} \sum_{k \neq 0} \frac{\Gamma(\chi_k) \zeta(\chi_k)}{\Gamma(\frac{\chi_k-1}{2})} A^{\frac{-\chi_k}{2}} n^{\chi_k/2} \\ = \log_4 n - \frac{\gamma}{2 \log 2} - \frac{1}{\log 2} + \frac{3}{2} + \frac{\log \pi}{\log 2} - \frac{\log A}{2 \log 2} + \psi(\log_4 n), \end{aligned}$$

with a tiny periodic function $\psi(x)$ of period 1.

REFERENCES

- [1] N. G. De Bruijn, D. E. Knuth, and S. O. Rice. The average height of planted plane trees. In R. C. Read, editor, *Graph Theory and Computing*, pages 15–22. Academic Press, 1972.
- [2] P. Flajolet and A. M. Odlyzko. Singularity analysis of generating functions. 3:216–240, 1990.
- [3] P. Flajolet and H. Prodinger. Register allocation for unary-binary trees. *SIAM J. Comput.*, 15(3):629–640, 1986.
- [4] Philippe Flajolet and Robert Sedgewick. *Analytic combinatorics*. Cambridge University Press, Cambridge, 2009.
- [5] Clemens Heuberger, Helmut Prodinger, and Stephan Wagner. The height of multiple edge plane trees. *Aequationes Math.*, 90(3):625–645, 2016.
- [6] Hana Kim and Richard P. Stanley. A refined enumeration of hex trees and related polynomials. *European J. Combin.*, 54:207–219, 2016.
- [7] H. Prodinger. Philippe Flajolet’s early work in combinatorics. 2:1 (2022) Article S1H1.:11 pages, 2022.
- [8] Helmut Prodinger. Introduction to Philippe Flajolet’s work on the register function and related topics. <https://www.math.tugraz.at/~prodinger/flajolet-introduction-register.pdf>, 7 pages, 2025.
- [9] Neil J. A. Sloane. The on-line encyclopedia of integer sequences. *Notices Amer. Math. Soc.*, 65(9):1062–1074, 2018.

HELMUT PRODINGER, MATHEMATICS DEPARTMENT, STELLENBOSCH UNIVERSITY, 7602 STELLENBOSCH, SOUTH AFRICA.

Email address: `hprodinger@sun.ac.za`