

A Logic for Reasoning About LF Specifications

Gopalan Nadathur

Mary Southern

April 12, 2022

Abstract

We present a logic named \mathcal{L}_{LF} whose intended use is to formalize properties of specifications developed in the dependently typed lambda calculus LF. The logic is parameterized by the LF signature that constitutes the specification. Atomic formulas correspond to typing derivations relative to this signature. The logic includes a collection of propositional connectives and quantifiers. Quantification ranges over expressions that denote LF terms and LF contexts. Quantifiers of the first variety are qualified by simple types that describe the functional structure associated with the variables they bind; deeper, dependency related properties are expressed by the body of the formula. Context-level quantifiers are qualified by *context schemas* that identify patterns of declarations out of which actual contexts may be constructed. The semantics of variable-free atomic formulas is articulated via the derivability in LF of the judgements they encode. Propositional constants and connectives are understood in the usual manner and the meaning of quantifiers is explicated through substitutions of expressions that adhere to the type qualifications. The logic is complemented by a proof system that enables reasoning that is sound with respect to the described semantics. The main novelties of the proof system are the provision for case-analysis style reasoning about LF judgements, support for inductive reasoning over the heights of LF derivations and the encoding of LF meta-theorems. The logic is motivated by the paradigmatic example of type assignment in the simply-typed lambda calculus and the proof system is illustrated through the formalization of a proof of type uniqueness for this calculus.

1 Introduction

The Edinburgh Logical Framework of LF has been proposed as a vehicle for formalizing rule-based presentation of object systems and has also been successfully used in many such formalizations. Two aspects of LF are critical to its use in such applications: its basis in a regime of dependent types and its ability to support a meta-level treatment of binding. The typical deployment of LF in this context involves the presentation of a signature that represents the object system. Exploiting their parameterization by terms, types are then used to encode relations between objects. Inhabitation of such a type, demonstrated through the derivation of a typing judgement, provides witness to the validity of a relation of interest. The meta-level encoding of binding is exhibited in the form of typing judgements in which typing contexts that scope over types and terms play an important role.

The objective of this paper is to develop a framework for formalizing and proving properties of specifications that are constructed in the manner described above. This framework is intended to provide a formal counterpart to the informal style of reasoning about object systems based on their rule-based description as is seen, for example, in [14]. To fit this role, the framework would need to be based on a logic that allows for the expression of complex properties that are built on top of derivability judgements whose validity is determined by the rules describing the system. Moreover, the framework would need to accommodate case analysis and inductive forms

of reasoning, thereby enabling the least-fixed point treatment of the rules that is typically assumed in the informal arguments.

Towards delivering on the mentioned objective, we describe here a logic in which properties of LF specifications can be formalized. The logic is parameterized by an LF signature that describes the particular object system that is of interest. The atomic formulas in the logic then comprise typing judgements that assert the inhabitation of types relative to LF contexts. Complex formulas can be constructed from these judgements using a collection of propositional constants and connectives and quantifiers. The quantification that is permitted ranges over expressions that denote both LF terms and LF contexts. A critical issue that must be addressed in this context is the qualification of quantifiers to limit their scope to meaningful subclasses of expressions. For term quantifiers, we address this issue by using a special class of simple types that are referred to as *arity types*. Such a type limits attention to terms that satisfy a particular functional structure, leaving deeper, dependency related properties to the domain of the formula that the quantifier ranges over. For context quantifiers, we introduce a new kind of types that we call *context schemas*. These types, which are motivated by the *regular worlds* used with Twelf developments [11, 15], identify patterns of declarations out of which the actual contexts that instantiate the quantifiers must be constructed. The validity of atomic formulas that are devoid of free variables is determined, as might be expected, by the derivability of the LF judgements that they represent. Propositional symbols are understood in the usual manner. An especially interesting part of the semantics is the treatment of quantifiers: these are interpreted via the substitution of expressions that adhere to the type qualifications that adorn them.

Perhaps the most significant contribution of this paper is a proof system that can be used to establish the validity of formulas in the logic that is described. This proof system is oriented around sequents that intuitively encapsulate states in the development of a proof. A basic part of this system is a set of rules that encode the meanings of the propositional symbols and quantifiers. The truly innovative part of the system is its treatment of atomic formulas that embodies their interpretation as typing judgements in LF. Included in this part is a mechanism for analyzing an atomic assumption formula via the parameterizing signature and the declarations in the LF context that is a part of the formula. The rules also build in the capability to reason by induction on the heights of LF derivations and to utilize meta-theorems about LF derivability. We show in this paper that the proof system is both coherent and sound. While we illustrate the logic and the proof system, we leave the demonstration of their effectiveness to other work, e.g. see [1, 18].

The rest of the paper is organized as follows. In the next section, we present the notions related to LF that we need in other parts of this work. A major part of this presentation is the treatment of *simultaneous hereditary substitution*, a concept that is important to much of the technical development in this paper and that has also not been treated in prior work. Section 3 then presents the logic \mathcal{L}_{LF} and illustrates its use in formalizing the type uniqueness property for the simply-typed lambda calculus, a paradigmatic example for such work. Section 4 describes the notion of a sequent and develops a collection of proof rules for deriving sequents. The main burden of the technical work in this section is the demonstration of the coherence and soundness of the rules that are presented. Section 5 illustrates the proof system by showing how it can be used to encode the informal argument outlined in Section 3. We conclude the paper in Section 6 by placing the ideas in this paper in the context of other work related to reasoning about LF specifications.

2 LF and the Formalization of Object Systems

The methodology for modeling object systems in LF relies on adequacy theorems that, in turn, depend on canonical $\beta\eta$ -forms for terms. The original presentation of LF [5] includes terms in both canonical and non-canonical form. Such a presentation simplifies the treatment of substitution but at the price of complicating arguments concerning adequacy and LF derivability. In light of this an alternative treatment of LF has been proposed that admits only terms that are in β -normal form and that are well-typed only if they are additionally in η -long form [6, 21]. We use this presentation of LF, called *canonical LF*, as the basis for our work. The first subsection below recalls this presentation and develops notions related to it that will be used in the later parts of this paper. Towards motivating the development of the logic that is the main content of this paper, we then discuss the use of LF in representing object systems and in reasoning about them at an informal level. The section concludes with an identification of meta-theorems related to derivability in LF that will be used in articulating proof rules in our logic.

2.1 Canonical LF

Our presentation of canonical LF, henceforth referred to simply as LF, differs from that in [6] in two respects. First, we elide the subordination relation in typing judgements since it is orthogonal to the thrust of this paper. Second, we treat substitution independently of LF typing judgements and we also extend the notion to include the simultaneous replacement of multiple variables. The elaboration below builds in these ideas.

Kinds	$K ::= \text{Type} \mid \Pi x:A. K$
Canonical Type Families	$A, B ::= P \mid \Pi x:A. B$
Atomic Type Families	$P ::= a \mid P M$
Canonical Terms	$M, N ::= R \mid \lambda x. M$
Atomic Terms	$R ::= c \mid x \mid R M$
Signatures	$\Sigma ::= \cdot \mid \Sigma, c : A \mid \Sigma, a : K$
Contexts	$\Gamma ::= \cdot \mid \Gamma, x : A$

Figure 2.1: The Syntax of LF Expressions

2.1.1 The Syntax

The syntax of LF expressions is described in Figure 2.1. The primary interest is in three categories of expressions: kinds, types which are indexed by kinds, and terms which are indexed by types. In these expressions, λ and Π are binding or abstraction operators. Relative to these operators, we assume the principle of equivalence under renaming that is applied as needed. We also assume as understood the notions of free and bound variables that are usual to expressions involving such operators. To ensure the absence of β -redexes, terms are stratified into *canonical* and *atomic* forms. A similar stratification is used with types that is exploited by the formation rules to force all well-typed terms to be in η -long form. We use x and y to represent term-level variables, which are bound by abstraction operators or in the contexts that are associated with terms. Further, we use c and d for term-level constants, and a and b for type-level constants, both of which are typed in signatures. The expression $A_1 \rightarrow A_2$ is used as an alternative notation for the type family

$\Pi x:A_1. A_2$ when x does not appear free in A_2 . An atomic term has the form $(h M_1 \dots M_n)$ where h is a variable or a constant. We refer to h as the *head symbol* of such a term.

2.1.2 Simultaneous Hereditary Substitution

We will need to consider substitution into LF expressions when explicating typing and other logical notions related to these expressions. To preserve the form of these expressions, it is necessary to build β -reduction into the application of such substitutions. An important consideration in this context is that substitution application must be a terminating operation. Towards ensuring this property, substitutions are indexed by types that are eventually intended to characterize the functional structure of expressions.

Definition 2.1. *The collection of expressions that are obtained from the constant o using the binary infix constructor \rightarrow constitute the arity types. Corresponding to each canonical type A , there is an arity type called its erased form and denoted by $(A)^-$ and given as follows: $(P)^- = o$ and $(\Pi x:A_1. A_2)^- = (A_1)^- \rightarrow (A_2)^-$.*

Definition 2.2. *A substitution θ is a finite set of the form $\{\langle x_1, M_1, \alpha_1 \rangle, \dots, \langle x_n, M_n, \alpha_n \rangle\}$, where, for $1 \leq i \leq n$, x_i is a distinct variable, M_i is a canonical term and α_i is an arity type.¹ Given such a substitution, $\text{dom}(\theta)$ denotes the set $\{x_1, \dots, x_n\}$ and $\text{rng}(\theta)$ denotes the set $\{M_1, \dots, M_n\}$.*

Given a substitution θ and an expression E that is a kind, a type, a canonical term or a context, we wish the expression $E[\theta]$ notionally to denote the application of θ to E . However, such an application is not guaranteed to exist. We therefore use the expression $E[\theta] = E'$ to indicate when it is defined and has E' as a result. The key part of defining this relation is that of articulating its meaning when E is a canonical term. This is done in Figure 2.2 via rules for deriving this relation. These rules use an auxiliary definition of substitution into an atomic term which accounts for any normalization that is necessitated by the replacement of a variable by a term. The different categories of rules in this figure are distinguished by being preceded by a box containing the judgement form they relate to. The extension of this definition to the case where E is a kind or a type corresponds essentially to the application of the substitution to the terms that appear within E . This idea is made explicit for types in Figure 2.3 and its elaboration for kinds is similar. A substitution is meaningfully applied to a context only when it does not replace variables to which the context assigns types and when a replacement does not lead to inadvertent capture. When these conditions are satisfied, the substitution distributes to the types that are assigned to the variables as the rules in Figure 2.4 make clear.

We define a measure on substitutions that is useful in showing that their application terminates.

Definition 2.3. *The size of an arity type is the number of occurrences of \rightarrow in it. The size of a substitution is the largest of the sizes of the arity types in each of its triples.*

The following theorem can be proved by induction first on the sizes of substitutions and then on the structures of expressions; it would first be proved simultaneously for canonical and atomic terms and then extended to types, kinds and contexts.

Theorem 2.4. *For any context, kind, type or canonical term E and any substitution θ , it is decidable whether there is an E' such that $E[\theta] = E'$ is derivable. Moreover, there is at most one E' for which it is derivable. Similarly, for any atomic term R and substitution θ , it is decidable*

¹Note that by a systematic abuse of notation, n may be less than m in a sequence written in the form s_m, \dots, s_n , in which case the empty sequence is denoted. In this particular instance, a substitution can be an empty set of triples.

$$\boxed{M[\theta] = M'}$$

$$\frac{R[\theta]^r = R'}{R[\theta] = R'}$$

$$\frac{R[\theta]^r = M' : \alpha'}{R[\theta] = M'}$$

$$\frac{x \text{ not free in } \text{dom}(\theta) \cup \text{rng}(\theta) \quad M[\theta] = M'}{(\lambda x. M)[\theta] = \lambda x. M'}$$

$$\boxed{R[\theta]^r = M' : \alpha'}$$

$$\frac{\langle x, M, \alpha \rangle \in \theta}{x[\theta]^r = M : \alpha}$$

$$\frac{R[\theta]^r = \lambda x. M' : \alpha' \rightarrow \alpha'' \quad M[\theta] = M'' \quad M'[\{\langle x, M'', \alpha' \rangle\}] = M'''}{(R M)[\theta]^r = M''' : \alpha''}$$

$$\boxed{R[\theta]^r = R'}$$

$$\frac{}{c[\theta]^r = c}$$

$$\frac{x \notin \text{dom}(\theta)}{x[\theta]^r = x}$$

$$\frac{R[\theta]^r = R' \quad M[\theta] = M'}{(R M)[\theta]^r = R' M'}$$

Figure 2.2: Applying Substitutions to Terms

$$\frac{}{a[\theta] = a} \quad \frac{P[\theta] = P' \quad M[\theta] = M'}{(P M)[\theta] = (P' M')}$$

$$\frac{x \text{ not free in } \text{dom}(\theta) \cup \text{rng}(\theta) \quad A_1[\theta] = A'_1 \quad A_2[\theta] = A'_2}{(\Pi x:A_1. A_2)[\theta] = \Pi x:A'_1. A'_2}$$

Figure 2.3: Applying Substitutions to Types

$$\frac{}{\cdot[\theta] = \cdot} \quad \frac{x \text{ not free in } \text{dom}(\theta) \cup \text{rng}(\theta) \quad \Gamma[\theta] = \Gamma' \quad A[\theta] = A'}{(\Gamma, x : A)[\theta] = \Gamma', x : A'}$$

Figure 2.4: Applying Substitutions to Contexts

whether there is an R' or an M' and α' such that $R[\theta]^r = R'$ or $R[\theta]^r = M' : \alpha'$ is derivable, at most one of these judgements is derivable and that too for a unique R' , respectively, M' and α' .

The following property has an obvious proof by induction on the structure of the expression.

Theorem 2.5. *If E is a kind, a type or a canonical term none of whose free variables is a member of $\text{dom}(\theta)$, then $E[\theta] = E$ has as derivation. If R is an atomic term none of whose free variables is a member of $\text{dom}(\theta)$ then $R[\theta]^r = R$ has a derivation.*

Simultaneous hereditary substitution enjoys a permutation property that is similar to the one described in [6] for unitary substitution.

Theorem 2.6. *Let θ_1 be the substitution $\{\langle x_1, M_1, \alpha_1 \rangle, \dots, \langle x_n, M_n, \alpha_n \rangle\}$. Further, let θ_2 be the substitution $\{\langle y_1, N_1, \beta_1 \rangle, \dots, \langle y_m, N_m, \beta_m \rangle\}$ where y_1, \dots, y_m are variables that are distinct from x_1, \dots, x_n and that do not appear free in M_1, \dots, M_n . Finally, suppose that for each i , $1 \leq i \leq m$, there is some N'_i such that $N_i[\theta_1] = N'_i$ has a derivation and let $\theta_3 = \{\langle y_1, N'_1, \beta_1 \rangle, \dots, \langle y_m, N'_m, \beta_m \rangle\}$. For every kind, type and canonical term E , E_1 and E_2 such $E[\theta_1] = E_1$ and $E[\theta_2] = E_2$ have derivations, there must be an E' such that $E_2[\theta_1] = E'$ and $E_1[\theta_3] = E'$ have derivations.*

Proof. The proof proceeds by a primary induction on the sum of the sizes of θ_1 and θ_2 and a secondary induction on the derivation of $E[\theta_2] = E_2$. We omit the details which are similar to those for Lemma 2.10 in [6]. \square

$$\frac{c : \alpha \in \Theta}{\Theta \vdash_{\text{at}}^r c : \alpha} \quad \frac{x : \alpha \in \Theta}{\Theta \vdash_{\text{at}}^r x : \alpha} \quad \frac{\Theta \vdash_{\text{at}}^r R : \alpha' \rightarrow \alpha \quad \Theta \vdash_{\text{at}} M : \alpha'}{\Theta \vdash_{\text{at}}^r R M : \alpha}$$

$$\frac{\{x : \alpha_1\} \uplus \Theta \vdash_{\text{at}} M : \alpha_2}{\Theta \vdash_{\text{at}} \lambda x. M : \alpha_1 \rightarrow \alpha_2} \quad \frac{\Theta \vdash_{\text{at}}^r R : o}{\Theta \vdash_{\text{at}} R : o}$$

Figure 2.5: Arity Typing for Canonical Terms

While the application of a substitution to an LF expression may not always exist, this is guaranteed to be the case when certain arity typing constraints are satisfied as we describe below.

Definition 2.7. *An arity context Θ is a set of unique assignments of arity types to (term) constants and variables; these assignments are written as $x : \alpha$ or $c : \alpha$. Given two arity contexts Θ_1 and Θ_2 , we write $\Theta_1 \uplus \Theta_2$ to denote the collection of all the assignments in Θ_1 and the assignments in Θ_2 to the constants or variables not already assigned a type in Θ_1 . The rules in Figure 2.5 define the arity typing relation denoted by $\Theta \vdash_{\text{at}} M : \alpha$ between a term M and an arity type α relative to an arity context Θ . A kind or type E is said to respect an arity context Θ under the following conditions: if E is Type; if E is an atomic type and for each canonical term M appearing in E there is an arity type α such that $\Theta \vdash_{\text{at}} M : \alpha$ is derivable; and if E has the form $\Pi x:A. E'$ and A respects Θ and E' respects $\{x : (A)^-\} \uplus \Theta$. A context Γ is said to respect Θ if for every $x : A$ appearing in Γ it is the case that A respects Θ . A substitution θ is arity type preserving with respect to Θ if for every $\langle x, M, \alpha \rangle \in \theta$ it is the case that $\Theta \vdash_{\text{at}} M : \alpha$ is derivable. Associated with a substitution θ is the arity context $\{x : \alpha \mid \langle x, M, \alpha \rangle \in \theta\}$ that is denoted by $\text{ctx}(\theta)$.*

Theorem 2.8. *Let θ be a substitution that is arity type preserving with respect to Θ and let Θ' denote the arity context $\text{ctx}(\theta) \uplus \Theta$.*

1. If E is a canonical type or kind that respects the arity context Θ' , then there must be an E' that respects Θ and that is such that $E[[\theta]] = E'$ is derivable.
2. If M is a canonical term such that $\Theta' \vdash_{\text{at}} M : \alpha$ is derivable, then there must be an M' such that $M[[\theta]] = M'$ and $\Theta \vdash_{\text{at}} M' : \alpha$ are derivable.
3. If R is an atomic term such that $\Theta' \vdash_{\text{at}}^r R : \alpha$ is derivable, then either there is an atomic term R' such that $R[[\theta]] = R'$ and $\Theta \vdash_{\text{at}}^r R' : \alpha$ are derivable or there is a canonical term M such that $R[[\theta]] = M : \alpha$ and $\Theta \vdash_{\text{at}} M : \alpha$ are derivable.

Proof. The first clause in the theorem is an easy consequence of the second. We prove clauses (2) and (3) simultaneously by induction first on the sizes of substitutions and then on the structure of terms. The argument proceeds by considering the cases for the term structure, first proving (3) and then using this in proving (2). \square

We will often consider expressions and substitutions that satisfy the arity typing requirements of the theorem above, which then guarantees that the applications of the substitutions have results. We introduce a notation that is convenient in this situation: we will write $E[[\theta]]$ to denote the unique E' such that $E[[\theta]] = E'$ has a derivation whenever such a derivation is known to exist.

Definition 2.9. *Two substitutions θ_1 and θ_2 are said to be arity type compatible relative to the arity context Θ if θ_2 is type preserving with respect to Θ and θ_1 is type preserving with respect to $\text{ctx}(\theta_2) \uplus \Theta$. The composition of two such substitutions, written as $\theta_2 \circ \theta_1$, is the substitution*

$$\{\langle x, M', \alpha \rangle \mid \langle x, M, \alpha \rangle \in \theta_1 \text{ and } M[[\theta_2]] = M' \text{ has a derivation}\} \cup \{\langle y, N, \beta \rangle \mid \langle y, N, \beta \rangle \in \theta_2 \text{ and } y \notin \text{dom}(\theta_1)\}$$

By Theorem 2.8 there must be an M' for which $M[[\theta_2]] = M'$ has a derivation for each $\langle x, M, \alpha \rangle \in \theta_1$. Moreover such an M' must be unique. Thus, the composition described herein is well-defined. Note also that the composition must also be arity type preserving with respect to Θ .

Theorem 2.10. *Let θ_1 and θ_2 be substitutions that are arity type compatible relative to Θ and let Θ' denote the arity context $\text{ctx}(\theta_2 \circ \theta_1) \uplus \Theta$.*

1. If E is a canonical kind, type or context that respects Θ' and E' and E'' are, respectively, canonical types or kinds such that $E[[\theta_1]] = E'$ and $E'[[\theta_2]] = E''$ have derivations, then $E[[\theta_2 \circ \theta_1]] = E''$ has a derivation.
2. If M is a canonical term such that, for some arity type α , $\Theta' \vdash_{\text{at}} M : \alpha$ is derivable and M' and M'' are canonical terms such that $M[[\theta_1]] = M'$ and $M'[[\theta_2]] = M''$ have derivations, then $M[[\theta_2 \circ \theta_1]] = M''$ has a derivation.
3. If R is a canonical term such that, for some arity type α , $\Theta' \vdash_{\text{at}} R : \alpha$ is derivable and
 - (a) M' and M'' are canonical terms such that $R[[\theta_1]]^r = M' : \alpha$ and $M'[[\theta_2]] = M''$ have derivations, then $R[[\theta_2 \circ \theta_1]]^r = M'' : \alpha$ has a derivation;
 - (b) R' and M'' are, respectively, an atomic and a canonical term such that $R[[\theta_1]]^r = R'$ and $R'[[\theta_2]]^r = M'' : \alpha$ have derivations then $R[[\theta_2 \circ \theta_1]]^r = M'' : \alpha$ has a derivation;
 - (c) R' and R'' are atomic terms such that $R[[\theta_1]]^r = R'$ and $R'[[\theta_2]]^r = R''$ have derivations, then $R[[\theta_2 \circ \theta_1]]^r = R''$ has a derivation.

Proof. Clause 1 of the theorem follows easily from an induction on the structure of the canonical type or kind, assuming the property stated in clause 2.

We prove clauses 2 and 3 together. These clauses are premised on the existence of a derivation corresponding to the application of the substitution θ_1 to either M or R . The argument is by induction on the size of this derivation and it proceeds by considering the cases for the last rule in the derivation.

We consider first the cases where the derivation is for $M[\theta_1] = M'$; the clause in the theorem relevant to these cases is 2. An easy argument using the induction hypothesis yields the desired conclusion when M is of the form $\lambda x. M_1$. In the case that M is an atomic term, there is a shorter derivation for $M[\theta_1]^r = M' : \alpha$ or $M[\theta_1]^r = M'$. In the first case, the induction hypothesis, specifically clause 3(a), allows us to conclude that $M[\theta_2 \circ \theta_1] = M''$ has a derivation. In the second case, M' must be an atomic term and there must therefore be a derivation for $M'[\theta_2]^r = M'' : \alpha$ or $M'[\theta_2]^r = M''$. Using the induction hypothesis, specifically clause 3(b) or 3(c), we can again conclude that there must be a derivation for $M[\theta_2 \circ \theta_1] = M''$.

We consider next the cases for the last rule when the derivation is for $R[\theta_1]^r = M' : \alpha$.

- If M is a variable x such that $\langle x, M', \alpha \rangle \in \theta_1$, it must be the case that $\langle x, M'', \alpha \rangle \in \theta_2 \circ \theta_1$. Hence there must be a derivation for $M[\theta_2 \circ \theta_1]^r = M'' : \alpha$.
- Otherwise M must be of the form $(R_1 M_2)$ where $R_1[\theta_1] = \lambda x. M_3 : \alpha' \rightarrow \alpha$, $M_2[\theta_1] = M_4$, and $M_3[\{\langle x, M_4, \alpha' \rangle\}] = M'$ have derivations for suitable choices for M_3 , α' and M_4 . We note first that $(ctx(\theta_2 \circ \theta_1)) \uplus \Theta = ctx(\theta_1) \uplus (ctx(\theta_2) \uplus \Theta)$. Then, by the assumptions of the theorem and Theorem 2.8, it follows that there must be terms M'_3 and M'_4 such that $M_3[\theta_2] = M'_3$ and $M_4[\theta_2] = M'_4$ have derivations. We see by using the induction hypothesis with respect to the derivation for $R_1[\theta_1] = \lambda x. M_3 : \alpha' \rightarrow \alpha$ that there must be a derivation for $R_1[\theta_2 \circ \theta_1] = \lambda x. M'_3 : \alpha' \rightarrow \alpha$. Using the induction hypothesis again with respect to the derivation for $M_2[\theta_1] = M_4$, we see that there must be a derivation for $M_2[\theta_2 \circ \theta_1] = M'_4$. By Theorem 2.6 it follows that $M'_3[\{\langle x, M'_4, \alpha' \rangle\}] = M''$ has a derivation and, hence that $(R_1 M_2)[\theta_2 \circ \theta_1]^r = M'' : \alpha$ has one too.

Finally we consider the cases for the last rule when the derivation is for $R[\theta_1]^r = R'$. The argument when R is a constant is trivial. The case when R is a variable follows almost as immediately using the definition of $\theta_2 \circ \theta_1$. The only remaining case is when R is of the form $(R_1 M_2)$ and R' is $(R'_1 M'_2)$ where $R_1[\theta_1]^r = R'_1$ and $M_2[\theta_1] = M'_2$ have shorter derivations for suitable terms R'_1 and M'_2 . We then have two subcases to consider with respect to the application of θ_2 to $(R'_1 M'_2)$:

- There is a derivation for $(R'_1 M'_2)[\theta_2]^r = (R''_1 M''_2)$ where R''_1 and M''_2 are terms such that $R'_1[\theta_2]^r = R''_1$ and $M'_2[\theta_2]^r = M''_2$ have derivations; note that the relevant clause in this case is 3(c) and R'' is $(R''_1 M''_2)$. The induction hypothesis lets us conclude that $R_1[\theta_2 \circ \theta_1]^r = R''_1$ and $M_2[\theta_2 \circ \theta_1]^r = M''_2$ have derivations. Hence, $(R_1 M_2)[\theta_2 \circ \theta_1]^r = (R''_1 M''_2)$ must have a derivation.
- There is a derivation for $(R'_1 M'_2)[\theta_2]^r = M'' : \alpha$. In this case, for suitable choices for M_3 , α' and M_4 , there must be derivations for $R'_1[\theta_2]^r = \lambda x. M_3 : \alpha' \rightarrow \alpha$, $M'_2[\theta_2] = M_4$ and $M'_3[\{\langle x, M_4, \alpha' \rangle\}] = M''$. The induction hypothesis now lets us conclude that there are derivations for $R_1[\theta_2 \circ \theta_1]^r = \lambda x. M_3 : \alpha' \rightarrow \alpha$ and $M_2[\theta_2 \circ \theta_1] = M_4$. It then follows easily that there must be a derivation for $(R_1 M_2)[\theta_2 \circ \theta_1]^r = M'' : \alpha$.

□

The erased form of a type is invariant under substitution. This is the content of the theorem below whose proof is straightforward.

Theorem 2.11. *For any type A and substitution θ , if $A[\theta] = A'$ has a derivation, then $(A)^- = (A')^-$.*

2.1.3 Wellformedness Judgements

Canonical LF includes seven judgements: $\vdash \Sigma$ **sig** that ensures that the constants declared in a signature are distinct and their type or kind classifiers are well-formed; $\vdash_{\Sigma} \Gamma$ **ctx** that ensure that the variables declared in a signature are distinct and their type classifiers are well-formed in the preceding declarations and well-formed signature Σ ; $\Gamma \vdash_{\Sigma} K$ **kind** that determines that a kind K is well-formed with respect to a well-formed signature and context pair; $\Gamma \vdash_{\Sigma} A$ **type** and $\Gamma \vdash_{\Sigma} P \Rightarrow K$ that check, respectively, the formation of a canonical and atomic type relative to a well-formed signature, context and kind triple; and $\Gamma \vdash_{\Sigma} M \Leftarrow A$ and $\Gamma \vdash_{\Sigma} R \Rightarrow A$ that ensure, respectively, that a canonical and atomic term are well-formed with respect to a well-formed signature, context and canonical type triple. Figure 2.6 presents the rules for deriving these judgements. In the rules *CANON_KIND_PI* and *CANON_TERM_LAM* we assume x to be a variable that does not appear free in Γ . The formation rule for type and term level application, i.e. *ATOM_FAM_APP* and *ATOM_TERM_APP*, require the substitution of a term into a kind or a type. Use is made towards this end of hereditary substitution. The index for such a substitution is obtained by erasure from the type established for the term.

The judgement forms other than $\vdash \Sigma$ **sig** that are described above are parameterized by a signature that remains unchanged in the course of their derivation. In the rest of this paper we will assume a fixed signature that has in fact been verified to be well-formed at the outset. The judgement forms require some of their other components to satisfy additional restrictions. For example, judgements of the forms $\Gamma \vdash_{\Sigma} M \Leftarrow A$ and $\Gamma \vdash_{\Sigma} R \Rightarrow A$ require that Σ , Γ and A be well-formed as an ensemble. To be coherent, the rules in Figure 2.6 must ensure that in deriving a judgement that satisfies these requirements, it is necessary only to consider the derivation of judgements that also accord with these requirements. The fact that they possess this property can be verified by an inspection of their structure, using the observation that will be made in Theorem 2.17 that hereditary substitution preserves the property of being well-formed for kinds and types.

Arity typing judgements for terms approximate LF typing judgements as made precise below.

Definition 2.12. *The arity context induced by the signature Σ and context Γ is the collection of assignments that includes $x : (A)^-$ for each $x : A \in \Gamma$ and $c : (A)^-$ for each $c : A \in \Sigma$. When the context Γ is irrelevant or empty, we shall refer to the arity context as the one induced by just Σ .*

Theorem 2.13. *Let Θ be the arity context induced by the signature Σ and context Γ . If $\vdash_{\Sigma} \Gamma$ **ctx** then Γ respects Θ . If $\Gamma \vdash_{\Sigma} K$ **kind** or $\Gamma \vdash_{\Sigma} A$ **type** then, respectively, K or A respect Θ . If $\Gamma \vdash_{\Sigma} M \Leftarrow A$ is derivable, then $\Theta \vdash_{at} M : (A)^-$ must also be derivable. If $\Gamma \vdash_{\Sigma} R \Rightarrow A$ is derivable, then $\Theta \vdash_{at} R : (A)^-$ must also be derivable.*

Proof. The last two parts of the theorem are proved simultaneously by induction on the size of the derivation of $\Gamma \vdash_{\Sigma} M \Leftarrow A$ and $\Gamma \vdash_{\Sigma} R \Rightarrow A$. The first two parts follows from them, again by induction on the derivation size. \square

$\boxed{\vdash \Sigma \text{ sig}}$

$$\frac{}{\vdash \cdot \text{ sig}} \text{SIG_EMPTY} \quad \frac{\vdash \Sigma \text{ sig} \quad \cdot \vdash_{\Sigma} A \text{ type} \quad c \text{ does not appear in } \Sigma}{\vdash \Sigma, c : A \text{ sig}} \text{SIG_TERM}$$

$$\frac{\vdash \Sigma \text{ sig} \quad \cdot \vdash_{\Sigma} K \text{ kind} \quad a \text{ does not appear in } \Sigma}{\vdash \Sigma, a : K \text{ sig}} \text{SIG_FAM}$$

$\boxed{\vdash_{\Sigma} \Gamma \text{ ctx}}$

$$\frac{}{\vdash_{\Sigma} \cdot \text{ ctx}} \text{CTX_EMPTY} \quad \frac{\vdash_{\Sigma} \Gamma \text{ ctx} \quad \Gamma \vdash_{\Sigma} A \text{ type} \quad x \text{ does not appear free in } \Gamma}{\vdash_{\Sigma} \Gamma, x : A \text{ ctx}} \text{CTX_TERM}$$

$\boxed{\Gamma \vdash_{\Sigma} K \text{ kind}}$

$$\frac{}{\Gamma \vdash_{\Sigma} \text{Type kind}} \text{CANON_KIND_TYPE} \quad \frac{\Gamma \vdash_{\Sigma} A \text{ type} \quad \Gamma, x : A \vdash_{\Sigma} K \text{ kind}}{\Gamma \vdash_{\Sigma} \Pi x : A. K \text{ kind}} \text{CANON_KIND_PI}$$

$\boxed{\Gamma \vdash_{\Sigma} A \text{ type}}$

$$\frac{\Gamma \vdash_{\Sigma} P \Rightarrow \text{Type}}{\Gamma \vdash_{\Sigma} P \text{ type}} \text{CANON_FAM_ATOM} \quad \frac{\Gamma \vdash_{\Sigma} A_1 \text{ type} \quad \Gamma, x : A_1 \vdash_{\Sigma} A_2 \text{ type}}{\Gamma \vdash_{\Sigma} \Pi x : A_1. A_2 \text{ type}} \text{CANON_FAM_PI}$$

$\boxed{\Gamma \vdash_{\Sigma} P \Rightarrow K}$

$$\frac{a : K \in \Sigma}{\Gamma \vdash_{\Sigma} a \Rightarrow K} \text{ATOM_FAM_CONST}$$

$$\frac{\Gamma \vdash_{\Sigma} P \Rightarrow \Pi x : A. K_1 \quad \Gamma \vdash_{\Sigma} M \Leftarrow A \quad K_1[\{\{x, M, (A)^-\}\}] = K}{\Gamma \vdash_{\Sigma} P M \Rightarrow K} \text{ATOM_FAM_APP}$$

$\boxed{\Gamma \vdash_{\Sigma} M \Leftarrow A}$

$$\frac{\Gamma \vdash_{\Sigma} R \Rightarrow P}{\Gamma \vdash_{\Sigma} R \Leftarrow P} \text{CANON_TERM_ATOM} \quad \frac{\Gamma, x : A_1 \vdash_{\Sigma} M \Leftarrow A_2}{\Gamma \vdash_{\Sigma} \lambda x. M \Leftarrow \Pi x : A_1. A_2} \text{CANON_TERM_LAM}$$

$\boxed{\Gamma \vdash_{\Sigma} R \Rightarrow A}$

$$\frac{x : A \in \Gamma}{\Gamma \vdash_{\Sigma} x \Rightarrow A} \text{ATOM_TERM_VAR} \quad \frac{c : A \in \Sigma}{\Gamma \vdash_{\Sigma} c \Rightarrow A} \text{ATOM_TERM_CONST}$$

$$\frac{\Gamma \vdash_{\Sigma} R \Rightarrow \Pi x : A_1. A_2 \quad \Gamma \vdash_{\Sigma} M \Leftarrow A_1 \quad A_2[\{\{x, M, (A_1)^-\}\}] = A}{\Gamma \vdash_{\Sigma} R M \Rightarrow A} \text{ATOM_TERM_APP}$$

Figure 2.6: The Formation Rules for LF

$tp : \text{Type}$	$of_empty : of\ empty\ unit$
$unit : tp$	
$arr : tp \rightarrow tp$	$of_app : \Pi E_1:tm. \Pi E_2:tm. \Pi T_1:tp. \Pi T_2:tp.$ $\Pi D_1:of\ E_1\ (arr\ T_1\ T_2). \Pi D_2:of\ E_2\ T_1.$ $of\ (app\ E_1\ E_2)\ T_2$
$tm : \text{Type}$	
$empty : tm$	
$app : tm \rightarrow tm \rightarrow tm$	$of_lam : \Pi R:tm \rightarrow tm. \Pi T_1:tp. \Pi T_2:tp.$ $\Pi D:(\Pi x:tm. \Pi y:of\ x\ T_1. of\ (R\ x)\ T_2).$ $of\ (lam\ T_1\ (\lambda x. R\ x))\ (arr\ T_1\ T_2)$
$lam : tp \rightarrow (tm \rightarrow tm) \rightarrow tm$	
$of : tm \rightarrow tp \rightarrow \text{Type}$	
$eq : tp \rightarrow tp \rightarrow \text{Type}$	$refl : \Pi T:tp. eq\ T\ T$

Figure 2.7: An LF Specification for the Simply-Typed Lambda Calculus

2.2 Formalizing Object Systems in LF

A key use of LF is in formalizing systems that are described through relations between objects that are specified through a collection of inference rules. In the paradigmatic approach, each such relation is represented by a dependent type whose term arguments are encodings of objects that might be in the relation in question. The inference rules translate in this context into term constructors for the type representing the relation. We illustrate these ideas through an encoding of the typing relation for the simply-typed λ -calculus (STLC), a running example for this paper.

We assume the reader to be familiar with the types and terms in the STLC and also with the rules that define its typing relation. Figure 2.7 presents an LF signature that serves as an encoding of this system. This encoding uses the higher-order abstract syntax approach to treating binding. The specification introduces two type families, tp and tm to represent the simple types and λ -terms. Additionally, for each expression form in the object system, it includes a constant that produces a term of type tp or tm ; as should be apparent from the declarations, we have assumed an object language whose terms are constructed from a single constant of atomic type that is represented by the LF constant $empty$ and whose type is represented by the LF constant $unit$. This signature also provides a representation of two relations over object language expressions: typing between terms and types and equality between types. Specifically, the type-level constants of and eq are included towards this end. The rules defining the relations of interest in the object system are encoded by constants in the signature. The types associated with these constants ensure that well-formed terms of atomic type that are formed using the constants correspond to derivations of the relation in the object language that is represented by the type.

One of the purposes for constructing a specification is to use it to prove properties about the object system. For example, we may want to show that when a type can be associated with a term in the STLC, it must be unique. Based on our encoding, this property can be stated as the following about typing derivations in LF:

For any terms M_1, M_2, E, T_1, T_2 , if there are LF derivations for $\vdash_{\Sigma} E \Leftarrow tm, \vdash_{\Sigma} T_1 \Leftarrow tp, \vdash_{\Sigma} T_2 \Leftarrow tp, \vdash_{\Sigma} M_1 \Leftarrow of\ E\ T_1$ and $\vdash_{\Sigma} M_2 \Leftarrow of\ E\ T_2$, then there must be a term M_3 such than there is a derivation for $\vdash_{\Sigma} M_3 \Leftarrow eq\ T_1\ T_2$.

To prove this property, we would obviously need to unpack its logical structure. We would also need to utilize an understanding of LF in analyzing the hypothesized typing derivations corresponding to the STLC typing judgements. Considering the case where E is an abstraction will lead us to actually wanting to prove a more general property:

For any terms M_1, M_2, E, T_1, T_2 and contexts Γ , if there are LF derivations for the judgements $\Gamma \vdash_{\Sigma} E \Leftarrow tm$, $\Gamma \vdash_{\Sigma} T_1 \Leftarrow tp$, $\Gamma \vdash_{\Sigma} T_2 \Leftarrow tp$, $\Gamma \vdash_{\Sigma} M_1 \Leftarrow of E T_1$ and $\Gamma \vdash_{\Sigma} M_2 \Leftarrow of E T_2$, then there must be a term M_3 such that there is a derivation for $\vdash_{\Sigma} M_3 \Leftarrow eq T_1 T_2$.

Now, this property is not provable without some constraints on the form of contexts. In this example, it suffices to prove it when Γ is restricted to being of the form

$$(x_1 : tm, y_1 : of x_1 T y_1, \dots, x_n : tm, y_n : of x_n T y_n).$$

In completing the argument, we would need to use properties of LF derivability. A property that would be essential in this case is the finiteness of LF derivations, which enables us to use an inductive argument.

The objective in this paper is to provide a formal mechanism for carrying out such analysis. We do this by describing a logic that is suitable for this purpose. One of the requirements of this logic is that it should permit the expression of the kinds of properties that arise in the process of reasoning. Beyond this, it should further be possible to complement the statement of properties with inference rules that permit the encoding of interesting and sound forms of reasoning.

2.3 Meta-Theoretic Properties about LF Derivability

Our reasoning system will need to embody an understanding of derivability in LF. We describe some properties related to this notion here that will be useful in this context. The first three theorems, which express structural properties about derivations, have easy proofs. The fourth theorem states a substitutivity property for wellformedness judgements. This theorem is proved in [6].

Theorem 2.14. *If \mathcal{D} is a derivation for $\Gamma \vdash_{\Sigma} K \text{ kind}$, $\Gamma \vdash_{\Sigma} A \text{ type}$ or $\Gamma \vdash_{\Sigma} M \Leftarrow A$, then, for any variable x that is fresh to the judgement and for any A' such that $\Gamma \vdash_{\Sigma} A' \text{ type}$ is derivable, there is a derivation, respectively, for $\Gamma, x : A' \vdash_{\Sigma} K \text{ kind}$, $\Gamma, x : A' \vdash_{\Sigma} A \text{ type}$ or $\Gamma, x : A' \vdash_{\Sigma} M \Leftarrow A$ that has the same structure as \mathcal{D} .*

Theorem 2.15. *If \mathcal{D} is a derivation for judgements $\Gamma, x : A' \vdash_{\Sigma} K \text{ kind}$, $\Gamma, x : A' \vdash_{\Sigma} A \text{ type}$ or $\Gamma, x : A' \vdash_{\Sigma} M \Leftarrow A$ and x is a variable that does not appear free in K , A , or M and A respectively, then there must be a derivation that has the same structure as \mathcal{D} for $\Gamma \vdash_{\Sigma} K \text{ kind}$, $\Gamma \vdash_{\Sigma} A \text{ type}$ or $\Gamma \vdash_{\Sigma} M \Leftarrow A$, respectively.*

Theorem 2.16. *If x does not appear in A_2 then $\Gamma_1, y : A_2, x : A_1, \Gamma_3$ is a well-formed context with respect to a signature Σ whenever $\Gamma_1, x : A_1, y : A_2, \Gamma_3$ is. Further, if there is a derivation \mathcal{D} for $\Gamma, x : A_1, y : A_2, \Gamma_2 \vdash_{\Sigma} K \text{ kind}$, $\Gamma, x : A_1, y : A_2, \Gamma_2 \vdash_{\Sigma} A \text{ type}$ or $\Gamma, x : A_1, y : A_2, \Gamma_2 \vdash_{\Sigma} M \Leftarrow A$, then there must be a derivation that has the same structure as \mathcal{D} for $\Gamma, y : A_2, x : A_1, \Gamma_2 \vdash_{\Sigma} K \text{ kind}$, $\Gamma, y : A_2, x : A_1, \Gamma_2 \vdash_{\Sigma} A \text{ type}$ or $\Gamma, y : A_2, x : A_1, \Gamma_2 \vdash_{\Sigma} M \Leftarrow A$, respectively.*

Theorem 2.17. *Assume that $\vdash_{\Sigma} \Gamma_1, x_0 : A_0, \Gamma_2 \text{ ctx}$ and $\Gamma_1 \vdash_{\Sigma} M_0 \Leftarrow A_0$ have derivations, and let θ be the substitution $\{\langle x_0, M_0, (A_0)^{\neg} \rangle\}$. Then there is a Γ'_2 such that $\Gamma_2[\theta] = \Gamma'_2$ and $\vdash_{\Sigma} \Gamma_1, \Gamma'_2 \text{ ctx}$ have derivations. Further,*

1. *if $\Gamma_1, x_0 : A_0, \Gamma_2 \vdash_{\Sigma} K \text{ kind}$ has a derivation, then there is a K' such that $K[\theta] = K'$ and $\Gamma_1, \Gamma'_2 \vdash_{\Sigma} K' \text{ kind}$ have derivations;*
2. *if $\Gamma_1, x_0 : A_0, \Gamma_2 \vdash_{\Sigma} A \text{ type}$ has a derivation, then there is an A' such that $A[\theta] = A'$ and $\Gamma_1, \Gamma'_2 \vdash_{\Sigma} A' \text{ type}$ have derivations; and*

3. if $\Gamma_1, x_0 : A_0, \Gamma_2 \vdash_{\Sigma} M \Leftarrow A$ has a derivation (for some well-formed type A), there is an A' and an M' such that $A[[\theta]] = A'$, $M[[\theta]] = M'$, and $\Gamma_1, \Gamma_2' \vdash_{\Sigma} M' \Leftarrow A'$ have derivations.

The reasoning system will need to build in a means for analyzing typing derivations of the form $\Gamma \vdash_{\Sigma} M \Leftarrow A$. This analysis will be driven by the structure of the type A . The decomposition when A is of the form $\Pi x_1 : A_1. A_2$ has an obvious form. The development below, culminating in Theorem 2.20, provides the basis for the analysis when A is an atomic type.

Lemma 2.18. *Let Γ be a context such that $\vdash_{\Sigma} \Gamma \text{ ctx}$ has a derivation and let Θ be the arity context induced by Σ and Γ . Suppose that $\Pi y_1 : A_1. \dots \Pi y_n : A_n. A$ is a type associated with a (term) constant or variable by Σ or Γ , or that $\Pi y_1 : A_1. \dots \Pi y_n : A_n. K$ is a kind associated with a (type) constant by Σ , where the y_i s are distinct variables. Then, for $1 \leq i \leq n$, A_i and $\Pi y_i : A_i. \dots \Pi y_n : A_n. A$ or, respectively, $\Pi y_i : A_i. \dots \Pi y_n : A_n. K$ respect the arity context $\{y_1 : (A_1)^-, \dots, y_{i-1} : (A_{i-1})^-\} \uplus \Theta$. Further, A or, respectively, K respects the arity context $\{y_1 : (A_1)^-, \dots, y_n : (A_n)^-\} \uplus \Theta$.*

Proof. Since Σ and Γ are well-formed by assumption, depending on the case under consideration, either $\Gamma \vdash_{\Sigma} \Pi y_1 : A_1. \dots \Pi y_n : A_n. A \text{ type}$ or $\cdot \vdash_{\Sigma} \Pi y_1 : A_1. \dots \Pi y_n : A_n. K \text{ kind}$ must have a derivation. The desired conclusions now follow from Theorem 2.13 and Definition 2.7. \square

Lemma 2.19. *Let Γ_1 be a context such that $\vdash_{\Sigma} \Gamma_1 \text{ ctx}$ has a derivation, let Θ be the arity context induced by Σ and Γ_1 , and let θ be a substitution that is arity type preserving with respect to Θ . Further, let x_0 be a variable that is neither bound in Γ_1 nor a member of $\text{dom}(\theta)$, let A_0 and M_0 be such that $\Gamma_1 \vdash_{\Sigma} A_0 \text{ type}$ and $\Gamma_1 \vdash_{\Sigma} M_0 \Leftarrow A_0$ are derivable and let $\theta' = \theta \cup \{\langle x_0, M_0, (A_0)^- \rangle\}$.*

1. θ' is arity type preserving with respect to Θ .
2. Let Γ_2 be a context that respects an arity context Θ' such that $\text{ctx}(\theta') \uplus \Theta \subseteq \Theta'$ and let Γ_2' be a context such that $\Gamma_2[[\theta]] = \Gamma_2'$, and $\vdash_{\Sigma} \Gamma_1, x_0 : A_0, \Gamma_2' \text{ ctx}$ have derivations. Then there is a context Γ_2'' such that the following hold:

- (a) $\Gamma_2'[[\{\langle x_0, M_0, (A_0)^- \rangle\}]] = \Gamma_2''$, $\Gamma_2[[\theta']] = \Gamma_2''$ and $\vdash_{\Sigma} \Gamma, \Gamma_2'' \text{ ctx}$ have derivations;
- (b) if K is a kind that also respects Θ' and K' is a kind such that there are derivations for $K[[\theta]] = K'$ and $\Gamma_1, x_0 : A_0, \Gamma_2' \vdash_{\Sigma} K' \text{ kind}$, then there is a kind K'' such that $K'[[\{\langle x_0, M_0, (A_0)^- \rangle\}]] = K''$, $K[[\theta']] = K''$ and $\Gamma_1, \Gamma_2' \vdash_{\Sigma} K'' \text{ kind}$ are derivable; and
- (c) if A is a type that also respects Θ' and A' is a type such that there are derivations for $A[[\theta]] = A'$ and $\Gamma_1, x_0 : A_0, \Gamma_2' \vdash_{\Sigma} A' \text{ type}$, then there is a type A'' such that $A'[[\{\langle x_0, M_0, (A_0)^- \rangle\}]] = A''$, $A[[\theta']] = A''$ and $\Gamma_1, \Gamma_2' \vdash_{\Sigma} A'' \text{ type}$ have derivations.

Proof. Since $\Gamma_1 \vdash_{\Sigma} M_0 \Leftarrow A_0$ has a derivation, it follows from Theorem 2.13 that $\{\langle x_0, M_0, (A_0)^- \rangle\}$ is type preserving with respect to Θ . It then follows from the assumptions in the lemma that θ' is in fact $\{\langle x_0, M_0, (A_0)^- \rangle\} \circ \theta$ and type preserving with respect to Θ . The various observations in clause 2 now follow from Theorems 2.10 and 2.17. \square

Theorem 2.20. *Let Γ be a context such that $\vdash_{\Sigma} \Gamma \text{ ctx}$ has a derivation.*

1. $\Gamma \vdash_{\Sigma} R \Rightarrow A'$ has a derivation if
 - (a) R is of the form $(c M_1 \dots M_n)$ for some $c : \Pi y_1 : A_1. \dots \Pi y_n : A_n. A \in \Sigma$ or of the form $(x M_1 \dots M_n)$ for some $x : \Pi y_1 : A_1. \dots \Pi y_n : A_n. A \in \Gamma$,
 - (b) there is a sequence of types A'_1, \dots, A'_n such that, for $1 \leq i \leq n$, there are derivations for $A_i[[\{\langle y_1, M_1, (A_1)^- \rangle, \dots, \langle y_{i-1}, M_{i-1}, (A_{i-1})^- \rangle\}]] = A'_i$ and $\Gamma \vdash_{\Sigma} M_i \Leftarrow A'_i$, and

(c) $A[\{\langle y_1, M_1, (A_1)^- \rangle, \dots, \langle y_n, M_n, (A_n)^- \rangle\}] = A'$ and $\Gamma \vdash_\Sigma A'$ **type** have derivations.

2. $\Gamma \vdash_\Sigma R \Rightarrow A'$ has a derivation of height h only if

(a) R is of the form $(c M_1 \dots M_n)$ for some $c : \Pi y_1:A_1. \dots \Pi y_n:A_n. A \in \Sigma$ or of the form $(x M_1 \dots M_n)$ for some $x : \Pi y_1:A_1. \dots \Pi y_n:A_n. A \in \Gamma$,

(b) there is a sequence of types A'_1, \dots, A'_n such that, for $1 \leq i \leq n$, there is a derivation for $A_i[\{\langle y_1, M_1, (A_1)^- \rangle, \dots, \langle y_{i-1}, M_{i-1}, (A_{i-1})^- \rangle\}] = A'_i$ and a derivation of height less than h for $\Gamma \vdash_\Sigma M_i \Leftarrow A'_i$, and

(c) $A[\{\langle y_1, M_1, (A_1)^- \rangle, \dots, \langle y_n, M_n, (A_n)^- \rangle\}] = A'$ and $\Gamma \vdash_\Sigma A'$ **type** have derivations.

Proof. At the outset, we should check the coherence of clauses 1(b) and 2(b) in the theorem statement by verifying that, for $1 \leq i \leq n$, it is the case that $\Gamma \vdash_\Sigma A'_i$ **type** has a derivation. Towards this end, we first note that there must be a derivation for $\Gamma, y_1 : A_1, \dots, y_{i-1} : A_{i-1} \vdash_\Sigma A_i$ **type** since Σ and Γ are well-formed. The desired conclusion then follows from using Lemma 2.19 repeatedly and observing, via Theorem 2.11, that erasure is preserved under substitution.

We now introduce some notation that will be useful in the arguments that follow. We will use Θ to denote the arity context induced by Σ and Γ . Further, for $1 \leq i \leq n+1$, we will write θ_i for the substitution $\{\langle y_1, M_1, (A_1)^- \rangle, \dots, \langle y_{i-1}, M_{i-1}, (A_{i-1})^- \rangle\}$. An observation that we will make use of below is that if for $1 \leq j < i$ it is the case that $\Gamma \vdash_\Sigma M_j \Leftarrow A'_j$ has a derivation, then θ_i is type preserving with respect to Θ . This is an easy consequence of Theorems 2.13 and 2.11.

Proof of (1). We will consider explicitly only the case where R is $(c M_1 \dots M_n)$; the argument for the case when R is $(x M_1 \dots M_n)$ is similar. We will show for $1 \leq i \leq n+1$ that, under the conditions assumed for M_1, \dots, M_{i-1} , there is a type A''_i such that $(\Pi y_i:A_i. \dots \Pi x_n:A_n. A)[\theta_i] = A''_i$, $\Gamma \vdash_\Sigma A''_i$ **type** and $\Gamma \vdash_\Sigma (c M_1 \dots M_{i-1}) \Rightarrow A''_i$ have derivations. The desired conclusion follows from noting that A' must be A''_{n+1} because the result of substitution application is unique.

The claim is proved by induction on i . Consider first the case when i is 1. Since $\theta_1 = \emptyset$, A''_1 is $\Pi y_1:A_1. \dots \Pi y_n:A_n. A$. The wellformedness of Σ ensures that $\Gamma \vdash_\Sigma A''_1$ **type** has a derivation and we get a derivation for $\Gamma \vdash_\Sigma c \Rightarrow A'$ by using an *ATOM_TERM_CONST* rule.

Let us then assume the claim for i and show that it must also hold for $i+1$. By the hypothesis, there is an A''_i of the form $\Pi y_i:A'_i. A''$ where A'' is such that $\Pi y_{i+1}:A_{i+1}. \dots \Pi x_n:A_n. A[\theta_i] = A''$ has a derivation. Since $\Gamma \vdash_\Sigma A''_i$ **type** has a derivation, so must $\Gamma, y_i : A'_i \vdash_\Sigma A''$ **type**. By Lemma 2.18, $\Pi y_{i+1}:A_{i+1}. \dots \Pi x_n:A_n. A$ respects the arity context $\{y_1 : (A_1)^-, \dots, y_i : (A_i)^-\} \uplus \Theta$. Since there are derivations for $\Gamma \vdash_\Sigma M_j \Leftarrow A'_j$ for $1 \leq j < i$, θ_i is type preserving over Θ . We now invoke Lemma 2.19 to conclude that there is a term A''' such that there are derivations for $A''[\{\langle y_i, M_i, (A'_i)^- \rangle\}] = A'''$, $\Pi y_{i+1}:A_{i+1}. \dots \Pi x_n:A_n. A[\theta_{i+1}] = A'''$, and $\Gamma \vdash_\Sigma A'''$ **type**. By the hypothesis, there is a derivation for $\Gamma \vdash_\Sigma c M_1 \dots M_{i-1} \Rightarrow \Pi y_i:A'_i. A''$. Using an *ATOM_TERM_APP* rule together with this derivation and the ones for $\Gamma \vdash_\Sigma M_i \Leftarrow A'_i$, and $A''[\{\langle y_i, M_i, (A'_i)^- \rangle\}] = A'''$, we get a derivation for $\Gamma \vdash_\Sigma (c M_1 \dots M_i) \Rightarrow A'''$. Letting A''_{i+1} be A''' we see that all the requirements are satisfied.

Proof of (2). We prove the claim by induction on the height of the derivation of $\Gamma \vdash_\Sigma R \Rightarrow A'$. We consider the cases for the last rule used in the derivation. If this rule is *ATOM_TERM_VAR* or *ATOM_TERM_CONST*, the argument is straightforward. The only case to be considered further, then, is that when the rule is *ATOM_TERM_APP*.

In this case, we know that R must be of the form $(R' M')$ where there is a shorter derivation for $\Gamma \vdash_\Sigma R' \Rightarrow B'$ for some type B' . From the induction hypothesis, it follows that R' has the form $(c M_1 \dots M_n)$ or $(x M_1 \dots M_n)$ for some $c : \Pi y_1:A_1. \dots \Pi y_n:A_n. B \in \Sigma$ or $x : \Pi y_1:A_1. \dots \Pi y_n:A_n. B \in \Gamma$ and that there must be a sequence of types A'_1, \dots, A'_n that, together

with the terms M_1, \dots, M_n satisfy the requirements stated in clause 2(b). Moreover, B' must be such that $B[\theta_{n+1}] = B'$ and $\Gamma \vdash_{\Sigma} B' \text{ type}$ have derivations. Since the rule is an *ATOM_TERM_APP*, B' must have the structure of an abstracted type. From this it follows that B must be of the form $\Pi y_{n+1}:A_{n+1}. A$ and, correspondingly, B' must be of the form $\Pi y_{n+1}:A'_{n+1}. A''$ where $A_{n+1}[\theta_{n+1}] = A'_{n+1}$ and $A[\theta_{n+1}] = A''$ have derivations. Noting that the type of c or x is really of the form $\Pi y_1:A_1. \dots \Pi y_{n+1}:A_{n+1}. A$ it follows from Lemma 2.18 that A respects the arity context $\{y_1 : (A_1)^-, \dots, y_{n+1} : (A_{n+1})^-\} \uplus \Theta$. Also, since $\Gamma \vdash_{\Sigma} B' \text{ type}$ has a derivation, it must be the case that $\Gamma, y_{n+1} : A'_{n+1} \vdash_{\Sigma} A'' \text{ type}$ has one. Since the derivation concludes with a *ATOM_TERM_APP* rule, it must be the case that $\Gamma \vdash_{\Sigma} M' \Leftarrow A'_{n+1}$ and $A''[\{\langle y_{n+1}, M', (A'_{n+1})^- \rangle\}] = A'$ have shorter derivations than the one for $\Gamma \vdash_{\Sigma} R \Rightarrow A'$. Since θ_{n+1} is type preserving with respect to Θ , we may now use Lemma 2.19 and Theorem 2.11 to conclude that $A[\theta_{n+1} \cup \{\langle y_{n+1}, M_2, (A_{n+1})^- \rangle\}] = A'$ and $\Gamma \vdash_{\Sigma} A' \text{ type}$ have derivations. Renaming M' to M_{n+1} we see that all the requirements of clause 2 are satisfied. \square

Theorem 2.20 gives us an alternative means for deriving judgements of the form $\Gamma \vdash_{\Sigma} R \Leftarrow P$, in the process dispensing with judgements of the form $\Gamma \vdash_{\Sigma} R \Rightarrow A$. Note also that in *analyzing* judgements of the form $\Gamma \vdash_{\Sigma} R \Leftarrow P$, it is necessary to consider only *shorter* derivations for subterms of R . This observation will be used in developing a means for arguing inductively on the heights of LF derivations.

A property similar to that in Theorem 2.20 can be observed for wellformedness judgements for atomic types. Theorem 2.21 presents a version that suffices for this paper. A proof of this theorem can be constructed based essentially on the one for Theorem 2.20.

Theorem 2.21. *Let Γ be a context such that $\vdash_{\Sigma} \Gamma \text{ ctx}$ is derivable. Then $\Gamma \vdash_{\Sigma} P \Rightarrow K'$ has a derivation if and only if there is an $a : \Pi y_1:A_1. \dots \Pi y_n:A_n. K \in \Sigma$ such that*

1. P is of the form $(a M_1 \dots M_n)$;
2. there is a sequence of types A'_1, \dots, A'_n such that, for $1 \leq i \leq n$, there are derivations for $A_i[\{\langle y_1, M_1, (A_1)^-, \dots, \langle y_{i-1}, M_{i-1}, (A_{i-1})^- \rangle\}] = A'_i$ and $\Gamma \vdash_{\Sigma} M_i \Leftarrow A'_i$; and
3. $K[\{\langle y_1, M_1, (A_1)^-, \dots, \langle y_n, M_n, (A_n)^- \rangle\}] = K'$ and $\Gamma \vdash_{\Sigma} K' \text{ kind}$ have derivations.

3 A Logic for Expressing Properties of LF Specifications

We turn now to the task of designing a logic in which we can express properties of an object system that has been specified in LF. The discussions in Section 2.2 suggest a possible structure for such a logic. The logic would be parameterized by an LF signature that has been determined to be well-formed at the outset. The basic building blocks for the properties that are to be described would be typing judgements. More specifically, the logic would use such judgements as its atomic formulas and would interpret them using LF derivability. More complex formulas would be then be constructed using logical connectives and quantifiers over LF terms. As the example in Section 2.2 illustrates, it would be necessary to also permit a quantification over LF contexts.

To develop an actual logic based on these ideas, we need to describe a more precise correspondence between LF typing judgements and atomic formulas. The judgement forms that need to be considered in this context are those for typing canonical and atomic terms, i.e., the $\Gamma \vdash_{\Sigma} M \Leftarrow A$ and $\Gamma \vdash_{\Sigma} R \Rightarrow A$ forms. The main judgement form is in fact the first one: the second form serves mainly to explicate judgements of the first kind when the type is atomic and, as we have noted already, Theorem 2.20 provides the basis for circumventing such an explicit treatment through a

special “focused” typing rule. In light of this, it suffices to describe an encoding of only the first judgement form. The judgement in the LF setting assumes the wellformedness of the context Γ and the type A . In the logic, the context and, therefore, also the type can be dynamically determined by instantiations for context variables. To deal with this situation, we will build the wellformedness of Γ and A into the interpretation of the encoding of the judgement. There is, however, an aspect of the wellformedness checking that we would like to extract into a static pre-processing phase. The LF typing rules combine the checking of canonicity of terms with the determination of inhabitation that relies on the semantically more meaningful aspect of dependencies in types. To allow the focus in the logic to be on the latter aspect, we will build the former into a wellformedness criterion for formulas using arity types.

Another aspect that needs further consideration is the representation of LF contexts in atomic formulas. To support typing derivations that use the *CANON_TERM_LAM* rule, this representation must allow for the explicit association of types with variables. These variables may appear free in the terms and types in the atomic formula. However, their interpretation in this context must be different from the variables that are bound by quantifiers: in particular, these variables cannot be instantiated and each of them must be treated as being distinct within the atomic formula. The necessary treatment of these variables can be realized by representing them by *nominal constants* in the style of [4, 19]. Context expressions must, in addition, allow for an unspecified part whose exact extent is to be determined by instantiation of an external context quantifier. To support this ability, we will allow context variables to appear in these expressions. However, as observed in Section 2.2, we would like to be able to restrict the instantiation of such variables to blocks of declarations adhering to specified forms. To impose such constraints, the logic will permit context variables to be typed by *context schemas* that are motivated by regular world descriptions used in the Twelf system [11, 15].

In the rest of this section, we present a logic called \mathcal{L}_{LF} that substantiates the ideas outlined above. The first two subsections present the well-formed formulas of \mathcal{L}_{LF} and identify their intended meaning. The end result of this discussion is a means for describing properties of a specification given by an LF signature and for assessing the validity of such properties. The third subsection illuminates this capability through a collection of examples. The last subsection observes the counterpart in \mathcal{L}_{LF} of the property of irrelevance of the particular names that are chosen for the variables bound by the context in an LF judgement. The particular expression of this property takes the form of the invariance of validity of formulas under permutations of nominal constants.

3.1 The Formulas of the Logic

We begin by considering the representation of LF terms and types in \mathcal{L}_{LF} . Figure 3.1 presents the syntax of these expressions. As with LF syntax, we use c and d to represent term level constants, a and b to represent type level constants and x and y to represent term-level variables. We also use n to represent a special category of symbols called the nominal constants. LF terms and types are obviously a subset of the expressions presented here. Going the other way, there are two main additions to the LF counterparts in the collection of expressions described here. First, nominal constants may be used in constructing terms. Second, as we shall soon see, variables may be bound not only by term and type level abstractions but also by formula level quantifiers.

The logic \mathcal{L}_{LF} is parameterized by an LF style signature Σ that assigns kinds to type-level constants and types to term-level ones. This signature is assumed to be well-formed in the sense described in Section 2. The logic also assumes as given a set \mathcal{N} of nominal constants, each specified with an arity type, with a countably infinite supply of such constants for each arity type α . The types of the nominal constants are fixed once and for all by \mathcal{N} . This allows us to treat sets of such

Terms	$M, N ::= R \mid \lambda x. M$
Atomic Terms	$R ::= c \mid x \mid n \mid R M$
Types	$A ::= P \mid \Pi x:A_1. A_2$
Atomic Types	$P ::= a \mid P M$

Figure 3.1: Terms and Types in \mathcal{L}_{LF}

constants ambiguously as collections whose elements are of the form $n : \alpha$ or simply n .

$$\begin{array}{c}
\frac{a : K \in \Sigma}{\Theta \vdash_{ak}^p a : K} \qquad \frac{\Theta \vdash_{ak}^p P : \Pi x:A. K \quad \Theta \vdash_{at} M : (A)^-}{\Theta \vdash_{ak}^p P M : K} \\
\frac{\Theta \vdash_{ak}^p P : \text{Type}}{\Theta \vdash_{ak} P \text{ type}} \qquad \frac{\Theta \vdash_{ak} A_1 \text{ type} \quad \{x : (A_1)^-\} \uplus \Theta \vdash_{ak} A_2 \text{ type}}{\Theta \vdash_{ak} \Pi x:A_1. A_2 \text{ type}}
\end{array}$$

Figure 3.2: Arity Kinding for Canonical Types

As explained earlier, expressions in \mathcal{L}_{LF} will be expected to satisfy typing constraints that check for canonicity. At the term level, these constraints will be realized through arity typing relative to a suitable arity context. At the type level, we must additionally ensure that (type) constants have been supplied with an adequate number of arguments. We make these notions precise below; we assume the obvious extension of erasure to types in \mathcal{L}_{LF} here and elsewhere.

Definition 3.1. *The typing relation between an arity context, a term and an arity type that is described in Definition 2.7 is extended to the present context by permitting terms to contain nominal constants and by allowing arity contexts to contain assignments to such constants. The rules in Figure 3.2 define an arity kinding property denoted by $\Theta \vdash_{ak} A$ type for a type A relative to an arity context Θ . In these rules, Σ is the signature parameterizing \mathcal{L}_{LF} . We will often need to refer to the arity context induced by Σ . We call this the initial constant context and we reserve the symbol Θ_0 to denote it.*

Hereditary substitution extends naturally to the terms and types in \mathcal{L}_{LF} by treating nominal constants like other constants. The following theorem relating to such substitutions has an obvious proof.

Theorem 3.2. *If θ is type preserving with respect to Θ and $ctx(\theta) \uplus \Theta \vdash_{ak} A$ type and $A[\theta] = A'$ have derivations, then $\Theta \vdash_{ak} A'$ type has a derivation.*

Block Declarations	$\Delta ::= \cdot \mid \Delta, y : A$
Block Schema	$\mathcal{B} ::= \{x_1 : \alpha_1, \dots, x_n : \alpha_n\} \Delta$
Context Schema	$\mathcal{C} ::= \cdot \mid \mathcal{C}, \mathcal{B}$

Figure 3.3: Block Schemas and Context Schemas

The logic allows for quantifiers over LF contexts. In the intended interpretation, such quantifiers are meant to be instantiated with context expressions that assign LF types to nominal constants. However, it will be necessary to be able to constrain the possible instantiations in real applications.

This ability is supported by typing context quantifiers using *context schemas* whose structure is presented in Figure 3.3. In essence, a context schema comprises a collection of *block schemas*. A block schema consists of a header of variables annotated with arity types and a body of declarations associating types with variables. Each variable in the header and that is assigned a type in the body of a block schema is required to be distinct. A block is intended to serve as a template for generating a sequence of bindings for nominal constants through an instantiation process that will be made clear in the next subsection. A context expression corresponding to a context schema is to be obtained by some number of instantiations of its block schemas. Block and context schemas are required to satisfy typing constraints towards ensuring that the context expressions generated from them will be well-formed in the manner required by the logic. These constraints are represented by the typing judgements $\vdash \mathcal{B}$ *blk schema* and $\vdash \mathcal{C}$ *ctx schema*, respectively, that are defined by the rules in Figure 3.4.

$$\begin{array}{c}
\frac{}{\Theta \vdash_{dec} \cdot \Rightarrow \Theta} \qquad \frac{\Theta \vdash_{dec} \Delta \Rightarrow \Theta' \quad y \text{ is not assigned by } \Theta' \quad \Theta' \vdash_{ak} A \text{ type}}{\Theta \vdash_{dec} \Delta, y : A \Rightarrow \Theta' \cup \{y : (A)^-\}} \\
\\
\frac{x_1, \dots, x_n \text{ are distinct variables} \quad \Theta_0 \cup \{x_1 : \alpha_1, \dots, x_n : \alpha_n\} \vdash_{dec} \Delta \Rightarrow \Theta'}{\vdash \{x_1 : \alpha_1, \dots, x_n : \alpha_n\} \Delta \text{ blk schema}} \\
\\
\frac{}{\vdash \cdot \text{ ctx schema}} \qquad \frac{\vdash \mathcal{C} \text{ ctx schema} \quad \vdash \mathcal{B} \text{ blk schema}}{\vdash \mathcal{C}, \mathcal{B} \text{ ctx schema}}
\end{array}$$

Figure 3.4: Wellformedness Judgements for Block and Context Schemas

$$\begin{array}{ll}
\textbf{Context Expressions} & G ::= \cdot \mid \Gamma \mid G, n : A \\
\textbf{Formulas} & F ::= \{G \vdash M : A\} \mid \top \mid \perp \mid F_1 \supset F_2 \mid F_1 \wedge F_2 \mid \\
& F_1 \vee F_2 \mid \Pi \Gamma : \mathcal{C}.F \mid \forall x : \alpha.F \mid \exists x : \alpha.F
\end{array}$$

Figure 3.5: The Formulas of \mathcal{L}_{LF}

We are finally in a position to describe the formulas of \mathcal{L}_{LF} . The syntax of these formulas is presented in Figure 3.5. The symbol Γ is used in these formulas to represent context variables. Atomic formulas, which represent LF typing judgements, have the form $\{G \vdash M : A\}$. The context in these formulas is constituted by a sequence of type associations with nominal constants, possibly preceded by a context variable. Included in the collection are the logical constants \top and \perp and the familiar connectives for constructing more complex formulas. Universal and existential quantification over term variables is also permitted and these are written as $\forall x : \alpha.F$ and $\exists x : \alpha.F$, respectively. Such quantification is indexed, as might be expected, by arity types. The collection also includes universal quantification over context variables that is typed by context schemas, written as $\Pi \Gamma : \mathcal{C}.F$. We assume the usual principle of equivalence under renaming with respect to the term and context quantifiers and apply them as needed.

A formula F is determined to be well-formed or not relative to an arity context Θ and a collection of context variables Ξ . This judgement is written concretely as $\Theta; \Xi \vdash F$ *fmla* and the rules defining it are presented in Figure 3.6. At the top-level, formulas are expected to be *closed*, i.e., to not have any free term or context variables. More specifically, we expect $\mathcal{N} \cup \Theta_0; \emptyset \vdash F$ *fmla* to be derivable for such formulas. The analysis within the scope of term and context quantifiers augments these sets in the expected way. For context quantifiers, this analysis must also check

$$\begin{array}{c}
\frac{}{\Theta; \Xi \vdash \cdot \text{ context}} \quad \frac{\Gamma \in \Xi}{\Theta; \Xi \vdash \Gamma \text{ context}} \\
\frac{\Theta; \Xi \vdash G \text{ context} \quad n : (A)^- \in \Theta \quad \Theta \vdash_{ak} A \text{ type}}{\Theta; \Xi \vdash G, n : A \text{ context}} \\
\frac{\Theta; \Xi \vdash G \text{ context} \quad \Theta \vdash_{ak} A \text{ type} \quad \Theta \vdash_{at} M : (A)^-}{\Theta; \Xi \vdash \{G \vdash M : A\} \text{ fmla}} \\
\frac{}{\Theta; \Xi \vdash \top \text{ fmla}} \quad \frac{}{\Theta; \Xi \vdash \perp \text{ fmla}} \quad \frac{\Theta; \Xi \vdash F_1 \text{ fmla} \quad \Theta; \Xi \vdash F_2 \text{ fmla}}{\Theta; \Xi \vdash F_1 \bullet F_2 \text{ fmla}} \quad \bullet \in \{\supset, \wedge, \vee\} \\
\frac{\vdash \mathcal{C} \text{ ctx schema} \quad \Theta; \Xi \cup \{\Gamma\} \vdash F \text{ fmla}}{\Theta; \Xi \vdash \Pi \Gamma : \mathcal{C}.F \text{ fmla}} \quad \frac{\{x : \alpha\} \uplus \Theta; \Xi \vdash F \text{ fmla}}{\Theta; \Xi \vdash Qx : \alpha.F \text{ fmla}} \quad Q \in \{\forall, \exists\}
\end{array}$$

Figure 3.6: The Wellformedness Judgement for Formulas

that the annotating context schema is well-formed. An atomic formula $\{G \vdash M : A\}$ is deemed well-formed if its components G , M and A are well-formed and if M can be assigned the erased form of A as its arity type. The context expression G is well-formed if any context variable used in it is bound in the overall formula and if the types assigned to nominal constants in the explicit part of G are well-formed and such that their erased forms match the arity types of the nominal constants they are assigned to. Note that these types may use nominal constants without paying attention to dependency ordering; assessing whether they are used in a manner that respects this ordering is a part of the meaning of the atomic formula.

The following theorem, whose proof is obvious, shows that the wellformedness judgement for formulas continues to hold under the augmentation of the two contexts that parameterize it.

Theorem 3.3. *If $\Theta; \Xi \vdash F$ fmla has a derivation and $\Theta \subseteq \Theta'$ and $\Xi \subseteq \Xi'$, then $\Theta'; \Xi' \vdash F$ fmla also has a derivation.*

3.2 The Interpretation of Formulas

A key component to understanding the meanings of formulas is understanding the interpretation of the quantifiers over term and context variables. These quantifiers are intended to range over closed expressions of the relevant categories. For a quantifier over a term variable, this translates concretely into closed terms of the relevant arity type. For a quantifier over a context variable, we must first explain when a context expression satisfies a context schema.

We do this by describing the relation of “being an instance of” between a closed context expression G and a context schema \mathcal{C} . This relation is indexed by a nominal constant context \mathbb{N} that is a subset of \mathcal{N} and a *term variables context* Ψ that identifies a finite collection of such variables together with their arity types: in combination with the constants in Θ_0 , these collections, circumscribe the symbols that can be used in the declarations in the context expressions.² The relation is written as $\mathbb{N}; \Psi \vdash \mathcal{C} \rightsquigarrow_{cs} G$ and it is defined by the rules in Figure 3.7. This relation is defined via the repeated use of a “one-step” instantiation relation written as $\mathbb{N}; \Psi \vdash \mathcal{C} \rightsquigarrow_{cs}^1 G$; note that by G, G' we mean a context expression that is obtained by adding the bindings corresponding to G' in

²In determining closed instances of context schemas, \mathbb{N} will be \mathcal{N} and Ψ will be the empty set. The more general form for this relation, which includes a parameterization by these sets, will be useful in later sections.

$$\begin{array}{c}
\frac{\mathbb{N} \vdash \Delta \rightsquigarrow_{dec} G \bowtie \theta \quad n : (A)^- \in \mathbb{N} \quad A[\theta] = A'}{\mathbb{N} \vdash \Delta, y : A \rightsquigarrow_{dec} G, n : A' \bowtie \theta \cup \{\langle y, n, (A)^- \rangle\}} \\
\\
\frac{\mathbb{N} \vdash \Delta \rightsquigarrow_{dec} G' \bowtie \theta \quad \{\mathbb{N} \cup \Psi \cup \Theta_0 \vdash_{at} t_i : \alpha_i \mid 1 \leq i \leq n\} \quad G'[\{\langle x_i, t_i, \alpha_i \rangle \mid 1 \leq i \leq n\}] = G}{\mathbb{N}; \Psi \vdash \{x_1 : \alpha_1, \dots, x_n : \alpha_n\} \Delta \rightsquigarrow_{bs} G} \\
\\
\frac{\mathbb{N}; \Psi \vdash \mathcal{B} \rightsquigarrow_{bs} G \quad \mathbb{N}; \Psi \vdash \mathcal{C} \rightsquigarrow_{cs}^1 G}{\mathbb{N}; \Psi \vdash \mathcal{C}, \mathcal{B} \rightsquigarrow_{cs}^1 G} \\
\\
\frac{\mathbb{N}; \Psi \vdash \mathcal{C} \rightsquigarrow_{cs} G \quad \mathbb{N}; \Psi \vdash \mathcal{C} \rightsquigarrow_{cs}^1 G'}{\mathbb{N}; \Psi \vdash \mathcal{C} \rightsquigarrow_{cs} G, G'} \\
\\
\frac{}{\mathbb{N}; \Psi \vdash \mathcal{C} \rightsquigarrow_{cs} \cdot}
\end{array}$$

Figure 3.7: Instantiating a Context Schema

front of those in G . The definition of the one-step instantiation relation for context schemas uses an auxiliary judgement $\mathbb{N}; \Psi \vdash \mathcal{B} \rightsquigarrow_{bs} G$ that denotes the relation of “being an instance of” between a block schema and a context expression fragment. This relation holds when the context expression is obtained by generating a sequence of bindings for nominal constants from \mathbb{N} using the body of the block schema and then instantiating the variables in the header of the block schema with terms of the right arity types. The former task is realized through the relation $\mathbb{N} \vdash \Delta \rightsquigarrow_{dec} G \bowtie \theta$ that holds between a block of declarations Δ , a context expression G that is obtained by replacing the variables assigned in Δ with suitable nominal constants, and a substitution θ that corresponds to this replacement. We assume here and elsewhere that the application of a hereditary substitution to a sequence of declarations corresponds to its application to the type in each assignment.

Theorem 3.4. *Let \mathcal{C} and G be a context schema and a context expression such that $\vdash \mathcal{C}$ ctx schema and $\mathbb{N}; \Psi \vdash \mathcal{C} \rightsquigarrow_{cs} G$ are derivable. Then for any arity context Θ such that $\mathbb{N} \cup \Psi \cup \Theta_0 \subseteq \Theta$, it is the case that $\Theta; \emptyset \vdash G$ context has a derivation.*

Proof. We first show that for any block declaration Δ and any arity context Θ such that $\mathbb{N} \subseteq \Theta$, if $\Theta \vdash_{dec} \Delta \Rightarrow \Theta'$ and $\mathbb{N} \vdash \Delta \rightsquigarrow_{dec} G' \bowtie \theta'$ are derivable for some Θ' and θ' , then (a) θ' is type preserving with respect to Θ , (b) Θ' is $ctx(\theta') \uplus \Theta$, and (c) each binding in G' is of the form $n : A$ where $n : (A)^- \in \Theta$ and $\Theta \vdash_{ak} A$ type has a derivation. This claim is proved by induction on the derivation of $\Theta \vdash_{dec} \Delta \Rightarrow \Theta'$; properties (a) and (b) are included in the claim because they are useful together with Theorem 3.2 in showing property (c) in the induction step. Next we show, through an easy inductive argument, that if $\Theta_0 \cup \{x_1 : \alpha_1, \dots, x_n : \alpha_n\} \vdash_{dec} \Delta \Rightarrow \Theta'$ has a derivation and Θ is such that $\mathbb{N} \cup \Psi \cup \Theta_0 \subseteq \Theta$, then, for some Θ'' , it is the case that $\{x_1 : \alpha_1, \dots, x_n : \alpha_n\} \uplus \Theta \vdash_{dec} \Delta \Rightarrow \Theta''$ has a derivation. Using Theorem 3.2 with these two observations, we can show easily that if $\vdash \{x_1 : \alpha_1, \dots, x_n : \alpha_n\} \Delta$ blk schema and $\mathbb{N}; \Psi \vdash \{x_1 : \alpha_1, \dots, x_n : \alpha_n\} \Delta \rightsquigarrow_{bs} G$ have derivations then for each binding of the form $n : A$ in G it is the case that $n : (A)^- \in \Theta$ and $\Theta \vdash_{ak} A$ type. The theorem follows easily from this observation. \square

In defining validity for formulas, we will need to consider substitutions for context and term variables. Context variables substitutions have the form $\{G_1/\Gamma_1, \dots, G_n/\Gamma_n\}$ where, for $1 \leq i \leq n$, Γ_i is a context variable and G_i is a context expression. If σ is such a substitution, we will write $dom(\sigma)$ to denote the set $\{\Gamma_1, \dots, \Gamma_n\}$. Further, the application of σ to a formula F , which is denoted by $F[\sigma]$, will correspond to the replacement of the free occurrences of the variables $\Gamma_1, \dots, \Gamma_n$

in F by the corresponding context expressions, renaming bound context variables appearing in F away from those appearing in G_1, \dots, G_n . For term variables, the replacement must also ensure the transformation of the resulting expression to normal form. Towards this end, we adapt hereditary substitution to formulas. The application of this substitution simply distributes over quantifiers and logical symbols, respecting the scopes of quantifiers through the necessary renaming. The application to the atomic formula $\{G \vdash M : A\}$ also distributes to the component parts. We have already discussed the application to terms and types. The application to context expressions leaves context variables unaffected and simply distributes to the types in the explicit bindings. Note that no check is mandated in the process for clashes in the names of nominal constants appearing in the context expression being substituted into and the substitution terms. In this respect, this application is unlike that to LF contexts that is defined in Figure 2.4.

Theorem 3.5. *Let Θ be an arity context and let Ξ be a collection of context variables.*

1. *If θ is a term variables substitution that is arity type preserving with respect to Θ and F is a formula such that there is a derivation for $\text{ctx}(\theta) \uplus \Theta; \Xi \vdash F$ fmla, then there is a unique formula F' such that $F[\theta] = F'$ has a derivation. Moreover, for this F' it is the case that $\Theta; \Xi \vdash F'$ fmla is derivable.*
2. *If $\sigma = \{G_1/\Gamma_1, \dots, G_n/\Gamma_n\}$ is a context variables substitution which is such that all judgements in the collection $\{\Theta; \Xi \setminus \{\Gamma_1, \dots, \Gamma_n\} \vdash G_i \text{ context} \mid 1 \leq i \leq n\}$ are derivable and F is a formula such that there is a derivation for $\Theta; \Xi \vdash F$ fmla, then there is a derivation for $\Theta; \Xi \setminus \{\Gamma_1, \dots, \Gamma_n\} \vdash F[\sigma]$ fmla.*

Proof. The first clause follows from an induction on the derivation of $\text{ctx}(\theta) \uplus \Theta; \Xi \vdash F$ fmla, using Theorems 2.4 and 2.8 in the atomic case to ensure the appropriate arity typing judgements will be derivable under the substitution θ . The second clause follows from an induction on the derivation of $\Theta; \Xi \vdash F$ fmla, using the assumption that $\Theta; \Xi \setminus \{\Gamma_1, \dots, \Gamma_n\} \vdash G_i$ context is derivable to ensure wellformedness under the substitution σ in the atomic case. \square

Following the notation introduced after Theorem 2.8, if F and θ are a formula and a substitution that together satisfy the requirements of the first part of the theorem, we will write $F[\theta]$ to denote the F' for which $F[\theta] = F'$ is derivable. As is implicit in the preceding discussion, term and context variables substitutions may introduce new nominal constants. If θ is a term variables substitution, we will write $\text{supp}(\theta)$ to denote the collection of such constants that appear in the terms in $\text{rng}(\theta)$. Similarly, if σ is the context variables substitution $\{G_1/\Gamma_1, \dots, G_n/\Gamma_n\}$, we will write $\text{supp}(\sigma)$ to denote the collection of nominal constants that appear in G_1, \dots, G_n .

A closed atomic formula of the form $\{G \vdash M : A\}$ is intended to encode an LF judgement of the form $\Gamma \vdash_{\Sigma} M \Leftarrow A$. In this encoding, nominal constants that appear in terms represent free variables for which bindings appear in the context in LF judgements. To substantiate this interpretation, the rules *CANON_KIND_PI*, *CANON_FAM_PI* and *CANON_TERM_LAM* must introduce fresh nominal constants into contexts in typing derivations and they must replace bound variables appearing in terms and types with these constants. We use this interpretation to define validity for closed atomic formulas with one further qualification: unlike in the LF judgement, for the atomic formula we must also ascertain the wellformedness of the context and the type. This notion of validity is then extended to all closed formulas by recursion on formula structure.

Definition 3.6. *Let F be a formula such that $\mathcal{N} \cup \Theta_0; \emptyset \vdash F$ fmla is derivable.*

- *If F is $\{G \vdash M : A\}$, then it is valid exactly when all of $\vdash_{\Sigma} G$ ctx, $G \vdash_{\Sigma} A$ type, and $G \vdash_{\Sigma} M \Leftarrow A$ are derivable in LF, under the interpretation of nominal constants as variables*

bound in a context and with the modification of the rules `CANON_KIND_PI`, `CANON_FAM_PI` and `CANON_TERM_LAM` to introduce fresh nominal constants into contexts and to instantiate the relevant bound variables in kinds, types and terms with these constants.

- If F is \top it is valid and if it is \perp it is not valid.
- If F is $F_1 \supset F_2$, it is valid if F_2 is valid in the case that F_1 is valid.
- If F is $F_1 \wedge F_2$, it is valid if both F_1 and F_2 are valid.
- If F is $F_1 \vee F_2$, it is valid if either F_1 or F_2 is valid.
- If F is $\Pi \Gamma : C.F$, it is valid if $F[\{G/\Gamma\}]$ is valid for every G such that $\mathcal{N}; \emptyset \vdash_{\mathcal{C}} G \rightsquigarrow_{cs} G$ is derivable.
- If F is $\forall x : \alpha.F$, it is valid if $F[\{\{x, M, \alpha\}\}]$ is valid for every M such that $\mathcal{N} \cup \Theta_0 \vdash_{at} M : \alpha$ is derivable.
- If F is $\exists x : \alpha.F$, it is valid if $F[\{\{x, M, \alpha\}\}]$ is valid for some M such that $\mathcal{N} \cup \Theta_0 \vdash_{at} M : \alpha$ is derivable.

Theorems 3.4 and 3.5 ensure the coherence of this definition.

3.3 Understanding the Notion of Validity

In the examples we consider below, we assume an instantiation of \mathcal{L}_{LF} based on the signature presented in Section 2.2. Obviously, any LF typing judgement based on that signature is expressible in the logic. Moreover, the corresponding formula will be valid exactly when the typing judgement is derivable in LF. Thus, the formulas $\{\cdot \vdash \text{empty} : tm\}$, $\{\cdot \vdash (\text{lam unit } (\lambda x.x)) : tm\}$ and $\{n : tm \vdash n : tm\}$ are all valid. Similarly, the formulas $\exists d : o. \{\cdot \vdash d : (\text{of empty unit})\}$ and

$$\exists d : o. \{\cdot \vdash d : (\text{of } (\text{lam unit } (\lambda x.x)) (\text{arr unit unit}))\}$$

are valid but the formula $\exists d : o. \{\cdot \vdash d : (\text{of } (\text{lam unit } (\lambda x.x)) \text{unit})\}$ is not. Note that the arity type associated with the quantified variable in each of these formulas provides only a rough constraint on the instantiation needed to verify the validity of the formula; to do this, the instance must also satisfy LF typeability requirements represented by formula that appears within the scope of the quantifier.

Wellformedness conditions for formulas ensure only that the terms appearing within formulas satisfy canonicity requirements, i.e. that these terms are in β -normal form and that variables and constants are applied to as many arguments as they can take. Arity typing does not distinguish between terms in different expression categories. For example, the formula

$$\exists d : o. \{\cdot \vdash d : (\text{of } (\text{lam empty } (\lambda x.x)) (\text{arr unit unit}))\}$$

is well-formed but not valid. An alternative design choice, with equivalent consequences from the perspective of the valid properties that can be expressed in the logic, might have been to let the fact that `lam` is ill-applied to `empty` to impact on the wellformedness of the formula. The wellformedness conditions do not also enforce a distinctness requirement for bindings in a context. Thus, the formula $\{n : tm, n : tp \vdash \text{empty} : tm\}$ is well-formed. However, it is not valid because $\vdash_{\Sigma} n : tm, n : tp \text{ ctx}$ is not derivable in LF under the described interpretation for nominal constants. An implication of these observations is that a naive form of weakening does not hold with respect to the

encoding of LF derivability in \mathcal{L}_{LF} ; additional conditions similar to this described in Theorem 2.14 must be verified for this principle to apply.

To provide a more substantive example of the kinds of properties that can be expressed in \mathcal{L}_{LF} , let us consider the formal statement of the property of uniqueness of type assignments for the STLC. As noted in Section 2.2, this property is best described in a form that considers typing expressions in contexts that have a particular kind of structure. That structure can be formalized in \mathcal{L}_{LF} by a context schema comprising the single block

$$\{t : o\}x : tm, y : of\ x\ t.$$

Let us denote this context schema by c . Observe that a context that instantiates this schema will not provide a variable that can be used to construct an atomic term of type tp . Thus, the strengthening property for expressions representing types that is expressed by the formula

$$\Pi \Gamma : c. \forall t : o. \{\Gamma \vdash t : tp\} \supset \{\cdot \vdash t : tp\}.$$

should hold. We can in fact easily show this formula to be valid by using Theorem 2.20 and an induction on the height of the derivation for $\{G \vdash t : tp\}$ for a closed term t and a closed instance G of c . Using the validity of this formula, we can also easily argue that the following formula that expresses a strengthening property pertaining to the equality of types is also valid:

$$\Pi \Gamma : c. \forall d : o. \forall t_1 : o. \forall t_2 : o. \{\Gamma \vdash d : eq\ t_1\ t_2\} \supset \{\cdot \vdash d : eq\ t_1\ t_2\}.$$

The property of uniqueness of type assignments for the STLC can be expressed through the following formula:

$$\begin{aligned} \Pi \Gamma : c. \forall e : o. \forall t_1 : o. \forall t_2 : o. \forall d_1 : o. \forall d_2 : o. \\ \{\Gamma \vdash e : tm\} \supset \{\Gamma \vdash t_1 : tp\} \supset \{\Gamma \vdash t_2 : tp\} \supset \\ \{\Gamma \vdash d_1 : of\ e\ t_1\} \supset \{\Gamma \vdash d_2 : of\ e\ t_2\} \supset \exists d_3 : o. \{\cdot \vdash d_3 : eq\ t_1\ t_2\}. \end{aligned}$$

This formula can be seen to be valid using the strengthening property just described if we can establish the validity of the formula

$$\begin{aligned} \Pi \Gamma : c. \forall e : o. \forall t_1 : o. \forall t_2 : o. \forall d_1 : o. \forall d_2 : o. \\ \{\Gamma \vdash e : tm\} \supset \{\Gamma \vdash t_1 : tp\} \supset \{\Gamma \vdash t_2 : tp\} \supset \\ \{\Gamma \vdash d_1 : of\ e\ t_1\} \supset \{\Gamma \vdash d_2 : of\ e\ t_2\} \supset \exists d_3 : o. \{\Gamma \vdash d_3 : eq\ t_1\ t_2\}. \end{aligned}$$

To show this, it suffices to argue that, for a closed context expression G that instantiates the schema c and for closed expressions d_1 , d_2 , e , t_1 , and t_2 , if the formulas $\{G \vdash e : tm\}$, $\{G \vdash t_1 : tp\}$, $\{G \vdash t_2 : tp\}$, $\{G \vdash d_1 : of\ e\ t_1\}$ and $\{G \vdash d_2 : of\ e\ t_2\}$ are valid, then there must be a closed expression d_3 such that $\{G \vdash d_3 : eq\ t_1\ t_2\}$ is also valid. Such an argument can be constructed by induction on the height of the LF derivation of $G \vdash_{\Sigma} d_1 \leftarrow of\ e\ t_1$, which we analyze using Theorem 2.20 in the manner discussed earlier. There are essentially four cases to consider, corresponding to whether the head symbol of d_1 is *of_empty*, *of_app*, *of_lam*, or a nominal constant that is assigned the type $(of\ n\ t_1)$ in G where n is also a nominal constant that is bound in G . In the last case, we use the fact that the validity of $\{G \vdash d_1 : of\ e\ t_1\}$ implies that $\vdash_{\Sigma} G\ ctx$ is derivable to conclude the uniqueness of n and, hence, of the typing. The argument when d_1 is *of_empty* has an obvious form. The argument when d_1 has *of_app* or *of_lam* as its head symbol will invoke the induction hypothesis. In the case where the head symbol is *of_lam*, we will need to consider a shorter derivation of a typing judgement in which the context has been enhanced. However, we will be able to use the induction hypothesis by observing that the enhancements to the context conform to the constraints imposed by the context schema. Note that the form of d_1 also constrains the form of e in all the cases, a fact that is used implicitly in the analysis outlined.

3.4 Nominal Constants and Invariance Under Permutations

The particular choices for bound variable names in the kinds, types and terms that comprise LF expressions are considered irrelevant. This understanding is built in concretely through the notion of α -conversion that renders equal expressions that differ only in the names used for such variables. Typing derivations transform expressions with bound variables into ones where variables are ostensibly free but in fact bound implicitly in the associated contexts. The lack of importance of name choices is reflected in this case in an invariance in the validity of typing judgements under a suitable renaming of variables appearing in the judgements. In a situation where context variables are represented by nominal constants, this property can be expressed via an invariance of formula validity under permutations of nominal constants as we describe here. We begin with a definition of the notions of permutations of nominal constants and their applications to expressions.

Definition 3.7. *A permutation of the nominal constants is an arity type preserving bijection from \mathcal{N} to \mathcal{N} that differs from the identity map at only a finite number of constants. The permutation that maps n_1, \dots, n_m to n'_1, \dots, n'_m , respectively, and is the identity everywhere else is written as $\{n'_1/n_1, \dots, n'_m/n_m\}$. The support of a permutation $\pi = \{n'_1/n_1, \dots, n'_m/n_m\}$, denoted by $\text{supp}(\pi)$, is the collection of nominal constants $\{n_1, \dots, n_m\}$ or, identically, $\{n'_1, \dots, n'_m\}$. Every permutation π has an obvious inverse that is written as π^{-1} .*

Definition 3.8. *The application of a permutation π to an expression E of a variety of kinds is described below and is denoted in all cases by $\pi.E$. If E is a term, type, or kind then the application consists of replacing each nominal constant n that appears in E with $\pi(n)$. If E is a context then the application of π to E replaces each explicit binding $n : A$ in E with $\pi(n) : \pi.A$. If E is an LF judgement \mathcal{J} then the permutation is applied to each component of the judgement in the way described above. If E is a formula then the permutation is applied to its component parts. The application of π to a term variables substitution $\{\langle x_1, M_1, \alpha_1 \rangle, \dots, \langle x_n, M_n, \alpha_n \rangle\}$ yields the substitution $\{\langle x_1, \pi.M_1, \alpha_1 \rangle, \dots, \langle x_n, \pi.M_n, \alpha_n \rangle\}$. The application of π to a context variables substitution $\{G_1/\Gamma_1, \dots, G_n/\Gamma_n\}$ yields $\{\pi.G_1/\Gamma_1, \dots, \pi.G_n/\Gamma_n\}$.*

The following theorem expresses the property of interest concerning LF judgements cast in the form relevant to \mathcal{L}_{LF} .

Theorem 3.9. *Let LF judgements and derivations be recast in the form discussed earlier in this section: variables that are bound in a context are represented by nominal constants and the rules CANON_KIND_PI, CANON_FAM_PI and CANON_TERM_LAM introduce fresh nominal constants into contexts and replace variables in kinds, types and terms with these constants. In this context, let \mathcal{J} be an LF judgement which has a derivation. Then for any permutation π , $\pi.\mathcal{J}$ is derivable. Moreover, the structure of this derivation is the same as that for \mathcal{J} .*

Proof. This proof is by induction on the derivation for \mathcal{J} . Perhaps the only observation worthy of note is that the freshness of nominal constants used in CANON_KIND_PI, CANON_FAM_PI, and CANON_TERM_LAM rules is preserved under permutations of nominal constants. \square

The above observation underlies the main theorem of this section.

Theorem 3.10. *Let F be a closed formula and let π be a permutation. Then F is valid if and only if $\pi.F$ is valid.*

Proof. Noting that π^{-1} is also a permutation and that $\pi^{-1}.\pi.F$ is F , it suffices to prove the claim in only one direction. We do this by induction on the structure of F .

The desired result follows easily from Theorem 3.9 and the relationship of validity to LF derivability when F is atomic. The cases where F is \top or \perp are trivial and the ones in which F is $F_1 \supset F_2$, $F_1 \wedge F_2$ or $F_1 \vee F_2$ are easily argued with recourse to the induction hypothesis and by noting that the permutation distributes to the component formulas.

In the case where F is $\Pi \Gamma : \mathcal{C}.F'$, we first note that if $\mathcal{N}; \emptyset \vdash \mathcal{C} \rightsquigarrow_{cs} G$ has a derivation then $\mathcal{N}; \emptyset \vdash \mathcal{C} \rightsquigarrow_{cs} \pi^{-1}.G$ must also have one. From this and the validity of F it follows that $F'[\{(\pi^{-1}.G)/\Gamma\}]$ must be valid. Moreover, $F'[\{(\pi^{-1}.G)/\Gamma\}]$ has the same structural complexity as F' . Hence, by the induction hypothesis, $\pi.(F'[\{(\pi^{-1}.G)/\Gamma\}])$ is valid. Noting that this formula is the same as $(\pi.F')[\{G/\Gamma\}]$ and that $\Pi \Gamma : \mathcal{C}.\pi.F'$ is identical to $\pi.(\Pi \Gamma : \mathcal{C}.F')$, the validity of $\pi.F$ easily follows.

Suppose that F has the form $\forall x : \alpha.F'$. We observe here that if $\mathcal{N} \cup \Theta_0 \vdash_{at} M : \alpha$ has a derivation then $\mathcal{N} \cup \Theta_0 \vdash_{at} \pi^{-1}.M : \alpha$ has one too and that $\pi.(F'[\{\langle x, \pi^{-1}.M, \alpha \rangle\}])$ is the same formula as $(\pi.F')[\{\langle x, M, \alpha \rangle\}]$. Using the definition of validity, the induction hypothesis and the fact that permutation distributes to the component formula together with the above observations, we may easily conclude that $\pi.F$ is valid.

Finally, suppose that F is of the form $\exists x : \alpha.F'$. Here we note that if $\mathcal{N} \cup \Theta_0 \vdash_{at} M : \alpha$ has a derivation then $\mathcal{N} \cup \Theta_0 \vdash_{at} \pi.M : \alpha$ has one too and that $\pi.(F'[\{\langle x, M, \alpha \rangle\}])$ is the same formula as $(\pi.F')[\{\langle x, \pi.M, \alpha \rangle\}]$. Using the definition of validity and the induction hypothesis, it is now easy to conclude that $\pi.F$ must be valid. \square

4 A Proof System for the Logic

In this section we describe a proof system that provides a formal mechanism for demonstrating validity for formulas in \mathcal{L}_{LF} . This proof system is oriented around sequents that represent assumption and conclusion formulas augmented with devices that capture additional aspects of states that arise in the process of reasoning. The syntax for sequents is more liberal than is meaningful at the outset, and this is rectified by imposing wellformedness requirements on them. We associate a semantics with sequents that is consistent with their intended use. We then present a collection of proof rules that can be used to derive sequents. These rules belong to two broad categories. The first category comprises rules that embody logical aspects such as the meanings of sequents and of the logical symbols that appear in formulas. A key aspect of \mathcal{L}_{LF} is that its atomic formulas represent the notion of derivability in LF that is also open to analysis. The second category of proof rules builds in capabilities for such analysis. An important property for our proof rules is that they should require the proofs only of well-formed sequents in constructing derivations for well-formed sequents. One concern in their presentation is therefore to check that they satisfy this property. At a more substantive level, the proof rules must support a reasoning process that is both sound and effective. We focus in this paper on the issue of soundness, leaving the demonstration of effectiveness to other work, e.g. see [18]. In the proofs of soundness, we will assume the wellformedness of sequents, as guaranteed by the complementary consideration.

The first subsection below presents the sequents underlying the proof system and identifies a semantics with them. The remaining subsections develop the collection of proof rules. In Section 4.2, we present a collection of rules that encapsulate the meanings of the logical symbols and also certain structural aspects of sequents. We then turn to the rules that internalize aspects of LF derivability that are intrinsic to the understanding of the atomic formulas. Section 4.3 develops rules for analyzing atomic formulas. An important component of these rules is the interpretation of typing judgements involving atomic types via the particular LF specification that parameterizes the logic: this interpretation leads, in particular, to a case analysis rule for such atomic formulas that

$$\begin{array}{c}
\frac{}{\mathbb{N}; \Psi \vdash \mathcal{C}[\cdot] \text{ ctx-ty}} \qquad \frac{\mathbb{N}; \Psi \vdash \mathcal{C} \rightsquigarrow_{cs}^1 G \quad \mathbb{N}; \Psi \vdash \mathcal{C}[\mathcal{G}] \text{ ctx-ty}}{\mathbb{N}; \Psi \vdash \mathcal{C}[\mathcal{G}; G] \text{ ctx-ty}} \\
\\
\frac{}{\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\cdot] \rightsquigarrow_{csty} \cdot} \qquad \frac{\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[\mathcal{G}] \in \Xi \quad (\mathcal{N} \setminus \mathbb{N}_\Gamma) \subseteq \mathbb{N}}{\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\cdot] \rightsquigarrow_{csty} \Gamma} \\
\\
\frac{\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\mathcal{G}] \rightsquigarrow_{csty} G \quad \mathbb{N}; \Psi \vdash \mathcal{C} \rightsquigarrow_{cs}^1 G'}{\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\mathcal{G}] \rightsquigarrow_{csty} G, G'} \qquad \frac{\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\mathcal{G}] \rightsquigarrow_{csty} G}{\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\mathcal{G}; G'] \rightsquigarrow_{csty} G, G'}
\end{array}$$

Figure 4.1: Well-Formed Context Variable Types and their Instantiations

appear as assumptions in a sequent. Section 4.4 presents rules that enable reasoning based on induction on the heights LF derivations. In the final subsection, we introduce proof rules that encode meta-theorems concerning LF derivability that often find use in reasoning about LF specifications.

4.1 The Structure of Sequents

A sequent in our proof system is characterized by a collection of assumption formulas and a conclusion or goal formula. The formulas may contain free term and context variables that are to be interpreted as being implicitly universally quantified over the sequent and, therefore, its proof. We find it useful also to identify with the sequent a collection of nominal constants that circumscribes the ones that appear in its formulas.

The nominal constants and term variables that appear in the sequent have arity types associated with them. Context variables are also typed and their types are, in spirit, based on context schemas. However, subproofs may require a partial elaboration of a context variable and the types associated with such variables accommodates this possibility. More specifically, these types have the form $\mathcal{C}[G_1; \dots; G_n]$ where \mathcal{C} is a context schema and G_1, \dots, G_n are context expressions. Such a type is intended to represent the collection of context expressions obtained by interspersing G_1, \dots, G_n with instantiations of the context schema \mathcal{C} and possibly prefixed by a context variable of suitable type that represents a yet to be elaborated sequence of declarations. Additionally, context variables are annotated with a collection of nominal constants that express the constraint that the elaborations of these variables must not use names in these collections; the ability to express such constraints is an essential part of the mechanism for analyzing typing judgements involving abstractions as we will see later in this section.

The ideas pertaining to context variable typing are made precise through the following definition.

Definition 4.1. *A context variable type is an expression of the form $\mathcal{C}[\mathcal{G}]$ where \mathcal{C} is a context schema such that $\vdash \mathcal{C}$ ctx schema is derivable and \mathcal{G} represents a sequence of context blocks given as follows:*

$$\mathcal{G} ::= \cdot \mid \mathcal{G}; n_1 : A_1, \dots, n_k : A_k.$$

Such a type is said to be well-formed with respect to a nominal constant set $\mathbb{N} \subseteq \mathcal{N}$ and a term variables context Ψ if it is the case that the relation $\mathbb{N}; \Psi \vdash \mathcal{C}[\mathcal{G}]$ ctx-ty that is defined by the rules in Figure 4.1 holds; intuitively, if \mathcal{G} is a listing of blocks obtained by instantiating block schemas comprising \mathcal{C} . A context variables context is a collection of associations with context variables of sets of nominal constants and context variable types, each written in the form $\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[\mathcal{G}]$. Given a context variables context Ξ , we write Ξ^- for the set $\{\Gamma \mid \Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[\mathcal{G}] \in \Xi\}$, i.e., the collection of context variables assigned types by Ξ . A context expression G is said to be an instance of a

context type $\mathcal{C}[\mathcal{G}]$ relative to \mathbb{N} , Ψ and Ξ if the relation $\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\mathcal{G}] \rightsquigarrow_{\text{csty}} G$, that is also defined in Figure 4.1, holds.

The following theorem, whose proof is straightforward, shows that an instance of a well-formed context variable type is a well-formed context relative to the relevant arity context and context variable collection.

Theorem 4.2. *If $\mathbb{N}; \Psi \vdash \mathcal{C}[\mathcal{G}]$ ctx-ty and $\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\mathcal{G}] \rightsquigarrow_{\text{csty}} G$ have derivations then so does $\mathbb{N} \cup \Psi; \Xi \vdash G$ context.*

The following theorem, whose proof is also straightforward, provides us a means that we will often use for adjusting contexts that parameterize the relevant wellformedness judgements.

Theorem 4.3. *Let $\mathbb{N} \subseteq \mathbb{N}'$, $\Psi \subseteq \Psi'$, and $\Xi \subseteq \Xi'$.*

1. *If $\mathbb{N}; \Psi \vdash \mathcal{C}[\mathcal{G}]$ ctx-ty is derivable, then so is $\mathbb{N}'; \Psi' \vdash \mathcal{C}[\mathcal{G}]$ ctx-ty.*
2. *If $\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\mathcal{G}] \rightsquigarrow_{\text{csty}} G$ is derivable, then so is $\mathbb{N}'; \Psi'; \Xi' \vdash \mathcal{C}[\mathcal{G}] \rightsquigarrow_{\text{csty}} G$.*

The wellformedness judgements in Figure 4.1 are preserved under meaningful substitutions as the theorem below explicates.

Theorem 4.4. *Let θ be an arity type preserving substitution with respect to $\mathcal{N} \cup \Theta_0 \cup \Psi$ and let $\mathbb{N}; \text{ctx}(\theta) \uplus \Psi \vdash \mathcal{C}[\mathcal{G}]$ ctx-ty have a derivation. Then*

1. *there must be a derivation for $\mathbb{N} \cup \text{supp}(\theta); \Psi \vdash \mathcal{C}[\mathcal{G}[\theta]]$ ctx-ty, and*
2. *if $\mathbb{N}; \text{ctx}(\theta) \uplus \Psi; \Xi \vdash \mathcal{C}[\mathcal{G}] \rightsquigarrow_{\text{csty}} G$ also has a derivation, there must be a derivation for $\mathbb{N} \cup \text{supp}(\theta); \Psi; \Xi \vdash \mathcal{C}[\mathcal{G}[\theta]] \rightsquigarrow_{\text{csty}} G[\theta]$.*

Proof. Since θ is arity type preserving with respect to $\mathcal{N} \cup \Theta_0 \cup \Psi$, using Theorem 2.8 we see that if there is a derivation for $\mathbb{N} \cup (\text{ctx}(\theta) \uplus \Psi) \cup \Theta_0 \vdash_{\text{at}} t : \alpha$ then $t[\theta]$ must be well-defined and there must be a derivation for $(\mathbb{N} \cup \text{supp}(\theta)) \cup \Psi \cup \Theta_0 \vdash_{\text{at}} t[\theta] : \alpha$. An induction on the derivation of $\mathbb{N}; \text{ctx}(\theta) \uplus \Psi \vdash \mathcal{C}[\mathcal{G}]$ ctx-ty using these observations allows us to confirm the first part of the theorem. Now suppose that there is a derivation for $\mathbb{N}; \text{ctx}(\theta) \uplus \Psi \vdash \mathcal{C} \rightsquigarrow_{\text{cs}}^1 G'$. By an induction on this derivation using the facts observed earlier, it can be concluded that there must be a derivation for $\mathbb{N} \cup \text{supp}(\theta); \Psi \vdash \mathcal{C} \rightsquigarrow_{\text{cs}}^1 G'[\theta]$. The second part of the theorem follows by another obvious induction from this. \square

We now define the notion of sequents that underlies our proof system.

Definition 4.5. *A sequent, written as $\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$, is a judgement that relates a finite subset \mathbb{N} of \mathcal{N} , a term variables context Ψ , a context variables context Ξ , a finite set Ω of assumption formulas and a conclusion or goal formula F . The sequent is well-formed if (a) for each $\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[\mathcal{G}]$ in Ξ it is the case that $\mathbb{N}_\Gamma \subseteq \mathbb{N}$ and that $\mathbb{N} \setminus \mathbb{N}_\Gamma; \Psi \vdash \mathcal{C}[\mathcal{G}]$ ctx-ty is derivable and (b) for each formula F' in $\{F\} \cup \Omega$ it is the case that $\mathbb{N} \cup \Psi \cup \Theta_0; \Xi \vdash F'$ fmla is derivable. Given a well-formed sequent $\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$, we will refer to \mathbb{N} as its support set, to Ψ as its term variables context and Ξ as its context variables context, and we will denote the collection of variables assigned types by Ψ and Ξ by $\text{dom}(\Psi)$ and $\text{dom}(\Xi)$, respectively. We will use a comma to denote set union in representing sequents, writing Ω, F_1, \dots, F_n to denote the set $\Omega \cup \{F_1, \dots, F_n\}$.*

We will need to consider substitutions for term variables in sequents. We will require legitimate substitutions to not use the nominal constants in the support set of the sequent; this restriction will be part of a mechanism for controlling dependencies in context declarations. We will further require substitutions to satisfy arity typing constraints for their applications to be well-defined. These considerations are formalized below in the notion of substitution compatibility.

Definition 4.6. *A pair $\langle \theta, \Psi' \rangle$ comprising a substitution and a term variables context is said to be substitution compatible with a well-formed sequent $\mathcal{S} = \mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$ if*

1. θ is arity type preserving with respect to the context $\mathcal{N} \cup \Theta_0 \cup \Psi'$,
2. $\text{supp}(\theta) \cap \mathbb{N} = \emptyset$, and
3. for any variable x , if $x : \alpha \in \Psi$ and $x : \alpha' \in \text{ctx}(\theta) \uplus \Psi'$, then $\alpha = \alpha'$.

The application of a substitution may introduce new nominal constants into a sequent. When this happens, substitutions for the term variables in the resulting sequent must be permitted to contain these constants. We use the technique of raising to realize this requirement [7]. The following definition is useful in formalizing this idea.

Definition 4.7. *Let Ψ be the term variables context $\{x_1 : \alpha_1, \dots, x_m : \alpha_m\}$, let n_1, \dots, n_k be a listing of the elements of a finite collection of the nominal constants \mathbb{N} , and let β_1, \dots, β_k be the arity types associated with these constants. Then a version of Ψ raised over \mathbb{N} is a set $\{y_1 : \gamma_1, \dots, y_m : \gamma_m\}$ where, for $1 \leq i \leq m$, y_i is a distinct variable that is also different from the variables in $\{x_1, \dots, x_m\}$ and γ_i is $\beta_1 \rightarrow \dots \rightarrow \beta_k \rightarrow \alpha_i$. Further, the raising substitution associated with this version is the set $\{(x_i, (y_i \ n_1 \ \dots \ n_k), \alpha_i) \mid 1 \leq i \leq m\}$.*

The basis for using raising in the manner described is the content of the following theorem. We say here and elsewhere that an arity context Θ is compatible with \mathcal{N} if the types that Θ assigns to nominal constants are identical to their assignments in \mathcal{N} .

Theorem 4.8. *Let θ be a substitution that is arity type preserving with respect to an arity context Θ that is compatible with \mathcal{N} . Further, let Ψ be a version of $\text{ctx}(\theta)$ raised over some listing of a collection \mathbb{N} of nominal constants and let θ_r be the associated raising substitution. Then there is a substitution θ' with $\text{supp}(\theta') = \text{supp}(\theta) \setminus \mathbb{N}$ and $\text{ctx}(\theta') = \Psi$ that is arity type preserving with respect to Θ and such that for any E for which $\text{ctx}(\theta) \uplus \Theta \vdash_{ak} E$ type or, for some arity type α , $\text{ctx}(\theta) \uplus \Theta \vdash_{at} E : \alpha$ has a derivation, it is the case that $E[\![\theta_r]\!] [\![\theta']\!] = E[\![\theta]\!]$.*

Proof. Each of the substitutions involved in the expression $E[\![\theta_r]\!] [\![\theta']\!] = E[\![\theta]\!]$ will have a result under the conditions described, thereby justifying the use of the notation introduced after Theorem 2.8. Now, let n_1, \dots, n_k be the listing of the constants in \mathbb{N} in the raising substitution, let $\alpha_1, \dots, \alpha_k$ be the respective types of these constants, let $\langle x, t, \alpha \rangle$ be a tuple in θ and let $\langle x, (y \ n_1 \ \dots \ n_k), \alpha \rangle$ be the tuple corresponding to x in θ_r . Let x_1, \dots, x_k be a listing of distinct variables that do not appear in t and let t' be the result of replacing n_i by x_i in t , for $1 \leq i \leq k$. We construct θ' by including in it the substitution $\langle y, \lambda x_1. \dots \lambda x_k. t', \alpha_1 \rightarrow \dots \rightarrow \alpha_k \rightarrow \alpha \rangle$ for each case of the kind considered. It is easy to see that $\text{supp}(\theta') = \text{supp}(\theta) \setminus \mathbb{N}$ and that θ_r and θ' are arity type compatible with respect to $\Theta \cup \mathcal{N}$. The remaining part of the theorem follows from noting that $\theta = \theta' \circ \theta_r$ and using Theorem 2.10. \square

The following definition formalizes the application of a term substitution to a well-formed sequent when the conditions of substitution compatibility are met. We assume here and elsewhere

that the application of a term substitution to a set of formulas distributes to each member of the set, its application to a context variables context distributes to each context variable type in the context and its application to a context variable type $\mathcal{C}[\mathcal{G}]$ distributes to each context block in \mathcal{G} .

Definition 4.9. Let \mathcal{S} be the well-formed sequent $\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$ and let $\langle \theta, \Psi' \rangle$ be substitution compatible with \mathcal{S} . Further, let Ψ'' be a version of $(\Psi \setminus \text{ctx}(\theta)) \cup \Psi'$ raised over $\text{supp}(\theta)$ and let θ_r be the corresponding raising substitution. Then the application of θ to \mathcal{S} relative to Ψ' , denoted by $\mathcal{S}[\![\theta]\!]_{\Psi'}$, is the sequent $\mathbb{N} \cup \text{supp}(\theta); \Psi''; \Xi[\![\theta]\!][\![\theta_r]\!]; \Omega[\![\theta]\!][\![\theta_r]\!] \longrightarrow F[\![\theta]\!][\![\theta_r]\!]$.

The definition and notation above are obviously ambiguous since they depend on the particular choices of Ψ'' and θ_r . We shall mean $\mathcal{S}[\![\theta]\!]_{\Psi'}$ to denote any one of the sequents so determined, referring to Ψ'' and θ_r as the raised context and the raising substitution associated with the application of the substitution where disambiguation is needed. Note also that the definition assumes that the application of the substitutions θ and θ_r to the relevant context variable types and formulas is well-defined. We show this to be the case in the theorem below.

Theorem 4.10. Let $\langle \theta, \Psi' \rangle$ be substitution compatible with the well-formed sequent \mathcal{S} . Then $\mathcal{S}[\![\theta]\!]_{\Psi}$ is well-defined and it yields a well-formed sequent.

Proof. As in Definition 4.9, let $S = \mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$, let Ψ'' be the raised context and let θ_r be the corresponding raising substitution. To prove the theorem, it suffices to show the following:

1. for each $\Gamma \uparrow \mathbb{N}_{\Gamma} : \mathcal{C}[\mathcal{G}]$ in Ξ it is the case that $\mathbb{N}_{\Gamma} \subseteq \mathbb{N} \cup \text{supp}(\theta)$ and there is a derivation for the judgement $(\mathbb{N} \cup \text{supp}(\theta)) \setminus \mathbb{N}_{\Gamma}; \Psi'' \vdash \mathcal{C}[\mathcal{G}[\![\theta]\!][\![\theta_r]\!]] \text{ ctx-ty}$, and
2. for each formula F' in $\{F\} \cup \Omega$ it is the case that $(\mathbb{N} \cup \text{supp}(\theta)) \cup \Psi'' \cup \Theta_0; \Xi^- \vdash F'[\![\theta]\!][\![\theta_r]\!] \text{ fmla}$ is derivable.

Implicit here is the requirement that the relevant substitutions are well-defined.

We first show (1). Since \mathcal{S} is well-formed, it must be the case that $\mathbb{N}_{\Gamma} \subseteq \mathbb{N}$ so the first part of this requirement obviously holds. The wellformedness of \mathcal{S} implies that $\mathbb{N} \setminus \mathbb{N}_{\Gamma}; \Psi \vdash \mathcal{C}[\mathcal{G}] \text{ ctx-ty}$ has a derivation. The substitution compatibility of $\langle \theta, \Psi' \rangle$ with \mathcal{S} implies that $\text{ctx}(\theta)$ and $\Psi' \setminus \text{ctx}(\theta)$ agree with Ψ on the type assignments to the variables that are common to them, from which it follows that $\Psi \subseteq \text{ctx}(\theta) \uplus ((\Psi \setminus \text{ctx}(\theta)) \cup \Psi')$. Theorem 4.3 now allows us to conclude that there must be a derivation for $(\mathbb{N} \setminus \mathbb{N}_{\Gamma}; \text{ctx}(\theta) \uplus ((\Psi \setminus \text{ctx}(\theta)) \cup \Psi')) \vdash \mathcal{C}[\mathcal{G}] \text{ ctx-ty}$. From the substitution compatibility of $\langle \theta, \Psi' \rangle$ with \mathcal{S} it also follows that θ must be type preserving with respect to $(\mathbb{N} \cup \text{supp}(\theta)) \cup \Theta_0 \cup ((\Psi \setminus \text{ctx}(\theta)) \cup \Psi')$ and therefore with respect to $\mathcal{N} \cup \Theta_0 \cup ((\Psi \setminus \text{ctx}(\theta)) \cup \Psi')$ and, because $\text{supp}(\theta)$ is disjoint from \mathbb{N} , that $(\mathbb{N} \setminus \mathbb{N}_{\Gamma}) \cup \text{supp}(\theta) = (\mathbb{N} \cup \text{supp}(\theta)) \setminus \mathbb{N}_{\Gamma}$. Using Theorem 4.4 we now determine that $(\mathbb{N} \cup \text{supp}(\theta)) \setminus \mathbb{N}_{\Gamma}; (\Psi \setminus \text{ctx}(\theta)) \cup \Psi' \vdash \mathcal{C}[\mathcal{G}[\![\theta]\!]] \text{ ctx-ty}$ has a derivation. From Definitions 4.7 and 4.9, it follows that θ_r is type preserving with respect to $\mathcal{N} \cup \Theta_0 \cup \Psi''$, that $\text{ctx}(\theta_r) = (\Psi \setminus \text{ctx}(\theta)) \cup \Psi'$, and that $\text{supp}(\theta_r) \subseteq (\mathbb{N} \cup \text{supp}(\theta)) \setminus \mathbb{N}_{\Gamma}$. Thus, using Theorem 4.4 again, we determine that $(\mathbb{N} \cup \text{supp}(\theta)) \setminus \mathbb{N}_{\Gamma}; \Psi'' \vdash \mathcal{C}[\mathcal{G}[\![\theta]\!][\![\theta_r]\!]] \text{ ctx-ty}$ has a derivation.

We now consider requirement (2). Let F' be a formula in $\{F\} \cup \Omega$. As we have seen already, θ is type preserving with respect to the arity context $(\mathbb{N} \cup \text{supp}(\theta)) \cup \Theta_0 \cup ((\Psi \setminus \text{ctx}(\theta)) \cup \Psi')$. Since \mathcal{S} is well-formed, there must be a derivation for $\mathbb{N} \cup \Psi \cup \Theta_0; \Xi^- \vdash F' \text{ fmla}$. Since $\text{ctx}(\theta)$ and Ψ agree on the type assignments to common variables, we can conclude, using Theorem 3.3, that there must be a derivation for $\text{ctx}(\theta) \uplus ((\mathbb{N} \cup \text{supp}(\theta)) \cup ((\Psi \setminus \text{ctx}(\theta)) \cup \Psi') \cup \Theta_0); \Xi^- \vdash F' \text{ fmla}$. From Theorem 3.5, it then follows that $(\mathbb{N} \cup \text{supp}(\theta)) \cup (\Psi \setminus \text{ctx}(\theta)) \cup \Psi' \cup \Theta_0; \Xi^- \vdash F'[\![\theta]\!] \text{ fmla}$ has a derivation. This last judgement may be rewritten as $\text{ctx}(\theta_r) \uplus ((\mathbb{N} \cup \text{supp}(\theta)) \cup \Theta_0); \Xi^- \vdash F'[\![\theta]\!] \text{ fmla}$ from which, using Theorem 3.3, it follows that $\text{ctx}(\theta_r) \uplus ((\mathbb{N} \cup \text{supp}(\theta)) \cup \Psi'' \cup \Theta_0); \Xi^- \vdash F'[\![\theta]\!] \text{ fmla}$

has a derivation. Noting that θ_r is type preserving with respect to $\text{supp}(\theta) \cup \Psi''$ and hence with respect to $((\mathbb{N} \cup \text{supp}(\theta)) \cup \Psi'' \cup \Theta_0)$, we may use Theorem 3.5 to conclude that there must be a derivation for $(\mathbb{N} \cup \text{supp}(\theta)) \cup \Psi'' \cup \Theta_0; \Xi^- \vdash F'[\![\theta]\!]_{[\![\theta_r]\!]}$ *fmla*. \square

We will also need to consider the application of context variables substitutions to sequents. We require such substitutions to respect the types associated with the variables. In contrast to term variables substitutions, context variables substitutions are not permitted to introduce new term variables into the sequent and they may use nominal constants that are already present in the sequent. We formalize these ideas in the definition of appropriateness for such substitutions.

Definition 4.11. *Let Ξ be a context variables context and let σ be the context variables substitution $\{G_1/\Gamma_1, \dots, G_n/\Gamma_n\}$. We write Ξ_σ to denote the result of trimming Ξ to exclude associations for the variables in $\text{dom}(\sigma)$. We say that σ is appropriate for a context variables context Ξ with respect to a term variables context Ψ and a collection of nominal constants \mathbb{N} if, for $1 \leq i \leq n$, it is the case that $\Gamma_i \uparrow \mathbb{N}_i : \mathcal{C}_i[\mathcal{G}_i] \in \Xi$ and, further, $\mathbb{N}; \Psi \vdash \mathcal{C}_i[\mathcal{G}_i]$ *ctx-ty* and $\text{supp}(\sigma) \setminus \mathbb{N}_i; \Psi; \Xi_\sigma \vdash \mathcal{C}_i[\mathcal{G}_i] \rightsquigarrow_{\text{csty}} G_i$ have derivations. The substitution σ is additionally said to cover Ξ if $\Xi_\sigma = \emptyset$. We say that σ is appropriate for a well-formed sequent $\mathcal{S} = \mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$ if it is appropriate for Ξ with respect to Ψ and \mathbb{N} and that it covers \mathcal{S} if it covers Ξ .*

Context types are unaffected by context variables substitutions. Context expressions may be impacted by such substitutions but, for the right kind of substitutions, they continue to be instances of relevant context types. This is made precise in the following theorem.

Theorem 4.12. *Let σ be a context variables substitution that is appropriate for Ξ with respect to Ψ and \mathbb{N} and let $\text{supp}(\sigma) \subseteq \mathbb{N}$. If $\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\mathcal{G}] \rightsquigarrow_{\text{csty}} G$ has a derivation, then $\mathbb{N}; \Psi; \Xi_\sigma \vdash \mathcal{C}[\mathcal{G}] \rightsquigarrow_{\text{csty}} G[\sigma]$ also has a derivation.*

Proof. By induction on the derivation of $\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\mathcal{G}] \rightsquigarrow_{\text{csty}} G$. The only case that perhaps needs explicit consideration is that where G is a context variable Γ for which σ substitutes the context expression G' . In this case \mathcal{G} must be empty and there must be an assignment of the form $\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[\mathcal{G}']$ in Ξ for some \mathbb{N}_Γ and \mathcal{G}' . Further, by the appropriateness of σ , there must be a derivations for $\mathbb{N}; \Psi \vdash \mathcal{C}[\mathcal{G}']$ *ctx-ty* and $\text{supp}(\sigma) \setminus \mathbb{N}_\Gamma; \Psi; \Xi_\sigma \vdash \mathcal{C}[\mathcal{G}'] \rightsquigarrow_{\text{csty}} G_i$. Using Theorem 4.3 and the assumption that $\text{supp}(\sigma) \subseteq \mathbb{N}$, it follows that $\mathbb{N}; \Psi; \Xi_\sigma \vdash \mathcal{C}[\mathcal{G}'] \rightsquigarrow_{\text{csty}} G_i$ has a derivation. An easy induction on the first derivation suffices to show that if $\mathbb{N}; \Psi; \Xi_\sigma \vdash \mathcal{C}[\mathcal{G}'] \rightsquigarrow_{\text{csty}} G_i$ and $\mathbb{N}; \Psi \vdash \mathcal{C}[\mathcal{G}']$ *ctx-ty* have derivations then there must be one for $\mathbb{N}; \Psi; \Xi_\sigma \vdash \mathcal{C}[\cdot] \rightsquigarrow_{\text{csty}} G_i$. \square

We formalize the application of a context variables substitution to a well-formed sequent when the conditions of appropriateness are met in the definition below. Note that context substitutions can also introduce new nominal constants, and we use the technique of raising once again to permit these constants to be used in substitutions for term variables in the resulting sequent. We assume here and elsewhere that the application of a context variables substitution to a set of formulas distributes to each member of the set.

Definition 4.13. *Let \mathcal{S} be a well-formed sequent $\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$ and σ be appropriate for \mathcal{S} . Further let Ψ' be a version of Ψ raised over $\text{supp}(\sigma)$ and let θ_r be the corresponding raising substitution. Then the application of σ to \mathcal{S} , denoted by $\mathcal{S}[\sigma]$, is the sequent*

$$\mathbb{N} \cup \text{supp}(\sigma); \Psi'; \Xi_\sigma[\![\theta_r]\!]; \Omega[\sigma][\![\theta_r]\!] \longrightarrow F[\sigma][\![\theta_r]\!].$$

Once again, we must justify the use in the above definition of the notation introduced after Theorem 2.8. This is part of the content of the following theorem.

Theorem 4.14. *Let \mathcal{S} be a well-formed sequent and let σ be a context variables substitution that is appropriate for \mathcal{S} . Then $\mathcal{S}[\sigma]$ is a well-defined and it yields a well-formed sequent.*

Proof. Let \mathcal{S} be $\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$ and let $\mathcal{S}[\sigma]$ be $\mathbb{N} \cup \text{supp}(\sigma); \Psi'; \Xi_\sigma[[\theta_r]]; \Omega[\sigma][[\theta_r]] \longrightarrow F[\sigma][[\theta_r]]$, where θ_r and Ψ' are the raising substitution and the raised term variables context as per Definition 4.13. Using an argument similar to that in the proof of Theorem 4.10, we can show that $(\mathbb{N} \cup \text{supp}(\sigma)) \setminus \mathbb{N}_\Gamma; \Psi' \vdash (\mathcal{C}[\mathcal{G}]][[\theta_r]]$ ctx-ty must have a derivation for each $\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[\mathcal{G}]$ in Ξ_σ . From the wellformedness of \mathcal{S} and the appropriateness of σ for \mathcal{S} , it is easy to see that $(\mathbb{N} \cup \text{supp}(\sigma)) \cup \Psi \cup \Theta_0; \Xi_\sigma^- \vdash F'[\sigma]$ fmla must have a derivation for each $F' \in \Omega \cup \{F\}$. Noting that the variables assigned types by Ψ' are distinct from those assigned types by Ψ and that $\text{ctx}(\theta_r) = \Psi$, we may use Theorems 3.3 and 3.5 to conclude that $F'[\sigma][[\theta_r]]$ is well-defined and that $(\mathbb{N} \cup \text{supp}(\sigma)) \cup \Psi' \cup \Theta_0; \Xi_\sigma^- \vdash F'[\sigma][[\theta_r]]$ fmla has a derivation. The theorem follows easily from these observations. \square

Term and context substitutions will often be used in tandem to produce instances of sequents. In this context, we shall say that θ and σ are proper for the sequent \mathcal{S} if there is a term variables context Ψ such that $\langle \theta, \Psi \rangle$ is substitution compatible with \mathcal{S} and σ is appropriate for $\mathcal{S}[[\theta]]_\Psi$.

We now define validity for sequents. For a sequent with empty term variables and context variables contexts, we base the definition on the validity of closed formulas. For a sequent that has formulas with free term and context variables, this is done by considering all their relevant substitution instances.

Definition 4.15. *A well-formed sequent of the form $\mathbb{N}; \emptyset; \emptyset; \Omega \longrightarrow F$, referred to as a closed sequent, is valid if either some formula in Ω is not valid or F is valid. A well-formed sequent \mathcal{S} of the form $\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$ is valid if for every term substitution θ that is type preserving with respect to $\mathcal{N} \cup \Theta_0$ and such that $\Psi = \text{ctx}(\theta)$ and $\langle \theta, \emptyset \rangle$ is substitution compatible with \mathcal{S} , and for every context substitution σ that is appropriate for and covers $\mathcal{S}[[\theta]]_\emptyset$, it is the case that $\mathcal{S}[[\theta]]_\emptyset[\sigma]$ is valid. Note that each such $\mathcal{S}[[\theta]]_\emptyset[\sigma]$ will be a well-formed and closed sequent in these circumstances and we shall refer to it as the closed instance of \mathcal{S} identified by θ and σ .*

The following theorem, whose proof is obvious, provides the basis for using our proof system for determining the validity of formulas.

Theorem 4.16. *Let F be a formula such that $\mathcal{N} \cup \Theta_0; \emptyset \vdash F$ fmla is derivable and let \mathbb{N} be the set of nominal constants that appear in F . Then the sequent $\mathbb{N}; \emptyset; \emptyset; \emptyset \longrightarrow F$ is well-formed. Moreover, F is valid if and only if $\mathbb{N}; \emptyset; \emptyset; \emptyset \longrightarrow F$ is.*

We will need in later discussions to consider the application of permutations of the nominal constants to sequents. We make this notion precise below.

Definition 4.17. *We extend the application of a permutation π of nominal constants to a variety of expressions E associated with sequents, denoting the result as before by $\pi.E$. The application of π to a set \mathbb{N} of nominal constants yields the set $\{n' \mid n \in \mathbb{N} \text{ and } \pi(n) = n'\}$. The application of π to a context variable type distributes to the constituent block instances, its application to an association of the form $\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[\mathcal{G}]$ with context variables distributes to \mathbb{N}_Γ and $\mathcal{C}[\mathcal{G}]$, and its application to a collection of formulas or a context variables context distributes to the members of the collection. Finally, the application of π to a sequent $\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$ yields the sequent $\pi.\mathbb{N}; \Psi; \pi.\Xi'; \pi.\Omega' \longrightarrow \pi.F'$.*

The following theorem notes that wellformedness of sequents is preserved under these permutations and that permutations commute in a suitable sense with term and context variables substitutions. The proof of the theorem is routine even if tedious; we omit the details.

$$\begin{array}{c}
\frac{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F_1}{\mathbb{N}; \Psi; \Xi; \Omega, F_2 \longrightarrow F_1} \text{ wk} \qquad \frac{\mathbb{N}; \Psi; \Xi; \Omega, F_2, F_2 \longrightarrow F_1}{\mathbb{N}; \Psi; \Xi; \Omega, F_2 \longrightarrow F_1} \text{ cont} \\
\\
\frac{\begin{array}{c} \Xi = \{\Gamma_i \uparrow (\mathbb{N}_i \setminus \mathbb{N}') : \mathcal{C}_i[\mathcal{G}_i] \mid \Gamma_i \uparrow \mathbb{N}_i : \mathcal{C}_i[\mathcal{G}_i] \in \bar{\Xi}\} \\ \{(\mathbb{N}, \mathbb{N}') \setminus \mathbb{N}_i; (\Psi, \Psi') \vdash \mathcal{C}_i[\mathcal{G}_i] \text{ ctx-ty} \mid \Gamma_i \uparrow \mathbb{N}_i : \mathcal{C}_i[\mathcal{G}_i] \in \Xi'\} \\ \mathbb{N}, \mathbb{N}'; \Psi, \Psi'; \bar{\Xi}, \Xi'; \Omega \longrightarrow F \end{array}}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F} \text{ ctx-str} \\
\\
\frac{\begin{array}{c} \Xi = \{\Gamma_i \uparrow (\mathbb{N}_i \setminus \mathbb{N}') : \mathcal{C}_i[\mathcal{G}_i] \mid \Gamma_i \uparrow \mathbb{N}_i : \mathcal{C}_i[\mathcal{G}_i] \in \bar{\Xi}\} \\ \{(\mathbb{N} \setminus \mathbb{N}_i); \Psi \vdash \mathcal{C}_i[\mathcal{G}_i] \text{ ctx-ty} \mid \Gamma_i \uparrow \mathbb{N}_i : \mathcal{C}_i[\mathcal{G}_i] \in \Xi\} \\ \{\mathbb{N} \cup \Theta_0 \cup \Psi; \{\Gamma \mid \Gamma \uparrow \mathbb{N} : \mathcal{C}[\mathcal{G}] \in \Xi\} \vdash F' \text{ fmla} \mid F' \in \Omega \cup \{F\}\} \\ \mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F \end{array}}{\mathbb{N}, \mathbb{N}'; \Psi, \Psi'; \bar{\Xi}, \Xi'; \Omega \longrightarrow F} \text{ ctx-wk}
\end{array}$$

Figure 4.2: The Structural Rules

Theorem 4.18. *Let π be a permutation of the nominal constants. If \mathcal{S} is a well-formed sequent then so is $\pi.\mathcal{S}$. Further, if $\langle \theta, \Psi \rangle$ is substitution compatible with $\pi.\mathcal{S}$ then $\langle \pi^{-1}.\theta, \Psi \rangle$ is substitution compatible with \mathcal{S} and $(\pi.\mathcal{S})[\theta]_{\Psi} = \pi.(\mathcal{S}[\pi^{-1}.\theta]_{\Psi})$. Finally, if σ is a context variables substitution that is appropriate for $\pi.\mathcal{S}$ then $\pi^{-1}.\sigma$ is appropriate for \mathcal{S} and $(\pi.\mathcal{S})[\sigma] = \pi.(\mathcal{S}[\pi^{-1}.\sigma])$.*

Theorem 4.19 provides a counterpart to Theorem 3.10 at the level of sequents. Its proof, whose details we again omit, uses the earlier result and the observation about substitutions contained in Theorem 4.18.

Theorem 4.19. *If π is a permutation of the nominal constants and \mathcal{S} is a well-formed closed sequent that is valid, then $\pi.\mathcal{S}$ is also a closed, valid sequent.*

4.2 The Core Proof Rules

The base for our proof system is provided by a collection of rules that embody the interpretation of the logical symbols and also certain aspects of the meanings of sequents. We present these rules in this subsection: we first identify some structural rules, we then consider axioms and the cut rule, and we finally introduce rules for the logical symbols.

4.2.1 Structural Rules

This subcollection of rules is presented in Figure 4.2. These rules can be subcategorized into those that allow for weakening and contracting the assumption set in a sequent and those that permit the weakening and strengthening of the support set, the term variables context, and the context variables context. Rules of the second subcategory encode the fact that vacuous (well-formed) extensions to the bindings manifest in a sequent will not impact its validity. The strengthening and weakening rules for contexts include premises that force modifications to context variable types and the satisfaction of typing judgements that are necessary to ensure the well-formedness of the sequents in any application of the rule.

The following theorem shows that these rules require the proof of only well-formed sequents in constructing a proof of a well-formed sequent.

Theorem 4.20. *The following property holds for each rule in Figure 4.2: if the conclusion sequent is well-formed, the premises expressing typing conditions have derivations and the conditions expressed by the other, non-sequent premises are satisfied, then all the sequent premises must be well-formed.*

Proof. The claim is obvious for the *wk*, *cont*, and *ctx-wk* rules. This leaves only the *ctx-str* rule. The wellformedness requirements as they pertain to the formulas in $\Omega \cup \{F\}$ in the premise sequent in this rule follow easily from Theorem 3.3. The second premise ensures that the context types in Ξ' in the rule meet the necessary requirements. For the remaining context types, we use the relationship between $\bar{\Xi}$ and Ξ and Theorem 4.3 to derive the needed property from the wellformedness of the conclusion sequent. \square

The following lemma will be useful in showing the soundness of the structural rules.

Lemma 4.21. *Let $\mathcal{S} = \mathbb{N}; \Psi; \bar{\Xi}; \Omega \longrightarrow F$ and $\mathcal{S}' = \mathbb{N}'; \Psi'; \Xi'; \Omega \longrightarrow F$ be well-formed sequents such that $\mathbb{N}' \subseteq \mathbb{N}$, $\Psi' \subseteq \Psi$, and there is some subset $\bar{\Xi}$ for the context variables context Ξ where*

$$\bar{\Xi}' = \{ \Gamma_i \uparrow (\mathbb{N}_i \setminus (\mathbb{N} \setminus \mathbb{N}')) : \mathcal{C}_i[\mathcal{G}_i] \mid \Gamma_i \uparrow \mathbb{N}_i : \mathcal{C}_i[\mathcal{G}_i] \in \bar{\Xi} \}.$$

Then \mathcal{S} is valid if and only if \mathcal{S}' is valid.

Proof. We will make use in the proof of the easily confirmed fact that a closed and well-formed sequent that differs from another such sequent only in its support set is valid exactly when the other sequent is valid. We note now that if θ and σ are proper for \mathcal{S} and σ' is a restriction of σ to $\text{dom}(\bar{\Xi})$, then θ and σ' must be proper for \mathcal{S}' . From this it follows easily that corresponding to each closed instance of \mathcal{S} there must be a closed instance of \mathcal{S}' that differs at most in its support set. Thus, if \mathcal{S}' is valid, \mathcal{S} must be too.

In the “only if” direction, we note first that a term substitution θ and a context substitution σ that identify a closed instance of \mathcal{S}' may not be proper for \mathcal{S} because θ and σ use nominal constants from $\mathbb{N} \setminus \mathbb{N}'$ in the terms they substitute for variables. To rectify this problem, we use a permutation π that swaps the nominal constants in $\mathbb{N} \setminus \mathbb{N}'$ with ones that do not appear in \mathbb{N} , θ or σ , leaving the other nominal constants unaffected. It is easily seen that $\pi.\theta$ and $\pi.\sigma$ identify a closed instance of \mathcal{S}' and are proper with respect to \mathcal{S} . These substitutions can now be extended to substitutions θ' and σ' that substitute closed expressions of the appropriate kinds for the variables in $\Psi \setminus \Psi'$ and $\bar{\Xi} \setminus \bar{\Xi}'$, respectively, and that thereby identify a closed instance of \mathcal{S} . This closed instance is valid by assumption. Further, it differs from $\mathcal{S}'[[\pi.\theta]]_\emptyset[[\pi.\sigma]]$ only in its support set. Thus, the latter sequent must also be valid. Noting that $\mathcal{S}' = \pi.\mathcal{S}'$ and using Theorem 4.18, we see that $\mathcal{S}'[[\pi.\theta]]_\emptyset[[\pi.\sigma]]$ is the same sequent as $\pi.\mathcal{S}'[[\theta]]_\emptyset[[\sigma]]$. Using Theorem 4.19, we now conclude that $\mathcal{S}'[[\theta]]_\emptyset[[\sigma]]$ is valid. Since this argument is independent of the choice of θ and σ , it follows that \mathcal{S}' must be valid if \mathcal{S} is. \square

The soundness of the structural rules is the content of the following theorem.

Theorem 4.22. *The following property holds for every instance of each of the rules in Figure 4.2: if the premises expressing typing judgements are derivable, the conditions described in the other non-sequent premises are satisfied and the premise sequent is valid, then the conclusion sequent must also be valid.*

Proof. The argument is obvious for the *wk* and *cont* rules. For the *ctx-str* and *ctx-wk* rules, we make use of Lemma 4.21. \square

$$\frac{F' \in \Omega \quad \pi \text{ is a permutation of nominal constants such that } \text{supp}(\pi) \subseteq \mathbb{N} \quad \Xi \vdash F' \equiv_{\pi} F}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F} \text{ id}$$

$$\frac{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F_2 \quad \mathbb{N}; \Psi; \Xi; \Omega, F_2 \longrightarrow F_1 \quad \mathbb{N} \cup \Theta_0 \cup \Psi; \text{dom}(\Xi) \vdash F_2 \text{ fmla}}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F_1} \text{ cut}$$

Figure 4.3: The Axiom and the Cut Rule

4.2.2 The Axiom and the Cut Rule

The *cut* rule codifies the notion of lemmas: if we can show the validity of a formula relative to a given assumption set, then this formula can be included in the assumptions to simplify the reasoning process. The *id* rule recognizes the validity of a sequent in which the conclusion formula appears in the assumption set. In its simplest form, the *id* rule would require the conclusion formula to be included as is in the assumption set. It is possible, and also pragmatically useful, to generalize this form to allow also for a permutation of nominal constants in the formulas in the process of matching. However, this has to be done with care to ensure that identity under the considered permutations continues to hold even after later instantiations of term and context variables appearing in the formulas. The following definition presents a notion of equivalence for formulas under permutations that encodes this care.

Definition 4.23 (Formula Equivalence). *The equivalence of two context expressions G_1 and G_2 with respect to a context variables context Ξ and a permutation π , written $\Xi \vdash G_2 \equiv_{\pi} G_1$, is a relation defined by the following three clauses:*

1. $\Xi \vdash \cdot \equiv_{\pi} \cdot$ holds for any Ξ and π .
2. $\Xi \vdash \Gamma \equiv_{\pi} \Gamma$ holds if for some $\Gamma' \uparrow \mathbb{N} : \mathcal{C}[\mathcal{G}] \in \Xi$ it is the case that $\Gamma = \Gamma'$ and $\text{supp}(\pi) \subseteq \mathbb{N}$.
3. If $G_1 = (G'_1, n_1 : A_1)$ and $G_2 = (G'_2, n_2 : A_2)$ then $\Xi \vdash G_2 \equiv_{\pi} G_1$ holds if $\pi.n_2$ is identical to n_1 , $\pi.A_2$ is the same type as A_1 , and $\Xi \vdash G'_2 \equiv_{\pi} G'_1$ holds.

Two atomic formulas $\{G' \vdash M' : A'\}$ and $\{G \vdash M : A\}$ are considered equivalent with respect to Ξ and π if $\Xi \vdash G' \equiv_{\pi} G$ holds, $\pi.M'$ and M are the same terms, and $\pi.A'$ and A are the same types. Two arbitrary formulas F' and F are considered equivalent with respect to Ξ and π if their component parts are so equivalent, allowing, of course, for a renaming of variables bound by quantifiers. We denote this equivalence by the expression $\Xi \vdash F' \equiv_{\pi} F$.

The *id* and the *cut* rules are presented in Figure 4.3. The *id* rule limits the permutations to be considered to ones that rename only nominal constants appearing in the support set of the sequent. The *cut* rule includes a premise that ensures the wellformedness of the cut formula.

Theorem 4.24. *The following property holds of the *id* and the *cut* rule: if the conclusion sequent is well-formed, the premises expressing typing conditions have derivations and the conditions expressed by the other, non-sequent premises are satisfied, then the premise sequents must be well-formed.*

Proof. The requirement is vacuously true for the *id* rule and it has an obvious proof for the *cut* rule. \square

In showing the soundness of the *id* rule, we will need the observation contained in the two lemmas below that the equivalence of formulas modulo permutations is preserved under the kinds of substitutions that have to be considered in determining the validity of sequents.

Lemma 4.25. *Suppose that $\Xi \vdash F_2 \equiv_\pi F_1$ holds for some formulas F_1 and F_2 and some Ξ and π of the right kinds. If θ is a term substitution such that $\text{supp}(\theta) \cap \text{supp}(\pi) = \emptyset$ and both $F_2[\theta] = F_2'$ and $F_1[\theta] = F_1'$ have derivations for some F_1' and F_2' , then $\Xi[\theta] \vdash F_2' \equiv_\pi F_1'$ holds.*

Proof. Let E be a type or term on which the application of the substitution θ is defined and let π be a permutation of the nominal constants. We observe then that $\pi.(E[\theta])$ is the same type or term as $(\pi.E)[\pi.\theta]$. This observation, which was implicit in Theorem 4.18, can be established by induction on the derivation of $E[\theta] = E'$ for the relevant E' . If $\text{supp}(\theta)$ and $\text{supp}(\pi)$ are disjoint as is assumed in this lemma, it further follows that $\pi.(E[\theta])$ is the same term or type as $(\pi.E)[\theta]$.

The lemma itself is proved by induction on the structure of F_1 or, equivalently, F_2 . Most of the cases are straightforward, the only one perhaps needing explicit consideration being that when F_1 and F_2 are the atomic formulas $\{G_1 \vdash M_1 : A_1\}$ and $\{G_2 \vdash M_2 : A_2\}$, respectively. In this case, by assumption, $\pi.M_2$ and M_1 are the same terms, $\pi.A_2$ and A_1 are the same types and $\Xi \vdash G_2 \equiv_\pi G_1$ holds. Using the observation at the beginning of the proof, we see that $\pi.(M_2[\theta])$ is the same term as $M_1[\theta]$ and $\pi.(A_2[\theta])$ is the same type as $A_1[\theta]$. Thus, the lemma would follow if we can show that $\Xi \vdash G_2[\theta] \equiv_\pi G_1[\theta]$ holds. However, this is easily done by an induction on the structure of G_1 . The argument is obvious when G_1 is \cdot or a context variable. If G_1 is of the form $(G_1', n_1 : A_1)$, then it must be the case that G_2 has the form $(G_2', n_2 : A_2)$, where $\pi.n_2$ is the same nominal constant as n_1 , $\pi.A_2$ is equal to A_1 up to renaming of bound variables, and $\Xi \vdash G_2' \equiv_\pi G_1'$ holds. Drawing again the observation at the beginning of the proof, we may conclude that $\pi.A_2[\theta]$ is the same type as $A_1[\theta]$. From the induction hypothesis, we know that $\Xi \vdash G_2'[\theta] \equiv_\pi G_1'[\theta]$ holds. It is now evident that $\Xi \vdash G_2[\theta] \equiv_\pi G_1[\theta]$ must hold. \square

Lemma 4.26. *Let σ be an appropriate substitution for Ξ with respect to some term variables context Ψ and set of nominal constants \mathbb{N} , and let $\Xi \vdash F' \equiv_\pi F$ hold. Then $\Xi_\sigma \vdash F'[\sigma] \equiv_\pi F[\sigma]$ must also hold.*

Proof. The key observation underlying the proof is that if G_1 and G_2 are two context expressions such that $\Xi \vdash G_1 \equiv_\pi G_2$ holds and σ is a context substitution that is appropriate for Ξ with respect to Ψ and \mathbb{N} , then $\Xi_\sigma \vdash G_1[\sigma] \equiv_\pi G_2[\sigma]$ must hold. This fact can be verified by an induction on the structure of G_1 . The only complex case is that where G_1 is a context variable Γ for which σ substitutes the context expression G' . Here, Ξ must contain an association of the form $\Gamma \uparrow \mathbb{N}' : \mathcal{C}[\mathcal{G}]$, with $\text{supp}(\pi) \subseteq \mathbb{N}'$ and with there being a derivation for $\text{supp}(\sigma) \setminus \mathbb{N}' ; \Psi ; \Xi_\sigma \vdash \mathcal{C}[\mathcal{G}] \rightsquigarrow_{\text{csty}} G'$. By an obvious inductive argument on the derivation, we can show that $\Xi_\sigma \vdash G' \equiv_\pi G'$ must hold, as we need to do in this case.

The lemma itself is proved by an obvious induction on the structure of F . In the case where F is the atomic formula $\{G \vdash M : A\}$, we make use of the above observation. \square

We can now show the soundness of the *id* and *cut* rules.

Theorem 4.27. *The following property holds for every instance of the *id* and *cut* rules: if the premises expressing typing judgements are derivable, the conditions described in the other non-sequent premises are satisfied and all the premise sequents are valid, then the conclusion sequent must also be valid.*

Proof. Let θ and σ identify a closed instance of the conclusion of the *id* rule. The requirements that θ and σ must satisfy and the assumption that the premise conditions $\text{supp}(\pi) \subseteq \mathbb{N}$ and $\Xi \vdash F' \equiv_\pi F$

hold allow us to use Lemmas 4.25 and 4.26 to conclude that $\emptyset \vdash F'[[\theta]][\sigma] \equiv_{\pi} F[[\theta]][\sigma]$ holds. Now, for any context variables context Ξ' , permutation π' , and closed formulas G and H it is easily seen that if $\Xi' \vdash G \equiv_{\pi'} H$ holds then H must be the same formula as $\pi'.G$. We may therefore conclude that the closed instance of the conclusion of the *id* rule identified by θ and σ is valid. Since the argument was independent of the choice of θ and σ , we conclude that the theorem must be true for this case.

For the *cut* rule, we note that, given the wellformedness premise pertaining to the formula F_2 , any substitutions θ and σ that identify a closed instance of the conclusion sequent must also identify closed instances of the two premise sequents. It is easy to see now that the validity of the latter two sequents must entail the validity of the former. We conclude that the theorem holds for this rule as well by noting that the argument is again independent of the choice of θ and σ . \square

4.2.3 Rules for the Logical Symbols

Figure 4.4 presents proof rules that facilitate reasoning based on the meanings of the logical symbols that are permitted in formulas. That we may focus attention only on well-formed sequents in the context of these rules is the content of the following theorem.

Theorem 4.28. *The following property holds of the rules in Figure 4.4: if the conclusion sequent is well-formed, the premises expressing typing conditions have derivations and the conditions expressed by the other, non-sequent premises are satisfied, then the premise sequents must be well-formed.*

Proof. The argument pertaining to \top , \perp and all the propositional connectives is straightforward, leaving us only to argue for the quantifier rules.

The considerations for the rules \forall -R and \exists -L are similar and we therefore discuss only the case for the former rule in detail. In showing that the context variables context for the premise sequent satisfies the wellformedness conditions, we observe that the corresponding context for the conclusion sequent does by assumption and we then use Theorem 4.3. In showing that the assumption formulas in the premise sequent satisfy the required conditions, we again use the fact that these formulas in the conclusion sequent do and then invoke Theorem 3.3. Thus, it only remains to show that F' satisfies the conditions required of it. From the wellformedness of $\forall x : \alpha.F$ relative to the conclusion sequent, we see that $\mathbb{N} \cup \Theta_0 \cup \Psi, x : \alpha; \Xi^- \vdash F \text{ fmla}$ has a derivation. The substitution $\{x, y, n_1 \dots n_m, \alpha\}$ is obviously type preserving with respect to the arity typing context $\mathbb{N} \cup \Theta_0 \cup (\Psi, y : \alpha_1 \rightarrow \dots \rightarrow \alpha_m \rightarrow \alpha)$. We may therefore invoke Theorem 3.5 to conclude that the judgement $\mathbb{N} \cup \Theta_0 \cup \Psi, y : \alpha_1 \rightarrow \dots \rightarrow \alpha_m \rightarrow \alpha; \Xi^- \vdash F' \text{ fmla}$ has a derivation.

The considerations for the rules \forall -L and \exists -R are also similar and so we discuss only the case for \forall -L in detail. Most of the wellformedness requirements for the premise sequent in fact follow trivially from the fact they hold for the conclusion sequent. The only perhaps nontrivial part is showing that F'_1 is well-formed. From the wellformedness of the conclusion sequent, we know that $\mathbb{N} \cup \Theta_0 \cup \Psi \cup \{x : \alpha\}; \Xi^- \vdash F_1 \text{ fmla}$ has a derivation. Since $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{at} t : \alpha$ has a derivation, $\{x, t, \alpha\}$ is an arity type preserving substitution with respect to $\mathbb{N} \cup \Theta_0 \cup \Psi$. We can now invoke Theorem 3.5 to conclude that there must be a derivation for $\mathbb{N} \cup \Theta_0 \cup \Psi; \Xi^- \vdash F'_1 \text{ fmla}$.

We consider the case for Π -R next. Here we must show that the addition $\Gamma' \uparrow \emptyset : \mathcal{C}[\cdot]$ to the context variables context in the premise sequent satisfies the wellformedness requirements. That it does follows easily from the observations that $\vdash \mathcal{C}$ *ctx schema* must be derivable because the formula $\Pi \Gamma : \mathcal{C}.F$ is well-formed (relative to the conclusion sequent) and that $\mathbb{N}; \Psi \vdash \mathcal{C}[\cdot]$ *ctx-ty* has a trivial derivation. The wellformedness of all the other associations in the context variables context follows immediately from the wellformedness of the conclusion sequent and the wellformedness of the formulas in Ω can be argued similarly, additionally using Theorem 3.3. Finally, using the fact

$$\begin{array}{c}
\frac{}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \top} \top\text{-R} \quad \frac{}{\mathbb{N}; \Psi; \Xi; \Omega, \perp \longrightarrow F} \perp\text{-L} \\
\frac{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F_1 \quad \mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F_2}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F_1 \wedge F_2} \wedge\text{-R} \quad \frac{\mathbb{N}; \Psi; \Xi; \Omega, F_i \longrightarrow F \quad i \in \{1, 2\}}{\mathbb{N}; \Psi; \Xi; \Omega, F_1 \wedge F_2 \longrightarrow F} \wedge\text{-L}_i \\
\frac{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F_i \quad i \in \{1, 2\}}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F_1 \vee F_2} \vee\text{-R}_i \quad \frac{\mathbb{N}; \Psi; \Xi; \Omega, F_1 \longrightarrow F \quad \mathbb{N}; \Psi; \Xi; \Omega, F_2 \longrightarrow F}{\mathbb{N}; \Psi; \Xi; \Omega, F_1 \vee F_2 \longrightarrow F} \vee\text{-L} \\
\frac{\mathbb{N}; \Psi; \Xi; \Omega, F_1 \longrightarrow F_2}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F_1 \supset F_2} \supset\text{-R} \quad \frac{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F_1 \quad \mathbb{N}; \Psi; \Xi; \Omega, F_2 \longrightarrow F}{\mathbb{N}; \Psi; \Xi; \Omega, F_1 \supset F_2 \longrightarrow F} \supset\text{-L} \\
\frac{\mathbb{N} = \{n_1 : \alpha_1, \dots, n_m : \alpha_m\} \quad y \notin \text{dom}(\Psi) \quad F[\{\langle x, y \ n_1 \dots n_m, \alpha \rangle\}] = F' \quad \mathbb{N}; \Psi \cup \{y : (\alpha_1 \rightarrow \dots \rightarrow \alpha_m \rightarrow \alpha)\}; \Xi; \Omega \longrightarrow F'}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \forall x : \alpha. F} \forall\text{-R} \\
\frac{\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{at} t : \alpha \quad F_1[\{\langle x, t, \alpha \rangle\}] = F'_1 \quad \mathbb{N}; \Psi; \Xi; \Omega, F'_1 \longrightarrow F_2}{\mathbb{N}; \Psi; \Xi; \Omega, \forall x : \alpha. F_1 \longrightarrow F_2} \forall\text{-L} \\
\frac{\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{at} t : \alpha \quad F[\{\langle x, t, \alpha \rangle\}] = F' \quad \mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F'}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \exists x : \alpha. F} \exists\text{-R} \\
\frac{\mathbb{N} = \{n_1 : \alpha_1, \dots, n_m : \alpha_m\} \quad y \notin \text{dom}(\Psi) \quad F_1[\{\langle x, y \ n_1 \dots n_m, \alpha \rangle\}] = F'_1 \quad \mathbb{N}; \Psi \cup \{y : (\alpha_1 \rightarrow \dots \rightarrow \alpha_m \rightarrow \alpha)\}; \Xi; \Omega, F'_1 \longrightarrow F_2}{\mathbb{N}; \Psi; \Xi; \Omega, \exists x : \alpha. F_1 \longrightarrow F_2} \exists\text{-L} \\
\frac{\mathbb{N}; \Psi; \Xi, \Gamma' \uparrow \emptyset : \mathcal{C}[\cdot]; \Omega \longrightarrow F[\{\Gamma'/\Gamma\}] \quad \Gamma' \notin \text{dom}(\Xi)}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \Pi \Gamma : \mathcal{C}.F} \Pi\text{-R} \\
\frac{\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\cdot] \rightsquigarrow_{csty} G \quad \mathbb{N}; \Psi; \Xi; \Omega, F_1[\{\langle G/\Gamma \rangle\}] \longrightarrow F_2}{\mathbb{N}; \Psi; \Xi; \Omega, \Pi \Gamma : \mathcal{C}.F_1 \longrightarrow F_2} \Pi\text{-L}
\end{array}$$

Figure 4.4: The Logical Rules

that $\mathbb{N} \cup \Theta_0 \cup \Psi; \Xi^- \vdash \Pi \Gamma : \mathcal{C}.F$ *fmla* has a derivation, we can easily show that there must be a derivation for $\mathbb{N} \cup \Theta_0 \cup \Psi; \Xi^- \cup \{\Gamma'\} \vdash F[\{\Gamma'/\Gamma\}]$ *fmla*.

The only remaining case is that of $\Pi\text{-L}$. Moreover, the only requirement whose validation is nontrivial in this case is that there is a derivation for $\mathbb{N} \cup \Theta \cup \Psi; \Xi^- \vdash F_1[\{\langle G/\Gamma \rangle\}]$ *fmla*. There is a trivial derivation for $\mathbb{N}; \Psi \vdash \mathcal{C}[\cdot]$ *ctx-ty* and, from one of the premises, we know that there is a derivation for $\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\cdot] \rightsquigarrow_{csty} G$. But then, from Theorem 4.2, it follows that $\mathbb{N}; \Psi \vdash G$ *context* has a derivation. By the well-formedness of the conclusion sequent, $\mathbb{N} \cup \Theta \cup \Psi; \Xi^- \vdash \Pi \Gamma : \mathcal{C}.F_1$ *fmla* has a derivation, from which it follows that $\mathbb{N} \cup \Theta \cup \Psi; \Xi^- \cup \{\Gamma\} \vdash F_1$ *fmla* must also have one. We now use Theorem 3.5 to reach the desired conclusion. \square

The proof of soundness of the rules $\Pi\text{-L}$, $\forall\text{-L}$ and $\exists\text{-R}$ will use inversions in the order of application of substitutions that are justified by the three lemmas below.

Lemma 4.29. *Let θ_1 and θ_2 be term variables substitutions that are arity type preserving with respect to Θ and $\text{ctx}(\theta_1) \uplus \Theta$, respectively, and such that the variables in $\text{dom}(\theta_2)$ are distinct from*

those in $\text{dom}(\theta_1)$ and do not appear free in the terms in $\text{rng}(\theta_1)$. Further, let θ'_2 be the substitution

$$\{\langle x, M', \alpha \rangle \mid \langle x, M, \alpha \rangle \in \theta_2 \text{ and } M[\theta_1] = M'\}.$$

If F is a formula such that $\text{ctx}(\theta_2) \uplus (\text{ctx}(\theta_1) \uplus \Theta); \Xi^- \vdash F$ *fm*la is derivable for some context variables context Ξ , then both $F[\theta_2][\theta_1]$ and $F[\theta_1][\theta'_2]$ are defined and are in fact the same formulas.

Proof. We note first that the substitution θ'_2 is well-defined by Theorem 2.8. The lemma is proved by an induction on the structure of formulas, making use eventually of Theorem 2.6. \square

Lemma 4.30. *Let σ_1 and σ_2 be context variables substitutions such that the variables substituted for by σ_2 are distinct from those substituted for by σ_1 and they also do not appear in the context expressions in the range of σ_1 . Further, let σ'_2 be the context variables substitution*

$$\{G'/\Gamma \mid G/\Gamma \in \sigma_2 \text{ and } G[\sigma_1] = G'\}.$$

Then, for any formula F , $F[\sigma_2][\sigma_1]$ and $F[\sigma_1][\sigma'_2]$ represent identical formulas.

Proof. Using an inductive argument, we first show the property for context expressions and then extend it to formulas. \square

Lemma 4.31. *Let σ be a context variables substitution appropriate for Ξ with respect to $\text{ctx}(\theta) \cup \Psi$ and \mathbb{N} , let θ be an arity type preserving substitution with respect to $\mathbb{N} \cup \Theta_0 \cup \Psi$, and let F be a formula such that $\mathbb{N} \cup \Theta_0 \cup (\text{ctx}(\theta) \uplus \Psi); \Xi^- \vdash F$ *fm*la is derivable. Then, if σ' be the substitution $\{G'/\Gamma \mid G/\Gamma \in \sigma \text{ and } G[\theta] = G'\}$, $F[\sigma][\theta]$ and $F[\theta][\sigma']$ denote the same formulas.*

Proof. Note first that for any $G/\Gamma \in \sigma$ it must be the case that $\mathbb{N} \cup \Theta_0 \cup (\text{ctx}(\theta) \uplus \Psi); \Xi_\sigma^- \vdash G$ context is derivable; this follows from the appropriateness property for σ , using Theorem 4.2. It is easily seen from this that $G[\theta]$ must be defined and, hence, that the substitution σ' is well-defined. Using Theorem 3.5, it also follows that $\mathbb{N} \cup \Theta_0 \cup (\text{ctx}(\theta) \uplus \Psi); \Xi^- \vdash F[\sigma]$ *fm*la is derivable. Using Theorem 3.5 again, it follows that $F[\sigma][\theta]$ must be defined. Finally, by Theorem 2.8, $F[\theta]$ must be defined. This establishes the coherence of the lemma statement. Its proof itself is based on an induction on the structure of the formula F , noting that term substitutions leave context variables unchanged. \square

The soundness of the logical rules is the content of the following theorem.

Theorem 4.32. *The following property holds for every instance of each of the rules in Figure 4.4: if the premises expressing typing judgements are derivable, the conditions described in the other non-sequent premises are satisfied and all the premise sequents are valid, then the conclusion sequent must also be valid.*

Proof. The argument is straightforward for the rules pertaining to the logical constants and propositional connectives. We therefore focus on the quantifier rules in the rest of the proof. The wellformedness of sequents ensures that the applications of substitutions to formulas and terms must be defined in actual instances of these rules. In light of this, we will freely use the notation introduced after Theorems 2.8 in the discussions below.

The arguments for the \forall -R and \exists -L rules are similar, so we present only the argument for the \forall -R rule. Suppose that the claim is false in this case. Then there must be substitutions θ and σ that identify a closed instance of the conclusion sequent that is not valid. We may assume without loss of generality that $y \notin \text{dom}(\theta)$; if this is not true at the outset, since $y \notin \text{dom}(\Psi)$, we may drop the tuple pertaining to y from θ to get another term substitution on which to base the argument. Now, in the

situation under consideration, all the formulas in $\Omega[\theta][\sigma]$ must be valid and $(\forall x : \alpha.F)[\theta][\sigma]$ must not be valid. The latter means that there is some term t for which $\mathcal{N} \cup \Theta_0 \vdash_{at} t : \alpha$ has a derivation that is such that $F[\theta][\sigma][\{\langle x, t, \alpha \rangle\}]$ is not valid; note that x is a variable that does not appear in θ or σ . Using Lemma 4.31, we see that this is the same formula as $F[\theta][\{\langle x, t, \alpha \rangle\}][\sigma]$. Using Theorem 4.8, we can conclude that there must be a term t' that is devoid of nominal constants from \mathbb{N} for which the typing judgement $\mathcal{N} \cup \Theta_0 \vdash_{at} t' : \alpha_1 \rightarrow \dots \rightarrow \alpha_m \rightarrow \alpha$ is derivable and that is such that $F[\theta][\{\langle x, t, \alpha \rangle\}][\sigma]$ is the same formula as $F[\theta][\{\langle x, y \ n_1 \dots n_m, \alpha \rangle\}][\{\langle y, t', \alpha \rangle\}][\sigma]$. Since x does not appear in θ and y is not in $\text{dom}(\theta)$, we can conclude using Lemma 4.29 that the latter formula is the same as $F[\{\langle x, y \ n_1 \dots n_m, \alpha \rangle\}][\theta][\{\langle y, t', \alpha_1 \rightarrow \dots \rightarrow \alpha_m \rightarrow \alpha \rangle\}][\sigma]$. Using Theorem 2.10, we see further that this formula is the same as $F[\{\langle x, y \ n_1 \dots n_m, \alpha \rangle\}][\theta']$, where θ' is the substitution $\theta \cup \{\langle y, t', \alpha_1 \rightarrow \dots \rightarrow \alpha_m \rightarrow \alpha \rangle\}$. We observe at this point that θ' and σ are substitutions that identify a closed instance of the premise sequent: since t' is a closed term of the right type that is devoid of the nominal constants in \mathbb{N} , it follows easily that θ' and σ must be “proper” for the premise sequent if θ and σ are proper for the conclusion sequent, the formulas in $\Omega[\theta'][\sigma]$ are identical to the ones in $\Omega[\theta][\sigma]$ and hence all closed, and the formula $F[\{\langle x, y \ n_1 \dots n_m, \alpha \rangle\}][\theta'][\sigma]$ is identical to $F[\theta][\sigma][\{\langle x, t, \alpha \rangle\}]$ and hence also closed. However, we now have a contradiction since this closed instance of the premise sequent must also be one that is not valid.

The arguments for the \forall -L and \exists -R rules are also similar, so we present only the one for the former rule. Let θ and σ be substitutions that identify a closed instance of the conclusion sequent. Now, if any formula in $(\Omega \cup \forall x : \alpha.F_1)[\theta][\sigma]$ is not valid, then the conclusion sequent must be valid. Let us therefore assume that all these formulas are valid. Our task is to show that then $F_2[\theta][\sigma]$ must also be valid. Clearly, θ and σ also identify a closed instance of the premise sequent. If we can show that the assumption formula $F_1[\{\langle x, t, \alpha \rangle\}][\theta][\sigma]$ in this sequent is valid, then all its assumption formulas would be valid, thereby enabling us to conclude that $F_2[\theta][\sigma]$ is also valid, as desired. Using Lemmas 4.29 and 4.31, we see that the assumption formula in question is the same as $F_1[\theta][\sigma][\{\langle x, t[\theta], \alpha \rangle\}]$. Since $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{at} t : \alpha$ has a derivation, by Theorem 2.8 there must be one for $\mathbb{N} \cup \Theta_0 \vdash_{at} t[\theta] : \alpha$. But this means that $F_1[\theta][\sigma][\{\langle x, t[\theta], \alpha \rangle\}]$ is a closed instance of $\forall x : \alpha.F_1[\theta][\sigma]$ and must therefore be valid by assumption.

Let us consider the case for Π -R next. If the claim is false, then there must be substitutions θ and σ that identify a closed instance of the conclusion sequent that is not valid. For this to happen, it must be the case that all the formulas in $\Omega[\theta][\sigma]$ are valid and the formula $\Pi \Gamma : \mathcal{C}.F[\theta][\sigma]$ is not valid. From the latter, it follows that for some context expression G it is the case that $\mathcal{N}; \emptyset \vdash \mathcal{C} \rightsquigarrow_{cs} G$ is derivable and $F[\theta][\sigma][\{G/\Gamma\}]$ is not valid. But now consider the substitutions θ and σ' where $\sigma' = \sigma \cup \{G/\Gamma'\}$. It is easily checked that these are proper for the premise sequent. Further, $F[\theta][\sigma']$ is the same formula as $F[\theta][\sigma][\{G/\Gamma\}]$ and, since $\Gamma' \notin \text{dom}(\Xi)$, $\Omega[\theta][\sigma']$ is the same collection of formulas as $\Omega[\theta][\sigma]$. Thus θ and σ' identify a closed instance of the premise sequent that is not valid, thereby contradicting the assumption of falsity of the theorem.

The only remaining case is that of Π -L. Let θ and σ be substitutions that identify a closed instance of the conclusion sequent. If any formula in $(\Omega \cup \{\Pi \Gamma : \mathcal{C}.F\})[\theta][\sigma]$ is not valid, then the conclusion sequent must be valid. Let us therefore assume that all these formulas are valid. We must show that $F_2[\theta][\sigma]$ is then also valid. By assumption, there is a derivation for $\mathbb{N}; \Psi; \Xi \vdash \mathcal{C}[\cdot] \rightsquigarrow_{csty} G$. Using Theorem 4.3, it follows that there must be a derivation for $\mathcal{N}; \Psi; \Xi \vdash \mathcal{C}[\cdot] \rightsquigarrow_{csty} G$. Using Theorems 4.4 and 4.12 we can show from this that $\mathcal{N}; \emptyset; \emptyset \vdash \mathcal{C}[\cdot] \rightsquigarrow_{csty} G[\theta][\sigma]$ is derivable; to invoke Theorem 4.12, we would need σ to be appropriate for $\Xi[\theta]$ with respect to \emptyset and \mathcal{N} but this is ensured by the properness of θ and σ with respect to the conclusion sequent. By Theorem 4.2, there must be a derivation for $\mathcal{N}; \emptyset \vdash \mathcal{C} \rightsquigarrow_{cs} G[\theta][\sigma]$. Clearly, θ and σ must identify a closed instance of the premise sequent. Moreover, all the assumption formulas in this instance must be

valid: we have assumed this to be the case for the formulas in $\Omega[\theta][\sigma]$ and $F_1[G/\Gamma][\theta][\sigma]$ must be valid because $\Pi\Gamma : \mathcal{C}.F[\theta][\sigma]$ is valid and $\mathcal{N};\emptyset \vdash \mathcal{C} \rightsquigarrow_{cs} G[\theta][\sigma]$ is derivable. But then, by the assumption of validity of the premise sequent, it must be the case that $F_2[\theta][\sigma]$ is valid. \square

4.3 Proof Rules that Interpret Atomic Formulas

We now present rules that build in an analysis of atomic formulas based on the understanding that they represent LF typing judgements. When an atomic formula appears as the conclusion of a sequent, the analysis takes an obvious form: the derivability of the sequent can be based on that of a sequent in which the conclusion judgement has been unfolded using an LF rule. The treatment of an atomic assumption formula is more complex: we must consider all the ways in which this formula could be valid in assessing the validity of the sequent. This “case analysis” can be driven by the structure of the type in the formula. When the type is of the form $\Pi x:A. B$, the term must be an abstraction and there is exactly one way in which a purported typing derivation could have concluded. When the type is atomic, Theorem 2.20 provides us information about the different cases that need to be considered. However, a complicating factor is that the formula may contain term and context variables in it and we must consider also all the possible instantiations of these variables that could made the judgement true in the analysis.

The analysis of atomic assumption formulas is obviously somewhat intricate and we devote the first part of this subsection to its discussion. We then use the resulting understanding to develop proof rules for atomic formulas.

4.3.1 Analyzing an Atomic Assumption Formula with an Atomic Type

Consider an atomic formula $\{G \vdash R : P\}$ where, obviously, P has the form $(a M_1 \dots M_n)$. Let us suppose initially that this formula is closed. In this case, for the formula to be valid, R would need to have as a head a constant declared in Σ or a nominal constant assigned a type in G . If the arguments of R do not satisfy the constraints imposed by the type associated with the head, then the typing judgement will not be derivable and hence we can conclude that the sequent is in fact valid. On the other hand, if the arguments of R do satisfy the required constraints, then Theorem 2.20 gives us a means for decomposing the given typing judgement into ones pertaining to M_1, \dots, M_n . The validity of the given sequent can therefore be reduced to the validity of a sequent that results from replacing the atomic formula under consideration by ones that represent the mentioned typing judgements.

In the general case, the formula $\{G \vdash R : P\}$ may not be closed. This could happen in two different ways. First, the context expression may have a part that is yet to be determined, i.e., G may be of the form $\Gamma, n_1 : A_1, \dots, n_m : A_m$ where Γ has a set of names \mathbb{N} and a context variable type of the form $\mathcal{C}[G_1; \dots; G_\ell]$ associated with it in the sequent. Second, the expressions in the atomic formula and the context variable type may contain variables in them that are bound in the term variables context. To articulate a proof rule around the atomic formula in this situation, it is necessary to develop a means for analyzing the formula in a way that pays attention to the validity of the sequent under all acceptable instantiations of the context and term variables.

The analysis that we describe proceeds in two steps. We first describe a finite way to consider elaborations of the context variable that make explicit all the heads that need to be considered for the term in an analysis of the closed instances of the atomic formula. This process yields a finite collection of pairs comprising a sequent in which the context variable may have been partially instantiated, and a specifically identified possibility for the head that is either drawn from the signature or that appears explicitly in the context; the intent here, which is verified in Lemma 4.39,

is that considering just the second components of these pairs as the heads of the term in the typing judgement will suffice for a complete analysis based on Theorem 2.20. The second step actually carries out the analysis in each of these cases, using the idea of unification in the application of Theorem 2.20 to accommodate all possible closed instantiations of the term variables in the sequent.

Elaborating Context Variables and Identifying Head Possibilities We first note that context expressions may have implicit and explicit parts, the former being subject to elaboration via context substitutions.

Definition 4.33. *Let $\mathcal{S} = \mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$ be a well-formed sequent. If G is a context expression appearing in \mathcal{S} then it must be of either the form $n_1 : A_1, \dots, n_m : A_m$ or of the form $\Gamma, n_1 : A_1, \dots, n_m : A_m$ where Γ is a context variable with an associated declaration $\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[G_1; \dots; G_n]$ in Ξ . In the latter case, we say that G has an implicit part relative to \mathcal{S} that is given by $\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[G_1; \dots; G_n]$. Further, we refer to $n_1 : A_1, \dots, n_m : A_m$ in the former case and to the sequence formed by listing the bindings in G_1, \dots, G_n followed by $n_1 : A_1, \dots, n_m : A_m$ in the latter case as the explicit bindings in G relative to \mathcal{S} .*

Let Γ be a context variable that has the type $\mathcal{C}[G_1; \dots; G_\ell]$. Closed instances of Γ are then generated by interspersing G_1, \dots, G_ℓ with blocks of declarations generated from the block schema comprising \mathcal{C} . In determining possibilities for the head of R from the implicit part of G in an atomic formula of the form $\{G \vdash R : P\}$, we need to consider an elaboration of G with only one such block; of course, for a complete analysis, we will need to consider all the possibilities for such an elaboration. The function *AddBlock* defined below formalizes such an elaboration, returning a modified sequent and a potential head for the term in the typing judgement. Note that in an elaboration based on a block schema of the form $\{x_1 : \alpha_1, \dots, x_n : \alpha_n\} y_1 : A_1, \dots, y_k : A_k$, it would be necessary to consider a choice of nominal constants for the schematic variables y_1, \dots, y_k . The function is parameterized by such a choice. We must also accommodate all possible instantiations for the variables x_1, \dots, x_n , subject to the proviso that these instantiations do not use nominal constants that appear in a later part of the context expression. This is done by introducing new term variables for x_1, \dots, x_n and by raising such variables over the nominal constants that are not prohibited from appearing in the instantiations; to support the latter requirement, the function is parameterized by a collection of nominal constants. Finally, we observe that the elaboration process may introduce new nominal constants into the sequent, necessitating a raising of the term variables over the new constants.

Definition 4.34. *Let \mathcal{S} be the well-formed sequent $\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$, let $\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[G_1; \dots; G_n]$ be an assignment in Ξ , and let $\mathcal{B} = \{x_1 : \alpha_1, \dots, x_n : \alpha_n\} y_1 : A_1, \dots, y_k : A_k$ be one of the block schemas comprising \mathcal{C} . Further, let $\mathbb{N}' \subseteq (\mathbb{N} \setminus \mathbb{N}_\Gamma)$ be a collection of nominal constants, let ns be a list n_1, \dots, n_k of distinct nominal constants that are also different from the constants in \mathbb{N}' and that are such that, for $1 \leq i \leq k$, $n_i : (A_i)^- \in (\mathcal{N} \setminus \mathbb{N}_\Gamma)$. Finally, for $0 \leq j \leq n$, let \mathbb{N}_j be the collection of nominal constants assigned types in G_1, \dots, G_j . Then, letting*

1. Ψ'_j be a version of Ψ raised over $\{n_1, \dots, n_k\} \setminus \mathbb{N}$ and θ'_j be the associated raising substitution,
2. A'_1, \dots, A'_k be the types A_1, \dots, A_k with the schematic variables y_1, \dots, y_k replaced with the names n_1, \dots, n_k ,
3. Ψ''_j be a version of $\{x_1 : \alpha_1, \dots, x_n : \alpha_n\}$ raised over $\mathbb{N}' \cup \mathbb{N}_j \cup (\{n_1, \dots, n_k\} \setminus \mathbb{N})$ with the new variables chosen to be distinct from those in Ψ'_j , θ''_j be the associated raising substitution, and G be the context expression $n_1 : A'_1 \llbracket \theta''_j \rrbracket, \dots, n_k : A'_k \llbracket \theta''_j \rrbracket$, and

4. Ξ'_j be the context variables context

$$(\Xi \setminus \{\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[G_1; \dots; G_n]\})[\theta'_j] \cup \{\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[G_1[\theta'_j]; \dots; G_j[\theta'_j]; G; G_{j+1}[\theta'_j]; \dots; G_n[\theta'_j]]\},$$

for $0 \leq j \leq n$ and $1 \leq i \leq k$, $\text{AddBlock}(\mathcal{S}, \Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[G_1; \dots; G_n], \mathcal{B}, ns, \mathbb{N}', j, i)$ is defined to be the tuple

$$\langle \mathbb{N} \cup ns; \Psi'_j \cup \Psi''_j; \Xi'_j; \Omega[\theta'_j] \longrightarrow F[\theta'_j], n_i : A'_i[\theta''_j] \rangle.$$

Note that the conditions in the definition ensure that all the substitutions involved in it will have a result, thereby permitting us to use the notation introduced after Theorem 2.8.

The elaboration just described is parameterized by the choice of nominal constants for the variables assigned types in the block schema. In identifying the choices that have to be considered, it is useful to partition the members of $(\mathcal{N} \setminus \mathbb{N}_\Gamma)$ into two sets: those that appear in the support set of the sequent whose elaboration is being considered and those that do not. It is necessary to consider all possible assignments that satisfy arity typing constraints from the first category. From the second category, as we shall soon see, it suffices to consider exactly one representative assignment. Note also that we may insist that the nominal constant in each assignment of the block be distinct; if this is not the case, the sequent is easily seen to be valid. The function *NamesLsts* defined below embodies these ideas. The function is parameterized by a sequence of arity types corresponding to the declarations in the block schema, a collection of “known” nominal constants that are available for use in an elaboration of the block schema and a collection of nominal constants that are already bound in the context expressions and hence must not be used again.

Definition 4.35. Let *tys* be a sequence of arity types and let \mathbb{N}_o and \mathbb{N}_b be finite sets of nominal constants. Further, let *nil* denote an empty sequence and $x :: xs$ denote a sequence that starts with *x* and continues with the sequence *xs*. Then the collection of name choices for *tys* relative to \mathbb{N}_o and away from \mathbb{N}_b is denoted by $\text{NamesLsts}(tys, \mathbb{N}_o, \mathbb{N}_b)$ and defined by recursion on *tys* as follows:

$$\text{NamesLsts}(tys, \mathbb{N}_o, \mathbb{N}_b) = \begin{cases} \{\text{nil}\} & \text{if } tys = \text{nil} \\ \{n :: nl \mid n : \alpha \in \mathcal{N}, n \in \mathbb{N}_o \setminus \mathbb{N}_b, \text{ and} \\ \quad nl \in \text{NamesLsts}(tys', \mathbb{N}_o, \mathbb{N}_b \cup \{n\})\} \cup \\ \{n :: nl \mid n \text{ is the first nominal constant} & \text{if } tys = \alpha :: tys' \\ \quad \text{such that } n : \alpha \in \mathcal{N} \text{ and } n \notin \mathbb{N}_o \cup \mathbb{N}_b, \\ \quad \text{and } nl \in \text{NamesLsts}(tys', \mathbb{N}_o, \mathbb{N}_b \cup \{n\})\} \end{cases}$$

We assume in this definition the existence of an ordering on the nominal constants that allows us to select the first of these constants that satisfies a criterion of interest.

We can now identify a finite collection of elaborations of the implicit part of a context expression that must be considered in the analysis of an assumption formula of the form $\{G \vdash R : P\}$ that appears in a sequent \mathcal{S} . We do this below through the definition of the function *ImplicitHeads*.

Definition 4.36. Let \mathcal{S} be the well-formed sequent $\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$, let *G* be a context expression appearing in a formula in \mathcal{S} that has an implicit part relative to \mathcal{S} that is given by $\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[G_1; \dots; G_n]$, and let $\mathcal{B} = \{x_1 : \alpha_1, \dots, x_n : \alpha_n\}y_1 : A_1, \dots, y_k : \alpha_k$ be one of the block schemas comprising \mathcal{C} . Further, let \mathbb{N}_b be the collection of nominal constants assigned types by the explicit bindings of *G* relative to \mathcal{S} and let $\mathbb{N}_o = (\mathbb{N} \setminus \mathbb{N}_\Gamma) \setminus \mathbb{N}_b$. Finally, let $\text{AllBlocks}(\mathcal{S}, \Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[G_1; \dots; G_n], \mathcal{B})$ denote the set

$$\{AddBlock(\mathcal{S}, \Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[G_1; \dots; G_n], \mathcal{B}, ns, \mathbb{N}_o, j, i) \mid \\ 0 \leq j \leq n, 1 \leq i \leq k, ns \in NamesLsts(((A_1)^-, \dots, (A_k)^-), \mathbb{N}_o, \mathbb{N}_\Gamma \cup \mathbb{N}_b)\}.$$

If $\{\mathcal{B}_1, \dots, \mathcal{B}_m\}$ is the collection of block schemas comprising \mathcal{C} , then the implicit heads in G relative to \mathcal{S} is defined to be the set

$$\bigcup \{AllBlocks(\mathcal{S}, \Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[G_1; \dots; G_n], \mathcal{B}) \mid \mathcal{B} \in \{\mathcal{B}_1, \dots, \mathcal{B}_m\}\}.$$

This set is denoted by $ImplicitHeds(\mathcal{S}, G)$.

The complete set of heads and corresponding (elaborated) sequents that must be considered in the analysis of an atomic formula of the form $\{G \vdash R : P\}$ is identified through the function $Heds$ that is defined below.

Definition 4.37. Let \mathcal{S} be a well-formed sequent and let G be a context expression appearing in a formula in \mathcal{S} . Let $NewHds$ be the set $ImplicitHeds(\mathcal{S}, G)$ if G has an implicit part relative to \mathcal{S} and the empty set otherwise. Then the heads in G relative to \mathcal{S} is defined to be the set

$$\{\langle \mathcal{S}, c : A \rangle \mid c : A \in \Sigma\} \cup \{\langle \mathcal{S}, n : A \rangle \mid n : A \text{ is an explicit binding in } G \text{ relative to } \mathcal{S}\} \cup NewHds.$$

This set is denoted by $Heds(\mathcal{S}, G)$.

The first property that we observe of the elaboration process described is that it requires us to consider only well-formed sequents.

Lemma 4.38. Let $\mathcal{S} = \mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$ be a well-formed sequent and let $\{G \vdash R : P\}$ be an atomic formula in Ω . Then for each $(\mathcal{S}', h : A) \in Heds(\mathcal{S}, G)$ it must be the case that \mathcal{S}' is a well-formed sequent. Further, if \mathcal{S}' is $\mathbb{N}'; \Psi'; \Xi'; \Omega' \longrightarrow F'$, it must be the case that $(\mathbb{N}' \cup \Theta_0 \cup \Psi') \vdash_{ak} A$ type is derivable.

Proof. The claim is not immediately obvious only when $(\mathcal{S}', h : A) \in ImplicitHeds(\mathcal{S}, G)$. For these cases, it suffices to show that every pair generated by $AddBlock$ satisfies the requirements of the lemma. However, this is easily argued. The main observation—that gets used twice—is that if Ψ_2 is a version of Ψ_1 raised over some collection of nominal constants \mathbb{N}_2 with θ being the associated raising substitution, and $\Theta \cup \Psi_1 \vdash_{ak} A'$ type holds for some arity context Θ that is disjoint from Ψ_1 and Ψ_2 , then $\Theta \cup \Psi_2 \cup \mathbb{N}_2 \vdash_{ak} A'[\theta]$ type also holds. \square

We want next to show that the collection of pairs of sequents and heads identified by the elaboration process are sufficient for the analysis of validity for a sequent with an assumption formula of the form $\{G \vdash R : P\}$. The following lemma provides the basis for this observation. We obviously do to identify all possible elaborations because we consider only a single representative for a “new name” for a binding in a block instance. However, we show that we cover all elaborations up to a permutation; Theorem 4.19 then ensures that this is sufficient.

Lemma 4.39. Let $\mathcal{S} = \mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F$ be a well-formed sequent and let $\{G \vdash R : P\}$ be a formula in Ω . Further, let θ and σ be term and context variables substitutions that identify a closed instance of \mathcal{S} and that are such that $\{G \vdash R : P\}[\theta][\sigma]$ is valid. If the term $R[\theta] = (h M_1 \dots M_n)$, then there is a pair $\langle \mathcal{S}', h' : A' \rangle$ in $Heds(\mathcal{S}, G)$ such that

1. there is a formula $\{G' \vdash R' : P'\}$ amongst the assumption formulas of \mathcal{S}' with $h' : A'$ appearing in either Σ or in the explicit bindings in G' relative to \mathcal{S}' , and

2. there is a closed instance of \mathcal{S}' identified by term and context variables substitutions θ' and σ' and a permutation π such that $\pi.h' = h$, $\pi.\{G \vdash R : P\} \llbracket \theta' \rrbracket [\sigma'] = \{G \vdash R : P\} \llbracket \theta \rrbracket [\sigma]$, and $\pi.\mathcal{S}' \llbracket \theta' \rrbracket [\sigma'] = \mathcal{S} \llbracket \theta \rrbracket [\sigma]$.

Proof. Since $\{G \vdash R : P\} \llbracket \theta \rrbracket [\sigma]$ is valid, it must be the case that there are LF derivations for $\vdash_{\Sigma} G \llbracket \theta \rrbracket [\sigma]$ **ctx**, $G \llbracket \theta \rrbracket [\sigma] \vdash_{\Sigma} P \llbracket \theta \rrbracket [\sigma]$ **type**, and $G \llbracket \theta \rrbracket [\sigma] \vdash_{\Sigma} R \llbracket \theta \rrbracket [\sigma] \Leftarrow P \llbracket \theta \rrbracket [\sigma]$. Using Theorem 2.20 together with the fact that $R \llbracket \theta \rrbracket [\sigma] = (h M_1 \dots M_n)$, we see that, for an appropriate A , $h : A$ must be a member of Σ or it must appear in $G \llbracket \theta \rrbracket [\sigma]$. Our argument distinguishes two ways that this could happen: it could be because $h : A$ is a member of Σ or it is an instance of a declaration in the explicit part of G or because it is introduced into the context $G \llbracket \theta \rrbracket [\sigma]$ by the substitution σ .

The first collection of cases is easily dealt with: essentially, we pick h' , \mathcal{S}' , θ' and σ' to be identical to h , \mathcal{S} , θ and σ , respectively, and we let π be the identity permutation. The requirements of the lemma then follow easily from the definition of the *Heads* function.

In the cases that remain, G must have the form $\Gamma, n_1^G : A_1^G, \dots, n_p^G : A_p^G$ for some context variable Γ that has the set of names \mathbb{N}_{Γ} and the type $\mathcal{C}[G_1; \dots; G_{\ell}]$ assigned to it in Ξ and h must be introduced by the substitution that σ makes for Γ as the i^{th} binding, for some i , in a block of declarations resulting from instantiating one of the block schemas constituting \mathcal{C} . Let us suppose the relevant block schema is \mathcal{B} and it has the form $\{x_1 : \alpha_1, \dots, x_n : \alpha_n\}(y_1 : B_1, \dots, y_k : B_k)$. Moreover, let us suppose that this block of declarations appears in $G \llbracket \theta \rrbracket [\sigma]$ somewhere between the instances of G_j and G_{j+1} , for some j between 0 and ℓ . We may, without loss of generality, assume x_1, \dots, x_n to be distinct from the variables assigned types by Ψ . We can then visualize the block introducing h as $(n_1 : B'_1 \llbracket \theta^h \rrbracket, \dots, n_k : B'_k \llbracket \theta^h \rrbracket)$ for some n_1, \dots, n_k of the requisite types, for some types B'_1, \dots, B'_k which are the types B_1, \dots, B_k with the schematic variables of the schema replaced by these names, and for a closed substitution θ^h whose domain is x_1, \dots, x_n and, since $\vdash_{\Sigma} G \llbracket \theta \rrbracket [\sigma]$ **ctx** is derivable, whose support does not contain the nominal constants in \mathbb{N}_{Γ} , n_1^G, \dots, n_p^G , or those that are assigned a type in G_{j+1}, \dots, G_{ℓ} . It follows from this that if we can associate the type $\mathcal{C}[G_1 \llbracket \theta \rrbracket; \dots; G_j \llbracket \theta \rrbracket; n_1 : B'_1 \llbracket \theta^h \rrbracket, \dots, n_k : B'_k \llbracket \theta^h \rrbracket; G_{j+1} \llbracket \theta \rrbracket; \dots; G_{\ell} \llbracket \theta \rrbracket]$ with Γ , then the context expression that σ substitutes for Γ can still be generated from the changed type. The key to our showing that the requirements of the lemma are met in these cases will be to establish that $\text{Heads}(\mathcal{S}, G)$ contains a sequent and head pair such that the type of Γ is elaborated to a form from which the above type can be obtained, up to a permutation of nominal constants, by a well-behaved substitution and the head is identified as the i^{th} item in the introduced block of declarations.

Towards this end, let us consider the tuple $\langle \mathcal{S}'', h'' : A'' \rangle$ that is generated by

$$\text{AddBlock}(\mathcal{S}, \Gamma \uparrow \mathbb{N}_{\Gamma} : \mathcal{C}[G_1; \dots; G_{\ell}], \mathcal{B}, (n_1, \dots, n_k), \mathbb{N}_o, j, i),$$

where \mathbb{N}_o is the collection of nominal constants obtained by leaving out of \mathbb{N} the constants in \mathbb{N}_{Γ} and the constants that appear amongst the explicit bindings of G relative to \mathcal{S} . In this case, \mathcal{S}'' will have the form $\mathbb{N}''; \Psi''; \Xi''; \Omega'' \longrightarrow F''$ with the following properties. First, \mathbb{N}'' will be identical to $\mathbb{N} \cup \{n_1, \dots, n_k\}$. Second, Ψ'' will comprise two disjoint parts $\Psi_r^{\mathcal{S}}$ and $\Psi_r^{\mathcal{B}}$, where $\Psi_r^{\mathcal{S}}$ is a version of Ψ raised over the nominal constants in $\{n_1, \dots, n_k\}$ that are not members of \mathbb{N} with a corresponding raising substitution θ_r^{Ψ} , and $\Psi_r^{\mathcal{B}}$ is a version of $\{x_1, \dots, x_n\}$ raised over all the nominal constants in $\mathbb{N} \cup \{n_1, \dots, n_k\}$ except the ones that are assigned a type in G_{j+1}, \dots, G_{ℓ} or that appear in n_1^G, \dots, n_p^G with the corresponding raising substitution $\theta_r^{\mathcal{B}}$. Third, Ξ'' will be

$$\hat{\Xi} \cup \{\Gamma \uparrow \mathbb{N}_{\Gamma} : \mathcal{C}[G_1 \llbracket \theta_r^{\Psi} \rrbracket; \dots, G_j \llbracket \theta_r^{\Psi} \rrbracket; n_1 : B'_1 \llbracket \theta_r^{\mathcal{B}} \rrbracket, \dots, n_k : B'_k \llbracket \theta_r^{\mathcal{B}} \rrbracket; G_{j+1} \llbracket \theta_r^{\Psi} \rrbracket, \dots, G_{\ell} \llbracket \theta_r^{\Psi} \rrbracket]\}$$

where $\hat{\Xi} = (\Xi \setminus \{\Gamma \uparrow \mathbb{N}_{\Gamma} : \mathcal{C}[G_1; \dots; G_{\ell}]\}) \llbracket \theta_r^{\Psi} \rrbracket$. Finally, each formula in $\Omega'' \cup \{F''\}$ is obtained by applying the raising substitution θ_r^{Ψ} to a corresponding one in $\Omega \cup \{F\}$. Using Theorem 4.8 we

observe that, because $\text{supp}(\theta)$ is disjoint from the set \mathbb{N} , there is a (closed) raising substitution θ_r with $\text{ctx}(\theta_r) = \Psi_r^{\mathcal{S}}$ whose support is disjoint from the set $\mathbb{N} \cup \{n_1, \dots, n_k\}$ and which is such that $\Omega''[\theta_r] = \Omega[\theta]$, $F''[\theta_r] = F[\theta]$, $\hat{\Xi}[\theta_r]$ is equal to $(\Xi \setminus \{\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[G_1; \dots; G_\ell]\})[\theta]$, and, for each q , $1 \leq q \leq \ell$, $G_q[\theta_r^\Psi][\theta_r] = G_q[\theta]$. Using Theorem 4.8 again, we see that there is a (closed) substitution θ_r^h with $\text{ctx}(\theta_r^h) = \Psi_r^{\mathcal{B}}$ whose support is disjoint from $\mathbb{N} \cup \{n_1, \dots, n_k\}$ and that is such that, for $1 \leq q \leq k$, it is the case that $B'_q[\theta_r^{\mathcal{B}}][\theta_r^h] = B'_q[\theta^h]$. Based on all these observations, it is easy to see that if we let $\theta'' = \theta_r \cup \theta_r^h$, then $\langle \theta'', \emptyset \rangle$ is substitution compatible with \mathcal{S}'' and $\mathcal{S}''[\theta'']_\emptyset$ is identical to $\mathcal{S}[\theta]_\emptyset$ except for the fact that the type associated with Γ in its context variables context is $\mathcal{C}[G_1[\theta]; \dots; G_j[\theta]; n_1 : B'_1[\theta^h], \dots, n_k : B'_k[\theta^h]; G_{j+1}[\theta]; \dots; G_\ell[\theta]]$. By the earlier observation, σ is appropriate for $\mathcal{S}''[\theta'']_\emptyset$ and, in fact $\mathcal{S}''[\theta'']_\emptyset[\sigma] = \mathcal{S}[\theta]_\emptyset[\sigma]$. Noting also that $h'' : A''$ must, by the definition of *AddBlock*, be $n_i : B'_i[\theta_r^{\mathcal{B}}]$, if $\text{Heads}(\mathcal{S}, G)$ includes in it a pair obtained by this particular call to *AddBlock*, then we can pick \mathcal{S}' to be \mathcal{S}'' , h' to be h'' , A' to be A'' , θ' to be θ'' , σ' to be σ and π to be the identity permutation to satisfy the requirements of the lemma.

We are, of course, not assured that there will be a pair in $\text{Heads}(\mathcal{S}, G)$ corresponding to the use of *AddBlock* with exactly the arguments considered above. Specifically, the sequences of nominal constants that are considered for the block instance may not include n_1, \dots, n_k . However, we know that some sequence n'_1, \dots, n'_k will be considered that is identical to n_1, \dots, n_k except for constants in identical locations in the two sequences that are not drawn from \mathbb{N} . Since the constants in any sequence must be distinct, it follows easily that we can describe a permutation π' on the nominal constants that is the identity map on \mathbb{N} and that maps n'_1, \dots, n'_k to n_1, \dots, n_k . It can also be seen then that $\text{Heads}(\mathcal{S}, G)$ will include a tuple $\langle \mathcal{S}''', h''' : A''' \rangle$ such that $\pi'.\mathcal{S}''' = \mathcal{S}''$, $\pi'.h''' = h''$, and $\pi'.A''' = A''$. Picking \mathcal{S}' to be \mathcal{S}''' , h' to be h''' , A' to be A''' , θ' to be $\pi'^{-1}.\theta''$, σ' to be $\pi'^{-1}.\sigma''$, π to be π' and using Theorem 4.18, we can once again see that the requirements of the lemma are met. \square

Generating a Covering Set of Premise Sequents We have, at this stage, a way for identifying all the possible heads to consider for R in analyzing an atomic assumption formula $\{G \vdash R : P\}$ in a sequent \mathcal{S} . However, we still need a systematic approach for considering all the term and context substitutions that yield closed instances of \mathcal{S} in which the term component of F has the relevant head. We now turn to this task. Rather than identifying the closed instances immediately, we will think of taking a step in this direction that also allows us to reduce the typing judgement represented by F based on the typing rule for the LF judgement it represents. The first step in this direction will be to determine a substitution that makes the head of R identical to the one it needs to be in its closed form. We introduce a particular form of unification towards this end. The next step will be to reduce the sequent based on the observations in Theorem 2.20. The eventual proof rule will then combine the identification of relevant heads using $\text{Heads}(\mathcal{S}, G)$, the solving of a unification problem based on each such head, and the reduction of the sequent.

We begin this development by describing the relevant notion of unification.

Definition 4.40. A unification problem \mathcal{U} is a tuple $\langle \mathbb{N}, \Psi, \mathcal{E} \rangle$ in which \mathbb{N} is a collection of nominal constants, Ψ is a term variables context, and \mathcal{E} is a set of equations $\{E_1 = E'_1, \dots, E_n = E'_n\}$ where, for $1 \leq i \leq n$, either $\mathbb{N} \cup \Psi \cup \Theta_0 \vdash_{\text{ak}} E_i$ type and $\mathbb{N} \cup \Psi \cup \Theta_0 \vdash_{\text{ak}} E'_i$ type have derivations or there is an arity type α such that $\mathbb{N} \cup \Psi \cup \Theta_0 \vdash_{\text{at}} E_i : \alpha$ and $\mathbb{N} \cup \Psi \cup \Theta_0 \vdash_{\text{at}} E'_i : \alpha$ have derivations. A solution to such a unification problem is a pair $\langle \theta, \Psi' \rangle$ of a substitution and a term variables context such that

1. θ is type preserving with respect to $\mathcal{N} \cup \Theta_0 \cup \Psi'$,
2. $\text{supp}(\theta) \cap \mathbb{N} = \emptyset$,

3. for any x if $x : \alpha \in \Psi$ and $x : \alpha' \in \text{ctx}(\theta) \uplus \Psi'$ then $\alpha = \alpha'$, and
4. for each i , $1 \leq i \leq n$, expressions E_i and E'_i from $E_i = E'_i$ are such that $E_i[\theta] = E'_i[\theta]$.

Note that the typing constraints validate the use of the notation $E_i[\theta]$ and $E'_i[\theta]$.

Given an atomic term R , we can determine its instances that have a particular head h through the unification of R with h applied to a sequence of fresh variables. We will use this idea to narrow down the set of instances of a sequent that must be considered once we have determined what the head of the term in an atomic goal of the form $\{G \vdash R : P\}$ must be. However, we must first build into our notion of a fresh variable the ability to instantiate it with nominal constants appearing in the sequent. We do this below by using the mechanism of raising.

Definition 4.41. Let Ψ be a term variables context and let \mathbb{N} be a finite subset of \mathcal{N} . Further, let n_1, \dots, n_k be a listing of the nominal constants in \mathbb{N} and let $\alpha_1, \dots, \alpha_k$ be the respective types of these constants. Then, for any variable z that does not appear in Ψ , $z : \alpha_1 \rightarrow \dots \rightarrow \alpha_k \rightarrow \beta$ is said to be a variable of arity type β away from Ψ and raised over \mathbb{N} . Moreover, $(z \ n_1 \ \dots \ n_k)$ is said to be the generalized variable term corresponding to z .

The following lemma now formalizes the described refinement of the sequent.

Lemma 4.42. Let $\mathcal{S} = \mathbb{N}; \Psi; \Xi; \Omega, \{G \vdash R : P\} \longrightarrow F'$ be a well-formed sequent. Further, let θ be a term substitution that together with a context substitution σ identifies a closed instance of \mathcal{S} and is such that $R[\theta] = (h \ M_1 \ \dots \ M_n)$ and $P[\theta] = P'[\{\langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_n, M_n, (A_n)^- \rangle\}]$ for a head h that Σ or $G[\theta][\sigma]$ assigns the type $\Pi x_1:A_1. \dots \Pi x_n:A_n. P'$. Finally for $1 \leq i \leq n$ let $z_i : \alpha'_i$ be a distinct variable of type $(A_i)^-$ away from Ψ and raised over \mathbb{N} , and let t_i be the generalized variable term corresponding to z_i . Then $\langle \theta, \emptyset \rangle$ is a solution to the unification problem

$$\langle \mathbb{N}, \Psi \cup \{z_1 : \alpha'_1, \dots, z_n : \alpha'_n\}, \\ \{P = P'[\{\langle x_1, t_1, (A_1)^- \rangle, \dots, \langle x_n, t_n, (A_n)^- \rangle\}], R = (h \ t_1 \ \dots \ t_n)\} \rangle.$$

Proof. As θ and σ identify a closed instance of \mathcal{S} it must be that $\langle \theta, \emptyset \rangle$ is substitution compatible with \mathcal{S} . But then, $\langle \theta, \emptyset \rangle$ satisfies the first three clauses of the definition for a solution. The assumptions $R[\theta] = (h \ M_1 \ \dots \ M_n)$ and $P[\theta] = P'[\{\langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_n, M_n, (A_n)^- \rangle\}]$ ensure that the final clause is also satisfied, implying that $\langle \theta, \emptyset \rangle$ solves the given unification problem. \square

The reduction of a sequent based on an atomic assumption formula is the content of the next definition. Note that the type and the term in the formula are required to be atomic for this reduction to be applicable and the type of the head of the term determines the reduction.

Definition 4.43. Let $\mathcal{S} = \mathbb{N}; \Psi; \Xi; \Omega, F \longrightarrow F'$ be a well-formed sequent, where F is the formula $\{G \vdash h \ M_1 \ \dots \ M_n : P\}$ for some h that is assigned the type $A = \Pi x_1:A_1. \dots \Pi x_n:A_n. P'$ in Σ or the explicit bindings in G relative to \mathcal{S} . The sequent obtained by decomposing the assumption formula F based on the type A is $\mathbb{N}; \Psi; \Xi; \Omega, \{G \vdash M_1 : A'_1\}, \dots, \{G \vdash M_n : A'_n\} \longrightarrow F'$ where, for $1 \leq i \leq n$, A'_i is $A_i[\{\langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_{i-1}, M_{i-1}, (A_{i-1})^- \rangle\}]$. This sequent is denoted by $\text{ReduceSeq}(\mathcal{S}, F)$. Note that the well-formedness of \mathcal{S} justifies the use of the notation $A_i[\{\langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_{i-1}, M_{i-1}, (A_{i-1})^- \rangle\}]$.

The following lemma expresses the soundness of the idea of reducing a sequent. Additionally, it identifies a measure with atomic formulas that diminishes with the replacements effected by a reduction step; this property will be useful in showing soundness for an induction rule that we present in Section 4.4.

Lemma 4.44. *Let $\mathcal{S} = \mathbb{N}; \Psi; \Xi; \Omega, F \longrightarrow F'$ be a well-formed sequent with F an atomic formula of the form $\{G \vdash R : P\}$. Further, let h be assigned the type $\prod x_1:A_1. \dots \prod x_n:A_n. P'$ in either Σ or the explicit bindings in G relative to \mathcal{S} and for each i , $1 \leq i \leq n$, let $z_i : \alpha'_i$ be a distinct variable of type $(A_i)^-$ away from Ψ and raised over \mathbb{N} , and let t_i be the generalized variable term corresponding to z_i . Finally, let \mathcal{U} be the unification problem*

$$\langle \mathbb{N}, \Psi \cup \{z_1 : \alpha'_1, \dots, z_n : \alpha'_n\}, \\ \{P = P'[\{\langle x_1, t_1, (A_1)^- \rangle, \dots, \langle x_n, t_n, (A_n)^- \rangle\}], R = (h \ t_1 \ \dots \ t_n)\} \rangle.$$

Then any solution to \mathcal{U} is substitution compatible with \mathcal{S} . Further, for any $\langle \theta, \Psi_\theta \rangle$ that is a solution to \mathcal{U} and θ_r that is a raising substitution associated with the application of θ to \mathcal{S} relative to Ψ_θ , there must be terms M_1, \dots, M_n such that the following hold:

1. $R[\theta][\theta_r]$ is $(h \ M_1 \ \dots \ M_n)$.
2. For any θ' and σ' identifying a closed instance of $\mathcal{S}[\theta]_{\Psi_\theta}$, if $\{G \vdash R : P\}[\theta][\theta_r][\theta'][\sigma']$ is valid and there is a derivation for $(G \vdash_\Sigma R \Leftarrow P)[\theta][\theta_r][\theta'][\sigma']$ of height k , then for each i , $1 \leq i \leq n$, letting $A'_i = A_i[\{\langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_{i-1}, M_{i-1}, (A_{i-1})^- \rangle\}]$, it must be the case that $(\{G[\theta][\theta_r] \vdash M_i : A'_i\})[\theta'][\sigma']$ is valid and that there is a derivation of height less than k for $(G[\theta][\theta_r] \vdash_\Sigma M_i \Leftarrow A'_i)[\theta'][\sigma']$.
3. There is an \mathcal{S}' such that $\mathcal{S}' = \text{ReduceSeq}(\mathcal{S}[\theta]_{\Psi_\theta}, F[\theta][\theta_r])$ and \mathcal{S}' is valid only if $\mathcal{S}[\theta]_{\Psi_\theta}$ is.

Proof. A straightforward examination of Definitions 4.6 and 4.40 suffices to verify that solutions to \mathcal{U} must be substitution compatible with \mathcal{S} . Any solution $\langle \theta, \Psi_\theta \rangle$ to the unification problem \mathcal{U} must be such that $R[\theta] = (h \ t_1 \ \dots \ t_n)[\theta]$. From this it follows that $R[\theta][\theta_r] = (h \ t_1 \ \dots \ t_n)[\theta][\theta_r]$. Since h is unaffected by substitutions, it is easy to see that $(h \ t_1 \ \dots \ t_n)[\theta][\theta_r] = (h \ (t_1[\theta][\theta_r]) \ \dots \ (t_n[\theta][\theta_r]))$. Picking M_i to be the term $t_i[\theta][\theta_r]$ for each i , $1 \leq i \leq n$, we see that clause (1) in the lemma is satisfied.

For the second clause we note first that the typing judgements in question must be closed and hence the consideration is meaningful. Consider an arbitrary closed instance of $\mathcal{S}[\theta]_{\Psi_\theta}$ identified by θ' and σ . If $F[\theta][\theta_r][\theta'][\sigma]$ is a valid formula then using the definition of validity as well as clause (1) in the lemma we can extract a derivation for $\vdash_\Sigma G[\theta][\theta_r][\theta'][\sigma]$ **ctx** and a derivation of height k for $G[\theta][\theta_r][\theta'][\sigma] \vdash_\Sigma (h \ M_1 \ \dots \ M_n)[\theta'] \Leftarrow P[\theta][\theta_r][\theta']$. Application of Theorems 2.20 and 2.6 are then sufficient to conclude that there is a derivation of height less than k for $(G[\theta][\theta_r] \vdash_\Sigma M_i \Leftarrow A'_i)[\theta'][\sigma]$. For us to be able to conclude that, for each i , $1 \leq i \leq n$, the formula $(\{G[\theta][\theta_r] \vdash M_i : A'_i\})[\theta'][\sigma]$ is valid, it only remains to show that there is a derivation for $G[\theta][\theta_r][\theta'][\sigma] \vdash_\Sigma A'_i[\theta']$ **type**. However, this has been done in the proof of Theorem 2.20.

We finish by proving the third clause. From clause (1) we know that $R[\theta][\theta_r]$ will be of the form $(h \ M_1 \ \dots \ M_n)$, thus by the definition of *ReduceSeq* there must exist an \mathcal{S}' such that $\mathcal{S}' = \text{ReduceSeq}(\mathcal{S}[\theta]_{\Psi_\theta}, F[\theta][\theta_r])$. Suppose \mathcal{S}' is valid. Consider an arbitrary closed instance of $\mathcal{S}[\theta]_{\Psi_\theta}$ identified by θ' and σ . These same θ' and σ also identify a closed instance of \mathcal{S}' as these sequents only differ in that the assumption formula $F[\theta][\theta_r]$ has been replaced with the collection of reduced formulas $\{\{G[\theta][\theta_r] \vdash M_i : A'_i\} \mid 1 \leq i \leq n\}$. If any formula in the set of assumption formulas of $\mathcal{S}[\theta]_{\Psi_\theta}[\theta'][\sigma]$ were not valid then this instance would be vacuously valid, so suppose all such formulas are valid. Then in particular, $F[\theta][\theta_r][\theta'][\sigma]$ must be valid and thus by clause (2), for each i , $1 \leq i \leq n$, the formula $\{G[\theta][\theta_r] \vdash M_i : A'_i\}[\theta'][\sigma]$ will be valid. But then all of the assumption formulas of $\mathcal{S}'[\theta'][\sigma]$ must be valid and since this is a closed instance of a valid sequent we can conclude that the goal formula $F'[\theta][\theta_r][\theta'][\sigma]$ is valid. Therefore any closed instance of $\mathcal{S}[\theta]_{\Psi_\theta}$ will be valid, and clause (3) in the lemma is satisfied. \square

Lemmas 4.42 and 4.44 yield the following possibility for analyzing the derivability of a sequent with $\{G \vdash R : P\}$ as an atomic assumption formula relative to an identified head for R : use the unification problem identified in Lemma 4.42 to generate a collection of term substitutions and analyze the derivability of the reduced sequent under these substitutions. Unfortunately, this would not be a very effective strategy if we have to treat each unifier separately. Towards dealing with this issue, we introduce the idea of a covering set of solutions to a unification problem. The next three definitions culminate in a formulation of this notion.

Definition 4.45. *The restriction of a substitution θ to the term variables context Ψ is the substitution $\{\langle x, M, \alpha \rangle \mid \langle x, M, \alpha \rangle \in \theta \text{ and } x : \alpha' \in \Psi\}$. This substitution is denoted by $\theta|_{\Psi}$.*

Definition 4.46. *Let Ψ , Ψ_1 and Ψ_2 be term variables contexts, and let θ_1 and θ_2 be substitutions that are arity type preserving with respect to $\mathcal{N} \cup \Theta_0 \cup \Psi_1$ and $\mathcal{N} \cup \Theta_0 \cup \Psi_2$, respectively. Then $\langle \theta_2, \Psi_2 \rangle$ is said to cover $\langle \theta_1, \Psi_1 \rangle$ relative to Ψ if there exists a pair $\langle \theta_3, \Psi_3 \rangle$ of a substitution and a term variables context such that*

1. θ_3 is type preserving with respect to $\mathcal{N} \cup \Theta_0 \cup \Psi_3$,
2. for any $x : \alpha \in \Psi_2$, if $x : \alpha' \in \text{ctx}(\theta_3) \uplus \Psi_3$ then $\alpha = \alpha'$, and
3. The substitutions $\theta_1|_{\Psi}$ and $(\theta_3 \circ \theta_2)|_{\Psi}$ are identical.

Note that the second condition ensures that $\mathcal{N} \cup \Theta_0 \cup ((\Psi_2 \setminus \text{ctx}(\theta_3))) \cup \Psi_3$ determines a valid arity context and that θ_2 and θ_3 are arity type compatible with respect to this context. Thus, the composition of θ_2 and θ_3 in the third condition is well-defined.

Definition 4.47. *A collection S of solutions to a unification problem $\mathcal{U} = \langle \mathbb{N}, \Psi, \mathcal{E} \rangle$ is said to be covering set of solutions for \mathcal{U} if every solution to \mathcal{U} is covered by some solution in S relative to Ψ .*

The following lemma provides the basis for using the reduced forms generated by just a covering set of solutions for the relevant unification problem in analyzing the derivability of the sequent.

Lemma 4.48. *Let S be the well-formed sequent $\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F'$ and let θ_1 and σ identify a closed instance S' of S . Further, let $\langle \theta_2, \Psi_2 \rangle$ be substitution compatible with S and such that it covers $\langle \theta_1, \emptyset \rangle$ relative to Ψ . Then there is a term substitution θ that together with σ identifies a closed instance of $S[[\theta_2]]_{\Psi_2}$ that is valid if and only if S' is.*

Proof. We argue below that, under the assumptions of the lemma, there is a substitution θ_3 and a term variables context Ψ_3 such that $\langle \theta_3, \Psi_3 \rangle$ is substitution compatible with $S[[\theta_2]]_{\Psi_2}$ and for any term M such that $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{\text{at}} M : \alpha$ is derivable it is the case that $M[[\theta_2]][[\theta_{2r}]] [[\theta_3]] = M[[\theta_1]]$, where θ_{2r} is the raising substitution associated with the application of θ_2 to S relative to Ψ_2 . It follows from this that the formulas and context types appearing in $S[[\theta_2]]_{\Psi_2} [[\theta_3]]_{\Psi_3}$ must be identical to the ones in $S[[\theta_1]]_{\emptyset}$. It is then easily seen that θ_3 can be extended into a substitution θ that together with σ identifies a closed instance of $S[[\theta_2]]_{\Psi_2}$ whose formulas are identical to those of S' . The lemma is an immediate consequence.

Since $\langle \theta_2, \Psi_2 \rangle$ covers $\langle \theta_1, \emptyset \rangle$ relative to Ψ we know that there exists a pair $\langle \theta'_3, \Psi_3 \rangle$ satisfying the conditions of Definition 4.46. We claim that we may further assume of θ'_3 that (1) $\text{ctx}(\theta'_3) = (\Psi \setminus \text{ctx}(\theta_2)) \cup \Psi_2$, and (2) $\text{supp}(\theta'_3) \cap \mathbb{N} = \emptyset$. Condition (1) may initially be violated because there are tuples in θ'_3 for variables that are not assigned types by $(\Psi \setminus \text{ctx}(\theta_2)) \cup \Psi_2$ or because $\text{ctx}(\theta'_3)$ does not span all of Ψ_2 . The first is easily fixed by dropping such tuples and the second by adding tuples of the form $\langle x, x, \alpha \rangle$ to θ'_3 and, if needed, the type assignment $x : \alpha$ to Ψ_3 . As for the second condition,

suppose that it is violated at first. Consider then a permutation π of nominal constants that swaps the constants in $\text{supp}(\theta'_3) \cap \mathbb{N}$ with ones that are not contained in $\text{supp}(\theta_1) \cup \text{supp}(\theta_2) \cup \mathbb{N}$ and is the identity map on all other constants. By the choice of π and the fact that $\langle \theta_1, \emptyset \rangle$ and $\langle \theta_2, \Psi_2 \rangle$ are both substitution compatible with \mathcal{S} , it is the case that $\pi.M$ is identical to M for any M such that $\langle x, M, \alpha \rangle$ is a member of θ_1 or θ_2 . Now, we could replace θ'_3 with $\pi.\theta'_3$ if the latter substitution also satisfies the conditions needed of it by Definition 4.46. It is easy to check that $\pi.\theta'_3$ satisfies the first two of these conditions. From the fact that $\theta'_3|_{\Psi \setminus \text{ctx}(\theta_2)} = \theta_1|_{\Psi \setminus \text{ctx}(\theta_2)}$, it follows that the tuples in $\theta'_3|_{\Psi \setminus \text{ctx}(\theta_2)}$ will be unaffected by the permutation and hence $\pi.\theta'_3|_{\Psi \setminus \text{ctx}(\theta_2)} = \theta_1|_{\Psi \setminus \text{ctx}(\theta_2)}$. Thus, to conclude that $\pi.\theta'_3 \circ \theta_2|_{\Psi} = \theta_1|_{\Psi}$, it suffices to show that for any tuple of the form $\langle x, M, \alpha \rangle$ that belongs to θ_2 , if it is the case that $\langle x, M[\theta'_3], \alpha \rangle$ belongs to θ_1 , then $M[\theta'_3] = M[\pi.\theta'_3]$. A simple inductive argument establishes the fact that $\pi.M[\theta'_3] = (\pi.M)[\pi.\theta'_3]$. The desired conclusion follows by utilizing the observation that $M[\theta'_3]$ and M are unaffected by the permutation.

We now use Theorem 4.8 to obtain a “raised” version of θ'_3 that together with Ψ_3 will constitute the pair $\langle \theta_3, \Psi_3 \rangle$ that we desired at the outset. Specifically, the theorem allows us to conclude that there is a substitution θ_3 satisfying the following properties:

1. $\text{supp}(\theta_3)$ is disjoint from $\mathbb{N} \cup \text{supp}(\theta_2)$,
2. $\text{ctx}(\theta_3)$ is identical to the raised version of $(\Psi \setminus \text{ctx}(\theta_2)) \cup \Psi_2$ corresponding to the raising substitution θ_{2r} ,
3. θ_3 is arity type preserving with respect to $\mathcal{N} \cup \Theta \cup \Psi_3$, and
4. for every term M such that $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{at} M : \alpha$, $M[\theta_2][\theta'_3] = M[\theta_2][\theta_{2r}][\theta_3]$.

The argument for the first three of these properties is obvious. For the last property, we observe, using Theorem 2.8, that $(\mathbb{N} \cup \text{supp}(\theta_2)) \cup \Theta_0 \cup ((\Psi \setminus \text{ctx}(\theta_2)) \cup \Psi_2) \vdash_{at} M[\theta_2] : \alpha$ has a derivation under the condition described; Theorem 4.8 can then be invoked in an obvious way. It follows immediately from the first three properties that $\langle \theta_3, \Psi_3 \rangle$ is substitution compatible with $\mathcal{S}[\theta_2]_{\Psi_2}$. It therefore only remains to show that for every M such that $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{at} M : \alpha$, it is the case that $M[\theta_1] = M[\theta_2][\theta_{2r}][\theta_3]$. An easy inductive argument shows that for any M of the kind described and any θ , if $M[\theta] = M'$ is derivable exactly when $M[\theta|_{\Psi}] = M'$ is derivable. It follows from this that $M[\theta_1] = M[\theta_2][\theta'_3]$. Property (4) then yields the desired result. \square

We now use the observations in Lemmas 4.39, 4.42, 4.44 and 4.48 to describe a complete analysis of the derivability of a sequent around an atomic assumption formula.

Definition 4.49. *Let \mathcal{S} be the well-formed sequent $\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F'$, let $F = \{G \vdash R : P\}$ be a formula in Ω and let $h : \prod x_1 : A_1. \dots \prod x_n : A_n. P'$ be a type assignment in Σ or in the explicit bindings in G . Further, for $1 \leq i \leq n$, let $z_i : \alpha_i$ be a distinct variable of type $(A_i)^-$ away from Ψ and raised over \mathbb{N} , and let t_i be the generalized variable term corresponding to z_i . Finally, let \mathcal{U} be the unification problem*

$$\langle \mathbb{N}, \Psi \cup \{z_1 : \alpha_1, \dots, z_n : \alpha_n\}, \\ \{P = P'[\{\langle x_1, t_1, (A_1)^-\rangle, \dots, \langle x_n, t_n, (A_n)^-\rangle\}], R = (h \ t_1 \ \dots \ h_n)\} \rangle$$

and let C be a covering set of solutions for \mathcal{U} . Then the analysis of \mathcal{S} based on F and h is denoted by $\text{Cases}(\mathcal{S}, F, h : A)$ and is given by the set of sequents

$$\{\text{ReduceSeq}(\mathcal{S}[\theta]_{\Psi_\theta}, F') \mid \langle \theta, \Psi_\theta \rangle \in C \text{ and } F' \text{ is the formula in } \mathcal{S}[\theta]_{\Psi_\theta} \text{ resulting from } F\}.$$

If \mathcal{S} is a well-formed sequent and $F = \{G \vdash R : P\}$ is an assumption formula in \mathcal{S} , then the complete analysis of \mathcal{S} based on F is the set of sequents

$$\bigcup \{ \text{Cases}(\mathcal{S}', F', h : A) \mid (\mathcal{S}'; h : A) \in \text{Heads}(\mathcal{S}, G) \\ \text{and } F' \text{ is the formula in } \mathcal{S}' \text{ resulting from } F \}.$$

This collection is denoted by $\text{AllCases}(\mathcal{S}, F)$. Note that the notations $\text{Cases}(\mathcal{S}, F, h : A)$ and $\text{AllCases}(\mathcal{S}, F)$ are both ambiguous—for instance, the first notation leaves out mention of the covering set of solutions that plays a role in generating the set it denotes. We will assume them to denote any of the set of sequents that can be generated in the respective ways described in this definition.

The following lemmas will be useful in showing that the proof rule that we will present using AllCases is well-defined and sound.

Lemma 4.50. *If \mathcal{S} is a well-formed sequent and F is an atomic assumption formula of the form $\{G \vdash R : P\}$ that appears in \mathcal{S} , then every sequent in $\text{AllCases}(\mathcal{S}, F)$ is well-formed.*

Proof. By Lemma 4.38 we know that every $\langle \mathcal{S}', h : A \rangle \in \text{Heads}(\mathcal{S}, G)$ is such that \mathcal{S}' is a well-formed sequent. Further, if \mathcal{S}' is the sequent $\mathbb{N}'; \Psi'; \Xi'; \Omega' \longrightarrow F'$, then $\mathbb{N}' \cup \Theta_0 \cup \Psi' \vdash_{ak} A$ type must have a derivation. Let F'' be the formula from \mathcal{S}' corresponding to F and let us denote the unification problem that is considered by $\text{Cases}(\mathcal{S}', F', h : A)$ by \mathcal{U} . Then, by Theorem 4.10, the application of any solution for \mathcal{U} to \mathcal{S}' must be well-formed. Now, if $\{G' \vdash h M_1 \dots M_n : P'\}$ is a formula that is well-formed with respect to $\mathbb{N} \cup \Theta_0 \cup \Psi$ and Ξ^- where $h : \prod x_1 : A_1. \dots \prod x_n : A_n. P''$ appears in Σ or the explicit bindings of G' , then it is easily argued that, for $1 \leq i \leq n$,

$$\{G \vdash M_i : A_i \llbracket \{ \langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_{i-1}, M_{i-1}, (A_{i-1})^- \rangle \} \rrbracket\}$$

is also a well-formed formula with respect to $\mathbb{N} \cup \Theta_0 \cup \Psi$ and Ξ^- . It follows from this that each sequent in $\text{Cases}(\mathcal{S}', F', h : A)$ must be well-formed. Since this is true regardless of the pair $\langle \mathcal{S}', h : A \rangle$ that is considered, it must be the case that all the sequents in $\text{AllCases}(\mathcal{S}, F)$ are well-formed. \square

Lemma 4.51. *Let \mathcal{S} be a well-formed sequent and let F be an atomic assumption formula of the form $\{G \vdash R : P\}$ in \mathcal{S} . If all the sequents in $\text{AllCases}(\mathcal{S}, F)$ are valid then \mathcal{S} must be valid.*

Proof. Consider an arbitrary closed instance of the sequent \mathcal{S} identified by θ and σ . By Lemma 4.39 and Theorem 4.19, there is a pair $\langle \mathcal{S}', h' : A' \rangle$ in $\text{Heads}(\mathcal{S}, G)$ that has in it an atomic assumption formula F' of the form $\{G' \vdash R' : P'\}$ and that is such that if every closed instance of \mathcal{S}' identified by substitutions θ' and σ' in which the head of $R' \llbracket \theta' \rrbracket$ is identical to h' is valid then $\mathcal{S} \llbracket \theta \rrbracket_{\theta} \llbracket \sigma \rrbracket$ must be valid. By Lemmas 4.42, 4.44, and 4.48, there is a sequent \mathcal{S}'' in $\text{Cases}(\mathcal{S}', F', h' : A')$ that is such that if \mathcal{S}'' is valid then the mentioned instance of \mathcal{S}' must be valid. Finally, since $\text{AllCases}(\mathcal{S}, F)$ collects all these sequents for each pair in $\text{Heads}(\mathcal{S}, G)$, it follows that the validity of the sequents in it ensures the validity of every closed instance of \mathcal{S} and, hence, of \mathcal{S} . \square

4.3.2 Proof Rules that Introduce Atomic Formulas

Figure 4.5 presents rules for introducing atomic formulas that internalize their understanding as encodings of typing derivations in LF. The rule for introducing an atomic goal formula with an atomic type has a proviso on the context expression. This proviso ensures that the corresponding LF context is well-formed if the assumption formulas of the sequent instance are valid. This rule can be complemented by others, such as ones that check (closed) contexts for wellformedness or that check closed goal formulas for LF derivability. We do not discuss these alternatives explicitly

here, but their soundness should be easy to verify. The treatment of an atomic assumption and goal formula with a type corresponding to an abstraction is identical: a nominal constant of the right type that does not appear in the support set of the sequent is picked and used for the abstracted variable as per the LF rule for typing abstractions, changing the association with any relevant context variable to ensure that only those of its instantiations are considered that do not use the chosen nominal constant.

$$\begin{array}{c}
\frac{AllCases(\mathbb{N}; \Psi; \Xi; \Omega, \{G \vdash R : P\} \longrightarrow F, \{G \vdash R : P\})}{\mathbb{N}; \Psi; \Xi; \Omega, \{G \vdash R : P\} \longrightarrow F} \text{ atm-app-L} \\
\\
\begin{array}{c}
h : \Pi x_1 : A_1. \dots \Pi x_n : A_n. P \in \Sigma \text{ or the explicit bindings in } G \\
G \text{ appears as the context of some atomic formula in } \Omega \\
P[\{\langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_n, M_n, (A_n)^- \rangle\}] = P' \\
\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \\
\{G \vdash M_i : A_i[\{\langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_{i-1}, M_{i-1}, (A_{i-1})^- \rangle\}] \mid 1 \leq i \leq n\} \\
\hline
\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \{G \vdash h M_1 \dots M_n : P'\} \text{ atm-app-R}
\end{array} \\
\\
\begin{array}{c}
n \text{ is new to } \mathbb{N} \text{ and has type } (A_1)^- \\
\Xi' = \begin{cases} (\Xi \setminus \{\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[\mathcal{G}]\}) \cup \{\Gamma \uparrow (\mathbb{N}_\Gamma, n : (A_1)^-) : \mathcal{C}[\mathcal{G}]\} & \text{if } \Gamma \text{ appears in } G \\ \Xi & \text{otherwise} \end{cases} \\
\mathbb{N}, n : (A_1)^-; \Psi; \Xi'; \Omega, \{G, n : A_1 \vdash M[\{\langle x, n, (A_1)^- \rangle\}] : A_2[\{\langle x, n, (A_1)^- \rangle\}]\} \longrightarrow F \\
\hline
\mathbb{N}; \Psi; \Xi; \Omega, \{G \vdash \lambda x. M : \Pi x : A_1. A_2\} \longrightarrow F \text{ atm-abs-L}
\end{array} \\
\\
\begin{array}{c}
n \text{ is new to } \mathbb{N} \text{ and has type } (A_1)^- \\
\Xi' = \begin{cases} (\Xi \setminus \{\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[\mathcal{G}]\}) \cup \{\Gamma \uparrow (\mathbb{N}_\Gamma, n : (A_1)^-) : \mathcal{C}[\mathcal{G}]\} & \text{if } \Gamma \text{ appears in } G \\ \Xi & \text{otherwise} \end{cases} \\
\mathbb{N}, n : (A_1)^-; \Psi; \Xi'; \Omega \longrightarrow \{G, n : A_1 \vdash M[\{\langle x, n, (A_1)^- \rangle\}] : A_2[\{\langle x, n, (A_1)^- \rangle\}]\} \\
\hline
\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \{G \vdash \lambda x. M : \Pi x : A_1. A_2\} \text{ atm-abs-R}
\end{array}
\end{array}$$

Figure 4.5: Rules that Introduce Atomic Formulas

As usual, we show that these rules preserve the wellformedness of sequents and are also sound.

Theorem 4.52. *The following property holds for each rule in Figure 4.5: if the conclusion sequent is well-formed, the premises expressing typing conditions have derivations and the conditions expressed by the other, non-sequent premises are satisfied, then all the sequent premises must be well-formed.*

Proof. Lemma 4.50 ensures that the property holds for the *atm-app-L* rule. For the *atm-app-R* rule, we note first that the wellformedness of the conclusion sequent immediately assures us that the context variables contexts and the assumption formulas of each of the n premise sequents satisfy the conditions required of them for the wellformedness of the corresponding sequent. This leaves us needing to show only that

$$\mathbb{N} \cup \Theta_0 \cup \Psi; \Xi \vdash \{G \vdash M_i : A_i[\{\langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_{i-1}, M_{i-1}, (A_{i-1})^- \rangle\}]\} \text{ fmla}$$

has a derivation for $1 \leq i \leq n$. Without loss of generality, we may assume that the variables x_1, \dots, x_n are chosen to be distinct and also different from the ones in Ψ . For $1 \leq i \leq (n+1)$,

let θ_i denote the term substitution $\{\langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_{i-1}, M_{i-1}, (A_{i-1})^- \rangle\}$. From the wellformedness of the conclusion sequent and the premise conditions for the rule we easily see that there must be derivations for $\mathbb{N} \cup \Theta_0 \cup \Psi \cup \text{ctx}(\theta_i) \vdash_{\text{ak}} A_i$ type for $1 \leq i \leq n$ and $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{\text{at}} h M_1 \dots M_n : (P[\theta_{n+1}])^-$, where $h : (\Pi x_1:A_1. \dots \Pi x_n:A_n. P)^-$ is in $\mathbb{N} \cup \Theta_0 \cup \Psi$. Using an induction on i and invoking Theorems 2.11 and 3.2, we can show from this that, for $1 \leq i \leq n$, it must be the case that θ_i is type preserving with respect to $\mathbb{N} \cup \Theta_0 \cup \Psi$, that $A_i[\theta_i]$ is well-defined, and that $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{\text{ak}} A_i[\theta_i]$ type and $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{\text{at}} M_i : (A_i[\theta_i])^-$ have derivations. The desired conclusion is an easy consequence of this.

We consider the two remaining cases, for *atm-abs-L* and *atm-abs-R*, simultaneously. Using the newness of n and the wellformedness of the conclusion sequent, it is easily shown that the context variables context, all the formulas in Ω and, in the case of *atm-abs-L*, the formula F all satisfy the conditions required of them for the premise sequent to be well-formed. This leaves it only to be shown that

$$(\mathbb{N} \cup \{n : (A_1)^-\}) \cup \Theta_0 \cup \Psi; \Xi'^- \vdash \{G, n : A_1 \vdash M[\{\langle x, n, (A_1)^- \rangle\}] : A_2[\{\langle x, n, (A_1)^- \rangle\}]\} \text{ fmla}$$

has a derivation. Noting that $\Xi'^- = \Xi^-$ and the wellformedness of the conclusion sequent, we can easily see that $(\mathbb{N} \cup \{n : (A_1)^-\}) \cup \Theta_0 \cup \Psi; \Xi'^- \vdash G$ context must have a derivation. What remains to be shown, then, is that $(\mathbb{N} \cup \{n : (A_1)^-\}) \cup \Theta_0 \cup \Psi \vdash_{\text{ak}} A_2[\{\langle x, n, (A_1)^- \rangle\}]$ type and $(\mathbb{N} \cup \{n : (A_1)^-\}) \cup \Theta_0 \cup \Psi \vdash_{\text{at}} M[\{\langle x, n, (A_1)^- \rangle\}] : (A_2[\{\langle x, n, (A_1)^- \rangle\}])^-$ have derivations. However, this is also easily argued for using the facts that, by the wellformedness of the conclusion sequent, $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{\text{ak}} \Pi x:A_1. A_2$ type and $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{\text{at}} \lambda x. M : (\Pi x:A_1. A_2)^-$ must have derivations and that $\{\langle x, n, (A_1)^- \rangle\}$ is an arity type preserving substitution; this argument would invoke Theorems 2.11 and 3.2 at relevant points. \square

Theorem 4.53. *The following property holds for every instance of each of the rules in Figure 4.5: if the premises expressing typing judgements are derivable, the conditions described in the other non-sequent premises are satisfied and the premise sequents are valid, then the conclusion sequent must also be valid.*

Proof. Lemma 4.51 verifies that the property holds for the *atm-app-L* rule. Consider a closed instance of the conclusion sequent of the *atm-app-R* rule that is identified by the substitutions θ and σ . If any formula in $\Omega[\theta][\sigma]$ is not valid, then this instance would be valid. We may therefore assume that all these formulas are valid and our task is then to show that the formula $\{G \vdash h M_1 \dots M_n : P'\}[\theta][\sigma]$ must also be valid under the assumptions of the theorem. Since G appears as the context in one of the formulas in Ω , $\vdash_{\Sigma} G[\theta][\sigma]$ ctx must have a derivation. Clearly, the substitutions θ and σ also identify closed instances of the premise sequents all of whose assumption formulas are valid. From the validity of these instances, the definition of substitution application and Theorem 2.6, it follows that, for $1 \leq i \leq n$, the formulas $\{G[\theta][\sigma] \vdash M_i[\theta] : A_i[\theta][\{\langle x_1, M_1[\theta], (A_1)^- \rangle, \dots, \langle x_{i-1}, M_{i-1}[\theta], (A_{i-1})^- \rangle\}]\}$ must be valid. Their validity means that, for $1 \leq i \leq n$, the LF judgements

$$G[\theta][\sigma] \vdash_{\Sigma} M_i[\theta] \Leftarrow A_i[\theta][\{\langle x_1, M_1[\theta], (A_1)^- \rangle, \dots, \langle x_{i-1}, M_{i-1}[\theta], (A_{i-1})^- \rangle\}]$$

must have derivations; these judgements are coherent because the validity of the atomic formula also assures us of the wellformedness of the corresponding types. From the assumption concerning h and the definition of substitution application, it follows that $h : \Pi x_1:A_1[\theta]. \dots \Pi x_n:A_n[\theta]. P[\theta]$ is a member of Σ or $G[\theta][\sigma]$. We may now invoke Theorem 2.20, part (1), to conclude that there must be derivations for $G[\theta][\sigma] \vdash_{\Sigma} P[\theta][\{\langle x_1, M_1[\theta], (A_1)^- \rangle, \dots, \langle x_n, M_n[\theta], (A_n)^- \rangle\}]$ type and

$$G[\theta][\sigma] \vdash_{\Sigma} h M_1[\theta] \dots M_n[\theta] \Leftarrow A_i[\theta][\{\langle x_1, M_1[\theta], (A_1)^- \rangle, \dots, \langle x_{i-1}, M_{i-1}[\theta], (A_{i-1})^- \rangle\}];$$

the wellformedness of the type is established in the proof. Using Theorem 2.6, we see that the type $P'[\theta][\{\langle x_1, M_1[\theta], (A_1)^- \rangle, \dots, \langle x_n, M_n[\theta], (A_n)^- \rangle\}]$ is identical to $P'[\theta]$. By the definition of substitution ($h M_1[\theta] \dots M_n[\theta]$) is the same term as $(h M_1 \dots M_n)[\theta]$. Thus, there are derivations for all three judgements determining the validity of $\{G \vdash h M_1 \dots M_n : P'\}[\theta][\sigma]$, implying that it must be valid.

We consider next the case for the *atm-abs-L* rule. Let θ and σ identify a closed instance of the conclusion sequent. We must show that this instance is valid provided the assumptions of the theorem are satisfied. We may assume without loss of generality that $\text{supp}(\theta)$ and $\text{supp}(\sigma)$ does not include the nominal constant n . If this condition is initially violated, we may consider instead variants of these substitutions under a permutation that swaps n with a nominal constant that does not appear in $\text{supp}(\theta) \cup \text{supp}(\sigma) \cup \mathbb{N}$, show the validity of the closed instance under these variants and then invoke Theorem 4.19. If any formula in $(\Omega, \{G \vdash \lambda x. M : \Pi x:A_1. A_2\})[\theta][\sigma]$ is not valid then the conclusion sequent must be valid. Thus it only remains for us to show the validity of $F[\theta][\sigma]$ when all these formulas are valid. From the assumption about θ and σ , it is easy to see that they must also determine a closed instance of the premise sequent. Using the definition of validity and the assumption of newness for n , it is easy to see that the validity of $(\lambda x. M : \Pi x:A_1. A_2)[\theta][\sigma]$ entails that there must be derivations for all three judgements that determine the validity of $(\{G, n : A_1 \vdash M[\{\langle x, n, (A_1)^- \rangle\}] : A_2[\{\langle x, n, (A_1)^- \rangle\}]\})[\theta][\sigma]$. From this it follows that all the assumption formulas in the instance of the premise sequent under consideration must be valid. From the assumed validity of this sequent, it then follows that $F[\theta][\sigma]$ must be valid, as we needed to show.

The only case left to be considered is that of the *atm-abs-R* rule. Let θ and σ identify a closed instance of the conclusion sequent. Our task is to show that the instance they identify is valid under the assumptions of the theorem. As before, we may assume that $\text{supp}(\theta)$ and $\text{supp}(\sigma)$ do not contain n . These substitutions must then identify a closed instance of the premise sequent. The conclusion sequent would be trivially valid if any formula in $\Omega[\theta][\sigma]$ is not valid. Thus, we only need to show that $\lambda x. M : \Pi x:A_1. A_2[\theta][\sigma]$ when all these formulas are valid. Noting that the assumption formulas in the relevant instance of the premise sequent are exactly $\Omega[\theta][\sigma]$, its validity implies the validity of $\{G, n : A_1 \vdash M[\{\langle x, n, (A_1)^- \rangle\}] : A_2[\{\langle x, n, (A_1)^- \rangle\}]\}[\theta][\sigma]$. Using the definitions of validity and LF derivability, it is easy to see that the validity of this formula implies the derivability of all three judgements that determine the validity of $\Omega[\theta][\sigma]$. Thus, this formula must be valid, thereby concluding the proof. \square

4.4 Proof Rules for Induction on the Heights of LF Derivations

The idea we use to build in a means for reasoning on the heights of LF derivations is borrowed from the Abella proof assistant [2, 3]. This idea is based on an annotation scheme that serves to determine when an atomic formula represents a typing derivation in LF that has a height less than that of the corresponding LF derivation represented by an atomic formula that appears in a formula being proved and, hence, when a property in which this atomic formula appears negatively can be assumed to hold in an inductive argument. The first part of this subsection describes these annotations and defines a refined semantics for sequents that pays attention to their intended meaning. The second part introduces an induction rule which uses annotations. The last part presents auxiliary forms of the rules for atomic formulas as well as the *id* rule that are applicable to annotated formulas; these forms are needed to make effective use of the induction rule.

4.4.1 Extending Formula Syntax with Annotations

We use a denumerable collection of annotations that go in pairs: @ and *, @@ and **, and so on. We will write @ⁿ (resp. *ⁿ) to denote a sequence in which the character @ (resp. *) is repeated n times. We use $F^{\text{@}^i}$ on an atomic formula F to indicate that it has a certain height and F^{*^i} to indicate that it has a strictly smaller height; we will explain what a height means shortly. This height annotation is decreased whenever we decompose a derivation into sub-derivations based on its structure, as is done in the *atm-app-L* rule of the previous section.

To understand the meaning of the annotations recall that an atomic formula $\{G \vdash M : A\}$, is valid if then there are LF derivations for $\vdash_{\Sigma} G$ ctx, $G \vdash_{\Sigma} A$ type and $G \vdash_{\Sigma} M \Leftarrow A$. It is the height of the typing judgement $G \vdash_{\Sigma} M \Leftarrow A$ that is the basis for our scheme for inductive reasoning. Thus, when we talk of the height of an atomic formula, we mean the height of the derivation of this typing judgement. In particular, the valid closed instances of the annotated atomic formula $\{G \vdash M : A\}^{\text{@}^i}$ are the ones for which the corresponding instances of $G \vdash_{\Sigma} M \Leftarrow A$ have derivations of height up to some particular size m , while the closed instances of the relatedly annotated formula $\{G' \vdash M' : A'\}^{*^i}$ will be valid only if the corresponding instances of $G' \vdash_{\Sigma} M' \Leftarrow A'$ have derivations of a height strictly smaller than m . Having available a denumerable collection of pairs of such annotations allows us to simultaneously relate the heights of different pairs of atomic formulas in this manner. We may of course also want to consider atomic formulas without any annotations. We use the notation $\{G \vdash M : A\}^{\text{Ann}}$ to denote a formula which may be unannotated ($\{G \vdash M : A\}$) or have an annotation ($\{G \vdash M : A\}^{\text{@}^n}$ or $\{G \vdash M : A\}^{*^n}$). Note that only the syntax of atomic formulas is extended with these annotations.

Definition 4.54. *A formula F containing annotations is well-formed with respect to Θ and Ξ if the formula F' obtained by erasing all annotations in F is such that $\Theta; \Xi \vdash F'$ fmla holds. We overload notation by denoting this property also by $\Theta; \Xi \vdash F$ fmla. A sequent \mathcal{S} containing annotations is well-formed if the sequent \mathcal{S}' obtained from \mathcal{S} by erasing all annotations is well-formed by virtue of Definition 4.5.*

We now refine the semantics of sequents to take into account the annotations on formulas. The basis for doing so is provided by an association between annotations and actual heights.

Definition 4.55. *A height assignment Υ maps each annotation @ⁱ to a natural number m_i . Given a particular Υ , a height assignment that is identical to Υ except that @ⁱ is mapped to m is denoted by $\Upsilon[\text{@}^i \leftarrow m]$.*

Definition 4.56. *We define validity only for closed annotated formulas that are well-formed, i.e., for formulas F such that $\mathcal{N} \cup \Theta_0; \emptyset \vdash F$ fmla is derivable. For such formulas, we first define formula validity with respect to a height assignment Υ , written $\Upsilon \models F$ valid, as follows:*

- $\Upsilon \models \{G \vdash M : A\}^{\text{Ann}}$ valid holds if $\vdash_{\Sigma} G$ ctx and $G \vdash_{\Sigma} A$ type are derivable, and
 - $G \vdash_{\Sigma} M \Leftarrow A$ has a derivation, if Ann is the empty annotation,
 - $G \vdash_{\Sigma} M \Leftarrow A$ has a derivation of height less than or equal to $\Upsilon(\text{@}^i)$, if Ann is @ⁱ, and
 - $G \vdash_{\Sigma} M \Leftarrow A$ has a derivation of height less than $\Upsilon(\text{@}^i)$, if Ann is *ⁱ.
- $\Upsilon \models \top$ valid holds.
- $\Upsilon \models \perp$ valid does not hold.
- $\Upsilon \models F_1 \supset F_2$ valid holds if $\Upsilon \models F_2$ valid holds in the case that $\Upsilon \models F_1$ valid holds.

- $\Upsilon \vDash F_1 \wedge F_2$ valid holds if both $\Upsilon \vDash F_1$ valid and $\Upsilon \vDash F_2$ valid hold.
- $\Upsilon \vDash F_1 \vee F_2$ valid holds if either $\Upsilon \vDash F_1$ valid or $\Upsilon \vDash F_2$ valid holds.
- $\Upsilon \vDash \Pi \Gamma : \mathcal{C}.F$ valid holds if $\Upsilon \vDash F[\{G/\Gamma\}]$ valid holds for every context expression G such that $\mathcal{N}; \emptyset \vdash_{cs} G$ is derivable.
- $\Upsilon \vDash \forall x : \alpha.F$ valid holds if $\Upsilon \vDash F[\{\langle x, M, \alpha \rangle\}]$ valid holds for every M such that $\mathcal{N} \cup \Theta_0 \vdash_{at} M : \alpha$ is derivable.
- $\Upsilon \vDash \exists x : \alpha.F$ valid holds if $\Upsilon \vDash F[\{\langle x, M, \alpha \rangle\}]$ valid holds for some M such that $\mathcal{N} \cup \Theta_0 \vdash_{at} M : \alpha$ is derivable.

As with Definition 3.6, the coherence of this definition is assured by Theorems 3.4 and 3.5.

The validity of a sequent containing annotations corresponds to the validity of each of its closed instances relative to every height assignment. In formalizing this idea, we assume the adaptation of the notions of the compatibility of term substitutions (Definition 4.6), the appropriateness of context substitutions (Definition 4.11) and the applications of these substitutions (Definitions 4.9 and 4.13) to annotated sequents that is obtained by ignoring the annotations on formulas. Further, we refer to every instance of an annotated well-formed sequent that is determined by term and context substitutions as in Definition 4.15 as one of its closed instances; the wellformedness of these closed instances follows easily from the results of Section 4.1.

Definition 4.57. *A well-formed sequent of the form $\mathbb{N}; \emptyset; \emptyset; \Omega \longrightarrow F$ is valid with respect to a height assignment Υ if $\Upsilon \vDash F$ valid holds whenever $\Upsilon \vDash F'$ valid holds for every $F' \in \Omega$. A well-formed sequent \mathcal{S} is valid with respect to Υ if every closed instance of \mathcal{S} is valid with respect to Υ and it is valid, without qualifications, under the extended semantics if it is valid with respect to every height assignment.*

While height assignments associate heights with every annotation, the assignments to only a finite subset of annotations matter in determining the validity of a formula or sequent. This observation is an immediate consequence of the following lemma.

Lemma 4.58. *If F is a formula in which the annotations $@^i$ and $*^i$ do not occur, then $\Upsilon \vDash F$ valid holds for a height assignment Υ if and only if $\Upsilon[@^i \leftarrow m] \vDash F$ valid holds for every choice of m . Similarly, if \mathcal{S} is a sequent in which the annotations $@^i$ and $*^i$ do not occur, then \mathcal{S} is valid with respect to Υ if and only if \mathcal{S} is valid with respect to $\Upsilon[@^i \leftarrow m]$ for every choice of m .*

Proof. The first clause is shown by a straightforward induction on the formation of F where in the base case we know that no atomic formula is annotated by $@^i$ or $*^i$ and therefore the m_i assigned to that annotation cannot play a role in determining its validity. The second clause follows from the first using the definition of validity for sequents of Definition 4.57. \square

Our ultimate interest is in determining that formulas devoid of annotations are valid in the sense articulated in Definition 3.6. This also means that we are eventually interested in the validity of sequents devoid of annotations in the sense described in Definition 4.15. However, in building in capabilities for inductive reasoning, we will consider rules that will introduce annotated formulas into sequents. Establishing the soundness of the resulting proof system requires us to refine the definition of validity for sequents to the one presented in Definition 4.57. That this is acceptable from the perspective of our eventual goal is the content of the following theorem.

$$\frac{\mathbb{N}; \Psi; \Xi; \Omega, \mathcal{Q}_1.F_1 \supset \dots \supset \mathcal{Q}_{k-1}.F_{k-1} \supset \mathcal{Q}_k.\{G \vdash M : A\}^{*i} \supset \mathcal{Q}_{k+1}.F_{k+1} \supset \dots \supset F_n \longrightarrow \mathcal{Q}_1.F_1 \supset \dots \supset \mathcal{Q}_{k-1}.F_{k-1} \supset \mathcal{Q}_k.\{G \vdash M : A\}^{\textcircled{i}} \supset \mathcal{Q}_{k+1}.F_{k+1} \supset \dots \supset F_n}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \mathcal{Q}_1.F_1 \supset \dots \supset \mathcal{Q}_{k-1}.F_{k-1} \supset \mathcal{Q}_k.\{G \vdash M : A\} \supset \mathcal{Q}_{k+1}.F_{k+1} \supset \dots \supset F_n} \text{ind}$$

Figure 4.6: The Induction Proof Rule

Theorem 4.59. *A well-formed sequent that is devoid of annotations is valid in the sense of Definition 4.57 if and only if it is valid in the sense of Definition 4.15.*

Proof. Using an argument similar to that for Lemma 4.58, we show that a closed instance of a sequent devoid of annotations is valid in the sense of Definition 4.15 if and only if it is valid by virtue of Definition 4.56 with respect to any height assignment. The theorem is an immediate consequence. \square

Our attention henceforth will be on sequents that potentially contain annotated formulas. For this reason, absent qualifications, we shall interpret “wellformedness” by virtue of Definition 4.54 and “validity” by virtue of Definitions 4.56 and 4.57. As we show below, the proof rules that we have discussed up to this point that apply to non-atomic formulas and that are different from the *id* rule lift in an obvious way to the situation where formulas carry annotations. Similarly, the *id* rule still applies as before when the formula that is the focus of the rule is unannotated, and no change is needed to the rules for atomic formulas when these formulas are unannotated. The extension of the *id* rule and the rules for atomic formulas to the situation when the focus formula is annotated needs some care. We discuss this matter after the presentation of the induction rule.

Theorem 4.60. *The following properties hold for the lifted forms of the proof rules in Figure 4.2, Figure 4.4, and the cut proof rule from Figure 4.3, and to the *id* rule and the rules from Figure 4.5 when the formula they pertain to are unannotated:*

1. *If the conclusion sequent is well-formed, the premises expressing typing conditions have derivations and the conditions expressed by the other, non-sequent premises are satisfied, then the premise sequents must be well-formed.*
2. *If the premises expressing typing judgements are derivable, the conditions described in the other non-sequent premises are satisfied and all the premise sequents are valid, then the conclusion sequent must also be valid.*

Proof. The definition of wellformedness for annotated sequents is based on the original notion via the erasure of annotations. Hence, the first claim follows immediately from the earlier results for unannotated sequents. The second claim can be proved relative to an arbitrary height assignment; this then generalizes to all possible height assignments. For the proof rules not relating to atomic formulas, the heights which may be assigned to annotations can play no role in the soundness argument as there are no atomic formulas being interpreted within the proof. For the atomic proof rules the claim is restricted to the case where the formulas being analyzed in the rules are not annotated, and thus again the height assignment has no impact on the previously presented argument for soundness. \square

4.4.2 The Induction Proof Rule

The induction rule is presented in Figure 4.6. In this rule, \mathcal{Q}_j , for $1 \leq j < n$, represents a sequence of context and universal term quantifiers. There is a proviso on the rule: the annotations $@^i$ and $*^i$ must be fresh, i.e., they must not already appear in the sequent that is the conclusion of the rule.

Given the definition of well-formedness for formulas and sequents that contain annotations, the following theorem has an obvious proof.

Theorem 4.61. *If the conclusion sequent of an instance of the ind rule is well-formed, then the premise sequent must be well-formed.*

A derivation of the premise sequent of the induction rule provides a schema for constructing a concrete, valid derivation for any closed instance of the conclusion sequent based on the height m of the assumption derivation in LF. Since this proof will be general with respect to the choice of m it ensures that the property holds for all natural numbers, and so the sequent without an annotated atomic formula will be valid. The soundness of this rule is shown by formalizing these ideas as a meta-level argument using induction on natural numbers. The following two lemmas are useful towards this end.

Lemma 4.62. *Let $F_1, \dots, F_{k-1}, F_{k+1}, \dots, F_n$ be formulas in which the annotations $@^i$ and $*^i$ do not appear, let $F^{\textcircled{i}}$ be a closed well-formed formula of the form*

$$\mathcal{Q}_1.F_1 \supset \dots \supset \mathcal{Q}_{k-1}.F_{k-1} \supset \mathcal{Q}_k.\{G \vdash M : A\}^{\textcircled{i}} \supset \mathcal{Q}_{k+1}.F_{k+1} \supset \dots \supset F_n$$

*in which, for $1 \leq j < n$, \mathcal{Q}_j denotes a sequence of context or universal term quantifiers, and let F^{*i} be the (obviously closed and well-formed) formula*

$$\mathcal{Q}_1.F_1 \supset \dots \supset \mathcal{Q}_{k-1}.F_{k-1} \supset \mathcal{Q}_k.\{G \vdash M : A\}^{*i} \supset \mathcal{Q}_{k+1}.F_{k+1} \supset \dots \supset F_n$$

*For any height assignment Υ and natural number m , if it is the case that $\Upsilon[@^i \leftarrow l] \models F^{\textcircled{i}}$ valid holds for every $l < m$, then $\Upsilon[@^i \leftarrow m] \models F^{*i}$ valid must also hold.*

Proof. We will prove the lemma by an induction on the number of top-level quantifiers and implications in $F^{\textcircled{i}}$ prior to the occurrence of the formula $\{G \vdash M : A\}^{\textcircled{i}}$ in it; the argument will implicitly make use of the closed and well-formed nature of the formulas under consideration.

In the base case, $F^{\textcircled{i}}$ is a formula of the form $\{G \vdash M : A\}^{\textcircled{i}} \supset F''$, where F'' is such that the annotations $@^i$ and $*^i$ do not appear in it. The formula F^{*i} correspondingly has the form $\{G \vdash M : A\}^{*i} \supset F''$. We may assume here that $\Upsilon[@^i \leftarrow m] \models \{G \vdash M : A\}^{*i}$ valid holds because the lemma is obviously true if it does not. From this, it easily follows that there is an l less than m for which $\Upsilon[@^i \leftarrow l] \models \{G \vdash M : A\}^{\textcircled{i}}$ valid holds. From the assumption about $F^{\textcircled{i}}$, it follows then that $\Upsilon[@^i \leftarrow l] \models F''$ valid holds. But then, by Lemma 4.58, it follows that $\Upsilon[@^i \leftarrow m] \models F''$ valid must also hold, from which we conclude easily that $\Upsilon[@^i \leftarrow m] \models F^{*i}$ valid holds.

The cases in which $F^{\textcircled{i}}$ begins with a context and universal term quantifier are similar, so we consider only the latter explicitly. In this case, $F^{\textcircled{i}}$ is a formula of the form $\forall x : \alpha.F'^{\textcircled{i}}$ and F^{*i} is correspondingly a formula of the form $\forall x : \alpha.F'^{*i}$, where F'^{*i} is a formula that is identical to $F'^{\textcircled{i}}$ except that the annotation on the formula $\{G \vdash M : A\}$ within it is changed to $*^i$. Note that $F'^{\textcircled{i}}$ and F'^{*i} must be formulas such that any of their substitution instances taken in pairs meet the descriptions in the lemma statement and all the substitution instances of $F'^{\textcircled{i}}$ must be less complex than $F^{\textcircled{i}}$. Now consider a term t such that $\mathcal{N} \cup \Theta_0 \vdash_{at} t : \alpha$. By the definition of validity and the assumption in the lemma, $\Upsilon[@^i \leftarrow l] \models F'^{\textcircled{i}}[\{\langle x, t, \alpha \rangle\}]$ valid must hold for any

$l < m$. But then, by the induction hypothesis, $\Upsilon[\textcircled{i} \leftarrow m] \models F'^{*^i}[\{\langle x, t, \alpha \rangle\}]$ **valid** must also hold. Since this is true regardless of the choice of t , it follows from the definition of validity that $\Upsilon[\textcircled{i} \leftarrow m] \models \forall x : \alpha. F'^{*^i}$ **valid**, i.e. $\Upsilon[\textcircled{i} \leftarrow m] \models F'^{*^i}$ **valid**, must hold.

Finally, suppose that $F^{\textcircled{i}}$ is a formula of the form $F'' \supset F'^{\textcircled{i}}$ and that F'^{*^i} is correspondingly a formula of the form $F'' \supset F'^{*^i}$, where F'^{*^i} differs from $F'^{\textcircled{i}}$ only in that the annotation on the occurrence of the formula $\{G \vdash M : A\}$ within it is changed to $*^i$. Note that $F'^{\textcircled{i}}$ and F'^{*^i} must be formulas that, as a pair, meet the descriptions in the lemma statement and $F'^{\textcircled{i}}$ must also be less complex than $F^{\textcircled{i}}$. We may assume in this case that $\Upsilon[\textcircled{i} \leftarrow m] \models F''$ **valid** holds because the lemma has a trivial proof if it does not. Using Lemma 4.58, we then see that $\Upsilon[\textcircled{i} \leftarrow l] \models F''$ **valid** must hold for any $l < m$. From the assumption in the lemma, it then easily follows that it must be the case that $\Upsilon[\textcircled{i} \leftarrow l] \models F'^{\textcircled{i}}$ **valid** holds for any $l < m$. Using the induction hypothesis, we may now conclude that $\Upsilon[\textcircled{i} \leftarrow m] \models F'^{*^i}$ **valid** holds, from which it follows easily that $\Upsilon[\textcircled{i} \leftarrow m] \models F'^{*^i}$ **valid** must hold. \square

Lemma 4.63. *Let F be a closed formula of the form*

$$\mathcal{Q}_1.F_1 \supset \dots \supset \mathcal{Q}_{k-1}.F_{k-1} \supset \mathcal{Q}_k.\{G \vdash M : A\} \supset \mathcal{Q}_{k+1}.F_{k+1} \supset \dots \supset F_n$$

in which the annotations \textcircled{i} or $^i$ do not occur and let $F^{\textcircled{i}}$ correspondingly be the formula*

$$\mathcal{Q}_1.F_1 \supset \dots \supset \mathcal{Q}_{k-1}.F_{k-1} \supset \mathcal{Q}_k.\{G \vdash M : A\}^{\textcircled{i}} \supset \mathcal{Q}_{k+1}.F_{k+1} \supset \dots \supset F_n$$

*If $\Upsilon[\textcircled{i} \leftarrow m] \models F^{\textcircled{i}}$ **valid** holds for every natural number m , then $\Upsilon \models F$ **valid** must also hold.*

Proof. This lemma is also proved by an induction on the number of top-level quantifiers and implications in F prior to the occurrence of the formula $\{G \vdash M : A\}$ in it. The argument again implicitly uses the information that the F and $F^{\textcircled{i}}$ are closed and well-formed.

In the base case, F is a formula of the form $\{G \vdash M : A\} \supset F''$, where F'' is a formula that is devoid of the annotations \textcircled{i} and $*^i$, and $F^{\textcircled{i}}$ is correspondingly the formula $\{G \vdash M : A\}^{\textcircled{i}} \supset F''$. If $\Upsilon[\textcircled{i} \leftarrow m] \models \{G \vdash M : A\}$ **valid** does not hold for any m , then $G \vdash_{\Sigma} M \Leftarrow A$ must not have a derivation and hence $\Upsilon \models \{G \vdash M : A\}$ **valid** must not hold. It is easy to see that the lemma must be true in this situation. On the other hand, if $\Upsilon[\textcircled{i} \leftarrow m] \models \{G \vdash M : A\}$ **valid** does hold for some m , then, by the assumptions of the lemma, $\Upsilon[\textcircled{i} \leftarrow m] \models F''$ **valid** must also hold. But then, by Lemma 4.58, $\Upsilon \models F''$ **valid** must hold and, hence, so also should $\Upsilon \models F$ **valid**.

The argument in the inductive cases follows a pattern that is very similar to that in the proof of Lemma 4.62. We avoid repeating the details that should be obvious at this point. \square

Theorem 4.64. *If the conclusion sequent of an instance of the ind rule is well-formed, the premise sequent is valid and the requirement of non-occurrence of the annotations \textcircled{i} and $*^i$ is satisfied, then the conclusion sequent of the rule instance must be valid.*

Proof. Let θ and σ be substitutions that identify a closed instance of the conclusion sequent and let Υ be a height assignment. The instance of the conclusion sequent would obviously be valid with respect to Υ if any formula in $\Omega[\theta][\sigma]$ is not valid with respect to Υ , so let us assume that they are all in fact so valid. Our task then is to show that the instance of the goal formula in the conclusion sequent determined by θ and σ is valid with respect to Υ . Using a harmless overloading of notation, we will identify this instance as the formula

$$\mathcal{Q}_1.F_1 \supset \dots \supset \mathcal{Q}_{k-1}.F_{k-1} \supset \mathcal{Q}_k.\{G \vdash M : A\} \supset \mathcal{Q}_{k+1}.F_{k+1} \supset \dots \supset F_n,$$

and refer to it as F . We will also use the notation $F^{\textcircled{i}}$ and F^{*i} below to denote the formulas that differ from F only in that the distinguished occurrence of the subformula $\{G \vdash M : A\}$ is annotated with \textcircled{i} and $*^i$, respectively.

Our objective is to show that $\Upsilon \vDash F$ **valid** holds. We will do this by showing that the judgement $\Upsilon[\textcircled{i} \leftarrow m] \vDash F^{\textcircled{i}}$ **valid** holds for every m and then invoking Lemma 4.63. In proving the latter, we will use the observation that θ and σ identify a closed instance also of the premise sequent of the rule and the assumption that all instances of the premise sequent are valid. The argument is carried out by (strong) induction on m . Given the structure of the formula $F^{\textcircled{i}}$ and the fact that no typing derivation in LF has height 0, it is obvious that $\Upsilon[\textcircled{i} \leftarrow 0] \vDash F^{\textcircled{i}}$ **valid** holds. Now assume that $\Upsilon[\textcircled{i} \leftarrow l] \vDash F^{\textcircled{i}}$ **valid** holds for every l less than m . By Lemma 4.62, it follows then that $\Upsilon[\textcircled{i} \leftarrow m] \vDash F^{*i}$ **valid** holds. Since every formula in $\Omega[\theta][\sigma]$ is valid with respect to Υ by assumption, it follows by virtue of Lemma 4.58 and the proviso that the annotations \textcircled{i} and $*^i$ does not appear in the formulas of Ω that every assumption formula in the instance of the premise sequent identified by θ and σ must be true under the height assignment $\Upsilon[\textcircled{i} \leftarrow m]$. That $\Upsilon[\textcircled{i} \leftarrow m] \vDash F^{\textcircled{i}}$ **valid** must hold is now a consequence of the validity of this instance of the premise sequent. \square

4.4.3 Additional Proof Rules that Interpret Annotations

With the addition of annotations to the syntax, we may consider additional rules for introducing annotated atomic formulas. The form that these rules take must pay attention to the heights of the LF derivations that these formulas encode; this is essential to being able to make effective use of the induction rule. In this spirit, our proof system includes the variants of the existing rules for atomic formulas that are shown in Figure 4.7. In the *atm-app-L** and *atm-abs-L** rules, *Ann* must be either \textcircled{i} or $*^i$. In the *atm-app-L** rule, Ω' denotes the formulas that result from Ω and $\{G_1 \vdash M_1 : A_1\}, \dots, \{G_k \vdash M_k : A_k\}$ denote the formulas that result from $\{G \vdash R : P\}$ in the reduction of each of the instances of the conclusion sequent that are considered by *AllCases*. Note that the \textcircled{i} annotation on the assumption formula that is the focus of the rule is replaced by the $*^i$ annotation in the assumption formulas in the premise sequent, as is needed for the hypothesis added by the induction rule to be useful.

The *id* rule described in Section 4.2.2 requires the assumption and conclusion formulas that are matched to be equivalent in terms of validity. With the inclusion of annotations, it is possible to weaken this requirement. For example, the assumption formula $\{G \vdash M : A\}^*$ ensures the validity of a sequent in which the conclusion formula is $\{G \vdash M : A\}^{\textcircled{i}}$. Similarly, the presence of either of these formulas as an assumption in a sequent suffices to ensure its validity if its conclusion formula is $\{G \vdash M : A\}$. These observations can be expanded to include non-atomic formulas with the proviso that the polarity of the occurrence of the formula must be paid attention to. For example, it is the validity of $\{G \vdash M : A\}^{\textcircled{i}} \supset F$ that implies that $\{G \vdash M : A\}^* \supset F$ is valid, and the validity of both of these formulas is implied by the validity of $\{G \vdash M : A\} \supset F$. We make these thoughts precise through a definition of comparative strengths of formulas.

Definition 4.65. *We first identify an ordering over annotations: $*^i$ is stronger than \textcircled{i} which, in turn, is stronger than no annotation. Next we define a strength relation between atomic formulas relative to a context variables context Ξ and a permutation π of nominal constants: $\{G \vdash M : A\}^{\text{Ann}}$ is at least as strong as $\{G' \vdash M' : A'\}^{\text{Ann}'}$ in this context if $\Xi \vdash \{G \vdash M : A\} \equiv_{\pi} \{G' \vdash M' : A'\}$ holds and *Ann* is stronger than or identical to *Ann'*. Finally, we extend this strength relation to arbitrary formulas. The formula $F_2 \supset F_2'$ is at least as strong as $F_1 \supset F_1'$ with respect to Ξ and π if F_1 is at least as strong as F_2 with respect to Ξ and π^{-1} and F_2' is at least as strong as F_1' with*

$$\begin{array}{c}
\mathcal{CS} = \text{AllCases}(\mathbb{N}; \Psi; \Xi; \Omega, \{G \vdash R : P\} \longrightarrow F, \{G \vdash R : P\}) \\
\frac{\{ \mathbb{N}'; \Psi'; \Xi'; \Omega', \{G_1 \vdash M_1 : A_1\}^{*i}, \dots, \{G_k \vdash M_k : A_k\}^{*i} \longrightarrow F' \\
\quad | \quad \mathbb{N}'; \Psi'; \Xi'; \Omega', \{G_1 \vdash M_1 : A_1\}, \dots, \{G_k \vdash M_k : A_k\} \longrightarrow F' \in \mathcal{CS} \}}{\mathbb{N}; \Psi; \Xi; \Omega, \{G \vdash R : P\}^{\text{Ann}} \longrightarrow F} \text{ atm-app-L}^* \\
\\
h : \prod x_1 : A_1. \dots \prod x_n : A_n. P \in \Sigma \text{ or the explicit bindings in } G \\
G \text{ appears as the context of some atomic formula in } \Omega \\
P[\{\langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_n, M_n, (A_n)^- \rangle\}] = P' \\
\frac{\{ \mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \\
\quad \{G \vdash M_i : A_i[\{\langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_{i-1}, M_{i-1}, (A_{i-1})^- \rangle\}]^{*i} \mid 1 \leq i \leq n\}}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \{G \vdash h M_1 \dots M_n : P'\}^{\text{@}^i}} \text{ atm-app-R}^* \\
\\
\Xi' = \begin{cases} (\Xi \setminus \{\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[\mathcal{G}]\}) \cup \{\Gamma \uparrow (\mathbb{N}_\Gamma, n : (A_1)^-) : \mathcal{C}[\mathcal{G}]\} & \text{if } \Gamma \text{ appears in } G \\ \Xi & \text{otherwise} \end{cases} \\
\mathbb{N}, n : (A_1)^-; \Psi; \Xi'; \\
\frac{\Omega, \{G, n : A_1 \vdash M[\{\langle x, n, (A_1)^- \rangle\}] : A_2[\{\langle x, n, (A_1)^- \rangle\}]^{*i} \longrightarrow F}{\mathbb{N}; \Psi; \Xi; \Omega, \{G \vdash \lambda x. M : \prod x : A_1. A_2\}^{\text{Ann}} \longrightarrow F} \text{ atm-abs-L}^* \\
\\
\Xi' = \begin{cases} (\Xi \setminus \{\Gamma \uparrow \mathbb{N}_\Gamma : \mathcal{C}[\mathcal{G}]\}) \cup \{\Gamma \uparrow (\mathbb{N}_\Gamma, n : (A_1)^-) : \mathcal{C}[\mathcal{G}]\} & \text{if } \Gamma \text{ appears in } G \\ \Xi & \text{otherwise} \end{cases} \\
\mathbb{N}, n : (A_1)^-; \Psi; \Xi'; \\
\frac{\Omega \longrightarrow \{G, n : A_1 \vdash M[\{\langle x, n, (A_1)^- \rangle\}] : A_2[\{\langle x, n, (A_1)^- \rangle\}]^{*i} \}}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \{G \vdash \lambda x. M : \prod x : A_1. A_2\}^{\text{@}^i}} \text{ atm-abs-R}
\end{array}$$

Figure 4.7: Rules that Introduce Atomic Formulas with Annotations

respect to Ξ and π . For all other non-atomic formulas, F_2 is at least as strong as F_1 with respect to Ξ and π if their components satisfy the same relation, under a possibly extended Ξ in the case of context quantification, allowing for renaming of variables bound by quantifiers. The “at least as strong as” relation for formulas relative to Ξ and π is represented by the judgement $\Xi \vdash F_2 \succeq_\pi F_1$.

The strength relation between formulas is preserved by term and context variables substitutions under some provisos.

Lemma 4.66. *Suppose that $\Xi \vdash F_2 \succeq_\pi F_1$ holds for suitable F_1, F_2, Ξ and π . Then,*

1. *if θ is a term substitution such that $\text{supp}(\theta) \cap \text{supp}(\pi) = \emptyset$ and both $F_2[\theta] = F'_2$ and $F_1[\theta] = F'_1$ have derivations for some F'_1 and F'_2 , then $\Xi[\theta] \vdash F_2[\theta] \succeq_\pi F_1[\theta]$ holds; and*
2. *if σ is an appropriate substitution for Ξ with respect to some term variables context Ψ and set of nominal constants \mathbb{N} then $\Xi_\sigma \vdash F_2[\sigma] \succeq_\pi F_1[\sigma]$ holds.*

Proof. Both clauses can be shown to be true by an inductive argument on the structure of F_2 , using the definition of substitution and invoking Lemmas 4.25 and 4.26 in the case of atomic formulas. \square

$$\frac{F_2 \in \Omega \quad \pi \text{ is a permutation of nominal constants such that } \text{supp}(\pi) \subseteq \mathbb{N} \quad \Xi \vdash F_2 \succeq_\pi F_1}{\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow F_1} \textit{id}$$

Figure 4.8: The Axiom Rule Generalized to Annotated Formulas

The main property for the strength definition that we will use is that validity of a closed formula is ensured by that of another closed formula that is at least as strong as it is.

Lemma 4.67. *Let F_1 and F_2 be closed well-formed formulas, let Ξ be a context variables context and let π be a permutation of nominal constants such that $\Xi \vdash F_2 \succeq_\pi F_1$ holds. If F_2 is valid with respect to a height assignment Υ , then F_1 must also be valid with respect to Υ .*

Proof. The proof by an induction on the structure of F_2 using the definition of $\Xi \vdash F_2 \succeq_\pi F_1$. The argument takes an obvious form in the inductive cases and we consider only the case for atomic formulas, where the annotations become important, in detail. In this case, F_2 and F_1 must, respectively, be of the form $\{G_2 \vdash M_2 : A_2\}^{Ann_2}$ and $\{G_1 \vdash M_1 : A_1\}^{Ann_1}$, where Ann_2 is an annotation that is stronger than or identical to Ann_1 , and $\Xi \vdash \{G_2 \vdash M_2 : A_2\} \equiv_\pi \{G_1 \vdash M_1 : A_1\}$ must hold. Since the formulas are closed, the last observation leads to the conclusion that $\pi.(\{G_2 \vdash M_2 : A_2\})$ and $\{G_1 \vdash M_1 : A_1\}$ are identical. Using Theorem 3.9, it follows from this that if there are derivations for $\vdash_\Sigma G_2 \textit{ ctx}$, $G_2 \vdash_\Sigma A_2 \textit{ type}$ and $G_2 \vdash_\Sigma M_2 \Leftarrow A_2$, then there must be derivations that have the *same structure* for $\vdash_\Sigma G_1 \textit{ ctx}$, $G_1 \vdash_\Sigma A_1 \textit{ type}$ and $G_1 \vdash_\Sigma M_1 \Leftarrow A_1$, respectively. The claim in the lemma can now be verified using the definition of validity. \square

Figure 4.8 presents a generalized version of the *id* rule based on the discussions above. We show the preservation of wellformedness and the soundness properties for the new rules below.

Theorem 4.68. *The following properties hold for every instance of the rules in Figures 4.7 and 4.8:*

1. *If the conclusion sequent is well-formed, the premises expressing typing conditions have derivations and the conditions expressed by the other, non-sequent premises are satisfied, then all the sequent premises must be well-formed.*
2. *If the conclusion sequent is well-formed, the premises expressing typing judgements are derivable, the conditions described in the other non-sequent premises are satisfied and the premise sequent is valid, then the conclusion sequent must also be valid.*

Proof. The truth of the first clause can be verified by using Theorems 4.24 and 4.52 and noting that the wellformedness of sequents with annotations is determined by the wellformedness of the sequent that results from erasing all the annotations.

The truth of the second clause for the new *id* rule follows easily from Lemmas 4.66 and 4.67 and the definition of validity for sequents. For the rules in Figure 4.7, we adapt the argument for Theorem 4.53 to the new semantics. The refinement to the semantics to take into account the heights of derivations for typing judgements associated with atomic formulas does not impact the consideration of any formula other than the one introduced by the rule and the ones in the premise sequents associated with it. For the formula introduced by the *atm-app-L** rule, we use the fact verified in Lemma 4.44 that if the typing judgement that determines the height of a closed instance of the formula $\{G \vdash R : P\}$ has a derivation of height k , then the typing judgements that determine the heights of the corresponding closed instances of the new assumption formulas in the reduced

form of the sequent must have derivations of height less than k . For the formula introduced by the $atm\text{-app}\text{-}R^*$ rule, we use the observation contained in Theorem 2.20 that the typing judgement that determines the height of a closed instance of the formula $\{G \vdash h M_1 \dots M_n : P'\}$ must have a derivation of height at most $(k + 1)$ if the typing judgement associated with the corresponding instances of the goal formulas of the premise sequents have height at most k . Finally, for the formula introduced by the $atm\text{-abs}\text{-}L^*$ and $atm\text{-abs}\text{-}R^*$ rules, we use the easily verified fact that an LF judgement of the form $\Gamma \vdash_{\Sigma} \lambda x. M \Leftarrow \Pi x:A_1. A_2$ has a derivation of height $(k + 1)$ if and only if the judgement $\Gamma, x : A_1 \vdash_{\Sigma} M \Leftarrow A_2$ has a derivation of height k . \square

4.5 Proof Rules Encoding LF Meta-Theorems

We now present proof rules that encode the meta-theorems discussed in Section 2.3 are often used in informal arguments about the properties of LF specifications. The strengthening, permutation and instantiation meta-theorems can be encoded as axioms, i.e., as proof rules that have no premise sequents although they may have side-conditions that determine applicability. Given the interpretation of atomic formulas, the weakening rule, which introduces a new type assignment into the LF context, requires the type to be checked for wellformedness. This checking is built into the proof rule that encodes it via suitable premise sequents.

To determine the premise sequents for the weakening rule, we use the process of “type decomposition” that is embodied in the following definition.

Definition 4.69. *The decomposition of an LF type A with respect to a collection of nominal constants \mathbb{N} , a context variables context Ξ and a context expression G , denoted by $Decompose(\mathbb{N}, \Xi, G, A)$, is defined by recursion on the type A as follows:*

1. *If A is a type of the form $(a M_1 \dots M_n)$ where $a : \Pi x_1:A_1. \dots \Pi x_n:A_n$. $Type \in \Sigma$, then $Decompose(\mathbb{N}, \Xi, G, A)$ is the collection*

$$\{\langle \mathbb{N}, \Xi, \{G \vdash M_i : A_i[\langle \langle x_1, M_1, (A_1)^- \rangle, \dots, \langle x_{i-1}, M_{i-1}, (A_{i-1})^- \rangle]\} \mid 1 \leq i \leq n \rangle\}.$$

2. *If $A = \Pi x:A_1. A_2$, then letting G' be $G, n : A_1$, \mathbb{N}' be $\mathbb{N} \cup \{n\}$, and Ξ' be the set*

$$\{\Gamma \uparrow \mathbb{N}'_{\Gamma} : \mathcal{C}[\mathcal{G}] \mid \Gamma \uparrow \mathbb{N}_{\Gamma} : \mathcal{C}[\mathcal{G}] \in \Xi \text{ and } \mathbb{N}'_{\Gamma} \text{ is } \mathbb{N}_{\Gamma} \cup \{n\} \text{ if } \Gamma \text{ occurs in } G \text{ and } \mathbb{N}_{\Gamma} \text{ otherwise}\}$$

for some nominal constant $n : (A_1)^- \in \mathcal{N} \setminus \mathbb{N}$, $Decompose(\mathbb{N}, \Xi, G, A)$ is the collection $Decompose(\mathbb{N}, \Xi, G, A_1) \cup Decompose(\mathbb{N}', \Xi', G', A_2[\langle \langle x, n, (A_1)^- \rangle \rangle])$.

We shall refer to the set $\bigcup\{\mathbb{N}' \mid \langle \mathbb{N}', \Xi', F' \rangle \in Decompose(\mathbb{N}, \Xi, G, A)\} \setminus \mathbb{N}$ as the new name set of $Decompose(\mathbb{N}, \Xi, G, A)$; we circumvent the ambiguity in this identification by assuming that new names are chosen in the same way each time.

The definition above is abusive of notation because it uses the convention discussed after Theorem 2.8 for denoting the result of applying a substitution without the certainty that such a result will exist. However, this abuse is harmless: the result will exist in all the situations that we will use $Decompose$. The following lemma describes the conditions under which we will use this function and shows the harmlessness of the abuse in addition to establishing additional properties of $Decompose$ that we will need in justifying its use.

Lemma 4.70. *Let Ξ be a context variables context, Ψ a term variables context, and \mathbb{N} a set of nominal constants such that for each $\Gamma_i \uparrow \mathbb{N}_i : \mathcal{C}_i[\mathcal{G}_i]$ in Ξ the judgement $\mathbb{N} \setminus \mathbb{N}_i; \Psi \vdash \mathcal{C}_i[\mathcal{G}_i]$ ctx-ty has a derivation. Further, let G be a context expression and B be a type such that there are derivations*

for $\mathbb{N} \cup \Theta_0 \cup \Psi; \Xi^- \vdash G$ context and $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{ak} B$ type. Then $Decompose(\mathbb{N}, \Xi, G, B)$ must be defined. Moreover, letting $\bar{\mathbb{N}}$ be the new name set of $Decompose(\mathbb{N}, \Xi, G, B)$, the following properties hold of the set $Decompose(\mathbb{N}, \Xi, G, B)$ identifies:

1. For each $\langle \mathbb{N}', \Xi', F' \rangle \in Decompose(\mathbb{N}, \Xi, G, B)$ it is the case that

- (a) $\Gamma_i \uparrow \mathbb{N}_i : \mathcal{C}_i[\mathcal{G}_i] \in \Xi$ if and only if $\Gamma_i \uparrow \mathbb{N}'_i : \mathcal{C}_i[\mathcal{G}_i] \in \Xi'$ for some \mathbb{N}'_i such $\mathbb{N}' \setminus \mathbb{N} \supseteq \mathbb{N}'_i \setminus \mathbb{N}_i$,
- (b) each member $\Gamma_i \uparrow \mathbb{N}'_i : \mathcal{C}_i[\mathcal{G}_i]$ of Ξ' is such that $\mathbb{N}' \setminus \mathbb{N}'_i; \Psi \vdash \mathcal{C}_i[\mathcal{G}_i]$ ctx-ty has a derivation,
- (c) there is a derivation for $\mathbb{N}' \cup \Theta_0 \cup \Psi; \Xi'^- \vdash F'$ fmla.

2. For any closed term substitution θ with $dom(\theta) = \Psi$ and closed context substitution σ with $dom(\sigma) = \Xi$ and $supp(\sigma)$ disjoint from $\bar{\mathbb{N}}$, if it is the case that $G[\theta][\sigma]$ and $B[\theta]$ are defined and $\vdash_{\Sigma} G[\theta][\sigma]$ ctx has a derivation, and, further, that $F'[\theta][\sigma]$ is defined and is a valid formula for every $\langle \mathbb{N}', \Xi', F' \rangle \in Decompose(\mathbb{N}, \Xi, G, B)$, then there must be a derivation for the LF judgement $G[\theta][\sigma] \vdash_{\Sigma} B[\theta]$ type.

Proof. We must first determine the coherence of the lemma by checking that $G[\theta][\sigma]$, $B[\theta]$ and $F'[\theta][\sigma]$ are closed expressions for each of the substitutions θ , σ and formulas F' considered in clause (3). This must be the case because the substitutions are closed, all the free variables in B are contained in Ψ , and all the free variables in F' and G must be contained in Ψ and Ξ .

The lemma is now proved by induction on the structure of the type B . We consider the cases for this structure below.

B is the atomic type $(a M_1 \dots M_n)$.

Since $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{ak} B$ type is derivable, Σ must assign a kind of the form $\Pi x_1:A_1. \dots \Pi x_n:A_n$. Type to a and there must be derivations for $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{at} M_i : (A_i)^-$ for $1 \leq i \leq n$. From the wellformedness of Σ it follows that $\mathbb{N} \cup \Theta_0 \cup \Psi \cup \{x_1 : (A_1)^-, \dots, x_{i-1} : (A_{i-1})^-\} \vdash_{ak} A_i$ type has a derivation. We may now use Theorem 3.2 to conclude that $A_i[\{\langle x_1, M_1, (A_1)^-\rangle, \dots, \langle x_{i-1}, M_{i-1}, (A_{i-1})^-\rangle\}]$ is defined, thereby ensuring that $Decompose(\mathbb{N}, \Xi, G, B)$ is defined in this case.

We now turn to showing the various clauses. Clause 1(a) is trivial and clause 1(b) follows from the assumptions of the lemma. For the remainder of the argument, let us write A'_i to denote the type $A_i[\{\langle x_1, M_1, (A_1)^-\rangle, \dots, \langle x_{i-1}, M_{i-1}, (A_{i-1})^-\rangle\}]$, for $1 \leq i \leq n$. Theorem 3.2 yields the further observation that there are derivations for $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{ak} A'_i$ type. We also note that $Decompose(\mathbb{N}, \Xi, G, B)$ is the set $\{\langle \mathbb{N}, \Xi, \{G \vdash M_i : A'_i\} \mid 1 \leq i \leq n \rangle\}$. Now, for clause 1(c), it suffices to show that $\mathbb{N} \cup \Theta_0 \cup \Psi; \Xi^- \vdash G$ context, $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{ak} A'_i$ type, and $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{at} M_i : (A'_i)^-$ have derivations. The first is true by assumption and the latter two follow from earlier observations and the fact that erasure is preserved under substitution (Theorem 2.11).

This leaves only clause (2). The validity of $\{G[\theta][\sigma] \vdash M_i[\theta] : A'_i[\theta]\}$, for $1 \leq i \leq n$, implies that there must be LF derivations for $G[\theta][\sigma] \vdash_{\Sigma} M_i[\theta] \Leftarrow A'_i[\theta]$. Using Theorem 2.6, we see that $A'_i[\theta]$ is identical to $A_i[\theta][\{\langle x_1, M_1[\theta], (A_1)^-\rangle, \dots, \langle x_{i-1}, M_{i-1}[\theta], (A_{i-1})^-\rangle\}]$; we assume here that the variables x_1, \dots, x_n are chosen so as to not be members of $dom(\theta)$ or to appear free in the terms in $rng(\theta)$, as is possible to ensure through renaming. Using Theorems 2.21 and 2.11, we can conclude now that there must be a derivation for $G[\theta][\sigma] \vdash_{\Sigma} (a M_1[\theta] \dots M_n[\theta]) \Rightarrow \text{Type}$. The claim follows easily from this by noting that $B[\theta]$ is identical to $(a M_1[\theta] \dots M_n[\theta])$.

B is the type $\Pi x:A_1. A_2$.

We first show that the parameters of the two recursive uses of $Decompose$ satisfy what is required of them for the lemma statement to apply. The assumption that $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{ak} \Pi x:A_1. A_2$ type has a derivation implies that $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{ak} A_1$ type and $\mathbb{N} \cup \Theta_0 \cup \Psi \cup \{x : (A_1)^-\} \vdash_{ak} A_2$ type have

derivations. From the latter, using Theorem 3.2, we see that $A_2[\{\langle x, n, (A_1)^- \rangle\}]$ must be defined and that $\mathbb{N} \cup \Theta_0 \cup \Psi \vdash_{ak} A_2[\{\langle x, n, (A_1)^- \rangle\}]$ type. must have a derivation for any nominal constant n that is assigned the type $(A_1)^-$. Given these observations, it is straightforward to verify that the required conditions are indeed satisfied.

The induction hypothesis now yields the conclusion that $Decompose(\mathbb{N}, \Xi, G, B)$ is defined. A similar observation applies to clauses 1(a), 1(b) and 1(c). For clause 2, we observe first that substitutions θ and σ are “good” for this case only if they are good for the recursive calls; hence we may use them without specific reference to the context. We then observe that if $B[\theta]$ is defined, then so must $A_1[\theta]$ and $A_2[\theta]$ be. Now, if can show that, for the n chosen in the definition of $Decompose$, it is the case that $(G, n : A_1)[\theta][\sigma]$ is defined and that $\vdash_{\Sigma} (G, n : A_1)[\theta][\sigma]$ **ctx** has a derivation, we can invoke the induction hypothesis relative to the second recursive call to $Decompose$ to conclude that $(G, n : A_1)[\theta][\sigma] \vdash_{\Sigma} A_2[\theta]$ **type** must have a derivation and, hence, that $G[\theta][\sigma] \vdash_{\Sigma} \Pi x : A_1.A_2. [\theta]$ **type** has one, as we desired to show. That $(G, n : A_1)[\theta][\sigma]$ is defined follows from the assumption that $G[\theta][\sigma]$ defined and the earlier observation that $A_1[\theta]$ is. The induction hypothesis invoked relative to the first recursive call to $Decompose$ allows us to conclude that there is a derivation for $G[\theta][\sigma] \vdash_{\Sigma} A_1[\theta]$ **type**. By assumption, $\vdash_{\Sigma} G[\theta][\sigma]$ **ctx** has a derivation. Noting that $(G, n : A_1)[\theta][\sigma]$ is the same expression as $(G[\theta][\sigma], n : A_1[\theta])$, we see that $\vdash_{\Sigma} (G, n : A_1)[\theta][\sigma]$ **ctx** would have a derivation if we could be sure that n is not assigned a type in $G[\theta][\sigma]$. But this must be the case: the way the n is chosen ensures that G cannot have such a binding and the condition that $supp(\sigma)$ is disjoint from $\overline{\mathbb{N}}$ ensures that σ cannot introduce one. \square

$$\begin{array}{c}
\begin{array}{c}
n \text{ does not appear in } M, A, \text{ or the explicit bindings in } G \\
\text{if } \Gamma_i \uparrow \mathbb{N}_i : \mathcal{C}_i[\mathcal{G}_i] \in \Xi \text{ and } \Gamma_i \text{ appears in } G, \text{ then } n \in \mathbb{N}_i \\
\{N'; \Psi; \Xi'; \Omega \longrightarrow F' \mid (N', \Xi', F') \in Decompose(\mathbb{N}, \Xi, G, B)\} \\
\hline
\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \{G \vdash M : A\}^{Ann} \supset \{G, n : B \vdash M : A\}^{Ann} \quad LF-wk
\end{array} \\
\\
\begin{array}{c}
n \text{ does not appear in } M, A, \text{ or the explicit bindings in } G \\
\hline
\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \{G, n : B \vdash M : A\}^{Ann} \supset \{G \vdash M : A\}^{Ann} \quad LF-str
\end{array} \\
\\
\begin{array}{c}
G = G_1, n_1 : A_1, n_2 : A_2, G_2 \\
G' = G_1, n_2 : A_2, n_1 : A_1, G_2 \\
n_1 \text{ does not appear in } A_2 \\
\hline
\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \{G \vdash M : A\}^{Ann} \supset \{G' \vdash M : A\}^{Ann} \quad LF-perm
\end{array} \\
\\
\begin{array}{c}
G' = n : B, n_1 : A_1, \dots, n_m : A_m \quad \{A_i[\{\langle n, N, (B)^- \rangle\}] = A'_i \mid 1 \leq i \leq m\} \\
G'' = G, n_1 : A'_1, \dots, n_m : A'_m \\
M[\{\langle n, N, (B)^- \rangle\}] = M' \quad A[\{\langle n, N, (B)^- \rangle\}] = A' \\
\hline
\mathbb{N}; \Psi; \Xi; \Omega \longrightarrow \{G, G' \vdash M : A\} \supset \{G \vdash N : B\} \supset \{G'' \vdash M' : A'\} \quad LF-inst
\end{array}
\end{array}$$

Figure 4.9: Rules that Encode Meta-Theoretic Properties of LF

Figure 4.9 presents the proof rules which encode the content of the LF meta-theorems. The symbol *Ann* in the first three rules, which encode weakening, strengthening, and context permu-

tation, stands for no annotation, $@^i$ or $*^i$ for some i , used in the same manner throughout the rule instance. Annotations are permitted in these rules to reflect the fact that the corresponding meta-theorems guarantee the preservation of the structure, and thus height, of LF derivations.

We show that these rules are well-defined by demonstrating that they satisfy the preservation of wellformedness property for sequents and that they are sound.

Theorem 4.71. *The following property holds for every instance of the rules in Figure 4.9: if the conclusion sequent is well-formed, the premises expressing typing conditions have derivations and the conditions expressed by the other, non-sequent premises are satisfied, then the premise sequents must be well-formed.*

Proof. The theorem holds vacuously for the rules *LF-str*, *LF-perm*, and *LF-inst*. For *LF-wk*, Lemma 4.70 yields the conditions that are required of Ξ' and F' for each of the premise sequents, and the wellformedness of the conclusion sequent yields the condition required of the formulas in Ω . \square

Theorem 4.72. *The following property holds for every instance of each of the rules in Figure 4.9: if the premises expressing typing judgements are derivable, the conditions described in the other non-sequent premises are satisfied and all the premise sequents are valid, then the conclusion sequent must also be valid.*

Proof. To establish the theorem, we must show the validity of each closed instance of the conclusion sequent under the conditions described. In the case of the *LF-wk*, *LF-str* and *LF-perm*, the notion of validity must also be relativized to a height assignment. Since closed instances of atomic formulas are interpreted via the derivability of LF judgements, we may use the meta-theorems presented in Section 2.3 in this argument. The details are entirely straightforward in the case of the rules *LF-str*, *LF-perm* and *LF-inst*, and we therefore avoid their presentation here. The argument is more involved for the rule *LF-wk*: the interpretation of atomic formulas includes the wellformedness of contexts and types and we must show that the premises of this rule suffice to ensure this holds for the extension of the context embodied in the consequent of the goal formula. We therefore present the proof for this case in more detail below.

Let θ and σ be substitutions that identify a closed instance of the conclusion sequent of this rule. As noted above, we must show that this instance is valid with respect to any height assignment Υ if the assumptions of the lemma are satisfied. Let $\overline{\mathbb{N}}$ be the new name set of *Decompose* (\mathbb{N}, Ξ, G, B). We show what is required initially under the assumption that $\text{supp}(\theta)$ and $\text{supp}(\sigma)$ are disjoint from $\overline{\mathbb{N}}$; we will explain later how this condition can be obviated.

We claim that θ and σ identify closed instances of all the premise sequents under the above assumption. Such a sequent must have the form $\mathbb{N}'; \Psi; \Xi'; \Omega \longrightarrow F'$ where $\mathbb{N}' \subseteq \overline{\mathbb{N}} \cup \mathbb{N}$ and, by Lemma 4.70, each $\Gamma_i \uparrow \mathbb{N}'_i : \mathcal{C}_i[\mathcal{G}_i] \in \Xi'$ corresponds to a $\Gamma_i \uparrow \mathbb{N}_i : \mathcal{C}_i[\mathcal{G}_i] \in \Xi$ with $\mathbb{N}' \setminus \mathbb{N} \supseteq \mathbb{N}'_i \setminus \mathbb{N}_i$. Since $\Xi'^- = \Xi^-$, we only need to show that θ and σ are proper with respect to this sequent to be sure they identify a closed instance of it. The compatibility of $\langle \theta, \emptyset \rangle$ with the conclusion sequent implies that $\text{supp}(\theta) \cap \mathbb{N} = \emptyset$. Since we also have $\text{supp}(\theta) \cap \overline{\mathbb{N}} = \emptyset$, it follows that $\text{supp}(\theta)$ is disjoint from the support set of the shown sequent and hence $\langle \theta, \emptyset \rangle$ must be substitution compatible with it. Using Theorems 4.71 and 4.10, we see that the context types in the context variables context that results from applying θ to the sequent must be well-formed relative to it. Since the application of a term variables substitution does not remove any entries from a context variables context, there must be an association in $\Xi'[\theta]$ for every variable in $\text{dom}(\sigma)$. To determine that σ is appropriate for the sequent, it only remains to be shown that there is a derivation for $\text{supp}(\sigma) \setminus \mathbb{N}'_i; \emptyset; \emptyset \vdash \mathcal{C}_i[\mathcal{G}_i[\theta]] \rightsquigarrow_{\text{csty}} G_i$ for each context expression G_i that σ substitutes for a variable Γ_i . The appropriateness of σ for the

corresponding instance of the conclusion sequent tells us that $\text{supp}(\sigma) \setminus \mathbb{N}_i; \emptyset; \emptyset \vdash \mathcal{C}_i[\mathcal{G}_i[\theta]] \rightsquigarrow_{\text{csty}} G_i$ has a derivation. We would get the desired conclusion from this if $\text{supp}(\sigma) \setminus \mathbb{N}'_i = \text{supp}(\sigma) \setminus \mathbb{N}_i$. But this must be the case: since $\mathbb{N}' \setminus \mathbb{N} \supseteq \mathbb{N}'_i \setminus \mathbb{N}_i$ it follows that $\overline{\mathbb{N}} \supseteq \mathbb{N}'_i \setminus \mathbb{N}_i$ and we know that $\text{supp}(\sigma) \cap \overline{\mathbb{N}} = \emptyset$.

We turn now to the main task, that of showing that the instance of the conclusion sequent is valid respect to the height assignment Υ . For this, it suffices to show that if all the formulas in $\Omega \cup \{\{G \vdash M : A\}^{Ann}\}[\theta][\sigma]$ are valid with respect to Υ , then $\{G, n : B \vdash M : A\}^{Ann}[\theta][\sigma]$ is also so valid. Using Theorem 2.14, the interpretation of atomic formulas, and the fact that σ cannot introduce a binding for n into G if it is appropriate for the instance of the conclusion sequent resulting from applying θ , we see that this would be the case if $G[\theta][\sigma] \vdash_{\Sigma} B[\theta]$ type has a derivation. To show that this is so, we use Lemma 4.70. The wellformedness of the conclusion sequent ensures that the parameters of the call to *Decompose* satisfy the requirements of the lemma. The substitutions θ and σ also meet the conditions required of them by clause 2: they are closed substitutions with domains identical to Ψ and Ξ^- , respectively, $G[\theta][\sigma]$ and $B[\theta]$ must obviously be defined, and $\vdash_{\Sigma} G[\theta][\sigma]$ ctx must have a derivation because $\{G \vdash M : A\}^{Ann}[\theta][\sigma]$ is valid by assumption. Thus, the lemma would allow us to conclude that $G[\theta][\sigma] \vdash_{\Sigma} B[\theta]$ type has a derivation if $F'[\theta][\sigma]$ is valid for every $\langle \mathbb{N}', \Xi', F' \rangle \in \text{Decompose}(\mathbb{N}, \Xi, G, B)$. But this must be the case. These are the goal formulas of closed instances of the premise sequents that we have assumed to be valid. Further, the assumption formulas of these sequents are $\Omega[\theta][\sigma]$, which are valid with respect to Υ by assumption. It follows that each $F'[\theta][\sigma]$ must be valid with respect to Υ . Since these formulas do not contain annotations, it is easy to see that they must be valid without qualification.

It remains only to dispense with the assumption concerning $\text{supp}(\theta)$ and $\text{supp}(\sigma)$. If these substitutions do not satisfy the assumption at the outset, then we may consider their variants θ' and σ' that are obtained through a permutation that swaps nominal constants in the set $\overline{\mathbb{N}}$ with ones that do not appear in $\overline{\mathbb{N}} \cup \mathbb{N} \cup \text{supp}(\theta) \cup \text{supp}(\sigma)$. This permutation will leave the conclusion sequent unaffected and will ensure that $\text{supp}(\theta')$ and $\text{supp}(\sigma')$ are disjoint from $\overline{\mathbb{N}}$. We may then use the earlier argument with respect to θ' and σ' and conclude the proof by invoking Theorem 4.18 and a version of Theorem 4.19 relativized to validity with respect to height assignments that can be easily shown to be true. \square

5 An Example Development: Uniqueness of Typing

We illustrate the proof system that has been developed in Section 4 by showing how a formal proof can be provided within it for the type uniqueness property of the STLC, the running example in this paper. We have discussed how this property can be expressed within \mathcal{L}_{LF} in Section 3.3. To recall, it is the situation in which the LF signature parameterizing \mathcal{L}_{LF} is the one presented in Figure 2.7 and c denotes the context schema comprising the single block

$$\{t : o\}x : tm, y : of\ x\ t,$$

the formula expressing the property of interest is the following:

$$\begin{aligned} & \Pi \Gamma : c. \forall e : o. \forall t_1 : o. \forall t_2 : o. \forall d_1 : o. \forall d_2 : o. \\ & \quad \{\Gamma \vdash e : tm\} \supset \{\Gamma \vdash t_1 : tp\} \supset \{\Gamma \vdash t_2 : tp\} \supset \\ & \quad \{\Gamma \vdash d_1 : of\ e\ t_1\} \supset \{\Gamma \vdash d_2 : of\ e\ t_2\} \supset \exists d_3 : o. \{ \cdot \vdash d_3 : eq\ t_1\ t_2 \}. \end{aligned}$$

Following the earlier discussion, we will actually construct a proof for the sequent $\emptyset; \emptyset; \emptyset; \emptyset \longrightarrow F$ where F is the formula

$$\begin{aligned} & \Pi \Gamma : c. \forall e : o. \forall t_1 : o. \forall t_2 : o. \forall d_1 : o. \forall d_2 : o. \\ & \quad \{\Gamma \vdash e : tm\} \supset \{\Gamma \vdash t_1 : tp\} \supset \{\Gamma \vdash t_2 : tp\} \supset \\ & \quad \{\Gamma \vdash d_1 : of\ e\ t_1\} \supset \{\Gamma \vdash d_2 : of\ e\ t_2\} \supset \exists d_3 : o. \{\Gamma \vdash d_3 : eq\ t_1\ t_2\}. \end{aligned}$$

We can then obtain a proof from the sequent representing the formula that we really want to prove using a proof of the strengthening lemma for equality of types identified in Section 3.3. We will not discuss a proof of the strengthening lemma or how exactly the use of lemmas plays out within our proof system. Suffice it to say that the former should hold no mysteries after we have discussed the proof of the sequent $\emptyset; \emptyset; \emptyset; \emptyset \longrightarrow F$ and the latter involves the use of the *cut* rule in the manner that we would expect. Also, our discussion of proofs will be in a “backwards,” proof-construction style: given a target or goal sequent, we will identify a collection of new goal sequents from whose proofs we can construct a proof of the original one by using rules of the proof system.

To get to the details, the informal proof of the formula F was based on an induction of the height of the typing derivation associated with the formula $\{\Gamma \vdash d_1 : of\ e\ t_1\}$ that appears as an antecedent of an implication in F . We can mimic this process in the formal proof through the use of the *ind* rule, that would yield as a new goal the task of constructing a proof for the sequent $\emptyset; \emptyset; \emptyset; \{F^*\} \longrightarrow F^\circledast$; we use the notation F^* and F^\circledast to denote formulas that are identical to F except that the occurrence of the formula $\{\Gamma \vdash d_1 : of\ e\ t_1\}$ within them is annotated with a $*$ and an \circledast , respectively. We may now think of deriving this sequent by using a sequence of rules for introducing logical symbols into its goal formula, leaving us with the task of constructing a proof for the following sequent:

$$\begin{aligned} & \cdot; (e : o, t_1 : o, t_2 : o, d_1 : o, d_2 : o); \Gamma \uparrow \emptyset : c[\cdot]; \\ & \{F^*, \{\Gamma \vdash e : tm\}, \{\Gamma \vdash t_1 : tp\}, \{\Gamma \vdash t_2 : tp\}, \{\Gamma \vdash d_1 : of\ e\ t_1\}^\circledast, \{\Gamma \vdash d_2 : of\ e\ t_2\}\} \\ & \quad \longrightarrow \exists d_3 : o. \{\Gamma \vdash d_3 : eq\ t_1\ t_2\} \end{aligned}$$

At this point in the informal proof we used case analysis to identify the different ways in which the formula $\{\Gamma \vdash d_1 : of\ e\ t_1\}$ may have been derived. This effect is realized in our proof system by considering the application of the *atm-app-L* rule with respect to the assumption formula $\{\Gamma \vdash d_1 : of\ e\ t_1\}^\circledast$. Doing so leads to the conclusion that we can complete the proof if we can construct proofs for the following four sequents:

1. $(n : o, n_1 : o, n_2 : o); (t_2 : o, d_2 : o, r : o \rightarrow o, t : o, u : o, a : o \rightarrow o \rightarrow o); \Gamma \uparrow \emptyset : c[\cdot];$
 $\{F^*, \{\Gamma \vdash lam\ t\ (\lambda x. r\ x) : tm\}, \{\Gamma \vdash arr\ t\ u : tp\}, \{\Gamma \vdash t_2 : tp\},$
 $\{\Gamma \vdash d_2 : of\ (lam\ t\ (\lambda x. r\ x))\ t_2\}, \{G, n : tm \vdash r\ n : tm\}^*, \{\Gamma \vdash t : tp\}^*,$
 $\{\Gamma \vdash u : tp\}^*, \{G, n_1 : tm, n_2 : of\ n_1\ t \vdash a\ n_1\ n_2 : of\ (r\ n_1)\ u\}^*\}$
 $\longrightarrow \exists d_3 : o. \{\Gamma \vdash d_3 : eq\ (arr\ t\ u)\ t_2\}$
2. $\cdot; (t_1 : o, t_2 : o, d_2 : o, m : o, n : o, u : o, a_1 : o, a_2 : o); \Gamma \uparrow \emptyset : c[\cdot];$
 $\{F^*, \{\Gamma \vdash app\ m\ n : tm\}, \{\Gamma \vdash t_1 : tp\}, \{\Gamma \vdash t_2 : tp\},$
 $\{\Gamma \vdash d_2 : of\ (app\ m\ n)\ t_2\}, \{\Gamma \vdash m : tm\}^*, \{\Gamma \vdash n : tm\}^*, \{\Gamma \vdash t_1 : tp\}^*,$
 $\{\Gamma \vdash u : tp\}^*, \{\Gamma \vdash a_1 : of\ M\ (arr\ u\ t_1)\}^*, \{\Gamma \vdash a_2 : of\ n\ u\}^*\}$
 $\longrightarrow \exists d_3 : o. \{\Gamma \vdash d_3 : eq\ t_1\ t_2\}$
3. $\cdot; (t_2 : o, d_2 : o); \Gamma \uparrow \emptyset : c[\cdot];$
 $\{F^*, \{\Gamma \vdash empty : tm\}, \{\Gamma \vdash unit : tp\}, \{\Gamma \vdash t_2 : tp\}, \{\Gamma \vdash d_2 : of\ empty\ t_2\}\}$
 $\longrightarrow \exists d_3 : o. \{\Gamma \vdash d_3 : eq\ unit\ t_2\}$
4. $(n : o, n_1 : o); (t_1 : o \rightarrow o \rightarrow o, t_2 : o \rightarrow o \rightarrow o, d_2 : o \rightarrow o \rightarrow o);$
 $\Gamma \uparrow \emptyset : c[n : tm, n_1 : of\ n\ (t_1\ n\ n_1)];$

$$\{F^*, \{\Gamma \vdash n : tm\}, \{\Gamma \vdash t_1 n n_1 : tp\}, \{\Gamma \vdash t_2 n n_1 : tp\}, \{\Gamma \vdash d_2 n n_1 : of n (t_2 n n_1)\}\} \\ \longrightarrow \exists d_3 : o. \{\Gamma \vdash d_3 : eq (t_1 n n_1) (t_2 n n_1)\}$$

The first of these sequents covers all of the cases where the head of d_1 is the constant of_lam , the second where it is of_app , the third where it is of_empty , and the fourth where it is a bound variable that appears in the context that instantiates Γ .

In the first of these cases, the informal proof was based on considering the derivation of the main LF typing judgement associated with $\{\Gamma \vdash d_2 : of (lam t (\lambda x. r x)) t_2\}$. Given the structure of the type, such a derivation will exist only when the head of d_2 is the constant of_lam . In the proof system, this analysis is realized by considering the use of the $atm-app-L$ rule relative to this formula, leading to the conclusion that we could complete a proof for this subcase if we are able to provide a proof for the sequent

$$(n : o, n_1 : o, n_2 : o, n_3 : o, n_4 : o, n_5 : o); \\ (r : o \rightarrow o, t : o, u : o, a : o \rightarrow o \rightarrow o, u_2 : o, a_2 : o \rightarrow o \rightarrow o); \Gamma \uparrow \emptyset : c[\cdot]; \\ \{F^*, \{\Gamma \vdash lam t (\lambda x. r x) : tm\}, \{\Gamma \vdash arr t u : tp\}, \{\Gamma \vdash arr t u_2 : tp\}, \{\Gamma, n : tm \vdash r n : tm\}^*, \\ \{\Gamma \vdash t : tp\}^*, \{\Gamma \vdash u : tp\}^*, \{\Gamma, n_1 : tm, n_2 : of n_1 t \vdash a n_1 n_2 : of (r n_1) u\}^*, \\ \{\Gamma, n_3 : tm \vdash r n_3 : tm\}, \{\Gamma \vdash t : tp\}, \{\Gamma \vdash u_2 : tp\}, \\ \{\Gamma, n_4 : tm, n_5 : of n_1 t \vdash a_2 n_4 n_5 : of (r n_4) u_2\}\} \\ \longrightarrow \exists d_3 : o. \{\Gamma \vdash d_3 : eq (arr t u) (arr t u_2)\}$$

The informal proof now uses the induction hypothesis based on the strictly smaller derivation that is assumed to exist for the main typing judgement associated with the assumption formula $\{\Gamma, n_1 : tm, n_2 : of n_1 t \vdash a n_1 n_2 : of (r n_1) u\}$. In the formal proof, this effect is realized by using the formula F^* together with the other assumption formulas in the sequent. Specifically, we would consider a sequence of uses of the $\Pi-L$ and $\forall-L$, $\supset-L, id$ and wk rules that would leave us needing to prove a sequent in which the consequent of a relevant instance of the formula F has been added to the set of assumption formulas. To actually carry out this process, it would be necessary to check that the context expression $\Gamma, n_1 : tm, n_2 : of n_1 t$ has the structure needed to satisfy the context schema represented by c , and we would need to instantiate the quantified variable e in F with $(r n_1)$, t_1 with u_1 , t_2 with u_2 , d_1 with $(a n_1 n_2)$, and d_2 with $(a_2 n_1 n_2)$. We leave it to the reader to check that these requirements can be satisfied and to elaborate the process by which the task can be reduced to that of constructing a proof for the sequent

$$(n : o, n_1 : o, n_2 : o, n_3 : o, n_4 : o, n_5 : o); \\ (r : o \rightarrow o, t : o, u : o, a : o \rightarrow o \rightarrow o, u_2 : o, a_2 : o \rightarrow o \rightarrow o, d_3 : o \rightarrow o \rightarrow o \rightarrow o \rightarrow o \rightarrow o); \\ \Gamma \uparrow \emptyset : c[\cdot]; \\ \{F^*, \{\Gamma \vdash lam t (\lambda x. r x) : tm\}, \{\Gamma \vdash arr t u : tp\}, \{\Gamma \vdash arr t u_2 : tp\}, \{\Gamma, n : tm \vdash r n : tm\}^*, \\ \{\Gamma \vdash t : tp\}^*, \{\Gamma \vdash u : tp\}^*, \{\Gamma, n_1 : tm, n_2 : of n_1 t \vdash a n_1 n_2 : of (r n_1) u\}^*, \\ \{\Gamma, n_3 : tm \vdash r n_3 : tm\}, \{\Gamma \vdash t : tp\}, \{\Gamma \vdash u_2 : tp\}, \\ \{\Gamma, n_4 : tm, n_5 : of n_1 t \vdash a_2 n_4 n_5 : of (r n_4) u_2\}, \\ \{\Gamma, n_1 : tm, n_2 : of n_1 t \vdash d_3 n n_1 n_2 n_3 n_4 n_5 : eq u u_2\}\} \\ \longrightarrow \exists d_3 : o. \{\Gamma \vdash d_3 : eq (arr t u) (arr t u_2)\}$$

The key observation now is that the main typing judgement corresponding to the new assumption formula in this sequent is derivable only if the head of d_3 is an instance of $refl$ and u and u_2 are instantiated with identical types. This reasoning step is reflected in the formal proof in the employment, again, of the $atm-app-L$ with respect to this assumption formula, which results in the conclusion that it suffices to construct a proof for the sequent

$$\begin{aligned}
& (n : o, n_1 : o, n_2 : o, n_3 : o, n_4 : o, n_5 : o); \\
& (r : o \rightarrow o, t : o, u : o, a : o \rightarrow o \rightarrow o, a_2 : o \rightarrow o \rightarrow o); \Gamma \uparrow \emptyset : c[\cdot]; \\
& \{F^*, \{\Gamma \vdash \text{lam } t (\lambda x. r \ x) : tm\}, \{\Gamma \vdash \text{arr } t \ u : tp\}, \{\Gamma \vdash \text{arr } t \ u_2 : tp\}, \{\Gamma, n : tm \vdash r \ n : tm\}^*, \\
& \quad \{\Gamma \vdash t : tp\}^*, \{\Gamma \vdash u : tp\}^*, \{\Gamma, n_1 : tm, n_2 : \text{of } n_1 \ t \vdash a \ n_1 \ n_2 : \text{of } (r \ n_1) \ u\}^*, \\
& \quad \{\Gamma, n_3 : tm \vdash r \ n_3 : tm\}, \{\Gamma \vdash t : tp\}, \{\Gamma \vdash u : tp\}, \\
& \quad \{\Gamma, n_4 : tm, n_5 : \text{of } n_1 \ t \vdash a_2 \ n_4 \ n_5 : \text{of } (r \ n_4) \ u\}, \{\Gamma, n_1 : tm, n_2 : \text{of } n_1 \ t \vdash u : tp\}\} \\
& \quad \longrightarrow \exists d_3 : o. \{\Gamma \vdash d_3 : \text{eq } (\text{arr } t \ u) (\text{arr } t \ u)\}
\end{aligned}$$

It is easy to see that this sequent would be derivable if the goal formula in it is replaced with the one obtained by instantiating the existentially quantified variable d_3 with the term $(\text{refl } (\text{arr } t \ u))$. However, that suffices to complete the proof of this subcase: we simply use the \exists - R below that derivation to get one for the sequent that is really of interest.

The analysis in the three remaining cases all begin with a step similar to that in the case just considered, that of analyzing the assumption formula corresponding to the other posited STLC typing derivation for the term. Since the first derivation has fixed the top level structure of the term, this analysis will identify only one possibility, i.e., that the final step in that derivation is the same as that in the analysis of the first derivation. In the case corresponding to sequent identified by the label 2, the argument will again employ the inductive hypothesis by identifying a suitable instance of the formula F^* . In each of the cases, the conclusion of the proof for that case will involve the instantiation of the existential goal, and decomposition of the resulting atomic goal formula. We leave it to the reader to fill out the details.

6 Related Work and Conclusion

We have described a logic called \mathcal{L}_{LF} in this paper that provides a means for formalizing properties of LF specifications. The atomic formulas in this logic encode typing judgements relative to an LF signature that parameterizes the logic. Propositional connectives can be used over these formulas to describe more complex properties and the logic also supports quantification over variables denoting LF contexts and terms. We have complemented the logic by presenting a proof system that can be used to establish the validity of formulas that express properties of the specified systems. In addition to providing rules that interpret the logical symbols, the proof system builds in the capability to carry out a case-analysis style reasoning over typing judgements in LF and to reason inductively based on the heights of LF derivations. The logic also enables the encoding of generic mechanisms for reasoning about LF derivability, a fact that we have illustrated by describing proof rules that embody a collection of commonly used meta-theorems about LF. In work that builds on the results of this paper, we have developed a proof assistant called Adelfa that provides mechanized support for the proof system and that we have used to demonstrate its effectiveness in reasoning [16, 18].

There have been other efforts directed at building a capability for reasoning about LF specifications. However, the approaches taken within these efforts have differed from the one that we have explored in this paper. One prominent approach is that embedded in what might be called the “Twelf family” of systems. The first realization of this approach is the Twelf system itself [10]. This system uses LF once again to formalize properties of systems described in it. More specifically, the properties that are of interest are described by other LF types that have a functional structure. The validity of these properties is then demonstrated by constructing inhabitants of the types that represent them and exhibiting the totality of these inhabitants as functions via means that are external to the encoding calculus. This approach has achieved much success but it is also limited by the fact that the property that need to be expressed must be transformed into a form that is encodable by a function type, i.e., by formulas that have a $\forall\exists$ quantifier structure. Another

drawback of the approach is that there isn't an explicit proof to be extracted at the end of a development for a property that has ostensibly been demonstrated. The logic M_2^+ has been enunciated towards mitigating this issue [15], and it appears possible to mechanically relate the “reasoning” embodied in a Twelf development to a proof in this logic [20]. While some of the proof rules in M_2^+ bear a resemblance to the ones in \mathcal{L}_{LF} , there are significant differences in the specific treatment especially of inductive reasoning.

The Twelf system and the M_2^+ logic differ in another marked way from the logic that we have described in this paper: they do not provide an explicit means for quantifying over contexts. It is possible to parameterize a development by a context description, but it is one fixed context that then permeates the development. As an example, it is not possible to express the strengthening lemma pertaining to the equality of types in the STLC that we discussed in Section 3.3. The Beluga system [13] alleviates this problem by using a richer version of type theory that allows for an explicit treatment of contexts as its basis [9]. Beluga is based on a computational view of reasoning that is similar in many respects to the philosophy underlying Twelf: one writes dependently-typed recursive functions and the type system ensures that the admitted functions are ones that are total and hence embody “proofs” of the properties expressed by the types. This system is more expressive than Twelf because the type system is more expressive, but the structure of the formulas that encode properties is similarly limited. A recent development that appears related to this line of work is that of COCON, a Martin-Löf style type theory that embeds LF within a rich dependently typed calculus that supports recursion [12]. It would be interesting to compare the reasoning capabilities that result from this kind of a combination with what is possible to achieve with an approach like ours. We leave a further exploration of this issue to future work.

Another approach that has been explored for reasoning about LF specifications is based on their translation to a predicate logic form. A particular exemplar of this approach is one that translates LF specifications into specifications in the logic of hereditary Harrop formulas [8], to then be reasoned about using the Abella system [17]. This approach has the virtue that benefit can be derived from any of the (generic) reasoning capabilities that have been developed for the host system. In the mentioned example, several such advantages are derived from the expressiveness of the logic underlying Abella [4]: it is possible to define relations between contexts, to treat binding notions explicitly in the reasoning process through the ∇ -quantifier, and to use inductive (and co-inductive) definitions in the reasoning logic. There are, however, some drawbacks to the translation-based approach. Perhaps the most significant problem is that the proof steps that can be taken under it are determined by the logic of the host system, and this may allow for more possibilities than are sensible in the LF context. One way to overcome this difficulty is to design macro proof steps that capture the natural process of reasoning about LF specifications. In this respect, one important outcome of the work that we have described here, especially for the structure that we have developed for case analysis, might be an understanding of how one might build within proof assistants such as Abella a targeted capability for reasoning about LF specifications.

Acknowledgements

This paper is based upon work supported by the National Science Foundation under Grant No. CCF-1617771. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

References

- [1] The Adelfa system. <http://adelfa.cs.umn.edu/>.
- [2] D. Baelde, K. Chaudhuri, A. Gacek, D. Miller, G. Nadathur, A. Tiu, and Y. Wang. Abella: A system for reasoning about relational specifications. *Journal of Formalized Reasoning*, 7(2), 2014.
- [3] A. Gacek. *A Framework for Specifying, Prototyping, and Reasoning about Computational Systems*. PhD thesis, University of Minnesota, 2009.
- [4] A. Gacek, D. Miller, and G. Nadathur. Nominal abstraction. *Information and Computation*, 209(1):48–73, 2011.
- [5] R. Harper, F. Honsell, and G. Plotkin. A framework for defining logics. *Journal of the ACM*, 40(1):143–184, 1993.
- [6] R. Harper and D. R. Licata. Mechanizing metatheory in a logical framework. *Journal of Functional Programming*, 17(4–5):613–673, July 2007.
- [7] D. Miller. Unification under a mixed prefix. *Journal of Symbolic Computation*, 14(4):321–358, 1992.
- [8] D. Miller and G. Nadathur. *Programming with Higher-Order Logic*. Cambridge University Press, June 2012.
- [9] A. Nanevski, F. Pfenning, and B. Pientka. Contextual model type theory. *ACM Trans. on Computational Logic*, 9(3):1–49, 2008.
- [10] F. Pfenning and C. Schürmann. System description: Twelf — A meta-logical framework for deductive systems. In H. Ganzinger, editor, *16th Conf. on Automated Deduction (CADE)*, number 1632 in LNAI, pages 202–206, Trento, 1999. Springer.
- [11] F. Pfenning and C. Schürmann. *Twelf User’s Guide*, Dec. 2002. Available from <http://www.cs.cmu.edu/~twelf/guide-1-4>.
- [12] B. Pientka, A. Abel, F. Ferreira, D. Thibodeau, and R. Zucchini. COCON: Computation in contextual type theory. Available from <http://arxiv.org/abs/1901.03378>, 2019.
- [13] B. Pientka and J. Dunfield. Beluga: A framework for programming and reasoning with deductive systems (system description). In J. Giesl and R. Hähnle, editors, *Fifth International Joint Conference on Automated Reasoning*, number 6173 in LNCS, pages 15–21, 2010.
- [14] B. C. Pierce. *Types and Programming Languages*. MIT Press, 2002.
- [15] C. Schürmann. *Automating the Meta Theory of Deductive Systems*. PhD thesis, Carnegie Mellon University, Oct. 2000.
- [16] M. Southern. *A Framework for Reasoning About LF Specifications*. PhD thesis, University of Minnesota, May 2021.

- [17] M. Southern and K. Chaudhuri. A two-level logic approach to reasoning about typed specification languages. In V. Raman and S. P. Suresh, editors, *34th International Conference on Foundation of Software Technology and Theoretical Computer Science (FSTTCS 2014)*, volume 29 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 557–569, Dagstuhl, Germany, 2014.
- [18] M. Southern and G. Nadathur. Adelfa: A system for reasoning about LF specifications. In *International Workshop on Logical Frameworks and Meta-Languages: Theory and Practice (LFMTP)*, 2021.
- [19] A. Tiu. A logic for reasoning about generic judgments. In A. Momigliano and B. Pientka, editors, *Int. Workshop on Logical Frameworks and Meta-Languages: Theory and Practice (LFMTP'06)*, volume 173 of *ENTCS*, pages 3–18, 2006.
- [20] Y. Wang and G. Nadathur. Towards extracting explicit proofs from totality checking in Twelf. In *LFMTP 2013 - Proceedings of the 2013 ACM SIGPLAN Workshop on Logical Frameworks and Meta-Languages*, pages 55–66, Oct. 2013.
- [21] K. Watkins, I. Cervesato, F. Pfenning, and D. Walker. A concurrent logical framework I: Judgments and properties. Technical Report CMU-CS-02-101, Carnegie Mellon University, 2003. Revised, May 2003.