

Local available quantum correlations of X states: The symmetric and anti-symmetric cases

Hermann L. Albrecht Q.,^{1,*} David M. Bellorin R.,¹ and Douglas F. Mundarain²

¹*Departamento de Física, Universidad Simón Bolívar, AP 89000, Caracas 1080, Venezuela.*

²*Departamento de Física, Universidad Católica del Norte, Casilla, 1280 Antofagasta, Chile.*

Abstract

Local available quantum correlations (LAQC), as defined by Mundarain et al., are analyzed for two subsets of 2-qubit X states. We start by studying X-states that are symmetric under the exchange of subsystems, that is, those with the same non-null local Bloch vector. We also analyze the subset of states that are anti-symmetric under subsystem exchange, that is, those that have non-null local Bloch vectors with an equal magnitude but opposite direction. We present various examples and compare the obtained results to concurrence as an entanglement measure and with quantum discord. We have also included markovian decoherence, with the analysis of amplitude damping decoherence for Werner states. As was previously observed for depolarization and phase damping decoherence, LAQC did not exhibit sudden death behavior for Werner states under amplitude damping decoherence.

I. INTRODUCTION

Quantum correlations and their quantification are of central importance in Quantum Information Theory (QIT). Entanglement [1] was seen as the main quantum resource until the development of Quantum Discord [2, 3] in 2001. Since then, the establishment of other quantum correlations and their quantifiers [4] has become a very active field of research, both within Information Theory and for their use in Quantum Computation and other applied specialties.

Local measurements are of the utmost importance to properly define correlations, as these latter must quantify the ability of a local observer to infer the results of another one from his results. This was shown to be the case in the abovementioned Quantum Discord

$$D_A(\rho_{AB}) \equiv \min_{\{\Pi_i^A\}} \{I(\rho_{AB}) - I[(\Pi^A \otimes \mathbb{1}_2)\rho_{AB}]\}, \quad (1)$$

which is based on comparing the quantum Mutual Information, defined for the original state ρ_{AB} as

$$I(\rho_{AB}) \equiv S(\rho_A) + S(\rho_B) - S(\rho_{AB}), \quad (2)$$

where ρ_A and ρ_B are the reduced operators, i.e. marginals, with a corresponding postmeasurement state in the absence of readout, where the measurement is performed locally on subsystem A. This latter state is referred to as a classical-quantum (or A-classical) state,

$$\begin{aligned} \rho_{AB}^{cq} &= \sum_i p_i |i\rangle\langle i| \otimes \rho_B^i \\ &= \sum_i p_i \Pi_i^{(A)} \otimes \rho_B^i, \end{aligned} \quad (3)$$

and all such 2-qubit states constitute a set, denoted by Ω_o . An analogous set, denoted by Ω'_o , is readily defined

when the local measurement is performed on the B subsystem. That is,

$$D_B(\rho_{AB}) \equiv \min_{\{\Pi_i^B\}} \{I(\rho_{AB}) - I[(\mathbb{1}_2 \otimes \Pi^B)\rho_{AB}]\}. \quad (4)$$

Quantum correlations defined using the set Ω_o (or Ω'_o) are called Discords and are in general not symmetric so that $D_A(\rho_{AB}) \neq D_B(\rho_{AB})$.

Another set of quantum correlations and quantifiers was developed by considering classical states, that is, postmeasurement states in the absence of readout whose measurements were performed locally on both subsystems:

$$\rho_{AB}^c = \sum_{i,j} p_{ij} \Pi_i^{(A)} \otimes \Pi_j^{(B)}. \quad (5)$$

A special subset of such states are product states,

$$\rho_{AB}^\Pi = \sum_{i,j} p_i \Pi_i^{(A)} \otimes p_j \Pi_j^{(B)} = \rho^{(A)} \otimes \rho^{(B)}, \quad (6)$$

also referred to as uncorrelated states. For these, the coefficients p_{ij} in eq. (5) need to be factorizable, $p_{ij} = p_i p_j$. We denote the sets of classical and product 2-qubit states as Ω_c and Ω_p , respectively.

Wu et al. considered general quantum correlations defined in terms of local bipartite measurements in [5] and, in a brief final appendix, outlined symmetric quantum correlations in relation to mutual information. Mundarain et al. in [6] developed the so-called Local Available Quantum Correlations (LAQC) by focusing on a slightly different version of those symmetric correlations, defining them in terms of mutual information of local bipartite measurements on the so-called complementary basis of a previously determined optimal computational basis.

This work is focused on determining the LAQC's quantifier for two particular sets of the so-called X states [7]

$$\rho^X = \begin{pmatrix} \rho_{11} & 0 & 0 & \rho_{14} \\ 0 & \rho_{22} & \rho_{23} & 0 \\ 0 & \rho_{32} & \rho_{33} & 0 \\ \rho_{41} & 0 & 0 & \rho_{44} \end{pmatrix}. \quad (7)$$

* Corresponding Author: albrecht@usb.ve

This family of 2-qubit states has been studied extensively [8] and are a widely used set of 2-qubit density matrices in QIT. That is because any arbitrary 2-qubit state ρ_{AB} can be mapped to such ρ^X , preserving its main characteristics, e.g. quantum correlations [9, 10].

Experimentally, this type of 2-qubit states has been achieved with cold trapped atoms and ions [11] and in non-linear crystals [12]. Also, X states appear naturally in spin chain systems when the reduced matrix of two neighboring spins is studied [13, 14] and any pure 2-qubit spin- $\frac{1}{2}$ state evolves to an X state via decoherence due to magnetic field fluctuations [8].

Concurrence is a *bona fide* entanglement measure introduced by Wootters [15] as

$$\mathcal{C} \equiv \max\{0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4\}, \quad (8)$$

where $\{\lambda_i\}$ are the decreasing ordered eigenvalues of $\mathbf{R} = \sqrt{\sqrt{\rho} \tilde{\rho} \sqrt{\rho}}$, with $\tilde{\rho} = (\sigma_y \otimes \sigma_y) \rho^* (\sigma_y \otimes \sigma_y)$, ρ^* the complex conjugate of ρ , and σ_y the corresponding Pauli matrix. For X states, a direct calculation shows that Concurrence takes a much simpler expression [7]:

$$\mathcal{C}_X = \frac{1}{2} \max\{0, \mathcal{C}_1, \mathcal{C}_2\} \quad (9)$$

where

$$\mathcal{C}_1 \equiv 2(|\rho_{14}| - \sqrt{\rho_{22} \rho_{33}}), \quad \mathcal{C}_2 \equiv 2(|\rho_{23}| - \sqrt{\rho_{11} \rho_{44}}).$$

A closed analytical expression of the quantum discord of X states has not been achieved yet. In 2010, Ali et al. [16] claimed to have found such an expression, but several counterexamples have been found since then that shows that their formula is not correct [17–19]. For a subgroup of this family of 2-qubit states, where $\rho_{22} = \rho_{33}$, some alternatives have been proposed [8]. In this article, we use the results of Li et al. [20], which we briefly present in an appendix.

We start by briefly reviewing the procedure for determining the quantifier of local available quantum correlations [6]. Then, we proceed to study its calculation for X states in two distinct subsets: when there is the symmetry under exchange of subsystems, that is when both subsystems have the same non-null local Bloch vector, and when the states are anti-symmetric under subsystem exchange, that is when both subsystems have local Bloch vectors with the same magnitude but opposite direction. We analyze various examples and include the amplitude damping decoherence of a Werner state [21].

II. LOCAL AVAILABLE QUANTUM CORRELATIONS OF 2-QUBITS

In [6], the starting point for Mundarain et al. is that a density operator ρ can always be written in terms of

different basis. For 2-qubits, we have that

$$\begin{aligned} \rho_{AB} &= \sum_{klmn} \rho_{kl}^{mn} |km\rangle\langle ln| \\ &= \sum_{ijpq} R_{ip}^{jq} |B(i, j)\rangle\langle B(p, q)| \end{aligned} \quad (10)$$

where $k, l, m, n \in \{0, 1\}$, $\{|km\rangle\}$ is the standard computational basis, that is the basis of eigenvectors of σ_z , which is local, and $\{|B(i, j)\rangle\}$ is another local basis that is equivalent under local unitary transformations to the former one. This equivalence can be stated as $|B(i, j)\rangle = \mathbb{U}_A \otimes \mathbb{U}_B |ij\rangle$, with $\mathbb{U}_A, \mathbb{U}_B \in \mathbf{U}(2)$. Any such basis can be used as the new computational one, where we now have the basis of eigenvector of $\sigma_{\hat{\mathbf{u}}} \equiv \vec{\sigma} \cdot \hat{\mathbf{u}}$, with $\vec{\sigma}$ the vector whose components are the Pauli matrices and $\hat{\mathbf{u}} \in \mathbb{E}^3$ is a generic unitary vector. The choosing of such $\hat{\mathbf{u}}$ can depend on various conditions and/or requirements of the system at hand.

For classical states, it is said that there exists a local basis for which ρ_{AB}^c (5) is diagonal. Therefore, one can define $X_\rho \in \Omega_c$ as the classical state induced by a measurement which minimizes

$$S(\rho_{AB} || X_\rho) = \min_{\Omega_c} S(\rho_{AB} || \chi_\rho), \quad (11)$$

where $S(\rho || \chi) = -\text{Tr}(\rho \log_2 \chi) - S(\rho)$ is the relative entropy and

$$\begin{aligned} \chi_\rho &= \sum_{ij} R_{ij} |B(i, j)\rangle\langle B(i, j)|, \\ R_{ij} &= \langle B(i, j) | \rho_{AB} | B(i, j) \rangle. \end{aligned} \quad (12)$$

The minimization in (11) is equivalent to determining the coefficients $\{R_{ij}^{opt}\}$ when $\chi_\rho = X_\rho$. Such coefficients are associated with the optimal basis $\{|B(i, j)\rangle^{opt}\}$ which will then serve as the new computational basis, in whose terms local available quantum correlations are then defined.

To determine the optimal basis, we start by defining the most general orthonormal basis for each subsystem:

$$\begin{aligned} |\mu_0^{(n)}\rangle &= \cos\left(\frac{\theta_n}{2}\right) |0^{(n)}\rangle + \sin\left(\frac{\theta_n}{2}\right) \exp^{i\phi_n} |1^{(n)}\rangle \\ |\mu_1^{(n)}\rangle &= -\sin\left(\frac{\theta_n}{2}\right) |0^{(n)}\rangle + \cos\left(\frac{\theta_n}{2}\right) \exp^{i\phi_n} |1^{(n)}\rangle, \end{aligned} \quad (13)$$

where $n = 1$ denotes subsystem A and $n = 2$, subsystem B. The classical correlations quantifier defined in this context in [6] is given by

$$\mathcal{C}(\rho) \equiv S(X_\rho || \Pi_{X_\rho}) = I(X_\rho) \quad (14)$$

where Π_{X_ρ} is the product state nearest to X_ρ and this relative entropy can be written as the Mutual Information of state X_ρ , as shown by Modi et al. [22]. Since the

mutual information (2) may be written as

$$I(\rho_{AB}) = \sum_{i,j} P_{\theta,\phi}(i_A, j_B) \times \log_2 \left[\frac{P_{\theta,\phi}(i_A, j_B)}{P_{\theta,\phi}(i_A)P_{\theta,\phi}(j_B)} \right], \quad (15)$$

where $P_{\theta,\phi}(i_A, j_B) = \langle \mu_i^{(1)}, \mu_j^{(2)} | \rho_{AB} | \mu_i^{(1)}, \mu_j^{(2)} \rangle$ are the probability distributions corresponding to ρ_{AB} and $P_{\theta,\phi}(i_A) = \langle \mu_i^{(1)} | \rho_A | \mu_i^{(1)} \rangle$, $P_{\theta,\phi}(j_B) = \langle \mu_j^{(2)} | \rho_B | \mu_j^{(2)} \rangle$ the ones corresponding to its marginals ρ_A and ρ_B , the required minimization of the relative entropy (11) yields a minima for the classical correlations quantifier defined in (14). It is straightforward to realize that this probability distributions are directly related to the $\{R_{ij}^{opt}\}$ coefficients when $\left\{ \left| \mu_i^{(1)}, \mu_j^{(2)} \right\rangle \right\}$ is the optimal computational basis.

Once this new computational basis $\left\{ |0^{(m)}\rangle_{opt}, |1^{(m)}\rangle_{opt} \right\}$ has been determined, the state ρ_{AB} is rewritten and the complementary basis defined in terms of a new general orthonormal basis:

$$\begin{aligned} |u_0^{(m)}\rangle &= \frac{1}{\sqrt{2}} \left(|0^{(m)}\rangle_{opt} + \exp^{i\Phi_m} |1^{(m)}\rangle_{opt} \right), \\ |u_1^{(m)}\rangle &= \frac{1}{\sqrt{2}} \left(|0^{(m)}\rangle_{opt} - \exp^{i\Phi_m} |1^{(m)}\rangle_{opt} \right). \end{aligned} \quad (16)$$

In this generic basis, the corresponding probability distributions,

$$P(i_A, j_B, \Phi_1, \Phi_2) = \langle u_i^A, u_j^B | \rho_{AB} | u_i^A, u_j^B \rangle, \quad (17)$$

and the marginal probability distributions are determined. The maximization of $I(\Phi_1, \Phi_2)$ (15) corresponds to the LAQC quantifier:

$$\mathcal{L}(\rho_{AB}) \equiv \max_{\{\Phi_1, \Phi_2\}} I(\Phi_1, \Phi_2). \quad (18)$$

III. LAQC OF X STATES

We start by using the Bloch representation of 2-qubit states, also referred to as the Fano form or Fano-Bloch representation [23], as it allows to directly describe the so-called canonical X states in terms of five independent real parameters:

$$\rho^X = \frac{1}{4} \left(\mathbb{1}_4 + x_3 \sigma_3 \otimes \mathbb{1}_2 + \mathbb{1}_2 \otimes y_3 \sigma_3 + \sum_{n=1}^3 T_n \sigma_n \otimes \sigma_n \right), \quad (19)$$

where σ_n are the Pauli matrices, $x_3 = \text{Tr}[\rho(\sigma_3 \otimes \mathbb{1}_2)]$, $y_3 = \text{Tr}[\rho(\mathbb{1}_2 \otimes \sigma_3)]$ y $T_n = \text{Tr}[\rho(\sigma_n \otimes \sigma_n)]$.

An important set of (19) is comprised of those for which their local Bloch vectors are of equal magnitude ($|x_3| = |y_3|$). Among these, Bell Diagonal states [24] constitute a special subset where $|x_3| = |y_3| = 0$ and their LAQC were already analyzed in [25]. For $|x_3| = |y_3| \neq 0$, a further classification will prove useful: symmetric and antisymmetric X states, where $x_3 = y_3$ and $x_3 = -y_3$, respectively.

It is also worth noting that single parameter mixed states between a Bell state $\{|\psi^\pm\rangle, |\phi^\pm\rangle\}$ and an element of the computational basis $\{|i, j\rangle\}$ belong to this set of X states. These can be thought of as analogous to what Werner states [21] are to Bell Diagonal ones. That is, as a one-parameter subset of the latter.

A. Symmetric X States

The set of symmetric X states ρ^{X_s} is invariant under subsystem exchange $\mathbf{A} \leftrightarrow \mathbf{B}$ since their local Bloch parameters are equal, $x_3 = y_3$. Such states arise when studying amplitude decoherence of Werner states. They also turn out as the ground state within the approximation of nanoelectric LC-circuits as two coupled harmonic oscillators presented in [26].

We start our analysis by determining the previously defined coefficients $R_{ij}(\theta_1, \theta_2, \phi_1, \phi_2)$ (12), obtaining

$$\begin{aligned} R_{00} &= \frac{1}{4} + \frac{1}{4}(\cos \theta_1 + \cos \theta_2)x_3 \\ &\quad + \Lambda \sum_{m=0}^1 \cos[\phi_1 + (-1)^m \phi_2][T_1 - (-1)^m T_2] \\ &\quad + \frac{1}{4} \cos \theta_1 \cos \theta_2 T_3, \\ R_{01} &= \frac{1}{4} + \frac{1}{4}(\cos \theta_1 - \cos \theta_2)x_3 \\ &\quad - \Lambda \sum_{m=0}^1 \cos[\phi_1 + (-1)^m \phi_2][T_1 - (-1)^m T_2] \\ &\quad - \frac{1}{4} \cos \theta_1 \cos \theta_2 T_3, \\ R_{10} &= \frac{1}{4} - \frac{1}{4}(\cos \theta_1 - \cos \theta_2)x_3 \\ &\quad - \Lambda \sum_{m=0}^1 \cos[\phi_1 + (-1)^m \phi_2][T_1 - (-1)^m T_2] \\ &\quad - \frac{1}{4} \cos \theta_1 \cos \theta_2 T_3, \\ R_{11} &= \frac{1}{4} - \frac{1}{4}(\cos \theta_1 + \cos \theta_2)x_3 \\ &\quad + \Lambda \sum_{m=0}^1 \cos[\phi_1 + (-1)^m \phi_2][T_1 - (-1)^m T_2] \\ &\quad + \frac{1}{4} \cos \theta_1 \cos \theta_2 T_3. \end{aligned} \quad (20)$$

where

$$\Lambda \equiv \frac{1}{2} \prod_{i=1}^2 \sin\left(\frac{\theta_i}{2}\right) \cos\left(\frac{\theta_i}{2}\right). \quad (21)$$

Since symmetric X states are invariant under subsystem exchange $\mathbf{A} \leftrightarrow \mathbf{B}$, the angles θ_i and ϕ_i must respect this symmetry in R_{ij} (20). Therefore, only two angles are sufficient to determine the optimal computational base. By defining $\theta_1 = \theta_2 = \theta$ and $\phi_1 = \phi_2 = \phi$ as the respective common angles, the previous expressions are readily simplified:

$$\begin{aligned} R_{00}(\theta, \phi) &= \frac{1}{4} [1 + 2 \cos \theta x_3 + \sin^2 \theta (\cos^2 \phi T_1 \\ &\quad + \sin^2 \phi T_2) + \cos^2 \theta T_3] \\ R_{01}(\theta, \phi) &= \frac{1}{4} [1 - \sin^2 \theta (\cos^2 \phi T_1 + \sin^2 \phi T_2) \\ &\quad - \cos^2 \theta T_3] = R_{10}(\theta, \phi) \\ R_{11}(\theta, \phi) &= \frac{1}{4} [1 - 2 \cos \theta x_3 + \sin^2 \theta (\cos^2 \phi T_1 \\ &\quad + \sin^2 \phi T_2) + \cos^2 \theta T_3] \end{aligned} \quad (22)$$

The corresponding coefficients $\{R_{i_A}(\theta, \phi), R_{j_B}(\theta, \phi)\}$ for the reduced matrices are given by

$$\begin{aligned} R_{0_A}(\theta) &= \frac{1}{2} (1 + x_3 \cos \theta) = R_{0_B}(\theta), \\ R_{1_A}(\theta) &= \frac{1}{2} (1 - x_3 \cos \theta) = R_{1_B}(\theta). \end{aligned} \quad (23)$$

The necessary minimization of the classical correlations quantifier (14) leads to three main cases to be considered:

- $\theta = 0, \phi \in [0, 2\pi]$:

$$\begin{aligned} R_{00}(\theta = 0, \phi) &= \frac{1}{4} [1 + T_3 + 2x_3], \\ R_{01}(\theta = 0, \phi) &= \frac{1}{4} [1 - T_3] = R_{10}, \\ R_{11}(\theta = 0, \phi) &= \frac{1}{4} [1 + T_3 - 2x_3]. \end{aligned} \quad (24)$$

- $\theta = \frac{\pi}{2}, \phi = 0$:

$$\begin{aligned} R_{00}\left(\theta = \frac{\pi}{2}, \phi = 0\right) &= \frac{1}{4} [1 + T_1] = R_{11}, \\ R_{01}\left(\theta = \frac{\pi}{2}, \phi = 0\right) &= \frac{1}{4} [1 - T_1] = R_{10}. \end{aligned} \quad (25)$$

- $\theta = \frac{\pi}{2}, \phi = \frac{\pi}{2}$:

$$\begin{aligned} R_{00}\left(\theta = \frac{\pi}{2}, \phi = \frac{\pi}{2}\right) &= \frac{1}{4} [1 + T_2] = R_{11}, \\ R_{01}\left(\theta = \frac{\pi}{2}, \phi = \frac{\pi}{2}\right) &= \frac{1}{4} [1 - T_2] = R_{10}. \end{aligned} \quad (26)$$

From the previous expressions, it is direct to see that ϕ is determined by the relation between T_1 and T_2 , since for $\theta = 0$ any $\phi \in [0, 2\pi]$ complies with the minimization. Lets define

$$\alpha_{\pm} \equiv T_3 \pm 2x_3 \quad \text{and} \quad T_m \equiv \min(|T_1|, |T_2|). \quad (27)$$

Next, we rewrite (24), (25), and (26) so we can focus our attention on θ :

- $\theta = 0$:

$$\begin{aligned} R_{00}(\theta = 0) &= \frac{1}{4} [1 + \alpha_+], \\ R_{01}(\theta = 0) &= \frac{1}{4} [1 - T_3] = R_{10}, \\ R_{11}(\theta = 0) &= \frac{1}{4} [1 + \alpha_-]. \end{aligned} \quad (28)$$

- $\theta = \frac{\pi}{2}$:

$$\begin{aligned} R_{00}\left(\theta = \frac{\pi}{2}\right) &= \frac{1}{4} [1 + T_m] = R_{11}, \\ R_{01}\left(\theta = \frac{\pi}{2}\right) &= \frac{1}{4} [1 - T_m] = R_{10}. \end{aligned} \quad (29)$$

The general expression for the quantifier of the classical correlations (14) is then given by

$$\mathcal{C}(\rho^{X_s}) = \sum_{i,j} R_{i,j}(\theta, \phi) \log_2 \left[\frac{R_{i,j}(\theta, \phi)}{R_i(\theta)R_j(\theta)} \right]. \quad (30)$$

The minimization can be achieved either by analyzing the coefficients $R_{i,j}(\theta, \phi)$ or the above expression. As we will see in our examples, graphical methods will be more efficient for states with one or two independent parameters since it will be straightforward to visualize the case that minimizes (30).

After the θ and ϕ angles have been determined, the state ρ^{X_s} is rewritten in the optimal computational basis as to determine the probability distributions $P^{(\theta, \phi)}(i_A, j_B, \Phi)$ (17) as well as the marginal distributions $P_A^{(\theta, \phi)}(i_A, \Phi)$ and $P_B^{(\theta, \phi)}(i_B, \Phi)$. For each of the above cases, the following results are obtained:

- $\theta = 0, \phi = 0$:

$$\begin{aligned} P^{(0,0)}(0, 0, \Phi) &= \frac{1}{4} (1 + T_1 \cos^2 \Phi + T_2 \sin^2 \Phi), \\ &= P^{(0,0)}(1, 1, \Phi), \\ P^{(0,0)}(1, 0, \Phi) &= \frac{1}{4} (1 - T_1 \cos^2 \Phi - T_2 \sin^2 \Phi), \\ &= P^{(0,0)}(0, 1, \Phi), \end{aligned} \quad (31)$$

and

$$\begin{aligned} P_A^{(0,0)}(0) &= P_A^{(0,0)}(1) = \frac{1}{2}, \\ P_B^{(0,0)}(0) &= P_B^{(0,0)}(1) = \frac{1}{2}. \end{aligned} \quad (32)$$

- $\theta = \frac{\pi}{2}, \phi = 0$:

$$\begin{aligned}
P^{(\frac{\pi}{2}, 0)}(0, 0, \Phi) &= \frac{1}{4}(1 + T_2 \sin^2 \Phi + T_3 \cos^2 \Phi \\
&\quad - 2x_3 \cos \Phi), \\
P^{(\frac{\pi}{2}, 0)}(1, 0, \Phi) &= \frac{1}{4}(1 - T_2 \sin^2 \Phi - T_3 \cos^2 \Phi), \\
&= P^{(\frac{\pi}{2}, 0)}(0, 1, \Phi), \\
P^{(\frac{\pi}{2}, 0)}(1, 1, \Phi) &= \frac{1}{4}(1 + T_2 \sin^2 \Phi + T_3 \cos^2 \Phi \\
&\quad + 2x_3 \cos \Phi).
\end{aligned} \tag{33}$$

and

$$\begin{aligned}
P_A^{(\frac{\pi}{2}, 0)}(0) &= P_B^{(\frac{\pi}{2}, 0)}(0) = \frac{1}{2}(1 - x_3 \cos \Phi), \\
P_A^{(\frac{\pi}{2}, 0)}(1) &= P_B^{(\frac{\pi}{2}, 0)}(1) = \frac{1}{2}(1 + x_3 \cos \Phi).
\end{aligned} \tag{34}$$

- $\theta = \frac{\pi}{2}, \phi = \frac{\pi}{2}$:

$$\begin{aligned}
P^{(\frac{\pi}{2}, \frac{\pi}{2})}(0, 0, \Phi) &= \frac{1}{4}(1 + T_1 \sin^2 \Phi + T_3 \cos^2 \Phi \\
&\quad - 2x_3 \cos \Phi), \\
P^{(\frac{\pi}{2}, \frac{\pi}{2})}(1, 0, \Phi) &= \frac{1}{4}(1 - T_1 \sin^2 \Phi - T_3 \cos^2 \Phi), \\
&= P^{(\frac{\pi}{2}, \frac{\pi}{2})}(0, 1, \Phi), \\
P^{(\frac{\pi}{2}, \frac{\pi}{2})}(1, 1, \Phi) &= \frac{1}{4}(1 + T_1 \sin^2 \Phi + T_3 \cos^2 \Phi \\
&\quad + 2x_3 \cos \Phi).
\end{aligned} \tag{35}$$

and again we have that

$$\begin{aligned}
P_A^{(\frac{\pi}{2}, \frac{\pi}{2})}(0) &= P_B^{(\frac{\pi}{2}, \frac{\pi}{2})}(0) = \frac{1}{2}(1 - x_3 \cos \Phi), \\
P_A^{(\frac{\pi}{2}, \frac{\pi}{2})}(1) &= P_B^{(\frac{\pi}{2}, \frac{\pi}{2})}(1) = \frac{1}{2}(1 + x_3 \cos \Phi).
\end{aligned} \tag{36}$$

To determine the LAQC's quantifier (18), the above expressions need to be maximized for $\Phi \in [0, 2\pi]$. We summarize next each case and present the corresponding expression of $\mathcal{L}(\rho^{X_s})$:

- $\theta = 0, \phi = 0$: The angle Φ depends on the maximum between $|T_1|$ and $|T_2|$. Therefore, analogous to T_m , we define

$$T_M \equiv \max(|T_1|, |T_2|) \tag{37}$$

and write the corresponding probability distributions:

$$\begin{aligned}
P^{(0, 0)}(0, 0) &= \frac{1}{4}(1 + T_M) = P^{(0, 0)}(1, 1), \\
P^{(0, 0)}(1, 0) &= \frac{1}{4}(1 - T_M) = P^{(0, 0)}(0, 1).
\end{aligned} \tag{38}$$

For $T_M = |T_1|$, $\Phi = \{0, \pi, 2\pi\}$, while for $T_M = |T_2|$, $\Phi = \{\frac{\pi}{2}, \frac{3\pi}{2}\}$. With these results, from eqs. (15) and (18) we have that

$$\begin{aligned}
\mathcal{L}(\rho^{X_s}) &= \frac{1 + T_M}{2} \log_2(1 + T_M) \\
&\quad + \frac{1 - T_M}{2} \log_2(1 - T_M).
\end{aligned} \tag{39}$$

- $\theta = \frac{\pi}{2}, \phi = 0$: Recalling that we previously defined $\alpha_{\pm} \equiv T_3 \pm 2x_3$ (27), we have two distinct cases, $\Phi = \{0, \pi, 2\pi\}$, and $\Phi = \{\frac{\pi}{2}, \frac{3\pi}{2}\}$:

– For $\Phi = \{0, 2\pi\}$ we have that

$$\begin{aligned}
P^{(\frac{\pi}{2}, 0)}(0, 0) &= \frac{1}{4}(1 + \alpha_-), \\
P^{(\frac{\pi}{2}, 0)}(1, 0) &= \frac{1}{4}(1 - T_3) \\
&= P^{(\frac{\pi}{2}, 0)}(0, 1), \\
P^{(\frac{\pi}{2}, 0)}(1, 1) &= \frac{1}{4}(1 + \alpha_+).
\end{aligned} \tag{40}$$

and

$$\begin{aligned}
P_A^{(\frac{\pi}{2}, 0)}(0) &= P_B^{(\frac{\pi}{2}, 0)}(0) = \frac{1}{2}(1 - x_3), \\
P_A^{(\frac{\pi}{2}, 0)}(1) &= P_B^{(\frac{\pi}{2}, 0)}(1) = \frac{1}{2}(1 + x_3).
\end{aligned} \tag{41}$$

For $\Phi = \pi$ we obtain equivalent expressions, with $P^{(\frac{\pi}{2}, 0)}(0, 0, \Phi = \pi) = P^{(\frac{\pi}{2}, 0)}(1, 1, \Phi = 0)$, $P^{(\frac{\pi}{2}, 0)}(1, 1, \Phi = \pi) = P^{(\frac{\pi}{2}, 0)}(0, 0, \Phi = 0)$, $P_A^{(\frac{\pi}{2}, 0)}(i_A, \Phi = 0) = P_B^{(\frac{\pi}{2}, 0)}(i_B, \Phi = \pi)$, and $P_B^{(\frac{\pi}{2}, 0)}(i_B, \Phi = 0) = P_A^{(\frac{\pi}{2}, 0)}(i_A, \Phi = \pi)$. With these results, we have that

$$\begin{aligned}
\mathcal{L}(\rho^{X_s}) &= \frac{1 + \alpha_-}{4} \log_2 \left[\frac{1 + \alpha_-}{(1 - x_3)^2} \right] \\
&\quad + \frac{1 - T_3}{2} \log_2 \left(\frac{1 - T_3}{1 - x_3^2} \right) \\
&\quad + \frac{1 + \alpha_+}{4} \log_2 \left[\frac{1 + \alpha_+}{(1 + x_3)^2} \right].
\end{aligned} \tag{42}$$

– For $\Phi = \frac{\pi}{2}, \frac{3\pi}{2}$ we have that

$$\begin{aligned}
P^{(\frac{\pi}{2}, 0)}(0, 0) &= \frac{1}{4}(1 + T_2) = P^{(\frac{\pi}{2}, 0)}(1, 1), \\
P^{(\frac{\pi}{2}, 0)}(1, 0) &= \frac{1}{4}(1 - T_2) = P^{(\frac{\pi}{2}, 0)}(0, 1).
\end{aligned} \tag{43}$$

and

$$\begin{aligned}
P_A^{(\frac{\pi}{2}, 0)}(0) &= P_A^{(\frac{\pi}{2}, 0)}(1) = \frac{1}{2}, \\
P_B^{(\frac{\pi}{2}, 0)}(0) &= P_B^{(\frac{\pi}{2}, 0)}(1) = \frac{1}{2}.
\end{aligned} \tag{44}$$

Therefore, we have that in this case the LAQC quantifier is given by

$$\mathcal{L}(\rho^{X_s}) = \frac{1+T_2}{2} \log_2(1+T_2) + \frac{1-T_2}{2} \log_2(1-T_2). \quad (45)$$

- $\theta = \frac{\pi}{2}, \phi = \frac{\pi}{2}$: As was the case for $\theta = \frac{\pi}{2}, \phi = 0$, we have two distinct cases with $\Phi = \{0, \pi, 2\pi\}$, and $\Phi = \{\frac{\pi}{2}, \frac{3\pi}{2}\}$. For the first case, the same expressions as before are obtained and therefore the LAQCs quantifier is again given by (42). For $\Phi = \{\frac{\pi}{2}, \frac{3\pi}{2}\}$, we obtain analogous expressions as before but with T_1 instead of T_2 , so that the LAQC quantifier is now given by:

$$\mathcal{L}(\rho^{X_s}) = \frac{1+T_1}{2} \log_2(1+T_1) + \frac{1-T_1}{2} \log_2(1-T_1). \quad (46)$$

1. Amplitude Damping Decoherence of Werner states

Werner states, ρ_w , can be written as:

$$\rho_w = z |\Psi^-\rangle\langle\Psi^-| + \frac{1-z}{4} \mathbb{1}_4, \quad (47)$$

where $z \in [0, 1]$ and $|\Psi^-\rangle = \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle)$ is the singlet, one of the four maximally entangled 2-qubit states known as Bell states. In a previous work [25], we analyzed the behaviour of LAQC for such states under Markovian decoherence, considering Depolarizing and Phase Damping channels. Amplitude Damping decoherence was not included due the resulting state no longer belonging to the Bell Diagonal set, but rather to the symmetric X states.

Within the Kraus operators formalism [27], the effect of the environment for a 2-qubit state is described via

$$\rho \longrightarrow \rho' = \sum_{i,j} \left(\mathbf{E}_i^{(A)} \otimes \mathbf{E}_j^{(B)} \right) \rho \left(\mathbf{E}_i^{(A)} \otimes \mathbf{E}_j^{(B)} \right)^\dagger \quad (48)$$

where the type of interaction for each subsystem may not be the same or even the parameter describing a common type of interaction might be different. In our present analysis, we study an equal interaction parameter for both subsystems.

Amplitude damping decoherence [28] describes the process of energy dissipation into the environment, like the T_1 process in NMR, for example. The Kraus operators for this type of one-way decay are

$$\mathbf{E}_0^{(AD)} = \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{1-p} \end{pmatrix}, \quad \mathbf{E}_1^{(AD)} = \begin{pmatrix} 0 & \sqrt{p} \\ 0 & 0 \end{pmatrix}, \quad (49)$$

where p can be thought of as the probability of losing a single quantum of energy, i.e. $|1\rangle \rightarrow |0\rangle$.

The state $\rho_w^{(AD)}$ resulting from applying the above Kraus operators via (48) is a symmetric X state and has the following non-null Bloch parameters:

$$x_3 = y_3 = p, \quad T_1 = T_2 = -(1-p)z, \quad T_3 = p^2 - (1-p)^2 z. \quad (50)$$

Since $T_1 = T_2$, we are left with only two possible expressions for $\mathcal{C}(\rho_w^{(AD)})$ (30). For $\theta = 0$, we have that

$$\begin{aligned} \mathcal{C}_0^{wAD} &= \frac{1}{4} (1-p)^2 (1-z) \log_2(1-z) \\ &+ \frac{1}{4} [(1-z)(1+p^2) + 2(1+z)p] \\ &\quad \times \log_2 [(1-z)(1+p^2) + 2(1+z)p] \\ &+ \frac{1}{2} (1-p) [(1-z)p + z + 1] \\ &\quad \times \log_2 [(1-z)p + z + 1] \\ &- (p+1) \log_2(1+p) \end{aligned} \quad (51)$$

and for $\theta = \frac{\pi}{2}$,

$$\begin{aligned} \mathcal{C}_{\frac{\pi}{2}}^{wAD} &= \frac{1}{2} [1 + (1-p)z] \log_2 [1 + (1-p)z] \\ &+ \frac{1}{2} [1 - (1-p)z] \log_2 [1 - (1-p)z]. \end{aligned} \quad (52)$$

In Figure 1 we present a graphical comparison of \mathcal{C}_0^{wAD}




Figure 1. Graphical comparison of the classical correlation function for $\theta = 0$ (grid) and $\theta = \frac{\pi}{2}$ (solid gray).

(grid) and $\mathcal{C}_{\frac{\pi}{2}}^{wAD}$ (solid gray). From that figure we can immediately conclude that the computational basis, that is $\theta = 0$ and $\phi = 0$, is the optimal one and the classical correlations quantifier is the one given in (51).

With the optimal computational basis defined, we can now turn our attention to our previous results and directly use (39). Since $T_1 = T_2 = T_M = -(1-p)z$, we have that the LAQC quantifier (18) is given by

$$\begin{aligned} \mathcal{L}(\rho_w^{(AD)}) &= \frac{1}{2} [1 + (1-p)z] \log_2 [1 + (1-p)z] \\ &+ \frac{1}{2} [1 - (1-p)z] \log_2 [1 - (1-p)z]. \end{aligned} \quad (53)$$




Figure 2. LAQC (above), QD (middle), and Concurrence (below) for a Werner state under amplitude damping decoherence.

A simple calculation shows that the above expression is only zero when $p = 0 \forall z$ and $z = 0 \forall p$.

The Concurrence (9) for an X state with Bloch parameters (50) is given by

$$C = \max [0, C_2], \quad \text{with} \quad C_2 = \frac{1-p}{2} \left[2z - \sqrt{1-z} \sqrt{(1+p)^2 - (1-p)^2 z} \right], \quad (54)$$

while its Quantum Discord is given by (A.17) using the corresponding Bloch parameters (50).

In Figure 2 we present the graphical behaviour of LAQC, QD, and concurrence. As expected, Werner states exhibit the so called ‘Entanglement Sudden Death’ [29], while LAQC only vanishes asymptotically, as does

the QD. That was already observed for two other decoherent interactions (depolarization and phase damping) in [25], therefore LAQC appears to be more robust as a quantum resource than entanglement, at least for this family of BD states. In Figure 3 we can observe the difference between QD and LAQC for this type of states. That is, we show the surface

$$S(z, p) = D_A(\rho_w^{(AD)}) - \mathcal{L}(\rho_w^{(AD)}). \quad (55)$$




Figure 3. Difference between QD and LAQC for a Werner state under amplitude damping decoherence.

2. A single parameter symmetric X state

As previously stated, by using Bell states and vectors of the computational basis, a single parameter X state can be attained such that $|x_3| = |y_3|$. As an example of such a symmetric X state we define the following density operator:

$$\rho_s = F |\Psi^-\rangle\langle\Psi^-| + (1-F) |00\rangle\langle 00|, \quad (56)$$

whose non-zero Bloch parameters are given by

$$x_3 = y_3 = 1 - F, \quad T_1 = T_2 = -F, \quad T_3 = 1 - 2F. \quad (57)$$

For this state the $R_{ij}(\theta, \phi)$ coefficients (22) are given by

$$\begin{aligned} R_{00}(\theta) &= \frac{1}{4}(1-F)(1 + \cos \theta)^2, \\ R_{10}(\theta) &= \frac{1}{8}[1 + 3F - (1-F) \cos(2\theta)] = R_{01}(\theta), \\ R_{11}(\theta) &= \frac{1}{4}(1-F)(1 - \cos \theta)^2 \end{aligned} \quad (58)$$

and marginal coefficients $R_i(\theta)$

$$\begin{aligned} R_0(\theta) &= \frac{1}{2}[1 + (1-F) \cos \theta], \\ R_1(\theta) &= \frac{1}{2}[1 - (1-F) \cos \theta]. \end{aligned} \quad (59)$$

The classical correlation function for $\theta = 0$ is then

$$\begin{aligned} C_0^s &= 2 - F - F \log_2(2 - F) \\ &\quad + (1-F) \log_2 \left[\frac{1-F}{(2-F)^2} \right]. \end{aligned} \quad (60)$$

For $\theta = \frac{\pi}{2}$, the corresponding function is

$$\mathcal{C}_{\frac{\pi}{2}}^s = \frac{1}{2} \left[(1-F) \log_2(1-F) + (1+F) \log_2(1+F) \right] \quad (61)$$




Figure 4. Graphical comparison of the classical correlation function for $\theta = 0$ (black) and for $\theta = \frac{\pi}{2}$ (gray).

Although a direct analysis of the above functions leads to conclude that the angles $\theta = 0$ and $\phi = 0$ are the ones that minimize (30), we can verify this graphically (Figure 4). Since the computational basis is optimal, we proceed to determine the probability distributions (31) and the marginals (32), obtaining

$$\begin{aligned} P_{00}(F) &= \frac{1}{4}(1-F) = P_{11}(F), \\ P_{01}(F) &= \frac{1}{4}(1+F) = P_{10}(F), \end{aligned} \quad (62)$$

and $P_0 = P_1 = \frac{1}{2}$. With all these results we have that the LAQC quantifier (18) is given by

$$\begin{aligned} \mathcal{L}(\rho_s) &= \frac{1}{2} [1+F] \log_2 [1+F] \\ &\quad + \frac{1}{2} [1-F] \log_2 [1-F] \end{aligned} \quad (63)$$

It is straightforward to verify that the Concurrence for (56) is

$$\mathcal{C}_s = F, \quad (64)$$

while its Quantum Discord, computed using the results presented in [20] and discussed in the appendix, is given by (A.17) using the corresponding Bloch parameters (57).

In Figure 5 the abovementioned quantifiers are presented. As can be observed, $\mathcal{L}(\rho_s) < D_A(\rho_s) < \mathcal{C}_s$ for $F \in (0, 1)$ and are only equal at $F = 0$ and $F = 1$.




Figure 5. LAQC (continuous black line), QD (continuous gray line), and Concurrence (dash-dot line) for state (56).

B. Anti-symmetric X States

We now focus on studying antisymmetric X states, i.e. ρ^X whose local Bloch vectors have equal norm but opposing direction. In the Fano-Bloch representation (19), for such states we have that $x_3 = -y_3$. The $R_{ij}(\theta_1, \theta_2, \phi_1, \phi_2)$ coefficients (12) previously defined are readily determined:

$$\begin{aligned} R_{00} &= \frac{1}{4} + \frac{1}{4}(\cos \theta_1 - \cos \theta_2)x_3 \\ &\quad + \Lambda \sum_{m=0}^1 \cos[\phi_1 + (-1)^m \phi_2][T_1 - (-1)^m T_2] \\ &\quad + \frac{1}{4} \cos \theta_1 \cos \theta_2 T_3, \\ R_{01} &= \frac{1}{4} + \frac{1}{4}(\cos \theta_1 + \cos \theta_2)x_3 \\ &\quad - \Lambda \sum_{m=0}^1 \cos[\phi_1 + (-1)^m \phi_2][T_1 - (-1)^m T_2] \\ &\quad - \frac{1}{4} \cos \theta_1 \cos \theta_2 T_3, \\ R_{10} &= \frac{1}{4} - \frac{1}{4}(\cos \theta_1 + \cos \theta_2)x_3 \\ &\quad - \Lambda \sum_{m=0}^1 \cos[\phi_1 + (-1)^m \phi_2][T_1 - (-1)^m T_2] \\ &\quad - \frac{1}{4} \cos \theta_1 \cos \theta_2 T_3, \\ R_{11} &= \frac{1}{4} - \frac{1}{4}(\cos \theta_1 - \cos \theta_2)x_3 \end{aligned} \quad (65)$$

$$\begin{aligned}
& + \mathbf{\Lambda} \sum_{m=0}^1 \cos[\phi_1 + (-1)^m \phi_2] [T_1 - (-1)^m T_2] \\
& + \frac{1}{4} \cos \theta_1 \cos \theta_2 T_3.
\end{aligned}$$

where $\mathbf{\Lambda}$ is as defined in (21).

As with the symmetric case, a relation between the angles θ_i and ϕ_i is expected. A direct calculation interchanging local angles θ_i and ϕ_i leads to

$$\begin{aligned}
R_{ii}(\theta_1, \theta_2, \phi_1, \phi_2) &= R_{ii}(\theta_2 + \pi, \theta_1 + \pi, \phi_1, \phi_2), \\
R_{ij}(\theta_1, \theta_2, \phi_1, \phi_2) &= R_{ij}(\theta_2, \theta_1, \phi_1, \phi_2), \quad i \neq j.
\end{aligned} \tag{66}$$

Therefore, as was done for the symmetric case, we can define θ and ϕ as the respective common local angles and the expressions in (65) are readily simplified:

$$\begin{aligned}
R_{00}(\theta, \phi) &= \frac{1}{4} [1 + \sin^2 \theta (\cos^2 \phi T_1 + \sin^2 \phi T_2) \\
&\quad + \cos^2 \theta T_3] = R_{11}(\theta, \phi), \\
R_{01}(\theta, \phi) &= \frac{1}{4} [1 + 2 \cos \theta x_3 - \sin^2 \theta (\cos^2 \phi T_1 \\
&\quad + \sin^2 \phi T_2) - \cos^2 \theta T_3], \\
R_{10}(\theta, \phi) &= \frac{1}{4} [1 - 2 \cos \theta x_3 - \sin^2 \theta (\cos^2 \phi T_1 \\
&\quad + \sin^2 \phi T_2) - \cos^2 \theta T_3].
\end{aligned} \tag{67}$$

The corresponding coefficients $\{R_{i_A}(\theta, \phi), R_{j_B}(\theta, \phi)\}$ for the reduced matrices are given by

$$\begin{aligned}
R_{0_A}(\theta) &= \frac{1}{2} (1 + x_3 \cos \theta) = R_{1_B}(\theta), \\
R_{1_A}(\theta) &= \frac{1}{2} (1 - x_3 \cos \theta) = R_{0_B}(\theta).
\end{aligned} \tag{68}$$

The necessary minimization to determine the optimal computational base leads again to three main cases to be considered:

- $\theta = 0, \phi \in [0, 2\pi]$:

$$\begin{aligned}
R_{00}(\theta = 0, \phi) &= \frac{1}{4} [1 + T_3] = R_{11}, \\
R_{01}(\theta = 0, \phi) &= \frac{1}{4} [1 - T_3 + 2x_3], \\
R_{10}(\theta = 0, \phi) &= \frac{1}{4} [1 - T_3 - 2x_3].
\end{aligned} \tag{69}$$

- $\theta = \frac{\pi}{2}, \phi = 0$:

$$\begin{aligned}
R_{00}\left(\theta = \frac{\pi}{2}, \phi = 0\right) &= \frac{1}{4} [1 + T_1] = R_{11}, \\
R_{01}\left(\theta = \frac{\pi}{2}, \phi = 0\right) &= \frac{1}{4} [1 - T_1] = R_{10}.
\end{aligned} \tag{70}$$

- $\theta = \frac{\pi}{2}, \phi = \frac{\pi}{2}$:

$$\begin{aligned}
R_{00}\left(\theta = \frac{\pi}{2}, \phi = \frac{\pi}{2}\right) &= \frac{1}{4} [1 + T_2] = R_{11}, \\
R_{01}\left(\theta = \frac{\pi}{2}, \phi = \frac{\pi}{2}\right) &= \frac{1}{4} [1 - T_2] = R_{10}.
\end{aligned} \tag{71}$$

We use α_{\pm} and T_m (27) to rewrite the previous expressions and center our attention on θ :

- $\theta = 0$:

$$\begin{aligned}
R_{00}(\theta = 0) &= \frac{1}{4} [1 + T_3] = R_{11}, \\
R_{01}(\theta = 0) &= \frac{1}{4} [1 - \alpha_-], \\
R_{11}(\theta = 0) &= \frac{1}{4} [1 - \alpha_+].
\end{aligned} \tag{72}$$

- $\theta = \frac{\pi}{2}$:

$$\begin{aligned}
R_{00}\left(\theta = \frac{\pi}{2}\right) &= \frac{1}{4} [1 + T_m] = R_{11}, \\
R_{01}\left(\theta = \frac{\pi}{2}\right) &= \frac{1}{4} [1 - T_m] = R_{10}.
\end{aligned} \tag{73}$$

The general expression for the quantifier of the classical correlations (14) is given by

$$\mathcal{C}(\rho^{X_{as}}) = \sum_{i,j} R_{i,j}(\theta, \phi) \log_2 \left[\frac{R_{i,j}(\theta, \phi)}{R_i(\theta) R_j(\theta)} \right], \tag{74}$$

which is completely analogous to (30). The minimization of the above expression can again be done either by directly minimizing it for θ and ϕ or by comparing the coefficients $R_{i,j}(\theta, \phi)$.

After the angles θ and ϕ have been determined, the state $\rho^{X_{as}}$ is rewritten in the optimal computational basis as was done previously for the symmetric case and the probability distributions $P^{(\theta, \phi)}(i_A, j_B, \Phi)$ (17) as well as the marginal distributions $P_A^{(\theta, \phi)}(i_A, \Phi)$ and $P_B^{(\theta, \phi)}(i_B, \Phi)$ are determined. Since the maximization procedure for these probability distributions is analogous to the one performed for the symmetric case, with some of the results even being the same, we present the corresponding analysis in a more succinct manner. For each pair (θ, ϕ) defining the optimal computational basis, the following results are obtained:

- $\theta = 0, \phi = 0$:

$$\begin{aligned}
P^{(0,0)}(0, 0, \Phi) &= \frac{1}{4} (1 + T_1 \cos^2 \Phi + T_2 \sin^2 \Phi), \\
&= P^{(0,0)}(1, 1, \Phi), \\
P^{(0,0)}(1, 0, \Phi) &= \frac{1}{4} (1 - T_1 \cos^2 \Phi - T_2 \sin^2 \Phi), \\
&= P^{(0,0)}(0, 1, \Phi),
\end{aligned} \tag{75}$$

and

$$\begin{aligned}
P_A^{(0,0)}(0) &= P_A^{(0,0)}(1) = \frac{1}{2}, \\
P_B^{(0,0)}(0) &= P_B^{(0,0)}(1) = \frac{1}{2}.
\end{aligned} \tag{76}$$

These are exactly the same expressions that we obtained for the symmetric case in (31) and (32).

Thus, they will lead to same expression for the LAQC quantifier. Using T_M (37), we again have that the LAQC quantifier is given, as in (39), by

$$\begin{aligned} \mathcal{L}(\rho^{X_{as}}) &= \frac{1+T_M}{2} \log_2(1+T_M) \\ &\quad + \frac{1-T_M}{2} \log_2(1-T_M). \end{aligned} \quad (77)$$

• $\theta = \frac{\pi}{2}, \phi = 0$:

$$\begin{aligned} P^{(\frac{\pi}{2},0)}(0,0,\Phi) &= \frac{1}{4}(1+T_2 \sin^2 \Phi + T_3 \cos^2 \Phi), \\ &= P^{(\frac{\pi}{2},0)}(1,1,\Phi), \\ P^{(\frac{\pi}{2},0)}(1,0,\Phi) &= \frac{1}{4}(1-T_2 \sin^2 \Phi - T_3 \cos^2 \Phi \\ &\quad + 2x_3 \cos \Phi), \\ P^{(\frac{\pi}{2},0)}(0,1,\Phi) &= \frac{1}{4}(1-T_2 \sin^2 \Phi - T_3 \cos^2 \Phi \\ &\quad - 2x_3 \cos \Phi). \end{aligned} \quad (78)$$

and

$$\begin{aligned} P_A^{(\frac{\pi}{2},0)}(0) &= P_B^{(\frac{\pi}{2},0)}(1) = \frac{1}{2}(1-x_3 \cos \Phi), \\ P_A^{(\frac{\pi}{2},0)}(1) &= P_B^{(\frac{\pi}{2},0)}(0) = \frac{1}{2}(1+x_3 \cos \Phi). \end{aligned} \quad (79)$$

As with the symmetric case, we have two distinct cases, $\Phi = \{0, \pi, 2\pi\}$ and $\Phi = \{\frac{\pi}{2}, \frac{3\pi}{2}\}$. For $\Phi = \{0, 2\pi\}$ we have that

$$\begin{aligned} P^{(\frac{\pi}{2},0)}(0,0) &= \frac{1}{4}(1+T_3) \\ &= P^{(\frac{\pi}{2},0)}(1,1), \\ P^{(\frac{\pi}{2},0)}(1,0) &= \frac{1}{4}(1-\alpha_-), \\ P^{(\frac{\pi}{2},0)}(0,1) &= \frac{1}{4}(1-\alpha_+). \end{aligned} \quad (80)$$

and

$$\begin{aligned} P_A^{(\frac{\pi}{2},0)}(0) &= P_B^{(\frac{\pi}{2},0)}(1) = \frac{1}{2}(1-x_3), \\ P_A^{(\frac{\pi}{2},0)}(1) &= P_B^{(\frac{\pi}{2},0)}(0) = \frac{1}{2}(1+x_3). \end{aligned} \quad (81)$$

For $\Phi = \pi$ we obtain equivalent expressions with an analogue change in ordering as in the symmetric case. With these results, we have that

$$\begin{aligned} \mathcal{L}(\rho^{X_{as}}) &= \frac{1-\alpha_-}{4} \log_2 \left[\frac{1-\alpha_-}{(1+x_3)^2} \right] \\ &\quad + \frac{1+T_3}{2} \log_2 \left(\frac{1+T_3}{1-x_3^2} \right) \\ &\quad + \frac{1-\alpha_+}{4} \log_2 \left[\frac{1-\alpha_+}{(1-x_3)^2} \right]. \end{aligned} \quad (82)$$

For $\Phi = \frac{\pi}{2}, \frac{3\pi}{2}$, we obtain the same results as for the symmetric case and the LAQC quantifier is therefore given by the same expression as in (45):

$$\begin{aligned} \mathcal{L}(\rho^{X_{as}}) &= \frac{1+T_2}{2} \log_2(1+T_2) \\ &\quad + \frac{1-T_2}{2} \log_2(1-T_2). \end{aligned} \quad (83)$$

• $\theta = \frac{\pi}{2}, \phi = \frac{\pi}{2}$: In this case we obtain analogous results as for $\theta = \frac{\pi}{2}$ and $\phi = 0$ by exchanging $T_2 \rightarrow T_1$ in (78):

$$\begin{aligned} P^{(\frac{\pi}{2},0)}(0,0,\Phi) &= \frac{1}{4}(1+T_1 \sin^2 \Phi + T_3 \cos^2 \Phi), \\ &= P^{(\frac{\pi}{2},0)}(1,1,\Phi), \\ P^{(\frac{\pi}{2},0)}(1,0,\Phi) &= \frac{1}{4}(1-T_1 \sin^2 \Phi - T_3 \cos^2 \Phi \\ &\quad + 2x_3 \cos \Phi), \\ P^{(\frac{\pi}{2},0)}(0,1,\Phi) &= \frac{1}{4}(1-T_1 \sin^2 \Phi - T_3 \cos^2 \Phi \\ &\quad - 2x_3 \cos \Phi). \end{aligned} \quad (84)$$

and again we have that

$$\begin{aligned} P_A^{(\frac{\pi}{2},\frac{\pi}{2})}(0) &= P_B^{(\frac{\pi}{2},\frac{\pi}{2})}(1) = \frac{1}{2}(1-x_3 \cos \Phi), \\ P_A^{(\frac{\pi}{2},\frac{\pi}{2})}(1) &= P_B^{(\frac{\pi}{2},\frac{\pi}{2})}(0) = \frac{1}{2}(1+x_3 \cos \Phi). \end{aligned} \quad (85)$$

As before, we have two distinct cases with $\Phi = \{0, \pi, 2\pi\}$, and $\Phi = \{\frac{\pi}{2}, \frac{3\pi}{2}\}$. For the first case, the same expressions as before are obtained and therefore the LAQCs quantifier is again given by (82). For $\Phi = \{\frac{\pi}{2}, \frac{3\pi}{2}\}$, we obtain analogous expressions as before but with T_1 instead of T_2 :

$$\begin{aligned} \mathcal{L}(\rho^{X_{as}}) &= \frac{1+T_1}{2} \log_2(1+T_1) \\ &\quad + \frac{1-T_1}{2} \log_2(1-T_1). \end{aligned} \quad (86)$$

1. A single parameter antisymmetric X state

In [30], Verstraete et al. introduced a single parameter mixed 2-qubit state, defined as:

$$\rho_v = F |\Phi^+\rangle\langle\Phi^+| + (1-F) |01\rangle\langle 01| \quad (87)$$

where $|\Phi^+\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$. It is straightforward to verify that this state is an anti-symmetric X state, with Bloch parameters

$$\begin{aligned} x_3 = -y_3 = 1-F, \quad T_1 = -T_2 = F, \\ T_3 = 2F-1, \end{aligned} \quad (88)$$

and all others equal to zero. The $R_{ij}(\theta, \phi)$ coefficients (67) are given by

$$\begin{aligned} R_{00}(\theta, \phi) &= \frac{1}{4} [1 + F \cos(2\phi) \sin^2 \theta \\ &\quad + (2F - 1) \cos^2 \theta] = R_{11}(\theta, \phi), \\ R_{10}(\theta) &= \frac{1}{4} [1 - F \cos(2\phi) \sin^2 \theta \\ &\quad - (2F - 1) \cos^2 \theta - 2(1 - F) \cos \theta], \\ R_{01}(\theta) &= \frac{1}{4} [1 - F \cos(2\phi) \sin^2 \theta \\ &\quad - (2F - 1) \cos^2 \theta + 2(1 - F) \cos \theta], \end{aligned} \quad (89)$$

and the marginal coefficients $R_i(\theta)$ are

$$\begin{aligned} R_{0_A}(\theta) &= \frac{1}{2} [1 + (1 - F) \cos \theta] = R_{1_B}(\theta), \\ R_{1_A}(\theta) &= \frac{1}{2} [1 - (1 - F) \cos \theta] = R_{0_B}(\theta). \end{aligned} \quad (90)$$

The classical correlation function for $\theta = 0$ is given by

$$\begin{aligned} C_0^v &= 2 - F - F \log_2(2 - F) \\ &\quad + (1 - F) \log_2 \left[\frac{1 - F}{(2 - F)^2} \right], \end{aligned} \quad (91)$$

and for $\theta = \frac{\pi}{2}$, $\phi = 0, \frac{\pi}{2}$ by

$$\begin{aligned} C_{\frac{\pi}{2}}^v &= \frac{1}{2} [(1 - F) \log_2(1 - F) \\ &\quad + (1 + F) \log_2(1 + F)] \end{aligned} \quad (92)$$

These are the same functions (60) and (61) that we obtained for state (56). As before, the computational basis is the optimal one and this leads to equivalent expressions for the probability distributions (75) and the marginals (76) as before

$$\begin{aligned} P_{00}(F) &= \frac{1}{4} [1 - F \cos(2\Phi)] = P_{11}(F), \\ P_{01}(F) &= \frac{1}{4} [1 + F \cos(2\Phi)] = P_{10}(F), \end{aligned} \quad (93)$$

and $P_0 = P_1 = \frac{1}{2}$. These distributions are maximized for $\Phi = n \frac{\pi}{2}$, $n = 0, 1, 2, 3$, and the LAQC quantifier (18) is given by

$$\begin{aligned} \mathcal{L}(\rho_v) &= \frac{1}{2} [1 + F] \log_2 [1 + F] \\ &\quad + \frac{1}{2} [1 - F] \log_2 [1 - F], \end{aligned} \quad (94)$$

which is the same expression found for the symmetric case (63).

The Concurrence for state (87) is again given by (64) and the QD is the same as in the one-parameter symmetric X state presented in (56). Therefore, the corresponding graph is the same as the one in Figure 5.

IV. CONCLUSIONS

We have studied the local available quantum correlations (LAQC) [6] for 2-qubit X states that have a local Bloch vector with equal magnitude. Such states can be regarded as symmetric and antisymmetric X states wether their local Bloch vector is parallel or anti-parallel, respectively. We determined all possible expressions for the classical correlations and LAQC quantifiers in terms of the Bloch parameters of the state. As examples of interesting symmetric states we have analyzed the (markovian) amplitude damping decoherence of Werner states and found that LAQC do not suffer sudden death under this quantum channel. Also, examples of one parameter X states were studied, both symmetric and antisymmetric. For such states, there are less LAQC than entanglement (as measured via concurrence) as well as quantum discord. The robustness of this quantum correlation under markovian decoherence is to be determined.

ACKNOWLEDGMENTS

This work was partially funded by the *2020 BrainGain Venezuela* grant awarded to H. Albrecht by the *Physics without Frontiers* program of the ICTP. Albrecht and Bellorin would also like to thank the support given by the research group *GID-30, Teoría de Campos y Óptica Cuántica*, at the Universidad Simón Bolívar, Venezuela.

Appendix: Quantum Discord of 2-qubit Symmetric and Antisymmetric X states

In [20], Li et al. presented an analytic expression for the quantum discord of X states with parallel local Bloch vectors. Their analysis allows to compute the classical correlations $\mathcal{C}_D(\rho^X)$ when you rewrite (1) as

$$D_A(\rho^X) = I(\rho^X) - \mathcal{C}_D(\rho^X), \quad (A.1)$$

where

$$\mathcal{C}_D(\rho^X) \equiv \min_{\{\Pi_i^A\}} I[(\Pi^A \otimes \mathbb{1}_2) \rho_{AB}]. \quad (A.2)$$

Lets start by defining monotonically decreasing function f for $x \in [0, 1]$:

$$f(x) = -\frac{1+x}{2} \log_2(1+x) - \frac{1-x}{2} \log_2(1-x), \quad (A.3)$$

so that the von Neumann entropy of the reduced matrices of ρ^X is given by

$$S(\rho_A^X) = 1 + f(x_3), \quad S(\rho_B^X) = 1 + f(y_3). \quad (A.4)$$

Since the eigenvalues of ρ^X (19) are given

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{4} \left[1 - T_3 \pm \sqrt{(x_3 - y_3)^2 + (T_1 + T_2)^2} \right], \\ \lambda_{3,4} &= \frac{1}{4} \left[1 + T_3 \pm \sqrt{(x_3 + y_3)^2 + (T_1 - T_2)^2} \right], \end{aligned} \quad (A.5)$$

the mutual information (2) is then

$$I(\rho^X) = 2 + f(x_3) + f(y_3) + \sum_{i=1}^4 \lambda_i \log_2 \lambda_i. \quad (\text{A.6})$$

As for $\mathcal{C}_D(\rho^X)$, it is given by

$$\mathcal{C}_D(\rho^X) = 1 - f(x_3) - \min\{S_1, S_2, S_3\}, \quad (\text{A.7})$$

where

$$S_1 \equiv - \sum_{i=1}^4 \rho_{ii}^X \log_2 \left[\frac{2\rho_{ii}^X}{1 - (-1)^i y_3} \right] \quad (\text{A.8})$$

and

$$\begin{aligned} \rho_{11}^X &= \frac{1}{4}(1 + x_3 \pm y_3 \pm T_3), \\ \rho_{33}^X &= \frac{1}{4}(1 - x_3 \pm y_3 \mp T_3), \end{aligned} \quad (\text{A.9})$$

and

$$S_{2,3} \equiv 1 + f\left(\sqrt{x_3^2 + T_{1,2}^2}\right). \quad (\text{A.10})$$

For X states where $|x_3| = |y_3|$, we have that $f(x_3) = f(y_3)$ so that

$$I(\rho^{X_s}) = 2 + 2f(x_3) + \sum_{i=1}^4 \lambda_i \log_2 \lambda_i, \quad (\text{A.11})$$

where

$$\begin{aligned} \lambda_{1,2}^s &= \frac{1}{4}[1 - T_3 \pm (T_1 + T_2)], \\ \lambda_{3,4}^s &= \frac{1}{4}\left[1 + T_3 \pm \sqrt{4x_3^2 + (T_1 - T_2)^2}\right], \end{aligned} \quad (\text{A.12})$$

for $x_3 = y_3$ (symmetric) and

$$\begin{aligned} \lambda_{1,2}^{as} &= \frac{1}{4}\left[1 - T_3 \pm \sqrt{4x_3^2 + (T_1 + T_2)^2}\right], \\ \lambda_{3,4}^{as} &= \frac{1}{4}[1 + T_3 \pm (T_1 - T_2)], \end{aligned} \quad (\text{A.13})$$

for $x_3 = -y_3$ (anti-symmetric). As for S_1 , we have that for ρ^{X_s}

$$\begin{aligned} \rho_{11}^{X_s} &= \frac{1}{4}(1 + 2x_3 + T_3), & \rho_{22}^{X_s} &= \frac{1}{4}(1 - T_3), \\ \rho_{33}^{X_s} &= \frac{1}{4}(1 - T_3), & \rho_{44}^{X_s} &= \frac{1}{4}(1 - 2x_3 + T_3), \end{aligned} \quad (\text{A.14})$$

and

$$\begin{aligned} \rho_{11}^X &= \frac{1}{4}(1 + T_3), & \rho_{22}^X &= \frac{1}{4}(1 + 2x_3 - T_3), \\ \rho_{33}^X &= \frac{1}{4}(1 - 2x_3 - T_3), & \rho_{44}^X &= \frac{1}{4}(1 + T_3), \end{aligned} \quad (\text{A.15})$$

for $\rho^{X_{as}}$.

As for the examples presented in this paper, it is straightforward to verify that for all three cases (50), (56), and (87),

$$\min\{S_1, S_2, S_3\} = S_2 = S_3, \quad (\text{A.16})$$

so that

$$\begin{aligned} D_A(\rho_{AB}) &= 3 + f(x_3) + f\left(\sqrt{x_3^2 + T_2^2}\right) \\ &+ \sum_{i=1}^4 \lambda_i \log_2 \lambda_i, \end{aligned} \quad (\text{A.17})$$

where the Bloch parameters were given in (50), (57), and (88), respectively.

-
- [1] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki. Quantum entanglement. *Rev. Mod. Phys.*, 81:865–942, 6 2009. arXiv:quant-ph/0702225.
- [2] H. Ollivier and W.H. Zurek. Quantum discord: A measure of the quantumness of correlations. *Phys. Rev. Lett.*, 88:017901–017904, 12 2001. arXiv:quant-ph/0105072.
- [3] L. Henderson and V. Vedral. Classical, quantum and total correlations. *J. Phys. A*, 34(35):6899–6905, aug 2001. arXiv:quant-ph/0105028.
- [4] K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral. The classical-quantum boundary for correlations: Discord and related measures. *Rev. Mod. Phys.*, 84:1655–1707, Nov 2012. arXiv:1112.6238.
- [5] S. Wu, Z. Ma, Z. Chen, and S. Xia. Reveal quantum correlation in complementary bases. *Sci. Rep.*, 4:4036–4041, 2 2015. arXiv:1301.6838.
- [6] D.F. Mundarain and M.L.L. de Guevara. Local available quantum correlations. *Quantum Inf Process*, 14:4493–4510, 10 2015.
- [7] T. Yu and J. H. Eberly. Evolution from entanglement to decoherence of bipartite mixed “x” states. *Quantum Info. Comput.*, 7(5):459–468, 2007. arXiv:quant-ph/0503089.
- [8] N. Quesada, A. Al-Qasimi, and D.F.V. James. Quantum properties and dynamics of x states. *J Mod Opt*, 59(15):1322–1329, 2012. arXiv:1207.3689.
- [9] P. E. M. F. Mendonça, M. A. Marchioli, and D. Galetti. Entanglement universality of two-qubit x-states. *Ann. Phys.*, 351:79–103, 2014. arXiv:1407.3021.
- [10] S.R. Hedemann. X states of the same spectrum and entanglement as all two-qubit states. *Quantum Inf Process*, 17(11):293, 2018. arXiv:1802.03038.
- [11] C. Monroe, D. M. Meekhof, B. E. King, W. M. Itano, and D. J. Wineland. Demonstration of a fundamental quantum logic gate. *Phys. Rev. Lett.*, 75:4714–4717, 12 1995.
- [12] P. G. Kwiat, K. Mattle, Y. Weinfurter, A. Zeilinger, A. V. Sergienko, and Y. Shih. New high-intensity source of polarization-entangled photon pairs. *Phys. Rev. Lett.*, 75:4337–4341, 12 1995.
- [13] T. Werlang, C. Trippe, G.A.P. Ribeiro, and G. Rigolin.

- Quantum correlations in spin chains at finite temperatures and quantum phase transitions. *Phys. Rev. Lett.*, 105:095702–095705, 2010. arXiv:1006.3332.
- [14] J. S. Pratt. Universality in the entanglement structure of ferromagnets. *Phys. Rev. Lett.*, 93:237205–237208, Dec 2004. arXiv:quant-ph/0411125.
- [15] W.K. Wootters. Entanglement of formation of an arbitrary state of two qubits. *Phys. Rev. Lett.*, 80:2245–2248, 3 1998. arXiv:quantph/9709029.
- [16] M. Ali, A. R. P. Rau, and G. Alber. Quantum discord for two-qubit x states. *Phys. Rev. A*, 81:042105, Apr 2010.
- [17] Q. Chen, C. Zhang, S. Yu, X. X. Yi, and C. H. Oh. Quantum discord of two-qubit x states. *Phys. Rev. A*, 84:042313, Oct 2011.
- [18] X.-M. Lu, J. Ma, Z. Xi, and X. Wang. Optimal measurements to access classical correlations of two-qubit states. *Phys. Rev. A*, 83:012327, Jan 2011.
- [19] Y. Huang. Quantum discord for two-qubit x states: Analytical formula with very small worst-case error. *Phys. Rev. A*, 88:014302, Jul 2013.
- [20] B. Li, Z.-X. Wang, and S.-M. Fei. Quantum discord and geometry for a class of two-qubit states. *Phys. Rev. A*, 83:022321, Feb 2011.
- [21] R.F. Werner. Quantum states with einstein-podolsky-rosen correlations admitting a hidden-variable model. *Phys. Rev. A*, 40:4277–4281, Oct 1989.
- [22] K. Modi, T. Paterek, W. Son, V. Vedral, and M. Williamson. Unified view of quantum and classical correlations. *Phys. Rev. Lett.*, 104:080501, Feb 2010. arXiv:0911.5417.
- [23] U. Fano. Pairs of two-level systems. *Rev. Mod. Phys.*, 55:855–874, Oct 1983.
- [24] R. Horodecki and M. Horodecki. Information-theoretic aspects of inseparability of mixed states. *Phys. Rev. A*, 54:1838–1843, Sep 1996.
- [25] H.L. Albrecht Q., M.I. Caicedo S., and D.F. Mundarain. Local available quantum correlations for bell diagonal states and markovian decoherence. *Rev. Mex. Fis.*, 64:662—670, Nov-Dec 2018. arXiv:1803.02426.
- [26] A.K. Fedorov, E.O. Kiktenko, O.V. Man’ko, and V.I. Man’ko. Tomographic discord for a system of two coupled nanoelectric circuits. *Phys. Scr.*, 90(5):055101, 4 2015. arXiv:1409.5265.
- [27] K. Kraus. General state changes in quantum theory. *Ann. Phys.*, 64(2):311–335, 6 1971.
- [28] M.A. Nielsen and I.L. Chuang. *Quantum Computation and Quantum Information: 10th Anniversary Edition*. Cambridge University Press, 2010.
- [29] T. Yu and J. H. Eberly. Sudden death of entanglement. *Science*, 323(5914):598–601, 2009. arXiv:0910.1396.
- [30] F. Verstraete and H. Verschelde. Optimal teleportation with a mixed state of two qubits. *Phys. Rev. Lett.*, 90:097901, Mar 2003. arXiv:quant-ph/0303007.