

RECIPROCITY SHEAVES AND LOGARITHMIC MOTIVES

SHUJI SAITO

CONTENTS

Introduction	1
1. Preliminaries	5
2. Purity with reduced modulus	10
3. Review on higher local symbols	14
4. Logarithmic cohomology of reciprocity sheaves	17
5. Invariance of logarithmic cohomology under blowups	22
6. Relation with logarithmic sheaves with transfers	27
References	30

ABSTRACT. We connect two developments aiming at extending Voevodsky’s theory of motives over a field in such a way to encompass non- \mathbf{A}^1 -invariant phenomena. One is theory of *reciprocity sheaves* introduced by Kahn-Saito-Yamazaki. Another is theory of the triangulated category $\mathbf{logDM}^{\text{eff}}$ of *logarithmic motives* launched by Binda, Park and Østvær. We prove that the Nisnevich cohomology of reciprocity sheaves is representable in $\mathbf{logDM}^{\text{eff}}$.

INTRODUCTION

We fix once and for all a perfect base field k . The main purpose of this paper is to connect two developments aiming at extending Voevodsky’s theory of motives over k in such a way to encompass non- \mathbf{A}^1 -invariant phenomena. One is theory of *reciprocity sheaves* introduced by Kahn-Saito-Yamazaki ([6] and [7]) and developed in [16] and [2]. Voevodsky’s theory is based on the category \mathbf{PST} of *presheaves with transfers*, defined as the category of additive presheaves of abelian groups on the category \mathbf{Cor} of finite correspondences: \mathbf{Cor} has the same objects as the category \mathbf{Sm} of separated smooth schemes of finite type over k and morphisms in \mathbf{Cor} are finite correspondences. Let $\mathbf{NST} \subset \mathbf{PST}$ be the full subcategory of Nisnevich sheaves, i.e. those objects $F \in \mathbf{PST}$ whose restrictions F_X to the small étale site

The author is supported by the JSPS KAKENHI Grant (20H01791).

$X_{\acute{e}t}$ over X are Nisnevich sheaves for all $X \in \mathbf{Sm}$. Voevodsky proved that \mathbf{NST} is a Grothendieck abelian category and defined the triangulated category \mathbf{DM}^{eff} of effective motives as the localization of the derived category $D(\mathbf{NST})$ of complexes in \mathbf{NST} with respect to an \mathbf{A}^1 -weak equivalence (see [10, Def. 14.1]). It is equipped with a functor $M : \mathbf{Sm} \rightarrow \mathbf{DM}^{\text{eff}}$ associating the motive $M(X)$ of $X \in \mathbf{Sm}$.

Let $\mathbf{HI}_{\text{Nis}} \subset \mathbf{NST}$ be the full subcategory consisting of \mathbf{A}^1 -invariant objects, namely such $F \in \mathbf{NST}$ that the projection $\pi_X : X \times \mathbf{A}^1 \rightarrow X$ induces an isomorphism $\pi_X^* : F(X) \simeq F(X \times \mathbf{A}^1)$ for any $X \in \mathbf{Sm}$. We say that $F \in \mathbf{HI}_{\text{Nis}}$ is strictly \mathbf{A}^1 -invariant if π_X induces isomorphisms

$$\pi_X^* : H_{\text{Nis}}^i(X, F_X) \simeq H_{\text{Nis}}^i(X \times \mathbf{A}^1, F_{X \times \mathbf{A}^1}) \quad \text{for all } i \geq 0.$$

The following theorem plays a fundamental role in Voevodsky's theory.

Theorem 0.1. (Voevodsky [17]) *Any $F \in \mathbf{HI}_{\text{Nis}}$ is strictly \mathbf{A}^1 -invariant and we have a natural isomorphism*

$$(0.1.1) \quad H_{\text{Nis}}^i(X, F_X) \simeq \text{Hom}_{\mathbf{DM}^{\text{eff}}}(M(X), L^{\mathbf{A}^1}F[i]) \quad \text{for } X \in \mathbf{Sm},$$

where $L^{\mathbf{A}^1} : D(\mathbf{NST}) \rightarrow \mathbf{DM}^{\text{eff}}$ is the localization functor.

Notice that there are interesting and important objects of \mathbf{NST} which do not belong to \mathbf{HI}_{Nis} . Such examples are given by the sheaves Ω^i of (absolute or relative) differential forms, and the p -typical de Rham-Witt sheaves $W_m \Omega^i$ of Bloch-Deligne-Illusie, and smooth commutative k -groups schemes with a unipotent part (seen as objects of \mathbf{NST}), and the complexes $R\varepsilon_* \mathbb{Z}/p^r(n)$ in case $\text{ch}(k) = p > 0$, where $\mathbb{Z}/p^r(n)$ is the étale motivic complex of weight n with \mathbb{Z}/p^r coefficients and ε is the change of site functor from the étale to the Nisnevich topology. For such examples, (0.1.1) fails to hold since $\pi_X : X \times \mathbf{A}^1 \rightarrow X$ induces an isomorphism $M(X \times \mathbf{A}^1) \simeq M(X)$ in \mathbf{DM}^{eff} but the maps induced on cohomology of those sheaves are not isomorphism.

The category $\mathbf{RSC}_{\text{Nis}}$ of reciprocity sheaves is a full abelian subcategory of \mathbf{NST} that contains \mathbf{HI}_{Nis} as well as the non- \mathbf{A}^1 -invariant objects mentioned above. Heuristically, its objects satisfy the property that for any $X \in \mathbf{Sm}$, each section $a \in F(X)$ “has bounded ramification at infinity” and the objects of \mathbf{HI}_{Nis} are special reciprocity sheaves with the property that every section $a \in F(X)$ has “tame” ramification at infinity¹. Slightly more exotic examples of reciprocity sheaves are given by the sheaves Conn^1 (in case $\text{ch}(k) = 0$), whose sections over X are rank 1-connections, or Lisse_ℓ^1 (in case $\text{ch}(k) = p > 0$), whose sections on X are the lisse $\overline{\mathbb{Q}}_\ell$ -sheaves of rank 1. Since $\mathbf{RSC}_{\text{Nis}}$ is an abelian category equipped with a lax symmetric monoidal structure by

¹This heuristic viewpoint is manifested in [13, Th. 2].

[11], many more interesting examples can be manufactured by taking kernels, quotients and tensor products (see [2, §11.1] for more examples).

The main purpose of this article is to establish the formula (0.1.1) for all $F \in \mathbf{RSC}_{\text{Nis}}$ in a new category which enlarges \mathbf{DM}^{eff} (see (0.2.2)). It is the triangulated category $\mathbf{logDM}^{\text{eff}}$ of *logarithmic motives* introduced by Binda, Park and Østvær in [1]. Let \mathbf{lSm} be the category of log smooth and separated fs log schemes of finite type over k and \mathbf{lCor} be the category with the same objects as \mathbf{lSm} and whose morphisms are log finite correspondences (see [1, Def. 2.1.1]). Let $\mathbf{PSh}^{\text{ltr}}$ be the category of additive presheaves of abelian groups on \mathbf{lCor} and $\mathbf{Shv}_{d\text{Nis}}^{\text{ltr}} \subset \mathbf{PSh}^{\text{ltr}}$ be the full subcategory consisting of those \mathcal{F} whose restrictions to \mathbf{lSm} are dividing Nisnevich sheaves (see [1, Def. 3.1.4]). It is shown in [1, §4 and Pr. 4.6.6] that $\mathbf{Shv}_{d\text{Nis}}^{\text{ltr}}$ is a Grothendieck abelian category, and $\mathbf{logDM}^{\text{eff}}$ is defined as the localization of the derived category $D(\mathbf{Shv}_{d\text{Nis}}^{\text{ltr}})$ of complexes in $\mathbf{Shv}_{d\text{Nis}}^{\text{ltr}}$ with respect to a $\overline{\square}$ -weak equivalence, where $\overline{\square}$ is \mathbf{P}^1 with the log-structure associated to the effective divisor $\infty \hookrightarrow \mathbf{P}^1$ (see [1, Def. 5.2.1]²). It is equipped with a functor $M : \mathbf{lSm} \rightarrow \mathbf{logDM}^{\text{eff}}$ associating the logarithmic motive $M(\mathfrak{X})$ of $\mathfrak{X} \in \mathbf{lSm}$.

Now we can state the main result of this paper.

Theorem 0.2. (Theorems 6.1.1 and 6.3) *There exists an exact and fully faithful functor*

$$(0.2.1) \quad \mathcal{L}og : \mathbf{RSC}_{\text{Nis}} \rightarrow \mathbf{Shv}_{d\text{Nis}}^{\text{ltr}} : F \rightarrow F^{\text{log}} = \mathcal{L}og(F)$$

such that F^{log} for $F \in \mathbf{RSC}_{\text{Nis}}$ is strictly $\overline{\square}$ -invariant in the sense [1, Def. 5.2.2]³. For $X \in \mathbf{Sm}$ we have a natural isomorphism

$$(0.2.2) \quad H_{\text{Nis}}^i(X, F_X) \simeq \text{Hom}_{\mathbf{logDM}^{\text{eff}}}(M(X, \text{triv}), L^{\overline{\square}} F^{\text{log}}[i]),$$

where $L^{\overline{\square}} : D(\mathbf{Shv}_{d\text{Nis}}^{\text{ltr}}) \rightarrow \mathbf{logDM}^{\text{eff}}$ is the localization functor and (X, triv) is the log-scheme with the trivial log-structure.

We remark (see Remark 5.5) that for $F = \Omega^i$, $F^{\text{log}}(\mathfrak{X})$ for $\mathfrak{X} \in \mathbf{lSm}$ whose underlying scheme is smooth, agrees with the sheaf of logarithmic differential forms of \mathfrak{X} at least assuming $\text{ch}(k) = 0$ ⁴.

²In fact it is defined in loc.cite. as the localization of the homotopy category of complexes in $\mathbf{Shv}_{d\text{Nis}}^{\text{ltr}}$ with respect to a $\overline{\square}$ -local descent model structure.

³It is an logarithmic analogue of Voevodsky's strict \mathbf{A}^1 -invariance.

⁴The assumption is necessary to use [13, Cor. 6.8] proved in case $\text{ch}(k) = 0$. We expect that it is removed by using a forthcoming work of K. Rülling extending [13, Cor. 6.8] to the case $\text{ch}(k) > 0$.

We now explain the organization of the paper.

In §1 we discuss some preliminaries and fix the notation. We recall the definitions and basic properties of *modulus (pre)sheaves with transfers* from [4], [5], [7] and [16]. It is a generalization of Voevodsky's (pre)sheaves with transfers to a version with modulus. The category $\underline{\mathbf{MCor}}$ of *modulus correspondences* is introduced. Its objects are pairs $\mathcal{X} = (\overline{X}, D)$, where \overline{X} is a separated scheme of finite type over k equipped with an effective Cartier divisor D such that the interior $\overline{X} - D = X$ is smooth. The morphisms are finite correspondences on the interiors satisfying some admissibility and a properness condition. Let $\underline{\mathbf{MPST}}$ be the category of additive presheaves of abelian groups on $\underline{\mathbf{MCor}}$. A full subcategory $\underline{\mathbf{MNST}} \subset \underline{\mathbf{MPST}}$ of *Nisnevich sheaves* is defined and there is a functor (see §1(20))

$$\underline{\omega}^{\mathbf{CI}} : \mathbf{RSC}_{\text{Nis}} \rightarrow \underline{\mathbf{MNST}}.$$

For every $F \in \mathbf{RSC}_{\text{Nis}}$ and $X \in \mathbf{Sm}$, it provides an exhaustive filtration on the group $F(X)$ of sections over X which measures depth of ramification along a boundary of a partial compactification of X : For $(\overline{X}, D) \in \underline{\mathbf{MCor}}$ with $\overline{X} - D = X$, we get the subgroups $\tilde{F}(\overline{X}, D) \subset F(X)$ with $\tilde{F} = \underline{\omega}^{\mathbf{CI}}F$ such that $\tilde{F}(\overline{X}, D_1) \subset \tilde{F}(\overline{X}, D_2)$ if $D_1 \leq D_2$.

In §2 we prove as a key technical input an analogue of Zariski-Nagata's purity theorem ([18, X 3.4]) for $\tilde{F}(\overline{X}, D)$ as above. It asserts the exactness of the sequence

$$0 \rightarrow \tilde{F}(\overline{X}, D) \rightarrow F(X) \rightarrow \bigoplus_{\xi \in D^{(0)}} \frac{F(\overline{X}_{|\xi}^h - \xi)}{\tilde{F}(\overline{X}_{|\xi}^h, \xi)},$$

in case $\overline{X} \in \mathbf{Sm}$ and D is reduced simple normal crossing divisor, where $D^{(0)}$ is the set of the irreducible components of D and $\overline{X}_{|\xi}^h$ is the henselization of \overline{X} at ξ . In [14], this result is generalized to the case where D may not be reduced under the assumption that \overline{X} admits a smooth compactification.

In §3 we review *higher local symbols* for reciprocity sheaves constructed in [14]. It is an effective tool with which one can decide when a given element of $F(X)$ with $F \in \mathbf{RSC}_{\text{Nis}}$ and $X \in \mathbf{Sm}$ belongs to $\tilde{F}(\overline{X}, D)$ as above. The construction of the pairing depends on push-forward maps for cohomology of reciprocity sheaves constructed in [2] (which means that Theorem 0.2.2 depends on the result of [2]).

In §4, we prove the following result: Let $\underline{\mathbf{MCor}}_{ls}^{\text{fin}}$ be the subcategory of $\underline{\mathbf{MCor}}$ whose objects are pairs (X, D) such that $X \in \mathbf{Sm}$ and the reduced parts D_{red} of D is a SNCD on X and whose morphisms

are modulus correspondences satisfying a finiteness conditions instead of the properness condition (see §1(5)). Then, for $F \in \mathbf{RSC}_{\text{Nis}}$, the association

$$\tilde{F}^{\log} : (X, D) \rightarrow \underline{\omega}^{\mathbf{CI}} F(X, D_{\text{red}})$$

gives a presheaf on $\mathbf{MCor}_{l_s}^{\text{fin}}$, which gives rise to a cohomology theory $H_{\log}^i(-, \tilde{F}^{\log})$ on $\mathbf{MCor}_{l_s}^{\text{fin}}$, called *the i -th logarithmic cohomology with coefficient F* (see Definition 4.4). The higher local symbols for F plays a fundamental role in the proof of the result .

In §5, we prove the invariance of logarithmic cohomology under blowups: Let $\Lambda_{l_s}^{\text{fin}}$ be the subcategory of $\mathbf{MCor}_{l_s}^{\text{fin}}$ whose objects are the same as $\mathbf{MCor}_{l_s}^{\text{fin}}$ and whose morphisms are those $\rho : (Y, E) \rightarrow (X, D)$ where $E = \rho^* D$ and ρ are induced by blowups of X in smooth centers $Z \subset D$ which are normal crossing to D (see the beginning of the section). Then, for $F \in \mathbf{RSC}_{\text{Nis}}$ and $\rho : \mathcal{Y} \rightarrow \mathcal{X}$ in $\Lambda_{l_s}^{\text{fin}}$, we have

$$\rho^* : H_{\log}^i(\mathcal{X}, F) \cong H_{\log}^i(\mathcal{Y}, F) \quad \text{for } \forall i \geq 0.$$

In §6, we prove Theorem 0.2.2, which is a formal consequence of the theorems in §4 and §5.

Acknowledgements. The author would like to thank Kay Rülling and F. Binda for many valuable discussions and comments. He is also grateful to A. Merici to whom he owes crucial ideas for §2.

1. PRELIMINARIES

We fix once and for all a perfect base field k . In this section we recall the definitions and basic properties of modulus sheaves with transfers from [4] and [16] (see also [7] for a more detailed summary).

- (1) Denote by \mathbf{Sch} the category of separated schemes of finite type over k and by \mathbf{Sm} the full subcategory of smooth schemes. For $X, Y \in \mathbf{Sm}$, an integral closed subscheme of $X \times Y$ that is finite and surjective over a connected component of X is called a *prime correspondence from X to Y* . The category \mathbf{Cor} of finite correspondences has the same objects as \mathbf{Sm} , and for $X, Y \in \mathbf{Sm}$, $\mathbf{Cor}(X, Y)$ is the free abelian group on the set of all prime correspondences from X to Y (see [17]). We consider \mathbf{Sm} as a subcategory of \mathbf{Cor} by regarding a morphism in \mathbf{Sm} as its graph in \mathbf{Cor} .

Let \mathbf{PST} be the category of additive presheaves of abelian groups on \mathbf{Cor} whose objects are called *presheaves with transfers*. Let $\mathbf{NST} \subseteq \mathbf{PST}$ be the category of Nisnevich sheaves

with transfers and let

$$a_{\mathbf{Nis}}^V : \mathbf{PST} \rightarrow \mathbf{NST}$$

be Voevodsky's Nisnevich sheafification functor, which is an exact left adjoint to the inclusion $\mathbf{NST} \rightarrow \mathbf{PST}$. Let $\mathbf{HI} \subseteq \mathbf{PST}$ be the category of \mathbf{A}^1 -invariant presheaves and put $\mathbf{HI}_{\mathbf{Nis}} = \mathbf{HI} \cap \mathbf{NST} \subseteq \mathbf{NST}$.

- (2) Let \mathbf{Sm}^{pro} be the category of k -schemes X which are essentially smooth over k , i.e. X is a limit $\varprojlim_{i \in I} X_i$ over a filtered set I , where X_i is smooth over k and all transition maps are étale. Note $\text{Spec } K \in \mathbf{Sm}^{\text{pro}}$ for a function field K over k thanks to the assumption that k is perfect. We define $\mathbf{Cor}^{\text{pro}}$ whose objects are the same as \mathbf{Sm}^{pro} and morphisms are defined as [13, Def. 2,2]. We extend $F \in \mathbf{PST}$ to a presheaf on $\mathbf{Cor}^{\text{pro}}$ by $F(X) := \varprojlim_{i \in I} F(X_i)$ for X as above.
- (3) We recall the definition of the category \mathbf{MCor} from [4, Definition 1.3.1]. A pair $\mathcal{X} = (X, D)$ of $X \in \mathbf{Sch}$ and an effective Cartier divisor D on X is called a *modulus pair* if $M - |M_\infty| \in \mathbf{Sm}$. Let $\mathcal{X} = (X, D_X)$, $\mathcal{Y} = (Y, D_Y)$ be modulus pairs and $\Gamma \in \mathbf{Cor}(X - D_X, Y - D_Y)$ be a prime correspondence. Let $\bar{\Gamma} \subseteq X \times Y$ be the closure of Γ , and let $\bar{\Gamma}^N \rightarrow X \times Y$ be the normalization. We say Γ is *admissible* (resp. *left proper*) if $(D_X)_{\bar{\Gamma}^N} \geq (D_Y)_{\bar{\Gamma}^N}$ (resp. if $\bar{\Gamma}$ is proper over X). Let $\mathbf{MCor}(\mathcal{X}, \mathcal{Y})$ be the subgroup of $\mathbf{Cor}(X - D_X, Y - D_Y)$ generated by all admissible left proper prime correspondences. The category \mathbf{MCor} has modulus pairs as objects and $\mathbf{MCor}(\mathcal{X}, \mathcal{Y})$ as the group of morphisms from \mathcal{X} to \mathcal{Y} .
- (4) Let $\mathbf{MCor}_{ls} \subset \mathbf{MCor}$ be the full subcategory of $(X, D) \in \mathbf{MCor}$ with $X \in \mathbf{Sm}$ and $|D|$ a normal crossing divisor on X .
- (5) Let $\mathbf{MCor}^{\text{fin}} \subset \mathbf{MCor}$ be the full subcategory of the same objects such that $\mathbf{MCor}^{\text{fin}}(\mathcal{X}, \mathcal{Y})$ are generated by all admissible *finite* prime correspondences, where finite prime correspondences are defined by replacing the left properness in (3) by finiteness. We also define $\mathbf{MCor}_{ls}^{\text{fin}} \subset \mathbf{MCor}_{ls}^{\text{fin}} \cap \mathbf{MCor}_{ls}$.
- (6) There is a canonical pair of adjoint functors $\lambda \dashv \underline{\omega}$:

$$\lambda : \mathbf{Cor} \rightarrow \mathbf{MCor} \quad X \mapsto (X, \emptyset),$$

$$\underline{\omega} : \mathbf{MCor} \rightarrow \mathbf{Cor} \quad (X, D) \mapsto X - |D|,$$

- (7) There is a full subcategory $\mathbf{MCor} \subset \mathbf{MCor}$ consisting of *proper modulus pairs*, where a modulus pair (X, D) is *proper* if X is

proper. Let $\tau : \mathbf{MCor} \hookrightarrow \underline{\mathbf{MCor}}$ be the inclusion functor and $\omega = \underline{\omega}\tau$.

- (8) Let \mathbf{MPST} (resp. $\underline{\mathbf{MPST}}$) be the category of additive presheaves of abelian groups on \mathbf{MCor} (resp. $\underline{\mathbf{MCor}}$) whose objects are called *modulus presheaves with transfers*. For $\mathcal{X} \in \mathbf{MCor}$, let $\mathbb{Z}_{\text{tr}}(\mathcal{X}) = \underline{\mathbf{MCor}}(-, \mathcal{X})$ be the representable object of $\underline{\mathbf{MPST}}$. We sometime write \mathcal{X} for $\mathbb{Z}_{\text{tr}}(\mathcal{X})$ for simplicity.
- (9) By the same manner as (2), the category $\underline{\mathbf{MCor}}^{\text{pro}}$ is defined and $F \in \underline{\mathbf{MPST}}$ is extended to a presheaf on $\underline{\mathbf{MCor}}^{\text{pro}}$ (see [13, §3.7]).
- (10) The adjunction $\lambda \dashv \underline{\omega}$ induce a string of 4 adjoint functors $(\lambda_! = \underline{\omega}^!, \lambda^* = \underline{\omega}_!, \lambda_* = \underline{\omega}^*, \underline{\omega}_*)$ (see [4, Pr. 2.3.1]):

$$\begin{array}{ccc} & \xleftarrow{\underline{\omega}^!} & \\ & \underline{\mathbf{MPST}} & \xrightarrow{\underline{\omega}_!} \mathbf{PST} \\ & \xleftarrow{\underline{\omega}^*} & \\ & \xrightarrow{\underline{\omega}_*} & \end{array}$$

where $\underline{\omega}_!, \underline{\omega}_*$ are localisations and $\underline{\omega}^!$ and $\underline{\omega}^*$ are fully faithful.

- (11) The functor τ yields a string of 3 adjoint functors $(\tau_!, \tau^*, \tau_*)$:

$$\begin{array}{ccc} & \xrightarrow{\tau_!} & \\ & \mathbf{MPST} & \xleftarrow{\tau^*} \underline{\mathbf{MPST}} \\ & \xrightarrow{\tau_*} & \end{array}$$

where $\tau_!, \tau_*$ are fully faithful and τ^* is a localisation; $\tau_!$ has a pro-left adjoint $\tau^!$, hence is exact (see [4, Pr. 2.4.1]). We will denote by $\underline{\mathbf{MPST}}^\tau$ the essential image of $\tau_!$ in $\underline{\mathbf{MPST}}$.

- (12) The modulus pair $\overline{\square} := (\mathbf{P}^1, \infty)$ has an interval structure induced by the one of \mathbf{A}^1 (see [7, Lem. 2.1.3]). We say $F \in \underline{\mathbf{MPST}}$ is $\overline{\square}$ -invariant if $p^* : F(\mathcal{X}) \rightarrow F(\mathcal{X} \otimes \overline{\square})$ is an isomorphism for any $\mathcal{X} \in \mathbf{MCor}$, where $p : \mathcal{X} \otimes \overline{\square} \rightarrow \mathcal{X}$ is the projection. Let \mathbf{CI} be the full subcategory of $\underline{\mathbf{MPST}}$ consisting of all $\overline{\square}$ -invariant objects and $\mathbf{CI}^\tau \subset \underline{\mathbf{MPST}}$ be the essential image of \mathbf{CI} under $\tau_!$.
- (13) Recall from [7, Theorem 2.1.8] that \mathbf{CI} is a Serre subcategory of $\underline{\mathbf{MPST}}$, and that the inclusion functor $i^{\overline{\square}} : \mathbf{CI} \rightarrow \underline{\mathbf{MPST}}$ has a left adjoint $h_0^{\overline{\square}}$ and a right adjoint $h_{\overline{\square}}^0$ given for $F \in \underline{\mathbf{MPST}}$ and $\mathcal{X} \in \mathbf{MCor}$ by

$$h_0^{\overline{\square}}(F)(\mathcal{X}) = \text{Coker}(i_0^* - i_1^* : F(\mathcal{X} \otimes \overline{\square}) \rightarrow F(\mathcal{X})),$$

$$h_{\overline{\square}}^0(F)(\mathcal{X}) = \text{Hom}(h_0^{\overline{\square}}(\mathcal{X}), F).$$

For $\mathcal{X} \in \mathbf{MCor}$, we write $h_0^{\overline{\square}}(\mathcal{X}) = h_0^{\overline{\square}}(\mathbb{Z}_{\text{tr}}(\mathcal{X})) \in \mathbf{CI}$, and by abuse of notation, we let $h_0^{\overline{\square}}(\mathcal{X})$ denote also for $\tau_! h_0^{\overline{\square}}(\mathcal{X}) \in \mathbf{CI}^\tau$.

- (14) For $F \in \underline{\mathbf{MPST}}$ and $\mathcal{X} = (X, D) \in \underline{\mathbf{MCor}}$, write $F_{\mathcal{X}}$ for the presheaf on the small étale site $X_{\text{ét}}$ over X given by $U \rightarrow F(\mathcal{X}_U)$ for $U \rightarrow X$ étale, where $\mathcal{X}_U = (U, D|_U) \in \underline{\mathbf{MCor}}$. We say F is a Nisnevich sheaf if so is $F_{\mathcal{X}}$ for all $\mathcal{X} \in \underline{\mathbf{MCor}}$ (see [4, Section 3]). We write $\underline{\mathbf{MNST}} \subset \underline{\mathbf{MPST}}$ for the full subcategory of Nisnevich sheaves and put

$$\underline{\mathbf{MNST}}^{\tau} = \underline{\mathbf{MNST}} \cap \underline{\mathbf{MPST}}^{\tau}, \quad \mathbf{CI}_{\text{Nis}}^{\tau} = \mathbf{CI}^{\tau} \cap \underline{\mathbf{MNST}}^{\tau}.$$

By [4, Prop. 3.5.3] and [5, Theorem 2], the inclusion functor $i_{\text{Nis}} : \underline{\mathbf{MNST}} \rightarrow \underline{\mathbf{MPST}}$ has an exact left adjoint $\underline{a}_{\text{Nis}}$ such that $\underline{a}_{\text{Nis}}(\underline{\mathbf{MPST}}^{\tau}) \subset \underline{\mathbf{MNST}}^{\tau}$. The functor $\underline{a}_{\text{Nis}}$ has the following description: For $F \in \underline{\mathbf{MPST}}$ and $\mathcal{Y} \in \underline{\mathbf{MCor}}$, let $F_{\mathcal{Y}, \text{Nis}}$ be the usual Nisnevich sheafification of $F_{\mathcal{Y}}$. Then, for $(X, D) \in \underline{\mathbf{MCor}}$ we have

$$\underline{a}_{\text{Nis}} F(X, D) = \varinjlim_{f: Y \rightarrow X} F_{(Y, f^*D), \text{Nis}}(Y)$$

where the colimit is taken over all proper maps $f : Y \rightarrow X$ that induce isomorphisms $Y - |f^*D| \xrightarrow{\sim} X - |D|$.

- (15) By [5, Pr. 6.2.1], $\underline{\omega}^*$ and $\underline{\omega}_!$ from (10) respect $\underline{\mathbf{MNST}}$ and \mathbf{NST} and induce a pair of adjoint functors (which for simplicity we write $\underline{\omega}_!$ and $\underline{\omega}^*$). Moreover, we have

$$\underline{\omega}_! \underline{a}_{\text{Nis}} = a_{\text{Nis}}^V \underline{\omega}_!.$$

By [7, Lem. 2.3.1] and [5, Pr. 6.2.1a)], for $F \in \mathbf{PST}$, we have $F \in \mathbf{HI}$ (resp $F \in \mathbf{HI}_{\text{Nis}}$) if and only if $\underline{\omega}^* F \in \mathbf{CI}^{\tau}$ (resp $\underline{\omega}^* F \in \mathbf{CI}_{\text{Nis}}^{\tau}$).

- (16) We say that $F \in \underline{\mathbf{MPST}}$ is *semi-pure* if the unit map

$$u : F \rightarrow \underline{\omega}^* \underline{\omega}_! F$$

is injective. For $F \in \underline{\mathbf{MPST}}$ (resp. $F \in \underline{\mathbf{MNST}}$), let $F^{sp} \in \underline{\mathbf{MPST}}$ (resp. $F^{sp} \in \underline{\mathbf{MNST}}$) be the image of $F \rightarrow \underline{\omega}^* \underline{\omega}_! F$ (called the semi-purification of F . See [16, Lem. 1.30]). For $F \in \underline{\mathbf{MPST}}$ we have

$$\underline{a}_{\text{Nis}}(F^{sp}) \simeq (\underline{a}_{\text{Nis}} F)^{sp}.$$

This follows from the fact that $\underline{a}_{\text{Nis}}$ is exact and commutes with $\underline{\omega}^* \underline{\omega}_!$. For $F \in \underline{\mathbf{MPST}}^{\tau}$ we have $F^{sp} \in \underline{\mathbf{MPST}}^{\tau}$ since τ is exact and $\underline{\omega}^* \underline{\omega}_! \tau_! = \tau_! \omega^* \omega_!$.

- (17) Let $\mathbf{CI}_{\text{Nis}}^{\tau, sp} \subset \mathbf{CI}^{\tau}$ be the full subcategory of semipure objects and consider the full subcategory

$$\mathbf{CI}_{\text{Nis}}^{\tau, sp} = \mathbf{CI}^{\tau, sp} \cap \underline{\mathbf{MNST}}^{\tau} \subset \mathbf{CI}_{\text{Nis}}^{\tau}.$$

By [16, Th. 0.1 and 0.4], we have $\underline{a}_{\mathbf{Nis}}(\mathbf{CI}^{\tau, sp}) \subset \mathbf{CI}_{\mathbf{Nis}}^{\tau, sp}$.

- (18) We write $\mathbf{RSC} \subseteq \mathbf{PST}$ for the essential image of \mathbf{CI} under $\omega_!$ (which is the same as the essential image of $\mathbf{CI}^{\tau, sp}$ under $\underline{\omega}_!$ since $\omega_! = \underline{\omega}_! \tau_!$ and $\omega_! F = \underline{\omega}_! F^{sp}$). Put $\mathbf{RSC}_{\mathbf{Nis}} = \mathbf{RSC} \cap \mathbf{NST}$. The objects of \mathbf{RSC} (resp. $\mathbf{RSC}_{\mathbf{Nis}}$) are called reciprocity presheaves (resp. sheaves). By [16, Th. 0.1], we have

$$(1.0.1) \quad a_{\mathbf{Nis}}^V(\mathbf{RSC}) \subset \mathbf{RSC}_{\mathbf{Nis}}.$$

We have $\mathbf{HI} \subseteq \mathbf{RSC}$ and it contains also smooth commutative group schemes (which may have non-trivial unipotent part), and the sheaf Ω^i of Kähler differentials, and the de Rham-Witt sheaves $W\Omega^i$ (see [6] and [7]).

- (19) \mathbf{NST} is a Grothendieck abelian category by [17, Lem. 3.1.6] and we can make $\mathbf{RSC}_{\mathbf{Nis}}$ its full sub-abelian category as follows: We define the kernel (resp. cokernel) of a map $\phi : F \rightarrow G$ in $\mathbf{RSC}_{\mathbf{Nis}}$ to be that of ϕ as a map in \mathbf{NST} . Here we need (1.0.1) to ensure that the cokernel of ϕ in \mathbf{NST} stays in $\mathbf{RSC}_{\mathbf{Nis}}$. By definition, a sequence $0 \rightarrow F \rightarrow G \rightarrow H \rightarrow 0$ is exact in $\mathbf{RSC}_{\mathbf{Nis}}$ if and only if it is exact in \mathbf{NST} .

- (20) By [7, Prop. 2.3.7] we have a pair of adjoint functors:

$$(1.0.2) \quad \mathbf{CI} \begin{array}{c} \xleftarrow{\omega^{\mathbf{CI}}} \\ \xrightarrow{\omega_!} \end{array} \mathbf{RSC},$$

where $\omega^{\mathbf{CI}} = h_{\square}^0 \omega^*$ and it is fully faithful. It induces a pair of adjoint functors:

$$(1.0.3) \quad \mathbf{CI}^{\tau} \begin{array}{c} \xleftarrow{\underline{\omega}^{\mathbf{CI}}} \\ \xrightarrow{\underline{\omega}_!} \end{array} \mathbf{RSC},$$

where $\underline{\omega}^{\mathbf{CI}} = \tau_! h_{\square}^0 \omega^*$ and it is fully faithful. Indeed, let $F = \tau_! \hat{F}$ for $\hat{F} \in \mathbf{CI}$ and $G \in \mathbf{RSC}$. In view of (13) and the exactness and full faithfulness of $\tau_!$, we have

$$\begin{aligned} \mathrm{Hom}_{\mathbf{CI}^{\tau}}(F, \tau_! h_{\square}^0 \omega^* G) &\simeq \mathrm{Hom}_{\mathbf{CI}}(\hat{F}, h_{\square}^0 \omega^* G) \simeq \\ \mathrm{Hom}_{\mathbf{MPST}}(\hat{F}, \omega^* G) &\simeq \mathrm{Hom}_{\mathbf{MPST}}(\tau_! \hat{F}, \underline{\omega}^* G) \simeq \mathrm{Hom}_{\mathbf{RSC}}(\omega_! F, G). \end{aligned}$$

In view of (15), (1.0.3) induce pair of adjoint functors:

$$(1.0.4) \quad \mathbf{CI}_{\mathbf{Nis}}^{\tau, sp} \begin{array}{c} \xleftarrow{\omega^{\mathbf{CI}}} \\ \xrightarrow{\underline{\omega}_!} \end{array} \mathbf{RSC}_{\mathbf{Nis}},$$

2. PURITY WITH REDUCED MODULUS

For $F \in \underline{\mathbf{MPST}}$, we put

$$F_{-1} = \text{Ker} \left(\underline{\text{Hom}}_{\underline{\mathbf{MPST}}}((\mathbf{P}^1 - 0, \infty), F) \xrightarrow{i_1^*} F \right),$$

$$F_{-1}^{(1)} = \text{Ker} \left(\underline{\text{Hom}}_{\underline{\mathbf{MPST}}}((\mathbf{P}^1, 0 + \infty), F) \xrightarrow{i_1^*} F \right),$$

Note that if $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$, we have for $\mathcal{X} \in \underline{\mathbf{MCor}}$

(2.0.1)

$$F_{-1}^{(1)}(\mathcal{X}) = \text{Hom}_{\underline{\mathbf{MPST}}}(h_{0, \text{Nis}}^{\square, \text{sp}}(\mathbf{P}^1, 0 + \infty)^0, \underline{\text{Hom}}_{\underline{\mathbf{MPST}}}(\mathbb{Z}_{\text{tr}}(\mathcal{X}), F)),$$

$$F_{-1}(\mathcal{X}) = \varinjlim_n \text{Hom}_{\underline{\mathbf{MPST}}}(h_{0, \text{Nis}}^{\square, \text{sp}}(\mathbf{P}^1, n \cdot 0 + \infty)^0, \underline{\text{Hom}}_{\underline{\mathbf{MPST}}}(\mathbb{Z}_{\text{tr}}(\mathcal{X}), F)),$$

where

$$h_{0, \text{Nis}}^{\square, \text{sp}}(\mathbf{P}^1, n \cdot 0 + \infty)^0 = \text{Coker} \left(\mathbb{Z} = \mathbb{Z}_{\text{tr}}(\text{Spec } k, \emptyset) \xrightarrow{i_1} h_{0, \text{Nis}}^{\square, \text{sp}}(\mathbf{P}^1, n \cdot 0 + \infty) \right).$$

Definition 2.1. For $e_1, \dots, e_r \in \{0, 1\}$, put

$$\tau^{(e_1, \dots, e_r)} F = \tau^{(e_r)} \dots \tau^{(e_1)} F,$$

where

$$\tau^{(0)} F = F_{-1} \quad \text{and} \quad \tau^{(1)} F = F_{-1}/F_{-1}^{(1)}.$$

The existence of retractions in the following lemma was suggested by A. Merici. It implies $\tau^{(e_1, \dots, e_r)} F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$ if $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$.

Lemma 2.2. For $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$, the inclusion $F_{-1}^{(1)} \rightarrow F_{-1}$ admits a retraction $s_F : F_{-1} \rightarrow F_{-1}^{(1)}$ such that for any map $\phi : F \rightarrow G$ in $\mathbf{CI}_{\text{Nis}}^{\tau, sp}$, the following diagram is commutative:

$$\begin{array}{ccc} F_{-1} & \xrightarrow{s_F} & F_{-1}^{(1)} \\ \downarrow \phi & & \downarrow \phi \\ G_{-1} & \xrightarrow{s_F} & G_{-1}^{(1)} \end{array}$$

In particular $\tau^{(1)} F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$ if $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$.

Proof. In view of (2.0.1), this follows from [2, Lem. 2.4]. \square

Theorem 2.3. Let $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$. Let $\mathcal{X} = \text{Spec } K\{t_1, \dots, t_n\}$ and $D = \{t_1^{e_1} \dots t_n^{e_n} = 0\} \subset \mathcal{X}$ with $e_1, \dots, e_n \in \{0, 1\}$. For a subset $I \subset [1, n]$ let $i_{\mathcal{H}} : \mathcal{H} \hookrightarrow \mathcal{X}$ be the closed immersion defined by $\{t_i = 0\}_{i \in I}$ and $D_{\mathcal{H}} = \{ \prod_{j \in [1, n] - I} t_j^{e_j} = 0 \} \subset \mathcal{H}$. Then

$$(2.3.1) \quad R^\nu i_{\mathcal{H}}^! F_{(\mathcal{X}, D)} = 0 \quad \text{for } \nu \neq q := |I|,$$

and there is an isomorphism

$$(2.3.2) \quad (\tau^{(e_I)} F)_{(\mathcal{H}, D_{\mathcal{H}})} \simeq R^q i_{\mathcal{H}}^! F_{(\mathcal{X}, D)} \quad \text{with } e_I = (e_i)_{i \in I} \in \mathbb{Z}_{\geq 0}^q.$$

Proof. The proof is divided into two steps.

Step 1: We prove (2.3.1) and (2.3.2) in case $q = |I| = 1$.

For $\nu = 0$ (2.3.1) follows from the semipurity of F . Thus it suffices to show (2.3.1) only for $\nu > 1$. Let $J = \{j \in [1, n] \mid e_j \neq 0\}$ and $r = |J|$. If $\dim(\mathcal{X}) = 0$, the assertion is trivial. If $r = 0$, the assertion follows from [16, Cor. 8.6(3)]. Assume $r > 0$ and $\dim(\mathcal{X}) \geq 1$, and proceed by the double induction on r and $\dim(\mathcal{X})$. Without loss of generality, we may assume

$$(\spadesuit) \quad e_1 \neq 0, \text{ and } \mathcal{H} = \{t_1 = 0\} \text{ if } \mathcal{H} \subset |D|.$$

Let $\iota : \mathcal{Z} \hookrightarrow \mathcal{X}$ be the closed immersion defined by $\{t_1 = 0\}$ and $D_{\mathcal{Z}} = \{t_2^{e_2} \cdots t_r^{e_r} = 0\} \subset \mathcal{Z}$ and $D' = \{t_2^{e_2} \cdots t_r^{e_r} = 0\} \subset \mathcal{X}$. By [16, Lem. 7.1], we have an exact sequence sheaves on \mathcal{X}_{Nis} :

$$0 \rightarrow F_{(\mathcal{X}, D')} \rightarrow F_{(\mathcal{X}, D)} \rightarrow \iota_* (F_{-1}^{(e_1)})_{(\mathcal{Z}, D_{\mathcal{Z}})} \rightarrow 0,$$

which gives rise to a long exact sequence of sheaves on \mathcal{H}_{Nis} :

$$(2.3.3) \quad \cdots \rightarrow R^\nu i_{\mathcal{H}}^! F_{(\mathcal{X}, D')} \rightarrow R^\nu i_{\mathcal{H}}^! F_{(\mathcal{X}, D)} \rightarrow R^\nu i_{\mathcal{H}}^! \iota_* (F_{-1}^{(e_1)})_{(\mathcal{Z}, D_{\mathcal{Z}})} \rightarrow \cdots.$$

By the induction hypothesis, $R^\nu i_{\mathcal{H}}^! F_{(\mathcal{X}, D')} = 0$ for $\nu > 1$. In case $\mathcal{H} \neq \mathcal{Z}$, we have a Cartesian diagram of closed immersions

$$\begin{array}{ccc} \mathcal{H} \cap \mathcal{Z} & \xrightarrow{\iota'} & \mathcal{H} \\ i_{\mathcal{H} \cap \mathcal{Z}} \downarrow & & \downarrow i_{\mathcal{H}} \\ \mathcal{Z} & \xrightarrow{\iota} & \mathcal{X} \end{array}$$

and we have an isomorphism

$$R^\nu i_{\mathcal{H}}^! \iota_* (F_{-1}^{(e_1)})_{(\mathcal{Z}, D_{\mathcal{Z}})} \simeq \iota'_* R^\nu i_{\mathcal{H} \cap \mathcal{Z}}^! (F_{-1}^{(e_1)})_{(\mathcal{Z}, D_{\mathcal{Z}})}.$$

By the induction hypothesis, $R^\nu i_{\mathcal{H} \cap \mathcal{Z}}^! (F_{-1}^{(e_1)})_{(\mathcal{Z}, D_{\mathcal{Z}})} = 0$ for $\nu > 1$ noting $F_{-1}^{(e_1)} \in \mathbf{CI}_{\text{Nis}}^{T, sp}$ by Lemma 2.2. So the desired vanishing follows from (2.3.3). Moreover, the assumptions (\spadesuit) and $\mathcal{H} \neq \mathcal{Z}$ imply that $\mathcal{H} \not\subset |D|$. Then (2.3.2) (with $q = 1$) follows from [16, Lem. 7.1(2)].

In case $\mathcal{Z} = \mathcal{H}$, we have

$$R^\nu i_{\mathcal{H}}^! \iota_* (F_{-1}^{(e_1)})_{(\mathcal{Z}, D_{\mathcal{Z}})} = R^\nu \iota^! \iota_* (F_{-1}^{(e_1)})_{(\mathcal{Z}, D_{\mathcal{Z}})},$$

which vanishes for $\nu > 0$. Hence (2.3.3) gives the desired vanishing together with an exact sequence:

$$0 \rightarrow (F_{-1}^{(e_1)})_{(\mathcal{H}, D_{\mathcal{H}})} \xrightarrow{\delta} R^1 i_{\mathcal{H}}^! F_{(\mathcal{X}, D')} \rightarrow R^1 i_{\mathcal{H}}^! F_{(\mathcal{X}, D)} \rightarrow 0.$$

By [16, Lem. 7.1(2)] we have an isomorphism

$$(F_{-1})_{(\mathcal{H}, D_{\mathcal{H}})} \simeq R^1 i_{\mathcal{H}}^! F_{(\mathcal{X}, D')}$$

through which δ is identified with the map induced by the canonical map $F_{-1}^{(e_1)} \rightarrow F_{-1}$. This proves the desired isomorphism (2.3.2) in case $\mathcal{Z} = \mathcal{H}$ and completes Step 1.

Step 2: We prove the theorem by the induction on q assuming $q > 0$. Let $I = \{i_1, \dots, i_q\} \subset [1, n]$ and $\mathcal{Y} \subset \mathcal{X}$ be the closed subscheme defined by $\{t_{i_1} = 0\}$. Let $i_{\mathcal{Y}} : \mathcal{Y} \hookrightarrow \mathcal{X}$ and $i_{\mathcal{H}, \mathcal{Y}} : \mathcal{H} \rightarrow \mathcal{Y}$ be the induced closed immersions. By Step 1 we have $R^\nu i_{\mathcal{Y}}^! F_{(\mathcal{X}, D)} = 0$ for $\nu \neq 1$ and we have an isomorphism

$$(\tau^{(e_{i_1})} F)_{(\mathcal{Y}, D_{\mathcal{Y}})} \simeq R^1 i_{\mathcal{Y}}^! F_{(\mathcal{X}, D)} \quad \text{with } D_{\mathcal{Y}} = \{t_1^{e_1} \cdots t_{i_1}^{e_{i_1}} \cdots t_n^{e_n} = 0\} \subset \mathcal{Y}.$$

Note $\tau^{(e_{i_1})} F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$ by Lemma 2.2. Thus, by the induction hypothesis, we have $R^\nu i_{\mathcal{H}, \mathcal{Y}}^! \tau^{(e_{i_1})} F_{(\mathcal{Y}, D_{\mathcal{Y}})} = 0$ for $\nu \neq q - 1$. By the spectral sequence

$$E_2^{a,b} = R^b i_{\mathcal{H}, \mathcal{Y}}^! R^a i_{\mathcal{Y}}^! F_{(\mathcal{X}, D)} \Rightarrow R^{a+b} i_{\mathcal{H}}^! F_{(\mathcal{X}, D)},$$

we get the desired vanishing (2.3.1) and an isomorphism

$$\begin{aligned} R^q i_{\mathcal{H}}^! F_{(\mathcal{X}, D)} &\simeq R^{q-1} i_{\mathcal{H}, \mathcal{Y}}^! R^1 i_{\mathcal{Y}}^! F_{(\mathcal{X}, D)} \simeq R^{q-1} i_{\mathcal{H}, \mathcal{Y}}^! (\tau^{(e_{i_1})} F)_{(\mathcal{Y}, D_{\mathcal{Y}})} \\ &\simeq (\tau^{(e_{i_2}, \dots, e_{i_q})} (\tau^{(e_{i_1})} F))_{(\mathcal{H}, D_{\mathcal{H}})} \simeq (\tau^{(e_{i_1}, e_{i_2}, \dots, e_{i_q})} F)_{(\mathcal{H}, D_{\mathcal{H}})}, \end{aligned}$$

where the third isomorphism holds by the induction hypothesis. This completes the proof of the theorem. \square

We say $\mathcal{X} = (X, D) \in \mathbf{MCor}$ reduced if so is D . The following corollaries 2.4 and 2.5 are immediate consequences of Theorem 2.3.

Corollary 2.4. *Take $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$ and $(X, D) \in \mathbf{MCor}_{ls}$ reduced. Let $x \in X^{(n)}$ with $K = k(x)$ and let $\mathcal{X} = X_{|x}^h$ be the henselization of X at x . Then*

$$H_x^i(X_{\text{Nis}}, F_{(X, D)}) = 0 \quad \text{for } i \neq n.$$

Choosing an isomorphism

$$\varepsilon : \mathcal{X} \simeq \text{Spec } K\{t_1, \dots, t_n\}$$

such that $D|_{\mathcal{X}} = \{t_1^{e_1} \cdots t_n^{e_n} = 0\} \subset \mathcal{X}$ with $e_1, \dots, e_n \in \{0, 1\}$, there exists an isomorphism depending on ε :

$$\theta_\varepsilon : \tau^{(e_1, e_2, \dots, e_n)} F(x) \simeq H_x^n(X_{\text{Nis}}, F_{(X, D)}).$$

Corollary 2.5. *For $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$ and $\mathcal{X} = (X, D) \in \underline{\mathbf{MCor}}_{l_s}$ reduced, the following sequence is exact:*

$$0 \rightarrow F(X, D) \rightarrow F(X - D, \emptyset) \rightarrow \bigoplus_{\xi \in D^{(0)}} \frac{F(X|_{\xi}^h - \xi, \emptyset)}{F(X|_{\xi}^h, \xi)}.$$

The idea of deducing the following corollary from the above is due to A. Merici.

Corollary 2.6. *Let $\mathcal{X} = (X, D) \in \underline{\mathbf{MCor}}_{l_s}$ be reduced.*

(1) *Assume given an exact sequence in $\underline{\mathbf{MNST}}$:*

$$(2.6.1) \quad 0 \rightarrow H \xrightarrow{\phi} G \xrightarrow{\psi} F$$

such that $F, G, H \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$ and that $\underline{\omega}_! \psi$ is surjective in \mathbf{NST} . If X is henselian local,

$$0 \rightarrow H(\mathcal{X}) \rightarrow G(\mathcal{X}) \rightarrow F(\mathcal{X}) \rightarrow 0$$

is exact.

(2) *Let $\gamma : F \rightarrow G$ be a map in $\mathbf{CI}_{\text{Nis}}^{\tau, sp}$ such that $\underline{\omega}_! \gamma$ is an isomorphism. Then $F(\mathcal{X}) \rightarrow G(\mathcal{X})$ is an isomorphism.*

(3) *For $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$, the unit map $u : F \rightarrow \underline{\omega}^{\mathbf{CI}} \underline{\omega}_! F$ induces an isomorphism $F(\mathcal{X}) \cong \underline{\omega}^{\mathbf{CI}} \underline{\omega}_! F(\mathcal{X})$.*

Proof. To show (1), it suffices to show the surjectivity of $G(\mathcal{X}) \rightarrow F(\mathcal{X})$. Let $\eta \in X$ be the generic point and consider the following commutative diagram of the Cousin complexes:

$$\begin{array}{ccccccc} 0 & \longrightarrow & H(\mathcal{X}) & \longrightarrow & H(\eta) & \longrightarrow & \bigoplus_{x \in X^{(1)}} H_x^1(X, H_{\mathcal{X}}) \longrightarrow \bigoplus_{y \in X^{(2)}} H_y^2(X, H_{\mathcal{X}}) \\ & & \downarrow & & \downarrow \phi(\eta) & & \downarrow H_x^1(\phi) \qquad \downarrow H_y^2(\phi) \\ 0 & \longrightarrow & G(\mathcal{X}) & \longrightarrow & G(\eta) & \longrightarrow & \bigoplus_{x \in X^{(1)}} H_x^1(X, G_{\mathcal{X}}) \longrightarrow \bigoplus_{y \in X^{(2)}} H_y^2(X, G_{\mathcal{X}}) \\ & & \downarrow & & \downarrow \psi(\eta) & & \downarrow H_x^1(\psi) \qquad \downarrow H_y^2(\psi) \\ 0 & \longrightarrow & F(\mathcal{X}) & \longrightarrow & F(\eta) & \longrightarrow & \bigoplus_{x \in X^{(1)}} H_x^1(X, F_{\mathcal{X}}) \longrightarrow \bigoplus_{y \in X^{(2)}} H_y^2(X, F_{\mathcal{X}}) \end{array}$$

By Corollary 2.4, the horizontal sequences are exact. By the assumption, $\psi(\eta)$ is surjective. By a diagram chase we are reduced to showing the following.

Claim 2.6.1. (i) For $x \in X^{(1)}$, the sequence

$$H_x^1(X, H_{\mathcal{X}}) \rightarrow H_x^1(X, G_{\mathcal{X}}) \rightarrow H_x^1(X, F_{\mathcal{X}})$$

is exact.

(ii) For $y \in X^{(2)}$, $H_y^2(\phi)$ is injective.

To show (i), by Corollary 2.4, it suffices to show the exactness of $\tau^{(e)}H \rightarrow \tau^{(e)}G \rightarrow \tau^{(e)}F$ for $e \in \{0, 1\}$. The case $e = 0$ follows from the left exactness of the endofunctor $\underline{\text{Hom}}_{\underline{\text{MPST}}}(\mathcal{X}, -)$ on $\underline{\text{MNST}}$ for any $\mathcal{X} \in \underline{\text{MCor}}$. We have a commutative diagram

$$\begin{array}{ccccc} \tau^{(1)}H & \xrightarrow{\phi} & \tau^{(1)}G & \xrightarrow{\psi} & \tau^{(1)}F \\ \uparrow p_H \downarrow s_H & & \uparrow p_G \downarrow s_G & & \uparrow p_F \downarrow s_F \\ \tau^{(0)}H & \xrightarrow{\phi} & \tau^{(0)}G & \xrightarrow{\psi} & \tau^{(0)}F \end{array}$$

where p_* are the projections and s_* is a right inverse of p_* coming from the retractions from Lemma 2.2. We have

$$\phi \circ p_H = p_G \circ \phi, \quad \psi \circ p_G = p_F \circ \psi, \quad \phi \circ s_H = s_G \circ \phi, \quad \psi \circ s_G = s_F \circ \psi.$$

By a diagram chase, the case $e = 1$ follows from the case $e = 0$.

To show (ii), by Corollary 2.4, it suffices to show the injectivity of $\tau^{(\underline{e})}H \rightarrow \tau^{(\underline{e})}G$ for $\underline{e} \in \{(0, 0), (0, 1), (1, 0), (1, 1)\}$. The case $\underline{e} = (0, 0)$ follows from the same left exactness as above, and the other cases from this case thanks to Lemma 2.2.

To show (2), we may assume \mathcal{X} is henselian local. Then it follows from (1). (3) follows from (2) since $\underline{\omega}_1 u$ is an isomorphism. This completes the proof of the corollary. \square

3. REVIEW ON HIGHER LOCAL SYMBOLS

In this section we recall from [14] the higher local symbols for reciprocity sheaves, which is a fundamental tool to prove Theorem 4.2, one of the main theorems of this paper.

3.1. Let X be an excellent separate scheme of pure dimension d . We denote by $X^{(c)}$ the set of points of codimension c . Let $x, y \in X$ be two points. We write

$$x > y :\iff \overline{\{x\}} \supset \overline{\{y\}}, \text{ i.e. } , y \in \overline{\{x\}}.$$

A *specialization chain* (or just *chain*) in X is a sequence

$$x = (x_1, \dots, x_n) \quad \text{with } x_1 > x_2 > \dots > x_n.$$

We say a specialization chain x is *maximal* or a *Parsin chain* if $n = d$ and $x_i \in X^{(i)}$ ⁵. We denote by $c(X)$ the set of specialization chains in

⁵The notation is different from [14] in which a specialization chain $x = (x_1, \dots, x_n)$ satisfies $x_n > \dots > x_1$ and x is maximal if the closures of x_i are of dimension i .

X and

$$\text{mc}(X) := \{x \in c(X) \mid x \text{ is maximal}\}.$$

If $d = 0$, the only element in $\text{mc}(X)$ is the empty chain.

We say a specialization chain $x = (x_1, \dots, x_n)$ is a Q -chain with break at r if $0 \leq r \leq n = d - 1$ and $x_i \in X^{(i)}$ for $i \leq r$ and $x_i \in X^{(i+1)}$ for $i > r$. We denote

$$Q_r(X) := \{Q\text{-chains with break at } r \text{ in } X\}.$$

For $x \in Q_r(X)$, we denote by $B(x)$ the set of all $y \in X$ such that

$$x(y) := (x_1, \dots, x_r, y, x_{r+1}, \dots, x_{d-1}) \in \text{mc}(X).$$

For a specialization chain $x = (x_1, \dots, x_n)$ in X , we let $\mathcal{O}_{X,x}^h$ be the henselization of X along x as defined in [3, Def. 1.6.2]: If $n = 1$, $\mathcal{O}_{X,x}^h$ is the henselization of $\mathcal{O}_{X,x}$. If $n > 1$, let $y = (x_2, \dots, x_n)$ and assume that we have defined $R = \mathcal{O}_{X,y}^h$. Let T be the set of all prime ideals of R lying over x_1 . Then we define $\mathcal{O}_{X,x}^h$ as the product of the henselization of R at \mathfrak{p} for all $\mathfrak{p} \in T$. By definition, $\mathcal{O}_{X,x}^h$ is a finite product of henselian local rings and we let $\mathfrak{m}_x \subset \mathcal{O}_{X,x}^h$ denote the product of its maximal ideals and $K_{X,x}^h$ denote its total fraction ring.

For $x \in Q_r(X)$ and $x(y)$ with $y \in B(x)$ as above, we have a natural inclusion of rings

$$(3.1.1) \quad \iota_y : K_{X,x}^h \rightarrow K_{X,x(y)}^h.$$

In what follows in this section, we fix $F \in \mathbf{RSC}_{\text{Nis}}$ and write $\tilde{F} = \omega_{\text{Nis}}^{\mathbf{CI}} F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$ (cf. (1.0.4)). We also fix a function field K over the base field k . Let X be an integral K -scheme of dimension d . Recall from [14] that we have a collection of bilinear pairings

$$(3.1.2) \quad \{(-, -)_{X/K,x} : F(K(X)) \otimes K_d^M(K_{X,x}^h) \rightarrow F(K)\}_{x \in \text{mc}(X)},$$

where we recall $K_{X,x}^h$ is a finite product of fields and $K_n^M(K_{X,x}^h)$ for an integer $n > 0$ denotes the product of the Milnor K -groups of the fields. We also note that $\mathcal{O}_{X,x}^h$ is a finite product of henselian dvr's and let $K_n^M(\mathcal{O}_{X,x}^h)$ denote the product of the Milnor K -groups of the dvr's.

For a local ring R and an ideal $I \subset R$, let $K_n^M(R, I) \subset K_n^M(R)$ denote the subgroup generated by symbols

$$\{1 + a, b_1, \dots, b_{n-1}\} \quad \text{with } a \in I, b_i \in R^\times.$$

The following properties hold for all $a \in F(K(X))$:

(HS1) Let $X \hookrightarrow X'$ be an open immersion where X' is an integral K -scheme of dimension d . Then we have $(a, \beta)_{X/K,x} = (a, \beta)_{X'/K,x}$ for all $\beta \in K_d^M(K_{X,x}^h)$.

(HS2) Let $x = (x_1, \dots, x_d) \in \text{mc}(X)$ and $X_1 \subset X$ be the closure of x_1 and $\delta = (x_2, \dots, x_d) \in \text{mc}(X_1)$. Then for all $\beta \in K_d^M(K_{X,x}^h)$

$$(a, \beta)_{X/K,x} = \begin{cases} \beta \cdot \text{Tr}_{E/K}(a), & \text{if } d = 0 \text{ and } E = K(X); \\ (a|_{X_1}, \partial_{x_1}\beta)_{X_1/K,\delta}, & \text{if } d \geq 1 \text{ and } a \in F(\mathcal{O}_{X,x_1}); \end{cases}$$

where $a|_{X_1} \in F(K(X_1))$ is the restriction of a , and

$$\partial_{x_1} : K_d^M(K_{X,x}^h) \rightarrow K_{d-1}^M(K_{X_1,\delta}^h)$$

is the tame symbol coming from the fact that $\mathcal{O}_{X,x}^h$ is a product of dvr 's and $K_{X_1,\delta}^h$ is identified with the product of its residue fields.

(HS3) Let $D \subset X$ be an effective Cartier divisor such that $(X, D) \in \mathbf{MCor}^{\text{pro}}$ (cf. §1(9)). Let $I_D \subset \mathcal{O}_X$ be the ideal sheaf of D . Assume $a \in \tilde{F}(X, D)$. Then, for all $x \in \text{mc}(X)$, we have

$$(a, \beta)_{X/K,x} = 0 \text{ for } \forall \beta \in K_d^M(\mathcal{O}_{X,x}^h, I_D \mathcal{O}_{X,x}^h).$$

(HS4) Let $x \in Q_r(X)$ with $0 \leq r \leq d-1$. Then $(a, \iota_y(\beta))_{X/K,x(y)} = 0$ for almost all $y \in B(x)$. If $r < d-1$ or X is projective over K ,

$$\sum_{y \in B(x)} (a, \iota_y(\beta))_{X/K,x(y)} = 0, \text{ for all } \beta \in K_d^M(K_{X,x}^h),$$

where $\iota_y : K_{X,x}^h \rightarrow K_{X,x(y)}^h$ is the natural map.

(HS5) Let $f : Y \rightarrow X$ be a finite surjective K -morphism between two integral K -schemes. Let $x \in \text{mc}(X)$ and $y \in \text{mc}(Y)$ with $f(y) = x$. Then we have

$$(f^*a, \beta)_{Y/K,y} = (a, \text{Nm}_{y/x}(\beta))_{X/K,x} \text{ (} a \in F(K(X)), \beta \in K_d^M(K_{Y,y}^h)\text{),}$$

$$(f_*b, \alpha)_{X/K,x} = \sum_{\substack{z \in \text{mc}(X') \\ f(z)=x}} (b, i_{x/z}\alpha)_{Y/K,z} \text{ (} b \in F(K(X')), \alpha \in K_d^M(K_{X,x}^h)\text{),}$$

where $\text{Nm}_{y/x} : K_d^M(K_{Y,y}^h) \rightarrow K_d^M(K_{X,x}^h)$ is the norm map on Milnor K -theory and $i_{x/z} : K_d^M(K_{X,x}^h) \rightarrow K_d^M(K_{Y,z}^h)$ is the inclusion map.

Now assume that $X \in \mathbf{Sm}$ of dimension d and $D \subset X$ be a reduced SNCD on X . For a function field K over k and a scheme Z over k , write $Z_K = Z \otimes_k K$ with $\phi_Z : Z_K \rightarrow Z$ the projection. If Z is integral, we denote by $K(Z)$ the function field of Z_K . We quote the following result from [14, Pr. 8.4].

Proposition 3.2. *Let $F \in \mathbf{RSC}_{\text{Nis}}$ and $a \in F(X - |D|)$. Assume that there exists an open subset $U \subset X$ which contains all generic points of D such that the following condition holds: For any function field K over k and any $x = (x_1, \dots, x_d) \in \text{mc}(U_K)$ with $x_1 \in D_K^{(0)}$, we have*

$$(\phi_X^*(a), \beta)_{U_K/K, x} = 0 \quad \text{for } \forall \beta \in K_d^M(\mathcal{O}_{X_K, x}^h, \mathbf{m}_x).$$

Then we have $a \in F(X, D)$.

4. LOGARITHMIC COHOMOLOGY OF RECIPROCITY SHEAVES

For $\mathcal{X} = (X, D) \in \underline{\mathbf{MCor}}_{ls}$, we write $\mathcal{X}_{\text{red}} = (X, |D|) \in \underline{\mathbf{MCor}}_{ls}$. We say $\mathcal{X} = (X, D) \in \underline{\mathbf{MCor}}_{ls}$ is reduced if $\mathcal{X} = \mathcal{X}_{\text{red}}$.

Definition 4.1. Let $F \in \underline{\mathbf{MPST}}$.

- (1) We say that F is *log-semipure* if for any $\mathcal{X} \in \underline{\mathbf{MCor}}_{ls}$, the map $F(\mathcal{X}_{\text{red}}) \rightarrow F(\mathcal{X})$ is injective. Note that if F is semipure, F is log-semipure.
- (2) We say that F is *logarithmic* if it is log-semipure and satisfies the condition that for $\mathcal{X}, \mathcal{Y} \in \underline{\mathbf{MCor}}_{ls}$ with \mathcal{X} reduced and $\alpha \in \underline{\mathbf{MCor}}^{\text{fin}}(\mathcal{Y}, \mathcal{X})$, the image of $\alpha^* : F(\mathcal{X}) \rightarrow F(\mathcal{Y})$ is contained in $F(\mathcal{Y}_{\text{red}}) \subset F(\mathcal{Y})$, where $\mathcal{Y}_{\text{red}} = (Y, E_{\text{red}})$ for $\mathcal{Y} = (Y, E)$.

Let $\underline{\mathbf{MPST}}_{\text{log}}$ be the full subcategory of $\underline{\mathbf{MPST}}$ consisting of logarithmic objects and put $\underline{\mathbf{MNST}}_{\text{log}} = \underline{\mathbf{MNST}} \cap \underline{\mathbf{MPST}}_{\text{log}}$.

Theorem 4.2. *Any $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$ is logarithmic, i.e. $\mathbf{CI}_{\text{Nis}}^{\tau, sp} \subset \underline{\mathbf{MNST}}_{\text{log}}$.*

We need a preliminary for the proof of the theorem.

Lemma 4.3. *Let $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$. Let $\mathbf{A}_K^n = \text{Spec } K[x_1, \dots, x_n]$ be the affine space over a function field K over k and $\mathcal{X} = \text{Spec } K\{x_1, \dots, x_n\}$ be the henselization of \mathbf{A}_K^n at the origin. Let $L_i = \{x_i = 0\} \subset \mathbf{A}^n$ and $\mathcal{L}_i = L_i \times_{\mathbf{A}_K^n} \mathcal{X}$ for $i \in [1, n]$. For an integer $0 < r \leq n$, the natural map*

$$K\{x_{r+1}, \dots, x_n\}[x_1, \dots, x_r] \rightarrow K\{x_1, \dots, x_n\}$$

induces a map in $\underline{\mathbf{MCor}}_{ls}$:

$$\rho_r : (\mathcal{X}, \mathcal{L}_1 + \dots + \mathcal{L}_r) \rightarrow (\mathbf{A}_S^r, \{x_1 \cdots x_r = 0\}) \simeq (\mathbf{A}^1, 0)^{\otimes r} \otimes (S, \emptyset),$$

where $S = \text{Spec } K\{x_{r+1}, \dots, x_n\}$. It induces

$$(4.3.1) \quad \rho_r^* : F(\mathbf{A}_S^r, \{x_1 \cdots x_r = 0\}) \rightarrow F(\mathcal{X}, \mathcal{L}_1 + \dots + \mathcal{L}_r)$$

Then $F(\mathcal{X}, \mathcal{L}_1 + \dots + \mathcal{L}_r)$ is generated by the image of ρ_r^ and*

$$F(\mathcal{X}, \mathcal{L}_1 + \dots + \mathcal{L}_i + \dots + \mathcal{L}_r) \quad \text{for } i = 1, \dots, r.$$

Proof. For $\mathcal{Y} \in \underline{\mathbf{M}}\mathbf{Cor}$, let $F^{\mathcal{Y}} \in \underline{\mathbf{M}}\mathbf{PST}$ be defined by $F^{\mathcal{Y}}(\mathcal{Z}) = F(\mathcal{Y} \otimes \mathcal{Z})$. By [2, Lem. 1.5(3)], $F^{\mathcal{Y}} \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$ for $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$. We prove the lemma by the induction on r . The case $r = 1$ holds since by [16, Lem. 7.1 and Lem 5.9], ρ_1 induces an isomorphism

$$F^{(\mathbf{A}^1, 0)}(S)/F^{(\mathbf{A}^1, \emptyset)}(S) \xrightarrow{\simeq} F(\mathcal{X}, \mathcal{L}_1)/F(\mathcal{X}).$$

By definition $\mathcal{L}_1 = \text{Spec } K\{x_2, \dots, x_n\}$ and we have a map in $\underline{\mathbf{M}}\mathbf{Cor}$:

$$(\mathcal{X}, \mathcal{L}_1 + \dots + \mathcal{L}_r) \rightarrow (\mathbf{A}^1, 0) \otimes (\mathcal{L}_1, \mathcal{L}_1 \cap (\mathcal{L}_2 + \dots + \mathcal{L}_r))$$

induced by the natural map $K\{x_2, \dots, x_n\}[x_1] \rightarrow K\{x_1, \dots, x_n\}$. By [16, Lem. 7.1 and Lem 5.9], it induces an isomorphism

$$F^{(\mathbf{A}^1, 0)}(\mathcal{L}_1, E)/F^{(\mathbf{A}^1, \emptyset)}(\mathcal{L}_1, E) \xrightarrow{\simeq} F(\mathcal{X}, \mathcal{L}_1 + \dots + \mathcal{L}_r)/F(\mathcal{X}, \mathcal{L}_2 + \dots + \mathcal{L}_r)$$

with $E = \mathcal{L}_1 \cap (\mathcal{L}_2 + \dots + \mathcal{L}_r)$. By the induction hypothesis, $F^{(\mathbf{A}^1, 0)}(\mathcal{L}_1, E)$ is generated by $F^{(\mathbf{A}^1, 0)}(\mathcal{L}_1, E_j)$ with $E_j = \mathcal{L}_1 \cap (\mathcal{L}_2 \cdots + \mathcal{L}_j \overset{\vee}{+} \cdots \mathcal{L}_r)$ for $j = 2, \dots, r$ together with the image of the map

$$(F^{(\mathbf{A}^1, 0)})^{(\mathbf{A}^1, 0)^{\otimes r-1}}(S) = F^{(\mathbf{A}^1, 0)^{\otimes r}}(S) \rightarrow F^{(\mathbf{A}^1, 0)}(\mathcal{L}_1, E)$$

induced by

$$(\mathcal{L}_1, E) \rightarrow (\mathbf{A}_S^{r-1}, \{x_2 \cdots x_r = 0\}) \simeq (\mathbf{A}^1, 0)^{\otimes r-1} \otimes (S, \emptyset)$$

coming from the map $K\{x_{r+1}, \dots, x_n\}[x_2, \dots, x_r] \rightarrow K\{x_2, \dots, x_d\}$. This proves the lemma. \square

Proof of Theorem 4.2: By Corollary 2.6(3), we may assume $F = \underline{\omega}^{\mathbf{CI}G}$ for $G \in \mathbf{RSC}_{\text{Nis}}$. Take $\mathcal{X} = (X, D), \mathcal{Y} = (Y, E) \in \underline{\mathbf{M}}\mathbf{Cor}_{\text{ts}}$ with \mathcal{X} reduced and let $\alpha \in \underline{\mathbf{M}}\mathbf{Cor}^{\text{fm}}(\mathcal{Y}, \mathcal{X})$ be an elementary correspondence. We need to show that $\alpha^*(F(\mathcal{X})) \subset F(\mathcal{Y}_{\text{red}})$. The question is Nisnevich local over X and Y . Hence we may assume $(X, D) = (\mathcal{X}, \mathcal{L}_1 + \dots + \mathcal{L}_r) \in \underline{\mathbf{M}}\mathbf{Cor}^{\text{pro}}$ under the notation from Lemma 4.3. If $r = 0$, we have $\alpha \in \underline{\mathbf{M}}\mathbf{Cor}((Y, \emptyset), (X, \emptyset))$ by the assumption $\alpha \in \underline{\mathbf{M}}\mathbf{Cor}^{\text{fm}}(\mathcal{Y}, \mathcal{X})$ so that

$$\alpha^*(F(\mathcal{X})) = \alpha^*(F(X, \emptyset)) \subset F(Y, \emptyset) \subset F(\mathcal{Y}_{\text{red}}).$$

Assume $r > 0$ and proceed by the induction on r . By Lemma 4.3, we may assume then

$$(X, D) = \mathcal{M} := (\mathbf{A}^1, 0)^{\otimes r} \otimes (S, \emptyset) \text{ for } S \in \mathbf{Sm}.$$

On the other hand, by Corollary 2.5, we have an exact sequence

$$0 \rightarrow F(Y, E_{\text{red}}) \rightarrow F(Y - E_{\text{red}}, \emptyset) \rightarrow \bigoplus_{\xi \in E^{(0)}} \frac{F(Y|_{\xi}^h - \xi, \emptyset)}{F(Y|_{\xi}^h, \xi)}.$$

Hence we may replace Y with its Nisnevich neighborhood of a generic point ξ of E . Using the assumption that k is perfect, we may then assume the following condition (\spadesuit). Recall that α is by definition an integral closed subschem of $(Y - E) \times (X - D)$ finite surjective over $Y - E$ and its closure $\bar{\alpha}$ in $Y \times X$ is finite surjective over Y .

- (\spadesuit) X and Y are irreducible and α is smooth. The normalization Y' of $\bar{\alpha}$ is smooth and $E' := E \times_Y Y' \subset Y'$ is irreducible and E'_{red} is smooth.

Let $g : Y' \rightarrow Y$ and $f : Y' \rightarrow X$ be the induced maps. We have $E' = g^*E \geq f^*D$ as Cartier divisors on Y' by the modulus condition for α . Hence these maps induce

$$F(X, D) \xrightarrow{f^*} F(Y', E') \xrightarrow{g^*} F(Y, E).$$

We claim that $\alpha^* : F(X, D) \rightarrow F(Y, E)$ agrees with this map. Indeed, this follows from the equality

$$\Gamma_f \circ {}^t \Gamma_g = \alpha \in \mathbf{Cor}(Y - E, X - D),$$

where ${}^t \Gamma_g \in \mathbf{Cor}(Y - E, Y' - E')$ is the transpose of the graph of g and $\Gamma_f \in \mathbf{Cor}(Y' - E', X - D)$ is the graph of f . By definition this follows from the equality

$${}^t \Gamma_g \times_{Y' - E'} \Gamma_f = \alpha \subset (Y - E) \times (X - D)$$

which one can check easily noting $Y' \rightarrow \bar{\alpha}$ is an isomorphism over α assumed to be smooth. Then we get a commutative diagram

$$\begin{array}{ccccc} & & F(Y', E'_{\text{red}}) & & \\ & & \downarrow \hookrightarrow & & \\ & & F(Y', E'_{\text{red}} \times_Y Y') & \xrightarrow{g^*} & F(Y, E'_{\text{red}}) \\ & & \downarrow \hookrightarrow & & \downarrow \hookrightarrow \\ F(X, D) & \xrightarrow{f^*} & F(Y', E') & \xrightarrow{g^*} & F(Y, E) \end{array}$$

where the top inclusion comes from $E'_{\text{red}} \times_Y Y' \geq E'_{\text{red}}$ as Cartier divisors on Y' thanks to the sempurity of F (cf. §1(16)). Hence it suffices to show $f^*(F(X, D)) \subset F(Y', E'_{\text{red}})$. By replacing (Y, E) with (Y', E') , we may now assume that α is induced by a morphism $f : Y \rightarrow X = \mathbf{A}^r \times S$. Then α factors in \mathbf{MCor} as

$$(Y, E) \rightarrow (\mathbf{A}^1, 0)^{\otimes r} \otimes (Y, \emptyset) \rightarrow (\mathbf{A}^1, 0)^{\otimes r} \otimes (S, \emptyset).$$

where the first map is induced by the map

$$i = (pr_{\mathbf{A}^r} \circ f, id_Y) : Y \rightarrow \mathbf{A}^r \times Y,$$

and the second induced by

$$id_{\mathbf{A}^r} \times (pr_S \circ f) : \mathbf{A}^r \times Y \rightarrow \mathbf{A}^r \times S.$$

Note that i is a section of the projection $\mathbf{A}^r \times Y \rightarrow Y$. Thus we are reduced to showing $i^*(F((\mathbf{A}^1, 0)^{\otimes r} \otimes (Y, \emptyset))) \subset F(Y, E_{\text{red}})$. By Proposition 3.2 this follows from the following.

Claim 4.3.1. Assume $F = \omega^{\mathbf{CI}}G$ for some $G \in \mathbf{RSC}_{\text{Nis}}$. Take $a \in F((\mathbf{A}^1, 0)^{\otimes r} \otimes (Y, \emptyset))$. Let K be a function field over k . After replacing Y by an open subset containing ξ , we have

$$(i^*(a)_K, \gamma)_{Y_K/K, \delta} = 0 \text{ for } \forall \gamma \in K_e^M(\mathcal{O}_{Y_K, \delta}^h, \mathfrak{m}_\delta)$$

and for any $\delta = (\xi, \delta_1, \dots, \delta_{e-1}) \in \text{mc}(Y_K)$, where $e = \dim(Y)$ and $\xi \in E$ is the generic point, and

$$(-, -)_{Y_K/K, \delta} : F(K(Y)) \otimes K_d^M(K_{Y_K, \delta}^h) \rightarrow F(K)$$

is from (3.1.2).

Proof. Write

$$\mathbf{A}^r \times Y = \text{Spec } A[x_1, \dots, x_r], \quad (\mathbf{A}^1, 0)^{\otimes r} \otimes (Y, \emptyset) = (\mathbf{A}_Y^r, \{x_1 \cdots x_r = 0\}).$$

After replacing Y by an open subset containing ξ , we can write

$$i(Y) = \bigcap_{1 \leq i \leq r} \{x_i - u_i \pi^{m_i} = 0\} \text{ with } m_i \in \mathbb{Z}_{\geq 0}, u_i \in A^\times.$$

Let $\delta = (\xi, \delta_1, \dots, \delta_{e-1})$ be as in the claim and put $\delta' = (\delta_1, \dots, \delta_{e-1}) \in \text{mc}((E_{\text{red}})_K)$. Let $X_K = \mathbf{A}^r \times Y_K$ and z_j for $1 \leq j \leq r$ be the generic point of

$$Z_j = \bigcap_{1 \leq i \leq j} \{x_i - u_i \pi^{m_i} = 0\} \subset X_K,$$

and w_j for $1 \leq j \leq r$ be the generic point of

$$W_j = \{\pi = x_1 = \cdots = x_j = 0\} = F \cap Z_j \text{ with } F = \{\pi = 0\} \subset X_K.$$

The section i induces isomorphisms

$$(4.3.2) \quad Y_K \simeq Z_r \text{ and } (E_{\text{red}})_K \simeq W_r.$$

Let $\sigma = (\eta_1, w_1, \dots, w_r, i(\delta')) \in \text{mc}(X_K)$, where η_1 is the generic point of $D_1 = \{x_1 = 0\} \subset X_K$ and $i(\delta') \in \text{mc}(W_r)$ is the image of δ' under (4.3.2). Let $\tilde{\sigma}$ be the chain $(w_r, i(\delta'))$ in X_K . By the definition of henselization along chains, the projection $X = \mathbf{A}^r \times Y \rightarrow Y$ induces $\iota : \mathcal{O}_{Y_K, \delta}^h \rightarrow \mathcal{O}_{X_K, \tilde{\sigma}}^h$. Take any $\gamma \in K_e^M(\mathcal{O}_{Y_K, \delta}^h, \mathfrak{m}_\delta)$ and put

$$(4.3.3) \quad \beta = \{\iota(\gamma), \frac{u_1 \pi^{m_1} - x_1}{u_1 \pi^{m_1}}, \dots, \frac{u_r \pi^{m_r} - x_r}{u_r \pi^{m_r}}\} \in K_d^M(K_{X_K, \tilde{\sigma}}^h) \text{ (} d = e + r\text{),}$$

For $a \in F((\mathbf{A}^1, 0)^{\otimes r} \otimes (Y, \emptyset))$ and its restriction $a_K \in F((\mathbf{A}^1, 0)^{\otimes r} \otimes (Y_K, \emptyset))$, we have⁶

$$\begin{aligned} 0 &= (a_K, \beta)_{X_K/K, \sigma} = - \sum_{\substack{\tau \in X_K^{(1)} \\ \tau > w_1, \tau \neq \eta_1}} (a_K, \beta)_{X_K/K, (\tau, w_1, \dots, w_r, i(\delta'))} \\ &= -(a_K, \beta)_{X_K/K, (z_1, w_1, \dots, w_r, i(\delta'))} \\ &= \pm((a_K)|_{Z_1}, \beta_1)_{Z_1/K, (w_1, \dots, w_r, i(\delta'))}, \\ \beta_1 &= \left\{ \iota_1(\gamma), \frac{u_2 \pi^{m_2} - x_2}{u_2 \pi^{m_2}}, \dots, \frac{u_r \pi^{m_r} - x_r}{u_r \pi^{m_r}} \right\} \in K_{d-1}^M(K_{Z_1, \tilde{\sigma}_1}^h) \end{aligned}$$

where $\tilde{\sigma}_1$ is the chain $(w_r, i(\delta'))$ in Z_1 and $\iota_1 : \mathcal{O}_{Y_K, \delta}^h \rightarrow \mathcal{O}_{Z_1, \tilde{\sigma}_1}^h$ is induced by $Z_1 \hookrightarrow X_K \rightarrow Y_K$. The first equality follows from §3 (HS3) applied to $D_1 \subset X_K$ noting that β lies in $K_d^M(\mathcal{O}_{X_K, \sigma}^h, \mathfrak{m}_\sigma)$ since $(u_1 \pi^{m_1} - x_1)/u_1 \pi^{m_1} \in 1 + x_1 \mathcal{O}_{X_K, \eta_1}$. The second follows from (HS4) applied to $x = (w_1, \dots, w_r, i(\delta')) \in Q_0(X_K)$. The third equality holds since z_1 is the unique $\tau \in X_K^{(1)} - \{\eta_1\}$ such that $\tau > w_1$ and $(a_K, \beta)_{X_K/K, (\tau, w_1, \dots, w_r, i(\delta'))}$ may not vanish, which follows from (HS2) noting $\iota(\gamma)|_F = 0$. Finally the last equality follows from (HS2). We further get

$$\begin{aligned} 0 &= ((a_K)|_{Z_1}, \beta_1)_{Z_1/K, (w_1, w_2, \dots, w_r, i(\delta'))} = - \sum_{\substack{\tau \in Z_1^{(1)} \\ \tau > w_2, \tau \neq w_1}} ((a_K)|_{Z_1}, \beta_1)_{Z_1/K, (\tau, w_2, \dots, w_r, i(\delta'))} \\ &= -((a_K)|_{Z_1}, \beta_1)_{Z_1/K, (z_2, w_2, \dots, w_r, i(\delta'))} \\ &= \pm((a_K)|_{Z_2}, \beta_2)_{Z_2/K, (w_2, \dots, w_r, i(\delta'))}, \\ \beta_2 &= \left\{ \iota_2(\gamma), \frac{u_3 \pi^{m_3} - x_3}{u_3 \pi^{m_3}}, \dots, \frac{u_r \pi^{m_r} - x_r}{u_r \pi^{m_r}} \right\} \in K_{d-1}^M(K_{Z_2, \tilde{\sigma}_2}^h), \end{aligned}$$

where $\tilde{\sigma}_2$ is the chain $(w_r, i(\delta'))$ in Z_2 and $\iota_2 : \mathcal{O}_{Y_K, \delta}^h \rightarrow \mathcal{O}_{Z_2, \tilde{\sigma}_2}^h$ is induced by $Z_2 \hookrightarrow X_K \rightarrow Y_K$. The above equalities hold by the same arguments as above except that for the third equality, there are a priori two $\tau \in Z_1^{(1)} - \{w_1\}$ with $\tau > w_2$ for which $((a_K)|_{Z_1}, \beta_1)_{Z_1/K, (\tau, w_2, \dots, w_r, i(\delta'))}$ may not vanish. One is z_2 and another is the generic point η_2 of $Z_1 \cap D_2$ with $D_2 = \{x_2 = 0\} \subset X_K$, but $((a_K)|_{Z_1}, \beta_1)_{Z_1/K, (\eta_2, w_2, \dots, w_r, i(\delta'))} = 0$. Indeed, $(a_K)|_{Z_1} \in F(\text{Spec}(\mathcal{O}_{Z_1, \eta_2}), \eta_2)$ since Z_1 and D_2 intersect transversally in X_K . Hence the vanishing follows from (HS3) applied to $Z_1 \cap D_2 \subset Z_1$

⁶By abuse of notation, for $\rho \in \text{mc}(X_K)$ containing $\tilde{\sigma}$ as a sub-chain, β in $(a_K, \beta)_{X_K/K, \rho}$ denotes the image of β from (4.3.3) under the natural map $K_d^M(K_{X_K, \tilde{\sigma}}^h) \rightarrow K_d^M(K_{X_K, \rho}^h)$. The same convention for β_1 and β_2 below.

noting $\beta_1 \in K_d^M(\mathcal{O}_{Z_1, (\eta_2, \bar{\sigma}_1)}, \mathfrak{m}_{\eta_2})$. Repeating the same arguments, we finally get

$$0 = ((a_K)|_{Z_r}, \iota_r(\gamma))_{Z_r/K, (w_r, i(\delta'))} = ((a_K)|_{Y_K}, \gamma)_{Y_K/K, \delta},$$

where $\iota_r : \mathcal{O}_{Y_K, \delta}^h \rightarrow \mathcal{O}_{Z_r, (w_r, i(\delta'))}^h$ is induced by $Z_r \hookrightarrow X_K \rightarrow Y_K$ and the second equality follows from (4.3.2). This completes the proof of the claim and Theorem 4.2. \square

Definition 4.4. For $F \in \underline{\mathbf{MNST}}_{\log}$ and an integer $i \geq 0$, consider the association

$$H_{\log}^i(-, F) : \underline{\mathbf{MCor}}_{ls}^{\text{fin}} \rightarrow \mathbf{Ab}; (X, D) \rightarrow H^i(X_{\text{Nis}}, F_{(X, D_{\text{red}})}).$$

By the definition this gives a presheaf on $\underline{\mathbf{MCor}}_{ls}^{\text{fin}}$, which we call *the i -th logarithmic cohomology with coefficient F* .

5. INVARIANCE OF LOGARITHMIC COHOMOLOGY UNDER BLOWUPS

Let the notation be as in §4. Let $\Lambda_{ls}^{\text{fin}}$ be the subcategory of $\underline{\mathbf{MCor}}_{ls}^{\text{fin}}$ whose objects are the same as $\underline{\mathbf{MCor}}_{ls}^{\text{fin}}$ and whose morphisms are those $\rho : (Y, E) \rightarrow (X, D)$ where $E = \rho^*D$ and ρ are induced by blowups of X in smooth centers $Z \subset D$ which are normal crossing to D in the following sense: For any point x of D , there exists a system z_1, \dots, z_d of regular parameters of X at x satisfying the conditions:

- Locally at x , $Z = \{z_1 = \dots = z_r = 0\}$ with $r = \text{codim}_X Z$.
- Locally at x , $|D| = \{\prod_{j \in J} z_j = 0\}$ for some $J \subset \{1, \dots, r\}$.

Theorem 5.1. For $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$ and $\rho : \mathcal{Y} \rightarrow \mathcal{X}$ in $\Lambda_{ls}^{\text{fin}}$, we have

$$(5.1.1) \quad \rho^* : H_{\log}^i(\mathcal{X}, F) \cong H_{\log}^i(\mathcal{Y}, F) \quad \text{for } \forall i \geq 0.$$

Proof. Writing $\mathcal{Y} = (Y, E)$ and $\mathcal{X} = (X, D)$, ρ is induced by a blowup $\rho : Y \rightarrow X$ in a smooth center $Z \subset D$ normal crossing to D . First we prove the theorem in case $i = 0$. We may assume that D is reduced and $E = \rho^*D$. Then ρ is invertible in $\underline{\mathbf{MCor}}$ so that $\rho^* : F(\mathcal{X}) \cong F(\mathcal{Y})$. Since this factors through $F(Y, E_{\text{red}})$, we get (5.1.1) for $i = 0$.

To show (5.1.1) for $i > 0$, it now suffices to prove $R^i \rho_* F_{(Y, E_{\text{red}})} = 0$. By [8, Lem. 9], Nisnevich locally around a point of Z , (X, D) is isomorphic to

$$(\mathbf{A}^c, L_1 + \dots + L_r) \otimes \mathcal{W} \quad \text{with } \mathcal{W} = (W, W^\infty) \in \underline{\mathbf{MCor}}_{ls},$$

where $\mathbf{A}^c = \text{Spec } k[t_1, \dots, t_c]$ with $c = \text{codim}_z(Z, X)$ and $L_i = V(t_i)$ for $i = 1, \dots, r$ with $1 \leq r \leq c$, and Z corresponds to $0 \times W$. Hence the theorem follows from the following proposition. \square

Proposition 5.2. *Let $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$ and $\mathcal{W} = (W, W^\infty) \in \mathbf{MCor}_{ls}$. Let $\mathbf{A}^n = \text{Spec } k[t_1, \dots, t_n]$ and put $L_i = V(t_i)$ for $1 \leq i \leq n$. Let $\rho : Y \rightarrow \mathbf{A}^n$ be the blow-up at the origin $0 \in \mathbf{A}^n$ and $\tilde{L}_i \subset Y$ be the strict transforms of L_i for $i = 1, \dots, r$ and $E = \rho^{-1}(0) \subset Y$. For any $1 \leq r \leq n$, we have*

$$(5.2.1) \quad R^i \rho_{W*} F_{(Y, \tilde{L}_1 + \dots + \tilde{L}_r + E) \otimes \mathcal{W}} = 0 \quad \text{for } i \geq 1,$$

where $\rho_W := \rho \times \text{id}_W : Y \times W \rightarrow \mathbf{A}^2 \times W$.

Lemma 5.3. *Proposition 5.2 holds for $n = 2$.*

Proof. We can assume W is henselian local. The case $r = 1$ is proved in [2, Lem. 2.13] and we show the case $r = 2$.⁷ Put $D = L_1 + L_2$. By the case $i = 0$ of Theorem 5.1, we get

$$(5.3.1) \quad F_{(\mathbf{A}^2, D) \otimes \mathcal{W}} \cong \rho_{W*} F_{(Y, \tilde{L}_1 + \tilde{L}_2 + E) \otimes \mathcal{W}}.$$

Set

$$\mathcal{F} := F_{(Y, \tilde{L}_1 + \tilde{L}_2 + E) \otimes \mathcal{W}}.$$

Since $R^i \rho_{W*} \mathcal{F}$ for $i \geq 1$ is supported in $0 \times W$ we have

$$R^i \rho_{W*} \mathcal{F} = 0 \iff H^0(\mathbf{A}_W^2, R^i \rho_{W*} \mathcal{F}) = 0,$$

where $\mathbf{A}_W^2 = \mathbf{A}^2 \times W$, and

$$H^j(\mathbf{A}_W^2, R^i \rho_{W*} \mathcal{F}) = 0, \quad \text{for all } i, j \geq 1.$$

By (5.3.1) and [2, Lem. 2.11]

$$H^i(\mathbf{A}_W^2, \rho_{W*} \mathcal{F}) = H^i(\mathbf{A}_W^2, F_{(\mathbf{A}^2, D) \otimes \mathcal{W}}) = 0.$$

Thus the Leray spectral sequence yields

$$H^0(\mathbf{A}_W^2, R^i \rho_{W*} \mathcal{F}) = H^i(Y \times W, \mathcal{F}), \quad i \geq 0,$$

and we have to show, that this group vanishes for $i \geq 1$. We can write

$$\mathbf{A}^2 = \text{Spec } k[x, y] \quad \text{and} \quad L_1 = V(x), \quad L_2 = V(y) \subset \mathbf{A}^2.$$

Then we have

$$Y = \text{Proj } k[x, y][S, T]/(xT - yS) \subset \mathbf{A}^2 \times \mathbf{P}^1.$$

Denote by

$$\pi_0 : Y \hookrightarrow \mathbf{A}^2 \times \mathbf{P}^1 \rightarrow \mathbf{P}^1 = \text{Proj } k[S, T]$$

the morphism induced by projection and let $\pi : Y \times W \rightarrow \mathbf{P}_W^1$ be its base change. Then π_0 induces an isomorphism $E \simeq \mathbf{P}^1$, and we have

$$(5.3.2) \quad \tilde{L}_1 = \pi_0^{-1}(0), \quad \tilde{L}_2 = \pi_0^{-1}(\infty).$$

⁷The following argument is adopted from [2, Lem. 2.13], but the present case is easier.

Set $s = S/T = x/y$ and write

$$\mathbf{P}^1 \setminus \{\infty\} = \mathbf{A}_s^1 := \operatorname{Spec} k[s], \quad \mathbf{P}^1 \setminus \{0\} = \operatorname{Spec} k[\frac{1}{s}].$$

Set $U := \mathbf{A}_s^1 \times W$ and $V := (\mathbf{P}^1 \setminus \{0\}) \times W$ and

$$\mathcal{U} := (\mathbf{A}_s^1, 0) \otimes \mathcal{W}, \quad \mathcal{V} := (\mathbf{P}^1 \setminus \{0\}, \infty) \otimes \mathcal{W}.$$

We have

$$\pi^{-1}(U) = \mathbf{A}_y^1 \times U, \quad \pi^{-1}(V) = \mathbf{A}_x^1 \times V,$$

and the restriction of π to these open subsets is given by projection. Furthermore, $E \times W \subset Y$ is defined by $y = 0$ on $\pi^{-1}(U)$ and by $x = 0$ on $\pi^{-1}(V)$. In view of (5.3.2), we have

$$(5.3.3) \quad \mathcal{F}|_{\pi^{-1}(U)} = F_{(\mathbf{A}_y^1, 0) \otimes \mathcal{U}}, \quad \mathcal{F}|_{\pi^{-1}(V)} = F_{(\mathbf{A}_x^1, 0) \otimes \mathcal{V}}.$$

Thus [2, Lem. 2.11] yields

$$R^j \pi_* \mathcal{F} = 0 \quad \text{for } j \geq 1,$$

and it remains to show

$$(5.3.4) \quad H^i(\mathbf{P}_W^1, \pi_* \mathcal{F}) = 0 \quad \text{for } i \geq 1.$$

where $\mathbf{P}_W^1 = \mathbf{P}^1 \times W$. For this consider the map

$$a_0 : Y \rightarrow \mathbf{A}_x^1 \times \mathbf{P}^1$$

which is the closed immersion $Y \hookrightarrow \mathbf{A}^2 \times \mathbf{P}^1$ followed by the projection $\mathbf{A}^2 \rightarrow \mathbf{A}_x^1$. Let $a : Y \times W \rightarrow \mathbf{A}_x^1 \times \mathbf{P}^1 \times W$ be its base change. In view of (5.3.3), the map a induces a morphism in $\underline{\mathbf{M}}\mathbf{Cor}$:

$$\alpha : (Y, \tilde{L}_1 + \tilde{L}_2 + E) \otimes \mathcal{W} \rightarrow (\mathbf{A}_x^1, 0) \otimes (\mathbf{P}^1, \infty) \otimes \mathcal{W},$$

which is an isomorphism over $(\mathbf{A}_x^1, 0) \otimes (\mathbf{P}^1 \setminus \{0\}, \infty) \otimes \mathcal{W}$. Setting

$$F_1 := \underline{\mathbf{H}}\mathbf{om}(\mathbb{Z}_{\operatorname{tr}}(\mathbf{A}_x^1, 0), F) \in \mathbf{CI}_{\operatorname{Nis}}^{\tau, sp},$$

it induces a map of Nisnevich sheaves on \mathbf{P}_W^1 :

$$\pi_*(\alpha^*) : F_{1, (\mathbf{P}^1, \infty) \otimes \mathcal{W}} \rightarrow \pi_* \mathcal{F},$$

which becomes an isomorphism over $(\mathbf{P}^1 - \{0\}) \times W$. Hence (5.3.4) follows from

$$H^i(\mathbf{P}_W^1, F_{1, (\mathbf{P}^1, \infty) \otimes \mathcal{W}}) = 0 \quad \text{for } i \geq 1,$$

which follows from [16, Th. 0.6]. \square

Lemma 5.4. *Let $N > 2$ be an integer and assume that Proposition 5.2 holds for $n < N$. Let $(X, D) \in \underline{\mathbf{M}}\mathbf{Cor}_{ls}$ and $Z \subset X$ be a smooth integral closed subscheme with $2 \leq \operatorname{codim}(Z, Y) =: c < N$. Assume*

$$D = D_1 + \cdots + D_r + D' \quad \text{with } r \leq c,$$

where D_1, \dots, D_r are distinct and reduced irreducible components of D containing Z and D' is an effective divisor on X such that none of the component of D' contains Z and Z is transversal to $|D'|$. Let $\rho : Y \rightarrow X$ be the blow-up of X in Z and $\tilde{D}_i, \tilde{D}' \subset Y$ be the strict transforms of D_i and D' respectively and $E_Z = \rho^{-1}(Z)$. Then, for all $\mathcal{W} = (W, W^\infty) \in \underline{\mathbf{MCor}}_{1s}$,

$$R^i \rho_{W*} F_{(Y, \tilde{D}_1 + \dots + \tilde{D}_r + E_Z + \tilde{D}') \otimes \mathcal{W}} = 0 \text{ for } i \geq 1,$$

where $\rho_W : Y \times W \rightarrow X \times W$ denotes the base change of ρ .

Proof. ⁸ The question is Nisnevich local around the points in $Z \times W$. Let $z \in Z \times W$ be a point and set $A := \mathcal{O}_{X \times W, z}^h$. For $V \subset Y \times W$ we denote by $V_{(z)} := V \times_{X \times W} \text{Spec } A$. By assumption we find a system of local parameters t_1, \dots, t_m of A , such that

$$\begin{aligned} (D_i \times W)_{(z)} &= V(t_i) \text{ for } 1 \leq i \leq r, \quad (Z \times W)_{(z)} = V(t_1, \dots, t_c), \\ (D' \times W)_{(z)} &= V(t_{c+1}^{e_{c+1}} \cdots t_{m_0}^{e_{m_0}}) \text{ with } c+1 \leq m_0 \leq m, \\ (X \times W^\infty)_{(z)} &= V(t_{m_0+1}^{e_{m_0+1}} \cdots t_{m_1}^{e_{m_1}}) \text{ with } m_0 \leq m_1 \leq m. \end{aligned}$$

Letting K be the residue field of A , we can choose a ring homomorphism $K \hookrightarrow A$ which is a section of $A \rightarrow K$. Then we obtain an isomorphism

$$K\{t_1, \dots, t_m\} \xrightarrow{\sim} A.$$

Let $\rho_1 : \widetilde{\mathbf{A}}^c \rightarrow \mathbf{A}^c$ be the blow-up in 0. By the above

$$\rho_W : (Y, \tilde{D}_1 + \dots + \tilde{D}_r + E_Z + \tilde{D}') \otimes \mathcal{W} \rightarrow (X, D) \otimes \mathcal{W}$$

is Nisnevich locally around z isomorphic over k to the morphism

$$(\widetilde{\mathbf{A}}^c, \tilde{L}_1 + \dots + \tilde{L}_r + E) \otimes \mathcal{W}' \rightarrow (\mathbf{A}^c, L_1 + \dots + L_r) \otimes \mathcal{W}',$$

$$(\mathcal{W}' = (\mathbf{A}_K^{m-c}, (\prod_{i=c+1}^{m_1} t_i^{e_i})))$$

induced by a map $(\widetilde{\mathbf{A}}^c, \tilde{L}_1 + \dots + \tilde{L}_r + E) \rightarrow (\mathbf{A}^c, L_1 + \dots + L_r)$ as in Proposition 5.2. Hence the statement follows from the proposition for $n = c < N$. \square

Proof of Proposition 5.2. The proof is by induction on $n \geq 2$. The case $n = 2$ follows from Lemma 5.3. Assume $n > 2$ and the theorem is proven for \mathbf{A}^m with $m < n$. In case $r = 1$, Proposition 5.2 is proved in [2, Th. 2.12]. Assume $r \geq 2$. Let $Z := L_1 \cap L_2 \subset \mathbf{A}^n$ and $\tilde{Z} \subset Y$ be the strict transform of Z . Denote by $\rho' : Y' \rightarrow Y$ the blow-up of Y in \tilde{Z} and $\tilde{L}'_i, E' \subset Y'$ be the strict transforms of \tilde{L}, E respectively and

⁸The proof is adopted from [2, Lem. 2.14].

$E'' = (\rho')^{-1}(\tilde{Z})$. Note that $\tilde{Z} = \tilde{L}_1 \cap \tilde{L}_2$ intersecting transversally with $\tilde{L}_3 + \cdots + \tilde{L}_r + E$ and $\text{codim}(\tilde{Z}, Y) = 2$. Hence, by Lemma 5.4

$$R^i \rho'_{W*} F_{(Y', \tilde{L}'_1 + \cdots + \tilde{L}'_r + E' + E'') \otimes \mathcal{W}} = 0 \text{ for } i \geq 1.$$

Since Theorem 5.1 has been proved for $i = 0$, we have

$$\rho'_* F_{(Y', \tilde{L}'_1 + \cdots + \tilde{L}'_r + E' + E'') \otimes \mathcal{W}} = F_{(Y, \tilde{L}_1 + \cdots + \tilde{L}_r + E) \otimes \mathcal{W}}.$$

Hence we obtain

$$(5.4.1) \quad R^i \rho_{W*} F_{(Y, \tilde{L}_1 + \cdots + \tilde{L}_r + E) \otimes \mathcal{W}} = R^i (\rho \rho')_{W*} F_{(Y', \tilde{L}'_1 + \cdots + \tilde{L}'_r + \tilde{E} + E') \otimes \mathcal{W}}.$$

Denote by $\sigma : \hat{Y} \rightarrow \mathbf{A}^n$ the blow-up in Z and $\hat{L}_i \subset \hat{Y}$ be the strict transform of L_i and $\Xi = \sigma^{-1}(Z)$. By Lemma 5.4 we get

$$(5.4.2) \quad R^i \sigma_{W*} F_{(\hat{Y}, \hat{L}_1 + \cdots + \hat{L}_r + \Xi) \otimes \mathcal{W}} = 0 \text{ for } i \geq 1.$$

Denote by $\sigma' : \hat{Y}' \rightarrow \hat{Y}$ the blow-up in $\hat{Z} = \sigma^{-1}(0) \subset \Xi$ and $\hat{L}'_i, \Xi' \subset \hat{Y}'$ be the strict transforms of \hat{L}_i, Ξ respectively and $\Xi'' = \sigma'^{-1}(\hat{Z})$. Note that $\tilde{Z} \subset \hat{L}_3 \cap \cdots \cap \hat{L}_n \cap \Xi$ and $\text{codim}(\tilde{Z}, \hat{Y}) = n - 1$ and \tilde{Z} intersects transversally with $\hat{L}_1 + \hat{L}_2$. Thus by Lemma 5.4 and the case $i = 0$ of Theorem 5.1, we obtain

$$(5.4.3) \quad R \sigma'_{W*} F_{(\hat{Y}', \hat{L}'_1 + \cdots + \hat{L}'_r + \Xi' + \Xi'') \otimes \mathcal{W}} = F_{(\hat{Y}, \hat{L}_1 + \cdots + \hat{L}_r + \Xi) \otimes \mathcal{W}}.$$

Finally, by [2, Lem. 2.15], there is an isomorphism of $\mathbf{A}^n \times W$ -schemes

$$(5.4.4) \quad (\hat{Y}', \hat{L}'_1, \dots, \hat{L}'_r, \Xi', \Xi'') \cong (Y', \tilde{L}'_1, \dots, \tilde{L}'_r, E', E'').$$

Altogether we obtain for $i \geq 1$

$$\begin{aligned} R^i \rho_{W*} F_{(Y, \tilde{L}_1 + \cdots + \tilde{L}_r + E) \otimes \mathcal{W}} &= R^i (\rho \rho')_{W*} F_{(Y', \tilde{L}'_1 + \cdots + \tilde{L}'_r + E' + E'') \otimes \mathcal{W}}, & \text{by (5.4.1),} \\ &= R^i (\sigma \sigma')_{W*} F_{(\hat{Y}', \hat{L}'_1 + \cdots + \hat{L}'_r + \Xi' + \Xi'') \otimes \mathcal{W}}, & \text{by (5.4.4),} \\ &= R^i \sigma_{W*} F_{(\hat{Y}, \hat{L}_1 + \cdots + \hat{L}_r + \Xi) \otimes \mathcal{W}}, & \text{by (5.4.3),} \\ &= 0, & \text{by (5.4.2).} \end{aligned}$$

This completes the proof of the proposition. \square

Remark 5.5. For simplicity, we write

$$H_{\log}^i(-, F) = H_{\log}^i(-, \underline{\omega}^{\mathbf{CI}} F) \text{ for } F \in \mathbf{RSC}_{\text{Nis}}.$$

By [13, Cor. 6.8], if $\text{ch}(k) = 0$ and $F = \Omega^i$, we have

$$H_{\log}^i(-, \Omega^i) = H^i(X, \Omega^i(\log |D|)) \text{ for } (X, D) \in \mathbf{MCor}_{ls}.$$

Hence $H_{\log}^i(-, F)$ for $F \in \mathbf{RSC}_{\text{Nis}}$ is a generalization of cohomology of sheaves of logarithmic differentials.

6. RELATION WITH LOGARITHMIC SHEAVES WITH TRANSFERS

In this section we use the same notations as [1].

Let \mathbf{lSm} be the category of log smooth and separated fs log schemes of finite type over the base field k and $\mathbf{SmlSm} \subset \mathbf{lSm}$ be the full subcategory consisting of objects whose underlying schemes are smooth over k . Let \mathbf{lCor} be the category with the same objects as \mathbf{lSm} and whose morphisms are log correspondences defined in [1, Def. 2.1.1]. Let $\mathbf{lCor}_{\mathbf{SmlSm}} \subset \mathbf{lCor}$ be the full subcategory consisting of all objects in \mathbf{SmlSm} .

Let \mathbf{PSh}^{ltr} be the category of additive presheaves of abelian groups on \mathbf{lCor} and $\mathbf{Shv}_{dNis}^{ltr} \subset \mathbf{PSh}^{ltr}$ be the full subcategory consisting of those \mathcal{F} whose restrictions to \mathbf{lSm} are dividing Nisnevich sheaves (see [1, Def. 3.1.4]). It is shown in [1, §4 and Pr. 4.6.6] that $\mathbf{Shv}_{dNis}^{ltr}$ is a Grothendieck abelian category and there is an equivalence of categories

$$(6.0.1) \quad \mathbf{Shv}_{dNis}^{ltr} \simeq \mathbf{Shv}_{dNis}^{ltr}(\mathbf{SmlSm}),$$

where the right hand side denotes the full subcategory of the category $\mathbf{PSh}^{ltr}(\mathbf{SmlSm})$ of additive presheaves of abelian groups on $\mathbf{lCor}_{\mathbf{SmlSm}}$ consisting of those \mathcal{F} whose restrictions to \mathbf{SmlSm} are dividing Nisnevich sheaves.

Now we construct a functor

$$(6.0.2) \quad \mathcal{L}og : \underline{\mathbf{MNST}}_{\log} \rightarrow \mathbf{Shv}_{dNis}^{ltr}.$$

For $\mathfrak{X} = (X, \mathcal{M}) \in \mathbf{SmlSm}$, we put $\mathfrak{X}^{MP} = (X, \partial\mathfrak{X})$, where $\partial\mathfrak{X} \subset X$ is the closed subscheme consisting of the points where the log-structure \mathcal{M} is not trivial. By [12, Theorem III.1.11.12], $\partial\mathfrak{X}$ with reduced structure is a normal crossing divisor on X so that we can view \mathfrak{X}^{MP} as an object of $\underline{\mathbf{MCor}}_{ls}$. For $F \in \underline{\mathbf{MPST}}_{\log}$ and $\mathfrak{X} \in \mathbf{SmlSm}$, we put

$$(6.0.3) \quad F^{\log}(\mathfrak{X}) = F(\mathfrak{X}^{MP}).$$

Take $\mathfrak{Y} \in \mathbf{SmlSm}$ and $\alpha \in \mathbf{lCor}(\mathfrak{Y}, \mathfrak{X})$. By [1, Def. 2.1.1 and Rem. 2.1.1(iv)], we have

$$\alpha \in \underline{\mathbf{MCor}}^{\text{fin}}((Y, n \cdot \partial\mathfrak{Y}), (X, \partial\mathfrak{X})) \text{ for some } n > 0,$$

where $n \cdot \partial\mathfrak{Y} \hookrightarrow Y$ is the n -th thickening of $\partial\mathfrak{Y} \hookrightarrow Y$. By the assumption $F \in \underline{\mathbf{MPST}}_{\log}$, the induced map

$$F^{\log}(\mathfrak{X}) = F(\mathfrak{X}^{MP}) \xrightarrow{\alpha^*} F(Y, n \cdot \partial\mathfrak{Y})$$

factors through $F^{\log}(\mathfrak{Y}) = F(Y, \partial\mathfrak{Y}) \subset F(Y, n \cdot \partial\mathfrak{Y})$ and we get a map

$$\alpha^{*\log} : F^{\log}(\mathfrak{X}) \rightarrow F^{\log}(\mathfrak{Y}).$$

Moreover, for a map $\gamma : F \rightarrow G$ in $\underline{\mathbf{MPST}}_{\log}$, the diagram

$$\begin{array}{ccc} F^{\log}(\mathfrak{X}) & \xrightarrow{\gamma} & G^{\log}(\mathfrak{X}) \\ \downarrow \alpha^{*\log} & & \downarrow \alpha^{*\log} \\ F^{\log}(\mathfrak{Y}) & \xrightarrow{\gamma} & G^{\log}(\mathfrak{Y}) \end{array}$$

is obviously commutative. Hence the assignment $\mathcal{X} \rightarrow F^{\log}(\mathcal{X})$ gives an object F^{\log} of $\mathbf{PSh}^{ltr}(\mathbf{SmlSm})$ and we get a functor

$$(6.0.4) \quad \mathcal{L}og : \underline{\mathbf{MPST}}_{\log} \rightarrow \mathbf{PSh}^{ltr}(\mathbf{SmlSm}) ; F \rightarrow F^{\log}.$$

By the definitions of sheaves ([4, Def. 1] and [1, Def. 3.1.4]) and [4, Pr. 1.9.2], this induces a functor

$$\underline{\mathbf{MNST}}_{\log} \rightarrow \mathbf{Shv}_{dNis}^{ltr}(\mathbf{SmlSm})$$

which induces the desired functor (6.0.2) using (6.0.1). By the construction, for $F \in \underline{\mathbf{MNST}}_{\log}$ and $\mathfrak{X} \in \mathbf{SmlSm}$ with $\mathcal{X} = \mathfrak{X}^{MP} \in \underline{\mathbf{MCor}}_{ls}$, we have

$$(6.0.5) \quad H_{\text{Nis}}^i(X, F_{\mathcal{X}}) = H_{s\text{Nis}}^i(\mathfrak{X}, F^{\log}) (F^{\log} = \mathcal{L}og(F)),$$

where the right hand side is the cohomology for the strict Nisnevich topology (see [1, Def. 4.3.1]).

Theorem 6.1. *For $F \in \mathbf{CI}_{\text{Nis}}^{\tau, sp}$, $F^{\log} = \mathcal{L}og(F) \in \mathbf{Shv}_{dNis}^{ltr}$ is strictly $\overline{\square}$ -invariant in the sense [1, Def. 5.2.2]. For $\mathfrak{X} \in \mathbf{SmlSm}$ with $\mathcal{X} = \mathfrak{X}^{MP} \in \underline{\mathbf{MCor}}_{ls}$, we have a natural isomorphism*

$$(6.1.1) \quad H_{\text{Nis}}^i(X, F_{\mathcal{X}}) \simeq \text{Hom}_{\mathbf{logDM}^{\text{eff}}}(M(\mathfrak{X}), F^{\log}[i]),$$

where $\mathbf{logDM}^{\text{eff}}$ is the triangulated category of logarithmic motives defined in [1, Def. 5.2.3].

Proof. We have isomorphisms

$$H_{\text{Nis}}^i(X, F_{\mathcal{X}}) \stackrel{(6.0.5)}{\simeq} H_{s\text{Nis}}^i(\mathfrak{X}, F^{\log}) \stackrel{(*1)}{\simeq} \varinjlim_{\mathfrak{Y} \in \mathfrak{X}_{div}^{Sm}} H_{s\text{Nis}}^i(\mathfrak{X}, F^{\log}) \stackrel{(*2)}{\simeq} H_{d\text{Nis}}^i(\mathfrak{X}, F^{\log}),$$

where (*2) comes from [1, Th. 5.1.8] and (*1) is a consequence of Theorem 5.1 in view of (6.0.5). Hence the strict $\overline{\square}$ -invariance of F^{\log} follows from [16, Th. 0.6]. Finally (6.1.1) follows from [1, Pr. 5.2.8]. \square

Now we consider the composite functor

$$\mathcal{L}og' : \mathbf{RSC}_{\text{Nis}} \xrightarrow{\omega^{\text{CI}}} \mathbf{CI}_{\text{Nis}}^{\tau, sp} \xrightarrow{\mathcal{L}og} \mathbf{Shv}_{dNis}^{ltr}.$$

Lemma 6.2. *$\mathcal{L}og$ and $\mathcal{L}og'$ have the same essential image.*

Proof. This follows directly from the construction and Corollary 2.6(3). \square

In what follows, we let

$$(6.2.1) \quad \mathcal{L}og : \mathbf{RSC}_{\text{Nis}} \rightarrow \mathbf{Shv}_{d\text{Nis}}^{\text{ltr}} : F \rightarrow F^{\log}$$

denote $\mathcal{L}og'$ defined as above. By (6.0.3), we have

$$(6.2.2) \quad F^{\log}(X, \text{triv}) = F(X) \text{ for } F \in \mathbf{RSC}_{\text{Nis}}, X \in \mathbf{Sm},$$

where (X, triv) denotes the log-scheme with the trivial log structure.

Theorem 6.3. *$\mathcal{L}og$ is exact and fully faithful.*

Proof. First we prove the full faithfulness. The faithfulness follows from (6.2.2). Let $F, G \in \mathbf{RSC}_{\text{Nis}}$ and $\gamma : F^{\log} \rightarrow G^{\log}$ be a map in $\mathbf{Shv}_{d\text{Nis}}^{\text{ltr}}$. By (6.2.2) it induces maps $\gamma_X : F(X) \rightarrow G(X)$ for all $X \in \mathbf{Sm}$. They are compatible with the action of \mathbf{Cor} since by [1, Rem 2.1.3(3)],

$$\mathbf{Cor}(Y, X) = \mathbf{lCor}(Y, \text{triv}), (X, \text{triv}) \text{ for } X, Y \in \mathbf{Sm}.$$

Thus γ_X for $X \in \mathbf{Sm}$ give a map $\gamma_{\mathbf{RSC}_{\text{Nis}}} : F \rightarrow G$ in $\mathbf{RSC}_{\text{Nis}}$. To see $\mathcal{L}og(\gamma_{\mathbf{RSC}_{\text{Nis}}}) = \gamma$, it suffices by (6.0.1) to show that $\mathcal{L}og(\gamma_{\mathbf{RSC}_{\text{Nis}}})$ and γ induce the same map $F^{\log}(\mathfrak{X}) \rightarrow G^{\log}(\mathfrak{X})$ for $\mathfrak{X} \in \mathbf{SmlSm}$. If \mathfrak{X} has the trivial log-structure, this follows immediately from the construction of $\gamma_{\mathbf{RSC}}$. The general case follows from this in view of the commutative diagram

$$\begin{array}{ccc} F^{\log}(\mathfrak{X}) & \xrightarrow{\gamma} & G^{\log}(\mathfrak{X}) \\ \downarrow j^* & & \downarrow j^* \\ F^{\log}(X \setminus \partial\mathfrak{X}, \text{triv}) & \xrightarrow{\gamma} & G^{\log}(X \setminus \partial\mathfrak{X}, \text{triv}) \end{array}$$

where j^* are induced by the natural map $(X \setminus \partial\mathfrak{X}, \text{triv}) \rightarrow \mathfrak{X}$ of log-schemes and injective by the construction and the semipurity of $\underline{\omega}^{\text{CI}}F$. This completes the proof of the full faithfulness.

Next we show the exactness of $\mathcal{L}og$. It suffices to show the following.

Claim 6.3.1. Given an exact sequence $0 \rightarrow F \rightarrow G \rightarrow H \rightarrow 0$ in $\mathbf{RSC}_{\text{Nis}}$, the induced sequence

$$0 \rightarrow F^{\log}(\mathfrak{X}) \rightarrow G^{\log}(\mathfrak{X}) \rightarrow H^{\log}(\mathfrak{X}) \rightarrow 0$$

is exact for every $\mathfrak{X} \in \mathbf{SmlSm}$ with X henselian local.

Indeed, by the definition of $\mathcal{L}og$, this is reduced to the exactness of

$$0 \rightarrow \underline{\omega}^{\text{CI}}F(\mathfrak{X}^{MP}) \rightarrow \underline{\omega}^{\text{CI}}G(\mathfrak{X}^{MP}) \rightarrow \underline{\omega}^{\text{CI}}H(\mathfrak{X}^{MP}) \rightarrow 0,$$

which follows from Corollary 2.6(2). This completes the proof of Theorem 6.3. \square

REFERENCES

- [1] F. Binda, D. Park, P.A. Østvær, *Triangulated categories of logarithmic motives over a field*, <https://arxiv.org/abs/2004.12298>.
- [2] F. Binda, K. Rülling, S. Saito, *On the cohomology of reciprocity sheaves*, <https://arxiv.org/abs/2010.03301>.
- [3] K. Kato, S. Saito, *Global class field theory of arithmetic schemes*, Contemporary Math. **55**, Part I (1986), 255–331.
- [4] B. Kahn, H. Miyazaki, S. Saito, T. Yamazaki, *Motives with modulus, I: Modulus sheaves with transfers*, to appear in *Epjournal de Geometrie Algebrique* (2021).
- [5] B. Kahn, H. Miyazaki, S. Saito, T. Yamazaki, *Motives with modulus, II: Modulus sheaves with transfers for proper modulus pairs*, to appear in *Epjournal de Geometrie Algebrique* (2021).
- [6] B. Kahn, S. Saito, T. Yamazaki, *Reciprocity Sheaves, I*, *Compositio Mathematica*, **9**, volume 151 (2016), 1851–1898.
- [7] B. Kahn, S. Saito, T. Yamazaki, *Reciprocity Sheaves, II*, to appear in *Homology, Homotopy and Applications* (2021).
- [8] S. Kelly, S. Saito, *Smooth blowups square for motives with modulus*, to appear in *Bulletin of the Polish Academy of Sciences - Mathematics*. (2020).
- [9] A. Merici, S. Saito, *Cancellation theorems for reciprocity sheaves*, <https://arxiv.org/abs/2001.07902>.
- [10] C. Mazza, V. Voevodsky, C. Weibel, *Lecture Notes on Motivic Cohomology*, Clay Mathematics Monographs, 2006
- [11] K. Rülling, R. Sugiyama, T. Yamazaki, *Tensor structures in the theory of modulus presheaves with transfers*, preprint, 2019, <https://arxiv.org/abs/1911.05291>.
- [12] A. Ogus, *Lectures on Logarithmic Geometry*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, (2018).
- [13] K. Rülling, S. Saito, *Reciprocity sheaves and their ramification filtrations*, to appear in *J. Inst. Math. Jussieu*. (2021).
- [14] Rülling, Kay and Saito, Shuji, *Higher local symbols and Zariski-Nagata's purity for reciprocity sheaves*, in preparation.
- [15] K. Rülling, T. Yamazaki, *Suslin homology of relative curves with modulus*, *J. London Math. Soc.* **93**, issue 2 (2016), 567–589.
- [16] S. Saito, *Purity of reciprocity sheaves*, in *Adv. Math.* **365** (2020), 107067.
- [17] V. Voevodsky, *Triangulated categories of motives over a field*, in E. Friedlander, A. Suslin, V. Voevodsky *Cycles, transfers and motivic cohomology theories*, *Ann. Math. Studies* **143**, Princeton University Press, 2000, 188–238.
- [18] A. Grothendieck, *Cohomologie locale des faisceaux cohérent et théorèmes de Lefschetz locaux et globaux (SGA2)*, *Lecture Notes in Math.*, *Advanced Studies in Pure Math.* **2**, North-Holland Publishing Co., Amsterdam, Paris, 1968.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO,
 3-8-1 KOMABA, TOKYO 153-8941, JAPAN
Email address: sshuji@msb.biglobe.ne.jp