

Never Go Full Batch (in Stochastic Convex Optimization)

Idan Amir* Yair Carmon† Tomer Koren‡ Roi Livni§

November 16, 2021

Abstract

We study the generalization performance of *full-batch* optimization algorithms for stochastic convex optimization: these are first-order methods that only access the exact gradient of the empirical risk (rather than gradients with respect to individual data points), that include a wide range of algorithms such as gradient descent, mirror descent, and their regularized and/or accelerated variants. We provide a new separation result showing that, while algorithms such as stochastic gradient descent can generalize and optimize the population risk to within ε after $O(1/\varepsilon^2)$ iterations, full-batch methods either need at least $\Omega(1/\varepsilon^4)$ iterations or exhibit a dimension-dependent sample complexity.

1 Introduction

Stochastic Convex Optimization (SCO) is a fundamental problem that received considerable attention from the machine learning community in recent years [26, 14, 3, 10, 1]. In this problem, we assume a learner that is provided with a finite sample of convex functions drawn i.i.d. from an unknown distribution. The learner’s goal is to minimize the expected function. Owing to its simplicity, it serves as an almost ideal theoretical model for studying generalization properties of optimization algorithms ubiquitous in practice, particularly *first-order methods* which utilize only first derivatives of the loss rather than higher-order ones.

One prominent approach for SCO—and learning more broadly—is to consider the *empirical risk* (the average objective over the sample) and apply a first-order optimization algorithm to minimize it. The problem of learning is then decoupled into controlling the optimization error over the empirical risk (training error) and bounding the difference between the empirical error and the expected error (generalization error).

In convex optimization, the convergence of different first-order methods has been researched extensively for many years (e.g., [24, 23, 4]), and we currently have a very good understanding of this setting in terms of upper as well lower bounds on worst-case complexity. However, in SCO where the generalization error must also be taken into account, our understanding is still lacking. In fact, this is one of the few theoretical learning models where the optimization method affects not only the optimization error but also the generalization error (distinctively from models such as PAC learning and generalized linear models). In particular, it has been shown [26, 14] that some minima

*Department of Electrical Engineering, Tel Aviv University; idanamir@mail.tau.ac.il.

†Blavatnik School of Computer Science, Tel Aviv University; ycarmon@tauex.tau.ac.il .

‡Blavatnik School of Computer Science, Tel Aviv University and Google Research; tkoren@tauex.tau.ac.il.

§Department of Electrical Engineering, Tel Aviv University; rlivni@tauex.tau.ac.il.

of the empirical risk may obtain large generalization error, while other minima have a vanishingly small generalization error. To put differently, learning in SCO is not only a question of minimizing the empirical risk, but also a question of *how* one minimizes it. However, the results of [26, 14] leave open the question of whether concrete optimization also have different generalization properties.

Towards better understanding, Amir et al. [1] recently studied the generalization properties of full-batch gradient descent (GD), where each step is taken with respect to the gradient of the empirical risk. For GD (and a regularized variant thereof), they gave a lower bound on the generalization error as a function of iteration number, which is strictly larger than the well-known optimal rate obtained by stochastic gradient descent (SGD), where each step is taken with respect to the gradient at a sampled example. Notably, the lower bound of [1] precisely matches the dimension-independent stability-based upper bound recently shown for full-batch GD by Bassily et al. [3]. The separation between full-batch GD and SGD is the first evidence that not only abstract Empirical Risk Minimizers may fail to generalize in SCO, but in fact also basic methods such as GD could be prone to such overfitting. A natural question is, then, whether overfitting is inherent to *full-batch* algorithms, that minimize the objective only through access to the exact empirical risk, or whether this suboptimality can be remedied by adding regularization, noise, smoothing, or any other mechanism for improving the generalization of GD.

In this work we present and analyze a model of *full-batch* optimization algorithms for SCO. Namely, we focus on algorithms that access the empirical risk only via a first-order oracle that computes the *exact* (full-batch) gradient of the empirical loss, rather than directly accessing gradients with respect to individual samples. Our main result provides a negative answer to the question above by significantly generalizing and extending the result of Amir et al. [1]: we show that *any* optimization method that uses full-batch gradients needs at least $\Omega(1/\varepsilon^4)$ iterations to minimize the expected loss to within ε error. This is in contrast with the empirical loss, which can be minimized with only $O(1/\varepsilon^2)$ steps.

Comparing SGD and GD in terms of the sample size n , we see that SGD converges to an optimal generalization error of $O(1/\sqrt{n})$ after $O(n)$ iterations, whereas a full-batch method must perform $\Omega(n^2)$ iterations to achieve the same $O(1/\sqrt{n})$ test error. We emphasize that we account here for the *oracle complexity*, which coincides with the iteration complexity in the case of gradient methods. In terms of individual gradients calculations, while SGD uses at most $O(n)$ gradient calculations (one sample per iteration), a full-batch method will perform $\Omega(n^3)$ calculations (n samples per iteration).

The above result is applicable to a wide family of full-batch learning algorithms: regularized GD (with any *data-independent* regularization function), noisy GD, GD with line-search or adaptive step sizes, GD with momentum, proximal methods, coordinate methods, and many more. Taken together with upper bound of Bassily et al. [3], we obtain a sharp rate of $\Theta(1/\varepsilon^4)$ for the generalization-complexity of full-batch methods. Surprisingly, this rate is achieved by standard GD (with an unusual step-size choice of $\eta = \Theta(\varepsilon^3)$), and it cannot be improved by adding regularization of any sort, nor by adding noise or any other form of implicit/explicit bias.

1.1 Related work

This work extends and generalizes the results of Amir et al. [1] who proved generalization lower bounds for GD (and a specific instance of regularized GD). Our work shows that in fact *any* full-batch method will suffer from similar lower bounds. Our construction builds upon the one used in [1], which in turn builds upon previous constructions [3, 26]. However, our arguments and proofs here are more challenging, as we need to reason about a general family of algorithms, and not about a specific algorithm whose trajectory can be analyzed directly. Our developments also build on

ideas from the literature on oracle complexity lower bounds in optimization [23, 24, 28, 7, 11, 8]. In particular, we first prove our result in the simplified setting of algorithms constrained to the span of observed gradients [23, 24] and subsequently lift it to general algorithms using a random high-dimensional embedding technique proposed by Woodworth and Srebro [28] and later refined in [7, 11]. However, while these works lower bound what we call the empirical risk, we lower bound the generalization error. This requires us to develop a somewhat different argument for how the span of the gradients evolve during the optimization: in prior work, the algorithm learns the component of the solution coordinate by coordinate, whereas in our work the true (generalizing) solution is present in the observed gradients from the first query, but spurious sampling artifacts drown it out.

Empirical studies (outside of the scope of SCO) support the claim that generalization capabilities degrade with the increase of the batch size. Specifically, Zhu et al. [31] indicates that SGD outperforms GD in terms of generalization. The works of Keskar et al. [20] and Hoffer et al. [18] exhibit a similar phenomenon in which small-batch SGD generalizes better than large-batch SGD with the same iteration budget. We provide the first *theoretical* evidence for this phenomenon for *convex* losses. Several theoretical studies explore the convergence of stochastic methods that use mini-batches [9, 21, 29]. Note that this setting differs from ours, as they assume access to mini-batches sampled *without replacement* whereas full-batch means we reuse the same (full) batch with each gradient step. The work of Wu et al. [30] also explores the separation between GD and SGD and interprets mini-batch SGD as a noisy version of GD. They propose a modified algorithm with noise injected to the full-batch gradients. Interestingly, the noise production requires access to the sample-points. Our work shows that (in SCO) this is unavoidable: namely, no data-independent noise can be used to improve generalization.

Several other works study the generalization performance of GD [27, 15, 19, 22]. The work of Soudry et al. [27], for example, examines GD on unregularized logistic regression problems. They show that, in the limit, GD converges to a well-generalizing solution by arguing about the bias of the algorithm. Interestingly, both our and their results require slow-training, beyond what is required for empirical error optimization. Another work that highlights the slow convergence of GD is that of Bassily et al. [3]. They were the first to address uniform stability of (non-smooth) GD and SGD, and provided tight bounds. Stability entails generalization, hence our results lead to stability lower bounds for any full-batch method. Consequently, we extend the lower bounds for GD in the work of Bassily et al. [3] to a wider class. It might be thought that the instability argument of Bassily et al. [3] can be used to obtain similar generalization lower bounds—however, we note that their techniques also prove instability of SGD (which does generalize). Hence, instability does not immediately imply, in this setting, lack of generalization.

Finally, we note that under smoothness and strong convexity, it is well known that improved rates can be obtained. Specifically, using the stability bound of Bousquet and Elisseeff [5] one can show that we can achieve generalization error of $O(1/\sqrt{n})$ after $O(n)$ iterations if the population risk is $O(1)$ -strongly convex. The arguments of Hardt et al. [17] imply generalization bound to instances where every sample risk is $O(\sqrt{n})$ smooth. Our result implies that, even though these special families of functions enjoy appealing learning rates, in general it is impossible to obtain better rates by strong-convexifying or smoothing problem instances via first-order full-batch oracle queries.

2 Problem Setup and Main Results

We study the standard setting of stochastic convex optimization. In this setting, a learning problem is specified by a fixed domain $\mathcal{W} \subseteq \mathbb{R}^d$ in d -dimensional Euclidean space, and a loss function $f : \mathcal{W} \times \mathcal{Z} \rightarrow \mathbb{R}$, which is both convex and L -Lipschitz with respect to its first argument (that is, for any $z \in \mathcal{Z}$ the function $f(w; z)$ is L -Lipschitz and convex with respect to w). In particular, throughout the paper, our construction consists of 1-Lipschitz functions and we will focus on a fixed domain \mathcal{W} defined to be the unit Euclidean ball in \mathbb{R}^d , namely $\mathcal{W} = \{w : \|w\|_2 \leq 1\}$.

We also assume that there exists an unknown distribution D over parameters z and the goal of the learner is to optimize the *true risk* (or *true loss*, or *population risk*) defined as follows:

$$F(w) := \mathbb{E}_{z \sim D} [f(w; z)], \quad (1)$$

We assume that a sample $S = \{z_1, \dots, z_n\}$ is drawn from the distribution D , and the learner has to output $w_S \in \mathcal{W}$ (the exact access the learner has to the sample, and how w_S may depend on S is discussed below). We require the solution to be ε -optimal in expectation for some parameter $\varepsilon > 0$, i.e.,

$$\mathbb{E}_{S \sim D^n} [F(w_S)] - \min_{w^* \in \mathcal{W}} F(w^*) \leq \varepsilon.$$

As discussed, the standard setting assumes that the learner has direct access to the i.i.d. sample, as well as to the gradients of the loss function (i.e., a first-order oracle). In this work, though, we focus on a specific family *full-batch* methods. Hence, the optimization process is described as follows: First, an i.i.d. sample $S = (z_1, \dots, z_n)$ is drawn from D . Then, the learner is provided with access only to the *empirical risk* via a *full-batch first-order oracle* which we define next.

Full-batch first-order oracle. Consider a *fixed* sample $S = (z_1, \dots, z_n)$ of size n , drawn i.i.d. from D . The *empirical risk* over the sample S is

$$F_S(w) = \frac{1}{n} \sum_{i=1}^n f(w; z_i).$$

Then, a *full-batch first-order oracle* is a procedure that, given input $w \in \mathcal{W}$, outputs

$$\mathcal{O}(w) := (\nabla F_S(w); F_S(w)).$$

where $\nabla F_S(w)$ is an empirical risk sub-gradient of the form

$$\nabla F_S(w) = \frac{1}{n} \sum_{i=1}^n \nabla f(w; z_i), \quad (2)$$

and each sub-gradient $\nabla f(w, z_i)$ is computed by the oracle as a function of w and z_i (that is, independently of z_j for $j \neq i$).

We emphasize that the sample is fixed throughout the optimization, so that the oracle computes the gradient of the *same* empirical risk function at every call, hence the name *full-batch*. Note that the subgradient with respect to a single data point, i.e., $\nabla f(w; z_i)$, is not accessible through this oracle, which only returns the average gradient over the sample S .

Notice that our definition above is slightly narrower than a general sub-gradient oracle for the empirical risk due to the requirement that the sub-gradients $\nabla f(w, z_i)$ are chosen independently of z_j for $j \neq i$ – since we provide here with a lower bound, this restriction strengthens our result. We make this restriction to avoid some degenerate constructions (that in fact can even be used to fail SGD if the gradient at z_i may depend on the whole sample), which are of no practical implications.

Full-batch first-order algorithm. A *full-batch* (first-order) method is naturally defined as any algorithm that has access to the optimization objective—namely the empirical risk F_S —only via the full-batch first order oracle. In particular, if w_t is the t 'th query of the algorithm to the full-batch oracle then w_t has to be of the form

$$w_t = Q_t(\mathcal{O}(w_0), \dots, \mathcal{O}(w_{t-1})), \quad (3)$$

where $Q_t : (\mathbb{R}^{d+1})^t \rightarrow \mathcal{W}$ is a fixed (possibly randomized) mapping. At the end of the process the algorithm outputs w_S . We study the algorithm's oracle complexity, which is the number of iterations T the algorithm performs before halting. Therefore, we assume without loss of generality that $w_S = w_T$, i.e., the algorithm's output is its T 'th query.

2.1 Main result

In this section we establish our main result, which provides a generalization lower-bound for full-batch first order algorithms. The complete proof is provided in Section 5.

Theorem 1. *Let $\varepsilon > 0$ and $n, T \in \mathbb{N}$; there exists $d = \text{poly}(2^n, T, 1/\varepsilon)$ such that the following holds. For any full-batch first-order algorithm with oracle complexity at most T , there exists a 1-Lipschitz convex function $f(w; z)$ in \mathcal{W} , the unit-ball in \mathbb{R}^d , and a distribution D over \mathcal{Z} such that, for some universal constant $c > 0$:*

$$\mathbb{E}_{S \sim D^n} [F(w_S)] \geq \min_{w^* \in \mathcal{W}} F(w^*) + \varepsilon + \Omega(\min\{1 - c\varepsilon^2\sqrt{T}, 0\}). \quad (4)$$

An immediate consequence of Theorem 1 is that in order to obtain less than ε true risk we need at least $T = \Omega(1/\varepsilon^4)$ iterations.

For simplicity, we state and prove the lower bound in Theorem 1 for the class of first-order full-batch algorithms defined above. However, our constructions readily generalize to *local full-batch oracles* that provide a complete description of F_S in an arbitrarily small neighborhood of the query point [23, 16]. Such oracles subsume second-order oracles, and consequently our generalization lower bounds hold also for second-order full-batch algorithms.

2.2 Discussion

Theorem 1 suggests that full-batch first-order algorithms are inferior to other types of first-order algorithms that operate with access to individual examples, such as SGD. Importantly, this separation is achieved not in terms of the *optimization* performance but in terms of the *generalization* performance. In light of this result, we next discuss and revisit the role of the optimization algorithm in the context of SCO. In particular, we wish to discuss the implications to what are perhaps the two most prominent full-batch optimization methods, GD and regularized-GD, and in turn compare them.

Gradient descent. Perhaps the simplest example of a full-batch method is (projected) GD: GD is an iterative algorithm that at each iteration performs an update step

$$w_t = \Pi_{\mathcal{W}}[w_{t-1} - \eta \nabla F_S(w_t)],$$

where \mathcal{W} is a convex set on which we project the iterated step. The output of GD is normally taken to be $w_S = \frac{1}{T} \sum w_t$ (or a randomly chosen w_t). Notice, that each step requires one call to a full batch oracle, and a single projection operation. The convergence analysis of GD to the

optimal solution of the *empirical risk* has been widely studied. Specifically, if n is the sample-size, it is known that with $\eta = O(1/\sqrt{n})$ and $T = O(n)$, GD converges to a minimizer of F_S that is $O(1/\sqrt{n})$ -sub optimal. For the *exact* variant of GD depicted above, the generalization performance was analyzed in the work of Amir et al. [1] that showed that with $T = O(n)$ steps, GD will suffer $\Omega(1/\sqrt[4]{n})$ generalization error. Theorem 1 extends the above result to any variant of GD (dynamic learning-rate, noisy GD, normalized GD, etc.).

Regularized gradient descent. We would also like to discuss the implication of Theorem 1 with respect to *regularized* variants of GD that operate on the regularized empirical risk

$$\hat{F}(w) = \lambda r(w) + F_S(w).$$

The main motivation of introducing the regularization term r is to avoid overfitting, and a popular choice for r is the Euclidean norm $r(w) = \|w\|_2^2$. This choice leads to the following update rule for GD:

$$w_{t+1} = \Pi_{\mathcal{W}} [(1 - \eta_t) \cdot (2\lambda w_t) - \eta_t \nabla F_S(w_t)],$$

Again, this update can be implemented using a single first-order full-batch oracle call that computes the quantity $\nabla F_S(w_t)$. More generally, for any data-independent r , GD on \hat{F} is a full-batch algorithm¹. When r is the Euclidean norm, the minimizer of \hat{F} is known to enjoy (with choice $\lambda = O(1/\sqrt{n})$), an optimal generalization error of $O(1/\sqrt{n})$ [5, 26]. This demonstrates the power of regularization and how it can provably induce generalization. Nevertheless, Theorem 1 still applies to any optimization method over \hat{F} . Since optimization of \hat{F} (the regularized empirical risk) to $O(1/\sqrt{n})$ -precision can be done via a full-batch method, and with less than $O(n)$ calls, we observe that there are methods that minimize the regularized-empirical risk but, due to Theorem 1 do not reach the optimal generalization error.

The role of regularization. Finally, in light of Theorem 1 let us compare the different variants of GD and regularized GD that *do* generalize well, in order to sharpen our understanding of the role of regularization in generalization. The conclusion of Theorem 1 is that any full-batch method that generalizes well performs at least $O(n^2)$ steps. For regularized GD, with ℓ_2 regularization, $O(n^2)$ are indeed sufficient. In particular, with $O(n^2)$ iterations we can find a solution that has $O(1/n)$ empirical error. Any such solution would enjoy a generalization error of $O(1/\sqrt{n})$ [26]. For GD, Bassily et al. [3] showed that $O(n^2)$ iterations would also suffice to achieve $O(1/\sqrt{n})$ error. This is achieved by tuning the learning rate to $\eta = O(1/n^{3/2})$. Notice that this improvement does not require any type of added regularization.

To summarize, both GD and regularized GD with optimal parameters require $\Theta(n^2)$ iterations to attain the optimal $O(1/\sqrt{n})$ generalization error. Overall then, explicitly adding regularization is not necessary nor does it improve the convergence rate. One might be tempted to believe that tuning the learning rate in GD induces *implicitly* some sort of regularization. For example, one might imagine that GD can be biased towards minimal norm solution, which might explain redundancy of regularizing by this norm. However, this turns out also to be false: Dauber et al. [10] showed how GD (with any reasonable choice of learning rate) can diverge from the minimal norm solution. In fact, for any regularization term r , one can find examples where GD does not converge to the regularized solution. Thus, even though GD and regularized-GD are comparable algorithms in terms of generalization and oracle complexity, they are distinct in terms of the solutions they select.

¹Note that we are not concerned with the computational cost of computing $\nabla r(w_t)$ since it does not factor into oracle complexity.

3 Technical Overview

In this section we give an overview of our construction and approach towards proving Theorem 1. For the sake of exposition, we will describe here a slightly simpler construction which proves the main result only for algorithms that remain in the span of the gradients. In more detail, let us examine the family of iterative algorithms of the form

$$w_t \in \text{span}\{\nabla F_S(w_0), \nabla F_S(w_1), \dots, \nabla F_S(w_{t-1})\} \cap \mathcal{W}, \quad (5)$$

where \mathcal{W} is the unit ball and $\nabla F_S(w_t)$ is full-batch oracle response to query w_t as defined in (2) above. Well-studied algorithms such as GD and GD with standard ℓ_2 norm regularization fall into this category of algorithms.

To extend the lower bound to algorithms not restricted the gradient span we refine the simpler construction and apply well-established techniques of random embedding in high-dimensional space. We discuss these modifications briefly in the end of this section and provide the full details in Sections 4 and 5 below.

3.1 A simpler construction

Let us fix $n, d \geq 1$ and parameters $z = (\alpha, \varepsilon, \gamma) \in \{0, 1\}^d \times \mathbb{R} \times \mathbb{R}^2 = \mathcal{Z}$, such that $\alpha \in \{0, 1\}^d$, $\varepsilon > 0$ and $\gamma_1, \gamma_2 > 0$. Define the hard instance $f_{(6)} : \mathbb{R}^{d+2} \times \mathcal{Z} \rightarrow \mathbb{R}$ as follows:

$$f_{(6)}(w; (\alpha, \varepsilon, \gamma)) = g_\gamma(w; \alpha) + \gamma_1 v_\alpha \cdot w + \varepsilon w \cdot e_{d+2} + r(w), \quad (6)$$

where g_γ, v_α and r are

- $g_\gamma(w; \alpha) := \sqrt{\sum_{i \in [d]} \alpha(i) h_\gamma^2(w(i))}$ with $h_\gamma(a) := \begin{cases} 0 & a \geq -\gamma_2; \\ a + \gamma_2 & a < -\gamma_2, \end{cases}$
- $r(w) := \max\{0, \max_{i \in [d+1]} \{w(i)\}\}$,
- $v_\alpha(i) := \begin{cases} -\frac{1}{2n} & \text{if } \alpha(i) = 0; \\ +1 & \text{if } \alpha(i) = 1; \\ 0 & \text{if } i \in \{d+1, d+2\}, \end{cases}$

and e_{d+2} is the $(d+2)$ 'th standard basis vector. The distribution we will consider is uniform over α . That is, we draw $\alpha \in \{0, 1\}^d$ uniformly at random and pick the function $f_{(6)}(w; (\alpha, \varepsilon, \gamma))$.

The parameters γ_1 and γ_2 of the construction should be thought of as arbitrarily small. In particular, the term $\gamma_1 v_\alpha \cdot w$ in Eq. (6) should be thought of as negligible, and the first term, g_γ , is roughly

$$g_\gamma(w; \alpha) \approx \sqrt{\sum_{i \in [d]} \alpha(i) (\max\{-w(i), 0\})^2}.$$

Another useful property of the construction is the population risk $F(w) = \mathbb{E}_{z \sim D} f_{(6)}(w; z)$ is minimized at $w^* \approx -e_{d+2}$, with expected loss $F(w^*) \approx -\varepsilon$. However, as we will see, the choice of the perturbation vector v_α and the term $r(w)$ hinder the learner from observing this coordinate and; the first $\Omega(\varepsilon^{-4})$ queries are constrained to a linear subspace where all the points have a high generalization error due to the expectation of the first term g_γ .

3.2 Analysis

We next state the main lemmas we use, with proofs deferred to Appendix A. Given a sample S , let us denote $\bar{v} = \frac{1}{n} \sum_{\alpha \in S} v_\alpha$, and

$$\text{span}^1\{u_1, u_2, \dots\} := \text{span}\{u_1, u_2, \dots\} \cap \mathcal{W}.$$

Additionally, given a fixed sample we write

$$\mathcal{I}(S) = \{i : \alpha(i) = 0 \forall \alpha \in S\} \cup \{d+1\}$$

for the set of coordinates $i \in [d]$ such that $\alpha(i) = 0$ for every α in the sample S , plus the coordinate $d+1$.

Lemma 2. *Let $\gamma_1 \leq \frac{1}{2T}$, $\gamma_2 = \frac{2\gamma_1}{\varepsilon}$, and suppose that the sample S satisfies $|\mathcal{I}(S)| > T$. Then there exists a first-order full-batch oracle such that for any algorithm that adheres to*

$$w_t \in \text{span}^1\{\nabla F_S(w_0), \nabla F_S(w_1), \dots, \nabla F_S(w_{t-1})\}, \quad (7)$$

with respect to $f(w; (\alpha, \varepsilon, \gamma))$ defined in Eq. (6), we have

$$w_t \in \text{span}^1_{i \in \mathcal{I}_t(S)} \left\{ \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i \right\} \text{ for all } t \in [T],$$

where $\mathcal{I}_t(S)$ is the set of the $t+1$ largest coordinates in $\mathcal{I}(S)$.

We next observe that in any span of the form $\{\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i\}_{i \in \mathcal{I}_T(S)}$ such that $|\mathcal{I}_T(S)| \leq T$, we cannot find a solution with better risk than 0. On the other hand, note that for $\bar{w} = -e_{d+2}$, we have that

$$f_{(6)}(\bar{w}; (\alpha, \varepsilon, \gamma)) = -\varepsilon.$$

In other words, our lower bound stems from the following result:

Lemma 3. *For sufficiently small $\gamma_1 \leq 2n\varepsilon\gamma_2$, $\gamma_2 \leq \varepsilon/\sqrt{4T}$, and any vector $\|\bar{v}\| \leq \sqrt{d}$, any output*

$$w_S \in \text{span}^1_{i \in \mathcal{I}_T(S)} \left\{ \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i \right\},$$

satisfies

$$\frac{1}{2} \sqrt{\sum_{i \in [d]} h_\gamma^2(w_S(i)) + \varepsilon w_S(e_{d+2})} \geq \min\left\{1 - 2\varepsilon^2\sqrt{T}, 0\right\} - \frac{1}{2}\varepsilon. \quad (8)$$

Lower bound proof sketch for span-restricted algorithms of the form (5). First, observe that the probability of an arbitrary index i to satisfy $\alpha(i) = 0$ for all $\alpha \in S$ is $(1/2)^n$. Therefore, $|\mathcal{I}(S)| - 1$, the number of indexes that hold this from the possible d , is distributed as a binomial with d experiments and success probability $p = 2^{-n}$. Using elementary probability arguments one can show that for sufficiently large d we have $|\mathcal{I}(S)| > T$ with high probability; see Claim 1 in the appendix. This implies that the conditions of Lemmas 2 and 3 hold w.h.p. To conclude, we relate the LHS of Eq. (8) to the expected risk

$$F(w) = \mathbb{E}_{\alpha \sim D} [f_{(6)}(w; (\alpha, \varepsilon, \gamma))] = \mathbb{E}_{\alpha \sim D} [g_\gamma(w; \alpha)] + \gamma_1 \cdot \mathbb{E}_{\alpha \sim D} [v_\alpha] \cdot w + \varepsilon w \cdot e_{d+2} + r(w).$$

As $g_\gamma(w; \alpha)$ is convex w.r.t. α (since $\alpha(i) = \alpha^2(i)$) we can apply Jensen's inequality with $\mathbb{E}_{\alpha \sim D}[\alpha(i)] = \frac{1}{2}$ to obtain:

$$\mathbb{E}_{\alpha \sim D} [g_\gamma(w_S; \alpha)] \geq \frac{1}{2} \sqrt{\sum_{i \in [d]} h_\gamma^2(w_S(i))}.$$

Applying the Cauchy-Schwarz inequality to the second term while also using the facts that $\|v_\alpha\| \leq \sqrt{d}$ and that w_S is in the unit ball, we get:

$$\gamma_1 \mathbb{E}_{\alpha \sim D} [v_\alpha] \cdot w \geq -\gamma_1 \mathbb{E}_{\alpha \sim D} [\|v_\alpha\| \cdot \|w\|] \geq -\gamma_1 \sqrt{d}.$$

For sufficiently small γ_1 this term is negligible, and since $r(w) \geq 0$ we get that the expected risk is approximately the LHS term in Eq. (8). Lastly, recalling that $F(-e_{d+2}) = -\varepsilon$ we get that

$$F(w_S) - \min_{w \in \mathcal{W}} F(w) \geq \frac{1}{2}\varepsilon + \min\{1 - 2\varepsilon^2\sqrt{T}, 0\} \quad \text{w.h.p.}$$

The same lower bound (up to a constant) also holds in expectation by the law of total expectation. Our distribution is supported on 5-Lipschitz convex functions, so that re-parametrizing $\frac{1}{10}\varepsilon \rightarrow \varepsilon$ as well as $f_{(6)}$ yields the claimed lower bound (4) for the case of span-restricted algorithms. \blacksquare

3.3 Handling general full-batch algorithms

The above construction establishes an $\Omega(1/\varepsilon^4)$ oracle complexity lower bound on any algorithm whose iterates lie in the span of the previous gradients. While this covers a large class of algorithms, techniques like preconditioning [12], coordinate methods [25] and randomized smoothing [13] do not satisfy this assumption. In fact, a trivial algorithm that always outputs $-e_{d+2}$ will solve the hard instance (6) in a single iteration.

To address general algorithms, we employ a well-established technique in optimization lower bounds [28, 7, 11] wherein we embed a hard instance $f(w; z)$ for span-constrained algorithms in a random high-dimensional space. More concretely, we draw a random orthogonal matrix $U \in \mathbb{R}^{d' \times d}$ ($U^\top U = I_{d \times d}$) and consider the $d' > d$ -dimensional instance $f_U(w; z) = f(U^\top w; z)$ along with its corresponding empirical objective $F_{S,U}(w) = \frac{1}{n} \sum_{i \in [n]} f_U(w; z_i)$. Roughly speaking, we show that for a general algorithm operating with the appropriate subgradient oracle for $F_{S,U}$ the iterate w_t is approximately in the span of $\{\nabla F_{S,U}(w_0), \dots, \nabla F_{S,U}(w_{t-1})\}$ in the sense that the component of w_t outside that span is nearly orthogonal to the columns of U . Consequently, the response of the oracle to the query w_t at iteration t is, with high probability, identical to the information it would return if queried with the projection of w_t to the span of the previously observed gradients. This reduces, in a sense, the problem back to the span-restricted setting described above.

For the embedding technique to work, we must robustify the hard instance construction so that small perturbations around points in the span of previous gradients do not “leak” additional information about the embedding U . To do that we make a fairly standard modification to the component $r(w)$ in (6) (known as Nemirovski's function [11, 6]), replacing it with $\max\{0, \max_{i \in [d]} \{w(i) + i\gamma'\}, w(d+1) + \gamma''\}$, where γ', γ'' are small offset coefficients that go to zero as the embedding dimension d' tends to infinity. We provide the full construction and the proof of Theorem 1 in Sections 4 and 5.

4 The Full Construction

As explained above, the key difference between the simplified construction $f_{(6)}$ and the full construction with which we prove Theorem 1 is that we modify the Nemirovski function term $r(w)$ in order to make it robust to queries that are nearly within a certain linear subspace. In particular, we bias the different terms in the maximization defining $r(w)$ so as to control the index of the coordinate attaining the maximum. For ease of reference, we now provide a self-contained definition of our full construction with the modified Nemirovski function.

Fix $n, d \geq 1$ and parameters $z = (\alpha, \varepsilon, \gamma) \in \{0, 1\}^d \times \mathbb{R} \times \mathbb{R}^3 = \mathcal{Z}$ are such that $\alpha \in \{0, 1\}^d$, $\varepsilon > 0$ and $\gamma_1, \gamma_2, \gamma_3 > 0$. Define the hard instance $f_{(9)} : \mathbb{R}^{d+2} \times \mathcal{Z} \rightarrow \mathbb{R}$ as follows:

$$f_{(9)}(w; (\alpha, \varepsilon, \gamma)) = g_\gamma(w; \alpha) + \gamma_1 v_\alpha \cdot w + \varepsilon w \cdot e_{d+2} + r(w), \quad (9)$$

where g_γ, v_α and r are

- $g_\gamma(w; \alpha) := \sqrt{\sum_{i \in [d]} \alpha(i) h_\gamma^2(w(i))}$ with $h_\gamma(a) := \begin{cases} 0 & a \geq -\gamma_2; \\ a + \gamma_2 & a < -\gamma_2, \end{cases}$
- $r(w) := \max\{0, \max_{i \in [d+1]} \{w(i) + \sigma_i\}\}$ with $\sigma_i := \begin{cases} i \cdot \frac{\gamma_1 \gamma_3}{4dn} & \text{if } i \in [d]; \\ 2\gamma_3 & \text{if } i = d + 1. \end{cases}$
- $v_\alpha(i) := \begin{cases} -\frac{1}{2n} & \text{if } \alpha(i) = 0; \\ +1 & \text{if } \alpha(i) = 1; \\ 0 & \text{if } i \in \{d + 1, d + 2\}, \end{cases}$

and e_i is the i 'th standard basis vector in \mathbb{R}^{d+2} . We consider a distribution D over α that is distributed uniformly over $\{0, 1\}^d$; that is, we draw $\alpha \in \{0, 1\}^d$ uniformly at random and pick the function $f_{(9)}(w; (\alpha, \varepsilon, \gamma))$. The rest of the parameters are set throughout the proof as follows:

$$\gamma_1 = \frac{\varepsilon \gamma_2}{4}, \quad \gamma_2 = \frac{\varepsilon}{T \sqrt{d}}, \quad \gamma_3 = \frac{\varepsilon}{16}. \quad (10)$$

With this choice of distribution D as well our choice of parameters we obtain, since $\|v_\alpha\| \leq \sqrt{d}$ and by our choice of γ_1 (as well as Jensen's inequality and $r(\cdot) \geq 0$):

$$F(w) = \mathbb{E}_{\alpha \sim D} [f_{(9)}(w; (\alpha, \varepsilon, \gamma))] \geq \frac{1}{2} \sqrt{\sum_{i \in [d]} h_\gamma^2(w(i))} + \varepsilon w(d+2) - \frac{\varepsilon}{4}. \quad (11)$$

Notice that we also have that for a choice $w^* = -e_{d+2}$, since $r(w^*) = 2\gamma_3$:

$$F(w^*) = -\varepsilon + \frac{\varepsilon}{8} = -\frac{7\varepsilon}{8} \quad (12)$$

Our development makes frequent use of the following notation from Section 3:

$$\mathcal{I}(S) = \{i : \alpha(i) = 0 \forall \alpha \in S\} \cup \{d + 1\}, \quad \mathcal{I}_t(S) = t \text{ largest elements in } \mathcal{I}(S), \quad \text{and } \bar{v} = \frac{1}{n} \sum_{\alpha \in S} v_\alpha.$$

We begin with the following lemma, which is a robust version of Lemma 2 in Section 3. The proof is provided in Appendix B.1.

Lemma 4. *Suppose that $w_0 = 0$. Consider $f_{\mathfrak{g}}(w; (\alpha, \varepsilon, \gamma))$ with parameters as in Eq. (10). Suppose S is a sample such that $|\mathcal{I}(S)| > t + 1$. Assume that w is such that*

$$w = w_t + q,$$

where

$$w_t \in \text{span}_{i \in \mathcal{I}_t(S)}^1 \left\{ \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i \right\}, \text{ and } \|q\|_\infty \leq \min \left\{ \frac{\gamma_2}{3}, \frac{\gamma_1 \gamma_3}{16dn} \right\}. \quad (13)$$

Then,

$$\nabla F_S(w) = \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i,$$

for some $i \in \mathcal{I}_{t+1}(S)$, where $\mathcal{I}_t(S)$ is the set of the $t + 1$ largest coordinates in $\mathcal{I}(S)$.

The following corollary states that the gradient oracle's answers are resilient to small perturbation of the query (as long as they are in vicinity of the "right" subspace): the proof is provided in Appendix B.2:

Corollary 5. *Assume that w is such that*

$$w = w_t + q,$$

where

$$w_t \in \text{span}_{i \in \mathcal{I}_t(S)}^1 \left\{ \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i \right\}, \text{ and } \|q\|_\infty \leq \frac{1}{4\sqrt{d}} \min \left\{ \frac{\gamma_2}{3}, \frac{\gamma_1 \gamma_3}{16dn} \right\}. \quad (14)$$

Then,

$$\nabla F_S(w) = \nabla F_S(\Pi_{t+1}(w)), \quad F_S(w) = F_S(\Pi_{t+1}(w)),$$

where Π_t is a projection onto $\text{span}_{i \in \mathcal{I}_t(S)} \{ \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i \}$.

5 Proof of Theorem 1

To prove Theorem 1 we embed the construction of Section 4 into a random, higher-dimensional space. More formally, let $f_{\mathfrak{g}}(w; z)$ be as in Eq. (9), and, for $d_2 \geq d$, let $U \in \mathbb{R}^{d_2 \times d}$ be an orthogonal matrix, i.e., such that $U^\top U = I_{d \times d}$. We consider then the objective function over \mathbb{R}^{d_2} :

$$f_U(w; z) = f_{\mathfrak{g}}(U^\top w; (\alpha, \varepsilon, \gamma)).$$

Given a sample S we use the notation

$$F_{S,U}(w) = \frac{1}{n} \sum_{i=1}^n f_U(w; z_i), \quad (15)$$

for the empirical error and

$$F_U(w) = \mathbb{E}_{\alpha \sim D} [f_U(w; (\alpha, \varepsilon, \gamma))], \quad (16)$$

for the expected error, where, as before, D is such that the coordinates of α are i.i.d. Bernoulli(1/2) and the parameters $\gamma_1, \gamma_2, \gamma_3$ are fixed as in Eq. (10). We start with the following claim:

Lemma 6. Fix a deterministic full-batch first-order algorithm, and a sample S such that $|\mathcal{I}(S)| > 2T$. Let $U \in \mathbb{R}^{d_2 \times d}$ be a random orthogonal matrix, then for some

$$d = O((T+1)2^n) \quad \text{and} \quad d_2 = \tilde{O}\left(\frac{d^3 T n^2}{\varepsilon^6}\right),$$

we have that with probability at least 0.99 (over the draw of U):

$$F_U(w_S) - F_U(w^*) \geq \min\left\{1 - 2\varepsilon^2\sqrt{2T}, 0\right\} + \frac{1}{8}\varepsilon, \quad (17)$$

where F_U is as in Eq. (16).

Before we proceed with the proof of Lemma 6, we explain how Theorem 1 follows. Fix a full-batch algorithm A and let D be a distribution as in Lemma 6. Let $\mathcal{X}_{U,S,\xi}$ be an indicator for the event that Eq. (17) holds, where U is the random orthogonal matrix, S is the sample and ξ is the random seed of the algorithm A , which is independent of U and S . Then by Lemma 6 we have that

$$\mathbb{E}_{S,\xi} \left[\mathbb{E}_U[\mathcal{X}_{U,S,\xi} \mid |\mathcal{I}(S)| > 2T] \right] > 0.99. \quad (18)$$

The next claim follows a standard concentration inequality and shows that the event $|\mathcal{I}(S)| > 2T$ is indeed probable; the proof is provided in Section 5.2.

Claim 1. Suppose $d \geq \max\{16, 4T\}2^n$. Then with probability at least 3/4, it holds that $|\mathcal{I}(S)| > 2T$.

From Claim 1 and Eq. (18), we can conclude that:

$$\mathbb{E}_{S,\xi} \mathbb{E}_U[\mathcal{X}_{U,S,\xi}] > 2/3.$$

By changing order of expectation we conclude that there exists a matrix U such that w.p. at least 2/3 (over the sample S as well as the random bits of the algorithm A) the lower bound (17) holds. Theorem 1 now follows from

$$\mathbb{E}[F_U(w_S)] - F_U(w^*) \geq \mathbb{E}\left[\mathcal{X}_{U,S,\xi} \left(\min\left\{1 - 2\varepsilon^2\sqrt{2T}, 0\right\} + \frac{1}{8}\varepsilon \right)\right] \geq \frac{2}{3} \min\left\{1 - 2\varepsilon^2\sqrt{2T}, 0\right\} + \frac{1}{12}\varepsilon.$$

We are left with proving Claim 1 and Lemma 6, which we do in Sections 5.1 and 5.2.

5.1 Proof of Lemma 6

We start by defining inductively a chain of algorithms A_0, \dots, A_T act as intermediaries between A and a full-batch first order oracle for F_S . A_t should be thought of as an arbitrator between A and the oracle, where at each iteration i , it receives a query $w_i^{(t)}$ from A , submits some query to the oracle, and returns some answer $\mathcal{O}_{A_t}(w_i^{(t)})$ to A (not necessarily the oracle's answer). We will build the chain A_0, \dots, A_T in such a way that $A_0 \equiv A$, while A_T forces queries to stay in the span of the gradients. We will then relate the error of A_0 to the error of A_T by bounding the probabilities that they observe different information from the oracle.

We formally define A_t is as follows.

- For each $i \leq t$, algorithm A_t receives the query point $w_i^{(t)}$, then the algorithm defines $v_i^{(t)} = \Pi_{2i}(U^\top w_i^{(t)})$, where $\Pi_i \in \mathbb{R}^{(d+2) \times (d+2)}$ is the Euclidean projection onto

$$\text{span}^1_{j \in \mathcal{I}_i(S)} \{\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_j\}.$$

The algorithm then inputs the query $v_i^{(t)}$ to a full-batch first-order oracle for F_S , receives $(\nabla F_S(v_i^{(t)}), F_S(v_i^{(t)}))$ and provides A with

$$\mathcal{O}_{A_t}(w_i^{(t)}) = (U \nabla F_S(v_i^{(t)}), F_S(v_i^{(t)})) = (U \nabla F_S(\Pi_{2i}(U^\top w_i^{(t)})), F_S(\Pi_{2i}(U^\top w_i^{(t)}))).$$

- For $i > t$, algorithm A_t behaves like a standard full-batch oracle. Namely, it receives a query $w_i^{(t)}$, defines $v_i^{(t)} = U^\top w_i^{(t)}$, queries the oracle with it, receives $(\nabla F_S(v_i^{(t)}), F_S(v_i^{(t)}))$ and provides A with

$$\mathcal{O}_{A_t}(w_i^{(t)}) = (U \nabla F_S(v_i^{(t)}), F_S(v_i^{(t)})) = (U \nabla F_S(U^\top w_i^{(t)}), F_S(U^\top w_i^{(t)})).$$

Notice that A_0 is the algorithm A interacting with a valid full-batch first order oracle for $F_{S,U}$ defined in Eq. (15). In particular, at each iteration A_0 provides, as required from a full-batch first-order oracle:

$$\begin{aligned} U \nabla F_S(U^\top w_i^{(0)}) &= U \cdot \frac{1}{n} \sum_{z \in S} \nabla f_{\mathcal{G}}(U^\top w_i^{(0)}; z) = \frac{1}{n} \sum_{z \in S} U \nabla f_{\mathcal{G}}(U^\top w_i^{(0)}; z) \\ &= \frac{1}{n} \sum_{z \in S} \nabla f_U(w_i^{(0)}; z) = \nabla F_{S,U}(w_i^{(0)}). \end{aligned}$$

At the other extreme the algorithm A_T is an algorithm that only queries points in

$$\text{span}^1_{i \in \mathcal{I}_{2T}(S)} \{\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i\}.$$

We obtain then, by Lemma 3 as well as Eqs. (11) and (12) that for every $v_i^{(T)}$:

$$\begin{aligned} F_U(Uv_i^{(T)}) &= F(v_i^{(T)}) \\ &\geq \frac{1}{2} \sqrt{\sum_{j \in [d]} h_\gamma^2(v_i^{(T)}(j))} + \varepsilon v_i^{(T)} \cdot e_{d+2} - \frac{\varepsilon}{4} \end{aligned} \quad (\text{Eq. (11)})$$

$$\geq \min \left\{ 1 - 2\varepsilon^2 \sqrt{2T}, 0 \right\} - \frac{3}{4} \varepsilon \quad (\text{Lemma 3})$$

$$\geq F_U(-Ue_{d+2}) + \min \left\{ 1 - 2\varepsilon^2 \sqrt{2T}, 0 \right\} + \frac{1}{8} \varepsilon. \quad (\text{Eq. (12)})$$

Denote by P_t the probability that algorithm A_t outputs a sequence $v_i^{(t)}$ such that for some i :

$$F_U(Uv_i^{(t)}) - F(-Ue_{d+2}) \geq \min \left\{ 1 - 2\varepsilon^2 \sqrt{2T}, 0 \right\} + \frac{1}{8} \varepsilon. \quad (19)$$

In particular, we have argued so far that $P_t = 0$.

Next, for two vectors v, v' , let us write $v \equiv v'$ if

$$\nabla F_S(v) = \nabla F_S(v'), \text{ and } F_S(v) = F_S(v').$$

Now, suppose we run A_t, A_{t-1} and we observe at step t a vector $w_t^{(t-1)}$ such that $v_t^{(t-1)} \equiv v_t^{(t)}$. Notice that in that case the output of A_t and the output of A_{t-1} is completely identical. Indeed, up to step t the two algorithms are identical and at step t they provide the same response to A (and after that they again behave identically). Thus,

$$P_T \geq D_{t-1} - \Pr(v_t^{(t-1)} \neq v_t^{(t)}).$$

Rearranging terms and iteratively applying the formula above we obtain:

$$\begin{aligned} P_0 &\leq P_T + \sum_{t=1}^T \Pr(v_t^{(t-1)} \neq v_t^{(t)}) \\ &\leq \sum_{t=1}^T \Pr(U^\top w_t^{(t)} \neq \Pi_{2t} U^\top w_t^{(t)}) \\ &\leq \sum_{t=1}^T \Pr\left(\|(1 - \Pi_{2t-1})U^\top w_t^{(t)}\|_\infty \geq \frac{\varepsilon^3}{2^{12}d^2Tn}\right). \end{aligned} \quad (\text{Corollary 5 \& Eq. (10)})$$

The result now follows from the next lemma, whose proof we defer to Appendix C.

Lemma 7. *Let \mathcal{S}_k be a fixed k -dimensional subspace in \mathbb{R}^{d_1} . Let $U \in \mathbb{R}^{d_2 \times d_1}$, $d_2 \geq d_1$ be a random orthogonal matrix. Let $\Pi \in \mathbb{R}^{d_1 \times d_1}$ be the orthogonal projection on \mathcal{S}_k . Let w be a random unit vector that is deterministic conditional on $U\Pi$. Then*

$$\Pr(\|(1 - \Pi)U^\top w\|_\infty > c) \leq 2d_2 e^{-\frac{d_1 c^2}{2} \cdot (d_2 - k + 1)}.$$

To apply the lemma, we set:

$$\mathcal{S}_t = \text{span}_{i \in \mathcal{I}_t(S)} \{\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i\},$$

and we want to show that $w_i^{(t)}$ is deterministic conditional on $U\Pi_{2t-1}$. To see that, note that throughout the interaction between A and A_t , given $w_i^{(t)}$ with $i < t$, A_t calculates $v_i^{(t)} = \Pi_{2i} U^\top w_i^{(t)} = \Pi_{2i} \Pi_{2t-1} U^\top w_i^{(t)} = \Pi_{2i} (U\Pi_{2t-1})^\top w_i^{(t)}$, which is determined given $U\Pi_{2t-1}$. In turn, A_t returns to A the vector:

$$U \nabla F_S(v_i^{(t)}).$$

By Lemma 4, since $v_i^{(t)} \in \mathcal{S}_{2i}$ we have that $\nabla F_S(v_i^{(t)}) \in \mathcal{S}_{2i+1} \subseteq \mathcal{S}_{2t-1}$, for $i < t$, hence:

$$U \nabla F_S(v_i^{(t)}) = U \Pi_{2t-1} \nabla F_S(v_i^{(t)}).$$

Which is again determined by $v_i^{(t)}$ and $U\Pi_{2t-1}$. We conclude that $w_t^{(t)}$ is deterministic if we condition on $U\Pi_{2t-1}$. Overall we obtain:

$$P_0 \leq 2Td_2 e^{-\frac{d^3 T^2 n^2}{2^{25} \varepsilon^6} \cdot (d_2 - 2T)}.$$

Thus, for some

$$d_2 \leq O\left(\frac{d^3 T^2 n^2}{\varepsilon^6} \log(Td_2)\right) = \tilde{O}\left(\frac{d^3 T^2 n^2}{\varepsilon^6}\right),$$

we obtain that $A = A_0$ satisfies Eq. (17) with probability at least 0.99.

5.2 Proof of Claim 1

The probability that a given index $i \in [d]$ is such that $\alpha(i) = 0$ for all $\alpha \in S$ is 2^{-n} . Thus, the expected number of such indices is $\mu = 2^{-n}d$ and the standard deviation is $\sigma = \sqrt{2^{-n}(1 - 2^{-n})d} \leq \sqrt{\mu}$. By an application of Chebyshev’s inequality we obtain

$$\begin{aligned} \Pr(|\mathcal{I}(S)| \leq 2T) &\leq \Pr(|\mathcal{I}(S)| \leq \frac{1}{2}\mu) && (d \geq 4T \cdot 2^n) \\ &\leq \Pr(|\mathcal{I}(S)| \leq \mu - 2\sqrt{\mu}) && (\mu \geq 16 \text{ for } d \geq 16 \cdot 2^n) \\ &\leq \Pr(|\mathcal{I}(S)| \leq \mu - 2\sigma) && (\sigma \leq \sqrt{\mu}) \\ &\leq \frac{1}{4}. && (\text{Chebyshev's inequality}) \end{aligned}$$

Acknowledgements

This work has received support from the Israeli Science Foundation (ISF) grant no. 2549/19 and grant no. 2188/20, from the Len Blavatnik and the Blavatnik Family foundation, from the Yandex Initiative in Machine Learning, and from an unrestricted gift from Google. Any opinions, findings, and conclusions or recommendations expressed in this work are those of the author(s) and do not necessarily reflect the views of Google.

References

- [1] I. Amir, T. Koren, and R. Livni. SGD generalizes better than GD (and regularization doesn’t help). In *Conference on Learning Theory*, 2021. [1](#), [2](#), [6](#)
- [2] K. Ball et al. An elementary introduction to modern convex geometry. *Flavors of Geometry*, 31:1–58, 1997. [26](#)
- [3] R. Bassily, V. Feldman, C. Guzmán, and K. Talwar. Stability of stochastic gradient descent on nonsmooth convex losses. In *Advances in Neural Information Processing Systems*, 2020. [1](#), [2](#), [3](#), [6](#)
- [4] L. Bottou, F. E. Curtis, and J. Nocedal. Optimization methods for large-scale machine learning. *SIAM Review*, 60(2):223–311, 2018. [1](#)
- [5] O. Bousquet and A. Elisseeff. Stability and generalization. *Journal of Machine Learning Research*, 2:499–526, 2002. [3](#), [6](#)
- [6] S. Bubeck, Q. Jiang, Y. T. Lee, Y. Li, A. Sidford, et al. Complexity of highly parallel non-smooth convex optimization. *Advances in Neural Information Processing Systems*, 2019. [9](#)
- [7] Y. Carmon, J. C. Duchi, O. Hinder, and A. Sidford. Lower bounds for finding stationary points I. *Mathematical Programming*, pages 1–50, 2019. [3](#), [9](#)
- [8] Y. Carmon, A. Jambulapati, Q. Jiang, Y. Jin, Y. T. Lee, A. Sidford, and K. Tian. Acceleration with a ball optimization oracle. In *Advances in Neural Information Processing Systems*, 2020. [3](#)
- [9] A. Cotter, O. Shamir, N. Srebro, and K. Sridharan. Better mini-batch algorithms via accelerated gradient methods. In *Advances in Neural Information Processing Systems*, 2011. [3](#)

- [10] A. Dauber, M. Feder, T. Koren, and R. Livni. Can implicit bias explain generalization? stochastic convex optimization as a case study. In *Advances in Neural Information Processing Systems*, 2020. 1, 6
- [11] J. Diakonikolas and C. Guzmán. Lower bounds for parallel and randomized convex optimization. In *Conference on Learning Theory*, 2019. 3, 9
- [12] J. Duchi, E. Hazan, and Y. Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of Machine Learning Research*, 12(7), 2011. 9
- [13] J. C. Duchi, P. L. Bartlett, and M. J. Wainwright. Randomized smoothing for stochastic optimization. *SIAM Journal on Optimization*, 22(2):674–701, 2012. 9
- [14] V. Feldman. Generalization of ERM in stochastic convex optimization: The dimension strikes back. In *Advances in Neural Information Processing Systems*, 2016. 1, 2
- [15] S. Gunasekar, J. D. Lee, N. Srebro, and D. Soudry. Implicit bias of gradient descent on linear convolutional networks. *Advances in Neural Information Processing Systems*, 2018. 3
- [16] C. Guzmán and A. Nemirovski. On lower complexity bounds for large-scale smooth convex optimization. *Journal of Complexity*, 31(1):1–14, 2015. 5
- [17] M. Hardt, B. Recht, and Y. Singer. Train faster, generalize better: Stability of stochastic gradient descent. In *International Conference on Machine Learning*, 2016. 3
- [18] E. Hoffer, I. Hubara, and D. Soudry. Train longer, generalize better: closing the generalization gap in large batch training of neural networks. In I. Guyon, U. von Luxburg, S. Bengio, H. M. Wallach, R. Fergus, S. V. N. Vishwanathan, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, 2017. 3
- [19] Z. Ji and M. Telgarsky. The implicit bias of gradient descent on nonseparable data. In *Conference on Learning Theory*, 2019. 3
- [20] N. S. Keskar, D. Mudigere, J. Nocedal, M. Smelyanskiy, and P. T. P. Tang. On large-batch training for deep learning: Generalization gap and sharp minima. In *International Conference on Learning Representations*, 2017. 3
- [21] M. Li, T. Zhang, Y. Chen, and A. J. Smola. Efficient mini-batch training for stochastic optimization. In S. A. Macskassy, C. Perlich, J. Leskovec, W. Wang, and R. Ghani, editors, *International Conference on Knowledge Discovery and Data Mining*, 2014. 3
- [22] Y. Li, E. X. Fang, H. Xu, and T. Zhao. Implicit bias of gradient descent based adversarial training on separable data. In *International Conference on Learning Representations*, 2019. 3
- [23] A. S. Nemirovski and D. B. Yudin. *Problem complexity and method efficiency in optimization*. 1983. 1, 3, 5
- [24] Y. Nesterov. *Introductory lectures on convex optimization: A basic course*, volume 87. Springer, 2003. 1, 3
- [25] Y. Nesterov. Efficiency of coordinate descent methods on huge-scale optimization problems. *SIAM Journal on Optimization*, 22(2):341–362, 2012. 9

- [26] S. Shalev-Shwartz, O. Shamir, N. Srebro, and K. Sridharan. Stochastic convex optimization. In *Conference on Learning Theory*, 2009. 1, 2, 6
- [27] D. Soudry, E. Hoffer, M. S. Nacson, S. Gunasekar, and N. Srebro. The implicit bias of gradient descent on separable data. *The Journal of Machine Learning Research*, 19(1):2822–2878, 2018. 3
- [28] B. Woodworth and N. Srebro. Tight complexity bounds for optimizing composite objectives. In *Advances in Neural Information Processing Systems*, 2016. 3, 9
- [29] B. Woodworth, J. Wang, A. Smith, B. McMahan, and N. Srebro. Graph oracle models, lower bounds, and gaps for parallel stochastic optimization. In *Advances in Neural Information Processing Systems*, 2018. 3
- [30] J. Wu, W. Hu, H. Xiong, J. Huan, V. Braverman, and Z. Zhu. On the noisy gradient descent that generalizes as sgd. In *International Conference on Machine Learning*, 2020. 3
- [31] Z. Zhu, J. Wu, B. Yu, L. Wu, and J. Ma. The anisotropic noise in stochastic gradient descent: Its behavior of escaping from sharp minima and regularization effects. In K. Chaudhuri and R. Salakhutdinov, editors, *International Conference on Machine Learning*, 2019. 3

Appendix

A Proofs for Section 3

A.1 Proof of Lemma 2

To define the first-order full-batch oracle fulfilling the lemma, it suffices to define a sub-gradient oracle for the functions $r(w) = \max\{0, \max_{i \in [d+1]} \{w(i)\}\}$ and g_γ (all the other components of the construction are differentiable). To that end, when $\max_{i \in [d+1]} \{w(i)\} \geq 0$ we let $\nabla r(w) = e_{i_w}$ where i_w is the largest index for which $r(w) = w(i_w)$; when $\max_{i \in [d+1]} \{w(i)\} < 0$ the function is differentiable with $\nabla r(w) = 0$. For, g_γ , we simply set $\nabla g_\gamma(w, \alpha) = 0$ for any w where $g_\gamma(w, \alpha)$ is not differentiable.

We prove that $w_t \in \text{span}_{i \in \mathcal{I}_t(S)}^1 \{\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i\}$ for all $t \leq T$ by induction on t . The base case $t = 0$ is trivial since $w_0 = 0$ is the only vector in the span of the empty set. Moreover, we have

$$\begin{aligned}
 \nabla F_S(w_0) = \nabla F_S(0) &= \frac{1}{n} \sum_{\alpha \in S} \nabla g_\gamma(0; \alpha) + \gamma_1 \bar{v} + \varepsilon e_{d+2} + \nabla r(0) \\
 &= \gamma_1 \bar{v} + \varepsilon e_{d+2} + \nabla r(0) && (\nabla g_\gamma(0; \alpha) = 0) \\
 &= \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_{d+1}. && (\nabla r(0) = e_{d+1})
 \end{aligned}$$

For the induction step we assume that $w_t \in \text{span}_{i \in \mathcal{I}_t(S)}^1 \{\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i\}$. Therefore w_t takes the form:

$$w_t = \sum_{i \in \mathcal{I}_t(S)} \beta_i (\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i). \tag{20}$$

for some $\beta_i \in \mathbb{R}$. Observe that $\bar{v}(i) = 0$ for $i \in \{d+1, d+2\}$, and $\bar{v}(i) = -\frac{1}{2n}$ for any $i \in \mathcal{I}(S)$, and also that $\bar{v}(i) \in (\frac{1}{2n}, 1]$ for $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$. Denote $\bar{\beta} = \sum_{i \in \mathcal{I}(S)} \beta_i$, therefore

$$w_t(i) = \begin{cases} -\frac{\gamma_1}{2n}\bar{\beta} + \beta_i & \text{if } i \in \mathcal{I}_t(S) \setminus \{d+1\}; \\ -\frac{\gamma_1}{2n}\bar{\beta} & \text{if } i \in \mathcal{I}(S) \setminus \mathcal{I}_t(S); \\ \gamma_1 \bar{v}(i) \bar{\beta} & \text{if } i \notin \{\mathcal{I}(S) \cup \{d+2\}\}; \\ \beta_i & \text{if } i = d+1; \\ \varepsilon \bar{\beta} & \text{if } i = d+2. \end{cases} \quad (21)$$

Consider then the sub-gradient at w_t ,

$$\nabla F_S(w_t) = \frac{1}{n} \sum_{\alpha \in S} \nabla g_\gamma(w_t; \alpha) + \gamma_1 \bar{v} + \varepsilon e_{d+2} + \nabla r(w_t). \quad (22)$$

We examine the first and last terms in Eq. (22). For the first term we present the following claim:

Claim 2. *Suppose w_t is of the form (20). For sufficiently small $\gamma_1 \leq \frac{\varepsilon \gamma_2}{2}$ we have*

$$\frac{1}{n} \sum_{\alpha \in S} \nabla g_\gamma(w_t; \alpha) = 0.$$

Proof. Since $\alpha(i) = 0$ for $i \in \mathcal{I}(S)$ we get,

$$\frac{1}{n} \sum_{\alpha \in S} \nabla g_\gamma(w_t; \alpha) = \frac{1}{n} \sum_{\alpha \in S} \nabla \left(\sqrt{\sum_{i \notin \{\mathcal{I}(S) \cup \{d+2\}\}} \alpha(i) h_\gamma^2(w_t(i))} \right). \quad (23)$$

Assume that $\gamma_1 \leq \frac{\varepsilon \gamma_2}{2}$ implies $w_t(i) > -\gamma_2$ for $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$. Given the assumption, note that $h_\gamma(w_t(i)) = 0$ for any $w_t(i) \geq -\gamma_2$, then together with Eq. (23) we conclude the claim. We proceed by proving that this assumption holds.

Using Eq. (21) implies that $w_t(i) = \gamma_1 \bar{v}(i) \bar{\beta}$ for $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$. Observe that for $\bar{\beta} \geq 0$ it follows immediately that $w_t(i) \geq 0 > -\gamma_2$. Then, for $\bar{\beta} < 0$, we assume by contradiction that $w_t(i) \leq -\gamma_2$ for some $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$. Therefore, we get that $\gamma_1 \bar{v}(i) \bar{\beta} \leq -\gamma_2$, or alternatively that

$$\bar{\beta} \leq -\frac{\gamma_2}{\gamma_1 \bar{v}(i)}. \quad (24)$$

In addition, from Eq. (21) it is clear that

$$\begin{aligned} \|w_t\| &\geq \|\varepsilon \bar{\beta} e_{d+2}\| && \text{(Eq. (21))} \\ &= \varepsilon(-\bar{\beta}) && (\bar{\beta} < 0) \\ &\geq \frac{\varepsilon \gamma_2}{\gamma_1 \bar{v}(i)} && \text{(Eq. (24))} \\ &\geq \frac{\varepsilon \gamma_2}{\gamma_1} && (\bar{v}(i) \in (\frac{1}{2n}, 1] \text{ for } i \notin \{\mathcal{I}(S) \cup \{d+2\}\}) \\ &\geq 2. && (\gamma_1 \leq \frac{\varepsilon \gamma_2}{2}) \end{aligned}$$

However, we have $\|w_t\|_2 \leq 1$ since the domain \mathcal{W} is the unit Euclidean ball. ■

For the last term in Eq. (22), we use the claim below:

Claim 3. Suppose w_t is of the form (20). For $|\mathcal{I}(S)| > T$ and $\gamma_1 \leq \frac{1}{2T}$ we get that

$$\nabla r(w_t) = e_i,$$

for some $i \in \mathcal{I}_{t+1}(S)$.

Proof. First, note that $\nabla r(w_t) = e_i$ for some $i \in \mathcal{I}(S)$ implies $i \in \mathcal{I}_{t+1}(S)$, because $w_t(i) = w_t(j)$ for every $i, j \in \mathcal{I}(S) \setminus \mathcal{I}_t(S)$, and therefore our choice of $r(w)$ guarantees that either $i^* \in \mathcal{I}_t(S)$ or i^* is the largest element in $\mathcal{I}(S) \setminus \mathcal{I}_t(S)$; both cases imply $i^* \in \mathcal{I}_{t+1}(S)$. To show that $\nabla r(w_t) = e_i$ for some $i \in \mathcal{I}(S)$, we split the analysis to two cases.

Case 1: $\bar{\beta} \geq 0$. Observe that if $\beta_i = 0$ for all $i \in \mathcal{I}_t(S)$, then $w_t = 0$ which coincides with the base of the induction. Assume that not all $\beta_i = 0$. Then, we next show that for sufficiently small $\gamma_1 \leq \frac{1}{2T}$ we get that

$$i^* := \arg \max_{i \in [d+1]} \{w_t(i)\} \in \mathcal{I}(S). \quad (25)$$

To see that Eq. (25) holds, note that since $\bar{\beta} \geq 0$, then for $i' = \arg \max_{i \in \mathcal{I}_t(S)} \beta_i$ we have that $\beta_{i'} > 0$. Observe that $\gamma_1 \bar{v}(i) \bar{\beta} \leq \gamma_1 \bar{\beta}$ for $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$ due to the fact that $\bar{v}(i) \leq 1$. Suppose $\bar{\beta} > 0$, so that for any $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$, again by Eq. (21)

$$\begin{aligned} w_t(i) &= \gamma_1 \bar{v}(i) \bar{\beta} \leq \gamma_1 \bar{\beta} = 2\gamma_1 \bar{\beta} - \gamma_1 \bar{\beta} \\ &< \frac{1}{T} \bar{\beta} - \frac{\gamma_1}{2n} \bar{\beta} && (\gamma_1 \leq \frac{1}{2T}) \\ &\leq \frac{1}{|\mathcal{I}_t(S)|} \bar{\beta} - \frac{\gamma_1}{2n} \bar{\beta} && (|\mathcal{I}_t(S)| \leq t \leq T) \\ &= \frac{1}{|\mathcal{I}_t(S)|} \sum_{j \in \mathcal{I}_t(S)} \beta_j - \frac{\gamma_1}{2n} \bar{\beta} \\ &\leq \max_{j \in \mathcal{I}_t(S)} \beta_j - \frac{\gamma_1}{2n} \bar{\beta} \\ &= \beta_{i'} - \frac{\gamma_1}{2n} \bar{\beta} \\ &= w_t(i'). \end{aligned}$$

And therefore $i^* \in \mathcal{I}(S)$. Similarly, when $\bar{\beta} = 0$ it follows immediately that for any $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$ we get

$$w_t(i) = 0 < \beta_{i'} = w_t(i'),$$

which implies again that $i^* \in \mathcal{I}(S)$; our choice of $\nabla r(w)$ guarantees that $\nabla r(w_t) = e_{i^*}$.

Case 2: $\bar{\beta} < 0$. From Eq. (21), note that since $w_t(i) < 0$ for $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$, then

$$i^* := \arg \max_{i \in [d+1]} \{w_t(i)\} \in \mathcal{I}(S), \quad (26)$$

as long as $\mathcal{I}(S) \setminus \mathcal{I}_t(S) \neq \emptyset$ (Namely, $t < |\mathcal{I}(S)|$). This is justified by the assumption that $|\mathcal{I}(S)| > T$. Therefore, $\nabla r(w_t) = e_i$ for some $i \in \mathcal{I}(S)$. \blacksquare

Combining both observations in Claims 2 and 3 and plugging it into Eq. (22) we obtain,

$$\nabla F_S(w_t) = \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i,$$

for some $i \in \mathcal{I}_{t+1}(S)$. We conclude that,

$$\begin{aligned} w_{t+1} &\in \text{span}\{\nabla F_S(w_0), \nabla F_S(w_1), \dots, \nabla F_S(w_t)\} \\ &= \text{span}_{i \in \mathcal{I}_{t+1}(S)}\{\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i\}, \end{aligned}$$

as $w_t \in \text{span}\{\nabla F_S(w_0), \nabla F_S(w_1), \dots, \nabla F_S(w_{t-1})\} = \text{span}_{i \in \mathcal{I}_t(S)}\{\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i\}$. This completes the induction.

A.2 Proof of Lemma 3

By assumption, for some sequence $\{\beta_i\}_{i \in \mathcal{I}_T(S)}$ where $\beta_i \in \mathbb{R}$, the algorithm output w_S takes the form

$$w_S = \sum_{i \in \mathcal{I}_T(S)} \beta_i (\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i).$$

Observe that the solution w_S must remain in the unit ball, thus

$$\varepsilon \left| \sum_{i \in \mathcal{I}_T(S)} \beta_i \right| \leq \|w_S\|_2 \leq 1. \quad (27)$$

Denote $\mathcal{I}_T^- = \{i \in \mathcal{I}_T(S) : \beta_i \leq 0\}$ as the non-positive subset in $\mathcal{I}_T(S)$. Recalling that $\bar{v} \cdot e_{d+2} = 0$, we have

$$\begin{aligned} \frac{1}{2} \sqrt{\sum_{i \in [d]} h_\gamma^2(w_S(i))} + \varepsilon w_S \cdot e_{d+2} &= \frac{1}{2} \sqrt{\sum_{i \in \mathcal{I}_T^-} h_\gamma^2(w_S(i))} + \varepsilon^2 \bar{\beta} \\ &\geq \frac{1}{2} \sqrt{\sum_{i \in \mathcal{I}_T^-} h_\gamma^2(w_S(i))} + \varepsilon^2 \sum_{i \in \mathcal{I}_T^-} \beta_i \\ &\geq \frac{1}{2} \sqrt{\sum_{i \in \mathcal{I}_T^-} h_\gamma^2(\beta_i + \gamma_2)} + \varepsilon^2 \sum_{i \in \mathcal{I}_T^-} \beta_i. \end{aligned} \quad (28)$$

Where the last inequality holds since $h^2 \gamma(\cdot)$ is non-increasing, together with Eq. (27) and the fact that $\gamma_1 \leq 2n\varepsilon\gamma_2$

$$w_S(i) = \beta_i - \frac{\gamma_1}{2n} \sum_{j \in \mathcal{I}_T(S)} \beta_j \leq \beta_i + \frac{\gamma_1}{2n\varepsilon} \leq \beta_i + \gamma_2,$$

for $i \in \mathcal{I}_T^- \setminus \{d+1\}$. Clearly, this also holds for $i = d+1$, as $w_S(d+1) = \beta_{d+1} \leq \beta_{d+1} + \gamma_2$. Using the reverse triangle inequality, and the observations that $|\mathcal{I}_T^-| \leq T$ and that $|h_\gamma(a) - a| \leq \gamma_2$ for any $a \leq \gamma_2$, we get

$$\sqrt{\sum_{i \in \mathcal{I}_T^-} h_\gamma^2(\beta_i + \gamma_2)} \geq \sqrt{\sum_{i \in \mathcal{I}_T^-} (\beta_i + \gamma_2)^2} - \sqrt{\sum_{i \in \mathcal{I}_T^-} (h_\gamma(\beta_i + \gamma_2) - (\beta_i + \gamma_2))^2} \geq \sqrt{\sum_{i \in \mathcal{I}_T^-} \beta_i^2} - 2\gamma_2 \sqrt{T}.$$

Plugging this into Eq. (28),

$$\begin{aligned} \frac{1}{2} \sqrt{\sum_{i \in [d]} h_\gamma^2(w_S(i))} + \varepsilon w_S \cdot e_{d+2} &\geq \frac{1}{2} \sqrt{\sum_{i \in \mathcal{I}_T^-} \beta_i^2} + \varepsilon^2 \sum_{i \in \mathcal{I}_T^-} \beta_i - \gamma_2 \sqrt{T} \\ &\geq \left(\frac{1}{2} - \varepsilon^2 \sqrt{T}\right) \sqrt{\sum_{i \in \mathcal{I}_T^-} \beta_i^2} - \frac{1}{2} \varepsilon, \end{aligned} \quad (\gamma_2 \leq \varepsilon / \sqrt{4T})$$

where the second inequality follows from $\|u\|_1 \leq \sqrt{d} \|u\|_2$ and $|\mathcal{I}_T^-| \leq T$. To conclude, note that

$$\sqrt{\sum_{i \in \mathcal{I}_T(S)} \beta_i^2} - \frac{\gamma_1}{2n} \left| \sum_{i \in \mathcal{I}_T(S)} \beta_i \right| \leq \|w_S\|_2 \leq 1,$$

and we obtain the required result for $\gamma_1 \leq 2n\varepsilon$.

B Proofs for Section 5

B.1 Proof of Lemma 4

The proof mirrors that of Lemma 2 and comprises of using induction on t to show for any $t \leq T$ and w as in Eq. (13), we have $\nabla F_S(w) = \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i$ for $i \in \mathcal{I}_{t+1}(S)$. As in the proof of Lemma 2, we choose a sub-gradient oracle for $r(w)$ such that all sub-gradients are standard basis vectors, and if multiple coordinate achieve the maximum defining r , the largest one is selected.

The basis of the induction is $t = 0$, where we observe that for $\|q\|_\infty \leq \gamma_2/3$ we have that $\nabla g_\gamma(q; \alpha) = 0$. Note that $\|q\|_\infty \leq \gamma_3/3$ for $\gamma_1 < \frac{8n}{3}$, and therefore $\nabla r(q) = e_{d+1}$ due to the fact that $\sigma_i + \gamma_3/3 < \sigma_{d+1} - \gamma_3/3$ for any $i \in [d]$. Hence

$$\begin{aligned} \nabla F_S(w) &= \frac{1}{n} \sum_{\alpha \in S} \nabla g_\gamma(q; \alpha) + \gamma_1 \bar{v} + \varepsilon e_{d+2} + \nabla r(q) \\ &= \gamma_1 \bar{v} + \varepsilon e_{d+2} + \nabla r(q) && (\nabla g_\gamma(q; \alpha) = 0) \\ &= \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_{d+1} && (\nabla r(q) = e_{d+1}) \\ &= \nabla F_S(w_0). && (w_0 = 0) \end{aligned}$$

For the induction step we assume that $w_t \in \text{span}_{i \in \mathcal{I}_t(S)}^1 \{ \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i \}$. Therefore w_t takes the form

$$w_t = \sum_{i \in \mathcal{I}_t(S)} \beta_i (\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i).$$

for some $\beta_i \in \mathbb{R}$. Observe that $\bar{v}(i) = 0$ for $i \in \{d+1, d+2\}$, that $\bar{v}(i) = -\frac{1}{2n}$ for any $i \in \mathcal{I}(S)$, and that $\bar{v}(i) \in (\frac{1}{2n}, 1]$ for $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$. Denote $\bar{\beta} = \sum_{i \in \mathcal{I}_t(S)} \beta_i$, therefore

$$w_t(i) = \begin{cases} -\frac{\gamma_1}{2n} \bar{\beta} + \beta_i & \text{if } i \in \mathcal{I}_t(S) \setminus \{d+1\}; \\ -\frac{\gamma_1}{2n} \bar{\beta} & \text{if } i \in \mathcal{I}(S) \setminus \mathcal{I}_t(S); \\ \gamma_1 \bar{v}(i) \bar{\beta} & \text{if } i \notin \{\mathcal{I}(S) \cup \{d+2\}\}; \\ \beta_i & \text{if } i = d+1; \\ \varepsilon \bar{\beta} & \text{if } i = d+2. \end{cases} \quad (29)$$

Consider then the gradient at w_t ,

$$\nabla F_S(w_t) = \frac{1}{n} \sum_{\alpha \in S} \nabla g_\gamma(w_t; \alpha) + \gamma_1 \bar{v} + \varepsilon e_{d+2} + \nabla r(w_t). \quad (30)$$

We examine the first and last terms in Eq. (30). For the first term we repeat and modify the analysis in Claim 2 to handle the noise vector q .

Claim 4. *Suppose that $w = w_t + q$ where w_t satisfies Eq. (20) and $\|q\|_\infty \leq \gamma_2/3$. Then, we have*

$$\frac{1}{n} \sum_{\alpha \in S} \nabla g_\gamma(w_t; \alpha) = \frac{1}{n} \sum_{\alpha \in S} \nabla g_\gamma(w; \alpha) = 0.$$

Proof. Since $\alpha(i) = 0$ for $i \in \mathcal{I}(S)$ we get,

$$\frac{1}{n} \sum_{\alpha \in S} \nabla g_\gamma(w_t; \alpha) = \frac{1}{n} \sum_{\alpha \in S} \nabla \left(\sqrt{\sum_{i \notin \{\mathcal{I}(S) \cup \{d+2\}\}} \alpha(i) h_\gamma^2(w_t(i))} \right). \quad (31)$$

Assume that $\gamma_1 \leq \frac{\varepsilon \gamma_2}{2}$ implies $w_t(i) > -\gamma_2/2$ for $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$. Under this assumption we also have $w_t(i) + q(i) > -\gamma_2/2 - \gamma_2/3 > -\gamma_2$, and hence, since $h_\gamma(w) = 0$ for any $w \geq -\gamma_2$, then together with Eq. (31) we conclude the claim. From here the proof continues as in Claim 2. \blacksquare

For the last term in Eq. (30), we use the following claim, which is the noise-robust counterpart of Claim 3.

Claim 5. *Suppose that $w = w_t + q$ where w_t satisfies Eq. (20), $\|q\|_\infty \leq \frac{\gamma_1 \gamma_3}{8n}$ and $|\mathcal{I}(S)| > t$. Then*

$$\nabla r(w) = e_i,$$

for some $i \in \mathcal{I}_{t+1}(S)$.

Proof. For the proof, we will show that there is always $i^* \in \mathcal{I}(S)$ such that:

$$w_t(i^*) + \sigma_{i^*} > \max_{i \notin \{\mathcal{I}(S) \cup \{d+2\}\}} \{w_t(i) + \sigma_i\} + \frac{\gamma_1 \gamma_3}{4n}. \quad (32)$$

Combined with the assumption $\|q\|_\infty \leq \frac{\gamma_1 \gamma_3}{8n}$, we immediately conclude that $\nabla r(w_t + q) = e_{i'}$ for some $i' \in \mathcal{I}(S)$. We may further conclude that $i' \in \mathcal{I}_{t+1}(S)$, because $w_t(i) = w_t(j)$ for every $i, j \in \mathcal{I}(S) \setminus \mathcal{I}_t(S)$, and therefore our choice of $r(w)$ guarantees that either. We now turn to show Eq. (32) following the lines of the proof of Claim 3 but taking the noise vector q into account. We split the analysis to five cases.

Case 1a: $\bar{\beta} > 0$ and $\beta_{d+1} + \gamma_3 \geq \gamma_1 \bar{\beta}$. By Eq. (29), for any $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$ we have

$$\begin{aligned} w_t(d+1) + \sigma_{d+1} &= \beta_{d+1} + 2\gamma_3 \\ &\geq \gamma_1 \bar{v}(i) \bar{\beta} + \gamma_3 && (\bar{v}(i) \leq 1 \text{ for any } i \in [d+1]) \\ &> \gamma_1 \bar{v}(i) \bar{\beta} + \frac{\gamma_1 \gamma_3}{2n} && (\gamma_1 < 2n) \\ &\geq w_t(i) + \sigma_i + \frac{\gamma_1 \gamma_3}{4n}, && (\sigma_i \leq \frac{\gamma_1 \gamma_3}{4n}) \end{aligned}$$

so Eq. (32) holds with $i^* = d+1$.

Case 1b: $\bar{\beta} > 0$ and $\beta_{d+1} + \gamma_3 < \gamma_1 \bar{\beta}$. For $\gamma_1 \leq \frac{1}{4T} \leq 1$ we get,

$$\beta_{d+1} + \gamma_3 < \gamma_1 \bar{\beta} \leq \beta_{d+1} + \sum_{i \in \mathcal{I}_t(S) \setminus \{d+1\}} \beta_i.$$

In other words,

$$\gamma_3 < \sum_{i \in \mathcal{I}_t(S) \setminus \{d+1\}} \beta_i. \quad (33)$$

Thus, for any, $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$:

$$\begin{aligned} w_t(i) + \sigma_i + \frac{\gamma_1 \gamma_3}{4n} &= \gamma_1 \bar{v}(i) \bar{\beta} + i \cdot \frac{\gamma_1 \gamma_3}{4dn} + \frac{\gamma_3 \gamma_1}{4n} \\ &\leq \gamma_1 \bar{\beta} + \frac{\gamma_1 \gamma_3}{2n} && (\bar{v}(i) \leq 1 \text{ and } i \leq d) \\ &\leq \gamma_1 \bar{\beta} + \frac{\gamma_3}{2T} && (\gamma_1 \leq \frac{n}{T}) \\ &< 2\gamma_1 \bar{\beta} + \frac{1}{2T} \sum_{j \in \mathcal{I}_t(S) \setminus \{d+1\}} \beta_j - \gamma_1 \bar{\beta} && (\text{Eq. (33)}) \\ &< \frac{1}{2T} \bar{\beta} + \frac{1}{2T} \sum_{j \in \mathcal{I}_t(S) \setminus \{d+1\}} \beta_j - \frac{\gamma_1}{2n} \bar{\beta} && (\gamma_1 \leq \frac{1}{4T} \text{ and } \bar{\beta} > 0) \\ &\leq \frac{1}{2} \max_{j \in \mathcal{I}_t(S)} \beta_j + \frac{1}{2} \max_{j \in \mathcal{I}_t(S) \setminus \{d+1\}} \beta_j - \frac{\gamma_1}{2n} \bar{\beta} && (|\mathcal{I}_t(S)| \leq T) \\ &\leq \beta_{i'} - \frac{\gamma_1}{2n} \bar{\beta} && (i' := \arg \max_{i \in \mathcal{I}_t(S)} \beta_i) \\ &\leq w_t(i') + \sigma_{i'} && (\sigma_i \geq 0 \text{ for any } i \in [d+1]) \end{aligned}$$

Case 2: $\bar{\beta} \leq -\gamma_3$. For any $i' \in \mathcal{I}(S) \setminus \mathcal{I}_t(S)$ (non-empty by the assumption $|\mathcal{I}(S)| > t$) and for any $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$ we have

$$\begin{aligned} w_t(i') + \sigma_{i'} &= -\frac{\gamma_1}{2n} \bar{\beta} + i' \cdot \frac{\gamma_1 \gamma_3}{4dn} && (\text{Eq. (29)}) \\ &> \frac{\gamma_1 \gamma_3}{2n} && (\bar{\beta} \leq -\gamma_3) \\ &\geq \gamma_1 \bar{v}(i) \bar{\beta} + \frac{\gamma_1 \gamma_3}{2n} && (\bar{v}(i) > 0 \text{ for } i \notin \{\mathcal{I}(S) \cup \{d+2\}\}) \\ &= \gamma_1 \bar{v}(i) \bar{\beta} + \frac{\gamma_1 \gamma_3}{4n} + \frac{\gamma_1 \gamma_3}{4n} \\ &\geq w_t(i) + \sigma_i + \frac{\gamma_1 \gamma_3}{4n}. && (\sigma_i = i \cdot \frac{\gamma_1 \gamma_3}{4dn} \leq \frac{\gamma_1 \gamma_3}{4n}) \end{aligned}$$

Case 3a: $-\gamma_3 < \bar{\beta} \leq 0$ and $\beta_{d+1} > -\frac{3}{2}\gamma_3$. For any $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$ we get,

$$\begin{aligned} w_t(d+1) + \sigma_{d+1} &= \beta_{d+1} + 2\gamma_3 && (\text{Eq. (29)}) \\ &> \frac{1}{2}\gamma_3 && (\beta_{d+1} > -\frac{3}{2}\gamma_3) \\ &\geq \gamma_1 \bar{v}(i) \bar{\beta} + \frac{1}{2}\gamma_3 && (\bar{v}(i) > 0 \text{ for } i \notin \{\mathcal{I}(S) \cup \{d+2\}\}) \\ &\geq \gamma_1 \bar{v}(i) \bar{\beta} + \frac{\gamma_1 \gamma_3}{2n}, && (\gamma_1 \leq n) \end{aligned}$$

and Eq. (32) holds as in the previous case.

Case 3b: $-\gamma_3 < \bar{\beta} \leq 0$ and $\beta_{d+1} \leq -\frac{3}{2}\gamma_3$. Note that this case implies

$$\frac{\gamma_3}{2} < \sum_{i \in \mathcal{I}_t(S) \setminus \{d+1\}} \beta_i. \quad (34)$$

Hence, there exists $i' \in \mathcal{I}_t(S) \setminus \{d+1\}$ such that $\beta_{i'} > \frac{\gamma_3}{2(|\mathcal{I}_t(S)|-1)} \geq \frac{\gamma_3}{2T}$ and we get that for any $i \notin \{\mathcal{I}(S) \cup \{d+2\}\}$,

$$\begin{aligned} w_t(i') + \sigma_{i'} &= \beta_{i'} - \frac{\gamma_1}{2n} \bar{\beta} + i' \cdot \frac{\gamma_1 \gamma_3}{2dn} && \text{(Eq. (29))} \\ &> \frac{\gamma_3}{2T} && (\beta_{i'} > \frac{\gamma_3}{2T} \text{ and } \bar{\beta} \leq 0) \\ &\geq \gamma_1 \bar{v}(i) \bar{\beta} + \frac{\gamma_3}{2T} && (\bar{v}(i) > 0 \text{ for } i \notin \{\mathcal{I}(S) \cup \{d+2\}\}) \\ &\geq \gamma_1 \bar{v}(i) \bar{\beta} + \frac{\gamma_1 \gamma_3}{2n}, && (\gamma_1 \leq \frac{n}{T}) \end{aligned}$$

and Eq. (32) holds as in the previous two cases. \blacksquare

Combining both observations in Claims 4 and 5 and plugging it into Eq. (30) we obtain,

$$\nabla F_S(w) = \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i,$$

for some $i \in \mathcal{I}_{t+1}(S)$.

B.2 Proof of Corollary 5

We assume $w = w_t + q$ with w_t as in Lemma 4 and some q yet to be defined. By standard vector decomposition we know that $\Pi_{t+1}(w)$ can be expressed as

$$\Pi_{t+1}(w) = w_t + \tilde{q}, \quad (35)$$

for some \tilde{q} . Observe that,

$$\begin{aligned} \|\tilde{q}\|_\infty &\leq \|\tilde{q}\|_2 && (\|\cdot\|_\infty \leq \|\cdot\|_2) \\ &= \|\Pi_{t+1}(w) - w_t\|_2 && \text{(Eq. (35))} \\ &\leq \|\Pi_{t+1}(w) - w\|_2 + \|w - w_t\|_2 && \text{(triangle inequality)} \\ &\leq 2\|w - w_t\|_2 && (\|\Pi_{t+1}(w) - w\|_2 \leq \|w_t - w\|_2) \\ &= 2\|q\|_2 && (w = w_t + q) \\ &\leq 2\sqrt{d+2}\|q\|_\infty. && (\|u\|_2 \leq \sqrt{d}\|u\|_\infty \text{ for any } u \in \mathbb{R}^d) \end{aligned}$$

Thus, for $\Pi_{t+1}(w)$ to satisfy the conditions of Lemma 4 in Eq. (13) it suffices that

$$\|q\|_\infty \leq \frac{1}{4\sqrt{d}} \min\left\{\frac{\gamma_2}{3}, \frac{\gamma_1 \gamma_3}{8dn}\right\}.$$

In this case, we can apply Lemma 4 on $\Pi_{t+1}(w)$ and get that

$$\nabla F_S(\Pi_{t+1}(w)) = \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i,$$

for some $i \in \mathcal{I}_{t+1}(S)$. Denote this index by i_{Π}^* , then by the definition of $\nabla F_S(\Pi_{t+1}(w))$ it holds

$$\begin{aligned} i_{\Pi}^* &= \arg \max_{i \in \mathcal{I}_{t+1}(S)} \{\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i \cdot \Pi_{t+1}(w) + \sigma_i\} \\ &= \arg \max_{i \in \mathcal{I}_{t+1}(S)} \{(\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i) \cdot \Pi_{t+1}(w) + \sigma_i\}, \end{aligned}$$

where the last equality is trivial since $\gamma_1 \bar{v} + \varepsilon e_{d+2}$ is independent of the argument i we try to maximize. On the other hand, applying again Lemma 4 on $w = w_t + q$ we get,

$$\nabla F_S(w) = \gamma_1 \bar{v} + \varepsilon e_{d+2} + e_{i^*},$$

for some $i^* \in \mathcal{I}_{t+1}(S)$. Similarly to i_{Π}^* we have

$$i^* = \arg \max_{i \in \mathcal{I}_{t+1}(S)} \{(\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i) \cdot w + \sigma_i\}.$$

Observe that by orthogonal vector decomposition we get that $w = \Pi_{t+1}(w) + w^\perp$ where w^\perp is orthogonal to $\Pi_{t+1}(w)$. Therefore, $(\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i) \cdot w^\perp = 0$ for any $i \in \mathcal{I}_{t+1}(S)$. Consequently we have,

$$i^* = i_{\Pi}^* = \arg \max_{i \in \mathcal{I}_{t+1}(S)} \{(\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_i) \cdot \Pi_{t+1}(w) + \sigma_i\}. \quad (36)$$

And this implies that $\nabla F_S(w) = \nabla F_S(\Pi_{t+1}(w))$. In addition, from the proof of Claim 5 in Lemma 4 we deduce that $g_\gamma(w; \alpha) = g_\gamma(\Pi_{t+1}(w); \alpha) = 0$ for $\alpha \in S$. Also from Eq. (36) we know that $r(w) = w_{i^*} + \sigma_{i^*}$ and $i^* = i_{\Pi}^* \in \mathcal{I}_{t+1}(S)$. Therefore,

$$\begin{aligned} F_S(w) &= \frac{1}{n} \sum_{\alpha \in S} g_\gamma(w; \alpha) + \gamma_1 \bar{v} \cdot w + \varepsilon e_{d+2} \cdot w + r(w) \\ &= (\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_{i_{\Pi}^*}) \cdot w + \sigma_{i_{\Pi}^*} \\ &= (\gamma_1 \bar{v} + \varepsilon e_{d+2} + e_{i_{\Pi}^*}) \cdot \Pi_{t+1}(w) + \sigma_{i_{\Pi}^*} \quad (w = \Pi_{t+1}(w) + w^\perp) \\ &= F_S(\Pi_{t+1}(w)). \end{aligned}$$

C Proof of Lemma 7

Let $[s_1, \dots, s_k]$ be an orthonormal basis for \mathcal{S}_k , and $[s_{k+1}, \dots, s_d]$ an orthonormal basis of \mathcal{S}_k^\perp the orthogonal subspace of \mathcal{S}_k , and S be a matrix whose i th column is s_i . Note that S and S^\top are both orthogonal matrices, i.e., $S^\top S = S S^\top = I$. Now consider the following process of generating an orthogonal matrix U : we pick a random orthogonal matrix $\tilde{U} \in \mathbb{R}^{d_2 \times d}$, and then let $U = \tilde{U} S^\top$. Notice that because random orthogonal matrices are invariant to multiplication by an orthogonal matrix then U is indeed a random orthogonal matrix.

Next, note that the matrix $U\Pi$ determines the first k columns of \tilde{U} . Specifically, for each $i \leq k$:

$$U\Pi s_i = U s_i = \tilde{U} S^\top s_i = \tilde{U} e_i = \tilde{U}_i. \quad (37)$$

Also note that

$$\begin{aligned} \|(1 - \Pi)U^\top w\|_\infty &= \|S S^\top (1 - \Pi) \tilde{U}^\top w\|_\infty \\ &\leq \|S S^\top (1 - \Pi) \tilde{U}^\top w\|_2 \\ &\leq \|S^\top (1 - \Pi) \tilde{U}^\top w\|_2 \quad (S \text{ is orthogonal matrix}) \\ &\leq \sqrt{d} \|S^\top (1 - \Pi) \tilde{U}^\top w\|_\infty. \end{aligned}$$

Now we consider the matrix $Q = \tilde{U}S^\top(1 - \Pi)S$, and we want to calculate its columns. Observe that for $i \leq k$ we have that

$$Q_i = Qe_i = \tilde{U}S^\top(1 - \Pi)Se_i = \tilde{U}S^\top(1 - \Pi)s_i = 0.$$

and for $i > k$:

$$Q_i = Qe_i = \tilde{U}S^\top(1 - \Pi)Se_i = \tilde{U}S^\top s_i = \tilde{U}e_i = \tilde{U}_i.$$

In particular we have that

$$(Qw)_i = \begin{cases} 0 & i \leq k \\ \tilde{U}_i^\top w & i > k. \end{cases}$$

Thus, conditioning on $U\Pi = V$, we have that for the fixed vector w , by Eq. (37):

$$\begin{aligned} & \Pr\left(\|S^\top(1 - \Pi)S\tilde{U}^\top w\|_\infty > \frac{c}{\sqrt{d}} \mid \tilde{U}\Pi = V\right) \\ &= \Pr\left(\|S^\top(1 - \Pi)S\tilde{U}^\top w\|_\infty > \frac{c}{\sqrt{d}} \mid \tilde{U}_i = Vs_i, i = 1, \dots, k\right) \\ &= \Pr\left(\max_{j>k}\{\|Q^\top w\|_\infty\} > \frac{c}{\sqrt{d}} \mid \tilde{U}_i = Vs_i, i = 1, \dots, k\right) \\ &= \Pr\left(\max_{j>k}\{|\tilde{U}_j^\top w|\} > \frac{c}{\sqrt{d}} \mid \tilde{U}_i = Vs_i, i = 1, \dots, k\right) \\ &\leq \sum_{j=k+1}^{d_2} \Pr\left(|\tilde{U}_j^\top w| > \frac{c}{\sqrt{d}} \mid \tilde{U}_i = Vs_i, i = 1, \dots, k\right) \\ &\leq d_2 \Pr\left(|\tilde{U}_{k+1}^\top w| > \frac{c}{\sqrt{d}} \mid \tilde{U}_i = Vs_i, i = 1, \dots, k\right) \end{aligned}$$

where the last equality is due to symmetry. Since \tilde{U}_{k+1} is sampled uniformly from the sphere orthogonal to $\tilde{U}_1, \dots, \tilde{U}_k$ and w is determined by them, we may apply a standard concentration inequality for the inner product of a randomly sampled vector in \mathbb{R}^{d_2-k} against a constant vector $\|w\| \leq 1$ [see, e.g., 2, Lemma 2.2], yielding

$$\Pr\left(|\tilde{U}_{k+1}^\top w| > \frac{c}{\sqrt{d}} \mid \tilde{U}_i = Vs_i, i = 1, \dots, k\right) \leq 2e^{-\frac{c^2}{2d} \cdot (d_2 - k + 1)}.$$

Taking expectation over V , we conclude the proof.