

Extraction of Product and Higher Moment Weak Values: Applications in Quantum State Reconstruction

Sahil,^{1,*} Sohail,^{2,†} Subhrajit Modak,^{3,‡} Sibasish Ghosh,^{1,§} and Arun Kumar Pati^{2,¶}

¹*Optics & Quantum Information Group, The Institute of Mathematical Sciences, HBNI, CIT Campus, Taramani, Chennai 600113, India.*

²*Quantum Information & Computation Group, Harish-Chandra Research Institute, HBNI, Allahabad 211019, India.*

³*Department of Physical Sciences, Indian Institute of Science Education and Research Mohali, Sector 81, S. A. S. Nagar, Manauli PO 140306, Punjab, India.*

Weak measurements introduced by Aharonov, Albert and Vaidman (AAV) can provide informations about the system with minimal back action. Weak values of product observables (commuting) or higher moments of an observable are informationally important in the sense that they are useful to resolve some paradoxes, realize strange quantum effects, construct density matrices, etc. In this work, we show that it is possible to access the higher moment weak values of an observable using weak values of that observable with pairwise orthogonal post-selections. Although the higher moment weak values of an observable are inaccessible with Gaussian pointer states, our method allows any pointer state. We have calculated product weak values in a bipartite system for any given pure and mixed pre selected states. Such product weak values can be obtained using only the measurements of local weak values (which are defined as single system weak values in a multi-partite system). As an application, we use higher moment weak values and product weak values to reconstruct unknown quantum states of single and bipartite systems, respectively. Robustness of our method which occurs due to inappropriate choices of quantum observables and noisy post-selections is also discussed here. Our method can easily be generalized to the multi-partite systems.

I. INTRODUCTION

In quantum systems, measurement of product observables of two or more can contain information about quantum correlations and quantum dynamics. In the recent past, many theoretical and experimental works have been performed regarding the measurements of product observables using weak measurements. They have been used to resolve the Hardy's Paradox [1] with experimental verification [2], EPR-Bohm experiment [3], direct measurement of a density matrix [4], reconstruction of entangled quantum states [5]. It has also been reported that a strange quantum effect namely the "Quantum Cheshire Cats" where the properties of a quantum particle can be disembodied (e.g., photon's position and polarization degrees of freedoms can be separated from each other) can be realized using product weak values [6]. See also the references [7] and [8] for the experimental test of the existence of Quantum Cheshire Cats and the exchange of grins between two such Quantum Cheshire Cats, respectively. Weak measurements of product observables also play an important role in understanding the quantum mechanics such as Bell tests [9–11], nonlocality via post-selection [12].

Weak measurements in particular weak values are known to provide useful informations with simple experimental setups. The "weak value" was first introduced by Aharonov, Albert, and Vaidman (AAV) [13]. It was inspired by the two-time formulation of the quantum-mechanical system [14]. The mechanism of AAV method is that they took the von Neumann mea-

surement scheme (VMS) [15] one step forward by considering the coupling coefficient very small i.e., weak followed by a strong measurement in succession. This formulation is characterized by the pre- and postselected states of the system. By preparing a system initially in the state $|\psi\rangle$ and post-selecting in the state $|\phi\rangle$, as a result we obtain the weak value of any observable A which is defined as

$$\langle A_w \rangle_\psi^\phi = \frac{\langle \phi | A | \psi \rangle}{\langle \phi | \psi \rangle}. \quad (1)$$

This is a complex number and the spatial and momentum displacements of the pointer state give the real and imaginary parts of that weak value [16, 17], respectively. By this way, we get the full knowledge of the complex weak value. One of the exciting and interesting features of weak value is that it can lie outside the max-min range of the eigenvalues of the operator of interest. Simultaneously, measurement disturbance is quite small which gives one to perform further measurements or simultaneous measurement of multiple observables.

Weak measurements have been proven useful in understanding quantum systems such as for the direct measurement of the wave function of a quantum system [18], to calculate slow- and fast-light effects in birefringent photonic crystals [19], the confirmation of the Heisenberg-Ozawa uncertainty relationship [20], for detecting tiny spatial shifts [21, 22]. Weak value can also be used to measure non-Hermitian operators [23, 24], in hot thermometry [25], to detect entanglement universally in a two-qubit system [26].

Product weak values: The observables used in (VMS) are of simple kinds. Direct measurement of product observables is extremely difficult whether it is strong or weak. This difficulty arises from the fact that the measurement interaction in VMS couples two different observables to a single pointer [16]. To overcome this, an approach using multi particle interaction Hamiltonian was proposed by Resch and Steinberg

* sahilmd@imsc.res.in

† sohail@hri.res.in

‡ modoksuvrajit@gmail.com

§ sibasish@imsc.res.in

¶ akpati@hri.res.in

(RS) [27] applying AAV method. Namely, they have used the Hamiltonian of the form $H = g_1 A \otimes p_x + g_2 B \otimes p_y$ with gaussian pointer states. Here A and B are two observables of the system. p_x and p_y are the pointer's momentum degrees of freedom in two different directions x and y, respectively. g_1 and g_2 are coupling coefficients between the system and the pointer for two different directions, respectively. By performing a second order expansion in the two-dimensional pointer displacement $\langle XY \rangle$, they showed that it is possible to extract the real part of the product weak value for the case of commuting observables $[A, B] = 0$, namely $Re[\langle (AB)_w \rangle_\psi^\phi] = 2\langle XY \rangle / (g_1 g_2 t^2) - Re[\langle (A_w)_w \rangle_\psi^\phi * \langle (B_w)_w \rangle_\psi^\phi]$, where 't' is the interaction time. For imaginary part of the product weak value, one needs to look into $\langle XP_y \rangle$. Note that, the weak value of a tensor product observable in a bipartite system (we will call it product weak value in a bipartite system) can also be calculated according to the RS method as the local observables are commuting. It's generalised version i.e., the product weak values $\langle (AB)_w \rangle_\psi^\phi$ and it's higher orders i.e., $\langle (ABC \dots)_w \rangle_\psi^\phi$ using N pointers' correlations can be found in Ref. [28].

Summary of the present work: In this work, we show that by introducing a unique orthogonal state to the given post selection, it is possible to extract the higher moment weak values $\langle (A^n)_w \rangle_\psi^\phi$ in a single system as well as the product weak values for the given pure and mixed pre selected states in a bipartite system separately. We show application of our results to reconstruct quantum states of single and bipartite systems. We have used higher moment weak values to reconstruct a pure state of a single system and product weak values of a bipartite system to reconstruct unknown pure and mixed states of that bipartite system. Recently, Pan *et al.* [29] have used product weak values of projection operators to reconstruct an unknown bipartite pure state. They have considered entangled pointer states as well as the modular values of local projection operators and the modular values of sum of the local projection operators [30] to evaluate product weak values. We, for the first time show that such product weak values can be realized locally for the case of both pure and mixed states. Also, to reconstruct the states of single and bipartite systems, we have generalized the measurement of projection operators to arbitrary observables. Our method can be generalized to the multipartite systems. Finally we show that our methods of "extraction of product and higher moment weak values" are robust against the errors which occur due to the inappropriate choices of system observables and unsharp post-selections.

This paper is organized as follows. In sec. II we provide the formulation of our method. In sec. III we apply our method to reconstruct quantum states of single as well as bipartite systems separately. We show the robustness of our method in sec. IV and finally conclude in sec. V.

II. FORMULATION

The following identity which is sometimes referred as Vaidman's formula [31] will be used to derive the main results of

this paper

$$A|\phi\rangle = \langle A \rangle_\phi |\phi\rangle + \langle \Delta A \rangle_\phi |\phi_A^\perp\rangle, \quad (2)$$

where A is an Hermitian operator and $|\phi\rangle$ is any quantum state vector in the Hilbert space \mathcal{H} . The state vector $|\phi_A^\perp\rangle$ is orthogonal to $|\phi\rangle$, $\langle A \rangle_\phi = \langle \phi|A|\phi\rangle$ and $\langle \Delta A \rangle_\phi = \langle \phi_A^\perp|A|\phi\rangle$. For the derivation see Appendix A

A. Higher moment weak values

If $|\psi\rangle$, A and $|\phi\rangle$ are the pre-selected state, the observable and the post-selected state, respectively, then the weak value of the observable A is given by

$$\langle A_w \rangle_\psi^\phi = \left(\frac{\langle \psi|A|\phi\rangle}{\langle \psi|\phi\rangle} \right)^* = \langle A \rangle_\phi + \langle \Delta A \rangle_\phi \frac{\langle \phi_A^\perp|\psi\rangle}{\langle \phi|\psi\rangle}, \quad (3)$$

where we have used Eq. (2). A similar expression was considered in Ref. [32] to explain the origin of the complex and anomalous nature of a weak value. The Eq. (3) for the expression of the weak value will be useful for deriving the following result.

Result 1:- The weak value of the operator A^2 which we call the "second moment weak value" has the following expression

$$\langle (A^2)_w \rangle_\psi^\phi = \langle A \rangle_\phi \left(\langle A_w \rangle_\psi^\phi - \langle A_w \rangle_\psi^{\phi_A^\perp} \right) + \langle A_w \rangle_\psi^\phi \langle A_w \rangle_\psi^{\phi_A^\perp}, \quad (4)$$

where $\langle A_w \rangle_\psi^\phi$ and $\langle A_w \rangle_\psi^{\phi_A^\perp}$ are the weak values of the operator A for the given pre selection $|\psi\rangle$ with two post-selections $|\phi\rangle$ and $|\phi_A^\perp\rangle$, respectively.

Proof.

$$\begin{aligned} \langle (A^2)_w \rangle_\psi^\phi &= \left(\frac{\langle \psi|A^2|\phi\rangle}{\langle \psi|\phi\rangle} \right)^* \\ &= \left(\frac{1}{\langle \psi|\phi\rangle} \langle \psi|A[\langle A \rangle_\phi |\phi\rangle + \langle \Delta A \rangle_\phi |\phi_A^\perp\rangle] \right)^* \\ &= \left(\langle A \rangle_\phi \frac{\langle \psi|A|\phi\rangle}{\langle \psi|\phi\rangle} + \langle \Delta A \rangle_\phi \frac{\langle \psi|A|\phi_A^\perp\rangle}{\langle \psi|\phi\rangle} \right)^* \\ &= \langle A \rangle_\phi \frac{\langle \phi|A|\psi\rangle}{\langle \phi|\psi\rangle} + \langle \Delta A \rangle_\phi \frac{\langle \phi_A^\perp|A|\psi\rangle}{\langle \phi_A^\perp|\psi\rangle} \frac{\langle \phi_A^\perp|\psi\rangle}{\langle \phi|\psi\rangle}. \end{aligned} \quad (5)$$

From Eq. (3), using $\langle \Delta A \rangle_\phi \frac{\langle \phi_A^\perp|\psi\rangle}{\langle \phi|\psi\rangle} = \langle A_w \rangle_\psi^\phi - \langle A \rangle_\phi$ in Eq. (5), we will obtain Eq. (4). \square

In the similar way we obtain all the higher moment weak values which take the general form as

$$\langle A_w^n \rangle_\psi^\phi = \langle A \rangle_\phi \left(\langle A_w^{n-1} \rangle_\psi^\phi - \langle A_w^{n-1} \rangle_\psi^{\phi_A^\perp} \right) + \langle A_w^{n-1} \rangle_\psi^{\phi_A^\perp} \langle A_w \rangle_\psi^\phi, \quad (6)$$

for $n = 1, 2, \dots$.

Now consider the second moment weak value i.e., Eq. (4), where $\langle A_w \rangle_\psi^\phi$ and $\langle A_w \rangle_\psi^{\phi^\perp}$ are extractable from the one and the same experimental set-up for the post-selection of $|\phi\rangle$ and $|\phi^\perp\rangle$, respectively as these two states are orthogonal to each other. Note that, although the post-selection can be realized here in one and the same measurement set-up, nevertheless, in order to actually find out the weak values $\langle A_w \rangle_\psi^\phi$ and $\langle A_w \rangle_\psi^{\phi^\perp}$, measurements of phase-space displacements for the two post-selected states $|\phi\rangle$ and $|\phi^\perp\rangle$ need to be performed. $\langle A \rangle_\phi$ is the average value of A for the post selected state $|\phi\rangle$. Extraction of higher moment weak values becomes extremely simple in two dimensional Hilbert space discussed in the following.

Two dimensional case: In two dimensional Hilbert space, there are only two pairwise orthogonal post-selections which occur at the same time and hence both the weak values with orthogonal post-selections can be extracted simultaneously. Particularly in this dimension, we find that only by knowing the weak values $\langle A_w \rangle_\psi^\phi$ and $\langle A_w \rangle_\psi^{\phi^\perp}$, we are able to obtain all the higher moment weak values without any further complications in comparison to Ref. [33]. So the number of measurements is being reduced considerably than the earlier proposal [33].

Higher dimensional case: In higher dimensional Hilbert space, to extract second moment weak value of the observable A , one can perform the projective measurements $\{|\phi\rangle\langle\phi|, |\phi^\perp\rangle\langle\phi^\perp|, I - |\phi\rangle\langle\phi| - |\phi^\perp\rangle\langle\phi^\perp|\}$ for post-selections.

It can be shown that the n -th moment weak value i.e., $\langle A_w^n \rangle_\psi^\phi$, consists n number of different weak values, namely $\langle A_w \rangle_\psi^\phi, \langle A_w \rangle_\psi^{\phi^\perp}, \dots, \langle A_w \rangle_\psi^{(\phi^\perp)^{n-1}}$. Here $|(\phi^\perp)^{n-1}\rangle = |((\phi^\perp)^\perp)^\perp \dots (n-1) \text{ times}\rangle = \frac{1}{\langle \Delta A \rangle_{(\phi^\perp)^{n-2}}} (A - \langle A \rangle_{(\phi^\perp)^{n-2}}) |(\phi^\perp)^{n-2}\rangle$. Now, if n is even, then there exist pairwise orthogonal post-selected states i.e., $(|\phi\rangle, |\phi^\perp\rangle), (|(\phi^\perp)^\perp\rangle, |((\phi^\perp)^\perp)^\perp\rangle), \dots, (|(\phi^\perp)^{n-2}\rangle, |(\phi^\perp)^{n-1}\rangle)$.

So, it is possible to obtain weak values $\langle A_w \rangle_\psi^\phi$ and $\langle A_w \rangle_\psi^{\phi^\perp}$ simultaneously for the first pair of post-selected states, so on and so forth. So, effectively the total number of measurements to be performed according to the AAV method to extract the n -th moment weak value is $n/2$. For odd n , the number of measurements is $(n+1)/2$. Note that, all the measurements here are to be done by only changing the post-selections while keeping the observable A fixed in the AAV method.

Result 2:- The product weak value in Eq. (7) can be realized via local weak values as

$$\langle (A \otimes B)_w \rangle_{\psi_{AB}}^{\phi_A \phi_B} = \langle A \rangle_{\phi_A} \left(\langle B_w^{local} \rangle_{\psi_{AB}}^{\phi_A \phi_B} - \langle B_w^{local} \rangle_{\psi_{AB}}^{\phi_A^\perp \phi_B} \right) + \langle A_w^{local} \rangle_{\psi_{AB}}^{\phi_A \phi_B} \langle B_w^{local} \rangle_{\psi_{AB}}^{\phi_A^\perp \phi_B} \quad (9)$$

where $\langle A_w^{local} \rangle_{\psi_{AB}}^{\phi_A \phi_B}$, $\langle B_w^{local} \rangle_{\psi_{AB}}^{\phi_A \phi_B}$ and $\langle B_w^{local} \rangle_{\psi_{AB}}^{\phi_A^\perp \phi_B}$ are the ‘‘local’’ weak values and $|\phi_A^\perp\rangle = \frac{1}{\langle \Delta A \rangle_{\phi_A}} (A - \langle A \rangle_{\phi_A}) |\phi_A\rangle$ is given by (2) for the subsystem A.

Once we extract the n -th moment weak value, then all the lower moment weak values can be calculated from the data of n -th moment weak value.

Previously, it was thought that with Gaussian pointer states, one is not able to extract second or higher moment weak values. In Ref. [33], the authors have demonstrated that by employing two-dimensional pointer states endowed with orbital angular momentum (OAM) in the RS method [27], it is possible to extract weak values of the higher moments of single-particle operators. Essentially, the two dimensional OAM states are not factorizable in two different directions for non-zero orbital angular momentum (in fact, OAM states with zero angular momentum correspond to two dimensional Gaussian pointer states) and this is the key factor to retrieve higher moment weak values. In doing that one needs to engineer OAM states with higher winding numbers, or superpositions of OAM states to obtain the higher moment weak values. This procedure can become difficult as the moments increase. One needs to prepare pointer states with different combination of orbital angular momentum [33]. Moreover, there are several disadvantages of the RS method from experimental perspective which we have discussed later in this section.

As an application, we will use the higher moment weak values to reconstruct an unknown pure state of a single system. See Appendix B for the derivation of the second or higher moment weak values for the mixed pre selected state case.

B. Product weak values

Product weak values for pure pre-selected state: The product weak value of the observable $A \otimes B$ in a bipartite system is given by

$$\langle (A \otimes B)_w \rangle_{\psi_{AB}}^{\phi_A \phi_B} = \frac{\langle \phi_A \phi_B | (A \otimes B) | \psi_{AB} \rangle}{\langle \phi_A \phi_B | \psi_{AB} \rangle}, \quad (7)$$

where the pre selection is the bipartite pure state $|\psi_{AB}\rangle$ and the post-selection is a product state $|\phi_A \phi_B\rangle = |\phi_A\rangle \otimes |\phi_B\rangle$.

Consider the ‘‘local weak value’’ of the operator A

$$\begin{aligned} \langle A_w^{local} \rangle_{\psi_{AB}}^{\phi_A \phi_B} &= \frac{\langle \phi_A \phi_B | (A \otimes I_B) | \psi_{AB} \rangle}{\langle \phi_A \phi_B | \psi_{AB} \rangle} \\ &= \langle A \rangle_{\phi_A} + \langle \Delta A \rangle_{\phi_A} \frac{\langle \phi_A^\perp \phi_B | \psi_{AB} \rangle}{\langle \phi_A \phi_B | \psi_{AB} \rangle}, \end{aligned} \quad (8)$$

where we have used Eq. (2) in the subsystem A. Eq. (8) will be used to derive the following result.

Proof.

$$\begin{aligned}
\langle (A \otimes B)_w \rangle_{\psi_{AB}}^{\phi_A \phi_B} &= \left(\frac{\langle \psi_{AB} | (A \otimes B) | \phi_A \phi_B \rangle}{\langle \psi_{AB} | \phi_A \phi_B \rangle} \right)^* \\
&= \left(\frac{\langle \psi_{AB} | (\langle A \rangle_{\phi_A} |\phi_A\rangle + \langle \Delta A \rangle_{\phi_A} |\phi_A^\perp\rangle) \otimes B | \phi_B \rangle}{\langle \psi_{AB} | \phi_A \phi_B \rangle} \right)^* \\
&= \langle A \rangle_{\phi_A} \frac{\langle \phi_A \phi_B | (I_A \otimes B) | \phi_{AB} \rangle}{\langle \phi_A \phi_B | \psi_{AB} \rangle} + \langle \Delta A \rangle_{\phi_A} \frac{\langle \phi_A^\perp \phi_B | (I_A \otimes B) | \phi_{AB} \rangle}{\langle \phi_A^\perp \phi_B | \psi_{AB} \rangle} \frac{\langle \phi_A^\perp \phi_B | \psi_{AB} \rangle}{\langle \phi_A \phi_B | \psi_{AB} \rangle} \\
&= \langle A \rangle_{\phi_A} \langle B_w^{local} \rangle_{\psi_{AB}}^{\phi_A \phi_B} + \langle B_w^{local} \rangle_{\psi_{AB}}^{\phi_A^\perp \phi_B} \left(\langle A_w^{local} \rangle_{\psi_{AB}}^{\phi_A \phi_B} - \langle A \rangle_{\phi_A} \right),
\end{aligned}$$

where we have used Eq. (2) in the second line for the subsystem A and Eq. (8) in the fourth line. After the manipulation, we have Eq. (9). \square

Product weak values in terms of local weak values: We have obtained a product weak value using only local weak values. Note that the local weak values $\langle B_w^{local} \rangle_{\psi_{AB}}^{\phi_A \phi_B}$ and $\langle B_w^{local} \rangle_{\psi_{AB}}^{\phi_A^\perp \phi_B}$ can be measured in the same experimental setup as the post-selected states $|\phi_A\rangle$ and $|\phi_A^\perp\rangle$ are orthogonal to each other. We need another measurement setup for $\langle A_w^{local} \rangle_{\psi_{AB}}^{\phi_A \phi_B}$. So, effectively the total number of measurements to be performed according to the AAV method to extract the product weak value $\langle (A \otimes B)_w \rangle_{\psi_{AB}}^{\phi_A \phi_B}$ is only two. In experiment, local weak values like $\langle B_w^{local} \rangle_{\psi_{AB}}^{\phi_A \phi_B}$ can be realized as

$$\langle B_w^{local} \rangle_{\psi_{AB}}^{\phi_A \phi_B} = \frac{\langle \phi_B \phi_A | (I_A \otimes B) | \psi_{AB} \rangle}{\langle \phi_B \phi_A | \psi_{AB} \rangle} \quad (10)$$

$$= \frac{\langle \phi_B | B | \psi_B^{\phi_A} \rangle}{\langle \phi_B | \psi_B^{\phi_A} \rangle}, \quad (11)$$

where $|\psi_B^{\phi_A}\rangle = \langle \phi_A | \psi_{AB} \rangle$ is an unnormalized state for the

subsystem B. That is, we first measure the projection operator $\Pi_{\phi_A} = |\phi_A\rangle \langle \phi_A|$ on the subsystem A, then the state of the subsystem B becomes $\langle \phi_A | \psi_{AB} \rangle / \sqrt{\langle \psi_B^{\phi_A} | \psi_B^{\phi_A} \rangle}$ which we consider to be pre-selected state for the subsystem B. This pre-selected state is unknown as $|\psi_{AB}\rangle$ is unknown. So, for each given projector related to the subsystem A, there exists an unknown pre-selected state of the subsystem B. The observable is B and the post-selection is $|\phi_B\rangle$. So from Eq. (11), we see that the local weak value $\langle B_w^{local} \rangle_{\psi_{AB}}^{\phi_A \phi_B}$ can be realized according to the AAV method related to the subsystem B.

Product weak values for mixed pre-selected state: If the pre selection of the bipartite system is a mixed state ρ_{AB} then, the product weak value of the observable $A \otimes B$ is given by

$$\langle (A \otimes B)_w \rangle_{\rho_{AB}}^{\phi_A \phi_B} = \frac{\langle \phi_A \phi_B | (A \otimes B) \rho_{AB} | \phi_A \phi_B \rangle}{\langle \phi_A \phi_B | \rho_{AB} | \phi_A \phi_B \rangle}. \quad (12)$$

Note that generalization for mixed state of the Eq. (9) is not straightforward.

Result 3:- The product weak value in Eq. (12) can be realized via local weak values as

$$\begin{aligned}
\langle (A \otimes B)_w \rangle_{\rho_{AB}}^{\phi_A \phi_B} &= \langle A \rangle_{\phi_A} \langle B_w^{local} \rangle_{\rho_{AB}}^{\phi_A \phi_B} + \frac{\langle \Delta A \rangle_{\phi_A}}{2p(\rho_{AB}, \phi_A \phi_B)} \sum_{i=1}^m \left(\lambda_A^i \langle B_w^{local} \rangle_{\rho_{AB}}^{i_A \phi_B} p(\rho_{AB}, i_A \phi_B) \right. \\
&\quad \left. + \lambda_A^{i'} \langle B_w^{local} \rangle_{\rho_{AB}}^{i'_A \phi_B} p(\rho_{AB}, i'_A \phi_B) \right), \quad (13)
\end{aligned}$$

where $\{\lambda_A^i, |i_A\rangle\}$ and $\{\lambda_A^{i'}, |i'_A\rangle\}$ satisfy the spectral decomposition for the normal operators $|\phi_A\rangle \langle \phi_A^\perp| + |\phi_A^\perp\rangle \langle \phi_A|$ and $|\phi_A\rangle \langle \phi_A^\perp| - |\phi_A^\perp\rangle \langle \phi_A|$, respectively. $p(\rho_{AB}, \phi_A \phi_B) = \langle \phi_A \phi_B | \rho_{AB} | \phi_A \phi_B \rangle$ is the probability of obtaining the post-selected state $|\phi_A \phi_B\rangle$ for the given pre selected state ρ_{AB} and ‘ m ’ is the dimension of the subsystem A.

Proof.

$$\begin{aligned}
\langle (A \otimes B)_w \rangle_{\rho_{AB}}^{\phi_A \phi_B} &= \frac{\langle \phi_A \phi_B | (A \otimes B) \rho_{AB} | \phi_A \phi_B \rangle}{\langle \phi_A \phi_B | \rho_{AB} | \phi_A \phi_B \rangle} \\
&= \langle A \rangle_{\phi_A} \frac{\langle \phi_A \phi_B | (I_A \otimes B) \rho_{AB} | \phi_A \phi_B \rangle}{\langle \phi_A \phi_B | \rho_{AB} | \phi_A \phi_B \rangle} + \langle \Delta A \rangle_{\phi_A} \frac{\langle \phi_A^\perp \phi_B | (I_A \otimes B) \rho_{AB} | \phi_A \phi_B \rangle}{\langle \phi_A \phi_B | \rho_{AB} | \phi_A \phi_B \rangle}, \quad (14)
\end{aligned}$$

where we have used Eq. (2). Now $\langle \phi_A^\perp \phi_B | (I_A \otimes B) \rho_{AB} | \phi_A \phi_B \rangle = \text{Tr} [(|\phi_A\rangle \langle \phi_A^\perp| \otimes |\phi_B\rangle \langle \phi_B|) (I_A \otimes B) \rho_{AB}]$ can be calculated as

$$\langle \phi_A^\perp \phi_B | (I_A \otimes B) \rho_{AB} | \phi_A \phi_B \rangle + \langle \phi_A \phi_B | (I_A \otimes B) \rho_{AB} | \phi_A^\perp \phi_B \rangle = \text{Tr} [(|\phi_A\rangle \langle \phi_A^\perp| + |\phi_A^\perp\rangle \langle \phi_A|) \otimes |\phi_B\rangle \langle \phi_B| (I_A \otimes B) \rho_{AB}], \quad (15)$$

where $|\phi_A\rangle \langle \phi_A^\perp| + |\phi_A^\perp\rangle \langle \phi_A|$ is a normal operator satisfying $XX^\dagger = X^\dagger X$ (where X is any operator) and hence can be written in spectral decomposition

$$|\phi_A\rangle \langle \phi_A^\perp| + |\phi_A^\perp\rangle \langle \phi_A| = \sum_i^m \lambda_A^i |i_A\rangle \langle i_A|, \quad (16)$$

where $\{|i_A\rangle\}$ is the set of eigenvectors with eigenvalues $\{\lambda_A^i\}$. Using Eq. (16) in (15), we have

$$\begin{aligned} \langle \phi_A^\perp \phi_B | (I_A \otimes B) \rho_{AB} | \phi_A \phi_B \rangle + \langle \phi_A \phi_B | (I_A \otimes B) \rho_{AB} | \phi_A^\perp \phi_B \rangle &= \sum_i^m \lambda_A^i \langle i_A \phi_B | (I_B \otimes B) \rho_{AB} | i_A \phi_B \rangle \\ &= \sum_i^m \lambda_A^i \langle B_w^{local} \rangle_{\rho_{AB}}^{i_A \phi_B} p(\rho_{AB}, i_A \phi_B), \end{aligned} \quad (17)$$

where we have used the definition of the weak value $\langle B_w \rangle_{\rho_{AB}}^{i_A \phi_B}$ of the observable B for the given pre and post-selections ρ_{AB} and $|i_A \phi_B\rangle$, respectively. $p(\rho_{AB}, i_A \phi_B) = \langle i_A \phi_B | \rho_{AB} | i_A \phi_B \rangle$ is the probability of obtaining the product state $|i_A \phi_B\rangle$. Now similarly,

$$\langle \phi_A^\perp \phi_B | (I_A \otimes B) \rho_{AB} | \phi_A \phi_B \rangle - \langle \phi_A \phi_B | (I_A \otimes B) \rho_{AB} | \phi_A^\perp \phi_B \rangle = \sum_i^m \lambda_A^i \langle B_w^{local} \rangle_{\rho_{AB}}^{i'_A} p(\rho_{AB}, i'_A \phi_B), \quad (18)$$

where we have used the fact that $|\phi_A\rangle \langle \phi_A^\perp| - |\phi_A^\perp\rangle \langle \phi_A|$ is also a normal operator with the spectral decomposition

$$|\phi_A\rangle \langle \phi_A^\perp| - |\phi_A^\perp\rangle \langle \phi_A| = \sum_i^m \lambda_A^i |i'_A\rangle \langle i'_A|. \quad (19)$$

Now adding two equations (17) and (18), we have

$$\langle \phi_A^\perp \phi_B | (I_A \otimes B) \rho_{AB} | \phi_A \phi_B \rangle = \frac{1}{2} \sum_i^m \left(\lambda_A^i \langle B_w^{local} \rangle_{\rho_{AB}}^{i_A \phi_B} p(\rho_{AB}, i_A \phi_B) + \lambda_A^i \langle B_w^{local} \rangle_{\rho_{AB}}^{i'_A \phi_B} p(\rho_{AB}, i'_A \phi_B) \right). \quad (20)$$

Using Eq. (20) in (14), we obtain Eq. (13). □

Product weak values in terms of local weak values: The product weak value $\langle (A \otimes B)_w \rangle_{\rho_{AB}}^{\phi_A \phi_B}$ consists the local weak values $\langle B_w^{local} \rangle_{\rho_{AB}}^{\phi_A \phi_B}$, $\{\langle B_w^{local} \rangle_{\rho_{AB}}^{i_A \phi_B}\}_{i=1}^m$ and $\{\langle B_w^{local} \rangle_{\rho_{AB}}^{i'_A \phi_B}\}_{i=1}^m$. Here ‘ m ’ is the dimension of the subsystem A. Note that, $\{\langle B_w^{local} \rangle_{\rho_{AB}}^{i_A \phi_B}\}_{i=1}^m$ can be measured in the same experimental setup according to the AAV method as the post-selections $\{|i_A\rangle\}$ form complete set of orthogonal basis. Similarly $\{\langle B_w^{local} \rangle_{\rho_{AB}}^{i'_A \phi_B}\}_{i=1}^m$ can also be measured within the same experiment with the complete set of post-selections $\{|i'_A\rangle\}$ according to the AAV method. So, the total number of measurements to be performed according to the AAV method to extract $\langle (A \otimes B)_w \rangle_{\rho_{AB}}^{\phi_A \phi_B}$ is only three. Local weak values can be realized in the same way as discussed above in Eq. (11).

Comparison with previous works: In the Ref. [16, 27], authors have shown that it is also possible to extract the product weak values by obtaining local weak values of the observables separately as well as by looking into different pointers’ position and momentum correlations i.e., statistical averages of these different degrees of freedom. There are some cases where polarization degrees of freedom (polarization correlations) was considered instead of position and momentum variables [2]. For such cases statistical averages of those polarization degrees of freedom can be difficult to obtain or number of measurements will be large. Instead, our approach shows that one needs to obtain only local weak values. Our approach is not necessarily restricted with Gaussian pointer states. But in most of the previous works [16, 27, 34], different type of pointer states have been used with certain conditions.

Our methods are easy to perform in experiments due to their

local realizations. While in the previous works [16, 27, 33, 34] i.e., extraction of higher moment weak values and product weak values, there are N pointers' correlations to be measured. So the scalability of their methods face significant challenges. Moreover, from an experimental perspective, their schemes are hard to implement as the resources required to observe the correlations are of second order in terms of the interaction coefficient.

In the next section, we discuss about the applications of our results to reconstruct unknown quantum states in a single and bipartite systems.

III. QUANTUM STATE TOMOGRAPHY

In the following, we discuss some methods of quantum state tomography of a single and bipartite system using higher moments weak values and product weak values.

A. State reconstruction of a single system

1. Pure state

As an application of higher moment weak values we reconstruct a pure state. The method of quantum state reconstruction using weak values was introduced by Lundeen *et al.* [18] as follows. Any state can be written in computational basis $\{|i\rangle\}$ as

$$|\psi\rangle = \sum_{i=0} \alpha_i |i\rangle, \quad (21)$$

where $\alpha_i = \langle i|\psi\rangle$. Now the weak value of a projection operator $\Pi_i = |i\rangle\langle i|$ is given by

$$\langle (\Pi_i)_w \rangle_\psi^b = \frac{\langle b|i\rangle \langle i|\psi\rangle}{\langle b|\psi\rangle}, \quad (22)$$

with $\langle b|i\rangle \neq 0$. where $|b\rangle$ is a post-selection. So using Eq. (22), we finally construct the pure state Eq. (21)

$$|\psi\rangle = \sum_i \frac{\langle (\Pi_i)_w \rangle_\psi^b}{\langle b|i\rangle} |i\rangle. \quad (23)$$

The complex number $\langle b|\psi\rangle$ is not taken into account as it corresponds to the global phase factor after normalization. So to measure a pure state, we need to obtain weak values of the projection operators Π_i with pre selection $|\psi\rangle$ and post-selection $|b\rangle$, respectively.

Instead of measuring weak values of the projection operators individually, we want to use the higher moment weak values of the observable which satisfies spectral decomposition with those projection operators and using those higher moment weak values we will obtain weak values of the projection operators. Let the observable be

$$A = \sum_i a_i \Pi_i, \quad (24)$$

where a_i are the eigenvalues of the observable A . Now the weak value and higher moment weak values of the observable are

$$\langle A_w \rangle_\psi^b = \sum_i a_i \langle (\Pi_i)_w \rangle_\psi^b, \quad (25)$$

$$\langle A_w^n \rangle_\psi^b = \sum_i a_i^n \langle (\Pi_i)_w \rangle_\psi^b, \quad (26)$$

where ' n ' is any positive integer. Eqs. (25) and (26) can be solved to obtain the weak values of projection operators for different ' n '. For example in three dimensional Hilbert Space, we require only up to second moment weak values because one can use the completeness relation for the projection operators with pre selection $|\psi\rangle$ and post-selection $|b\rangle$

$$1 = \sum_i \langle (\Pi_i)_w \rangle_\psi^b. \quad (27)$$

In Appendix C, we explicitly show how to solve the above equations to obtain the weak values of the projection operators.

From (C1), the highest moment weak value is $\langle A_w^{d-1} \rangle_\psi^b$ and as we have discussed in (II A) that, extracting the highest moment weak value is enough to calculate all the lower moments weak values. *Hence the total number of measurements needed to reconstruct a pure state is $d/2$ if the dimension d is even and $(d-1)/2$ if the dimension d is odd* (see II A for detail discussion).

To compare with Lundeen *et al.* [18] and Wu [35], the number of measurement operators which is the complete set of projection operators with a fixed post-selection is $(d-1)$. Moreover, we measure only one system operator A , but post-selections are to be changed, while, in their method, there are $(d-1)$ system operators (projection operators) to be measured according to the AAV method.

Alternative:– The weak value of an observable C with pre and post-selections $|\psi\rangle$ and $|0\rangle$, respectively is

$$\langle C_w \rangle_\psi^0 = \frac{\langle 0|C|\psi\rangle}{\langle 0|\psi\rangle}. \quad (28)$$

Inserting the identity operator $I = \sum_i |i\rangle\langle i|$ in numerator of the right hand side of Eq. (28), we have

$$\langle C_w \rangle_\psi^0 - C_{00} = \sum_{i=1} C_{0i} \frac{\alpha_i}{\alpha_0}, \quad (29)$$

where $C_{0i} = \langle 0|C|i\rangle$. Like Eq. (29), we have to measure a set of observables to obtain the values of α_1/α_0 , $\alpha_2/\alpha_0, \dots, \alpha_{(d-1)}/\alpha_0$ (see Appendix C). The value of α_0 can be obtained from the normalization condition. Here, the number of measurement operators is $(d-1)$. This method will be used to reconstruct an unknown quantum pure state of a bipartite system.

2. Mixed state

Measurement of a mixed state of a quantum system was also introduced by Lundeen *et al.* [4] using product weak values of two non commuting projection operators. It was later

simplified by Shengjun Wu [35] where weak values of complete set of projection operators with complete set of post-selected states have been used. Here, we develop another method by using the weak values of arbitrary observables which can be thought as the generalization of the reference [35]. This part is added here because we will use the same procedure while dealing with bipartite mixed state reconstruction using product weak values.

Let the unknown mixed state be the pre-selected state then the weak value of the observable C with post-selection $|j\rangle$ is

$$\langle C_w \rangle_\rho^j = \frac{\langle j|C\rho|j\rangle}{\langle j|\rho|j\rangle}. \quad (30)$$

Inserting the identity operator $I = \sum_i |i\rangle \langle i|$, we have

$$p(\rho, j) \langle C_w \rangle_\rho^j = \sum_i C_{ji} \rho_{ij}, \quad (31)$$

where $p(\rho, j) = \langle j|\rho|j\rangle$ is the probability of obtaining the basis state $|j\rangle$ as a post-selection and $C_{ji} = \langle j|C|i\rangle$, $\rho_{ij} = \langle i|\rho|j\rangle$. To obtain the j -th column of the density matrix ρ from Eq. (31), we need to measure a set of arbitrary observables according to the AAV method to get a set of equations like (31) (see Appendix D).

To compare with the work by Lundeen *et al.* [4] where each matrix element is directly obtainable via sequential measurements of two non-commuting projection operators in the AAV method. Different combinations of position and momentum correlations are to be measured where the correlations are of second order in terms of the interaction coefficient. Our method is more efficient as we only need to measure $(d-1)$ arbitrary single observables according to AAV method. We do not discard any post-measurements data and thus reduces the number of experimental runs (see Appendix D).

Weak measurement methods have several key advantages for state reconstruction of a quantum system over the standard schemes [4]. For example, the state disturbance is minimum and thus it is possible for characterization of the state of a system during a physical process in an experiment. Unlike standard schemes, global reconstruction is not required by our methods as states can be determined locally i.e., each matrix element can directly be accessed. Standard scheme typically requires $\mathcal{O}(d^2)$ measurements, while our method require $\mathcal{O}(d-1)$ measurements for mixed state reconstruction.

B. State reconstruction of a bipartite system

1. Pure state

In the above, we introduced reconstruction of an unknown pure state of a given system using single observable (or projection) weak values. For the reconstruction of a bipartite state, one needs to measure product weak values namely the weak values of the tensor product observables. In standard scheme (VMS), measurement of product observables can not be realized directly as it requires the interaction Hamiltonian of the two distant subsystems to be of the form $H \propto (A \otimes B)$

which implies an instantaneous interaction between the two distant subsystems (a relativistic constraint). In this section, we use our version of product weak values (9) in a bipartite system to reconstruct a pure state following the same method of Eq. (29) as we saw for the pure state case in a single quantum system.

The pure state of a bipartite system can be written in computational basis as

$$|\psi_{AB}\rangle = \sum_{ij} \alpha_{ij} |i_A\rangle \otimes |j_B\rangle, \quad (32)$$

where $\alpha_{ij} = \langle ij|\psi_{AB}\rangle$ and $|ij\rangle = |i_A\rangle \otimes |j_B\rangle$. The product weak value of the observable $C_A \otimes C_B$ in a bipartite system is given by

$$\langle (C_A \otimes C_B)_w \rangle_{\psi_{AB}}^{00} = \frac{\langle 00|(C_A \otimes C_B)|\psi_{AB}\rangle}{\langle 00|\psi_{AB}\rangle}, \quad (33)$$

where $|\psi_{AB}\rangle$ is the bipartite pre selected state which is to be reconstructed. $|00\rangle = |0_A\rangle \otimes |0_B\rangle$ is the post-selected state. Now inserting identity operator $I = \sum_{ij} |ij\rangle \langle ij|$ of the joint Hilbert space in Eq. (33), we have

$$\langle (C_A \otimes C_B)_w \rangle_{\psi_{AB}}^{00} - [C_A]_{00} [C_B]_{00} = \sum_{i,j \neq (0,0)} [C_A]_{0i} [C_B]_{0j} \frac{\alpha_{ij}}{\alpha_{00}}, \quad (34)$$

where $[C_A]_{0i} [C_B]_{0j} = \langle 00|C_A \otimes C_B|ij\rangle$ and $\alpha_{00} \neq 0$. Again we have to solve a matrix equation using Eq. (34) for a set of product operators to obtain the values $\frac{\alpha_{ij}}{\alpha_{00}}$ (see Appendix E).

We have found using the matrix equation (E1) that we do not require to measure all the product weak values. For example, consider a two-qubit system where

$$\begin{aligned} C_A^{(1)} \otimes C_B^{(1)} &= I^A \otimes \sigma_x^B, & C_A^{(2)} \otimes C_B^{(2)} &= \sigma_x^A \otimes I^B \\ C_A^{(3)} \otimes C_B^{(3)} &= \sigma_x^A \otimes \sigma_x^B, \end{aligned} \quad (35)$$

then the square matrix in (E1) becomes an identity matrix having nonzero determinant and hence all the coefficients can be determined. *This way, the number of measurements of product weak values can be reduced.* In this particular case, we need only one product weak value i.e., $\langle (\sigma_x^A \otimes \sigma_x^B)_w \rangle_{\psi_{AB}}^{00}$ to be measured and according to Eq. (9), it can be calculated using local weak values $\langle (\sigma_x^B)_w \rangle_{\psi_{AB}}^{00}$, $\langle (\sigma_x^B)_w \rangle_{\psi_{AB}}^{10}$ and $\langle (\sigma_x^A)_w \rangle_{\psi_{AB}}^{00}$. So the total number of local weak values is only three to reconstruct two-qubit pure state. Note that, the local weak value $\langle (\sigma_x^B)_w \rangle_{\psi_{AB}}^{10}$ can be calculated by using $\langle (\sigma_x^B)_w \rangle_{\psi_{AB}}^{00}$ with the completeness relation for $|0\rangle$ and $|1\rangle$.

To compare with the method of Pan *et al.* [29], our method is experimentally simple because it does not depend on the nature of the pointer's state (entangled or product) and locally measurable (using local weak values only) while in their method, the use of entangled pointer's states are necessary (which might not be an easy task to perform) and local modular values as well as modular values of the sum of the local

operators are required. For certain cases, some of the probability amplitudes with the entangled pointer's states are considered to be sufficiently small. Complicated situations may arise for higher dimensions and multi-partite systems because of the entangled pointer states. Our method can be generalized both in higher dimensions and multi-partite systems with local weak value measurements only. In the method of Pan *et al.* [29], The number of product weak values is $(m-1)(n-1)$ and each product weak value consists one modular value of the sum of the two local projectors as well as two local projector modular values. Here 'm' and 'n' are the number of dimensions of the subsystems A and B, respectively. *In our method, there are $(d-1)$ (where $d = mn$) numbers of product weak values and each product weak value can be extracted with only two numbers of AAV type weak measurements.* But as we have seen for the case of two-qubit system (35), we do not need to calculate $(d-1)$ numbers of product weak values all the time. For example, effectively we need only two local weak values to reconstruct the pure state of the two-qubit system. Note that, in Ref. [29] for two-qubit system, the total number of measurements is three in which one is the modular value of the sum of two local projectors and two local projector modular values. So, in most of the cases, it is possible to reduce the number of product weak values considerably in our method of state reconstruction.

2. Mixed state

To reconstruct a mixed state of a bipartite system, we will use the method of Eq. (31). Now, let the pre-selection of the system be ρ_{AB} which is unknown and post-selection be any computational basis state $|kl\rangle$. Then the product weak value of the operator $C_A \otimes C_B$ is given by

$$\langle (C_A \otimes C_B)_w \rangle_{\rho_{AB}}^{kl} = \frac{\langle kl | (C_A \otimes C_B) \rho_{AB} | kl \rangle}{\langle kl | \rho_{AB} | kl \rangle}. \quad (36)$$

Now inserting the identity operator $I = \sum_{i,j} |ij\rangle \langle ij|$ in Eq. (36), we have

$$p(\rho_{AB}, kl) \langle (C_A \otimes C_B)_w \rangle_{\rho_{AB}}^{kl} = \sum_{i,j} [C_A]_{ki} [C_B]_{lj} [\rho_{AB}]_{ij,kl}, \quad (37)$$

where $[C_A]_{ki} [C_B]_{lj} = \langle kl | C_A \otimes C_B | ij \rangle$, $[\rho_{AB}]_{ij,kl} = \langle ij | \rho_{AB} | kl \rangle$ and $p(\rho_{AB}, kl) = \langle kl | \rho_{AB} | kl \rangle$ is the probability of the successful post-selection $|kl\rangle$. So to obtain the kl -th column of the density matrix ρ_{AB} , we have to form a matrix equation using Eq. (37) for a set of product operators (see Appendix F).

Clearly, mixed state reconstruction is more resource intensive than the pure state case in a bipartite system. The number of product weak values to be calculated here is $(d-1)$ and each product weak value can be extracted with only three numbers of AAV type weak measurements (see II B). Here $d = mn$, where 'm' and 'n' are dimensions of the subsystems A and B, respectively. *We will get advantage of using matrix equation (F1) where for some cases we do not require to calculate*

all the product weak values as we have seen for the case of bipartite pure state reconstruction.

It is important to note that, our method of calculating product weak values for pure and mixed states in a bipartite system can also be applied for projection operators and hence one can reconstruct pure state using the state reconstruction method of Ref. [29]. For mixed state reconstruction, one should look to the Ref. [35] by considering bipartite system conditions.

Full knowledge of the state of a quantum system is always crucial to understand a system better and for controlling quantum technologies. In particular, the measurement of bipartite (multi-partite) states are useful for information transfer, cryptography protocols, etc. They are also used to study nonlocality, quantum discord, entanglement entropy, etc. We have shown the application of product and higher moment weak values as quantum state reconstruction of a single and bipartite systems only. The calculations of product weak values of a bipartite system are even more fascinating because of their local realizations. We can have applications of product weak values to extract informations about multi-partite systems for future technologies. Product weak values with local realization can find it's applications in quantum steering, to perform some nonlocal tasks, etc.

IV. ROBUSTNESS

In AAV method, the coupling between the system and the pointer is extremely small and hence the state collapse is avoided. During the process, any resolution is insufficient to distinguish the different eigenvalues of the observable. Nevertheless, by performing the experiment many times on identically prepared systems, it is possible to reduce the uncertainty in the mean pointer displacement to any arbitrary precision.

There are other type of errors which are inevitable due to the inappropriate choices of system observables and unsharp post-selections. Here, we show that our methods of "extraction of product and higher moment weak values" are robust against them.

(i) *Error in choice of observable:* In experiment, let's say, we want to measure a spin-1/2 observable according to the AAV method but due to some technical difficulties, we are unable to measure the actual spin-1/2 observable (slightly changed θ and ϕ , where θ and ϕ define a point on the bloch sphere). Now let 'A' be the correct observable while 'A^e' be the erroneous one such that $|A - A^e| \leq \delta$, where $|X| = \text{Tr} \sqrt{X^\dagger X}$ is the trace norm of a square matrix X. So the error occurring in the weak value is given by

$$\begin{aligned} \Delta(\rho, A, \phi) &= \left| \langle A_w \rangle_\rho^\phi - \langle A_w^e \rangle_\rho^\phi \right| = \frac{|\langle \phi | (A - A^e) \rho | \phi \rangle|}{\langle \phi | \rho | \phi \rangle} \\ &\leq \frac{|\langle \phi | (A - A^e) \rho | \phi \rangle|}{m}, \end{aligned} \quad (38)$$

where m is the minimum of the probabilities for all the possible choices of rank-one post-selections with a given pre-selection. Now consider the spectral decomposition $A - A^e = \sum_i \lambda_i |i\rangle \langle i|$ where $\{|i\rangle\}$ is the complete set of orthogonal ba-

sis. Then

$$\begin{aligned} \Delta(\rho, A, \phi) &\leq \frac{1}{m} \left| \sum \lambda_i \langle \phi|i \rangle \langle i|\rho|\phi \rangle \right| \\ &\leq \frac{1}{m} \sum_i |\lambda_i| |\langle \phi|i \rangle \langle i|\rho|\phi \rangle|. \end{aligned} \quad (39)$$

Note that $|A - A^e| = \sum_i |\lambda_i|$ and since $0 \leq \rho \leq \mathbb{1}$, it can easily be shown that $|\langle \phi|i \rangle \langle i|\rho|\phi \rangle| \leq 1$. Hence

$$\Delta(\rho, A, \phi) \leq \frac{1}{m} \sum_i |\lambda_i| = \frac{|A - A^e|}{m} \leq \frac{\delta}{m}. \quad (40)$$

(ii) *Noisy post-selection:* Now we consider another type of error which is common in experiments is due to the unsharp post-selections. Let us assume that the unsharp post-selection is a mixture of the true post-selection $|\phi\rangle$ with probability $(1 - \epsilon)$ and noise state σ with probability ϵ

$$\Phi^\epsilon = (1 - \epsilon) |\phi\rangle \langle \phi| + \epsilon \sigma, \quad (41)$$

where ϵ is a sufficiently small positive quantity. Then the difference between the perturbed and true weak values is

$$\langle A_w \rangle_\rho^{\Phi^\epsilon} - \langle A_w \rangle_\rho^\phi \approx \epsilon \left[\frac{\text{Tr}(\sigma A \rho) - \langle A_w \rangle_\rho^\phi \text{Tr}(\sigma \rho)}{\langle \phi|\rho|\phi \rangle} \right]. \quad (42)$$

So in both the cases (Eqs. (40) and (42)), the weak values are robust. Now it is not hard to realize that product and higher moment weak values are also robust. The only thing we need to do is to replace the observable A by A^2 for a single system and $C_A \otimes C_B$ for a bipartite system in Eqs. (40) and (42). Hence the weak values which we have used to reconstruct the state of a single and bipartite systems are also robust.

V. CONCLUSION

We have derived the methods of extracting higher moment weak values and product weak values using Vaidman's formula. Such higher moment weak values are calculated using only the weak values of that observable with pairwise orthogonal post-selections. Two dimensional Hilbert space becomes

the simplest case for extracting the higher moment weak values. Our methods turn out to be simple from experimental perspective as we don't need to measure the N pointer's correlations as required in the previous works. Previously, it was thought that with Gaussian pointers' states, it is not possible to obtain the higher moment weak values but we have shown that instead of looking for different pointer states (e.g., OAM states) to obtain the higher moment weak values, we can use Vaidman's formula. To extract the product weak values in a bipartite system, we have again used the Vaidman's formula in one of the subsystems. The product weak values can be calculated using only local weak values. The key factor for such local realization is that the action of the local operator on the local post-selected state is equivalent to the superposition of that post-selected state and a unique orthogonal state to that given post-selected state. Our method can be used to verify Hardy's Paradox, to confirm the existence of quantum cheshire cats, to perform EPR-Bohm experiment, to realize non-locality via post-selections, etc.

As an application, we have shown how to reconstruct quantum states of a single and bipartite systems separately. We have used higher moment weak values to reconstruct an unknown pure state of a single system. The number of measurements are nearly half of the measurements required in previous works. Mixed state reconstruction has been shown using arbitrary observables rather than projection operators. We have used product weak values for reconstruction of pure and mixed states in a bipartite system. Such reconstructions become simply feasible in experiment using only the measurements of local weak values. In the previous works, projection measurement operators were the central for direct quantum state tomography. But we have generalized it to any arbitrary observables for both single and bipartite systems. Comparisons between the previous works and our work have been considered from various perspective (e.g., number of measurements according to the AAV method and experimental feasibility). Further, we have shown that our methods are robust against the errors which are inevitable due to the inappropriate choices of system observables and unsharp post-selections. Our method can easily be extended to multi-partite systems.

Acknowledgment: Sahil is thankful to the QIC group at HRI, Allahabad, for making an arrangement for visiting the group during which, part of the work was done. SM acknowledges financial support from the Visiting Postdoctoral Programme of IMSc, Chennai.

[1] Y. Aharonov, *et al.*, Phys. Lett. A 301 (2001) 130
[2] J. S. Lundeen and A. M. Steinberg, Phys. Rev. Lett. 102, 020404 (2009); K. Yokota, T. Yamamoto, M. Koashi, and N. Imoto, New J. Phys. 11, 033011 (2009).
[3] Y. Aharonov, D. Rohrlich, Quantum Paradoxes: Quantum Theory for the Perplexed, Wiley-VCH; Y. Aharonov *et al.*, Phys. Rev. A 87 (2012) 014105.
[4] Jeff S. Lundeen and Charles Bamber, Phys. Rev. Lett. 108, 070402 (2012); G. S. Thekkadath, et

al., Phys. Rev. Lett. 117, 120401 (2016).
[5] Wei-Wei Pan *et al.*, Phys. Rev. Lett. 123, 150402(2019); Yusuf Turek 2020 J. Phys. Commun. 4 075007.
[6] Yakir Aharonov *et al.*, 2013 New J. Phys. 15 113015.
[7] Tobias Denkmayr *et al.*, 10.1038/ncomms5492.
[8] Debmalya Das and Arun Kumar Pati 2020 New J. Phys. 22 063032.
[9] B. L. Higgins, *et al.*, Phys. Rev. A 91, 012113 (2015).
[10] S. Marcovitch and B. Reznik, arXiv:1005.3236.

- [11] Y. Aharonov *et al.*, Annals of Physics 355(2015)258–268
- [12] Y. Aharonov, and E. Cohen, in Quantum Nonlocality and Reality, arXiv:1504.03797
- [13] Y. Aharonov, D. Z. Albert, and L. Vaidman, Phys. Rev. Lett. 60, 1351 (1988).
- [14] Y. Aharonov, P. G. Bergmann, and J. L. Lebowitz, Phys. Rev. 134, B1410 (1964).
- [15] J. von Neumann *Mathematical Foundations of Quantum Mechanics*. Princeton Univ. Press, Princeton, 1955
- [16] J.S. Lundeen and K.J. Resch, Physics Letters A 334 (2005) 337–344.
- [17] Richard Jozsa, Phys. Rev. A 76, 044103 (2007).
- [18] Jeff S. Lundeen, et al., Nature 474, 188–191(2011).
- [19] D.R. Solli *et al.*, Phys. Rev. Lett. 92 (2004) 043601.
- [20] L. A. Rozema *et al.*, Phys. Rev. Lett. 109, 100404 (2012).
- [21] Omar S. Magana Loaiza *et al.*, Phys. Rev. Lett. 112, 200401 (2014).
- [22] O. Hosten and P. Kwiat, Science 319, 787 (2008).
- [23] Arun Kumar Pati, Uttam Singh, and Urbasi Sinha Phys. Rev. A 92, 052120 (2015).
- [24] Gaurav Nirala *et al.*, Phys. Rev. A 99, 022111 (2019).
- [25] Arun Kumar Pati *et al.*, Phys. Rev. A 102, 012204 (2020).
- [26] Suchetana Goswami *et al.*, Phys. Rev. A 99, 012327 (2019).
- [27] K. Resch and A. Steinberg, Phys. Rev. Lett. 92, 130402, 443(2004).
- [28] K.J. Resch, J. Opt. B 6 (2004) 482.
- [29] Wei-Wei Pan *et al.*, Phys. Rev. Lett. 123, 150402 (2019).
- [30] Y. Kedem and L. Vaidman, Phys. Rev. Lett. 105, 230401 (2010).
- [31] Lior Goldenberg and Lev Vaidman, American Journal of Physics 64, 1059 (1996).
- [32] Arun Kumar Pati, Junde Wu, arXiv:1410.5221.
- [33] G. Puentes, N. Hermosa, and J. P. Torres, Phys. Rev. Lett. 109, 040401 (2012).
- [34] Hirokazu Kobayashi, Graciana Puentes and Yutaka Shikano, Phys. Rev. A 86, 053805 (2012).
- [35] Shengjun Wu, Scientific Reports volume 3, 1193 (2013).

VI. APPENDICES

Appendix A

Let A be a Hermitian operator and $|\phi\rangle$ be a quantum state vector in a Hilbert space \mathcal{H} . Then it is always possible to decompose $A|\phi\rangle$ as [31]

$$A|\phi\rangle = \alpha|\phi\rangle + \beta|\phi_A^\perp\rangle, \quad (\text{A1})$$

where α and β can be any complex numbers. $|\phi_A^\perp\rangle$ is an orthonormal state to $|\phi\rangle$. The state $|\phi_A^\perp\rangle$ is unique and can be determined using Gram-Schmidt orthonormalization process (GSOP) as given below.

Let $A|\phi\rangle$ and $|\phi\rangle$ are two non-orthogonal state vectors in the same Hilbert space. Using GSOP we find the unnormalized $|\phi_A^\perp\rangle_{un}$

$$|\phi_A^\perp\rangle_{un} = A|\phi\rangle - \frac{\langle\phi|A|\phi\rangle}{\langle\phi|\phi\rangle}|\phi\rangle = (A - \langle A \rangle_\phi)|\phi\rangle. \quad (\text{A2})$$

Here $|\phi\rangle$ is normalized. So the normalized orthogonal state

vector to $|\phi\rangle$ is

$$|\phi_A^\perp\rangle = \frac{|\phi_A^\perp\rangle_{un}}{\sqrt{\langle\phi_A^\perp|\phi_A^\perp\rangle_{un}}}. \quad (\text{A3})$$

From Eqs. (A2) and (A3) we find $\alpha = \langle\phi|A|\phi\rangle = \langle A \rangle_\phi$ and $\beta = \langle\phi_A^\perp|A|\phi\rangle = \langle\Delta A\rangle_\phi$. Where $\langle A \rangle_\phi$ and $\langle\Delta A\rangle_\phi$ are the average and standard deviation of the observable A in the state $|\phi\rangle$, respectively. So the Eq. (A1) becomes

$$A|\phi\rangle = \langle A \rangle_\phi|\phi\rangle + \langle\Delta A\rangle_\phi|\phi_A^\perp\rangle. \quad (\text{A4})$$

Eq. (A4) is sometimes referred as Vaidman's formula.

Appendix B

For a given mixed pre selected state ρ and post-selected state $|\phi\rangle$, the second moment weak value of the observable A is given by

$$\begin{aligned} \langle(A^2)_w\rangle_\rho^\phi &= \left(\frac{\langle\phi|\rho A^2|\phi\rangle}{\langle\phi|\rho|\phi\rangle}\right)^* \\ &= \langle A \rangle_\phi \frac{\langle\phi|A\rho|\phi\rangle}{\langle\phi|\rho|\phi\rangle} + \langle\Delta A\rangle_\phi \frac{\langle\phi_A^\perp|A\rho|\phi\rangle}{\langle\phi|\rho|\phi\rangle}. \end{aligned} \quad (\text{B1})$$

Now $\langle\phi_A^\perp|A\rho|\phi\rangle = \text{Tr}(|\phi\rangle\langle\phi_A^\perp|A\rho)$ can be calculated as

$$\langle\phi_A^\perp|A\rho|\phi\rangle + \langle\phi|A\rho|\phi_A^\perp\rangle = \text{Tr}[(|\phi\rangle\langle\phi_A^\perp| + |\phi_A^\perp\rangle\langle\phi|)A\rho], \quad (\text{B2})$$

where $(|\phi\rangle\langle\phi_A^\perp| + |\phi_A^\perp\rangle\langle\phi|)$ is a normal operator satisfying $XX^\dagger = X^\dagger X$ (where X is any operator) and hence can be written in spectral decomposition

$$|\phi\rangle\langle\phi_A^\perp| + |\phi_A^\perp\rangle\langle\phi| = \sum_i^d \lambda_i |i\rangle\langle i|, \quad (\text{B3})$$

where $|i\rangle$ is the eigenvector with eigenvalue λ_i and 'd' is the dimension of the Hilbert space. Using Eq. (B3) in (B2), we have

$$\begin{aligned} \langle\phi_A^\perp|A\rho|\phi\rangle + \langle\phi|A\rho|\phi_A^\perp\rangle &= \sum_i^d \lambda_i \langle i|A\rho|i\rangle \\ &= \sum_i^d \lambda_i \langle A_w \rangle_\rho^i p(\rho, i), \end{aligned} \quad (\text{B4})$$

where $p(\rho, i) = \langle i|\rho|i\rangle$ is the probability of obtaining the eigenvector $|i\rangle$ and $\langle A_w \rangle_\rho^i$ is the weak value of the observable A for the given pre and post-selections ρ and $|i\rangle$, respectively. Now similarly,

$$\langle\phi_A^\perp|A\rho|\phi\rangle - \langle\phi|A\rho|\phi_A^\perp\rangle = \sum_i^d \lambda_i' \langle A_w \rangle_\rho^{i'} p(\rho, i'), \quad (\text{B5})$$

where we have used the fact that $|\phi\rangle\langle\phi_A^\perp| - |\phi_A^\perp\rangle\langle\phi|$ is also a normal operator with the spectral decomposition

$$|\phi\rangle\langle\phi_A^\perp| - |\phi_A^\perp\rangle\langle\phi| = \sum_i^d \lambda'_i |i'\rangle\langle i'|. \quad (\text{B6})$$

Now adding two equations (B4) and (B5), we have

$$\langle\phi_A^\perp|A\rho|\phi\rangle = \frac{1}{2} \sum_i^d \left(\lambda_i \langle A_w \rangle_\rho^i p(\rho, i) + \lambda'_i \langle A_w \rangle_\rho^{i'} p(\rho, i') \right). \quad (\text{B7})$$

So the final expression of Eq. (B1) is

$$\langle(A^2)_w\rangle_\rho^\phi = \langle A \rangle_\phi \langle A_w \rangle_\rho^\phi + \frac{\langle\Delta A\rangle_\phi}{2p(\rho, \phi)} \sum_i^d \left(\lambda_i \langle A_w \rangle_\rho^i p(\rho, i) + \lambda'_i \langle A_w \rangle_\rho^{i'} p(\rho, i') \right), \quad (\text{B8})$$

where $p(\rho, \phi) = \langle\phi|\rho|\phi\rangle$. The n -th moment weak value is

$$\langle(A^n)_w\rangle_\rho^\phi = \langle A \rangle_\phi \langle A_w^{n-1} \rangle_\rho^\phi + \frac{\langle\Delta A\rangle_\phi}{2p(\rho, \phi)} \sum_i^d \left(\lambda_i \langle A_w^{n-1} \rangle_\rho^i p(\rho, i) + \lambda'_i \langle A_w^{n-1} \rangle_\rho^{i'} p(\rho, i') \right). \quad (\text{B9})$$

Note that, $\{\langle A_w \rangle_\rho^i\}_{i=1}^d$ can be measured in the same experimental setup according to the AAV method as the post-selections $\{|i\rangle\}$ form a complete set of orthogonal basis. Similarly, $\{\langle A_w \rangle_\rho^{i'}\}_{i'=1}^d$ can also be measured within the same experiment with the complete set of post-selections $\{|i'\rangle\}$ according to the AAV method. So, to extract the second moment weak value, one needs only three AAV type measurements and these are $\langle A_w \rangle_\rho^\phi$, $\{\langle A_w \rangle_\rho^i\}_{i=1}^d$ and $\{\langle A_w \rangle_\rho^{i'}\}_{i'=1}^d$. Number of measurements will increase for mixed state case than when the system is prepared in the pure state.

Appendix C

For the measurement of a pure state of a single system, we have followed a method slightly different from Lundeen *et al.* [18]. Namely we consider higher moment weak values as discussed in the main text. Here we show how to complete the process. A is an observable having spectral decomposition (24), is measured according to AAV method. Now consider Eq. (25) and (26) with $n=1, \dots, d-1$, where d is the dimension of the Hilbert space. So, for different 'n', we obtain a matrix equation

$$\begin{pmatrix} 1 \\ \langle A_w \rangle_\psi^b \\ \vdots \\ \langle A_w^{d-1} \rangle_\psi^b \end{pmatrix} = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ a_0 & a_1 & & a_{d-1} \\ \vdots & & \ddots & \\ a_0^{d-1} & a_1^{d-1} & \cdots & a_{d-1}^{d-1} \end{pmatrix} \begin{pmatrix} \langle(\Pi_0)_w\rangle_\psi^b \\ \langle(\Pi_1)_w\rangle_\psi^b \\ \vdots \\ \langle(\Pi_{d-1})_w\rangle_\psi^b \end{pmatrix} \quad (\text{C1})$$

The solution of the above equation exists if the determinant of the square matrix is non-zero. The weak values of the projection operators are then extracted from (C1) using the higher moment weak values of A .

Alternative:— To obtain the values of α_i/α_0 , we have to solve the matrix equation using the Eq. (29) for a set of arbitrary

observables $C^{(n)}$, $n = 1, \dots, (d-1)$. So

$$\begin{pmatrix} \langle C_w^{(1)} \rangle_\psi^0 - C_{00}^{(1)} \\ \langle C_w^{(2)} \rangle_\psi^0 - C_{00}^{(2)} \\ \vdots \\ \langle C_w^{(d-1)} \rangle_\psi^0 - C_{00}^{(d-1)} \end{pmatrix} = \begin{pmatrix} C_{01}^{(1)} & C_{02}^{(1)} & \cdots & C_{0(d-1)}^{(1)} \\ C_{01}^{(2)} & C_{02}^{(2)} & & C_{0(d-1)}^{(2)} \\ \vdots & & \ddots & \\ C_{01}^{(d-1)} & C_{02}^{(d-1)} & \cdots & C_{0(d-1)}^{(d-1)} \end{pmatrix} \begin{pmatrix} \frac{\alpha_1}{\alpha_0} \\ \frac{\alpha_2}{\alpha_0} \\ \vdots \\ \frac{\alpha_{d-1}}{\alpha_0} \end{pmatrix} \quad (\text{C2})$$

The solution exists if the determinant of the square matrix above is non-zero. It may be noted here that there will always exist at least one set of observables $\{C^{(n)}\}_{n=1}^{d-1}$ for which the determinant of the coefficient matrix on the right hand side of (C2) is non-zero. Now we obtain the values of α_i/α_0 and then from Eq. (21), we have

$$|\psi\rangle = \alpha_0 \left(|0\rangle + \sum_{i=1}^{d-1} \frac{\alpha_i}{\alpha_0} |i\rangle \right). \quad (\text{C3})$$

The value of α_0 can be obtained from the normalization condition. If the determinant of the square matrix (C2) is zero with the post-selection $|0\rangle$, then one should look for the other post-selections such as $|1\rangle, |2\rangle$ and so on and hence solve the corresponding matrix equations.

Appendix D

The j -th column of the density matrix with post-selection $|j\rangle$ can be obtained using Eq. (31) for a set of arbitrary ob-

servables $C^{(n)}$, $n = 1, \dots, d$ as

$$p(\rho, j) \begin{pmatrix} \langle C_w^{(1)} \rangle_\rho^j \\ \langle C_w^{(2)} \rangle_\rho^j \\ \vdots \\ \langle C_w^{(d)} \rangle_\rho^j \end{pmatrix} = \begin{pmatrix} C_{j0}^{(1)} & C_{j1}^{(1)} & \cdots & C_{j(d-1)}^{(1)} \\ C_{j0}^{(2)} & C_{j1}^{(2)} & & C_{j(d-1)}^{(2)} \\ \vdots & \vdots & \ddots & \vdots \\ C_{j0}^{(d)} & C_{j1}^{(d)} & \cdots & C_{j(d-1)}^{(d)} \end{pmatrix} \begin{pmatrix} \rho_{0j} \\ \rho_{1j} \\ \vdots \\ \rho_{(d-1)j} \end{pmatrix} \quad (\text{D1})$$

With $C^{(1)} = I$, an identity operator. The j' -th column can be calculated measuring the same set of observables with post-selection $|j'\rangle$. Remember that post-selections are complete set of projection operators and hence can be measured simultaneously for each observables $C^{(n)}$. So measuring an observable $C^{(n)}$ according to AAV method, we obtain all the weak values for a complete set of post selections $\{|j\rangle\}$. We do not need to

$$\begin{pmatrix} \langle (C_A^{(1)} \otimes C_B^{(1)})_w \rangle_{\psi_{AB}}^{00} - [C_A^{(1)}]_{00} [C_B^{(1)}]_{00} \\ \langle (C_A^{(2)} \otimes C_B^{(2)})_w \rangle_{\psi_{AB}}^{00} - [C_A^{(2)}]_{00} [C_B^{(2)}]_{00} \\ \vdots \\ \langle (C_A^{(d-1)} \otimes C_B^{(d-1)})_w \rangle_{\psi_{AB}}^{00} - [C_A^{(d-1)}]_{00} [C_B^{(d-1)}]_{00} \end{pmatrix} = \begin{pmatrix} [C_A^{(1)}]_{00} [C_B^{(1)}]_{01} & [C_A^{(1)}]_{01} [C_B^{(1)}]_{00} & \cdots & [C_A^{(1)}]_{0(d-1)} [C_B^{(1)}]_{0(d-1)} \\ [C_A^{(2)}]_{00} [C_B^{(2)}]_{01} & [C_A^{(2)}]_{01} [C_B^{(2)}]_{00} & & [C_A^{(2)}]_{0(d-1)} [C_B^{(2)}]_{0(d-1)} \\ \vdots & \vdots & \ddots & \vdots \\ [C_A^{(d-1)}]_{00} [C_B^{(d-1)}]_{01} & [C_A^{(d-1)}]_{01} [C_B^{(d-1)}]_{00} & \cdots & [C_A^{(d-1)}]_{0(d-1)} [C_B^{(d-1)}]_{0(d-1)} \end{pmatrix} \begin{pmatrix} \frac{\alpha_{01}}{\alpha_{00}} \\ \frac{\alpha_{10}}{\alpha_{00}} \\ \vdots \\ \frac{\alpha_{(d-1)(d-1)}}{\alpha_{00}} \end{pmatrix} \quad (\text{E1})$$

Solution exists if the determinant of the square matrix in (E1) is nonzero and hence we calculate all the unknown coefficients α_{ij}/α_{00} . Then Eq. (31) becomes

$$|\psi_{AB}\rangle = \alpha_{00} \left(|00\rangle + \sum_{i,j \neq (0,0)} \frac{\alpha_{ij}}{\alpha_{00}} |ij\rangle \right). \quad (\text{E2})$$

Again, the value of α_{00} can be found using normalization condition. If the determinant of the square matrix becomes zero

perform any measurement for the last column as one can get it using normalization and Hermiticity condition of the density matrix.

Appendix E

In the main text, we discussed about how to develop an equation (34) which contains all the probability amplitudes of a pure state in a bipartite system of dimension $d = mn$, where ‘ m ’ and ‘ n ’ are the dimensions of subsystems A and B, respectively. Now, our task is to obtain all the unknown complex coefficients (probability amplitudes) in a pure state. To do so, we develop a matrix equation using Eq. (34) for a set of arbitrary product observables $C_A^{(n)} \otimes C_B^{(n)}$, $n = 1, \dots, (d-1)$

with the post selection $|00\rangle$ then one should look for the other post-selections such as $|01\rangle$, $|10\rangle$ and so on and hence solve the corresponding matrix equations.

Appendix F

The kl -th column of the density matrix with post-selection $|kl\rangle$ can be obtained using Eq. (37) for a set of arbitrary product observables $C_A^{(n)} \otimes C_B^{(n)}$, $n = 1, \dots, d$

$$p(\rho_{AB}, kl) \begin{pmatrix} \langle (C_A^{(1)} \otimes C_B^{(1)})_w \rangle_{\rho_{AB}}^{kl} \\ \langle (C_A^{(2)} \otimes C_B^{(2)})_w \rangle_{\rho_{AB}}^{kl} \\ \vdots \\ \langle (C_A^{(d)} \otimes C_B^{(d)})_w \rangle_{\rho_{AB}}^{kl} \end{pmatrix} = \begin{pmatrix} [C_A^{(1)}]_{k0} [C_B^{(1)}]_{l0} & [C_A^{(1)}]_{k0} [C_B^{(1)}]_{l1} & \cdots & [C_A^{(1)}]_{k(d-1)} [C_B^{(1)}]_{l(d-1)} \\ [C_A^{(2)}]_{k0} [C_B^{(2)}]_{l0} & [C_A^{(2)}]_{k0} [C_B^{(2)}]_{l1} & \cdots & [C_A^{(2)}]_{k(d-1)} [C_B^{(2)}]_{l(d-1)} \\ \vdots & \vdots & \ddots & \vdots \\ [C_A^{(d)}]_{k0} [C_B^{(d)}]_{l0} & [C_A^{(d)}]_{k0} [C_B^{(d)}]_{l1} & \cdots & [C_A^{(d)}]_{k(d-1)} [C_B^{(d)}]_{l(d-1)} \end{pmatrix} \begin{pmatrix} [\rho_{AB}]_{00,kl} \\ [\rho_{AB}]_{01,kl} \\ \vdots \\ [\rho_{AB}]_{(d-1)(d-1),kl} \end{pmatrix} \quad (\text{F1})$$

With $C_A^{(1)} \otimes C_B^{(1)} = I_A \otimes I_B$, an identity operator in the joint Hilbert space. Solution exists for non zero determinant of the above square matrix. The $k'l'$ -th column can be calculated measuring the same set of observables with post-selection $|k'l'\rangle$ and by solving the corresponding matrix equations. Remember that post-selections are complete set of projection operators and hence can be measured simultaneously

for each product observables $C_A^{(n)} \otimes C_B^{(n)}$. So measuring an observable $C_A^{(n)} \otimes C_B^{(n)}$ according to AAV method, we obtain all the weak values for a complete set of post selections $\{|kl\rangle\}$. One does not need to perform a measurement for the last column as it can be obtained from the normalization and Hermiticity condition of the density matrix.