

ON THE FUNCTIONAL GRAPH OF THE POWER MAP OVER FINITE GROUPS

CLAUDIO QURESHI AND LUCAS REIS

ABSTRACT. In this paper we study the description of the digraph associated with the power map over finite groups. Our main motivation comes from the nice description of such digraphs in the case of cyclic groups. In particular, we derive results on abelian groups, and also on flower groups, which are introduced in this paper. The class of flower groups includes many non abelian groups such as dihedral and generalized quaternion groups, and the projective general linear group of order two over a finite field. In particular, we provide improvements on past works.

1. INTRODUCTION

Given a pair (f, S) of a finite set S and a map $f : S \rightarrow S$, we can associate with it a graph $\mathcal{G}(f/S)$, called the functional graph of f over S . This is the directed graph with vertex set $V = \{s \mid s \in S\}$ and directed edges $\{s \rightarrow f(s) \mid s \in S\}$. The functional graph encodes the dynamics of f over S . For instance, the f -orbit $\{s, f(s), f^{(2)}(s), \dots\}$ of an element $s \in S$ is described by a path in $\mathcal{G}(f/S)$. Moreover, an element $s \in S$ is f -periodic if and only if it belongs to a cycle of $\mathcal{G}(f/S)$. One of the motivation of studying finite dynamical systems is due to their applications such as integer factorization methods in cryptography [12, 22] and pseudo-random number generators [3]. The description of $\mathcal{G}(f/S)$ has been considered for many algebraic structures S and well behaved maps f . In many cases, the functional graphs turns out to have many remarkable properties such as regularity on the indegrees and symmetries on its cycles that allow us to obtain a partial or complete description of their structure. See [5, 9, 10, 11, 13, 14, 15, 18, 19, 21] for a rich source of results regarding these issues and [8] for a survey including some applications of dynamical systems over finite fields.

When $S = G$ is a finite group, it is natural to consider the power map $\varphi_t : G \rightarrow G$ with $\varphi_t(g) = g^t$ (or tg if G is written additively), where t is an integer. When G is cyclic, the graph $\mathcal{G}(\varphi_t/G)$ is completely described in [13]. The latter can be extended to abelian groups, mainly using the results of [16] as we show in Section 2.3. For non abelian groups, only results for special families of groups are known. More specifically, in [1], [2] and [4] the cases where G is a dihedral, a generalized quaternion and some semidirect product of cyclic groups are explored, respectively. However the digraphs are not explicitly determined there. Instead, only partial descriptions of such graph are given like the distribution of indegrees, cycle lengths and number of cycles. Further results of this kind are given in [7] for special classes of finite groups.

2010 *Mathematics Subject Classification.* Primary 20D60, Secondary 05C20.
Key words and phrases. functional graph, finite groups, power map.

In this paper we introduce the class of *flower groups*, which contains the dihedral and generalized quaternion groups, and also some semidirect product of cyclic groups. Our main result is the description of the functional graph structure of the power map over flower groups under some mild conditions (Theorems 3.9 and 3.11).

Our paper is organized as follows. In Section 2 we introduce the basic notation and operations on graphs and review the structure theorem for the functional graph of the power map on cyclic groups. Then, using results from [16], we show how to extend this structure theorem for finite abelian groups. In the last part of this section we provide some general results on the functional graph of the power map on arbitrary finite groups that is further used to prove our main results. In Section 3 we introduce the notion of flower groups, obtain some properties about these groups and also prove our main results. In Section 4 we apply our main results to the explicit description of functional graphs associated to some non abelian groups.

2. PRELIMINARIES

This section provides some background material and minor results. We start by fixing some basic notation and reviewing the structural description of the functional graph of the power map on cyclic groups. Then we show how to extend this structural description for abelian groups. At the end of this section we provide some general results on the functional graph of the power map over arbitrary finite groups.

Along the paper we use the letters G, H to denote finite groups and C to denote cyclic groups. For a positive integer m , C_m denotes the cyclic group of order m . We use multiplicative notation for the operation of these groups unless otherwise specified. The ring of integers modulo d is denoted by \mathbb{Z}_d and their multiplicative group of units is denoted by \mathbb{Z}_d^* . The Euler totient function is denoted by $\varphi(d)$ and the (multiplicative) order of $t \in \mathbb{Z}_d^*$ is denoted by $o_d(t)$. For a group G , the centralizer of $S \subseteq G$ is denoted by $C_G(S)$ and the center of G is denoted by $Z(G) = C_G(G)$.

2.1. Basics on functional graphs. We use the same terminology as in [13] and [16]. It is a well known fact that the connected components of functional graphs are composed of a cycle and each vertex of this cycle is the root of a tree (the direction of the edges is from leaves to the root). Several interesting classes of functions over finite fields or finite structures studied in the literature present certain regularity that is reflected in some symmetry conditions on their functional graphs. We denote by $\text{Cyc}(m, T)$ a directed graph composed by an m -cycle (i.e. a cycle of length m) where every node in this cycle is the root of a tree isomorphic to T . When every connected component of a functional graph $\mathcal{G}(f/S)$ is of this form we say that this functional graph is *regular*, in this case there are integers m_1, \dots, m_k and rooted trees T_1, \dots, T_k such that $\mathcal{G}(f/S) = \bigoplus_{i=1}^k \text{Cyc}(m_i, T_i)$, where the circled plus symbol denotes a disjoint union of graphs. When the rooted tree T has a unique vertex we write $T = \bullet$. The m -cycle $\text{Cyc}(m, \bullet)$ is denoted by $\text{Cyc}(m)$ and the rooted tree T with a loop in the root $\text{Cyc}(1, T)$ is denoted by $\{T\}$. The notation $G = k \times H$ means that $G = \bigoplus_{i=1}^k H_i$ with each H_i isomorphic to the graph H . A forest is a disjoint union of rooted trees. Given a forest $G = \bigoplus_{i=1}^k T_i$, we denote by $\langle G \rangle$ the rooted tree with k children where these children are roots of trees isomorphic to T_1, \dots, T_k . We consider the empty graph \emptyset with the property that $\langle \emptyset \rangle = \bullet$.

Given rooted trees $T_1 = \langle G_1 \rangle, \dots, T_k = \langle G_k \rangle$ where each G_i is a forest we define $\sum_{i=1}^k T_i := \langle \bigoplus_{i=1}^k G_i \rangle$ (i.e. the sum of rooted trees is a new rooted tree which is obtained by identifying all the roots). For a tree $T = \langle G \rangle$ and $k \in \mathbb{Z}^+$ we denote $k \cdot T = \langle k \times G \rangle$ (note that $k \cdot T \neq k \times T$ if $k > 1$).

The vertices belonging to the cycles of the functional graph $\mathcal{G}(f/S)$ correspond to the periodic points of f . Given a point $x_0 \in S$, the least natural number $\delta = \delta(x_0)$ such that $f^{(\delta)}(x_0)$ is a periodic point is called the *preperiod* of x_0 (the periodic points corresponds to points x_0 with $\delta(x_0) = 0$). Regarding the functional graph $\mathcal{G}(f/S)$, the number $\delta(x_0)$ equals the depth of x_0 in the rooted tree which it belongs to (i.e. the distance of x_0 to the root).

There is a simple way to associate with each non-increasing finite sequence of positive integers $\mathbf{v} = (\nu_1, \nu_2, \dots, \nu_d)$ (i.e. $\nu_1 \geq \nu_2 \geq \dots, \nu_d \geq 1$) a rooted tree $T_{\mathbf{v}}$ defined recursively as follows:

$$(1) \quad \begin{cases} T_{\mathbf{v}}^0 = \bullet, \\ G_{\mathbf{v}}^k = \nu_k \times T_{\mathbf{v}}^{k-1} \oplus \bigoplus_{i=1}^{k-1} (\nu_i - \nu_{i+1}) \times T_{\mathbf{v}}^{i-1} \text{ and } T_{\mathbf{v}}^k = \langle G_{\mathbf{v}}^k \rangle \text{ for } 1 \leq k < d, \\ G_{\mathbf{v}} = (\nu_d - 1) \times T_{\mathbf{v}}^{d-1} \oplus \bigoplus_{i=1}^{d-1} (\nu_i - \nu_{i+1}) \times T_{\mathbf{v}}^{i-1} \text{ and } T_{\mathbf{v}} = \langle G_{\mathbf{v}} \rangle. \end{cases}$$

Trees associated with non-increasing sequences as above are called *elementary trees*, see [16] for more details on elementary trees. Figure 1 shows the inductive process to construct $T_{\mathbf{v}}$ for a 4-term sequence.




FIGURE 1. This figure (taken from [13]) illustrates the inductive definition of $T_{\mathbf{v}}$ for a 4-term sequence $\mathbf{v} = (\nu_1, \nu_2, \nu_3, \nu_4)$. A node labelled by a rooted tree T indicates that it is the root of a tree isomorphic to T .

Elementary trees appear in a wide range of context when we describe the dynamics of different classes of maps, see for example [5, 10, 13, 14, 15, 16, 20]. They will play a key role also in our description of the functional graph associated with the power map over flower groups. An important special case of elementary trees $T_{\mathbf{v}}$ is when the sequence \mathbf{v} is a multiplicative chain (i.e. when each term of the sequence divides the previous term), these are the trees that appear in the functional graph of power maps on abelian groups.

A rooted tree T is *homogeneous* if it can be written as $T = \langle \bigoplus_{i=0}^{d-1} n_i \times T_i \rangle$ where T_i is a rooted tree with depth i for $0 \leq i < d$ (i.e. all its subtrees with the same depth are isomorphic). In particular elementary trees are homogeneous since the subtrees $T_{\mathbf{v}}^i$ of $T_{\mathbf{v}}$ associated with the d -term sequence \mathbf{v} has depth i , for $0 \leq i < d$.

The following operation will be useful to describe trees in the functional graph associated with the power map on some finite non abelian groups.

Definition 2.1. Let $T = \langle \bigoplus_{i=0}^{d-1} n_i \times T_i \rangle$ be an homogeneous rooted tree where each T_i has depth i for $0 \leq i < d$ and S be a rooted tree. For $0 \leq j < d$, the j -sum of T and S is given by $T +_j S = \langle \bigoplus_{i \neq j} n_i \times T_i \oplus (n_j - 1) \times T_j \oplus (T_j + S) \rangle$.

We note that $T +_j S$ is obtained by replacing one of the depth- j -subtree T_j of T with $T_j + S$. The resulting rooted tree will be not an homogeneous tree in general.

2.2. The power map on finite cyclic groups. Let \mathcal{C}_n be the (multiplicative) cyclic group of order n and $\varphi_t : \mathcal{C}_n \rightarrow \mathcal{C}_n$ be the power map $x \rightarrow x^t$. The trees attached to the cyclic points in $\mathcal{G}(\varphi_t/\mathcal{C}_n)$ are described by the iterated gcd of n relative to t (also called ν -series in [13]), denoted by $\text{gcd}_t(n)$, and defined as follows¹: $\text{gcd}_t(n) = (1)$ if $\text{gcd}(t, n) = 1$, otherwise $\text{gcd}_t(n) = (\nu_1, \dots, \nu_D)$ where

$$\nu_1 = \text{gcd}(t, n), \quad \nu_{i+1} = \text{gcd}\left(t, \frac{n}{\nu_1 \cdots \nu_i}\right), \text{ for } i \geq 1,$$

and D is the least positive integer such that $\nu_{D+1} = 1$. It is easy to see that if $\nu := \nu_1 \cdots \nu_D$ then $\nu \mid n$, $\text{gcd}_t(n) = \text{gcd}_t(\nu)$ and $\omega := n/\nu$ is the greatest divisor of n that is relatively prime with t (see [13] for more details). The following proposition gives an explicit description for the functional graph $\mathcal{G}(\varphi_t/\mathcal{C}_n)$.

Proposition 2.2 ([15], Proposition 2.1). *Let $n = \nu\omega$, where ω is the greatest divisor of n that is relatively prime with t . Let $\text{gcd}_t(\nu) = (\nu_1, \nu_2, \dots, \nu_D)$ be the iterated gcd of ν relative to t and $T_{\text{gcd}_t(\nu)}$ be the elementary tree associated with this sequence. Then \mathcal{C}_n has exactly ω elements that are φ_t -periodic and the following isomorphism formula holds:*

$$(2) \quad \mathcal{G}(\varphi_t/\mathcal{C}_n) = \bigoplus_{d \mid \omega} \left(\frac{\varphi(d)}{\text{o}_d(t)} \times \text{Cyc}(\text{o}_d(t), T_{\text{gcd}_t(\nu)}) \right).$$

Moreover, the tree $T_{\text{gcd}_t(\nu)}$ has ν vertices and depth D .

2.3. The power map on finite abelian groups. Next we show how formula (2) can be extended to general abelian groups. Note that if \mathbf{u} and \mathbf{v} are non-increasing finite sequences which differ only possibly in the last terms that are all equal to 1, then the corresponding elementary trees are equal (i.e. $T_{\mathbf{u}} = T_{\mathbf{v}}$). In these case we say that these sequences are *equivalent*. The product $\mathbf{u}\mathbf{v}$ of two non-increasing sequences is the coordinatewise product (substituting the smallest sequence by other equivalent in such a way that both sequences have the same length). For what follows, \otimes denotes the *tensor product* of digraphs.

Lemma 2.3 ([16], Lemma 3.4). *For any maps of finite sets $f : X \rightarrow X$ and $g : Y \rightarrow Y$ we have the following graph isomorphism:*

$$\mathcal{G}(f \times g/X \times Y) \cong \mathcal{G}(f/X) \otimes \mathcal{G}(g/Y),$$

where $f \times g : X \times Y \rightarrow X \times Y$ is the map $f \times g(x, y) = (f(x), g(y))$.

¹In [13] the iterated gcd of n relative to t is called ν -series generated by n and t , denoted by $n(t)$, and defined only in the case when each prime divisor of n divides t . In order to avoid possible confusion we decide to use the alternative notation $\text{gcd}_t(n)$.

Lemma 2.4 ([16], Lemma 3.5). *Let $r \in \mathbb{Z}^+$ and T be a rooted tree. The following isomorphism holds:*

$$\text{Cyc}(r, T) \cong \text{Cyc}(r) \otimes \{T\}.$$

Lemma 2.5 ([16], Prop. 2.10). *For any non-decreasing sequences \mathbf{u}, \mathbf{v} we have $\{T_{\mathbf{u}}\} \otimes \{T_{\mathbf{v}}\} = \{T_{\mathbf{uv}}\}$.*

Lemma 2.6. *Let r_1, \dots, r_k be positive integers. We have*

$$\bigotimes_{i=1}^k \text{Cyc}(r_i) = \frac{r_1 r_2 \cdots r_k}{\text{lcm}(r_1, r_2, \dots, r_k)} \times \text{Cyc}(\text{lcm}(r_1, r_2, \dots, r_k)).$$

Proof. Consider the maps $s_i : \mathbb{Z}_{r_i} \rightarrow \mathbb{Z}_{r_i}$, $s_i(x) = x + 1$. Then $\text{Cyc}(r_i) = \mathcal{G}(s_i/\mathbb{Z}_{r_i})$ and $\bigotimes_{i=1}^k \text{Cyc}(r_i) = \mathcal{G}(s_1 \times \cdots \times s_k, \mathbb{Z}_{r_1} \times \cdots \times \mathbb{Z}_{r_k})$ which is a union of disjoint cycles. Each one of these cycles is in correspondence with the cosets of $H := \langle (1, 1, \dots, 1) \rangle$ in $\mathbb{Z}_{r_1} \times \cdots \times \mathbb{Z}_{r_k}$ and each one of these cosets has $|H| = \text{lcm}(r_1, r_2, \dots, r_k)$ elements. \square

Now consider any (multiplicative) abelian group G and the map $\varphi_t : G \rightarrow G$, $\varphi_t(g) := g^t$. By the fundamental theorem of finite abelian groups, there is an isomorphism $\eta : G \rightarrow \mathcal{C}_{r_1} \times \cdots \times \mathcal{C}_{r_k}$. Since $\eta \circ \varphi_t = \varphi_t \circ \eta$ we have that η induces an isomorphism between the functional graph of φ_t over G and over the direct product of cyclic groups, so we can assume $G = \mathcal{C}_{r_1} \times \cdots \times \mathcal{C}_{r_k}$. If \mathbf{d} and $\boldsymbol{\omega}$ are k -term sequences, $\mathbf{d} \mid \boldsymbol{\omega}$ means $d_i \mid \omega_i$ for every $1 \leq i \leq k$.

Proposition 2.7. *Let G be an abelian group and write $G = \mathcal{C}_{r_1} \times \cdots \times \mathcal{C}_{r_k}$, where \mathcal{C}_r denotes a cyclic group of order r . Let $r_i = \nu_i \omega_i$ where ω_i is the greatest divisor of r_i that is relatively prime with t , $\boldsymbol{\nu} := (\nu_1, \dots, \nu_k)$, $\boldsymbol{\omega} := (\omega_1, \dots, \omega_k)$ and $\text{gcd}_t(\boldsymbol{\nu}) := \prod_{i=1}^k \text{gcd}_t(\nu_i)$. For $\mathbf{d} = (d_1, \dots, d_k)$ define $\varphi(\mathbf{d}) := \prod_{i=1}^k \varphi(d_i)$ and $\text{o}_{\mathbf{d}}(t) = \text{lcm}\{\text{o}_{d_i}(t) : 1 \leq i \leq k\}$. Then G has exactly $\prod_{i=1}^k \omega_i$ elements that are φ_t -periodic and the following isomorphism formula holds:*

$$(3) \quad \mathcal{G}(\varphi_t/G) = \bigoplus_{\mathbf{d} \mid \boldsymbol{\omega}} \frac{\varphi(\mathbf{d})}{\text{o}_{\mathbf{d}}(t)} \times \text{Cyc}(\text{o}_{\mathbf{d}}(t), T_{\text{gcd}_t(\boldsymbol{\nu})})$$

Proof. If formula (3) holds, the number of φ_t -periodic points in G is $\sum_{\mathbf{d} \mid \boldsymbol{\omega}} \varphi(\mathbf{d}) = \prod_{i=1}^k \left(\sum_{d_i \mid \omega_i} \varphi(d_i) \right) = \prod_{i=1}^k \omega_i$. To prove the formula we use Proposition 2.2 and Lemma 2.3 to obtain:

$$\mathcal{G}(\varphi_t/G) = \bigotimes_{i=1}^k \mathcal{G}(\varphi_t/\mathcal{C}_{r_i}) = \bigotimes_{i=1}^k \bigoplus_{d_i \mid \omega_i} \left(\frac{\varphi(d_i)}{\text{o}_{d_i}(t)} \times \text{Cyc}(\text{o}_{d_i}(t), T_{\text{gcd}_t(\nu_i)}) \right).$$

By the commutativity of the tensor product and Lemma 2.4 we have:

$$\begin{aligned} \mathcal{G}(\varphi_t/G) &= \bigoplus_{\mathbf{d} \mid \boldsymbol{\omega}} \bigotimes_{i=1}^k \left(\frac{\varphi(d_i)}{\text{o}_{d_i}(t)} \times \text{Cyc}(\text{o}_{d_i}(t), T_{\text{gcd}_t(\nu_i)}) \right) \\ &= \bigoplus_{\mathbf{d} \mid \boldsymbol{\omega}} \left(\prod_{i=1}^k \frac{\varphi(d_i)}{\text{o}_{d_i}(t)} \right) \times \left(\bigotimes_{i=1}^k \text{Cyc}(\text{o}_{d_i}(t)) \right) \otimes \left(\bigotimes_{i=1}^k T_{\text{gcd}_t(\nu_i)} \right) \end{aligned}$$

Finally, using Lemmas 2.5, 2.6 and again Lemma 2.4 we obtain the desired formula. \square

Example 2.8. Consider the power map φ_{14} over the group $\mathbb{Z}_{91}^* \cong \mathcal{C}_6 \times \mathcal{C}_{12}$ of invertible elements modulo 91. Figure 2 shows the functional graph of this map. In this case $k = 2$, $r_1 = 6$, $r_2 = 12$, $\nu = (2, 4)$, $\omega = (3, 3)$ and $\gcd_{14}(\nu) = \gcd_{14}(2) \cdot \gcd_{14}(4) = (2) \cdot (2, 2) = (4, 2)$. Then,

$$\begin{aligned} \mathcal{G}(\varphi_{14}/G) &= \bigoplus_{\substack{d_1|3 \\ d_2|3}} \frac{\varphi(d_1)\varphi(d_2)}{\text{lcm}\{o_{d_1}(14), o_{d_2}(14)\}} \times \text{Cyc}(\text{lcm}\{o_{d_1}(14), o_{d_2}(14)\}, \mathcal{T}_{(4,2)}) \\ &= \{\mathcal{T}_{(4,2)}\} \oplus 4 \times \text{Cyc}(2, \mathcal{T}_{(4,2)}). \end{aligned}$$




FIGURE 2. The graph $\{\mathcal{T}_{(4,2)}\} \oplus 4 \times \text{Cyc}(2, \mathcal{T}_{(4,2)})$.

2.4. Some results on the power map over finite groups. Let G be a finite group and let d be a divisor of $|G|$. We denote by $G[d]$ the group generated by the elements $g \in G$ such that $g^d = 1$. In the next lemma we consider the factorization $|G| = \nu\omega$ where ω is the greatest divisor of $|G|$ that is relatively prime with t . We note that if $g \in G$ verifies $g^{t^n} = 1$ then the order of g is a divisor of ν since $\gcd(t^n, |G|)$ divides ν . In particular, the tree attached to 1 in $\mathcal{G}(\varphi_t/G)$, is contained in $G[\nu]$.

Definition 2.9. An element $g \in G$ is φ_t -periodic if there exists a positive integer n such that $g^{t^n} = \varphi_t^{(n)}(g) = g$. Moreover, the *preperiod* of $g \in G$ under φ_t is the least integer $i \geq 0$ such that $\varphi_t^{(i)}(g)$ is φ_t -periodic.

Proposition 2.10. Let G be a finite group with identity 1 and consider the power map $\varphi_t : G \rightarrow G$ with $g \mapsto g^t$. Let $|G| = \nu\omega$ where ω is the greatest divisor of $|G|$ that is relatively prime with t . If $h \in G$ is φ_t -periodic and h is in the centralizer of $G[\nu]$, then the rooted tree attached to h in $\mathcal{G}(\varphi_t/G)$ is isomorphic to the rooted tree attached to 1 in $\mathcal{G}(\varphi_t/G)$.

Proof. Suppose that s is the least positive integer such that $\varphi_t^{(s)}(h) = h$, i.e., $h^{t^s} = h$. Let \mathcal{T}_h and \mathcal{T}_1 be the rooted trees attached to h and 1 in $\mathcal{G}(\varphi_t/G)$, respectively. Set $\tau : \mathcal{T}_1 \rightarrow \mathcal{T}_h$ with $\tau(g) = gh^{t^{\delta(g)(s-1)}}$, where $\delta(g)$ is the preperiod of g under φ_t . As previously remarked, each element of \mathcal{T}_1 is in $G[\nu]$.

Claim. The map τ is well defined and preserves the preperiod, i.e., $\tau(g) \in \mathcal{T}_h$ and $\delta(\tau(g)) = \delta(g)$ for every $g \in \mathcal{T}_1$.

We proceed by induction on $n = \delta(g)$. If $n = 0$, then $g = 1$ and $\tau(1) = h \in \mathcal{T}_h$ with $\delta(h) = 0$. If $n = 1$, then $g \neq 1$, $g^t = 1$ and $\tau(g) = gh^{t^{s-1}}$. In particular, since h is in the centralizer of $G[\nu]$ and $g \in \mathcal{T}_1$, we have that $\varphi_t(\tau(g)) = g^t h^{t^s} = h$. Since $g \neq 1$, it follows that $\tau(g) = gh^{t^{s-1}} \neq h^{t^{s-1}}$, the unique φ_t -periodic element in the set $\varphi_t^{-1}(h)$. The latter implies that $\tau(g) \in \mathcal{T}_h$ and $\delta(\tau(g)) = 1$. Suppose that

$\tau(g) \in \mathcal{T}_h$ and $\delta(\tau(g)) = \delta(g)$ whenever $g \in \mathcal{T}_1$ and $\delta(g) = n$ for some $n \geq 1$ and let $g \in \mathcal{T}_1$ with $\delta(g) = n + 1$. Since $n \geq 1$, we necessarily have that $\varphi_t(g) \in \mathcal{T}_1$ and $\delta(\varphi_t(g)) = n$. By induction hypothesis, $\tau(\varphi_t(g)) \in \mathcal{T}_h$ and $\delta(\tau(\varphi_t(g))) = n$. Since $\mathcal{T}_1 \subseteq G[\nu]$, h is in the centralizer of $G[\nu]$ and $n(s-1) \equiv (n+1)(s-1) + 1 \pmod{s}$ we have that

$$(4) \quad \tau(\varphi_t(g)) = g^t h^{t^{n(s-1)}} = \left(gh^{t^{(n+1)(s-1)}}\right)^t = \varphi_t(\tau(g)).$$

In particular, $\tau(g)$ is in the set $\varphi_t^{-1}(f)$ for some element $f \in \mathcal{T}_h$ of preperiod n . Since $n \geq 1$, the latter implies that $\tau(g)$ is in \mathcal{T}_h and has preperiod $n + 1$. The proof of the claim is complete.

From the claim, τ is well defined and preserves the preperiod under φ_t . By the same reasoning, the map $\tau^* : \mathcal{T}_h \rightarrow \mathcal{T}_1$ that sends the element $g \in \mathcal{T}_h$ to the element $gh^{-t^{\delta(g)(s-1)}}$ is well defined and preserves the preperiod under φ_t . It is direct to verify that τ and τ^* are the compositional inverses of each other, hence τ is a bijection. By Equation (4), we have that $\tau \circ \varphi_t = \varphi_t \circ \tau$ and then τ preserves adjacency. \square

Corollary 2.11. *Let $h \in G$ be a φ_t -periodic element. If $h \in Z(G)$ then the rooted tree attached to h in $\mathcal{G}(\varphi_t/G)$ is isomorphic to the rooted tree attached to 1 in $\mathcal{G}(\varphi_t/G)$.*

3. ON FLOWER GROUPS

Given a group G , a cyclic subgroup H is a μ -subgroup of G if H is not contained in any cyclic subgroup of G other than H itself. We have the following definition.

Definition 3.1. Let G be a finite noncyclic group and let $S = \{C_1, C_2, \dots, C_k\}$ be the collection of its μ -subgroups. The group G is a flower group if there exists a subgroup C_0 of G such that $C_i \cap C_j = C_0$ for $1 \leq i < j \leq k$. The subgroup C_0 is the *pistil* of G and the elements of S are the *petals* of G . We define the type of G as $(c_0; c_1, \dots, c_k)$ where $c_i := |C_i|$ for $0 \leq i \leq k$.

In the following proposition we provide equivalent conditions for a group G to be a flower group.

Proposition 3.2. *For a finite noncyclic group G and a cyclic subgroup C_0 of G , the following are equivalent:*

- (i) G is a flower group with pistil C_0 ;
- (ii) for any $g \in G \setminus C_0$, there exists a unique μ -subgroup C such that $g \in C$ and $C_0 \subseteq C$;
- (iii) there exist $k \geq 1$ and distinct μ -subgroups C_1, \dots, C_k of G such that $C_i \cap C_j = C_0$ for any $1 \leq i < j \leq k$, verifying

$$\sum_{i=1}^k |C_i| - (k-1)|C_0| = |G|.$$

Proof. Let S be the collection of the μ -subgroups of G . It is direct to verify that S covers G , i.e., $G = \bigcup_{C \in S} C$. For the implication (i) \rightarrow (ii), we observe that if $g \in G$ belongs to two distinct μ -subgroups C_1, C_2 of G , then $g \in C_1 \cap C_2 = C_0$. The implication (ii) \rightarrow (iii) follows by a simple counting argument. It remains to

prove that (iii) \rightarrow (i). If (iii) holds, it follows that G is the disjoint union of the sets $C_i \setminus C_0$ and C_0 . In particular, since each C_i is a μ -subgroup of G and S covers G , we have that $S = \{C_1, \dots, C_k\}$. Therefore, G is a flower group with pistil C_0 . \square

In the following proposition we provide some basic properties of flower groups.

Proposition 3.3. *Let G be a flower group with pistil C_0 and center $Z(G)$. Then the following hold:*

- (i) *If $C = \langle g \rangle$ is a μ -subgroup of G then $C \subseteq C_G(g)$. In particular, $C_0 \subseteq Z(G)$.*
- (ii) *If H is any noncyclic subgroup of G , H is a flower group with pistil $H \cap C_0$;*

Proof. (i) Since $g \in C_G(g)$ we have $C = \langle g \rangle \subseteq C_G(g)$. Observe that every element of G is contained in at least one μ -subgroup of G . Therefore, the intersection of all the μ -subgroups is contained in the intersection of all the centralizers, i.e., $C_0 \subseteq Z(G)$.

- (ii) It suffices to prove that the μ -subgroups of H are of the form $H \cap C$ with C a μ -subgroup of G . Let C' be a μ -subgroup of H . Since C' is a cyclic subgroup of G , it is contained in a μ -subgroup C of G . We have that $C' \subseteq H \cap C$ and $H \cap C$ is a cyclic subgroup of H , hence $C' = H \cap C$. \square

The following result provides a special class of flower groups G whose pistil C_0 coincides with $Z(G)$.

Proposition 3.4. *Let G be a finite non abelian group such that for every element $g \in G \setminus Z(G)$, there exists a unique μ -subgroup C_g^* of the centralizer $C_G(g)$ containing g and $Z(G)$. Then G is a flower group with pistil $Z(G)$ and the set of petals of G equals $\{C_g^* \mid g \in G \setminus Z(G)\}$.*

Proof. We observe that, for every $g \in G \setminus Z(G)$, the inclusion $Z(G) \subseteq C_g^*$ holds and, in particular, $Z(G)$ is cyclic. From Proposition 3.2, it suffices to prove that for every $g \in G \setminus Z(G)$, the group C_g^* is the unique μ -subgroup of G that contains g . The latter follows from our hypothesis and the fact that any μ -subgroup of G containing g is necessarily a μ -subgroup of $C_G(g)$. \square

We observe that the condition of Proposition 3.4 is satisfied if, for instance, $C_G(g)$ is cyclic for every $g \in G \setminus Z(G)$. The groups satisfying the latter are fully characterized in [6].

3.1. Flower groups under the power map. In this part we prove that the functional graph induced by the power map on flower groups depends only on the type of the group and provide a description of the structure of this graph.

Definition 3.5. Let G be a flower group of type $(c_0; c_1, \dots, c_k)$ and petals C_1, \dots, C_k . A compatible system of generators for G is (g_1, \dots, g_k) where g_i is a generator of C_i for $1 \leq i \leq k$ and $g_i^{c_i/c_0} = g_j^{c_j/c_0}$ for $1 \leq i < j \leq k$.

We obtain the following result.

Lemma 3.6. *Every flower group has a compatible system of generators.*

Proof. Let G be a flower group with pistil C_0 and petals C_1, \dots, C_k and let $c_i := |C_i|$ for $1 \leq i \leq k$. Let g_1 be a generator of C_1 and consider generators h_i of C_i for $2 \leq i \leq k$. Since $h_i^{c_i/c_0}$ and $g_1^{c_1/c_0}$ are both generators of C_0 we have that

$h_i^{c_i f/c_0} = g_1^{c_1/c_0}$ for some integer f with $\gcd(f, c_0) = 1$. Consider a positive integer f_i such that $f_i \equiv f \pmod{c_0}$ and $\gcd(f_i, |G|) = 1$ (for example, using Dirichlet's Theorem on arithmetic progressions we can take a prime f_i such that $f_i \equiv f \pmod{c_0}$ and $f_i > |G|$). If we set $g_i = h_i^{f_i}$ for $2 \leq i \leq k$, we have that g_i is a generator of C_i satisfying $g_i^{c_i/c_0} = h_i^{c_i f_i/c_0} = g_1^{c_1/c_0}$. \square

The following proposition entails that the isomorphism class of the functional graph induced by power maps on flower groups, depend only on their type.

Proposition 3.7. *Let G and H be two flower groups of the same type, then there is a graph isomorphism $\psi : \mathcal{G}(\varphi_t/G) \rightarrow \mathcal{G}(\varphi_t/H)$ such that $\psi(1_G) = \psi(1_H)$.*

Proof. Let C_0 and C'_0 be the pistils of G and H , respectively. Let $S = \{C_1, \dots, C_k\}$ and $S' = \{C'_1, \dots, C'_k\}$ be the set of petal of G and H , respectively. Since G and H are flower groups of the same type, with no loss of generality, we may assume that $c_i := |C_i| = |C'_i|$ for $0 \leq i \leq k$. By Lemma 3.6 we consider a compatible system of generators (g_1, \dots, g_k) and (h_1, \dots, h_k) for G and H , respectively. We note that $g_0 := g_1^{c_1/c_0}$ and $h_0 := h_1^{c_1/c_0}$ are generators of C_0 and C'_0 , respectively. For each i with $1 \leq i \leq k$ we consider the group isomorphism $\psi_i : C_i \rightarrow C'_i$ such that $\psi(g_i) = h_i$ and define $\psi : G \rightarrow H$ such that $\psi|_{C_i} = \psi_i$. To prove that this map is well defined it suffices to prove that the maps ψ_i and ψ_j coincides in C_0 for $i < j \leq k$. This last assertion follows from the fact that $\psi_i(g_0) = \psi_i(g_i^{c_i/c_0}) = h_i^{c_i/c_0} = h_0$ for $1 \leq i \leq k$ and that g_0 is a generator of C_0 . It is clear that ψ is a bijection and since $\varphi_t \psi = \psi \varphi_t$ in each petal C_i , this also happens globally and ψ induces an isomorphism between the functional graphs $\mathcal{G}(\varphi_t/G)$ and $\mathcal{G}(\varphi_t/H)$. \square

Next we provide a description of the functional graph induced by the power map on flower groups. We start with the following important auxiliary result.

Lemma 3.8. *Let G be a flower group with petals C_1, \dots, C_k and pistil C_0 . Then for any $1 \leq i \leq k$, $\varphi_t^{-1}(C_i \setminus C_0) \subseteq C_i \setminus C_0$. In particular, for an element $g \in C_i \setminus C_0$ that is φ_t -periodic, we have that the tree attached to g in $\mathcal{G}(\varphi_t/G)$ is isomorphic to the tree attached to g in $\mathcal{G}(\varphi_t/C_i)$ and the cycle containing g in $\mathcal{G}(\varphi_t/G)$ comprises vertices from $C_i \setminus C_0$.*

Proof. Suppose that $h^t \in C_i \setminus C_0$. Hence $h \notin C_0$ and then $h \in C_j \setminus C_0$ for some j with $1 \leq j \leq k$. Since $C_i \cap C_j \setminus C_0 \neq \emptyset$, we conclude that $j = i$. The remaining statement follows directly by the inclusion $\varphi_t^{-1}(C_i \setminus C_0) \subseteq C_i \setminus C_0$. \square

We obtain the following result.

Theorem 3.9. *Let t be a positive integer and G be a flower group of type $(c_0; c_1, \dots, c_k)$. Set $c_i = \nu_i \cdot \omega_i$ in a way that ω_i is the greatest divisor of c_i that is relatively prime with t . Then there exists a rooted tree $\mathcal{T}_t(G)$ such that the functional graph $\mathcal{G}(\varphi_t/G)$ of the map $\varphi_t : G \rightarrow G$ with $g \mapsto g^t$ is isomorphic to*

$$\left(\bigoplus_{i=1}^k \bigoplus_{\substack{d_i | \omega_i \\ d_i \nmid \omega_0}} \frac{\varphi(d_i)}{o_{d_i}(t)} \times \text{Cyc}(o_{d_i}(t), \mathcal{T}_{\gcd_t(\nu_i)}) \right) \oplus \left(\bigoplus_{d_0 | \omega_0} \frac{\varphi(d_0)}{o_{d_0}(t)} \times \text{Cyc}(o_{d_0}(t), \mathcal{T}_t(G)) \right),$$

where $\mathcal{T}_t(G)$ equals the tree attached to $1 \in G$ in $\mathcal{G}(\varphi_t/G)$ and has $\sum_{i=1}^k \nu_i - (k-1)\nu_0$ nodes.

Proof. Let C_0 be the pistil of G and C_1, \dots, C_k be the petals such that $|C_i| = c_i$ for $1 \leq i \leq k$. Fix $1 \leq i \leq k$ and let P_i be the set of φ_t -periodic elements of $C_i \setminus C_0$. From Proposition 2.2, P_i has $\omega_i - \omega_0$ elements, corresponding to the cycle decomposition $\bigoplus_{\substack{d_i | \omega_i \\ d_i \nmid \omega_0}} \frac{\varphi(d_i)}{o_{d_i}(t)} \times \text{Cyc}(o_{d_i}(t))$ in $\mathcal{G}(\varphi_t/G)$. From Proposition 2.2 and Lemma 3.8, to each element of P_i is attached a rooted tree isomorphic to $\mathcal{T}_{\text{gcd}_t(\nu_i)}$; this yields the first component in our statement. It remains to consider the set P_0 of φ_t -periodic elements of C_0 . From Lemma 2.2, P_0 has ω_0 elements, corresponding to the cycle decomposition $\bigoplus_{d_0 | \omega_0} \frac{\varphi(d_0)}{o_{d_0}(t)} \times \text{Cyc}(o_{d_0}(t))$ in $\mathcal{G}(\varphi_t/G)$. From Proposition 3.3, C_0 is in the center of G and then, by Proposition 2.10, the trees attached to the elements of P_0 are all isomorphic to the tree attached to the element $1 \in G$ in $\mathcal{G}(\varphi_t/G)$. The statement about the number of nodes in such tree follows by a counting argument. \square

3.2. The rooted tree $\mathcal{T}_t(G)$. Theorem 3.9 provided a description of the functional graph of the power map φ_t on a flower group G . This description is complete except for the term $\mathcal{T}_t(G)$ which is the rooted tree associated with $1 \in G$. In this part we provide some properties of these trees and completely describe it under mild conditions.

By proposition 3.7, the tree $\mathcal{T}_t(G)$ only depends on the type of the flower group G . If G has type $(c_0; c_1, \dots, c_k)$ we denote $\mathcal{T}_t(c_0; c_1, \dots, c_k) := \mathcal{T}_t(G)$.

It is convenient to extend the definition of $\mathcal{T}_t(c_0; c_1, \dots, c_k)$ for positive integers c_0, c_1, \dots, c_k with $c_0 \mid c_i$, $1 \leq i \leq k$, even if there is no flower group of type $(c_0; c_1, \dots, c_k)$. We consider set $V = \bigcup_{i=0}^k \{i\} \times \mathbb{Z}_{c_i}$. Note that if $x \in \mathbb{Z}_{c_i}$ for some i , $1 \leq i \leq k$, verifies $x \equiv 0 \pmod{c_i/c_0}$ then there is a unique $x' \in \mathbb{Z}_{c_0}$ such that $x \equiv x'c_i/c_0 \pmod{c_i}$ and we define the identification function $f : V \rightarrow V$ given by

$$f(i, x) = \begin{cases} (i, x) & \text{if } i = 0 \text{ or } x \not\equiv 0 \pmod{\frac{c_i}{c_0}}; \\ (0, x') & \text{if } i \neq 0 \text{ and } x \equiv x' \cdot \frac{c_i}{c_0} \pmod{c_i}. \end{cases}$$

From the latter, we have the following definition.

Definition 3.10. The pseudo-flower group $F = F(c_0; c_1, \dots, c_k)$ is the quotient set V/f i.e. its elements are of the form $\overline{(i, x)} := \{(j, y) \in V : f(j, y) = f(i, x)\}$. The set $V_0 := \{\overline{(i, x)} : x \in \mathbb{Z}_{c_0}\}$ is the pistil of F and the sets $V_i := \{\overline{(i, x)} : x \in \mathbb{Z}_{c_i}\}$ with $1 \leq i \leq k$ are the petals of F .

Note that, for each i , $0 \leq i \leq k$, the sets V_i have a natural structure of cyclic groups given by $\overline{(i, x)} + \overline{(i, y)} := \overline{(i, x + y)}$ and that this sum is well defined in F . Indeed, if $f(i, x_1) = f(j, x_2) = (0, x')$ and $f(i, y_1) = f(j, y_2) = (0, y')$ with $0 \leq j < i \leq k$ we have that $x \equiv x' \cdot \frac{c_i}{c_0} \pmod{c_i}$, $y \equiv y' \cdot \frac{c_i}{c_0} \pmod{c_i}$ and $x + y \equiv (x' + y') \cdot \frac{c_i}{c_0} \pmod{c_i}$, thus $f(i, x_1 + y_1) = (0, x' + y')$. In a similar way we have that $f(j, x_2 + y_2) = (0, x' + y')$ (if $j = 0$ it follows directly from the fact that $x_2 = x'$ and $y_2 = y'$). Then, $f(i, x_1 + y_1) = f(j, x_2 + y_2)$ for $0 \leq j < i \leq k$. It is clear that each V_i is a cyclic group (isomorphic to \mathbb{Z}_{c_i}) for $0 \leq i \leq k$ and they satisfy $\bigcup_{i=1}^k V_i = F$ and $V_i \cap V_j = V_0$ for $1 \leq i < j \leq k$ but F is not a genuine flower group since the sum is not defined for every pair of elements of F . Nevertheless, we can define the power map $\varphi_t : F \rightarrow F$ given by $\varphi_t(x) = tx$. The

group V_0 is called the pistil of F and the groups V_1, \dots, V_k are called the petals of F .

It is direct to verify that if G is a flower group of type $(c_0; c_1, \dots, c_k)$ and (g_1, \dots, g_k) is a compatible system of generators for G , the map $\psi : G \rightarrow F(c_0; c_1, \dots, c_k)$ given by $g_i^j \mapsto (i, j)$ for $1 \leq i \leq k$ and $j \geq 0$ is well defined (using $(i, c_i/c_0) = (0, 1)$ for $1 \leq i \leq k$), bijective and satisfies $\varphi_t \psi = \psi \varphi_t$. Thus, the induced map $\tilde{\psi} : \mathcal{G}(\varphi_t/G) \rightarrow \mathcal{G}(\varphi_t/F(c_0; c_1, \dots, c_k))$ is a graph isomorphism. Since $\psi(1) = (0, 0)$, the tree $\mathcal{T}_t(c_0; c_1, \dots, c_k)$ is isomorphic to the tree attached to $(0, 0)$ in $\mathcal{G}(\varphi_t/F(c_0; c_1, \dots, c_k))$.

Now we have defined the rooted tree $\mathcal{T}_t(c_0; c_1, \dots, c_k)$ for every c_0, c_1, \dots, c_k satisfying $c_0 \mid c_i$ even if there is no flower group of type $(c_0; c_1, \dots, c_k)$. In the case that there is a flower group G of type $(c_0; c_1, \dots, c_k)$, this tree coincides with the tree attached to $1 \in G$ in the functional graph $\mathcal{G}(\varphi_t/G)$.

The following proposition brings us a method which allows to simplify the tree structure of $\mathcal{T}_t(c_0; c_1, \dots, c_k)$.

Theorem 3.11. *Let c_0, c_1, \dots, c_k be positive integers with $k \geq 2$ and $c_0 \mid c_i$ for $1 \leq i \leq k$. Then, the following holds:*

- i) $\mathcal{T}_t(c_0; c_1, \dots, c_k) = \mathcal{T}_t(c_0; c_{\theta(1)}, \dots, c_{\theta(k)})$ for every permutation θ of the set $\{1, 2, \dots, k\}$.
- ii) If $\gcd(c_0, t) = 1$, then $\mathcal{T}_t(c_0; c_1, \dots, c_k) = \sum_{i=1}^k \mathcal{T}_{\gcd_t(c_i)}$.
- iii) If $\gcd(t, \frac{c_k}{c_0}) = 1$ then $\mathcal{T}_t(c_0; c_1, \dots, c_{k-1}, c_k) = \mathcal{T}_t(c_0; c_1, \dots, c_{k-1})$.
- iv) If $c_k \mid t$ then $\mathcal{T}_t(c_0; c_1, \dots, c_{k-1}, c_k) = \mathcal{T}_t(c_0; c_1, \dots, c_{k-1}) + \langle (c_k - c_0) \times \bullet \rangle$.
- v) $\mathcal{T}_t(c_0; c_1) = \mathcal{T}_{\gcd_t(c_1)}$.

Proof. Consider the pseudo-flower group $F = F(c_0; c_1, \dots, c_k)$ with pistil V_0 and petals V_1, \dots, V_k and the power map $\varphi_t : F \rightarrow F$. We split the proof into cases:

- i) Item i) follows directly from the definition of F .
- ii) We observe that, if $\gcd(c_0, t) = 1$, then every element of V_0 is φ_t -periodic. In particular, the tree attached to $(0, 0) \in F$ contains, in addition to the root $(0, 0)$, only elements of the set $F \setminus V_0 = \bigcup_{i=1}^k V_i \setminus V_0$. Such tree is the union of the rooted trees associated with $(0, 0) \in F$ in each of the functional graphs $\mathcal{G}(\varphi_t/V_i)$, that is, $\sum_{i=1}^k \mathcal{T}_{\gcd_t(c_i)}$.
- iii) We first note that $\gcd(t, c_k/c_0) = 1$ implies $\varphi_t(V_k \setminus V_0) \subseteq V_k \setminus V_0$. Indeed, if $(k, x) \in V_k \setminus V_0$ then $\frac{c_k}{c_0} \nmid x$ and, since $\gcd\left(t, \frac{c_k}{c_0}\right) = 1$, we also have $\frac{c_k}{c_0} \nmid tx$. Hence, $\varphi_t((k, x)) = (k, tx) \in V_k \setminus V_0$. The inclusion $\varphi_t(V_k \setminus V_0) \subseteq V_k \setminus V_0$ implies that there are no points of the tree attached to $(0, 0)$ in $V_i \setminus V_0$ and it is equal to $\mathcal{T}_t(c_0; c_1, \dots, c_{k-1})$.
- iv) We observe that if $c_k \mid t$ then $\varphi_t(V_k) = \{\overline{(0, 0)}\}$. Set $F^* = (F \setminus V_k) \cup V_0$. The tree attached to $(0, 0)$ in $\mathcal{G}(\varphi_t/F)$ can be obtained as the union of the tree attached to $(0, 0)$ in $\mathcal{G}(\varphi_t/F^*)$ (which equals $\mathcal{T}_t(c_0; c_1, \dots, c_{k-1})$) and the tree $\langle (c_k - c_0) \times \bullet \rangle$ (corresponding to the $c_k - c_0$ points of $V_k \setminus V_0$ mapping to $(0, 0)$ by φ_t), both trees with the same root, then $\mathcal{T}_t(c_0; c_1, \dots, c_{k-1}, c_k) = \mathcal{T}_t(c_0; c_1, \dots, c_{k-1}) + \langle (c_k - c_0) \times \bullet \rangle$.

- v) We note that $F(c_0; c_1) = V_1$ is a cyclic group of order c_1 and then the equality $\mathcal{T}_t(c_0; c_1) = \mathcal{T}_{\gcd_t(c_1)}$ follows from Proposition 2.2. \square

4. APPLICATIONS

In this section we provide some applications of Theorem 3.9 to the explicit description of the functional graph of the power map on certain classes of finite groups. For clarity and organization, we consider them separately.

4.1. Generalized quaternions. In [2] the authors obtain an implicit description of the digraph associated to power maps over generalized quaternion groups, such as distribution of indegrees and cycle lengths. Here we provide a more explicit description of these digraphs. We start by showing that such groups are flower groups.

Lemma 4.1. *For $n \geq 2$, the generalized quaternion*

$$Q_{4n} = \langle a, b \mid a^{2n} = 1, a^n = b^2, bab^{-1} = a^{-1} \rangle,$$

of order $4n$ is a flower group with pistil $C_0 = \{1, a^n\}$. Moreover, the set of petals of Q_{4n} comprises n cyclic groups of order 4 and one cyclic group of order $2n$.

Proof. We observe that every element of Q_{4n} is written uniquely as $a^i b^j$, where $j = 0, 1$ and $0 \leq i \leq 2n - 1$. Therefore, the μ -subgroups of Q_{4n} are the groups $C_i = \langle a^i b \rangle = \{1, a^i b, a^n, a^{n+i} b\}$ with $1 \leq i \leq n$, each of order 4 and $C_{n+1} = \langle a \rangle$, which has order $2n$. From this fact the result follows directly. \square

We obtain the following corollary.

Corollary 4.2. *Fix t an integer, let Q_{4n} be as in Lemma 4.1 and set $2n = \nu \cdot \omega$ in a way that ω is the greatest divisor of $2n$ that is relatively prime with t . Let α be the integer such that $n/2^\alpha$ is odd. Then the functional graph $\mathcal{G}(\varphi_t/Q_{4n})$ of the map $\varphi_t : Q_{4n} \rightarrow Q_{4n}$ with $g \mapsto g^t$ is isomorphic to one of the following graphs:*

$$\left(\bigoplus_{\substack{d \mid \omega \\ d > 2}} \frac{\varphi(d)}{\text{O}_d(t)} \times \text{Cyc}(\text{O}_d(t), \mathcal{T}_{\gcd_t(\nu)}) \right) \oplus \left(k_t n \times \text{Cyc} \left(\frac{2}{k_t} \right) \right) \oplus (2 \times \{\mathcal{T}_{\gcd_t(\nu)}\}),$$

if t is odd, where $k_t = 2$ if $t \equiv 1 \pmod{4}$ and $k_t = 1$ if $t \equiv 3 \pmod{4}$, or

$$\left(\bigoplus_{\substack{d \mid \omega \\ d > 1}} \frac{\varphi(d)}{\text{O}_d(t)} \times \text{Cyc}(\text{O}_d(t), \mathcal{T}_{\gcd_t(\nu)}) \right) \oplus \{T_0\},$$

if t is even, where $T_0 = \begin{cases} T_{\gcd_t(\nu)} + \langle 2n \times \bullet \rangle & \text{if } t \equiv 0 \pmod{4}; \\ T_{\gcd_t(\nu)} + \alpha \langle 2n \times \bullet \rangle & \text{if } t \equiv 2 \pmod{4}. \end{cases}$

Proof. By Lemma 4.1, the type of Q_{4n} is $(c_0; c_1, \dots, c_{n+1})$ with $c_0 = 2$, $c_i = 4$ for $1 \leq i \leq n$ and $c_{n+1} = 2n$. Let C_0 be the pistil and C_1, \dots, C_{n+1} be the petals of Q_{4n} with $|C_i| = c_i$ for $1 \leq i \leq n+1$. Set $c_i = \nu_i \cdot \omega_i$ such that ω_i is the greatest divisor of c_i that is relatively prime with t . In all the cases we apply Theorem 3.9 to obtain the graph structure of $\mathcal{G}(\varphi_t/Q_{4n})$ except for the tree $\mathcal{T}_t(Q_{4n})$ which equals T_0 when $t \equiv 0 \pmod{2}$. We split the proof into cases.

- a) If $t \equiv 1 \pmod{2}$ we have $\gcd(t, \frac{c_i}{c_0}) = 1$ for $1 \leq i \leq n$ and by Theorem 3.11 (repeated application of i. and iii. and using v. to finish) we obtain $\mathcal{T}_t(Q_{4n}) = T_{\gcd_t(\nu)}$.
- b) If $t \equiv 0 \pmod{4}$ we have $c_i \mid t$ for $1 \leq i \leq n$ and by Theorem 3.11 (repeated application of i. and iv. and using v. to finish) we obtain $T_0 = T_{\gcd_t(\nu)} + \langle 2n \times \bullet \rangle$.
- c) If $t \equiv 2 \pmod{4}$ and $1 \leq i \leq n$, the power map φ_t maps the two points of $C_i \setminus C_0$ in the unique point of order 2 in C_0 and the hanging tree of 1 in $\mathcal{G}(\varphi_t/C_{n+1})$ is $T_{\gcd_t(\nu)}$. Then, the tree T_0 can be obtained from $T_{\gcd_t(\nu)}$ by replacing the subtree T whose node corresponds to the point of order 2 in C_{n+1} with $T + \langle 2n \times \bullet \rangle$, that is, $T_0 = T_{\gcd_t(\nu)} + \alpha \langle 2n \times \bullet \rangle$ where α is the depth T . To conclude we observe that α equals the greatest integer k for which there exists $x \in \mathbb{Z}$ such that $t^k x \equiv n \pmod{2n}$. This last equation has a solution if and only if $\gcd(t^k, 2n) \mid n$. If $e_2(m)$ denotes the exponent of 2 in the prime decomposition of m , the latter is equivalent to $\min\{k \cdot e_2(t), 1 + e_2(n)\} \leq e_2(n)$, i.e., $k \cdot e_2(t) \leq e_2(n)$. Since $e_2(t) = 1$ we have $\alpha = e_2(n)$ as desired. \square

We provide two numerical examples, showing the applicability of Corollary 4.2.

Example 4.3. Consider the group $Q_{24}(n = 6)$ and the map $\varphi_3 : Q_{24} \rightarrow Q_{24}$ with $\varphi_3(g) = g^3$. From Corollary 4.2, we obtain that $\mathcal{G}(\varphi_3/Q_{24})$ is isomorphic to $\text{Cyc}(2, \mathcal{T}_3) \oplus (2 \times \text{Cyc}(1, \mathcal{T}_3)) \oplus (6 \times \text{Cyc}(2))$; see Figure 3.




FIGURE 3. The graph $\text{Cyc}(2, \mathcal{T}_3) \oplus (2 \times \text{Cyc}(1, \mathcal{T}_3)) \oplus (6 \times \text{Cyc}(2))$.

Example 4.4. Consider the group $Q_{48}(n = 12)$ and the map $\varphi_{10} : Q_{48} \rightarrow Q_{48}$ with $\varphi_{10}(g) = g^{10}$. From Corollary 4.2, we obtain that $\mathcal{G}(\varphi_{10}/Q_{48})$ is isomorphic to $2 \times \{\mathcal{T}_{(2,2,2)}\} \oplus \{\mathcal{T}_{(2,2,2)} +_2 \langle 24 \times \bullet \rangle\}$; see Figure 4.




FIGURE 4. The graph $2 \times \{\mathcal{T}_{(2,2,2)}\} \oplus \{\mathcal{T}_{(2,2,2)} +_2 \langle 24 \times \bullet \rangle\}$.

4.2. Semidirect products of cyclic groups. Let $\mathcal{C}_n = \langle a \rangle$ and $\mathcal{C}_m = \langle b \rangle$ be two cyclic groups. Every homomorphism $\phi : \mathcal{C}_m \rightarrow \text{Aut}(\mathcal{C}_n)$ is defined by $\phi(b)(a) = a^s$ for some integer s with $s^n \equiv 1 \pmod{m}$ and its associated semidirect product is denoted by $\mathcal{C}_n \rtimes_s \mathcal{C}_m$.

In [4], the authors explored the digraph associated to the power map on such groups in the case where n is a prime number. The following lemma provides a class of such groups that are also flower groups.

Lemma 4.5. *Let $m, n, s > 1$ be integers such that $s^m \equiv 1 \pmod{n}$ and $\gcd\left(n, \frac{s^j-1}{s-1}\right) = 1$ for $1 \leq j < m$. Then the semi-direct product*

$$\mathcal{C}_n \rtimes_s \mathcal{C}_m = \langle a, b \mid b^n = a^m = 1, aba^{-1} = b^s \rangle,$$

is a flower group of order mn with pistil $C_0 = \{1\}$. Moreover, the petals of $\mathcal{C}_n \rtimes_s \mathcal{C}_m$ comprises n cyclic groups of order m and one cyclic group of order n .

Proof. We observe that every element of $G := \mathcal{C}_n \rtimes_s \mathcal{C}_m$ is written uniquely as $b^i a^j$ with $0 \leq i < n$ and $0 \leq j < m$. It is direct to verify that, for any $0 \leq i, i_0 < n$ and $0 \leq j, j_0 < m$ we have that

$$(b^i a^j) \cdot (b^{i_0} a^{j_0}) = b^{i+i_0 \cdot s^j} a^{j+j_0}.$$

In particular, we obtain that

$$(5) \quad (b^i a^j)^t = b^{i \cdot \frac{s^{jt}-1}{s^j-1}} a^{jt}, \quad t > 0.$$

We claim that for every element $g = b^i a^j \in G$ with $0 \leq i < n$, $1 \leq j < m$ there is an integer i_0 such that $(b^{i_0} a)^j = b^i a^j$. Indeed, by Equation 5, it suffices to take i_0 such that $i_0 \left(\frac{s^j-1}{s-1}\right) \equiv i \pmod{n}$ and the existence of such i_0 is guaranteed by the fact that $\gcd\left(n, \frac{s^j-1}{s-1}\right) = 1$ for every $1 \leq j < m$. Thus, the μ -subgroups are $\langle b \rangle$ (of order n) and $\langle b^i a \rangle$ with $0 \leq i < n$ (of order m). Since their union covers G and $(n-1) + (m-1)n = mn - 1 = |G| - 1$ we conclude that they are the μ -subgroups with pairwise trivial intersection which implies that G is a flower group. \square

Remark 4.6. The conditions on s, m and n in Lemma 4.5 are fulfilled if, for instance, the order of s modulo p equals m for every prime divisor p of n .

Remark 4.7. For every positive integer $n \geq 2$, the pair $(s, m) = (n-1, 2)$ satisfies the conditions of Lemma 4.5. In this case, the corresponding group is just the dihedral group \mathcal{D}_{2n} of order $2n$.


The following corollary is a direct application of Lemma 4.5 and Theorem 3.9.

Corollary 4.8. *Fix t a positive integer and let $m, n, s > 1$ and $\mathcal{C}_n \rtimes_s \mathcal{C}_m$ be as in Lemma 4.5. Write $n = \nu_1 \omega_1$ in a way that ω_1 is the greatest divisor of n that is relatively prime with t and write $m = \nu_2 \omega_2$ in the same way. Then, for $n_1 = 1$ and $n_2 = n$, the functional graph $\mathcal{G}(\varphi_t / \mathcal{C}_n \rtimes_s \mathcal{C}_m)$ of the map φ_t over $\mathcal{C}_n \rtimes_s \mathcal{C}_m$ is isomorphic to*

$$\left(\bigoplus_{i=1,2} \bigoplus_{\substack{d_i \mid \omega_i \\ d_i \neq 1}} \frac{n_i \cdot \varphi(d_i)}{o_{d_i}(t)} \times \text{Cyc}(o_{d_i}(t), \mathcal{T}_{\gcd_t(\nu_i)}) \right) \oplus \{ \mathcal{T}_{\gcd_t(\nu_1)} + n \cdot \mathcal{T}_{\gcd_t(\nu_2)} \}.$$

We provide a numerical example, showing the applicability of Corollary 4.8.

Example 4.9. Consider the group $G = \mathcal{C}_{35} \rtimes_8 \mathcal{C}_4$ as in Lemma 4.5 and let $\varphi_{10} : G \rightarrow G$ with $\varphi_{10}(g) = g^{10}$. From Corollary 4.8, we have that $\mathcal{G}(\varphi_{10}/G)$ is isomorphic to $\text{Cyc}(6, \mathcal{T}_{(5)}) \oplus \{ \mathcal{T}_{(5)} + 35 \cdot \mathcal{T}_{(2,2)} \}$. Figure 5 shows a picture of this graph.


 FIGURE 5. The graph $\text{Cyc}(6, \mathcal{T}_{(5)}) \oplus \{\mathcal{T}_{(5)} + 35 \cdot \mathcal{T}_{(2,2)}\}$.

4.3. The projective general linear group. Fix $q = p^s$ a prime power and let \mathbb{F}_q be the finite field with q elements. The Projective General Linear group of order 2 over \mathbb{F}_q is the quotient group $\mathbb{G}_q := \text{PGL}(2, q) = \frac{\text{GL}(2, q)}{\mathbb{F}_q^* \cdot I}$, where $\text{GL}(2, q)$ is the group of the 2×2 non-singular matrices with entries in \mathbb{F}_q and I is the 2×2 identity matrix. It is well known that \mathbb{G}_q has order $q^3 - q$. We shall prove that \mathbb{G}_q is, in fact, a flower group. For this, we need the following machinery.

Definition 4.10. For $A \in \text{GL}(2, q)$ such that $[A] \neq [I]$, A is of **type 1** (resp. 2, 3 or 4) if its eigenvalues are distinct and in \mathbb{F}_q (resp. equal and in \mathbb{F}_q , symmetric and in $\mathbb{F}_{q^2} \setminus \mathbb{F}_q$ or not symmetric and in $\mathbb{F}_{q^2} \setminus \mathbb{F}_q$).

We observe that elements of type 3 appear only when q is odd. Moreover, the types of A and $\lambda \cdot A$ are the same for any $\lambda \in \mathbb{F}_q^*$. For this reason, we say that $[A]$ is of type t if A is of type t . The number of elements of each type and the structure of the centralizers is well known and we display them in Table 1.

p	type of g	order of g	$C_{\mathbb{G}_q}(g)$	# elements
2	1	> 2	$\cong C_{q-1}$	$q(q+1)(q-2)/2$
	2	2	$\cong C_2^s$	$q^2 - 1$
	4	> 2	$\cong C_{q+1}$	$q^2(q-1)/2$
> 2	1	> 2	$\cong C_{q-1}$	$q(q+1)(q-3)/2$
	1	2	$\cong D_{2(q-1)}$	$q(q+1)/2$
	2	p	$\cong C_p^s$	$q^2 - 1$
	3	2	$\cong D_{2(q+1)}$	$q(q-1)/2$
	4	> 2	$\cong C_{q+1}$	$q(q-1)^2/2$

TABLE 1. Element structure in $\mathbb{G}_q = \text{PGL}(2, q)$, where $q = p^s$. Here C_n and D_{2n} denote the cyclic group of order n and the dihedral group of order $2n$, respectively.

We obtain the following result.

Proposition 4.11. *For any prime power $q = p^s$, the projective general linear group $\mathbb{G}_q = \text{PGL}(2, q)$ is a flower group with pistil $C_0 = \{[I]\}$. Moreover, for $q \geq 3$, the set of petals of $\text{PGL}(2, q)$ comprises $\frac{q(q+1)}{2}$ cyclic groups of order $q-1$, $\frac{q(q-1)}{2}$ cyclic groups of order $q+1$ and $\frac{q^2-1}{p-1}$ cyclic groups of order p .*

Proof. It is well known that the center of \mathbb{G}_q equals $\{[I]\}$. By Proposition 3.4, it suffices to prove that every non identity element $g \in \mathbb{G}_q$ is contained in a unique μ -subgroup of $C_G(g)$. This is trivially verified if $C_G(g)$ is cyclic or has prime exponent.




FIGURE 6. The graph $10 \times \text{Cyc}(2, \mathcal{T}_{(2)}) \oplus 6 \times \text{Cyc}(4) \oplus \{15 \cdot \mathcal{T}_{(2,2)} + 10 \cdot \mathcal{T}_{(2)}\}$.

Otherwise, $C_G(g)$ is a dihedral group, the order of g is 2 and q is odd (see Table 1). In this case, we observe that $n = q \pm 1$ is even and the center of \mathcal{D}_{2n} equals $\{1, h^{\frac{n}{2}}\}$ where h is an element of order n . Hence $g = h^{\frac{n}{2}}$ and it is direct to verify that such element is contained in a unique μ -subgroup of \mathcal{D}_{2n} , namely the group of order n generated by h . Thus \mathbb{G}_q is a flower group with pistil $\{[I]\}$ whose set of petals comprises the cyclic subgroups of orders $p, q+1$ and $q-1$. A detailed account in Table 1 provides the number of such subgroups, according to their orders. \square

The following corollary is an immediate application of Proposition 4.11 and Theorems 3.9 and 3.11.

Corollary 4.12. *Fix t a positive integer, let $q = p^s \geq 4$ be a prime power and $\mathbb{G}_q = \text{PGL}(2, q)$. Write $q-1 = \nu_1 \omega_1$ in a way that ω_1 is the greatest divisor of $q-1$ that is relatively prime with t and write $q+1 = \nu_2 \omega_2$ in the same way. Then, for $d_1 = \frac{q(q+1)}{2}$ and $d_2 = \frac{q(q-1)}{2}$, the functional graph $\mathcal{G}(\varphi_t/\mathbb{G}_q)$ of the map $\varphi_t : \mathbb{G}_q \rightarrow \mathbb{G}_q$ with $g \mapsto g^t$ is isomorphic to*

$$\left(\bigoplus_{i=1,2} \bigoplus_{\substack{d_i | \omega_i \\ d_i \neq 1}} \frac{d_i \cdot \varphi(d_i)}{o_{d_i}(t)} \times \text{Cyc}(o_{d_i}(t), \mathcal{T}_{\text{gcd}_t(\nu_i)}) \right) \oplus \mathcal{G},$$

$$\text{where } \mathcal{G} = \begin{cases} \left\{ \frac{q(q+1)}{2} \cdot \mathcal{T}_{\text{gcd}_t(\nu_1)} + \frac{q(q-1)}{2} \cdot \mathcal{T}_{\text{gcd}_t(\nu_2)} + \frac{q^2-1}{p-1} \cdot \mathcal{T}_{(p)} \right\} & \text{if } p \mid t; \\ \left\{ \frac{(q^2-1)}{o_p(t)} \times \text{Cyc}(o_p(t)) \oplus \left\{ \frac{q(q+1)}{2} \cdot \mathcal{T}_{\text{gcd}_t(\nu_1)} + \frac{q(q-1)}{2} \cdot \mathcal{T}_{\text{gcd}_t(\nu_2)} \right\} \right\} & \text{if } p \nmid t. \end{cases}$$

We provide a numerical example, showing the applicability of the previous corollary.

Example 4.13. Consider the map $\varphi_2 : \text{PGL}(2, 5) \rightarrow \text{PGL}(2, 5), g \mapsto g^2$. From Corollary 4.12, we obtain that $\mathcal{G}(\varphi_2/\text{PGL}(2, 5))$ is isomorphic to

$$10 \times \text{Cyc}(2, \mathcal{T}_{(2)}) \oplus 6 \times \text{Cyc}(4) \oplus \{15 \cdot \mathcal{T}_{(2,2)} + 10 \cdot \mathcal{T}_{(2)}\}; \text{ see Figure 6.}$$

ACKNOWLEDGMENTS

Part of this paper were developed during a pleasant stay by the second author at Universidad de la República, supported by PEDECIBA.

REFERENCES

- [1] U. Ahmad. The power digraphs associated with generalized dihedral groups. *Discrete Math. Algorithms Appl.* 7(4): 1550057 (2015).
- [2] U. Ahmad and M. Moeen. The digraphs arising by the power maps of generalized Quaternion groups. *J. Algebra Appl.* 16(9): 1750179 (2017).
- [3] L. Blum, M. Blum and M. Shub. A simple unpredictable pseudo-random number generator. *SIAM J. Comput.* 15(2): 364–383 (1986).

- [4] G. Deng and J. Zhao. Digraph from power mapping on noncommutative groups. *J. Algebra Appl.* 19(5): 2050084 (2020).
- [5] T. A. Gassert. Chebyshev action on finite fields. *Discr. Math.* 315: 83–94 (2014).
- [6] S.M. Jafarian Amiri and H. Rostami. Finite groups whose all proper centralizers are cyclic. *Bull. Iranian Math. Soc.* 43(3): 755–762 (2017).
- [7] M. Larson. Power maps in finite groups. *Integers* 19: #A58 (2019).
- [8] R. Martins, D. Panario and C. Qureshi. A Survey on Iterations of Mappings over Finite Fields. In: *Combinatorics and finite fields: Difference sets, polynomials, pseudorandomness and applications*. Vol. 23 Radon Series on Computational and Applied Mathematics, De Gruyter, Berlin (2019).
- [9] G.L. Mullen and T.P. Vaughan. Cycles of linear permutations over a finite field. *Linear Algebra Appl.* 108: 63–82 (1988).
- [10] D. Panario and L. Reis. The functional graph of linear maps over finite fields and applications. *Des. Codes Cryptogr.* 87(2), 437–453 (2019).
- [11] A. Peinado, F. Montoya, J. Munoz and A. J. Yuste. Maximal periods of $x^2 + c$ in \mathbb{F}_q . In: *International Symposium on Applied Algebra, Algebraic Algorithms and Error-Correcting Codes, Springer* pp. 219–228 (2001).
- [12] J. M. Pollard. A Monte Carlo method for factorization. *BIT* 15(3): 331–334 (1975).
- [13] C. Qureshi and D. Panario. Rédei actions on finite fields and multiplication map in cyclic groups. *SIAM J. on Discr. Math.* 29: 1486–1503 (2015).
- [14] C. Qureshi and D. Panario. The graph structure of Chebyshev polynomials over finite fields and applications. *Des. Codes Cryptogr.* 87(2), 393–416 (2019).
- [15] C. Qureshi, D. Panario and R. Martins. Cycle structure of iterating Rédei functions. *Adv. Math. Comm.* 11(2): 397–407 (2017).
- [16] C. Qureshi and L. Reis. Dynamics of the a -map over residually finite Dedekind domains. *J. Num. Theory* 204: 134–154 (2019).
- [17] L. Reis. Moebius-like maps on irreducible polynomials and rational transformations. *J. Pure Appl. Algebra* 224(1): 169–180 (2020).
- [18] T. Rogers. The graph of the square mapping on the prime fields. *Discr. Math.* 144: 317–324 (1996).
- [19] R. A. H. Toledo, Linear Finite Dynamical Systems. *Commun. Algebra.* 33 (9): 2977–2989 (2005).
- [20] S. Ugolini. Functional graphs of rational maps induced by endomorphisms of ordinary elliptic curves over finite fields. *Periodica Math. Hungarica* 77.2: 237–260 (2018).
- [21] T. Vasiga and J. Shallit. On the iteration of certain quadratic maps over $GF(p)$. *Discr. Math.* 277: 219–240 (2004).
- [22] M. Wiener and R. Zuccherato. Faster attacks on elliptic curve cryptosystems. *International workshop on selected areas in cryptography. Springer, Berlin, Heidelberg* 190–200 (1998).

INSTITUTO DE MATEMÁTICA Y ESTADÍSTICA RAFAEL LAGUARDIA, FACULTAD DE INGENIERÍA,
UNIVERSIDAD DE LA REPÚBLICA, MONTEVIDEO 11300, URUGUAY.

Email address: cqureshi@gmail.com

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE MINAS GERAIS, BELO HORIZONTE, MG, 30270-901, BRAZIL.

Email address: lucasreismat@gmail.com