

# Closed forms of Zassenhaus formula

Léonce Dupays<sup>1,2</sup> 

<sup>1</sup> Department of Physics and Materials Science, University of Luxembourg, L-1511 Luxembourg, G. D. Luxembourg

<sup>2</sup> Donostia International Physics Center, E-20018 San Sebastián, Spain

E-mail: leonce.dupays@uni.lu

**Abstract.** Zassenhaus formula is used in a wide range of fields in physics or mathematics, from fluid dynamics to differential geometry. The non commutativity of the elements of the algebra implies that the exponential of a sum of operators cannot be splitted in the product of exponential of operators. The exponential of the sum can then be decomposed as the product of the exponentials multiplied by a supplementary term which form is generally an infinite product of exponentials. However, for some special commutators, closed forms can be found. We propose a closed form for Zassenhaus formula when the commutator of the operators X and Y satisfy the relation  $[X, Y] = uX + vY + c\mathbb{1}$ .

Zassenhaus formula then reduces to the closed form

$$e^{X+Y} = e^X e^Y e^{g_r(u,v)[X,Y]} = e^X e^{g_c(u,v)[X,Y]} e^Y = e^{g_l(u,v)[X,Y]} e^X e^Y, \quad (1)$$

a left-sided, centered and right-sided formula are found, with respective arguments,

$$g_r(u, v) = g_c(v, u)e^u = g_l(v, u) = \frac{u(e^{u-v} - e^u) + v(e^u - 1)}{vu(u-v)}. \quad (2)$$

## 1. Introduction

Zassenhaus formula found recently a regain of interest [1, 2] due to its numerous applications in various fields such as Soliton physics, Dirac's monopole problem, quantum lattices of spins or even fluids dynamics [3, 4, 5, 6]. It is also studied for mathematical fundamental purpose such as disentangling exponential operators [7, 8] or in differential geometry [9]. This formula is the dual formula of the very famous Baker-Campbell-Hausdorff (BCH) formula which aims at composing two exponential of operators.

When X and Y do not commute, the expansion of  $e^{X+Y}$  is hard to find. Some approached forms have been proposed such as the Trotter-Suzuki decomposition [10]. Recently, some new methods have been suggested, such as using a recurrence [11]. However, for some special commutators, the expansion reduces to a closed form, the most famous one being Glauber formula, known as  $e^{X+Y} = e^X e^Y e^{-\frac{1}{2}[X,Y]}$  for  $[X, [X, Y]] = 0$  and  $[Y, [X, Y]] = 0$ . But if many articles are looking for closed forms of the BCH formula [12, 13, 14, 15, 16] or if some exotic forms of BCH have been found [17] it seems that Zassenhaus formula has been forgotten. Here we offer a closed form of Zassenhaus formula for the special commutator  $[X, [X, Y]] = uX + vY + c\mathbb{1}$ .

## 2. A closed form of Zassenhaus formula

Zassenhaus theorem states that  $X, Y$  being the generators of a Lie algebra.  $e^{X+Y}$  can be uniquely decomposed in the following way :

$$e^{X+Y} = e^X e^Y \prod_{n=2}^{\infty} e^{C_n(X,Y)} = e^X e^Y e^{C_2(X,Y)} e^{C_3(X,Y)} \dots e^{C_n(X,Y)} \dots \quad (3)$$

where  $C_n(X, Y)$  is a Lie polynomial in  $X$  and  $Y$  of degree  $n$ .  
The first terms are expressed in this way,

$$e^{X+Y} = e^X e^Y e^{-\frac{1}{2}[X,Y]} e^{\frac{1}{6}(2[Y,[X,Y]]+[X,[X,Y]])} e^{-\frac{1}{24}([[[X,Y],X],X]+3[[X,Y],X],Y)+3[[X,Y],Y],Y)} \dots \quad (4)$$

In the case of the specific commutators,  $[X, Y] = uX + vY + c\mathbb{1}$ , the multiple commutators in the argument of the exponentials all reduce to a constant multiplied by  $[X, Y]$ . Equation (3) may be recasted in the form  $e^{X+Y} = e^X e^Y e^{\mu[X,Y]}$  where  $\mu$  is to be determined. We denote the adjoint operator  $ad_A$  of an operator  $A$ , defined by recurrence from  $ad_A^m B = [A, ad_A^{m-1} B]$  with  $ad_A^0 B = B$  so that it is possible to rewrite  $e^A B e^{-A} = e^{ad_A} B$ . A first method to determine this constant is to express the Zassenhaus formula in an integral form, introducing a new variable  $t$  in the exponential, as could be noticed in [10]

$$e^{t(X+Y)} = e^{tX} e^{tY} \exp \left( \int_0^t ds e^{-s ad_Y} \left( e^{-s ad_X} - \mathbb{1} \right) Y \right). \quad (5)$$

This formula can be demonstrated in differentiating  $e^{t(X+Y)} = e^{tX} e^{tY} e^{\eta}$  with respect to the variable  $t$  and noticing that  $\frac{\partial \eta}{\partial t} = e^{-t ad_Y} (e^{-t ad_X} - \mathbb{1}) Y$ . It is then useful to notice that for the special commutator  $[X, Y] = uX + vY + c\mathbb{1}$ ,

$$e^{-s ad_Y} \left( e^{-s ad_X} - \mathbb{1} \right) Y = \frac{e^{s(u-v)} - e^{su}}{v} [X, Y]. \quad (6)$$

Inserting (6) in (5) allows to perform the integration, then choosing  $t = 1$  gives the closed expression for the Zassenhaus formula

$$e^{(X+Y)} = e^X e^Y \exp(g_r(u, v)[X, Y]). \quad (7)$$

With the right-sided coefficient  $g_r(u, v)$ ,

$$g_r(u, v) = \frac{u(e^{u-v} - e^u) + v(e^u - 1)}{uv(u-v)}. \quad (8)$$

Equations (7) and (8) are our main result.

## 3. Left-sided Zassenhaus formula with a second demonstration

It is possible to derive (7) and (8) using another derivation, constructing the Zassenhaus formula from a closed form of BCH formula. We will derive here a left-sided expression of the Zassenhaus formula for the special commutator  $[X, Y] = uX + vY + c\mathbb{1}$  with coefficient  $g_l(u, v)$

$$\exp(X + Y) = \exp(g_l(u, v)[X, Y]) \exp(X) \exp(Y). \quad (9)$$

Van-Brunt and Visser [12] found a closed form BCH formula for the commutator  $[X, Y] = uX + vY + c\mathbb{1}$ ,

$$\exp(X) \exp(Y) = \exp(X + Y + f(u, v)[X, Y]). \quad (10)$$

With  $f$ , the Van-Brunt and Visser function [12]

$$f(u, v) = \frac{ue^u(e^v - 1) - ve^v(e^u - 1)}{uv(e^u - e^v)}. \quad (11)$$

Inserting (10) in (9) one can find

$$\exp(X + Y) = \exp(g_l(u, v)[X, Y]) \exp(X + Y + f(u, v)[X, Y]). \quad (12)$$

In order to determine  $g_l(u, v)$  one can apply Van-Brunt Visser formula (10) directly to (12), and make the equality hold. It is useful to proceed to the change of variables  $\tilde{X} = g_l(u, v)[X, Y]$ ,  $\tilde{Y} = X + Y + f(u, v)[X, Y]$ , and to compute the commutator

$$[\tilde{X}, \tilde{Y}] = g_l(u, v)(u - v)[X, Y] \quad (13)$$

$$= g_l(u, v)(u - v)(uX + vY + c\mathbb{1}). \quad (14)$$

Furthermore, the commutator can also be written

$$[\tilde{X}, \tilde{Y}] = \tilde{u}\tilde{X} + \tilde{v}\tilde{Y} + \tilde{c}\mathbb{1}, \quad (15)$$

$$= \tilde{u}g_l(u, v)[X, Y] + \tilde{v}(X + Y + f(u, v)[X, Y]) + \tilde{c}\mathbb{1}, \quad (16)$$

$$= \{\tilde{u}g_l(u, v)u + \tilde{v} + f(u, v)\tilde{v}u\}X + \{\tilde{u}g_l(u, v)v + \tilde{v} + f(u, v)\tilde{v}v\}Y \\ + \{\tilde{u}g_l(u, v)c + \tilde{v}f(u, v)c + \tilde{c}\}\mathbb{1}. \quad (17)$$

This leads to the system of equations

$$g_l(u, v)(u - v)u = \{\tilde{u}g_l(u, v) + f(u, v)\tilde{v}\}u + \tilde{v}, \quad (18)$$

$$g_l(u, v)(u - v)v = \{\tilde{u}g_l(u, v) + f(u, v)\tilde{v}\}v + \tilde{v}, \quad (19)$$

$$g_l(u, v)(u - v)c = \{\tilde{u}g_l(u, v) + f(u, v)\tilde{v}\}c + \tilde{c}. \quad (20)$$

That is satisfied for  $\tilde{u} = u - v$ ,  $\tilde{v} = 0$ ,  $\tilde{c} = 0$ .

Hence,

$$\exp(X + Y) = \exp(g_l(u, v)[X, Y]) \exp(X + Y + f(u, v)[X, Y]), \quad (21)$$

$$= \exp\{X + Y + [g_l(u, v) + f(u, v) + f(u - v, 0)(u - v)g_l(u, v)][X, Y]\}. \quad (22)$$

To satisfy the previous equality,  $g_l(u, v)$  may take the form

$$g_l(u, v) = \frac{v(e^{v-u} - e^v) + u(e^v - 1)}{uv(v - u)}. \quad (23)$$

We notice that  $g_l(u, v) = g_r(v, u)$ . The same demonstration can also be employed to find a right-sided formula confirming the previously found result (8).

#### 4. Commutating the generators of the algebra

If  $X, Y$  are elements of an algebra which verify the commutation relationship  $[X, Y] = uX + vY + c\mathbb{1}$ , we can find the commutation relation between the generators of the algebra  $e^X$  and  $e^Y$ , and look for an expression of the form  $e^X e^Y = e^Y e^X e^{\gamma[X, Y]}$  where  $\gamma$  is to be determined. As  $X + Y = Y + X$  one can use (7) for both commutators  $[X, Y] = uX + vY + c\mathbb{1}$  and  $[Y, X] = \alpha Y + \beta X + \delta\mathbb{1} = -uX - vY - c\mathbb{1}$  so that

$$e^{X+Y} = e^X e^Y e^{g_r(u,v)[X,Y]}, \quad (24)$$

$$e^{Y+X} = e^Y e^X e^{g_r(\alpha,\beta)[Y,X]} = e^Y e^X e^{-g_r(-v,-u)[X,Y]}. \quad (25)$$

From the previous equalities:

$$e^X e^Y = e^Y e^X e^{-(g_r(-v,-u)+g_r(u,v))[X,Y]}. \quad (26)$$

$$-g_r(-v,-u) - g_r(u,v) = \frac{1}{vu(u-v)} \{v(e^{-v}-1)(e^u-1) + u(e^{-v}-1)(1-e^u)\}. \quad (27)$$

### 5. Properties of $g_r(u,v)$

In order to compute  $g_r(u,v)$  in the special cases  $u=0$ ,  $v=0$  or  $u=v=0$  De l'Hôpital's rule may be used, we compute

$$g_r(0,v) = -\frac{e^{-v}-1+v}{v^2}, \quad g_r(u,0) = \frac{e^u(1-u)-1}{u^2}, \quad g_r(0,0) = -\frac{1}{2}. \quad (28)$$

For a commutator of the form  $[X,Y] = c\mathbb{1}$  one can express  $e^{X+Y} = e^X e^Y e^{-\frac{1}{2}[X,Y]}$  which is nothing but Glauber formula. Finally the case  $u=v$  is also interesting

$$g_r(u,u) = \frac{u+1-e^u}{u^2}. \quad (29)$$

Special forms of  $g_l$  and  $g_c$  are directly found from  $g_r$ .

### 6. A centered formula

It is also possible to look for a centered formula with coefficient  $g_c(u,v)$  so that

$$e^{X+Y} = e^X e^{g_c(u,v)[X,Y]} e^Y. \quad (30)$$

Starting from,

$$e^{X+Y} = e^{g_l(u,v)[X,Y]} e^X e^Y, \quad (31)$$

and setting  $\tilde{X} = g_l(u,v)[X,Y]$  and  $\tilde{Y} = X$  so that the commutator reads  $[\tilde{X},\tilde{Y}] = -v\tilde{X}$ . Applying the result of (26), equation (31) becomes,

$$\begin{aligned} e^{X+Y} &= e^X e^{g_l(u,v)[X,Y]} e^{-(g_r(0,v)+g_r(-v,0))[\tilde{X},\tilde{Y}]} e^Y, \\ &= e^X e^{g_l(u,v)[X,Y]} e^{v(g_r(0,v)+g_r(-v,0))g_l(u,v)[X,Y]} e^Y. \end{aligned} \quad (32)$$

Where,

$$g_c(u,v) = \{1 + v(g_r(0,v) + g_r(-v,0))\} g_l(u,v), \quad (33)$$

$$= e^{-v} g_l(u,v). \quad (34)$$

## 7. Application to the Lie algebra $\{a^{\dagger 2}, a^2, a^{\dagger}a\}$

One can apply the formula to the commutators  $[a^{\dagger 2}, a^{\dagger}a] = -2a^{\dagger 2}$ ,  $[a^2, a^{\dagger}a] = 2a^2$

$$e^{a^{\dagger 2} + a^{\dagger}a} = e^{a^{\dagger 2}} e^{a^{\dagger}a} e^{g_r(-2,0)[X,Y]} = e^{a^{\dagger 2}} e^{a^{\dagger}a} e^{\frac{3e^{-2}-1}{4}a^{\dagger 2}}, \quad (35)$$

$$e^{a^2 + a^{\dagger}a} = e^{a^{\dagger 2}} e^{a^{\dagger}a} e^{g_r(2,0)[X,Y]} = e^{a^2} e^{a^{\dagger}a} e^{-\frac{e^2+1}{4}a^2}. \quad (36)$$

## 8. A third derivation of the formula

In [11] the authors developed a technique to compute recursively the exponents of the Zassenhaus formula. But it can be shown that if X and Y verify the above mentioned commutation relation, their work can be used to recover the relation (8).

### 8.1. Computing the Zassenhaus formula exponents with a recurrence

To find the terms of Zassenhaus formula by recurrence, Casas et al. [11] introduce a differentiable formula with respect to  $t$  that recovers Zassenhaus formula (3) for  $t = 1$

$$e^{t(X+Y)} = e^{tX} e^{tY} e^{t^2 C_2} e^{t^3 C_3} \dots \quad (37)$$

And the following compositions are considered

$$R_1(t) = e^{-tY} e^{-tX} e^{t(X+Y)}. \quad (38)$$

So that for each  $n \geq 2$ ,

$$R_n(t) = e^{-t^n C_n} \dots e^{-t^2 C_2} e^{-tY} e^{-tX} e^{t(X+Y)} = e^{-t^n C_n} R_{n-1}(t). \quad (39)$$

The authors [11] define the functions  $F_n = \left(\frac{d}{dt} R_n(t)\right) R_n(t)^{-1}$  that allows to determine the polynomial  $C_n$  through the recurrence relations

$$F_n(t) = e^{-t^n ad_{C_n}} \left( F_{n-1}(t) - \frac{t^{n-1}}{(n-1)!} F_{n-1}^{(n-1)}(0) \right), \quad (40)$$

$$C_n = \frac{1}{n!} F_{n-1}^{(n-1)}(0), \quad (41)$$

with the  $n^{th}$  derivative noted as  $F^{(n)} = \frac{\partial^n F}{\partial t^n}$ . Furthermore, still from [11], one can get:

$$F_1(t) = e^{-tad_Y} \left( e^{-tad_X} - \mathbb{1} \right) Y. \quad (42)$$

It is easily seen that from the knowledge of  $F_1$ ,  $C_2$  is known and so does  $F_2$ , etc. A recurrence scheme to determine de coefficients  $C_n$  is then established.

## 9. Impact of the closed form $[X, Y] = uX + vY + c\mathbb{1}$ on the recurrence

This recurrence scheme can be exploited in order to find directly the  $C_n$  without any recurrence if the commutator between X and Y is appropriate. If X and Y are two elements of an algebra such that their commutator verify  $[X, Y] = uX + vY + c\mathbb{1}$ . It is then obvious that  $[X, [X, Y]] = v[X, Y]$  and  $[Y, [X, Y]] = -u[X, Y]$ . And  $F_1$  can be written,

$$F_1(t) = e^{-tad_Y} \left( e^{-tad_X} - \mathbb{1} \right) Y = \frac{e^{tu}(e^{-tv} - \mathbb{1})}{v} [X, Y]. \quad (43)$$

Let's demonstrate by recurrence that  $F_n$  is always proportional to the commutator  $[X, Y]$ , so that  $F_n = \beta_n(t)[X, Y]$  where  $\beta_n$  is a function depending on  $t$ . The initialization is clear since  $F_1(t) = \frac{e^{tu}(e^{-tv}-1)}{v}[X, Y]$ .

We wish to express  $F_{n+1}(t)$  defined by

$$F_{n+1}(t) = e^{-t^{n+1}ad_{C_{n+1}}} \left( F_n(t) - \frac{t^n}{(n)!} F_n^{(n)}(0) \right). \quad (44)$$

But according to the recurrence hypothesis,

$$C_{n+1} = \frac{1}{(n+1)!} \left( \frac{\partial^n}{\partial t^n} \beta_n \right) (0)[X, Y]. \quad (45)$$

So that the adjoint operator is commuting with the term it acts on, leading to the simplification

$$F_{n+1}(t) = \left[ \beta_n(t) - \frac{t^n}{n!} \left( \frac{\partial^n}{\partial t^n} \beta_n \right) (0) \right] [X, Y], \quad (46)$$

so that,

$$\beta_{n+1}(t) = \beta_n(t) - \frac{t^n}{n!} \left( \frac{\partial^n}{\partial t^n} \beta_n \right) (0). \quad (47)$$

The  $F_n$  being of the form  $F_n = \beta_n[X, Y]$ , one can express directly the coefficients  $C_n$ .

### 9.1. Expression of the closed Zassenhaus formula

It is now possible to find a closed form of Zassenhaus formula in summing up all the  $C_n$ . The coefficients  $C_{n+1}$  can be expressed  $\forall n \geq 1$ ,

$$C_{n+1} = \frac{1}{(n+1)!} \left( \frac{\partial^n}{\partial t^n} \beta_n \right) (0)[X, Y]. \quad (48)$$

According to (47),

$$\frac{\partial^{n+1}}{\partial t^{n+1}} \beta_{n+1} = \frac{\partial^{n+1}}{\partial t^{n+1}} \beta_n. \quad (49)$$

From them on  $\forall n \geq 1$ ,

$$\begin{aligned} C_{n+1} &= \frac{1}{(n+1)!} \left( \frac{\partial^n}{\partial t^n} \beta_n \right) (0)[X, Y] = \frac{1}{(n+1)!} \left( \frac{\partial^n}{\partial t^n} \beta_1 \right) (0)[X, Y], \\ &= \frac{1}{(n+1)!} \left( \frac{\partial^n}{\partial t^n} \frac{e^{tu}(e^{-tv}-1)}{v} \right) (0)[X, Y]. \end{aligned} \quad (50)$$

Using a Cauchy product one can develop in whole series  $\frac{e^{tu}(e^{-tv}-1)}{v}$ .

$$\frac{e^{tu}(e^{-tv}-1)}{v} = \frac{1}{v} \sum_{k=0}^{\infty} \sum_{s=0}^k \left( \frac{u^s (-v)^{k-s}}{s! (k-s)!} \right) t^k - \frac{1}{v} \sum_{k=0}^{\infty} \frac{u^k}{k!} t^k. \quad (51)$$

Hence,

$$\frac{\partial^n}{\partial t^n} \left( \frac{e^{tu}(e^{-tv} - \mathbb{1})}{v} \right) = \sum_{k=0}^{\infty} \frac{k!}{(k-n+1)!} \left[ \frac{1}{v} \left( \sum_{s=0}^k \left( \frac{u^s (-v)^{k-s}}{s! (k-s)!} \right) - \frac{u^k}{k!} \right) \right] t^{k-n}, \quad (52)$$

and,

$$\begin{aligned} \frac{\partial^n}{\partial t^n} \left( \frac{e^{tu}(e^{-tv} - \mathbb{1})}{v} \right) (0) &= n! \left( \frac{1}{v} \left( \sum_{s=0}^n \left( \frac{u^s (-v)^{n-s}}{s! (n-s)!} \right) - \frac{u^n}{n!} \right) \right), \\ &= \frac{1}{v} ((u-v)^n - u^n). \end{aligned} \quad (53)$$

This leads to  $\forall n \geq 1$

$$C_{n+1} = \frac{1}{(n+1)!} \frac{1}{v} ((u-v)^n - u^n). \quad (54)$$

Hence, one can get the series,

$$\begin{aligned} \sum_{n=2}^{\infty} C_n &= \sum_{n=1}^{\infty} C_{n+1} = \sum_{n=1}^{\infty} \frac{1}{(n+1)!} \frac{1}{v} ((u-v)^n - u^n) = \frac{1}{v(u-v)} \sum_{n=1}^{\infty} \frac{(u-v)^{n+1}}{(n+1)!} - \frac{1}{vu} \sum_{n=1}^{\infty} \frac{u^{n+1}}{(n+1)!}, \\ &= \frac{1}{v(u-v)} \sum_{n=2}^{\infty} \frac{(u-v)^n}{n!} - \frac{1}{vu} \sum_{n=2}^{\infty} \frac{u^n}{n!} = \frac{1}{v(u-v)} (e^{u-v} - 1 - (u-v)) - \frac{1}{vu} (e^u - 1 - u), \end{aligned} \quad (55)$$

and obtain the closed form of the Zassenhaus formula given by equations (7) and (8).

## Conclusion

We demonstrated a new closed form of Zassenhaus formula for a special commutator, this closed form can be expressed in three forms, a right-sided, left-sided and centered formula. This formula proves to be useful in quantum mechanics, for instance to disentangle operators. We hope this formula can be useful in a broad range of contexts, notably in quantum mechanics.

## Acknowledgement

It is a great pleasure to thank Jean-Christophe Pain to have sparked the interest of the author in closed forms of Baker-Campbell-Hausdorff formula. It is also a pleasure to thank Adolfo del Campo and Iñigo Luis Egusquiza for bright comments. The author would like to acknowledge DIPC for financial support at early stage of this work.

## References

- [1] Kimura T 2017 *Progress of Theoretical and Experimental Physics* **2017** ISSN 2050-3911 URL <http://dx.doi.org/10.1093/ptep/ptx044>
- [2] Wang L, Gao Y and Jing N 2019 *Frontiers of Mathematics in China* **14** 421–433 ISSN 1673-3576 URL <http://dx.doi.org/10.1007/s11464-019-0760-1>
- [3] Tumer S N and Pashaev O K 2017 q-viscous burgers' equation: Dynamical symmetry, shock solitons and q-semiclassical expansion (*Preprint* [1707.01737](https://arxiv.org/abs/1707.01737))
- [4] Soloviev M 2016 *Physical Review D* **94** ISSN 2470-0029 URL <http://dx.doi.org/10.1103/PhysRevD.94.105021>
- [5] Braz F, Rodrigues F, de Souza S and Rojas O 2016 *Annals of Physics* **372** 523–543 ISSN 0003-4916 URL <http://dx.doi.org/10.1016/j.aop.2016.07.007>

- [6] Geiser J 2012 Embedded zassenhaus expansion to operator splitting schemes: Theory and application in fluid dynamics (*Preprint* [1204.0380](#))
- [7] Sridhar R and Jagannathan R 2002 On the q-analogues of the zassenhaus formula for disentangling exponential operators (*Preprint* [math-ph/0212068](#))
- [8] Quesne C 2004 *International Journal of Theoretical Physics* **43** 545–559 ISSN 0020-7748 URL <http://dx.doi.org/10.1023/B:IJTP.0000028885.42890.f5>
- [9] Nishimura H 2013 The baker-campbell-hausdorff formula and the zassenhaus formula in synthetic differential geometry (*Preprint* [1304.4667](#))
- [10] Masuo S 1985 *Journal of Mathematical Physics* **26** 601–612
- [11] Casas F, Murua A and Nadinic M 2012 *Computer Physics Communications* **183** 2386–2391 ISSN 0010-4655 URL <http://dx.doi.org/10.1016/j.cpc.2012.06.006>
- [12] Van-Brunt A and Visser M 2015 *Journal of Physics A: Mathematical and Theoretical* **48** 225207 ISSN 1751-8121 URL <http://dx.doi.org/10.1088/1751-8113/48/22/225207>
- [13] Matone M 2016 *The European Physical Journal C* **76** ISSN 1434-6052 URL <http://dx.doi.org/10.1140/epjc/s10052-016-4455-4>
- [14] Matone M 2015 *Journal of High Energy Physics* **2015** ISSN 1029-8479 URL [http://dx.doi.org/10.1007/JHEP05\(2015\)113](http://dx.doi.org/10.1007/JHEP05(2015)113)
- [15] Van-Brunt A and Visser M 2018 *Mathematics* **6** 135 ISSN 2227-7390 URL <http://dx.doi.org/10.3390/math6080135>
- [16] Pain J C 2012 *Journal of Physics A: Mathematical and Theoretical* **46** 035304 ISSN 1751-8121 URL <http://dx.doi.org/10.1088/1751-8113/46/3/035304>
- [17] Mostovoy J, Perez-Izquierdo J M and Shestakov I P 2016 A non-associative baker-campbell-hausdorff formula (*Preprint* [1605.00953](#))