

Extending Properly n -REA Sets¹

Peter Gerdes

Mathematics Department, Indiana University, Bloomington, Rawles Hall, 831 East 3rd St., Bloomington, IN 47405 USA
gerdes@invariant.org

Peter Cholak*

Mathematics Department, University of Notre Dame du Lac, 255 Hurley Building, Notre Dame, IN 46556, USA
cholak@nd.edu

Keywords: REA sets, recursively enumerable and above, Turing degree, recursively enumerable, computability theory, recursion theory

1. Introduction

In [2] Jockusch and Shore introduce the n -REA sets. A 1-REA set is just an r.e. set, while an $n + 1$ -REA set is a set of the form $A \oplus W_e^A$ where A is n -REA. Despite the fact that the n -REA sets are merely the result of iterating the operation of adding an r.e. set, many basic questions remain even when n is finite.

One natural property to investigate about n -REA sets is when they are properly n -REA.

Definition 1.1. A set B is properly n -REA if B is n -REA and B isn't Turing equivalent to any m -REA set with $m < n$.

It is evident that for all $n \in \omega$ there are properly n -REA sets. But it's obviously not the case that for every n -REA set A there is a properly $n + 1$ -REA set of the form $A \oplus W_e^A$, i.e., not every n -REA set can be extended to a properly $n + 1$ -REA set. For instance, the empty set is a 1-REA set but can't be extended to a properly 2-REA set.

A natural hypothesis is that any properly n -REA set can be extended to a properly $n + 1$ -REA set. This represents the most optimistic possible hypothesis about when n -REA sets can be extended to properly $n + 1$ -REA sets.

Hypothesis 1.2. If A is a properly n -REA set then there is a properly $n + 1$ -REA set (Turing equivalent to a set²) of the form $A \oplus W_e^A$. In such cases, we say A can be extended to a properly $n + 1$ -REA set.

Further evidence for this hypothesis comes in the form of a result by Soare and Stob [5] who demonstrate that for any r.e. set $W_e >_T \mathbf{0}$ there is a set 1-REA in W_e but not of r.e. degree. This establishes that every properly 1-REA set can be extended to a properly 2-REA set. This result was extended to $n = 2$ by Cholak and Hinman when they established the following result (recast using the above definitions) in [1].

Theorem 1.3. *If X is properly 2-REA then there is a REA(X) set $W_e^X \oplus X$ which is properly 3-REA.*

In other words any 2-REA set which isn't of r.e. degree can be extended to a 3-REA set not of 2-REA degree. However, despite the suggestive evidence we show that the above attractive hypothesis in fact fails by proving the following theorem.

¹Thanks to an anonymous referee for many helpful comments and editing suggestions.

*Cholak was partially supported by a Focused Research Group grant from the National Science Foundation of the United States, DMS-1854136.

²We add this caveat because, in the argument below, we adopt a slightly different form for n -REA sets that makes the argument more convenient.

Theorem 1.4. *There is a properly 3-REA set A which can't be extended to a properly 4-REA set. The set A can also be taken to be Δ_2^0 .*

Note that, if A isn't properly 3-REA then it is evident that no set of the form $A \oplus W_e^A$ will be properly 4-REA. Our strategy, roughly speaking, will be to build a 3-REA set A such that enumeration into the first and second components of A (where the first component of A is the r.e. part of A and the second component is the part of A r.e. in the first component) doesn't result in further changes to W_i^A . The construction takes the form of a finite injury argument, with the complexity arising from the difficulty of showing that the requirements eventually succeed. However, before we present the main construction, we first review some notation and define the n -REA sets in [section 2](#). Note that in this section we also introduce the idea that n -REA sets can be viewed as $n + 1$ column sets (with the 0-th column empty) produced by an r.e. set of axioms.

Readers familiar with n -REA sets may wish to jump ahead to [section 3](#), where we describe the requirements that our construction will meet and delineate some basic conditions that our construction will satisfy.

2. Background

2.1. Notational Conventions

We largely adopt the standard notation seen in [3] which we briefly review. The use of $\Phi_{i,s}(X; y)$ is denoted by $u[\Phi_{i,s}(X; y)]$, the e -th set r.e. in X by W_e^X and we write $y \searrow_s X$ ($y \searrow_s \bar{X}$) to indicate y enters (leaves) X at stage s . We let $\langle\langle x_0, \dots, x_n \rangle\rangle$ ($\langle x, y \rangle \stackrel{\text{def}}{=} \frac{1}{2}(x + y)(x + y + 1) + y$) denote a canonical bijection of $\omega^{<\omega}$ (ω^2) with ω and define $A \oplus B$, $\bigoplus_{n \in S} X_n$ and $X^{[n]}/X^{[<n]}$ standardly³. We write $\sigma \perp \tau$ ($\sigma \upharpoonright \tau$) to indicate that two strings/partial functions/etc. are incompatible (compatible).

We also adopt some less common notational conventions and assumptions. We let the variables χ, ξ, η range over $\{0, 1, \uparrow\}^{<\omega}$ (binary partial functions with finite domain) with extension denoted by \succ . By identifying sets with their characteristic functions (so A_s is really a $\{0, 1, \uparrow\}^{<\omega}$) we gain the ability for our approximations to a set A to take no position on whether $x \in A$. This turns out to be helpful in giving well-behaved stagewise approximations to n -REA sets.

With this notation in place, we briefly review n -REA sets and describe how we identify such sets with $n + 1$ (non-trivial) column sets. The reader familiar with n -REA sets may wish to skip ahead to [section 3](#) after familiarizing themselves with the notion of an A supported approximation in [definition 2.6](#).

2.2. α -REA Sets

In [2] Jockusch and Shore introduce the α -REA sets for any ordinal notation $\alpha \in \mathcal{O}_1$. In this paper, we will only be concerned with finite values of α so we adopt the following definition (equivalent up to 1-degree for finite α).

Definition 2.1. Given a computable function f , we define the n -REA operator $\mathcal{J}_f^n(X)$ (or $\mathcal{J}_e^n(X)$ where e is an index for f), $n \in \omega$ via the following inductive definition (where we let $W_{f(m)}^Z = \emptyset$ if $f(m) \uparrow$).

$$\begin{aligned} \widehat{\mathcal{J}}_f^0(X) &= X \\ \widehat{\mathcal{J}}_f^{m+1}(X) &= W_{f(m)}^{\mathcal{J}_f^m(X)} \\ \mathcal{J}_f^n(X) &= \bigoplus_{m < n+1} \widehat{\mathcal{J}}_f^m(X) \end{aligned} \tag{1}$$

³That is, $A \oplus B \stackrel{\text{def}}{=} \{y \mid y = 2x \wedge x \in A \vee y = 2x + 1 \wedge x \in B\}$ and $\bigoplus_{n \in S} X_n \stackrel{\text{def}}{=} \{(n, x) \mid n \in S \wedge x \in X_n\}$

Furthermore, C is an n -REA(X) set just if $C = \mathcal{J}_f^n(X)$ for some computable function f and C is an n -REA set if it is an n -REA(\emptyset) set. We define X_e^n (written just as X_e when n is understood) to be $\mathcal{J}_e^n(\emptyset)$.

We further note that we can describe the construction of n -REA(X) with $n \in \omega$ sets via the enumeration of ‘axioms’ defined as follows.

Definition 2.2. An **axiom** is a pair $\langle \sigma \rightarrow y \rangle$ with $\sigma \in \{0, 1, \uparrow\}^{<\omega}$, $y \in \omega$. An **REA axiom** is an axiom that further satisfies $y = \langle m, z \rangle$ with $m > 0$ and $\text{dom } \sigma \subset \omega^{[<m]}$. Finally, an **n -REA axiom** is an REA axiom with $m \leq n$.

We think of the axiom $\langle \sigma \rightarrow y \rangle$ as an instruction to put y into a set Z provided $\sigma \prec Y$. Thus, regarding y as coding an element of $2^{<\omega}$, an r.e. set of axioms defines (if compatible) a computable functional. An REA axiom $\langle \sigma \rightarrow \langle m, z \rangle \rangle$ is then an instruction to put z into $X^{[m]}$ provided $\sigma \prec X^{[<m]}$ (as this is equivalent to $\sigma \prec X$).

We now argue that for $n \in \omega$ we may identify n -REA operators \mathcal{J}_e^n (and hence n -REA(X) sets) with r.e. sets of n -REA axioms and effectively translate between r.e. indices for sets of n -REA axioms and indices for n -REA operators.

Lemma 2.3. *If \mathcal{A} is a r.e. set of n -REA axioms ($n \in \omega$) then the operator \mathcal{J}_A^n defined by*

$$\langle l, y \rangle \in \mathcal{J}_A^n(X) \iff (l = 0 \wedge y \in X) \vee \exists [\sigma \prec \mathcal{J}_A^n(X)] [\langle \sigma \rightarrow \langle l, y \rangle \rangle \in A] \quad (2)$$

is an n -REA operator. Conversely, given an n -REA operator \mathcal{J}_f^n there is a r.e. set A_f such that $\mathcal{J}_{A_f}^n = \mathcal{J}_f^n$. Furthermore, we can effectively translate between r.e. indices of sets of n -REA axioms and indices of n -REA operators.

Proof. We first prove that \mathcal{J}_A^n is an n -REA operator. Note that, for $0 < l \leq n$, $\mathcal{J}_A^n(X)^{[l]}$ is determined by $\mathcal{J}_A^n(X)^{[<l]}$ and X so $\mathcal{J}_A^n(X)$ is well defined. Furthermore, the above equation explicitly defines $\mathcal{J}_A^n(X)^{[l]}$ from $\mathcal{J}_A^n(X)^{[<l]}$ and l via a (uniformly) Σ_1^0 formula. Thus, by an application of the s-m-n theorem [4] there is a computable function f satisfying [definition 2.1](#).

Given \mathcal{J}_e^n we simply enumerate axioms $\langle \sigma \rightarrow \langle m + 1, x \rangle \rangle$ into \mathcal{A} when $\Phi_{f(m)}(\sigma; x) \downarrow$ for $m < n$. The uniformity claim is evident from the proof. \square

In light of this result we adopt the following notation.

Notation 2.4. $\mathcal{A}(J)$ denotes a canonical enumeration of an r.e. set of axioms corresponding to the n -REA operator J . When X_e is an n -REA set we write $\mathcal{A}(X_e)$ for $\mathcal{A}(\mathcal{J}_e^n)$ (remember $X_e \stackrel{\text{def}}{=} \mathcal{J}_e^n(\emptyset)$)

In various cases we’ll further specify which of the many potential enumerations of axioms we mean to specify with $\mathcal{A}(X_e)$. For instance, we’ll explicitly define the enumeration of axioms for the n -REA sets we explicitly construct and rely on [lemma 2.3](#) to produce the corresponding n -REA set. We adopt the following definition to assist us in defining the axioms we enumerate in these construction.

Definition 2.5. The axiom $\langle \sigma' \rightarrow y \rangle$ **depends** on σ if $\sigma \prec \sigma'$. We say the axiom $\langle \sigma \rightarrow y \rangle$ (element y) is enumerated dependent on δ to mean we enumerate $\langle \sigma \cup \delta \rightarrow y \rangle$ $\langle \delta \rightarrow y \rangle$ into \mathcal{A} .

We will call an axiom $\langle \sigma \rightarrow y \rangle$ compatible (incompatible) with a partial function τ just if the partial function σ determining when the axiom applies is compatible (incompatible) with τ . We now argue that we can use an axiom set to approximate an n -REA set in well-behaved ways.

Definition 2.6. If \mathcal{A} is a set of n -REA axioms we say that $\chi \in \{0, 1, \uparrow\}^{<\omega}$ is \mathcal{A} supported just if for all m, x with $\chi(\langle m, x \rangle) \downarrow$ all of the following hold

- If $m = 0$ then $\chi(\langle m, x \rangle) = X(x)$.

- If $m > n$ then $\chi(\langle m, x \rangle) = 0$.
- If $0 < m \leq n$ then $\chi(\langle m, x \rangle) = 1 \iff (\exists \sigma, x)[\langle \sigma \rightarrow \langle m, x \rangle \rangle \in A \wedge \sigma \prec \chi]$

If χ is an $\mathcal{A}_s(J)$ supported approximation we say that χ is an s -supported⁴ approximation to $J(X)$.

Note that the notion of being s -supported is relative to a choice of a canonical enumeration of axioms for $J(X)$. Also, it's worth observing the following point.

Remark 2.7. Given any set of axioms \mathcal{A} and finite domain $D \subset \omega$ there is a unique \mathcal{A} -supported χ with domain D .

It turns out that for any n -REA operator J we can effectively find an enumeration of axioms \mathcal{A}_s and a computable functional Y_s^X with $\lim_{s \rightarrow \infty} \text{dom } Y_s = \omega$, such that Y_s^X is always an \mathcal{A}_s supported approximation to $J(X)$ and infinitely often $Y_s^X \prec J(X)$. A proof of this claim can be found in the appendix to give the reader an idea how this can work. However, we don't actually need the full strength of this result and will instead directly specify an approximation to the n -REA sets we are constructing which we will ensure has the following property (where A here is the n -REA set being constructed).

Condition 1. $(\exists^\infty s)A_s \prec A$.

In fact, our approximation to A (which we define below) will turn out to satisfy the stronger condition that there are infinitely many stages such that $(\forall t > s)A_s \prec A_t$. Note that, this implies that our approximation A_s is a Δ_2^0 approximation to A .

3. Construction Overview

We adopt the following notation for easier manipulation of n -REA sets defined in terms of columns pursuant to [definition 2.1](#).

Definition 3.1.

- $A \oplus B \stackrel{\text{def}}{=} \bigoplus_{n \in \omega} A^{[n]} \oplus B^{[n]}$.
- For A an n -REA(X) set we let $A \upharpoonright_{[s]} = \{\langle k, x \rangle \mid k \leq n \wedge x < s\}$.
- $l_s > 4$ is a number chosen large⁵ at the end of stage s .

Note that if both A and B are both n -REA then so is $A \oplus B$. Using these conventions we can now describe the broad outline of our proof of [theorem 1.4](#).

To establish the desired claim we'll need to build a properly 3-REA set A that can't be extended to a properly 4-REA set. In other words, for every $i \in \omega$ the set $A \oplus W_i^A$ must not be properly 4-REA. So in addition to the set A we'll build a sequence of 3-REA sets $A \oplus Y_i$ Turing equivalent to the sets $A \oplus W_i^A$ (note that by using $A \oplus Y_i$ rather than just Y_i we avoid the need for unnecessarily copying changes to A over to Y_i). To verify these equivalences we'll also need computable functionals so, in addition to the 3-REA set A we'll also build sets Y_i and functionals Γ_i, Θ to satisfy all of the following requirements where X_e denotes the e -th 2-REA set.

⁴Technically, being s -supported is relative to the choice of an enumeration of axioms but we will only use this notation when there is no ambiguity about the enumeration of the set $\mathcal{A}(J)$.

⁵Specifically, we will need that l_s is large enough that if x is mentioned at or before stage s then $l_s > \langle k, x + 2 \rangle, 0 \leq k \leq 3$ so that l_s is large enough to see the next two elements in each column.

$$\begin{aligned} \mathcal{P}_i: \quad & \Gamma_i(A \oplus W_i^A) = Y_i \wedge \Theta(Y_i) = W_i^A \\ \mathcal{R}_{j,e}: \quad & \Phi_j(A) \neq X_e \vee \Phi_j(X_e) \neq A \end{aligned}$$

We now observe that satisfying the above requirements is sufficient to prove the claimed theorem. However, the impatient reader may wish to jump ahead to the informal description of how $\mathcal{R}_{j,e}$ is met in [section 3.1](#). Note that the general method there will be familiar to anyone familiar with the methods in [\[1, 5\]](#) and broadly resembles the construction of properly n -r.e. sets (with some extra bookkeeping). Readers eager to jump ahead to the unique challenges and features of this construction should turn to [section 3.3](#) for an informal discussion or to the full construction in [section 4](#).

First we observe that our choice to avoid index profusion by using j as the index for both computations in $\mathcal{R}_{j,e}$ is harmless.

Lemma 3.2. *Suppose A and $Y_i, i \in \omega$ satisfy the requirements \mathcal{P}_i and $\mathcal{R}_{j,e}$ for all i, j, e then A is properly 3-REA but can't be extended to a properly 4-REA set.*

Proof. If \mathcal{P}_i is satisfied for all i it follows that the 3-REA set $A \oplus \overline{Y_i}$ is Turing equivalent to $A \oplus W_i^A$. It only remains to demonstrate that A is properly 3-REA.

Suppose, for a contradiction, that $A \equiv_T X_e$ for some 2-REA set X_e . Then either $A^{[3]} = \emptyset$, making A literally 2-REA, or $A^{[3]} \neq \emptyset$. In the first case $\mathcal{R}_{j,e}$ would fail when j is an index for the identity function and e an index for the empty set. In the second case suppose that the Turing equivalence is witnessed by $\Phi_{j_0}(A) = X_e$ and $\Phi_{j_1}(X_e) = A$ and z is the least element in $A^{[3]}$. Now let $\Phi_j(Z)$ be the functional which asks if $\langle 3, z \rangle \in Z$ and if not computes $\Phi_{j_1}(Z)$ and if so computes $\Phi_{j_0}(Z)$. Hence, if A is equivalent to some 2-REA set X_e it follows that some $\mathcal{R}_{j,e}$ isn't satisfied. \square

We will build A or Y_i by enumerating axioms into \mathcal{A} and $\mathcal{A}(Y_i)$ respectively which, by the remarks above, uniquely determines the corresponding sets. We will define approximations $A_s, Y_{i,s}$ to the sets A, Y_i and adopt the convention that an attempt to enumerate x into A (or Y_i) at stage s means enumerating x dependent on a large initial segment of the prior columns as defined by $A_s, Y_{i,s}$.

We now formally specify our approximation A_s and specify some properties it will be constructed to have.

Property 1.

- a) A_s is the unique s -supported approximation (i.e., \mathcal{A}_s -supported) with $\text{dom } A_s = \omega \upharpoonright_{[l_s]}$.
- b) If x is enumerated into $A^{[k]}$ at stage s then $x \notin A_{s-1}^{[k]}$ and x is enumerated dependent on $A_s^{[k']}$ for $k' < k$ (hence on prior columns of A up to height l_s).
- c) Axioms are only enumerated into \mathcal{A} at even stages and at most one axiom is enumerated per stage.

To avoid the potential circularity induced by the interaction of [definition 3.1](#) and [property 1b](#) we queue any requests made to enumerate elements made during stage s until after l_s is chosen at the end of the stage. Using $A_s^{[<n]}$ and l_s we can build axioms satisfying [property 1b](#) for elements queued for enumeration into $A^{[n]}$ which, in turn, lets us determine $A_s^{[<n+1]}$. Working inductively, this lets us identify a set of axioms for elements enumerated during stage s which satisfy both constraints.

We will also treat W_i^A as the result of an enumeration of axioms \mathcal{A}_s^i and define an approximation $W_{i,s}^A$ that satisfies the following properties.

Property 2.

- a) No axioms are enumerated into \mathcal{A}^i at even stages.
- b) At any stage s there is at most one i and x such that an axiom enumerating some x into W_i^A is enumerated and $i, x < s$.
- c) If an axiom enumerating x into W_i^A is enumerated at stage s then x is enumerated dependent on $A_s \upharpoonright_{[i]}$.
- d) $W_{i,s}^{A_s}$ denotes the unique finite partial function with domain $\omega \upharpoonright_{l_s}$ that is \mathcal{A}_s^i -supported.
- e) W_i^A is the set built by applying the axioms \mathcal{A}^i to the set A .

Property 2c ensures that only axioms which agree with our current approximation A_s to A are enumerated at stage s and are canceled by any change in A_s . Note, these properties are only possible to meet, as we will now verify, because of **condition 1**.

Lemma 3.3. *For all i there is an (uniformly specified) enumeration \mathcal{A}_s^i and an effective approximation $W_{i,s}^{A_s}$ satisfying **property 2**.*

Proof. We can assume we start with an enumeration of axioms $\widehat{\mathcal{A}}_s^i$ such that $x \in W_i^A$ iff $(\exists \langle \sigma \rightarrow x \rangle \in \mathcal{A}^i)(\sigma \prec A)$ and satisfies the usual rules for the enumeration of an A -r.e. set such as the enumeration of at most one element a stage etc. We now define \mathcal{A}_s^i in terms of the sets \mathcal{A}_{s-1}^i , $i < s$.

If s is even we do nothing. So assume s is odd and let $t < s$ be the minimal value such that we can find $i, y < s$, $|\sigma| < s$ such that $\langle \sigma \rightarrow y \rangle \in \mathcal{A}_t^i$, $\sigma \prec A_{s-1}$ and $W_{i,s-1}^{A_{s-1}}(y) = 0$. If no such values can be found do nothing. If they are found enumerate the axiom $\langle A_s \rightarrow y \rangle$ into \mathcal{A}_s^i . By construction, we clearly satisfy **properties 2a to 2d**. It remains to show that **property 2e** holds. By **condition 1** and **property 1c** we can assume that infinitely often we have an odd stage s with $A_{s-1} \prec A_s \prec A$, ensuring that eventually every element in W_i^A gets an axiom enumerating it. \square

We now give an overview of how each requirement operates.

3.1. Overview of $\mathcal{R}_{j,e}$

The basic approach to meeting $\mathcal{R}_{j,e}$ is to use the fact that A is 3-REA to change our minds about the behavior of some initial segment of A more times than X_e is able to track. We now give an informal description of the process which we depict graphically in **fig. 1**. In the figure stages at which a well-behaved back and forth computation exists are indicated by shaded/hatched regions. The region of A which is either hatched or shaded indicates the use of a computation of the shaded region of X_e , which in turn extends the use of a computation of the shaded region of A . We describe such computations as well-behaved when (at stage s) an s -supported region of A (hatched and shaded) computes (via Φ_j) X_e on (at least) an s -supported region (shaded region) which in turn computes (via Φ_j) $A(\langle 3, c \rangle)$. We leave cells blank if empty — unless they've just been canceled by an enumeration into a prior column, in which case we place a 0 in the cell. Finally, we circle locations which conflict with a commitment made by Φ_j at a prior stage.

We pick a value c at some stage s_{-1} which we hold out of $A^{[3]}$. If we never see a stage s_0 and a well-behaved computation from A to X_e and back then we are done. If we do see such a well-behaved computation then at stage $s_0 + 1$ we enumerate c into $A^{[3]}$ dependent on some large value staying out of $A^{[2]}$.

If at some stage $s_1 > s_0 + 1$ we again see a well-behaved back and forth computation (meaning X_{e,s_1} disagrees with X_{e,s_0} on the region shaded at stage s_0) we enumerate an element b into $A^{[2]}$ canceling c from A_{s_1+1} . This restores the computation from A to X_e seen at stage s_0 . Note that the only way we can again see a well-behaved back and forth




Figure 1. Meeting $\mathcal{R}_{j,e}$

computation (given we restrain enumeration into A) is if our approximation to X_e restores the (used part of the) state it had at s_0 by enumerating an element into $X_e^{[1]}$ to cancel any changes.

Now suppose that at some stage s_2 we again see a well-behaved back and forth computation. We respond by enumerating an element into $A^{[1]}$ at stage $s_2 + 1$ canceling the enumeration made at stage s_1 into $A^{[2]}$. This restores the computation from A to X_e seen at stage s_1 . But now X_e is unable to match this change as it can't cancel the element enumerated into $X_e^{[1]}$ and, as we will prove in [section 3.1.1](#), the fact that A computed an s -supported approximation to X_e at s_1 means this element was predicted by A to be out of X_e at s_1 . This ensures that, provided we restrain modifications to A , we never again see a well-behaved Φ_j computation from A to X_e and back, guaranteeing that $\mathcal{R}_{j,e}$ is satisfied.

We now formalize the argument made in this sketch and demonstrate that if the equivalence of A and X_e is witnessed by Φ_j then we will see infinitely many computations we've been calling well-behaved.

3.1.1. Positive Change Property

We now show that REA sets have the following positive change property. That is, the only way an approximation to an REA set can change is by enumerating a new element. We prove the results in this section for n -REA sets for arbitrary n , as this result will help us understand the strategy used to satisfy \mathcal{P}_i .

Lemma 3.4. *Suppose χ, χ' are s, s' -supported approximations to an n -REA(Z) set Y with $s < s'$ and $\text{dom} \chi' \supseteq \text{dom} \chi$. If $\chi(\langle m, x \rangle) = 1$ but $\chi'(\langle m, x \rangle) = 0$ then $\chi \upharpoonright_{m-1} \chi'$. Moreover, there is a y and $m' < m$ with $\chi(\langle m', y \rangle) = 0$ but $\chi'(\langle m', y \rangle) = 1$.*

Proof. To establish the first claim suppose that the hypotheses of the lemma are satisfied. By [definition 2.6](#) there is some axiom $\langle \sigma \rightarrow y \rangle \in \mathcal{A}_s Y$ with $y = \langle m, x \rangle$ and $\sigma \prec \chi$. As $\text{dom} \sigma \subset \omega^{[m-1]}$ if $\chi' \upharpoonright_{m-1} \chi$ then $\sigma \prec \chi'$ as $\text{dom} \chi' \supseteq \text{dom} \chi$. But then by [definition 2.6](#) we would have $\chi'(y) = 1$, contrary to the assumptions of the lemma.

We now prove the second claim by induction. As $Y^{[1]}$ is r.e., the claim obviously holds for $m = 2$. Now suppose the claim holds for any $m' \leq m$ and that the hypotheses of the lemma are satisfied for $m + 1$. Thus, $\chi \upharpoonright_m \chi'$. If there is some $y = \langle m', x \rangle$ with $m' < m + 1$ and $\chi(y) = 0$ and $\chi'(y) = 1$, we are done. If not there must be a y with $\chi(y) = 1$ and $\chi'(y) = 0$. The result now follows by the inductive hypothesis applied to $y = \langle m', x \rangle$. \square

We now give a slightly modified version of the above lemma that is specifically phrased in terms of undoing a previous change.

Lemma 3.5. *Suppose χ, χ', χ'' are s, s', s'' -supported approximations to $\mathcal{J}_e^n(Z)$ with $s < s' < s''$, $\chi'' \succ \chi$ and $\chi' \upharpoonright_m \chi$ then $\chi'' \upharpoonright_{m-1} \chi'$.*

Proof. Note that WLOG we may assume $\text{dom } \chi'' \supset \text{dom } \chi'$ by restricting χ' to the domain of χ . With this assumption made, the pair χ, χ' satisfy the hypotheses of lemma 3.4 or χ', χ'' do. As $\chi'' \succ \chi$ whichever way we have $\chi'' \upharpoonright_{m-1} \chi'$. \square

We now define a predicate which holds at x, s if it appears that A_s computes enough of X_e for X_e to correctly compute the value of $A_s(x)$. Note that, in what follows, we will make use of an approximation $X_{e,s}$ to X_e . However, somewhat surprisingly, all that matters about this approximation is that $X_{e,s}$ be s -supported and that, infinitely often, it's domain includes any finite subset. As such, we can simply take $X_{e,s}$ to be the unique s -supported approximation such that $\text{dom } X_{e,s}^{[1]} = \text{dom } X_{e,s}^{[2]} = l_s$ even if this approximation isn't even guaranteed to be (pointwise) infinitely often correct.

Definition 3.6. We define $\Psi_{j,e}(x, s)$ (written $\Psi(x, s)$ when j, e is clear from context) to hold just when there is some s -supported approximation ξ to X_e satisfying

- $\Phi_{j,s}(A_s) \succ \xi$
- $\Phi_{j,s}(\xi) \succ A_s \upharpoonright_{x+1}$

We note that if $\mathcal{R}_{j,e}$ fails we can always wait for a stage at which our approximation witnesses this failure.

Lemma 3.7. *If $\Phi_j(A) = X_e \wedge \Phi_j(X_e) = A$, $A_t \prec A$ then for any x there are infinitely many stages at which $\Psi(x, s)$ holds and is witnessed by A_s, ξ where $A \succ A_s \succ A_t$.*

Proof. Suppose A_t is as in the lemma. We first note that it is enough to show one such $s > t$ exists, since further witnesses may be generated by applying this result. Moreover, we may presume that $x \in \text{dom } A_t$ since, as the approximation to A is infinitely often correct, we can simply wait for some later stage at which $x \in \text{dom } A_t$ and $A_t \prec A$ then argue as below.

Since the reductions are total we can find ξ, χ such that $\xi \prec X_e$ is an $\mathcal{A}(X_e)$ supported approximation to X_e satisfying $(\forall y \leq x)(\Phi_j(\xi; y) = A(y) = A_t(y))$ and χ is an \mathcal{A} supported approximation to A satisfying $A \succ \chi \succ A_t$ with $\Phi_j(\chi) \succ \xi$. Now let $t' > t$ be large enough that every axiom needed to ensure ξ, χ are $\mathcal{A}X_e, \mathcal{A}$ supported has been enumerated by stage t' , $\Phi_{j,t'}(\xi; x) = A(x) = A_t(x)$, $\Phi_{j,t'}(\chi) \succ \xi$ and the domain of ξ is contained in $\{l, x \mid l, x \leq t'\}$. Finally, choose $s > t'$ so that $A_s \prec A$ and $\text{dom } A_s \supseteq \text{dom } \chi \cup \text{dom } A_t$ which guarantees $A_s \succ \chi$. But now note that all the conjuncts in definition 3.6 are satisfied so $\Psi(x, t)$ holds. \square

Using this definition we can make precise the idea of an approximation changing its mind.

Definition 3.8. An increasing sequence of $k + 1$ stages s_0, s_1, \dots, s_k is k -flipflopping (for Φ_j, X_e) at x denoted $\Psi_{j,e}^k(x, s_0, s_1, \dots, s_k)$ just if

- (1) $(\forall l \leq k - 1)(A_{s_l}(x) \neq A_{s_{l+1}}(x))$
- (2) $(\forall l \leq k - 1)(\Psi_{j,e}(x, s_l))$ is witnessed by some approximation ξ_l to X_e .
- (3) $(\forall l \leq k - 2)(A_{s_l} \prec A_{s_{l+2}})$

An element x is k -flipflopping at stage s (for X_e, Φ_j), denoted $\Psi_{j,e}^k(x, s)$ just if there is an increasing sequence of stages $s_0, s_1, \dots, s_{k-1}, s$ such that $\Psi^k(x, s_0, s_1, \dots, s_{k-1}, s)$.

When j, e are clear from context we will omit mentioning them.

We can now provide the framework for diagonalizing against X_e .

Lemma 3.9. *If $\Psi_{j,e}^3(x, s)$ then $\Psi_{j,e}(x, s)$ fails to hold.*

Proof. Let $s_3 = s$ and suppose, for a contradiction, that both $\Psi_{j,e}^3(x, s_0, s_1, s_2, s_3)$ and $\Psi_{j,e}(x, s_3)$ hold and are witnessed by the approximations $\xi_l, l \leq 4$ to X_e . By part 3 of [definition 3.8](#) and [definition 3.6](#) it follows that $(\forall l \leq k-2)(\xi_l \prec \xi_{l+2})$.

As $A_{s_0}(x) \downarrow \neq A_{s_1}(x) \downarrow$ we have $\xi_0 \upharpoonright_2 \xi_1$ by [lemma 3.5](#) we have $\xi_2 \upharpoonright_1 \xi_1$ and as $A_{s_1}(x) \downarrow \neq A_{s_2}(x) \downarrow$ we can apply [lemma 3.5](#) again to infer that $\xi_3 \upharpoonright_0 \xi_2$. But s -supported approximations to an n -REA(Z) can't disagree on column 0. Contradiction. \square

With this in mind we can now reiterate the basic strategy we would use to meet $\mathcal{R}_{j,e}$ if there were no other requirements. Choose some x and hold it out of $A^{[3]}$ until we observe a computation witnessing $\Psi(x, s)$, i.e., a computation of an s -supported approximation ξ of X_e which computes $x \notin A^{[3]}$. This stage becomes s_0 in the lemma above and ξ becomes ξ_0 . Now enumerate x into $A^{[3]}$ dependent on a very large initial segment of $A^{[2]}$ and wait until we again see $\Psi(x, s)$ with a witness whose domain contains the domain of ξ_0 . This new stage becomes s_1 and this new witness becomes ξ_1 . Enumerate an element into $A^{[2]}$ depending on a large initial segment of $A^{[1]}$ canceling the enumeration of x and now wait until we again see $\Psi(x, s)$ with $A_s \succ A_{s_0}$ and a witness whose domain extends that of ξ_1 . Repeat the cancellation one last time via an enumeration into $A^{[1]}$ for a guaranteed win. At each point at which we wait on a computation we restrain any elements from being enumerated into any initial segment appearing in a computation we've used in this process. The lemma above ensures that this process ends in a victory.

3.2. Overview of \mathcal{P}_i

To meet \mathcal{P}_i we must construct sets Y_i and functionals that witness $Y_i \oplus A \equiv_{\mathbf{T}} A \oplus W_i^A$. As we control the construction of the sets Y_i we will simply settle on a particular way of coding W_i^A into Y_i so we can share a single functional Θ computing W_i^A from Y_i . As elements can both enter and leave (our approximation to) W_i^A during the construction our coding mechanism must allow Y_i to change it's mind about the value of $W_i^A(x)$.

To this end we use the x -th column of $Y_i^{[3]}$ to encode whether or not x is in W_i^A . To guess that $x \in W_i^A$ we place $\langle x, 0 \rangle$ into $Y_i^{[3]}$, to revoke that guess we place $\langle x, 1 \rangle$ into $Y_i^{[3]}$, to guess x in W_i^A we place $\langle x, 2 \rangle$ into $Y_i^{[3]}$ and so on. To avoid any need to memorize the particular coding convention we define the following notation.

Definition 3.10.

$$[x]_i^s = \langle x, k \rangle \text{ where } k \text{ is the least element of } \omega \text{ such that } Y_{i,s-1}^{[3]}(\langle x, k \rangle) \downarrow = 0$$

When used without stage it indicates the least element of the form $\langle x, k \rangle$ not yet enumerated into $Y_i^{[3]}$.

On the other hand, as we don't control the set W_i^A , the only way we can take full advantage of any change in W_i^A is to build functionals Γ_i so that it always tries to map $A_s = A_s \upharpoonright_{l_s}$ to $Y_{i,s} \upharpoonright_s$ whenever $i < s$ but is reset whenever \mathcal{P}_i is reinitialized.

Definition 3.11.

$$(1) \Theta_s(Z; x) = \begin{cases} 1 & \text{if } 1 + \max\{k \mid Z(\langle 3, \langle x, k \rangle \rangle) \downarrow = 1 \vee k = -1\} \equiv 0 \pmod{2} \\ 0 & \text{otherwise} \end{cases}$$

- (2) Let s' be the last stage with $s' \leq s$ at which \mathcal{P}_i is reinitialized (injured). Then $\Gamma_{i,s}(X \oplus Z; y) \downarrow = Y_{i,t}(y)$ where t is the least stage satisfying $s \geq t \geq s', t > y, Y_{i,t}(y) \downarrow, X \succ A_t$ and $Z \succ W_{i,t}^A \upharpoonright_{y+1}$. If there is no such t then $\Gamma_{i,s}(X \oplus Z; y) \uparrow$

Note that **condition 1** ensures that $\Gamma_i(A \oplus W_i^A)$ is total as for every y there is a stage $s > y$ at which $A_s \prec A$ and $W_{i,s+1}^A \upharpoonright_{y+1} = W_i^A \upharpoonright_{y+1}$. Also, observe that if $x > s$ then Γ_i makes no commitments about x prior to stage s and thus any commitments made about x will be dependent on at least the initial l_s elements of each column of A .

We build Y_i by enumerating axioms much like we do for A but, remember it is only the columnwise sum $Y_i \oplus A$ which we build to be 3-REA not Y_i itself (this merely saves us the trouble of copying every enumeration into A over to Y_i) Thus, the axioms enumerating elements into Y_i can depend not only on prior columns of Y_i but on prior columns of A as well. We omit the straightforward modifications to the definition of axiom to allow for this dependence. As with A we will simply indicate which elements we wish to enumerate into Y_i and rely on the following properties to uniquely define what axiom is enumerated and how our approximation to Y_i is affected.

Property 3.

- $Y_{i,s}$ is the unique partial function with $\text{dom } Y_{i,s} = \{\langle k, x \rangle \mid k < l_s + 4 \wedge x < l_s\}$ such that $Y_{i,s} \oplus A_s$ is an s -supported approximation (i.e., $\mathcal{A}_s(Y_i \oplus A)$ -supported) (hence, $k > 3, k < l_s + 4, x < l_s \implies Y_{i,s}(\langle k, x \rangle) \downarrow = 0$).
- If x is enumerated into $Y_i^{[k]}$ at stage s then x is enumerated dependent on $A_s^{[k']}$ and $Y_{i,s}^{[k']}$ for $k' < k$ (hence on prior columns of both A and Y_i up to height l_s).

Our construction will ensure that the following condition is satisfied.

Condition 2. For all s , $\Theta_s(Y_{i,s}; x) = W_{i,s}^A(x)$ and $\Gamma_{i,s}(A_s \oplus W_{i,s}^A; x) = Y_{i,s}(x)$ whenever both sides are defined. Moreover, Γ_i is well-defined.

To this end, unlike the functional Γ_i , we don't reset the set of axioms enumerated into $\mathcal{A}(Y_i)$ when the module implementing \mathcal{P}_i is reinitialized since W_i^A isn't reinitialized.

3.3. Requirement Interaction

Note that we can regard $A \oplus W_i^A$ as (equivalent to) a 4-REA set so the same considerations about an $n + 1$ -REA set avoiding equivalence with an n -REA from **section 3.1** apply but now it's our job to ensure equivalence. Obviously, if we could leave $A^{[1]}$ empty (or even computable) then we could trivially meet \mathcal{P}_i simply by enumerating $[x]_i^s$ into $Y_i^{[3]}$ whenever we see z enter W_i^A (as we could wait until $A^{[1]}$ had settled) and simply copying $A^{[2]}, A^{[3]}$ into $Y_i^{[1]}$ and $Y_i^{[2]}$ respectively. However, meeting requirements of the form $\mathcal{R}_{j,e}$ forces us to make (non-computable) enumerations into all three columns of A .

Specifically, an opponent building W_i^A could try to duplicate the kind of argument we gave in **section 3.1**. If such an opponent could arrange for the approximation to $A \oplus W_i^A$ to flip-flop on some value z 4 times (waiting for new Γ_i, Θ computations each time) then we would have no way to change $Y_i \oplus A$ to meet \mathcal{P}_i . This situation is loosely depicted in **fig. 2** where the shaded region is meant to represent the region below l_s (and thus use and domain of Θ and Γ_i) while the hatched region in stage t_4 represents the use of the Γ_i computation which disagrees with Y_i .

For instance, our opponent might start by reserving some value z to be kept out of W_i^A . By waiting until we've committed to the behavior of our functionals on the appropriate initial segments our opponent could force a change to Y_i by enumerating z into W_i^A with some large use and wait until some stage at which our functionals again witness (enough of) the equivalence $A \oplus W_i^A \equiv_T Y_i \oplus A$. The proof of **theorem 1.3** from [1] shows that our opponent can ensure that sometimes we're forced to enumerate some y into $Y_i^{[2]}$ (to cancel a previous enumeration into $Y_i^{[3]}$) in




Figure 2. Piggybacking off $\mathcal{R}_{j,e}$

response to some requirement $\mathcal{R}_{j,e}$ enumerating c into $A^{[3]}$. That is, we can't avoid getting to stage t_2 as depicted in fig. 2.

3.4. Safety Via Agreement

If \mathcal{P}_i is of higher priority than $\mathcal{R}_{j,e}$ we need to be able to meet $\mathcal{R}_{j,e}$ in a way that doesn't allow an opponent to force an injury to \mathcal{P}_i . Our strategy will be to modify the way (the module responsible for meeting) $\mathcal{R}_{j,e}$ operates to ensure that when $\mathcal{R}_{j,e}$ enumerates b into $A^{[2]}$ we aren't forced to enumerate any element into $Y_i^{[1]}$, i.e., we avoid ever reaching the stage labeled t_3 in fig. 2. If we can arrange this, it removes our opponents 1 column advantage on us which will prevent $\mathcal{R}_{j,e}$ from injuring \mathcal{P}_i (the final enumeration of a into $A^{[1]}$ will thus keep $A \oplus Y_i$ and $A \oplus W_i^A$ in lock step).

The key idea here is that b 's entry into $A^{[2]}$ can only force us to enumerate an element into $Y_i^{[1]}$ if there is some (small) value y with $y \notin Y_{i,t_1-1}$ but $y \in Y_{i,t_3-1}$. In other words we will wait to enumerate b until the second (and first) column of Y_i are the same as they were immediately before c was enumerated.

As the enumeration of c into $A^{[3]}$ at stage s_0 will generally force a change in $Y_i^{[2]}$ and since we can't control W_i^A we can't ensure this condition is met for any particular value c . Instead, we make multiple attempts to meet $\mathcal{R}_{j,e}$ with the first attempt enumerating c_0 into $A^{[3]}$, the second attempt c_1 and so on and argue that for some $c = c_k$ we either meet $\mathcal{R}_{j,e}$ without canceling c or that we eventually see the agreement needed to allow safe enumeration of b into $A^{[2]}$.

Specifically, at stage s_{-1} we initialize (the module for) $\mathcal{R}_{j,e}$ choosing c_0 large and setting $c_k = c_0 + k$ (in the full construction $A^{[3]}$ will be partitioned between the various requirements to avoid collision). At stages $t_k = s_k - 1$ we observe an active stage for $\mathcal{R}_{j,e}$ and, if we haven't yet found agreement (i.e. some c_n which we can now cancel without changing the first two columns of any set Y_i), we respond at stage s_k by enumerating (only) c_k into $A^{[3]}$.

There are two critical aspects of this strategy. First, we ensure that c_{k+1} is small relative to the stage at which c_k is first enumerated so that each time we enumerate some c_k we are extending the use of Γ_i . Hence, commitments made about Γ_i after stage s_{-1} are canceled when we enumerate any c_k into $A^{[3]}$. Second, the fact that we choose c_0 large means that if $z \in W_{i,s_{-1}}^A$ then no enumeration of c_k will cancel z (and thus no reason to change how Y_i codes z 's

membership in W_i^A). Note that it will also be the case that each time we enumerate some c_k we will remove⁶ any z enumerated into W_i^A since stage s_{-1} . Thus, the only real constraint prior commitments impose on us while trying to produce the desired agreement is that we must remove $[z]_i^{s-1}$ from $Y_i^{[3]}$ *if present* whenever we enumerate any c_k into $A^{[3]}$.

The main difficulty in the construction will be to argue that we can always arrange our enumerations into Y_i during the intervals (s_{k-1}, s_k) so that we eventually find some c_n we can cancel without changing the first two columns of any Y_i . Stated in terms of the following definition we need to show that we can find k', k such that $s_{k'} - 1 \approx s_k - 1$. That is, there are active stages for $\mathcal{R}_{j,e}$ (or there are only finitely many such stages) $s_{k'} - 1 < s_k - 1$ such that canceling $c_{k'}$ by enumerating $b_{k'}$ at stage⁷ s_k (which requires us to return all Y_i to a state compatible with that at $s_{k'} - 1$) won't require changes in the first two columns of Y_i (the third column is returned to it's earlier state for free by the enumeration of $b_{k'}$).

Definition 3.12. We define

$$s \approx^i t \stackrel{\text{def}}{\iff} Y_{i,s}^{[\leq 2]} \upharpoonright_{l_s} \setminus Y_{i,t}^{[\leq 2]} \upharpoonright_{l_t}$$

$$s \approx t \stackrel{\text{def}}{\iff} (\forall i)(s \approx^i t)$$

We say that s, t **agree** just when $s \approx t$.

For future use we also adopt the following terminology (reflecting the fact that if t is accessible at stage s then there is some possible enumeration of elements into the sets Y_i that would allow some later stage $s' > s$ to satisfy $t \approx s'$).

Definition 3.13. A stage t is **accessible** at a stage $s \geq t$ just if for all i

$$Y_{i,s}^{[\leq 1]} \upharpoonright_{l_s} \succ Y_{i,t}^{[\leq 1]} \upharpoonright_{l_t}$$

A stage is (***i*-canceled**) **canceled** just if it isn't (*i*-accessible) accessible.

We also adopt the definition below. Note that active stages are the stages at which $\mathcal{R}_{j,e}$ takes action in response to seeing a longer Φ_j back and forth computation on it's own accord rather than merely responding to some enumeration into W_i^A .

Definition 3.14. A stage s_m at which $\mathcal{R}_{j,e}$ enumerates an element into $A^{[3]}$ or $A^{[2]}$ is a ($\mathcal{R}_{j,e}$) **active stage**. A stage t_k is called a ($\mathcal{R}_{j,e}$) **preactive** stage just if $t_k + 1$ is an active stage.

3.5. Simplified Agreement

Let's consider how (the module for) $\mathcal{R}_{j,e}$ might enact this strategy with respect to a single requirement \mathcal{P}_i assuming only a single value z enters or leaves W_i^A . On this assumption, we'll describe a winning strategy for $\mathcal{R}_{j,e}$ that enumerates at most two elements c_0, c_1 into $A^{[3]}$. This strategy is visualized in [figs. 3](#) and [4](#). Note that, while the figures depict c_0 and $[z]_i^{s-1}$ as being on the same level but in reality $[z]_i^{s-1}$ is smaller than c_0 which is why we shade a larger region of Y_i (where the shaded region indicates the region below l_s at the given stage).

If z doesn't enter W_i^A in the interval $[s_{-1}, s_0)$ when we enumerate c_0 we get $s_0 - 1 \approx^i s_1 - 1$ for free since there was never any need to cancel any element in $Y_i^{[3]}$ (this case isn't depicted).

⁶Remember, that we've used the fact that we will ensure $A_s \prec A$ infinitely often to allow us to set the 'use' of a computation placing $z \in W_i^A$ at stage s to l_s .

⁷Note that in full construction we assume that no elements enter any W_i^A at stages s_k so we need not worry about changes between $s_k - 1$ and s_k .




Figure 3. Satisfying \mathcal{R}_{je} respecting \mathcal{P}_i on z




Figure 4. Alternate behavior after stage s_0

In fig. 3 we see that z enters W_i^A by stage $s_0 - 1$ forcing us to cancel $[z]_i^{s_0-1}$ at stage s_0 . Note that when z first enters W_i^A we become free to code that entry into Y_i since the enumeration changes W_i^A on the use of $\Gamma_i (A \oplus W_i^A; [z]_i^{s_0-1})$. However, in the figure z isn't enumerated into W_i^A during $[s_0, s_1)$. We've circled the value which blocks agreement at stage $s_1 - 1$ with the earlier stage $s_0 - 1$. But now observe that at stage $s_2 - 1$ agreement is achieved for free since we had no need to cancel any elements at stage s_1 (with c_1 playing the role of c). As we will see later, even if we'd had other intervening stages as long as they didn't enumerate any elements into $Y_i^{[1]}$ small at stage $s_1 - 1$ we could cancel their contributions. Note that when we enumerate b_1 into $A^{[2]}$ it automatically removes $[z]_i^{s_0-1}$ from $Y_i^{[3]}$ as this was enumerated during the interval (s_1, s_2) (remember $A \oplus Y_i$ is 3-REA not Y_i).

In fig. 4 we depict the case where z is enumerated into W_i^A at some stage $s'_0 \in (s_0, s_1)$. At stage s'_0 (we can't wait until $s_1 - 1$) we are faced with a choice. We could simply enumerate $[z]_i^{s'_0-1}$ into $Y_i^{[3]}$ again but then the value we enumerated into $Y_i^{[2]}$ at stage s_0 would prevent us from achieving agreement between stages $s_0 - 1$ and $s_1 - 1$. So instead we choose to cancel the value enumerated into $Y_i^{[2]}$ and restore the state of Y_i at stage $s_0 - 1$ producing agreement.




Figure 5. Satisfying $\mathcal{R}_{j,e}$, \mathcal{P}_i when b_0 removes z from W_i^A

So far our examples have always shown z entering W_i^A before the enumeration of c_0 as this is the most challenging case to achieve agreement. However, it's useful to consider the case to illustrate two important points. First, to demonstrate why it's only agreement in the first two columns of Y_i which matters not agreement about $z \in W_i^A$ and why later enumeration into W_i^A doesn't threaten agreement once it's been achieved. This situation is depicted in fig. 5.

In this scenario z doesn't enter W_i^A before, at s_0 , we enumerate c_0 into $A^{[3]}$. As such, we don't enumerate $[z]_i^{s_0-1} = [z]_i^{s_0}$ before s_0 either so it need not be canceled at stage s_0 . Hence, at stage s_0 we already have agreement with stage $s_0 - 1$, i.e., $s_0 \approx^i s_0 - 1$. However, we may have to wait to see a $\mathcal{R}_{j,e}$ expansionary stage and during that time we may see z enter W_i^A . But, note that, once we've achieved agreement we can always respond to such an entry by enumerating $[z]_i$ (labeled above as $[z]_i^{s_0-1}$) into $Y_i^{[3]}$ without jeopardizing that agreement. Thus, at stage $s_1 - 1$ when we finally see a $\mathcal{R}_{j,e}$ expansionary stage we have $s_1 - 1 \approx^i s_0 - 1$.

Now, remember, that it's not actually Y_i that we've committed to building as a 3-REA set but $A \oplus Y_i$ (the columnwise sum). Hence, the enumeration of b_0 into $A^{[2]}$ doesn't just cancel c_0 but also $[z]_i^{s_0-1}$ (in fig. 4 we glossed over need to reenumerate this element when b_0 was enumerated). Thus, at s_1 we comply with the commitments made for \mathcal{P}_i at $s_0 - 1$. Of course, z could still decide to finally enter W_i^A at this point but, since z wasn't in W_i^A at stage $s_0 - 1$, we are no longer committed to keeping $[z]_i^{s_0-1}$ out of $Y_i^{[3]}$ allowing us to preserve the computation required by \mathcal{P}_i . Finally, if we enumerate a , we again rely on the fact that it's $A \oplus Y_i$ that's 3-REA to allow a to cancel both $[z]_i^{s_0-1}$ and b_0 returning both sides to stages compatible with those they had at stage $s_1 - 1$. Note that if, instead, z had been out of W_i^A at stage $s_1 - 1$ and then entered W_i^A after the enumeration of a we could respond by enumerating $[z]_i^{s_0-1}$ just as we did when z entered W_i^A after the enumeration of b_0 but before a .

3.6. General Agreement

In section 3.5 we demonstrated how to meet $\mathcal{R}_{j,e}$ without injuring a single requirement \mathcal{P}_i on the assumption that only a single value z ever enters or leaves W_i^A . However, in the general case we must meet $\mathcal{R}_{j,e}$ without injuring any

of the finitely many higher priority requirements $\mathcal{P}_{i_0}, \mathcal{P}_{i_1}, \mathcal{P}_{i_2}, \dots$. We must also accommodate arbitrarily many elements entering and leaving the sets $W_{i_n}^A$.

3.6.1. Multiple elements entering some $W_{i_0}^A$

First we explain the relatively easy task of dealing with multiple elements entering and leaving the sets $W_{i_n}^A$. By the discussion in [section 3.4](#) the enumeration of c_k cancels any commitments we've made about Γ_i after stage s_{-1} . Hence, if $z \geq s_{-1}$ then [definition 3.11](#) means that at s_{-1} we've yet to make any commitment about z (i.e. about the value of Γ_i on $[z]_i^t$). This leaves us with finitely many values we must be concerned with entering or leaving any W_i^A when trying to achieve agreement. Note that if we ever see $z < z'$ enter W_i^A after stage s_{-1} this cancels any commitments we've made regarding z' allowing us to freely code changes in $W_i^A(z')$ by enumerating elements into $Y_i^{[3]}$. Morally speaking, this means it's only the least value of z entering W_i^A in the interval $[s_k, s_{k+1} - 1]$ which matters so we need only tweak the strategy above to deal with multiple values entering/leaving some W_i^A .

More specifically, we eventually find $k' < k$ such that the least $z \notin W_{i, s_{k'} - 1}^A$ which is in some $W_{i, s_{k'} - 1}^A, k' \leq k'' < k$ also is in $W_{i, s_k - 1}^A$ or eventually all $z \notin W_{i, s_{k'} - 1}^A$ aren't in $W_{i, s_k - 1}^A$. In the later case we have $s_k - 1 \approx^i s_{k+1} - 1$ as we are in the same situation depicted in [fig. 3](#). In the former case, as in [fig. 4](#), we ensure $s_{k'} - 1 \approx^i s_k - 1$ by making enumerations into $Y_i^{[1]}$ when we see a new least $z \notin W_{i, s_{k'} - 1}^A$ enter W_i^A that cancels all changes since the last stage $s_{k''} - 1$ with $z \in W_{i, s_{k''} - 1}^A$ and making whatever corrections needed to code the state of $z' > z$ by enumeration into $Y_i^{[3]}$. Note that no problem is introduced if, in response to the entry of z' into W_i^A at stage $s \in (s_{k-1}, s_k)$ we make an enumeration into $Y_i^{[1]}$ to cancel all changes between $s_{k'} - 1$ and s . Such enumerations may prevent ever achieving agreement with some $s_{k''} - 1, k' < k'' < k$ (as we can never remove an element from $Y_i^{[1]}$) but that's not a problem since the canceled stages are ones in which no $z'' < z$ enters W_i^A .

3.6.2. Multiple \mathcal{P}_i requirements

This leaves us with the harder problem of dealing with multiple requirements $\mathcal{P}_0, \mathcal{P}_1, \dots, \mathcal{P}_{\langle j, e \rangle}$ at the same time.

Unlike the case with multiple values entering a single set, W_i^A here entry into W_i^A doesn't cancel any commitments made about $\Gamma_{i'}, i' \neq i$. Thus, when we see z enter W_i^A we must immediately decide how to code this change in Y_i (by enumerating into $Y_i^{[3]}$ or cancelling a removal from it by enumerating into $Y_i^{[1]}$) without yet knowing what elements may later enter some $W_{i'}^A$. We now describe how we handle this problem under the simplifying assumption that there is only a single value z which may enter any set W_i^A and $z \notin W_{i, s_{-1}}^A$ ⁸ where $i \leq \langle j, e \rangle$. As there is only a single value of z in play we will only ever need to make use of a single location $[z]_i$ to code whether or not $z \in W_i^A$ (i.e. if we see z leave W_i^A we always remove $[z]_i$ from $Y_i^{[3]}$) and WLOG we can assume that value is independent of i and just write $[z]$.

We formulate the problem slightly more generally as trying to meet $\mathcal{R}_{j, e}$ while respecting the $n + 1$ requirements $\mathcal{P}_{i_0}, \mathcal{P}_{i_1}, \dots, \mathcal{P}_{i_n}$ (i.e., assuming that agreement is automatic for i not in $\{i_0, \dots, i_n\}$) and then argue that we can reduce this to the problem of respecting n requirements. Since we've assumed that $z \notin W_{i_j, s_{-1}}^A, j \leq n$ at each stage s_k where we enumerate c_k into $A^{[3]}$ this has the effect of removing z from $W_{i_j, s_k}^A, j \leq n$. To ensure that the sets Y_{i_j} reflect this we enumerate l_t into $Y_{i_j}^{[2]}$ if $z \in W_{i_j, s_k - 1}^A$ where t is the stage at which $[z]$ entered $Y_{i_j}^{[1]}$ in the interval (s_{k-1}, s_k) . Thus, at each stage s_k z isn't in any of the sets $W_{i_j}^A$ nor are the elements $[z]$. Note that, in the case where $z \notin W_{i_j, s_k - 1}^A$ for all $j \leq n$ we don't enumerate any elements into Y_{i_j} at stage s_k meaning that we automatically have $s_k - 1 \approx s_k$ and, by the remarks at the end of [section 3.5](#), once established we can maintain this agreement until stage $s_{k+1} - 1$ as we only need to enumerate elements into the third column of the sets Y_{i_j} to respect the requirements $\mathcal{P}_{i_0}, \mathcal{P}_{i_1}, \dots, \mathcal{P}_{i_n}$.

⁸If $z \in W_{i, s_{-1}}^A$ then z won't leave W_i^A as a result of any enumeration we make in trying to meet $\mathcal{R}_{j, e}$ ensuring that coding the status of z isn't in tension with our attempt to meet $\mathcal{R}_{j, e}$.




Figure 6. Agreement Respecting \mathcal{P}_{i_0} and \mathcal{P}_{i_1}




Figure 7. Alternative agreements respecting \mathcal{P}_{i_0} and \mathcal{P}_{i_1}

What we must do is describe a strategy for reacting to the enumeration of z into $W_{i_j}^A$ during the interval (s_k, s_{k+1}) (the interval between c_k and c_{k+1} entering A ^[3]) that ensures we will eventually produce agreement. The key factor that determines how we act to try and produce agreement during the interval (s_k, s_{k+1}) is which of the sets $W_{i_0}^A, W_{i_1}^A, \dots, W_{i_n}^A$ we see z enter first. To this end, define u_m^j so that $(s_{u_m^j}, s_{u_m^j+1})$ is the m -th interval on which z first enters $W_{i_j}^A$.

Our argument is essentially a priority argument in which we give highest priority to producing agreement between intervals (s_k, s_{k+1}) in which the first set z enters is $W_{i_0}^A$. We give second highest priority to producing agreement between intervals in which z first enters $W_{i_1}^A$ and so on. Specifically, during the interval $(s_{u_0^0}, s_{u_0^0+1})$ we will attempt to produce agreement with the stages $s_{u_0^0+1}-1, s_{u_1^0+1}-1, \dots, s_{u_{m-1}^0+1}-1$ ignoring any intervening intervals in which z first entered some other r.e. set in our list. However, as when we service a higher priority requirement in a finite injury argument, when we first see z enter $W_{i_0}^A$ we abandon our attempts to produce agreement on intervals where z first enters some other r.e. set. Thus, during the interval $(s_{u_m^1}, s_{u_m^1+1})$ we will only attempt to produce agreement with the stages $s_{u_{m'}^1+1}-1, s_{u_{m'+1}^1+1}-1, \dots, s_{u_{m-1}^1+1}-1$ where m' is the least value such that there is no k with $s_{u_{m'}^1+1}-1 < s_{u_k^0} < s_{u_{m-1}^1+1}-1$. Similarly, in the interval $(s_{u_m^2}, s_{u_m^2+1})$ we only try to produce agreement with the stages $s_{u_{m'}^2+1}-1$ which have occurred since the last interval in which z first entered $W_{i_0}^A$ or $W_{i_1}^A$ and so on. Of course, if we already have agreement with some earlier stage s_k-1 (e.g. because z didn't enter any of our r.e. sets in the previous interval) we simply maintain it regardless of which set is entered first.

Thus, if s is the first stage in (s_k, s_{k+1}) at which we see z enter some $W_{i_j}^A$ with there is a (potentially empty) list of earlier stages⁹ $s_{k_0^s+1}-1, \dots, s_{k_r^s+1}-1$ which we will try to produce agreement with. If this list is non-empty then we immediately¹⁰ and enumerate $l_t, t = s_{k_r^s+1}$ into $Y_{i_j}^{[1]}$ to cancel all elements enumerated into Y_{i_j} at intervening stages. This enumeration also cancels the element enumerated into $Y_{i_j}^{[2]}$ at stage t and thus returns $[z]$ into $Y_{i_j}^{[3]}$.

This has the effect of ensuring that $s_{k_0^s+1}-1 \approx^{ij} s_{k_1^s+1}-1 \approx^{ij} \dots \approx^{ij} s_{k_r^s+1}-1 \approx^{ij} s$. Note that if we are only trying to respect a single requirement \mathcal{P}_{i_0} this strategy ensures we always achieve agreement. Either z doesn't enter $W_{i_0}^A$ during the interval (s_{-1}, s_0) in which case we achieve the free win mentioned above or during the interval (s_0, s_1) then the list of prior stages we are attempting to agree with is non-empty ensuring that $s_0-1 \approx s_1-1$.

Now we notice that the argument we just gave depends only on having a sequence of intervals (t'_q, t_q) with $t_{q-1} \leq t'_q \leq t_q$ such that at every stage t_q we have $z \notin W_{i_j}^A, j \leq n, [z] \notin Y_{i_j}^{[3]}$, all elements enumerated into Y_{i_j} between t_q and t'_{q+1} are larger l_{t_q} and that, if z enters $W_{i_j}^A$ in (t'_q, t_q) for some $j \leq n$ then enumerating l_{t_q} into $Y_{i_j}^{[1]}$ restores $Y_{i_j} \upharpoonright [l_{t_q-1}]$ to the state it had at stage t_q-1 . Finally, we observe that if $(t'_q, t_q), q \leq m$ is such that $t_q = u_{q+m'}^j, q+m' \leq m$ where, as above, z doesn't first enter any $W_{i_e}^A, e < j$ during any interval between $u_{m'}^j$ and u_m^j and t'_q is the stage at which z enters $W_{i_j}^A$ in the interval ending at stage t_q satisfies these properties. That is, $t_q-1, q < m-m'$ is the sequence of stages that we try to achieve agreement with when we see z first enter $W_{i_j}^A$ during the interval ending with $t_{m-m'}$. Thus, when we see z first enter $W_{i_j}^A$ in an interval we can then run the same strategy on the indexes in $\{i_0, \dots, i_n\} - \{i_j\}$ on these reduced intervals.

The notation gets a bit heavy in that last discussion so to make this concrete consider [fig. 6](#) where we give an example of trying to produce agreement while respecting the two requirements \mathcal{P}_{i_0} and \mathcal{P}_{i_1} . In the figure we've only shown the final stage of intervals and we've boxed (including the double box) the stages which correspond to intervals in which z first enters $W_{i_0}^A$. Note that, on the intervals ending at stage s_1, s_2 where z first enters $W_{i_1}^A$ we have $s_1-1 \approx^{i_1} s_2-1$ and the behavior of Y_{i_0} looks just like the strategy we would follow if we were dealing with only the requirement \mathcal{P}_{i_0} . However, when we hit the interval ending at s_3 in which z first enters $W_{i_0}^A$ this attempt is injured so in the interval ending s_4, Y_{i_1} starts afresh and doesn't try to create agreement with stage s_2-1 .

⁹For instance, if z first enters $W_{i_0}^A$ and $(s_k, s_{k+1}) = (s_{u_m^0}, s_{u_m^0+1})$ then $k_q^s = u_q^0, q < m$ and if z first enters $W_{i_1}^A$ and $(s_k, s_{k+1}) = (s_{u_m^1}, s_{u_m^1+1})$ then $k_q^s = u_{m'+q}^1, m' + q < m$.

¹⁰Stage $s+1$ since we interleave stages at which enumeration into A and sets r.e. in A are seen.

Furthermore, note that at the boxed stages (the last stages of intervals at which z first enters $W_{i_0}^A$) agreement works as if the unboxed stages don't exist. Thus, since we have $s_0 - 1 \approx^{i_0} s_3 - 1 \approx^{i_0} s_5 - 1$ just as in the case where we dealt with only a single requirement \mathcal{P}_{i_1} we eventually achieve agreement on that index as well ensuring $s_3 - 1 \approx s_5 - 1$. Note that in [fig. 7](#) we give two alternate ways that the interval ending at stage s_3 could have played out. On the left we see that if we had a long enough consecutive sequence of intervals in which z first entered Y_{i_1} that too would have produced agreement while, on the right, we see how we would have achieved agreement between the double boxed stages had z entered Y_{i_1} during the interval (s_2, s_3) after z entered Y_{i_0} .

Finally, one might wonder how, even though a smaller value of z entering $W_{i_j}^A$ subsumes larger values entering the same set how we can handle the case where we first see the larger values enter $W_{i_j}^A$ when we don't yet know if the smaller value will later enter. We can handle this the same way we do with multiple sets where seeing 0 enter $W_{i_0}^A$ first can injure seeing 1 first enter $W_{i_0}^A$ and so forth. Though, by the discussion above, we need not worry about the case where the larger number enters during an interval after the smaller number.

Now that we've outlined the general approach we give the full construction. The complexity involved in this construction largely reflects the difficulty involved in doing accurate bookkeeping for the approach described in this section in the case where we must deal both with achieving agreement with respect to multiple values of i and multiple elements entering and leaving the associated r.e. sets.

4. Full Construction

4.1. General Framework

The construction takes the form of a finite injury argument with the module \mathcal{P}_i tasked with meeting \mathcal{P}_i having priority $2i$ and the module $\mathcal{R}_{j,e}$ tasked with meeting $\mathcal{R}_{j,e}$ having priority $2\langle j, e \rangle + 1$. Modules of the form \mathcal{P}_i merely define Γ_i as specified in [definition 3.11](#) and thus have no direct effect on any set or module. In contrast, any time the module $\mathcal{R}_{j,e}$ acts all lower priority modules are reinitialized.

At any even stage s at most one requirement of the form $\mathcal{R}_{j,e}$ with priority at most $2s - 1$ acts while (at most one per stage) enumerations into W_i^A occur only at odd stages. In response to an enumeration into W_i^A we give each requirement (in decreasing order of priority) the chance to act and claim the enumeration (injuring lower priority requirements) and enumerate elements into Y_i in response. We apply the following rule for unclaimed enumerations.

Rule 4.1. If z enters W_i^A at stage s but goes unclaimed then we enumerate $[z]_i^s$ into $Y_i^{[3]}$.

4.2. Module $\mathcal{R}_{j,e}$

We break up the description of the module $\mathcal{R}_{j,e}$ into two parts. First, we describe the behavior of the module $\mathcal{R}_{j,e}$ at even stages when it may act of its own volition to modify A and possibly the sets Y_i . Then, later, we describe action of the module $\mathcal{R}_{j,e}$ at odd stages in response to an enumeration into a set W_i^A .

4.2.1. Even Stages

The module $\mathcal{R}_{j,e}$ tasked with meeting $\mathcal{R}_{j,e}$ has four states, $\uparrow, 0, 1, 2, 3$ where \uparrow indicates $\mathcal{R}_{j,e}$ has yet to execute after (re)initialization and the later numbers (roughly) indicate the number of times we've changed our mind about whether $c \in A^{[3]}$. We denote the state of the module $\mathcal{R}_{j,e}$ at the end of stage s by $\mathcal{S}_s(\mathcal{R}_{j,e})$ and assume that the state remains the same at stage $s + 1$ unless otherwise noted. Note that, in state 1 we've enumerated at least one potential value for c_k for c into $A^{[3]}$ but we may not yet have enumerated the ultimate value c takes into $A^{[3]}$. If $\mathcal{R}_{j,e}$ is injured at stage s we set $\mathcal{S}_s(\mathcal{R}_{j,e}) = \uparrow$.

We describe the behavior $\mathcal{R}_{j,e}$ at stage s where we may assume that the following all hold.

- s is even, $\langle j, e \rangle < s$ and no higher priority requirement has chosen to act (injuring $\mathcal{R}_{j,e}$) at stage s , i.e., we execute this module at stage s .

- $s_{-1} \leq s$ is the first even stage after the most recent reinitialization of $\mathcal{R}_{j,e}$.
- The element $c_k, k > 0$ is defined to be $\langle \langle j, e \rangle, v + k \rangle$ if $c_0 = \langle \langle j, e \rangle, v \rangle$.
- c_k is enumerated into $A^{[3]}$ at stage $s_k = t_k + 1 > s_{-1}$ and m is the smallest value for which s_m is undefined and less than s (thus c_m is the next element waiting to be enumerated into $A^{[3]}$).
- $b_k, k < m$ are values chosen large at stage s_k satisfying $b_k \notin A_{s_k}^{[n]}$ and $b_k \notin Y_i^{[n]}$ for any i and $n \leq 2$. Note that this implies $l_{s_k-1} < b_k < l_{s_k}$.

Procedure 1 (Even Stages of $\mathcal{R}_{j,e}$).

Case $\mathcal{S}_{s-1}(\mathcal{R}_{j,e}) = \uparrow$: We act by choosing $c_0 = \langle \langle j, e \rangle, v \rangle$ where v is chosen large and setting $\mathcal{S}_s(\mathcal{R}_{j,e}) = 0$.

Case $\mathcal{S}_{s-1}(\mathcal{R}_{j,e}) = 0$: If $\Psi(c_0, s-1)$ holds then we act by enumerating c_0 into $A^{[3]}$, executing **procedure 2** and setting $\mathcal{S}_s(\mathcal{R}_{j,e}) = 1$

Case $\mathcal{S}_{s-1}(\mathcal{R}_{j,e}) = 1$: We execute the following steps.

Step 1: If $\Psi(c_m, s-1)$ fails to hold end the stage without acting. Otherwise the module acts by executing the subsequent steps.

Step 2: If there is $k < m$ with $s_k - 1 \approx s - 1$ perform the following steps and end the stage.

Step 2a: Choose $a \notin A^{[1]}$ large.

Step 2b: Set $c = c_k, b = b_k, \hat{s}_0 = s_k - 1, \hat{s}_1 = s - 1$.

Step 2c: Enumerate b into $A^{[2]}$ and set $\mathcal{S}_s(\mathcal{R}_{j,e}) = 2$

Step 3: If there is no such k we instead enumerate c_m into $A^{[3]}$ and execute **procedure 2**

Case $\mathcal{S}_{s-1}(\mathcal{R}_{j,e}) = 2$: If $\Psi(c, s-1)$ the module acts by enumerating a into $A^{[1]}$ and setting $\mathcal{S}_{s-1}(\mathcal{R}_{j,e}) = 3, \hat{s}_2 = s - 1$.

Case $\mathcal{S}_{s-1}(\mathcal{R}_{j,e}) = 3$: Once in state 3 the module never acts again.

We now specify the procedure we referenced above when we enumerated c_k . As we want to maintain **condition 2** (correctness of our functionals) at all active stages we must return Y_i , by potentially enumerating an element into $Y_i^{[2]}$, to a state compatible with $Y_{i,s-1}$. However, what, if any, element must be enumerated into $Y_i^{[2]}$ at s_k is determined by how we choose to respond to enumerations into W_i^A so we hand that task of keeping track of that value to the machinery which responds to those enumerations. Specifically, for each $i \in \omega$ we define (relative to $\mathcal{R}_{j,e}$) a marker $\mathcal{L}_i \in \omega \cup \{\uparrow\}$ with initial value $\mathcal{L}_{i,s-1-1} = \uparrow$. At active stages we'll just trust that we've placed this marker on the right value and do the following.

Procedure 2 (Resetting Y_i). Execute the following steps when called:

Step 1: For every $i < \langle j, e \rangle + 1$ enumerate \mathcal{L}_i into $Y_i^{[2]}$ if $\mathcal{L}_i \downarrow$ and update \mathcal{L}_i to \uparrow .

Step 2: If $i > \langle j, e \rangle$ or $x \geq s_{-1}$ then **at the end of the stage** enumerate $[x]_i^s$ into $Y_i^{[3]}$ if we would otherwise have $\Theta(Y_{i,s}; x) \downarrow \neq W_{i,s}^{A_s}(x) \downarrow$.

Note that unless $i, x < s$ no elements have ever been enumerated into W_i^A so we need take no action to maintain agreement between $W_{i,s}^{A_s}$ and $\Theta(Y_{i,s})$.

4.2.2. Odd Stages

It's now our task to ensure that there are two preactive stages t, t' that agree in the sense $t \approx t'$. Since we may injure lower priority modules we have no obligation to undo enumerations into $Y_i, i \geq \langle j, e \rangle + 1$ at active stages as we do with $Y_i, i < \langle j, e \rangle + 1$. Thus, we can simply leave enumerations into $Y_i, i \geq \langle j, e \rangle + 1$ to lower priority modules (or leave them unclaimed) without concern they will block us from achieving the desired agreement. On the other hand, if $z > s_{-1}$ then the use of Γ_i for any element coding the status of z is large enough that we can redefine it every time some c_m is enumerated into $A^{[3]}$ by $\mathcal{R}_{j,e}$. Finally, once $\mathcal{R}_{j,e}$ enters state 2 we are too far along in the process for changes in $Y_i^{[3]}$ to migrate all the way to the first column in time to cause a problem. We only claim those finitely many enumerations which don't fall into these easy cases.

Rule 4.2. If $z \setminus_{s} W_i^{A_s}, i < \langle j, e \rangle + 1$ then $\mathcal{R}_{j,e}$ claims z just if $\mathcal{S}_{s-1}(\mathcal{R}_{j,e}) \in \{0, 1\}$ and $z < s_{-1}$.

Our active management in these cases is complicated by the fact that we may only modify the elements in Y_i coding the status of $z < s_{-1}$ immediately in response to the enumeration of z . Our basic approach to deal with this problem is to use the order in which elements are enumerated into the sets W_i^A during an interval $[s_k, s_{k+1} - 1]$ to determine our response. However, our approach is only easily described in a recursive fashion. To that end we abstract away from the particular rule we will use to respond to enumeration and define the notion of a strategy. A strategy is a procedure with persistent state (i.e., a coroutine) that tells us how best to respond to an enumeration of z into the set W_i^A given some finite set S of stages we are trying to produce agreement with (or force future agreement).

We now define the notion of a procedure whose task is to tell us how to respond to an enumeration of z into W_i^A at stage s .

Definition 4.1. A strategy \mathcal{S} is a computable function $\mathcal{S}(s, z, i)$ (subroutine) with persistent state which enumerates a finite set of elements for entry into Y_i and recommends an update to \mathcal{L}_i . The persistent state of $\mathcal{S}(s, z, i)$ consists of a number of variables which $\mathcal{S}(s, z, i)$ updates and whose values persist across calls. In addition to the explicitly given arguments and persistent state we also assume that $\mathcal{S}(s, z, i)$ has access to a complete history of enumerations/recommendations made¹¹ prior to its execution.

We will describe the behavior of a strategy by giving a procedure by which it updates its persistent internal variables (initially set to \uparrow) and leave it to the pedantic reader to translate such a description into a fully formal object. We also need terminology for the stage (if any) in the intervals (s_{k-1}, s_k) at which control is first passed to a particular strategy as follow.

Definition 4.2. A stage s is an **initial enumeration** for \mathcal{S} (relative to $\mathcal{R}_{j,e}$) if \mathcal{S} is executed at stage s in response to the enumeration of z into W_i^A and if there is some k such that $s_{k-1} < s < s_k$ and \mathcal{S} was not executed during the interval (s_{k-1}, s) . In such cases we also say that the initial enumeration for \mathcal{S} during the interval (s_{k-1}, s_k) was into W_i^A .

Our goal is to prove that, given any value of s_{-1} we can effectively produce a strategy \mathcal{S} which, if followed whenever any $z < s_{-1}$ enters some $W_i^A, i \leq \langle j, e \rangle$, guarantees that after some finite number of preactive stages we'll succeed in producing agreement.

Specifically, we'll show that for any $h : \omega \mapsto \omega$ with compact support there is a strategy \mathcal{S}^h such that \mathcal{S}^h is capable of producing agreement while respecting the requirements \mathcal{P}_i for $i \in \text{supp } h$ provided that the only elements z entering or leaving W_i^A satisfy $h(i) > z$. We will then argue that if for every h' with $\int h' < \int h$ (relative to the counting measure) $\mathcal{S}^{h'}$ is guaranteed to win then so is \mathcal{S}^h . As when h is the zero function, there are no values of i, z with $h(i) > z$. \mathcal{S}^0 clearly satisfies the above assumption the desired result will follow by induction on $\int h$.

First, however, we describe the procedure \mathcal{S}^h . In this description, we style variables which persist across calls in Fraktur, e.g., τ , and adopt the computer science convention that we specify temporary values via 'let' while we update a variable in the strategy's internal state with 'set'.

¹¹Including enumerations/updates that some calling strategy has already committed to recommending but haven't yet been made.

Definition 4.3. The strategy \mathcal{S}^h for $h \in \omega^\omega$ with $\text{supp } h = I = \{i_0, \dots, i_{N-1}\}$ is defined to be the strategy with the following persistent internal variables (all of which are initialized to \uparrow) which executes [procedure 3](#) when called as¹² $\mathcal{S}^h(s, z, i_p)$.

- $m_r^h \in \omega, r < N$ which records the greatest non-canceled (i.e. accessible) preactive stage t_k at which initial enumeration for \mathcal{S}^h was made into $W_{i_r}^A$. Note that we may assume that m_r actually records the value of k rather than t_k behind the seems and uses k to return t_k so that we may assign t_{m+1} to m_r even when that stage hasn't yet happened.
- $\mathfrak{S}_r^h, r < N$ holds the current state of the substrategy tasked with producing agreement in response to initial enumerations made into $W_{i_r}^A$.
- $\mathfrak{b}_{i_r}^h, r < N$ identifies the value, if any, needed to cancel the values in Y_{i_r} coding the status of $h(i_r) - 1$ (the only value \mathcal{S}^h is directly responsible for) at the next active stage. \mathcal{S}^h will recommend that \mathfrak{c}_{i_r} be set to the least of $\mathfrak{b}_{i_r}^h$ and the recommendations of any sub-strategies.

In what follows we drop the superscript h from persistent variables when clear from context.

Procedure 3. Suppose that,

- $h \in \omega^\omega \wedge \text{supp } h = I = \{i_0 < i_1 < \dots < i_N\}$.
- m is largest with $t_m + 1 = s_m < s$.
- $z < h(i_p), p \leq N$
- $z \searrow_{>s} W_{i_p}^A$

then $\mathcal{S}^h(s, z, i_p)$ behaves as follows when executed at stage s . Note that $\mathcal{S}^h(s, z, i_p)$ always returns the current value of \mathfrak{b}_{i_p} , when it ceases execution as it's recommendation for \mathfrak{c}_{i_p} .

Step 1: If we project¹³ $t_m \approx s$ Enumerate $[z]_{i_p}^s$ into Y_{i_p} and exit any strategy (even parent strategies calling this one). There is no need to recommend an update to \mathfrak{c}_{i_p} , since we are guaranteed agreement.

Step 2: If s is an initial enumeration for \mathcal{S}^h set $\mathfrak{b}_{i_r}, r < N$ to be undefined and execute the following steps:

Step 2a: Set $\tau = p$ and for all $r, N > r > p$ set m_r and \mathfrak{S}_r to \uparrow .

Step 2b: Let n be such that $t_n = m_p$ and set m_p equal to t_{m+1} (yes, in the future). For all $r < p$ if $m_r = \uparrow$ set $m_r = t_{m+1}$.

Step 2c: If \mathfrak{S}_p is undefined set $\mathfrak{S}_p = \mathcal{S}^{h_p}$ with all persistent variables initialized to \uparrow where

$$h_p(i) \stackrel{\text{def}}{=} \begin{cases} h(i) - 1 & \text{if } i = i_p \\ h(i) & \text{otherwise.} \end{cases} \quad (3)$$

Step 2d: If m_p was undefined at the start of execution then do the following. If $z = h(i_p) - 1$ mark $[z]_{i_p}^s$ for enumeration into Y_{i_p} and set \mathfrak{b}_{i_p} to a large value returning that as our recommendation for \mathfrak{c}_{i_p} . If $z < h(i_p) - 1$ then execute $\mathfrak{S}_p(s, z, i_p) = \mathcal{S}^{h_p}(s, z, i_p)$ and let our recommendation for \mathfrak{c}_{i_p} be whatever was returned by $\mathfrak{S}_p(s, z, i_p)$. In either case, cease all further execution and return to caller.

Only if both \mathfrak{S}_p and m_p were defined at the start of execution will we pass beyond this point.

¹²That is, $\mathcal{S}^{\text{arg}_0}(\text{arg}_1, \text{arg}_2, \text{arg}_3)$ follows the steps in [procedure 3](#) with $h = \text{arg}_0, s = \text{arg}_1, z = \text{arg}_2, i_p = \text{arg}_3$.

¹³Projecting on the (valid) assumption that all recommendations for enumeration committed to by strategies executing before this one during stage s will be made.

Step 2e: Enumerate the least $q \notin Y_{i_p}^{[1]}$ satisfying $l_n < q < l_{n+1}$ into $Y_{i_p}^{[1]}$. This has the effect of returning Y_{i_p} to the state it was in at stage m_p .

Note that, if we've previously enumerated a $q' \in (l_n, l_{n+1})$ at some stage $t \geq t_n + 1$ it is enough to enumerate $q < l_t$ so we need not worry about running out of values but for simplicity we'll assume¹⁴ that we always have $l_n < q < l_{n+1}$.

Step 2f: $\boxed{\text{If } (\exists x < h(i_p) - 1) (x \in W_{i_p, t_n}^{A_{i_p}} \setminus W_{i_p, s-1}^{A_{i_p}})} \text{ place } \mathcal{A}_{i_p, t_n} \text{ into } Y_{i_p}^{[2]}$.

This has the effect of canceling (as if we were the active stage s_n) the elements in $Y_{i_p}^{[3]}$ which code the membership of those $x < h(i_p) - 1$. Note that this enumeration may be canceled by a sub-strategy at this very stage.

Step 2g: $\boxed{\text{If } h(i_p) - 1 \in W_{i_p, t_n}^{A_{i_p}} \setminus W_{i_p, s-1}^{A_{i_p}}}$ set $\mathfrak{b}_{i_p}^h$ to $\mathfrak{b}_{i_p, t_n}^h$ as at the next active stage s_{m+1} we must be prepared to cancel the enumerations into $Y_{i_p}^{[2]}$ we restored by the enumeration of q .

Step 3: $\boxed{\text{If } p \neq \tau \vee z + 1 < h(i_p)}$ Execute $\mathfrak{S}_\tau(z, i_p)$ (updating the internal state) and pass on the set of elements to be enumerated unmodified. Let b be the recommendation for \mathcal{A}_{i_p} , returned by the sub-strategy just executed.

Step 4: Let $V = \{\langle \sigma_0 \rightarrow \langle 3, y_0 \rangle \rangle, \langle \sigma_1 \rightarrow \langle 3, y_1 \rangle \rangle, \dots\}$ be the set of axioms projected to be active at the end of stage s such that y_k has the form $\langle h(i_p) - 1, x \rangle$ but $y_k \notin Y_{i, s-1}^{[3]}$. If V is empty then set $\mathfrak{b}_{i_p}^h$ to \uparrow . Otherwise, let t be the least stage at which one these axioms is enumerated and t_p the least preactive stage after t . If there is no such t_p , i.e., $t_p = t_{m+1}$, then leave $\mathfrak{b}_{i_p}^h$ unchanged. Otherwise, set $\mathfrak{b}_{i_p}^h$ to $\mathfrak{b}_{i_p, t_p}^h$.

Step 5: $\boxed{\text{If } p = \tau \wedge z = h(i_p) - 1}$ Check if, absent this step we are on track to have $\Theta(Y_{i_p, s}; h(i_p) - 1) \downarrow \neq W_{i_p, s}^{A_{i_p}}(h(i_p) - 1) \downarrow$. If not enumerate $[h(i_p) - 1]_{i_p}^s$ into Y_{i_p} and if $\mathfrak{b}_{i_p}^h$ is undefined set $\mathfrak{b}_{i_p}^h$ to be large.

Step 6: Return to caller with recommendation for \mathcal{A}_{i_p} , equal to the minimum of b and $\mathfrak{b}_{i_p}^h$, (always passing through any requests for enumeration).

We can now give the rule for responding to a claimed enumeration using the following specification

$$h^{j,e}(i) \stackrel{\text{def}}{=} \begin{cases} s-1 & \text{if } i < \langle j, e \rangle + 1 \\ 0 & \text{otherwise} \end{cases}$$

We presume that we initialize a strategy of the form $\mathcal{S}^{h^{j,e}}$ at stage s_{-1} and adopt the following rule.

Rule 4.3. If $z \searrow_s W_i^{A_s}$, and $\mathcal{R}_{j,e}$ claims z then execute $\mathcal{S}^{h^{j,e}}(s, z, i)$ and follow it's recommendations.

Note that, whenever we say things like “execute $\mathcal{S}^{h^{j,e}}(S, z, i)$ ” there is an ambiguity between the procedure which might be instantiated multiple times and the particular instance. However, we'll see in part 4 of lemma 5.3 that in any context each strategy will only ever be instantiated once so there is no risk of confusion.

¹⁴An assumption that will be vindicated by the calculation of an explicit bound on the number of preactive stages before agreement and thus on the total number of strategy executions which in turn gives us an explicit bound on the number of values we may need to enumerate between l_n and l_{n+1} .

5. Verification

We now demonstrate that the construction given above satisfies the requirements. In what follows we will frequently prove results about the strategy \mathcal{S}^h by either via induction or descent on strategies so we remind the reader that relation between strategies which holds whenever \mathcal{S}^h calls a strategy \mathcal{S}^{h_r} is a well-founded relation. To see this note that under the counting measure on ω , the integral is a functional from ω^ω to ω and for each r we have $\int h > \int h_r$.

To facilitate these arguments we've defined \mathcal{S}^h so that when we initially execute \mathcal{S}^{h_r} the configuration \mathcal{S}^{h_r} sees (i.e., Y_i) resembles that seen by \mathcal{S}^h (and indeed present at stage s_{-1}). However, to make use of this we need yet more notation.

Definition 5.1. If the strategy \mathfrak{S} is executed at stage s then let $Y_{i,s}^{\mathfrak{S}-}$ denote the state of Y_i at the start of execution of \mathfrak{S} (including any enumerations scheduled by strategies prior to this point) and let $Y_{i,s}^{\mathfrak{S}}$ denote the state of Y_i at the end of execution of \mathfrak{S} . When \mathfrak{S} instantiates \mathcal{S}^h we also denote these values by $Y_{i,s}^{h-}$ and $Y_{i,s}^h$ respectively.

Note that we'll continue to use notation like t_k which is defined relative to a requirement $\mathcal{R}_{j,e}$ without making the dependence on j, e explicit when it is obvious from context which requirement $\mathcal{R}_{j,e}$ is relevant.

5.1. Correct Computations

Our main goal in this section is to prove that [condition 2](#) holds. However, we first must prove that $Y_{i,s_k} \succ Y_{i,s_{-1}} \upharpoonright_{[s_{-1}]}$. But we need a number of utility results first, some of which depend on the very fact to be proved. We therefore present the lemma below but hold off on a proof until we can gather up all the inductive hypotheses needed.

Lemma 5.2. Suppose that $\mathcal{R}_{j,e}$ isn't initialized between s_{-1} and s_k and $\mathcal{S}_{s_k}(\mathcal{R}_{j,e}) < 2$ then

- (1) If $i \leq \langle j, e \rangle$ and $x < s_{-1}$ we have $Y_{i,s_{-1}}^{[3][x]} \prec Y_{i,s_k}^{[3][x]}$.
- (2) If s_k is the greatest active stage less than s and $Y_{i,s}^h$ is the result of enumerating $\mathcal{C}_{i,s}$ into $Y_{i,s}$, $i \leq \langle j, e \rangle$, $x < s_{-1}$ then $Y_{i,s}^{[3][x]} \succ Y_{i,s_{-1}}^{[3][x]}$.

We now prove our utility results, some of which assume that [lemma 5.2](#) holds at earlier stages.

Lemma 5.3. Suppose that \mathfrak{S} instantiates \mathcal{S}^h and receives an initial enumeration of z into W_i^A at stage $t \in (s_k, s_{k+1})$ (note that s_{k+1} is possibility infinite) then

- (1) If x is enumerated into W_i^A at stage $s \in (s_k, s_{k+1})$ then \mathfrak{S} is executed to handle this enumeration iff $x < h(i')$.
- (2) At no stage in (s_k, s_{k+1}) does any strategy other than \mathfrak{S} enumerate elements into $Y_i^{[3][h(i)-1]}$.
- (3) At no stage in (s_k, s_{k+1}) does any strategy not called from \mathfrak{S} enumerate elements into $Y_i^{[3][z]}$.
- (4) No other strategy implementing \mathcal{S}^h executes during (s_k, s_{k+1}) .
- (5) If \mathfrak{S} executes stage [step 2e](#) enumerating q with $l_n < q < l_{n+1}$ into $Y_i^{[1]}$ then no $q' < l_n$ has been enumerated since stage t_n .

If we further assume that [lemma 5.2](#) holds for $s_{k'}, k' \leq k$ and $x < h(i')$ then we may also conclude.

- (6) For all $k' \leq k$, $x < s_{-1}$ and $s \in (s_{k'}, s_{k'+1})$ if $(\forall x' \leq x) (x' \notin W_{i',s}^A \setminus W_{i',s_{-1}}^A)$ then $Y_{i',s_{-1}}^{[3][x]} \prec Y_{i',s}^{[3][x]}$.
- (7) For all i' , x if $x < h(i')$ then $Y_{i',s_{-1}}^{[3][x]} \prec Y_{i',t}^{h-[3][x]}$. In other words, initial enumerations to strategies start with a blank slate.

Proof. We prove these claims by descent. For claims 1, 2, 3 and 4 we observe they clearly hold for $h = h^{j^e}$ and we now prove that these claims also hold for the strategies $\mathfrak{S}_\tau = \mathcal{S}^{h_\tau}$ called from \mathfrak{S} .

To see that 1 holds note that the only time \mathfrak{S} doesn't pass on z is when $z = h(i) - 1$ and that by eq. (3) this is the only element which isn't below h_τ . To see that 2 (and, thus, 3) holds note that only step 5 ever enumerates elements and then only if $z = h(i_p) - 1$. \mathfrak{S}_τ is passed any element $x < h_r(i')$ as the only time x received by \mathfrak{S} and \mathfrak{S} only enumerates into $Y_i^{[\leq 2]}$ before executing \mathfrak{S}_τ for the first time.

To see that 5 holds it is enough to observe that if the first time after reinitialization that \mathfrak{S} runs is $s \in (s_{k'}, s_{k'+1})$ (with $s_{k'+1}$ possibly infinite) then neither \mathfrak{S} nor any strategy called from it will enumerate elements below $l_{s_{k'}}$. Then observe that if we ever run \mathfrak{S}_r with $r < p$ we reinitialize \mathfrak{S}_r so that if we had run some strategy that wasn't called by \mathfrak{S} but enumerated a value below l_{t_n} we would have reinitialized \mathfrak{S} after t_n contradicting the choice of q . The conclusion follows by observing that, after stage t_n , the first time \mathfrak{S} is run it enumerates q before any sub-strategy it calls enumerates anything.

To prove 6 note that the base case holds by the assumption of lemma 5.2 and we can ignore any stages at which no enumeration into $W_{i'}^A$ is claimed by $\mathcal{R}_{j,e}$. If $\mathcal{S}^{h'}(s, y, i')$ is executed at s then $h'(i') - 1 \geq y > x$ so direct enumeration can't cause disagreement. This leaves only enumeration into $Y_{i'}^{[\leq 2]}$ as a threat. However, when this happens step 2e restores the state as of t_n and by the inductive assumption (with t_n replacing s and s_{n-1} replacing $s_{k'}$) we can conclude that if no $x' \leq x$ are in $W_{i',s}^A \setminus W_{i',s-1}^A$ then $Y_{i',s-1}^{[3][x]} \prec Y_{i',s}^{[3][x]}$. Conversely, if there is such an x' then step 2f enumerates \mathcal{L}_{i',t_n} into $Y_{i'}^{[2]}$ which, by 5 ensures that $Y_{i',s-1}^{[3][x]} \prec Y_{i',s}^{[3][x]}$ by the assumption of lemma 5.2.

With item 6 established we can now proceed to prove 7 in exactly the same manner. Note that 7 clearly holds if $h = h^{j^e}$ by our assumption that lemma 5.2 holds at s_k . For the inductive case we need merely replace the assumption that no $x' \leq x$ are in $W_{i',s}^A \setminus W_{i',s-1}^A$ with the assumption that $\mathcal{S}^{h'}$ is executing before the initial enumeration for \mathcal{S}^h noting that by 1 if $\mathcal{S}^{h'}(s, y, i')$ executes before the initial enumeration for \mathcal{S}^h in (s_k, s_{k+1}) then $h'(i') - 1 > x$. \square

We now return and provide the promised proof.

Lemma 5.2. Suppose that $\mathcal{R}_{j,e}$ isn't initialized between s_{-1} and s_k and $\mathcal{S}_{s_k}(\mathcal{R}_{j,e}) < 2$ then

- (1) If $i \leq \langle j, e \rangle$ and $x < s_{-1}$ we have $Y_{i,s-1}^{[3][x]} \prec Y_{i,s_k}^{[3][x]}$.
- (2) If s_k is the greatest active stage less than s and $Y'_{i,s}$ is the result of enumerating $\mathcal{L}_{i,s}$ into $Y_{i,s}$, $i \leq \langle j, e \rangle$, $x < s_{-1}$ then $Y'_{i,s}^{[3][x]} \succ Y_{i,s-1}^{[3][x]}$.

Proof. We first prove 2 on the assumption that 1 holds at each $s_{k'}, k' \leq k$.

By lemma 5.3 it is enough to argue that if $z = h(i) - 1$ then $\mathfrak{b}_{i,s}^h$ (and thus $\mathcal{L}_{i,s} \leq \mathfrak{b}_{i,s}^h$) cancels any changes made by $\mathcal{S}^h(s, z', i)$, $z' \leq z$ to $Y'_{i,s}^{[3][z]}$. After all, item 7 ensures that the claim holds true at all stages prior to such an enumeration, item 7 ensures that it is true when \mathcal{S}^h receives it's initial enumeration, stages at which enumeration occurs into some $W_{i'}^A$, $i' \neq i$ don't alter $Y'_{i,s}^{[3]}$, $\mathfrak{b}_{i,s}^h$ or $\mathcal{L}_{i,s}$. But the combination of steps 4 and 5 and our assumption that 1 holds at all $s_{k'}, k' \leq k$ ensure that $\mathfrak{b}_{i,s}^h$ is small enough to cancel any active axioms causing $Y'_{i,s}^{[3][z]}$ to disagree with $Y_{i,s-1}^{[3][z]}$ or step 2d when it is an initial enumeration and m_i is undefined.

We now note that 2 is trivial if $k = -1$ and otherwise follows by induction on k by application of 2 and the fact that procedure 1 executes procedure 2 when it enumerates c_k . \square

We are now in a position to prove condition 2 holds. We break the proof up into two lemmas.

Lemma 5.4. For all s, i , $\Theta(Y_{i,s})$ is compatible with $W_{i,s}^{A_s}$. Moreover, if $W_{i,s}^{A_s}(x) \uparrow$ then $\Theta(Y_{i,s}; x) = 0$.

Proof. The moreover claim is trivial since if $x > l_s$ then we've taken no action regarding x at or before stage s . We let i be arbitrary and proceed by induction to prove the main claim. Trivially, the claim holds if $s = 0$ so we assume that the claim holds for all stages prior to s and argue the lemma holds at s . If no enumeration happens at s into either A or W_i^A the result is immediate. So we consider the following (disjoint) cases.

Case $x \searrow_s W_i^A$: This case can be broken into two subcases.

Case x is claimed by some $\mathcal{R}_{j,e}$: In this case, there is some \mathcal{S}^h executing at s such that $h(i) - 1 = x$. **Step 5** plus the fact that no elements are enumerated into either $Y_i^{[\leq 2]}$ or $Y_i^{[3][x]}$ after **step 5** during stage s establish the claim.

Case x is unclaimed: In this case, the result follows immediately by **rule 4.1**.

Case $c_k \searrow_s A^{[3]}$: Suppose $\mathcal{R}_{j,e}$ is responsible for the numeration of c_k and consider the following subcases.

Case $x \in W_{i,s-1}^{A_{s-1}}$: In this case, $x \in W_{i,s}^{A_s}$ as any enumeration below l_{s-1} would have injured $\mathcal{R}_{j,e}$. Similarly, $Y_{i,s-1}^{[3][x]} \subset Y_{i,s}^{[3][x]}$. But as we only enumerate elements into $Y_i^{[3][x]}$ when it appears that the lemma would fail no such elements are ever enumerated at or after $s-1$ so the claim holds by virtue of the inductive hypothesis applied at stage $s-1$.

Case $x \notin W_{i,s-1}^{A_{s-1}}$: Note that if $x \geq s-1 \vee i > \langle j, e \rangle$ then the claim follows immediately via the action of **step 2**. So assume $x < s-1 \wedge i \leq \langle j, e \rangle$. Since x is enumerated into W_i^A after stage $s-1$ it is (by induction) enumerated dependent on $A^{[3]}(c_k) = 0$. Thus, $x \notin W_{i,s}^{A_s}$ and the result follows by **lemma 5.2**.

Case $b = b_k \searrow_s A^{[2]}$: Suppose $\mathcal{R}_{j,e}$ is responsible for the numeration of b . **Step 2** ensures that there is some k with $s_k < s$ such that $s_k - 1 \approx s - 1$. But the enumeration of b (chosen large at stage s_k) cancels all enumerations into $Y_i^{[3]}$ after stage $s_k - 1$ and as $s_k - 1 \approx s - 1$ this ensures that $Y_{i,s} \succ Y_{i,s_k}$. As the enumeration of b also cancels any enumeration of x into W_i^A after stage $s_k - 1$ the result follows from the application of the inductive hypothesis to stage s_k .

Case $a \searrow_s A^{[1]}$: Suppose $\mathcal{R}_{j,e}$ is responsible for the numeration of a . The enumeration of a cancels all enumerations into $Y_i^{[3]}$ since the stage t at which b was enumerated as well as any enumeration of x into W_i^A after stage $t - 1$. As no enumerations were made into $Y_i^{[\leq 2]} \upharpoonright_{[t]}$ since stage $t - 1$ the state at stage $t - 1$ is restored and the claim follows from the inductive hypothesis applied at stage $t - 1$.

□

Now to prove the corresponding claim for Γ_i .

Lemma 5.5. For all s, i , $\Gamma_i(A_s \oplus W_{i,s}^{A_s}) \prec Y_{i,s}$ and Γ_i is well-defined.

Note that here Γ_i really refers to whatever version of the functional is left uninjured at the end of stage s since we will reinitialize each Γ_i finitely many times.

Proof. By **definition 3.11** it is enough to show that Γ_i is well-defined. Specifically, we must show that if $t < s$ and \mathcal{P}_i is not injured during the interval $(t, s]$ then

$$A_s \oplus W_{i,s}^{A_s} \succ A_t \oplus W_{i,t}^{A_t} \tag{4a}$$

$$\implies Y_{i,t} \upharpoonright_t \prec Y_{i,s} \tag{4b}$$

We assume the claim holds for all $s' < s$ and prove it also holds at s . This is trivial if no axioms are enumerated at stage s so, suppose, for contradiction, that some element is enumerated at stage s , \mathcal{P}_i is not injured during the interval $(t, s]$ but that eq. (4) fails. We consider the following cases.

Case $z \searrow_s W_i^A$: We break this case up into two subcases.

Case z is unclaimed: In this case, only rule 4.1 enumerates elements at s and all such elements are larger than z . But as no elements are enumerated into $Y_i^{[\leq 2]}$ it follows that the claim must have failed at stage t itself which is ruled out by the inductive hypothesis.

Case z is claimed by $\mathcal{R}_{j,e}$: In this case, $z < s_{-1}$ (as defined in the description of $\mathcal{R}_{j,e}$) and $\mathcal{S}_s(\mathcal{R}_{j,e}) < 2$. Note that we must have $z \notin W_{i,s-1}^{A_{s-1}}$ or it couldn't enter at stage s . So if $t < s_{-1}$ then there must have been some $z' \in A_{s-1}$, $z' < l_t$ which entered during the interval (t, s_{-1}) : rendering eq. (4a) false. If $t < s_k < s$ for some active stage s_k then the enumeration of c_k at stage s_k renders eq. (4a) false. If not then $z \notin W_{i,s}^{A_t}$ contradicting eq. (4a).

Case $q \searrow_s A$: Suppose that $\mathcal{R}_{j,e}$ enumerates q . Note that if $i > \langle j, e \rangle$ and $\mathcal{R}_{j,e}$ takes any action then \mathcal{P}_i is injured giving us the desired conclusion trivially. Thus, we can assume $i \leq \langle j, e \rangle$ and consider the following subcases.

Case $q = c_k \searrow_s A^{[3]}$: We must have $t < s_{-1}$ or $c_{k'}$ makes eq. (4a) false where $s_{k'}$ is the least active stage greater than t . If $A_t \oplus W_{i,t}^{A_t} \neq A_{s-1} \oplus W_{i,s-1}^{A_{s-1}}$ then that disagreement would persist until stage s making eq. (4a) false. But lemma 5.2 lets us conclude that $Y_{i,s-1} \upharpoonright_t \prec Y_{i,s}$ so $Y_{i,t} \upharpoonright_t \prec Y_{i,s}$.

Case $q \searrow_s A^{[\leq 2]}$: This follows from the inductive assumption plus lemma 5.2 which ensures that if $t < s_{-1}$ then eq. (4b) holds. If $t \geq s_{-1}$ then the enumeration of some c_k or q itself renders eq. (4a) false.

□

We can now prove the desired proposition.

Proposition 5.6. *Condition 2 is satisfied, i.e., Θ and Γ_i are well-defined and correct at every stage.*

Proof. Immediate from lemmas 5.4 and 5.5

□

5.2. Winning Strategies

In this section we seek to show that $\mathcal{R}_{j,e}$ only acts finitely many times. To prove this we need to show that our strategies eventually guarantee we find agreement between stages. But the strategy \mathcal{S}^h doesn't execute \mathcal{S}^{hr} on consecutive intervals $[s_k, s_{k+1}]$. It may, instead, skip a number of active stages during which \mathcal{S}^h executes some other sub-strategy only to return to \mathcal{S}^{hr} much later so we need some way to talk about intervals which behave as if we had run \mathcal{S}^{hr} on consecutive intervals.

To that end, we make the following definitions (these are implicitly relative to a particular module $\mathcal{R}_{j,e}$).

Definition 5.7. Say that t is an h successor of $s < t$ (relative to the module $\mathcal{R}_{j,e}$), just if t is an initial enumeration for \mathcal{S}^h and letting $\widehat{Y}_i = Y_{i,s}^h$ we have

$$(\forall i \leq \langle j, e \rangle)(\forall x < h(i)) \left(Y_{i,s}^{[\leq 2]} \upharpoonright_{[s]} \prec \widehat{Y}_i^{[\leq 2]} \wedge Y_{i,s}^{[3][x]} \prec \widehat{Y}_i^{[3][x]} \right)$$

Informally speaking, t is an h successor of s just if for all $i \leq \langle j, e \rangle$ the state of Y_i when \mathcal{S}^h receives its initial enumeration during t agrees with that of $Y_{i,s}$ excepting only elements \mathcal{S}^h can ignore as too large for its concern. Using this notion we can now define a notion of a sequence of intervals that look as if they are consecutive as far as \mathcal{S}^h is concerned.

Definition 5.8. Say that a sequence $[\widehat{t}_0, \widehat{s}_0], [\widehat{t}_1, \widehat{s}_1] \dots, [\widehat{t}_n, \widehat{s}_n]$ of intervals is \mathcal{S}^h -**virtually consecutive** (or just h -virtually consecutive) if

- (1) \mathcal{S}^h is not reinitialized during $(\widehat{t}_0, \widehat{s}_n)$
- (2) At stage \widehat{t}_0 \mathcal{S}^h is in its initialized state, i.e., hasn't been executed since it was last reinitialized.
- (3) \widehat{s}_k is the least active stage greater than \widehat{t}_k
- (4) For all $k < n$, either \widehat{t}_{k+1} is an h -successor of \widehat{s}_k or $\widehat{t}_{k+1} = \widehat{s}_{k+1}$ and no elements are enumerated during the interval ending at \widehat{s}_{k+1} and started at the prior active stage.
- (5) Every interval $\widehat{t}_k, \widehat{s}_k$ satisfying the above between \widehat{t}_0 and \widehat{s}_n appears in the above list.

Note that if the sequence is genuinely consecutive, i.e., $\widehat{t}_{k+1} = \widehat{s}_k + 1$ for all k , then it is trivially h -virtually consecutive. Now let's define what it means for a strategy to be successful.

Definition 5.9. Say that the strategy \mathcal{S}^h is a **winning strategy** if there is a number N_h (the winning number for h) such that if $[\widehat{t}_0, \widehat{s}_0], \dots, [\widehat{t}_{N_h-1}, \widehat{s}_{N_h-1}]$ is an h virtually consecutive sequence of intervals of length N_h then, for **any** acceptable enumeration of elements into the sets $W_i^A, i \in \text{supp } h$ satisfying $x < h(i)$ there are distinct $k, k' < N_h$ if the recommendations of \mathcal{S}^h are followed we have $\widehat{s}_k - 1 \approx \widehat{s}_{k'} - 1$.

We leave the task of formalizing the notion of relative to any acceptable enumeration to the pedantic reader but take the intended content to be clear. In particular, a winning strategy is guaranteed to produce agreement in a bounded number of intervals even if we permute the indices to r.e. sets. With this notion in place we can finally demonstrate that our strategies are winning strategies.

Lemma 5.10. *Suppose that for all $h_r, r < |\text{supp } h|$ the strategy \mathcal{S}^{h_r} is a winning strategy with winning number N_{h_r} then \mathcal{S}^h is a winning strategy.*

Proof. Suppose that $[\widehat{t}_0, \widehat{s}_0], [\widehat{t}_1, \widehat{s}_1] \dots, [\widehat{t}_{N-1}, \widehat{s}_{N-1}]$ is a sequence of h virtually consecutive intervals with $N = \prod_{r < |\text{supp } h|} (N_{h_r} + 2)$ with $z_k \searrow_{\widehat{s}_k} W_k^A$. We argue there are $k, k' < N$ with $\widehat{t}_k \approx \widehat{t}_{k'}$.

We now note that there must be a subsequence of length at least $N_{h_r} + 2$ on which we always execute the same substrategy but never reinitialize that strategy. Specifically, we observe that there is an $\widehat{i} \in \text{supp } h$ and a subsequence $\widehat{t}_{n_k}, k < N_{h_r} + 2$ such that $i_{n_k} = \widehat{i}$ and for all n' with $n_k < n' < n_{k+1}$ we have $i_{n'} > \widehat{i}$. This follows since if this isn't witnessed by i_0 then there must be a subsequence of length at least $N = \prod_{1 < r < |\text{supp } h|} (N_{h_r} + 2)$.

As the support of h is finite we must eventually find such a subsequence. Call this a good \widehat{i} subsequence and let $\widehat{r} = \left\{ y \in \text{supp } h \wedge y < \widehat{i} \right\}$, i.e., $\mathcal{S}^{h_{\widehat{r}}}$ is the strategy called (if necessary) by \mathcal{S}^h in an interval in which the initial enumeration was into $W_{\widehat{i}}^A$. Thinking of this in terms of a finite injury construction this subsequence represents the stages at which the \widehat{i} strategy gets attention without intervening injury.

Suppose that for some $k < N_{h_{\widehat{r}}} + 1$ we fail to enumerate any $x < h(i) - \delta_i(i)$ into W_i^A during $(\widehat{t}_{n_k}, \widehat{s}_{n_k})$. We claim that $\widehat{s}_{n_k} - 1 \approx \widehat{s}_{n_{k+1}} - 1$. By h -virtual connectivity when \mathcal{S}^h begins execution at stage $\widehat{t}_{n_{k+1}}$, $Y_i^{[\leq 2]}$ is compatible with its state at stage $\widehat{s}_{n_k} - 1$. Note that, by **definition 5.9**, we may assume that only $x < h(i)$ are enumerated into any W_i^A .

Thus, in any interval the first element is enumerated at stage \widehat{t}_k . Thus, when \mathcal{S}^h executes at stage $\widehat{t}_{n_{k+1}}$ the action of **step 2e** cancels any elements enumerated both at stage \widehat{s}_{n_k} and by any ancestor strategies executing **step 2f** (which must occur earlier in that stage).

As no $x < h(i) - \delta_i(i)$ was enumerated during the interval $(\widehat{t}_{n_k}, \widehat{s}_{n_k})$ **step 2f** never enumerates any elements and as no parent strategy enumerates any elements into the first two columns of Y_i after executing its child that agreement persists until the end of stage, and, indeed, the end of stage $\widehat{s}_{n_{k+1}} - 1$.

Thus, we can assume that we have a subsequence of length $[\widehat{t}_{n_k}, \widehat{s}_{n_k}]$, $k < N_{h_{r'}} + 1$ with $\tau_{n_k} = r'$ of length $N_{h_{r'}} + 1$ during which we never reinitialize $\mathfrak{S}_{r'}$ and during each interval $(\widehat{t}_{n_k}, \widehat{s}_{n_k})$ there is a \widehat{t}_{n_k} at which we execute $\mathcal{S}^{h_{r'}}$. To verify the claim we first show that the sequence $[\widehat{t}_{n_k}, \widehat{s}_{n_k}]$, $k < N_{h_{r'}}$ is $h_{r'}$ virtually consecutive. Note that $i_{r'} = \widehat{i}$ here and that we may presume that this is the first such subsequence, i.e., \widehat{t}_{n_0} is the first execution of $\mathcal{S}^{h_{r'}}$ since last reinitialization.

This leaves only part 4 of **definition 5.8** to be verified but this follows by **lemma 5.2**. Now suppose that for some k the only element enumerated during the interval $(\widehat{t}_{n_k}, \widehat{s}_{n_k})$ with $k < N_{h_{r'}}$ into W_i^A . By the same reasoning above we'll see agreement at $\widehat{s}_{n_{k+1}}$ since this means that for \widehat{i} we behave as above and the inductive hypothesis handles enumeration into W_i^A , $i \neq \widehat{i}$. Thus, we can assume that in each interval $(\widehat{t}_{n_k}, \widehat{s}_{n_k})$ we see some $x < h(\widehat{i})$ enumerated into W_i^A . In this case, when that x is enumerated the operation of **step 2e** by $\mathcal{S}^{h_{r'}}$ cancels any element enumerated at stage \widehat{t}_{n_k} during **step 2f**. Thus, the situation with respect to the first two columns of the sets Y_i is no different than if enumeration was restricted only to those x entering W_i^A where $x < h_{r'}(i)$. □

We can now prove our desired result.

Proposition 5.11. $\mathcal{R}_{j,e}$ acts only finitely many times.

Proof. Suppose not. The only possibility is that some module $\mathcal{R}_{j,e}$ is initialized at some stage s_{-1} and never subsequently reinitialized by the action of any higher priority module. But this can only happen if at all active stages s_k , $k > 0$ we have $\mathcal{S}_{s_k}(\mathcal{R}_{j,e}) = 1$.

But, by **lemma 5.10** we have that $\mathcal{S}^{h^{j,e}}$ is a winning strategy. Consider the sequence (\widehat{t}_k, s_k) where \widehat{t}_k is the first stage subsequent to s_k at which $\mathcal{R}_{j,e}$ claims an enumeration. As enumerations that go unclaimed or are claimed by modules with lower priority don't enumerate elements that are small relative to the last stage at which $\mathcal{R}_{j,e}$ acts into the first two columns of any Y_i it is trivial that this sequence is $h^{j,e}$ virtually contiguous. It thus follows that there are k, k' with $s_k - 1 \approx s_{k'} - 1$ and thus at stage $s_{k'}$ we have $\mathcal{S}_{s_{k'}}(\mathcal{R}_{j,e}) = 2$ contrary to our assumption. Hence every module acts only finitely often. □

5.3. Putting It Together

Lemma 5.12. Every requirement of the form $\mathcal{R}_{j,e}$ is satisfied.

Proof. Assume, for a contradiction, $\mathcal{R}_{j,e}$ isn't satisfied (i.e., $\Phi_j(A) = X_e \wedge \Phi_j(X_e) = A$) and that t_0 is the last stage at which the module $\mathcal{R}_{j,e}$ acts. If $s > t_0$ then $\mathcal{S}_{t_0}(\mathcal{R}_{j,e}) = \mathcal{S}_s(\mathcal{R}_{j,e})$ and $A_{t_0} \prec A_s$ since such a change would require $\mathcal{R}_{j,e}$ act at that stage. Hence, $A_{t_0} \prec A$.

If the variable c is defined for the $\mathcal{R}_{j,e}$ at stage t_0 (i.e., if $\mathcal{S}_{t_0}(\mathcal{R}_{j,e}) \geq 2$) we leave that value in place but if not set $c = c_m$ where c_m is the least m such that c_m hasn't been enumerated into $A^{[3]}$. Note that c_0 must be defined at t_0 since if $\mathcal{S}_{t_0}(\mathcal{R}_{j,e}) = \uparrow$ then $\mathcal{R}_{j,e}$ acts at stage $t_0 + 1$

By [lemma 3.7](#) let $s > t_0$ be a stage at which $\Psi(c, s)$ holds. But if $\mathcal{S}_s(\mathcal{R}_{j,e}) < 3$ then we have $\mathcal{R}_{j,e}$ acts at stage $s + 1$. On the other hand if $\mathcal{S}_s(\mathcal{R}_{j,e}) = 3$ then we have $\Psi^3(c, \hat{s}_0, \hat{s}_1, \hat{s}_2, s)$. However, this is exactly what [lemma 3.9](#) denies. Contradiction! \square

Before we can complete the proof of our main theorem we must fulfill the commitment we made in [condition 1](#) and verify that $\Gamma_i(A \oplus W_i^A)$ is total.

Lemma 5.13. *There are infinitely many stages s such that $A_s \prec A_t$ for all $t > s$ and thus $A_s \prec A$. Thus, [condition 1](#) hold and $\Gamma_i(A \oplus W_i^A)$ is total.*

Note that, this result entails that A is Δ_2^0 .

Proof. To verify the primary claim note that if $\mathcal{R}_{j,e}$ acts at stage s it injures lower priority modules and when reinitialized those modules never enumerate an element into any column of A below l_{s+1} . Thus, if s_k is last stage that any module with priority $\leq k$ acts we have $A_t \succ A_{s_k}$ for all $t > s_k$ and $A_{s_k} \prec A$.

[Condition 1](#) follows immediately and the totality of $\Gamma_i(A \oplus W_i^A)$ was observed to follow from this fact in the discussion immediately following [definition 3.11](#). \square

We can now easily complete the proof of our main theorem.

Theorem 1.4. *There is a properly 3-REA set A which can't be extended to a properly 4-REA set. The set A can also be taken to be Δ_2^0 .*

Proof. By [lemma 5.12](#) to show that A is properly 3-REA but can't be expanded to a 4-REA set it is enough to verify that the requirements of the form \mathcal{P}_i is satisfied. Clearly $\Theta(Y_i)$ is total for all i and $\Gamma_i(A \oplus W_i^A)$ is total by [lemma 5.13](#). Hence, it's enough to show that these functionals are correct at every stage. But by [proposition 5.6](#) this holds as long as no module for \mathcal{P}_i is injured infinitely many times. However, as only modules of the form $\mathcal{R}_{j,e}$ are responsible for injuries this is immediate from [proposition 5.11](#).

The fact that A can be taken to be Δ_2^0 follows from [lemma 5.13](#). \square

References

- [1] Peter A. Cholak and Peter G. Hinman. Iterated relative recursive enumerability. *Archive for Mathematical Logic*, 33(5):321–346, 1994.
- [2] Carl G. Jockusch and Richard A. Shore. Pseudo-jump operators. ii: Transfinite iterations, hierarchies and minimal covers. *The Journal of Symbolic Logic*, 49(04):1205–1236, 1984.
- [3] Piergiorgio Odifreddi. *Classical recursion theory: The theory of functions and sets of natural numbers*, volume 1. Elsevier, 1992.
- [4] H. Rogers. *Theory of recursive functions and effective computability*. MIT Press, Cambridge, Mass, 1987.
- [5] Robert I. Soare and Michael Stob. Relative recursive enumerability. In *Studies in Logic and the Foundations of Mathematics*, volume 107, pages 299–324. Elsevier, 1982.