

DISTINGUISHED STRATA IN A REDUCTIVE GROUP

G. LUSZTIG

INTRODUCTION

0.1. Let G be a connected reductive algebraic group over \mathbf{C} , with Weyl group W . In [L15] we have defined a partition of G into finitely many strata. The set of strata is in bijection with the image of a map $cl(W) \rightarrow \text{Irr}(W)$ where $cl(W)$ is the set of conjugacy classes in W and $\text{Irr}(W)$ is a set of representatives for the isomorphism classes of irreducible representations of W over \mathbf{Q} . The set of strata can be viewed as an enlargement of the set of unipotent classes of G (a unipotent class of G is contained in exactly one stratum) which is built from the sets of unipotent classes in groups like G but in all characteristics. According to [BC76], the classification of unipotent classes of G can be reduced to the classification of a much smaller set of unipotent classes, the distinguished ones. We will show that, similarly, the classification of strata of G can be essentially reduced to the classification of a much smaller set, that of distinguished strata. It turns out that the distinguished strata of G are indexed by a subset $CL_{dist}(W)$ of the set of elliptic conjugacy classes of W which may be called distinguished conjugacy classes.

0.2. Let $\mathcal{P} = \{2, 3, 5, \dots\}$ be the set of prime numbers. For any $r \in \mathcal{P}$ let $G^{(r)}$ be a connected reductive group over an algebraically closed field of characteristic r of the same type as G . We set $G^{(0)} = G$. For $r \in \{0\} \cup \mathcal{P}$ let $\mathcal{U}^{(r)}$ be the set of unipotent classes of $G^{(r)}$. By the Springer correspondence (extended in [L84] to small characteristic) there is a natural imbedding $\iota^{(r)} : \mathcal{U}^{(r)} \rightarrow \text{Irr}(W)$ whose image is denoted by $\mathbf{S}^{(r)}(W)$; it is known that $\mathbf{S}^{(0)}(W) \subset \mathbf{S}^{(r)}(W)$. Let

$$\mathbf{S}(W) = \cup_{r \in \{0\} \cup \mathcal{P}} \mathbf{S}^{(r)}(W) = \cup_{r \in \mathcal{P}} \mathbf{S}^{(r)}(W) \subset \text{Irr}(W).$$

In [L15] (where the notation $\mathcal{S}_2(W)$ is used instead of $\mathbf{S}(W)$) it is shown that $\mathbf{S}(W)$ depends only on W as a Coxeter group, not on the underlying root datum (but it is not clear whether $\mathbf{S}(W)$ makes sense for a finite non-crystallographic Coxeter group).

In [L15] we have defined for any $r \in \{0\} \cup \mathcal{P}$ a surjective map $\kappa^{(r)} : G^{(r)} \rightarrow \mathbf{S}(W)$ whose fibres are called the *strata* of $G^{(r)}$; each stratum is a union of conjugacy

Supported by NSF grant DMS-1855773 and by a Simons Fellowship

classes of the same dimension, independent of r and, according to [C20], is locally closed in $G^{(r)}$. If $g \in G^{(r)}$ is unipotent, then $\kappa^{(r)}(g)$ is the same as the image of the conjugacy class of g under $\iota^{(r)}$. It follows that for any $E \in \mathbf{S}(W)$, there exists $r \in \{0\} \cup \mathcal{P}$ such that the stratum $(\kappa^{(r)})^{-1}(E)$ contains some unipotent element.

0.3. An element of $G^{(r)}$ is said to be distinguished if it is not contained in a Levi subgroup of a proper parabolic subgroup of $G^{(r)}$ (see [BC76]). Let $\mathcal{U}_{dist}^{(r)}$ be the set of unipotent classes in $G^{(r)}$ in which some/any element is distinguished. Let $\mathbf{S}_{dist}^{(r)}(W) = \iota^{(r)}(\mathcal{U}_{dist}^{(r)})$.

We say that $E \in \mathbf{S}(W)$ is distinguished if $E \in \cup_{r \in \{0\} \cup \mathcal{P}} \mathbf{S}_{dist}^{(r)}(W)$ or equivalently if there exists $r \in \{0\} \cup \mathcal{P}$ such that the stratum $(\kappa^{(r)})^{-1}(E)$ contains some distinguished unipotent element. Let $\mathbf{S}_{dist}(W)$ be the set of distinguished elements of $\mathbf{S}(W)$. In §1 we will show:

(a) $E \in \mathbf{S}(W)$ is distinguished if and only if there exists $r \in \{0\} \cup \mathcal{P}$ such that the stratum $(\kappa^{(r)})^{-1}(E)$ contains some distinguished (not necessarily unipotent) element of $G^{(r)}$.

A stratum of $G^{(r)}$ (with $r \in \{0\} \cup \mathcal{P}$) is said to be distinguished if it is of the form $(\kappa^{(r)})^{-1}(E)$ where $E \in \mathbf{S}_{dist}(W)$ (such a stratum need not contain a distinguished element).

0.4. For $C \in cl(W)$ let $m(C)$ be the dimension of the 1-eigenspace of some/any $w \in C$ on the reflection representation of W .

We shall write $\Phi : cl(W) \rightarrow \mathbf{S}(W)$ for what in [L15] is denoted by $'\Phi$ (a surjective map). In [L15] it is shown that

(a) for any $E \in \mathbf{S}(W)$ there is a unique $C_E \in \Phi^{-1}(E)$ which is as elliptic as possible, that is $m(C_E) \leq m(C)$ for any $C \in \Phi^{-1}(E)$;

thus $E \mapsto C_E$ is a cross section of the surjective map Φ . The following variant of (a) will be verified in §1.

(b) for any $E \in \mathbf{S}(W)$ there is a unique $C'_E \in \Phi^{-1}(E)$ which is as non-elliptic as possible, that is $m(C'_E) \geq m(C)$ for any $C \in \Phi^{-1}(E)$.

Let $CL(W)$ be the image of the map $E \mapsto C'_E$, $\mathbf{S}(W) \rightarrow cl(W)$. Note that Φ restricts to a bijection $CL(W) \xrightarrow{\sim} \mathbf{S}(W)$. Under this bijection, the subset $\mathbf{S}_{disc}(W)$ of $\mathbf{S}(W)$ corresponds to a subset $CL_{dist}(W)$ of $CL(W)$. The conjugacy classes of W contained in CL_{dist} are said to be distinguished. The following result will be proved in §1.

(c) Let $C \in CL(W)$. We have $C \in CL_{dist}(W)$ if and only if C is elliptic (that is, $m(C) = 0$).

0.5. For $r \in \{0\} \cup \mathcal{P}$ let $\psi^{(r)} : cl(W) \rightarrow \mathcal{U}^{(r)}$ be the surjective map defined in [L11a]. (In the case $r = 0$, an alternative definition of this map is given in [Y20]. Let $\Phi^{(r)} : cl(W) \rightarrow \text{Irr}(W)$ be the composition of $\psi^{(r)}$ with $\iota^{(r)} : \mathcal{U}^{(r)} \rightarrow \text{Irr}(W)$. From the explicit description of $\psi^{(r)}$ in [L12] and the explicit description of $CL_{dist}(W)$ given in this paper we see that

(a) if $C \in CL_{dist}(W)$ then $\Phi^{(r)}(C)$ is independent of r . Hence, by the definition of Φ in [L15, 4.1], we have $\Phi(C) = \Phi^{(r)}(C)$ for all r .

0.6. Notation. A bipartition is a sequence $\lambda_* = (\lambda_1, \lambda_2, \lambda_3, \dots)$ in \mathbf{N} such that $\lambda_i = 0$ for i large and $\lambda_1 \geq \lambda_3 \geq \lambda_5 \geq \dots$, $\lambda_2 \geq \lambda_4 \geq \lambda_6 \geq \dots$. We write $|\lambda_*| = \lambda_1 + \lambda_2 + \lambda_3 + \dots$. Let BP be the set of bipartitions.

Let T be the set of all $\lambda_* \in BP$ such that $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \lambda_4 \geq \dots$. Let R be the set of all $\lambda_* \in T$ such that λ_i is even for any i . Let P be the set of all $\lambda_* \in T$ such that $\lambda_1 = \lambda_2, \lambda_3 = \lambda_4, \lambda_5 = \lambda_6, \dots$. For $\lambda_* \in T$ and $j > 0$ we set $\mu_j(\lambda_*) = \#\{k \geq 1; \lambda_k = j\}$. For $m \in \mathbf{N}$ let $BP^m = \{\lambda_* \in BP, |\lambda_*| = m\}$, $T^m = T \cap BP^m$. Let T_{ev} (resp. R_{ev}) be the subset of T (resp. R) consisting of the λ_* with an even number of > 0 terms. For a, b in \mathbf{Z} we write $a \gg b$ instead $a - b \geq 2$.

1. PROOF OF 0.3(A), 0.4(B), 0.4(C)

1.1. In this section we prove 0.3(a), 0.4(b), 0.4(c). To do this we can assume that G is almost simple. It is also enough to consider only one such G in each isogeny class. The case where G is of classical (resp. exceptional) type is considered in 1.2-1.9 (resp. 1.10-1.15).

1.2. Assume first that $G = SL_n(\mathbf{C}), n \geq 2$. In this case $\mathbf{S}^{(r)}(W) = \text{Irr}(W) = \mathbf{S}(W)$ for any r and the map Φ is a bijection $cl(W) \xrightarrow{\sim} \mathbf{S}(W)$. In this case 0.4(b) is obvious and we have $CL(W) = cl(W)$. Also 0.3(a) is immediate (an element is distinguished if and only if is regular unipotent times a central element). Note that $CL_{dist}(W)$ consists of a single element: the class of the Coxeter element; thus 0.4(c) holds.

1.3. Until the end of 1.5 we assume that $G = Sp_{2n}(\mathbf{C}), n \geq 2$. Then $\mathbf{S}(W) = \mathbf{S}^{(2)}(W)$ and Φ becomes a map $cl(W) \rightarrow \mathbf{S}^{(2)}(W)$. By [L12] we have bijections

(a) $\mathcal{U}^{(2)} \leftrightarrow$ set of all pairs (c_*, ϵ) where $c_* \in T^{2n}$ is such that $\mu_j(c_*) \in 2\mathbf{N}$ for any odd j and $\epsilon : \{j \in 2\mathbf{N} + 2; \mu_j(c_*) \in 2\mathbf{N} + 2\} \rightarrow \{0, 1\}$,

(b) $cl(W) \leftrightarrow (R \times P)^{2n} := \{(r_*, p_*) \in R \times P; |r_*| + |p_*| = 2n\}$.

Via (a),(b), $(\iota^{(2)})^{-1}\Phi : cl(W) \rightarrow \mathcal{U}^{(2)}$ becomes the map

(c) $(r_*, p_*) \mapsto (c_*, \epsilon)$ where $\mu_j(c_*) = \mu_j(r_*) + \mu_j(p_*)$ for $j > 0$ and for any $j \in 2\mathbf{N} + 2$ such that $\mu_j(c_*) \in 2\mathbf{N} + 2$, we have $\epsilon(j) = 1$ if $j = r_i$ for some i and $\epsilon(j) = 0$, otherwise.

Via (b), the map $cl(W) \rightarrow \mathbf{N}, C \mapsto m(C)$ becomes the map which to any $(r_*, p_*) \in (R \times P)^{2n}$ associates $|p_*|/2$. To prove 0.4(b) we must show that, if (c_*, ϵ) (as in (a)) is given, then there is a unique $(r_*, p_*) \in (R \times P)^{2n}$ which maps to it (as in (c)) and has $|p_*|$ maximum possible. Thus,

(d) for $j \in 2\mathbf{N} + 2$ such that $\mu_j(c_*) \in 2\mathbf{N} + 2$, we must have that $\mu_j(r_*) = 2, \mu_j(p_*) = \mu_j(c_*) - 2$ (if $\epsilon(j) = 1$) and $\mu_j(r_*) = 0, \mu_j(p_*) = \mu_j(c_*)$ (if $\epsilon(j) = 0$); for $j \in 2\mathbf{N} + 2$ such that $\mu_j(c_*) \in 2\mathbf{N} + 1$, we must have that $\mu_j(r_*) = 1, \mu_j(p_*) = \mu_j(c_*) - 1$; for $j \in 2\mathbf{N} + 2$ such that $\mu_j(c_*) = 0$, we must have that $\mu_j(r_*) =$

$0, \mu_j(p_*) = 0$; for $j \in 2\mathbf{N} + 1$, we must have that $\mu_j(r_*) = 0, \mu_j(p_*) = \mu_j(c_*)$.

This proves 0.4(b) in our case.

Note that:

(e) $CL(W)$ is (via (b)) the set of all $(r_*, p_*) \in (R \times P)^{2n}$ such that $\mu_j(r_*) \leq 2$ for any $j > 0$.

1.4. As in [L15] we have a bijection

(a) $\text{Irr}(W) \leftrightarrow BP^n$.

Using [L15, §3], we see that

(b) when $r \neq 2$, the subset $\iota^{(r)}(\mathcal{U}_{dist}^{(r)})$ of $\text{Irr}(W)$ becomes via (a) the subset of BP^n consisting of sequences of the form $(a_1 > a_2 > \dots > a_s > 0, 0, 0, \dots)$.

By [LS12, 6.2], the set $\mathcal{U}_{dist}^{(2)}$ can be identified via 1.3(a) with the subset of $\mathcal{U}^{(2)}$ consisting of

(c) all (c_*, ϵ) (as in 1.3(a)) such that $\mu_j(c_*) = 0$ for odd j , $\mu_j(c_*) \leq 2$ for even j and $\epsilon(j) = 1$ whenever j is even and $\mu_j(c_*) = 2$.

Using [L15, §3] we see that the subset $\iota^{(2)}(\mathcal{U}_{dist}^{(2)})$ of $\text{Irr}(W)$ becomes via (a) the subset of BP^n formed by the sequences $(c_1/2, c_2/2, c_3/2, \dots)$ for various (c_*, ϵ) as in (c). This is the same as the set of all $(a_1 \geq a_2 \geq a_3 \geq \dots) \in BP^n$ such that there are no consecutive equalities between the non zero a_i . This set contains the set (b). It follows that

(d) $\mathbf{S}_{dist}^{(r)}(W) \subset \mathbf{S}_{dist}^{(2)}(W)$ for any r .

Under our bijection $CL_{dist}(W) \leftrightarrow \mathbf{S}_{dist}(W)$, the set of (c_*, ϵ) as in (c) corresponds to the set of $(r_*, p_*) \in (R \times P)^{2n}$ such that $\mu_j(r_*) \leq 2$ for $j > 0$ and $p_* = (0, 0, 0, \dots)$; this is the same as the set of all $(r_*, p_*) \in (R \times P)^{2n}$ which correspond to elliptic conjugacy classes in W . This implies (by (d)) that 0.4(c) holds in our case.

1.5. Let $g \in G^{(r)}$ be a distinguished element. To prove 0.3(a) it is enough to show that $\kappa^{(r)}(g) \in \mathbf{S}_{dist}(W)$. If $r = 2$ then g is unipotent and the result is clear. Thus we can assume that $r \neq 2$. Using [L15, §3] we see that under the bijection 1.4(a), $\kappa^{(r)}(g)$ corresponds to a bipartition of the form

(a) $((a_1 + b_1)/2, (a_2 + b_2)/2, (a_3 + b_3)/2, \dots)$

where

$$a_1 = \nu_1/2, a_2 = \nu_2/2, a_3 = \nu_3/2, \dots, a_s = \nu_s/2, a_{s+1} = 0, a_{s+2} = 0, \dots,$$

$$b_1 = \nu'_1/2, b_2 = \nu'_2/2, b_3 = \nu'_3/2, \dots, b_t = \nu'_t/2, b_{t+1} = 0, b_{t+2} = 0, \dots,$$

and $\nu_1 > \nu_2 > \nu_3 > \dots > \nu_s, \nu'_1 > \nu'_2 > \nu'_3 > \dots > \nu'_t$ are even integers ≥ 2 with $\sum_k \nu_k + \sum_k \nu'_k = 2n$.

Clearly $(a_1 + b_1)/2 \geq (a_2 + b_2)/2 \geq (a_3 + b_3)/2 \geq \dots$. If $(a_i + b_i)/2 = (a_{i+2} + b_{i+2})/2$ then $a_i = a_{i+2}, b_i = b_{i+2}$ hence $a_{i+2} = 0, b_{i+2} = 0$ and $(a_{i+2} + b_{i+2})/2 = 0$. Thus (a) corresponds to an element of $\mathbf{S}_{dist}^{(2)}(W)$. This proves 0.3(a) in our case.

1.6. We now assume that $G = SO_{2n+1}(\mathbf{C}), n \geq 3$. Then the arguments in 1.3, 1.4 can be used word by word in the present case except that 1.4(b) must be replaced by the following statement:

(a) When $r \neq 2$, the subset $\iota^{(r)}(\mathcal{U}_{dist}^{(r)})$ of $\text{Irr}(W)$ becomes via 1.4(a) the subset of BP^n consisting of sequences (c_1, c_2, c_3, \dots) where

$$c_1 = (\nu_1 - 1)/2, c_2 = (\nu_2 + 1)/2, c_3 = (\nu_3 - 1)/2, c_4 = (\nu_4 + 1)/2, \dots, c_{2s+1} = (\nu_{2s+1} - 1)/2, c_{2s+2} = 0, c_{2s+3} = 0, \dots$$

and $\nu_1 > \nu_2 > \nu_3 > \dots > \nu_{2s+1}$ are odd integers ≥ 1 with sum $2n + 1$.

(Note that $c_1, c_2, \dots) \in P^n$, with no two successive equalities between its nonzero terms hence it corresponds to an element in $\mathbf{S}_{dist}^{(2)}(W)$.)

The argument in 1.5 also continues to hold except that 1.5(a) must be replaced by

$$(b) ((a_1 + b_1)/2, (a_2 + b_2)/2, (a_3 + b_3)/2, \dots)$$

where

$$a_1 = (\nu_1 - 1)/2, a_2 = (\nu_2 + 1)/2, a_3 = (\nu_3 - 1)/2, a_4 = (\nu_4 + 1)/2, \dots, a_{2s+1} = (\nu_{2s+1} - 1)/2, a_{2s+2} = 0, a_{2s+3} = 0, \dots,$$

$$b_1 = (\nu'_1 + 1)/2, b_2 = (\nu'_2 - 1)/2, b_3 = (\nu'_3 + 1)/2, b_4 = (\nu'_4 - 1)/2, \dots, b_{2t} = (\nu'_{2t} - 1)/2, b_{2t+1} = 0, b_{2t+2} = 0, \dots,$$

and $\nu_1 > \nu_2 > \nu_3 > \dots > \nu_{2s+1}$, $\nu'_1 > \nu'_2 > \nu'_3 > \dots > \nu'_{2t}$ are odd integers ≥ 1 with $\sum_k \nu_k + \sum_k \nu'_k = 2n + 1$.

We have $a_1 \geq a_2 \geq a_3 \geq \dots$ and $b_1 \geq b_2 \geq b_3 \geq \dots$ hence $(a_1 + b_1)/2 \geq (a_2 + b_2)/2 \geq (a_3 + b_3)/2 \geq \dots$. If $(a_i + b_i)/2 = (a_{i+2} + b_{i+2})/2$ then $a_i = a_{i+2}, b_i = b_{i+2}$ hence $a_i = b_i = 0$.

Thus (a) corresponds to an element of $\mathbf{S}_{dist}^{(2)}(W)$. This proves 0.3(a) in our case.

1.7. Until the end of 1.9 we assume that $G = SO_{2n}(\mathbf{C}), n \geq 4$. We have $\mathbf{S}(W) = \mathbf{S}^{(2)}(W)$ and Φ becomes a map $cl(W) \rightarrow \mathbf{S}^{(2)}(W)$. Now each of

$$cl(W), \text{Irr}(W), \mathbf{S}^{(2)}(W), \mathcal{U}^{(2)}$$

has a natural involution induced by conjugation by an element in the non-identity component of the full orthogonal group. We then have partitions

$$cl(W) = cl(W)' \sqcup cl(W)'', \text{Irr}(W) = \text{Irr}(W)' \sqcup \text{Irr}(W)'',$$

$$\mathbf{S}^{(2)}(W) = \mathbf{S}^{(2)}(W)' \sqcup \mathbf{S}^{(2)}(W)'', \mathcal{U}^{(2)} = \mathcal{U}^{(2)'} \sqcup \mathcal{U}^{(2)''},$$

where $()'$ denotes the set of fixed points of the involution and $()''$ denotes its complement.

By [L12] we have bijections

(a) (set of orbits of the involution on $\mathcal{U}^{(2)''}$) \leftrightarrow (set of all pairs (c_*, ϵ) as in 1.3(a) such that $\epsilon = 0$ and $\mu_j(c_*) = 0$ for j odd, $\mu_j(c_*)$ even for j even),

(b) (set of orbits of the involution on $cl(W)''$) \leftrightarrow (set of all pairs $(r_*, p_*) \in (R \times P)^{2n}$ such that $r_* = (0, 0, \dots)$ and $\mu_j(p_*) = 0$ for j odd),

(c) $\mathcal{U}^{(2)'}$ \leftrightarrow (set of all pairs (c_*, ϵ) as in 1.3(a) which are not as in (a) and are such that $c_* \in T_{ev}$),

(d) $cl(W)'$ \leftrightarrow (set of all pairs $(r_*, p_*) \in (R \times P)^{2n}$ which are not as in (b) and are such that $r_* \in R_{ev}$).

Now $(\iota^{(2)})^{-1}\Phi$ induces the (surjective) map $cl(W)' \rightarrow \mathcal{U}^{(2)'}$ which by (c),(d) becomes the map $(r_*, p_*) \mapsto (c_*, \epsilon)$ given by the same rule as in 1.3(c). The same proof as in 1.3 shows that if (c_*, ϵ) (as in (c)) is given, then there is a unique (r_*, p_*) (as in (d)) which maps to it and has $|p_*|$ maximum possible. This implies that 0.4(b) holds for any $E \in \mathbf{S}(W) = \mathbf{S}^{(2)}(W)$ that is contained in $\mathbf{S}^{(2)}(W)'$. If $E \in \mathbf{S}^{(2)}(W)''$ then $\Phi^{-1}(E)$ consists of a single element so that 0.4(b) holds automatically for such E . This proves 0.4(b) in our case.

In our case

(e) the set $CL(W)$ becomes the set of pairs (r_*, p_*) as in 1.3(e) such that $r_* \in R_{ev}$ and with each pair as in (b) repeated twice.

1.8. As in 1.4(a) we have a bijection

(a) $\text{Irr}(W)' \leftrightarrow$ set of all $(a_1, a_2, a_3, \dots) \in BP^n$ such that $a_1 - a_2, a_3 - a_4, \dots$ are not all zero.

Using [L15, §3] we see that

(b) when $r \neq 2$, the subset $\iota^{(r)}(\mathcal{U}_{dist}^{(r)})$ of $\text{Irr}(W)'$ becomes via (a) the subset of BP^n consisting of sequences of the form $((\nu_1 + 1)/2, (\nu_2 - 1)/2, (\nu_3 + 1)/2, \dots, (\nu_{2s} - 1)/2, 0, 0, 0, \dots)$ where $\nu_1 > \nu_2 > \dots > \nu_{2s}$ are odd ≥ 1 .

By [LS12, 6.2], the set $\mathcal{U}_{dist}^{(2)}$ can be identified via 1.7(a) with the set consisting of

(c) all (c_*, ϵ) (as in 1.3(a)) where $c_* \in T_{ev}$ and such that $\mu_j(c_*) = 0$ for odd j , $\mu_j(c_*) \leq 2$ for even j and $\epsilon(j) = 1$ whenever j is even and $\mu_j(c_*) = 2$.

Using [L15, §3] we see that the subset $\iota^{(2)}(\mathcal{U}_{dist}^{(2)})$ of $\text{Irr}(W)'$ becomes via (a) the subset of BP^n formed by the sequences $((c_1 + 2)/2, (c_2 - 2)/2, (c_3 + 2)/2, (c_4 - 2)/2, \dots, (c_{2s} - 2)/2, 0, 0, 0, \dots)$ for various (c_*, ϵ) as in (c) with $c_* = (c_1 \geq c_2 \geq \dots \geq c_{2s} > 0, 0, 0, \dots)$.

If $\nu_1 > \nu_2 > \dots > \nu_{2s}$ are odd ≥ 1 then

$$\begin{aligned} & ((\nu_1 + 1)/2, (\nu_2 - 1)/2, (\nu_3 + 1)/2, \dots, (\nu_{2s} - 1)/2, 0, 0, 0, \dots) \\ &= ((c_1 + 2)/2, (c_2 - 2)/2, (c_3 + 2)/2, (c_4 - 2)/2, \dots, (c_{2s} - 2)/2, 0, 0, 0, \dots) \end{aligned}$$

where $c_1 = \nu_1 - 1, c_2 = \nu_2 + 1, c_3 = \nu_3 - 1, \dots, c_{2s} = \nu_{2s} + 1$ are even and $c_1 \geq c_2 \gg c_3 \geq c_4 \gg \dots \geq c_{2s} \gg 0$. From this we see that $\iota^{(r)}(\mathcal{U}_{dist}^{(r)}) \subset \iota^{(2)}(\mathcal{U}_{dist}^{(2)})$ for any r , that is

(d) $\mathbf{S}_{dist}^{(r)}(W) \subset \mathbf{S}_{dist}^{(2)}(W)$ for any r .

Under our bijection $CL_{dist}(W) \leftrightarrow \mathbf{S}_{dist}(W)$, the set of (c_*, ϵ) as in (c) corresponds to the set of (r_*, p_*) as in 1.7(d) such that $\mu_j(r_*) \leq 2$ for $j > 0$ and $p_* = (0, 0, 0, \dots)$; this is the same as the set of all (r_*, p_*) in 1.7(e) which correspond to elliptic conjugacy classes in W . This implies (by (d)) that 0.4(c) holds in our case.

1.9. Let $g \in G^{(r)}$ be a distinguished element. To prove 0.3(a) it is enough to show that $\kappa^{(r)}(g) \in \mathbf{S}(W)$. If $r = 2$ then g is unipotent and the result is clear. Thus

we can assume that $r \neq 2$. Using [L15, §3] we see that under the bijection 1.8(a), $\kappa^{(r)}(g)$ corresponds to a bipartition of the form

$$(a) \left((a_1 + b_1)/2, (a_2 + b_2)/2, (a_3 + b_3)/2, \dots \right)$$

where

$$a_1 = (\nu_1 + 1)/2, a_2 = (\nu_2 - 1)/2, a_3 = (\nu_3 + 1)/2, a_4 = (\nu_4 - 1)/2, \dots, a_{2s} = (\nu_{2s} - 1)/2, a_{2s+1} = 0, a_{2t+2} = 0, \dots,$$

$$b_1 = (\nu'_1 + 1)/2, b_2 = (\nu'_2 - 1)/2, b_3 = (\nu'_3 + 1)/2, b_4 = (\nu'_4 - 1)/2, \dots, b_{2t} = (\nu'_{2t} - 1)/2, b_{2t+1} = 0, b_{2t+2} = 0, \dots,$$

and $\nu_1 > \nu_2 > \nu_3 > \dots > \nu_{2s}$, $\nu'_1 > \nu'_2 > \nu'_3 > \dots > \nu'_{2t}$ are odd integers ≥ 1 with $\sum_k \nu_k + \sum_k \nu'_k = 2n$. We can assume that $t \leq s$. We have

$$\left((a_1 + b_1)/2, (a_2 + b_2)/2, (a_3 + b_3)/2, \dots \right) = \left((c_1 + 2)/2, (c_2 - 2)/2, (c_3 + 2)/2, (c_4 - 2)/2, \dots, (c_{2s} - 2)/2, 0, 0, 0, \dots \right)$$

where $c_1 = \nu_1 + \nu'_1$, $c_2 = \nu_2 + \nu'_2$, \dots , $c_{2t} = \nu_{2t} + \nu'_{2t}$, $c_{2t+1} = \nu_{2t+1} - 1$, $c_{2t+2} = \nu_{2t+2} + 1$, \dots , $c_{2s} = \nu_{2s} + 1$. (The terms c_{2t+1}, \dots, c_{2s} are missing if $t = s$.) Note that c_1, c_2, \dots, c_{2s} are even, nonzero and $c_1 \geq c_2 \geq c_3 \geq c_4 \geq \dots \geq c_{2s}$ with no consecutive equalities. Thus (a) corresponds to an element of $\mathbf{S}_{dist}^{(2)}(W)$. This proves 0.3(a) in our case.

1.10. In this remainder of this section we assume that G is simple of exceptional type. In this case 0.4(b) can be deduced from tables in [L15]. In 1.11-1.15 we describe in each case, using notation of [L15, §4] the bijection between $CL(W)$ and $\mathbf{S}(W)$ in the form $C \leftrightarrow E$ (to which we add "dist" when $E \in \mathbf{S}_{dist}^{(r)}(W)$ for all r or "dist" _{r} when $E \in \mathbf{S}_{dist}^{(r)}(W)$ for a single r). Here we also use the description of distinguished unipotent elements in [LS12]. Now 0.3(a), 0.4(c) can be verified in each case using the definitions.

1.11. Type G_2 .

$$[G_2] \leftrightarrow 1_0, \text{ dist}$$

$$[A_2] \leftrightarrow 2_1, \text{ dist}$$

$$[A_1 + \tilde{A}_1] \leftrightarrow 2_2, \text{ dist}_3$$

$$[\tilde{A}_1] \leftrightarrow 1_3$$

$$[A_1] \leftrightarrow 1_3$$

$$[A_0] \leftrightarrow 1_6$$

1.12. Type F_4 .

$$[F_4] \leftrightarrow 1_0, \text{ dist}$$

$$[B_4] \leftrightarrow 4_1, \text{ dist}$$

$$[F_4(a_1)] \leftrightarrow 9_2, \text{ dist}$$

$$[B_3] \leftrightarrow 8_3$$

$$[C_3] \leftrightarrow 8_3$$

$$[D_4(a_1)] \leftrightarrow 12_4, \text{ dist}$$

$$[A_3 + \tilde{A}_1] \leftrightarrow 16_5, \text{ dist}_2$$

$$[A_3] \leftrightarrow 9_6$$

$$[B_2 + A_1] \leftrightarrow 9_6$$

$$\begin{aligned}
[\tilde{A}_2 + \tilde{A}_2] &\leftrightarrow 6_6, \text{ dist}_2 \\
[A_2 + \tilde{A}_1] &\leftrightarrow 4_7 \\
[\tilde{A}_2 + A_1] &\leftrightarrow 4_7 \\
[B_2] &\leftrightarrow 4_8 \\
[\tilde{A}_2] &\leftrightarrow 8_9 \\
[A_2] &\leftrightarrow 8_9 \\
[A_1 + \tilde{A}_1] &\leftrightarrow 9_{10} \\
[2A_1] &\leftrightarrow 4_{13} \\
[A_1] &\leftrightarrow 2_{16} \\
[\tilde{A}_1] &\leftrightarrow 2_{16} \\
[A_0] &\leftrightarrow 1_{24}
\end{aligned}$$

1.13. Type E_6 .

$$\begin{aligned}
[E_6] &\leftrightarrow 1_0, \text{ dist} \\
[E_6(a_1)] &\leftrightarrow 6_1, \text{ dist} \\
[D_5] &\leftrightarrow 20_2 \\
[E_6(a_2)] &\leftrightarrow 30_3, \text{ dist} \\
[A_5] &\leftrightarrow 15_4 \\
[D_5(a_1)] &\leftrightarrow 64_4 \\
[A_4 + A_1] &\leftrightarrow 60_5 \\
[D_4] &\leftrightarrow 24_6 \\
[A_4] &\leftrightarrow 81_6 \\
[D_4(a_1)] &\leftrightarrow 80_7 \\
[A_3 + A_1] &\leftrightarrow 60_8 \\
[2A_2 + A_1] &\leftrightarrow 10_9 \\
[A_3] &\leftrightarrow 81_{10} \\
[A_2 + 2A_1] &\leftrightarrow 60_{11} \\
[2A_2] &\leftrightarrow 24_{12} \\
[A_2 + A_1] &\leftrightarrow 64_{13} \\
[A_2] &\leftrightarrow 30_{15} \\
[3A_1] &\leftrightarrow 15_{16} \\
[2A_1] &\leftrightarrow 20_{20} \\
[A_1] &\leftrightarrow 6_{25} \\
[A_0] &\leftrightarrow 1_{36}
\end{aligned}$$

1.14. Type E_7 .

$$\begin{aligned}
[E_7] &\leftrightarrow 1_0, \text{ dist} \\
[E_7(a_1)] &\leftrightarrow 7_1, \text{ dist} \\
[E_7(a_2)] &\leftrightarrow 27_2, \text{ dist} \\
[E_7(a_3)] &\leftrightarrow 56_3, \text{ dist} \\
[E_6] &\leftrightarrow 21_3 \\
[E_6(a_1)] &\leftrightarrow 120_4 \\
[D_6] &\leftrightarrow 35_4 \\
[A_7] &\leftrightarrow 189_5, \text{ dist}
\end{aligned}$$

$$\begin{aligned}
[A_6] &\leftrightarrow 105_6 \\
[D_6(a_1)] &\leftrightarrow 210_6 \\
[D_5 + A_1] &\leftrightarrow 168_6 \\
[E_7(a_4)] &\leftrightarrow 315_7, \text{ dist} \\
[D_5] &\leftrightarrow 189_7 \\
[E_6(a_2)] &\leftrightarrow 405_8 \\
[D_6(a_2)] &\leftrightarrow 280_8 \\
[(A_5 + A_1)'] &\leftrightarrow 70_9 \\
[A_5''] &\leftrightarrow 216_9 \\
[D_5(a_1) + A_1] &\leftrightarrow 378_9 \\
[D_5(a_1)] &\leftrightarrow 420_{10} \\
[A_4 + A_2] &\leftrightarrow 210_{10} \\
[A_4 + A_1] &\leftrightarrow 512_{11} \\
[A_5'] &\leftrightarrow 105_{12} \\
[D_4 + A_1] &\leftrightarrow 84_{12} \\
[A_4] &\leftrightarrow 420_{13} \\
[A_3 + A_2 + A_1] &\leftrightarrow 210_{13} \\
[A_3 + A_2] &\leftrightarrow 378_{14} \\
[D_4] &\leftrightarrow 105_{15} \\
[D_4(a_1) + A_1] &\leftrightarrow 405_{15} \\
[A_3 + A_2] &\leftrightarrow 84_{15} \\
[(A_3 + 2A_1)'] &\leftrightarrow 216_{16} \\
[D_4(a_1)] &\leftrightarrow 315_{16} \\
[(A_3 + A_1)'''] &\leftrightarrow 280_{17} \\
[2A_2 + A_1] &\leftrightarrow 70_{18} \\
[(A_3 + A_1)'] &\leftrightarrow 189_{20} \\
[A_3] &\leftrightarrow 210_{21} \\
[2A_2] &\leftrightarrow 168_{21} \\
[A_2 + 3A_1] &\leftrightarrow 105_{21} \\
[A_2 + 2A_1] &\leftrightarrow 189_{22} \\
[A_2 + A_1] &\leftrightarrow 120_{25} \\
[(4A_1)'] &\leftrightarrow 15_{28} \\
[A_2] &\leftrightarrow 56_{30} \\
[(3A_1)'''] &\leftrightarrow 35_{31} \\
[(3A_1)'] &\leftrightarrow 21_{36} \\
[2A_1] &\leftrightarrow 27_{37} \\
[A_1] &\leftrightarrow 7_{46} \\
[A_0] &\leftrightarrow 1_{63}
\end{aligned}$$

1.15. Type E_8 .

$$\begin{aligned}
[E_8] &\leftrightarrow 1_0, \text{ dist} \\
[E_8(a_1)] &\leftrightarrow 8_1, \text{ dist} \\
[E_8(a_2)] &\leftrightarrow 35_2, \text{ dist} \\
[E_8(a_4)] &\leftrightarrow 112_3, \text{ dist}
\end{aligned}$$

$$\begin{aligned}
[E_7] &\leftrightarrow 84_4 \\
[E_8(a_5)] &\leftrightarrow 210_4, \text{ dist} \\
[D_8] &\leftrightarrow 560_5, \text{ dist} \\
[E_7(a_1)] &\leftrightarrow 567_6 \\
[E_8(a_3)] &\leftrightarrow 700_6, \text{ dist} \\
[D_7] &\leftrightarrow 400_7, \\
[E_8(a_7)] &\leftrightarrow 1400_7, \text{ dist} \\
[E_8(a_6)] &\leftrightarrow 1400_8, \text{ dist} \\
[E_7(a_2)] &\leftrightarrow 1344_8 \\
[E_6 + A_1] &\leftrightarrow 448_9 \\
[D_8(a_2)] &\leftrightarrow 3240_9, \text{ dist}_2 \\
[D_7(a_1)] &\leftrightarrow 1050_{10} \\
[A_8] &\leftrightarrow 2240_{10}, \text{ dist} \\
[E_7(a_3)] &\leftrightarrow 2268_{10} \\
[E_6(a_1) + A_1] &\leftrightarrow 4096_{11} \\
[D_8(a_3)] &\leftrightarrow 1400_{11}, \text{ dist}_3 \\
[A_7'] &\leftrightarrow 175_{12} \\
[E_6] &\leftrightarrow 525_{12} \\
[D_7(a_2)] &\leftrightarrow 4200_{12} \\
[D_6] &\leftrightarrow 972_{12} \\
[E_6(a_1)] &\leftrightarrow 2800_{13} \\
[A_7 + A_1] &\leftrightarrow 4536_{13}, \text{ dist}_2 \\
[A_7'] &\leftrightarrow 6075_{14} \\
[A_6 + A_1] &\leftrightarrow 2835_{14} \\
[D_5 + A_2] &\leftrightarrow 840_{14} \\
[A_6] &\leftrightarrow 4200_{15} \\
[D_6(a_1)] &\leftrightarrow 5600_{15} \\
[E_8(a_8)] &\leftrightarrow 4480_{16}, \text{ dist} \\
[D_5 + A_1] &\leftrightarrow 3200_{16} \\
[E_7(a_4)] &\leftrightarrow 7168_{17} \\
[D_6(a_2)] &\leftrightarrow 4200_{18} \\
[E_6(a_2) + A_1] &\leftrightarrow 3150_{18} \\
[(A_5 + A_1)'] &\leftrightarrow 2016_{19} \\
[D_5(a_1) + A_2] &\leftrightarrow 1344_{19} \\
[D_5] &\leftrightarrow 2100_{20} \\
[A_4 + A_3] &\leftrightarrow 420_{20} \\
[E_6(a_2)] &\leftrightarrow 5600_{21} \\
[D_4 + A_3] &\leftrightarrow 4200_{21} \\
[(A_5 + A_1)'] &\leftrightarrow 3200_{22} \\
[D_5(a_1) + A_1] &\leftrightarrow 6075_{22} \\
[A_4 + A_2 + A_1] &\leftrightarrow 2835_{22} \\
[A_4 + A_2] &\leftrightarrow 4536_{23} \\
[A_4 + 2A_1] &\leftrightarrow 4200_{24}
\end{aligned}$$

$$\begin{aligned}
[D_4 + A_2] &\leftrightarrow 168_{24} \\
[D_5(a_1)] &\leftrightarrow 2800_{25} \\
[A_4 + A_1] &\leftrightarrow 4096_{26} \\
[(2A_3)'] &\leftrightarrow 840_{26} \\
[D_4 + A_1] &\leftrightarrow 700_{28} \\
[D_4(a_1) + A_2] &\leftrightarrow 2240_{28} \\
[A_3 + A_2 + A_1] &\leftrightarrow 1400_{29} \\
[A_4] &\leftrightarrow 2268_{30} \\
[(2A_3)'] &\leftrightarrow 3240_{31} \\
[D_4(a_1) + A_1] &\leftrightarrow 1400_{32} \\
[A_3 + A_2] &\leftrightarrow 972_{32} \\
[(A_3 + 2A_1)'] &\leftrightarrow 1050_{34} \\
[D_4] &\leftrightarrow 525_{36} \\
[2A_2 + 2A_1] &\leftrightarrow 175_{36} \\
[D_4(a_1)] &\leftrightarrow 1400_{37} \\
[A_3 + A_1] &\leftrightarrow 1344_{38} \\
[2A_2 + A_1] &\leftrightarrow 448_{39} \\
[2A_2] &\leftrightarrow 700_{42} \\
[A_2 + 3A_1] &\leftrightarrow 400_{43} \\
[A_3] &\leftrightarrow 567_{46} \\
[A_2 + 2A_1] &\leftrightarrow 560_{47} \\
[A_2 + A_1] &\leftrightarrow 210_{52} \\
[(4A_1)'] &\leftrightarrow 50_{56} \\
[A_2] &\leftrightarrow 112_{63} \\
[3A_1] &\leftrightarrow 84_{64} \\
[2A_1] &\leftrightarrow 35_{74} \\
[A_1] &\leftrightarrow 8_{91} \\
[A_0] &\leftrightarrow 1_{120}
\end{aligned}$$

2. COMPLEMENTS

2.1. Let I be the set of simple reflections of W . For any $J \subset I$ let W_J be the subgroup of W generated by J . We can view W_J as the Weyl group of a Levi subgroup L_J of a parabolic subgroup of G . Thus $CL(W_J), CL_{dist}(W_J)$ are defined in terms of L_J . The following results can be deduced from the results in §1.

(a) *If $C \in CL(W_J)$ and $C' \in cl(W)$ is defined by $C \subset C'$, then $C' \in CL(W)$. Conversely, if $C' \in CL(W)$ then there exists $J \subset I$ and $C \in CL_{dist}(W_J)$ such that $C \subset C'$.*

2.2. Let $r \in \{0\} \cup \mathcal{P}$. We state two properties similar to 0.4(a),(b).

(a) For any $E \in \mathbf{S}^{(r)}(W)$ there is a unique $C_E^{(r)} \in (\Phi^{(r)})^{-1}(E)$ which is as elliptic as possible, that is $m(C_E^{(r)}) \leq m(C)$ for any $C \in (\Phi^{(r)})^{-1}(E)$;

(b) For any $E \in \mathbf{S}^{(r)}(W)$ there is a unique $C'_E{}^{(r)} \in (\Phi^{(r)})^{-1}(E)$ which is as non-elliptic as possible, that is $m(C'_E{}^{(r)}) \leq m(C)$ for any $C \in (\Phi^{(r)})^{-1}(E)$.

Now (a) is proved in [L12]; the proof of (b) is similar. Let $CL^{(r)}(W)$ be the image of the map $E \mapsto C'_E, \mathbf{S}^{(r)}(W) \rightarrow cl(W)$. Note that $\Phi^{(r)}$ restricts to a bijection $CL^{(r)}(W) \xrightarrow{\sim} \mathbf{S}^{(r)}(W)$. Let $CL_{dist}^{(r)}(W)$ be the subset of $CL^{(r)}(W)$ corresponding to $\mathbf{S}_{disc}^{(r)} \subset \mathbf{S}^{(r)}(W)$ under this bijection. We have the following analogue of 0.4(c).

(c) *Let $C \in CL^{(r)}(W)$. We have $C \in CL_{dist}^{(r)}(W)$ if and only if C is elliptic.*

Note that $CL_{dist}(W) = \cup_r CL_{dist}^{(r)}(W)$.

2.3. For any semisimple element $s \in G = G^{(0)}$ let $E_s = j_{W_s}^W(\text{sign})$ (j -induction) where W_s is the Weyl group of the connected centralizer of s viewed as a subgroup of W . We have $E_s \in \mathbf{S}(W)$; the subset of $\mathbf{S}(W)$ formed by the E_s for various s as above is denoted by $\text{Irr}_{ss}(W)$.

Now let $E \in SS_{dist}(W)$ and let $C \in CL_{dist}(W)$ be the corresponding conjugacy class. According to 0.5(a) we have $E = \Phi^{(0)}(C)$ hence $E = \iota^{(0)}(\gamma)$ where $\gamma = \psi^{(0)}(C) \in \mathcal{U}^{(0)}$. (The elements of γ need not be distinguished.) From [L11b] there exists a semisimple element $s \in G^{(0)}$ such that $E = E_s$; thus, we have $SS_{dist}(W) \subset \text{Irr}_{ss}(W)$. (We can assume that G is almost simple; then the statement in the previous sentence holds for $E = \Phi^{(0)}(C)$ for any elliptic $C \in cl(W)$ with a single exception in type E_8 ; but that exception is not distinguished, see 1.15.) Note that s belongs to the stratum of $G^{(0)}$ that contains γ .

Thus, to $E \in \mathbf{S}_{dist}(W)$ (or $C \in CL_{dist}(W)$) one can associate a collection of reflection subgroups W_s (for various s as above). In the case where γ (as above) is in $\mathcal{U}_{dist}^{(0)}$ this collection includes the parabolic subgroup of W associated by Kostant to an element of γ . Thus the collection of subgroups that we have defined can be viewed as an extension of the Kostant parabolic subgroups.

Now, [L11a, 4.4(b)] can be reformulated as saying that the minimum length of an element in $C \in CL_{dist}(W)$ is equal to the dimension of the centralizer of s (as above) in G modulo the centre of G .

2.4. Let K be a maximal compact subgroup of $G^{(0)}$. We show:

(a) *Let X be a stratum of $G^{(0)}$ and let E be the corresponding element of $\mathbf{S}(W)$. We have $X \cap K \neq \emptyset$ if and only if $E \in \text{Irr}_{ss}(W)$.*

By the results in 2.3, we have $E \in \text{Irr}_{ss}(W)$ if and only if X contains a semisimple element of $G^{(0)}$. This last condition is clearly satisfied when $X \cap K \neq \emptyset$. Conversely, assume that X contains a semisimple element s of $G^{(0)}$. It is well known that we can find $s' \in K$ such that the connected centralizers of s and s' are conjugate. It follows that $E_s = E_{s'}$ hence s, s' belongs to the same stratum. Since $s \in X$ we have $s' \in X$ so that $X \cap K \neq \emptyset$. This proves (a).

We show:

(b) *Let X, E be as in (a). If E is distinguished then $X \cap K \neq \emptyset$.*

By (a) it is enough to show that $E \in \text{Irr}_{ss}(W)$. This follows from $SS_{dist}(W) \subset \text{Irr}_{ss}(W)$, see 2.3.

REFERENCES

- [BC76] P.Bala and R.Carter, *Classes of unipotent elements of simple algebraic groups*, Math. Proc. Camb. Phil. Soc. **79** (1976), 461-475.
- [C20] G.Carnovale, *Lusztig's strata are locally closed*, Arch.der Math. **115** (2020), 23-26.
- [LS12] M.Liebeck and G.Seitz, *Unipotent and nilpotent elements in simple algebraic groups and Lie algebras*, Math. Surveys and Monogr. 180, Amer. Math. Soc., Providence, R.I., 2012.
- [L84] G.Lusztig, *Intersection cohomology complexes on a reductive group*, Inv. Math. **75** (1984), 205-272.
- [L11a] G.Lusztig, *From conjugacy classes in the Weyl group to unipotent classes*, Represent.Th. **15** (2011), 494-530.
- [L11b] G.Lusztig, *On C -small conjugacy classes in a reductive group*, Transfor.Groups **16** (2011), 807-825.
- [L12] G.Lusztig, *From conjugacy classes in the Weyl group to unipotent classes, II*, Represent.Th. **16** (2012), 189-211.
- [L15] G.Lusztig, *On conjugacy classes in a reductive group*, Representations of Reductive Groups, Progr.in Math. 312, Birkhäuser, 2015, pp. 333-363.
- [Y20] Z.Yun, *Minimal reduction type and the Kazhdan-Lusztig map*, arxiv:2010.13642.

DEPARTMENT OF MATHEMATICS, M.I.T., CAMBRIDGE, MA 02139