

# Sets of Marginals and Pearson-Correlation-based CHSH Inequalities for a Two-Qubit System

Yuwen Huang and Pascal O. Vontobel  
Department of Information Engineering  
The Chinese University of Hong Kong  
hy018@ie.cuhk.edu.hk, pascal.vontobel@ieee.org

**Abstract**—Quantum mass functions (QMFs), which are tightly related to decoherence functionals, were introduced by Loeliger and Vontobel [IEEE Trans. Inf. Theory, 2017, 2020] as a generalization of probability mass functions toward modeling quantum information processing setups in terms of factor graphs.

Simple quantum mass functions (SQMFs) are a special class of QMFs that do *not explicitly* model classical random variables. Nevertheless, classical random variables appear *implicitly* in an SQMF if some marginals of the SQMF satisfy some conditions; variables of the SQMF corresponding to these “emerging” random variables are called classicable variables. Of particular interest are jointly classicable variables.

In this paper we initiate the characterization of the set of marginals given by the collection of jointly classicable variables of a graphical model and compare them with other concepts associated with graphical models like the sets of realizable marginals and the local marginal polytope.

In order to further characterize this set of marginals given by the collection of jointly classicable variables, we generalize the CHSH inequality based on the Pearson correlation coefficients, and thereby prove a conjecture proposed by Pozsgay *et al.* A crucial feature of this inequality is its nonlinearity, which poses difficulties in the proof.

## I. INTRODUCTION

Graphical models like factor graphs [1]–[3] have been used to represent various statistical models. In the following, we will call a factor graph consisting only of non-negative real-valued local functions a standard factor graph (S-FG). S-FGs have many applications, in particular in communications and coding theory (see, e.g., [4], [5]) and statistical mechanics (see, e.g., [6]). In these applications, factor graphs frequently represent the factorization of the joint probability mass functions (PMFs) of all the relevant random variables. Quantities of interest can then be obtained by exactly or approximately computing marginals of this joint PMF and suitably processing these marginals.

Factor graphs have also been used to represent quantum-mechanical probabilities [7], [8]. In contrast to S-FGs, these factor graphs consist of complex-valued local functions satisfying some constraints. In the following, we will call such factor graphs quantum-probability factor graphs (Q-FGs). A Q-FG is typically used to represent the factorization of the joint quantum mass function (QMF) as introduced in [7].

In this paper, we first discuss similarities and differences between PMFs and QMFs. Some of the features of QMFs will then motivate the study that is carried out in the rest of this paper.

## II. PMFs vs. QMFs

In this section, we highlight some similarities and crucial differences between PMFs and QMFs. First, we consider a classical setup. In particular, we assume that we are interested in a graphical model that represents the joint PMF  $P_{Y_1, \dots, Y_n}(y_1, \dots, y_n)$ , where  $Y_1, \dots, Y_n$  are some random variables of interest taking value in some alphabets  $\mathcal{Y}_1, \dots, \mathcal{Y}_n$ .<sup>1</sup> (In a typical application, we might have observed  $Y_1 = y_1, \dots, Y_{n-1} = y_{n-1}$  and would like to estimate  $Y_n$  based on these observations.) In most applications, the PMF  $P_{Y_1, \dots, Y_n}(y_1, \dots, y_n)$  does *not* have a “nice” factorization in terms of simple factors. However, frequently, with the introduction of suitable auxiliary variables  $x_1, \dots, x_m$  taking values in some alphabets  $\mathcal{X}_1, \dots, \mathcal{X}_m$ , respectively, there is a function  $p(\mathbf{x}, \mathbf{y})$ , where  $\mathbf{x} := (x_1, \dots, x_m)$  and  $\mathbf{y} := (y_1, \dots, y_n)$ , such that

$$\begin{aligned} p(\mathbf{x}, \mathbf{y}) &\in \mathbb{R}_{\geq 0} \quad (\text{for all } \mathbf{x}, \mathbf{y}), \\ \sum_{\mathbf{x}, \mathbf{y}} p(\mathbf{x}, \mathbf{y}) &= 1, \\ \sum_{\mathbf{x}} p(\mathbf{x}, \mathbf{y}) &= P_{\mathbf{Y}}(\mathbf{y}) \quad (\text{for all } \mathbf{y}), \end{aligned}$$

and such that  $p(\mathbf{x}, \mathbf{y})$  has a “nice” factorization. (For example, in a hidden Markov model, the joint PMF of the observations does not have a “nice” factorization, but the joint PMF of the hidden state process and the observations has a “nice” factorization.) Note that the function  $p(\mathbf{x}, \mathbf{y})$  can, thanks to its properties, be considered as a joint PMF of some random variables  $X_1, \dots, X_m, Y_1, \dots, Y_n$ .

Second, we consider a quantum-mechanical setup. We assume, again, that we are interested in a graphical model representing the joint PMF  $P_{Y_1, \dots, Y_n}(y_1, \dots, y_n)$ , where  $Y_1, \dots, Y_n$  are some random variables of interest taking values in some alphabets  $\mathcal{Y}_1, \dots, \mathcal{Y}_n$ . Such random variables can, for example, represent the measurements obtained when running some quantum-mechanical experiment, and we might be interested in estimating  $Y_n$  based on the observations

<sup>1</sup>For simplicity, in the following all alphabets will be finite.

$Y_1 = y_1, \dots, Y_{n-1} = y_{n-1}$ . As in the classical case, the PMF  $P_{Y_1, \dots, Y_n}(y_1, \dots, y_n)$  usually does *not* have a “nice” factorization in terms of simple factors. Moreover, standard physical modeling of quantum-mechanical systems shows that introducing a function  $p(\mathbf{x}, \mathbf{y})$  as defined above does usually *not* help toward obtaining a function with a “nice” factorization. However, in many quantum-mechanical setups of interest, with the introduction of suitable auxiliary variables  $x_1, \dots, x_m, x'_1, \dots, x'_m$  taking values in some alphabets  $\mathcal{X}_1, \dots, \mathcal{X}_m, \mathcal{X}'_1, \dots, \mathcal{X}'_m$  (with  $\mathcal{X}'_i = \mathcal{X}_i, i \in \{1, \dots, m\}$ ), there is a function  $q(\mathbf{x}, \mathbf{x}', \mathbf{y})$ , called quantum mass function (QMF) [7], such that

$$\begin{aligned} q(\mathbf{x}, \mathbf{x}', \mathbf{y}) &\in \mathbb{C} \quad (\text{for all } \mathbf{x}, \mathbf{x}', \mathbf{y}), \\ \sum_{\mathbf{x}, \mathbf{x}', \mathbf{y}} q(\mathbf{x}, \mathbf{x}', \mathbf{y}) &= 1, \\ q(\mathbf{x}, \mathbf{x}', \mathbf{y}) &\text{ is a PSD kernel in } (\mathbf{x}, \mathbf{x}') \text{ for every } \mathbf{y}, \\ \sum_{\mathbf{x}, \mathbf{x}'} q(\mathbf{x}, \mathbf{x}', \mathbf{y}) &= P_Y(\mathbf{y}) \quad (\text{for all } \mathbf{y}), \end{aligned}$$

and such that  $q(\mathbf{x}, \mathbf{x}', \mathbf{y})$  has a “nice” factorization. The major difference between  $p(\mathbf{x}, \mathbf{y})$  and  $q(\mathbf{x}, \mathbf{x}', \mathbf{y})$  is the fact that the former takes value in  $\mathbb{R}_{\geq 0}$ , whereas the latter takes value in  $\mathbb{C}$ . In particular,  $\sum_{\mathbf{y}} q(\mathbf{x}, \mathbf{x}', \mathbf{y})$  is in general not a PMF over  $(\mathbf{x}, \mathbf{x}')$ , thereby showing that  $\mathbf{x}, \mathbf{x}'$  *cannot* be considered as random variables. (See [7] for more details.)

In [8], the authors discussed an approach to QMFs where  $\mathbf{y}$  does *not* appear explicitly anymore, but “emerges” from a QMF. More precisely, they first introduced a simple quantum mass function (SQMF)  $q(\mathbf{x}, \mathbf{x}')$  that satisfies

$$\begin{aligned} q(\mathbf{x}, \mathbf{x}') &\in \mathbb{C}_{\geq 0} \quad (\text{for all } \mathbf{x}, \mathbf{x}'), \\ \sum_{\mathbf{x}, \mathbf{x}'} q(\mathbf{x}, \mathbf{x}') &= 1, \\ q(\mathbf{x}, \mathbf{x}') &\text{ is a PSD kernel in } (\mathbf{x}, \mathbf{x}'). \end{aligned}$$

Afterwards, they defined “classicable” variables.

**Definition 1.** Let  $\mathcal{I}$  be a subset of  $\{1, \dots, m\}$  and let  $\mathcal{I}^c := \{1, \dots, m\} \setminus \mathcal{I}$  be its complement. The variables  $\mathbf{x}_{\mathcal{I}}$  are called *jointly classicable* if the function

$$q(\mathbf{x}_{\mathcal{I}}, \mathbf{x}'_{\mathcal{I}}) := \sum_{\mathbf{x}_{\mathcal{I}^c}, \mathbf{x}'_{\mathcal{I}^c}} q(\mathbf{x}, \mathbf{x}')$$

is zero for all  $(\mathbf{x}_{\mathcal{I}}, \mathbf{x}'_{\mathcal{I}})$  satisfying  $\mathbf{x}_{\mathcal{I}} \neq \mathbf{x}'_{\mathcal{I}}$ .<sup>2</sup>

Note that if  $\mathbf{x}_{\mathcal{I}}$  are jointly classicable, then one can define the function  $p(\mathbf{x}_{\mathcal{I}}) := q(\mathbf{x}_{\mathcal{I}}, \mathbf{x}_{\mathcal{I}})$ , for which it is straightforward, thanks to the properties of SQMFs, to show that it is a PMF. It is in this sense that random variables  $y_1, \dots, y_n$  that were omitted when going from QMFs to SQMFs can “emerge” again.<sup>3</sup>

<sup>2</sup>It would be more precise to call this function  $q_{\mathcal{I}}$ . However, for conciseness, we drop the index  $\mathcal{I}$  as it can be inferred from the arguments.

<sup>3</sup>Note that there is a strong connection of SQMFs to the so-called decoherence functional [9], [10], and via this also to the consistent-histories approach to quantum mechanics [11]. However, the starting point of our investigations is quite different.

**Definition 2.** Let  $\mathcal{K}$  be a collection of subsets  $\mathcal{I}$  of  $\{1, \dots, m\}$  such that  $\mathbf{x}_{\mathcal{I}}$  is classicable.

**Example 3.** Consider the Q-FG  $N_4$  in Fig. 4, whose global function is an SQMF. In that Q-FG, the matrix  $\rho$  represents a PSD matrix and the matrices  $U_1, U_2$  are unitary matrices. One can show that for all choices of  $\rho, U_1$ , and  $U_2$ , the collection  $\mathcal{K}$  can be chosen to contain the sets  $\{1, 2\}, \{1, 4\}, \{2, 3\}$ , and  $\{3, 4\}$ .

Interestingly enough, the collection of functions  $\{p(\mathbf{x}_{\mathcal{I}})\}_{\mathcal{I} \in \mathcal{K}}$  is usually such that there is *no* PMF  $p(\mathbf{x})$  such that for every  $\mathcal{I} \in \mathcal{K}$ , the function  $p(\mathbf{x}_{\mathcal{I}})$  can be obtained as a marginal of  $p(\mathbf{x})$ .<sup>4</sup> In general, we can only guarantee that for two sets  $\mathcal{I}_1, \mathcal{I}_2 \in \mathcal{K}$  the following consistency constraint holds:

$$\sum_{\mathbf{x}_{\mathcal{I}_1 \setminus \mathcal{I}_2}} p(\mathbf{x}_{\mathcal{I}_1}) = \sum_{\mathbf{x}_{\mathcal{I}_2 \setminus \mathcal{I}_1}} p(\mathbf{x}_{\mathcal{I}_2}) \quad (\text{for all } \mathbf{x}_{\mathcal{I}_1 \cap \mathcal{I}_2}).$$

Let us comment on these special properties of  $\{p(\mathbf{x}_{\mathcal{I}})\}_{\mathcal{I} \in \mathcal{K}}$ :

- It turns out that these special properties of  $\{p(\mathbf{x}_{\mathcal{I}})\}_{\mathcal{I} \in \mathcal{K}}$  are at the heart of quantum mechanical phenomena like Hardy’s paradox [12] and the Frauchiger–Renner paradox [13].<sup>5</sup> In fact, the Q-FG  $N_4$  in Fig. 4 can be used to analyze Hardy’s paradox. On the side, note that  $N_4$  also captures the essence of Bell’s game [14].
- Interestingly, these special properties of  $\{p(\mathbf{x}_{\mathcal{I}})\}_{\mathcal{I} \in \mathcal{K}}$  are very similar to the properties of the beliefs in the local marginal polytope of an S-FG (see, e.g., [15]).<sup>6</sup>

The above observations motivate the systematic study of the collection  $\{p(\mathbf{x}_{\mathcal{I}})\}_{\mathcal{I} \in \mathcal{K}}$  for a given SQMF. Indeed, one key contribution of this paper is to study this collection for the Q-FG  $N_4$  in Fig. 4 and compare this collection with other objects that can be associated with this Q-FG.

### III. CONTRIBUTIONS






To better understand classicable variables’ marginals, we define the set  $\mathcal{M}(N_4)$ , which is the set of the marginals created by the classicable variables in the two-qubit system  $N_4$ , as shown in Fig. 4. One of our paper’s main topics is to fully characterize  $\mathcal{M}(N_4)$ . For comparison, we introduce  $\mathcal{LM}(\mathcal{K})$  (the local marginal polytope of the S-FG  $N_1$  in Fig. 1),  $\mathcal{M}(N_1)$  (the set of realizable marginals of  $N_1$ ),  $\mathcal{M}(N_2)$  (the set of realizable marginals of the Markov chain  $N_2$  in Fig. 2), and  $\mathcal{M}(N_3)$  (the set of realizable marginals of  $N_3$  in Fig. 3). We have the following results.

- We prove the Venn diagram in Fig. 6 by showing that each part in the diagram is non-empty. We can see that  $\mathcal{M}(N_3)$  and  $\mathcal{M}(N_4)$  are strict subsets of  $\mathcal{LM}(\mathcal{K})$ ; both  $\mathcal{M}(N_1)$  and  $\mathcal{M}(N_2)$  have marginals that are not in

<sup>4</sup>A similar observation is at the origin of the so-called “single-framework” rule in the consistent-histories approach to quantum mechanics.

<sup>5</sup>For a discussion of the latter in terms of SQMFs, see [8].

<sup>6</sup>Local marginal polytopes are of relevance, for example, when characterizing locally operating message-passing iterative algorithms like the sum-product algorithm [16], [17].

Fig. 1: The S-NFG  $\mathcal{N}_1$ .Fig. 2: The S-NFG  $\mathcal{N}_2$ .Fig. 3: The S-NFG  $\mathcal{N}_3$ .Fig. 4: The Q-NFG  $\mathcal{N}_4$ .Fig. 5: The Q-NFG representation of  $\beta_{i,j}(x_i, x_j)$ .Fig. 6: The Venn diagram for  $\mathcal{M}(\mathcal{N}_1)$ ,  $\mathcal{M}(\mathcal{N}_2)$ ,  $\mathcal{M}(\mathcal{K})$ ,  $\mathcal{M}(\mathcal{N}_4)$ , and  $\mathcal{LM}(\mathcal{K})$ .

$\mathcal{M}(\mathcal{N}_4)$ ; the set  $\mathcal{M}(\mathcal{N}_4)$  consists of marginals that are not compatible with any joint PMF.

- We generalize the Clauser-Horne-Shimony-Holt (CHSH) inequality [18] for Pearson correlation coefficients (PCCs), which resolves a conjecture proposed in [19]. Because PCCs are non-linear functions with respect to marginals, the inequality has a non-trivial proof. We suspect that the proof approach is applicable for proving other non-linear Bell inequalities. A violation of this inequality indicates that the associated marginals are not in  $\mathcal{M}(\mathcal{N}_3)$ .
- We illustrate Hardy's paradox, Bell's game, and the maximum quantum violation of the PCC-based CHSH inequality by the classifiable variables in  $\mathcal{N}_4$  in Fig. 4.

Besides these specific results, our paper is, more generally, about leveraging tools from factor graphs to understand certain quantities of interest in quantum information processing. In particular, given that factor graphs have been proven very

useful in classical information processing, but can also be used for doing quantum information processing, they allow one to understand and appreciate the similarities and the differences between classical and quantum information processing.

The rest of this paper is structured as follows. Section IV reviews some basics of S-FGs. In particular, Section IV-A proves the PCC-based CHSH inequality, and Section IV-B discusses the Markov chain in Fig. 2. Section V introduces  $\mathcal{N}_4$ , proves the Venn diagram in Fig. 6 and illustrates the maximum quantum violation of the PCC-based CHSH inequality. Many details are left out due to space constraints; a more detailed discussion is given in [20].

#### A. Basic Notations and Definitions

The sets  $\mathbb{R}$ ,  $\mathbb{R}_{\geq 0}$ ,  $\mathbb{R}_{> 0}$ , and  $\mathbb{C}$  denote the field of real numbers, the set of nonnegative real numbers, the set of positive real numbers, and the field of complex numbers, respectively. An overline denotes complex conjugation. For any statement  $S$ , by the Iverson's convention, the function  $[S]$  is defined to be  $[S] := 1$  if  $S$  is true and  $[S] := 0$  otherwise.

### IV. STANDARD NORMAL FACTOR GRAPHS (S-NFGS)

In this section, we review some basic concepts and properties of an S-NFG. The word "normal" refers to the fact that variables are arguments of only one or two local functions. We use an example to introduce the fundamental concepts of an S-NFG first.

**Example 4.** [1], [3] Consider the multivariate function

$$g_{\mathcal{N}_1}(x_1, \dots, x_4) := f_{1,2}(x_1, x_2) \cdot f_{1,4}(x_1, x_4) \cdot f_{3,2}(x_3, x_2) \cdot f_{3,4}(x_3, x_4),$$

where  $g_{\mathcal{N}_1}$ , the so-called global function, is defined to be the product of the so-called local functions  $f_{1,2}$ ,  $f_{1,4}$ ,  $f_{3,2}$  and  $f_{3,4}$ . We can visualize the factorization of  $g$  with the help of the S-FG  $\mathcal{N}_1$  in Fig. 1. Note that the S-FG  $\mathcal{N}_1$  consists of four function nodes  $f_{1,2}, \dots, f_{3,4}$  and four (full) edges with associated variables  $x_1, \dots, x_4$ .

For an S-NFG, a half edge is an edge incident on one function node only and a full edge is an edge incident on two function nodes.

**Definition 5.** The S-NFG  $\mathcal{N}(\mathcal{F}(\mathcal{N}), \mathcal{E}(\mathcal{N}), \mathcal{X}(\mathcal{N}))$  consists of:

- 1) The graph  $(\mathcal{F}(\mathcal{N}), \mathcal{E}(\mathcal{N}))$  with vertex set  $\mathcal{F}(\mathcal{N})$  and edge set  $\mathcal{E}(\mathcal{N})$ , where  $\mathcal{E}(\mathcal{N})$  consists of all full edges and half edges in  $\mathcal{N}$ . With some slight abuse of notation, an  $f \in \mathcal{F}(\mathcal{N})$  will denote a function node and the corresponding local function.
- 2) The alphabet  $\mathcal{X}(\mathcal{N}) := \prod_{e \in \mathcal{E}(\mathcal{N})} \mathcal{X}_e$ , where  $\mathcal{X}_e$  is the alphabet associated with the edge  $e \in \mathcal{E}(\mathcal{N})$ .

**Definition 6.** Given  $\mathcal{N}(\mathcal{F}(\mathcal{N}), \mathcal{E}(\mathcal{N}), \mathcal{X}(\mathcal{N}))$ , we make the following definitions:

- 1) For every function node  $f \in \mathcal{F}(\mathcal{N})$ , the set  $\partial f$  is the set of edges incident on  $f$ .
- 2) An assignment  $\mathbf{x} := (x_e)_{e \in \mathcal{E}(\mathcal{N})} \in \mathcal{X}(\mathcal{N})$  is called a configuration of the S-NFG.

- 3) The local function  $f$  associated with function node  $f \in \mathcal{F}(\mathbf{N})$  denotes an arbitrary mapping  $f : \prod_{e \in \partial f} \mathcal{X}_e \rightarrow \mathbb{R}_{\geq 0}$ .
- 4) The global function is  $g_{\mathbf{N}}(\mathbf{x}) := \prod_{f \in \mathcal{F}(\mathbf{N})} f(\mathbf{x}_{\partial f})$ .
- 5) The partition function is  $Z(\mathbf{N}) := \sum_{\mathbf{x}} g_{\mathbf{N}}(\mathbf{x})$ , where  $\sum_{\mathbf{x}}$  denotes  $\sum_{\mathbf{x} \in \mathcal{X}(\mathbf{N})}$ .
- 6) The PMF induced on  $\mathbf{N}$  is  $p_{\mathbf{N}}(\mathbf{x}) := g_{\mathbf{N}}(\mathbf{x})/Z(\mathbf{N})$ .
- 7) Let  $\mathcal{I}$  be a subset of  $\mathcal{E}(\mathbf{N})$  and let  $\mathcal{I}^c := \mathcal{E}(\mathbf{N}) \setminus \mathcal{I}$  be its complement. The marginal  $p_{\mathbf{N}, \mathcal{I}}(\mathbf{x}_{\mathcal{I}})$  is defined to be  $p_{\mathbf{N}, \mathcal{I}}(\mathbf{x}_{\mathcal{I}}) := \sum_{\mathbf{x}_{\mathcal{I}^c}} p_{\mathbf{N}}(\mathbf{x})$ .

**Definition 7.** Considering  $\mathbf{N} \in \{\mathbf{N}_1, \mathbf{N}_2, \mathbf{N}_3\}$ , we make the following definitions:

- 1) The alphabet  $\mathcal{X}_e$  is  $\mathcal{X}_e := \{0, 1\}$  for all  $e \in \mathcal{E}(\mathbf{N})$ .
- 2) The set  $\mathcal{K}$  is  $\mathcal{K} := \{\{1, 2\}, \{1, 4\}, \{2, 3\}, \{3, 4\}\}$ .
- 3) For  $\{i, j\} \in \mathcal{K}$ , the marginal  $p_{\mathbf{N}, i, j}$  is defined to be a  $|\mathcal{X}_e|$ -by- $|\mathcal{X}_e|$  matrix with the entry  $p_{\mathbf{N}, i, j}(x_i, x_j)$  and the marginal  $p_{\mathbf{N}, i}$  is defined to be a  $|\mathcal{X}_e|$ -by- $|\mathcal{X}_e|$  diagonal matrix with  $p_{\mathbf{N}, i}(x_i)$  being the  $x_i$ -th diagonal term.
- 4) The collection of matrices  $\beta$  is defined to be  $\beta := ((\beta_{i, j})_{\{i, j\} \in \mathcal{K}}, (\beta_i)_{i \in \mathcal{E}(\mathbf{N}_1)})$ . In particular, the matrix  $\beta_{i, j}$  is defined to be a  $|\mathcal{X}_e|$ -by- $|\mathcal{X}_e|$  matrix with entry  $\beta_{i, j}(x_i, x_j) \in \mathbb{R}_{\geq 0}$  and the matrix  $\beta_i$  is defined to be a  $|\mathcal{X}_e|$ -by- $|\mathcal{X}_e|$  diagonal matrix with  $\beta_i(x_i) \in \mathbb{R}_{\geq 0}$  being the  $x_i$ -th diagonal term.
- 5) The set of realizable marginals of  $\mathbf{N}$  is defined to be

$$\mathcal{M}(\mathbf{N}) := \left\{ \beta \left| \begin{array}{l} \text{there exists an } \mathcal{F}(\mathbf{N}) \text{ such that} \\ \beta_{i, j} = p_{\mathbf{N}, i, j}, \beta_i = p_{\mathbf{N}, i}, \{i, j\} \in \mathcal{K} \end{array} \right. \right\}.$$

- 6) The set  $\mathcal{LM}(\mathcal{K})$  is defined to be

$$\mathcal{LM}(\mathcal{K}) := \left\{ \beta \left| \begin{array}{l} 0 \leq \beta_{i, j}(x_i, x_j) \leq 1, \forall x_i, x_j, i, j \\ \sum_{x_j} \beta_{i, j}(x_i, x_j) = \beta_i(x_i), \forall x_i, i \\ \sum_{x_i} \beta_{i, j}(x_i, x_j) = \beta_j(x_j), \forall x_j, j \\ \sum_{x_i} \beta_i(x_i) = 1, \forall i \end{array} \right. \right\}.$$

The set  $\mathcal{LM}(\mathcal{K})$  is essentially the local marginal polytope of the S-NFG  $\mathbf{N}_1$  in Fig. 1. The definition of the local marginal polytope for an S-NFG is given in [15, Section 4.1.1].

- 7) For each  $\beta \in \mathcal{LM}(\mathcal{K})$  and  $\{i, j\} \in \mathcal{K}$ , each marginal  $\beta_{i, j}$  can be used to represent the PMF for two random variables  $Y_1, Y_2 \in \mathcal{X}_e$  by setting the probability  $\Pr(Y_1 = x_i, Y_2 = x_j) = \beta_{i, j}(x_i, x_j)$ ,  $x_i, x_j \in \mathcal{X}_e$ . The functions  $\text{Cov}(Y_1, Y_2)$ ,  $\text{Var}(Y_1)$ ,  $\text{Var}(Y_2)$  are defined to be the covariance of  $Y_1$  and  $Y_2$ , and the variances of  $Y_1$  and  $Y_2$ , respectively. When  $\text{Var}(Y_1), \text{Var}(Y_2) > 0$ , the PCC of  $Y_1$  and  $Y_2$  is defined to be  $\text{Corr}(\beta_{i, j}) := \text{Cov}(Y_1, Y_2) / \sqrt{\text{Var}(Y_1) \cdot \text{Var}(Y_2)}$ .

When there is no ambiguity, we use short-hands  $(\cdot)_{i, j}$ ,  $(\cdot)_i$ ,  $\sum_{x_i}$ , and  $\{\cdot\}_{x_i}$  for  $(\cdot)_{\{i, j\} \in \mathcal{K}}$ ,  $(\cdot)_{i \in \mathcal{E}(\mathbf{N}_1)}$ ,  $\sum_{x_i \in \mathcal{X}_e}$ , and  $\{\cdot\}_{x_i \in \mathcal{X}_e}$ , respectively.

Because  $\mathcal{LM}(\mathcal{K})$  is a convex set by definition, Carathéodory's theorem [21, Proposition B.6] states that each element in  $\mathcal{LM}(\mathcal{K})$  can be written as a convex combination of the vertices in  $\mathcal{LM}(\mathcal{K})$ . The full list of the vertices in  $\mathcal{LM}(\mathcal{K})$  is given in [20, Appendix A].

**Proposition 8.** For  $\{i, j\} \in \mathcal{K}$  and  $0 < \beta_i(0), \beta_j(0) < 1$ , the PCC  $\text{Corr}(\beta_{i, j})$  satisfies

$$\text{Corr}(\beta_{i, j}) = \frac{\det(\beta_{i, j})}{\sqrt{\det(\beta_i) \cdot \det(\beta_j)}}.$$

The requirement  $0 < \beta_i(0), \beta_j(0) < 1$  ensures that  $\det(\beta_i), \det(\beta_j) > 0$ , and thus  $\text{Corr}(\beta_{i, j})$  is well-defined.

*Proof.* See the proof of [20, Corollary 9].  $\blacksquare$

**Definition 9.** Suppose that  $\beta \in \mathcal{LM}(\mathcal{K})$  and  $0 < \beta_i(0) < 1$ ,  $i \in \mathcal{E}(\mathbf{N}_1)$ , we define

$$\text{CorrCHSH}(\beta) :=$$

$$\text{Corr}(\beta_{1, 2}) + \text{Corr}(\beta_{1, 4}) + \text{Corr}(\beta_{3, 2}) - \text{Corr}(\beta_{3, 4}).$$

#### A. Properties for $\mathbf{N}_3$

In this subsection, we prove inequalities with respect to  $\text{CorrCHSH}(\beta)$  for  $\beta \in \mathcal{M}(\mathbf{N}_3)$ . These inequalities genuinely are (nonlinear) Bell inequalities [22] in the usual sense. By definition, it holds that

$$\mathcal{M}(\mathbf{N}_1) \subseteq \mathcal{M}(\mathbf{N}_3), \quad \mathcal{M}(\mathbf{N}_2) \subseteq \mathcal{M}(\mathbf{N}_3),$$

so any inequality that holds for all  $\beta \in \mathcal{M}(\mathbf{N}_3)$  also holds for all  $\beta \in \mathcal{M}(\mathbf{N}_1) \cup \mathcal{M}(\mathbf{N}_2)$ .

**Theorem 10.** For any  $\beta \in \mathcal{M}(\mathbf{N}_3)$  such that  $0 < \beta_i(0) < 1$  for all  $i \in \mathcal{E}(\mathbf{N}_1)$ , we have

$$|\text{CorrCHSH}(\beta)| < 2\sqrt{2}.$$

*Proof.* We prove it by contradiction. On the one hand, the set  $\mathcal{M}(\mathbf{N}_3)$  consists of marginals for binary random variables only. On the other hand, to have  $\text{CorrCHSH}(\beta) = 2\sqrt{2}$  for some  $\beta \in \mathcal{M}(\mathbf{N}_3)$ , the PMF realizing  $\beta$  needs to be the joint PMF for random variables with alphabet size greater than two. For details, see the proof in [20, Appendix C].  $\blacksquare$

The main idea in the proof of Theorem 10 can be used to verify whether a proposed bound for a function with binary random variables is achievable. It is different from the idea in the proof of the upcoming Theorem 11.

**Theorem 11.** For any  $\beta \in \mathcal{M}(\mathbf{N}_3)$  such that  $0 < \beta_i(0) < 1$  for all  $i \in \mathcal{E}(\mathbf{N}_1)$ , we have  $|\text{CorrCHSH}(\beta)| \leq 5/2$ .

*Proof.* We give a proof sketch here. For details, see the proof in [20, Appendix E].

- Consider a subset of  $\mathcal{LM}(\mathcal{K})$  such that in this subset,  $0 < \beta_i(0) < 1$  for  $i \in \mathcal{E}(\mathbf{N}_1)$ , and the elements in  $\beta$  satisfy the original linear CHSH inequality. Denote this set as  $\mathcal{LM}_{\text{CHSH}}(\mathcal{K})$ . We have  $\mathcal{M}(\mathbf{N}_3) \subseteq \mathcal{LM}_{\text{CHSH}}(\mathcal{K})$ .
- Find a  $\beta^* \in \mathcal{M}(\mathbf{N}_3)$  such that  $\text{CorrCHSH}(\beta^*) = 5/2$ .
- We formulate an optimization problem where  $\text{CorrCHSH}(\beta)$  is maximized over  $\beta \in \mathcal{LM}_{\text{CHSH}}(\mathcal{K})$  such that  $\beta$  has a similar structure as  $\beta^*$ , e.g., having the same number of zero entries in  $(\beta_{i, j})_{i, j}$ . Note that this optimization problem has linear constraints only, which helps determine the optimal solution. We prove  $\text{CorrCHSH}(\beta) \leq 5/2$  in this case.

- We generalize the proof for all  $\beta \in \mathcal{LM}_{\text{CHSH}}(\mathcal{K})$ .
- The proof of  $\text{CorrCHSH}(\beta) \geq -5/2$  is similar. ■

Theorem 11 proves the conjecture stated in [19]. The key idea of the proof is that we consider  $\mathcal{LM}_{\text{CHSH}}(\mathcal{K})$  instead of  $\mathcal{M}(\mathbb{N}_3)$ . Suppose that we want to prove  $\text{CorrCHSH}(\beta) \leq 5/2$  for  $\beta \in \mathcal{M}(\mathbb{N}_3)$  directly. Because  $\mathcal{M}(\mathbb{N}_3)$  is a convex set, for any  $\beta \in \mathcal{M}(\mathbb{N}_3)$ , the marginal  $\beta_{i,j}$  can be written as a convex combination of some joint PMF for  $X_1, \dots, X_4$ , i.e.,  $\{p_{\mathbb{N}_3}(\mathbf{x})\}_{\mathbf{x}}$ , which makes the expression of  $\text{CorrCHSH}(\beta)$  non-trivial. By considering a superset of  $\mathcal{M}(\mathbb{N}_3)$ , i.e.,  $\mathcal{LM}_{\text{CHSH}}(\mathcal{K})$ , we can simplify  $\text{CorrCHSH}(\beta)$ . We suspect that this idea can be generalized in the proof of other non-linear Bell inequalities.

### B. Markov Chain in Fig. 2

In this subsection, we consider the Markov chain  $\mathbb{N}_2$  in Fig. 2.

**Theorem 12.** *For the Markov chain  $\mathbb{N}_2$  in Fig. 2, we have*

$$\text{Corr}(\beta_{3,4}) = \text{Corr}(\beta_{3,2}) \cdot \text{Corr}(\beta_{1,2}) \cdot \text{Corr}(\beta_{1,4}).$$

*Proof.* See [23, Corollary 19]. ■

**Corollary 13.** *For the Markov chain  $\mathbb{N}_2$  in Fig. 2, it holds that  $|\text{Corr}(\beta_{3,4})| \leq |\text{Corr}(\beta_{1,2})| \leq 1$ .*

*Proof.* It can be proven using Theorem 12 and  $|\text{Corr}(\beta_{i,j})| \leq 1$  for  $\{i, j\} \in \mathcal{K}$ . ■

We prove another variation of the PCC-based CHSH inequality for  $\mathbb{N}_2$ .

**Corollary 14.** *For the Markov chain  $\mathbb{N}_2$  in Fig. 2, we have*

$$|\text{Corr}(\beta_{1,2}) + \text{Corr}(\beta_{2,4}) + \text{Corr}(\beta_{1,3}) - \text{Corr}(\beta_{3,4})| \leq 2.$$

*Proof.* See the proof of [20, Proposition 20]. ■

## V. QUANTUM-PROBABILITY NORMAL FACTOR GRAPHS (Q-NFGs)

This section considers a quantum system represented by the Q-NFG  $\mathbb{N}_4$  in Fig. 4. Such Q-NFGs have been discussed thoroughly in [7], [8]. Note that in Fig. 4 and Fig. 5, the row index of a matrix is marked by a black dot. The details of  $\mathbb{N}_4$  are shown in [20, Definition 21].

**Proposition 15.** *For any  $\{i, j\} \in \mathcal{K}$ , the variables  $\tilde{x}_i$  and  $\tilde{x}_j$  are jointly classicable, which implies that the marginals  $q_{i,j}(\tilde{x}_i, \tilde{x}_j)$  and  $q_i(\tilde{x}_i)$  are non-negative real numbers for any  $\tilde{x}_i, \tilde{x}_j \in \mathcal{X}_e^2$ .*

*Proof.* It can be proven directly by Definition 1. ■

Then we define the set of realizable marginals of  $\mathbb{N}_4$  based on the jointly classicable variables  $\tilde{x}_i$  and  $\tilde{x}_j$  for all  $\{i, j\} \in \mathcal{K}$ .

**Definition 16.** *With  $\tilde{0} := (0, 0)$ ,  $\tilde{1} := (1, 1)$ , and  $\{i, j\} \in \mathcal{K}$ , the matrices  $\mathbf{q}_{i,j}$  and  $\mathbf{q}_i$  induced by  $q_{\mathbb{N}_4}$  are defined to be*

$$\mathbf{q}_{i,j} := \begin{pmatrix} q_{i,j}(\tilde{0}, \tilde{0}) & q_{i,j}(\tilde{0}, \tilde{1}) \\ q_{i,j}(\tilde{1}, \tilde{0}) & q_{i,j}(\tilde{1}, \tilde{1}) \end{pmatrix}, \quad \mathbf{q}_i := \begin{pmatrix} q_i(\tilde{0}) & 0 \\ 0 & q_i(\tilde{1}) \end{pmatrix}.$$

The set of realizable marginals of  $\mathbb{N}_4$  is defined to be the set  $\mathcal{M}(\mathbb{N}_4) := \{\beta \mid \beta_{i,j} = \mathbf{q}_{i,j}, \beta_i = \mathbf{q}_i, \{i, j\} \in \mathcal{K}\}$ , which is not the set of quantum correlations in the usual Bell nonlocality sense.

**Proposition 17.** *For any  $\beta \in \mathcal{M}(\mathbb{N}_4)$ , there exist matrices  $\rho$ ,  $U_1$ , and  $U_2$  such that*

$$\beta_{i,j}(x_i, x_j) = \text{Tr}((A_{i,x_i} \otimes B_{j,x_j}) \cdot \rho \cdot (A_{i,x_i} \otimes B_{j,x_j})^H), \quad (1)$$

for all  $x_i, x_j \in \mathcal{X}_e$  and  $\{i, j\} \in \mathcal{K}$ , where

$$A_{i,x_i} := E_{x_i} \cdot U_1^{[i=3]}, \quad B_{j,x_j} := E_{x_j} \cdot U_2^{[j=4]}, \\ E_{x_i}(y_i, y'_i) := [y_i = x_i] \cdot [y_i = y'_i], \quad x_i, y_i, y'_i \in \mathcal{X}_e.$$

Note that the set  $\{E_{x_i}\}_{x_i}$  denotes the measurement of a single qubit in the computational basis. Then we have

$$\sum_{x_i \in \mathcal{X}_e} A_{i,x_i}^H \cdot A_{i,x_i} = \sum_{x_j \in \mathcal{X}_e} B_{j,x_j}^H \cdot B_{j,x_j} = I, \quad \{i, j\} \in \mathcal{K},$$

which shows that both  $\{A_{i,x_i}\}_{x_i}$  and  $\{B_{j,x_j}\}_{x_j}$  are sets of measurement matrices with binary outcomes  $x_i$  and  $x_j$ , respectively.

*Proof.* It can be proven directly. ■

After closing the dashed box in Fig. 5, i.e., summing over the variables inside the box, we obtain (1).

**Proposition 18.** *There exists a  $\beta \in \mathcal{M}(\mathbb{N}_4)$  such that  $|\text{Corr}(\beta_{3,4})| > |\text{Corr}(\beta_{1,2})|$ .*

*Proof.* See [20, Proposition 26]. ■

Compared with Corollary 13, Proposition 18 implies that  $\mathcal{M}(\mathbb{N}_4)$  provides extra  $\beta$  that is not in  $\mathcal{M}(\mathbb{N}_2)$ .

**Theorem 19.** *The Venn diagram in Fig. 6 holds.*

*Proof.* See the proof of [20, Theorem 49]. ■

We make some remarks on the Venn diagram in Fig. 6:

- On the one hand, the set  $\mathcal{M}(\mathbb{N}_4)$  provides extra marginals that are not in  $\mathcal{M}(\mathbb{N}_3)$ . For example, by introducing entanglement in the quantum system, one can obtain a set of incompatible marginals (see, e.g., [20, Example 30]).
- On the other hand, the sets  $\mathcal{M}(\mathbb{N}_1)$ ,  $\mathcal{M}(\mathbb{N}_2)$ , and  $\mathcal{M}(\mathbb{N}_3)$  also consist of marginals that are not in  $\mathcal{M}(\mathbb{N}_4)$ .

**Proposition 20.** *For  $\beta \in \mathcal{M}(\mathbb{N}_4)$  satisfying  $0 < \beta_i(0) < 1$  for all  $i \in \mathcal{E}(\mathbb{N}_1)$ , we have  $|\text{CorrCHSH}(\beta)| \leq 2\sqrt{2}$ .*

*Proof.* See [19, Appendix B]. ■

**Proposition 21.** *Hardy's paradox [12] and Bell's game [14] can be illustrated via the classicable variables induced by the SQMF  $q_{\mathbb{N}_4}(\tilde{\mathbf{x}})$ . In Bell's game, we have  $\text{CorrCHSH}(\beta) = 2\sqrt{2}$ , which also realizes the maximum quantum violation of  $\text{CorrCHSH}(\beta)$  as proven in Proposition 20.*

*Proof.* See [20, Example 30, Proposition 31]. ■

## REFERENCES

- [1] F. R. Kschischang, B. J. Frey, and H.-. Loeliger, "Factor graphs and the sum-product algorithm," *IEEE Trans. Inf. Theory*, vol. 47, no. 2, pp. 498–519, Feb. 2001.
- [2] G. D. Forney, "Codes on graphs: normal realizations," *IEEE Trans. Inf. Theory*, vol. 47, no. 2, pp. 520–548, Feb. 2001.
- [3] H.-A. Loeliger, "An introduction to factor graphs," *IEEE Signal Process. Mag.*, vol. 21, no. 1, pp. 28–41, Jan. 2004.
- [4] H. Wymeersch, *Iterative Receiver Design*. Cambridge, U.K.: Cambridge Univ. Press, 2007.
- [5] T. Richardson and R. Urbanke, *Modern Coding Theory*. Cambridge, U.K.: Cambridge Univ. Press, 2008.
- [6] M. Mézard and A. Montanari, *Information, Physics and Computation*. Oxford, U.K.: Oxford Univ. Press, 2009.
- [7] H.-A. Loeliger and P. O. Vontobel, "Factor graphs for quantum probabilities," *IEEE Trans. Inf. Theory*, vol. 63, no. 9, pp. 5642–5665, Sep. 2017.
- [8] H. Loeliger and P. O. Vontobel, "Quantum measurement as marginalization and nested quantum systems," *IEEE Trans. Inf. Theory*, vol. 66, no. 6, pp. 3485–3499, Jun. 2020.
- [9] M. Gell-Mann and J. B. Hartle, "quantum mechanics in the light of quantum cosmology," in *Proc. Santa Fe Institute Workshop on Complexity, Entropy, and the Physics of Information*, May 1989.
- [10] H. F. Dowker and J. J. Halliwell, "Quantum mechanics of history: The decoherence functional in quantum mechanics," *Phys. Rev. D*, vol. 46, pp. 1580–1609, Aug. 1992.
- [11] R. B. Griffiths, *Consistent Quantum Theory*. Cambridge Univ. Press, 2002.
- [12] L. Hardy, "Quantum mechanics, local realistic theories, and Lorentz-invariant realistic theories," *Phys. Rev. Lett.*, vol. 68, pp. 2981–2984, May 1992.
- [13] D. Frauchiger and R. Renner, "Quantum theory cannot consistently describe the use of itself," *Nature Communications*, vol. 9, no. 3711, 2018.
- [14] N. Gisin, *Quantum Chance: Nonlocality, Teleportation and Other Quantum Marvels*. USA: Copernicus, 2014.
- [15] M. J. Wainwright and M. I. Jordan, "Graphical models, exponential families, and variational inference," *Foundation and Trends Machine Learning*, vol. 1, no. 1–2, pp. 1–305, 2008.
- [16] J. S. Yedidia, W. T. Freeman, and Y. Weiss, "Constructing free-energy approximations and generalized belief propagation algorithms," *IEEE Trans. Inf. Theory*, vol. 51, no. 7, pp. 2282–2312, Jul. 2005.
- [17] P. O. Vontobel, "Counting in graph covers: A combinatorial characterization of the Bethe entropy function," *IEEE Trans. Inf. Theory*, vol. 59, no. 9, pp. 6018–6048, Sep. 2013.
- [18] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, "Proposed experiment to test local hidden-variable theories," *Phys. Rev. Lett.*, vol. 23, pp. 880–884, Oct. 1969.
- [19] V. Pozsgay, F. Hirsch, C. Branciard, and N. Brunner, "Covariance Bell inequalities," *Phys. Rev. A*, vol. 96, p. 062128, Dec. 2017.
- [20] Y. Huang and P. O. Vontobel, "Sets of marginals and Pearson-correlation-based CHSH inequalities for a two-qubit system (extended version)," 2021. [Online]. Available: [https://staff.ie.cuhk.edu.hk/~vpascal/students/huang\\_vontobel\\_isit2021\\_long.pdf](https://staff.ie.cuhk.edu.hk/~vpascal/students/huang_vontobel_isit2021_long.pdf)
- [21] D. Bertsekas, *Nonlinear Programming*, ser. Athena scientific optimization and computation series. Athena Scientific, 2016.
- [22] A. Fine, "Hidden variables, joint probability, and the Bell inequalities," *Phys. Rev. Lett.*, vol. 48, pp. 291–295, Feb. 1982.
- [23] R. Mori, "Loop calculus for non-binary alphabets using concepts from information geometry," *IEEE Trans. Inf. Theory*, vol. 61, no. 4, pp. 1887–1904, Apr. 2015.