

Almost radial gauge

Andrzej Herdegen

Abstract. An almost radial gauge A^{ar} of the electromagnetic potential is constructed for which $x \cdot A^{\text{ar}}(x)$ vanishes arbitrarily fast in timelike directions. This potential is in the class introduced by Dirac with the purpose of forming gauge-invariant quantities in quantum electrodynamics. In quantum case the construction of smeared operators $A^{\text{ar}}(K)$ is enabled by a natural extension of the free electromagnetic field algebra introduced earlier (represented in a Hilbert space). The space of possible smearing functions K includes vector fields with the asymptotic spacetime behavior typical for scattered currents (the conservation condition in the whole spacetime need not be assumed). This construction is motivated by a possible application to the infrared problem in QED.

Mathematics Subject Classification (2010). Primary 81V10; Secondary 81T05.

Keywords. quantum electrodynamics, infrared problems, radial gauge.

1. Introduction

The long-range nature of interaction entails the so called infrared problems in electrodynamics, both classical and quantum [17]. Some of the quantum problems in this area are specific for this realm, but other reveal themselves already on the classical level. The identification of asymptotic charged fields and the scattering theory belong to the latter class.

The present author has long followed the idea that an appropriate choice of the electromagnetic gauge may relieve the scattering infrared problems. After preliminary results [12], it was shown recently that indeed the classical asymptotic problem for scattering of the Dirac field in electromagnetic time-dependent field, typical for the complete theory, disappears in certain gauges. The main constituent feature of this class is sufficiently fast vanishing of the product $x \cdot A(x)$ for x tending to timelike infinity (with arbitrarily fixed origin $x = 0$ in Minkowski space) [16]. The choice of a specific potential in this class has a large freedom in this classical context. However, we plan to extend our

analysis to quantum electrodynamics, and in that case a judicious choice of our gauge is of fundamental importance.

The first gauge which probably comes to mind is the radial gauge $x \cdot A^{\text{rg}}(x) = 0$, going also in literature by the names Fock-Schwinger or relativistic Poincaré gauge ([9], [23]; see also [11]). In classical electrodynamics this gauge is well defined and may be obtained by integration of the electromagnetic field tensor. In quantum theory the latter property would be an advantage, as no indefinite metric space would be needed. However, we shall indicate in the following that the quantum version of this gauge is singular, which explains the need for *a posteriori* regularization in attempts to construct perturbative calculus with the use of this gauge [19].

It is the purpose of the present article to construct an ‘almost radial gauge’, which may also be obtained from the electromagnetic field tensor alone, and for which $x \cdot A^{\text{ar}}(x)$ vanishes fast in timelike directions, but which at the same time admits quantization. Almost radial gauge turns out to be in the Dirac family of ‘gauge-invariant gauges’, which in standard formulation of QED may be constructed only after initial regularization and the use of indefinite Gupta-Bleuler metric (see the discussion of this problem in [24], p. 191). We choose a different approach. For the construction of the quantum field we use a natural extension of the algebra of the free field proposed earlier [13], which enables representation of infrared-singular fields. This extension admits the construction of the almost radial gauge, and allows moreover its well defined smearing with vector functions typically appearing in scattering. We hope to apply our construction to infrared problems in QED in future.

As we shall see, the construction of the almost radial gauge involves integration of the electromagnetic field along radial \mathbb{R} -axes. This bears some similarity to the integration of the field along spacelike semi-axes, which is the principle of the construction of string-localized potentials introduced by Mund, Schroer and Yngvason [21]. We shall discuss similarities and differences of the two constructions.

The plan of the article is as follows. In Section 2 we summarize our notation and mathematical tools to be used in the article. Section 3 contains a brief summary of the quantization of the standard free field algebra, and its extension mentioned above. The classical radial gauge, and its singularity in quantum case, are discussed in Section 4. Section 5 contains our main results: construction and evaluation of properties of the almost radial gauge, both classical and quantum, along the lines announced above. In Section 6 we compare our construction with the string-localized potentials. Proofs of some technical points are shifted to Appendix.

2. Preliminaries

We list here our notation, conventions and preliminary formulas. More extensive summary of these questions may be found in [15].

We consider the flat spacetime with a fixed origin, thus described by the Minkowski vector space M with the scalar product $x \cdot y$ with signature $(+, -, -, -)$. We choose physical units for which \hbar , c and an arbitrarily chosen length scale are all equal to 1. Then, in particular, $x \in M$ may be treated as dimensionless. We fix a unit timelike, future-pointing vector t and for a Minkowski vector x we write $x^0 = x \cdot t$ and $\mathbf{x} = x - (x \cdot t)t$. The 3-dimensional norm of \mathbf{x} is denoted $|\mathbf{x}|$, and we also write $|x| = (|x^0|^2 + |\mathbf{x}|^2)^{\frac{1}{2}}$. The tensor indices, which are often suppressed, are denoted by a, b etc. The Minkowski volume measure element is denoted dx . Our results do not depend on the choice of vector t .

2.1. Homogeneous functions on the future lightcone

We shall write l for any future-pointing lightlike (nonzero) vector and denote

$$C_+ = \{l \in M \mid l \cdot l = 0, l^0 > 0\}, \quad C_+^t = \{l \in C_+ \mid l^0 = 1\}.$$

For C^1 -functions $f(l)$ on C_+ , the intrinsic differentiation operators

$$L_{ab} = l_a \frac{\partial}{\partial l^b} - l_b \frac{\partial}{\partial l^a}$$

are the generators of the Lorentz transformations $f(l) \mapsto f_\Lambda(l) = f(\Lambda^{-1}l)$, where Λ is a Lorentz transformation of the Minkowski space. The derivative $\partial f / \partial l^a$ can only be given unique meaning if one determines a particular C^1 -extension of f to a neighborhood of C_+ . However, any two such extensions differ by a term $l^2 g(l)$, with a C^1 -function $g(l)$ in a neighborhood of C_+ . Thus on C_+ the derivatives $\partial f(l)$ obtained from these two extensions differ by $2l g(l)$. Therefore, if we define

$$\partial_a f(l) = (l^0)^{-1} L_{0a} f(l)$$

in any chosen reference frame, then $\partial_a f(l)$ is uniquely defined as an equivalence class with respect to the addition of terms proportional to l_a . If the definition of $f(l)$ is the restriction of a function naturally defined in some neighborhood of C_+ , then this function may also serve to define $\partial_a f(l)$.

For any measurable function $W(l)$ on C_+ , homogeneous of degree -2 , the integral over the set of null directions defined by

$$\int W(l) d^2l = \int_{C_+^t} W(l) d\Omega_t(l),$$

where $d\Omega_t(l)$ is the angle measure on the unit sphere, does not depend on the choice of the vector t . If W is C^1 , it follows that

$$\int L_{ab} W(l) d^2l = 0.$$

Another closely related identity is the following. Let $V^a(l)$ be a C^1 -vector function on C_+ , such that $l \cdot V(l) = 0$. Suppose $V(l)$ is extended to a neighborhood of the cone with the preservation of these properties. Then in each

Minkowski frame one has

$$L_{0a} \left(\frac{V^a}{l^0} \right) = \partial \cdot V(l),$$

so the rhs of this equation is both frame- and extension (preserving orthogonality)-independent. Moreover, if in addition $V(l)$ is homogeneous of degree -1 , then

$$\int \partial \cdot V(l) d^2l = 0. \quad (1)$$

If $f(l)$ is a scalar C^1 -function, homogeneous of degree 0, then the last identity and the earlier remarks on $\partial_a f(l)$ imply the ‘integration by parts’ identity

$$\int f(l) \partial \cdot V(l) d^2l = - \int V(l) \cdot \partial f(l) d^2l. \quad (2)$$

We end these remarks with the following definition. Let $F(l)$ be any continuous function, homogeneous of degree -1 . Then it is easily calculated that

$$\lim_{\epsilon \searrow 0} \frac{1}{2\pi} \int \delta(l \cdot l' - \epsilon l^0 l'^0) F(l') d^2l' = F(l),$$

independently of the frame in which the coordinates $l^0 l'^0$ are taken, where $\delta(\cdot)$ is the Dirac delta distribution. Therefore, we can interpret the lhs as the action of a distribution $\delta(l, l')$, homogeneous of degree -1 in each of the arguments:

$$\int \delta(l, l') F(l') d^2l' = F(l).$$

2.2. Fourier transforms

We use the following conventions for a function f on M and a function g on $\mathbb{R} \times C_+$:

$$\begin{aligned} \hat{f}(p) &= \frac{1}{2\pi} \int_M f(x) e^{ip \cdot x} dx, \\ \tilde{g}(\omega, l) &= \frac{1}{2\pi} \int_{\mathbb{R}} g(s, l) e^{i\omega s} ds \end{aligned} \quad (3)$$

2.3. Wave equation

Solutions of the homogeneous wave equation may be represented by

$$A(x) = -\frac{1}{2\pi} \int \dot{V}(x \cdot l, l) d^2l, \quad (4)$$

where¹ $\dot{V}(s, l)$ (of any algebraic type) is a function homogeneous of degree -2 . This representation is related to the Fourier representation

$$A(x) = \frac{1}{\pi} \int_M e^{-ix \cdot k} a(k) \operatorname{sgn}(k^0) \delta(k^2) dk$$

by

$$\tilde{\dot{V}}(\omega, l) = -\omega a(\omega l).$$

¹In what follows the overdot will always denote the differentiation with respect to the real parameter s ; for a choice of $V(s, l)$ see below.

In physical contexts of interest the function $\dot{V}(s, l)$ is of class C^N , for some $N \in \{0, 1, \dots\} \cup \{\infty\}$, and for l scaled² to $l^0 = 1$ and some $\varepsilon > 0$, the following bounds hold

$$|L_{a_1 b_2} \dots L_{a_k b_k} \dot{V}(s, l)| \leq \frac{\text{const}}{(1 + |s|)^{1+\varepsilon}}, \quad k = 0, 1, \dots, N. \quad (5)$$

With this condition $A(x)$ has null asymptotes given by

$$\lim_{R \rightarrow \infty} RA(x \pm Rl) = \pm [V(x \cdot l, l) - V(\pm\infty, l)], \quad (6)$$

where $V(s, l)$ is any primitive function of $\dot{V}(s, l) = \partial V(s, l)/\partial s$, determined up to an additive s -independent term. Independent of this choice is the difference

$$\Delta V(l) = V(+\infty, l) - V(-\infty, l) = \int_{\mathbb{R}} \dot{V}(s, l) ds.$$

In what follows two spaces of functions $V(s, l)$ will appear naturally: one characterized by $V(+\infty, l) = 0$, and another by $V(-\infty, l) = 0$, respectively. In addition, we shall also find convenient to define another choice, denoted by

$$\bar{V}(s, l) = \frac{1}{2} \int_{\mathbb{R}} \text{sgn}(s - \tau) \dot{V}(\tau, l) d\tau, \quad (7)$$

for which

$$\bar{V}(\pm\infty, l) = \pm \frac{1}{2} \Delta V(l).$$

Beside the existence of null asymptotes, condition (5) entails (with the use of estimate (78)) the following decay of $A(x)$ in spacetime:

$$\begin{aligned} |A(x)| &\leq \text{const} \int \frac{d\Omega_t(l)}{(1 + |x \cdot l|)^{1+\varepsilon}} = \int_{-1}^1 \frac{\text{const} du}{(1 + ||x^0| + |\mathbf{x}|u|)^{1+\varepsilon}} \\ &\leq \frac{\text{const}}{1 + |x^0| + |\mathbf{x}|} \left\{ \theta(-x^2) + \frac{\theta(x^2)}{(1 + |x^0| - |\mathbf{x}|)^\varepsilon} \right\}. \end{aligned} \quad (8)$$

We end this section by mentioning that the standard Pauli-Jordan function, a particular solution of the wave equation, has the representation

$$D(x) = \frac{1}{2\pi} \text{sgn}(x^0) \delta(x^2) = -\frac{1}{8\pi^2} \int \delta'(x \cdot l) d^2l. \quad (9)$$

This may be shown by writing the rhs as

$$-\frac{1}{8\pi^2} \frac{\partial}{\partial x^0} \int \frac{\delta(x \cdot l)}{l^0} d^2l = -\frac{1}{4\pi} \frac{\partial}{\partial x^0} \int_{-1}^1 \delta(x^0 + |\mathbf{x}|u) du = -\frac{1}{4\pi} \frac{\partial}{\partial x^0} \frac{\theta(-x^2)}{|\mathbf{x}|},$$

which gives the standard form.

²This scaling is to be understood in all further estimates, to appear below, containing s or l .

2.4. Free electromagnetic field

A Lorenz potential A of a free electromagnetic field F has the representation (4), where $\dot{V}(s, l)$ is a vector function satisfying the condition

$$l \cdot \dot{V}(s, l) = 0. \quad (10)$$

We assume that the condition (5) with $N \geq 1$ is satisfied, so $\Delta V(l)$ exists, and satisfies $l \cdot \Delta V(l) = 0$. Moreover, we demand in addition that

$$L_{[ab}\Delta V_{c]}(l) = 0,$$

which under the orthogonality condition is equivalent to the existence of a homogeneous of degree 0 function $\Phi(l)$, such that

$$\Delta V_a(l) = -\partial_a \Phi(l), \quad (11)$$

equality in the sense described in Section 2.1. Then A is of the type of radiation potentials of fields created in scattering processes. The corresponding electromagnetic field is then given by

$$F_{ab}(x) = -\frac{1}{2\pi} \int [l_a \ddot{V}_b(x \cdot l, l) - l_b \ddot{V}_a(x \cdot l, l)] d^2 l. \quad (12)$$

The gauge freedom within the space of Lorenz gauges is represented in terms of $\dot{V}(s, l)$ by the transformations

$$\dot{V}(s, l) \mapsto \dot{V}(s, l) + l \dot{\alpha}(s, l), \quad (13)$$

with $\alpha(s, l)$ homogeneous of degree -2 . The spacelike asymptotic behaviour of A (and F) depends on $\Delta V(l)$ and is given by

$$\lim_{R \rightarrow \infty} R A(x + Ry) = -\frac{1}{2\pi} \int \Delta V(l) \delta(y \cdot l) d^2 l, \quad y^2 < 0.$$

Fields with $\Delta V(l) \neq 0$ are usually termed as infrared-singular; their spacelike tail falls off as that of the Coulomb field of electric charge.

For future use we note the following representation

$$x \cdot A(x) = \frac{1}{2\pi} \int (\partial \cdot V)(x \cdot l, l) d^2 l; \quad (14)$$

here and in what follows we denote

$$(\partial_a V)(x \cdot l, l) = \frac{\partial}{\partial l^a} V(s, l)|_{s=x \cdot l}$$

(and similarly for L_{ab} and other functions in place of V). Representation (14) follows immediately from the relation

$$\frac{\partial}{\partial l} \cdot V(x \cdot l, l) = x \cdot \dot{V}(x \cdot l, l) + (\partial \cdot V)(x \cdot l, l)$$

by the use of identity (1).

Finally, it is important for what comes to relate representation (4) to the construction of the radiation potential of a conserved vector current $J(x)$,

$$\partial \cdot J(x) = 0.$$

Let $J(x)$ be such current, for which $\dot{V}(s, l)$, with

$$V(s, l) = \int J(x) \delta(s - x \cdot l) dx, \quad (15)$$

satisfies conditions (5). Then the radiation potential of this current has two equivalent forms

$$A[J](x) = 4\pi \int_M D(x - y) J(y) dy = -\frac{1}{2\pi} \int \dot{V}(x \cdot l, l) d^2l, \quad (16)$$

which is shown with the use of representation (9). We note that orthogonality (10) is then automatic, but in general

$$l \cdot V(s, l) = Q, \quad (17)$$

Q being the charge of the current.

3. Free quantum field

Quantization of the local free electromagnetic field may be described in terms of the algebra of electromagnetic potential elements $A(J)$ satisfying the relations

$$[A(J_1), A(J_2)] = i4\pi \int J_1(x) D(x - y) J_2(x) dx dy = i\{J_1, J_2\}, \quad (18)$$

where J_i are smooth, compactly supported conserved test currents (as discussed in [13] and [14]). After exponentiation to the Weyl form, this defines a net of C^* -algebras satisfying Einstein causality in the form of commutation for spacelike separated supports of J_i .³ This formulation is close to the discussion given in [22]. In more standard formulation one treats the free electromagnetic field as an operator valued distribution on the space of antisymmetric, smooth, compactly supported functions $\varphi^{ab}(x)$. Then the elements

$$F(\varphi) = A(J), \quad \text{where } J^a = 2\partial_b \varphi^{ab}, \quad (19)$$

generate the standard net of electromagnetic field, where now $F(\varphi)$ is considered as an element of the local algebra of any open set containing $\text{supp } \varphi$. However, as each conserved, compactly supported current J^a has a representation (19) with compactly supported⁴ φ^{ab} , the nets have the same elements, with some differences in the assignment to spacetime sets. Moreover, two important properties bind the two nets:

³More recently, description in terms of ‘intrinsic potential’ $A(J)$ has been used for an axiomatic construction of a general local electromagnetic algebra in [6]. In that case, according to this analysis, the commutators for spacelike separated, but topologically nontrivial regions need not vanish, but are in the center of the algebra.

⁴This more specific Poincaré lemma for compact supports follows from its standard form and the properties of cohomology of spheres, see e.g. [18], Lemma 17.27. Also, the proof in that reference easily implies that for $\text{supp } J$ contained in an open, star-shaped set, there exists φ with support contained in the same set. See also [6].

- (i) the nets are local with respect to each other, i.e.

$$[F(\varphi), A(J)] = 0$$

for spacelike separated supports of φ and J ;

- (ii) for each open star-shaped set the two local algebras assigned to this set are equal.

The first property is obvious from (18) and (19), and for the second see footnote 4.

The symplectic form on the rhs of (18) may be expressed in terms of functions $V(s, l)$ defined by (15). Substituting in this form representation (9) one obtains⁵

$$\begin{aligned} \{V_1, V_2\} &= \frac{1}{4\pi} \int [\dot{V}_1(s, l) \cdot V_2(s, l) - V_1(s, l) \cdot \dot{V}_2(s, l)] ds d^2l \\ &= i \int \overline{\tilde{V}_1(\omega, l)} \cdot \tilde{V}_2(\omega, l) \frac{d\omega}{\omega} d^2l, \end{aligned} \quad (20)$$

where for the Fourier transformed version we used (3). If one writes symbolically $A(J_i) = \int_M A(x) J_i(x) dx$, with $A(x)$ represented by (4) with quantum function $V^q(s, l)$, then

$$A(J_i) = \{V_i, V^q\}. \quad (21)$$

The quantization condition (18) takes now the form

$$\{ \{V_1, V^q\}, \{V_2, V^q\} \} = i \{V_1, V_2\}. \quad (22)$$

This condition is now extended to the space of all smooth, compactly supported functions $V(s, l)$, satisfying orthogonality⁶

$$l \cdot V(s, l) = 0, \quad (23)$$

which for $V(s, l)$ obtained from J reflects the fact that the current is chargeless. The form (20) is nondegenerate on the space of equivalence classes of such functions with respect to gauge transformations (13). This may be also interpreted as follows: the symplectic structure is defined and nondegenerate on the space of infrared-regular classical fields (12). This leads to the unique Weyl algebra satisfying the corresponding exponentiated condition on this space (see [4], Section 5.2.2.2). Consider the scalar product on the space of equivalence classes

$$(\tilde{V}_1, \tilde{V}_2) = \int_{\mathbb{R}_+} \left[-\overline{\tilde{V}_1(\omega, l)} \cdot \tilde{V}_2(\omega, l) \right] \frac{d\omega}{\omega} d^2l, \quad (24)$$

and the Hilbert space \mathcal{H} obtained by completion. The standard construction leads to the representation of the algebra on the Fock space based on \mathcal{H} . When restricted to the exponentiated elements $F(\varphi)$, this is the standard irreducible vacuum representation of the local net.

In interacting theory one needs to form elements $A(K)$, where the vector function K need not be conserved, and secondly, need not be compactly

⁵By a slight abuse of notation we use here the same symbol as in (18).

⁶This formulation of the quantization is in the spirit of ‘asymptotic quantization’ discussed by Bramson [3] and later by Ashtekar [1]).

supported. The first problem (non-conservation of K) is dealt by choosing A in some special gauge, and in local formulation of QED this leads to the Gupta-Bleuler indefinite metric. The second problem (non-compactness) is avoided in usual formulations by the introduction of a switching function $g(x)$ and the subsequent performance of the ‘adiabatic limit’ $g \rightarrow 1$. This programme of the solution of infrared problems is far from complete.⁷ Here we want to propose a construction avoiding both problems, with the use of a natural extension of the above algebra to the free infrared-singular fields, put forward in [13]. We briefly sketch the construction of this extension.

The symplectic structure (20) has a natural extension to a wider space. It is easy to see that the form remains well defined (absolutely integrable) and gauge invariant for general functions V_i subject to conditions (5) with $N = \infty$, (11) and (23). However, for this structure to be again consistently formulated on the space of all classical fields (12) (now also those infrared-singular when $\Delta V_i(l) \neq 0$), we have to make the choice of $V(s, l)$ for given $\dot{V}(s, l)$ unique. We make this choice guided by the following physical motivation (as discussed in [14]). Free fields appear in realistic physical theories as incoming or outgoing fields in scattering setting. We want to admit the possibility of testing these fields with currents infinitely extended in timelike directions.⁸ Formula (15) tells us that for $V(+\infty, l)$ ($V(-\infty, l)$) to be different from zero, the test current J must have a sufficiently slow (in fact, of $|x|^{-3}$ type) decay in future (past), respectively. Therefore, it seems reasonable to adopt the rule that

$$\begin{aligned} V(+\infty, l) &= 0 \quad \text{for ‘in’ case,} \\ V(-\infty, l) &= 0 \quad \text{for ‘out’ case,} \end{aligned} \tag{25}$$

consistently with the spacetime regions where the fields are tested: ‘in’ in the past, and ‘out’ in the future; the same rule is adopted for the quantum variables $V^q(s, l)$. Let us mention that all such extended currents are charge-free (see (17)), and that the symplectic form $\{V_1, V_2\}$ for such currents may be represented in the form

$$\{J_1, J_2\} = \frac{1}{2} \int_M (A[J_1] \cdot J_2 - A[J_2] \cdot J_1)(x) dx, \tag{26}$$

where $A[J_i](x)$ are radiation fields (16). For compactly supported J_i this agrees with the definition in (18); however, the latter is not absolutely integrable in general.

Relations (22), considered on the space of functions $V(s, l)$ satisfying (5) with $N = \infty$, (11), (23) and (25), define (after exponentiation) a C^* -algebra, an extension of the standard local algebra. Algebras for ‘in’ and ‘out’ cases are canonically isomorphic. For J_i with supports spacelike separated the usual commutation rule is fulfilled. Also, the form (26) shows that the timelike commutativity characteristic for massless fields extends as well: if $\text{supp } J_1$ is timelike separated from $\text{supp } J_2$, then the two fields commute, irrespective of the size of supports.

⁷But see a recent contribution to this programme [8].

⁸Extensive discussion of such test currents may be found in [14]; see also what follows.

The Poincaré group transformations (z, Λ) are represented by automorphisms $\alpha_{z, \Lambda}$ acting on the algebra by

$$\alpha_{z, \Lambda}[A(J)] = A(T_{z, \Lambda}J), \quad (27)$$

where $T_{z, \Lambda}$ is the natural representation of the Poincaré group on tensor functions. Expressed in terms of notation (21) this reads

$$\alpha_{z, \Lambda}[\{V, V^q\}] = \{T_{z, \Lambda}V, V^q\}, \quad (28)$$

where

$$[T_{z, \Lambda}V]_a(s, l) = \Lambda_a{}^b V_b(s - z \cdot l, \Lambda^{-1}l).$$

Translationally covariant, positive energy, irreducible representations of the extended algebra have been constructed in [13] and [14]. Also, there exist among them those for which rotations are implemented, although not Lorentz boosts. In these representations there are vector states with arbitrarily small energy content, but no vacuum vector state exists.

The integral in the Fourier transformed version of the symplectic form (20) is now in the principal value sense. Note also that $\tilde{V}(0, l) = (2\pi)^{-1} \Delta V(l)$. Therefore, elements $\{V_1, V^q\}$ with $\Delta V_1(l) \neq 0$, describing infrared-singular fields, have no representation in the vacuum representation of the local algebra, which is based on the scalar product (24). We shall see in Section 5 below, that our larger arena allows the construction of elements $A^{\text{ar}}(K)$ for appropriately extended functions K and almost radial gauge A^{ar} to be defined there.

For later comparison we write the symbolic distributional formula equivalent to (22):

$$[\dot{V}_a^q(s, l), \dot{V}_b^q(s', l')] = -i g_{ab} \frac{2\pi}{l^0 l'^0} \delta(l, l') \delta' \left(\frac{s}{l^0} - \frac{s'}{l'^0} \right). \quad (29)$$

The rhs has the correct homogeneity and is independent of the choice of the vector t . The metric tensor on the rhs could suggest indefinite metric, but this is not the case: this relation is smeared with (equivalence classes with respect to (13) of) functions $V_i(s, l)$ orthogonal to l . In local formulation of interacting theory one needs to smear $A(x)$ with non-conserved test vector fields, and then the usual Gupta-Bleuler formulation is needed.

Finally, let us note for future use that if both $V_i(s, l)$ satisfy one and the same of the conditions (25), then

$$\{V_1, V_2\} = \{\bar{V}_1, \bar{V}_2\}, \quad (30)$$

where $\bar{V}_i(s, l)$ are formed by the convention (7).

4. Radial gauge

As announced in Introduction, we are now looking for a suitable quantum gauge, in which $x \cdot A(x)$ vanishes (sufficiently fast) in timelike directions. We start by testing the radial gauge for that purpose, for classical field first.

If $F_{ab}(x) = \partial_a A_b(x) - \partial_b A_a(x)$, with A a C^1 -function, then without any further conditions the following identity holds

$$x^a F_{ab}(x) = (x \cdot \partial + 1)A_b(x) - \partial_b(x \cdot A(x)),$$

so

$$\lambda x^a F_{ab}(\lambda x) = \frac{\partial}{\partial \lambda} [\lambda A_b(\lambda x)] - \partial_b(x \cdot A(\lambda x)). \quad (31)$$

Integrating with respect to λ on $[0, 1]$ one finds

$$\int_0^1 \lambda x^a F_{ab}(\lambda x) d\lambda = A_b(x) - \partial_b S^{\text{rg}}(x) \equiv A_b^{\text{rg}}(x), \quad (32)$$

$$S^{\text{rg}}(x) = \int_0^1 x \cdot A(\lambda x) d\lambda,$$

which for any electromagnetic field F_{ab} defines the radial gauge with the property

$$x \cdot A^{\text{rg}}(x) = 0. \quad (33)$$

Potential (32) is the unique, up to the addition of a constant, regular solution of the condition (33). Indeed, any addition to $A_a^{\text{rg}}(x)$ of a gauge term $\partial_a G(x)$ respecting (33) must result from a homogeneous function $G(x)$, which has to be singular (not continuous) at the origin, if it is not constant.

The lhs of (32) shows immediately that for F —a free Maxwell field, the radial gauge is a Lorenz gauge. Let $A(x)$ be a Lorenz gauge given by (4) with (10). Then one finds

$$A^{\text{rg}}(x) = -\frac{1}{2\pi} \int \dot{V}^{\text{rg}}(x \cdot l, l) d^2 l, \quad (34)$$

with

$$\dot{V}_b^{\text{rg}}(s, l) = \dot{V}_b(s, l) + l_b \frac{\partial \cdot V(s, l) - \partial \cdot V(0, l)}{s}. \quad (35)$$

This is shown by noting that by (14) we have

$$x \cdot A(\lambda x) = \frac{1}{2\pi \lambda} \int (\partial \cdot V)(\lambda x \cdot l, l) d^2 l,$$

so

$$S^{\text{rg}}(x) = \frac{1}{2\pi} \int_0^1 \left(\int (\partial \cdot V)(\lambda x \cdot l, l) d^2 l \right) \frac{d\lambda}{\lambda}$$

and

$$\begin{aligned} \partial_b S^{\text{rg}}(x) &= \frac{1}{2\pi} \int_0^1 \int l_b (\partial \cdot \dot{V})(\lambda x \cdot l, l) d^2 l d\lambda \\ &= \frac{1}{2\pi} \int l_b \frac{(\partial \cdot V)(x \cdot l, l) - (\partial \cdot V)(0, l)}{x \cdot l} d^2 l \end{aligned}$$

(note that $S^{\text{rg}}(x)$ is well defined, since $\int (\partial \cdot V)(0, l) d^2 l = 0$).

For a classical field, with sufficiently regular $V(s, l)$, the radial gauge function (35) is regular and the representation (34) is well defined (e.g., for a smooth $V(s, l)$ both $V^{\text{rg}}(s, l)$ and $A^{\text{rg}}(x)$ are smooth), although with

marked differences as compared to A . In contrast to $\dot{V}(s, l)$, which satisfies (5), $\dot{V}^{\text{rg}}(s, l)$ decays only as s^{-1} for large $|s|$. This has two consequences: first, there is no null asymptotics of the form (6) for $A^{\text{rg}}(x)$; and second, the estimate (8) is replaced by a slower decay

$$|A^{\text{rg}}(x)| \leq \text{const} \int \frac{d\Omega(l)}{1 + |x \cdot l|} \leq \frac{\text{const}}{1 + |x^0| + |\mathbf{x}|} \log(2 + |\mathbf{x}|),$$

which is easily obtained with the use of estimate (78).

However, for the quantum version it follows immediately from the juxtaposition of formulas (35) and (29) that $V^{\text{rg}}(s, l)$, and consequently also $A^{\text{rg}}(x)$, are not properly defined, due to the appearance of a distributional quantity at a point: $V(0, l)$ (the commutator of $V(0, l)$ and $V(0, l')$ would yield $\delta(0)$). This, in our view, is the fundamental reason for the reported difficulties and a need to *a posteriori* regularization in attempts to establish Feynman rules in radial gauge (see [19] and references therein).

5. Almost radial gauge

The source of trouble in the radial gauge is the term $\partial \cdot V(0, l)$ in the function $\dot{V}^{\text{rg}}(s, l)$. Our plan is to modify the definition (32) so as to eliminate this term. This will only be possible with some smearing of the central, reference point. The result of this programme is the construction of an *almost radial gauge* potential $A_b^{\text{ar}}(x)$ in formulas (48), (49), (50) and (71) below.

The starting point for the construction is the following variant of the identity (31)

$$(\xi + 1)\overline{a\hat{x}}^a F_{ab}(x + \xi\overline{a\hat{x}}) = \frac{\partial}{\partial \xi} [(\xi + 1)A_b(x + \xi\overline{a\hat{x}})] - \partial_b^x (\overline{a\hat{x}} \cdot A(x + \xi\overline{a\hat{x}})), \quad (36)$$

where ξ is a real parameter and to abbreviate notation we introduced a usual affine space symbol

$$\overline{a\hat{x}} = x - a.$$

To obtain this identity we shift the argument of A and F in (31) by vector a , and write it with x replaced by y :

$$\lambda y^a F_{ab}(a + \lambda y) = \frac{\partial}{\partial \lambda} [\lambda A_b(a + \lambda y)] - \frac{\partial}{\partial y^b} (y \cdot A(a + \lambda y)).$$

Now the substitutions $y = \overline{a\hat{x}}$ and $\lambda = \xi + 1$ give the result.

5.1. Potential of fast decay

We start discussion with the case of $A_b(x)$ a C^1 -function of fast decay, together with its derivative. Multiplying (36) by $(-\frac{1}{2})\text{sgn}(\xi)$ and integrating with respect to ξ on \mathbb{R} we find

$$-\frac{1}{2} \int_{\mathbb{R}} \text{sgn}(\xi) (\xi + 1) \overline{a\hat{x}}^a F_{ab}(x + \xi\overline{a\hat{x}}) d\xi = A_b(x) - \partial_b^x S_k(a, x), \quad (37)$$

where

$$S(a, x) = -\frac{1}{2} \int_{\mathbb{R}} \vec{a}\vec{x} \cdot A(x + \xi\vec{a}\vec{x}) \operatorname{sgn}(\xi) d\xi. \quad (38)$$

It is easy to see that $S(a, x)$ is regular outside $x = a$, bounded by a constant, while $|\partial^x S(a, x)| \leq \operatorname{const}|x - a|^{-1}$. In order to remove the singularity we smear (37) with a real Schwartz function $\rho(a)$ on Minkowski space, such that

$$\int_M \rho(a) da = 1, \quad \int_M \rho(a) a^\alpha da = 0 \quad \text{for } 1 \leq |\alpha| \leq n, \quad (39)$$

for some arbitrarily chosen (large) $n \in \mathbb{N}$. The smearing of $\partial^x S(a, x)$ may be pulled under the differential sign and we obtain

$$\begin{aligned} \int_M S(a, x) \rho(a) da &= \frac{1}{2} \int_M A(y) \cdot \vec{y}\vec{x} \int_{\mathbb{R}} \rho(x + \xi^{-1}\vec{y}\vec{x}) |\xi|^{-5} d\xi dy \\ &= \frac{1}{2} \int_M A(y) \cdot \vec{y}\vec{x} \int_{\mathbb{R}} \rho(x + u\vec{y}\vec{x}) |u|^3 du dy. \end{aligned}$$

In the first step above we substituted $S(a, x)$ (38) and changed integration variable a to $y = x + \xi(x - a)$. In the second step we substituted $\xi = u^{-1}$.

We now introduce the vector function

$$r^a(x, z) = \frac{1}{2} z^a \int_{\mathbb{R}} \rho(x + uz) |u|^3 du, \quad (40)$$

which together with its derivatives with respect to x is estimated as in (79), and satisfies the distributional equation (see Appendix B)

$$\partial^z \cdot r(x, z) = \delta(z), \quad (41)$$

and define the **almost radial gauge** by

$$A_b^{\text{ar}}(x) = -\frac{1}{2} \int_M \rho(a) \int_{\mathbb{R}} \operatorname{sgn}(\xi) (\xi + 1) \vec{a}\vec{x}^a F_{ab}(x + \xi\vec{a}\vec{x}) d\xi da \quad (42)$$

$$= A_b(x) - \partial_b S^{\text{ar}}(x), \quad (43)$$

$$S^{\text{ar}}(x) = \int_M r(x, x - y) \cdot A(y) dy = \int_M r(x, z) \cdot A(x - z) dz. \quad (44)$$

Formulas (41), (43) and (44) together show that our almost radial gauge is in the class of generalized ‘gauge-invariant gauges’ first postulated by Dirac [7] in an attempt to obtain ‘physical’, out of local potentials (see also [24]). The generalization consists in nontrivial dependence of $r^a(x, x - y)$ on the first argument. Potential $A^{\text{ar}}(x)$ is a continuous function, decaying as $|x|^{-1}$ for large $|x|$, thus

$$|A^{\text{ar}}(x)| \leq \frac{\operatorname{const}}{1 + |x|}.$$

To estimate the radial component of A^{ar} assume for simplicity that the support of $\rho(a)$ is contained in a small ball $|a| \leq \delta$, and note that $\vec{a}\vec{x}^a x^b F_{ab} = \vec{a}\vec{x}^a a^b F_{ab}$. Then the contraction of (42) with x gives

$$|x \cdot A^{\text{ar}}(x)| \leq \delta \frac{\operatorname{const}}{1 + |x|}.$$

In the present case of a fast decaying function A the ‘almost radial’ property has only a global aspect, appearance of δ on the rhs. It is the case of free field discussed below, where this term has a fuller justification.

We note that a strict Dirac gauge with $r(0, x - y)$ replacing $r(x, x - y)$ in (44) is also possible, but it does not have the almost radial property.

5.2. Free classical field

We now assume that $F_{ab}(x)$ is a free classical field obtained from the potential (4), where $\dot{V}(s, l)$ satisfies all restrictions formulated there. The definition (44) may not be directly applied in this case, but we shall find that an extension of this definition may be obtained as a limit of a regularized expression. We admit infrared-singular fields, thus $\Delta V(l)$ may be different from zero. For the sake of the construction of almost radial gauge at the classical level, it will prove convenient to use the convention $\bar{V}(s, l)$ (7). The choice (25) will be restored at the stage of quantization.

The following auxiliary functions will be needed

$$\begin{aligned}\sigma(s, l) &= \frac{1}{2} \int_M \operatorname{sgn}(s - a \cdot l) \rho(a) da, \\ \zeta(s, l) &= \int_M \frac{\rho(a)}{s - a \cdot l} da = \int_{\mathbb{R}} \frac{\dot{\sigma}(\tau, l)}{s - \tau} d\tau, \\ \eta(s, l) &= \int_M \frac{\rho(a)a}{s - a \cdot l} da = - \int_{\mathbb{R}} \frac{\partial \sigma(\tau, l)}{s - \tau} d\tau\end{aligned}\quad (45)$$

(we assume $\sigma(s, l)$ is defined by the above formula also in a neighborhood of the cone, so $\partial \sigma(s, l)$ is also unique in this case). Properties of these functions are discussed in Appendix C. Let $\chi(x)$ be a Schwartz function, such that

$$\chi(-x) = \chi(x), \quad \chi(0) = 1,$$

and for $\delta > 0$ denote (cf. (44))

$$S_\delta^{\text{ar}}(x) = \int_M \chi(\delta z) r(x, z) \cdot A(x - z) dz, \quad (46)$$

with $A(x)$ as described above. Then the following limit exists

$$S^{\text{ar}}(x) = \lim_{\delta \searrow 0} S_\delta^{\text{ar}}(x) = \frac{1}{2\pi} \int [\eta(x \cdot l, l) - \zeta(x \cdot l, l)x] \cdot \bar{V}(x \cdot l, l) d^2l. \quad (47)$$

We give the proof of this fact in Appendix D. Gradient of this function is easily obtained

$$\partial_b S^{\text{ar}}(x) = \frac{1}{2\pi} \int \left\{ l_b \partial_s [(\eta(s, l) - \zeta(s, l)x) \cdot \bar{V}(s, l)] - \zeta(s, l) \bar{V}_b(s, l) \right\} \Big|_{s=x \cdot l} d^2l.$$

We use identity of the form (14) to transform

$$-\frac{1}{2\pi} x^a \int l_b \partial_s [\bar{V}_a(s, l) \zeta(s, l)] \Big|_{s=x \cdot l} d^2l = \frac{1}{2\pi} \int \partial_a [l_b \bar{V}^a(s, l) \zeta(s, l)] \Big|_{s=x \cdot l} d^2l.$$

Taking into account that $\partial_a[l_b\bar{V}^a\zeta] = \bar{V}_b\zeta + l_b\partial \cdot [\bar{V}\zeta]$, we find

$$\partial_b S^{\text{ar}}(x) = \frac{1}{2\pi} \int l_b \left\{ \partial_s [\eta(s, l) \cdot \bar{V}(s, l)] + \partial \cdot [\zeta(s, l) \bar{V}(s, l)] \right\} \Big|_{s=x \cdot l} d^2 l.$$

We can now extend the definition of the **almost radial gauge** to the free field:

$$A_b^{\text{ar}}(x) = A_b(x) - \partial_b S^{\text{ar}}(x) = -\frac{1}{2\pi} \int \dot{V}^{\text{ar}}(x \cdot l, l) d^2 l, \quad (48)$$

where

$$\dot{V}^{\text{ar}}(s, l) = \dot{\bar{V}}(s, l) + l \left\{ \partial_s [\eta(s, l) \cdot \bar{V}(s, l)] + \partial \cdot [\zeta(s, l) \bar{V}(s, l)] \right\} \quad (49)$$

$$= \dot{\bar{V}}(s, l) + l \left\{ \eta(s, l) \cdot \dot{\bar{V}}(s, l) + \zeta(s, l) \partial \cdot \bar{V}(s, l) \right\}, \quad (50)$$

the second form (50) obtained with the use of identity (80). Our almost radial gauge is a functional of the smearing function ρ (39), which serves to define functions ζ and η (45). Note that putting formally $\rho(a) = \delta(a)$ (which, precisely, is not allowed), one has $\zeta(s, l) = s^{-1}$, $\eta(s, l) = 0$ and (50) reproduces formula (35) with the term $(\partial \cdot V)(0, l)$ omitted (which, again, would result in singularity).

The almost radial gauge is again a Lorenz gauge satisfying the wave equation. It is also gauge invariant with respect to gauge transformations

$$A_b(x) \rightarrow A_b(x) + \partial_b \Lambda(x), \quad \Lambda(x) = -\frac{1}{2\pi} \int \bar{\alpha}(x \cdot l, l) d^2 l,$$

which correspond to the replacement $\bar{V}_b(s, l) \rightarrow \bar{V}_b(s, l) + l_b \bar{\alpha}(s, l)$. This replacement leaves $\dot{V}_b^{\text{ar}}(s, l)$ unchanged. To show this we put $l\bar{\alpha}(s, l)$ in place of $\bar{V}(s, l)$ in (50). Recall that to calculate $\partial \cdot [l\bar{\alpha}(s, l)]$ one has to extend $l\bar{\alpha}(s, l)$ to the neighborhood of the cone respecting orthogonality to l , e.g. by

$$l_a \bar{\alpha}(s, l) \rightarrow [l_a \bar{\alpha}(s, l)]_{\text{ext}} = [l_a - (l^0)^{-1} l^2 \delta_a^0] \bar{\alpha}(s, l),$$

and then on the cone $\partial \cdot [l\bar{\alpha}(s, l)]_{\text{ext}} = (l \cdot \partial + 2) \bar{\alpha}(s, l)$. In this way the change in (50) becomes

$$l\zeta(s, l)(s\partial_s + l \cdot \partial + 2) \bar{\alpha}(s, l) = 0,$$

where we used identity (81) and the homogeneity of degree -2 of $\bar{\alpha}(s, l)$.

Similarly as in the radial gauge case, $\dot{V}^{\text{ar}}(s, l)$ decays only as $1/s$ for $|s| \rightarrow \infty$, so $A^{\text{ar}}(x)$ has no null asymptotes and decays as

$$|A^{\text{ar}}(x)| \leq \frac{\text{const}}{1 + |x^0| + |\mathbf{x}|} \log(2 + |\mathbf{x}|). \quad (51)$$

On the other hand, using identity (81) we find

$$\begin{aligned} & x \cdot \dot{V}^{\text{ar}}(x \cdot l, l) \\ &= x \cdot \dot{\bar{V}}(x \cdot l, l) + (\partial \cdot \bar{V})(x \cdot l, l) + \eta(x \cdot l, l) \cdot [l \partial \cdot \bar{V}(x \cdot l, l) + x \cdot l \dot{\bar{V}}(x \cdot l, l)], \end{aligned}$$

and by identity (14) we obtain

$$x \cdot A^{\text{ar}}(x) = -\frac{1}{2\pi} \int \eta(s, l) \cdot [l \partial \cdot \bar{V}(s, l) + s \dot{\bar{V}}(s, l)]|_{s=x \cdot l} d^2 l. \quad (52)$$

The expression in square brackets in (52) is bounded by a constant, while $\eta(s, l)$ satisfies (82), so

$$\begin{aligned} |x \cdot A^{\text{ar}}(x)| &\leq \text{const} \int \frac{d\Omega(l)}{(1 + |x \cdot l|)^n} \\ &\leq \frac{\text{const}}{1 + |x^0| + |\mathbf{x}|} \left\{ \theta(-x^2) + \frac{\theta(x^2)}{(1 + |x^0| - |\mathbf{x}|)^{n-1}} \right\}. \end{aligned} \quad (53)$$

In timelike directions the radial product $x \cdot A^{\text{ar}}(x)$ vanishes arbitrarily fast (depending on the choice of the function $\rho(x)$, see (39)). Moreover, let us introduce the scaling $\rho_\delta(a) = \delta^{-4} \rho(a/\delta)$, which for $\delta \searrow 0$ contracts the function to the central point, and denote by A_δ^{ar} the corresponding potential. Then $\eta_\delta(s, l) = \eta(s/\delta, l)$ and for $x \cdot A_\delta^{\text{ar}}(x)$ the bound (53) holds with the replacement $x \rightarrow x/\delta$. Thus outside $x = 0$ this quantity vanishes with δ as

$$|x \cdot A_\delta^{\text{ar}}(x)| \leq \frac{\text{const} \delta}{\delta + |x^0| + |\mathbf{x}|} \left\{ \theta(-x^2) + \frac{\theta(x^2) \delta^{n-1}}{(\delta + |x^0| - |\mathbf{x}|)^{n-1}} \right\}.$$

5.3. Smearing with a test function

Let $K^b(x)$ be a vector Schwartz test function (no conservation condition) and denote

$$W(s, l) = \int_M K(x) \delta(s - x \cdot l) dx, \quad (54)$$

which is a C^∞ -function, fast decaying in s , together with all its derivatives. Then

$$A^{\text{ar}}(K) = \int_M A^{\text{ar}}(x) \cdot K(x) dx = -\frac{1}{2\pi} \int \dot{V}^{\text{ar}}(s, l) \cdot W(s, l) ds d^2 l.$$

Setting here (49) and integrating by parts (use (2) to transfer derivative ∂) one finds

$$\begin{aligned} A^{\text{ar}}(K) &= \frac{1}{2\pi} \int \bar{V}(s, l) \cdot \dot{V}_K(s, l) ds d^2 l \\ &= -\frac{1}{2\pi} \int \dot{V}(s, l) \cdot \bar{V}_K(s, l) ds d^2 l = \{\bar{V}_K, \bar{V}\}, \end{aligned} \quad (55)$$

where

$$\begin{aligned} \dot{V}_K(s, l) &= \dot{W}(s, l) + \eta(s, l) l \cdot \dot{W}(s, l) + \zeta(s, l) \partial(l \cdot W(s, l)), \\ &= \dot{W}(s, l) + \partial_s [\eta(s, l) l \cdot W(s, l)] + \partial[\zeta(s, l) l \cdot W(s, l)], \\ \bar{V}_K(s, l) &= \frac{1}{2} \int_{\mathbb{R}} \text{sgn}(s - \tau) \dot{V}_K(\tau, l) d\tau. \end{aligned} \quad (56)$$

It follows that

$$l \cdot \bar{V}_K(s, l) = 0, \quad (57)$$

$$\Delta V_K(l) = \partial \int_{\mathbb{R}} \zeta(s, l) l \cdot W(s, l) ds. \quad (58)$$

Orthogonality (57) is showed with the use of identity (81) as follows:

$$\begin{aligned} l \cdot \dot{V}_K(s, l) &= l \cdot \dot{W}(s, l) + (s\zeta(s, l) - 1)l \cdot \dot{W}(s, l) + \zeta(s, l)l \cdot \partial(l \cdot W(s, l)) \\ &= \zeta(s, l)(s\partial_s + l \cdot \partial)(l \cdot W(s, l)) = 0, \end{aligned}$$

the last equality by the homogeneity of $l \cdot W(s, l)$. Equality (58) is immediate from the second form of \dot{V}_K in (56).

We note that the function $\bar{V}_K(s, l)$ satisfies all demands (5) with $N = \infty$, (11) and (23) on a test function in the extended symplectic form (20), but in general—for non-conserved K —is outside the original test functions space.

5.4. More general smearing

Now we want to extend the smearing $A^{\text{ar}}(K)$ to C^1 -vector functions $K(x)$, with the spacetime asymptotic behavior characteristic for currents of scattered free particles or fields. We shall show that with the conditions on K made appropriately specific the function $W(s, l)$ (54) is of class C^1 and satisfies the bounds

$$|\dot{W}(s, l)| \leq \frac{\text{const}}{(1 + |s|)^{1+\varepsilon}}, \quad |\partial[l \cdot W(s, l)]| \leq \frac{\text{const}}{(1 + |s|)^\varepsilon}, \quad (59)$$

so the integrals in (55), with the definitions (56), are absolutely convergent for this extension. Moreover, $\bar{V}_K(s, l)$ thus formed satisfies conditions (5) with $N = 0$, (11) and (23). In addition, let $g(x)$ be a Schwartz function, with $g(0) = 1$ and denote $K_\delta(x) = g(\delta x)K(x)$. Then we shall find that irrespective of the shape of g the following extension of the identity (55) holds true

$$\begin{aligned} A^{\text{ar}}(K) &\equiv \lim_{\delta \searrow 0} A^{\text{ar}}(K_\delta) = \frac{1}{2\pi} \int \bar{V}(s, l) \cdot \dot{V}_K(s, l) ds d^2l \\ &= -\frac{1}{2\pi} \int \dot{V}(s, l) \cdot \bar{V}_K(s, l) ds d^2l = \{\bar{V}_K, \bar{V}\}. \end{aligned} \quad (60)$$

We first note that once the existence of $W(s, l)$ with the properties (59) is established, the existence of $\bar{V}_K(s, l)$ with the property (5) with $N = 0$, and orthogonality (57) are obviously satisfied.

To specify the class for which the above results hold it is convenient to characterize such class by splitting the function K into three C^1 -contributions: $K = K_1 + K_2 + K_3$, with the following properties:

$$\begin{aligned} K_1(x) &= x \kappa(x), \quad \text{supp } \kappa \subseteq \{x^2 \geq 0, |x^0| \geq \frac{1}{2}\}, \\ \kappa(\lambda x) &= \lambda^{-4} \kappa(x) \quad \text{for } \lambda \geq 1, |x^0| \geq 1, \end{aligned} \quad (61)$$

$$|\kappa(x)| \leq \frac{\text{const}}{(1 + |x|)^4}, \quad |\partial \kappa(x)| \leq \frac{\text{const}}{(1 + |x|)^5},$$

$$\text{supp } K_2 \subseteq \{x^2 \geq 0\}, \quad |K_2(x)| \leq \frac{\text{const}}{(1 + |x|)^3}, \quad \text{oscillatory part}, \quad (62)$$

$$|K_3(x)| \leq \frac{\text{const}}{(1 + |x|)^{3+\varepsilon}}, \quad |\partial K_3(x)| \leq \frac{\text{const}}{(1 + |x|)^{4+\varepsilon}}. \quad (63)$$

Condition (61) characterizes the dominating asymptotic behavior of particles, and the non-oscillatory part of asymptotic behavior of fields. The oscillatory contribution in case of fields is represented by K_2 , bounded by (62), and we shall characterize it more precisely below. The rest is represented by K_3 , conditions (63) characterizing the next to leading behaviour inside the cone, and decay of K outside. We prove our claims separately for each contribution K_i , except for property (11), which will be discussed at the end.

We first note that all $W_i(s, l)$, defined by (54) with K replaced by K_i , are absolutely convergent and bounded. Moreover, in case $i = 3$ one has

$$|W_3(s, l)|, |L_{ab}W_3(s, l)| \leq \frac{\text{const}}{(1 + |s|)^\varepsilon}, \quad |\dot{W}_3(s, l)| \leq \frac{\text{const}}{(1 + |s|)^{1+\varepsilon}};$$

all these statements are proved as in Lemma 20 and Theorem 21 in ref. [16]. Therefore, bounds (59) are satisfied for W_3 . Next, we observe that both

$$A^{\text{ar}}(x) \cdot K_3(x) \quad \text{and} \quad A^{\text{ar}}(x) \cdot K_1(x) = x \cdot A^{\text{ar}}(x) \kappa(x)$$

are absolutely integrable (see (51) and (53)), so the δ -regularization in (60) is not needed for these contributions. Thus in case $i = 3$ we obtain

$$\begin{aligned} A(K_3) &= -\frac{1}{2\pi} \int_M \int \dot{V}^{\text{ar}}(x \cdot l, l) d^2l K_3(x) dx \\ &= -\frac{1}{2\pi} \int \dot{V}^{\text{ar}}(s, l) \cdot W_3(s, l) ds d^2l, \end{aligned}$$

so the thesis for this contribution follows.

In case $i = 1$ we denote

$$U(s, l) = \int_M \kappa(x) \delta(s - x \cdot l) dx, \quad |U(s, l)| \leq \frac{\text{const}}{1 + |s|}$$

(the bound again by Lemma 20 in [16]), and observe that

$$\partial U(s, l) = -\dot{W}_1(s, l), \quad sU(s, l) = l \cdot W_1(s, l). \quad (64)$$

Moreover, we note that the inside of the lightcone (past and future) may be parametrized by $x = \lambda v$, $\lambda \in \mathbb{R}$ and v on the future hyperboloid $v^2 = 1$,

$v^0 \geq 0$, and then $dx = |\lambda|^3 d\lambda d\mu(v)$, where $d\mu(v) = d^3v/v^0$. It follows then from assumptions (61) that for $|s| \geq l^0$ one has

$$\begin{aligned} W_1(s, l) &= \int \operatorname{sgn}(\lambda) v \kappa(\operatorname{sgn}(\lambda)v) \delta(s - \lambda v \cdot l) d\lambda d\mu(v) \\ &= \operatorname{sgn}(s) \int \kappa(\operatorname{sgn}(s)v) \frac{v}{v \cdot l} d\mu(v), \\ l \cdot W_1(s, l) &= \operatorname{sgn}(s) \int \kappa(\operatorname{sgn}(s)v) d\mu(v). \end{aligned} \quad (65)$$

Therefore,

$$\dot{W}_1(s, l) = 0, \quad \partial[l \cdot W_1(s, l)] = 0 \quad \text{for } |s| \geq l^0.$$

We can now use (52) to obtain (we omit the arguments (s, l) for the sake of clarity)

$$\begin{aligned} A(K_1) &= -\frac{1}{2\pi} \int \eta \cdot [l \partial \cdot \bar{V} + s \dot{\bar{V}}] U ds d^2l \\ &= -\frac{1}{2\pi} \int [sU \partial_s (\bar{V} \cdot \eta) + (1 - s\zeta) \bar{V} \cdot \partial U] ds d^2l, \end{aligned}$$

where for the second equality we integrated ∂ by parts and used the equality $\partial(l \cdot \eta) = -s\dot{\zeta}$, which follows from identities (80) and (81). Performing now the substitutions (64) (for sU using the second of these relations in both places where it appears) and integrating ∂_s by parts, we obtain

$$A(K_1) = \frac{1}{2\pi} \int \bar{V} \cdot [\dot{W}_1 + \eta l \cdot \dot{W}_1 + \zeta \partial(l \cdot W_1)] ds d^2l,$$

which is the required result.

We turn to the case $i = 2$. The decay of K_2 is not sufficient to apply the proof presented in case $i = 3$. However, the oscillatory behavior of K_2 damps the integrals, and we now add further assumption that W_2 satisfies bounds as in case W_3 :

$$|W_2(s, l)|, |L_{ab}W_2(s, l)| \leq \frac{\text{const}}{(1 + |s|)^\varepsilon}, \quad |\dot{W}_2(s, l)| \leq \frac{\text{const}}{(1 + |s|)^{1+\varepsilon}}. \quad (66)$$

We denote

$$W_{2\delta}(s, l) = \int_M g(\delta x) K_2(x) \delta(s - x \cdot l) dx, \quad (67)$$

and assume further that

$$|W_{2\delta}(s, l)| \leq \frac{\text{const}}{(1 + |s|)^\varepsilon}. \quad (68)$$

With these assumptions we have

$$\begin{aligned} A(K_{2\delta}) &= \int \dot{V}^{\text{ar}}(s, l) \cdot W_{2\delta}(s, l) ds d^2l \\ &\rightarrow \int \dot{V}^{\text{ar}}(s, l) \cdot W_2(s, l) ds d^2l \quad \text{for } \delta \searrow 0, \end{aligned}$$

the limit by the dominated convergence theorem. This, together with the estimates (66), leads to the thesis. We show in Appendix E that our assumptions on W_2 are satisfied for a term of the type characteristic for the Dirac field.

Finally, we shall close the proof of our claims by showing that

$$\Delta V_K(l) = -\partial\Phi_K(l), \quad (69)$$

with

$$\begin{aligned} \Phi_K(l) &= \int [\kappa(v) + \kappa(-v)]\rho(a) \log \frac{|a \cdot l|}{v \cdot l} d\mu(v) da \\ &\quad - \int \zeta(s, l) l \cdot [W(s, l) - W(\operatorname{sgn}(s)\infty, l)] ds. \end{aligned} \quad (70)$$

First we observe that it is only W_1 that contributes to $W(\pm\infty, l)$, so by (65) we have

$$W(\pm\infty, l) = \pm \int \kappa(\pm v) \frac{v}{v \cdot l} d\mu(v), \quad l \cdot W(\pm\infty, l) = \pm \int \kappa(\pm v) d\mu(v) \equiv Q_{\pm}.$$

Differentiating the first line in (70) we obtain

$$\begin{aligned} \int [\kappa(v) + \kappa(-v)]\rho(a) \left[\frac{a}{a \cdot l} - \frac{v}{v \cdot l} \right] d\mu(v) da \\ = \eta(0, l)(Q_- - Q_+) - W(+\infty, l) + W(-\infty, l). \end{aligned}$$

Differentiation of the second line in (70) gives

$$- \int \partial[\zeta(s, l) l \cdot W(s, l)] ds + \int \partial\zeta(s, l) Q_{\operatorname{sgn}(s)} ds.$$

Using (80) in the second integral and summing the contributions we obtain

$$\partial\Phi_K(l) = -W(+\infty, l) + W(-\infty, l) - \int \partial[\zeta(s, l) l \cdot W(s, l)] ds,$$

which substituted into (69) gives the correct value. Note that the differential ∂ cannot be extracted outside the integral in the last formula, as without it the integrand is not integrable.

5.5. Quantum field

We have shown in classical case that $A^{\operatorname{ar}}(K) = \{\bar{V}_K, \bar{V}\}$. However, recall the identity (30), which holds in each of the cases (25), that is both for ‘in’ and ‘out’ algebra. Thus, on the algebraic level, in each of these two cases we propose to *define* the **quantum almost radial gauge** of the potential by

$$A^{\operatorname{ar}}(K) = \{V_K, V^{\operatorname{q}}\}, \quad (71)$$

whenever $V_K(s, l)$ defined by (56) and (25) falls in our extended symplectic space defined after (26). This condition demands stronger regularity properties of K than those assumed in Section 5.4, leading to classical version of (71). One can show that a class of such functions may be obtained by a modification of the currents J considered in [14], where the modification consists in demanding the continuity equation to be satisfied only asymptotically in

time. On the other hand, in concrete representations the scope of admitted test functions is wider, and the definition of the almost radial gauge may be also appropriately extended at the level of representation. We do not go into details here and leave the question to be decided in applications.

The almost radial gauge is a Lorenz gauge. For $K(x) = \partial F(x)$ one finds that $V_K(s, l) = W(s, l)\varpropto l$, which is in the zero equivalence class. Thus

$$A^{\text{ar}}(\partial F) = 0.$$

Moreover, for $K(x) = \square G(x)$ we have $W(s, l) = 0$, so

$$A^{\text{ar}}(\square G) = 0.$$

For test functions $K(x)$ in the class of conserved currents admitted in $A(K)$ in one of the cases ‘in’ or ‘out’ one has $l \cdot W(s, l) = 0$, so that $\dot{V}_K(s, l) = \dot{W}(s, l)$ and $V_K(s, l) = W(s, l)$. Therefore, in this case

$$A^{\text{ar}}(K) = A(K),$$

which reproduces the extended free algebra.

Finally, let us note the following important corollary. If K is a conserved, non-radiating current, such as the current of a free, charged classical particle, or of the free (classical or quantum) Dirac field, then $W(s, l) = W(l)$ is constant in s and $l \cdot W(l) = Q$, the charge of the current (see, e.g. ref. [12], discussion starting with Eq. (2.70)). Therefore, for such currents $V_K = 0$ and

$$A^{\text{ar}}(K) = 0.$$

5.6. Transformation and commutation properties

We recall that almost radial gauge is a functional of the function ρ serving to smear the point a from which the radial line to x is drawn. Let us make this dependence explicit by writing $A^{\text{ar}}(K; \rho)$. With Poincaré transformations of the extended algebra defined in (27), these elements have the covariance property given by

$$\alpha_{z, \Lambda} [A^{\text{ar}}(K; \rho)] = A^{\text{ar}}(T_{z, \Lambda} K; T_{z, \Lambda} \rho). \quad (72)$$

In this sense $A(K; \rho)$ may be regarded algebraically covariant, with unitarily implementable translations. For the proof of the covariance relation one notes that

$$\begin{aligned} W[T_{z, \Lambda} K](s, l) &= \Lambda W[K](s - z \cdot l, \Lambda^{-1} l), \\ \zeta[T_{z, \Lambda} \rho](s, l) &= \zeta[\rho](s - z \cdot l, \Lambda^{-1} l), \\ \eta[T_{z, \Lambda} \rho](s, l) &= \Lambda \eta[\rho](s - z \cdot l, \Lambda^{-1} l) + z \zeta[\rho](s - z \cdot l, \Lambda^{-1} l). \end{aligned}$$

Setting these relations into the formula (56) for V_K , and then using the result in (28) one arrives at the thesis.

Elements $A^{\text{ar}}(K; \rho)$ do not have compact localization even if the support of K is compact. However, the following remnant timelike locality with respect to A holds. Let the support of an admissible test vector function K and the support of a conserved test current J be timelike separated, and let

$V(s, l)$ for current J (15) have $\Delta V(l) = 0$ (in particular, this is always true for compactly supported J). Then

$$[A^{\text{ar}}(K; \rho), A(J)] = 0, \quad (73)$$

irrespective of the choice of ρ . To show this we first note that it is sufficient to show this when the two supports are placed inside the future and the past parts of the same lightcone, with any vertex point b . Next, we apply to the commutator translation automorphism $\alpha_{-b, \mathbb{1}}$. The commutator, without changing its value (being proportional to identity), takes now the form $[A^{\text{ar}}(K'; \rho'), A(J')]$, where K' and J' are similarly separated as K and J , but with the vertex of the cone in the origin. In consequence, the functions $W'(s, l)$ for K' (54), and $V'(s, l)$ for J' (15), are supported in $s \in (0, +\infty)$ or $s \in (-\infty, 0)$, each in a different of the two sets. Now, it follows from (56) that the support of $\dot{V}_{K'}(s, l)$ is not larger than the support of $W'(s, l)$, so the same conclusion is valid for the pair $\dot{V}_{K'}(s, l)$ and $V'(s, l)$. But from $\Delta V'(l) = 0$, together with (25), it follows that

$$\{V_{K'}, V'\} = \frac{1}{2\pi} \int_{\mathbb{R}} \dot{V}_{K'}(s, l) \cdot V'(s, l) \, ds \, d^2l,$$

so the thesis follows.

6. Almost radial gauge and string-localized fields: comparative remarks

Definition of the almost radial gauge was motivated by integration, on the classical level, of the electromagnetic potential along straight axes (42). Another construction based on integration along infinitely extended straight lines is the formalism of string-localized fields (potentials) developed by Mund, Schroer and Yngvason [21] (based on earlier works, as explained in this reference) and worked out later in many articles.⁹ The two constructions result from quite different motivations and aims; nevertheless, it seems interesting to make some comparison. For that purpose we briefly recall and formulate in our language the latter in case of electrodynamics.¹⁰ All electromagnetic fields are assumed to satisfy the homogeneous Maxwell equations.

The string-localized potentials in electrodynamics are distributions in position vectors x and spacelike unit vectors e defined by¹¹

$$A_b^{\text{sl}}(x, e) = \int_{\mathbb{R}_+} F_{ba}(x + \lambda e) e^\alpha \, d\lambda.$$

⁹See e.g. a recent discussion of an application to QED problems [20].

¹⁰Our perspective is different from the discussion of string-localized fields in [21], which starts from the unitary representations of the Poincaré group and engages the notion of modular localization [2]. Here we want to avoid the *a priori* assumption on the action of a representation of the Lorentz group.

¹¹There exists a version of this construction with lightlike e , see [10].

To obtain a vector potential as an x -distribution one needs a scalar test function $h(e)$ on the hyperboloid $e^2 = -1$, such that

$$\int h(e) d\nu(e) = 1, \quad (74)$$

where the integration is over the hyperboloid with the Lorentz invariant measure $d\nu(e)$. Then

$$A_b^{\text{sl}}(x; h) = \int A_b(x, e) h(e) d\nu(e)$$

is an h -dependent potential, commuting with the field $F_{ab}(y)$ at points y in spacelike position to the set of points $x + \lambda e$, $\lambda \in \mathbb{R}_+$, $e \in \text{supp } h$. In particular, if $\text{supp } h$ is contained in some double-cone spacelike separated from 0, then $A^{\text{sl}}(x; h)$ is localized in a spacelike cone.

This potential is in the Lorenz class, and using the representation (12) one can show that on classical level

$$A_b^{\text{sl}}(x; h) = -\frac{1}{2\pi} \int \dot{V}_b^{\text{sl}}(x \cdot l, l) d^2l,$$

where

$$\dot{V}^{\text{sl}}(s, l) = \dot{V}(s, l) - l n(l) \cdot \dot{V}(s, l) - lz(l) \cdot \int \frac{\dot{V}(\tau, l)}{s - \tau} d\tau,$$

$$n(l) = \int \frac{eh(e)}{e \cdot l} d\nu(e), \quad z(l) = \int eh(e) \delta(e \cdot l) d\nu(e),$$

and not to burden notation we do not indicate the dependence on h explicitly. For a vector Schwartz test function $K(x)$ we then find in analogy to (55)

$$A^{\text{sl}}(K; h) = \{V_K^{\text{sl}}, V\}, \quad (75)$$

where in place of (56), and keeping notation (54), we have

$$V_K^{\text{sl}}(s, l) = W(s, l) - n(l)l \cdot W(s, l) + z(l) \int \frac{l \cdot W(\tau, l)}{s - \tau} d\tau. \quad (76)$$

In contrast to $W(s, l)$, which is of fast decrease, $V_K^{\text{sl}}(s, l)$ vanishes in general only as s^{-1} , see the last term in (76). Therefore, the (exponentiated) formula (75) cannot give, upon quantization, an element in the local algebra, even if K is local. However, we have

$$\tilde{V}_K^{\text{sl}}(\omega, l) = \tilde{W}(\omega, l) - n(l)l \cdot \tilde{W}(\omega, l) + i\pi z(l) \text{sgn}(\omega l) \cdot \tilde{W}(\omega, l),$$

which is sufficient for elements (75) to be represented in the standard vacuum representation of the field F , in contrast to the almost radial gauge, which needs the extended algebra including infrared singular fields.

The formulation of a gauge theory in terms of string-localized potentials puts stress on their Poincaré covariance. In vacuum representation the action of the Poincaré automorphisms $\alpha_{z, \Lambda}$ of the observable algebra given by (27) is implemented by unitary representation $U(z, \Lambda)$. Then, in this representation, one also has

$$\alpha_{z, \Lambda}[A^{\text{sl}}(K; h)] = A^{\text{sl}}(T_{z, \Lambda}K; T_{0, \Lambda}h) = U(z, \Lambda)A^{\text{sl}}(K; h)U(z, \Lambda)^*. \quad (77)$$

At the algebraic level our transformation law (72) is analogous to (77). However, as mentioned before, in our case not all Lorentz transformations are implemented.

Let us now return again to the question of localization and commutation. String-localized fields located on spacelike separated strings commute (which remains true for slightly smeared strings); this is an important property in the formalism of these fields. Whether almost radial gauge may be localized in spacelike extended regions in some special cases and special points x is a subtle question, which we do not want to consider here; anyway, this cannot be a generic property of this gauge. On the other hand, it has a timelike commutation property (73). We have seen that this property depended on the fact that the support of $\dot{V}_K(s, l)$ was not greater than that of $W(s, l)$. The latter property does not hold for $V_K^{\text{sl}}(s, l)$ because of the last term in (76). Therefore, for string-localized fields timelike commutativity does not hold, in general.

It is interesting to note that the function $z(l)$ vanishes for even functions h , $h(-e) = h(e)$, which are admitted by the condition (74). For such functions the last term in (76) is absent and the elements $A^{\text{sl}}(K; h)$ acquire the timelike locality as formulated above for $A^{\text{ar}}(K; \rho)$. As here the algebra is the standard local electromagnetic algebra of the free field, this leads in fact to the following implication in the vacuum representation. If the support of the smearing function K is contained in an open double-cone with vertices x_1 and x_2 , $x_2^0 > x_1^0$, then $A^{\text{sl}}(K)$ belongs to the intersection of the von Neuman algebras connected with the following two regions: the inside of the future lightcone with the vertex in x_1 , and the inside of the past lightcone with the vertex in x_2 . This follows from the timelike duality for vacuum representation of massless fields, see [5]. Condition of evenness of h may be equivalently formulated as string integration along whole real axes, instead of semi-axes; the spacelike localization and commutation properties are then retained (with obvious modifications).¹²

One could also observe that the last term in (76) seems to worsen the infrared/adiabatic limit behavior: for K becoming infinitely extended so that $W(s, l)$ does not vanish in infinity, this term becomes meaningless. On the other hand, if the term is absent, then the algebra of elements (75) may be also built in the extended algebra of electromagnetic field described earlier.

Acknowledgement

I am grateful to Paweł Duch, Wojciech Dybalski and José M. Gracia-Bondía for reading the manuscript and for interesting correspondence.

¹²The relation with the notion of modular localization, as used in the string-localized formalism of [20], is then lost, or at least becomes indirect.

Appendix

A. Estimates

Here we note the following estimates. For $a, b, c, \alpha > 0$, one has

$$\int_0^c \frac{du}{(a+bu)^\alpha} \leq \begin{cases} \frac{\alpha}{\alpha-1} \frac{c}{a^{\alpha-1}(a+bc)}, & \alpha > 1, \\ \frac{1}{1-\alpha} \frac{c}{(a+bc)^\alpha}, & \alpha < 1, \\ \frac{2c}{a+bc} \log\left(e + \frac{bc}{a}\right), & \alpha = 1, \end{cases} \quad (78)$$

with Euler's number e . Cases $\alpha \leq 1$ may be found in [12], Appendix B, and are reproduced here for convenience of the reader. In case $\alpha = 1$ the result of integration is $\frac{1}{b} \log\left(1 + \frac{bc}{a}\right)$. For $bc \leq a$ this is bounded by $\frac{c}{a} \leq \frac{2c}{a+bc}$, while for $bc \geq a$ by $\frac{2c}{a+bc} \log\left(1 + \frac{bc}{a}\right)$.

B. Function $r^a(x, z)$

Here we discuss properties of function $r^a(x, z)$ defined in (40), and we start with its estimate. We denote by x_z and x_\perp the parallel and the perpendicular (in Euclidean metric) to z components of x , respectively. Then for each $k \in \mathbb{N}$ we have

$$|r^a(x, z)| \leq \text{const}(k) \frac{(1 + |x_z|)^3}{(1 + |x_\perp|)^k} \frac{1}{|z|^3}. \quad (79)$$

To show this we use $|\rho(x)| \leq \text{const}(1 + |x|)^{-k-4}$ to estimate the rhs of (40) by

$$\text{const} \int_{\mathbb{R}} \frac{|z| |u|^3 du}{(1 + |x_\perp| + |u|z + x_z)^{k+4}} \leq \frac{\text{const}}{|z|^3} \int_0^\infty \frac{(r + |x_z|)^3 dr}{(1 + |x_\perp| + r)^{k+4}},$$

which leads easily to (79).

Next, we prove identity (41). We observe that outside $z = 0$ the function $\int \rho(x + uz) |u|^3 du$ is smooth and homogeneous in z of degree -4 , so the distribution $\partial^z \cdot r(x, z)$ is concentrated in $z = 0$. We choose a C^1 -function $f(z)$ which together with its derivative decays at least as $|z|^{-2}$, denote $\hat{z} = z/|z|$ and write $dS(\hat{z})$ for the integration element on the (Euclidean) unit sphere. Then

$$\begin{aligned} - \int_M r(x, z) \cdot \partial f(z) dz &= - \lim_{\epsilon \searrow 0} \int_M r(x, z) \cdot \partial f(z) \theta(|z| - \epsilon) dz \\ &= \frac{1}{2} \lim_{\epsilon \searrow 0} \int_M \int_{\mathbb{R}} \rho(x + uz) |u|^3 du f(z) \delta(|z| - \epsilon) |z| dz \\ &= \frac{1}{2} \lim_{\epsilon \searrow 0} \int_{S^3} \int_{\mathbb{R}} \rho(x + r\hat{z}) |r|^3 dr f(\epsilon\hat{z}) dS(\hat{z}) = f(0) \int_M \rho(x + z) dz = f(0), \end{aligned}$$

where the third equality is obtained by the change of variables $r = \epsilon u$. This ends the proof.

C. Functions ζ and η

The functions ζ and η defined by (45) are smooth outside $l = 0$ and homogeneous of degree -1 with respect to their arguments, and satisfy the following identities

$$\partial\zeta(s, l) + \dot{\eta}(s, l) = 0, \quad (80)$$

$$s\zeta(s, l) = 1 + l \cdot \eta(s, l). \quad (81)$$

The Fourier transforms are

$$\tilde{\zeta}(\omega, l) = i\pi \operatorname{sgn}(\omega)\hat{\rho}(\omega l), \quad \tilde{\eta}(\omega, l) = \pi \operatorname{sgn}(\omega)(\partial\hat{\rho})(\omega l).$$

Conditions (39) are equivalent to

$$\hat{\rho}(0) = \frac{1}{2\pi}, \quad \partial^\alpha \hat{\rho}(0) = 0 \quad \text{for } 1 \leq |\alpha| \leq n,$$

which implies that $\tilde{\eta}(\omega, l)$ is $n - 1$ times continuously differentiable and

$$\partial_\omega^k \tilde{\eta}(\omega, l)|_{\omega=0} = 0, \quad k = 0, \dots, n - 1,$$

while $\partial_\omega^n \tilde{\eta}(\omega, l)$ is bounded and integrable. It follows that

$$|\eta(s, l)| \leq \frac{\operatorname{const}}{(1 + |s|)^n}. \quad (82)$$

D. Derivation of $S^{\text{ar}}(x)$ for free field

Here we prove formula (47). Formula (46) for $S_\delta^{\text{ar}}(x)$ may be interpreted as the definition (44) with $\chi(\delta z)A(x - z)$ substituted for $A(x - z)$. Equivalently, this is expression (38) with $\chi(-\delta\xi\vec{a}\vec{x})A(x + \xi\vec{a}\vec{x})$ substituted for $A(x + \xi\vec{a}\vec{x})$, and smeared with $\rho(a)$. We change integration variables from a to $z = a - x$, multiply the expression for convenience by 2π , and obtain

$$2\pi S_\delta^{\text{ar}}(x) = -\frac{1}{2} \int_M \int_{\mathbb{R}} \chi(\delta\xi z) \rho(x + z) \int z \cdot \dot{V}(x \cdot l - \xi z \cdot l, l) d^2l \operatorname{sgn}(\xi) d\xi dz.$$

We note that the integrand is absolutely integrable, so we can freely change the order of integration and leave the d^2l integral to be performed at the last stage. In the remaining integrals we change ξ to $\lambda = \delta\xi$ and reduce integration with respect to λ to positive semi-axis by reflecting $\lambda \rightarrow -\lambda$ from the negative semi-axis. The result may be written as

$$-\int_M \int_{\mathbb{R}_+} \chi(\lambda z) \rho(x + z) \frac{\partial}{\partial \lambda} \frac{z \cdot Z(x \cdot l, \delta^{-1}\lambda z \cdot l, l)}{z \cdot l} d\lambda dz, \quad (83)$$

where

$$Z(\tau, u, l) = \bar{V}(\tau, l) - \frac{1}{2} [\bar{V}(\tau + u, l) + \bar{V}(\tau - u, l)].$$

We note that $Z(\tau, u, l)$ is bounded, $Z(\tau, 0, l) = 0$, and

$$\lim_{\delta \rightarrow 0} Z(x \cdot l, \delta^{-1}\lambda z \cdot l, l) = \bar{V}(x \cdot l, l) \quad \text{almost everywhere,}$$

with the choice of $\bar{V}(s, l)$ according to the formula (7).

We now submit (83) to the following transformations:

- integration with respect to λ by parts;

- replacement of $\rho(x+z)$ by $\nu(x, z) = \frac{1}{2}[\rho(x+z) + \rho(x-z)]$, possible due to the evenness in z of the combined remaining factors;
- choice of variables w^a for vector z , such that $w^0 = z \cdot l$, $dz = dw^0 d^3 w_\perp$, with $w_\perp = (w^1, w^2, w^3)$.

In this way (83) takes the form

$$\int_{\mathbb{R}} \int_{\mathbb{R}_+} (w^0)^{-1} f(x, \lambda, w^0) \cdot Z(x \cdot l, \delta^{-1} \lambda w^0, l) d\lambda dw^0, \quad (84)$$

where

$$f(x, \lambda, w^0) = \int_{\mathbb{R}^3} \frac{\partial}{\partial \lambda} \chi(\lambda w) \nu(x, w) w dw_\perp.$$

It is easy to see that $f(x, \lambda, w^0)$ is a C^∞ -function, and $f(x, \lambda, 0) = 0$ due to the oddness of the integrand in w , so $(w^0)^{-1} f(x, \lambda, w^0)$ is also C^∞ . An easy calculation also shows that the latter function is absolutely integrable with respect to $d\lambda dw^0$. Therefore, we can take the $\delta \rightarrow 0$ limit under all integrations and the limit of (84) becomes

$$\bar{V}(x \cdot l, l) \cdot \int_{\mathbb{R}} \int_{\mathbb{R}_+} f(x, \lambda, w^0) (w^0)^{-1} d\lambda dw^0 = -\bar{V}(x \cdot l, l) \cdot \int_{\mathbb{R}^4} \frac{w \nu(x, w)}{w^0} dw,$$

in the latter form integral in the principal value sense. We can now restore back the abstract notation of vector z , and the function $\rho(x+z)$ in place of $\nu(x, z)$ (due to the evenness of $(w^0)^{-1} w$). We finally restore the integration variable $a = x+z$ and then

$$- \int_{\mathbb{R}^4} \frac{\nu(x, w) w}{w^0} dw = \int_M \frac{\rho(a)(a-x)}{x \cdot l - a \cdot l} da = \eta(x \cdot l, l) - x \zeta(x \cdot l, l).$$

E. Oscillatory asymptotic term

Here we make more specific assumptions on oscillating smearing function K_2 introduced in Section 5.4. Function K_2 specified below obviously satisfies (62), and we shall show that assumptions (66) and (68) are satisfied with $\varepsilon = 1$.

Let $K_2(x)$ be of the form (cf. [12])

$$K_2(x) = e^{i\mu\sqrt{x^2}} k(x), \quad \mu > 0,$$

where $k(x)$ is a C^2 -vector function, supported inside the lightcone and vanishing for $|x^0| \leq 1$. Moreover, we assume that in the parametrization $x = \lambda v$ introduced after (64) the following bounds are satisfied

$$|(\partial^\alpha k)(\lambda v)| \leq \frac{\text{const}}{\lambda^{3+|\alpha|} (v^0)^n}, \quad |\alpha| \leq 2. \quad (85)$$

where for simplicity we assume that n may be arbitrarily large; by a rather tedious control of the decay in v this could be substantially weakened, but we do not need this for our illustrative purposes.

For $W_{2\delta}$ formed as in (67) we have

$$W_{2\delta}(s, l) = \int e^{i\mu\lambda} k(\lambda v) g(\delta\lambda v) \delta(s - \lambda v \cdot l) |\lambda|^3 d\lambda d\mu(v).$$

We integrate with respect to λ , which gives

$$W_{2\delta}(s, l) = \int e^{i\mu\lambda} k(\lambda v) g(\delta\lambda v) |\lambda|^3 \frac{d\mu(v)}{v \cdot l}, \quad (86)$$

where from now on we put $\lambda = \lambda(s, v, l) = s/v \cdot l$. It is clear that $|W_{2\delta}(s, l)|$ is bounded by a constant independent of δ , and

$$\lim_{\delta \rightarrow 0} W_{2\delta}(s, l) = W_2(s, l) = \int e^{i\mu\lambda} k(\lambda v) |\lambda|^3 \frac{d\mu(v)}{v \cdot l}. \quad (87)$$

To proceed, we need an integral identity on the hyperboloid $v^2 = 1$, $v^0 > 0$. Consider operators

$$M_{ab} = v_a \frac{\partial}{\partial v^b} - v_b \frac{\partial}{\partial v^a}, \quad \delta_b = v^a M_{ab} = \frac{\partial}{\partial v^b} - v_b v \cdot \frac{\partial}{\partial v},$$

intrinsic differentiation operators in the hyperboloid. The measure $d\mu(v)$ is Lorentz invariant, and operators M_{ab} generate Lorentz transformations in the hyperboloid, so if $\chi(v)$ and $M_{ab}\chi(v)$ are absolutely integrable, then

$$\int M_{ab}\chi(v) d\mu(v) = 0.$$

Moreover, we have $M_{ab}v^a\chi(v) = (\delta_b - 3v_b)\chi(v)$. Therefore, if

$$|\chi(v)| \leq \frac{\text{const}}{(v^0)^{3+\varepsilon}}, \quad |\delta\chi(v)| \leq \frac{\text{const}}{(v^0)^{2+\varepsilon}},$$

then

$$\int (\delta - 3v)\chi(v) d\mu(v) = 0. \quad (88)$$

We note the identities and a bound

$$l \cdot \delta \frac{1}{v \cdot l} = 1, \quad e^{i\mu\lambda} = -\frac{i}{s\mu} l \cdot \delta e^{i\mu\lambda}, \quad l \cdot \delta \frac{v}{v \cdot l} = \frac{l}{v \cdot l}, \quad \frac{1}{v \cdot l} \leq \frac{2v^0}{l^0}.$$

Substituting the second of these relations in (86) and integrating by parts by (88) (our assumptions on decay of k and ∂k make it possible), we find

$$\begin{aligned} W_{2\delta}(s, l) &= \frac{i}{\mu s} \int e^{i\mu\lambda} g(\delta\lambda v) |\lambda|^3 \left(k(\lambda v) + \frac{\lambda}{v \cdot l} l \cdot \partial k(\lambda v) \right) d\mu(v), \\ &+ \frac{i\delta}{\mu s} \int e^{i\mu\lambda} |\lambda|^3 k(\lambda v) \frac{\lambda}{v \cdot l} l \cdot \partial g(\delta\lambda v) d\mu(v). \end{aligned}$$

We note that $|g(\delta\lambda v)|$ is bounded by a constant and tends to 1 for $\delta \rightarrow 0$, while $\delta|\lambda||v^0||\partial g(\delta\lambda v)|$ is also bounded by a constant and tends to zero in that limit. Thus using (85) we find that $|W_{2\delta}(s, l)| \leq \text{const}|s|^{-1}$, so the bound (68) and the first of the bounds (66) are satisfied with $\varepsilon = 1$. Moreover, taking the limit we find

$$W_2(s, l) = \frac{i}{\mu s} \int e^{i\mu\lambda} |\lambda|^3 \left(k(\lambda v) + \frac{|\lambda|}{v \cdot l} l \cdot \partial k(\lambda v) \right) d\mu(v).$$

Differentiating this formula with respect to l^a and s and using similar technique as above one proves the remaining bounds in (66) for $|s| \geq 1$. For $|s| \leq 2$ formula (87) should be used instead. We omit the details.

References

- [1] Ashtekar, A.: *Asymptotic Quantization*, Bibliopolis, Napoli, 1986
- [2] Brunetti, R., Guido, D. and Longo, R.: Modular localization and Wigner particles, *Rev. Math. Phys.* **14**, 759-786 (2002)
- [3] Bramson, B.D.: Physics in cone space, in *Asymptotic Structure of Space-Time*, ed. Esposito, F.P. and Witten, L., Plenum, New York, 1977, pp. 273-359
- [4] Bratteli, O. and Robinson, D.W.: *Operator Algebras and Quantum Statistical Mechanics 2*, 2nd. Ed., Springer, New York, 1996
- [5] Buchholz, D.: On the structure of local quantum fields with non-trivial interaction, in: *Proceedings of the international conference on operator algebras, ideals and their applications in theoretical physics, Leipzig 1977*, Teubner, Leipzig, 1978
- [6] Buchholz, D., Ciolli, F., Ruzzi, G. and Vasselli, E.: The universal C^* -algebra of the electromagnetic field, *Lett Math Phys* **106**, 269-285 (2016). <https://doi.org/10.1007/s11005-015-0801-y>
- [7] Dirac, P.A.M.: Gauge-invariant formulation of quantum electrodynamics, *Can. J. Phys.* **33**, 650-660 (1955)
- [8] Duch, P.: Infrared problem in perturbative quantum field theory, arxiv: 1906.00940 [math-ph]
- [9] Fock, V.A.: Die Eigenzeit in der klassischen und in der Quantenmechanik, *Phys. Zeit. Sowjetunion* **12** 404-425 (1937)
- [10] Gracia-Bondía, J.M., Mund, J. and Várilly, J.C.: The Chirality Theorem, *Ann. Henri Poincaré* **19** 843-874 (2018)
- [11] Jackson, J.D.: From Lorenz to Coulomb and other explicit gauge transformations, *Am. J. Phys.* **70**, 917-928 (2002)
- [12] Herdegen, A.: Long-range effects in asymptotic fields and angular momentum of classical field electrodynamics, *J. Math. Phys.* **36**, 4044-4086 (1995)
- [13] Herdegen, A.: Semidirect product of CCR and CAR algebras and asymptotic states in quantum electrodynamics, *J. Math. Phys.* **39**, 1788-1817 (1998)
- [14] Herdegen, A.: Infrared problem and spatially local observables in electrodynamics, *Ann. Henri Poincaré* **9**, 373-401 (2008). <https://doi.org/10.1007/s00023-008-0359-7>
- [15] Herdegen, A.: Asymptotic structure of electrodynamics revisited, *Lett. Math. Phys.* **107**, 1439-1470 (2017). <https://doi.org/10.1007/s11005-017-0948-9>
- [16] Herdegen, A.: Infrared problem vs gauge choice: scattering of classical Dirac field, *Ann. Henri Poincaré* (2021). <https://doi.org/10.1007/s00023-020-01015-y>
- [17] Jauch, J.M., Rohrlich, F.: *The Theory of Photons and Electrons*, 2nd Ed., Springer, Berlin, 1976
- [18] Lee, J.M.: *Introduction to Smooth Manifolds*, 2nd Ed., Springer, New York, 2013
- [19] Leupold, S.: Feynman rules in radial gauge, arXiv:hep-th/9609222 (1996)
- [20] Mund, J., Rehren, K.-H. and Schroer, B.: Gauss' Law and String-Localized Quantum Field Theory, *JHEP* **01** 001 (2020)
- [21] Mund, J., Schroer, B. and Yngvason, J.: String-Localized Quantum Fields and Modular Localization, *Commun. Math. Phys.* **268** 621-672 (2006)

- [22] Roepstorff, G.: Coherent Photon States and Spectral Condition, *Comm. Math. Phys.* **19** 301-314 (1970)
- [23] Schwinger, J.: On gauge invariance and vacuum polarization, *Phys. Rev.* **82**, 664-679 (1951)
- [24] Steinmann, O.: *Perturbative Quantum Electrodynamics and Axiomatic Field Theory*, Springer, Berlin, 2000

Andrzej Herdegen
Institute of Theoretical Physics
Jagiellonian University
ul. S. Łojasiewicza 11
30-348 Kraków
Poland
e-mail: herdegen@th.if.uj.edu.pl